



Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2017 March ; 26(3): 413–419. doi:10.1158/1055-9965.EPI-16-0630.

URINARY MELATONIN IN RELATION TO POSTMENOPAUSAL BREAST CANCER RISK ACCORDING TO MELATONIN 1 RECEPTOR STATUS

Elizabeth E. Devore¹, Erica T. Warner², A. Heather Eliassen^{1,3}, Susan B. Brown⁴, Andrew H. Beck⁵, Susan E. Hankinson^{1,3,4}, and Eva S. Schernhammer^{1,6}

¹Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts

²Clinical Translational Epidemiology Unit, Massachusetts General Hospital, Boston, Massachusetts

³Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts

⁴Division of Biostatistics and Epidemiology, University of Massachusetts Amherst, Amherst, Massachusetts

⁵Department of Pathology, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, Massachusetts

⁶Department of Epidemiology, Center for Public Health, Medical University of Vienna, Vienna, Austria

Abstract

Background—Urinary melatonin levels have been associated with a reduced risk of breast cancer in postmenopausal women, but this association might vary according to tumor melatonin 1 receptor (MT1R) expression.

Methods—We conducted a nested case-control study among 1,354 postmenopausal women in the Nurses' Health Study, who were cancer free when they provided first-morning spot urine samples in 2000–2002; urine samples were assayed for 6-sulfatoxymelatonin (aMT6s, a major metabolite of melatonin). Five-hundred fifty-five of these women developed breast cancer before May 31, 2012, and were matched to 799 control subjects. In a subset of cases, immunohistochemistry was used to determine MT1R status of tumor tissue. We used multivariable-adjusted conditional logistic regression to estimate the relative risk (RR) of breast cancer (with 95% confidence intervals [CI]) across quartiles of creatinine-standardized urinary aMT6s level, including by MT1R subtype.

Corresponding author: Elizabeth E. Devore, 181 Longwood Avenue, Room 448, Boston, Massachusetts 02115; phone: 617-525-2042; fax: 617-525-2008; nheed@channing.harvard.edu.

Conflict of interest statement: The authors declare no potential conflicts of interest.

Results—Higher urinary melatonin levels were suggestively associated with a lower overall risk of breast cancer (multivariable-adjusted RR=0.78, 95% CI=0.61–0.99, comparing quartile 4 vs. quartile 1; p-trend=0.08); this association was similar for invasive vs. in situ tumors (p-heterogeneity=0.12). There was no evidence that associations differed according to MT1R status of the tumor (e.g., p-heterogeneity for overall breast cancer=0.88).

Conclusion—Higher urinary melatonin levels were associated with reduced breast cancer risk in this cohort of postmenopausal women, and the association was not modified by MT1R subtype.

Impact—Urinary melatonin levels appear to predict the risk of breast cancer in postmenopausal women. However, future research should evaluate these associations with longer-term follow-up and among premenopausal women.

Keywords

Melatonin; breast cancer; melatonin 1 receptor; postmenopausal; epidemiology

INTRODUCTION

Light exposure during the biologic night has been hypothesized to increase risk of breast cancer(1). Melatonin, an indolamine hormone, is a molecular marker of the circadian system; it is entrained to the 24-hour environmental light-dark cycle, released by the pineal gland, and suppressed by light(2). In addition, melatonin has been shown to mediate numerous cell-signaling pathways involved in breast cancer, including estrogen-dependent pathways(3, 4). Some epidemiologic studies(5–7), although not all(8–11), have reported that lower levels of urinary melatonin are associated with greater risk of breast cancer—in line with the notion that light exposure at night, which suppresses melatonin secretion, promotes carcinogenesis.

Moreover, two melatonin receptors have been identified, melatonin 1 receptor (MT1R) and melatonin 2 receptor (MT2R), and MT1R has been found on the surface of breast tumors in cell culture(12–14). In rodents, MT1R overexpression has been linked to reduced breast tumor incidence(15), whereas age-related decline in MT1R expression has been shown to reduce the sensitivity of melatonin for these receptors, leading to enhanced tumor growth(16, 17). Indeed, experimental work indicates that melatonin’s anti-carcinogenic effects may be largely mediated by MT1R(18), but this potential mechanism has been understudied in human populations. In the Nurses’ Health Study, previous research found that higher urinary melatonin levels were strongly associated with lower risk of breast cancer based on 357 cases that developed over six years of follow up(7); however, the combined role of endogenous melatonin levels and MT1R in breast cancer has not been explored.

To extend our previous study, we evaluated the association of urinary melatonin and breast cancer risk in the Nurses’ Health Study, utilizing ~200 additional breast cancer cases with twice the length of follow-up. Furthermore, we examined whether this association differed according to MT1R status of the breast tumor.

MATERIALS AND METHODS

Study population

We conducted a nested case-control study within the Nurses' Health Study (NHS) cohort. The NHS cohort began in 1976, when 121,701 female nurses, who were 30–55 years old and living in the United States, returned an initial mailed questionnaire with information on breast cancer risk factors and major medical diagnoses. This information has been updated using similar mailed questionnaires every two years, and participation has exceeded 90% at every questionnaire cycle. Between 2000 and 2002, 18,643 participants provided spot urine samples and completed a short supplemental questionnaire, including information on date and time of urine collection, confirmation of first-morning urine, current weight, and recent postmenopausal hormone (PMH) use; 93% of urine samples were first-morning samples. These samples were returned on ice, by overnight mail, after which they were aliquoted and stored in nitrogen freezers (at –130 degrees Celsius) until they were assayed for melatonin.

Women were eligible for the present analysis if they provided a urine sample and had no history of cancer (except non-melanoma skin cancer) prior to urine collection; cases were women who developed breast cancer after urine collection and before May 31, 2012. These cases were matched either 1:1 (for postmenopausal cases with recent PMH use) or 1:2 (all other cases) to controls who did not develop breast cancer during the follow-up period. Matching factors included: birth year, menopausal status, recent PMH use (defined as use 3 months from the time of urine collection), timing of urine collection (month and time of day), and fasting status at urine collection. Women were considered postmenopausal if they reported natural menopause or bilateral oophorectomy, or if they reported hysterectomy without bilateral oophorectomy and were aged 56 years and a non-smoker, or aged 54 years and a current smoker (90% of women in the cohort had reached natural menopause by these ages). We focused on postmenopausal women in this study. The Institutional Review Board of Brigham and Women's Hospital approved this study, and informed consent was implied by voluntary return of mailed questionnaires and biological specimens.

Breast cancer case ascertainment

Participants self reported a breast cancer diagnosis on one of the biennial cohort questionnaires. All self-reported cases were asked for permission to review their medical records, which provided confirmation of their diagnosis. When medical records were unavailable, breast cancer cases were defined as probable and included in the analysis if corroborated by phone interview or written confirmation from the participant. In some cases, breast cancer was reported during death follow-up, when family members or the postal service reported a participant's death to study personnel. The National Death Index was reviewed after each questionnaire cycle to determine the status of women who were unresponsive to cohort questionnaires. In this cohort, self-reported breast cancer was >98% accurate compared to pathology reports(19).

Urinary melatonin assessment

Urine samples for breast cancer cases and controls were removed from the freezer simultaneously, handled identically, and each batch was shipped in one parcel. Three batches

were sent for laboratory analysis based on the biennial questionnaire cycle in which women reported incident breast cancer. The first batch (i.e., incident cases reported on the 2002 and 2004 biennial questionnaires and their matched controls) and second batch (i.e., incident cases reported on the 2006 biennial questionnaire and their matched controls) were sent to the Endocrine Core Laboratory of M. Wilson (Yerkes National Primate Center at Emory University, Atlanta, GA). The third batch (i.e., incident cases reported on the 2008 and 2010 biennial questionnaires and their matched controls) were sent to the Specialty Assay Research Core Laboratory of R. Carroll (Brigham and Women's Hospital, Boston, MA). 6-sulfatoxymelatonin (aMT6s) was assayed using the Buhlmann enzyme-linked immunosorbent assay (ELISA), with a lower limit of detection of 0.8 ng/mL; creatinine levels were also assayed, which enabled creatinine standardization to account for differences in urine concentration. Laboratory personnel were blinded to case-control status, and case-control samples were assayed together on the same day and in the same run; quality control samples were included in each batch. In the present analyses, within-batch coefficients of variation were: 9.5%, 10.3%, and 6.2% for aMT6s and 8.5%, 3.1%, and 1.1% for creatinine across the three batches, respectively.

The reproducibility of creatinine-adjusted aMT6s levels over three years has been previously established, using the collection and assessment method described for this cohort (intraclass correlation=0.72)(20).

MT1 receptor assessment

A detailed description of the breast tumor tissue block collection, microarrays (TMAs) construction, and immunohistochemical analyses performed can be found elsewhere(21–23). Briefly, we collected archived formalin-fixed, paraffin-embedded breast cancer blocks from incident breast cancer cases through 2006. Hematoxylin and eosin sections from those cases were reviewed to confirm the diagnosis, classify the cancer according to histological type and grade, and identify the area from which the cores for the TMAs would be taken. TMAs were constructed in the Dana-Farber Harvard Cancer Center Tissue Microarray Core Facility (Boston, Massachusetts). Three cores 0.6 mm in diameter were obtained from each breast cancer sample and inserted into the recipient TMA blocks.

Immunohistochemical staining for the MT1 receptor was performed on 5- μ m paraffin sections cut from TMA blocks, using the following antibody: Melatonin Receptor (MT2/MTNR1B) from LS Bio (Catalog #: LS-A930; Lot #: 7195/7196AP3-1; 1:1000 dilution). A maximum of three cores were reviewed for each tumor, and degree of MT1 receptor staining in the epithelium was assessed using image analysis software (Definiens, Carlsbad, CA). Mean staining across available cores was used to define the outcome; epithelial staining was scored as positive if the mean percentage of stained cells was \geq 50%; it was otherwise considered negative. This cut point was determined by maximizing the accuracy of image analysis software results compared to those of an expert pathologist, who manually reviewed one of the TMAs (n=126 participants, with concordance of MT1R positive and negative ratings=89% and 79%, respectively). Because there is no established cut point for MT1R positivity, we also defined alternative cut points based on the absolute distribution of mean

epithelial staining (1/3, >1/3 to 2/3, and >2/3 of cells positive) and relative distribution of mean epithelial staining (in tertiles).

Statistical analysis

Of the 18,643 cohort participants with urine samples, 606 women without a prior history of cancer developed incident breast cancer during the follow-up period; they were matched to 1,085 cancer-free controls. We excluded 279 women whose urine sample was not assayed for aMT6s, nine women whose aMT6s values were statistical outliers (according to the generalized extreme studentized deviate many-outlier detection approach(24)), and 23 women who were not postmenopausal at the time of urine collection. In addition, we excluded 26 women whose matched pair was removed based on these exclusions. The remaining 1,354 women (555 breast cancer cases and 799 matched controls, with 158 having MT1R subtype information) comprised our analytic sample.

For these analyses, aMT6s levels below the limit of detection (n=29) were conservatively set to 0.8 ng/mL (i.e., the lower limit of detection), and aMT6s levels were standardized to creatinine. Because absolute aMT6s levels assayed by the Wilson laboratory were consistently lower compared to the Carroll laboratory, we used a previously described method to recalibrate these measurements by accounting for batch-to-batch variability that was not explained by age and body-mass index (BMI) (i.e., the strongest predictors of aMT6s levels)(25, 26).

Using logistic regression models conditioned on matching factors, we estimated simple and multivariable-adjusted odds ratios to approximate relative risks [RR] of breast cancer (with corresponding 95% confidence intervals [CI]) across quartiles of creatinine-standardized urinary aMT6s levels (based on re-calibrated values). Our analyses extend a previous study of urinary melatonin and incident breast cancer in this cohort, which included follow-up time up until May 31, 2006(7); thus, to facilitate a comparison with previous findings, we also calculated effect estimates based on the second half of follow up (i.e., cases occurring between June 1, 2006 and May 31, 2012, and their matched controls). In the subset of case-control pairs for which MT1R subtyping was available in the cases, we obtained separate estimates for MT1R-positive vs. negative tumors (defined as a tumor with 50% vs. <50 percent of cells with MT1R staining) using similar models. We used batch-corrected quartiles based on the creatinine-standardized aMT6s distribution in the controls; the lowest quartile served as the reference category. Tests of linear trend were calculated across quartiles using the Wald statistic, and tests of heterogeneity were calculated across breast cancer subtypes (invasive vs. in situ and MT1R-positive vs. negative) using polytomous conditional logistic regression.

Multivariable-adjusted models were controlled for possible breast cancer risk factors: age at menarche (<12, 12, 13, 14 years), age at menopause (45, 46–50, 51–53, 54 years), parity (nulliparous, 1–2, 3–4, or 5 children), family history of breast cancer (mother or sister: yes, no), (BMI continuous in kg/m²), alcohol intake (0, 1–14.9, 15 g/day), smoking status (current: yes, no), history of benign breast disease (yes, no), PMH use (current: yes, no), and type and duration of PMH use (estrogen-only: 0, <5, 5 years; estrogen + progesterone: 0,

<5, 5 years). Antidepressant use (current: yes, no) and first morning urine (yes, no) variables were included in our models to reduce variation in urinary melatonin levels.

We conducted several secondary analyses. First, we repeated our overall analysis of urinary melatonin and incident breast cancer while restricting to women with first-morning urine samples. Second, we repeated our analysis restricting our sample to estrogen- and progesterone-positive cases (i.e., the majority of breast cancer cases in our sample) because melatonin might influence breast cancer development via hormone-dependent pathways(3, 4). Third, we re-evaluated the association between urinary melatonin and incident breast cancer while implementing a two-year lag between time of urine collection and breast cancer diagnosis to minimize the possibility of subclinical breast tumors affecting urinary melatonin levels. Finally, in analyses incorporating the MT1R status of breast tumors, we explored alternative categories for defining MT1R-positivity (low, medium, and high) based on the absolute and relative distributions of mean epithelial staining, as described above.

Statistical tests were two sided and were considered statistically significant at $p < 0.05$. All analyses were conducted using SAS version 9.3 (SAS Institute, Cary, NC).

RESULTS

When we examined the distribution of breast cancer risk factors, we observed several expected patterns in our sample: breast cancer cases were more likely to have a family history of breast cancer, and a history of benign breast disease compared to controls (Table 1). In addition, cases had a higher mean intake of alcohol relative to controls. Among 158 cases with MT1R subtyping, women with MT1R-positive tumors were less likely to be current smokers, recent PMH users, and estrogen-only PMH users for 5 years compared to women with MT1R-negative tumors; MT1R-positive cases also had higher alcohol consumption compared to MT1R-negative cases.

There were few meaningful differences in participant characteristics at the time of urine collection according to quartiles of urinary melatonin level; however, as expected, mean age and BMI decreased across increasing melatonin quartiles (Table 2).

In our main analyses, we observed a suggestive trend of decreasing risk of overall breast cancer (including invasive and in situ cases) across increasing melatonin quartiles during a mean of 4.0 years of follow up in simple and multivariable-adjusted models (p -trend=0.08 for both models) (Table 3). Point estimates were similar in simple vs. multivariable-adjusted models, with significantly reduced risk comparing quartile 4 vs. quartile 1 in the fully-adjusted model (i.e., the 95% CI for this comparison did not include the null value of 1.00) (RR=0.78 95% CI=0.61–0.99). When we examined invasive and in situ cases separately, there were similar trends with decreasing risk of breast cancer observed over increasing quartiles of melatonin (p -heterogeneity=0.12), although neither of these individual trends reached statistical significance (p -trend for final models=0.19 for invasive cases and 0.09 for in situ cases). When we evaluated these associations based on cases occurring during the second half of follow up (and their matched controls), we found that urinary melatonin levels were not related to risk of breast cancer during that period (multivariable-adjusted RR:

1.05, 95% CI: 0.66–1.67 comparing quartile 4 vs. quartile 1 of urinary melatonin, p-trend=0.75) (Supplemental Table).

In the subset of 158 case-control pairs with MT1R subtyping in the cases, we found a similar pattern of overall association between urinary melatonin and breast cancer (Table 4). In multivariable-adjusted models, associations were also similar when we separately examined MT1R-negative cases (RR=0.60, 95% CI=0.35–1.06 comparing tertile 3 vs. tertile 1, p-trend=0.07) and MT1R-positive cases (RR=0.76, 95% CI=0.39–1.51 comparing tertile 3 vs. tertile 1, p-trend=0.39) (p-heterogeneity=0.88). When we restricted to invasive cases only (there were not enough in situ cases to examine them separately), these results were largely unchanged, and there were no differences of association found comparing MT1R-positive vs. negative cases (p-heterogeneity=0.57).

In secondary analyses, our results were very similar when we restricted our analysis of urinary melatonin and incident breast cancer to women with first-morning urine samples (multivariable-adjusted estimate comparing quartile 4 vs. quartile 1 of urinary melatonin: RR=0.79, 95% CI=0.61–1.03, p-trend=0.09). Results were also similar when we examined estrogen- and progesterone-positive cases only (multivariable-adjusted estimate comparing quartile 4 vs. quartile 1 of urinary melatonin: RR=0.82, 95% CI=0.60–1.11, p-trend=0.19), and after excluding cases diagnosed within two years of urine collection (multivariable-adjusted estimate comparing quartile 4 vs. quartile 1 of urinary melatonin: RR=0.79, 95% CI=0.58–1.06, p-trend=0.22) (data not shown in tables). For analyses based on MT1R status of tumors, results were similar when we utilized alternative cut points for the definition of MT1R positivity. For example, use of cut points based on absolute values of the distribution yielded the following results: multivariable-adjusted estimates comparing tertile 3 vs. tertile 1 of urinary melatonin: RR=0.59, 95% CI=0.21–1.65, p-trend=0.41, for low MT1R positivity; RR=0.67, 95% CI=0.33–1.33, p-trend=0.14, for medium MT1R positivity; RR=1.00, 95% CI=0.44–2.29, p-trend=0.90, for high MT1R positivity; p-heterogeneity=0.80) (data not shown in tables).

DISCUSSION

Overall, we found a suggestion that higher levels of urinary melatonin were related to lower risk of breast cancer risk over twelve years of follow up, which is consistent with previous findings with shorter follow up in this cohort. However, this association appeared to be driven by results from the first half of follow up, as there was no association between melatonin levels and breast cancer risk during the second half of follow up. In a subset of women with MT1R subtyping, we found no evidence that the association between urinary melatonin and incident breast cancer differed comparing MT1R-positive vs. negative tumors. To our knowledge, this is the first epidemiologic study to evaluate this association according to MT1R subtype of the breast tumor.

Two recent meta-analyses reported that higher levels of urinary melatonin were associated with a reduced risk of breast cancer in women based on published studies(27, 28). The summary relative risk of one meta-analysis indicated an approximately 18% decreased risk comparing quantile 4 vs. quartile 1 of urinary melatonin, with borderline significance

(RR=0.82, 95% CI=0.68–0.99)(27); this is very similar to the 22% risk reduction that we identified comparing quartile 4 vs. quartile 1 in our present study (OR=0.78, 95% CI=0.61–0.99). Results of the second meta-analysis are also similar to our results, with an inverse association observed between urinary melatonin and breast cancer incidence, which appeared to be confined to postmenopausal women (RR=0.81 95% CI=0.70–0.92 for each 15ng/mg creatinine increase in aMT6s)(28); however, the effect estimate comparing quartile 4 vs. quartile 1 was not given, making it more difficult to compare these results to those of the present study.

Indeed, one of the studies included in these meta-analyses is our previous analysis of the association between urinary melatonin levels and breast cancer risk in the NHS cohort(7). With shorter follow up, our prior study identified a 38% lower risk of breast cancer among women in the quartile 4 vs. quartile 1 of urinary melatonin (RR=0.62, 95% CI=0.41–0.95); the p-trend was also highly significant (p=0.004), although the risk appeared to be equally lower in the top two quartiles. In addition, exclusion of breast cancer cases diagnosed within one to two years of urine collection somewhat attenuated the effect estimate and it became non-significant (RR=0.72, 95% CI=0.45–1.17, with a one-year lag; and RR=0.76, 95% CI=0.44–1.31, with a two-year lag). In the current study, we observed that this association was attenuated further and became null during the period six to twelve years after urine collection. Taken together, these results suggest that endogenous melatonin levels could influence breast cancer incidence in the shorter but not longer term.

With regard to the melatonin receptor, to date, few studies have assessed MT1R status of breast tumors in women. One study with MT1R information identified an association between MT1R positivity and survival among women with triple-negative breast cancer(29). These findings suggest that greater numbers of MT1R may be related to better outcomes in breast cancer, but more specific interpretation is difficult due to the lack of urinary melatonin assessment and restriction to triple-negative breast tumors in this population. More directly related to our analyses, a second study examined the association between melatonin receptor genes and breast cancer incidence in Chinese women, and results suggested that genotype might influence risk of breast cancer(30). Nonetheless, differences in genetic variation across different ethnic populations could limit the generalizability of these findings, and no additional studies have been undertaken in other ethnic populations. Although the current study does not suggest differences in the association between endogenous melatonin levels and breast cancer risk according to MT1R type, experimental evidence demonstrating an age-related decrease in sensitivity of these receptors for melatonin might explain this null finding(16, 17). Clearly, additional studies are needed to replicate our findings in postmenopausal women, as well as explore these associations in premenopausal women.

Our study is unique because it incorporates information on both urinary melatonin levels and MT1R subtype of breast tumors in women. An important limitation, however, is the relatively small number of women in our cohort (29%) with both of these measurements, which limits our power to detect differences in the association between urinary melatonin and breast cancer by MT1R subtype. Moreover, because of this small number of cases, we were unable to examine these associations for in situ cases separately. In addition, there are limitations with our measurements of urinary melatonin levels and MT1R subtyping.

Because urine was collected at only one point in time, it is possible that melatonin levels at that time may not reflect melatonin levels over time; however, previous research has established that urinary melatonin levels are highly correlated over three years in this cohort (intraclass correlation=0.72), which lends more credibility to the notion that one-time melatonin assessment might contain relevant information about longer-term melatonin levels(20). For MT1R subtyping, there is little established information on the optimal cut point for characterizing positivity of this tumor marker. Thus, in our study, we utilized several different approaches to this definition, including ones based on a comparison of pathologist vs. machine read values, as well as absolute and relative values of our distribution. Results were similar comparing these various approaches, and therefore this limitation is less likely to have affected our interpretation of results. Finally, we assessed MT1R expression of breast tumors because this receptor is thought to largely mediate melatonin's anti-cancer effects; however, it is possible that expression of MT2R and key downstream signaling proteins play an important role in mediating this association, and we have not assessed these tumor factors in our cohort(18).

In summary, higher melatonin levels were associated with reduced breast cancer risk in postmenopausal women—consistent with our previous finding based on shorter follow up in this cohort. Moreover, there was no difference in the association comparing MT1R-positive vs. negative breast tumors. Future research should evaluate these associations with longer-term follow up, a larger number of cases with MT1R subtyping, and among premenopausal women.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: This study was funded by the National Cancer Institute (P01 CA87969 to M.J. Stampfer), which also supports the Nurses' Health Study (UM1 CA186107 to M.J. Stampfer). Additional support was provided by the Centers for Disease Control and Prevention/National Institute of Occupational Safety and Health (R01 OH009803 to E.S. Schernhammer.).

We would like to thank our participants for their commitment to research in the Nurses' Health Study. We also would like to thank Meir J. Stampfer for his role in obtaining funding for research in this cohort.

References

1. Stevens RG, Brainard GC, Blask DE, Lockley SW, Motta ME. Breast cancer and circadian disruption from electric lighting in the modern world. *CA: a cancer journal for clinicians*. 2014; 64:207–18. [PubMed: 24604162]
2. Hardeland R, Cardinali DP, Srinivasan V, Spence DW, Brown GM, Pandi-Perumal SR. Melatonin--a pleiotropic, orchestrating regulator molecule. *Progress in neurobiology*. 2011; 93:350–84. [PubMed: 21193011]
3. Hill SM, Frasnich T, Xiang S, Yuan L, Duplessis T, Mao L. Molecular mechanisms of melatonin anticancer effects. *Integrative cancer therapies*. 2009; 8:337–46. [PubMed: 20050373]
4. Proietti S, Cucina A, Reiter RJ, Bizzarri M. Molecular mechanisms of melatonin's inhibitory actions on breast cancers. *Cellular and molecular life sciences : CMLS*. 2013; 70:2139–57. [PubMed: 23007844]

5. Schernhammer ES, Hankinson SE. Urinary melatonin levels and breast cancer risk. *Journal of the National Cancer Institute*. 2005; 97:1084–7. [PubMed: 16030307]
6. Schernhammer ES, Berrino F, Krogh V, Secreto G, Micheli A, Venturelli E, et al. Urinary 6-sulfatoxymelatonin levels and risk of breast cancer in postmenopausal women. *Journal of the National Cancer Institute*. 2008; 100:898–905. [PubMed: 18544743]
7. Schernhammer ES, Hankinson SE. Urinary melatonin levels and postmenopausal breast cancer risk in the Nurses' Health Study cohort. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2009; 18:74–9.
8. Schernhammer ES, Berrino F, Krogh V, Secreto G, Micheli A, Venturelli E, et al. Urinary 6-Sulphatoxymelatonin levels and risk of breast cancer in premenopausal women: the ORDET cohort. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2010; 19:729–37.
9. Brown SB, Hankinson SE, Eliassen AH, Reeves KW, Qian J, Arcaro KF, et al. Urinary melatonin concentration and the risk of breast cancer in Nurses' Health Study II. *American journal of epidemiology*. 2015; 181:155–62. [PubMed: 25587174]
10. Travis RC, Allen DS, Fentiman IS, Key TJ. Melatonin and breast cancer: a prospective study. *Journal of the National Cancer Institute*. 2004; 96:475–82. [PubMed: 15026473]
11. Wu AH, Stanczyk FZ, Wang R, Koh WP, Yuan JM, Yu MC. Sleep duration, spot urinary 6-sulfatoxymelatonin levels and risk of breast cancer among Chinese women in Singapore. *International journal of cancer Journal international du cancer*. 2013; 132:891–6. [PubMed: 22644618]
12. Jablonska K, Pula B, Zemla A, Owczarek T, Wojnar A, Rys J, et al. Expression of melatonin receptor MT1 in cells of human invasive ductal breast carcinoma. *Journal of pineal research*. 2013; 54:334–45. [PubMed: 23330677]
13. Rogelsperger O, Ekmekcioglu C, Jager W, Klimpfinger M, Konigsberg R, Krenbek D, et al. Coexpression of the melatonin receptor 1 and nestin in human breast cancer specimens. *Journal of pineal research*. 2009; 46:422–32. [PubMed: 19552766]
14. Dillon DC, Easley SE, Asch BB, Cheney RT, Brydon L, Jockers R, et al. Differential expression of high-affinity melatonin receptors (MT1) in normal and malignant human breast tissue. *American journal of clinical pathology*. 2002; 118:451–8. [PubMed: 12219788]
15. Collins A, Yuan L, Kiefer TL, Cheng Q, Lai L, Hill SM. Overexpression of the MT1 melatonin receptor in MCF-7 human breast cancer cells inhibits mammary tumor formation in nude mice. *Cancer letters*. 2003; 189:49–57. [PubMed: 12445677]
16. Hill SM, Cheng C, Yuan L, Mao L, Jockers R, Dauchy B, et al. Age-related decline in melatonin and its MT1 receptor are associated with decreased sensitivity to melatonin and enhanced mammary tumor growth. *Current aging science*. 2013; 6:125–33. [PubMed: 23895529]
17. Hill SM, Cheng C, Yuan L, Mao L, Jockers R, Dauchy B, et al. Declining melatonin levels and MT1 receptor expression in aging rats is associated with enhanced mammary tumor growth and decreased sensitivity to melatonin. *Breast cancer research and treatment*. 2011; 127:91–8. [PubMed: 20549340]
18. Hill SM, Belancio VP, Dauchy RT, Xiang S, Brimer S, Mao L, et al. Melatonin: an inhibitor of breast cancer. *Endocrine-related cancer*. 2015; 22:R183–204. [PubMed: 25876649]
19. Willett WC, Browne ML, Bain C, Lipnick RJ, Stampfer MJ, Rosner B, et al. Relative weight and risk of breast cancer among premenopausal women. *American journal of epidemiology*. 1985; 122:731–40. [PubMed: 4050766]
20. Schernhammer ES, Rosner B, Willett WC, Laden F, Colditz GA, Hankinson SE. Epidemiology of urinary melatonin in women and its relation to other hormones and night work. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2004; 13:936–43.
21. Tamimi RM, Baer HJ, Marotti J, Galan M, Galaburda L, Fu Y, et al. Comparison of molecular phenotypes of ductal carcinoma in situ and invasive breast cancer. *Breast cancer research : BCR*. 2008; 10:R67. [PubMed: 18681955]

22. Tamimi RM, Colditz GA, Hazra A, Baer HJ, Hankinson SE, Rosner B, et al. Traditional breast cancer risk factors in relation to molecular subtypes of breast cancer. *Breast cancer research and treatment*. 2012; 131:159–67. [PubMed: 21830014]
23. Collins LC, Botero ML, Schnitt SJ. Bimodal frequency distribution of estrogen receptor immunohistochemical staining results in breast cancer: an analysis of 825 cases. *American journal of clinical pathology*. 2005; 123:16–20. [PubMed: 15762275]
24. Rosner B. Percentage points for a generalized ESD many-outlier procedure. *Technometrics*. 1983; 25:165–72.
25. Rosner B, Cook N, Portman R, Daniels S, Falkner B. Determination of blood pressure percentiles in normal-weight children: some methodological issues. *American journal of epidemiology*. 2008; 167:653–66. [PubMed: 18230679]
26. Rice MS, Tworoger SS, Rosner BA, Pollak MN, Hankinson SE, Tamimi RM. Insulin-like growth factor-1, insulin-like growth factor-binding protein-3, growth hormone, and mammographic density in the Nurses' Health Studies. *Breast cancer research and treatment*. 2012; 136:805–12. [PubMed: 23085768]
27. Basler M, Jetter A, Fink D, Seifert B, Kullak-Ublick GA, Trojan A. Urinary excretion of melatonin and association with breast cancer: meta-analysis and review of the literature. *Breast care*. 2014; 9:182–7. [PubMed: 25177260]
28. Yang WS, Deng Q, Fan WY, Wang WY, Wang X. Light exposure at night, sleep duration, melatonin, and breast cancer: a dose-response analysis of observational studies. *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation*. 2014; 23:269–76.
29. Oprea-Ilies G, Haus E, Sackett-Lundeen L, Liu Y, McLendon L, Busch R, et al. Expression of melatonin receptors in triple negative breast cancer (TNBC) in African American and Caucasian women: relation to survival. *Breast cancer research and treatment*. 2013; 137:677–87. [PubMed: 23250547]
30. Deming SL, Lu W, Beeghly-Fadiel A, Zheng Y, Cai Q, Long J, et al. Melatonin pathway genes and breast cancer risk among Chinese women. *Breast cancer research and treatment*. 2012; 132:693–9. [PubMed: 22138747]

Table 1

Baseline characteristics (at urine collection in 2000) of breast cancer cases (n=555), including by MT1R subtype (n=158), and matched controls (n=799)

Characteristic	Controls (n=799)	All breast cancer cases (n=555)	MT1R negative cases (n=91)	MT1R positive cases (n=67)
Mean melatonin level, in ng/mg creatinine (SD)	29.8 (30.7)	26.6 (25.9)	29.1 (28.7)	24.2 (34.0)
Mean age, in years (SD) ^a	66.9 (6.8)	66.6 (6.8)	67.4 (7.1)	66.9 (6.5)
Mean age at menarche, in years (SD)	12.6 (1.4)	12.5 (1.4)	12.6 (1.3)	12.4 (1.4)
Mean age at menopause, in years (SD)	49.6 (4.7)	50.3 (5.0)	50.4 (4.3)	50.5 (4.7)
Mean number of births (SD)	3.2 (1.6)	3.1 (1.4)	3.2 (1.5)	3.1 (1.3)
Family history of breast cancer, %	16	23	23	21
Mean BMI, in kg/m ² (SD)	26.4 (5.2)	26.8 (5.0)	26.8 (4.7)	27.3 (4.8)
Mean alcohol intake, in g/day (SD)	5.3 (8.5)	6.3 (9.8)	5.8 (9.1)	6.8 (11.8)
Current smoker, %	4	6	9	3
History of benign breast disease, %	59	65	63	60
Recent PMH use, % ^a	56	66	73	66
Duration of PMH use (estrogen only), %				
None	58	63	54	72
<5 years	13	10	11	15
5 years	29	27	35	13
Duration of PMH use (estrogen + progesterone), %				
None	62	54	56	52
<5 years	15	14	6	15
5 years	23	32	38	33
First morning urine, %	95	93	97	97
ER+/PR+ tumor, %	n/a	74	78	75

^aThese are matching factors.

SD=standard deviation

Table 2

Baseline characteristics (at urine collection in 2000) of matched controls across quartiles of creatinine-standardized urinary melatonin level (n=799)

Characteristic	Quartiles of urinary melatonin levels (in ng/mg creatinine)			
	Quartile 1	Quartile 2	Quartile 3	Quartile 4
Mean melatonin level, in ng/mg creatinine (SD)	6.9 (3.0)	17.6 (3.4)	30.3 (4.3)	64.8 (44.2)
Mean age, in years (SD)	67.9 (6.6)	67.5 (6.6)	66.1 (6.9)	66.2 (6.9)
Mean age at menarche, in years (SD)	12.6 (1.4)	12.6 (1.3)	12.6 (1.3)	12.6 (1.4)
Mean age at menopause, in years (SD)	49.4 (4.4)	49.4 (5.3)	49.7 (4.2)	49.8 (4.9)
Mean number of births (SD)	3.3 (1.6)	3.2 (1.8)	3.2 (1.5)	3.3 (1.5)
Family history of breast cancer, %	14	17	13	19
Mean BMI, in kg/m ² (SD)	27.6 (5.8)	26.9 (5.4)	25.9 (4.9)	25.3 (4.1)
Mean alcohol intake, in g/day (SD)	5.5 (9.0)	4.9 (8.8)	5.9 (8.7)	4.9 (7.3)
Current smoker, %	7	3	3	4
History of benign breast disease, %	61	61	50	61
Recent PMH use, %	54	58	56	58
Duration of PMH use (estrogen only), %				
None	58	57	57	59
<5 years	13	15	12	12
5 years	29	28	31	29
Duration of PMH use (estrogen + progesterone), %				
None	62	61	62	63
<5 years	19	13	17	12
5 years	19	26	21	25
First morning urine, %	94	97	96	95

SD=standard deviation

Table 3

Relative risks (RR) and 95% confidence intervals of breast cancer across quartiles of creatinine-standardized urinary melatonin level (n=1,354)

	Quartiles of urinary melatonin levels (in ng/mg creatinine)				
	Quartile 1	Quartile 2	Quartile 3	Quartile 4	p-trend
Invasive + in situ cases combined					
No. of cases / no. of controls	162 / 200	148 / 199	136 / 201	109 / 199	
Simple RR ^a	1.00 (ref)	0.95 (0.76–1.19)	0.90 (0.72–1.13)	0.79 (0.62–1.01)	0.08
Multivariable RR ^b	1.00 (ref)	0.96 (0.77–1.20)	0.92 (0.73–1.16)	0.78 (0.61–0.99) ^c	0.08
Invasive cases only					
No. of cases / no. of controls	121 / 156	114 / 154	111 / 150	87 / 161	
Simple RR ^a	1.00 (ref)	0.97 (0.75–1.26)	0.97 (0.75–1.26)	0.80 (0.61–1.06)	0.15
Multivariable RR ^b	1.00 (ref)	0.99 (0.76–1.28)	1.01 (0.78–1.31)	0.80 (0.61–1.07)	0.19
In situ cases only					
No. of cases / no. of controls	41 / 44	34 / 45	25 / 51	22 / 38	
Simple RR ^a	1.00 (ref)	0.89 (0.57–1.41)	0.68 (0.42–1.12)	0.76 (0.45–1.28)	0.29
Multivariable RR ^b	1.00 (ref)	0.84 (0.52–1.35)	0.63 (0.38–1.06)	0.62 (0.36–1.07)	0.09

^aSimple conditional logistic regression model.

^bMultivariable conditional logistic regression model adjusted for the following: age at menarche (<12, 12, 13, 14 years), age at menopause (45, 46–50, 51–53, 54 years), parity (nulliparous, 1–2, 3–4, or 5 children), family history of breast cancer (mother or sister: yes, no), BMI (continuous in kg/m²), alcohol intake (0, 1–14.9, 15 g/day), smoking status (current: yes, no), history of benign breast disease (yes, no), antidepressant use (current: yes, no), PMH (current: yes, no), type and duration of PMH (estrogen only: 0, <5, 5 years; estrogen + progesterone: 0, <5, 5 years), and first morning urine (yes, no).

^cThis estimate is statistically significant because the 95% confidence interval does not include the null value of 1.00.

Table 4

Multivariable-adjusted relative risks (RR) and 95% confidence intervals of breast cancer, according to MTIR subtype, across tertiles of creatinine-standardized urinary melatonin level (n=403 [245 controls, 91 MTIR negative cases, 67 MTIR positive cases])

	Tertiles of urinary melatonin levels (in ng/mg creatinine)				
	Tertile 1	Tertile 2	Tertile 3	p-trend	p-heterogeneity
Invasive + in situ cases combined					
No. of cases / no. of controls	61 / 77	59 / 74	38 / 94		
MTIR negative + positive ^a	1.00 (ref)	1.01 (0.70–1.47)	0.67 (0.44–1.02)	0.05	
No. of cases / no. of controls	32 / 42	35 / 34	24 / 61		
MTIR negative ^b	1.00 (ref)	1.08 (0.64–1.81)	0.60 (0.35–1.06)	0.07	0.88
No. of cases / no. of controls	29 / 35	24 / 40	14 / 33		
MTIR positive ^b	1.00 (ref)	0.95 (0.53–1.70)	0.76 (0.39–1.51)	0.39	
Invasive cases only					
No. of cases / no. of controls	49 / 64	44 / 57	32 / 73		
MTIR negative + positive ^a	1.00 (ref)	0.98 (0.64–1.50)	0.72 (0.45–1.14)	0.14	
No. of cases / no. of controls	29 / 38	29 / 32	23 / 54		
MTIR negative ^b	1.00 (ref)	1.00 (0.57–1.75)	0.65 (0.37–1.17)	0.14	0.57
No. of cases / no. of controls	20 / 26	15 / 25	9 / 19		
MTIR positive ^b	1.00 (ref)	0.88 (0.40–1.95)	0.97 (0.39–2.42)	0.79	

^aMultivariable conditional logistic regression model adjusted for the following: age at menarche (<12, 12, 13, 14 years), age at menopause (45, 46–50, 51–53, 54 years), parity (nulliparous, 1–2, 3–4, or 5 children), family history of breast cancer (mother or sister: yes, no), BMI (continuous in kg/m²), alcohol intake (0, 1–14.9, 15 g/day), smoking status (current: yes, no), history of benign breast disease (yes, no), antidepressant use (current: yes, no), PMH use (current: yes, no), type and duration of PMH use (estrogen only: 0, <5, 5 years; estrogen + progesterone: 0, <5, 5 years), and first morning urine (yes, no).

^bMultivariable conditional polytomous logistic regression model with same covariates as listed above for conditional logistic regression.