

# CDC PUBLIC HEALTH GRAND ROUNDS

## Emerging Tickborne Diseases



March 21, 2017



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Expanding Diversity and Distribution of Tickborne Diseases



**Rebecca Eisen, PhD**

*Research Biologist*

Division of Vector-Borne Diseases

National Center for Emerging and Zoonotic Infectious Diseases



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# The Basics of Tickborne Diseases

- **All known tickborne infectious diseases are diseases of animals that can be transmitted to humans via a tick vector (e.g., zoonoses)**
  - Ticks can maintain the pathogens through transmission to their offspring
  - Ticks can acquire infection through feeding on infectious hosts
- **Humans are incidental hosts, infected by the bite of infected ticks**

# Ticks Can Transmit Diverse Types of Bacteria in the United States

## Bacterial Diseases (9)

## Pathogens (14)

## Tick Genera (5)

Anaplasmosis

*Anaplasma phagocytophilum*

*Ixodes* spp.

*Borrelia miyamotoi* disease

*Borrelia miyamotoi*

*Ixodes* spp.

Ehrlichiosis

*Ehrlichia chaffeensis*

*Ehrlichia ewingii*

*Ehrlichia muris eauclarensis*

*Amblyomma* spp.

*Ixodes* spp.

Lyme disease

*Borrelia burgdorferi*

*Borrelia mayonii*

*Ixodes* spp.

*Rickettsia parkeri* rickettsiosis

*Rickettsia parkeri*

*Amblyomma* spp.

Rocky Mountain spotted fever

*Rickettsia rickettsii*

*Dermacentor* spp.

*Rhipicephalus* spp.

Pacific Coast tick fever

*Rickettsia philipii*

*Dermacentor* spp.

Relapsing fever

*Borrelia hermsii*

*Borrelia parkeri*

*Borrelia turicatae*

*Ornithodoros* spp.

Tularemia

*Francisella tularensis*

*Amblyomma* spp.

*Dermacentor* spp.

# Other Types of Pathogens Ticks Can Transmit


Diseases (4)	Pathogens (4)	Tick Genera (3)
<b>Viruses</b>		
Colorado tick fever	Colorado tick fever virus (Coltivirus)	<i>Dermacentor spp.</i>
Heartland virus disease	Heartland virus (Phlebovirus)	<i>Amblyomma spp.</i>
Powassan encephalitis	Powassan virus (Flavivirus)	<i>Ixodes spp.</i>
<b>Protozoa</b>		
Babesiosis	<i>Babesia microti</i>	<i>Ixodes spp.</i>

# Overview of Trends

- **Majority of vector-borne diseases in the U.S. are tickborne diseases**
- **Increasing number of tickborne disease cases over time**
- **Expanding geographic range of tickborne cases**
- **Growing number of tickborne agents recognized to cause human disease**



# Cases of Nationally Notifiable Vector-borne Diseases Reported in the U.S., 2014



# Three Species Cause Majority of Human Diseases

- *Ixodes scapularis* (Blacklegged tick)
- *Amblyomma americanum* (Lone star tick)
- *Dermacentor variabilis* (American dog tick)



*Ixodes scapularis*



*Amblyomma americanum*



*Dermacentor variabilis*

# Distribution of *Ixodes scapularis* (Blacklegged Tick)



## Transmits agents that cause:

- Anaplasmosis
- Babesiosis
- *Borrelia miyamotoi* disease
- Ehrlichiosis
- Lyme disease
- Powassan encephalitis



Adult female



Adult male



Nymph



Larva

# Distribution of *Amblyomma americanum* (Lone star tick)



**Transmits agents that cause:**

- Ehrlichiosis
- Tularemia
- Heartland virus disease



**Adult female**



**Adult male**



**Nymph**



**Larva**

# Distribution of *Dermacentor variabilis* (American Dog Tick)



**Transmits agents that cause:**

- Rocky Mountain spotted fever
- Tularemia



**Adult female**



**Adult male**




**Nymph**




**Larva**

# Geographic Distribution of Nationally Notifiable Tickborne Diseases, 2015

**Lyme disease**



**Anaplasmosis**




**Babesiosis**



**Ehrlichiosis**



**Rocky Mountain Spotted Fever**




**Tularemia**




Each dot represents a reported case in the county of residence

# Cases of Lyme Disease Have Increased


Annual Reported Cases of Lyme Disease, 1992–2015




# Other Nationally Notifiable Tickborne Diseases Have Also Increased



# Expanding Numbers and Geographic Distribution of Lyme Disease Cases Mirrors Other Tickborne Diseases




**2001**



**2015**


# Discovery of Tickborne Pathogens as Causes of Human Disease By Year, 1909–1959



Year represents when tickborne pathogen was recognized as cause of human disease.

Adapted from: Paddock CD, Lane RS, Staples JE, Labruna MB. 2016. In: Mack A, Editor. Global health impacts of vector-borne diseases: workshop summary. National Academies Press. p. 221-257.

# Discovery of Tickborne Pathogens Has Accelerated, 1960–2016



Year represents when tickborne pathogen was recognized as cause of human disease.

Adapted from: Paddock CD, Lane RS, Staples JE, Labruna MB. 2016. In: Mack A, Editor. Global health impacts of vector-borne diseases: workshop summary. National Academies Press. p. 221-257.

# Explanation of Increasing Cases and Geographic Spread


- Improved diagnostics and clinical recognition
- Range expansion and population increases of ticks
- Lack of effective prevention strategies



A blood-fed *Amblyomma americanum*,  
“Lone star tick”

# Geographic Expansion of Ticks

## Locations Where *Ixodes scapularis* Recorded




- Established:**  $\geq 6$  or more ticks or  $\geq 1$  life stage recorded in a single year
- Reported:**  $< 6$  individuals of a single life stage recorded in a single year

Dennis DT, Nekomoto TS, Victor JC, et al. J Med Entomol. 1998 Sep;35(5):629-38.

Eisen RJ, Eisen L, Beard CB. J Med Entomol. 2016 Mar;53(2):349-86.


# Geographic Expansion of Vectors Matches Increases in Tickborne Disease Cases

## Changes in *I. scapularis* distribution, 1996–2015



**■ Established as of 1996**   **■ Reported as of 2015**  
**■ Established as of 2015**

## Time Period When County First Reached High Incidence of Lyme Disease



**■ 1993–1997**   **■ 2003–2007**  
**■ 1998–2002**   **■ 2008–2012**

Established defined as: 6 or more ticks; or 1 or more; Reported defined as: 6 or more ticks or >1 life stage recorded in a single year

Adapted from: Eisen RJ, Eisen L, Beard CB. *J Med Entomol.* 2016 Mar;53(2):349-86.

Kugeler KJ, Farley GM, Forrester JD, et al. *Emerg Infect Dis.* 2015 Aug;21(8):1455-7.

# Prevention in the Absence of Vaccines

- **Increasing human contact with ticks**
- **No human vaccines to prevent tickborne diseases in the U.S.**
  - No single effective, widely accepted method of preventing tickborne diseases
- **Prevention strategies include**
  - Personal protection
  - Environmental modification
  - Tick suppression

# Prevention Through Personal Protection

- **Avoid tick habitat**
- **Repel ticks**
  - Use 20–30% DEET on exposed skin and clothing
  - Wear permethrin-treated clothing




# Prevention Through Personal Protection

## ➤ Find and remove ticks from your body

- Bathe or shower as soon as possible after coming indoors
- Every day check for and remove ticks on body, pets and outdoor gear

## ➤ Tumble dry clothing

- Tumble dry clothes in a dryer on high heat for 10 minutes
- If the clothes are damp, additional time may be needed
- If the clothes require washing first, use hot water
  - ❑ Cold and medium temperature water will not kill ticks effectively



# Prevention Through Environmental Modification and Tick Suppression

## ➤ Reduce the numbers of host-seeking ticks

- Landscape management
- Kill host-seeking ticks with acaricides or biological agents



**Acaricide Treatment  
of Vegetation**



# Prevention Through Environmental Modification and Tick Suppression

- **Host reduction or exclusion**
  - Install deer-proof fencing
- **Reduce the numbers of ticks on hosts**
  - Acaricide treatment of rodents or deer

**Acaricide Treatment Station for Deer**



As deer feed, a pesticide (acaricide) is applied topically.

# Emerging Prevention Strategies

- **Reduce the number of infected hosts**
- **Rodent-targeted methods to reduce infection in ticks**
  - Vaccines
  - Antibiotics

**Bait Box for Rodents**



Attracted by the bait inside, rodents enter. Once inside, the rodent consumes the bait containing an oral vaccine or antibiotic.

# Looking Forward

- **Recognition of more tickborne diseases and tickborne agents**

# Looking Forward

- **Recognition of more tickborne diseases and tickborne agents**
- **Continuing range expansion of ticks and associated tickborne diseases**

# Looking Forward

- **Recognition of more tickborne diseases and tickborne agents**
- **Continuing range expansion of ticks and associated tickborne diseases**
- **Importance of co-infections**
  - Single species can carry multiple disease agents

# Looking Forward

- **Recognition of more tickborne diseases and tickborne agents**
- **Continuing range expansion of ticks and associated tickborne diseases**
- **Importance of co-infections**
  - Single species can carry multiple disease agents
- **Need for effective, widely-acceptable approaches to prevent tickborne diseases**

# Tickborne Spotted Fevers – Old and New



**Christopher D. Paddock, MD**

Medical Officer, Rickettsial Zoonoses Branch


Division of Vector-Borne Diseases

National Center for Emerging and Zoonotic Infectious Diseases

# A Long Time Ago, in A Valley Far, Far Away...



Image courtesy of Pete Ramberg



# Rocky Mountain Spotted Fever

## The First Recognized Tickborne Disease of Humans in the U.S.



Early RMSF patient with  
“spotted” rash

Years	Cases	Deaths	CFR
1880–1889	19	15	<b>79%</b>
1890–1899	99	70	<b>71%</b>
1900–1909	225	128	<b>57%</b>
1880–1909	343	213	<b>62%</b>




*Dermacentor andersoni*,  
the Rocky Mountain wood tick

CFR: Case fatality rate

Philip RN. 2000. Rocky Mountain spotted fever in Western Montana. Anatomy of a Pestilence. Bitter Root Valley Historical Society, Hamilton, MT.

RMSF: Rocky Mountain spotted fever


# Why is Rocky Mountain Spotted Fever So Deadly?



***Rickettsia rickettsii* bacteria infecting endothelial cells of small blood vessel**




**Petechial rash involving sole**



**Histologic section of lung from patient with fatal RMSF**


# The Favorable Impact of Doxycycline Therapy on Survival

- **Discovery of tetracycline-class drugs resulted in dramatic reduction in case-fatality rates**
- **Doxycycline is the drug of choice for patients of all ages with RMSF and other tickborne rickettsial diseases**
- **Doxycycline should be initiated immediately if RMSF is suspected**
  - Delay of therapy (> 5 days) is the most important predictor of fatal outcome




Doxycycline

# Incidence of Spotted Fever Group Rickettsioses (including RMSF) in the United States, 2008–2012



Incidence By County, 2008–2012  
Rate per million person-years



*Dermacentor variabilis*,  
the American dog tick

# The Brown Dog Tick Emerges as an Unexpected Vector of Rocky Mountain Spotted Fever (RMSF) in Arizona



*Rhipicephalus sanguineus*,  
the brown dog tick

- Not considered important in the epidemiology of RMSF in the U.S. until 2004
- Cause of epidemic levels of RMSF in several American Indian communities in Arizona
- Outbreaks precipitated by large populations of free-roaming, tick-infested dogs



# Community-Based Intervention Successfully Reduces the Number of Ticks and Cases of RMSF

- **Collaborative endeavor among tribal partners, IHS, private sector, Arizona DOH, USDA, Inter Tribal Council of Arizona, and CDC**
  - Yards treated with acaricide spray
  - Tick collars placed on dogs
- **Dramatic reduction in ticks on dogs and in the environment**
  - Ticks found on <1% of treated dogs vs. 64% of untreated dogs
- **43% reduction in cases of RMSF**



Placing tick collars on dogs

IHS: Indian Health Service

USDA: United States Department of Agriculture

Drexler N, Miller M, Gerding J, et al. *PLoS One*. 2014 Dec 5;9(12):e112368.

DHS: Department of Health Services

RMSF: Rocky Mountain spotted fever

# A Second Pathogen Emerges from the Past

## 1939

### OBSERVATIONS ON AN INFECTIOUS AGENT FROM *AMBLYOMMA MACULATUM*<sup>1</sup>

By R. R. PARKER, Director, Rocky Mountain Laboratory, GLEN M. KOHLS, Assistant Entomologist, United States Public Health Service, GEORGE W. COX, Executive Officer, Texas State Department of Public Health, and GORDON E. DAVIS, Bacteriologist, United States Public Health Service



*Amblyomma maculatum*,  
the Gulf Coast tick

## 2004


### *Rickettsia parkeri*: A Newly Recognized Cause of Spotted Fever Rickettsiosis in the United States

Christopher D. Paddock,<sup>1</sup> John W. Sumner,<sup>1</sup> James A. Comer,<sup>1</sup> Sherif R. Zaki,<sup>1</sup> Cynthia S. Goldsmith,<sup>1</sup> Jerome Goddard,<sup>2</sup> Susan L. F. McLellan,<sup>3</sup> Cynthia L. Tamminga,<sup>4</sup> and Christopher A. Ohl<sup>1,5</sup>

<sup>1</sup>Division of Viral and Rickettsial Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia; <sup>2</sup>Mississippi Department of Health, Jackson, Mississippi; <sup>3</sup>Infectious Diseases Section, Tulane University Health Sciences Center, New Orleans, Louisiana; <sup>4</sup>Division of Infectious Diseases, Portsmouth Naval Medical Center, Portsmouth, Virginia; and <sup>5</sup>Section on Infectious Diseases, Wake Forest University School of Medicine, Winston-Salem, North Carolina

(See the editorial commentary by Raoult on pages 812–3)

## 2014



# Rashes of Rocky Mountain Spotted Fever (RMSF) versus *Rickettsia parkeri* rickettsiosis




RMSF



*R. parkeri*  
rickettsiosis



# Distinctive Eschars of *Rickettsia parkeri* Rickettsiosis



Paddock CD, Sumner JW, Comer JA, et al. *Clin Infect Dis*. 2004 Mar 15;38(6):805-11.

Cragun WC, Bartlett BL, Ellis MW, et al. *Arch Dermatol*. 2010 Jun;146(6):641-8.

# RMSF versus *Rickettsia parkeri* rickettsiosis

Clinical Feature	RMSF (n = 398)	<i>R. parkeri</i> rickettsiosis (n = 21)
Fever	99%	100%
Headache	80%	86%
Any rash	92%	90%
Petechial rash	52%	14%
Pustular/vesicular rash	---	33%
Eschar	---	95%
Nausea/vomiting	66%	10%
Coma/seizure	27%	---
Death	8%	---

Paddock CD, Finley RW, Wright CS, et al. *Clin Infect Dis*. 2008 Nov 1;47(9):1188-96.

Cragun WC, Bartlett BL, Ellis MW, et al. *Arch Dermatol*. 2010 Jun;146(6):641-8.

Myers T, Lalani T, Dent M, et al. *Emerg Infect Dis*. 2013 May;19(5):778-80.

Ekenna O, Paddock CD, Goddard J. *J Miss State Med Assoc*. 2014 Jul;55(7):216-9.

# *Rickettsia parkeri* rickettsiosis: How Much is Out There?

Morbidity and Mortality Weekly Report

➤ **5 cases identified by one clinician in Mississippi during 2007–2012**

➤ **5 cases identified at one urgent care practice in Georgia during 2012–2014**


## Notes from the Field

### *Rickettsia parkeri* Rickettsiosis — Georgia, 2012–2014

Anne Straily, DVM<sup>1,2</sup>; Amanda Feldpausch, MPH<sup>3</sup>; Carl Ulbrich, DO<sup>1</sup>; Kiersten Schell<sup>4</sup>; Shannon Casillas, MPH<sup>3</sup>; Sherif R. Zaki, MD, PhD<sup>5</sup>; Amy M. Denison, PhD<sup>5</sup>; Marah Condit, MS<sup>2</sup>; Julie Gabel, DVM<sup>3</sup>; Christopher D. Paddock, MD<sup>2</sup>

During 2012–2014, five cases of *Rickettsia parkeri* rickettsiosis were identified by a single urgent care practice in Georgia, located approximately 40 miles southwest of Atlanta. Symptom onset occurred during June–October, and all patients had a known tick bite. Patients ranged in age from 27 to 72 years (median = 53 years), and all were male. The most commonly reported initial signs were erythema (n = 3) and swelling (n = 2) at the site of the bite. Two patients reported fever and a third patient reported a rash and lymphadenopathy without fever. Other symptoms included myalgia (n = 3), chills (n = 3), fatigue (n = 2), arthralgia (n = 2), and headache (n = 2). Eschar biopsy specimens were collected from each patient using a 4-mm or 5-mm punch and placed in 10% neutral buffered formalin or

FIGURE. Female (A) and male (C) Gulf Coast ticks (*Amblyomma maculatum*); (B) necrotic, ulcerated or scabbed lesion at the tick bite site, known as an inoculation eschar; and (D) immunohistochemical stain indicating the presence of a spotted fever group *Rickettsia* species in the tissue



# Percentage of Ticks Infected with *R. parkeri* Greater than Percentage of Ticks Infected with *R. rickettsii*

## Frequency of *R. rickettsii* in *D. variabilis* vs. *R. parkeri* in *A. maculatum*

### *Dermacentor variabilis*



Pathogen	Location, years	No. (% infected)
<i>R. rickettsii</i>	North Carolina, 1982	2,123 (0.05)
	Ohio, 1984–1989	12,631 (0.06)
	Maryland, 2002	392 (0)
	Tennessee, 2007–2008	555 (0)

### *Amblyomma maculatum*



<i>R. parkeri</i>	Florida, 2005–2007	128 (22)
	Mississippi, 2008–2012	698 (15)
	North Carolina, 2009–2010	101 (31)
	Virginia, 2010–2011	293 (53)

Paddock CD, Fournier PE, Sumner JW, et al. *Appl Environ Microbiol.* 2010 May;76(9):2689-96.

Pagac BB, Miller MK, Mazzei MC, et al. *Emerg Infect Dis.* 2014 Oct;20(10):1750-2.

Ferrari FA, Goddard J, Paddock CD, et al. *Emerg Infect Dis.* 2012 Oct;18(10):1705-7.

44 Varela-Stokes AS, Paddock CD, Engber B, et al. *Emerg Infect Dis.* 2011 Dec;17(12):2350-3.

Nadolny RM, Wright CL, Sonenshine DE, et al. *Ticks Tick Borne Dis.* 2014 Feb;5(1):53-7.

# Conclusions

- **Etiologic spectrum of tickborne rickettsioses in the U.S. has expanded during the past 15 years**
- **Rocky Mountain spotted fever and *R. parkeri* rickettsiosis share many clinical features but differ considerably in severity**
- **Doxycycline is the drug of choice for all tickborne rickettsioses and in patients of all ages**
  - Therapy should be initiated immediately, based on a presumptive diagnosis

# Tickborne Viruses: Emerging Public Health Concern



**Gregory D. Ebel, ScD**

*Associate Professor and Director*

Arthropod-borne and Infectious Diseases Laboratories

Colorado State University

# Worldwide Ticks Are Vectors For Diverse Array of Viral Pathogens and Diseases

Disease and Etiologic Agent	Signs and Symptoms	Geographic Location
<b>Colorado Tick Fever</b> Colorado Tick Fever	Flu-like symptoms (FLS), headache, rash	Western U.S.
<b>Crimean-Congo Hemorrhagic Fever</b> CCHF virus	FLS, hemorrhagic fever (HF)	Asia, Africa, and Europe
<b>Omsk Hemorrhagic Fever</b> Omsk hemorrhagic fever virus	FLS, HF	Southwestern Russia
<b>Kyasanur Forest Disease</b> Kyasanur forest disease virus	FLS, HF	Southern India, Saudi Arabia (aka Alkhurma disease in Saudi Arabia)
<b>Tick-Borne Encephalitis</b> Tick-borne encephalitis virus	FLS, encephalitis, HF	Temperate regions of Europe and northern Asia

# Ticks Are Highly Efficient Vectors

## ➤ Abundance

- One fully engorged female *Ixodid* yields several thousand offspring

## ➤ Host preference

- At least two stages of feeding (usually larvae and nymphs) on the same host



A female *Ixodes scapularis* (Blacklegged tick) engorged with a host blood meal

# Ticks Are Highly Efficient Vectors

## ➤ Longevity of ticks

- Once infected with a pathogen, ticks are likely to survive and to transmit pathogen to new host
  - Survival of pathogen's extrinsic incubation period

## ➤ Modification of the site of attachment

- Pharmacology of tick saliva



## ➤ Favorable gut environment

- Intracellular digestion of bloodmeal

# Emerging Tickborne Viruses in the United States

- **Powassan virus (*Flaviviridae: Flavivirus*)**
  - Isolated 1958, Powassan, Ontario
- **Sole North American representative of the tickborne encephalitis complex of the Flaviviruses**
- **Severe acute disease**
  - 10.5% case-fatality rate
- **Long-term sequelae**
  - 47% of cases experience hemiplegia, wasting, severe headaches, etc.
- **Transmission depends on ecological factors that are not well understood**


# Diverse Vectors and Transmission Cycles Maintain Powassan Virus



Deer tick-associated viruses appear to pose greatest public health risk

# Infection Among Deer Is Increasing

Serological Prevalence of POWV in Hunter-killed Deer, 1979–2010



POWV: Powassan virus

Nofchissey RA, Deardorff ER, Blevins TM, et al. *Am J Trop Med Hyg.* 2013 Jun;88(6):1159-62.

# Clinical Features of Powassan Virus Disease Consistent Through Time


**Demographic, Clinical Features and Outcomes of 8 Patients  
with Powassan Virus Encephalitis, 2013–2015**

Patient Age and Gender	Exposure	Fever	Rash	Gastro-intestinal	Outcome
82 M	Outdoors	Y	N	Vomiting	Death
74 M	Outdoors, gardening	Y	N	None	Residual Deficits
21 M	Outdoors	Y	Y	Vomiting	Improved
67 M	Outdoors	Y	Y	Vomiting, diarrhea	Improved
65 F	Tick bite	Y	N	Vomiting	Improved
52 M	Tick bite	Y	N	None	Residual Deficits
49 M	Tick bite	Y	N	None	Death
44 M	Outdoors, hunting	N	Y	None	Improved


Adapted from: Piantadosi A, Rubin DB, McQuillen DP, et al. *Clin Infect Dis*. 2016 Mar 15;62(6):707-13.

# Number and Geographic Distribution of Reported Powassan Virus Cases, 2006-2015


## Powassan Virus Neuroinvasive Disease Cases Reported by Year, 2006–2015




## Powassan Virus Neuroinvasive Disease Cases Reported by State, 2006–2015



# Powassan Virus Causes Neurological Damage Both Directly and via Inflammatory Processes



**Inflammatory changes within the perivascular and parenchymal portions of the brain**



**Direct neuronal injury of the Purkinje cells**

# Discovery of New Tickborne Viruses Is Ongoing

## ➤ Heartland virus (*Bunyaviridae: Phlebovirus*)

- First 2 cases in Missouri, 2009
- Additional 6 confirmed cases in Tennessee and Missouri, 2012–2013
- CDC continues to investigate additional cases
- Few fatalities
- Lone star tick
  - ▣ (*Amblyomma americanum*)
- Wildlife serologic studies suggest widespread distribution of virus




*Amblyomma americanum*,  
"Lone star tick"

# Discovery of Another Virus Likely to Be Tickborne

## ➤ Bourbon virus (*Orthomyxoviridae: Thogotovirus*)


- First case in Kansas, 2015
- Likely tickborne
  - ▣ *In vitro* studies support
- Fatal outcome



Numerous extracellular Bourbon virus virions with slices through strands of viral nucleocapsids

# Knowledge Gaps and Unmet Needs


- **As new viruses emerge, so do questions and concerns**
- **What determines human risk and likelihood of emergence as human threat? How best to intervene?**



**Forest cover in the U.S., East of the Mississippi, from colonization to the present**  
**Reforestation Drives Emergence**

# Ecology: How Do Tickborne Arboviruses Perpetuate in Nature?

- Modeling and experimental studies show multiple transmission routes required
- Model parameters obtainable and transmission cycles can be studied
- Niche complexity




Nonaka E, Ebel GD, Wearing HJ. *PLoS One*. 2010 Jul 23;5(7):e11745.

Nuttall PA, Labuda M. *Adv Virus Res*. 2003;60:233-72.

Davis S, Bent SJ. *J Theor Biol*. 2011 Jan 21;269(1):96-103.

# How Do Ticks, Hosts, and Pathogens Interact?

- **What factors influence tick vector competence at the molecular level?**
- **Most work covers tick salivary secretions**
  - Hundreds of protein and non-protein molecules
    - ❑ Affect hemostasis, inflammation, i.e., host immune response to tick feeding
    - ❑ Effect of pathogens poorly understood




*Ehrlichia microti*. Polyhedral clusters of rickettsiae (arrows) within hypertrophied salivary acinus

# Figuring Out Function of Tick Genes

## ➤ Tick genome sequence

- Extremely large
- Most genes duplicated
- Metagenomics also contributes to key phenotypes



Distribution of assembled contigs from cDNA library from the salivary glands of adult female *Hyalomma marginatum rufipes* ticks


# Exciting Time to Study Tickborne Diseases



**In vitro  
and in vivo  
systems to  
study tick-  
borne  
viruses**



**Ecology of ticks  
and their hosts**



# Barriers to Rapid Advances in Tickborne Virus Research

## ➤ Technical

- Containment and regulatory restrictions
- System complexity
- Long tick life cycle
  - ▣ One tick lifecycle per grant or funding cycle

## ➤ Environmental

- Tickborne viruses emerge slowly relative to mosquito-borne viruses
- Significance, impact, innovation

## ➤ Few researchers active



Note tick in tweezers

# Summary

- **Tick borne viruses are emerging health concerns**
- **Interesting, relevant and feasible questions present opportunities to the field**
- **Difficult technical and environmental barriers currently impede progress**

# Advances in Diagnosing Tickborne Diseases



**Bobbi S. Pritt, MD, MSc, DTM&H**

*Director, Clinical Parasitology Laboratory*

*Co-Director, Vector-borne Diseases Laboratory Services*

*Professor of Pathology and Laboratory Medicine*

*Division of Clinical Microbiology, Mayo Clinic*

# Primary Methods for Diagnosis of Tickborne Diseases

## ➤ Clinical evaluation

## ➤ Tickborne diseases must be considered in differential diagnosis

- Order correct laboratory tests
- Begin empirical antimicrobial therapy if indicated
- Especially if rickettsiosis, ehrlichiosis, or anaplasmosis is suspected
  - ❑ Infections may be rapidly fatal
  - ❑ Need to be treated quickly, often before test results are available

Rash of Rocky Mountain spotted fever



Eschar of *Rickettsia parkeri* rickettsiosis



# Indirect Laboratory Methods: Serology


## ➤ Principle:

- Detection of the host's immune response to organisms
- IgM or IgG class host antibodies in serum

## ➤ Uses:


- Method of choice for diagnosing many tickborne diseases
  - *Rickettsia*, *Ehrlichia* and *Anaplasma* species, tickborne viruses
- Testing for tickborne viruses uses laboratory-developed methods
  - Available primarily through state public health laboratories or CDC
- Not primary diagnostic choice for babesiosis
- Sensitivity varies by the time that the specimen is obtained during the course of infection

# Serology: IgM and IgG Patterns




# Direct Laboratory Methods for Diagnosis of Tickborne Diseases


## ➤ Microscopy



*Anaplasma phagocytophilum*  
morula




Relapsing fever  
*Borrelia* spp.  
spirochetes



*Babesia* spp.  
parasites  
within red  
blood cells

- Nucleic acid amplification tests (e.g., polymerase chain reaction)
- Culture not routinely used, except for tularemia

# Direct Methods - General Pattern



# Nucleic Acid Amplification Tests – What Can They Do?

- **One of the earliest methods for detecting some organisms**
- **Insensitive if nucleic acid is no longer detectable by the time the patient presents for evaluation**
- **Especially useful for *Anaplasma phagocytophilum*, *Ehrlichia* species and *Babesia* species (whole blood specimens)**
  - These organisms are usually present in high amounts during the symptomatic stage of infection
- **Variety of nucleic acid amplification methods available**
  - None are FDA-approved/cleared


# Polymerase Chain Reaction

- **Fluorescent probes are used to detect DNA as it is amplified**
  - Can be highly sensitive and specific
  - Semiquantitative detection
- **Melting curve analysis**
  - Uses nonspecific DNA-binding dyes or specific probes
  - Additional benefit of detecting mutations (and even new organisms)



# Example: Melting Temperature Analysis with Fluorescence Resonance Energy Transfer (FRET) Hybridization Probes

Real-time PCR targeting *groEL* (gene encoding the heat shock operon) of *Ehrlichia* spp.




Previous name:  
*Ehrlichia muris-like agent* (EMLA)

New accepted name:  
*Ehrlichia muris euclairensis*

# Example: Melting Temperature Analysis with Fluorescence Resonance Energy Transfer (FRET) Hybridization Probes

Real-time PCR targeting *groEL* (gene encoding the heat shock operon) of *Ehrlichia* spp.



Previous name:  
*Ehrlichia muris-like agent (EMLA)*

New accepted name:  
*Ehrlichia muris eauclairensis*

This same method allowed for detection of *Borrelia mayonii*

# Beyond Singleplex PCR

## ➤ **Multiplex molecular panels**

- Specific primers/probes to detect multiple bacterial, viral, and parasitic pathogens in a single test
  - ❑ Panels for tickborne disease under development
- Only detects targeted organisms, non-targeted organisms will be missed

## ➤ **Broad range sequencing**

- Targeted gene amplification with subsequent sequence identification
  - ❑ Common targets: 16S rRNA gene (bacteria), internal transcribed spacer (ITS) region (fungi)
  - ❑ No equivalent for viruses; instead, groups of closely related viruses can be targeted (e.g. flavivirus)


# Next Steps in Testing

## ➤ Metagenomics

- Amplification all nucleic acid in a specimen: bacterial, fungal, viral, parasitic and human
- Extensive pre- and post-processing steps used to select for targets of interest and remove nonrelevant nucleic acid
- Currently very expensive and time consuming; this is bound to change in the future!




# Recommended Reading




# Thank you

# Resources for Tickborne Diseases



[cdc.gov/lyme/resources/tickbornediseases.pdf](https://cdc.gov/lyme/resources/tickbornediseases.pdf)



[cdc.gov/mobile/applications/mobileframework/tickborne-diseases.html](https://cdc.gov/mobile/applications/mobileframework/tickborne-diseases.html)

[www.cdc.gov/ticks](https://www.cdc.gov/ticks)

# CDC PUBLIC HEALTH GRAND ROUNDS

## Emerging Tickborne Diseases



March 21, 2017



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention