

District of Columbia	40 (1)	0 (0)
Florida‡	1,131 (22)	26 (59)
Georgia	110 (2)	0 (0)
Hawaii	16 (<1)	0 (0)
Idaho	4 (<1)	0 (0)
Illinois	105 (2)	0 (0)
Indiana	51 (1)	0 (0)
Iowa	27 (1)	1 (2)
Kansas	22 (<1)	0 (0)
Kentucky	33 (1)	0 (0)
Louisiana	39 (1)	0 (0)
Maine	14 (<1)	0 (0)
Maryland	134 (3)	0 (0)
Massachusetts	126 (3)	1 (2)
Michigan	72 (1)	0 (0)
Minnesota	64 (1)	0 (0)
Mississippi	25 (1)	0 (0)
Missouri	36 (1)	0 (0)
Montana	9 (<1)	0 (0)
Nebraska	13 (<1)	0 (0)
Nevada	22 (<1)	1 (2)
New Hampshire	13 (<1)	0 (0)
New Jersey	189 (4)	0 (0)
New Mexico	10 (<1)	0 (0)
New York	1,020 (20)	3 (7)
North Carolina	100 (2)	0 (0)
North Dakota	3 (<1)	0 (0)
Ohio	85 (2)	0 (0)
Oklahoma	29 (1)	0 (0)
Oregon	48 (1)	0 (0)
Pennsylvania	179 (4)	0 (0)
Rhode Island	55 (1)	0 (0)
South Carolina	57 (1)	0 (0)
South Dakota	3 (<1)	0 (0)
Tennessee	61 (1)	0 (0)
Texas‡	322 (6)	4 (9)

Utah	22 (<1)	0 (0)
Vermont	13 (<1)	0 (0)
Virginia	115 (2)	0 (0)
Washington	72 (1)	0 (0)
West Virginia	11 (<1)	1 (2)
Wisconsin	56 (1)	0 (0)
Wyoming	2 (<1)	0 (0)

Territories	Symptomatic disease cases**	Presumptive viremic blood donors†
	(N=36,569) No. (%)	(N=325) No. (%)
American Samoa	132 (<1)	0 (0)
Puerto Rico	35,418 (97)	325 (100)
US Virgin Islands	1,109 (3)	0 (0)

** Includes reported confirmed and probable Zika virus disease cases per the [CSTE case definitions](#).

† Presumptive viremic blood donors are people who reported no symptoms at the time of donating blood, but whose blood tested positive when screened for the presence of Zika virus RNA by the blood collection agency. Some presumptive viremic blood donors develop symptoms after their donation or may have had symptoms in the past. These individuals may be reported as both Zika virus disease cases and presumptive viremic blood donors.

‡ Includes 217 cases in Florida and 6 cases in Texas that were acquired through presumed local mosquito-borne transmission.