



ZIKA 101

Updated June 12, 2017



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention



INTRODUCTION

What is Zika?

- Zika virus is spread to people primarily through the bite of an infected *Aedes* species mosquito (*Ae. aegypti* and *Ae. albopictus*).
- Many people infected with Zika virus won't have symptoms or will only have mild symptoms.
- Zika virus infection during pregnancy can cause microcephaly and other severe brain defects.



Where has Zika been found?

- Before 2015, Zika outbreaks occurred in Africa, Southeast Asia, and the Pacific Islands.
- Currently outbreaks are occurring in many countries and territories.




<http://www.cdc.gov/zika/geo/index.html>

SPREAD AND SYMPTOMS


How is Zika spread?

- Zika can be spread through
 - » Mosquito bites
 - » From a pregnant woman to her fetus
 - » Sex with an infected person
 - » Laboratory exposure
- Zika may be spread through blood transfusion.
- No reports of infants getting Zika through breastfeeding.




How does Zika affect people?

- Many people with Zika will not have symptoms or will only have mild symptoms.
- Symptoms last several days to a week.
- People usually don't get sick enough to go to the hospital.
- People very rarely die of Zika.



What are the symptoms?

- For people with symptoms, the most common symptoms of Zika are
 - » Fever
 - » Rash
 - » Headache
 - » Joint pain
 - » Conjunctivitis (red eyes)
 - » Muscle pain



ZIKA AND PREGNANCY

How can Zika affect pregnancies?

- Zika virus can pass from a pregnant woman to her fetus during pregnancy or around the time of birth.
- It is not known how often this happens.



How can Zika affect pregnancies?

- Infection during pregnancy can cause damage to the brain, microcephaly, and congenital Zika syndrome
- Linked to other problems, such as miscarriage, stillbirth, and birth defects
- No evidence that past infection will affect future pregnancies once the virus has cleared the body



How can Zika affect pregnancies?

- Congenital Zika syndrome
 - » Distinct pattern of birth defects in fetuses and infants of women infected during pregnancy
 - » Associated with 5 types of birth defects not seen or rarely seen with other infections during pregnancy
 - Severe microcephaly (small head size) resulting in a partially collapsed skull
 - Decreased brain tissue with brain damage
 - Damage to the back of the eye with a specific pattern of scarring and increased pigment
 - Limited range of joint motion, such as clubfoot
 - Too much muscle tone restricting body movement soon after birth



Assessing pregnant women for possible Zika exposure


- At each prenatal care visit, all pregnant women should be asked if they
 - » Traveled to or live in an area with risk of Zika
 - » Had sex without a condom with a partner who lives in or traveled to an area with risk of Zika



GUILLAIN- BARRÉ SYNDROME

Does Zika cause Guillain-Barré syndrome (GBS)?

- GBS is an uncommon sickness of the nervous system in which a person's own immune system damages the nerve cells, causing muscle weakness, and sometimes, paralysis.
- GBS is strongly associated with Zika but only a small proportion of people with recent Zika infection get GBS.
- CDC is continuing to investigate the link between GBS and Zika to learn more.



TESTING

How is Zika diagnosed?

- A doctor or other healthcare provider will ask about any recent travel and any signs and symptoms.
- A blood or urine test can confirm a Zika infection.



Who should be tested for Zika?

- Anyone who has or recently had Zika symptoms
 - » And lives in or traveled to any area with risk of Zika, or
 - » Had unprotected sex with a partner who lives in or traveled to any area with risk of Zika



Who should be tested for Zika?

- All pregnant women (regardless of symptoms) who
 - » Live in or recently traveled to an area with risk of Zika that has a CDC Zika travel notice, or
 - » Had unprotected sex with a partner who lives in or traveled to an area with risk of Zika that has a CDC Zika travel notice
- Pregnant women who live in or recently traveled to an area with risk of Zika but **without** a CDC Zika travel notice
 - » If they develop symptoms of Zika, or
 - » If their fetus has abnormalities on an ultrasound that may be related to Zika infection



Testing babies for Zika

CDC recommends laboratory testing for

- All infants born to mothers with laboratory evidence of Zika virus infection during pregnancy
- Infants who have abnormal clinical or neuroimaging findings suggestive of congenital Zika syndrome and a mother with a possible exposure to Zika virus, regardless of maternal Zika virus testing results



WHAT TO DO IF YOU GET INFECTED

How is Zika treated?

- There is no specific medicine or vaccine for Zika virus infection.
- Treat the symptoms
 - » Rest
 - » Drink fluids to prevent dehydration
 - » Do not take aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs)
 - » Take acetaminophen (Tylenol®) to reduce fever and pain



What to do if you have Zika

- Protect yourself from mosquito bites. During the first week of illness, Zika virus can be found in blood.
- The virus can be passed from an infected person to a mosquito through bites.
- An infected mosquito can spread the virus to other people.



SURVEILLANCE

Reporting of Zika in the United States

- Healthcare providers should report cases to their local, state, or territorial health department.
- State and territorial health departments are encouraged to report confirmed cases to CDC through ArboNET, the national surveillance system for arboviral diseases.
- Pregnant women with any lab evidence of possible Zika virus infection should be reported to the US Zika Pregnancy Registry.



For the most recent case counts, visit <https://www.cdc.gov/zika/geo/united-states.html>.

Zika Pregnancy Registries

US Zika Pregnancy Registry



Zika Active Pregnancy Surveillance System (Puerto Rico)



Proyecto Vigilancia de Embarazadas con Zika (Colombia)



US Zika Pregnancy Registry

- CDC established the US Zika Pregnancy Registry to collect information and learn more about pregnant women in the US with Zika and their infants.
- Data will be used to
 - » Update recommendations for clinical care
 - » Plan for services for pregnant women and families affected by Zika
 - » Improve prevention of Zika infection during pregnancy
- Zika Active Pregnancy Surveillance System is used in Puerto Rico.



<https://www.cdc.gov/zika/hc-providers/registry.html>

<https://www.cdc.gov/zika/public-health-partners/zapss.html>

PREVENTION

Protect from mosquito bites

Zika is primarily spread through the bite of an infected *Aedes aegypti* or *Ae. albopictus* mosquito. Take steps to protect yourself and others.




Control mosquitoes outside

- Here's what you can do to help control mosquitoes outside your home
 - » Once a week, empty and scrub, turn over, cover, or throw out items that hold water.
 - » Tightly cover water storage containers.
 - » Use larvicides to kill larvae in containers of water that cannot be emptied and will not be used for drinking.
 - » Use an outdoor insect spray made to kill mosquitoes in areas where they rest.
 - » If you have a septic tank, repair cracks or gaps.



Control mosquitoes inside

- Here's what you can do to help control mosquitoes inside your home:
 - » Use window and door screens.
 - » Use air conditioning when possible.
 - » Once a week, empty, scrub, turn over, or throw out items that hold water.
 - » If you have mosquitoes inside your home, use an indoor insect fogger or indoor insect spray.
 - When using insecticides, always follow label directions.



Wear insect repellent

- Use Environmental Protection Agency (EPA)-registered insect repellents.
 - » Use a repellent with one of the following active ingredients: DEET, picaridin, IR3535, or oil of lemon eucalyptus, para-menthane-diol, or 2-undecanone.
- Always follow the product label instructions.
- Do not spray repellent on the skin under clothing.
- If also using sunscreen, apply sunscreen before applying insect repellent.



Create a barrier between you and mosquitoes

- Cover up exposed skin
 - » Wear long-sleeved shirts and long pants.
- Treat clothing and gear
 - » Use permethrin* to treat clothing and gear or buy pre-treated items.
 - » See product information to learn how long the protection will last.
 - » Do not use permethrin products directly on skin.



* Permethrin is not effective in Puerto Rico.

Protect your family

- For babies and children
 - » Dress your child in clothing that covers arms and legs.
 - » For children older than 2 months, use insect repellent on exposed skin.
 - » Cover crib, stroller, and baby carrier with mosquito netting.



Protect your family

- Applying insect repellent on children
 - » Do not apply repellent onto hands, eyes, mouth, and cut or irritated skin.
 - » Adults: Spray onto your hands and then apply to a child's face.
 - » Do not use insect repellent on babies younger than 2 months old.
 - » Do not use products containing oil of lemon eucalyptus or para-menthane-diol on children younger than 3 years old.



PREVENTION

Preventing sexual transmission


About sexual transmission

- Zika can be passed through sex from a person who has Zika to his or her sex partners.
 - » Sex includes vaginal, anal, and oral sex and the sharing of sex toys.
 - » Zika can be passed through sex before symptoms start, during, and after symptoms end.
 - » It can be passed even if the infected person does not have symptoms at the time or never develops symptoms.
- Zika virus can stay in semen longer than in vaginal fluids, urine, and blood.



Protect your partner

- Not having sex eliminates the risk of getting Zika from sex.
- Condoms can reduce the chance of getting Zika from sex.
 - » Includes male and female condoms.
 - » Condoms should be used from start to finish, every time during vaginal, anal, and oral sex and the sharing of sex toys.



Protect your partner

- People with a partner who traveled to an area with risk of Zika can use condoms or not have sex
 - » If the traveler is female: For at least 8 weeks after return, or after start of symptoms or diagnosis
 - » If the traveler is male: For at least 6 months after return, or after start of symptoms or diagnosis
- People living in an area with risk of Zika can use condoms or not have sex.



During pregnancy

- Pregnant couples in which one or both partners live in or traveled to an area with risk of Zika should:
 - » Use condoms from start to finish every time they have sex (oral, vaginal, or anal) or not have sex during the pregnancy.
 - » Not share sex toys during the pregnancy.



If you're thinking about having a baby

Exposure from travel to or sex without a condom with someone who lives in or traveled to an area **with a CDC Zika travel notice**

Women

Wait at least 8 weeks after symptoms start or last possible exposure before trying to get pregnant.

Men

Wait at least 6 months after symptoms start or last possible exposure before trying to conceive with your partner.



If you're thinking about having a baby

Exposure from travel to or sex without a condom with someone who lives in or traveled to an area with risk of Zika but **without a CDC Zika travel notice**

Women

If no symptoms, talk with a healthcare provider about plans for pregnancy.

Wait at least 8 weeks after symptoms start before trying to get pregnant.

Men

If no symptoms, talk with a healthcare provider about plans for pregnancy.

Wait at least 6 months after symptoms start before trying to get pregnant.



If you're thinking about having a baby

People living in or frequently traveling to any area with risk of Zika

Women

If no symptoms, talk with a healthcare provider about plans for pregnancy.

Wait at least 8 weeks after symptoms start before trying to get pregnant.

Men

If no symptoms, talk with a healthcare provider about plans for pregnancy.

Wait at least 8 weeks after symptoms start before trying to get pregnant.



PREVENTION

Traveling

Travel guidance for pregnant women

- If you are pregnant, do not travel to areas with risk of Zika.
- If you must travel, talk to your doctor or other healthcare provider and strictly follow steps to prevent mosquito bites during the trip.



Protect yourself while traveling

- If you travel to an area with risk of Zika
 - » Strictly follow steps to prevent mosquito bites.
 - » Use condoms or do not have sex during the trip.



Protect yourself while traveling

- Stay in places with air conditioning and with window and door screens.
- Use a bed net if air conditioned or screened rooms are not available or if sleeping outdoors.



Protect yourself and others after travel

- Even if they do not feel sick, travelers returning from an area with risk of Zika should take steps to prevent mosquito bites for 3 weeks so they do not spread Zika to uninfected mosquitoes.



Do your homework before traveling

See the latest travel notices at:

wwwnc.cdc.gov/travel/page/zika-travel-information



WHAT CDC IS DOING

What is CDC doing?

- Activated Emergency Operations Center (EOC) to level 1
- Alerting healthcare providers and the public about Zika
- Posting travel guidance
- Monitoring infections among pregnant women to identify the long-term consequences of congenital Zika infection.
- Working with clinical experts and organizations to update guidance
- Researching factors that might affect birth defects in fetuses and babies, including the timing of Zika infection during pregnancy.
- Improving laboratory testing for Zika and providing state, tribal, local, and territorial health laboratories with diagnostic tests.





CDC is working with partners to

- Monitor and report cases of Zika.
- Conduct studies to learn more about the potential link between Zika and Guillain-Barré syndrome.
- Create action plans for state and local health officials to improve Zika preparedness.
- Publish and disseminate guidelines to inform testing and treatment of people with suspected or confirmed Zika.
- Working with partners to better understand the risk and spectrum of birth defects from Zika infection during pregnancy and risks for sexual transmission.



Zika in the United States

- Local mosquito-borne spread of Zika virus was identified in Miami-Dade County, Florida, and Brownsville, Texas.
 - » Pregnant women should consider postponing travel to Brownsville, Texas (currently a yellow area).
 - » CDC lifted the yellow area designation for Miami-Dade County on June 2, 2017.



Brownsville, Texas

Miami-Dade County, Florida

CDC'S Response to Zika



For more information, contact CDC
1-800-CDC-INFO (232-4636)
TTY: 1-888-232-6348 www.cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention