Weekly Report

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE PUBLIC HEALTH SERVICE

Prepared by the

COMMUNICABLE DISEASE CENTER

634-5131

For release November 1, 1963

ATLANTA, GEORGIA 30333

Vol. 12. No. 43

PROVISIONAL INFORMATION ON SELECTED NOTIFIABLE DISEASES IN THE UNITED STATES AND ON DEATHS IN SELECTED CITIES FOR WEEK ENDED OCTOBER 26, 1963

POLIOMYELITIS – Ten cases of poliomyelitis, all paralytic, were reported for the week ending October 26. Two cases each were reported from Connecticut, New Jersey, and Pennsylvania. Single cases were reported from Illinois, Wisconsin, South Carolina, and Florida.

The 10 cases bring the total thus far in 1963 to 352 cases (301 of which have been paralytic). This total figure remains well under one-half the number reported last year. Nineteen hundred and sixty-two represented the the fewest cases of poliomyelitis ever recorded in the United States.

POLIOMYELITIS (CUMULATED WEEKLY) 1ST THROUGH 43RD WEEK

	1963	1962	1961	1960	1959
Paralytic	301	574	741	1932	4908
Total	352	724	1142	2798	7491
POLIOMYEL	ITIS (SIX	WEEK TOTALS	38TH	THROUGH 43RD	WEEK
Paralytic	75	158	224	620	1453
Tatal	90	188	362	890	2017

EPIDEMIOLOGICAL REPORTS

Botulism - Conned Liver Poste - Conado and U.S.A.—
Three cases of type B botulism, including one death, all related to ingestion of a commercially canned liver paste, were reported from Montreal, Canada, for the week ending October 26. All victims had consumed Paragon label canned liver paste, processed by Delicious Food Products, Ltd., Montreal.

Shipments of this same product were made to the United States, to distributors in Elmhurst, New York, and Union, New Jersey. The New York shipment bore the Paragon label, while the New Jersey shipment carried the Epicure Club label. From these two distributors, shipments were made throughout the United States. The Food and Drug Administration and State and local health authorities have now accounted for most cans in commercial channels.

Table 1. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES

(Cumulative totals include revised and delayed reports through previous week)

		43rd Wee	k		Cumulative	
Disease	Ended	Ended	Median	Fir	rst 43 wee	ks
Discuse	October 26, 1963	October 27, 1962	1958 - 1962	1963	1962	Median 1958 - 1962
DECEMBER 1991 OF THE RESERVE OF THE	1703	1701		to almost two 2	THE PERSON OF	The second second
Aseptic meningitis	49	64		1,538	2,161	
rucellosis	8	6	9	311	343	616
aphuleria	9	18	24	215	366	557
incephalitis, infectious	26	35	49	1,307	1,578	1,578
epatitis, infectious and serum	841	970	863	35,923	45,517	31,455
ICASIPE	1,207	1,708	1,536	363,931	449,640	402,004
iciiIIIgococcal intectione	51	32	36	1,990	1,777	1,879
orioniyentis, total	10	22	118	352	724	2,805
raralviic	10	20	81	301	574	1,924
Nonparalytic.	-	1	19	35	107	590
Ullsbecified		and I display to	18	16	43	291
treptococcal sore throat		W. D. CO.	ALL MANY COLUMN			100
and Scarler forces	5,146	5,019		279,132	259,004	fir he pay hade.
Ctanus		8		225	238	
	6	2		245	241	100 250 100
	5	10	25	451	520	701
ipinus Lever, fick-horne.	al ambientos s			12772		
(Rocky Mountain spotted)	-	4	odi seed lints	170	209	Andreas de la
abies in Animals	54	44	45	3,137	3,121	3,121

Table 2. NOTIFIABLE DISEASES OF LOW FREQUENCY

		10. 1
Anthrax: Botulism:	Cum. 4 Psittacosis:	67
Malaria: Colo 1	34 Rabies in Man: 83 Smallpox:	-
- ague:	- Typhus, murine:	24

No cases of botulism, related to this product, have been reported in the United States, as of noon Wednesday, October 30.

In the Montreal outbreak, 2 of the cases were brothers, who made sandwiches from a can of the liver paste. The elder, a 17-year-old, consumed 3 sandwiches while his 16-year-old brother had but a few bites of another sandwich for supper on October 19.

The following day, the elder brother, who consumed the greater quantity, experienced numbness of his mouth, dizziness, diplopia, urinary retention, and weakness to the point that he was unable to walk. He retired for the evening and was found dead in bed the next morning, about 36 hours following the meal.

Early the morning of his brother's death, the younger victim began experiencing diarrhea, dizziness, and diplopia. He was immediately hospitalized and treated with bivalent antitoxin. He has since recovered uneventfully.

The can providing the liver paste for their sandwiches had been discarded and could not be located. However, Clostridium botulinum type B was demonstrated in remaining portions of the sandwiches.

Following Canadian publicity of the outbreak, it was learned that a 17-year-old male had been treated in Montreal with bivalent antitoxin September 2 for a case of diagnosed botulism. This patient had experienced headache, nausea, vomiting, dysphonia, diplopia and respiratory weakness. He was placed in a respirator. He recovered. The vehicle concerned was originally thought to be home preserved strawberry jam; subsequently, Paragon brand liver paste has been implicated.

In addition to the above cases, approximately 25 to 30 others were treated with antitoxin as suspect cases of botulism. They had consumed the same product and had experienced gastrointestinal, but no neurological, symptoms. These have not been officially reported as botulism cases.

No cases of botulism were reported from elsewhere in Canada.

The commercial product was removed from the shelves of the stores in the 6 provinces in which it was offered for sale: Newfoundland, New Brunswick, Nova Scotia, Prince Edward Island, Quebec, and Ontario. Laboratory tests on cans randomly selected from the shelves are in progress.

The processing plant is being investigated by Canadian Food and Drug officials. The plant has had no previous incidents of food poisoning and has allowed voluntary Canadian inspection annually for the past 2 decades.

The liver paste was made from pork liver.

(Reported by Dr. E.W.R. Best, Chief, Epidemiology Division, Department of National Health and Welfare, Ottawa,

Canada and the U.S. Food and Drug Administration.)

Editor's Note: The liver paste is the third botulism incident reported in North America this year to involve a commercially processed product. The first was the canned tuna fish (see MMWR, Vol. 12, p. 95) and the second was the smoked whitefish chubs (see MMWR, Vol. 12, pp. 329, 337 and 345).

Botulism - Smoked Fish Products - United States

The Food & Drug Administration, acting on the basis of its own findings and on the advice of a special advisory committee on botulism, has recommended the destruction of all products currently on the market which are made from Great Lakes fish or processed in establishments in the Great Lakes region. The FDA made two exceptions, with the provision that the criteria can be established with with certainty: "(1) That the fish was heated immediately after it was packaged in a manner that would assure a temperature of at least 180°F. for at least 30 minutes in the coldest part of the fish. Such fish should be marked 'Perishable – Keep Under Refrigeration', or, (2) that the fish was frozen immediately after packaging and has since been maintained continuously in a frozen condition."

The FDA based its decision on the 3 known outbreaks of Type E botulism which have occurred during the past 3 years from fresh water fish obtained from the Great Lakes, (See MMWR, Vol. 12, pp. 329, 337, and 345), and on subsequent laboratory tests which it performed on such fish under special conditions.

The FDA release stated further: "These outbreaks, including recent findings from the Tennessee and Michigan outbreaks, indicate that Clostridium botulinum type E spores are present in the incoming fish, the smoking establishments, or both. Bulk and packaged smoked fish packed by three Michigan firms, held deliberately by FDA laboratories without refrigeration for several days, showed evidence of type E botulinus toxin. . . .

"The organism is capable of germination, growth, and toxin formation at temperatures within the normal refrigeration zone, as low as 38°F., within 31-45 days in one case. Growth and toxin formation has not been reported at 36°F and below for longer periods. Existing conditions for shipping and handling packaged and bulk smoked fish products do not insure against growth in toxin formation if the organism is present. Freezing of products immediately after processing, and maintaining in the frozen condition until delivered to the consumers, will prevent development of the toxin."

Dengue-Like Illness - Puerto Rico

During the past week, 543 cases of dengue-like illness were officially reported to the Puerto Rico Department of Health, bringing the 1963 total to 16,771 through October 28.

Intensive day-to-day surveillance continues in Barriada Frailes Llanos and Barriada Marrero in the townshop of Guaynabo. Attack rates in these population groups are as follows:

	DENGUE-LIKE ILLNESS By week of onset*
number of cases	
257	
20-	BARRIADA MARRERO
15-	
10-	-111 ₀ -1
5"	
0	3 10 17 24 31 7 14 21 28 5 12 19 26
30-	
25 -	BARRIADA FRAILES LLANOS
20-	
15-	
10-	
5-	
week ending	3 10 17 24 31 7 14 21 28 5 12 19 26 AUGUST SEPTEMBER OCTOBER

Period	Area	Population	Dengue-like Illness	Attack Rate(%)
August 1 -	13/12/12/14	ST 12.1	LEL LOCK	Garage .
October 29	Guaynabo Township*	898	291	32.4
August 1 -	Barriada Frailes			art out
October 23	Llanos	488	173	35.5
August 1 -				
October 23	Barriada Marrero	209	80	38.3

^{*}From morbidity survey conducted on October 29.

The epidemic curves for Barriada Frailes Llanos and Barriada Marrero are shown (left). In one community the incidence of cases dropped abruptly following a peak during the week ending September 28, whereas new cases have continued to occur in the other area.

In addition, the total attack rate for females is significantly higher than the attack rate for males in Barriada Frailes Llanos; there is no significant difference in Barriada Marrero.

(Reported by Victor Gonzales, M.D., Director, Bureau of Health, Puerto Rico Department of Health, and a team from the Communicable Disease Center.)

(Continued on page 364)


INFANT DEATHS IN 108 CITIES

The weekly average number of infant deaths in 108 cities for the four-week period ending October 26 was 759 as compared with an expected weekly average of 764.

TOTAL DEATHS UNDER ONE YEAR OF AGE IN 108 CITIES

200		WEEK EN	4 Week	Weekly		
lates:	10/5	10/12	10/19	10/26	Total	Average
Observed	761	789	717	770	3,037	759
Expected	761	763	765	767	3,056	764
Excess	0	26	- 48	3	– 19	– 5

DEATHS UNDER ONE YEAR OF AGE IN IOS U.S. CITIES
Average number per week by four-week periods



(See table, page 363)

Table 3. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES FOR WEEKS ENDED

OCTOBER 26, 1963 AND OCTOBER 27, 1962

To the season of the	Polic	omyelitis			Po1	iomyeliti			Poliomy nonpara	velitis, ulytic	Aseptic Meningitis	
Area	43rd	week		lative 3 weeks	43rd	week	1 =	lative 43 weeks	/, 3 m d	week	43rd	week
196 1 40	1963	1962	1963	1962	1963	1962	1963	1962	1963	1962	1963	1962
UNITED STATES	10	 	252									
UNITED STATES		22	352	724	10	20	301	574	110E V	1	49	64
EW ENGLAND	2		7	7	2	_	7	7 -	_	1 :	2	1
New Hampshire			-			_	1 -	[_	1 -		1 -
Vermont	5	-	1	THE 2	8 -	-	1	-			1	-
Massachusetts	-	-	3	6	- A	-	3	6				1
Rhode Island	2		2	1	2		_ 2	1	-	-	1 -	-
IDDLE ATLANTIC	4	1	112	74	4	1		5.0]		_
New York	W 10	i	8	57	-	1	89 5	53 39		-	8 8	3
New Jersey	2	-	3	6	2	-	3	6	9 -	-	-	1 -
Pennsylvania	2	-	101	11	2	-	81	8	B 1-	-	-	2
AST NORTH CENTRAL	2	5	51	101	2	4	41	75	Luc-	_	8	5
Ohio		1	8	18	9.	1	4	16	ALE-III	-	. 2	2
IndianaIllinois	1	4	17	18 45	1	3	3 16	14 31		-	4	1
Michigan	er leg 4		15	14	100	-	15	11			2	2
Wisconsin	1	4	7	6	1	200	3	3		1	-	-
EST NORTH CENTRAL	-	2	5	38	-	2	5	27		100	1	5
Minnesota	The su	Jan Ber	4	7	-	-	4	7		-	1	5
Iowa Missouri		1 -	1000	7 10		1 -	-	3	100	- "	- us-	
North Dakota	E136311	1	11 1-1-11	5		1	- 5	5	100	1 2		1 1
South Dakota	100	-		1	-	1 2	11 1520	1	120		120	TA I
Nebraska	0.00		1	8	GII	-	1	8		-	-	_
Kansas	Description	- 1	had an		-	-	4.7	-	The same		-	-
OUTH ATLANTIC	2		62	56	2	-	53	49		-	6	4
Delaware	W-84	L	1	- 1	-	-	1			70.00	1	-
Maryland	× 1,444	19/01/1	3	1 2	//II	-	1	1		-	-	-
Virginia	1000	With the	18	8		20 PE 193	13	1 8	all to the	17000	ī	1
West Virginia	-		3	5	-	- 11	3	5	10 - DE	-	i	1
North Carolina		-	3	10	-	4.7-6	3	8		O. R. S. H.	P=3,	-
South Carolina	1	1	7 19	6 14	1	-	6	6	-	-	-	-
Florida	1		8	10	1		18 8	13	-	-	3	2
AST SOUTH CENTRAL		1	64	67		1	60	55		_	2	4
Kentucky	-		1	26	-	_	1	20		-	1	1
Tennessee	-	1	8	10	-	1	8	5	_		1	-
Alabama Mississippi	100	DE AUG	48 7	22	-		44	22 8		- III	160-	3
EST SOUTH CENTRAL	100	9 2	25 5	294 14	-	8	24	229		1		7
Louisiana		2	14	24	-	2	14	14 21	40.5	700	HTA.	
Oklahoma		3	-	20	-	3	17	15	-	2		2
Texas	-	2	6	236	-	2	6-	179	-	-	il do an	5
OUNTAIN	-	1	5	15	_	1	4	11	100		3	3
Montana	10- 00	N 11-151	W	4	_	-	-	3			1	-
Idaho	W- L	- 510	1	2	-	1111	1	1				-
Wyoming Colorado	Ben	= = 1	3 57	2	-	-	-	1	-	-		-
New Mexico	nkor		1	2	1	-	200	1		11.	1	3
Arizona	10.5	3.14	3	3		-	3	3	-		1	
Utah Nevada		1	10 2	2	-	1		2	-		-	:
Paragraph 1076 Balling	LITTLE STATE		THE R	History			- 3107	•	-			
PACIFIC		3	21	72		3	18	68	-		19	32
Washington		3	2	5		3	2	5		-	2	1
Oregon	7		2 17	5		201	1	5	575	115		21
Alaska			-	62			15	58	-		17	31
Hawaii	-	-	_ - _	-	=	-		Un 200	447	12.12	-	-
							+					+

Table 3. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES FOR WEEKS ENDED

OCTOBER 26, 1963 AND OCTOBER 27, 1962 - (Continued)

Went of resident	Brucel	losis	Diphth	eria	Encepha infect		infe	Hepati ectious a		m	Mea	sles
Area		Cumu- lative		Cumu- lative			1-11-		week	7		ALDE.
	43rd week	43 weeks	43rd week		s 43rd	week	Under 20 yr.	20 & over	To	tal	43rd	week
	1963	1963	1963	1963	1963	1962	1963	1963	1963	1962	1963	1962
UNITED STATES	8	311	9	215	26	35	432	367	841	970	1,207	1,70
NEW ENGLAND	100	1	_	8			65	44	112	91	88	7
Maine	7 32 7			-	-	-	25	11	36	37	4	315
New Hampshire	-		-	-	-		14	6	23	15	v-1	
Vermont	-	1	-	-	-	-	7	6	13	2	9	100
Massachusetts	-		-	6 2			14	5	19	32	19	
Rhode Island			-	-			5	14	19	1	17 39	
	1	8		22	1	7	80	7.0	1.50	176	000	
New York	1	4		22 13	1	4	45	73 49	153 94	176 90	233	1
New Jersey.	_	i	-	4	1		12	9	21	33	56	
Pennsylvania	10-	3	-	5	-	3	23	15	38	53	59	
EAST NORTH CENTRAL	3	39	1	24	3	3	71	51	126	148	210	5
Ohio	-	-		1	1	2	10	14	24	40	40	
Indiana	-	6	1	6		-	7	4	11	18	24	
Illinois	2	20		12	1	-	14	9	24	33	34	
Michigan	1	5 8	A 11	3 2	1	1 -	40	24	64	50 7	56 56	1 2
									- 3	146	27,544	0111
VEST NORTH CENTRAL	2	166 8	4 AT 1	39 15	3	1	23	13	43 5	51 18	36	lie a
Iowa	2	124	- 1	1			13	7	21	6	18	
Missouri	I	12		î	0.00	0 2	4	2	6	12	4	
North Dakota	-	-	N	2		-	\$1	1	1	4	8	-
South Dakota	-	10		12		6 DF=-	19	91 -	-	1	1	1.00
Nebraska		6	3 7 11	8	-		:	1	3	6	-	
Kansas	+1 (* - 1	6	7.5	2.1	3	and the	5	-	7	4	NN	
SOUTH ATLANTIC	1	19	1	50	6	2	54	42	97	106	144	
Delaware	41 07 1			8. 1	-	100	9	3 7	3 16	10	-	
Maryland	41 35 4 1	4 4	9 E T	1	-	100	9	1	10	12	19	-7-5
Virginia	1	9	11	< 20		2000	6	4	10	11	48	
West Virginia	J 1- J			1		1	13	8	21	20	57	10/91
North Carolina	1 5 71	4		2	-	1	20	10	30	34	5	100
South Carolina	21 3-1	3	1 10	17	-	-	1 2	-	1	4	4	-
GeorgiaFlorida	1 7 1	3	1	18 11	6		3	3 6	5 10	24	10	1000
	11 14 1	10		10	3	2	/0	.,		0.5	70	71.0
AST SOUTH CENTRAL	the set of	13 3	- [18	3	2	40	21	61	95 27	78 26	
Tennessee		6		3	3	- 1	16	10	26	32	52	55.4
Alabama	1 42	4	- 1	12		9.2	11	3	14	15		111/1-10
Mississippi	-	(- J	1	3	2. 11	- 1	4	2	6	21	1975	
EST SOUTH CENTRAL		35	7	45		8	23	25	53	50	64	
Arkansas	1 3-	8	-	2	-	1	1	4	5	7	3	
Louisiana	-	8	6	27	-	-	9	8	19	5		
Oklahoma. Texas	1 1	14	1	6 10		6	13	11	27	35	60	- 4
	104		7.2			1				7-1-1	HOUSE	100
Montana		9	1	5	2	1	11 2	9	41	50 6	87 55	2
Idaho			- 1						14	3	18	1
Wyoming	- 1	1	_	-	-		A45 1	1	1	4	7711-	
Colorado	-		-	- 3	-	I	2	1	6	20	2	
New Mexico	J (- 1)		- 9	2	1	-	5	3	8	4	NN	
Arizona	01 5 1	3	-	-	1	-	-	-	4	7	6	
Utah Nevada	3 111	5 -		4 4			2 -	2 -	4	3	6	- 60
		1.0			N				1 0			1 640
ACIFIC	1	21		4	8	12	65	89	155	203	267	3
Washington		3			1	2	10	16	26	24	79	1
Oregon	1	17		4	7	10	3 49	9 61	13 110	15 113	101	1
Alaska				-	4	10	2	3	5	43	52	15
Hawaii.		1	- 11	-		-	ī		í	8	12	
and the second s					1		1					

Morbidity and Mortality Weekly Report

Table 3. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES FOR WEEKS ENDED

OCTOBER 26, 1963 AND OCTOBER 27, 1962 - (Continued)

- cal (4.0)	Mening Infect	ococcal	Strepto Sore Th Scarlet	roat &	Tetanus	Tickborne Typhus (Rcky Mt.	Tularemia	Typhoi	d Fever	Rabie	s in Anir	nals
Area	43rd wk.	Cumu- lative 43 weeks	43rd	week	43rd wk.	Spotted)	43rd Wk.	43rd wk.	Cumu- lative 43 weeks	43rd	week	Cumu- lative 43 weeks
GO DELL'ES	1963	1963	1963	1962	1963	1963	1963	1963	1963	1963	1962	1963
UNITED STATES	51	1,990	5,146	5,019	4	Va la	6	5	451	54	44	3,137
et 1145 11	Table 1	as had	100	34 E	viii)							and of
NEW ENGLAND Maine	5 1	124 18	355 20	470 126			1 1 1		12	1	1 1	31
New Hampshire	4-75	4		3		-	I - b	_	-	1	_	15
Vermont	1	5	16	12	-	à -	- 1	-	- 1		-	13
Massachusetts	2	58	60	56		= -	- 1	/a []	6	0.00	- 15-11	1
Rhode Island	1	11 28	24 235	50 223			7 1		3	110011		
ATTORIE ATTIANTES	4	270	202	213	4		10 4		0.0	-		100
MIDDLE ATLANTIC New York	1	272 126	202 149	107	4			7 1	83 43	1	6	100 75
New Jersey	11/12	41	21	41		1 -	4 - do	-	5		-	-
Pennsylvania	3	105	32	65	-	-	- 1	-	35	-	2	25
EAST NORTH CENTRAL	8	309	457	370	-		2 - 1	1	53	5	-	479
Ohio	111	80	56	51		-	6 - 5	1	21	2		282
Indiana	1.0	45	99	82	3: 15-1	PT -		75 -	7	1		44
Illinois Michigan	5	60 92	63 160	58 94	5.11		1 : 1	1 1	10	1		67 44
Wisconsin	3	32	79	85	-	- 7	1 - 1	-	5	1		42
WEST NORTH CENTRAL	2	122	223	151	1		1 . 4.	1	26	19	15	819
Minnesota	1	24	8	6	1	2 - .	3 - 10		3	10	6	213
Iowa	-	7	50	64	2:	2 -	F - II	140-24	3	5	7	298
Missouri.	1	35 13	19 42	7 29			-	1	16	2	2	138
North Dakota	100	7	5	29				LOUI TO	1	9-4,0,7 1.1	1100	32 89
Nebraska.	-	25		5		-			1	1	11000	29
Kansas	= 1-	11	99	40	-	-	-	-	2	1	-	20
SOUTH ATLANTIC	6	363	545	439	3	K 1- 1-	1	2	62	11	4	450
Delaware	-	4	4	3	-	-		- 1	4	Line .		1
Maryland	-	51	29	12	3		4 - 1		11		na E	1
Dist. of Columbia Virginia	2	6 82	1 160	107		10 10		1 - 1	8	8	1	172
West Virginia	-	19	185	87	1 11			16 Tel	7	-	2	110
North Carolina	3	67	48	25	1	6 le 1 3	B - B	2	9	Paris I	5-17-0	14
South Carolina	-1-3-	20	42	38	-	3/17	4 - 4	(C	4	100 ACO		9
Georgia. Florida.	1	30 84	2 74	161	2		1	E -	17	3	1	73
THE	38	724 U	. ALC: 1		e 100	4X-10		pt- II			pulatica o	NEW TEX
EAST SOUTH CENTRAL	1	140	954	823	-	T 12 12 1		15-	63	4	4	243
Kentucky Tennessee	1	31 64	40 864	68 673	5	21.7			13 24	1 3	1 2	113 112
Alabama		23	32	13	-	3 - 13		10 201	11		1	18
Mississippi	II	22	18	69		-		1000-11	15	-	-	-
WEST SOUTH CENTRAL.	5	178	536	422		-	3	1	83	9	5	584
Arkansas	10	11	100			3E - 17	2	16.5	32	2	1	72
Louisiana	3	74	14	2		8 1- 12	1		25	1	2	44
Oklahoma. Texas	2	31 62	16 506	420		St Lin	1	1	6 20	2 4	1	53 415
DESCRIPTION OF STREET	314	70	0/0		x 4	3 1 1	3.		1.09	li-sam	(income)	300
MOUNTAIN	7	70	948 76	1,192 53	4 7				16	THE STATE OF	3	124
Idaho.		6	60	105	-	STATE OF THE				1 (150,000)		
Wyoming	3	7	81	72	3 - 17	1 - E	3		-			insta
Colorado	1	20	216	370	-			12 -31	6	arrain.	2	16
New Mexico	1	11	309 141	366 81	1 TE			(E. 7)	3 7			37
Utah	2	16	65	143			1. 1	1.4			1 -	57 3
Nevada	The second	3		2			- 7	T	-	-	-	11
PACIFIC	13	412	926	939		C.L.III	2	10 4	53	4	6	307
Washington	2	34	247	325	3 . 17	3 E T	4-1	12 - 3	3		Correct.	hei -
Oregon	2	31	12	23	2 Tab		- 1	7.12	2	1		12
California.	9	326 12	492 37	575 14	TE- IT		2		45 1	3	6	286
Hawaii		9	138	2			-	145	2	1000	1000	9
			2	8					-			

Table 4 (C). TOTAL DEATHS UNDER 1 YEAR OF AGE IN REPORTING CITIES

(Tables 4(A), 4(B), 4(C), and 4(D) will be published in sequence covering a four-week period.)

Area	550 b	For week	s ending	olum.	nelson-true on no because.		For weeks ending				
Area	10/5	10/12	10/19	10/26	Area	10/5	10/12	10/19	10/26		
EW ENGLAND:		1			SOUTH ATLANTIC:	3/17/19	1808/5	M. Y.	1579/		
Boston, Mass	13	18	12	13	Atlanta, Ga	8	17	9	8		
Bridgeport, Conn	7112211	4	5	3	Baltimore, Md	26	19	16	36		
Cambridge, Mass	1	1	THE P.		Charlotte, N.C.	4	1	3	4		
Fall River, Mass	3		2	1	Jacksonville, Fla	6	2	4	9		
Hartford, Conn	3	5	5	4	Miami, Fla	4	7	4	men.		
Lowell, Mass	2	3	2	1 - 1	Norfolk, Va	4	9	4			
Lynn, Mass	11 - 0-	1	1	3 1	Richmond, Va	12	8	9			
New Bedford, Mass New Haven, Conn	2	2	2	1	Savannah, Ga	2	2	3	o emili		
Providence, R.I	7	2	5	6	St. Petersburg, Fla Tampa, Fla	5 7	1 3	2 2	0.5		
Somerville, Mass.	A laborate	1	ī	-	Washington, D.C.	28	14	8	1		
Springfield, Mass	-	3	4	3	Wilmington, Del	6	1	6	1		
Waterbury, Conn	3	2	2	1	Translating Control of the Control o			it min			
Worcester, Mass	2	4	-	2	EAST SOUTH CENTRAL:		-				
					Birmingham, Ala	6	12	7	- SV		
IDDLE ATLANTIC:					Chattanooga, Tenn	13.U=14	2	6	2007		
Albany, N.Y	2	1	1	2	Knoxville, Tenn	3	3	4	1000		
Allentown, Pa	2	2	1	2	Louisville, Ky	3	7	18	9		
Buffalo, N.Y	12	9	6	8	Memphis, Tenn	15	8	9	1		
Camden, N.J.	3	7	3	8	Mobile, Ala	5	7	5			
Elizabeth, N.J	2	7	-	2	Montgomery, Ala	2	3	3			
Erie, Pa.	4	4	2	1	Nashville, Tenn.	9	9	3			
Jersey City, N.J.	6	3 6	7 7	1 7	SI DICTOR HE STREET BY BAR		1-60 P	PASSES.	594		
New York City, N.Y	87	88	85	117	WEST SOUTH CENTRAL:	and the second		,			
Paterson, N.J.	1	2	4	3	Austin, Tex	5	6	3 2			
Philadelphia, Pa	30	27	11	39	Baton Rouge, La Corpus Christi, Tex	4	5	2	100		
Pittsburgh, Pa.	12	18	11	7	Dallas, Tex	16	13	7			
Reading, Pa.	2	_	3	-	El Paso, Tex	3	5	7			
Rochester, N.Y	5	8	6	6	Fort Worth, Tex	6	4	5	PUR		
Schenectady, N.Y	- 100	-	1	- 1	Houston, Tex	17	23	19	1.		
Scranton, Pa	5 - 1	20	7716	3	Little Rock, Ark	10	19	5			
Syracuse, N.Y	1	2	6	5	New Orleans, La	13	16	19	1		
Trenton, N.J	7	2	5	4	Oklahoma City, Okla	3	4	6	1		
Utica, N.Y	10.65	1	2	By Terral	San Antonio, Tex	12	5	4			
Yonkers, N.Y		-	1	3	Shreveport, La	3	5	8			
AST MODELL OF MEDIA					Tulsa, Okla	2	4	6	100		
AST NORTH CENTRAL:	4	6	2	6	MOUNTA TNI-						
Akron, Ohio	4	2	2	2*	MOUNTAIN:	2	5	2			
Chicago, Ill	49	48	47	45	Colorado Springs, Colo			1	hipe		
Cincinnati, Ohio	6	13	6	9	Denver, Colo	21	8	2	wil.		
Cleveland, Ohio	4	21	24	15	Ogden, Utah		2	2	200		
Columbus, Ohio	6	9	16	12	Phoenix, Ariz	11	4	11	1:		
Dayton, Ohio	2	5	6	7	Pueblo, Colo	1	2	1	and a		
Detroit, Mich	25	30	19	11	Salt Lake City, Utah	3	4	3			
Evansville, Ind	2	2	1	1	Tucson, Ariz	6	2	3	1		
Flint, Mich	7	2	5	3	Charles Or marketing spiritely		entain s	sees at ID	N 153		
Fort Wayne, Ind.	1	3	-	1	PACIFIC:				* 193		
Gary, Ind		3	2	2	Berkeley, Calif	11111	D 16-12	2			
Grand Rapids, Mich	2	3	1	5	Fresno, Calif	4	6	4 ,	100		
Indianapolis, Ind	9	14	9	11	Glendale, Calif	1	1	MI COL			
Madison, Wis	3	4	2	2 5	Honolulu, Hawaii	1	8	6	124		
Milwaukee, Wis	5 2	5	4 2	5 2	Long Beach, Calif	41	7 35	32	2		
Peoria, Ill	2	3		5	Los Angeles, Calif	9	33	32 4	2		
Rockford, Ill	2	1	3	2	Oakland, Calif	-	2	1			
South Bend, Ind Toledo, Ohio	6	6	3	6	Pasadena, Calif	5	5	1	1		
Youngstown, Ohio	1	2	1	4	Portland, Oreg Sacramento, Calif	2	7	6			
and a second	HIA.		(FOLES)		San Diego, Calif	10	7	8			
ST NORTH CENTRAL:					San Francisco, Calif	2	6	14	1		
Des Moines, Iowa	3	4	4	1	San Jose, Calif		-	4			
Duluth, Minn	1	1			Seattle, Wash	8	3	2	1:		
Kansas City, Kans	7	3	2	4	Spokane, Wash	2	2	2			
Kansas City, Mo	9	4	3	9	Tacoma, Wash		3	2			
Lincoln, Nebr.		1	-	2							
Minneapolis, Minn	7	7	9	8	San Juan, P.R.	5	2	5	4		
Omaha, Nebr	5	13	5	2							
St. Louis, Mo	16	7	21	12							
St. Paul, Minn.	3	2	4	8	OCurrent Week Mortality for	- 108 Se	lected Cf	ties			

^{*}Estimate - based on average percent of divisional total.
Totals for previous weeks include reported correction

NOTE: All deaths by place of occurrence.

Bat Rabies - North Carolina

Since the first proved rabid bat was discovered in North Carolina in late July, 1963, approximately 100 residents of the State have been placed on an anti-rabies immunization regimen because of definite or probable exposure to bats. Most of the instances represent unprovoked attack by bats upon humans during the daytime hours.

In only 8 instances, however, have bats collected at the site of the attack been shown to be rabid by direct microscopic examination or by the fluorescent antibody technique.

The low incidence of proved bat rabies in the captured, peculiarly behaving bats suggests that another disease may be responsible in many cases. One possibility presently being considered is the effect of insecticides. Further investigation is in progress.

(Reported by Dr. Jacob Koomen, Assistant Secretary and State Health Director, North Carolina State Board of Health.)

Editor's Note: Since rabies virus is known to occur in salivary glands in animals without being present in their central nervous systems, it is imperative that all bona fide bat bites receive post exposure vaccine treatment unless the salivary glands, as well as brain tissues, are specifically examined and found to be negative.

INTERNATIONAL NOTES

One case of paralytic poliomyelitis was reported in Canada for the week ending October 19. This brings the Canadian total to 92 thus far in 1963.

This week's case was reported from Quebec, bringing that province's total to 85 so far in 1963.

For the comparable period in 1962, 52 cases of paralytic polio were reported from Quebec, whereas 87 cases were reported from the entire country. In 1961, for a comparable period, 99 cases of polio were reported in Quebec, and 165 cases throughout Canada.

(Reported in Weekly Report of Canadian Department of National Health and Welfare.)

In addition to the established procedures for reporting morbidity and mortality, the Communicable Disease Center welcomes accounts of interesting outbreaks or cases. Such accounts should be addressed to:

Lawrence K. Altman, M.D., Editor Morbidity and Mortality Weekly Report Communicable Disease Center Atlanta, Georgia 30333

> Library 81 CDC 7 61

HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
Communicable Disease Center
Atlanta, Georgia 30333

Official Business

MA I L

POSTAGE AND FEES PAID U. S. DEPARTMENT OF H. E. W.