This is an official CDC HEALTH ADVISORY

Distributed via the CDC Health Alert Network September 13, 2017, 1230 ET (12:30 PM ET) CDCHAN-00407

Rifampin/Penicillin-Resistant Strain of RB51 Brucella Contracted from Consumption of Raw Milk

Summary:

The Texas Department of State Health Services, with assistance from CDC, is investigating *Brucella* RB51 exposures and illnesses that may be connected to the purchase and consumption of raw (unpasteurized) milk from K-Bar Dairy in Paradise, Texas. Symptoms of brucellosis can include: fever, sweats, malaise, anorexia, headache, fatigue, muscle & joint pain, and potentially more serious complications (e.g., swelling of heart, liver, or spleen, neurologic symptoms).


Background

A person who drank raw milk from K-Bar Dairy in Paradise, Texas, has been hospitalized with brucellosis. Milk samples from the dairy have tested positive for a *Brucella* strain called RB51. People who consumed milk or milk products from this dairy from June 1, 2017, to August 7, 2017 are at an increased risk for brucellosis and should receive appropriate post-exposure prophylaxis (PEP). They are advised to consult with their health care providers regarding PEP care and possible diagnostic testing. Please note: the incubation period for brucellosis can range from five days to six months.

Recommendations

Brucella strain RB51 is resistant to rifampin and penicillin. A combination of doxycycline and trimethoprim/sulfamethoxazole for 21 days is the recommended first-line PEP regimen for RB51 exposure. There is no serological test available to detect RB51 infection. Blood culture is the recommended diagnostic test for exposed symptomatic individuals. When ordering blood cultures to diagnose brucellosis, please advise the laboratory that blood culture may grow *Brucella* and that appropriate laboratory¹ precautions should be observed. If brucellosis occurs despite prophylaxis, treatment regimens² should be selected based on antimicrobial susceptibility results. Please see the diagram below for information on developing an evaluation and treatment plan for exposed patients. Women who are pregnant or breastfeeding should consult with their health care provider.

Treatment Decision Tree for Patients Who Were Exposed to (Consumed) Raw Milk/Raw Milk Products from K-Bar Dairy June 1, 2017 - August 7, 2017



For More Information

Risks from Unpasteurized Dairy Products https://www.cdc.gov/brucellosis/exposure/unpasteurized-dairy-products.html

Exposure to RB51 through Raw Milk or Milk Products: How to Reduce Risk of Infection <u>https://www.cdc.gov/brucellosis/clinicians/rb51-raw-milk.html</u>

Symptoms of Brucellosis https://www.cdc.gov/brucellosis/symptoms/index.html

Brucellosis and Expecting Mothers https://www.cdc.gov/brucellosis/exposure/expecting-mothers.html

Raw Milk Questions and Answers https://www.cdc.gov/foodsafety/rawmilk/raw-milk-questions-and-answers.html

CDC-INFO https://www.cdc.gov/cdc-info/index.html or 1-800-232-4636

CDC Emergency Operations Center (EOC) 770-488-7100

Bacterial Special Pathogens Branch bspb@cdc.gov or 404-639-1711

References

¹Biosafety in Microbiological and Biomedical Laboratories <u>https://www.cdc.gov/biosafety/publications/bmbl5/bmbl.pdf</u>

²Treatment of Brucellosis (for alternative treatment guidance)

- Ariza J *et al.* 2007. Perspectives for the Treatment of Brucellosis in the 21st Century: The Ioannina Recommendations. PLoS Med. 4(12): e317. http://www.plosmedicine.org/article/info:doi/10.1371/journal.pmed.0040317
- Al-Tawfiq JA. 2008. Therapeutic options for human brucellosis. Expert Rev Anti Infect Ther. 6(1): 109-120. http://www.ncbi.nlm.nih.gov/pubmed/18251668
- Solera J. 2010. Update on brucellosis: therapeutic challenges. Intl J Antimicrob Agent. 36S, S18–S20. http://www.ncbi.nlm.nih.gov/pubmed/20692127

Brucellosis Reference Guide

https://www.cdc.gov/brucellosis/pdf/brucellosi-reference-guide.pdf

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages:

Health AlertRequires immediate action or attention; highest level of importanceHealth AdvisoryMay not require immediate action; provides important information for a specific incident or situationHealth UpdateUnlikely to require immediate action; provides updated information regarding an incident or situationHAN Info ServiceDoes not require immediate action; provides general public health information

##This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, HAN coordinators, and clinician organizations##