

Web Table 1. Predicted Differences^a in Body Mass Index, Waist Circumference, and Percent Body Fat From Ages 7–15 Years According to Tertiles (Low, Medium, and High^b) of Benzophenone-3 Concentrations Among Girls in the Breast Cancer and Environment Research Program, 2004–2015 (*n* = 1,017)

Age (Years) and Tertile Comparison	Predicted Difference in Body Mass Index		Predicted Difference in Waist Circumference, cm		Predicted Difference in Percent Body Fat, %	
	Difference	95% CI	Difference	95% CI	Difference	95% CI
7 years						
Medium to low	0.34	−0.18, 0.86	0.69	−0.87, 2.25	0.31	−1.13, 1.76
High to low	0.53	−0.01, 1.06	1.27	−0.33, 2.87	0.51	−0.97, 2.00
High to medium	0.19	−0.32, 0.70	0.58	−0.97, 2.12	0.20	−1.22, 1.62
8 years						
Medium to low	0.12	−0.40, 0.64	0.02	−1.39, 1.43	−0.12	−1.49, 1.25
High to low	0.29	−0.25, 0.82	0.34	−1.11, 1.79	−0.07	−1.48, 1.33
High to medium	0.17	−0.35, 0.68	0.32	−1.06, 1.71	0.04	−1.30, 1.39
9 years						
Medium to low	−0.04	−0.59, 0.51	−0.53	−1.96, 0.89	−0.45	−1.78, 0.87
High to low	0.09	−0.47, 0.66	−0.43	−1.90, 1.03	−0.53	−1.89, 0.83
High to medium	0.14	−0.40, 0.68	0.10	−1.30, 1.50	−0.08	−1.38, 1.22
10 years						
Medium to low	−0.16	−0.75, 0.43	−0.97	−2.50, 0.56	−0.69	−2.00, 0.62
High to low	−0.05	−0.66, 0.55	−1.06	−2.63, 0.51	−0.86	−2.20, 0.49
High to medium	0.10	−0.48, 0.68	−0.09	−1.59, 1.41	−0.16	−1.45, 1.12
11 years						
Medium to low	−0.22	−0.85, 0.42	−1.29	−2.94, 0.36	−0.84	−2.14, 0.47
High to low	−0.15	−0.80, 0.50	−1.54	−3.23, 0.16	−1.05	−2.39, 0.29
High to medium	0.06	−0.56, 0.69	−0.25	−1.87, 1.38	−0.21	−1.49, 1.07
12 years						
Medium to low	−0.22	−0.91, 0.46	−1.49	−3.26, 0.27	−0.89	−2.20, 0.42
High to low	−0.20	−0.91, 0.50	−1.86	−3.67, −0.05	−1.12	−2.46, 0.23
High to medium	0.02	−0.65, 0.69	−0.37	−2.10, 1.37	−0.23	−1.51, 1.06
13 years						
Medium to low	−0.18	−0.92, 0.56	−1.58	−3.46, 0.30	−0.85	−2.17, 0.48
High to low	−0.21	−0.96, 0.55	−2.04	−3.97, −0.10	−1.05	−2.41, 0.31
High to medium	−0.03	−0.75, 0.70	−0.46	−2.30, 1.39	−0.20	−1.51, 1.10
14 years						
Medium to low	−0.08	−0.88, 0.72	−1.55	−3.58, 0.48	−0.71	−2.07, 0.65
High to low	−0.16	−0.98, 0.66	−2.06	−4.14, 0.02	−0.86	−2.25, 0.54
High to medium	−0.08	−0.86, 0.70	−0.51	−2.50, 1.48	−0.14	−1.48, 1.19
15 years						
Medium to low	0.07	−0.79, 0.94	−1.41	−3.64, 0.83	−0.48	−1.91, 0.94
High to low	−0.07	−0.95, 0.82	−1.94	−4.23, 0.36	−0.53	−1.99, 0.93
High to medium	−0.14	−0.99, 0.71	−0.53	−2.73, 1.67	−0.05	−1.45, 1.35

CI, confidence interval.

^aModels included benzophenone-3 concentration tertiles, age, age squared, an interaction term between age and benzophenone-3 concentration tertiles, an interaction term between age squared and benzophenone-3 concentration tertiles, and an interaction term between race and age.

^bTertiles of benzophenone-3 (median, $\mu\text{g/g}$ creatinine): low, 5.3; medium, 28; high, 284.

Web Table 2. Predicted Differences^a in Body Mass Index, Waist Circumference, and Percent Body Fat From Ages 7–15 Years According to Tertiles (Low, Medium, and High^b) of Bisphenol A Concentrations Among Girls in the Breast Cancer and Environment Research Program, 2004–2015 (*n* = 1,017)

Age (Years) and Tertile Comparison	Predicted Difference in Body Mass Index		Predicted Difference in Waist Circumference, cm		Predicted Difference in Percent Body Fat, %	
	Difference	95% CI	Difference	95% CI	Difference	95% CI
7 years						
Medium to low	−0.01	−0.52, 0.50	0.62	−0.92, 2.15	−0.16	−1.58, 1.25
High to low	−0.34	−0.85, 0.17	−0.04	−1.57, 1.50	−1.31	−2.72, 0.10
High to medium	−0.33	−0.84, 0.18	−0.65	−2.18, 0.88	−1.15	−2.56, 0.26
8 years						
Medium to low	0.06	−0.45, 0.57	0.31	−1.07, 1.69	−0.09	−1.43, 1.25
High to low	−0.30	−0.81, 0.21	−0.46	−1.83, 0.92	−1.30	−2.64, 0.03
High to medium	−0.36	−0.87, 0.15	−0.77	−2.14, 0.60	−1.21	−2.54, 0.12
9 years						
Medium to low	0.12	−0.42, 0.66	0.10	−1.29, 1.50	−0.02	−1.32, 1.28
High to low	−0.27	−0.81, 0.27	−0.75	−2.15, 0.64	−1.28	−2.57, 0.01
High to medium	−0.39	−0.93, 0.14	−0.86	−2.25, 0.53	−1.26	−2.55, 0.04
10 years						
Medium to low	0.18	−0.40, 0.75	0.00	−1.50, 1.50	0.05	−1.23, 1.33
High to low	−0.24	−0.82, 0.33	−0.92	−2.42, 0.57	−1.24	−2.51, 0.04
High to medium	−0.42	−0.99, 0.15	−0.92	−2.42, 0.58	−1.29	−2.56, −0.01
11 years						
Medium to low	0.22	−0.40, 0.84	−0.01	−1.63, 1.61	0.12	−1.15, 1.40
High to low	−0.23	−0.85, 0.39	−0.97	−2.58, 0.64	−1.18	−2.46, 0.09
High to medium	−0.45	−1.07, 0.17	−0.96	−2.57, 0.66	−1.31	−2.58, −0.04
12 years						
Medium to low	0.26	−0.42, 0.93	0.08	−1.66, 1.81	0.20	−1.08, 1.48
High to low	−0.22	−0.88, 0.45	−0.89	−2.62, 0.84	−1.11	−2.39, 0.16
High to medium	−0.47	−1.14, 0.20	−0.97	−2.70, 0.76	−1.31	−2.59, −0.04
13 years						
Medium to low	0.28	−0.44, 1.00	0.26	−1.58, 2.11	0.27	−1.02, 1.57
High to low	−0.21	−0.93, 0.50	−0.69	−2.53, 1.15	−1.03	−2.32, 0.27
High to medium	−0.49	−1.21, 0.22	−0.95	−2.80, 0.89	−1.30	−2.59, −0.01
14 years						
Medium to low	0.29	−0.49, 1.07	0.55	−1.44, 2.54	0.35	−0.98, 1.68
High to low	−0.22	−1.00, 0.55	−0.36	−2.34, 1.62	−0.93	−2.25, 0.40
High to medium	−0.52	−1.29, 0.26	−0.91	−2.89, 1.07	−1.28	−2.61, 0.05
15 years						
Medium to low	0.30	−0.55, 1.15	0.94	−1.26, 3.13	0.43	−0.96, 1.83
High to low	−0.24	−1.08, 0.61	0.09	−2.09, 2.28	−0.81	−2.20, 0.58
High to medium	−0.54	−1.38, 0.31	−0.84	−3.04, 1.35	−1.24	−2.63, 0.15

CI, confidence interval.

^aModels included bisphenol A concentration tertiles, age, age squared, an interaction term between age and bisphenol A concentration tertiles, an interaction term between age squared and bisphenol A concentration tertiles, and an interaction term between race and age.

^bTertiles of bisphenol A (median, µg/g creatinine): low, 1.2; medium, 2.4; high, 5.8.

Web Table 3. Predicted Differences^a in Body Mass Index, Waist Circumference, and Percent Body Fat From Ages 7–15 Years According to Tertiles (Low, Medium, and High^b) of Daidzein Concentrations Among Girls in the Breast Cancer and Environment Research Program, 2004–2015 (*n* = 995)

Age (Years) and Tertile Comparison	Predicted Difference in Body Mass Index		Predicted Difference in Waist Circumference, cm		Predicted Difference in Percent Body Fat, %	
	Difference	95% CI	Difference	95% CI	Difference	95% CI
7 years						
Medium to low	0.38	−0.13, 0.90	0.36	−1.18, 1.91	1.07	−0.34, 2.49
High to low	0.30	−0.22, 0.82	0.49	−1.07, 2.05	0.91	−0.52, 2.34
High to medium	−0.08	−0.61, 0.44	0.13	−1.45, 1.70	−0.17	−1.62, 1.29
8 years						
Medium to low	0.39	−0.12, 0.91	0.80	−0.59, 2.19	1.16	−0.18, 2.50
High to low	0.33	−0.19, 0.85	0.66	−0.74, 2.06	0.99	−0.37, 2.34
High to medium	−0.06	−0.59, 0.46	−0.14	−1.55, 1.28	−0.17	−1.55, 1.20
9 years						
Medium to low	0.42	−0.12, 0.96	1.17	−0.23, 2.57	1.24	−0.06, 2.54
High to low	0.36	−0.19, 0.90	0.80	−0.61, 2.22	1.06	−0.25, 2.38
High to medium	−0.06	−0.62, 0.49	−0.37	−1.80, 1.07	−0.18	−1.51, 1.15
10 years						
Medium to low	0.47	−0.11, 1.05	1.47	−0.04, 2.97	1.32	0.03, 2.60
High to low	0.39	−0.19, 0.97	0.91	−0.61, 2.42	1.14	−0.16, 2.43
High to medium	−0.08	−0.67, 0.51	−0.56	−2.10, 0.98	−0.18	−1.49, 1.14
11 years						
Medium to low	0.54	−0.08, 1.16	1.70	0.07, 3.32	1.38	0.11, 2.66
High to low	0.43	−0.20, 1.06	0.98	−0.66, 2.62	1.21	−0.08, 2.50
High to medium	−0.11	−0.75, 0.52	−0.72	−2.38, 0.94	−0.17	−1.48, 1.14
12 years						
Medium to low	0.63	−0.04, 1.30	1.86	0.12, 3.59	1.44	0.16, 2.72
High to low	0.47	−0.21, 1.15	1.02	−0.74, 2.77	1.28	−0.01, 2.58
High to medium	−0.16	−0.85, 0.52	−0.84	−2.61, 0.94	−0.16	−1.47, 1.15
13 years						
Medium to low	0.74	0.02, 1.46	1.95	0.10, 3.80	1.50	0.20, 2.79
High to low	0.51	−0.22, 1.24	1.03	−0.84, 2.90	1.35	0.04, 2.66
High to medium	−0.23	−0.97, 0.51	−0.92	−2.81, 0.96	−0.15	−1.47, 1.18
14 years						
Medium to low	0.87	0.09, 1.65	1.98	−0.01, 3.97	1.55	0.22, 2.87
High to low	0.56	−0.23, 1.35	1.00	−1.01, 3.01	1.42	0.08, 2.76
High to medium	−0.31	−1.11, 0.49	−0.98	−3.01, 1.06	−0.13	−1.49, 1.23
15 years						
Medium to low	1.02	0.18, 1.87	1.93	−0.27, 4.14	1.59	0.20, 2.98
High to low	0.61	−0.24, 1.47	0.94	−1.27, 3.16	1.49	0.08, 2.89
High to medium	−0.41	−1.28, 0.46	−0.99	−3.24, 1.25	−0.10	−1.52, 1.32

CI, confidence interval.

^aModels included daidzein concentration tertiles, age, age squared, an interaction term between age and daidzein concentration tertiles, an interaction term between age squared and daidzein concentration tertiles, and an interaction term between race and age.

^bTertiles of daidzein (median, $\mu\text{g/g}$ creatinine): low, 26; medium, 98; high, 644.

Web Table 4. Predicted Differences^a in Body Mass Index, Waist Circumference, and Percent Body Fat From Ages 7–15 Years According to Tertiles (Low, Medium, and High^b) of Genistein Concentrations Among Girls in the Breast Cancer and Environment Research Program, 2004–2015 (*n* = 995)

Age (Years) and Tertile Comparison	Predicted Difference in Body Mass Index		Predicted Difference in Waist Circumference, cm		Predicted Difference in Percent Body Fat, %	
	Difference	95% CI	Difference	95% CI	Difference	95% CI
7 years						
Medium to low	0.36	−0.16, 0.87	0.43	−1.13, 1.98	0.39	−1.04, 1.81
High to low	0.19	−0.33, 0.71	0.37	−1.21, 1.94	0.18	−1.26, 1.62
High to medium	−0.16	−0.69, 0.36	−0.06	−1.63, 1.51	−0.20	−1.66, 1.25
8 years						
Medium to low	0.35	−0.16, 0.87	0.76	−0.63, 2.15	0.57	−0.77, 1.92
High to low	0.13	−0.39, 0.65	0.29	−1.11, 1.70	0.16	−1.20, 1.53
High to medium	−0.22	−0.75, 0.31	−0.47	−1.88, 0.95	−0.41	−1.79, 0.97
9 years						
Medium to low	0.37	−0.17, 0.91	1.07	−0.33, 2.47	0.74	−0.57, 2.04
High to low	0.10	−0.45, 0.65	0.26	−1.16, 1.68	0.18	−1.14, 1.50
High to medium	−0.27	−0.83, 0.28	−0.81	−2.24, 0.63	−0.56	−1.89, 0.78
10 years						
Medium to low	0.41	−0.17, 0.99	1.35	−0.16, 2.85	0.88	−0.41, 2.16
High to low	0.09	−0.49, 0.68	0.27	−1.26, 1.79	0.24	−1.06, 1.54
High to medium	−0.32	−0.91, 0.28	−1.08	−2.62, 0.46	−0.64	−1.96, 0.68
11 years						
Medium to low	0.47	−0.15, 1.10	1.59	−0.04, 3.21	1.00	−0.28, 2.28
High to low	0.11	−0.52, 0.75	0.31	−1.33, 1.96	0.33	−0.96, 1.63
High to medium	−0.36	−1.00, 0.28	−1.28	−2.94, 0.39	−0.66	−1.98, 0.65
12 years						
Medium to low	0.55	−0.12, 1.23	1.80	0.06, 3.54	1.09	−0.19, 2.37
High to low	0.16	−0.52, 0.84	0.40	−1.37, 2.16	0.46	−0.84, 1.77
High to medium	−0.39	−1.08, 0.30	−1.41	−3.19, 0.37	−0.63	−1.94, 0.69
13 years						
Medium to low	0.66	−0.07, 1.38	1.98	0.13, 3.84	1.16	−0.14, 2.46
High to low	0.24	−0.49, 0.97	0.52	−1.36, 2.39	0.63	−0.68, 1.95
High to medium	−0.42	−1.16, 0.32	−1.47	−3.36, 0.43	−0.53	−1.86, 0.81
14 years						
Medium to low	0.78	0.00, 1.57	2.14	0.14, 4.13	1.21	−0.12, 2.54
High to low	0.34	−0.45, 1.13	0.68	−1.33, 2.70	0.84	−0.51, 2.19
High to medium	−0.44	−1.24, 0.36	−1.46	−3.49, 0.58	−0.37	−1.73, 1.00
15 years						
Medium to low	0.93	0.08, 1.78	2.26	0.05, 4.46	1.23	−0.16, 2.63
High to low	0.47	−0.39, 1.33	0.88	−1.34, 3.11	1.09	−0.33, 2.50
High to medium	−0.46	−1.33, 0.41	−1.37	−3.62, 0.87	−0.15	−1.58, 1.28

CI, confidence interval.

^aModels included genistein concentration tertiles, age, age squared, an interaction term between age and genistein concentration tertiles, an interaction term between age squared and genistein concentration tertiles, and an interaction term between race and age.

^bTertiles of genistein (median, $\mu\text{g/g}$ creatinine): low, 12; medium, 45; high, 320.

Web Table 5. Predicted Differences^a in Body Mass Index, Waist Circumference, and Percent Body Fat From Ages 7–15 Years According to Tertiles (Low, Medium, and High^b) of the Sum of Parabens^c Concentrations Among Girls in the Breast Cancer and Environment Research Program (*n* = 937)

Age (Years) and Tertile Comparison	Predicted Difference in Body Mass Index		Predicted Difference in Waist Circumference, cm		Predicted Difference in Percent Body Fat, %	
	Difference	95% CI	Difference	95% CI	Difference	95% CI
7 years						
Medium to low	-0.01	-0.53, 0.50	0.09	-1.45, 1.63	-0.05	-1.48, 1.39
High to low	-0.03	-0.57, 0.51	-0.32	-1.91, 1.27	-0.06	-1.55, 1.44
High to medium	-0.02	-0.51, 0.48	-0.41	-1.90, 1.08	-0.01	-1.38, 1.36
8 years						
Medium to low	-0.03	-0.55, 0.50	-0.06	-1.43, 1.31	-0.10	-1.45, 1.26
High to low	0.01	-0.54, 0.55	-0.21	-1.65, 1.22	0.00	-1.42, 1.41
High to medium	0.03	-0.47, 0.53	-0.16	-1.49, 1.17	0.09	-1.20, 1.39
9 years						
Medium to low	-0.04	-0.59, 0.50	-0.21	-1.61, 1.19	-0.17	-1.49, 1.14
High to low	0.03	-0.54, 0.60	-0.14	-1.61, 1.33	0.00	-1.38, 1.38
High to medium	0.07	-0.45, 0.60	0.07	-1.29, 1.43	0.17	-1.09, 1.44
10 years						
Medium to low	-0.07	-0.66, 0.52	-0.38	-1.89, 1.14	-0.28	-1.58, 1.02
High to low	0.04	-0.58, 0.65	-0.09	-1.68, 1.50	-0.04	-1.40, 1.32
High to medium	0.11	-0.46, 0.67	0.28	-1.20, 1.76	0.23	-1.02, 1.48
11 years						
Medium to low	-0.10	-0.74, 0.53	-0.55	-2.20, 1.10	-0.41	-1.71, 0.89
High to low	0.03	-0.64, 0.69	-0.08	-1.81, 1.65	-0.14	-1.50, 1.22
High to medium	0.13	-0.48, 0.74	0.47	-1.14, 2.08	0.27	-0.98, 1.52
12 years						
Medium to low	-0.14	-0.83, 0.54	-0.73	-2.51, 1.04	-0.57	-1.87, 0.74
High to low	0.00	-0.72, 0.72	-0.10	-1.96, 1.76	-0.28	-1.65, 1.09
High to medium	0.14	-0.52, 0.80	0.63	-1.10, 2.36	0.28	-0.98, 1.54
13 years						
Medium to low	-0.19	-0.93, 0.55	-0.92	-2.82, 0.97	-0.75	-2.08, 0.58
High to low	-0.05	-0.83, 0.73	-0.15	-2.15, 1.85	-0.48	-1.87, 0.92
High to medium	0.14	-0.57, 0.86	0.77	-1.07, 2.62	0.27	-1.01, 1.55
14 years						
Medium to low	-0.25	-1.05, 0.56	-1.13	-3.18, 0.92	-0.96	-2.33, 0.41
High to low	-0.11	-0.96, 0.73	-0.23	-2.40, 1.93	-0.72	-2.16, 0.72
High to medium	0.14	-0.64, 0.91	0.89	-1.10, 2.88	0.24	-1.08, 1.56
15 years						
Medium to low	-0.31	-1.19, 0.56	-1.34	-3.62, 0.95	-1.20	-2.65, 0.24
High to low	-0.19	-1.11, 0.73	-0.35	-2.76, 2.07	-1.01	-2.53, 0.51
High to medium	0.12	-0.72, 0.96	0.99	-1.23, 3.21	0.19	-1.20, 1.58

CI, confidence interval.

^aModels included sum of parabens concentration tertiles, age, age squared, an interaction term between age and sum of parabens concentration tertiles, an interaction term between age squared and sum of parabens concentration tertiles, and an interaction term between race and age.

^bTertiles of sum of parabens (median, $\mu\text{g/g}$ creatinine): low, 23; medium, 92; high, 515.

^cSum of parabens: Paraben metabolites (methyl, ethyl, and propyl parabens) were summed based on molecular weight, expressed as propyl paraben (molecular weight 180.2 g/mol).

Web Table 6. Predicted Differences^a in Body Mass Index, Waist Circumference, and Percent Body Fat From Ages 7–15 Years According to Tertiles (Low, Medium, and High^b) of Triclosan Concentrations Among Normal Weight Girls (BMI<85th percentile) in the Breast Cancer and Environment Research Program, 2004–2015 (*n* = 683)

Age (Years) and Tertile Comparison	Predicted Difference in Body Mass Index		Predicted Difference in Waist Circumference, cm		Predicted Difference in Percent Body Fat, %	
	Difference	95% CI	Difference	95% CI	Difference	95% CI
7 years						
Medium to low	-0.02	-0.35, 0.31	0.11	-1.35, 1.57	0.07	-1.11, 1.25
High to low	-0.04	-0.38, 0.29	-0.69	-2.15, 0.77	-0.60	-1.81, 0.61
High to medium	-0.02	-0.35, 0.31	-0.80	-2.22, 0.62	-0.68	-1.87, 0.52
8 years						
Medium to low	-0.08	-0.38, 0.22	0.16	-0.90, 1.23	-0.02	-1.11, 1.08
High to low	-0.18	-0.48, 0.13	-0.36	-1.43, 0.70	-0.48	-1.60, 0.64
High to medium	-0.10	-0.39, 0.20	-0.53	-1.57, 0.51	-0.46	-1.57, 0.64
9 years						
Medium to low	-0.14	-0.45, 0.17	0.18	-0.82, 1.19	-0.12	-1.19, 0.95
High to low	-0.28	-0.59, 0.04	-0.08	-1.09, 0.94	-0.40	-1.50, 0.70
High to medium	-0.14	-0.45, 0.17	-0.26	-1.26, 0.74	-0.28	-1.36, 0.80
10 years						
Medium to low	-0.20	-0.55, 0.16	0.17	-0.97, 1.32	-0.25	-1.33, 0.84
High to low	-0.34	-0.71, 0.02	0.17	-1.00, 1.34	-0.37	-1.48, 0.74
High to medium	-0.15	-0.50, 0.21	0.00	-1.15, 1.14	-0.12	-1.22, 0.97
11 years						
Medium to low	-0.25	-0.66, 0.15	0.13	-1.18, 1.44	-0.38	-1.50, 0.74
High to low	-0.38	-0.80, 0.03	0.37	-0.97, 1.72	-0.38	-1.53, 0.77
High to medium	-0.13	-0.54, 0.28	0.24	-1.08, 1.57	0.00	-1.13, 1.13
12 years						
Medium to low	-0.31	-0.77, 0.14	0.05	-1.41, 1.51	-0.54	-1.71, 0.62
High to low	-0.39	-0.86, 0.08	0.54	-0.96, 2.03	-0.45	-1.64, 0.75
High to medium	-0.08	-0.54, 0.39	0.48	-0.98, 1.95	0.09	-1.09, 1.27
13 years						
Medium to low	-0.37	-0.89, 0.14	-0.06	-1.66, 1.54	-0.71	-1.94, 0.51
High to low	-0.36	-0.89, 0.16	0.66	-0.98, 2.29	-0.56	-1.81, 0.70
High to medium	0.01	-0.51, 0.53	0.72	-0.90, 2.33	0.16	-1.08, 1.39
14 years						
Medium to low	-0.43	-1.01, 0.15	-0.20	-1.99, 1.59	-0.90	-2.20, 0.40
High to low	-0.31	-0.90, 0.29	0.73	-1.09, 2.56	-0.72	-2.05, 0.62
High to medium	0.12	-0.46, 0.71	0.94	-0.87, 2.75	0.19	-1.13, 1.50
15 years						
Medium to low	-0.49	-1.15, 0.17	-0.38	-2.49, 1.73	-1.11	-2.53, 0.31
High to low	-0.22	-0.90, 0.45	0.77	-1.37, 2.92	-0.92	-2.37, 0.53
High to medium	0.27	-0.39, 0.94	1.15	-0.98, 3.28	0.19	-1.25, 1.62

CI, confidence interval.

^aModels included triclosan concentration tertiles (low, medium, and high), age (at exam, centered and estimated to the nearest tenth of a year), age squared (to allow for non-linearity), an interaction term between age and triclosan concentration tertiles, an interaction term between age squared and triclosan concentration tertiles, and an interaction term between race and age.

^bTertiles of triclosan (median, $\mu\text{g/g}$ creatinine): low, 3.7; medium, 15; high, 86.

Web Table 7. Predicted Differences^a in Body Mass Index, Waist Circumference, and Percent Body Fat From Ages 7–15 Years According to Tertiles of Enterolactone and 2,5-Dichlorophenol (Low, Medium, and High^b) Concentrations Among Girls in the Breast Cancer and Environment Research Program, 2004–2015 (*n* = 994)

Age (Years) and Tertile Comparison	Predicted Difference in Body Mass Index		Predicted Difference in Waist Circumference, cm		Predicted Difference in Percent Body Fat, %	
	Difference	95% CI	Difference	95% CI	Difference	95% CI
7 years						
Medium to low	0.17	−0.45, 0.79	0.71	−1.18, 2.61	0.80	−0.92, 0.36
High to low	0.84	0.00, 1.68	3.54	0.99, 6.08	3.12	0.80, 5.44
High to medium	0.67	0.04, 1.31	2.82	0.90, 4.74	2.32	0.57, 4.06
8 years						
Medium to low	0.54	−0.09, 1.16	1.34	−0.33, 3.02	1.50	−0.12, 3.12
High to low	1.46	0.62, 2.31	4.58	2.30, 6.86	4.24	2.05, 6.43
High to medium	0.93	0.29, 1.56	3.23	1.51, 4.96	2.74	1.09, 4.39
9 years						
Medium to low	0.84	0.19, 1.50	1.88	0.19, 3.57	2.05	0.47, 3.62
High to low	1.97	1.09, 2.86	5.47	3.18, 7.77	5.07	2.94, 7.19
High to medium	1.13	0.47, 1.80	3.59	1.86, 5.33	3.02	1.42, 4.62
10 years						
Medium to low	1.09	0.39, 1.79	2.32	0.51, 4.13	2.43	0.89, 3.98
High to low	2.37	1.43, 3.32	6.22	3.77, 8.66	5.59	3.50, 7.69
High to medium	1.28	0.57, 1.99	3.90	2.05, 5.75	3.16	1.59, 4.73
11 years						
Medium to low	1.27	0.52, 2.02	2.66	0.70, 4.62	2.66	1.12, 4.21
High to low	2.66	1.64, 3.68	6.81	4.17, 9.45	5.82	3.74, 7.90
High to medium	1.39	0.63, 2.15	4.15	2.16, 6.14	3.16	1.59, 4.72
12 years						
Medium to low	1.39	0.58, 2.20	2.91	0.81, 5.00	2.74	1.19, 4.29
High to low	2.84	1.74, 3.93	7.26	4.43, 10.08	5.75	3.66, 7.84
High to medium	1.44	0.62, 2.27	4.35	2.22, 6.48	3.01	1.44, 4.58
13 years						
Medium to low	1.45	0.58, 2.33	3.06	0.83, 5.29	2.65	1.09, 4.22
High to low	2.90	1.72, 4.08	7.55	4.54, 10.57	5.38	3.27, 7.50
High to medium	1.45	0.56, 2.33	4.49	2.23, 6.76	2.73	1.14, 4.31
14 years						
Medium to low	1.45	0.51, 2.40	3.12	0.72, 5.51	2.41	0.81, 4.02
High to low	2.85	1.58, 4.13	7.70	4.46, 10.94	4.71	2.55, 6.88
High to medium	1.40	0.45, 2.36	4.59	2.15, 7.02	2.30	0.67, 3.93
15 years						
Medium to low	1.39	0.37, 2.41	3.08	0.42, 5.73	2.02	0.33, 3.70
High to low	2.70	1.32, 4.07	7.70	4.14, 11.26	3.75	1.48, 6.01
High to medium	1.31	0.27, 2.34	4.62	1.95, 7.30	1.73	0.03, 3.43

CI, confidence interval.

^aModels included enterolactone/2,5-dichlorophenol tertiles (low, medium, and high), age (at exam, centered and estimated to the nearest tenth of a year), age squared (to allow for non-linearity), an interaction term between age and concentration tertiles, an interaction term between age squared and concentration tertiles, and an interaction term between race and age.

^b Categories were based on combinations of tertiles of 2,5-dichlorophenol and enterolactone. Low: low 2,5-dichlorophenol/high enterolactone; High: high 2,5-dichlorophenol/low enterolactone; Medium: all other 2,5-dichlorophenol/enterolactone.