

This slide set contains information about and data from the School Health Profiles or Profiles. It includes state data from 2016.

What is the School Health Profiles (Profiles)?

- Profiles is a system of surveys assessing school health policies and practices in states, territories, and large urban school districts
- Profiles surveys are conducted biennially (during even-numbered years) by state, territorial, and local education and health agencies with technical assistance from CDC
- Profiles questionnaires are administered to middle and high school principals and lead health education teachers

What is the School Health Profiles (Profiles)?

- Profiles is a system of surveys assessing school health policies and practices in states, territories, and large urban school districts
- Profiles surveys are conducted biennially (during even-numbered years) by state, territorial, and local education and health agencies with technical assistance from CDC
- Profiles questionnaires are administered to middle and high school principals and lead health education teachers

Topics Monitored by Profiles

- School health education requirements and content
- Physical education and physical activity
- Practices related to bullying and sexual harassment
- School health policies related to tobacco-use prevention and nutrition
- School-based health services
- Family engagement and community involvement
- School health coordination

Topics Monitored by Profiles


- School health education requirements and content
- Physical education and physical activity
- Practices related to bullying and sexual harassment
- School health policies related to tobacco-use prevention and nutrition
- School-based health services
- Family engagement and community involvement
- School health coordination

Profiles Participation, 1998 – 2016

	1998	2000	2002	2004	2006	2008	2010	2012	2014	2016
# of states	38	38	43	41	44	50	50	49	50	50
# of large urban school districts	10	13	14	13	16	21	20	19	21	21
# of territories	--	--	--	--	--	5	6	5	3	4
# of tribal governments	--	--	--	--	--	2	2	2	--	--
Total # of sites	48	51	57	54	60	78	78	75	74	75
Total % of sites with weighted data	77%	75%	77%	70%	82%	94%	99%	92%	96%	97%

This slide contains state, large urban school district, territory, and tribal government Profiles participation data from 1998 through 2016.

- In 1998, 38 states and 10 cities (including Washington, D.C.) participated in Profiles, for a total number of 48 states and cities. The percent of sites with weighted data in 1998 was 77%.
- In 2000, 38 states and 13 cities (including Washington, D.C.) participated in Profiles, for a total number of 51 states and cities. The percent of sites with weighted data in 2000 was 75%.
- In 2002, 43 states and 14 cities (including Washington, D.C.) participated in Profiles, for a total number of 57 states and cities. The percent of sites with weighted data in 2002 was 77%.
- In 2004, 41 states and 13 cities (including Washington, D.C.) participated in Profiles, for a total number of 54 states and cities. The percent of sites with weighted data in 2004 was 70%.
- In 2006, 44 states and 16 cities (including Washington, D.C.) participated in Profiles, for a total number of 60 states and cities. The percent of sites with weighted data in 2006 was 82%.
- In 2008, 50 states, 21 cities (including Washington, D.C.), 5 territories, and 2 tribes participated in Profiles, for a total number of 78 sites. The percent of sites with weighted data in 2008 was 94%.
- In 2010, 50 states, 20 cities (including Washington, D.C.), 6 territories, and 2 tribes participated in Profiles, for a total number of 78 sites. The percent of sites with weighted data in 2010 was 99%.
- In 2012, 49 states, 19 cities (including Washington, D.C.), 5 territories, and 2 tribes participated in Profiles, for a total number of 75 sites. The percent of sites with weighted data in 2012 was 92%.
- In 2014, 50 states, 21 cities (including Washington, D.C.), and 3 territories participated in Profiles, for a total number of 74 sites. The percent of sites with weighted data in 2014 was 96%.
- In 2016, 50 states, 21 cities (including Washington, D.C.), and 4 territories participated in Profiles, for a total number of 75 sites. The percent of sites with weighted data in 2016 was 97%.



This map illustrates states and large urban school districts that participated in the School Health Profiles in 2016. States and large urban school districts with weighted and unweighted results are shown.

States with weighted results are: Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming.

States with unweighted results are: Colorado and Iowa.

Surveyed large urban school districts with weighted results are: Baltimore, MD; Boston, MA; Broward County, FL; Chicago, IL; Cleveland, OH; DeKalb County, GA; Detroit, MI; District of Columbia; Duval County, FL; Fort Worth, TX; Houston, TX; Los Angeles, CA; Miami-Dade County, FL; New York City, NY; Oakland, CA; Orange County, FL; Palm Beach County, FL; Philadelphia, PA; San Diego, CA; San Francisco, CA; Shelby County, TN.

Surveyed territories with weighted results are: Guam, Northern Mariana Islands, Palau, and Puerto Rico.

Weighted data means that at least 70% of the principals or lead health education teachers in the sample completed the survey. Weighted data represent the state, large urban school district, territory, or tribal government. With weighted data, it is possible to say, for example, "X% of schools in our state offer students intramural activities or physical activity clubs."

Unweighted data represent only the schools that completed the questionnaire.

Uses of Profiles Data

Education and health officials use Profiles data to

- Describe school health policies and practices and compare them across jurisdictions
- Identify professional development needs
- Plan and monitor programs
- Support health-related policies and legislation
- Seek funding
- Garner support for future surveys

Uses of Profiles Data

Education and health officials use Profiles data to

- Describe school health policies and practices and compare them across jurisdictions
- Identify professional development needs
- Plan and monitor programs
- Support health-related policies and legislation
- Seek funding
- Garner support for future surveys

Examples of Uses of Profiles Data

- **District of Columbia (Office of the State Superintendent of Education)**
 - Used Profiles data to inform the modification and revamping of their 2016 Health Education Standards
- **Orange County Public Schools (Florida)**
 - Used Profiles data to support updates to the district's Local Wellness Policy, as well as to support updates to human sexuality lesson plans and resources for students in grades 6-12
- **Arkansas**
 - Used Profiles data to monitor the implementation of mandated requirements within districts

Examples of Uses of Profiles Data

Profiles Information

<http://www.cdc.gov/schoolhealthprofiles>

- Overview of the School Health Profiles
- Questionnaires and item rationales
- Comprehensive results
- Fact sheet
- Publications and journal articles
- Participation history
- Frequently asked questions about requesting data files
- Data file request form


For more information about Profiles go to <http://www.cdc.gov/schoolhealthprofiles>.

This web site contains

- Overview of the School Health Profiles
- Questionnaires and item rationales
- Comprehensive results
- Fact sheet
- Publications and journal articles
- Participation history
- Frequently asked questions about requesting data files
- Data file request form


HIV, STD, and Pregnancy Prevention

HIV, STD, and Pregnancy Prevention




This slide shows the percentage of secondary schools that taught 11 key HIV, STD, and pregnancy prevention topics in a required course during grades 6, 7, or 8 and during grades 9, 10, 11, or 12*. The values range from 7% to 66%. Alaska, Arizona, Idaho, Louisiana, Nebraska, North Dakota, South Dakota, Tennessee, Utah, range from 0% to 24%. Alabama, Arkansas, California, Florida, Georgia, Hawaii, Illinois, Indiana, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Missouri, Montana, Ohio, Pennsylvania, South Carolina, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. Connecticut, Delaware, Maryland, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, Rhode Island, West Virginia, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that assessed the ability of students to do 7 skills in a required course taught during grades 6, 7, or 8 and during grades 9, 10, 11, or 12*. The values range from 14% to 76%. Arizona, range from 0% to 24%. Alabama, Alaska, California, Florida, Hawaii, Idaho, Kansas, Louisiana, Maine, Massachusetts, Nebraska, North Dakota, South Dakota, Tennessee, Utah, Vermont, Washington, range from 25% to 49%. Arkansas, Connecticut, Georgia, Illinois, Indiana, Kentucky, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Virginia, Wisconsin, Wyoming, range from 50% to 74%. Delaware, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.


*During the current school year.




This slide shows the percentage of secondary schools in which those who teach sexual health education are provided with key materials for teaching sexual health education. The values range from 35% to 87%. Alaska, Arizona, Arkansas, Delaware, Hawaii, Illinois, Indiana, Kansas, Maine, Minnesota, Missouri, Montana, New Mexico, North Dakota, Ohio, Oregon, South Dakota, Utah, Vermont, West Virginia, Wyoming, range from 25% to 49%. Alabama, California, Connecticut, Florida, Georgia, Idaho, Kentucky, Louisiana, Massachusetts, Michigan, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, Rhode Island, South Carolina, Tennessee, Virginia, Washington, Wisconsin, range from 50% to 74%. Maryland, Mississippi, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on HIV prevention in a required course. The values range from 36% to 98%. Arizona, range from 25% to 49%. Alaska, Florida, Louisiana, Tennessee, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.



This slide shows the percentage of secondary schools that tried to increase student knowledge on pregnancy prevention in a required course. The values range from 32% to 90%. Arizona, range from 25% to 49%. Alabama, Alaska, California, Florida, Georgia, Hawaii, Louisiana, Michigan, South Dakota, Tennessee, range from 50% to 74%. Arkansas, Connecticut, Delaware, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on STD prevention in a required course. The values range from 35% to 94%. Arizona, range from 25% to 49%. Alabama, Alaska, Florida, Louisiana, Tennessee, range from 50% to 74%. Arkansas, California, Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that taught how HIV and other STDs are transmitted in a required course during grades 6, 7, or 8*. The values range from 18% to 93%. Arizona, range from 0% to 24%. Alabama, Alaska, Kansas, Louisiana, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Kentucky, Maine, Massachusetts, Michigan, Montana, Nebraska, New Mexico, Ohio, Pennsylvania, Vermont, range from 50% to 74%. Delaware, Illinois, Indiana, Maryland, Minnesota, Mississippi, Missouri, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Rhode Island, South Carolina, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how HIV and other STDs are transmitted in a required course during grades 9, 10, 11, or 12*. The values range from 49% to 100%. Arizona, range from 25% to 49%. Alaska, Louisiana, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught health consequences of HIV, other STDs, and pregnancy in a required course during grades 6, 7, or 8*. The values range from 18% to 91%. Arizona, range from 0% to 24%. Alabama, Alaska, Kansas, Louisiana, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Florida, Georgia, Hawaii, Idaho, Kentucky, Maine, Massachusetts, Michigan, Montana, Nebraska, New Mexico, Ohio, Pennsylvania, Vermont, range from 50% to 74%. Connecticut, Delaware, Illinois, Indiana, Maryland, Minnesota, Mississippi, Missouri, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Rhode Island, South Carolina, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught health consequences of HIV, other STDs, and pregnancy in a required course during grades 9, 10, 11, or 12*. The values range from 47% to 100%. Arizona, range from 25% to 49%. Alaska, Louisiana, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the benefits of being sexually abstinent in a required course during grades 6, 7, or 8*. The values range from 18% to 93%. Arizona, range from 0% to 24%. Alabama, Alaska, Louisiana, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Florida, Georgia, Hawaii, Idaho, Kansas, Kentucky, Maine, Massachusetts, Michigan, Nebraska, New Mexico, Ohio, Pennsylvania, Vermont, range from 50% to 74%. Connecticut, Delaware, Illinois, Indiana, Maryland, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Rhode Island, South Carolina, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the benefits of being sexually abstinent in a required course during grades 9, 10, 11, or 12*. The values range from 42% to 100%. Arizona, range from 25% to 49%. Alaska, Louisiana, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to access valid and reliable health information, products, or services related to HIV, other STDs, and pregnancy in a required course during grades 6, 7, or 8*. The values range from 13% to 86%. Arizona, range from 0% to 24%. Alabama, Alaska, Georgia, Hawaii, Kansas, Kentucky, Louisiana, Nebraska, Ohio, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Connecticut, Florida, Idaho, Illinois, Indiana, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, New Mexico, North Dakota, Oregon, Pennsylvania, South Carolina, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Delaware, Maryland, Nevada, New Hampshire, New Jersey, New York, North Carolina, Rhode Island, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to access valid and reliable health information, products, or services related to HIV, other STDs, and pregnancy in a required course during grades 9, 10, 11, or 12*. The values range from 36% to 100%. Arizona, range from 25% to 49%. Alaska, Louisiana, Nebraska, North Dakota, South Dakota, Utah, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the influences of family, peers, media, technology, and other factors on sexual risk behaviors in a required course during grades 6, 7, or 8*. The values range from 17% to 88%. Arizona, range from 0% to 24%. Alabama, Alaska, Kansas, Louisiana, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Florida, Georgia, Hawaii, Idaho, Illinois, Kentucky, Maine, Massachusetts, Michigan, Missouri, Montana, Nebraska, New Mexico, Ohio, Pennsylvania, Vermont, Washington, Wisconsin, Wyoming, range from 50% to 74%. Connecticut, Delaware, Indiana, Maryland, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Rhode Island, South Carolina, Utah, Virginia, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the influences of family, peers, media, technology, and other factors on sexual risk behaviors in a required course during grades 9, 10, 11, or 12*. The values range from 35% to 100%. Arizona, range from 25% to 49%. Alaska, Louisiana, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught communication and negotiation skills related to eliminating or reducing risk for HIV, other STDs, and pregnancy in a required course during grades 6, 7, or 8*. The values range from 15% to 87%. Arizona, range from 0% to 24%. Alabama, Alaska, Kansas, Kentucky, Louisiana, Nebraska, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Mexico, Ohio, Oregon, Pennsylvania, South Carolina, Vermont, Wisconsin, Wyoming, range from 50% to 74%. Delaware, Indiana, Maryland, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Rhode Island, Utah, Virginia, Washington, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught communication and negotiation skills related to eliminating or reducing risk for HIV, other STDs, and pregnancy in a required course during grades 9, 10, 11, or 12*. The values range from 36% to 100%. Arizona, range from 25% to 49%. Alaska, Louisiana, Nebraska, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught goal-setting and decision-making skills related to eliminating or reducing risk for HIV, other STDs, and pregnancy in a required course during grades 6, 7, or 8*. The values range from 17% to 87%. Arizona, range from 0% to 24%. Alabama, Alaska, Hawaii, Kansas, Kentucky, Louisiana, Nebraska, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Connecticut, Florida, Georgia, Idaho, Illinois, Maine, Massachusetts, Michigan, Minnesota, Missouri, Montana, New Mexico, North Dakota, Ohio, Oregon, Pennsylvania, South Carolina, Vermont, Washington, Wisconsin, Wyoming, range from 50% to 74%. Delaware, Indiana, Maryland, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, Rhode Island, Utah, Virginia, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught goal-setting and decision-making skills related to eliminating or reducing risk for HIV, other STDs, and pregnancy in a required course during grades 9, 10, 11, or 12*. The values range from 38% to 97%. Arizona, range from 25% to 49%. Alaska, Hawaii, Louisiana, Nebraska, North Dakota, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught influencing and supporting others to avoid or reduce sexual risk behaviors in a required course during grades 6, 7, or 8*. The values range from 17% to 84%. Arizona, range from 0% to 24%. Alabama, Alaska, Hawaii, Kansas, Kentucky, Louisiana, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Connecticut, Florida, Georgia, Idaho, Illinois, Maine, Massachusetts, Michigan, Missouri, Montana, Nebraska, New Mexico, Ohio, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, Wisconsin, Wyoming, range from 50% to 74%. Delaware, Indiana, Maryland, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, South Carolina, Utah, Virginia, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught influencing and supporting others to avoid or reduce sexual risk behaviors in a required course during grades 9, 10, 11, or 12*. The values range from 37% to 100%. Arizona, range from 25% to 49%. Alaska, Hawaii, Louisiana, Nebraska, North Dakota, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught efficacy of condoms in a required course during grades 6, 7, or 8*. The values range from 10% to 79%. Alabama, Arizona, South Dakota, Tennessee, Utah, range from 0% to 24%. Alaska, Arkansas, Florida, Georgia, Hawaii, Idaho, Indiana, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, South Carolina, Wyoming, range from 25% to 49%. California, Connecticut, Delaware, Illinois, Maine, Maryland, Massachusetts, Mississippi, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, Rhode Island, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. Nevada, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught efficacy of condoms in a required course during grades 9, 10, 11, or 12*. The values range from 35% to 99%. Alaska, Arizona, Louisiana, Utah, range from 25% to 49%. Florida, Hawaii, Idaho, Indiana, Kansas, Montana, Nebraska, North Dakota, South Dakota, Tennessee, Wyoming, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Georgia, Illinois, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the importance of using condoms consistently and correctly in a required course during grades 6, 7, or 8*. The values range from 7% to 68%. Alabama, Arizona, Kansas, Kentucky, South Dakota, Tennessee, Utah, range from 0% to 24%. Alaska, Arkansas, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Louisiana, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, South Carolina, Virginia, Wisconsin, Wyoming, range from 25% to 49%. California, Delaware, Maine, Maryland, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, Rhode Island, Vermont, Washington, West Virginia, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the importance of using condoms consistently and correctly in a required course during grades 9, 10, 11, or 12*. The values range from 33% to 98%. Alaska, Arizona, Louisiana, Nebraska, South Dakota, Utah, range from 25% to 49%. Alabama, Arkansas, Florida, Georgia, Hawaii, Idaho, Indiana, Kansas, Michigan, Mississippi, Missouri, Montana, New Mexico, North Dakota, Ohio, South Carolina, Tennessee, Virginia, Wyoming, range from 50% to 74%. California, Connecticut, Delaware, Illinois, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Nevada, New Hampshire, New Jersey, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to obtain condoms in a required course during grades 6, 7, or 8*. The values range from 0% to 56%. Alabama, Arizona, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Michigan, Missouri, Nebraska, North Dakota, Ohio, Pennsylvania, South Dakota, Tennessee, Utah, range from 0% to 24%. Alaska, Arkansas, California, Connecticut, Florida, Hawaii, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Montana, Nevada, New Hampshire, New Jersey, Oregon, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. Delaware, New Mexico, New York, North Carolina, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to obtain condoms in a required course during grades 9, 10, 11, or 12*. The values range from 19% to 94%. Utah, range from 0% to 24%. Alaska, Arizona, Georgia, Idaho, Indiana, Kansas, Louisiana, Missouri, Nebraska, North Dakota, South Dakota, Tennessee, range from 25% to 49%. Alabama, Arkansas, California, Florida, Hawaii, Illinois, Kentucky, Michigan, Minnesota, Mississippi, Montana, Nevada, New Mexico, North Carolina, Ohio, Pennsylvania, South Carolina, Virginia, Wisconsin, Wyoming, range from 50% to 74%. Connecticut, Delaware, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, Oregon, Rhode Island, Vermont, Washington, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to correctly use a condom in a required course during grades 6, 7, or 8*. The values range from 1% to 50%. Alabama, Alaska, Arizona, Arkansas, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Virginia, range from 0% to 24%. California, Connecticut, Delaware, Florida, Hawaii, Maine, Maryland, Massachusetts, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, Oregon, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New Mexico, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to correctly use a condom in a required course during grades 9, 10, 11, or 12*. The values range from 12% to 92%. Arizona, Louisiana, South Dakota, Utah, range from 0% to 24%. Alabama, Alaska, Arkansas, Georgia, Idaho, Indiana, Kansas, Michigan, Missouri, Montana, Nebraska, North Dakota, Ohio, Tennessee, Virginia, Wyoming, range from 25% to 49%. California, Florida, Hawaii, Illinois, Kentucky, Massachusetts, Minnesota, Mississippi, Nevada, New Mexico, North Carolina, Pennsylvania, South Carolina, Washington, West Virginia, Wisconsin, range from 50% to 74%. Connecticut, Delaware, Maine, Maryland, New Hampshire, New Jersey, New York, Oregon, Rhode Island, Vermont, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught methods of contraception other than condoms in a required course during grades 6, 7, or 8*. The values range from 7% to 73%. Alabama, Arizona, Kansas, Kentucky, Louisiana, South Dakota, Tennessee, Utah, range from 0% to 24%. Alaska, Arkansas, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, Rhode Island, South Carolina, Virginia, Wisconsin, Wyoming, range from 25% to 49%. California, Delaware, Maine, Maryland, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, Vermont, Washington, West Virginia, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught methods of contraception other than condoms in a required course during grades 9, 10, 11, or 12*. The values range from 36% to 99%. Alaska, Arizona, Louisiana, South Dakota, Utah, range from 25% to 49%. Alabama, Arkansas, Florida, Georgia, Hawaii, Idaho, Indiana, Kansas, Michigan, Missouri, Montana, Nebraska, North Dakota, Tennessee, Wyoming, range from 50% to 74%. California, Connecticut, Delaware, Illinois, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the importance of using a condom at the same time as another form of contraception to prevent both STDs and pregnancy in a required course during grades 6, 7, or 8*. The values range from 6% to 82%. Alabama, Arizona, Kansas, Kentucky, Louisiana, South Dakota, Tennessee, Utah, range from 0% to 24%. Alaska, Arkansas, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, South Carolina, Virginia, Wyoming, range from 25% to 49%. California, Maine, Maryland, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, range from 50% to 74%. Delaware, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the importance of using a condom at the same time as another form of contraception to prevent both STDs and pregnancy in a required course during grades 9, 10, 11, or 12*. The values range from 31% to 100%. Alaska, Arizona, Louisiana, Utah, range from 25% to 49%. Alabama, Arkansas, Florida, Georgia, Hawaii, Idaho, Indiana, Kansas, Michigan, Missouri, Montana, Nebraska, North Dakota, Ohio, South Carolina, South Dakota, Tennessee, Virginia, Wyoming, range from 50% to 74%. California, Connecticut, Delaware, Illinois, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to create and sustain healthy and respectful relationships in a required course during grades 6, 7, or 8*. The values range from 18% to 95%. Arizona, range from 0% to 24%. Alabama, Alaska, Louisiana, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Delaware, Florida, Georgia, Hawaii, Idaho, Kansas, Kentucky, Maine, Massachusetts, Michigan, Missouri, Montana, Nebraska, New Mexico, Ohio, Pennsylvania, Washington, Wyoming, range from 50% to 74%. Connecticut, Illinois, Indiana, Maryland, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Rhode Island, South Carolina, Utah, Vermont, Virginia, West Virginia, Wisconsin, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught how to create and sustain healthy and respectful relationships in a required course during grades 9, 10, 11, or 12*. The values range from 45% to 100%. Arizona, range from 25% to 49%. Alaska, Louisiana, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the importance of limiting the number of sexual partners in a required course during grades 6, 7, or 8*. The values range from 11% to 81%. Alabama, Arizona, range from 0% to 24%. Alaska, Florida, Georgia, Hawaii, Kansas, Kentucky, Louisiana, Nebraska, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Connecticut, Delaware, Idaho, Illinois, Indiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Mexico, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. New Jersey, New York, North Carolina, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught the importance of limiting the number of sexual partners in a required course during grades 9, 10, 11, or 12*. The values range from 40% to 100%. Alaska, Arizona, range from 25% to 49%. Hawaii, Louisiana, Nebraska, North Dakota, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught preventive care that is necessary to maintain reproductive and sexual health in a required course during grades 6, 7, or 8*. The values range from 11% to 81%. Alabama, Arizona, range from 0% to 24%. Alaska, Florida, Georgia, Hawaii, Kansas, Kentucky, Louisiana, Michigan, Montana, Nebraska, Ohio, Pennsylvania, South Dakota, Tennessee, range from 25% to 49%. Arkansas, California, Connecticut, Delaware, Idaho, Illinois, Indiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Nevada, New Jersey, New Mexico, North Dakota, Oregon, Rhode Island, South Carolina, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. New Hampshire, New York, North Carolina, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught preventive care that is necessary to maintain reproductive and sexual health in a required course during grades 9, 10, 11, or 12*. The values range from 35% to 99%. Arizona, range from 25% to 49%. Alaska, Florida, Hawaii, Louisiana, Montana, Nebraska, North Dakota, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught about sexual orientation in a required course during grades 6, 7, or 8*. The values range from 8% to 63%. Alabama, Alaska, Arizona, Georgia, Idaho, Kansas, Kentucky, Louisiana, Nebraska, South Carolina, South Dakota, Tennessee, Utah, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Hawaii, Illinois, Indiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New Jersey, New Mexico, New York, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught about sexual orientation in a required course during grades 9, 10, 11, or 12*. The values range from 13% to 94%. Utah, range from 0% to 24%. Alaska, Arizona, Georgia, Idaho, Kansas, Louisiana, Michigan, Missouri, Montana, Nebraska, North Dakota, Pennsylvania, South Carolina, South Dakota, Tennessee, Virginia, Wyoming, range from 25% to 49%. Alabama, Arkansas, California, Delaware, Florida, Hawaii, Illinois, Indiana, Kentucky, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New Mexico, North Carolina, Ohio, Washington, West Virginia, Wisconsin, range from 50% to 74%. Connecticut, Maine, Maryland, New Jersey, New York, Oregon, Rhode Island, Vermont, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught about gender roles, gender identity, or gender expression in a required course during grades 6, 7, or 8*. The values range from 10% to 66%. Alabama, Alaska, Arizona, Georgia, Idaho, Kansas, Kentucky, Louisiana, South Carolina, South Dakota, Tennessee, Utah, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Hawaii, Illinois, Indiana, Maine, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. Maryland, Massachusetts, Mississippi, New Jersey, New Mexico, New York, Rhode Island, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools that taught about gender roles, gender identity, or gender expression in a required course during grades 9, 10, 11, or 12*. The values range from 29% to 94%. Alaska, Arizona, Georgia, Idaho, Louisiana, Michigan, Missouri, Montana, Nebraska, North Dakota, South Carolina, South Dakota, Utah, Virginia, Wyoming, range from 25% to 49%. Alabama, Arkansas, California, Florida, Hawaii, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New Mexico, North Carolina, Ohio, Oregon, Pennsylvania, Tennessee, Washington, West Virginia, Wisconsin, range from 50% to 74%. Connecticut, Delaware, New Jersey, New York, Rhode Island, Vermont, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on HIV prevention. The values range from 10% to 57%. Alabama, Alaska, Arizona, Idaho, Indiana, Kansas, Kentucky, Missouri, Montana, Nebraska, North Dakota, Oregon, Pennsylvania, Rhode Island, South Dakota, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Louisiana, Maine, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maryland, Michigan, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

* During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on human sexuality. The values range from 9% to 55%. Alabama, Alaska, Arizona, Hawaii, Idaho, Indiana, Kansas, Kentucky, Louisiana, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, South Dakota, West Virginia, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Illinois, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia, Washington, Wisconsin, range from 25% to 49%. Maryland, Utah, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on pregnancy prevention. The values range from 9% to 48%. Alabama, Alaska, Arizona, Georgia, Idaho, Indiana, Kansas, Kentucky, Louisiana, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Utah, Virginia, Wisconsin, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Hawaii, Illinois, Maine, Maryland, Massachusetts, Michigan, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, South Carolina, Tennessee, Vermont, Washington, West Virginia, range from 25% to 49%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on STD prevention. The values range from 10% to 54%. Alabama, Alaska, Arizona, Idaho, Indiana, Kansas, Kentucky, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Virginia, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Louisiana, Maine, Massachusetts, Michigan, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, South Carolina, Tennessee, Utah, Vermont, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maryland, Mississippi, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools that provide curricula or supplementary materials that include HIV, STD, or pregnancy prevention information that is relevant to LGBTQ youth . The values range from 20% to 68%. Arizona, Louisiana, Utah, range from 0% to 24%. Alabama, Alaska, Arkansas, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Michigan, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Mexico, North Carolina, North Dakota, Ohio, Pennsylvania, South Carolina, South Dakota, Tennessee, Virginia, West Virginia, Wyoming, range from 25% to 49%. California, Connecticut, Delaware, Maine, Maryland, Massachusetts, Minnesota, Mississippi, New Jersey, New York, Oregon, Rhode Island, Vermont, Washington, Wisconsin, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on teaching students of different sexual orientations or gender identities . The values range from 7% to 46%. Alabama, Alaska, Arizona, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Missouri, Montana, Nebraska, North Carolina, North Dakota, Ohio, South Carolina, South Dakota, Tennessee, Utah, Virginia, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Maine, Maryland, Massachusetts, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, range from 25% to 49%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on aligning lessons and materials with the district scope and sequence for sexual health education. The values range from 12% to 59%. Alabama, Alaska, Arizona, Idaho, Indiana, Kansas, Kentucky, Montana, Nebraska, North Dakota, Ohio, Rhode Island, South Dakota, West Virginia, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Louisiana, Maine, Massachusetts, Michigan, Minnesota, Missouri, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, Pennsylvania, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. Maryland, Mississippi, Nevada, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on creating a comfortable and safe learning environment for students receiving sexual health education . The values range from 11% to 53%. Alabama, Alaska, Arizona, Idaho, Indiana, Kansas, Kentucky, Louisiana, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, South Dakota, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Maine, Massachusetts, Michigan, Minnesota, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maryland, Mississippi, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on connecting students to on-site or community-based sexual health services . The values range from 5% to 41%. Alabama, Alaska, Arizona, Arkansas, California, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Minnesota, Missouri, Montana, Nebraska, New Hampshire, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Connecticut, Delaware, Florida, Maryland, Massachusetts, Michigan, Mississippi, Nevada, New Jersey, New Mexico, New York, North Carolina, Vermont, range from 25% to 49%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on using a variety of effective instructional strategies to deliver sexual health education . The values range from 7% to 54%. Alabama, Alaska, Arizona, Idaho, Indiana, Kansas, Kentucky, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Louisiana, Maine, Massachusetts, Michigan, Minnesota, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, South Carolina, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maryland, Mississippi, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on building student skills in HIV, other STDs, and pregnancy prevention. The values range from 7% to 53%. Alabama, Alaska, Arizona, Idaho, Indiana, Kansas, Kentucky, Louisiana, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Virginia, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, South Carolina, Tennessee, Utah, Vermont, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maryland, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.



This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on assessing student knowledge and skills in sexual health education . The values range from 7% to 55%. Alabama, Alaska, Arizona, Idaho, Indiana, Kansas, Kentucky, Louisiana, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, South Dakota, Tennessee, Virginia, West Virginia, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Rhode Island, South Carolina, Utah, Vermont, Washington, Wisconsin, range from 25% to 49%. Maryland, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on understanding current district or school board policies or curriculum guidance regarding sexual health education. The values range from 9% to 63%. Alabama, Alaska, Arizona, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Missouri, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, Rhode Island, South Dakota, West Virginia, Wyoming, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Louisiana, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Oregon, South Carolina, Tennessee, Vermont, Virginia, Washington, Wisconsin, range from 25% to 49%. Maryland, Mississippi, Nevada, Utah, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.


*During the 2 years before the survey.

Practices Related to Safe and Supportive Environments


Practices Related to Safe and Supportive Environments




This slide shows the percentage of secondary schools that implement parent engagement strategies for all students. The values range from 24% to 64%. Wyoming, range from 0% to 24%. Alabama, Alaska, Arizona, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Louisiana, Maine, Michigan, Missouri, Montana, Nebraska, Nevada, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Virginia, Washington, Wisconsin, range from 25% to 49%. Arkansas, California, Kentucky, Maryland, Massachusetts, Minnesota, Mississippi, New Hampshire, New Jersey, New York, Rhode Island, Texas, Utah, Vermont, West Virginia, range from 50% to 74%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that implement school connectedness strategies. The values range from 63% to 94%. Alaska, Arizona, Idaho, Kansas, Maine, Montana, Nevada, New Mexico, North Dakota, Oklahoma, Pennsylvania, Rhode Island, South Dakota, Washington, Wyoming, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Indiana, Kentucky, Louisiana, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oregon, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, West Virginia, Wisconsin, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that prevent bullying and sexual harassment, including electronic aggression, among all students. The values range from 21% to 78%. Alaska, Wyoming, range from 0% to 24%. Arizona, California, Connecticut, Delaware, Georgia, Hawaii, Idaho, Kansas, Maine, Michigan, Missouri, Montana, Nebraska, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Dakota, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Alabama, Arkansas, Florida, Illinois, Indiana, Kentucky, Louisiana, Maryland, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New York, South Carolina, Tennessee, Vermont, range from 50% to 74%. New Jersey, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools with a gay/straight alliance or similar club. The values range from 9% to 61%. Alabama, Arizona, Arkansas, Georgia, Illinois, Kansas, Kentucky, Louisiana, Mississippi, Montana, Nebraska, North Dakota, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Wyoming, range from 0% to 24%. Alaska, California, Delaware, Florida, Hawaii, Idaho, Indiana, Maine, Maryland, Michigan, Minnesota, Missouri, Nevada, New Hampshire, New Jersey, New Mexico, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Connecticut, Massachusetts, New York, range from 50% to 74%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that identify “safe spaces” where LGBTQ youth can receive support from administrators, teachers, or other school staff. The values range from 48% to 91%. Alaska, North Dakota, South Dakota, Wyoming, range from 25% to 49%. Alabama, Arizona, Arkansas, Delaware, Georgia, Hawaii, Idaho, Illinois, Kansas, Kentucky, Louisiana, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, New Mexico, North Carolina, Ohio, Oklahoma, South Carolina, Tennessee, Texas, Virginia, West Virginia, Wisconsin, range from 50% to 74%. California, Connecticut, Florida, Indiana, Maine, Massachusetts, Minnesota, Nevada, New Hampshire, New Jersey, New York, Oregon, Pennsylvania, Rhode Island, Utah, Vermont, Washington, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that prohibit harassment based on a student's perceived or actual sexual orientation or gender identity. The values range from 78% to 99%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that encourage staff to attend professional development on safe and supportive school environments for all students, regardless of sexual orientation or gender identity. The values range from 46% to 89%. Arizona, Wyoming, range from 25% to 49%. Alabama, Alaska, Arkansas, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Michigan, Mississippi, Missouri, Montana, Nebraska, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. California, Connecticut, Florida, Maine, Maryland, Massachusetts, Minnesota, Nevada, New Hampshire, New Jersey, New York, Rhode Island, Vermont, range from 75% to 100%. Colorado and Iowa did not have weighted results.



This slide shows the percentage of secondary schools that facilitate access to providers not on school property who have experience in providing health services to LGBTQ youth. The values range from 31% to 72%. Alabama, Alaska, Arizona, Arkansas, Georgia, Idaho, Illinois, Indiana, Kansas, Louisiana, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, Wisconsin, Wyoming, range from 25% to 49%. California, Connecticut, Delaware, Florida, Hawaii, Kentucky, Maine, Maryland, Massachusetts, Minnesota, New Hampshire, New Jersey, New Mexico, New York, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, range from 50% to 74%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that facilitate access to providers not on school property who have experience in providing social and psychological services to LGBTQ youth. The values range from 36% to 79%. Alabama, Arizona, Arkansas, Georgia, Idaho, Illinois, Kansas, Louisiana, Mississippi, Montana, Nebraska, North Carolina, North Dakota, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Wyoming, range from 25% to 49%. Alaska, California, Connecticut, Delaware, Florida, Hawaii, Indiana, Kentucky, Maine, Maryland, Michigan, Minnesota, Missouri, Nevada, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oregon, Pennsylvania, Rhode Island, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. Massachusetts, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that provide curricula or supplementary materials and engage in 5 practices related to LGBTQ youth. The values range from 3% to 40%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Maine, Massachusetts, Minnesota, New Hampshire, New Jersey, New York, Rhode Island, Vermont, range from 25% to 49%. Colorado and Iowa did not have weighted results.

School Health Services


School Health Services



This slide shows the percentage of secondary schools that have a full-time registered nurse who provides health services to students. The values range from 2% to 99%. Alaska, California, Hawaii, Idaho, Michigan, Montana, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Utah, Wisconsin, range from 0% to 24%. Arizona, Kansas, Louisiana, Maine, Mississippi, Nebraska, North Carolina, Ohio, Washington, West Virginia, Wyoming, range from 25% to 49%. Florida, Georgia, Illinois, Kentucky, Maryland, Minnesota, New Mexico, Tennessee, range from 50% to 74%. Alabama, Arkansas, Connecticut, Delaware, Indiana, Massachusetts, Missouri, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Texas, Vermont, Virginia, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that have a part-time registered nurse who provides health services to students. The values range from 11% to 83%. Alabama, Alaska, Arizona, Delaware, Hawaii, Missouri, New Hampshire, New Jersey, New York, North Dakota, Rhode Island, South Carolina, range from 0% to 24%. Arkansas, Connecticut, Florida, Georgia, Idaho, Illinois, Indiana, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Montana, Oklahoma, Pennsylvania, South Dakota, Tennessee, Texas, Vermont, Virginia, range from 25% to 49%. California, Kansas, Louisiana, Maine, Mississippi, Nebraska, New Mexico, North Carolina, Ohio, Oregon, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Nevada, Utah, Washington, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that have a school-based health center* that offers health services to students. The values range from 6% to 52%. Alabama, Alaska, Arkansas, California, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Louisiana, Maine, Massachusetts, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, Washington, Wisconsin, Wyoming, range from 0% to 24%. Arizona, Connecticut, Delaware, Indiana, Kentucky, Maryland, Minnesota, New Jersey, New Mexico, Vermont, range from 25% to 49%. West Virginia, range from 50% to 74%. Colorado and Iowa did not have weighted results.


*A place on school campus where enrolled students can receive primary care.




This slide shows the percentage of secondary schools that provide students with on-site services or referrals to healthcare providers for 7 sexual health services. The values range from 8% to 53%. Alabama, Alaska, Arizona, Arkansas, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Louisiana, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Virginia, Wisconsin, range from 0% to 24%. California, Connecticut, Delaware, Kentucky, Maine, Maryland, Massachusetts, Minnesota, New Hampshire, New Mexico, Oregon, Rhode Island, Vermont, Washington, West Virginia, Wyoming, range from 25% to 49%. New York, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide HIV testing to students. The values range from 0% to 19%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide HIV treatment to students. The values range from 0% to 9%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide STD testing to students. The values range from 0% to 23%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide STD treatment to students. The values range from 0% to 19%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide pregnancy testing to students. The values range from 0% to 26%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Delaware, range from 25% to 49%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide condoms to students. The values range from 0% to 21%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide condom-compatible lubricants to students. The values range from 0% to 10%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide contraceptives other than condoms to students. The values range from 0% to 11%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.



This slide shows the percentage of secondary schools that provide prenatal care to students. The values range from 0% to 15%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide HPV vaccine administration to students. The values range from 0% to 29%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Rhode Island, range from 25% to 49%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide students with referrals* for HIV testing. The values range from 12% to 57%. Alabama, Alaska, Arizona, Georgia, Hawaii, Idaho, Illinois, Kansas, Louisiana, Nebraska, Nevada, Ohio, Tennessee, Texas, Virginia, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, New Hampshire, New Jersey, New Mexico, North Carolina, North Dakota, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for HIV treatment. The values range from 18% to 58%. Alabama, Arizona, Georgia, Hawaii, Louisiana, range from 0% to 24%. Alaska, Arkansas, California, Connecticut, Delaware, Florida, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New Mexico, New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for nPEP (non-occupational post-exposure prophylaxis for HIV). The values range from 16% to 58%. Alabama, Arizona, Georgia, Hawaii, Idaho, Louisiana, Tennessee, range from 0% to 24%. Alaska, Arkansas, California, Connecticut, Delaware, Florida, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New Mexico, New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for STD testing. The values range from 14% to 58%. Alabama, Alaska, Arizona, Georgia, Hawaii, Idaho, Kansas, Louisiana, Nevada, Ohio, Tennessee, Texas, Virginia, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Illinois, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, North Carolina, North Dakota, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for STD treatment. The values range from 11% to 57%. Alabama, Alaska, Arizona, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Louisiana, Nebraska, Nevada, Ohio, Tennessee, Texas, Virginia, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, New Hampshire, New Jersey, New Mexico, North Carolina, North Dakota, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for pregnancy testing. The values range from 15% to 60%. Alabama, Alaska, Arizona, Georgia, Hawaii, Idaho, Kansas, Louisiana, Nevada, Ohio, Tennessee, Virginia, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Illinois, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, North Carolina, North Dakota, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Texas, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New Mexico, New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for condoms. The values range from 10% to 54%. Alabama, Alaska, Arizona, Arkansas, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Louisiana, Michigan, Mississippi, Missouri, Nebraska, Nevada, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, range from 0% to 24%. California, Connecticut, Delaware, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Montana, New Hampshire, New Jersey, New Mexico, Oregon, South Dakota, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for condom-compatible lubricants. The values range from 8% to 53%. Alabama, Alaska, Arizona, Arkansas, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Louisiana, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Virginia, Wisconsin, range from 0% to 24%. California, Connecticut, Delaware, Kentucky, Maine, Maryland, Massachusetts, Minnesota, New Hampshire, New Mexico, Oregon, Vermont, Washington, West Virginia, Wyoming, range from 25% to 49%. New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for contraceptives other than condoms. The values range from 10% to 55%. Alabama, Alaska, Arizona, Arkansas, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Louisiana, Michigan, Mississippi, Missouri, Nebraska, Nevada, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, range from 0% to 24%. California, Connecticut, Delaware, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Montana, New Hampshire, New Jersey, New Mexico, Oregon, South Dakota, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for prenatal care. The values range from 11% to 58%. Alabama, Alaska, Arizona, Georgia, Hawaii, Idaho, Illinois, Kansas, Louisiana, Nevada, Ohio, Tennessee, Virginia, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, North Carolina, North Dakota, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Texas, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New York, Rhode Island, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for HPV vaccine administration. The values range from 18% to 66%. Alabama, Arizona, Florida, Georgia, Hawaii, range from 0% to 24%. Alaska, Arkansas, California, Connecticut, Delaware, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. New Mexico, New York, Rhode Island, West Virginia, range from 50% to 74%. Colorado and Iowa did not have weighted results. Utah did not ask this question.


*To any organizations or health care professionals not on school property.



This slide shows the percentage of secondary schools that do not require parental consent or parental notification when sexual or reproductive health services are provided by the school. The values range from 0% to 6%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that do not require parental consent or parental notification when sexual or reproductive health services are referred by the school. The values range from 0% to 20%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that do not require parental consent but may notify parents when sexual or reproductive health services are provided by the school*. The values range from 0% to 26%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Delaware, range from 25% to 49%. Colorado and Iowa did not have weighted results.

*Responses to this question are not indicative of compliance with states' minors' right to consent for confidential services laws.




This slide shows the percentage of secondary schools that do not require parental consent but may notify parents when sexual or reproductive health services are referred by the school*. The values range from 4% to 39%. Alabama, Alaska, Arizona, Arkansas, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Virginia, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. California, Delaware, Maine, Massachusetts, Minnesota, New Mexico, New York, Oregon, Rhode Island, Vermont, Washington, range from 25% to 49%. Colorado and Iowa did not have weighted results.

*Responses to this question are not indicative of compliance with states' minors' right to consent for confidential services laws.




This slide shows the percentage of secondary schools that identify and track students with chronic conditions* that may require daily or emergency management. The values range from 85% to 100%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*For example, asthma or food allergies.




This slide shows the percentage of secondary schools that have protocols that ensure students with a chronic condition that may require daily or emergency management are enrolled into private, state, or federally funded insurance programs if eligible. The values range from 44% to 81%. Alaska, North Dakota, range from 25% to 49%. Alabama, Arizona, Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Mexico, New York, North Carolina, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Delaware, Maryland, Massachusetts, New Hampshire, New Jersey, Oregon, Vermont, West Virginia, range from 75% to 100%. Colorado and Iowa did not have weighted results.



This slide shows the percentage of secondary schools that provide referrals to community-based medical care providers for students identified with chronic conditions or at risk for activity, diet, and weight-related chronic conditions. The values range from 27% to 100%. Alabama, Alaska, Arizona, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Michigan, Nevada, North Dakota, Ohio, Oklahoma, South Dakota, Tennessee, Utah, Virginia, Wisconsin, range from 25% to 49%. Arkansas, California, Connecticut, Delaware, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, North Carolina, Oregon, Pennsylvania, Rhode Island, South Carolina, Washington, West Virginia, Wyoming, range from 50% to 74%. New York, Texas, Vermont, range from 75% to 100%. Colorado and Iowa did not have weighted results.

School Health Coordination


School Health Coordination




This slide shows the percentage of secondary schools that had a school health council, committee, or team that offers guidance on the development of policies or coordinates activities on health topics. The values range from 36% to 82%. Alabama, Alaska, Arizona, California, Georgia, Idaho, Illinois, Nevada, North Carolina, North Dakota, Ohio, Oregon, South Dakota, Utah, Washington, range from 25% to 49%. Connecticut, Delaware, Florida, Hawaii, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Jersey, New Mexico, New York, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Virginia, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Arkansas, New Hampshire, Vermont, range from 75% to 100%. Colorado and Iowa did not have weighted results.

Health Education


Health Education



This slide shows the percentage of secondary schools that required students to take 2 or more health education courses. The values range from 9% to 85%. Alabama, Arizona, California, Florida, Kentucky, New Mexico, range from 0% to 24%. Alaska, Georgia, Hawaii, Kansas, Louisiana, Michigan, Mississippi, Nevada, Ohio, South Carolina, South Dakota, Tennessee, Utah, Washington, range from 25% to 49%. Arkansas, Connecticut, Delaware, Idaho, Illinois, Indiana, Maine, Maryland, Massachusetts, Missouri, Nebraska, New Hampshire, New York, North Carolina, North Dakota, Pennsylvania, Vermont, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Minnesota, Montana, New Jersey, Oregon, Rhode Island, Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.




This slide shows the pmong secondary schools that required a health education course, percentage that required students who fail the course to repeat it. The values range from 37% to 89%. California, Maryland, Massachusetts, New Jersey, South Carolina, Vermont, Virginia, West Virginia, range from 25% to 49%. Arizona, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Indiana, Kentucky, Louisiana, Maine, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New York, North Carolina, North Dakota, Oregon, Pennsylvania, Rhode Island, Tennessee, Utah, Washington, Wisconsin, Wyoming, range from 50% to 74%. Alabama, Alaska, Arkansas, Idaho, Kansas, Mississippi, New Mexico, Ohio, South Dakota, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.



This slide shows the percentage of secondary schools in which the lead health education teacher had professional preparation in health education or in health and physical education combined. The values range from 18% to 89%. Alaska, California, range from 0% to 24%. Alabama, Arizona, Florida, Hawaii, Missouri, New Mexico, South Carolina, Vermont, Washington, range from 25% to 49%. Connecticut, Delaware, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Mississippi, Montana, Nebraska, Nevada, New Hampshire, New York, North Carolina, North Dakota, Ohio, Oregon, South Dakota, Tennessee, Utah, Wisconsin, Wyoming, range from 50% to 74%. Arkansas, Georgia, Minnesota, New Jersey, Pennsylvania, Rhode Island, Virginia, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.


Tobacco-Use Prevention

Tobacco-Use Prevention




This slide shows the percentage of secondary schools that prohibit all tobacco use at all times in all locations*. The values range from 38% to 80%. Alaska, Idaho, Kansas, Kentucky, Missouri, Nebraska, Ohio, Tennessee, Wyoming, range from 25% to 49%. Alabama, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Indiana, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Dakota, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. North Carolina, Texas, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Prohibit the use of all tobacco, including cigarettes, smokeless tobacco, cigars, and pipes, by students, faculty and school staff, and visitors, in school buildings, outside on school grounds, on school buses or other vehicles used to transport students, and at off-campus, school-sponsored events, during school hours and non-school hours.




This slide shows the percentage of secondary schools that prohibit the use of electronic vapor products* by students during any school-related activity. The values range from 68% to 98%. Vermont, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Such as e-cigarettes, vape pipes, and hookah pens.




This slide shows the percentage of secondary schools that taught 19 key tobacco-use prevention topics in a required course*. The values range from 15% to 76%. Arizona, range from 0% to 24%. Alaska, California, Connecticut, Florida, Hawaii, Idaho, Illinois, Kansas, Maine, Massachusetts, Michigan, Minnesota, Nebraska, New Hampshire, New Mexico, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. Alabama, Arkansas, Delaware, Georgia, Indiana, Kentucky, Louisiana, Maryland, Mississippi, Missouri, Montana, Nevada, New Jersey, New York, North Carolina, South Dakota, range from 50% to 74%. West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.



This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on tobacco-use prevention. The values range from 8% to 57%. Alabama, Arizona, California, Georgia, Hawaii, Idaho, Indiana, Kansas, Maine, Massachusetts, Minnesota, Missouri, Nebraska, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Virginia, Washington, Wyoming, range from 0% to 24%. Alaska, Arkansas, Connecticut, Delaware, Florida, Illinois, Kentucky, Louisiana, Maryland, Michigan, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Tennessee, Utah, Vermont, West Virginia, Wisconsin, range from 25% to 49%. Mississippi, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.


*During the 2 years before the survey.



This slide shows the percentage of secondary schools that posted signs marking a tobacco-free school zone. The values range from 43% to 98%. Connecticut, range from 25% to 49%. Delaware, Hawaii, Idaho, Illinois, Kentucky, Maryland, Massachusetts, Michigan, Missouri, Nevada, New Jersey, New Mexico, Ohio, Pennsylvania, South Carolina, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Florida, Georgia, Indiana, Kansas, Louisiana, Maine, Minnesota, Mississippi, Montana, Nebraska, New Hampshire, New York, North Carolina, North Dakota, Oklahoma, Oregon, Rhode Island, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.


Nutrition

Nutrition




This slide shows the percentage of secondary schools that did not sell less healthy foods and beverages* in vending machines, canteen, or snack bar. The values range from 34% to 89%. Alabama, Georgia, Idaho, Indiana, Kansas, Louisiana, Minnesota, Missouri, Montana, Nebraska, Nevada, New York, North Dakota, South Carolina, South Dakota, Utah, Washington, Wisconsin, Wyoming, range from 25% to 49%. Arizona, California, Delaware, Florida, Illinois, Kentucky, Maine, Maryland, Massachusetts, Michigan, Mississippi, New Hampshire, New Jersey, New Mexico, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, Tennessee, Texas, Vermont, Virginia, range from 50% to 74%. Alaska, Arkansas, Connecticut, Hawaii, West Virginia, range from 75% to 100%. Colorado and Iowa did not have weighted results.


*Baked goods not low in fat, salty snacks not low in fat, chocolate candy, other kinds of candy, soda pop or fruit drinks that are not 100% juice, and sports drinks. During the current school year.



This slide shows the percentage of secondary schools in which students could not purchase chocolate candy from vending machines or at the school store, canteen, or snack bar. The values range from 68% to 98%. Louisiana, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase other kinds of candy from vending machines or at the school store, canteen, or snack bar. The values range from 54% to 98%. Georgia, Idaho, Louisiana, Utah, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Hawaii, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase salty snacks that are not low in fat* from vending machines or at the school store, canteen, or snack bar. The values range from 65% to 96%. Idaho, Louisiana, Michigan, New Jersey, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.


*Such as regular potato chips.



This slide shows the percentage of secondary schools in which students could not purchase cookies, crackers, cakes, pastries, or other baked goods that are not low in fat from vending machines or at the school store, canteen, or snack bar. The values range from 68% to 97%. Idaho, Louisiana, Minnesota, New Jersey, New York, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase soda pop or fruit drinks that are not 100% juice from vending machines or at the school store, canteen, or snack bar. The values range from 61% to 97%. Idaho, Indiana, Montana, Tennessee, Utah, Virginia, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Texas, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase sports drinks* from vending machines or at the school store, canteen, or snack bar. The values range from 39% to 92%. Montana, Nevada, South Carolina, range from 25% to 49%. Alabama, Arizona, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Dakota, Tennessee, Texas, Utah, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Alaska, Arkansas, California, Connecticut, Hawaii, Massachusetts, Rhode Island, Vermont, West Virginia, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Such as Gatorade.




This slide shows the percentage of secondary schools in which students could not purchase energy drinks* from vending machines or at the school store, canteen, or snack bar. The values range from 93% to 100%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.


*Such as Red Bull or Monster.



This slide shows the percentage of secondary schools in which students could not purchase foods or beverages containing caffeine from vending machines or at the school store, canteen, or snack bar. The values range from 65% to 99%. Idaho, Indiana, Missouri, Montana, Tennessee, Utah, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that allowed students to purchase fruits and vegetables from vending machines or at the school store, canteen, or snack bar. The values range from 3% to 39%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Florida, Michigan, New Hampshire, New Jersey, Rhode Island, Texas, range from 25% to 49%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that prohibit all forms of advertising and promotion of candy, fast food restaurants, or soft drinks in all locations*. The values range from 26% to 75%. Alabama, Arkansas, Delaware, Georgia, Illinois, Indiana, Kentucky, Michigan, Missouri, Montana, Nebraska, North Carolina, Ohio, Oklahoma, Oregon, South Carolina, South Dakota, Tennessee, Virginia, Wisconsin, Wyoming, range from 25% to 49%. Alaska, Arizona, California, Connecticut, Florida, Hawaii, Idaho, Kansas, Louisiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New Mexico, North Dakota, Pennsylvania, Rhode Island, Texas, Utah, Vermont, Washington, West Virginia, range from 50% to 74%. New York, range from 75% to 100%. Colorado and Iowa did not have weighted results.


*In school buildings; on school grounds, including on the outside of the school building, on playing fields, or other areas of the campus; on school buses or other vehicles used to transport students; and in school publications.




This slide shows the percentage of secondary schools that priced nutritious foods and beverages at a lower cost while increasing the price of less nutritious foods and beverages. The values range from 4% to 21%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results.



This slide shows the percentage of secondary schools that provided information to students or families on the nutrition and caloric content of foods available. The values range from 33% to 74%. Alaska, Arkansas, Hawaii, Idaho, Louisiana, Montana, Nevada, New Mexico, New York, North Dakota, Oregon, Utah, Wyoming, range from 25% to 49%. Alabama, Arizona, California, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, North Carolina, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that placed fruits and vegetables near the cafeteria cashier, where they are easy to access. The values range from 42% to 93%. Alaska, range from 25% to 49%. Alabama, Arizona, Arkansas, Georgia, Hawaii, Idaho, Illinois, Kansas, Louisiana, Missouri, Montana, Nevada, New Mexico, North Carolina, North Dakota, Oklahoma, Pennsylvania, South Dakota, Wisconsin, Wyoming, range from 50% to 74%. California, Connecticut, Delaware, Florida, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Nebraska, New Hampshire, New Jersey, New York, Ohio, Oregon, Rhode Island, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that allow students to have access to drinking water*. The values range from 45% to 87%. Illinois, New Jersey, Pennsylvania, Rhode Island, range from 25% to 49%. Alabama, Alaska, Arkansas, Connecticut, Delaware, Georgia, Idaho, Indiana, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, New Hampshire, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Arizona, California, Florida, Hawaii, Kansas, Nebraska, Nevada, Oklahoma, range from 75% to 100%. Colorado and Iowa did not have weighted results.


*Permit students to have a drinking water bottle with them during the school day in at least certain locations and offer a free source of drinking water in 5 locations.




This slide shows the percentage of secondary schools that offer fruits or non-fried vegetables when foods or beverages are offered at school celebrations. The values range from 19% to 62%. Kansas, Missouri, Nebraska, North Dakota, South Dakota, range from 0% to 24%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kentucky, Louisiana, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Montana, Nevada, New Jersey, New Mexico, New York, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. Delaware, Maine, New Hampshire, Vermont, range from 50% to 74%. Colorado and Iowa did not have weighted results.



This slide shows the percentage of secondary schools that prohibit school staff from giving students food or food coupons as a reward for good behavior or good academic performance. The values range from 12% to 69%. Alaska, Georgia, Idaho, Illinois, Louisiana, Montana, Nebraska, New York, Ohio, Oregon, South Dakota, Utah, Virginia, Washington, Wisconsin, Wyoming, range from 0% to 24%. Alabama, Arizona, California, Connecticut, Delaware, Florida, Hawaii, Indiana, Kansas, Kentucky, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, range from 25% to 49%. Arkansas, Massachusetts, New Hampshire, Texas, West Virginia, range from 50% to 74%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that prohibit less nutritious foods and beverages from being sold for fundraising purposes. The values range from 20% to 71%. Nebraska, range from 0% to 24%. Alabama, Alaska, Arizona, Arkansas, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. California, Connecticut, Massachusetts, New Hampshire, Rhode Island, South Carolina, West Virginia, range from 50% to 74%. Colorado and Iowa did not have weighted results.



This slide shows the percentage of secondary schools in which the lead health education teacher received professional development* on nutrition and dietary behavior. The values range from 16% to 65%. Alaska, California, Georgia, Hawaii, Idaho, Indiana, Missouri, Oregon, Rhode Island, Wyoming, range from 0% to 24%. Alabama, Arizona, Connecticut, Delaware, Florida, Illinois, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Montana, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Pennsylvania, South Carolina, South Dakota, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Arkansas, Mississippi, New Hampshire, Tennessee, range from 50% to 74%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the 2 years before the survey.




This slide shows the percentage of secondary schools that taught 20 key nutrition and dietary behavior topics in a required course*. The values range from 29% to 79%. Alaska, Arizona, California, Connecticut, Hawaii, Kansas, Maine, Massachusetts, Michigan, Minnesota, Oregon, South Carolina, Vermont, Washington, Wisconsin, range from 25% to 49%. Alabama, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kentucky, Louisiana, Maryland, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Pennsylvania, Rhode Island, South Dakota, Tennessee, Utah, Virginia, Wyoming, range from 50% to 74%. Arkansas, Mississippi, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.


*During the current school year.

Physical Education And Physical Activity

Physical Education and Physical Activity




This slide shows the percentage of secondary schools that have established, implemented, or evaluated a Comprehensive School Physical Activity Program. The values range from 1% to 14%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 6th grade*. The values range from 72% to 100%. Hawaii, Michigan, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 7th grade*. The values range from 51% to 100%. Hawaii, Michigan, Nevada, Oklahoma, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 8th grade*. The values range from 53% to 100%. Hawaii, Michigan, New Mexico, Oklahoma, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 9th grade*. The values range from 26% to 100%. Oklahoma, range from 25% to 49%. Arizona, Idaho, range from 50% to 74%. Alabama, Alaska, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 10th grade*. The values range from 13% to 100%. Kansas, Kentucky, North Carolina, Oklahoma, range from 0% to 24%. Arizona, Georgia, Michigan, Nebraska, South Carolina, South Dakota, range from 25% to 49%. Alabama, Arkansas, Florida, Hawaii, Idaho, Indiana, Maryland, Missouri, New Hampshire, New Mexico, North Dakota, Ohio, Oregon, Tennessee, Washington, West Virginia, Wyoming, range from 50% to 74%. Alaska, California, Connecticut, Delaware, Illinois, Louisiana, Maine, Massachusetts, Minnesota, Mississippi, Montana, Nevada, New Jersey, New York, Pennsylvania, Rhode Island, Texas, Utah, Vermont, Virginia, Wisconsin, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 11th grade*. The values range from 5% to 100%. California, Hawaii, Indiana, Kansas, Kentucky, Minnesota, Montana, Nebraska, North Carolina, North Dakota, Oklahoma, Virginia, range from 0% to 24%. Arizona, Delaware, Florida, Georgia, Idaho, Maine, Maryland, Michigan, Missouri, Ohio, Oregon, South Carolina, South Dakota, Tennessee, West Virginia, Wyoming, range from 25% to 49%. Alabama, Arkansas, Connecticut, Louisiana, Nevada, New Hampshire, New Mexico, Texas, Utah, Vermont, Washington, Wisconsin, range from 50% to 74%. Alaska, Illinois, Massachusetts, Mississippi, New Jersey, New York, Pennsylvania, Rhode Island, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 12th grade*. The values range from 6% to 100%. California, Hawaii, Indiana, Kansas, Kentucky, Minnesota, Montana, Nebraska, North Carolina, North Dakota, Oklahoma, Virginia, Wyoming, range from 0% to 24%. Arizona, Delaware, Florida, Georgia, Idaho, Maine, Maryland, Michigan, Missouri, Nevada, New Hampshire, New Mexico, Ohio, Oregon, South Carolina, South Dakota, Tennessee, Utah, Washington, West Virginia, Wisconsin, range from 25% to 49%. Alabama, Arkansas, Connecticut, Louisiana, Texas, Vermont, range from 50% to 74%. Alaska, Illinois, Massachusetts, Mississippi, New Jersey, New York, Pennsylvania, Rhode Island, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which at least 1 physical education teacher or specialist received professional development on physical education*. The values range from 39% to 98%. Alaska, range from 25% to 49%. Arizona, Hawaii, Idaho, Michigan, New Mexico, Oklahoma, Oregon, Rhode Island, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Pennsylvania, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.

*During the year before the survey.




This slide shows the percentage of secondary schools in which students participate in physical activity breaks in classrooms during the school day*. The values range from 28% to 82%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New York, North Dakota, Ohio, Pennsylvania, Rhode Island, South Carolina, South Dakota, Texas, Utah, Virginia, Washington, range from 25% to 49%. Arizona, Hawaii, Louisiana, Maine, Maryland, Nebraska, New Hampshire, New Jersey, New Mexico, North Carolina, Oklahoma, Oregon, Tennessee, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Alaska, Vermont, range from 75% to 100%. Colorado and Iowa did not have weighted results.


*Outside of physical education.



This slide shows the percentage of secondary schools that offer opportunities for all students to participate in intramural sports programs or physical activity clubs. The values range from 32% to 86%. Arkansas, Kansas, Nebraska, North Dakota, South Dakota, Wyoming, range from 25% to 49%. Alabama, Arizona, Delaware, Georgia, Idaho, Illinois, Indiana, Kentucky, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Montana, New Mexico, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, Tennessee, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. Alaska, California, Connecticut, Florida, Hawaii, Maine, Maryland, Massachusetts, Nevada, New Hampshire, New Jersey, New York, Rhode Island, Vermont, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that offer interscholastic sports to students. The values range from 69% to 95%. Hawaii, Louisiana, Maryland, Utah, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado and Iowa did not have weighted results.




This slide shows the percentage of secondary schools that offer opportunities for students to participate in physical activity before the school day*. The values range from 18% to 64%. Alabama, Connecticut, Delaware, Louisiana, Mississippi, North Carolina, Virginia, range from 0% to 24%. Arkansas, California, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Missouri, Nevada, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Vermont, Washington, West Virginia, Wyoming, range from 25% to 49%. Alaska, Arizona, Minnesota, Montana, Nebraska, North Dakota, Texas, Utah, Wisconsin, range from 50% to 74%. Colorado and Iowa did not have weighted results.

*Through organized physical activities or access to facilities or equipment for physical activity.



This slide shows the percentage of secondary schools that taught 13 key physical activity topics in a required course*. The values range from 37% to 86%. Alaska, Arizona, range from 25% to 49%. Alabama, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Nebraska, Nevada, New Hampshire, New Mexico, New York, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Arkansas, Indiana, Mississippi, Missouri, Montana, New Jersey, North Carolina, West Virginia, range from 75% to 100%. Colorado, Iowa, Oklahoma, and Texas did not have weighted results.

*During the current school year.



This slide shows the percentage of secondary schools that have a joint use agreement for shared use of physical activity facilities. The values range from 47% to 79%. Delaware, Indiana, Mississippi, Ohio, Oklahoma, range from 25% to 49%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Utah, range from 75% to 100%. Colorado and Iowa did not have weighted results.