



West Nile Virus Disease Cases* and Presumptive Viremic Blood Donors by State – United States, 2013

State	Neuroinvasive Disease Cases†	Non-neuroinvasive Disease Cases	Total cases	Deaths	Presumptive Viremic Blood Donors‡
Alabama	3	6	9	0	2
Arizona	50	12	62	6	6
Arkansas	16	2	18	1	6
California	237	142	379	15	68
Colorado	90	232	322	7	22
Connecticut	1	3	4	0	1
Delaware	3	0	3	0	0
District of Colombia	0	1	1	0	0
Florida	5	2		0	1
Georgia	4	6	10	0	3
Idaho	14	26	40	2	12
Illinois	86	31	117	12	21
Indiana	19	4	23	2	3
lowa	24	20	44	0	12
Kansas	34	57	91		35
Kentucky	1	2	3	1	0
Louisiana	34	20	<u>5</u>	4	5
Maryland	11	5	16	1	1
Massachusetts	7	1	8	0	2
Michigan	24	12	36	2	4
Minnesota	31	48	<u> </u>	3	26
Mississippi	27	18	45	<u>5</u>	5
Missouri	24	5	29	2	8
Montana	10	28	38	2	5
Nebraska	54	172	226	5	38
Nevada	8	3	11	<u>5</u> 1	5
New Hampshire	<u>8</u> 1	0	1	0	0
	10	2	12	2	3
New Jersey New Mexico	24	14	38	3	2
New York	18	14	32	0	9
North Carolina	3	0	3	1	0
North Dakota	64	61	125	2	15
	21	3		3	
Ohio			24	<u>3</u> 	6
Oklahoma	60 7	<u>29</u> 9	89	0	18 0
Oregon			<u>16</u>		-
Pennsylvania Phodo Island	6	5	11	1	0
Rhode Island	<u>1</u> 3	0	<u> </u>	0	<u>0</u> 2
South Carolina		4		0	
South Dakota	57	92	149	3	26
Tennessee	17	7	24	2	0
Texas	113	70	183	14	39
Utah	4	3	7	0	5
Vermont	1	1	2	0	0
Virginia	6	0	6	0	0
Washington	0	1	1	0	1
West Virginia	1	0	1	0	0
Wisconsin	17	4	21	2	9
Wyoming	16	25	41	1	5
Totals	1,267	1,202	2,469	119	431

^{*}Includes cases reported as meningitis, encephalitis, or acute flaccid paralysis.

‡Presumptive viremic blood donors (PVDs) are people who had no symptoms at the time of donating blood through a blood collection agency, but whose blood tested positive when screened for the presence of West Nile virus. Some PVDs develop symptoms after donation; of the 431 PVDs, 60 (14%) developed clinical illness and are also included as disease cases.

[†]Includes confirmed and probable cases.