



West Nile virus disease cases and presumptive viremic blood donors reported to ArboNET, United States, 2011

State	Neuroinvasive	Non-neuroinvasive	Total		Presumptive viremic
	disease cases*	disease cases	cases†	Deaths	blood donors‡
Alabama	5	0	5	1	0
Arizona	49	20	69	4	13
Arkansas	1	0	1	0	0
California	110	48	158	9	27
Colorado	2	5	7	0	1
Connecticut	8	1	9	0	0
Delaware	1	0	1	0	0
District of Columbia	10	5	15	0	0
Florida	20	4	24	2	3
Georgia	14	8	22	3	4
Idaho	1	2	3	0	0
Illinois	22	12	34	3	7
Indiana	7	2	9	1	3
Iowa	5	4	9	0	0
Kansas	4	0	4	1	0
Kentucky	4	1	5	1	2
Louisiana	6	4	10	0	3
Maryland	10	9	19	0	3
Massachusetts	5	1	6	0	1
Michigan	32	2	34	2	
Minnesota	1	<u>-</u> 1	2		0
Mississippi	31	21	 52	5	8
Missouri	6	4	10	0	2
Montana	1	0	1	0	0
Nebraska	14	15	29	0	9
Nevada	12	4	16	2	3
New Jersey	2	5	7	0	1
New Mexico	4	0	4	0	2
New York	28	16	44	2	8
North Carolina	2	0	2	0	0
North Dakota	<u>2</u> 1	3	4	0	0
Ohio	10	11	21	1	9
Oklahoma	1	0	1	0	0
Pennsylvania	5	1	6	0	0
Rhode Island	1	0	1	0	0
South Dakota	0	2	2	0	0
Tennessee	16	2	18	2	0
Texas	20	7	27	2	16
Utah	1	2	3	0	1
Vermont	1	0	<u>5</u> 1	0	1
Virginia	8	0 1	9	1	9
	2	0	2	0	0
West Virginia Wisconsin	2	<u>0</u> 1	3	0	0
	<u>2</u> 1	2	3	0	0
Wyoming	486	226	712	U	U

^{*}Includes cases reported as meningitis, encephalitis, or acute flaccid paralysis.

[†]Includes confirmed and probable cases.

[‡]Presumptive viremic blood donors (PVDs) are people who had no symptoms at the time of donating blood through a blood collection agency, but whose blood tested positive when screened for the presence of West Nile virus. Some PVDs develop symptoms after donation; of the 137 PVDs, 33 (24%) developed clinical illness and are also included as disease cases.