

Table 4 (Supplemental Table). Characteristics of travelers who were immunized for influenza before travel and those who were not immunized\*

	<b>Immunized for influenza, n=454</b>	<b>Not immunized for influenza, n=170</b>	<b>PRR (95% CI)</b>	<b>P value</b>
Median age, years (range)	49 (19-83)	42 (20-82)	N/A	0.0009
<b>Duration of travel</b>	<b>(n, row%)</b>			
≥14 days	181 (76)	56 (24)	1.07 (0.98, 1.18)	0.15
<14 days	270 (71)	110 (29)		
<b>Birth country</b>				
Foreign-born (n=108)	73 (68)	35 (32)	0.92 (0.80-1.05)	0.19
U.S.-born (n=515)	380 (74)	135 (26)		
<b>Travel Reason**</b>				
<i>VFR</i> (n=16)	9 (56)	7 (44)	0.77 (0.5-1.19)	0.13
<i>Tourism/Vacation</i> (n=313)	229 (73)	84 (27)	1.01 (0.91-1.13)	0.80
<i>Business</i> (n=90)	65 (72)	25 (28)	0.99 (0.86-1.14)	0.90
<i>Education</i> (n=13)	10 (77)	3 (23)	1.05 (0.78-1.43)	1
<i>Volunteer/Missionary/Aid Work</i> (n=64)	48 (75)	16 (25)	1.04 (0.89-1.21)	0.67
<b>Destination grouping by UN Human Development Index †</b>				
Low HD (n=163)	119 (73)	44 (27)	N/A	0.89
Medium HD (n=318)	229 (72)	89 (28)		
High HD (n=143)	106 (74)	37 (26)		
<b>Experienced a health problem during or after travel (n=398)</b>	288 (72)	110 (28)	0.99 (0.89-1.09)	0.77
<b>Respiratory Symptoms† (n=143)</b>	100 (70)	43 (30)	0.95 (0.84-1.07)	0.39

\* Travelers were considered immunized if they received influenza vaccine prior to the BATMN pre-travel consultation or at the time of the BATMN pre-travel consultation

\*\* Excludes those who indicated multiple purposes of travel (n=126).

¥ Based on UN human development ranking (high + very high is combined); if traveler went to  $\geq 1$  country, the lowest-ranked country was used

† Anybody who experienced cough, sore throat, OR runny nose during or after travel