HEALTH STATISTICS

FROM THE U. S. NATIONAL HEALTH SURVEY

Diabetes

reported in interviews

United States July 1957 - June 1959

Statistics on prevalence of diabetes and associated disability by age, sex, and medical care status. Based on data collected in household interviews during the period July 1957-June 1959.

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE Arthur S. Flemming, Secretary

> Public Health Service Leroy E. Burney, Surgeon General

Division of Public Health Methods William H. Stewart, M. D., Chief

U. S. NATIONAL HEALTH SURVEY

Forrest E. Linder, Ph. D., Director Theodore D. Woolsey, Assistant Director Alice M. Waterhouse, M. D., Chief Medical Advisor James E. Kelly, D. D. S., Dental Advisor Walt R. Simmons, Statistical Advisor O. K. Sagen, Ph. D., Chief, Special Studies Philip S. Lawrence, Sc. D., Chief, Health Interview Survey Margery R. Cunningham, Staff Assistant Robert T. Little, Chief, Automatic Data Processing

The U. S. National Health Survey is a continuing program under which the Public Health Service makes studies to determine the extent of illness and disability in the population of the United States and to gather related information. It is authorized by Public Law 652, 84th Congress.

CO-OPERATION OF THE BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies. For the Health Interview Survey the Bureau of the Census designed and selected the sample, conducted the household interviews, and processed the data in accordance with specifications established by the Public Health Service.

Public Health Service Publication No. 584-B21

CONTENTS

3414

_

	Page
Source and Classification of Data	1
Prevalence of Diabetes According to Sex and Age	2
Diabetes According to Medical Care Status	3
Disability Due to Diabetes	.3
Summary	4
Detailed Tables	6
Appendix I. Technical Notes on Methods Background of This Report Statistical Design of the Health Interview	12 12
Survey General Qualifications Reliability of Estimates	12 13 13
Appendix II. Definitions of Certain Terms Used in This Report Terms Relating to Chronic Conditions Terms Relating to Disability Chronic Activity and Mobility Limitation Demographic Terms	16 16 16 17 17
Appendix III Questionnaire	10

SYMBOLS AND NOTES	•
Data not available (three dashes)	
Category not applicable (three dots)	•••
Quantity is zero (1 dash)	-
Magnitude greater than zero but less than one-half of the unit used	0 or 0.0
Magnitude of the sampling error precludes showing separate estimates	(*)
NOTE: Due to rounding detailed figures within tables may not add to totals	

DIABETES

The need to find and control cases of diabetes, and thus to prevent or to postpone disability and deaths due to it-rather than excessive prevalence-makes this chronic disease one of the major health problems in the United States today. Diabetes mellitus ranks far below such chronic conditions as heart disease, and hypertension in number of cases and deaths, but it is nevertheless eighth among the 10 leading causes of death, and fifth among diseases usually regarded as chronic, exceeded only by diseases of heart, cancer, vascular lesions of the central nervous system, and general arteriosclerosis, in that order. For the year ending December 1958, there were 27,500 deaths caused by diabetes mellitus, with a rate of 15.9 per 100,000 population.¹

The introduction of insulin in 1922, and continued research since then to improve it and other therapeutic measures fortunately have changed the status of diabetes from a disease with a very serious outlook to one that, if diagnosed early, in many cases can be controlled, thus permitting the patient to live practically a normal life.² In order to locate the diabetic or the potential diabetic, large-scale mass screening programs have been organized and conducted over the past 20 years by Federal and local health agencies and such private organizations as the American Diabetes Association. The results of these efforts, and of such studies as that of Wilkerson and Krall³, have led investigators to the opinion that the number of unsuspected cases of diabetes in this country may be equal to the known cases or very nearly so.

The volume of medical literature on all phases of this disease is enormous. Much has been published on the statistics of diabetes, in terms of both the known and the undiscovered cases. Prevalence estimates based on the results of physical examinations and clinical records will obviously differ from estimates from the findings in household interview surveys. Objectives, techniques, definitions, and resulting estimates may vary considerably even between studies of the same general class and time period. It is to be expected that

This report was prepared by Louise E. Bollo, of the U. S. National Health Survey staff.

differences should exist between estimates obtained from records prepared by trained clinical examiners and those resulting from responses given by household members who may or may not be co-operative and well-informed.

Diabetes prevalence data included here can of course deal only with those cases: (1) which have been diagnosed; (2) about which the family has been informed; and (3) which are believed by the family to have been present in the year prior to the interview.

SOURCE AND CLASSIFICATION OF DATA

Data on diabetes presented in this report are based primarily on replies to four "illness-recall" questions:

- 1. Were you sick at any time last week or the week before?
- 2. Last week or the week before did you take any medicine or treatment for any condition?
- 3. At the present time do you have any ailments or conditions that have lasted for a long time? (If "No") Even though they don't bother you all the time?
- 4. Has anyone in the family...had any of these conditions during the past 12 months? (Interviewer reads list of major chronic conditions which includes "diabetes.")

Positive responses may come from any one or more of the four questions. The unduplicated positive replies serve as the basis for estimates of the prevalence of diabetes in the population. Because these estimates are based on a continuing sample of household interviews throughout a 24-month period July 1957-June 1959, the prevalence is actually the average number of cases of diabetes reported by the people during that period.

Additional questions are asked to obtain information about medical attention and disability. Replies to these are used to establish the extent to which diabetics have medical care and are or are not restricted or limited in their daily living.

Reports of "diabetes" were assumed to mean "diabetes mellitus," and were classified under category 260 (diabetes mellitus), of the International Classification of Diseases, 1955 Revision. It is possible that cases such as "renal diabetes," "bronze diabetes," or "diabetes insipidus" could have been reported only as "diabetes," but these conditions are too infrequent to have any appreciable effect on the results. However, if the respondent knew and reported such modifiers as "renal," "bronze," "insipidus," the appropriate codes for these would have been selected. Terms such as "sugar in urine," or "sugar in blood," were not counted as cases of diabetes unless diabetes was also mentioned.

Impairments and certain other conditions reported to be due to diabetes were recorded separately under their own code numbers, but the code 260 was also applied. Minor symptoms due to diabetes were coded only to the diabetes category. Code 260 was assigned only once per person with diabetes and is the only condition code included in these data. Therefore the total cases with mention of diabetes, complicated or uncomplicated, represent the total number of diabetics in the survey.

Terminology used by the household members did not constitute a problem because if they knew and reported that they had diabetes they used the single term "diabetes" in nearly all cases. The matter of self-diagnosis is considered to be a minor source of error because the layman is very unlikely to believe that he has diabetes unless he has been told by a medical person that he has this condition.

A description of the statistical design of the household survey and general qualifications of the data presented in this report is given in Appendix I. Particular attention is called to the section in Appendix I on Reliability of Estimates, which includes tables of sampling errors and instruction for their use. Explanations and definitions of special terms and concepts are presented in Appendix II.

PREVALENCE OF DIABETES ACCORDING TO SEX AND AGE

Based on U. S. National Health Survey data collected during the period July 1957-June 1959, the prevalence of diabetes in the United States is estimated to be 1½ million cases, a rate of 9 cases per 1,000 population. This figure probably represents a slight underestimate of the known cases since it does not include the diabetics living in old-age homes and other resident institutions.

The 1935-36 National Health Survey provided an estimate of roughly 3 known cases of diabetes per 1,000 population. On the basis of this rate a national estimate of 660,000 cases as of 1937 was made.⁴ Reasons for the large increase in cases of diabetes over the past 20-25 years undoubtedly include the continuing national concentration upon case-finding, mentioned earlier in this report, the increase in the population particularly of the older age groups, the increased longevity of diabetics, and the possibility that physicians are now diagnosing as diabetes quite mild cases, based on laboratory findings, which in 1935 would not have been called diabetes.

As indicative of the prevalence rates found in studies conducted by means of clinical examinations of samples of the population, the previously mentioned study of Wilkerson and Krall³ in Oxford, Massachusetts, in 1946 and 1947, and the studies conducted under the sponsorship of the Commission on Chronic Illness in Baltimore, Maryland,⁵ and Hunterdon County, New Jersey, (1953-54)⁶ may be cited. In Oxford, a prevalence rate of 17 per 1,000 population was found. The Baltimore and Hunterdon County studies yielded rates of 27 per 1,000 and 10 per 1,000, respectively. In each instance the prevalence rate included, of course, both the previously known cases and those discovered during the study.

The estimate of 9 cases per 1,000 population produced by the health-interview phase of the present National Health Survey is only slightly higher than the rates found in two other recent interview surveys: 8.7 per 1,000 in the 1955-57 Hagerstown, Maryland, study⁷ and 7.7 per 1,000 in the Kansas City Metropolitan Area Health Survey (Dec. 1954-Jan. 1957).⁸

A lower rate of 5.5 per 1,000 was reported in the California Health Survey (1954-55).⁹ Remein,¹⁰ who estimated the prevalence of known diabetes in the United States for the year 1958 to be 8.8 per 1,000 population, has commented upon this low California rate as follows: "The relationship between mortality and prevalence is by no means a sure one, but the mortality rate from diabetes in California is only two thirds that of the United States as a whole. It is therefore conceivable that the prevalence of known diabetes is considerably less in California than in the nation as a whole."

Diabetes is essentially a disease of middle and old age, as indicated in figure 1 and table 1. Two thirds of all diabetics reported in the survey were 55 years of age or older. Females outnumber males in every age group over 45, most noticeably at ages 65-74 (table A). The rate per 1,000 persons drops for both sexes after 74 years of age, but more sharply for females than for males of this age; it is probable that the exclusion of patients in resident institutions accounts for some of this drop.

The survey estimate of the prevalence of diabetes among persons in the younger age groups, particularly those under 25, should be interpreted with great caution. It is generally recognized that diabetes is rare among young people, but the survey figure of 0.9 per 1,000 persons under 25 years of age could be quite unrealistic because of errors of sampling.



Figure 1. Number of persons reported to have diabetes per 1,000 population by sex and age.

Table A. Prevalence of known cases of diabetes per 1,000 population and ratio of females to males as reported in interviews: United States, July 1957-June 1959

Age	Preva per l popul	lence ,000 ation	Ratio of	
	Fe- male	Male	to males.	
All ages	10.0	8.0	1.25	
0-24 25-44 45-54 55-64 65-74 75+	(*) 3.9 13.7 31.5 50.3 38.8	(*) 4.9 11.2 25.2 34.4 31.5	(*) 0.79 1.22 1.25 1.46 1.23	

DIABETES ACCORDING TO MEDICAL CARE STATUS

Table 1 shows that practically all cases had been attended by a physician at some time, and that a very large proportion (about 92 percent) were "under care" at the time of the interview (table B). Persons "under care" are those for whom a positive response was obtained to the question "Do you still take any medicine or treatment that the doctor prescribed for diabetes? Or follow any advice he gave?" The small percent of all diabetics who replied "no" were those who may not have fully understood the question, or those who may have been feeling well at the time and were risking the cessation of control measures.

Table B. Number and percent of cases of diabetes under medical care: United States, July.1957-June 1959

	Total cases	Cases under medical care		
`Age	di- abetes (in th	Number noușands)	Per- cent	
All ages	1,530	1,404	91.8	
0-24 25-44 45-54 55-64 65-74 75+	67 199 246 424 416 177	61 181 222 392 388 159	91.0 91.0 90.2 92.5 93.3 89.8	

DISABILITY DUE TO DIABETES

There are a number of factors inherent in the nature of this disease, its age and sex distribution and its potential for control, which are likely to make it a relatively minor cause of disability, especially when institutionalized cases are excluded. The survey findings point out the considerable extent to which many diabetics regard themselves as <u>not</u> restricted or limited because of this condition. It is possible, of course, that some who had disabling complications of diabetes—such as failing vision or nephritis—attributed their disability to the complication only, and not to the underlying diabetes.

The 1½ million cases of diabetes were responsible for less than 2 million work-loss days among the "usually working" population, with about 1 million of these days among males, and 705,000 among females. This low figure is not surprising, since the majority of diabetics, sick or well, are not members of the "usually working" population either because of their age, retirement status, or because they are females keeping house.

Ninety percent of the diabetics indicated that they had no chronic limitation of mobility due to diabetes. Even among diabetics over 65 years of

age (some 593,000 persons), 82 percent considered that this disease did not limit their ability to get around freely in the house or outside.

Seventy-five percent of the total reported that their diabetes did not affect the kinds or amounts of their day-to-day activities—working, keeping house, going to school, or whatever else that usually occupied them (table 2). Females over 65 years of age had the greatest amount of continuing limitation of this kind.

Table 3 indicates that 85 percent of all diabetics had had no bed-disability days, because of diabetes, in the 12 months prior to the interview.

Disability attributable to diabetes in terms of the <u>number of days</u> is shown in table 4. A day of restricted activity, by definition, is any day on which a person had to cut down on his usual activities for a whole day because of his condition whether or not he was confined to bed. Diabetes was responsible for about 30 million restrictedactivity days of which 13 million were bed-disability days. When distributed among all diabetics the average number of restricted-activity days per case per year due to diabetes was about 20, with the average number of bed-days being 8.5.

Figure 2 shows the distribution of bed-days due to diabetes among all diabetics by age and sex. Females 65 years of age and older experienced more bed disability than any of the other age-sex groups.



Figure 2. Average annual number of bed-disability days due to diabetes per diabetic by sex and age.

It is perhaps more realistic to consider the number of bed-days due to diabetes in relation only to those diabetics who had actually experienced bed disability since it has been shown in table 3 that 85 percent of all diabetics had had <u>no</u> bed-disability days. When averaged among the 15 percent who had had any bed-days (one or more), the number of bed-days in a year due to diabetes becomes 56.2, with persons over 65 experiencing about 80 days per year (table C).

Table C.	Number	of be	ed-days	per	year	due
to diat	oetes per	r dia	abetic,	and	per	dia-
betic	having	one	or mon	ce b	oed-d	ays:
United	States,	July	1957-Ju	ine]	L959	•

	Number of bed-days due to diabetes			
Age	Per diabetic	Per diabetic having 1+ bed-days		
All ages	8.5	56.2		
0-44 45-54 55-64 65-74 75+	2.9 8.8 7.1 10.8 14.5	16.4 58.2 52.9 78.6 79.9		

SUMMARY

The average prevalence of known cases of diabetes, based on an estimate from the U.S. National Health Survey household interview sample, was a 1½ million cases, corresponding to a rate of 9 per 1,000 population. The rate per 1,000 persons was higher among females than among males in each age group over 45, most noticeably at ages 65-74 where the prevalence rate for females was 50.3, and for males, 34.4.

Virtually all cases had been attended by a physician at some time and about 92 percent were still under care by, or instruction from, a physician at the time of the interview. 1

Ninety percent of the reported cases showed no chronic mobility limitation, and 75 percent were not chronically restricted as to daily activities. Eighty-five percent of the diabetics had had no bed-disability days in the year prior to interview because of diabetes. However, among those who had been confined to bed for one or more days because of this condition, the average annual number of such days was 56.2, with persons over 65 experiencing about 80 days per year. ¹National Office of Vital Statistics: <u>Vital</u> <u>Statistics of the United States, 1958</u>, Vols. I and II. In press.

²American Diabetes Association, Inc.,: <u>Facts</u> <u>About Diabetes</u>. New York, N.Y., 1956.

³Wilkerson, H. L., and Krall, L. P.: Diabetes in a New England Town. <u>J. A. M. A</u>. 135: 209-216. Sept. 27, 1947.

⁴National Health Survey, 1935-36. The Magnitude of the Chronic Disease Problem in the United States. <u>Preliminary Reports, Sickness and Medical</u> <u>Care Series</u>. Bulletin 6, Public Health Service, Washington, D. C., 1938.

⁵Commission on Chronic Illness in 1953-54: <u>Chronic Illness in a Large City</u>: The Baltimore Study (Chronic Illness in the United States, Vol. IV). Harvard University Press, Cambridge, Mass., 1959. ⁶Commission on Chronic Illness: <u>Chronic Illness in a Rural Area</u>: The Hunterdon Study (Chronic Illness in the United States, Vol. III). Harvard University Press, Cambridge, Mass., 1959.

⁷West, M. D., and Altenderfer, M. E.: <u>Illness</u> and <u>Medical Care in Hagerstown</u>, <u>Maryland</u>. <u>I</u>. <u>The</u> <u>Prevalence of Chronic Disease in 1955-57 as Measured by Household Interviews</u>. Public Health Service, Washington, D. C., Oct. 1958.

⁸Peterson, W. A.: Kansas City Metropolitan Area Health Survey. <u>Community Studies, Inc</u>., Kansas City, Mo., June 1959.

⁹State of California, Department of Public Health: <u>Health in California</u>. California State Printing Office, Sacramento. Calif.

¹⁰Remein, Q.: <u>A Current Estimate of the Prevalence of Diabetes Mellitus in the United States.</u> Annals of the New York Academy of Sciences, Vol. 82. Article 2, Pages 229-235. Sept. 25, 1959.

DETAILED TABLES

			Page
Table 1	1.	Average prevalence of diabetes according to medical care status as reported in interviews and average prevalence per 1,000 population by sex and age: United States, July 1957-June 1959	7
:	2.	Average number of diabetics and number and percent whose diabetes caused no chronic limitation of activity as reported in interviews by sex and age: United States, July 1958-June 1959	8
:	3.	Average number of diabetics and number and percent whose diabetes caused no bed disability as reported in interviews by sex and age: United States, July 1957-June 1959	9
L	4.	Average annual number of disability days associated with diabetes as reported in interviews and number of days per diabetic: United States, July 1957-June 1959	10
	5.	Average population used in obtaining rates shown in this publication by age and sex: United States, July 1957-June 1959	11

Table 1. Average prevalence of diabetes according to medical care status as reported in interviews and average prevalence per 1,000 population by sex and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

Sex and age	Total	Medically attended cases		Total	Medically attended cases	
·	cases	Total	Under care	cases	Total	Under care
Both sexes	Average	number i	n thousands	Rate p	er 1,000	population
All ages	1,530	1,525	1,404	9.0	9.0	8.3
0-24 25-44	67 199	66 198	61 181	0.9 4.4	0.9 4.3	0.8
45-54 5 <u>5</u> -64	246 424	246 423	222 392	12.4 28.4	12.4 28.3	11.2 26.3
65-74 75+	416 177	416 176	388 , 159	42.9 35.6	42.9 35.4	40.0 32.0
Male			•			
All ages	660	657	588	8.0	8.0	7.1
0-24 25-44	40 106	39 106	36 97	1.1 4.9	1.0 4.9	1.0 4.4
45-54 55-64	108 181	108 179	94 _ 163	11.2 25.2	11.2 24.9	9.7 22.7
65-74 75+	156 68	156 67	138 60	34.4 31.5	34.4 31.1	30.5 27.8
Female			_			
All ages	871	868	816	10.0	10.0	9.4
0-24 25-44	27 93	27 _ 92	26 84	0.7 3.9	0.7	0.7
45-54 55-64	139 244	138 244	129 229	13.7 31.5	13.6 31.5	12.7 29.6
65-74 75+	260 109	259 109	250 . 99	50.3 38.8	50.1 38.8	48.4 35.2

L

Table 2. Average number of diabetics and number and percent whose diabetes caused no chronic limitation of activity as reported in interviews by sex and age: United States, July 1958-June 1959

Data are based on household interviews during July 1958-June 1959. Data refer to the civilian noninstitutional population of the United States. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

1

S

Sex and age	Number of	Diabetics whose diabetes caused no chronic limitation of activity			
	diabetics (in tho	Number usands)	Percent		
Both sexes					
All ages	1,530	1,147	75.Ò		
0-24	67	59	88.1		
25-44	199	151	75.9		
45-54	246	191	77.6		
55-64	424	337	79.5		
65-74	416	296	71.2		
75+	177	112	63.3		
Male					
All ages	660	522	79.1		
0-24	40	36	90.0		
	106	85	80.2		
45-54	108	90	83.3		
55-64	181	147	81.2		
65-7475+	156	118	75.6		
	68	47	69.1		
Female	· .		·····		
All ages	871	624	71.6		
0-24	27	23	85.2		
25-44	93	66	71.0		
45-54	139	102	73.4		
	244	190	77.9		
65- 7 4	260	178	68.5		
7 5+	109	65	59.6		

Estimates in this table are based on the second year of data collection only. Therefore, the standard errors are 1.25 times the standard errors shown in Appendix 1.

Table 3. Average number of diabetics and number and percent whose diabetes caused no bed disability as reported in interviews by sex and ane: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

	Number of	Diabetics whose diabetes caused no bed disability			
Sex and age	diabetics (in th	diabetics Number (in thousands)			
Both sexes		· · · ·			
All ages	1,530	1,300	85.0		
0-24	67	53	79.1		
25-44	199	166	83.4		
45-54	246	209	85.0		
55-64	424	367	86.6		
65-74	416	359	86.3		
75 +	177	145	81.9		
Male					
All ages	660	-562	85.2		
0-24	40	32	80.0		
25-44	106	91	- 85.8		
45-54	108	91	84.3		
55-64	181	152	84.0		
65-74	156	138	88.5		
75 +	68	58	85.3		
Female					
All ages	871	738	84.7		
0-24	27	21	77.8		
25-44	93	76	81.7		
45-54	139	118	84.9		
55-64	244	215	88.1		
65-74	260	220	84.6		
7 5+	109	88	80.7		

ŗ

5

Table 4. Average annual number of disability days associated with diabetes as reported in interviews and number of days per diabetic: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1]

7

-1

Э

'n

۶

٦

	Disability days associated with diabetes						
Sex and age	Restricted	activity	Bed disa	bility			
	Average number (in thousands)	Per diabetic per year	Average number (in thousands)	Per diabetic per year			
Both sexes							
All ages	30,376	19.9	12,983	8.5			
0-44 45-54	3,173 5,539	11.9 22.5	772 2,154	2.9 8.8			
55-64 65-74	8,143 8,903	19.2 21.4	3,017 4,482	7.1 10.8			
75+	4,619	26.1	2,558	14.5			
Male	ł						
All ages	11,228	17.0	4,349	6.6			
0-44	. 932 2,428	6.3 22.5	303 916	2.1 8.5			
55-64 65-74	3,866 2,680	21.4 17.2	1,493 852	8.2 5.5			
75+	1,323	19.5	785	11.5			
Female	· · · ·	· ·					
All ages	19,148	22.0	8,634	9.9			
0-44 45-54	2,241 3,111	18.7 22.4	469 1,238	3.9 8.9			
55-64 65-74	4,277 6,223	17.5 23.9	1,524 3,630	6.2 14.0			
75+	3,296	30.2	1,774	16.3			

Table 5. Average population used in obtaining rates shown in this publication by age and sex: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 11

5

	Both sexes		Male		Female	
Age	Total	Persons with diabetes	Total	Persons with diabetes	Total	Persons with diabetes
	Average population in thousands					
All ages	169,835	1,530	82,633	660	87,202	871
0-24 25-44	74,826 45,579	67 199	37,233 21,854	40 106	37,593 23,725	· 27 93
45-54 55-64	19,833 14,930	246 424	9,675 7,183	108 181	10,157 7,747	139 244
65-74 75 +	9,698 4,969	416 177	4,530 2,157	156 68	5,167 2,812	260 109

NOTE: For official population estimates for more general use, see Bureau of the Census reports on the civilian population of the United States, in <u>Current Population Reports</u>: Series P-20, P-25, P-50, P-57, and P-60.

TECHNICAL NOTES ON METHODS

Background of This Report

This report on <u>Diabetes</u> is one of a series of statistical reports covering separate health-related topics which are prepared by the U.S. National Health Survey. The report is based on information collected in the continuing nationwide sample of households in the Health Interview Survey, which is a main aspect of the program.

The Health Interview Survey uses a questionnaire which, in addition to personal and demographic characteristics, elicits information on illnesses, injuries, chronic conditions, medical care, dental care, and hospitalization. As interview data relating to each of these various broad subject areas are tabulated and analyzed, separate reports are issued covering one or more specific topics. The present report contains data for 104 weeks of interviewing ending June 28, 1959.

The population covered by the sample for the Health Interview Survey is the civilian population living in the United States at the time of the household interview. Although the sample collection covers persons who are inmates of institutions, data for these persons are not included in the figures given in these reports. Also the sample does not include members of the Armed Forces, United States nationals living in foreign countries, and crews of vessels.

Statistical Design of the

Health Interview Survey

General plan.-The sampling plan of the survey follows a multistage probability design which permits a continuous sampling of the civilian population of the United States, This plan utilizes the 1,900 Primary Sampling Units consisting of counties, groups of contiguous counties, and Standard Metropolitan Statistical Areas into which the country has been divided. The first stage of the design consists of drawing a sample from these Primary Sampling Units (PSU's). During the first 18 months of the Health Interview Survey the sample size was 372 PSU's. This was increased to 500 PSU's in January 1959. However, the basic sampling design and methods of estimating remained unchanged during the two-year period covered by this report. The number of ratio estimating classes shown subsequently in this Appendix are those which applied to the first 18 months of the survey.

With no loss in general understanding, the remaining stages of the sampling can be telescoped and treated in this discussion as an ultimate stage. Within PSU's, then, ultimate stage units called segments are defined, also geographically, in such a manner that each segment contains an expected six households. Each week a random sample of about 120 segments is drawn. In the approximately 700 households in those segments, persons are interviewed concerning illnesses, injuries, chronic conditions, disability, and other factors related to health.

The household members included each week are a representative sample of the population so that samples for successive weeks can be combined into larger samples for, say, a calendar quarter, a year, or more. Thus the design permits both continuous measurement of characteristics of high incidence or prevalence in the population and, through the larger consolidated samples, more detailed analysis of less common characteristics and smaller categories. The continuous collection has administrative and operational advantages as well as technical assets, since it permits field work to be handled with an experienced, stable staff.

à

3

Sample size and geographic detail.—The national sample plan during the 24-month period ending June 1959 included approximately 235,000 persons from 73,000 households in 12,200 segments. The over-all sample was designed in such a fashion that tabulations can be provided for various geographic sections of the United States and for urban and rural sectors of the Nation.

<u>Collection of data</u>.—The field operations for the household survey are performed by the Bureau of the Census under specifications established by the Public Health Service. In accordance with these specifications the Bureau of the Census designs and selects the sample, conducts the field interviewing, and edits and codes the questionnaires. Tabulations are prepared by the Public Health Service using the Bureau of the Census electronic computers.

Estimating methods.—Each statistic produced by the survey is the result of two stages of ratio estimation. In the first of these, the factor is the ratio of the 1950 decennial U.S. total population count to the estimated population in 1950 of the U.S. National Health Survey's first-stage sample of PSU's. This factor is applied separately for more than 50 color-residence classes.

Later, ratios of sample-produced estimates of the population to official Bureau of the Census figures for current population in about 60 age-sex-color classes are computed, and serve as second-stage factors for ratio estimating.

The effect of the ratio estimating process is to make the sample more closely representative of the population by age, sex, color, and residence, thus reducing sampling variance.

As noted, each week's sample represents the population living during that week and characteristics of that population. Consolidation of samples over a time period such as a calendar quarter produces estimates of average characteristics of the United States population for that calendar quarter.

For prevalence statistics based on two years of data collection, figures are first calculated for each calendar quarter by averaging estimates for all weeks of interviewing in that quarter. Prevalence data based on eight quarters of interviewing are then obtained by averaging the eight quarterly figures.

For statistics measuring the number of occurrences during a specified time period, a similar computational procedure is used, but the statistics have a different interpretation. For the disability-day items, the interviewer asks for the respondent's experience over the two calendar weeks prior to the week of interview. Then, the estimated quarterly total for a statistic is simply 6.5 times the average two-week estimate produced by the 13 successive samples taken during the period. Thus, the experience of persons interviewed during a year-experience which actually occurred for each person in a two-calendar-week interval prior to week of interview-is treated in analysis as though it measured the total of such experience occurring in the year. For most statistics such interpretation leads to no significant bias. As noted earlier, the interviewing and estimation procedures are designed to reproduce the experience during the reference period of the questionnaire only for the population still living at the time of interview.

General Qualifications

ĸ

<u>Nonresponse</u>.—Data are adjusted for nonresponseby a procedure which imputes to persons in a household not interviewed the characteristics of persons in households which were interviewed in the same segment. The total noninterview rate is 5 percent; 1 percent is refusal, and the remainder is accounted for by other reasons, such as failure to find any household respondent after repeated trials.

The interview process.—The statistics presented in this report are based on replies secured by interviewing members of the sampled households. Each person, 18 years and over, available at the time of interview, is interviewed individually. Proxy respondents within the household are employed for children and for adults who are not available at the time of the interview, provided the respondent is closely related to the person about whom information is being obtained.

There are limitations to the accuracy of diagnostic and other information collected in household interviews. For diagnostic information the household respondent, can, at best, pass on to the interviewer only the information the physician has given to the family. For conditions not medically attended, diagnostic information is often no more than a description of symptoms. However, other types of facts such as the number of disability days caused by the condition can be obtained more accurately from household members than from any other source.

<u>Rounding of numbers</u>.—The original tabulations on which data in this report are based show all estimates to the nearest whole unit, All consolidations are made from these original tabulations before the numbers are rounded to the nearest thousand for the published tables. Derived statistics such as rates and percent distributions are computed after the estimates have been rounded. Rounding to thousands has been done throughout this report even though, because of sampling error, the estimates may not be accurate to that detail.

<u>Population figures.</u>—Some of the published tables include population figures for specified categories. Except for certain over-all totals by age and sex (which are independently estimated), these figures are based on the sample of households in the U.S. National Health Survey. They are given primarily for the purpose of providing denominators for rate computation, and for this purpose are more appropriate for use with the accompanying measures of health characteristics than other population data which may be available. In some instances they will permit users to recombine published data into classes more suitable to their specific needs. With the exception of the over-all totals by age and sex mentioned above, the population figures may in some cases differ from corresponding figures (which are derived from different sample surveys) published in reports of the Bureau of the Census. For population data for general use, see the official estimates presented in Bureau of the Census reports in the P-20, P-25, P-50, P-57, and P-60 series.

Reliability of Estimates

Since the estimates are based on a sample, they may differ somewhat from the figures that would have been obtained if a complete census had been taken using the same schedules, instructions, and interviewing personnel and procedures. As in any survey, the results are also subject to measurement error.

The standard error is primarily a measure of sampling variability, that is, the variations that might occur by chance because only a sample of the population is surveyed. As calculated for this report, the standard error also reflects part of the variation which arises in the measurement process. It does not include estimates of any biases which might lie in the data. The chances are about 68 out of 100 that an estimate from the sample differs from the value obtained from a complete census by less than the standard error. The chances are about 95 out of 100 that the difference is less than twice the standard error and about 99 out of 100 that it is less than 2½ times as large.

In order to derive standard errors which are applicable to a wide variety of health statistics and which can be prepared at a moderate cost, a number of approximations are required. As a result, tables I through III, included in this Appendix should be interpreted as providing an estimate of the standard error rather than as the precise standard error for any specific statistic.

The following guides will enable the reader to determine sampling errors for the statistics presented in this report:

- Approximate standard errors for estimates of the number of cases of a chronic condition, the number of disability days associated with a chronic condition, and the number of persons in a population group¹ are obtained from the appropriate columns of table I.
- 2. Approximate standard errors for percentage distributions of a chronic condition according to the number of bed-disability days or the extent of activity or mobility limitation associated with it are given in table II.
- 3. Approximate standard errors for prevalence estimates of a chronic condition per 1,000 persons in an age, sex, or color group or per 1,000 total population are obtained from table II. Since table II is set up for the estimation of the standard error of a rate per 100, the prevalence per 1,000 must first be converted to a percentage; table II is then entered with this percentage and the number of persons in the category (base of

¹The number of persons in an age, sex,or color group, or the total number of persons in the population is not subject to sampling error.

the percentage). The entry in the body of the table must then be multiplied by 10 to apply to the rate per 1,000 persons.

- 4. Approximate standard errors of percentage distributions of disability days associated with a chronic condition (not computed in this report) are given in table III.
- 5. A rough approximation of the standard errors for rates showing the average number of disability days per "persons with the condition² per year" is obtained by taking the square root of the sum of the square of the standard error of the numerator used in obtaining the rate divided by the numerator itself and the square of the standard error of the denominator used divided by the denominator itself, and then multiplying by the rate. This computation will normally give an overestimate of the true sampling error.

Example:

It is estimated that each person with diabetes has an average of 8.5 days of bed disability per year because of diabetes (table 4). The numerator of 12,983,000 bed-disability days (table 4) has a standard error of 1,367,000. The denominator of 1,530,000 persons with diabetes (table 1) has a standard error of 71,000. Using these numbers as shown below yields an answer of 1.0, the standard error of the estimated rate.

$$8.5 \times \sqrt{\left(\frac{1,367,000}{12,983,000}\right)^2 + \left(\frac{71,000}{1,500,000}\right)^2} = 1.0$$

²Note that where the rate refers to persons in a disease category, rule 5 applies, even if the group is further subdivided by age, sex, or color.

Table I. Standard errors of estimates of aggregates

(A1)	1 nu u	obers	shown	in	thousands)	1
			DIIIIIIIIIIIII	_		

Size of estimate	Persons with diabetes	Disability days
100	18	-
500	40	-
1,000	60	400
2,000	80	560
3,000	100	. 720
5,000	130	960
10,000	180	1,200
20,000	240	1,760
30,000	260	2,160
50,000	280	2,800
100,000	320	4,400
200,000	-	6,400
500,000	-	12,000
750,000	_	16,800
1,250,000	-	25,600

<u>Illustration of use of table 1.</u>—The estimated number of persons who have received medical attention for diabetes is 1,525,000 (table 1). Since this figure is not shown in the table, it is necessary to interpolate for the standard error. The standard error for an estimate of 1,000,000 is 60,000 and the standard error for an estimate of 2,000,000 is 80,000. Interpolation gives 71,000 as the standard error for 1,525,000.

When the	For e	stimat	ed perc	entages	of
percentage is: (in thousands)	2 or 98	5 or 95	10 or 90	25 or 75	50
	The e pe	e appro error (ercenta	ximate express ge poin	standar ed in its) is:	d
100 500 1,000 2,000	2.9 1.3 0.9 0.6	4.5 2.0 1.4 1.0	5.4 2.4 1.7 1.2	7.8 3.5 2.5 1.8	10.3 4.6 3.3 2.3
3,000 5,000 10,000 20,000	0.6 0.4 0.3 0.2	0.8 0.6 0.5 0.3	1.0 0.8 0.6 0.4	1.4 1.1 0.8 0.6	1.9 1.4 1.0 0.7

Table II. Standard errors of percentages based on persons with diabetes

<u>Illustration of use of table II.</u>—Of the 871,000 fe-males reported as having diabetes, 71.6 percent had no chronic limitation of activity (table 2). Since neither of these values can be read directly from the table, interpolation may be carried out as follows: for a base of 500,000 a statistic of 50 percent has a standard error of 4.6 percentage points and a statistic of 75 percent has a standard error of 3.5 percentage points. Interpolating, with a base of 500,000 an estimate of 71.6 percent would have a standard error of 3.6 percentage points. Corresponding calculations with a base of 1,000,000 produce a standard error of 2.6 percentage points. A final interpolation between these two results yields an estimate of 2.9 percentage points for a statistic of 71.6 percent with a base of 871,000. (Although interpolation has been carried out in two dimensions here to illustrate the use of the table, a simple scanning of the table will provide an approximate answer which will usually be sufficient.)

Table III. Standard errors of percentages based on disability days

When the base of the	For e	stimat	ed perc	entages	of
percentage is: (in thousands)	2 or 98	5 or 95	10 or 90	25`or 75	50
`````````````````````````````````	The e pe	e appro error ( ercenta	express ge poin	standar ed in its) is:	d
2,500 12,500 25,000 50,000	3.4 1.5 1.0 0.7	5.2 2.3 1.7 1.2	7.2 3.2 2.2 1.6	10.4 4.6 3.3 2.3	12.0 5.4 3.8 2.7
75,000 125,000 250,000 500,000	0.6 0.5 0.3 0.2	1.0 0.7 0.6 0.4	1.3 1.0 0.7 0.5	1.9 1.4 1.0 0.7	2.2 1.7 1.2 0.9

Illustration of use of table III. — Of the 30,376,000 restricted-activity days due to diabetes, 29.3 percent were for persons 65-74 years of age. Since neither of these values can be read directly from the table, interpolation may be carried out as follows: with a base of 25,000,000 a statistic of 25 percent has a standard error of 3.3 percentage points and a statistic of 50 percent has a standard error of 3.8 percentage points. Interpolating, a statistic of 29.3 percent with a base of 25,000,000 would have a standard error of 3.4 percent-age points. Corresponding calculations with a base of 50,000,000 produce a standard error of 2.4 percentage points. A final interpolation between these two results yields a, standard error of 3.2 percentage points for a statistic of 29.3 percent with a base of 30,376,000. (Although interpolation has been carried out in two dimensions here to illustrate the use of the table, a simple scanning of the table will provide an approximate answer which will be sufficient for most purposes.)

### APPENDIX II

### DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

#### Terms Relating to Chronic Conditions.

<u>Condition</u>—A condition is defined as any entry on the questionnaire which describes a departure from a state of physical or mental well-being. In the coding and tabulating process, conditions are first classified according to the type of disease, injury, or impairment, and then according to a number of other criteria such as whether they were medically attended, whether they resulted in disability, and whether they were acute or chronic. For the purposes of each published report or set of tables, only those conditions recorded on the questionnaire which satisfy certain stated criteria are included.

Conditions, except impairments, are coded by type according to the International Classification of Diseases, 1955 Revision, with certain modifications adopted to make the coding procedures more suitable for a household-interview-type survey. Impairments, consisting of selected chronic or permanent defects resulting from disease, injury, or congenital malformation, are coded according to a special supplementary classification which permits their classification by type and site, and also by etiology.

<u>Chronic condition</u>.—A condition is considered to be chronic if it is described by the respondent (1) in terms of one of the conditions on the "Check List of Chronic Conditions" or in terms of one of the impairments on the "Check List of Impairments" (Cards A and B, Appendix III), or (2) as having been first noticed more than three months before the interview. For this purpose, first noticed is defined as the time at which the person first felt sick or when he or his family was first told by a physician that he had a disease of which he was previously unaware. For a condition which is episodic in nature, the onset is always considered to be the original onset rather than the onset of the most recent episode.

Prevalence of a condition.—In general, the prevalence of a condition is the estimated number of cases existing in a population at a specific point in time or the average number existing during a specified period of time.

In the National Health Survey, the prevalence of a chronic condition is the number of cases reported to be present at the time of the interview or at any time during the 12 months prior to the interview. Estimates of the prevalence of chronic conditions may be restricted to cases which satisfy certain additional criteria. For example, only cases involving a day or more in bed during the past year or cases under care may be included.

<u>Medically attended condition</u>.—A condition is considered to be medically attended if a physician has been consulted about it either at its onset or at any time thereafter. Medical attention includes consultation either in person or by telephone for treatment or advice. Advice from the physician transmitted to the patient through the nurse is counted as well as visits to physicians in clinics or hospitals. If during the course of a single visit the physician is consulted about more than one condition for each of several patients, each condition of each patient is counted as medically attended.

Discussions of a child's condition by the physician and a responsible member of the household are considered as medical attention even if the child was not seen at that time.

For the purpose of this definition, the term "physician" includes doctors of medicine and osteopathic physicians.

<u>Condition under care</u>.—By under care is meant one or more of the following: (1) currently taking medicine or treatment prescribed by a physician, (2) observing a systematic course of diet or activity prescribed by a physician, (3) visiting the physician regularly for checking on the condition, or (4) under instruction from the physician to return if some particular thing happens.

Physician is again defined as a doctor of medicine or an osteopathic physician.

#### Terms Relating to Disability

Disability is a general term used to describe any reduction of a person's activity as a result of an acute or chronic condition. It is measured in numbers of disability days, and for persons with one or more chronic conditions, it is also expressed by the extent to which they are limited in their major activity and their mobility.

Disability days.—Disability days are classified according to whether they are days of restricted activity, days in bed, days in the hospital, days lost from work, or days lost from school. All hospital days are, by definition, days of bed disability; all days of bed disability are, by definition, days of restricted activity. The converse form of these statements is, of course, not true. Days lost from work and days lost from school are also days of restricted activity for the working and school-age populations. Hence, restricted activity is the most inclusive term used in describing disability days.

<u>Restricted-activity day.</u>—A restricted-activity day is one on which because of a specific illness or injury a person substantially reduces the amount of activity normal for that day. The type of reduction will vary with the age and occupation of the individual as well as with the day of the week or the season. Restricted activity thus covers a range from substantial reduction of normal activity to complete inactivity.

Bed-disability day.—A bed-disability day is one on which more than half the daylight hours were spent in bed because of a specific illness or injury. All hospital days for inpatients are considered to be days of bed disability even if the patient was not actually in bed at the hospital.

<u>Work-loss day</u>, —A work-loss day is a normal working day on which a person did not work at his job or business because of a specific illness or injury. The

number of days lost from work is determined only for persons 17 years of age or older.

#### Chronic Activity and Mobility Limitation

Chronic activity limitation, —Chronic activity limitation is ascertained for all persons with one or more chronic conditions. See Cards C, D, E, and F, Appendix III. The replies indicate to what extent the person (worker, housewife, child, or other) is limited by his chronic disease(s) with respect to his ability to carry on the major (or usual) activity of the group of which he is a member. The person may be completely limited; may be able to carry on the major activity but be limited in the kinds and amounts of his activities; or may not be limited in any of these ways.

<u>Chronic mobility limitation.</u>—Persons with one or more chronic conditions are also classified with respect to the extent to which their mobility is limited. See Card G, Appendix III. Degrees of limitation of  $mo_{\overline{1}}$  bility include being confined to the house all the time, or, if able to go outside, needing the help of another person or having trouble in getting about freely.

#### Demographic Terms

Age.—The age of the person on his last birthday recorded on the questionnaire in single years. Ages are then grouped in intervals suitable for the topic under discussion.

Usually working.—A term applied to an individual 17 years of age or older who was gainfully employed as a paid employee, a self-employed person, or as a worker in a family business for more than half of the 12 months prior to the interview. A person who does only volunteer or unpaid work—such as work in his own home or work for the church or community—is not considered to be gainfully employed.

# 

# QUESTIONNAIRE.

The items below show the exact content	and wording o	of the questionnair	e used in the household surv	ey. The actual
uestionnaire is designed for a household	as a unit and	includes additional	spaces for reports on more	than one person.

CONFI	DENTI	The N AL: would purpo	permit i ses of the	lealth Survey is aut identification of the e survey, and will n	horize indiv ot be o	d by Public Law vidual will be h lisclosed or rele	652 of eld str ased to	f the f rictly o othe	64th Congress confidentia rs for any o	s (70 Stat l, will be ther purpo	: 489; 42 U.S used only by ses (22 FR )	.C. 3 y pers 1687).	05). Al	l inform gaged in	ation whi and for	ch the
Form NHS- (4-18-58)	-2			U.S. DEPA BURI Acting a U.S. PU	RTMEN EAU OF 9 Collec BLIC H	TT OF COMMERC THE CENSUS ting Agent for the EALTH SERVICE	E	-			· · ·	1. Q	uestion of	haire		
				NATIONAL	. не	ALTH SUR	/EY					_				
			<u></u>										uc etion	Sauce .		
2. (e) Add	ress or	description	of location	• 		·	- 3.	Iden. Code	4. Sub- sample weight	5. Sample	6. PSU Number	7. S	egment	No.	8. Serial	No.
							_					L_			<u> </u>	
<u></u>	-		`~				- 9.	la thi	s house on a	fana or rar	ich?		• • • • •	·····	] Yeis 🗌	] No
								). Wha	t is the telep	hone numbe	r here?	11.	What is	the best	time to co	1117
(b) Type of ing qua	f lív- [ utets [	] Dwelliog u Other	nit (¢) Ni	me of Special Dwellin	ng Plac	e Code					Na phone					
12. Are the vocant	are any ), in thi	other living s building (o	quorters, partment)?	occupied or	🗖 Ye	No	-   ı.	4. Doe EN1	s anyone els FRANCE to g	aliving in at to his li	this building ving quarters?	use ¥( ?	0UR 	⊏	] Ye∎ []	יא [_
Ask at all	1	except a per-	nent house				+			<u> </u>	INSTRUCT	IONS				
13. Is ther to live	re ony c i la - si	ther building	on this p lor vacan	roperty for people t?	🗌 Ye	• M 🗋		lf "Ye: whethe listing	to question to one or mos is to be corr	e addition ected.	t 14 apply def al questionna	inition ires s	bould b	welling u a filled a	nit to dete and whethe	rmine rr the
		· · · · · · · · · · · · · · · · · · ·		· ·	r	15. RECORD OF	CALLS	AT H	OUSEHOLOS		<b></b>					1
		ltem	Date	· 1	Com.			om.	3	Com.	<u>                                     </u>		Com.		<u> </u>	Com.
Enti	re hous	ehold	Time		•											
indiv respon	idual idual	Col. No	Date Time					-								-
				<u>۱</u>		16. REASON	FOR N	ON-IN	TERVIEW							
TYPE:	·	<u> </u>		<u>`</u>	8		·		<u>c</u>	<u>{`</u>			Z		·	
	[] Re	fusal		Vacant - non-	scason	<b>•</b> L		emolis	bed .	Inte	rview not obta	ained f	for;			
		peated calls	-	Vacant - seas		A		sampl	e hy mistake	Col					. <u> </u>	
Resson:	C Te	mporarily ab	s <del>e</del> at	Armed Forces		watere		mple	.ed 10 600-	hec	LUSE:	-	• .			
		het (Specify)		Other (Specif	<b>7)</b>	~		ber (S	pecity)						•.	
Comments	00 Л0Л	-interview		I <del></del>					-							
17. Signat	ure of 1	ntervie wer		<u>.</u>								18.	Code			
Special in:	structio	as or aotes										Ļ				
1. (a) Wha	nt is the	nome of the	head of th	his household? (Enter means who live here?	name /lier	in first column)	sualiv I	ive he					Lastna	me		(1)
and	all per	sons staying	here who	have no usual place	of resid	ience elsewhere.	List the	ese	,	-		1	,			
peri (c) Da	sons in anv (ot	the prescrib her) lodgers	ed order.) er roomera	live hare? -	No	( Yes (List)						→				
(d) la t	here an	yone else wi	o lives he	rewholanow Temporarilvin	No	Yes (List)						<b>→</b> [	First n	me and i	nicial	
oho (e) la 4	ospitol	vone else at	vino here	now?	No	Yes (List)			<b></b>			<b>_</b>				
(f) De (	any of t	hese people	have a hor	me elsewhere? Yes (if not a	housel	old member, dele	·									
2. How ar	e you r	eloted to the	head of t	he household? (Enter	relatio	aship to head; for	example	.e:		<u> </u>			Relatio	nship	Head '	
											· · · ·	†	Age			ider year
3. How of	id were	you on your	last birthd	lay?							<u>.</u>		<b>□ v</b> h			gro
4. Race ((	beck of	e box for ea	h person)	<u> </u>			<u>.</u>			<u> </u>	-	+	<u> </u>	0 le		emale
6. Where	were yo	iu bern? (Re	cord state	or foreigo country)							<u>-</u>		Ste	te or for	ign count	:y)
lf 14 y 7. Are yo (Checi	venrs ol ou now v k one b	d or over, as merried, wide ox foc each p	k: wed, divo ersoa)	rced, separated or new	er mar	rind?					•		() Maa ♥ic	nied lowed	Dider 14 y	ivorced sparated ried
If 14 y	ears of	d or over, as	k;										Elem:		Under 14 ) 3 4`5 6	venra 78
8. Whot is (Circle	s the hi e highe	ighest grade nt grade com	you compl pleted or c	eted in school? check "None")		-							High: Colleg	1 2 e: 1 2	3 4 3 4 5 4	

•	lf Mal	e and	14 yea	urs old or ove	r, ask:	·					·····		Fem.	. or und. 14 yr
9.	(a) Die If "Ye	i you s," a	ever s sk:	erve in the Ar	med Forces (	of the U	nited States?					Yes_		<u>No</u>
	to) are you now in the armed Forces, not counting the reserves r If "Yes," delete this person from questionnaire)													
	(c) Was any of your service during a war or was it peace-time only?											- Wer		Peace- time onl
	li 🐂 (d) Du	z, n. ving '	ok: which v	war did you se	erve?				-			Spanis Americ	h — — —	🗍 WW-U
1	l "Pe	ice-ti	ime" or	ily, ask:	•	-					,	<b>WW-I</b>	<u> </u>	[] Korean
	(e) ₩a	a ony	of you	ar service her	ween June 2/	/, 1950 a	ind January 31	, 19557	· · ·			Yes		<u> </u>
10.	(a) W	het w	la or o ere ýo	ver, aak: J doing most (	of the past 17	2 months	-						Unde e	n 6 years
	(1 (1	for m for fe	ales or males	rer 16): worki over 16): wor	ng, looking fi king, looking	or work, for worl	or doing some k, keeping heu	thing else? se, or doing r	something e	1267		Lookin	g for work	£ .
1	() Soc	For cl nethis	hildren az else	6-16): going -	to school or a	doing so 50 years	mething else?	sk:		-	· ·	Going	to school	
	(b) A	<b>Ja</b> 70	u ratiri	d?	•••							Yes	ing else	
		Int	erviev	each adult p	erson for him	self for	mestions 11-2	6 and Tables	1. 11. and /	L. If		Respon	aded for a	elf
	ĸ	be	is at h	ome. Enter c	oluma aumbe	r of reap	ondent in each	1 column.	·, ··,			Col. No		respondent
	We cr	tni e	erested	in all kinds	of Illness, wi	hether s	erlous er net					- C.) Yes	-	№
11.	Were (c) W	you s hat w	as the	ony time LA3 matter?	T WEEK OR	THE WE	EK BEFORE	1.						
12.	(b) A	week	ng eise : or the	veek before	did you have	ony acc	idents or injur	ies, either at		(a) What were they?	· · ·	Yes		
home or away from home?											T Yes		. — [] No	
•••	Injury that happened before that time?													-
	(u) man werd these effects? (b) Anything else?											L		
14.	14. Loss week or real week before did you rake any mean mean mean and in the one of the second secon										Yes		N0	
15. AT THE PRESENT TIME do you have any aliaments or conditions that have lasted (0) What are they? for a long time? (if "No") Even though they don't bother you all the time? (b) Anything else?								Yes		. No				
_	Table I - ILLNESSES, IMPAIRMENTS AND ACCIDENTS													
_			Did	What did i	the doctor en;	v It	10 I - ILLING	25525, IMF 201 OF	U eye	What kind oftrouble	What part of the	body is	LAST W	EEK How
			700 8787	wastdi medical to	d he use any arms?	⁻ 1	sympton or a from q. 13 or	condition	trouble of any	la 117	affected?		OR THE	E- days,
1			talk to a	(If doctor	not talked to	- "No"	esk:	4	kind and	Ask only for:	Show in tollows for members lis	ng detsu ted below:	FORE d	id includ se ing
			doctor about	in col. (c) dent's des	- record resp cription)	póa-	What was the	cause of1	6 yrs. old ar	anemia rheumatism	Heod - (Skull, e face)	calp or	you to c down on	ut the 2 week-
			1	(If ill-effe	of earlies	.	(If accident or injury, slao ank: tumor (or cysts) Spine - (Upper, middle			middle or	your usu activitie	iol ands?		
ž	C01.			accidem,	record ill eff	ects.	fill Table A)			OR	bock		for as m as a day	uch /7
	No. 1	ion .		Erren					Can you	or (d-2) of:	Arm- (Shoulder,	upper,	Check	one
1	son	No.		occurring	during past 2	·			read ardin-	trouble condition disease	hand)	·	No	Yes
						-		. *	cery nows-	coupled	Leg - (nip, upp lower, an	er, Knee, kie, foot)		
				What part burt? Who	of the body v at kind of inj	ves ury			poper print	seeing or hearing;	ALSO If arm, leg, eye	. ar ear,	to Col.	_
				was it?	unything also	1			with glasses?	s part of the body; "ments!" or any	state whether C	HE or	(*))	
·				(A180, 11	1 Табіе лу	1				internal örgan				_   _
	(a)	<u>(b)</u>	(c)		(d-1)		(d	<u>+2)</u>	(d-3) X 🗔 Vee	(d-4) X	(4-5	) X	(e)	
1														Days
-				·	1	Cable ()	- HOSPITA	LIZATION	DURING F	AST 12 MONTHS	- · · · · · · · · · · · · · · · · · · ·			
			W	hen did	How many	To Ime	trvie wer		What did t	hey say at the hospital the co	ndition was ~	Were any	operation:	s performed
	C-1		++	e hos-	you in the	many	many	this	(lí "they"	did't say, ask):		the hespi	nal 7	noy ui
ų.	No.	tio	•	touth, Year)	not count-	- days	- days	still	What did i	the lost doctor you tolked to s	ay it was?	If "Yes"		ma of the
dana	per-	<b></b>			day you left?	the part	at the past 2 weeks?	hospital	(Show san	ne detail as in cols. (d-1)-(d-5	) of T.I)	opera	tion ?	
Libe						months	17	Sundéy night?	(If conditi	on from accideot or injury, fil	I TableA)	(b) Any of	ther opera	tions?
				-						-			<i>a</i> .	-
	(@)	-	<u> </u>	(c)	(4)	(e)	(1)	(a)		(2)		C Yes	(1)	10
			M	D:		or		L Ies				Y		•
1			1	~ .	Days	Days	Daye	□ N°		e				
							Τ/	ABLE A (/	Accidents	and Injuries)				
	Line M from	ło. 1	1. 1	that part of th	w body was l	burt? Wi	hat kind of inju	ury was it? A	laything el			Accid	eat happe 2 weeks	ned during
	4 =DIC	<u>_</u>			<u>.                                    </u>		. <u></u> .							
2	Then -	11d 14	hanne	n? V	(Free	er month	also if the we	ar is 1957 ai	1958)	Month			lent happe 2 weeks	ned during
-					(200				/					
3. 1	There	did f	he acci	ident hoppen?							-			

Æ

	At home (inside or ourside the house) (own home or someone else's)	*[] While in Armed Services
hicle		

Under 14 years at time of accident

Some other place

 Vers a car, truck, bus or other motor vehicle involved in the accident in any way?
 Yes
 No
 Vers you at work at your job or business when the accident happened?
 No

<ul> <li>16. Hos anyone in the family - yeu, your-, etchad any of these conditions DURING THE PAST 12 MONTHS? <ul> <li>(Read Card A, condition by condition; record any conditions mentioned in the column for the person)</li> </ul> </li> <li>17. Does anyone in the family have any of these conditions? <ul> <li>(Read Card B, condition by condition; record any conditions mentioned in the column for the person)</li> </ul> </li> <li>18. (a) LAST WEEK OR THE WEEK BEFORE did anyone in the family - you, your-, etctalk to a doctor or go to a doctor's affice or clinic? Anyone else? <ul> <li>(b) How many times during the past 2 weeks?</li> <li>(c) Where did you talk to the doctor?</li> <li>(d) How many times of - (home, affice, clinic, etc.)?</li> <li>(Record total number of times for each type of place) <ul> <li>(*Hospital clinic* excludes overnight stays)</li> </ul> </li> <li>19. (a) Last week or the week before did anyone in the family go to a dentist? Anyone else?</li> </ul></li></ul>	Yes     No       Yes     No       Yes     No       Yes     No       Place     Times       At home     Times       At office     Times       Company or industry     Over relephase       Over relephase
(Read Card A, condition by condition; record may conditions mentioned in the column for the person)         17. Does anyone in the family have any of these conditions? (Read Card B, condition by condition; record may conditions mentioned in the column for the person)         18. (a) LAST WEEK OR THE WEEK BEFORE did anyone in the family - you, your-, etctelk to a doctor or go to a doctor's office or clinic? Anyone else? (b) How many times during the post 2 weeks?         (c) Where did you tolk to the doctor? (d) How many times of the column office, clinic, etc.}?? (Record total aumhor to fuice of times for each type of place) ("Hospital clinic" excludes overnight stays)         19. (a) Last week or the week before did anyone in the family go to a dentist? Anyone else?	Ves No Ves No. of times Place Iines At home. At office
17. Does anyone in the family have any of these conditions?         (Read Card B, coadition by condition; tecord any conditions mentioned in the column for the person)         18. (a) LAST WEEK OR THE WEEK BEFORE did anyone in the family - you, your-, stetelk to a dector or go to a dector's affice or elinic? Anyone alse?         (b) How many times during the post 2 works?         (c) Where did you tolk to the doctor?         (d) How many times of the doctor?         (d) How many times of the doctor?         (e) Where did you tolk to the doctor?         (f) How many times for each type of place)         ("Hospical clinic" excludes overnight stays)         19. (a) Lest week or the week before did anyone in the family go to a dentist? Anyone else?	Yes     No       Yes     No       No. of times     No       At home.     Times       At home.
(Read Card B, condition by condition; record any conditions mentioned in the column for the person)         18. (a) LAST WEEK OR THE WEEK BEFORE did anyone in the family - you, you-, stctelk to a dector or go to a dector's office or clinic? Anyone else?         (b) How many times during the post 2 weeks?         (c) Mhere did you tolk to the dector?         (d) How many times for each type of place)         (*Hospital clinic" excludes overnight stays)         19. (a) Last week or the week of the dector did anyone in the family go to a dentist? Anyone else?	Ves N <u>Place</u> <u>times</u> At hoore. <u>toopial clinic</u> <u>Company or industry</u> <u>Over stelphace</u>
<ul> <li>18. (a) LAST WEEK OR THE WEEK BEFORE did anyone in the family - you, your-, etctelk to a dector up to a dector's affice or clinic? Anyone else?</li> <li>(b) How many times during the post 2 weeks?</li> <li>(c) Where did you tolk to the doctor?</li> <li>(d) How many times at - (home, affice, clinic, etc.)?</li> <li>(Record total aumber of times for each type of place)</li> <li>("Hospital clinic" excludes overnight stays)</li> <li>19. (a) Last week or the week before did anyone in the family go to a dentist? Anyone else?</li> </ul>	No. of times
(b) How many times during the past 2 weeks?     (c) Where did you talk to the doctor?     (d) How many times at - (home, affice, clinic, etc.)?     (Record task number of times for each type of place)     ("Hospical clinic" excludes overnight stays)     (9) Last week at the week before did anyone in the family go to a dentist? Anyone else?	No. of times           Place         Times           At boxes
<ul> <li>(c) Where did you talk to the doctor?</li> <li>(d) How many times at - (home, affice, clinic, etc.)?</li> <li>(Record task number of times for each type of place)</li> <li>("Hospical clinic" excludes overnight stays)</li> <li>19. (a) Last week or the week before did anyone in the family go to a dentist? Anyone else?</li> </ul>	Place     Times       At home
<ul> <li>(d) How mony times at (home, office, clinic, etc.)?</li> <li>(Record total number of times for each type of place)</li> <li>("Hospital clinic" excludes overnight stays)</li> <li>19. (a) Last week or the week before did anyone in the family go to a dentist? Anyone else?</li> </ul>	At boxe At office Hospital clinic Company or industry Over telephane
(Record total number of times for each type of place) ("Hospital clinic" excludes overnight stays) 19. (a) Last week or the week before did anyons in the family go to a dentist? Anyone else?	At office
("Hospital clinic" excludes overnight stays) 19. (a) Last week at the week before did anyone in the family go to a dentist? Anyone else?	Over telephone
19. (a) Last week at the week before did anyone in the family go to a dentist? Anyone else?	Other (Specify)
the second se	
(b) How many times during the past 2 weeks?	No. of times
20. How many times altogether in the post 12 months did you go to a dentist?	One Three Two Four or more None
21. (o) DURING THE PAST 12 MONTHS has anyone in the family been a patient in a hespital avernight or longer?	Yes (Table II)
(b) Haw many times were you in the hospital?	No. of times
22. (a) During the past 12 months has anyone in the family been a patient in a nursing home or sonitarium?	Yes (Table II) N
lf "Yes" (b) Haw many times were you in a nursing hame or sanitarium?	No. of times

How many of these         If 6 years old or over, sek:         Did you first notice         To DURING THE PAST 3 MONTHS         Did you first notice         How long notice         Do you         About hew         Ask after completing condition for each per condition for each per to be or before that just           In bed or of before         Did you first notice         Did you first notice         Did you first notice         Did you first notice         How long onte         Do you         About hew         Ask after completing condition for each per doys           In bed or the during the dary         II-Ster to before         Did sturt during the post         Did sturt during the post         Did sturt during the post         MONTHES or o dector the dary?         Did cover that this?         Sector of the this?         If sturt doctor         If sturt during the post         If sturt during the post         If sturt this?         If sturt doctor         If sturt doctor         If sturt doctor         If sturt doctor         If sturt doctor         If sturt doctor         If sturt is this cord doctor	Did you first notice To DURING THE PAST 3 MONTHS Later or before that fine? Viewer: DURING THE PAST 3 MONTHS Later viewer: DURING THE Later DURING THE Later DURING THE Later DURING THE Later DURING THE Later PAST 12 MONTHS or a before that during the past Checked, time?	Now long         Do you         Absut         Ask after completing last conditions for each present monty           site any medi-astic class         nonity         Please         If 1, 2         If         If <t< th=""><th>*1,* *2* *3*</th></t<>	*1,* *2* *3*
would you from months months months time? coodi- you work (going to (Co → 2 wecks, ssk): is on either which month? The full during past that one for? kept you to storesard. Seconds of any of the which month? The store of the which month? The store of the which month? The store of the storesard to storesard. Second of the which month? The store of the storesard to storesard. Second of the which month? The store of the storesard to storesard. Second of the storesard to storesard to storesard to storesard. Second of the storesard to store	mosths mosths time? condi (If during past tion (If during past (If during past (If during past 2 weeks, ask): roo (Co 2 weeks, ask): roo (Which month? of	fices prescribed kins read each because (s) (r) has cace for i kept you into the because (s) (r) north, in bed me which of any Piet nater Or, follow for all me which of the "Which" load that you double or mest stratement item this	col. · nek: nose k ot · o
working or a lob or busi- ness scrept for?     col. (n)     (m)     Which week, lost week, the week before?     coc of cards A or B, commer; commer; or wise     Out.1. for "No.")     he gove?     of the day?     fire you best. he gove?     wise of the gove?     fire you day?     he for best. he gove?     fire you day?     he gove?     of the day?     he gove?     fire you best. he gove?     he gove?     of the day?     he gove?     fire you best. he gove?     he gove?     fire you day?     he gove?     he gove?     fire you best. he gove?     he gove?     fire you best. he gove?     he gove?     fire you day?     he fire you he gove?     he gove?     fire you best. he gove?     fire you best. h	col. (n)) Which week, and of Cards for Instrueek or A or B, before? continue; other- wise STOP	(c)	d aad a d each a tament "I which tement t yau st. ow cd G) (u)
		Mos Yes Days	
Or         No         Or         Or         No         Or         No         Or         No         Or         No         No         Or         No         Or         No         Or         No         Or         No         Or         Or         No         Or         Or         No         Or         No         Or         No         Or         No         Or         Or         No         Or         No         Or         Or         Or         Or         Or	Before 2 with Before	Yra.         No         or         No           No Dr.         No ne         No         No	- 1

		For completed	hospitalizations	only:		• ,	
	heapitol you wave in? (Enter name, city or county and State)	Was any of the hospital hill pold for by any kind of insurance?	If "No" to col. (k), ask: Or, by any kind of plan that pays for hospitel costs?	If "No" to hoth cols. (k) and (l) Do you expect any of the hospital hill to be pold for by insurance or any plan of this kind?	What part of the haspital bill was (will be) taken care of by insurance?	Who carries the cost of pays the premium?,	this insurance—that is, <del>who</del>
٠İ	(j)	(k)	(1)	(m) .	(n)		(0)
		Yes (Skip to col.n)	Yes (Skip to col.a)	🖂 Yes	□ Under ½΄ □ ½ up to ½	Employer	Other (Specify)
Ľ		D No	Nº No	🗀 Na (Stop)	1 % or more	[_] Union, clubs, etc.	

POOTNOTES AND COMMENTS

21

· · ·

Card A	Gard C	Çard E	Card G
MATIONAL HEALTH SURVEY Check List of Chronic Conditions	NATIONAL HEALTH SURVEY For: Workers and other persons except Housewives and Children	NATIONAL HEALTH SURVEY For: Children from 6 to 16 years old and others going to school	NATIONAL HEALTH SURVEY
1. Asthma16. Kidney stones or other2. Any allergykidney trouble3. Tuberculosis17. Arthritis or rheumatism4. Chronic bronchitis18. Prostate trouble5. Repeated attacks of sinus trouble19. Diabetes6. Rheumatic fever20. Thyroid trouble or7. Hardening of the arteriesgoiter8. High blood pressure21. Epilepsy or convulsions9. Heart trouble22. Mental or nervous11. Trouble with varicose velnstrouble12. Hemorrhoids or piles23. Repeated trouble with13. Gallbladder or liver trouble24. Tumor or cancer15. Any other chronic25. Chronic skin troublestomach trouble26. Hernia or rupture	<ol> <li>Cannot work at all at present.</li> <li>Can work but limited in amount or kind of work.</li> <li>Can work but, limited in kind or amount of outside activities.</li> <li>Not limited in any of these ways.</li> </ol>	<ol> <li>Cannot go to school at all at present time.</li> <li>Can go to school but limited to certain types of schools or in school attendance.</li> <li>Can go to school but limited in other activities.</li> <li>Not limited in any of these ways.</li> </ol>	<ol> <li>Confined to the house all the time, except in emergencies.</li> <li>Can go outside but need the help of another person in getting around outside.</li> <li>Can go outside alone but have trouble in getting around freely.</li> <li>Not limited in any of these ways.</li> </ol>
-Card B	Card D	Card F	Card H
NATIONAL HEALTH SURVEY	NATIONAL HEALTH SURVEY	-NATIONAL REALTH SURVEY	NATIONAL HEALTH, SURVEY
Gheck List of impairments	For: Housewife	For: Children under 6 yeare old	Family income during past
<ol> <li>Deafness or serious trouble with hearing.</li> <li>Serious trouble with seeing, even with glasses.</li> <li>Condition present since birth, such as cleft palate or club foot.</li> <li>Stammering or other trouble with speech.</li> <li>Missing fingers, hand, or arm.</li> <li>Missing toes, foot, or leg.</li> <li>Cerebral palsy.</li> <li>Paralysis of any kind.</li> <li>Any permanent stiffness or deformity of the foot or leg,</li> </ol>	<ol> <li>Cannot keep house at all at present.</li> <li>Can keep house but limited in amount or kind of housework.</li> <li>Can keep house but limited in outside activities.</li> <li>Not limited in any of these ways.</li> </ol>	<ol> <li>Cannot take part at all in ordinary play with other children.</li> <li>Can play with other children but limited in amount or kind of play.</li> <li>Not limited in any of these ways.</li> </ol>	<ol> <li>under \$500 (Including loss)</li> <li>\$500 - \$999</li> <li>\$1,000 - \$1,999</li> <li>\$2,000 - \$2,999</li> <li>\$3,000 - \$3,999</li> <li>\$4,000 - \$4,999</li> <li>\$5,000 - \$6,999</li> <li>\$7,000 - \$9,999</li> <li>\$10,000 and over.</li> </ol>
fingers, arm, or back.			

[?] 22

ł