


INTERNATIONAL CIRCUMPOLAR SURVEILLANCE (ICS) SUMMARY REPORT



YEAR 2011 DATA

TABLE OF CONTENTS

	<u>Page</u>
Summary	1
Introduction	2
Goals	2
Methods	3
Results	
<i>Streptococcus pneumoniae</i>	8
<i>Haemophilus influenzae</i>	16
<i>Neisseria meningitidis</i>	20
Group A Streptococcus	22
Group B Streptococcus	25
Conclusions	28
Acknowledgements	28
Source	28
References	29
Participants	30

SUMMARY

International Circumpolar Surveillance (ICS) is a population-based surveillance system for invasive bacterial diseases established in the U.S. Arctic, Northern Canada, Greenland, Iceland, Norway, Finland, and Northern Sweden. Data collection began in 1999 and includes information on disease caused by *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Neisseria meningitidis*, and groups A and B *Streptococcus* (GAS, GBS). This report reviews the data collected for the year 2011.

Data on invasive disease with the organism *S. pneumoniae* are collected from all participating countries. A total of 1,803 cases of invasive pneumococcal disease were identified in 2011. Overall, rates of invasive *S. pneumoniae* were highest in individuals less than 2 years of age or in persons 65 years and older. Case fatality ratios ranged from 11-40%. Race and ethnicity data are collected only in N. Canada and the U.S. Arctic; rates of invasive pneumococcal disease in Northern Canadian Aboriginals and U.S. Arctic Native populations were 32 and 39 cases per 100,000 population, respectively, which are similar to the 2010 rates in these populations. Pneumonia and bacteremia were the most common clinical presentations; cigarette smoking alcohol abuse, and chronic lung disease were the most common risk factors. The most common *S. pneumoniae* serotype in Iceland was 14, in Greenland it was serotype 1, and in N. Canada and the U.S. Arctic the most common serotype was 7F.

Data on invasive disease due to *H. influenzae*, *N. meningitidis*, and groups A and B *Streptococcus* are currently collected in Greenland, Northern Canada, Northern Sweden and the U.S. Arctic; Norway also contributes data on cases of *N. meningitidis* and *H. influenzae*. A total of 127 *H. influenzae* cases, 43 *N. meningitidis* cases, 90 GAS cases, and 43 GBS cases were reported in 2011. In general, the highest rates of disease for all organisms occurred in N. Canada Aboriginal or Alaska Native persons less than two years of age and persons 65 and older.

Surveillance Organisms Reported by Country, ICS 2011 Data

Country	<i>S. pneumoniae</i> n (rate*)	<i>H. influenzae</i> n (rate*)	<i>N. meningitidis</i> n (rate*)	GAS n (rate*)	GBS n (rate*)
Finland	782 (14.5)	N/A	N/A	N/A	N/A
Greenland	6 (10.6)	0 (0)	0 (0)	1 (1.8)	0 (0)
Iceland	33 (10.4)	N/A	N/A	N/A	N/A
N. Canada	36 (23.4)	15 (9.7)	2 (1.3)	12 (7.8)	3 (1.9)
N. Sweden	76 (15)†	3 (1.2)‡	0 (0)‡	3 (1.2)‡	0 (0)‡
Norway	729 (14.8)	85 (1.7)	37 (0.8)	N/A	N/A
U.S. Arctic	141 (19.5)	24 (3.3)	2 (0.3)	74 (10.2)	40 (5.5)
Total	1,803 (14.9)	127 (2.1)	43 (0.7)	90 (7.6)	43 (3.6)

*Cases per 100,000

† Norbotten & Vasterbotten

‡Norbotten

INTRODUCTION

In January, 1999, the United States and Canada began international cooperative population-based surveillance for invasive *S. pneumoniae* by all laboratories serving residents of the North American Arctic. In January, 2000, this surveillance system expanded to include invasive diseases with the following organisms: *H. influenzae* (all types), *N. meningitidis*, GAS, and GBS. These pathogens were selected for ICS because rates of these diseases are elevated in indigenous peoples of the north, strains demonstrate resistance to commonly used antibiotics, they are routinely cultured in clinical laboratories, and clinically important serotypes of *S. pneumoniae*, *H. influenzae*, and *N. meningitidis* are vaccine preventable in infants and adults.

Denmark's autonomous region of Greenland joined ICS in 2000; Iceland, Norway (including Svalbard), and Finland joined in 2001; the Northern Sweden regions of Norbotten and Vasterbotten joined in 2003 and 2006, respectively.

GOALS

The goal of ICS is to establish an integrated network of hospital and public health facilities throughout the Arctic countries to monitor infectious diseases of concern. Linking public health facilities within Arctic nations will allow for the collection and sharing of uniform laboratory and epidemiological data that will describe the prevalence of infectious diseases in Arctic populations and assist in the formulation of prevention and control strategies.

The project, initiated in 1998, focused on establishing an ICS system for diseases caused by *S. pneumoniae*. This bacterium causes pneumonia, meningitis, and bacteremia in both the very young and the elderly. Once easily treated with antibiotics, this bacterium has become resistant to commonly used antibiotics. This is of great concern to the public health community and is increasingly a target for surveillance by many countries worldwide. A polysaccharide vaccine is available for use in persons two years of age and older. A conjugate vaccine for infants has been developed and is licensed for use in the U.S., Canada, and the European Union. The fact that diseases caused by *S. pneumoniae* were already being monitored by many public health authorities within the Arctic states made establishing a circumpolar surveillance system for this infection feasible. In addition, due to the availability of polysaccharide and conjugate vaccines, much of the morbidity and mortality caused by *S. pneumoniae* is currently preventable.

ICS objectives include:

- Identify key public health contacts within Arctic countries. These persons should be familiar with infectious disease surveillance systems in place (particularly surveillance systems for diseases caused by *S. pneumoniae*) in the member country. Through correspondence and working group meetings, the scope and gaps of the surveillance systems are determined.
- Determine the comparability of laboratory and data collection methods, and negotiate standard protocols and quality control programs.
- Share and report data in agreed upon formats.
- Form a working group of key laboratory and public health contacts to coordinate pneumococcal surveillance within their respective jurisdictions. This group meets on a regular basis to review problems, progress, compliance, report generation, and future plans.

- Form a steering committee of national Arctic health experts to coordinate new objectives and initiatives within ICS.

This program forms a framework through which surveillance of other infectious diseases as well as prevention and control programs can be added. Other infectious diseases of circumpolar community concern include: other invasive bacterial diseases (caused by *H. influenzae*, *N. meningitidis*, GAS, and GBS), tuberculosis, HIV, hepatitis, foodborne diseases (botulism, brucellosis), waterborne diseases, respiratory diseases of children such as those caused by respiratory syncytial virus, and chronic conditions related to infectious agents (hepatitis B virus and liver cancer, human papilloma virus and cervical cancer, *H. pylori* and stomach cancer), and climate sensitive diseases. In addition, the surveillance model developed by this program for infectious disease may be adapted to monitor other non-infectious human health priorities of community concern.

METHODS

ICS is coordinated by personnel at the Arctic Investigations Program (AIP), Centers for Disease Control and Prevention, in Anchorage, Alaska.

A case of invasive *S. pneumoniae*, *H. influenzae*, *N. meningitidis*, GAS or GBS is defined by the isolation of the bacteria from a normally sterile site, (including blood, cerebrospinal fluid, pleural fluid, peritoneal fluid or joint fluid) that has been taken from a resident of the surveillance area.

In the U.S. Arctic and Northern Canada, laboratory, demographic and clinical data are collected continually by ICS, while in Greenland, Iceland, Northern Sweden, Norway, and Finland, summary data are submitted to ICS in aggregate at the end of the year.

Surveillance System Description by Country/Region

The following table outlines the organisms reported and data provided by each country or region.

Data Provided by Country/Region, ICS 2011

Country	<i>S. pneumoniae</i>	<i>H. influenzae</i>	<i>N. meningitidis</i>	GAS	GBS	Serotype	Demographics	Race/Ethnicity	Risk Factors	Outcome	Vaccination Status	Antibiotic Susceptibility
Finland	X						X					X
Greenland	X	X	X	X	X	X	X			X		
Iceland	X					X	X			X		X
N. Canada	X	X	X	X	X	X	X	X	X	X	X	X
N. Sweden	X	X	X	X	X		X					X
Norway	X	X	X			X*	X			X	X	
U.S. Arctic	X	X	X	X	X	X	X	X	X	X	X	X

*Serotypes/serogroups provided for Hi/Nm only

Finland

- 23 district hospital laboratories participate in ICS.
 - Provide diagnostic microbiology services for all residents of Finland.
 - All invasive isolates of *S. pneumoniae* submitted to the National Public Health Institute (KTL) laboratory in Oulu.
- Antimicrobial susceptibility testing of *S. pneumoniae* isolates was performed by agar dilution method at district hospital laboratories as well as the KTL laboratory.
- Population estimates for 2011 were obtained from the website <http://www.stat.fi>

Greenland

- 15 district hospital laboratories participate in ICS.
 - Provide diagnostic microbiology services for all residents of Greenland.
 - All invasive isolates of *S. pneumoniae*, *H. influenzae*, *N. meningitidis*, GAS, and GBS submitted to reference laboratories in Nuuk and Copenhagen.
- Antimicrobial susceptibility testing of *S. pneumoniae* isolates was performed by agar dilution at the central laboratory at Queen Ingrid's Hospital in Nuuk.
- Serotyping was performed at the Statens Serum Institute in Copenhagen, Denmark, by the Quellung method.
- Clinical and demographic data for every case of invasive *S. pneumoniae*, *H. influenzae*, *N. meningitidis*, GAS, and GBS was collected by public health authorities at the end of the year and entered onto a standardized collection tool, the Bacterial Diseases Surveillance Form (BDSF), which is also used in Iceland, Northern Canada, and the U.S. Arctic.
- Population estimates for 2011 were obtained from the website <http://www.stat.gl>

Iceland

- 10 district hospital laboratories and one regional laboratory participate in ICS.
 - Provide diagnostic microbiology services for all residents of Iceland.
 - All invasive isolates of *S. pneumoniae* submitted to the reference hospital in Reykjavik.
- Antimicrobial susceptibility testing of *S. pneumoniae* isolates is performed by disc diffusion method at the Landspítali University Hospital (LUH) in Reykjavik and the laboratory at the regional hospital in Akureyri. All oxacillin resistant isolates are then analyzed by E test.
- Serotyping is performed at the LUH by coagglutination using antisera from Statens Serum Institute.
- Clinical and demographic data for every case of invasive *S. pneumoniae* was collected by public health authorities at the end of the year and entered onto the same collection form (BDSF) used in Greenland, Northern Canada, and the U.S. Arctic.
- Population estimates for 2011 were obtained from the website <http://www.statice.is>

Northern Canada

- 14 Canadian laboratories participate in ICS.
 - Provide diagnostic microbiology services for all residents of the Yukon Territory, Northwest Territories, Nunavut, Northern Quebec, and Northern Labrador.
 - Submit all invasive isolates of *S. pneumoniae*, *H. influenzae*, *N. meningitidis*, GAS, and GBS to one of two reference laboratories in Canada.
 - *S. pneumoniae*, *H. influenzae*, GAS, and GBS isolates are serotyped by the Quellung method using Statens Serum Institute antisera.
- Antimicrobial susceptibility of *S. pneumoniae*, GAS, and GBS isolates was tested by micro-broth dilution (according to NCCLS recommendations).
- Communicable disease consultants located within one of the five regions of Northern Canada provided clinical and demographic information on the same collection form (BDSF) used in Greenland, Iceland, and the U.S. Arctic.
- Population estimates for 2011 were obtained from the website <http://www.statcan.ca>

Northern Sweden

- 1 district laboratory participates in ICS.
 - Provides diagnostic microbiology services for all residents of Norrbotten and Vasterbotten counties.
 - The main reference laboratory is at the Swedish Institute for Infectious Disease Control in Stockholm.
 - Isolates are serotyped by the Quellung method.
- Antimicrobial susceptibility testing was by disc diffusion at the University Hospital in Umea and Sunderby Hospital in Lulea.
- Population estimates for 2011 were obtained from the website http://www.scb.se/default_2154.asp

Norway

- 33 district hospital laboratories participate in ICS.
 - Provide diagnostic microbiology services for all residents of Norway.
 - All invasive isolates of *S. pneumoniae* submitted to one of two reference laboratories in Oslo or Tromsö.
- Antimicrobial susceptibility testing of *S. pneumoniae* isolates is performed using the disc diffusion method at district hospital laboratories, the reference laboratory in Tromsö or the main national laboratory in Oslo.
- Serotyping is performed at the Statens Serum Institute in Denmark by the Quellung method.
- Population estimates for 2011 were obtained from the website <http://www.ssb.no>

U.S. Arctic


- 23 laboratories providing diagnostic services to residents of Alaska submitted to AIP isolates of *S. pneumoniae*, *H. influenzae*, *N. meningitidis*, GAS, and GBS cultured in blood, cerebrospinal fluid, or from other sterile sites.
 - *S. pneumoniae* and *H. influenzae* isolates are serotyped by the Quellung method using Statens Serum Institute antisera.
 - Serogroup testing of *N. meningitidis* was done using real-time PCR to detect the *ctaA* gene (capsule transport), as well as the genes required for serogroup-specific capsule biosynthesis.
- Antimicrobial susceptibility testing of *S. pneumoniae* isolates is performed at AIP by micro-broth dilution (according to NCCLS recommendations).
- Clinical and demographic information on each case-patient is recorded by AIP research nurses onto the same collection form (BDSF) used in Greenland, Iceland, and Northern Canada.
- Population estimates for 2011 were obtained from the website <http://www.labor.state.ak.us>

Quality Control

Streptococcus pneumoniae: Clinical laboratories forward isolates from patients with invasive pneumococcal disease to reference laboratories in their respective country. To ensure inter-laboratory comparability of *S. pneumoniae* serotyping and antimicrobial susceptibility testing between two reference laboratories in Canada (Alberta and Quebec) and one in the U.S. (Alaska), the ICS *S. pneumoniae* inter-laboratory quality control (QC) program was established in 1999. Statens Serum Institute, Copenhagen, Denmark, joined the program in 2004. Each reference laboratory is responsible for exporting one QC panel of seven *S. pneumoniae* isolates every other year to each of the other laboratories [1].

Neisseria meningitidis/*Haemophilus influenzae*: An interlaboratory quality control program for *Neisseria meningitidis* and *Haemophilus influenzae* was established in 2005. Participating laboratories include the National Microbiology Laboratory, Winnipeg, Manitoba, Canada; Arctic Investigations Program, Anchorage, Alaska, USA; Laboratoire Santé Publique du Québec, Sainte-Anne-de-Bellevue, Québec, Canada; National Centre for Streptococcus, Edmonton, Alberta, Canada; and Statens Serum Institut, Copenhagen, Denmark. Strain panels are distributed twice a year from the National Microbiology Laboratory or Arctic Investigations Program [2].

Participating Countries, ICS 2011



RESULTS

Streptococcus pneumoniae

Case Demographics

A total of 1,803 cases of invasive disease caused by *S. pneumoniae* were reported to ICS during 2011 by Finland, Greenland, Iceland, N. Canada, N. Sweden, Norway, and the U.S. Arctic. The highest rates of disease (23.4 per 100,000) occurred in N. Canada and the lowest in Iceland (10.4 per 100,000) with an overall rate for the ICS circumpolar region of 14.9 per 100,000; 55% of all cases occurred in males. The median age of cases overall was 61.5 years with the lowest median age in N. Canada (33 years) and the highest in N. Sweden (67 years). Case fatality ratios ranged from 11% in Norway to 40% in Greenland; the overall case fatality ratio was 13%.

Streptococcus pneumoniae Case Demographics, ICS 2011 Data

Country	Population	#		Sex		Median Age (min-max) yrs	Deaths n (CFR [†])
		Cases	Rate*	M (%)	F (%)		
Finland	5,401,267	782	14.5	456 (58)	59 (0-101)	‡	
Greenland	56,615	6	10.6	4 (67)	39 (7.2-53.4)	2 (40%) ^b	
Iceland	318,452	33	10.4	23 (70)	62 (1-93)	8 (24%)	
N. Canada	154,162	36	23.4	22 (61)	33 (0.2-79.4)	5 (14%) ^b	
N. Sweden	508,039 ^a	76	15	32 (42)	67 (1.5-95.5)	‡	
Norway	4,920,305	729	14.8	372 (51)	65.8 (0-96.8)	35 (11%) ^b	
U.S. Arctic	723,424	141	19.5	84 (60)	52 (0.2-87.8)	21 (15%)	
Total	12,082,264	1,803	14.9	993 (55)	61.5 (0-101)	71 (13%)	

*Number of cases per 100,000 per year

†Case fatality ratio

‡Case outcomes not reported from Finland, N. Sweden

^a Population for Norbotten and Vasterbotten

^b Case outcomes unknown in 1 case from Greenland, 1 case from N. Canada, 398 cases from Norway

Streptococcus pneumoniae by Age Category, ICS 2011 Data

Age		Finland	Greenland	Iceland	N. Canada	N. Sweden	Norway	U.S. Arctic
<2 yrs	Pop	121,578	1,753	9,826	5,686	9,979	124,613	22,194
	N (%) [*]	46 (6)	0 (0)	2 (6)	7 (19)	1 (1)	18 (3)	4 (3)
	Rate [†]	37.8	0	20.4	123.1	10	14.4	18
2-19 yrs	Pop	1,095,170	15,445	80,017	44,950	99,569	1,120,720	186,816
	N (%) [*]	62 (8)	2 (33)	2 (6)	8 (22)	2 (3)	35 (5)	14 (10)
	Rate [†]	5.7	13	2.5	17.8	2	3.1	7.5
20-64 yrs	Pop	3,204,879	35,470	189,436	94,672	293,361	2,932,729	455,553
	N (%) [*]	377 (48)	4 (67)	16 (49)	18 (50)	29 (38)	292 (40)	93 (66)
	Rate [†]	11.8	11.3	8.5	19	9.9	10	20.4
65+ yrs	Pop	979,640	3,947	39,173	8,854	105,130	742,243	58,861
	N (%) [*]	297 (38)	0 (0)	13 (39)	3 (8)	44 (58)	384 (53)	30 (21)
	Rate [†]	30.3	0	33.2	33.9	41.9	51.7	51
All ages	Pop	5,401,267	56,615	318,452	154,162	508,039	4,920,305	723,424
	N	782	6	33	36	76	729	141
	Rate [†]	14.5	10.6	10.4	23.4	15	14.8	19.5

*Proportion of total cases in each country/region

†Number of cases per 100,000 per year

When stratified by age, the highest rates of disease in most countries occurred either in those cases less than two years of age or in cases 65+ years of age. In Greenland, there were no cases in either of these age categories and in the U.S. Arctic, the highest rates were in the 20-64 years and 65+ years of age categories.

Seasonality

S. pneumoniae was diagnosed throughout the year in 2011 in each country. For all countries except Greenland, higher proportions of disease were seen in the first and second quarters of the year with declines during the third quarter and rising again during the fourth quarter. In Greenland, the highest proportion of cases was seen in the third quarter and the lowest in the second quarter of the year, however, due to the small number of cases, trends in seasonality cannot be described.

Race

Race and ethnicity data were collected in N. Canada and the U.S. Arctic. Rates of invasive pneumococcal disease were higher overall in Aboriginal and AK Native populations than in non-Aboriginal and non-AK Native populations. The highest rates of disease occurred in Aboriginal children less than 2 years of age in N. Canada and AK Native adults 65+ in the U.S. Arctic.

Streptococcus pneumoniae by Race and Age Categories, ICS 2011 Data

Age (yrs)	N. Canada*		U.S. Arctic		
	Aboriginal	Non-Aboriginal	AK Native	Non-AK Native	
<2	Population	3,928	1,758	6,518	15,676
	Cases (rate‡)	7 (178.2)	0 (0)	2 (30.7)	2 (12.8)
2-19	Population	35,444	9,506	49,769	137,047
	Cases (rate‡)	6 (16.9)	2 (21)	5 (10.1)	9 (6.6)
20-64	Population	45,212	49,460	77,168	378,385
	Cases (rate‡)	13 (28.8)	3 (6.1)	42 (54.4)	51 (13.5)
65+	Population	3,699	5,155	8,707	50,154
	Cases (rate‡)	2 (54.1)	1 (19.4)	7 (80.4)	23 (45.9)
All	Population	88,283	65,879	142,162	581,262
Ages	Cases (rate‡)	28 (31.7)	6 (9.1)	56 (39.4)	85 (14.6)

*Race unknown in 2 cases 20-64 years

‡Number of cases per 100,000 per year

Clinical Presentation

The primary clinical presentation was determined by a review of the discharge diagnoses in each patient's individual medical record associated with the invasive bacterial illness. In cases with multiple discharge diagnoses, the most serious diagnosis related to the pneumococcal infection was recorded as the primary clinical presentation. The most common clinical presentations associated with *S. pneumoniae* were pneumonia, bacteremia, and meningitis. In Greenland, N. Canada, and the U.S. Arctic the clinical presentation reported most often was pneumonia (100%, 47%, and 66%, respectively), in N. Sweden and Norway it was bacteremia (91% and 61%, respectively).

Clinical Presentation of Reported *Streptococcus pneumoniae* Cases, ICS 2011 Data

	Finland n (%)	Greenland n (%)	Iceland n (%)	N Canada n (%)	N Sweden** n (%)	Norway n (%)	US Arctic n (%)
Pneumonia*	0 (0)	6 (100)	0 (0)	17 (47)	0 (0)	195 (27)	93 (66)
Bacteremia	742 (95)	0 (0)	24 (73)	12 (33)	31 (91)	446 (61)	25 (18)
Meningitis	40 (5)	0 (0)	8 (24)	2 (6)	3 (9)	48 (7)	8 (6)
Empyema	0 (0)	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	3 (2)
Peritonitis	0 (0)	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	3 (2)
Septic arthritis	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	0 (0)	0 (0)
Endocarditis	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
Cellulitis	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	2 (1)
Pericarditis	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
Osteomyelitis	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
Epiglottitis	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
Other	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	39 (5)	1 (1)
Unknown	0 (0)	0 (0)	0 (0)	3 (8)	0 (0)	1 (<1)	0 (0)
Total Cases	782	6	33	36	34	729	141

*with bacteremia

**Clinical presentation data reported from Vasterbotten only

Risk Factors

N. Canada and the U.S. Arctic reported medical conditions or risk factors associated with *S. pneumoniae*. In adults 18 years and older, cigarette smoking and alcohol abuse were the most common associated conditions in N. Canada and the U.S. Arctic.

Streptococcus pneumoniae Risk Factor/Medical Conditions in Adults*, ICS 2011 Data†

	N. Canada n (%)	U.S. Arctic n (%)
Cigarette Smoking	8 (38)	59 (47)
Alcohol Abuse	7 (33)	39 (31)
Chronic Lung Disease and/or Asthma	2 (10)	32 (26)
Immunosuppressive Therapy	0 (0)	6 (5)
Diabetes	2 (10)	28 (22)
Injection Drug Use	1 (5)	2 (2)
Asplenia	0 (0)	0 (0)
Total Adult* Cases	21	125

*≥ 18 years

†Multiple risk factors may be reported per case

Vaccination Policy

In Finland, Iceland, N. Canada, Norway, and the U.S. Arctic, 23-valent pneumococcal polysaccharide vaccine (PS23) is recommended for persons over 60 years (Iceland) or over 65 years of age (Finland, N. Canada, Norway, U.S. Arctic), and to persons greater than two years of age (Finland, Iceland, Norway, U.S. Arctic) or greater than five years of age (N. Canada) with specific medical problems. The vaccine is only recommended for certain risk groups in N. Sweden. Three pneumococcal conjugate vaccines (PCV), PCV7, PCV10, and PCV13, for use in children under the age of 2 years have been used in ICS regions since 2001. Vaccines used and years introduced are presented in the table below.

Pneumococcal Vaccines Introduced, ICS 2011 Data

	Finland	Greenland	Iceland	N Canada	N Sweden	Norway	US Arctic
PCV7				2002-2006*	2007	2007	2001
PCV10	2010		2011	2009-2010*			
PCV13		2010		2010-2011*	2010	2011	2010
PS23	†	1996	†	1988	1994	†	1983

*PCV7 – N Quebec and Nunavut (2002), Newfoundland and Yukon (2005), Northwest Territory (2006); PCV10 – Nunavut (2010); PCV13 – N Quebec (2011)

†PS23 used but year of introduction unknown

Vaccination Status

Thirteen percent of *S. pneumoniae* cases in children less than 2 years of age with known vaccination status were vaccinated with a PCV in Norway; in N. Canada and the U.S. Arctic, 100% of children with known vaccine status were vaccinated. Only 4% of cases eligible for PS23 in Norway were vaccinated indicating much less frequent use of this vaccine than in N. Canada and the U.S. Arctic. Vaccine coverage data were not reported from Finland, Iceland and N. Sweden.

Streptococcus pneumoniae Case Vaccination Status for Pneumococcal Vaccine, ICS 2011 Data

	N. Canada	Norway	U.S. Arctic
Total cases eligible for PCV7 vaccine*	7	18	4
Vaccine status known in cases eligible for PCV7	6	15	2
Cases eligible for PCV7 vaccinated (%)[†]	6 (100%)	2 (13%)	2 (100%)
Total cases eligible for PS23 vaccine[‡]	3	383	30
Vaccine status known in cases eligible for PS23	1	73	18
Cases eligible for PS23 vaccinated (%)[†]	1 (100%)	3 (4%)	15 (83%)

*Children less than 2 years of age

†Percent of vaccine status known cases

‡Adults 65 years and older

Serotypes

The *S. pneumoniae* serotypes reported by ICS regions in 2011 are listed in the following table; yellow highlights the most common serotypes in each country.

Streptococcus pneumoniae Serotypes by Country, ICS 2011 Data

Serotype	Greenland n (% [†])	Iceland n (% [†])	N. Canada (% [†])	N. Sweden* n (% [†])	U.S. Arctic n (% [†])
1[‡]	2 (67)	0 (0)	0 (0)	0 (0)	1 (1)
3[‡]	0 (0)	2 (6)	1 (3)	5 (15)	12 (9)
4[‡]	0 (0)	2 (6)	1 (3)	0 (0)	0 (0)
6	0 (0)	0 (0)	0 (0)	6 (18)	0 (0)
6A[‡]	0 (0)	2 (6)	1 (3)	0 (0)	0 (0)
6C	0 (0)	1 (3)	1 (3)	0 (0)	6 (5)
7	0 (0)	0 (0)	0 (0)	2 (6)	0 (0)
7C	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
7F[‡]	1 (33)	2 (6)	6 (20)	0 (0)	28 (22)

Serotype	Greenland n (% †)	Iceland n (% †)	N. Canada (% †)	N. Sweden* n (% †)	U.S. Arctic n (% †)
8	0 (0)	0 (0)	0 (0)	2 (6)	11 (9)
9	0 (0)	0 (0)	0 (0)	1 (3)	0 (0)
9N	0 (0)	0 (0)	2 (7)	0 (0)	4 (3)
10A	0 (0)	0 (0)	4 (13)	0 (0)	2 (2)
11A	0 (0)	0 (0)	3 (10)	0 (0)	3 (2)
12F	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
14‡	0 (0)	6 (18)	0 (0)	0 (0)	0 (0)
15	0 (0)	0 (0)	0 (0)	1 (3)	0 (0)
15A	0 (0)	0 (0)	0 (0)	0 (0)	6 (5)
15B	0 (0)	0 (0)	1 (3)	0 (0)	2 (2)
15B/C	0 (0)	1 (3)	0 (0)	0 (0)	0 (0)
15C	0 (0)	0 (0)	1 (3)	0 (0)	1 (1)
16F	1 (0)	0 (0)	0 (0)	0 (0)	6 (5)
17F	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
18	0 (0)	0 (0)	0 (0)	1 (3)	0 (0)
18F	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)
19	0 (0)	0 (0)	0 (0)	1 (3)	0 (0)
19A‡	0 (0)	5 (15)	1 (3)	0 (0)	8 (6)
19F‡	1 (0)	3 (9)	0 (0)	0 (0)	2 (2)
20	0 (0)	1 (3)	0 (0)	0 (0)	3 (2)
22	0 (0)	2 (6)	0 (0)	6 (18)	0 (0)
22F	1 (0)	0 (0)	1 (3)	0 (0)	10 (8)
23	0 (0)	0 (0)	0 (0)	8 (24)	0 (0)
23A	0 (0)	1 (3)	0 (0)	0 (0)	4 (3)
23B	0 (0)	0 (0)	1 (3)	0 (0)	2 (2)
29	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
31	0 (0)	0 (0)	0 (0)	0 (0)	6 (5)
33	0 (0)	1 (3)	0 (0)	0 (0)	0 (0)
33A	0 (0)	0 (0)	1 (3)	0 (0)	1 (1)
33F	0 (0)	3 (9)	3 (10)	0 (0)	1 (1)
34	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
35	0 (0)	0 (0)	0 (0)	1 (3)	0 (0)
35B	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
35F	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)
38	0 (0)	1 (3)	0 (0)	0 (0)	3 (2)
42	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)
# Isolates Serotyped	6	33	30	34	129

*Serogroup level data from Vasterbotten only

†Percent of total isolates serotyped

‡Serotype included in PCV

Cases and Deaths with Vaccine Serotypes

For the countries reporting serotype data, more than 69% of *S. pneumoniae* cases in persons ≥ 2 years of age had a serotype that was included in the 23-valent polysaccharide vaccine. Twenty percent to 100% of *Strep pneumoniae* cases in children < 2 years of age had a serotype contained in the PCV13 vaccine. The proportion

of deaths with serotypes in the 23-valent polysaccharide vaccine related to *S. pneumoniae* ranged from 50% to 100%.

Proportion of Cases/Deaths with Invasive Pneumococcal Disease Vaccine Serotypes, ICS 2011 Data

	Greenland n/Denom* (%)	Iceland n/Denom* (%)	N. Canada n/Denom* (%)	U.S. Arctic n/Denom* (%)
Cases ≥ 2 years old with serotype in the 23-valent pneumococcal polysaccharide vaccine	5/6 (83)	24/28** (86)	19/25 (76)	86/125 (69)
Cases < 2 years old with serotype in the 13-valent pneumococcal conjugate vaccine	No cases	2/2 (100)	1/5 (20)	2/4 (50)
Deaths (all ages) for which the serotype was contained in the 23-valent pneumococcal vaccine	1 (50‡)	6 (100‡)	3 (60‡)	14 (70‡)

*Number of isolates serotyped by country by age group

**3 isolates tested to serogroup level only

‡Percentage of total death

†Outcomes not reported

Outcome

A total of 71 deaths associated with *S. pneumoniae* were reported to ICS in 2011. Overall, the highest case fatality ratio (CFR) occurred in persons 65+ years of age (17%). Finland and N. Sweden did not report outcome data.

Streptococcus pneumoniae Age-Specific Case-Fatality Ratios (CFR), ICS 2011 Data

		<2 years	2-19 years	20-64 years	65+ years	All Ages
Greenland	Deaths/Cases* (CFR)	0/0 (0%)	0/2 (0%)	2/3† (67%)	0/0 (0%)	2/5 (40%)
Iceland	Deaths/Cases* (CFR)	0/2 (0%)	0/2 (0%)	4/16 (25%)	4/13 (31%)	8/33 (24%)
N. Canada	Deaths/Cases* (CFR)	0/7 (0%)	0/8 (0%)	5/18 (28%)	0/2† (0%)	5/35 (14%)
Norway	Deaths/Cases* (CFR)	2/17† (12%)	1/25† (4%)	8/125† (6%)	24/164† (15%)	35/331 (11%)
U.S. Arctic	Deaths/Cases* (CFR)	0/4 (0%)	1/14 (7%)	12/93 (13%)	8/30 (27%)	21/141 (15%)
Total	Deaths/Cases* (CFR)	2/28 (7%)	2/51 (4%)	31/255 (12%)	36/209 (17%)	71/545 (13%)

*Cases with known outcome.

†Outcome unknown in (1) Norway case < 2 years; (10) Norway cases 2-19 years; (1) Greenland, (167) Norway cases 20-64 years; (1) N Canada, (220) Norway cases 65+ years

Antimicrobial Susceptibility

In 2011, antimicrobial susceptibility results were reported to ICS from Finland, Iceland, N. Canada, N. Sweden and the U.S. Arctic. The following table outlines for each country the antibiotics tested, the number of isolates reported tested for each antibiotic and the proportion (NS%) of isolates tested that showed any level of non-susceptibility.

***Streptococcus pneumoniae* Antibiotic Susceptibility Testing, ICS 2011 Data**

Antibiotic	Finland n (NS%)	Iceland n (NS%)	N. Canada n (NS%)	N. Sweden n (NS%)	U.S. Arctic n (NS%)
Ceftriaxone	-	33 (0%)	27 (0%)	-	129 (4%)
Chloramphenicol	-	33 (3%)	23 (0%)	-	129 (1%)
Clindamycin	-	-	26 (4%)	-	129 (3%)
Erythromycin	-	33 (24%)	25 (12%)	34 (6%)	129 (14%)
Levofloxacin	-	-	25 (0%)	-	129 (1%)
Penicillin	782 (2%)	33 (9%)	27 (0%)	34 (3%)	129 (16%)
Rifampin	-	-	-	-	129 (1%)
Tetracycline	-	33 (15%)	-	-	129 (6%)
TMP Sulfa*	-	33 (21%)	26 (12%)	33 (6%)	129 (10%)
Vancomycin	-	-	29 (0%)	-	129 (0%)

*Trimethoprim-sulfamethoxazole

Of the antibiotics tested, the largest numbers of isolates were resistant to penicillin, TMP-Sulfa and erythromycin. Results by country for each of these antibiotics are shown in the tables below. In the U.S. Arctic, 15 isolates comprised of 9 serotypes (3, 7F, 6C, 15A, 15B, 15C, 19A, 29, 35B) showed multi-drug resistance. The largest proportion (n = 6, 40%) of the multi-drug resistant isolates was serotype 19A and 5 of the 6 (83%) of isolates with this serotype were resistant to three or more antibiotics. In N. Canada, one serotype 19A isolate was resistant to four antibiotics. In Iceland, 5 isolates including serotypes 6C, 14, 19F, and 38 were resistant to more than one antibiotic. Finland and N. Sweden did not report serotypes.

***Streptococcus pneumoniae* Penicillin Susceptibility Results, ICS 2011 Data**

	# Tested	I* (%)	I* Serotypes (n)	R* (%)	R* Serotypes (n)
Finland	782	18 (2%)	†	1 (<1%)	†
Iceland	33	4 (12%)	19F (2), 14 (1), 38 (1)	0 (0%)	
N. Canada	22	0 (0%)		0 (0%)	
N. Sweden‡	83	1 (3%)	19 (1)	0 (0%)	
U.S. Arctic	129	17 (13%)	19A (5), 23A (4), 7F (3), 6C (2), 15A (1), 15B (1), 15C (1)	4 (3%)	19A (2), 29 (1), 35B (1)

*I=Intermediate resistance, R=Fully resistant

†Serotypes not reported

‡N. Sweden reported serogroups only

***Streptococcus pneumoniae* TMP-Sulfa Susceptibility Results, ICS 2011 Data**

	# Tested	I* (%)	I* Serotypes (n)	R* (%)	R* Serotypes (n)
Iceland	33	1 (3%)	15B/C (1)	7 (11%)	19F (2), 33F (2), 6A (1), 14 (1), 38 (1)
N. Canada	23	1 (4%)	33A (1)	2 (9%)	19A (1), 23B (1)
N. Sweden†	33	0 (0%)		2 (6%)	9 (1), 19 (1)
U.S. Arctic	129	5 (4%)	15A (1), 15B (1), 15C (1), 33A (1), 33F (1)	8 (6%)	19A (6), 6C (2)

*I=Intermediate resistance, R=Fully resistant

†N. Sweden reported serogroups only

***Streptococcus pneumoniae* Erythromycin Susceptibility Results, ICS 2011 Data**

	# Tested	I* (%)	I* Serotypes	R* (%)	R* Serotypes (n)
Iceland	33	0 (0%)		8 (24%)	14 (4), 19F (2), 6C (1), 38 (1)
N. Canada	25	0 (0%)		3 (12%)	19A (1), 33A (1), 33F (1)
N. Sweden†	34	1 (3%)	6 (1)	1 (3%)	23 (1)
U.S. Arctic	129	0 (0%)		18 (14%)	19A (5), 31 (4), 3 (1), 6C (2), 15A (1), 15B (1), 15C (1), 19F (1), 22F (1), 42 (1)

*I=Intermediate resistance, R=Fully resistant

†N. Sweden reported serogroups only

Quality Control

In 2011, two QC panels of seven *S. pneumoniae* isolates plus a control strain each were shipped and tested. Beginning in 2005, the MIC data was analyzed in two ways. The first comparison is based on MIC data provided by the distributing laboratory and is the method used historically in the QC program. The new methodology provides an analysis of MIC data for all the participating laboratories by using the modal MIC for each antibiotic-organism combination as the value to which the other data are compared. The modal MIC is the MIC most frequently reported. When two MIC values were reported with equal frequency, both were accepted as a modal value. Antibiotic-organism combinations for which there was no consensus on a modal MIC were excluded from the analysis. The rationale for initiating the second analysis recognizes that there is an allowable variation of one log₂ dilution inherent for any MIC testing system. This means that there is no one absolutely correct MIC value to which all others can be compared. The modal MIC may be a better representation of this ‘true’ value than any one laboratory can provide. For Panel 2011-A, overall serotyping correlation was 100%. The modal MIC comparison resulted in an overall correlation of 98.6% with individual participant correlation ranging from 75%-100%. Overall category agreement was 98.2% with individual participant agreement ranging from 93.7%-100%. For Panel 2011-B, overall serotyping correlation was 100%. The modal MIC comparison resulted in an overall correlation of 100%. Overall category agreement was 99.3% with individual participant agreement ranging from 98.4%-100%.

Conclusions

Streptococcus pneumoniae remains a major cause of invasive bacterial disease in circumpolar regions. Disease rates are highest in indigenous populations. All ICS regions have introduced either PCV10 or PCV13 into their vaccine schedules. Monitoring the impact of these conjugate vaccines will be important to confirm effectiveness and provide support for continuing immunization programs.

Haemophilus influenzae

Case Demographics

Greenland, N. Canada, N. Sweden (Norrbotten), Norway and the U.S. Arctic reported the occurrence of invasive *H. influenzae* disease during 2011. Greenland reported no cases. A total of 127 cases of invasive disease caused by *H. influenzae* were reported to ICS during 2011 by N. Canada, N. Sweden (Norrbotten), Norway and the U.S. Arctic. The highest rate of disease among regions reporting cases was in N. Canada (9.7 per 100,000) and the lowest in N. Sweden (1.2/100,000). Median age of cases was highest in N. Sweden (68 years) and lowest in N. Canada (1.8 years).

Haemophilus influenzae Case Demographics, ICS 2011 Data

Country	Population	# Cases	Rate*	Sex M (%)	Median Age (min-max) yrs	Deaths n (CFR†)
Greenland	56,615	0	0	No cases	No cases	No cases
N. Canada	154,162	15	9.7	6 (40)	1.8 (0.3-47.5)	3 (21%)‡
N. Sweden	248,545	3	1.2	0 (0)	68 (64-87)	‡
Norway	4,920,305	85	1.7	45 (53)	60.4 (0-95.6)	6 (10%)‡
U.S. Arctic	723,424	24	3.3	12 (50)	18.2 (0-101.1)	2 (8%)
Total	6,103,051	127	2.1	63 (50)	55.1 (0-101.1)	11 (11%)

*Number of cases per 100,000 per year

†Case fatality ratio

‡ Case outcome unknown in (1) N. Canada, (27) Norway cases; N. Sweden did not report case outcomes

When stratified by age, the highest rates of disease for N. Canada and the U.S. Arctic were in the <2 years age category; in N. Sweden and Norway, highest rate of disease was in the 65+ years age category.

Haemophilus influenzae by Age Category, ICS 2011 Data

Age		N. Canada	N. Sweden	Norway	U.S. Arctic
<2 yrs	Population	5,686	4,901	124,613	22,194
	Cases (%)*	8 (53)	0 (0)	2 (2)	12 (50)
	Rate†	140.7	0	1.6	54.1
2-19 yrs	Population	44,950	47,830	1,120,720	186,816
	Cases (%)*	2 (13)	0 (0)	8 (9)	0 (0)
	Rate†	4.5	0	0.7	0
20-64 yrs	Population	94,672	142,118	2,932,729	455,553
	Cases (%)*	5 (33)	1 (33)	40 (47)	6 (25)
	Rate†	5.3	0.7	1.4	1.3
65+ yrs	Population	8,854	53,696	742,243	58,861
	Cases (%)*	0 (0)	2 (67)	35 (41)	6 (25)
	Rate†	0	3.7	4.7	10.2
All ages	Population	154,162	248,545	4,920,305	723,424
	Cases	15	3	85	24
	Rate†	9.7	1.2	1.7	3.3

*Proportion of total cases in each country/region

†Number of cases per 100,000 per year

Race

Rates of disease were highest in N. Canada Aboriginal (203.7 per 100,000) and U.S. Arctic AK Native (168.8 per 100,000) cases less than two years of age. In the U.S. Arctic, rates of disease were higher in Native populations than in non-Native populations in all age categories.

***Haemophilus influenzae* by Race and Age Categories, ICS 2011 Data**

Age (yrs)	N. Canada		U.S. Arctic		
	Aboriginal	Non-Aboriginal	AK Native	Non-AK Native	
<2	Population	3,928	1,758	6,518	15,676
	Cases (rate†)	8 (203.7)	0 (0)	11 (168.8)	1 (6.4)
2-19	Population	35,444	9,506	49,769	137,047
	Cases (rate†)	2 (5.6)	0 (0)	0 (0)	0 (0)
20-64	Population	45,212	49,460	77,168	378,385
	Cases (rate†)	4 (8.9)	1 (2)	2 (2.6)	4 (1.1)
65+	Population	3,699	5,155	8,707	50,154
	Cases (rate†)	0 (0)	0 (0)	2 (23)	4 (8)
All	Population	88,283	65,879	142,162	581,262
Ages	Cases (rate†)	14 (15.9)	1 (1.5)	15 (10.6)	9 (1.6)

†Number of cases per 100,000 per year

Clinical Presentation

The primary clinical presentation was determined by a review of the discharge diagnoses in each patient's individual medical record associated with the invasive bacterial illness. In cases with multiple discharge diagnoses, the most serious diagnosis related to the infection was recorded as the primary clinical presentation. In N. Canada and the U.S. Arctic, the most common clinical presentation associated with *H. influenzae* was pneumonia (47% and 29% of reported cases, respectively). In Norway, the most common clinical presentation was bacteremia (46%). N. Sweden did not report clinical presentation data.

***Clinical Presentation of Reported Haemophilus influenzae* Cases, ICS 2011 Data**

	N. Canada n (%)	Norway n (%)	U.S. Arctic n (%)
Pneumonia*	7 (47)	21 (25)	7 (29)
Bacteremia	4 (27)	39 (46)	4 (17)
Meningitis	1 (7)	5 (6)	4 (17)
Cellulitis	1 (7)	0 (0)	2 (8)
Septic arthritis	0 (0)	0 (0)	1 (4)
Empyema	0 (0)	0 (0)	3 (13)
Endocarditis	0 (0)	0 (0)	1 (4)
Epiglottitis	0 (0)	0 (0)	1 (4)
Other/Unknown	2 (13)	20 (23)	1 (4)
Total	15	85	24

*with bacteremia

Risk Factors

Fifty percent of adult (≥ 18 years) cases of *H. influenzae* reported in the U.S. Arctic indicated diabetes as an associated risk factor; 25% indicated smoking, 17% alcohol abuse and 8% chronic lung disease or immune suppressive treatment as an associated risk factor. Forty percent of adult N. Canadian *H. influenzae* cases reported smoking or alcohol abuse as a risk factor and 20% reported chronic lung disease. N. Sweden and Norway did not report risk factor data.

Vaccination Status

The *H. influenzae* type b (Hib) conjugate vaccine is required as part of routine childhood vaccination in N. Canada, N. Sweden, Norway and the U.S. Arctic. One case of Hib was reported in N. Canada in children less than five years; the case had received Hib vaccine. Two cases of Hib were reported in the U. S. Arctic in children less than five years; both cases had received Hib vaccine. Data on vaccine status were not provided by N. Sweden and Norway.

***Haemophilus influenzae* Case Vaccination Status for Hib Vaccine, ICS 2011 Data**

	N. Canada	U.S. Arctic
Total cases* eligible for Hib vaccine†	8	12
Vaccine status known in cases* eligible for Hib vaccine	8	12
Cases* eligible for Hib vaccine vaccinated (%)‡	8 (100%)	9 (75%)

*All serotypes

†Children less than 5 years of age

‡Percent of vaccine status known cases

Serotypes

***Haemophilus influenzae* Serotypes by Country, ICS 2011 Data**

Serotype	N. Canada n (%)	Norway n (%)	U.S. Arctic n (%)
a	12 (86)	1 (1)	7 (30)
b	1 (2)	2 (2)	2 (9)
c	1 (2)	0 (0)	0 (0)
e	0 (0)	1 (1)	1 (4)
f	0 (0)	16 (19)	3 (13)
Non-typeable	0 (0)	65 (77)	10 (44)
Total*	14	85	23

*Number of isolates serotyped

The most common *H. influenzae* serotype in N. Canada was type a (86% of cases), in Norway, type f (19% of cases) and in the U.S. Arctic it was serotype a (30% of cases). Non-typeable cases also made up a large proportion of cases in Norway (77%) and the U.S. Arctic (44%). N. Sweden did not provide serotype data.

Outcome

Eleven deaths were associated with *H. influenzae* cases reported to ICS in 2011; three from N. Canada, two from the U.S. Arctic and six from Norway. N. Sweden did not provide outcome data.

Haemophilus influenzae Deaths by Country, ICS 2011 Data

	N. Canada	Norway	U.S. Arctic
No. Deaths	3	6	2
Case Fatality Ratio	21%*	10%*	8%
Min-max (yrs)	0.5-47.5	61-95.6	0.4-34.8
Serotypes (n)	a (3)	NT (6)	a (1), NT (1)
Clinical Presentation (n)	Bacteremia (2), unknown (1)	Bacteremia (4), pneumonia (1), meningitis (1)	Meningitis (1), pneumonia (1)

*Case outcome unknown in (1) N. Canada, (27) Norway cases

Quality Control

Two QC panels of five *H. influenzae* and *N. meningitidis* isolates were shipped and serotyped or serogrouped. There were two discrepant results, one of which was due to the primer set available that did not include all possible serogroups [2].

Conclusions

Widespread use of Hib conjugate vaccines has led to the virtual disappearance of Hib disease in these populations. Substantial replacement with other serotypes has not occurred. The proportion of disease caused by non-typeable organisms continues to be substantial.

Neisseria meningitidis

Case Demographics

Greenland, N. Canada, N. Sweden (Norrbotten), Norway and the U.S. Arctic reported the occurrence of *N. meningitidis* during 2011. A total of 43 cases of invasive disease caused by *N. meningitidis* were reported to ICS. N. Canada had the highest disease rate (1.3 per 100,000). Four deaths associated with *N. meningitidis* were reported from Norway and one from N. Canada.

Neisseria meningitidis Case Demographics, ICS 2011 Data

Country	Population	# Cases	Rate* Rate*	Sex M (%)	Median Age (min-max) yrs	Deaths n (CFR†)
Greenland	56,615	0	0	No cases	No cases	No cases
N. Canada	154,162	2	1.3	1 (50)	1.2 (0.4-2)	1 (50)
N. Sweden	248,545	2	0.8	1 (50)	54 (51-57)	‡
Norway	4,920,305	37	0.8	24 (65)	27.8 (0.6-91.3)	4 (14) ‡
U.S. Arctic	723,424	2	0.3	1 (50)	27.7 (13.8-41.5)	0 (0)
Total	6,103,051	43	0.7	27 (63)	27.8 (0.4-91.3)	5 (15)

*Number of cases per 100,000 per year

†Case fatality ratio

‡Outcome unknown in 9 Norway cases; N. Sweden did not report outcomes

The following table shows cases and rates stratified by age category.

Neisseria meningitidis by Age Category, ICS 2011 Data

Age		N. Canada	N. Sweden	Norway	U.S. Arctic
<2 yrs	Population	5,686	4,901	124,613	22,194
	Cases (%)*	1 (50)	0 (0)	6 (16)	0 (0)
	Rate†	17.6	0	4.8	0
2-19 yrs	Population	44,950	47,830	1,120,720	186,816
	Cases (%)*	1 (50)	0 (0)	11 (30)	1 (50)
	Rate†	2.2	0	1	0.5
20-64 yrs	Population	45,212	49,460	77,168	378,385
	Cases (%)*	0 (0)	2 (100)	12 (32)	1 (50)
	Rate†	0	1.4	0.4	0.2
65+ yrs	Population	8,854	53,696	742,243	58,861
	Cases (%)*	0 (0)	0 (0)	8 (22)	0 (0)
	Rate†	0	0	1.1	0
All ages	Population	154,162	248,545	4,920,305	723,424
	Cases	2	2	37	2
	Rate†	1.3	0.8	0.8	0.3

*Proportion of total cases in each country/region

†Number of cases per 100,000 per year

Race

In the U.S. Arctic, one case of *N. meningitidis* occurred in a non-AK Native person (rate 0.2/100,000) and one in an AK Native person (rate 0.7/100,000). In N. Canada, one Nm case occurred in a non-Aboriginal person (rate 1.5/100,000) and one in an Aboriginal person (rate 1.1/100,000).

Clinical Presentation

The clinical presentation reported for the two *N. meningitidis* cases both in N. Canada and the U.S. Arctic was bacteremia. In Norway, the most common presentation was meningitis (46%). N. Sweden did not report clinical presentations.

Clinical Presentation of Reported *Neisseria meningitidis* Cases, ICS 2011 Data

	N. Canada	Norway	U.S. Arctic
	n (%)	n (%)	n (%)
Pneumonia*	0 (0)	8 (22)	0 (0)
Bacteremia	2 (100)	9 (24)	2 (100)
Meningitis	0 (0)	17 (46)	0 (0)
Other	0 (0)	2 (5)	0 (0)
Unknown	0 (0)	1 (3)	0 (0)
Total	2	37	2

*with bacteremia

Serogroups

Forty of 43 cases of invasive *N. meningitidis* reported to ICS in 2011 included serogroup data. The table below lists serogroups by country. N. Sweden did not report serogroup data.

***Neisseria meningitidis* Serogroups by Country, ICS 2011 Data**

Serogroup	N. Canada	Norway	U.S. Arctic
	n (%)	n (%)	n (%)
B	2 (100)	10 (27)	1 (50)
C	0 (0)	4 (11)	1 (50)
W135	0 (0)	2 (5)	0 (0)
Y	0 (0)	20 (54)	0 (0)
Unknown	0 (0)	1 (3)	0 (0)
Total	2	37	2

Conclusions

Neisseria meningitidis is a relatively uncommon cause of invasive bacterial disease in the circumpolar area under surveillance, however, continued surveillance is warranted due to high morbidity and mortality associated with outbreaks.

Group A *Streptococcus*

Case Demographics

Greenland, N. Canada, N. Sweden (Norrbotten) and the U.S. Arctic each reported the occurrence of GAS during 2011. A total of 90 cases of invasive disease caused by GAS were reported to ICS. Among regions reporting cases, the rate of disease was highest in the U.S. Arctic (10.2 per 100,000) compared to the lowest in N. Sweden (1.2 per 100,000). Seven deaths were associated with GAS, all occurred in the U.S. Arctic.

Group A *Streptococcus* Case Demographics, ICS 2011 Data

Country	Population	# Cases	Rate*	Sex	Median Age (min-max) yrs	Deaths n (CFR†)
				M (%)		
Greenland	56,615	1	1.8	1 (100)	11.1 (1 case)	0 (0)
N. Canada	154,162	12	7.8	8 (67)	39 (0-79)	0 (0) ^a
N. Sweden	248,545	3	1.2	1 (33)	39 (7-79)	‡
U.S. Arctic	723,424	74	10.2	43 (58)	49.3 (0.7-90)	7 (9.5)
Total	1,182,746	90	7.6	53 (59)	47.4 (0-90)	7 (8.1)

*Number of cases per 100,000 per year

†Case fatality ratio

‡Outcomes not reported from N. Sweden

^aOutcome unknown in 1 case from N. Canada

When stratified by age, the highest rates of disease occurred in children <2 years in N. Canada (35.2 per 100,000) and in individuals 65+ years of age in the U.S. Arctic (27.2 per 100,000).

Group A *Streptococcus* by Age Category, ICS 2011 Data

Age		Greenland	N. Canada	N. Sweden	U.S. Arctic
<2 yrs	Population	1,753	5,686	4,901	22,194
	Cases (%) [*]	0 (0)	2 (17)	0 (0)	3 (4)
	Rate†	0	35.2	0	13.5
2-19 yrs	Population	15,445	44,950	47,830	186,816
	Cases (%) [*]	1 (100)	3 (25)	1 (33)	6 (8)
	Rate†	6.5	6.7	2.1	3.2
20-64 yrs	Population	35,470	94,672	142,118	455,553
	Cases (%) [*]	0 (0)	6 (50)	1 (33)	49 (66)
	Rate†	0	6.3	0.7	10.8
65+ yrs	Population	3,947	8,854	53,696	58,861
	Cases (%) [*]	0 (0)	1 (8)	1 (33)	16 (22)
	Rate†	0	11.3	1.9	27.2
All ages	Population	56,615	154,162	248,545	723,424
	Cases	1	12	3	74
	Rate†	1.8	7.8	1.2	10.2

*Proportion of total cases in each country/region

†Number of cases per 100,000 per year

Race

Race and ethnicity data were collected by N. Canada and the U.S. Arctic. The highest rates of disease occurred in N. Canada Aboriginal children <2 years and Alaska Native people 65 years and older in the U.S. Arctic.

Group A *Streptococcus* by Race and Age Categories, ICS 2011 Data

Age (yrs)		N. Canada		U.S. Arctic	
		Aboriginal	Non-Aboriginal	AK Native	Non-AK Native
<2	Population	3,928	1,758	6,518	15,676
	Cases (rate†)	2 (50.9)	0 (0)	2 (30.7)	1 (6.4)
2-19	Population	35,444	9,506	49,769	137,047
	Cases (rate†)	3 (8.5)	0 (0)	5 (10.1)	1 (0.7)
20-64	Population	45,212	49,460	77,168	378,385
	Cases (rate†)	5 (11.1)	1 (2)	24 (31.1)	25 (6.6)
65+	Population	3,699	5,155	8,707	50,154
	Cases (rate†)	1 (27)	0 (0)	6 (68.9)	10 (19.9)
All Ages	Population	88,283	65,879	142,162	581,262
	Cases (rate†)	11 (12.5)	1 (1.5)	37 (26)	37 (6.4)

†Number of cases per 100,000 per year

Clinical Presentation

The primary clinical presentation was determined by a review of the discharge diagnoses in each patient's individual medical record associated with the invasive bacterial illness. In cases with multiple discharge diagnoses, the most serious diagnosis related to the infection was recorded as the primary clinical presentation. The most common clinical presentation for GAS cases in Greenland, N. Canada and the U.S. Arctic was bacteremia, 100%, 50% and 27%, respectively. N. Sweden did not report clinical presentations.

Clinical Presentation of Reported group A *Streptococcus* Cases, ICS 2011 Data

	Greenland n (%)	N. Canada n (%)	U.S. Arctic n (%)
Bacteremia	1 (100)	6 (50)	20 (27)
Cellulitis*	0 (0)	3 (25)	18 (24)
Pneumonia*	0 (0)	0 (0)	12 (16)
Endocarditis	0 (0)	0 (0)	6 (8)
Septic arthritis	0 (0)	1 (8)	5 (7)
Necrotizing fasciitis	0 (0)	1 (8)	4 (5)
Empyema	0 (0)	1 (8)	3 (4)
Meningitis	0 (0)	0 (0)	1 (1)
Appendicitis	0 (0)	0 (0)	1 (1)
Epiglottitis	0 (0)	0 (0)	1 (1)
Other	0 (0)	0 (0)	3 (4)
Total	1	12	74

*with bacteremia

Risk Factors

Cigarette smoking, alcohol abuse and diabetes were the most common risk factors associated with adult (≥ 18 years) GAS cases in N. Canada and the U.S. Arctic. Greenland and N. Sweden did not report risk factor data.

Group A *Streptococcus* Risk Factor/Medical Conditions in Adults*, ICS 2011 Data†

	N. Canada	U.S. Arctic
	n (%)	n (%)
Cigarette Smoking	1 (14)	23 (35)
Alcohol Abuse	1 (14)	18 (27)
Chronic Lung Disease and/or Asthma	0 (0)	6 (9)
Immunosuppressive Therapy	0 (0)	1 (2)
Diabetes	1 (14)	14 (21)
Injection Drug Use	0 (0)	2 (3)
Asplenia	0 (0)	0 (0)
Total Adult* Cases	7	66

* ≥ 18 years

†Multiple risk factors may be reported per case

Outcome

Seven deaths in cases with GAS were reported from the U.S. Arctic (CFR 9.5%); two occurred in the 2-19 years old age category, four in the 20-64 years old age category and one in persons ≥ 65 . No deaths were reported in Greenland or N. Canada. N. Sweden did not report case outcome data.

Conclusions

These data suggest higher rates in indigenous populations, particularly in young children and older adults. Increased awareness of risk may help target improved treatment responses.

Group B *Streptococcus*

Case Demographics

Greenland, N. Canada, N. Sweden and the U.S. Arctic each reported the occurrence of GBS during 2011. A total of 43 cases of invasive disease caused by GBS were reported to ICS. Greenland and N. Sweden reported no cases. The rate of disease was highest in the U.S. Arctic (5.5 per 100,000) compared to N. Canada (2 per 100,000). Four deaths were associated with GBS in 2011.

Group B *Streptococcus* Case Demographics, ICS 2011 Data

Country	Population	#		Sex	Median Age (min-max) yrs	Deaths n (CFR†)
		Cases	Rate*	M (%)		
Greenland	56,615	0	0	No cases	No cases	No cases
N. Canada	154,162	3	2	2 (67)	0 (0-80.3)	0 (0)‡
N. Sweden	248,545	0	0	No cases	No cases	No cases
U.S. Arctic	723,424	40	5.5	17 (43)	60.3 (0-93.8)	4 (10)
Total	1,182,746	43	3.6	19 (44)	59.2 (0-93.8)	4 (9.5)

*Number of cases per 100,000 per year

†Case fatality ratio

‡Outcome unknown in one case

When stratified by age, the highest rates of disease occurred in cases less than two years of age in N. Canada and in persons 65 and older in the U.S. Arctic.

Group B *Streptococcus* by Age Category, ICS 2011 Data

Age	N. Canada	U.S. Arctic
<2 yrs	Population	5,686
	Cases (%)*	2 (67)
	Rate†	35.2
2-19 yrs	Population	44,950
	Cases (%)*	0 (0)
	Rate†	0
20-64 yrs	Population	94,672
	Cases (%)*	0 (0)
	Rate†	0
65+ yrs	Population	8,854
	Cases (%)*	1 (33)
	Rate†	11.3
All ages	Population	154,162
	Cases	3
	Rate†	2

*Proportion of total cases in each country/region

†Number of cases per 100,000 per year

There was one case of early-onset disease (cases less than 7 days old) in the U.S. Arctic (0.1 cases per 1,000 births) and two cases in N. Canada (0.7 cases per 1,000 births).

Race

Race and ethnicity data were collected in N. Canada and the U.S. Arctic. The overall rates of disease caused by GBS were higher in AK Native and Aboriginal people compared to non-Native and non-Aboriginal people. The highest rates of disease occurred in Aboriginal children less than 2 years old and AK Native persons 65 years and older.

Group B *Streptococcus* by Race and Age Categories, ICS 2011 Data

Age (yrs)	N. Canada		U.S. Arctic		
	Aboriginal	Non-Aboriginal	Native	Non-Native	
<2	Population	3,928	1,758	6,518	15,676
	Cases (rate*)	2 (50.9)	0 (0)	0 (0)	2 (12.8)
2-19	Population	35,444	9,506	49,769	137,047
	Cases (rate*)	0 (0)	0 (0)	0 (0)	0 (0)
20-64	Population	45,212	49,460	77,168	378,385
	Cases (rate*)	0 (0)	0 (0)	7 (9.1)	14 (3.7)
65+	Population	3,699	5,155	8,707	50,154
	Cases (rate*)	0 (0)	0 (0)	4 (45.9)	13 (25.9)
All	Population	88,283	65,879	142,162	581,262
Ages	Cases (rate*)	2 (2.3)	0 (0)	11 (7.7)	29 (5)

*Number of cases per 100,000 per year

Clinical Presentation

The primary clinical presentation was determined by a review of the discharge diagnoses in each patient's individual medical record associated with the invasive bacterial illness. In cases with multiple discharge diagnoses, the most serious diagnosis related to the infection was recorded as the primary clinical presentation. In the U.S. Arctic, cellulitis (35%) was the most common clinical presentation reported for cases of GBS in 2011 followed by bacteremia (25%) and pneumonia (10%). One case in N. Canada presented with septic arthritis and two with other clinical presentations.

Clinical Presentation of Reported group B *Streptococcus* Cases, ICS 2011 Data

	N. Canada	U.S. Arctic
	n (%)	n (%)
Cellulitis*	0 (0)	14 (35)
Bacteremia	0 (0)	10 (25)
Pneumonia*	0 (0)	4 (10)
Peritonitis	0 (0)	3 (7.5)
Endocarditis	0 (0)	2 (5)
Meningitis	0 (0)	2 (5)
Osteomyelitis	0 (0)	2 (5)
Septic abortion	0 (0)	1 (2.5)
Bursitis	0 (0)	1 (2.5)
Septic arthritis	1 (33)	0 (0)
Other	2 (67)	1 (2.5)
Total	3	40

*with bacteremia

Risk Factors

Twenty-six percent of GBS adult (≥ 18 years) cases reviewed in the U.S. Arctic indicated diabetes as a risk factor in 2011; 24% had chronic lung disease, 16% were smokers and/or reported alcohol abuse. No risk factors were reported in N. Canada.

Outcome

Four deaths in cases with GBS were reported in the U.S. Arctic (CFR 10%); all occurred in the 20-65 years age category.

Conclusions

Guidelines for universal screening of pregnant women for GBS carriage were established in 2002 which have resulted in decreases in early onset disease. Cases continue to occur in older age groups which warrants continued surveillance.

CONCLUSIONS

Monitoring rates of disease and levels of antimicrobial resistance in *S. pneumoniae*, *H. influenzae*, *N. meningitidis*, GAS and GBS via use of the ICS system is important in providing data on groups at risk for disease, measurement of effectiveness of prevention measures, and emerging challenges in serotype distribution and antimicrobial resistance. Efforts to expand ICS to include all circumpolar nations will continue.

ACKNOWLEDGMENTS

ICS is a cooperative project funded by the Office of Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, and by the Public Health Agency of Canada in Ottawa, Canada.

We would like to thank all individuals involved in ICS at participating laboratories and public health departments.

SOURCE

This report was prepared by:

Tammy Zulz, MPH
Surveillance Coordinator

Michael Bruce, MD MPH
Epidemiology Team Leader

Arctic Investigations Program
OID/NCEZID/DPEI
Centers for Disease Control and Prevention
Phone: (907) 729-3400
Fax: (907) 729-3429

REFERENCES

- [1] Reasonover A, Zulz T, Bruce MG, Bruden D, Jetté L, Kaltoft M, Lambertsen L, Parkinson A, Rudolph K, Lovgren M. The International Circumpolar Surveillance interlaboratory quality control program for *Streptococcus pneumoniae*, 1999 to 2008. *J Clin Microbiol.* 2011 Jan;49(1):138-43

- [2] Tsang RS, Rudolph K, Lovgren M, Bekal S, Lefebvre B, Lambertsen L, Zulz T, Bruce M. International circumpolar surveillance interlaboratory quality control program for serotyping *Haemophilus influenzae* and serogrouping *Neisseria meningitidis*, 2005 to 2009. *J Clin Microbiol.* 2012 Mar;50(3):651-6

PARTICIPANTS

International Circumpolar Surveillance (ICS) Participants, 2011

FINLAND

Reference Laboratory	National Public Health Institute (KTL) Laboratory, Oulu
Laboratories	Et.-Pohjanmaan sh-piiri, Seinäjoen sairaalan mikrobiol. lab. Etelä-Karjalan keskussairaalan kl.mikrobiologian laboratorio HY – Serobakteriologian laitos Jorvin sairaala, kliinisen mikrobiologian laboratorio KYS – Mikrobiologian laboratorio Kainuun keskussairaalan mikrobiologian laboratorio Kanta-Hämeen keskussairaalan mikrobiologian laboratorio Keski-Pohjanmaan keskussairaalan mikrobiologian laboratorio Keski-Suomen keskussairaalan mikrobiologian laboratorio Kymenlaakson keskussairaalan mikrobiologian laboratorio Lapin keskussairaalan mikrobiologian laboratorio Länsi-Pohjan keskussairaalan laboratorio Mikkelin keskussairaalan mikrobiologian laboratorio OYKS – Mikrobiologian laboratorio Oulun kiakonissalairoksen laboratorio Pohjois-Karjalan keskussairaalan mikrobiologian laboratorio Päijät-Hämeen keskussairaalan mikrobiologian laboratorio Rauman aluesairaalan laboratorio Satakunnan keskussairaalan mikrobiologian laboratorio Savonlinnan keskussairaalan laboratorio TAYS – Mikrobiologian laboratorio TYKS – Mikrobiologian laboratorio Vaasan keskussairaalan mikrobiologian laboratorio

GREENLAND

Reference Laboratories	Statens Serum Institute, Copenhagen, Denmark Centralab at Queen Ingrid's Hospital, Nuuk, Greenland
Laboratories	Nanortalik Hospital Qaqortoq Hospital Narsaq Hospital Paamiut Hospital Maniitsoq Hospital Sisimut Hospital Aasiaat Hospital Qasigiannuguit Hospital Ilulissat Hospital Qeqertarsuaq Hospital Uummannaq Hospital Upernavik Hospital Qaanaaq Hospital Ammassalik Hospital Ittoqqortoormiit Hospital

ICELAND

Reference Laboratory	Department of Microbiology, Landspítali University Hospital, Reykjavik
Laboratories	Akranes Hospital Isafjordur District Hospital Stykkisholmur Local Health Center St. Joseph's Hospital Hafnarfjorour Municipal Hospital of Vestmannaeyjar Akureyri Egilstadir Health Center Selfoss Health Center Sudurnes Health Center (Keflavik) Regional Hospital Neskaupstadur

NORTHERN CANADA

Laboratory Centre for Disease Control	Respiratory Division, Bureau of Infectious Diseases, Laboratory Centre for Disease Control, Ottawa
Reference Laboratories	National Centre for Streptococcus, Provincial Laboratory of Public Health, Edmonton, AB Laboratoire de Santé Publique du Québec, Montréal, QC National Centre for Meningococcus, Provincial Laboratory of Public Health, Winnipeg, MB
Laboratories	Whitehorse General Hospital, Whitehorse, YK Stanton Regional Health Board, Yellowknife, NT H.H. Williams Memorial Hospital, Hay River, NT Inuvik Regional Hospital, Inuvik, NT Baffin Regional Hospital, Iqaluit, NU Churchill Regional Health Authority, Churchill, MB Cadham Provincial Laboratory, Winnipeg, MB Ungava Tulattavik Health Centre, Kuujjuaq, QC Inulitsavik Hospital, Puvirnituk, QC Cree Health Board, Chisasibi, QC CSSSR, Chibougamou, QC Val d'Or Hospital, Val d'Or, QC Melville Hospital, Goose Bay, NL Newfoundland Public Health Laboratory, St. John's, NL
Public Health	Yukon Communicable Disease Control, Whitehorse, YK Health Protection Unit, Government of NWT, Yellowknife, NT JA Hildes Northern Medical Unit, Winnipeg, MB Régie Régionale de la Santé et des Services Sociaux, Kuujjuaq, QC Région Cri de la Baie James, Module de Santé Publique, Montreal, QC Communicable Disease Control, Health Laborador Corporation, Goose Bay, NL IMPAct Coordinator, Vaccine Evaluation Centre, Vancouver, BC

NORTHERN SWEDEN

Reference Laboratory	Department of Bacteriology, Swedish Institute for Infectious Disease Control, Stockholm
Laboratories	Department of Microbiology, Sunderby Hospital, Lulea

NORWAY

Reference Laboratory	Oslo/Tromsø
Laboratories	<p>Frederikstad, Østf. SSH Sarpsborg SH Akershus SSH, SiA Børum SH Aker SH Fürsts laborat, Oslo Dr. Willes med.lab. Radiumhospitalet Folkehelsa, vir.lab. Folkehelsa, bakt.lab. Forsv.mik.lab.Folk.h. Rikshospitalet, mik.lab. Ullevål SH, mik.lab. Lab. klin. mikrob. Oslo Lillehammer mik.lab Elverum mik.lab. Buskerud SSH, mik.lab. Vestfold SSH, mik.lab. Telelab Vest-Agder SSH, mik.lab. Rogaland SSH, mik.lab. Haukeland SH, mik.lab. Sogn-Fk. SSH, mik.lab. Ålesund FSH, mik.lab. Molde FSH, mik.lab. Trondheim RSH, mik.lab. Innherred SH, mik.lab. Namdal SH, mik.lab. Nordland SSH, mik.lab. Tromsø RSH, mik.lab. Kirkenes SH, mik.lab. Laboratorium INA/div. Haugesund, mik.lab.</p>

U.S. ARCTIC

Reference Laboratory	Arctic Investigations Program, National Center for Infectious Diseases, Centers for Disease Control and Prevention, Anchorage, AK
Laboratories	<p>Alaska Native Medical Center, Anchorage, AK Alaska Regional Hospital, Anchorage, AK Bartlett Regional Hospital, Juneau, AK Bassett Army Hospital, Fort Wainwright, AK Central Peninsula General Hospital, Soldotna, AK Cordova Community Medical Center, Cordova, AK Elmendorf Air Force Base Hospital, Anchorage, AK Fairbanks Memorial Hospital, Fairbanks, AK Kakanak Hospital, Dillingham, AK Ketchikan Regional Hospital, Ketchikan, AK Manilaq Medical Center, Kotzebue, AK Norton Sound Regional Hospital, Nome, AK Petersburg Medical Center, Petersburg, AK Providence Alaska Medical Center, Anchorage, AK Providence Island Medical Center, Kodiak, AK Samuel Simmonds Memorial Hospital, Barrow, AK Sitka Community Hospital, Sitka, AK South Peninsula Hospital, Homer, AK Southeast Area Regional Health Corporation, Sitka, AK State Public Health Laboratory, Division of Public Health, Department of Health and Social Services, Anchorage, AK Valdez Community Hospital, Valdez, AK Valley Hospital, Palmer, AK Wrangell General Hospital, Wrangell, AK Yukon-Kuskokwim Delta Regional Hospital, Bethel, AK</p>