

Acknowledgments

We thank Fernando Gil and Aldo López for the authorization for the development of the research in PVWR and VWRA, respectively. Thanks are due to the biologist Christian Carazas and the engineer José Junco for help in some sampling of snails; to Rosa Martínez for the use of the laboratory at Universidad Nacional Mayor de San Marcos, Peru; and to Airton Lobo for technical support. We are grateful to Sara Vanessa Brant for constructive suggestions in the earlier version of this work.

This project was supported financially by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) (process no. 23038.005297/2011-39) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil (scholarship to E.A.P.M. and L.F.V.F.).

About the Author

Mr. Pulido-Murillo is a PhD student from Peru working in the Universidade Federal de Minas Gerais, Brazil. His research interests focus on morphological and molecular study of zoonotic helminths.

References

- Chai JY, Darwin Murrell K, Lymbery AJ. Fish-borne parasitic zoonoses: status and issues. *Int J Parasitol.* 2005;35:1233–54. <http://dx.doi.org/10.1016/j.ijpara.2005.07.013>
- Pearson J. Family Heterophyidae Leiper, 1909. In: Bray RA, Gibson DI, Jones A, editors. *Keys to the Trematoda*. Vol. III. London: CAB International and Natural History Museum; 2008, p. 113–142.
- Keiser J, Utzinger J. Emerging foodborne trematodiasis. *Emerg Infect Dis.* 2005;11:1507–14. <http://dx.doi.org/10.3201/eid1110.050614>
- Chai JY, Jung BK. Fishborne zoonotic heterophyid infections: an update. *Food Water Parasitol.* 2017 [Epub ahead of print]. <https://doi.org/10.1016/j.fawpar.2017.09.001>
- Arya LK, Rathinam SR, Lalitha P, Kim UR, Ghatani S, Tandon V. Trematode fluke *Procerovum varium* as cause of ocular inflammation in children, South India. *Emerg Infect Dis.* 2016;22:192–200 <https://doi.org/10.3201/eid2202.150051>. <http://dx.doi.org/10.3201/eid2202.150051>
- Pinto HA, de Melo AL. A checklist of trematodes (Platyhelminthes) transmitted by *Melanoides tuberculata* (Mollusca: Thiaridae). *Zootaxa.* 2011;2799:15–28.
- Pointier JP, David P, Jarne P. The biological control of the snail hosts of schistosomes: the role of competitor snails and biological invasions. In: Toledo R, Fried B, editors. *Biomphalaria* snails and larval trematodes. New York: Springer; 2011. p. 215–38.
- Simone LRL. Land and freshwater mollusks of Brazil. São Paulo: Editora Gráfica Bernardi & Fundação de Amparo à Pesquisa do Estado de São Paulo; 2006.
- Scholz T, Aguirre-Macedo ML, Salgado-Maldonado G. Trematodes of the family Heterophyidae (Digenea) in Mexico: a review of species and new host and geographical records. *J Nat Hist.* 2001;35:1733–72. <http://dx.doi.org/10.1080/00222930152667087>
- Tkach VV, Littlewood DTJ, Olson PD, Kinsella JM, Swiderski Z. Molecular phylogenetic analysis of the Microphalloidea Ward, 1901 (Trematoda: Digenea). *Syst Parasitol.* 2003;56:1–15. <http://dx.doi.org/10.1023/A:1025546001611>

Address for correspondence: Hudson A. Pinto, Laboratório de Biologia de Trematoda, Department of Parasitology, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, C.P. 486, 30123-970, Belo Horizonte, Minas Gerais, Brazil; email: hudsonalves13@icb.ufmg.br

LETTER

Ceftriaxone-Resistant *Neisseria gonorrhoeae*, Canada, 2017

Alan R. Katz

Author affiliation: University of Hawaii, Honolulu, Hawaii, USA

DOI: <http://dx.doi.org/10.3201/eid2403.171892>

To the Editor: I read with great interest the report by Lefebvre et al. about a *Neisseria gonorrhoeae* isolate identified in Canada demonstrating a ceftriaxone MIC of 1 mg/L (1). The authors note: “As of October 15, 2017, only 5 ceftriaxone-resistant *Neisseria gonorrhoeae* isolates had been reported worldwide (MIC range 0.5–2 mg/L).” The authors cite published reports from Spain, Japan, Australia, and France.

I would like to clarify that additional *N. gonorrhoeae* isolates have been identified with ceftriaxone MICs ≥ 0.5 mg/L. Since 1987, as part of the Gonococcal Isolate Surveillance Project, the Centers for Disease Control and Prevention has been testing *N. gonorrhoeae* isolates for ceftriaxone susceptibility. During 1987–2016, the Centers for Disease

Control and Prevention identified and reported 5 isolates with ceftriaxone MICs of 0.5 mg/L in the United States. These isolates were found in San Diego, California (1987); Cincinnati, Ohio (1992 and 1993); Philadelphia, Pennsylvania (1997); and most recently, Oklahoma City, Oklahoma (2012) (2). Therefore, although the number of *N. gonorrhoeae* isolates with ceftriaxone MICs ≥ 0.5 mg/L identified globally to date has been small, these Gonococcal Isolate Surveillance Project findings should be acknowledged. Continued and enhanced global surveillance of gonococcal isolates for antimicrobial susceptibility testing is imperative.

References

- Lefebvre B, Martin I, Demczuk W, Deshaies L, Michaud S, Labbe AC, et al. Ceftriaxone-resistant *Neisseria gonorrhoeae*, Canada, 2017 [cited 2017 Nov 19]. 2018;24:381–3. <http://doi.org/10.3201/eid2402.171756>
- Centers for Disease Control and Prevention. Sexually transmitted diseases surveillance 2016 [cited 2017 Nov 19]. https://www.cdc.gov/std/stats16/CDC_2016_STDS_Report-for-508WebSep21_2017_1644.pdf

Address for correspondence: Alan R. Katz, University of Hawaii at Manoa, Office of Public Health Studies, Biomedical Sciences Bldg, Rm D104M, 1960 East-West Rd, Honolulu, HI 96822, USA; email: katz@hawaii.edu