

SUPPLEMENTAL MATERIAL

Table S1. Age-standardized prevalence of coronary heart disease and stroke by sex and birthplace, comparison of foreign-born with US-born by race/ethnicity, † National Health Interview Survey, 2006-2014.

	US-born-NH white	US-born NH-Black	US-born Hispanic	US-born NH-Asia	Mexico, Central America, Caribbean	South America	Europe	Africa	Indian Subcontinent	Asia
					% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)
Coronary heart disease										
Total	6.3(0.1)	6.6(0.1)	5.7(0.2)	4.7(0.4)	5.0(0.2)*	4.1(0.4)*	5.4(0.3)**	3.1(0.7)***	5.1(0.7)	3.4(0.2)**
Men	8.4(0.1)	6.9(0.2)	6.9(0.4)	5.9(0.8)	5.3(0.3)**	5.6(0.7)	6.7(0.5)**	4.4(1.2)*	7.0(1.0)	3.9(0.3)*
Women	4.6(0.1)	6.3(0.2)	4.6(0.3)	3.7(0.6)	4.8(0.2)	3.0(0.5)**	4.3(0.4)	1.6(0.6)***	2.7(0.8)	3.1(0.3)
Stroke										
Total	2.4(0)	4.2(0.1)	2.8(0.2)	1.9(0.3)	2.3(0.1)*	1.1(0.2)***	1.8(0.2)***	1.9(0.6)***	1.7(0.4)	1.9(0.2)
Men	2.5(0.1)	4(0.2)	2.7(0.3)	2.3(0.5)	2.3(0.2)	0.8(0.3)***	2.3(0.3)	0.8(0.3)***	2.0(0.5)	2.1(0.3)
Women	2.4(0.1)	4.3(0.2)	2.9(0.2)	1.6(0.3)	2.4(0.2)	1.4(0.3)***	1.4(0.2)***	2.9(1.0)	1.2(0.5)	1.7(0.2)

p-values (*<0.05, **<0.01, ***<0.001) .

† 1) for Mexico, Central America and Caribbean-born and South American-born, comparison was US-born Hispanics; 2) for Europe-born, comparison was US-born Non-Hispanic white; 3) for African-born, the comparison was US-born Non-Hispanic black; 4) for Asian-born and Indian Subcontinent-born, the comparison was US-born Asian.

Table S2. Odds ratios for coronary heart disease and stroke risk among foreign-born adults, National Health Interview Survey, 2006-2014.

		Compared with US-born race/ethnic group [†]	Model 1 [‡]	Model 2 [§]	Model 3 [¶]
CHD	Men	Mexico, Central America, Caribbean	0.88(0.74-1.04)	0.85(0.69-1.04)	0.89(0.72-1.11)
		South America	1.12(0.83-1.51)	1.29(0.90-1.85)	1.31(0.88-1.94)
		Europe	0.79(0.66-0.94)	0.81(0.67-0.97)	0.87(0.70-1.06)
		Africa	0.43(0.24-0.79)	0.73(0.37-1.43)	0.67(0.33-1.34)
		Indian Subcontinent	1.06(0.66-1.69)	1.60(0.99-2.58)	1.89(1.12-3.20)
		Asia	0.87(0.60-1.27)	0.74(0.50-1.11)	0.80(0.53-1.22)
CHD	Women	Mexico, Central America, Caribbean	1.38(1.16-1.65)	1.16(0.95-1.40)	1.13(0.92-1.37)
		South America	0.82(0.58-1.14)	0.83(0.57-1.22)	0.94(0.63-1.40)
		Europe	0.98(0.79-1.21)	1.00(0.79-1.26)	0.97(0.76-1.23)
		Africa	0.15(0.07-0.29)	0.30(0.13-0.68)	0.28(0.08-0.90)
		Indian Subcontinent	0.57(0.31-1.04)	0.61(0.26-1.42)	0.99(0.47-2.11)
		Asia	1.02(0.70-1.51)	0.80(0.50-1.26)	0.76(0.48-1.20)
Stroke	Men	Mexico, Central America, Caribbean	0.91(0.69-1.21)	0.86(0.61-1.19)	0.91(0.65-1.28)
		South America	0.39(0.20-0.77)	0.46(0.22-1.00)	0.53(0.24-1.19)
		Europe	0.92(0.68-1.24)	0.98(0.71-1.34)	0.98(0.71-1.36)
		Africa	0.20(0.08-0.51)	0.17(0.09-0.35)	0.29(0.14-0.59)
		Indian Subcontinent	0.74(0.37-1.47)	0.94(0.46-1.96)	0.97(0.44-2.12)
		Asia	1.28(0.76-2.14)	0.94(0.52-1.68)	0.87(0.48-1.60)
Stroke	Women	Mexico, Central America, Caribbean	1.05(0.84-1.30)	0.93(0.72-1.19)	0.97(0.75-1.26)
		South America	0.66(0.41-1.08)	0.72(0.41-1.25)	0.71(0.39-1.30)
		Europe	0.60(0.47-0.77)	0.59(0.45-0.77)	0.64(0.50-0.83)
		Africa	0.32(0.16-0.64)	0.70(0.33-1.49)	0.72(0.29-1.76)
		Indian Subcontinent	0.41(0.14-1.18)	0.53(0.16-1.76)	0.76(0.24-2.40)
		Asia	1.34(0.82-2.18)	1.15(0.67-1.98)	1.22(0.70-2.13)

†. Used US-born Hispanic as reference when the models included respondents from Mexico, Central America and Caribbean-born, and South America-born; Used US-born Non-Hispanic-white as reference with the model for Europe-born; Used US-born Non-Hispanic-black as reference with the model for Africa-born; Used US-born-Asian as reference with the models for Indian Subcontinent-born and Asian-born. ‡. Model 1 unadjusted. §. Model 2 adjusted for age. ¶. Model 3 adjusted for age, birthplace, sex, health insurance, education, body mass index, smoking status, hypertension, and diabetes.