

**DRAFT  
TOXICOLOGICAL PROFILE FOR  
CARBON MONOXIDE**

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Public Health Service  
Agency for Toxic Substances and Disease Registry

September 2009

## DISCLAIMER

The use of company or product name(s) is for identification only and does not imply endorsement by the Agency for Toxic Substances and Disease Registry.

This information is distributed solely for the purpose of pre dissemination public comment under applicable information quality guidelines. It has not been formally disseminated by the Agency for Toxic Substances and Disease Registry. It does not represent and should not be construed to represent any agency determination or policy.

## UPDATE STATEMENT

Toxicological profiles are revised and republished as necessary. For information regarding the update status of previously released profiles, contact ATSDR at:

Agency for Toxic Substances and Disease Registry  
Division of Toxicology and Environmental Medicine/Applied Toxicology Branch  
1600 Clifton Road NE  
Mailstop F-62  
Atlanta, Georgia 30333

This page is intentionally blank.

## FOREWORD

This toxicological profile is prepared in accordance with guidelines developed by the Agency for Toxic Substances and Disease Registry (ATSDR) and the Environmental Protection Agency (EPA). The original guidelines were published in the *Federal Register* on April 17, 1987. Each profile will be revised and republished as necessary.

The ATSDR toxicological profile succinctly characterizes the toxicologic and adverse health effects information for these toxic substances described therein. Each peer-reviewed profile identifies and reviews the key literature that describes a substance's toxicologic properties. Other pertinent literature is also presented, but is described in less detail than the key studies. The profile is not intended to be an exhaustive document; more comprehensive sources of specialty information are referenced.

The focus of the profiles is on health and toxicologic information. Each toxicological profile begins with a public health statement that describes, in nontechnical language, a substance's relevant toxicological properties. Following the public health statement is information concerning levels of significant human exposure and, where known, significant health effects. The adequacy of information to determine a substance's health effects is described in a health effects summary. Data needs that are of significance to protection of public health are identified by ATSDR and EPA.

Each profile includes the following:

- (A) The examination, summary, and interpretation of available toxicologic information and epidemiologic evaluations on a toxic substance to ascertain the levels of significant human exposure for the substance and the associated acute, subacute, and chronic health effects;
- (B) A determination of whether adequate information on the health effects of each substance is available or in the process of development to determine levels of exposure that present a significant risk to human health of acute, subacute, and chronic health effects; and
- (C) Where appropriate, identification of toxicologic testing needed to identify the types or levels of exposure that may present significant risk of adverse health effects in humans.

The principal audiences for the toxicological profiles are health professionals at the Federal, State, and local levels; interested private sector organizations and groups; and members of the public. We plan to revise these documents in response to public comments and as additional data become available. Therefore, we encourage comments that will make the toxicological profile series of the greatest use.

Comments should be sent to:


Agency for Toxic Substances and Disease Registry  
Division of Toxicology and Environmental Medicine  
1600 Clifton Road, N.E.  
Mail Stop F-62  
Atlanta, Georgia 30333

The toxicological profiles are developed under the Comprehensive Environmental Response, Compensation, and Liability Act of 1980, as amended (CERCLA or Superfund). CERCLA section 104(i)(1) directs the Administrator of ATSDR to "...effectuate and implement the health related authorities" of the statute. This includes the preparation of toxicological profiles for hazardous substances most commonly found at facilities on the CERCLA National Priorities List and that pose the most significant potential threat to human health, as determined by ATSDR and the EPA. Section 104(i)(3) of CERCLA, as amended, directs the Administrator of ATSDR to prepare a toxicological profile for each substance on the list. In addition, ATSDR has the authority to prepare toxicological profiles for substances not found at sites on the National Priorities List, in an effort to "...establish and maintain inventory of literature, research, and studies on the health effects of toxic substances" under CERCLA Section 104(i)(1)(B), to respond to requests for consultation under section 104(i)(4), and as otherwise necessary to support the site-specific response actions conducted by ATSDR.

This profile reflects ATSDR's assessment of all relevant toxicologic testing and information that has been peer-reviewed. Staffs of the Centers for Disease Control and Prevention and other Federal scientists have also reviewed the profile. In addition, this profile has been peer-reviewed by a nongovernmental panel and is being made available for public review. Final responsibility for the contents and views expressed in this toxicological profile resides with ATSDR.



Howard Frumkin, M.D., Dr. P.H.  
Director, National Center for Environmental  
Health/Agency for Toxic Substances and Disease  
Registry



Thomas R. Frieden, M.D., M.P.H.  
Administrator, Agency for Toxic Substances and  
Disease Registry

## QUICK REFERENCE FOR HEALTH CARE PROVIDERS

Toxicological Profiles are a unique compilation of toxicological information on a given hazardous substance. Each profile reflects a comprehensive and extensive evaluation, summary, and interpretation of available toxicologic and epidemiologic information on a substance. Health care providers treating patients potentially exposed to hazardous substances will find the following information helpful for fast answers to often-asked questions.

---

### *Primary Chapters/Sections of Interest*

**Chapter 1: Public Health Statement:** The Public Health Statement can be a useful tool for educating patients about possible exposure to a hazardous substance. It explains a substance's relevant toxicologic properties in a nontechnical, question-and-answer format, and it includes a review of the general health effects observed following exposure.

**Chapter 2: Relevance to Public Health:** The Relevance to Public Health Section evaluates, interprets, and assesses the significance of toxicity data to human health.

**Chapter 3: Health Effects:** Specific health effects of a given hazardous compound are reported by type of health effect (death, systemic, immunologic, reproductive), by route of exposure, and by length of exposure (acute, intermediate, and chronic). In addition, both human and animal studies are reported in this section.

**NOTE:** Not all health effects reported in this section are necessarily observed in the clinical setting. Please refer to the Public Health Statement to identify general health effects observed following exposure.

**Pediatrics:** Four new sections have been added to each Toxicological Profile to address child health issues:

<b>Section 1.6</b>	<b>How Can (Chemical X) Affect Children?</b>
<b>Section 1.7</b>	<b>How Can Families Reduce the Risk of Exposure to (Chemical X)?</b>
<b>Section 3.7</b>	<b>Children's Susceptibility</b>
<b>Section 6.6</b>	<b>Exposures of Children</b>

### **Other Sections of Interest:**

<b>Section 3.8</b>	<b>Biomarkers of Exposure and Effect</b>
<b>Section 3.11</b>	<b>Methods for Reducing Toxic Effects</b>

---

### *ATSDR Information Center*

**Phone:** 1-800-CDC-INFO (800-232-4636) or 1-888-232-6348 (TTY) **Fax:** (770) 488-4178  
**E-mail:** [cdcinfo@cdc.gov](mailto:cdcinfo@cdc.gov) **Internet:** <http://www.atsdr.cdc.gov>

The following additional material can be ordered through the ATSDR Information Center:

*Case Studies in Environmental Medicine: Taking an Exposure History*—The importance of taking an exposure history and how to conduct one are described, and an example of a thorough exposure history is provided. Other case studies of interest include *Reproductive and Developmental Hazards*; *Skin Lesions and Environmental Exposures*; *Cholinesterase-Inhibiting Pesticide Toxicity*; and numerous chemical-specific case studies.

*Managing Hazardous Materials Incidents* is a three-volume set of recommendations for on-scene (prehospital) and hospital medical management of patients exposed during a hazardous materials incident. Volumes I and II are planning guides to assist first responders and hospital emergency department personnel in planning for incidents that involve hazardous materials. Volume III—*Medical Management Guidelines for Acute Chemical Exposures*—is a guide for health care professionals treating patients exposed to hazardous materials.

*Fact Sheets (ToxFAQs)* provide answers to frequently asked questions about toxic substances.

---

### ***Other Agencies and Organizations***

*The National Center for Environmental Health (NCEH)* focuses on preventing or controlling disease, injury, and disability related to the interactions between people and their environment outside the workplace. Contact: NCEH, Mailstop F-29, 4770 Buford Highway, NE, Atlanta, GA 30341-3724 • Phone: 770-488-7000 • FAX: 770-488-7015.

*The National Institute for Occupational Safety and Health (NIOSH)* conducts research on occupational diseases and injuries, responds to requests for assistance by investigating problems of health and safety in the workplace, recommends standards to the Occupational Safety and Health Administration (OSHA) and the Mine Safety and Health Administration (MSHA), and trains professionals in occupational safety and health. Contact: NIOSH, 200 Independence Avenue, SW, Washington, DC 20201 • Phone: 800-356-4674 or NIOSH Technical Information Branch, Robert A. Taft Laboratory, Mailstop C-19, 4676 Columbia Parkway, Cincinnati, OH 45226-1998 • Phone: 800-35-NIOSH.

*The National Institute of Environmental Health Sciences (NIEHS)* is the principal federal agency for biomedical research on the effects of chemical, physical, and biologic environmental agents on human health and well-being. Contact: NIEHS, PO Box 12233, 104 T.W. Alexander Drive, Research Triangle Park, NC 27709 • Phone: 919-541-3212.

---

### ***Referrals***

*The Association of Occupational and Environmental Clinics (AOEC)* has developed a network of clinics in the United States to provide expertise in occupational and environmental issues. Contact: AOEC, 1010 Vermont Avenue, NW, #513, Washington, DC 20005 • Phone: 202-347-4976 • FAX: 202-347-4950 • e-mail: [AOEC@AOEC.ORG](mailto:AOEC@AOEC.ORG) • Web Page: <http://www.aoec.org/>.

*The American College of Occupational and Environmental Medicine (ACOEM)* is an association of physicians and other health care providers specializing in the field of occupational and environmental medicine. Contact: ACOEM, 25 Northwest Point Boulevard, Suite 700, Elk Grove Village, IL 60007-1030 • Phone: 847-818-1800 • FAX: 847-818-9266.

## CONTRIBUTORS

### CHEMICAL MANAGER(S)/AUTHOR(S):

Sharon Wilbur, M.A.  
Malcolm Williams, DVM, Ph.D.  
Robert Williams, Ph.D.  
Franco Scinicariello, M.D., M.P.H.  
ATSDR, Division of Toxicology and Environmental Medicine, Atlanta, GA

Julie M. Klotzbach, Ph.D.  
Gary L. Diamond, Ph.D.  
Mario Citra, Ph.D.  
SRC, Inc., North Syracuse, NY

### THE PROFILE HAS UNDERGONE THE FOLLOWING ATSDR INTERNAL REVIEWS:

1. Health Effects Review. The Health Effects Review Committee examines the health effects chapter of each profile for consistency and accuracy in interpreting health effects and classifying end points.
2. Minimal Risk Level Review. The Minimal Risk Level Workgroup considers issues relevant to substance-specific Minimal Risk Levels (MRLs), reviews the health effects database of each profile, and makes recommendations for derivation of MRLs.
3. Data Needs Review. The Applied Toxicology Branch reviews data needs sections to assure consistency across profiles and adherence to instructions in the Guidance.
4. Green Border Review. Green Border review assures the consistency with ATSDR policy.

This page is intentionally blank.

## PEER REVIEW

A peer review panel was assembled for carbon monoxide. The panel consisted of the following members:

1. Laurence Fechter, Ph.D., Senior Career Research Scientist, Jerry Pettis Memorial Veterans Medical Center, Loma Linda, California
2. Jerrold Leikin, M.D., Director of Medical Toxicology, NorthShore University Health System – OMEGA, Glenbrook Hospital, Glenview, Illinois
3. Stephen Thom, M.D., Ph.D., Professor, Emergency Medicine, University of Pennsylvania, Institute for Environmental Medicine, Philadelphia, Pennsylvania

These experts collectively have knowledge of carbon monoxide's physical and chemical properties, toxicokinetics, key health end points, mechanisms of action, human and animal exposure, and quantification of risk to humans. All reviewers were selected in conformity with the conditions for peer review specified in Section 104(I)(13) of the Comprehensive Environmental Response, Compensation, and Liability Act, as amended.

Scientists from the Agency for Toxic Substances and Disease Registry (ATSDR) have reviewed the peer reviewers' comments and determined which comments will be included in the profile. A listing of the peer reviewers' comments not incorporated in the profile, with a brief explanation of the rationale for their exclusion, exists as part of the administrative record for this compound.

The citation of the peer review panel should not be understood to imply its approval of the profile's final content. The responsibility for the content of this profile lies with the ATSDR.

This page is intentionally blank.

## CONTENTS

DISCLAIMER .....	ii
UPDATE STATEMENT .....	iii
FOREWORD .....	v
QUICK REFERENCE FOR HEALTH CARE PROVIDERS .....	vii
CONTRIBUTORS .....	ix
PEER REVIEW .....	xi
CONTENTS .....	xiii
LIST OF FIGURES .....	xvii
LIST OF TABLES .....	xix
1. PUBLIC HEALTH STATEMENT .....	1
1.1 WHAT IS CARBON MONOXIDE? .....	1
1.2 WHAT HAPPENS TO CARBON MONOXIDE WHEN IT ENTERS THE ENVIRONMENT? .....	2
1.3 HOW MIGHT I BE EXPOSED TO CARBON MONOXIDE? .....	3
1.4 HOW CAN CARBON MONOXIDE ENTER AND LEAVE MY BODY? .....	3
1.5 HOW CAN CARBON MONOXIDE AFFECT MY HEALTH? .....	3
1.6 HOW CAN CARBON MONOXIDE AFFECT CHILDREN? .....	4
1.7 HOW CAN FAMILIES REDUCE THE RISK OF EXPOSURE TO CARBON MONOXIDE? .....	4
1.8 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO CARBON MONOXIDE? .....	5
1.9 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH? .....	5
1.10 WHERE CAN I GET MORE INFORMATION? .....	6
2. RELEVANCE TO PUBLIC HEALTH .....	7
2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO CARBON MONOXIDE IN THE UNITED STATES .....	7
2.2 SUMMARY OF HEALTH EFFECTS .....	8
2.3 CARBON MONOXIDE DOSE-RESPONSE RELATIONSHIPS .....	17
2.4 MINIMAL RISK LEVELS (MRLs) .....	20
3. HEALTH EFFECTS .....	21
3.1 INTRODUCTION .....	21
3.2 DISCUSSION OF HEALTH EFFECTS .....	26
3.2.1 Death .....	30
3.2.2 Systemic Effects .....	33
3.2.3 Immunological and Lymphoreticular Effects .....	89
3.2.4 Neurological Effects .....	89
3.2.5 Reproductive Effects .....	95
3.2.6 Developmental Effects .....	96
3.2.7 Cancer .....	127
3.3 GENOTOXICITY .....	128
3.4 TOXICOKINETICS .....	128
3.4.1 Absorption .....	130
3.4.2 Distribution .....	133
3.4.3 Metabolism .....	136
3.4.4 Elimination and Excretion .....	142

3.4.5	Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models .....	144
3.5	MECHANISMS OF ACTION .....	162
3.5.1	Pharmacokinetic Mechanisms .....	162
3.5.2	Mechanisms of Toxicity .....	165
3.5.3	Animal-to-Human Extrapolations .....	171
3.6	TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS .....	171
3.7	CHILDREN'S SUSCEPTIBILITY .....	172
3.8	BIOMARKERS OF EXPOSURE AND EFFECT .....	175
3.8.1	Biomarkers Used to Identify or Quantify Exposure to Carbon Monoxide .....	176
3.8.2	Biomarkers Used to Characterize Effects Caused by Carbon Monoxide .....	176
3.9	INTERACTIONS WITH OTHER CHEMICALS .....	177
3.10	POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE .....	179
3.11	METHODS FOR REDUCING TOXIC EFFECTS .....	181
3.11.1	Reducing Peak Absorption Following Exposure .....	181
3.11.2	Reducing Body Burden .....	182
3.11.3	Interfering with the Mechanism of Action for Toxic Effects .....	183
3.12	ADEQUACY OF THE DATABASE .....	183
3.12.1	Existing Information on Health Effects of Carbon Monoxide .....	183
3.12.2	Identification of Data Needs .....	185
3.12.3	Ongoing Studies .....	191
4.	CHEMICAL AND PHYSICAL INFORMATION .....	195
4.1	CHEMICAL IDENTITY .....	195
4.2	PHYSICAL AND CHEMICAL PROPERTIES .....	195
5.	PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL .....	199
5.1	PRODUCTION .....	199
5.2	IMPORT/EXPORT .....	199
5.3	USE .....	199
5.4	DISPOSAL .....	200
6.	POTENTIAL FOR HUMAN EXPOSURE .....	201
6.1	OVERVIEW .....	201
6.2	RELEASES TO THE ENVIRONMENT .....	204
6.2.1	Air .....	204
6.2.2	Water .....	206
6.2.3	Soil .....	207
6.3	ENVIRONMENTAL FATE .....	207
6.3.1	Transport and Partitioning .....	207
6.3.2	Transformation and Degradation .....	208
6.3.2.1	Air .....	208
6.3.2.2	Water .....	208
6.3.2.3	Sediment and Soil .....	209
6.4	LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT .....	209
6.4.1	Air .....	209
6.4.2	Water .....	222
6.4.3	Sediment and Soil .....	223
6.4.4	Other Environmental Media .....	223
6.5	GENERAL POPULATION AND OCCUPATIONAL EXPOSURE .....	224
6.6	EXPOSURES OF CHILDREN .....	226
6.7	POPULATIONS WITH POTENTIALLY HIGH EXPOSURES .....	227

6.8	ADEQUACY OF THE DATABASE .....	229
6.8.1	Identification of Data Needs .....	229
6.8.2	Ongoing Studies .....	231
7.	ANALYTICAL METHODS.....	233
7.1	BIOLOGICAL MATERIALS.....	233
7.2	ENVIRONMENTAL SAMPLES .....	234
7.3	ADEQUACY OF THE DATABASE .....	240
7.3.1	Identification of Data Needs .....	240
7.3.2	Ongoing Studies .....	241
8.	REGULATIONS, ADVISORIES, AND GUIDELINES.....	243
9.	REFERENCES .....	247
10.	GLOSSARY .....	305
APPENDICES		
A.	ATSDR MINIMAL RISK LEVELS AND WORKSHEETS .....	A-1
B.	USER’S GUIDE.....	B-1
C.	ACRONYMS, ABBREVIATIONS, AND SYMBOLS.....	C-1
D.	INDEX .....	D-1

This page is intentionally blank.

## LIST OF FIGURES

2-1. Blood Carboxyhemoglobin Levels Corresponding to Adverse Health Effects of Carbon Monoxide in Humans .....	18
3-1. HH Human and Carbon Monoxide Rat Curves .....	93
3-2. HH Human and Carbon Monoxide Human .....	94
3-3. Summary of Carbon Monoxide Uptake, Distribution, and Elimination Pathways.....	131
3-4. Temporal Profile of Blood Carboxyhemoglobin (%) for Hypothetical Continuous Exposures to Air Carbon Monoxide at Concentrations of 25, 200, or 1,000 ppm .....	132
3-5. Oxyhemoglobin Dissociation Curve for Normal Human Blood Containing 50% Carboxyhemoglobin, and for Blood of Anemic Human .....	141
3-6. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance .....	146
3-7. Plot of Standardized Sensitivity Coefficients for Carboxyhemoglobin (COHb) Levels Predicted from the Coburn-Forster-Kane (CFK) Model for a 100 ppm Exposure to an Individual Engaged in Vigorous Exercise (Alveolar Ventilation Rate=30 L/Minute).....	150
3-8. Structure of the Smith et al. (1994) Model .....	152
3-9. Structure of the Hill et al. (1977) Maternal-Fetal Model.....	155
3-10. Structure of the Bruce et al. (2008) Model .....	159
3-11. Potential Signaling Pathways Modulated by Carbon Monoxide .....	169
3-12. Existing Information on Health Effects of Carbon Monoxide .....	184
6-1. Frequency of NPL Sites with Carbon Monoxide Contamination.....	202

This page is intentionally blank.

## LIST OF TABLES

2-1. Blood Carboxyhemoglobin Levels Corresponding to Adverse Health Effects of Carbon Monoxide .....	19
3-1. Carbon Monoxide Exposures and Carboxyhemoglobin Levels Associated with Health Effects from Selected Studies Representing the Lowest Adverse Effect Levels .....	22
3-2. Predicted Steady-State Blood Carboxyhemoglobin (COHb) Levels .....	25
3-3. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Respiratory Disease .....	37
3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease .....	51
3-5. Effects of Acute-Duration Carbon Monoxide Exposure on Cardiovascular Function in Healthy Subjects .....	60
3-6. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Cardiovascular Disease .....	67
3-7. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Hospital Admissions and Emergency Department Visits Related to Cardiovascular Disease .....	74
3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes .....	98
3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals .....	114
3-10. Genotoxicity of Carbon Monoxide .....	129
3-11. Post-Mortem Tissue Distribution of Carbon Monoxide in Humans .....	135
3-12. Hemoglobin and Myoglobin Binding Kinetics and Equilibrium Constants for Oxygen and Carbon Monoxide .....	139
3-13. Parameter Values for Coburn-Forster-Kane (CFK) Model .....	149
3-14. Parameter Values for the Smith et al. (1994) Model .....	153
3-15. Parameter Values for the Hill et al. (1977) Model .....	156
3-16. Parameter Values for the Bruce et al. (2008) Model .....	160
3-17. Heme Proteins Modulated by Carbon Monoxide .....	167
3-18. Ongoing Studies on the Health Effects of Carbon Monoxide .....	192
4-1. Chemical Identity of Carbon Monoxide .....	196

4-2. Physical and Chemical Properties of Carbon Monoxide .....	197
6-1. Carbon Monoxide Emissions (Thousands of Short Tons) from 1970 to 2008 .....	205
6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States .....	211
6-3. Average Annual Estimated Non Fatal Carbon Monoxide Exposure Cases in the United States Emergency Room Visits (2004–2006) .....	225
6-4. Carboxyhemoglobin Levels in the U.S. Population Based Upon Smoking Status.....	228
7-1. Analytical Methods for Determining Carbon Monoxide in Biological Samples .....	235
7-2. Analytical Methods for Determining Carbon Monoxide in Environmental Samples .....	236
8-1. Regulations, Advisories, and Guidelines Applicable to Carbon Monoxide .....	244

## 1. PUBLIC HEALTH STATEMENT

This public health statement tells you about carbon monoxide and the effects of exposure to it.

The Environmental Protection Agency (EPA) identifies the most serious hazardous waste sites in the nation. These sites are then placed on the National Priorities List (NPL) and are targeted for long-term federal clean-up activities. Carbon monoxide has been found in at least 12 of the 1,699 current or former NPL sites. Although the total number of NPL sites evaluated for this substance is not known, the possibility exists that the number of sites at which carbon monoxide is found may increase in the future as more sites are evaluated. This information is important because these sites may be sources of exposure and exposure to this substance may be harmful.

When a substance is released either from a large area, such as an industrial plant, or from a container, such as a drum or bottle, it enters the environment. Such a release does not always lead to exposure. You can be exposed to a substance only when you come in contact with it. You may be exposed by breathing, eating, or drinking the substance, or by skin contact.

If you are exposed to carbon monoxide, many factors will determine whether you will be harmed. These factors include the dose (how much), the duration (how long), and how you come in contact with it. You must also consider any other chemicals you are exposed to and your age, sex, diet, family traits, lifestyle, pregnancy status, and state of health.

### 1.1 WHAT IS CARBON MONOXIDE?

<b>Carbon monoxide is a gas</b>	Carbon monoxide is a colorless, nonirritating, odorless, and tasteless gas. It is found in both outdoor and indoor air.
---------------------------------	---

## 1. PUBLIC HEALTH STATEMENT

<b>Sources of carbon monoxide in the atmosphere</b>	<p>Carbon monoxide is made when carbon in fuel is not burned completely. Carbon monoxide is produced from both human-made and natural sources. The most important human-made source of carbon monoxide arises from the exhaust of automobiles.</p> <p>Inside homes, improperly adjusted gas appliances, furnaces, wood burning stoves, and fireplaces are a potential source of carbon monoxide (see Section 1.3).</p>
<b>Industrial uses</b>	<p>Carbon monoxide can be used in industry to synthesize many compounds such as acetic anhydride, polycarbonates, acetic acid, and polyketone.</p>

## 1.2 WHAT HAPPENS TO CARBON MONOXIDE WHEN IT ENTERS THE ENVIRONMENT?

<b>Converts to carbon dioxide</b>	<p>When carbon monoxide is released to the environment, it enters the air and remains in the atmosphere for an average of about 2 months.</p> <p>Eventually, carbon monoxide reacts with other compounds in the atmosphere and is converted to carbon dioxide.</p> <p>Microorganisms found in soil and water can also convert carbon monoxide to carbon dioxide.</p>
-----------------------------------	--

## 1. PUBLIC HEALTH STATEMENT

## 1.3 HOW MIGHT I BE EXPOSED TO CARBON MONOXIDE?

<b>Sources of exposure</b>	<p>All people are exposed to carbon monoxide at varying levels through inhalation of air. Places and times of the day that have a lot of vehicular traffic generally have higher levels of carbon monoxide as compared to areas of low traffic.</p> <p>You can be exposed to carbon monoxide from cigarette smoke whether as a smoker or from second-hand smoke.</p> <p>You can be exposed to carbon monoxide by using gas appliances or wood burning stoves and fireplaces.</p> <p>People are exposed to carbon monoxide inside of vehicles.</p>
----------------------------	---

## 1.4 HOW CAN CARBON MONOXIDE ENTER AND LEAVE MY BODY?

<b>Carbon monoxide enters and leaves the body</b>	<p>Carbon monoxide in the air rapidly enters all parts of the body, including blood, brain, heart, and muscles when you breathe.</p> <p>The carbon monoxide in your body leaves the body through your lungs, when you breathe out (exhale), but there is a delay in eliminating carbon monoxide.</p> <p>It takes about a full day for carbon monoxide to leave your body.</p>
---	---

## 1.5 HOW CAN CARBON MONOXIDE AFFECT MY HEALTH?

This section looks at studies concerning potential health effects in animal and human studies.

<b>Carbon monoxide can harm the heart, brain, and lungs</b>	<p>Breathing high levels of carbon monoxide can kill you.</p> <p>Breathing lower levels of carbon monoxide can permanently harm your heart and brain.</p> <p>Carbon monoxide can be more harmful to you if you have heart or lung disease.</p>
---	--

## 1. PUBLIC HEALTH STATEMENT

**1.6 HOW CAN CARBON MONOXIDE AFFECT CHILDREN?**

This section discusses potential health effects in humans from exposures during the period from conception to maturity at 18 years of age.

<p><b>Breathing carbon monoxide during pregnancy can harm your unborn child</b></p>	<p>Breathing high levels of carbon monoxide can lead to miscarriage.</p> <p>Breathing lower levels of carbon monoxide during pregnancy may harm the mental development of your child.</p>
---	---

**1.7 HOW CAN FAMILIES REDUCE THE RISK OF EXPOSURE TO CARBON MONOXIDE?**

<p><b>Reduce indoor air levels of carbon monoxide</b></p>	<p>The most dangerous levels of carbon monoxide usually occur in indoor air. High levels occur as a result of improperly installed or unvented appliances that burn natural gasoline, kerosene, or other fuels. These include stoves, furnaces, heaters, and generators. Make sure that all of your appliances are installed properly and have periodic maintenance performed by professional installers. Always follow the manufacturer's recommendations on installing and using these devices.</p> <p>Make certain wood burning heaters and fireplaces are properly vented.</p> <p>Never use a gas-powered generator or burn charcoal indoors, as this can quickly lead to dangerous levels of carbon monoxide in your home.</p> <p>Do not use older portable propane heaters in enclosed indoor settings, including campers and tents, as dangerous levels of carbon monoxide can build up. Look for portable heaters that contain an oxygen depletion sensor (ODS) and are safer to use when camping. If oxygen levels start to fall, the sensor automatically shuts down the heater before it can produce dangerous levels of carbon monoxide. Older generation heaters without an ODS are intended for outdoor use only and should not be used indoors.</p> <p>Do not let your car idle for long periods of time in your garage.</p>
---	---

## 1. PUBLIC HEALTH STATEMENT

<b>Avoid tobacco smoke</b>	You can reduce your exposure to carbon monoxide by avoiding smoke from cigarettes and cigars since the sidestream smoke contains carbon monoxide.
<b>Install carbon monoxide detectors in your home</b>	Carbon monoxide detectors can be purchased at home remodeling or hardware stores. It is important to understand that most smoke detectors do not detect carbon monoxide, so you should install carbon monoxide detectors in your home as well as smoke detectors.

### 1.8 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO CARBON MONOXIDE?

<b>Carbon monoxide exposure can be measured with a blood test</b>	Medical devices called carbon monoxide-oximeters that are found in clinical laboratories or hospitals can estimate the level of carbon monoxide in blood by a simple test.
---	--

### 1.9 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH?

The federal government develops regulations and recommendations to protect public health. Regulations can be enforced by law. The EPA, the Occupational Safety and Health Administration (OSHA), and the Food and Drug Administration (FDA) are some federal agencies that develop regulations for toxic substances. Recommendations provide valuable guidelines to protect public health, but cannot be enforced by law. The Agency for Toxic Substances and Disease Registry (ATSDR) and the National Institute for Occupational Safety and Health (NIOSH) are two federal organizations that develop recommendations for toxic substances.

Regulations and recommendations can be expressed as “not-to-exceed” levels. These are levels of a toxic substance in air, water, soil, or food that do not exceed a critical value. This critical value is usually based on levels that affect animals; they are then adjusted to levels that will help protect humans. Sometimes these not-to-exceed levels differ among federal organizations because they used different exposure times (an 8-hour workday or a 24-hour day), different animal studies, or other factors.

Recommendations and regulations are also updated periodically as more information becomes available. For the most current information, check with the federal agency or organization that provides it.

## 1. PUBLIC HEALTH STATEMENT

Some regulations and recommendations for carbon monoxide include the following:

<b>Levels in air set by EPA</b>	EPA established an environmental limit of 10 mg/m <sup>3</sup> (9 ppmv) of carbon monoxide in air averaged over 8 hours and not to be exceeded more than once per year.
<b>Levels in workplace air set by OSHA</b>	OSHA set a legal limit of 55 mg/m <sup>3</sup> (50 ppmv) for carbon monoxide in air averaged over an 8-hour work day.

**1.10 WHERE CAN I GET MORE INFORMATION?**

If you have any more questions or concerns, please contact your community or state health or environmental quality department, or contact ATSDR at the address and phone number below.

ATSDR can also tell you the location of occupational and environmental health clinics. These clinics specialize in recognizing, evaluating, and treating illnesses that result from exposure to hazardous substances.

Toxicological profiles are also available on-line at [www.atsdr.cdc.gov](http://www.atsdr.cdc.gov) and on CD-ROM. You may request a copy of the ATSDR ToxProfiles™ CD-ROM by calling the toll-free information and technical assistance number at 1-800-CDCINFO (1-800-232-4636), by e-mail at [cdcinfo@cdc.gov](mailto:cdcinfo@cdc.gov), or by writing to:

Agency for Toxic Substances and Disease Registry  
 Division of Toxicology and Environmental Medicine  
 1600 Clifton Road NE  
 Mailstop F-62  
 Atlanta, GA 30333  
 Fax: 1-770-488-4178

Organizations for-profit may request copies of final Toxicological Profiles from the following:

National Technical Information Service (NTIS)  
 5285 Port Royal Road  
 Springfield, VA 22161  
 Phone: 1-800-553-6847 or 1-703-605-6000  
 Web site: <http://www.ntis.gov/>

## 2. RELEVANCE TO PUBLIC HEALTH

### 2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO CARBON MONOXIDE IN THE UNITED STATES

Carbon monoxide is a colorless, odorless, non-irritating, and tasteless gas that is ubiquitous in the atmosphere. It arises from both natural and anthropogenic sources. It is produced as a primary pollutant during the incomplete combustion of fossil fuels and biomass. Carbon monoxide is also produced indirectly from the photochemical oxidation of methane and other volatile organic compounds (VOCs) in the atmosphere. Vegetation can emit carbon monoxide directly into the atmosphere as a metabolic byproduct, and the photooxidation of organic matter in surface waters (lakes, streams, rivers, oceans) and soil surfaces also results in the formation of carbon monoxide. Volcanic activity is an additional natural source of carbon monoxide in the atmosphere. Carbon monoxide is also produced endogenously in humans during the normal catabolism of hemoglobin (Hb). The vast majority of anthropogenic carbon monoxide emissions arise from gasoline-powered automobile usage, although the total amount of carbon monoxide emitted to the environment from this source has declined significantly over the past several decades due to the use of catalytic converters and other emission control devices that are standard equipment on modern passenger vehicles. However, there are numerous additional sources of carbon monoxide generation that while minor in terms of total tonnage released to the atmosphere, may have significance with regard to generation of carboxyhemoglobin (COHb) in exposed individuals.

The annual average outdoor carbon monoxide concentrations are roughly 0.12 parts per million by volume (ppmv) in the Northern Hemisphere and about 0.04 ppmv in the Southern Hemisphere. These levels are variable throughout the course of the year, with seasonal maximum levels occurring during late winter in both hemispheres and minimum levels being observed during late summer. Carbon monoxide concentrations are reported to range from a minimum of about 0.03 ppmv during summer in the Southern Hemisphere to about 0.20 ppmv at high latitudes in the Northern Hemisphere during winter. Urban locations with high automobile usage or a high volume of stationary emission sources such as refineries or power plants typically have greater atmospheric levels of carbon monoxide as compared to rural or remote sites. Carbon monoxide levels in indoor air are strongly influenced by the presence of various appliances and whether or not the occupants of the residence smoke tobacco products. Unvented kerosene and gas space heaters; leaking chimneys and furnaces; back-drafting from furnaces, gas water heaters, wood stoves, and fireplaces; gas stoves, generators and other gasoline-powered equipment; automobile exhaust from attached garages; and tobacco smoke all contribute to indoor air levels of carbon

## 2. RELEVANCE TO PUBLIC HEALTH

monoxide. Average levels in homes without gas stoves vary from 0.5 to 5 ppmv. Levels near properly adjusted gas stoves are often 5–15 ppmv and those near poorly adjusted stoves may be  $\geq 30$  ppmv.

Exposure of the general population to carbon monoxide occurs through inhalation of outdoor and indoor air. Populations living in urban areas with heavy vehicular traffic or stationary sources such as petroleum refineries, gas and coal burning power plants, petrochemical plants, and coke oven plants are more likely to be exposed to higher levels of carbon monoxide from ambient outdoor air. Occupational exposure for employees who work in these industries and other occupations that are subject to high levels of vehicular exhaust (such as taxi cab drivers and toll booth workers) are also likely to be exposed to higher levels. Industrial or in-home use of methylene chloride paint strippers in poorly ventilated areas can lead to high levels of carbon monoxide in blood since carbon monoxide is a metabolic byproduct of methylene chloride. Members of the public who smoke or work in smoke-filled environments such as restaurants, bars, and casinos where smoking is allowed are also exposed to higher levels of carbon monoxide than members of the population who do not smoke and are not frequently exposed to second-hand tobacco smoke. Section 6.5 discusses exposures to the general population and occupational exposures in greater detail.

### 2.2 SUMMARY OF HEALTH EFFECTS

Health effects associated with acute carbon monoxide poisoning have been extensively documented. In the last decade, growing evidence has revealed endogenous carbon monoxide (produced from catabolism of heme and other endogenous precursors) to be a cell signaling agent that contributes to the regulation of numerous physiological systems, including brain and muscle oxygen storage and utilization (myoglobin, neuroglobin), relaxation of vascular and extra-vascular smooth muscle, modulation of synaptic neurotransmission, anti-inflammation, anti-apoptosis, anti-proliferation, and anti-thrombosis (see Section 3.5.2, Mechanisms of Toxicity). Current toxicological and epidemiological research has focused on examining health effects of low-level carbon monoxide exposures that do not result in overt carbon monoxide poisoning and attempting to understand the connections between carbon monoxide toxicity and carbon monoxide physiology. This research has revealed that the heart and cardiovascular system, and brain and developing nervous system are particularly sensitive to carbon monoxide. These studies have also shown that people with ongoing cardiovascular and/or respiratory disease may be particularly vulnerable to carbon monoxide.

## 2. RELEVANCE TO PUBLIC HEALTH

***Modes of Action of Carbon Monoxide.*** Carbon monoxide exerts effects on cell metabolism through hypoxic and non-hypoxic modes of action. Both modes of action are thought to be largely (if not entirely) the result of the ability of carbon monoxide to bind to heme and alter function and/or metabolism of heme proteins. Formation of COHb decreases the O<sub>2</sub> carrying capacity of blood and impairs release of O<sub>2</sub> from Hb for its utilization in tissues. Through similar mechanisms, carbon monoxide decreases O<sub>2</sub> storage in muscle cells by binding to and displacing O<sub>2</sub> from myoglobin. Although all tissues are vulnerable to carbon monoxide-induced hypoxic injury, those having the highest O<sub>2</sub> demand are particularly vulnerable, including the brain and heart.

Most of the non-hypoxic mechanisms of action of carbon monoxide have been attributed to binding of carbon monoxide to heme in proteins other than Hb. Notable targets of carbon monoxide include components of several important physiological regulatory systems, including brain and muscle oxygen storage and utilization (myoglobin, neuroglobin); nitric oxide cell signaling pathway (e.g., nitric oxide synthase, guanylyl cyclase); prostaglandin cell signaling pathway (cyclooxygenase, prostaglandin H synthase); energy metabolism and mitochondrial respiration (cytochrome c oxidase, cytochrome c, NADPH oxidase); steroid and drug metabolism (cytochrome P450); cellular redox balance and reactive oxygen species (catalase, peroxidases); and various transcription factors (e.g., neuronal PAS domain protein, NPAS2).

***Endogenous Carbon Monoxide.*** In addition to inhalation exposure to carbon monoxide in air, internal exposures to carbon monoxide occur as a result of production of carbon monoxide from endogenous precursors (e.g., heme degradation, auto-oxidation of phenols, photo-oxidation of organic compounds, and lipid peroxidation of cell membrane lipids) and production of carbon monoxide from oxidative metabolism of exogenous precursors (e.g., carbon tetrachloride, dichloromethane and other dihalomethanes). The latter two metabolic sources of carbon monoxide result in a carbon monoxide body burden in the absence of exposure to exogenous carbon monoxide in air. Endogenous carbon monoxide production rate has been estimated to be approximately 0.42 mL carbon monoxide STPD/hour, 0.006 mL carbon monoxide/hour-kg body weight. However, numerous physiological and disease factors affect the rate of endogenous production of carbon monoxide, including the menstrual cycle, pregnancy, diseases, and stimuli that increase catabolism of Hb or other heme proteins, including hemolysis, hematomas, hemolytic anemias, thalassemia, and Gilbert's syndrome.

Growing evidence has revealed endogenous carbon monoxide to be a cell signaling agent that contributes to the regulation of numerous physiological systems, including brain and muscle oxygen storage and

## 2. RELEVANCE TO PUBLIC HEALTH

utilization (myoglobin, neuroglobin), relaxation of vascular and extra-vascular smooth muscle, modulation of synaptic neurotransmission, anti-inflammation, anti-apoptosis, anti-proliferation, and anti-thrombosis. This has potentially important implications for the understanding of carbon monoxide toxicology and dose-response relationships for the following reasons: (1) carbon monoxide modulation of physiological processes may underlie some aspects of the toxicity of exogenous carbon monoxide; (2) exogenous carbon monoxide may disrupt physiological regulation of those systems that are responsive to endogenous carbon monoxide (e.g., vascular resistance); and (3) exposures to exogenous carbon monoxide may affect carbon monoxide-mediated physiological responses at levels that approach those resulting from endogenous production. One implication of this is that the dose threshold for effects of exogenous carbon monoxide on carbon monoxide-modulated physiological systems may lie near or below ambient air carbon monoxide concentrations.

***Toxicokinetics.*** Inhaled carbon monoxide is rapidly and extensively absorbed into blood and distributes throughout the body. The distribution of carbon monoxide in the body largely reflects the binding of carbon monoxide to heme proteins (e.g., Hb, myoglobin). Measurements of total carbon monoxide concentrations in tissues obtained from human autopsies showed the highest concentrations in blood, spleen, lung, kidney, and skeletal muscle, with detectable levels also in brain and adipose tissue (Vreman et al. 2006). Higher concentrations of carbon monoxide in blood, heart, skeletal muscle, and spleen reflect the abundance of the major carbon monoxide binding proteins in these tissues. In blood, carbon monoxide rapidly distributes into erythrocytes where it exists primarily as a complex with Hb (COHb). Carbon monoxide in muscle exists primarily as a complex with myoglobin (COMb). Carbon monoxide in the maternal system distributes to fetal tissues where it binds to fetal Hb and other heme proteins. Steady-state fetal blood COHb concentrations are approximately 10–15% higher than maternal blood (fetal/maternal ratio=1.1–1.25) and fetal blood COHb elimination kinetics are slower than maternal (Hill et al. 1977; Longo 1977). Carbon monoxide binding to fetal Hb is analogously similar to maternal Hb.

Absorbed carbon monoxide is eliminated from the body by exhalation and oxidative metabolism. Oxidative metabolism of carbon monoxide has been estimated to be a relatively small fraction (<10%) of endogenous carbon monoxide elimination. Under most conditions, the dominant route of elimination of absorbed carbon monoxide is exhalation. The decline in blood %COHb following cessation of an inhalation exposure to carbon monoxide exhibits at least two kinetic phases. The fast phase is thought to reflect a combination of exhalation of carbon monoxide along with slower distribution of blood carbon monoxide to tissues that continues after cessation of exposure. The elimination half-time for the slow phase is approximately 100–300 minutes. The carbon monoxide elimination half-time increases with age,

## 2. RELEVANCE TO PUBLIC HEALTH

with the most pronounced increase occurring from age 2 to 20 years and is marginally longer in males compared to females. Exercise decreases carbon monoxide elimination half-time, although exercise and the increase in respiration would lead to increased CO exposure, if CO is still present in inspired air.

The importance of COHb as a potential biomarker of carbon monoxide exposure and hypoxia, particularly at low levels of exposure, has led to the development of a physiologically-based mechanistic model of carbon monoxide kinetics for predicting relationships between exposure and blood COHb levels. The model that has received the greatest attention and use in risk assessment and in clinical medicine is the Coburn-Forster-Kane (CFK) model. This model can be used to predict steady-state blood COHb levels that correspond to a given continuous inhalation exposure to carbon monoxide in a typical adult. The CFK model has been used to support discussions of the health effects of carbon monoxide in this Toxicological Profile, by providing a means for interconverting carbon monoxide exposure levels expressed in units of ppm or mg/m<sup>3</sup> and corresponding equivalent steady-state COHb% values (i.e., the COHb% that would be achieved with continuous exposure to the reported air carbon monoxide concentration). Predicted steady-state blood COHb levels corresponding to a range of carbon monoxide exposure concentrations are presented in the introduction to Section 3.2 (Table 3-1). Several other toxicokinetics models are described in Section 3.4.5.

***Acute Carbon Monoxide Poisoning.*** Carbon monoxide poisoning is one of leading causes of morbidity and mortality due to poisoning in the United States. The principal mechanism of many adverse effects of carbon monoxide exposure is COHb-induced tissue hypoxia; thus, tissues with high oxygen requirements (e.g., brain, heart) are the most sensitive to carbon monoxide-induced hypoxia. However, other non-hypoxic mechanisms (e.g., binding of carbon monoxide to other heme proteins, such as myoglobin and cytochrome c oxidase) and alterations in biological and physiological functions of endogenous carbon monoxide, likely contribute to the adverse effects of acute carbon monoxide poisoning.

The extent of injury from acute carbon monoxide exposure depends upon the concentration and duration of exposure and the underlying health status of the exposed individual. The most commonly reported signs and symptoms associated with acute carbon monoxide poisoning are due to effects on the central nervous system and the cardiovascular system; however, because carbon monoxide exposure has the potential to affect nearly all tissues, the clinical presentation of acute carbon monoxide poisoning includes a wide range of symptoms. The severity of carbon monoxide poisoning is typically categorized as mild, moderate, or severe, based on clinical presentation. Signs and symptoms of mild carbon monoxide poisoning include headache, nausea, vomiting, dizziness, and blurred vision; headache and dizziness are

## 2. RELEVANCE TO PUBLIC HEALTH

the most commonly reported symptoms. Because these symptoms mimic flu-like viral illnesses, mild carbon monoxide poisoning can easily be misdiagnosed. Symptoms associated with moderate carbon monoxide poisoning may include confusion, syncope, chest pain, dyspnea, weakness, tachycardia, tachypnea, and rhabdomyolysis. Effects of severe poisoning may be life-threatening, including cardiac arrhythmias, myocardial ischemia, cardiac arrest, hypotension, respiratory arrest, noncardiogenic pulmonary edema, seizures, and coma. In addition to the immediate-onset effects of exposure, delayed-onset development of neuropsychiatric impairment typically occurs from several days to approximately 3–4 weeks of exposure, with symptoms including inappropriate euphoria, impaired judgment, poor concentration, memory loss, cognitive and personality changes, psychosis, and Parkinsonism. Acute carbon monoxide poisoning during pregnancy has been associated with spontaneous abortion and fetal death; pregnancy outcome is likely to be dependent upon the severity of maternal poisoning and fetal age.

The relationship between the severity of clinical signs and symptoms of acute carbon monoxide poisoning and COHb levels is not well correlated. The poor correlation may be due to the length of time elapsed between cessation of exposure and measurement of COHb levels or to effects of supplemental oxygen treatment prior to COHb measurement. Generally, in healthy individuals, mild carbon monoxide poisoning that requires medical intervention are associated with COHb levels >20%. Fatalities due to carbon monoxide poisoning have been reported for a wide range of COHb levels (3–70%). Levels of COHb >50% are frequently fatal.

***Primary Targets of Low-level Carbon Monoxide Exposure.*** The primary targets of low-level exposures to carbon monoxide (i.e., those that result in blood COHb levels <20%) appear to include the heart and cardiovascular system, central nervous system, and the fetus and neonate. Adverse effects in the respiratory tract have also been observed in human clinical studies and in animal studies at higher exposures than those associated with effects on the cardiovascular system, central nervous system, and on development. A large body of epidemiologic studies also has provided evidence that ambient levels of carbon monoxide in air may contribute to respiratory morbidity and aggravation of ongoing respiratory disease (e.g., asthma; see Table 3-3). Epidemiologic studies have also examined possible associations between ambient air carbon monoxide concentrations and hematologic biomarkers of coagulation and inflammation. Although some studies have found significant associations, collectively, findings from these studies are inconclusive.

**Cardiovascular System.** Cardiovascular effects of inhalation exposures to carbon monoxide have been evaluated in controlled human clinical studies, epidemiology studies, and in various animal models

## 2. RELEVANCE TO PUBLIC HEALTH

(monkeys, dogs, rats, and rabbits). In general, these studies provide convincing evidence for adverse cardiovascular effects in association with carbon monoxide exposures that result in blood COHb levels of  $\geq 2.4\%$ , with effects occurring at the lowest levels in subjects who have compromised cardiovascular function (e.g., coronary artery disease).

Results of controlled clinical studies in patients with coronary artery disease show that acute-duration exposure to carbon monoxide at levels producing blood COHb levels between 2.4 and 5.9% exacerbates underlying cardiovascular disease, including enhanced myocardial ischemia and increased cardiac arrhythmias. In patients with exertional angina, carbon monoxide exposure exacerbated exercise-induced myocardial ischemia, including decreased time-to-onset of angina symptoms, increased duration of angina symptoms, decreased time-to-onset of ST-segment depression (EKG change indicative of myocardial ischemia), and decreased left ventricular ejection fraction. At the lowest blood COHb level evaluated in patients (i.e., COHb 2.4%), time-to-onset of angina symptoms and of ST-segment depression were significantly decreased by 4.2 and 5.1%, respectively.

Epidemiological studies of exposure to carbon monoxide and cardiovascular outcomes have yielded mixed results. In general, the weight of evidence suggests that risks of certain specific outcomes (hospitalizations and emergency room visits related to congestive heart failure, ischemic heart disease, myocardial infarction, and stroke) are associated with increasing ambient carbon monoxide concentrations. The interpretation of these associations is complicated by the possibility that ambient air carbon monoxide levels may be a surrogate measure for air pollution in general. However, the corroborated observations of associations between carbon monoxide exposure and outcomes related to ischemic heart disease is particularly provocative in the context of results of human clinical studies in which carbon monoxide-induced hypoxia exacerbated ischemia symptoms in patients with coronary artery disease. Mean ambient air carbon monoxide concentrations reported in studies that have found carbon monoxide-associated adverse cardiovascular outcomes have ranged from 0.5 to 10 ppm, with maximum values ranging from 2 to 50 ppm. These values correspond to approximate steady-state blood COHb levels of  $<2\%$  for the mean, and  $<10\%$  for the maximum.

Studies in animals have investigated adverse cardiovascular effects of carbon monoxide exposure over a much wider range of exposure conditions (e.g., exposure concentration and duration) and have evaluated additional outcome measures that are not possible to assess in humans. These studies provide further evidence of adverse cardiovascular effects of carbon monoxide exposure, including compensatory alterations in hemodynamics, cardiac hypertrophy, cardiac arrhythmias, and possibly atherosclerosis.

## 2. RELEVANCE TO PUBLIC HEALTH

Based on studies described above, a blood COHb concentration of 2.4% is identified as lowest-observed-adverse-effect level (LOAEL) for adverse cardiovascular outcomes in coronary artery disease patients. A no-observed-adverse-effect level (NOAEL) for this effect was not identified. The LOAEL for COHb can be converted to an equivalent human exposure concentration (for continuous exposure) that would yield the same steady-state blood COHb concentration (2.4%) by implementing the CFK model. The human equivalent exposure concentration is approximately 14 ppm.

**Developmental Effects.** Epidemiological studies have examined possible associations between exposure to ambient air carbon monoxide concentrations and various developmental outcomes, including pre-term birth, birth weight, congenital anomalies, and neonatal and infant death. Results of these studies have been mixed and, collectively do not provide strong evidence for developmental effects in association with exposures to ambient levels of carbon monoxide. In general, these studies have examined relatively low air carbon monoxide concentrations, typical of ambient levels (e.g., mean concentrations ranging from 0.5 to 3 ppm, with highest reported values <10 ppm). The air carbon monoxide measurements do not necessarily represent exposures that occurred during gestation or during any particular period of gestation.

Numerous studies on developmental effects of gestational and early postnatal exposure to carbon monoxide have been conducted in animals. In general, most studies evaluated effects of relatively low carbon monoxide concentrations (i.e.,  $\leq 300$  ppm), with exposure concentrations selected to produce maternal COHb levels typically associated with smoking (5–10%); however, studies did not consistently report maternal or fetal COHb levels. Studies in animals have examined effects of carbon monoxide exposure on numerous developmental outcomes, including several outcomes that have not been assessed in epidemiological studies (e.g., auditory and immune system development). Results of animal studies show adverse developmental effects of gestational and early postnatal carbon monoxide exposure, including decreased fetal weight, adverse central nervous system development, altered peripheral nervous system development, cardiac effects, altered sexual behavior, immunological effects, and hematological effects. In addition, some studies showed that developmental effects persisted beyond the postnatal period, although persistence of effects was not examined in all studies. The lowest LOAEL values for developmental effects were obtained in studies evaluating effects of carbon monoxide on the developing auditory system (i.e., LOAEL 12–25 ppm carbon monoxide); however, since other developmental outcomes were not assessed at this range of low carbon monoxide concentrations, it is not possible to determine if the developing auditory system is more sensitive to carbon monoxide exposure than other

## 2. RELEVANCE TO PUBLIC HEALTH

systems. Gestational and/or early postnatal exposure of rats to 25 ppm carbon monoxide produced morphological changes in the developing auditory system, including swelling, cytoplasmic vacuolization and atrophy of nerve terminals innervating inner hair cells, “distorted myelin” with vacuolization in the 8th cranial nerve at the level of the internal auditory canal, decreased immunoreactivity of the enzymes cytochrome oxidase, NADH-TR, and calcium-mediated myosin ATPase, and decreased immunostaining of neurofilament and myelin basic protein in the organ of corti. Exposure of rat pups during the early postnatal period decreased action potential amplitude of the 8th cranial nerve at  $\geq 12$  ppm carbon monoxide and decreased in otoacoustic emissions at  $\geq 50$  ppm carbon monoxide, with effects on action potential amplitude of the 8th nerve persisting through age 73 days.

Based on studies described above, a maternal exposure concentration of 12 ppm is identified as a LOAEL for adverse neurodevelopmental outcomes in rats. This LOAEL can be converted to an equivalent blood COHb level in the rat by implementation of the CFK model, as adapted for the rat. The time-averaged blood COHb level predicted for the 16-day exposure (22 hours/day) is 1.8%. The equivalent human exposure concentration (for continuous exposure) that would yield the same steady-state blood COHb concentration (1.82%) is approximately 10 ppm

**Central Nervous System.** As previously noted, acute exposure to high levels of carbon monoxide produces symptoms of central nervous system toxicity, including headache, dizziness, drowsiness, weakness, nausea, vomiting, confusion, disorientation, irritability, visual disturbances, convulsions, and coma. Lesions of the basal ganglia (primarily of the globus pallidus) and white matter have also been observed in magnetic resonance imaging (MRI) and computed tomography (CT) scans in association with acute carbon monoxide poisoning. Motor impairments consistent with damage to basal ganglia have been observed following carbon monoxide poisoning. Following acute-onset effects, delayed development of neuropsychiatric impairment may occur from several days to 3–4 weeks of exposure, with symptoms including inappropriate euphoria, impaired judgment, poor concentration, memory loss, cognitive and personality changes, psychosis, and Parkinsonism. Delayed neuropsychiatric impairment has been estimated to occur in up to 68% of patients with acute carbon monoxide poisoning. There is a poor correlation between initial symptom severity and the likelihood of developing of delayed neuropsychiatric impairment has not been established.

Based on an extensive database of acute carbon monoxide poisoning, it is generally accepted that central nervous system symptoms are associated with acute exposures that result in blood COHb levels  $\geq 20\%$ . However, despite extensive clinical experience, general consensus on the dose-response relationship for

## 2. RELEVANCE TO PUBLIC HEALTH

carbon monoxide-induced nervous system effects at blood COHb levels between 5 and 20% has not been achieved.

Based on studies described above, a blood COHb concentration of 20% is identified as a LOAEL for adverse neurological outcomes, although the LOAEL range may extend below this level to 5%. These levels, 5 and 20%, can be converted to equivalent human exposure concentrations (for continuous exposure) that would yield the same steady-state blood COHb concentration by implementing the CFK model. The human equivalent exposure concentrations are approximately 32 ppm (COHb=5%) and 160 ppm (COHb=20%).

**Respiratory Effects.** Although cardiopulmonary arrest is an end point of fatal carbon monoxide poisoning, results of controlled clinical studies in healthy subjects indicate that the respiratory tract does not appear to be a primary target organ for carbon monoxide toxicity. Brief exposure to carbon monoxide at levels >1,000 ppm may decrease ventilatory performance, although conflicting results have been reported. Epidemiological studies have examined possible associations between ambient air carbon monoxide concentrations and mortality. These studies have examined relatively low carbon monoxide concentrations (from a toxicological perspective) that are typical of ambient conditions of study period (mean concentrations ranging from 0.3 to 10 ppm with the highest values <30 ppm). Collectively, these studies have yielded mixed results, with some studies finding significant associations between increasing ambient air carbon monoxide concentrations and respiratory outcomes (e.g., exacerbation of asthma symptoms, hospitalizations and emergency room visits related to asthma), and few studies finding associations that persist after accounting for exposures to other air pollutants that also have been shown to contribute to respiratory disease risk (e.g., NO<sub>2</sub>, O<sub>3</sub>, particulate matter, and SO<sub>2</sub>). The lack of strong evidence for associations between ambient air carbon monoxide concentrations at <30 ppm and pulmonary function is also consistent with the results of human clinical studies. Studies conducted in animals provide supporting evidence that the respiratory tract does not appear to be a primary target organ for carbon monoxide. Most of these studies evaluated carbon monoxide exposures that produced much higher COHb concentrations (i.e., COHb >50%) than those evaluated in controlled clinical studies in humans.

**Hematological Effects.** Hematological effects of carbon monoxide include compensatory responses to tissue hypoxia resulting from binding of carbon monoxide to Hb (e.g., increased blood volume, Hb, hematocrit, and erythrocyte count and volume). Possible associations between ambient air carbon monoxide concentrations and biomarkers of coagulation and inflammation have been examined in

## 2. RELEVANCE TO PUBLIC HEALTH

epidemiological studies. Biomarkers examined have included inflammation markers, CRP, serum amyloid A (SAA), and white blood cell count (WBC); cell adhesion markers, E-selectin, von Willebrand factor, antigen (vWF), ICAM-1; and coagulation markers, fibrinogen, factor VII (FVII), prothrombin fragment 1+2, prothrombin time (PT), and activated partial thromboplastin time (APTT). Although some studies have found significant associations between environmental carbon monoxide exposures and alteration in plasma proteins, these studies do not distinguish between possible direct effects of carbon monoxide and/or other air pollutants directly on coagulation and immune systems from indirect effects that result in changes in blood biomarkers. Furthermore, findings across studies are inconsistent. Environmental carbon monoxide exposures were shown to correlate with elevated C reactive protein in one trial, but not another by the same group. Plasma fibrinogen was increased in one study, decreased in one study, and unchanged in two studies. Other studies have reported an elevation in soluble intercellular adhesion molecule-1, along with decreases in factor VII, serum albumin, and prothrombin time. Therefore, no conclusions can be drawn at this time on whether low-level environmental exposures may alter plasma inflammatory or coagulation markers. Nevertheless, direct effects are plausible, given mechanistic studies that have revealed evidence that endogenous carbon monoxide may participate in the regulation of thrombosis and immune function (see Section 3.5.2, Mechanisms of Toxicity).


### 2.3 CARBON MONOXIDE DOSE-RESPONSE RELATIONSHIPS

Epidemiological and clinical studies provide evidence for a progression of some of the adverse health effects of carbon monoxide in humans with increasing blood levels of COHb (Figure 2-1). The relationship shown in Figure 2-1 does not necessarily mean that these effects result directly from formation of COHb at the expense of decreasing O<sub>2</sub>Hb levels in blood (i.e., hypoxic mechanisms). Other important mechanisms, previously mentioned and described in greater detail in Section 3.5.2, (Mechanisms of Toxicity), may also contribute to these effects. COHb may serve as a biomarker for carbon monoxide body burden or carbon monoxide burdens in specific target tissues where nonhypoxic modes of actions of carbon monoxide exert effects.

An alternative presentation of the relationship between blood COHb levels and adverse health effects is provided in Table 2-1. This table shows the predicted relationship between blood COHb levels that

2. RELEVANCE TO PUBLIC HEALTH

**Figure 2-1. Blood Carboxyhemoglobin Levels Corresponding to Adverse Health Effects of Carbon Monoxide in Humans**



## 2. RELEVANCE TO PUBLIC HEALTH

**Table 2-1. Blood Carboxyhemoglobin Levels Corresponding to Adverse Health Effects of Carbon Monoxide**

Effect	COHb <sup>a</sup> (Percent)	Exposure <sup>b</sup> (ppm)
Endogenous production	<0.5	0
Typical level in nonsmoker	0.5–1.5	1–8
Increased risk of arrhythmias in coronary artery disease patients and exacerbation of asthma (epidemiological studies)	0.3–2 <sup>b</sup>	0.5–10 <sup>b</sup>
Neurodevelopmental effects on the auditory system in rats	2–4 <sup>b</sup>	12–25 <sup>b</sup>
Enhanced myocardial ischemia and increased cardiac arrhythmias in coronary artery disease patients	2.4–6	14–40
Decreased exercise stamina in healthy adults	5–8	30–50
Neurobehavioral/cognitive changes, including visual and auditory sensory effects (decreased visual tracking, visual and auditory vigilance, visual perception), fine and sensorimotor performance, cognitive effects (altered time discrimination, learning, attention level, driving performance), and brain electrical activity	5–20	30–160
Acute and delayed onset of neurological impairment (headache, dizziness, drowsiness, weakness, nausea, vomiting, confusion, disorientation, irritability, visual disturbances, convulsions, and coma), and pathology (basal ganglia lesions)	20–60	160–1000
High risk of death	>50	>600

<sup>a</sup>Reported value, unless otherwise denoted as predicted.

<sup>b</sup>Predicted from the Coburn-Forster-Kane (CFK) model (unless otherwise denoted as reported), with a rate of endogenous carbon monoxide production assigned a value of 0.006 mL CO/kg body weight and all other parameter values as noted in Table 3-13.

## 2. RELEVANCE TO PUBLIC HEALTH

roughly correspond to adverse health effects and their corresponding equivalent human exposure concentrations (HEC) that would result in the same-steady state blood COHb level. For example, in Table 2-1, a continuous exposure to approximately 14 ppm for a period exceeding 16 hours (i.e., sufficient to achieve steady state) would be expected to result in a blood COHb level of approximately 2.4%, the lower end of the range for cardiac effects in coronary artery disease patients. All predictions shown in Table 2-1 are based on the CKF model, described in Section 3.4.5 (Physiologically Based Pharmacokinetic/Pharmacodynamic Models).

#### 2.4 MINIMAL RISK LEVELS (MRLs)

Given the above considerations, MRLs for carbon monoxide are not proposed at this time. The rationale for this determination is as follows:

- (1) Growing evidence suggests that endogenous carbon monoxide production is physiologically regulated and plays a role in regulating various important physiological processes, including processes that may underlie adverse effects on the cardiovascular system, immune system, blood coagulation system, and nervous system that have been observed in human clinical studies, epidemiological studies, and animal studies.
- (2) Given the physiological role of endogenous carbon monoxide, it is likely that an exposure threshold for carbon monoxide actions, if one exists at all, would be at or near the endogenous production rate. Therefore, any exogenous source of carbon monoxide exposure would have the potential for exceeding the threshold and producing potentially adverse effects.
- (3) Although there may be an exposure level that can be tolerated with minimal risk of adverse effects, the currently available toxicological and epidemiological data do not identify such minimal risk levels. Experimental clinical studies and animal toxicology studies that identify the lowest LOAELs do not identify NOAELs. These LOAELs are relatively low: COHb 2.4% for cardiovascular effects in humans, exposure concentrations of  $\geq 12$  ppm in rats for developmental effects. Converting these to human equivalent exposure concentrations (i.e., levels of continuous exposure that would result in steady-state COHb concentration in blood estimated from the CFK model) yield corresponding LOAELs of 14 and 10 ppm, respectively. Application of appropriate uncertainty factors to these LOAELs (e.g., for extrapolation from a LOAEL, for extrapolation from animals to humans, and for extrapolation to sensitive subpopulations), would result in MRLs that are 30–100 times lower than the corresponding LOAELs (e.g., approximately 0.1–0.5 ppm). These values are within the range of ambient carbon monoxide concentrations in the United States and would result, even for acute (e.g., 14-day) exposures, in internal doses that would be similar to endogenous production of carbon monoxide.
- (4) Any exposure level determined to be of minimal risk at sea level would not necessarily be of minimal risk at higher altitudes (i.e., at lower O<sub>2</sub> partial pressures). This would apply, in particular, to modes of actions of carbon monoxide that involve competition between carbon monoxide and O<sub>2</sub> for heme binding sites. The latter would include hypoxic mechanisms of actions that appear to underlie adverse cardiovascular effects of carbon monoxide (e.g., exacerbation of exercise induced arrhythmias in patients who have coronary artery disease).

### 3. HEALTH EFFECTS

#### 3.1 INTRODUCTION

The primary purpose of this chapter is to provide public health officials, physicians, toxicologists, and other interested individuals and groups with an overall perspective on the toxicology of carbon monoxide. It contains descriptions and evaluations of toxicological studies and epidemiological investigations and provides conclusions, where possible, on the relevance of toxicity and toxicokinetic data to public health.

A glossary and list of acronyms, abbreviations, and symbols can be found at the end of this profile.

*Focus of the Health Effects Review.* The literature on the toxicology of carbon monoxide is immense. Relatively recent reviews of the literature have been prepared by various organizations, including WHO (1999). EPA (2000, 2009g) and numerous reviews have been compiled on the clinical toxicology and management of acute carbon monoxide poisoning. The review of the literature provided in this section is not intended to be comprehensive; rather, it is focused on the more important, recent developments in the toxicological and epidemiological assessment of carbon monoxide. Studies that have potential relevance to understanding the lower end of the dose-response relationship for carbon monoxide are emphasized. Typically, dose-response information is presented in Toxicological Profiles in Levels of Significant Exposure (LSE) tables and figures. However, this information on carbon monoxide has been represented in various types of units (e.g., exposure concentration, blood COHb), and end points are not easily assigned into categories of no-observed-adverse-effect-level (NOAEL) or lowest-observed-adverse-effect-level (LOAEL) (e.g., odds ratio from epidemiological studies). Therefore, in place of an LSE table, Table 3-1 provides an overview of exposures (ppm) and blood COHb levels (%) that have been associated with specific categories of health effects. Table 3-2 can be used to convert the reported ambient air carbon monoxide concentrations into equivalent predicted blood COHb levels. However, considerable uncertainty attends to these dose interconversions as well as their relevance to a particular outcome. Predicted COHb levels and corresponding exposure concentrations may have greater relevance to outcomes mediated through hypoxic mechanisms than to those mediated through nonhypoxic mechanisms. Furthermore, steady-state requires exposure durations of approximately 16–24 hours. Exposures to lower levels of carbon monoxide for longer durations and exposures to higher levels for shorter durations that achieve similar blood COHb levels may not yield equivalent responses.

The discussion of health effects has been limited to the inhalation exposure pathway. Carbon monoxide exists in the environment as a gas (Henry's law constant  $>50,000$  atm/mol fraction, 25 °C). As a result,

**Table 3-1. Carbon Monoxide Exposures and Carboxyhemoglobin Levels Associated with Health Effects from Selected Studies Representing the Lowest Adverse Effect Levels**

Study type	Subjects	Exposure (ppm)	COHb (%)	Effect	Comments	Reference
<b>Respiratory</b>						
Epidemiological	Asthmatic children	0.3–2	NR	Asthma	Exacerbation of asthma. Associations confounded by co-exposure to other air pollutants (NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , SO <sub>2</sub> ).	Park et al. 2005a; Rabinovitch et al. 2004; Rodriguez et al. 2007; Schildcrout et al. 2006; Silkoff et al. 2005; Slaughter et al. 2003; von Klot et al. 2002; Yu et al. 2000
<b>Cardiovascular</b>						
Clinical (acute exposure)	Coronary artery disease patients (63 male adult nonsmokers)	117 (1 hour)	2.4%	Myocardial ischemia	Decreased time-to-onset of angina and arrhythmia (ST-segment changes) during exercise.	Allred et al. 1989, 1991
Clinical (acute exposure)	Healthy adults (15 male adult nonsmokers)	NR	5.1%	Performance	Decreased maximal exercise duration and effort. No effect on heart rate or rhythm, cardiac perfusion, or blood pressure.	Adir et al. 1999
Epidemiological	General public	0.3–2	NR	Cardiovascular disease	Increased risk of congestive heart failure, ischemic heart disease, myocardial infarction, and stroke. Effect size more pronounced in elderly and people with ongoing respiratory or cardiovascular disease. Associations confounded by co-exposure to other air pollutants (NO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , SO <sub>2</sub> ).	Berger et al. 2006; D'Ippoliti et al. 2003; Hosseinpoor et al. 2005; Lanki et al. 2006; Lee et al. 2003b; Mann et al. 2002; Szyszkowicz 2007; von Klot et al. 2005
Animal study, chronic exposure	Rats	200 (20 hour/day, 72	NR	Cardiomegaly	Left and right ventricle weights increased by 20% (p<0.001) and 14% (p<0.001).	Sørhaug et al. 2006

**Table 3-1. Carbon Monoxide Exposures and Carboxyhemoglobin Levels Associated with Health Effects from Selected Studies Representing the Lowest Adverse Effect Levels**

Study type	Subjects	Exposure (ppm) week)	COHb (%)	Effect	Comments	Reference
<b>Neurological</b>						
Clinical, acute exposure	Healthy adults	NR	5–20	Neurobehavioral/cognitive changes.	Visual and auditory sensory effects (decreased visual tracking, visual and auditory vigilance, visual perception), fine and sensorimotor performance, cognitive effects (altered time discrimination, learning, attention level, driving performance), and brain electrical activity.	Beningus et al. 1994
Clinical, acute exposure	Carbon monoxide-poisoning	NR	40–60	Neurological impairment and pathology	Acute and delayed onset neurological impairment (headache, dizziness, drowsiness, weakness, nausea, vomiting, confusion, disorientation, irritability, visual disturbances, convulsions, and coma) and pathology (basal ganglia lesions).	Chambers et al. 2008; Dolan 1985; Ernst and Zibrak 1998; Hopkins et al. 2006; Kao and Nañagas 2006; Lo et al. 2007; Parkinson et al. 2002; Raub and Benignus 2002
Animal study, acute exposure	Rat	500–1,500		Neurosensory impairment	Potentiates noise-induced hearing loss, including noise-induced elevation of compound action potential threshold and auditory threshold shifts	Chen and Fechter 1999; Fechter et al. 1988; Young et al. 1987

**Table 3-1. Carbon Monoxide Exposures and Carboxyhemoglobin Levels Associated with Health Effects from Selected Studies Representing the Lowest Adverse Effect Levels**

Study type	Subjects	Exposure (ppm)	COHb (%)	Effect	Comments	Reference
<b>Developmental</b>						
Animal study, acute exposure	Neonatal rat (exposed post utero)	12–25	NR	Neurosensory development	Morphological changes in the developing auditory system, decreased 8 <sup>th</sup> cranial nerve action potentials and otoacoustic emissions in pups, which persisted to age 73 days	Lopez et al. 2003, 2008; Stockard-Sullivan et al. 2003; Webber et al. 2003
Animal study, acute exposure	Neonatal rats or mice (exposed <i>in utero</i> )	60–150		Neurosensory/neurobehavioral development	Decreased motor activity and response to stimulation, impaired righting reflexes, and impaired homing and memory acquisition behaviors.	De Salvia et al. 1995; Fechter and Annau 1977, 1980; Giustino et al. 1999; Mactutus and Fechter 1985; Singh 1986

COHb = blood carboxyhemoglobin; NR = not reported

## 3. HEALTH EFFECTS

**Table 3-2. Predicted Steady-State Blood Carboxyhemoglobin (COHb) Levels<sup>a</sup>**

Carbon monoxide exposure concentration (ppm)	Steady-state blood COHb (percent)
0.1	0.25
0.5	0.32
1	0.39
2	0.50
5	1.0
10	1.8
15	2.5
20	3.2
40	6.1
60	8.7
80	11
100	14
200	24
400	38
600	48
800	56
1,000	61

<sup>a</sup>Blood COHb levels are predicted from the Coburn-Forster-Kane (CFK) model (see Section 3.5.4 for a discussion of model and parameter values).

### 3. HEALTH EFFECTS

humans can be exposed to carbon monoxide from breathing and/or contact with carbon monoxide in air. No information is available on the dermal absorption or toxicity of carbon monoxide resulting from exposures to gaseous carbon monoxide. However, as is the case for other gases that are avidly absorbed from the lung (e.g., O<sub>2</sub>), during exposure to carbon monoxide in air, dermal absorption of carbon monoxide through intact skin would be expected to make a minor contribution to absorbed carbon monoxide, relative to inhalation. No information is available on the absorption or toxicity of carbon monoxide resulting from oral exposures to gaseous carbon monoxide. Although carbon monoxide can dissolve in water (23 mL carbon monoxide/L water, 20 °C), appreciable concentrations in drinking water would occur only at very high partial pressures of carbon monoxide in air, conditions in which inhalation would be the dominant absorption pathway. Therefore, the only relevant pathway of exposure to humans is the inhalation pathway, and oral and dermal exposures are not considered further.

#### 3.2 DISCUSSION OF HEALTH EFFECTS

This section of the Toxicological Profile summarizes results obtained from clinical cases of carbon monoxide poisoning, studies of controlled exposures conducted in humans, epidemiological studies of health outcomes associated with ambient air carbon monoxide concentrations, and experimental studies conducted in various animal models. Reported clinical studies are limited to exposures of acute duration. Animal studies have examined longer-duration exposures. Epidemiological studies have examined outcomes in the context of chronic exposures, or acute variations in exposure concentrations that occur during chronic exposures. Some studies (e.g., clinical studies) have reported carbon monoxide doses in terms of blood COHb levels, while most epidemiological studies and some animal studies have reported exposures concentrations (e.g., ppm, mg/m<sup>3</sup>). For comparability to human clinical studies, reported air carbon monoxide concentrations can be converted to corresponding equivalent steady-state COHb% values (i.e., the COHb% that would be achieved with continuous exposure to the reported air carbon monoxide concentration) model. This conversion has been made by application of the CFK model (see Section 3.4.5). Predicted steady-state blood COHb levels corresponding to a range of carbon monoxide exposure concentrations are presented in Table 3-2. This table can be used to convert the reported ambient air carbon monoxide concentrations into equivalent predicted blood COHb levels. Exposure concentrations are typically reported in units of ppm or mg/m<sup>3</sup>; in the Toxicological Profile, exposure units are presented in units of ppm. At standard temperature and pressure (e.g., 25 °C, 760 Torr), the conversion factor is approximately 1 (i.e., 1 mg/m<sup>3</sup> ≈ 0.87 ppm).

## 3. HEALTH EFFECTS

***Overview of Acute Carbon Monoxide Toxicity and Modes of Action.*** Carbon monoxide exerts effects on cell metabolism through both hypoxic and non-hypoxic mechanisms. Both types of effects are thought to be largely (but not entirely) the result of the ability of carbon monoxide to bind to heme and alter function and/or metabolism of heme proteins. Formation of COHb decreases the O<sub>2</sub> carrying capacity of blood and impairs release of O<sub>2</sub> from Hb for its utilization in tissues. Through similar mechanisms, carbon monoxide decreases O<sub>2</sub> storage in muscle cells by binding to and displacing O<sub>2</sub> from myoglobin. Although all tissues are vulnerable to carbon monoxide-induced hypoxic injury, those having the highest O<sub>2</sub> demand are particularly vulnerable, including the brain and heart. The developing fetus may also be a sensitive target of carbon monoxide, through hypoxic and/or non-hypoxic mechanisms (Carratu et al. 1993, 2000a, 2000b; De Salvia et al. 1995; Lopez et al. 2003, 2008; Stockard-Sullivan et al. 2003; Webber et al. 2003). Acute carbon monoxide poisoning is largely the result of tissue hypoxia. Signs and symptoms of carbon monoxide toxicity, in order of increasing severity include: (1) headache, nausea, dilation of cutaneous vasculature, vomiting, dizziness, and blurred vision; (2) confusion, syncope, chest pain, dyspnea, weakness, tachycardia, and tachynea rhabdomyolysis; and (3) palpitations, cardiac dysrhythmias, hypotension, myocardial ischemia, cardiac arrest, respiratory arrest, pulmonary edema, seizures, and coma (Kao and Nañagas 2006). Although binding of carbon monoxide to blood Hb is a primary component of the hypoxic mode of action of carbon monoxide, blood COHb levels have not been shown to be a reliable predictor of severity of acute toxicity, in part, due to time elapsed from removal from carbon monoxide exposure to COHb measurement and to the effects of emergency medical intervention (i.e., treatment with oxygen) on COHb levels prior to COHb measurement (Hampson and Hauff 2008). In general, typical levels of COHb in nonsmokers are <2% (Adams et al. 1988; Allred et al. 1991; Anderson et al. 1973; Hinderliter et al. 1989; Kleinman et al. 1989, 1998; Sheps et al. 1987, 1990). Of this, approximately 0.2–1.0% is derived from endogenous production of carbon monoxide (Coburn et al. 1963; Delivoria-Papadopoulos et al. 1974; Longo 1977). Levels ranging from 2 to 6% have been shown to exacerbate underlying cardiovascular disease, including enhanced myocardial ischemia and increased cardiac arrhythmias (Adams et al. 1988; Allred et al. 1989, 1991; Anderson et al. 1973; Kleinman et al. 1989, 1998; Leaf and Kleinman 1996b). In general, signs and symptoms of acute carbon monoxide poisoning can present at COHb levels ranging from 3 to 24% (Hampson and Hauff 2008; Kao and Nañagas 2006). This level overlaps with levels found immediately following cigarette smoking, up to approximately 10% (Kao and Nañagas 2006). More severe signs of carbon monoxide poisoning are poorly correlated with blood COHb, with loss of consciousness occurring at mean levels of 24.3% (range: 2–70; Hampson and Hauff 2008) and fatality at mean levels of 32.1% (range: 3.0–60; Hampson and Hauff 2008). Exposures that result in COHb levels >60–70% are usually fatal (Kao and Nañagas 2006). Persistent neurologic sequelae, which can be delayed in onset, can also occur, including memory loss,

## 3. HEALTH EFFECTS

impairments of concentration and language, affective disorders (e.g., depression), and Parkinsonism (Choi 2002; Klawans et al. 1982; Ringel and Klawans 1972), some of which may be related to pathologic changes in the brain (Lo et al. 2007; Gorman et al. 2003).

Most of the non-hypoxic mechanisms of action of carbon monoxide have been attributed to binding of carbon monoxide to heme in proteins other than Hb. Notable targets of carbon monoxide include components of several important physiological regulatory systems, such as brain and muscle oxygen storage and utilization (myoglobin, neuroglobin); nitric oxide cell signaling pathway (e.g., nitric oxide synthase, guanylyl cyclase); prostaglandin cell signaling pathway (cyclooxygenase, prostaglandin H synthase); energy metabolism and mitochondrial respiration (cytochrome c oxidase, cytochrome c, NADPH oxidase); steroid and drug metabolism (cytochrome P450), cellular redox balance, and reactive oxygen species (catalase, peroxidases); and various transcription factors (e.g., neuronal PAS domain protein, NPAS2). Endogenously produced carbon monoxide may participate in the physiological regulation of some, if not all, of these systems. This has potentially important implications for the understanding of carbon monoxide toxicology and dose-response relationships for the following reasons: (1) carbon monoxide modulation of physiological processes may underlie some aspects of the toxicity of exogenous carbon monoxide; (2) exogenous carbon monoxide may disrupt physiological regulation of those systems that are responsive to endogenous carbon monoxide (e.g., vascular resistance); and (3) exposures to exogenous carbon monoxide may affect carbon monoxide-mediated physiological responses at levels that approach those resulting from endogenous production. One implication of this is that the dose threshold for effects of exogenous carbon monoxide on carbon monoxide -modulated physiological systems may lie near or below ambient air carbon monoxide concentrations.

***Epidemiological Studies.*** Epidemiological studies of health outcomes associated with exposure to carbon monoxide fall into two major categories. Some studies have examined relationships between long-term average ambient air carbon monoxide concentrations and health outcomes, where the air carbon monoxide concentrations represent averages over years or decades. Other studies have examined associations between outcomes and air concentrations measured over relatively narrow time slices (e.g., 1–24-hour average or maxima). These *short-term* studies are particularly suited to outcomes that might be related to short-term elevations in carbon monoxide exposure in close temporal proximity to the outcome of interest, which might otherwise be lost in longer-term average carbon monoxide concentrations (e.g., change in heart rate, heart rate variability, arrhythmia). End points that have been assessed in epidemiological studies include mortality, morbidity, and rates of medical assistance. Examples of the latter category include rates of hospital admissions or emergency room visits recorded to

## 3. HEALTH EFFECTS

have been prompted by a given morbidity outcome, rates of activation of recorded activation of implantable cardioverter defibrillator (ICD) devices, and rates of rescue medication use by asthmatics. These studies are typically conducted using time-series analysis and Poisson regression models for estimated associations between carbon monoxide concentrations and outcome rates. Case-crossover designs have also been widely applied in air pollution epidemiology. This design offers many of the advantages of the case-control design (e.g., matching of individual cases and controls for potential co-variables and confounders), with the additional feature of each case serving as its own control, with each case assigned a *case exposure period*, usually within a few days of the outcome measurement, and a *control exposure period*, usually some time before or after the *case exposure period*. Another widely reported design in assessment of morbidity is the panel study in which members of the study cohort, usually a relatively small sample, are followed individually with respect to exposures and outcomes. Also reported are case-control studies, retrospective and cross-sectional cohort studies, and a few prospective studies of mortality.

Several important features of the epidemiological studies of ambient air carbon monoxide are of generic relevance to public health assessments. In particular, their utility for establishing dose-response relationships is limited by several factors:

- (1) Nearly all studies have relied on area monitoring for estimating exposure levels; this has the potential to produce errors in assigning exposure levels to individuals.
- (2) The temporal correspondence between monitored exposure levels and outcomes is highly uncertain. Outcomes that might be related to carbon monoxide exposure might occur in response to short-term elevations in carbon monoxide concentrations not captured in air monitoring data that are averaged over longer time periods (e.g., 24-hour average). Also, outcomes associated with a given exposure may have a latency period (e.g., myocardial infarction may occur days after the exposure that initiated the event). One approach to this problem has been to apply time-series techniques to quantify air concentration trends and to explore lag times between exposure concentration trends and the time of outcome.
- (3) Ambient air carbon monoxide concentrations tend to be strongly correlated with other air quality variables that can affect cardiovascular function, either directly or indirectly (e.g., through effects on the respiratory tract). Typical variables that correlate with air carbon monoxide concentrations include particulate matter (e.g., total suspended particulates, PM<sub>10</sub>, PM<sub>2.5</sub>), NO<sub>2</sub>, O<sub>3</sub>, and SO<sub>2</sub>. Ignoring co-variables that contribute to risk in regression models (e.g., single-pollutant models) can introduce upward bias in observed associations between air carbon monoxide concentrations and outcomes (e.g., inflate relative risk). On the other hand, assuming strongly correlated co-variables are independent in multi-pollutant models can introduce downward bias in the carbon monoxide association. Although the strength of the carbon monoxide association may be underestimated in multi-pollutant models, for the reasons noted above, persistence of significant associations between the measured outcome and air carbon monoxide concentrations, when co-pollutants are accounted for in multivariate models, provides stronger support for a contribution

### 3. HEALTH EFFECTS

of carbon monoxide to the outcome. Not all studies have explored multi-pollutant models; however, such models lend supporting evidence to the results of clinical studies that have shown carbon monoxide to exacerbate coronary vascular disease.

- (4) Outcomes investigated have tended to be serious, life-threatening events (e.g., cardiac arrhythmia, myocardial infarction, stroke, heart failure); these outcomes reflect the “late-stage” consequences of contributing pathophysiology that might be associated with exposures to lower levels of carbon monoxide for longer durations than indicated in the epidemiological studies.
- (5) Mean values of carbon monoxide exposure concentrations have ranged from 0.3 to 5 ppm with the highest values <30 ppm. These values correspond to predicted steady-state COHb levels of <1% for the mean and <5% for the highest values (predicted from the CFK model). At these COHb levels, it is possible that only highly susceptible individuals exhibit the serious hypoxia-related outcomes. For these and other reasons, epidemiologic studies have typically focused on highly susceptible populations. A typical design has been to examine associations between temporal trends in ambient air carbon monoxide concentrations and hospital admissions or emergency department visits for which the reported diagnosis was some form of cardiovascular and/or respiratory disease of impairment.

#### 3.2.1 Death

The Centers of Disease Control and Prevention (CDC) estimated that during the period 1999–2004, carbon monoxide was listed as a contributing cause of death on 16,447 death certificates in the United States, of which 2,631 deaths were categorized as unintentional and unrelated to fires (CDC 2007). Typically, fatal exposures to carbon monoxide produce coma, convulsions, and cardiorespiratory arrest (Raub et al. 2000; Wolf et al. 2008). Symptoms preceding death can include headache, dizziness, weakness, nausea, vomiting, mental confusion and visual disturbances, and loss of consciousness (Choi 2001; Raub et al. 2000). Clinical signs of life-threatening toxicity can include cardiac arrhythmia and myocardial ischemia, hypotension, pulmonary edema, and seizures (Kao and Nañagas 2006). Patients who have been resuscitated following carbon monoxide-induced cardiac arrest have a very low prognosis for survival (Hampson and Zmaeff 2001).

Although severe carbon monoxide toxicity is thought to primarily derive from hypoxia resulting from binding to and displacement of O<sub>2</sub> from heme proteins, including blood Hb, the relationship between blood COHb levels and signs indicative of life-threatening toxicity is highly uncertain (e.g., loss of consciousness, convulsions, coma, cardiopulmonary depression). In a survey of 1,407 cases of acute carbon monoxide poisoning patients (i.e., record of history of carbon monoxide exposure, signs and symptoms, blood COHb >2%, treatment with hyperbaric O<sub>2</sub>), the mean blood COHb concentration was 22.3% (95% confidence interval [CI]: 22.7–23.9, range: 2.1–72.3; Hampson and Hauff 2008). The group mean was only marginally lower than in patients who lost consciousness (24.3%, 95% CI: 23.3–

## 3. HEALTH EFFECTS

25.1, range: 2.1–73.3) or died (32.1%, 95% CI: 27.9–36.4, range: 3.0–60.0). Although survivors had lower mean COHb levels (23.1, 95% CI: 22.5–23.6) than that of fatalities, the upper end of the range exceeded that of fatalities (2.1–72.3%). The lower end of the range for COHb levels (2.1%) in this study reflects, at least in part, time delays between termination of exposure and measurement of blood COHb levels. However, that notwithstanding, this study suggests that blood COHb levels do not provide a reliable predictor of lethality. COHb levels can be converted to estimates of corresponding continuous air exposure levels by application of the CFK model. The mean for fatalities (32.1%) corresponds to a steady-state exposure (e.g., >500 minutes) to 300 ppm, or exposure to 1,000 ppm for approximately 80–90 minutes.

*Epidemiological Studies.* Epidemiological studies have examined possible associations between ambient air carbon monoxide concentrations and mortality. These studies have examined relatively low carbon monoxide concentrations (from a toxicological perspective) that are typical of ambient conditions of study period. Mean air carbon monoxide concentrations in most of the studies were <2 ppm, with maximum values <10 ppm. Epidemiologic studies of ambient carbon monoxide and mortality can be grouped into two categories: (1) studies that have examined the relationship between exposures to carbon monoxide estimated from short-term (e.g., daily or monthly average) ambient air carbon monoxide concentrations; and (2) studies that have based exposure estimates on long-term average air carbon monoxide concentrations. Although both types of studies have yielded mixed results, the larger, multi-city studies of long-term exposure have yielded estimates of carbon monoxide-associated mortality risk that are zero or less than zero (Jerrett et al. 2003; Lipfert et al. 2006b; Miller et al. 2007; Pope et al. 2002). Results from short-term exposures studies have been more varied, with some studies showing increased mortality risk when carbon monoxide is considered in single-pollutant models (percent increase in mortality: 0.2–0.6), which did not persist when the models are adjusted for co-pollutants (Burnett et al. 2004; Dominici et al. 2003a, 2003b; Samoli et al. 2007). This conclusion is supported by at least one meta-analysis of time-series published between 1985 and 2001 (Stieb et al. 2002), which estimated excess mortality risk (pooled meta estimate) for a 1.1 ppm increment in air carbon monoxide concentration to be 1.6 (95% CI: 1.1–2.1) based on a single-pollutant model and 0.7% (95% CI: -0.1–1.5), based on a multi-pollutant model that included NO<sub>2</sub>, O<sub>3</sub>, PM<sub>10</sub>, and SO<sub>2</sub> (both estimates from generalized additive model regression models). A study of survival among 1,073 carbon monoxide poisoning patients (followed from 1978 to 2005) found that survivors of carbon monoxide poisoning had an increased mortality (standardized mortality ratios [SMR]=1.9) (95% CI: 1.6–2.2; NIOSH Life Table analysis as reference; Hampson et al. 2009; Leikin and Wills 2009).

## 3. HEALTH EFFECTS

While these studies provide some evidence in support of a mortality risk associated with inhalation exposures to carbon monoxide, as previously discussed in the introduction to Section 3.2, their utility for establishing dose-response relationships for these effects are limited by several factors: (1) reliance on area monitoring for estimating exposure levels; and (2) relatively strong correlations between ambient air carbon monoxide concentrations and other air quality variables that may contribute to mortality risk. The discussion presented below focuses on outcomes of the most recent follow-ups of larger (e.g., multi-city) studies. Numerous single-city studies have also been reported; however, results of these studies are, in general, consistent with findings of multi-city studies.

Three multi-city prospective mortality studies have examined possible associations between long-term average ambient air carbon monoxide concentrations and mortality. A prospective study examined mortality in a cohort of 552,138 adults in 151 U.S. metropolitan areas, enrolled in the study in 1982 and followed through 1998 (Pope et al. 1995, 2002). A reanalysis of the study was reported in Jerrett et al. (2003). Based on estimated average ambient air carbon monoxide concentration during the period 1982–1998, increasing air carbon monoxide concentration was significantly associated with decreasing mortality from all-causes, cardiopulmonary, and lung cancer. The relative risk for cardiopulmonary death was estimated to be approximately 0.95 (95% CI: 0.9–0.99) per 1 ppm increase in carbon monoxide concentration (estimated from Figure 5 of Pope et al. 2002). A smaller prospective study focused on cardiovascular deaths among postmenopausal women between the ages of 50 and 79 years (Miller et al. 2007). Women (n=65,893) without previous history of cardiovascular disease were enrolled in the study from 36 U.S. metropolitan areas during the period 1994–1998 and deaths recorded up to mid-2003. Among the subset of the cohort that had complete data on ambient air pollutant levels from local monitoring stations (n=28,402), the hazard ratio for cardiovascular mortality was 0.92 (95% CI: 0.71–1.21) per 1 ppm increase in ambient air carbon monoxide concentration based on a single-pollutant model and 0.93 (95% CI: 0.67–1.30) based on a multi-pollutant model that adjusted for air concentrations of NO<sub>2</sub>, O<sub>3</sub>, PM<sub>2.5</sub>, PM<sub>10</sub>, and SO<sub>2</sub>. A prospective study of U.S. military veterans examined mortality outcomes in approximately 70,000 males who had been diagnosed with hypertension (Lipfert et al. 2000, 2006a, 2006b). Subjects were enrolled in the study in 1976 at the average age of 51 years and were followed through 2001 (Lipfert et al. 2006a). Mortality risk (all causes of death) was not significantly associated with ambient air carbon monoxide concentration. The relative risk was estimated to be 1.032 (95% CI: 0.954–1.117) per 1 ppm increase in air carbon monoxide concentration based on a single-pollutant model and 1.023 (95% CI: 0.939–1.115) after adjustment for NO<sub>2</sub> and O<sub>3</sub>.

### 3. HEALTH EFFECTS

A multi-city, cross-sectional study examined mortality in all U.S. counties (excluding those of Alaska) for the period 1969–1997 (Lipfert and Morris 2002). Risk estimates were made for various exposure periods ranging from 2 to 9 years and for age categories. Estimates of mortality risk (all causes) attributable to average ambient air carbon monoxide concentrations for the exposure periods were mostly either not significant or significantly negative. Elevated risk was observed in some exposure periods for the age group 15–44 years; however, this outcome was not consistently observed across time periods.

Several multi-city studies have found significant associations between increasing short-term average ambient air carbon monoxide concentration and increasing mortality risk when carbon monoxide is considered in single-pollutant models, with the associations attenuated when models are adjusted for other co-pollutants. A time-series study examined mortality in 82 U.S. cities during the period 1987–1994 (Dominici et al. 2003b; HEI 2005; Samet et al. 2000). In the Dominici et al. (2003b) analysis of these data, a 1 ppm increase in ambient air carbon monoxide concentration (lag 1 day) was associated with increased mortality (from all causes) of 0.46% (95% CI: 0.18–0.73). Mortality risk was not significant after adjustment for air concentration of PM<sub>10</sub>, alone, or PM<sub>10</sub> and NO<sub>2</sub>. A time-series study examined mortality in 12 Canadian cities during the period 1987–1994 (Burnett et al. 2004). Mortality risk (all non-accidental causes) was significantly associated with ambient air carbon monoxide concentration (lag 1 day) in single-pollutant models, with an increase of 0.68% (t=3.12) per 1.02 ppm increases in air carbon monoxide concentration. However, the association did not persist when adjusted for NO<sub>2</sub> (0.07%, t=0.30). A time-series analysis examined mortality in 19 European cities during the period 1990–1997 (Samoli et al. 2007). This study found a significant association between carbon monoxide and total non-accidental and cardiovascular mortality in single-pollutant models. The estimated effect size varied depending on the model applied. The effect size for total mortality based on a single-pollutant model ranged from 0.59% increase (95% CI: 0.41–1.79) to 1.20% (95% CI: 0.63–1.77) per 1 mg/m<sup>3</sup> increase in carbon monoxide concentration (0–1-day lag). The effect on cardiovascular mortality ranged from 0.8% increase (95% CI: 0.53–1.07) to 1.25 (95% CI: 0.30–2.21). Inclusion of black smoke or NO<sub>2</sub> in the models substantially decreased the estimated risk attributable to carbon monoxide, with the effect sizes no longer significant in some models.

#### 3.2.2 Systemic Effects

**Respiratory Effects.** Although cardiopulmonary arrest is an end point of fatal carbon monoxide poisoning, results of controlled clinical studies in healthy subjects indicate that the respiratory tract does not appear to be a primary target organ for carbon monoxide toxicity. Brief exposure to carbon monoxide

## 3. HEALTH EFFECTS

at levels >1,000 ppm may decrease ventilatory performance, although conflicting results have been reported (Chevalier et al. 1966; Fisher et al. 1969; Koike et al. 1991; Ren et al. 2001; Vesely et al. 2004). Epidemiological studies have examined possible associations between ambient air carbon monoxide concentrations and mortality. These studies have examined relatively low carbon monoxide concentrations (from a toxicological perspective) that are typical of ambient conditions of study period (mean concentrations ranging from 0.3 to 10 ppm with the highest values <30 ppm). Collectively, these studies have yielded mixed results, with some studies finding significant associations between increasing ambient air carbon monoxide concentrations and respiratory outcomes (e.g., exacerbation of asthma symptoms, hospitalizations and emergency room visits related to asthma), and few studies finding associations that persist after accounting for exposures to other air pollutants that also have been shown to contribute to respiratory disease risk (e.g., NO<sub>2</sub>, O<sub>3</sub>, particulate matter, and SO). The lack of strong evidence for associations between ambient air carbon monoxide concentrations at <30 ppm and pulmonary function is also consistent with the results of human clinical and animal studies. Studies conducted in animals provide supporting evidence that the respiratory tract does not appear to be a primary target organ for carbon monoxide (EPA 1991, 2000; Sørhaug et al. 2006).

***Clinical Studies.*** Few controlled clinical studies have evaluated adverse respiratory effects of carbon monoxide exposure. Available clinical studies have been conducted in small numbers (i.e., 4–12) of healthy subjects under acute exposure conditions (Chevalier et al. 1966; Fisher et al. 1969; Koike et al. 1991; Ren et al. 2001; Vesely et al. 2004). No controlled clinical studies evaluating respiratory effects of intermediate- or chronic-duration exposure in healthy subjects or of any duration exposure in patients with underlying respiratory diseases were identified.

Results of controlled clinical studies in healthy subjects indicate that brief exposure to carbon monoxide may decrease ventilatory performance, although conflicting results have been reported; however, based on results of available clinical studies, the respiratory tract does not appear to be a major target organ for carbon monoxide. Exposure of 10 healthy subjects to 5,000 ppm carbon monoxide for 2–4 minutes (COHb 4%) decreased inspiratory capacity by 7.5% ( $p<0.05$ ), total lung capacity decreased by 2.1% ( $p<0.02$ ), and mean resting diffusing capacity of lungs decreased by 7.6% ( $p<0.05$ ) (Chevalier et al. 1966). However, other studies have found no effects of acute carbon monoxide exposure on ventilatory function. No changes in ventilatory mechanics (vital capacity, functional residual capacity, airway conductance, lung volume, pulmonary resistance, dynamic lung compliance, diffusing capacity) were observed in four healthy subjects exposed to carbon monoxide producing blood COHb levels of 11–20% (Fisher et al. 1969). Exposure to 1,000 ppm carbon monoxide for 10–30 minutes (COHb ~10%;

## 3. HEALTH EFFECTS

10 subjects) (Ren et al. 2001) or 1,200 ppm carbon monoxide for 30–45 minutes (COHb 10.2%; 10 subjects) (Vesely et al. 2004) did not affect resting ventilation. During submaximal exercise challenge, acute exposure of 10 subjects to carbon monoxide producing blood COHb levels of 11–20% did not alter ventilation mechanics (Koike et al. 1991). The reason for these conflicting results is not apparent, although the small number of subjects evaluated in each study may have been a contributing factor.

***Epidemiological Studies.*** Epidemiological studies of exposure to carbon monoxide and respiratory outcomes fall into two major categories: (1) studies that have assessed respiratory morbidity in association with ambient air carbon monoxide concentrations; and (2) studies that have evaluated associations between ambient air carbon monoxide concentrations and the incidence of hospital admissions and/or emergency room visits related to respiratory disease (e.g., time-series studies, case-crossover). Collectively, these studies have yielded mixed results. Although some studies have found significant associations between increasing ambient air carbon monoxide concentrations and respiratory outcomes (e.g., exacerbation of asthma symptoms, hospitalizations and emergency room visits related to asthma), few studies have examined the robustness of the association in models that adjust for exposures to other air pollutants that also have been shown to contribute to respiratory disease risk (e.g., NO<sub>2</sub>, O<sub>3</sub>, particulate matter, and SO<sub>2</sub>). While these studies provide some evidence in support of adverse respiratory effects of inhalation exposures to carbon monoxide, as previously discussed in the introduction to Section 3.2, their utility for establishing dose-response relationships for these effects are limited by several factors: (1) reliance on area monitoring for estimating exposure levels; (2) uncertainty in knowledge of temporal correspondence between monitored exposure levels and outcomes; (3) relatively strong correlations between ambient air carbon monoxide concentrations and other air quality variables that can affect respiratory function; and (4) relatively low carbon monoxide exposures studied. Mean values of carbon monoxide exposure concentrations have ranged from 0.3 to 10 ppm with the highest values <30 ppm. These concentrations are well below those explored in experimental studies that found minimal or no effects of carbon monoxide on pulmonary function in association with acute exposures to carbon monoxide (i.e., ≥1,000 ppm). The relatively low carbon monoxide concentrations studied in epidemiological studies, together with concurrent exposures to other air pollutants that may have more pronounced effects on respiratory function, may have contributed to the mixed results of these studies and relatively weak associations with carbon monoxide that have been observed (e.g., odds ratios and relative risks very close to 1). The lack of strong evidence for associations between ambient air carbon monoxide concentrations at <30 ppm and pulmonary function is also consistent with the results of human clinical and animal studies.

## 3. HEALTH EFFECTS

The presentation of epidemiology is organized by major categories of respiratory outcomes for which the studies were designed to evaluate, which include pulmonary function, asthma and exacerbation of asthma symptoms, and hospital admissions and emergency room visits related to respiratory disease. Study conclusions are presented in the text, with selected supporting details presented in tabular form (Table 3-3). Although most studies explored various time lags between monitored air carbon monoxide concentrations and outcomes, as well as various sample strata, for the sake of brevity, only selected representative time lags (usually, those indicative of the strongest associations to carbon monoxide) are presented in the tables. Where co-pollutant models have been explored, these results are also presented.

***Studies of Pulmonary Function.*** Studies of possible associations between inhalation exposures to carbon monoxide and effects on pulmonary function have yielded mixed results (Chen et al. 1999; Fischer et al. 2002; Lagorio et al. 2006; Penttinen et al. 2001; Rabinovitch et al. 2004; Silkoff et al. 2005; Timonen et al. 2002). Several of these studies examined subjects who had ongoing lung disease (e.g., asthma, chronic obstructive lung disease) and who might be more sensitive to agents that affect pulmonary function (Rabinovitch et al. 2004; Silkoff et al. 2005; Timonen et al. 2002). Mean ambient air carbon monoxide concentrations studied ranged from 0.5 to 10 ppm, with the highest values <30 ppm (Lagorio et al. 2006). Only one of these studies explored multi-pollutant models and found that the association with carbon monoxide was substantially weakened when adjustments were made for air particle matter (Penttinen et al. 2001). Therefore, is it difficult to sort out the effects of carbon monoxide from those of other urban air pollutants that have strong correlations with air carbon monoxide concentrations and that also could have affected pulmonary function.

Studies of children have yielded mixed results that are further complicated by lack of assessment of models in which correlated concentrations of other air pollutants were included (Chen et al. 1999; Fischer et al. 2002; Rabinovitch et al. 2004; Timonen et al. 2002). A study conducted in the San Joaquin Valley, California, examined lung function in a cohort of 232 asthmatic children (age 6–11 years) found that long-term average exposures to carbon monoxide in infancy and childhood (age 0–6 years) was associated with significant decreasing forced expiratory volume in 1 second (FEV<sub>1</sub>)/forced vital capacity (FVC) and forced midexpiratory flow rate (FEF<sub>25–75</sub>)/FVC (Mortimer et al. 2008). Mean air carbon monoxide concentrations during the study period ranged from approximately 1–1.5 ppm, with the highest value <6 ppm. The effect sizes (per 1 ppm increase in air carbon monoxide concentration) were of approximately 2.5% decrease in FEV<sub>1</sub>/FVC and 4.8% decrease in FEF<sub>25–75</sub>/FVC. A panel study conducted in Denver, Colorado (1999–2002) examined lung function in asthmatic children (n=41–63, age

**Table 3-3. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Respiratory Disease**

Study	Design features	CO exposure	Effect size
<b>Lung function</b>			
Chen et al. (1999) Period: 1995–1996 Location: Taiwan, China	Outcome: Lung function (FVC, FEV <sub>1</sub> , FEV <sub>1</sub> /FVC, FEF <sub>25–75%</sub> , PEF) Design: Cross-sectional Sample: n=941, age 8–13 years	Avg time: 1 hour max Range: 0.4–3.6 ppm	Increment: NR β Coefficient (SE), lag 2 days for 24 hours avg CO: FVC (mL) -147.71 (64.48) FEV <sub>1</sub> (mL) -82.42 (60.95)
Fischer et al. (2002) Period: NR Location: Utrecht, Netherlands	Outcome: Lung function (FVC, FEV <sub>1</sub> , PEF, MMEF) Design: Panel study Sample: n=68, age 10–11 years	Avg time: 24 hours Mean: 0.80 ppm Range: 0.28–1,34 ppm	Increment: 100 μg/m <sup>3</sup> NO% increase (95% CI): 28.2 (6.9–53.9) No significant associations with lung function endpoints
Lagorio et al. (2006) Period: 1999 Location: Rome, Italy	Outcome: Lung function (FVC, FEV <sub>1</sub> ) Design: Time-series panel study Sample: Patients (age 50–80 years) diagnosed with COPD n=11; asthma, n=11; IHD, n=7	Avg time: 24 hours Mean: 6.5 ppm Range: 1.4–25.2 ppm	Increment: 1 mg/m <sup>3</sup> No significant associations with lung function endpoints
Park et al. (2005a) Period: 2002 Location: Incheon, Korea	Outcome: Lung function (PEF variability (>20%), mean PEF); Design: Panel Sample: n=64 bronchial asthmatics, age 16–75 years	Avg time: 24 hours Mean: 0.6 ppm SD range: 0.09–0.15 ppm	Increment: NR β Coefficient (SE): Mean PEF (L/min): -10.103 (2.7146)
Penttinen et al. (2001) Period: 1996–1997 Location: Helsinki, Finland	Outcome: Lung function (PEF) Design: Panel Sample: n=57 adults asthmatics	Avg time: 24 hours Median: 0.35 ppm Range: 0.09–0.96 ppm	Increment: 0.2 mg/m <sup>3</sup> β Coefficient (SE); lag 1 day: PEF Deviations (L/minute) CO: -1.08 (0.36) CO+PNC: -0.67 (0.64); 1
Mortimer et al. (2008) Period: 1989–2000 Location: San Joaquin Valley, California	Outcome: Lung function (FVC, FEV <sub>1</sub> , PEF, FEF <sub>25–75</sub> , FEV <sub>1</sub> /FVC, FEF <sub>25–75</sub> /FVC, FEF <sub>25</sub> , FEF <sub>75</sub> ) Design: Cohort Sample: n=232 asthmatic children (age 6–11 years)	Avg time: 8 hour max Mean: 1–1.5 ppm Range: max ~6 ppm	Increment: ~1 ppm β Coefficient (SE); FEV <sub>1</sub> /FVC: -0.0073 (0.0016) FEF <sub>25–75</sub> /FVC: -0.2179 (0.0446) % Decrease: FEV <sub>1</sub> /FVC: 2.5% FEF <sub>25–75</sub> /FVC: 4.8%

**Table 3-3. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Respiratory Disease**

Study	Design features	CO exposure	Effect size
Rabinovitch et al. (2004) Period: 1999–2002 Location: Denver, Colorado	Outcome: Lung function (FEV <sub>1</sub> , PEF) Design: Panel study Sample: n=41–63 asthmatics, age 6–12 years	Avg time: 24 hours Mean 1.0 ppm Range: 0.3, 3.5 ppm	Increment: 0.4 ppm No significant associations with lung function endpoints
Silkoff et al. (2005) Period: 1999–2001 Location: Denver, Colorado	Outcome: Lung function (FEV <sub>1</sub> , PEF) Design: Panel Sample: n=16–18, former smokers, age ≥40 years	Pollutant: CO Avg Time: 24 hours Mean: 1 ppm Range: 0.3–3.8 ppm	Increment: 1 ppm Significant association with decreasing FEV <sub>1</sub> in second winter season of study, but not in first (β coefficients reported graphically)
Timonen et al. (2002) Period: 1994 Location: Kuopio, Finland	Outcome: Lung function (FVC, FEV <sub>1</sub> , MMEF, AEFV) Design: Panel Sample: n=33 with respiratory symptoms, age 7–12	Avg time: 24 hours Mean: 0.5 ppm Range: 0.09–2.4 ppm	Increment: 0.32 mg/m <sup>3</sup> β Coefficient (SE), lag 2 days: Baseline FEV <sub>1</sub> (mL): 11.7 (SE: 5.77, p<0.05) Exercise ΔFEV <sub>1</sub> (%): 0.087 (0.26)
Asthma and exacerbation of asthma symptoms			
Hwang et al. (2005) Period: 2001 Location: Taiwan, China	Outcome: Asthma Design: Cross-sectional Sample: n=32,672, age 6–15 years	Avg time: year. Mean: 0.66 ppm Range: 0.42–0.96 ppm	Increment: 0.1 ppm Adjusted OR (95% CI) CO: 1.045 (1.017–1.074) CO+SO <sub>2</sub> : 1.066 (1.034–1.099) CO+PM <sub>10</sub> : 1.079 (1.047–1.112) CO+O <sub>3</sub> : 1.063 (1.1–1.474) CO+SO <sub>2</sub> +O <sub>3</sub> : 1.111 (1.074–1.15) CO+PM <sub>10</sub> + O <sub>3</sub> : 1.119 (1.084–1.155)
Park et al. (2005a) Period: 2002 Location: Incheon, Korea	Outcome: Asthma symptoms Design: Panel Sample: n=64 bronchial asthmatics, age 16–75 years	Avg time: 24 hours Mean: 0.6 ppm SD range: 0.09–0.15 ppm	Increment: NR No significant association with asthma symptoms
Rabinovitch et al. (2004) Period: 1999–2002 Location: Denver, Colorado	Outcome: Asthma symptoms, rescue inhaler use Design: Panel study Sample: n=41–63 asthmatics, age 6–12 years	Avg time: 24 hours Mean 1.0 ppm Range: 0.3, 3.5 ppm	Increment: 0.4 ppm β Coefficient (SE), 3-day moving avg: Asthma exacerbation: 1.012 (0.913–1.123) Rescue inhaler: 1.065 (1.001–1.133)

**Table 3-3. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Respiratory Disease**

Study	Design features	CO exposure	Effect size
Rodriguez et al. (2007) Period: 1996–2003 Location: Perth, Australia	Outcome: Respiratory symptoms Design: Panel Population: n=263, age 0–5 years	Avg time: 8 hours Mean 1.4 ppm Range: 0.01–8.0 ppm	Increment: NR OR (95%CI), wheeze/chest rattle, lag 5-days: lag 0 days: 1.089 (0.968–1.226) lag 5 days: 1.136 (1.016–1.26) lag 0–5 days: 1.035 (1.005–1.066)
Schildcrout et al. (2006) Period: 1993–1995 Location: 8 North American cities	Outcome: Asthma symptoms; rescue inhaler use Design: Panel Sample: 990 asthmatics, age 5–12 years	Avg time: 24 hours 10th–90th%: 0.4–2.4 ppm	Increment: 1.0 ppm OR (95% CI), asthma symptoms, lag 0 days: CO: 1.08 (1.01–1.14) CO+NO <sub>2</sub> 1.07 (1–1.14) CO+PM <sub>10</sub> : 1.08 (1.01–1.15) CO+SO <sub>2</sub> 1.07 (0.99–1.16)
Silkoff et al. (2005) Period: 1999–2001 Location: Denver, Colorado	Outcome: Asthma symptoms, rescue inhaler use Design: Panel Sample: n=16–18, former smokers, age ≥40 years	Avg time: 24 hours Mean: 1 ppm Range: 0.3–3.8 ppm	Increment: 1 ppm No significant associations with asthma symptoms or rescue medications
Slaughter et al. (2003) Period: 1993–1995 Location: Seattle, Washington	Outcome: Asthma severity; medication use Design: Panel Sample: n=133 asthmatics, age 5–13 years	Avg time: 24 hours Median: 1.47 ppm 25 <sup>th</sup> –75 <sup>th</sup> %: 0.23–1.87 ppm	Increment: 0.67 ppm OR (95% CI), asthma severity, lag 1 day: Asthma severity: 1.21 (1.08–1.35) Rescue inhaler use: 1.09 (1.03–1.16)
von Klot et al. (2002) Period 1996–1997 Location: Erfurt, Germany	Outcome: Asthma symptoms; medication use Design: Panel Sample: 53 asthmatics, age 37–77 years	Avg time: 24 hours Mean: 0.8 ppm Range: 0.3–2.3 ppm	Increment: 0.6 mg/m <sup>3</sup> OR (95% CI); wheezing prevalence, 5-day avg: CO: 0 1.13 (1.05–1.22) CO+MC <sub>0.01–2.5</sub> : 1.15 (1.04–1.27) CO+NC <sub>0.01–0.1</sub> : 1.09 (0.98–1.22)
Yu et al. (2000) Period: 1993–1995 Location: Seattle, Washington	Outcome: Asthma symptoms Design: Panel Sample: n=133 asthmatics, age 5–13 years	Avg time: 24 hours Mean 1.6 ppm Range: 0.65–4.18 ppm	Increment: 1.0 ppm OR (95% CI), lag=2 days: 1.26 (1.09–1.46)

**Table 3-3. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Respiratory Disease**

Study	Design features	CO exposure	Effect size
Hospital admissions and emergency room visits related to respiratory disease			
Burnett et al. (2001) Period: 1980–1994 Location: Toronto, Canada	Outcome: Asthma, acute bronchitis/bronchiolitis; croup; pneumonia HAs Design: Time series Sample: <2 years	Avg time: 1 hour Mean: 1.18 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 1.3–2.3 ppm	Increment: 1.9 ppm % Increase (t value), lag 0–1 day: CO: 19.20% (3.48) CO+O <sub>3</sub> : 14.30% (2.6)
Cakmak et al. (2006) Period: 1993–2000 Location: 10 Canadian cities	Outcome: Respiratory disease HAs Design: Time-series Sample: All ages	Avg time: 24 hours Mean: 0.8 ppm Range: 0.0–6.5 ppm	Increment: 0.8 ppm % Increase(95% CI), lag 2.8 days: CO: 0.60% (0.20, 1) CO+SO <sub>2</sub> +NO <sub>2</sub> +O <sub>3</sub> : -0.20% (-0.70–0.30)
Karr et al. (2006) Period: 1995–2000 Location: Southern California	Outcome: Acute bronchiolitis HAs Design: Case-crossover Sample: age ≤1 year	Avg time: 24 hours Mean: 17 ppm Range: 0.004–9.6 ppm	Increment: 1.36 ppm OR (95% CI), lag 1 day: All: 0.99 (0.96–1.02) Age 25–29 weeks: 0.86 (0.68–1.1)
Linn et al. (2000) Period 1992–1995 Location: Los Angeles, California	Outcome: Asthma, COPD, pulmonary HAs Design: Time-series Sample: age 0–29, ≥30 years	Avg time: 24 hours Mean: 1.0–2.1 ppm Range: 0.3–5.3 ppm	Increment: 1.0 ppm β (SE): Asthma, <30 years: 0.036 (0.016, p<0.05) Asthma ≥30 years: 0.028 (0.010, p<0.05) COPD, ≥30 years: 0.019 (0.007, p<0.05)
Lin et al. (2003) Period: 1981–1993 Location: Toronto, Canada	Outcome: Asthma HAs Design: Case-crossover Sample: age 6–12 years	Avg time: 24 hours Mean: 1.18 ppm Range: 0–6.1 ppm	Increment: 0.5 ppm OR (95% CI), lag 1 day: Males CO: 1.05 (1–1.11) Males CO+PM: 1.05 (0.99–1.11) Females CO: 1.00 (0.93–1.06) Females CO+PM: 1.00 (0.93–1.07)
Lin et al. (2005) Period: 1998–2001 Location: Toronto, Canada	Outcome: Respiratory infection HAs Design: Case-crossover Sample: age <15 years	Avg time: 24 hours Mean: 1.16 ppm Range: 0.38–2.45 ppm	Increment: 0.44 ppm OR (95% CI), lag 0–5 days: Males, CO: 1.13 (1.02–1.25) Males, CO+PM: 1.08 (0.97–1.20) Females, CO: 1.05 (0.93–1.18) Females, CO+PM: 1.02 (0.90–1.15)

**Table 3-3. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Respiratory Disease**

Study	Design features	CO exposure	Effect size
Peel et al. (2005) Period: 1993–2000 Location: Atlanta, Georgia	Outcome: Asthma; COPD, pneumonia EDVs Design: Time-series Sample: ages all, 2–18 years	Avg time: 1 hour maximum Mean: 1.8 ppm 10 <sup>th</sup> –90 <sup>th</sup> ‰: 0.5–3.4 ppm	Increment: 1.0 ppm RR (95%CI); lag 0–13 days: All respiratory: 1.066 (1.045–1.087) Asthma: 1.076 (1.047–1.105) Pneumonia: 1.045 (1.011–1.080) COPD: 1.032 (0.975–1.092), 1.026 (1.004–1.048; lag=0–2 days) No significant associations after adjustment for PM <sub>10</sub> , NO <sub>2</sub> , O <sub>3</sub>
Slaughter et al. (2005) Period: 1995–2001 Location: Spokane, Washington	Outcome: Respiratory, asthma; COPD, respiratory infection HAs and EDVs Design: Time-series Sample: All ages, adults	Avg time: 24 hours 5 <sup>th</sup> –95 <sup>th</sup> ‰: 1.25–3.05 ppm	Increment: 1.0 ppm RR (95% CI); lag 3 days: Respiratory EDV: 1.03 (1.00–1.06) Asthma EDV: 1.06 (1.00–1.11) COPD EDV, Adults: 1.01 (0.93–1.10) Respiratory HA: 0.99 (0.96–1.03) Asthma HA: 1.00 (0.91–1.11) COPD HA.: Adults: 0.97 (0.88–1.06)
Tolbert et al. (2007) Period: 1993–2004 Location: Atlanta, Georgia	Outcome: Asthma; COPD; URI, pneumonia; bronchiolitis EDVs Design: Time-series Sample: All ages	Avg time: 1 hour maximum Mean: 1.6 ppm Range: 0.1–7.7 ppm	Increment: 1.22 ppm RR (95% CI), lag 3 days: Respiratory diseases: 1.016 (1.009–1.022) Not significant after adjustment for O <sub>3</sub> , NO <sub>2</sub> , or PM <sub>10</sub> .
Yang et al. (2003) Period: 1986–1998 Location: Vancouver, Canada	Outcome: Respiratory disease HA) Design: Case-crossover Sample: ages <3; ≥65 years	Avg time: 24 hours Mean: 0.98 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.62–1.16 ppm	Increment: 0.54 ppm OR (95% CI), lag 1 day: <3 years, CO: 1.04 (1.01–1.07) <3 years, CO+O3: 1.04 (1.01–1.07) <3 years, CO+O3+NO2+ SO2: 1.02 (0.96–1.08) ≥65 years, CO: 1.02 (1.00–1.04) ≥65 years, CO+O3: 1.02 (1.00–1.04) ≥65 years, CO, O3, NO2, SO2: 0.96 (0.93–1.00)

**Table 3-3. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Respiratory Disease**

Study	Design features	CO exposure	Effect size
Zanobetti and Schwartz (2006) Period: 1995–1999 Location: Boston, Massachusetts	Outcome: Pneumonia HAs Design: Case-crossover Sample: All ages	Avg time: 24 hours 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.39–0.60 ppm	Increment: 0.475 ppm % Increase (95% CI): 5.45 (1.10, 9.51; lag=0 days) 5.12 (0.83, 9.16; lag=0–1 days)

AEFV = area under the expiratory flow-volume curve; Avg = average; CI = confidence interval; CO = carbon monoxide; COPD = chronic obstructive pulmonary disease; EDV = emergency department visit; FEF = forced expiratory flow; FEF25–75 = forced expiratory flow at 25–75% of vital capacity; FEV1 = forced expiratory volume at 1 second; FVC = forced vital capacity; HA = hospital admission, IHD = ischemic heart disease; MC = particle mass concentration; NC = particle number concentration; NR = not reported; PEF = peak expiratory flow, PNC = particle number concentration/count; RR = relative risk; SD = standard deviation; SE = standard error; URI = upper respiratory infection

## 3. HEALTH EFFECTS

6–12 years) and did not find a significant association between air carbon monoxide concentration and pulmonary function end points, FEV<sub>1</sub>, and peak expiratory flow (PEF) (Rabinovitch et al. 2004). The mean air carbon monoxide concentration in the study was 1.0 ppm (range: 0.3–3.5 ppm). A similar panel study conducted in Kuopio, Finland examined lung function in children (n=33, age 7–12 years) who were diagnosed with chronic respiratory disease symptoms (Timonen et al. 2002). Increasing air carbon monoxide concentration was associated with decreasing FVC and FEV<sub>1</sub>, when measured at rest, but not during exercise. Other air pollutants were also associated with decrements in baseline pulmonary function (particulate matter, NO<sub>2</sub>); however, multi-pollutant models were not explored. The air carbon monoxide concentrations in this study were similar to that in the Rabinovitch et al. (2004) study (mean 0.6 mg/m<sup>3</sup>, range: 0.1–2.8 mg/m<sup>3</sup>). A cross-sectional cohort study conducted in Taiwan, China (1995–1996) included asthmatic children (n=941, age 8–13 years; Chen et al. 1999). A significant association was found between increasing air carbon monoxide concentration and decreasing FVC; however, multi-pollutant models were not explored. Peak daily air carbon monoxide concentrations ranged from 0.4 to 3.6 ppm. Fischer et al. (2002) examined possible associations between carbon monoxide and pulmonary function in a panel of children (n=68, age 10–11 years) in The Netherlands. Increasing air carbon monoxide concentration was not associated with FVC, FEV<sub>1</sub>, PEF, or maximal midexpiratory flow (MMEF); however, it was associated with increasing levels of expired endogenous nitric oxide radical (NO•). Here again, multi-pollutant models were not explored. The mean air carbon monoxide concentration in this study was 0.9 mg/m<sup>3</sup> (range: 0.3–1.5 mg/m<sup>3</sup>).

Results of studies of pulmonary function in adults have also been mixed and conflicting (Lagorio et al. 2006; Penttinen et al. 2001; Silkoff et al. 2005). A panel study of adult nonsmoking asthmatics (n=57, mean age 53 years) conducted in Helsinki, Finland (1996–1997) found that increasing air carbon monoxide concentration was associated with decreasing PEF; however, the association was not significant when adjusted for particulate matter (Penttinen et al. 2001). The mean air carbon monoxide concentration was 0.4 mg/m<sup>3</sup> (range: 0.1–1.1 mg/m<sup>3</sup>). A panel study of adult former smokers, diagnosed with chronic obstructive pulmonary disease (COPD), conducted in Denver, Colorado (1999–2001, n=16–18, age ≥40 years) found somewhat conflicting results. A significant association between increasing air carbon monoxide concentration and decreasing FEV<sub>1</sub> was found in the second winter season of the study, but not in the first winter season of the study (Silkoff et al. 2005). Air carbon monoxide concentrations (ppm) were not significantly different in the two seasons (mean: 1.1–1.2; ranges: 0.3–3.8 and 0.4–2.8), although concentrations of other air pollutants (e.g., PM<sub>10</sub>, PM<sub>2.5</sub>, and NO<sub>2</sub>) were higher in the second winter season, suggesting that other air pollutants may have contributed to the associations attributed to carbon monoxide in single-pollutant models. The study conducted by Lagorio et al. (2006) is notable

## 3. HEALTH EFFECTS

because it included relatively high air carbon monoxide concentrations (mean range: 1.8 [spring]–10.6 [winter] ppm; range: 1.5–25.1 ppm). This panel study examined adults (n=29, age 18–80 years) in Rome, Italy (1999) who were diagnosed with asthma, COPD, or ischemic heart disease. Increasing air carbon monoxide concentration was not significantly associated with lung function parameters, and multi-pollutant models were not explored. A panel study conducted in Incheon, Korea examined lung function in asthmatics (n=57, ages 16–75 years) found a significant association between air carbon monoxide concentrations and decreasing PEF; however, multi-pollutant models were not explored (Park et al. 2005a). The mean air carbon monoxide concentration was 0.6 ppm.

*Studies of Asthma Incidence (or Prevalence) and Exacerbation of Asthma Symptoms.* A cross-sectional study examined asthma diagnoses among school-age children (n=32,672, age 6–15 years) in Taiwan, China (Hwang et al. 2005). The odds ratio for 0.1 ppm increases in air carbon monoxide concentration (annual average) was estimated to be 1.045 (95% CI: 1.017–1.074) and persisted after adjustment for O<sub>3</sub>, PM<sub>10</sub>, and/or SO<sub>2</sub> in multi-pollutant models. The mean air carbon monoxide concentration was 0.6 ppm (range: 0.4–0.9 ppm). Several studies have examined possible associations between ambient air carbon monoxide concentrations and asthma symptoms (e.g., coughing, wheezing, chest tightness, shortness of breath, inhaler use). Results of these studies have been mixed; however, collectively, these studies provide evidence for associations between increasing air carbon monoxide concentration and increasing severity of asthma (Park et al. 2005a; Rabinovitch et al. 2004; Rodriguez et al. 2007; Schildcrout et al. 2006; Silkoff et al. 2005; Slaughter et al. 2003; von Klot et al. 2002; Yu et al. 2000). One of the larger studies examined asthma symptoms in a panel of 990 children (ages 5–12 years) in eight North American cities (1993–1995, Schildcrout et al. 2006). A 1 ppm increase in air carbon monoxide concentration (lag=0 days) was associated with an odds ratio of 1.08 (95% CI: 1.01–1.14) for asthma symptoms. The association remained significant after adjusting for NO<sub>2</sub>, PM<sub>10</sub>, or SO<sub>2</sub> in two-pollutant models. Air carbon monoxide concentrations ranged from approximately 0.4 to 2.4 ppm (10<sup>th</sup>–90<sup>th</sup> percentile range). Significant associations with increasing carbon monoxide concentration were also found when rescue medication use was evaluated as the asthma symptom end point.

The above results are consistent with those of several smaller panel studies. A study conducted in Seattle, Washington examined asthma symptoms (Yu et al. 2000) and asthma severity (Slaughter et al. 2003) in a panel of 133 asthmatic children (age 5–13 years). The mean air carbon monoxide concentration was 1.6 ppm (range: 0.6–4.8 ppm; Yu et al. 2000). Both studies found significant associations between ambient air carbon monoxide and asthma symptoms. In the Yu et al. (2000) study, a 1 ppm increase in air carbon monoxide (lag=2 days) was associated with an odds ratio of 1.26 (95% CI: 1.09–1.46) for asthma

## 3. HEALTH EFFECTS

symptoms. In the Slaughter et al. (2003), a 0.67 ppm increase in air carbon monoxide concentration was associated with an odds ratio of 1.21 (95% CI: 1.08–1.35) for an increase in severity of asthma attack. A significant association with increasing carbon monoxide concentration was also found when rescue medication use was evaluated as the asthma severity end point (Slaughter et al. 2003). Rabinovitch et al. (2004) also found a significant association between increasing air carbon monoxide concentrations and increasing rescue medication use in a study of children in Denver, Colorado (1999–2002, n=41–63, age 6–12 years). A panel study of 263 children (age 0–5 years) in Perth, Australia, who were not diagnosed with asthma, also found a significant association between air carbon monoxide concentration and wheezing/chest rattle (Rodriguez et al. 2007).

Results of studies conducted in older asthmatic subjects have been mixed (Park et al. 2005a; Silkoff et al. 2005; von Klot et al. 2002). In a panel of 53 asthmatic adults (age 37–77 years) in Erfurt, Germany (1996–1997), a 0.6 mg/m<sup>3</sup> increase in air carbon monoxide concentration (5-day average) was associated with an odds ratio of 1.13 (95% CI: 1.05–1.22) for prevalence of wheezing; however, the association was attenuated when adjusted for ultrafine particle number concentration in air. In two other panel studies of asthmatics, one conducted in Incheon, Korea (2002, n=54, age 16–76 years; Park et al. 2005a) and one in Denver, Colorado (n=16 or 18, age ≥40 years, Silkoff et al. 2005) did not find significant associations between air carbon monoxide concentrations and asthma symptoms.

***Studies of Hospital Admissions and Emergency Room Visits Related to Respiratory Symptoms.*** A large number of studies have examined associations between ambient carbon monoxide concentrations and rates of hospital admissions (e.g., Burnett et al. 2001; Cakmak et al. 2006; Karr et al. 2006; Lin et al. 2003, 2005; Linn et al. 2000; Slaughter et al. 2005; Yang et al. 2003; Zanobetti and Schwartz 2006) or emergency room visits related to respiratory disease (e.g., Peel et al. 2005; Slaughter et al. 2005; Tolbert et al. 2007). Collectively, these studies provide evidence for associations between increasing carbon monoxide concentrations and increasing risk of respiratory symptoms that trigger the need for medical assistance. However, only a few studies have examined associations in multi-pollution models; therefore, relative contributions of other air pollutants to the reported outcomes are highly uncertain.

In studies that did evaluate multi-pollutant models, the associations between carbon monoxide concentrations and outcomes persisted in some studies (Burnett et al. 2001; Lin et al. 2003, Yang et al. 2003), but not in others (Lin et al. 2005; Peel et al. 2005; Tolbert et al. 2007). In a time-series analysis conducted in Toronto, Canada (1980–1994), Burnett et al. (2001) found a significant association between increasing air carbon monoxide concentrations and hospital admissions for asthma, acute bronchitis/

## 3. HEALTH EFFECTS

bronchiolitis, croup, and pneumonia in children (age <2 years). A 1.9 ppm increase in air carbon monoxide concentration (lag: 0–1 day) was associated with a 19.2% increase in hospital admissions, which remained elevated when adjusted for air O<sub>3</sub> concentration. A case-crossover study of hospital admissions of children (ages 6–12 years) for asthma in Toronto, Canada (1981–1993) found that a 0.5 ppm increase in air carbon monoxide concentration was associated with an odds ratio (lag: 1 day) of 1.05 (95% CI: 1.00–1.11) in males but not in females (odds ratio: 1.00, 95% CI: 0.93–1.06), and persisted in males after adjustment for particulate matter (Lin et al. 2003). Another case-crossover study directed at respiratory infections in children (age <15 years, males), also conducted in Toronto, Canada (1988–2001) estimated the odds ratio for a 0.44 ppm increase in air carbon monoxide concentration (lag: 0–5 days) to be 1.13 (95% CI: 1.01–1.25) and 1.08 (95% CI: 0.97–1.20) after adjustment for PM<sub>10</sub> and PM<sub>10-2.5</sub> (Lin et al. 2005). Odds ratios for females were not significant. Yang et al. (2003) conducted a case-crossover study of hospital admissions for various respiratory disease categories in Vancouver, Canada (1986–1998) and found that an increase in carbon monoxide concentration of 0.5 ppm was associated with odds ratios of 1.04 (95% CI: 1.01–1.07) for children (age <3 years) and 1.02 (95% CI: 1.00–1.04) for elderly adults (age ≥65 years). The associations persisted when adjusted for air O<sub>3</sub> concentration. In a time-series analysis of >400,000 emergency room visits for respiratory disease in Atlanta, Georgia (1993–2000), the estimated risk ratio for a 1 ppm increase in air carbon monoxide concentration (lag: 0–13 days) was 1.076 (95% CI: 1.047–1.105) for all ages and 1.019 (1.004–1.035) for asthma visits in children aged 2–18 years (Peel et al. 2005). The associations were not significant when adjusted for other air pollutants in two-pollutant models (e.g., PM<sub>10</sub>, NO<sub>2</sub>, or O<sub>3</sub>). This study also found a significant risk ratio for emergency room visits related to pneumonia (risk ratio: 1.045, 95% CI: 1.011–1.080; lag: 0–13 days). A revisit to the Atlanta, Georgia study, with the study period expended to 1993–2004 (Tolbert et al. 2007), found essentially similar results as reported in Peel et al. (2005). The risk ratio for all emergency room visits for all respiratory disease categories was 1.016 (95% CI: 1.009–1.022) per 1.22 ppm increase in 1-hour max carbon monoxide concentration (lag: 3 days); however, the association was attenuated in models that adjusted for O<sub>3</sub>, NO<sub>2</sub>, or PM<sub>10</sub>.

Studies that examined COPD admissions and emergency room visits yielded mixed results (Peel et al. 2005; Slaughter et al. 2005; Yang et al. 2003). In the Yang et al. (2003) time-series study, conducted in Vancouver, Canada (1994–1998), the risk ratio for a 0.5 ppm increase in air carbon monoxide concentration (lag: 0–6 days) was 1.14 (95% CI: 1.03–1.23) for COPD hospital admissions. On the other hand, significant associations between air carbon monoxide concentration and COPD hospital admissions or emergency room visits were not found in a time-series study conducted in Spokane, Washington (1995–2001; Slaughter et al. 2005). In the Peel et al. (2005) time-series study conducted in

## 3. HEALTH EFFECTS

Atlanta, Georgia (1993–2000), relative risk for emergency room visits for COPD was 1.03 (95% CI: 1.00–1.05) per 1 ppm increase in carbon monoxide concentration days, and 1.03 (95% CI: 0.98–1.09) for lag: 0–2 days, and 1.03 (95% CI: 0.98–1.09) for lag: 0–13 days. The air carbon monoxide concentrations in these studies were similar (10<sup>th</sup>–90<sup>th</sup> percentile range: approximately 1.25–4.3 ppm)

***Animal Studies.*** Most studies evaluating adverse respiratory effects of carbon monoxide in animals were conducted using carbon monoxide exposures that produced much higher COHb concentrations (i.e., COHb >50%) than those evaluated in controlled clinical studies in humans. Even at these high exposures, results of animal studies support those of controlled clinical studies indicating that the respiratory tract is not a major target organ for carbon monoxide. Animal studies published prior to 2000 have been reviewed in detail by the EPA (1991, 2000); therefore, these reviews largely form the basis of the following discussions for animals studies published prior to 2000.

Studies investigating adverse respiratory effects of carbon monoxide have evaluated the lung function and morphology, with most studies evaluating effects of brief exposure (i.e., a few minutes to <1 hour). In acute-duration studies, alterations in lung function (decreased compliance and increased airway resistance) were observed in rabbits and cats exposed to 8,000–28,400 ppm carbon monoxide for <1 hour (COHb >60%); however no changes in lung function were observed in dogs exposed to 8,000–14,000 ppm carbon monoxide for 10–20 minutes (COHb 18–59%) (EPA 1991). In an intermediate-duration study, continuous exposure of rats to 50 ppm carbon monoxide for 21 days (COHb not reported) did not alter the pulmonary pressure-flow relationship (Carraway et al. 2002). Results of these studies suggest that carbon monoxide-induced changes in pulmonary function occur only following exposure to high carbon monoxide concentrations.

Studies evaluating effects of acute-duration carbon monoxide exposure on lung morphology have reported conflicting results (EPA 1991). Minimal changes in type 2 epithelial cells (fragmentation of lamellar bodies) and in nonciliated bronchiolar cells (dilation of smooth endoplasmic reticulum and increased mitochondrial cristae) were reported in mice exposed to 50–90 ppm carbon monoxide for 1–5 days (COHb <10%) (Niden 1971); however, study results were not presented in detail and the toxicological significance of these finding is not known. In rabbits exposed to 8,000 ppm for 30 minutes (COHb 63%), examination of lung tissue by electron microscopy showed alveolar epithelial edema with detachment of the endothelium from the basement membrane (Fein et al. 1980). However, no morphological changes in lung tissue were observed following acute exposure (30 minute to 5 hours) of dogs and rabbits at carbon monoxide exposure producing blood COHb levels of 18–39% or following

## 3. HEALTH EFFECTS

intermediate-duration exposure (6 weeks) of rabbits to 200 ppm carbon monoxide (COHb 11.9–19%) (Fisher et al. 1969; Hugod 1980). Results of a recent study in rats exposed chronically to 200 ppm carbon monoxide (COHb 14.7%) for 20 hours/day for 72 weeks showed no histopathological changes to lung tissue (Sorhaug et al. 2006).

**Cardiovascular Effects.** Cardiovascular effects of inhalation exposures to carbon monoxide have been evaluated in controlled human clinical studies, epidemiology studies, and in various animal models (monkeys, dogs, rats, and rabbits). In general, these studies provide convincing evidence for adverse cardiovascular effects in association with carbon monoxide exposures that result in blood COHb levels of 2.4% and higher, with effects occurring at the lowest levels in subjects who have compromised cardiovascular function (e.g., coronary artery disease). Results of controlled clinical studies in patients with coronary artery disease show that acute-duration exposure to carbon monoxide at levels producing blood COHb levels between 2.4 and 5.9% exacerbates underlying cardiovascular disease, including enhanced myocardial ischemia and increased cardiac arrhythmias.

In healthy subjects, acute exposure to carbon monoxide at concentrations producing blood COHb levels of 3.35 and 20.5% resulted in compensatory cardiovascular responses and decreased exercise performance consistent with COHb-induced decreased O<sub>2</sub> carrying capacity of the blood; however, subjects did not exhibit exercise-induced adverse cardiovascular effects (e.g., evidence of myocardial ischemia or cardiac arrhythmias). Continuous exposure of healthy subjects, which included smokers, to carbon monoxide resulting in blood COHb levels of 2.4 and 5.1% produced P-wave deviations under resting conditions, although similar findings have not been reported in other studies.

Epidemiological studies of exposure to carbon monoxide and cardiovascular outcomes have yielded mixed results; although, in general, the weight of evidence suggests that risks of certain specific outcomes are associated with increasing ambient carbon monoxide concentrations (hospitalizations and emergency room visits related to congestive heart failure, ischemic heart disease, myocardial infarction, and stroke). The corroborated observations of associations between carbon monoxide exposure and outcomes related to ischemic heart disease is particularly provocative in the context of results of human clinical studies in which carbon monoxide-induced hypoxia exacerbated ischemia symptoms in patients with coronary artery disease. Mean ambient air carbon monoxide concentrations reported in studies that have found carbon monoxide-associated cardiovascular outcomes have ranged from 0.5 to 10 ppm, with maximum values ranging from 2 to 50 ppm. These values correspond to approximate steady-state blood COHb levels of <2% for the mean and <10% for the maximum. Studies conducted in animals provide additional

### 3. HEALTH EFFECTS

evidence for adverse cardiovascular effects of carbon monoxide exposure, including compensatory alterations in hemodynamics, cardiac hypertrophy, cardiac arrhythmias, and possibly atherosclerosis.

Studies in animals have investigated adverse cardiovascular effects of carbon monoxide exposure over a much wider range of exposure conditions (e.g., exposure concentration and duration) and have evaluated additional outcome measures that are not possible to assess in humans. These studies provide further evidence of adverse cardiovascular effects of carbon monoxide exposure, including compensatory alterations in hemodynamics, cardiac hypertrophy, cardiac arrhythmias, and possibly atherosclerosis.

#### *Human Clinical Studies*

*Studies of Patients with Cardiovascular Disease.* The effects of acute-duration carbon monoxide exposure on cardiovascular function have been evaluated in several controlled clinical studies in patients with underlying cardiovascular disease (Adams et al. 1988; Allred et al. 1989, 1991; Anderson et al. 1973; Dahms et al. 1993; Hinderliter et al. 1989; Kleinman et al. 1989, 1998; Leaf and Kleinman 1996b; Sheps et al. 1987, 1990) and in healthy subjects (Adir et al. 1999; Davies and Smith 1980; Hausberg and Somers 1997; Horvath et al. 1975; Kizakevich et al. 2000; Morse et al. 2008; Resch et al. 2005; Vogel and Gleser 1972; Zevin et al. 2001). In general, similar effects of acute-duration carbon monoxide exposure were not observed in studies on healthy study subjects, due to functional cardiovascular compensatory mechanisms (e.g., vasodilation and increased cardiovascular output) that protect against COHb-induced tissue hypoxia. Mechanisms contributing to cardiac effects of carbon monoxide in patients with coronary artery disease have not been conclusively established. Under conditions of cardiac ischemia, tissue hypoxia secondary to elevated COHb levels is thought to be a contributing factor. However, other direct cellular effects of carbon monoxide on cardiac muscle may also be important. These include modulation of coronary arteriole calcium-activated potassium channels. These channels are required for membrane hyperpolarization and flow-induced dilation, and are reversibly activated by carbon monoxide under normoxic conditions, but they are inhibited when hypoxic/ischemic conditions prevail (Wang and Wu 1997; Wang et al. 1997a, 1997b).

Controlled clinical studies conducted in small numbers of patients with underlying coronary artery disease have evaluated effects of carbon monoxide exposure on myocardial ischemia. Patients evaluated had a diagnosis of coronary artery disease with exertional angina (i.e., exercise-induced angina or stable angina) and/or ventricular arrhythmia. Most studies were conducted using a randomized, double-blind, crossover design, with cardiovascular function assessments conducted during exercise challenge, either

## 3. HEALTH EFFECTS

during or immediately following carbon monoxide exposure. Carbon monoxide exposures (concentration and duration) were designed to reach target blood COHb levels between 2 and 6%; exposure durations ranged from 1 to 4 hours via an environmental chamber with carbon monoxide concentrations ranging from 42 to 357 ppm. Effects of carbon monoxide exposure were evaluated by comparison to assessments conducted during room air exposure and at corresponding COHb concentrations (<1%). Additional experimental details for each study are summarized in Table 3-4.

Results of controlled clinical studies in patients with coronary artery disease show that acute-duration exposure to carbon monoxide at levels producing blood COHb levels between 2.4 and 5.9% exacerbates underlying cardiovascular disease, including enhanced myocardial ischemia and increased cardiac arrhythmias. In patients with exertional angina, carbon monoxide exposure exacerbated exercise-induced myocardial ischemia, including decreased time-to-onset of angina symptoms (Adams et al. 1988; Allred et al. 1989, 1991; Anderson et al. 1973; Kleinman et al. 1989, 1998; Leaf and Kleinman 1996b), increased duration of angina symptoms (Anderson et al. 1973; Kleinman et al. 1989), decreased time-to-onset of ST-segment depression (EKG change indicative of myocardial ischemia) (Allred et al. 1989, 1991; Kleinman et al. 1989), and decreased left ventricular ejection fraction (Adams et al. 1988). At the lowest blood COHb level evaluated in patients (i.e., COHb 2.4%), time-to-onset of angina symptoms and of ST-segment depression were significantly decreased by 4.2 and 5.1%, respectively (Allred et al. 1989, 1991). In contrast, no carbon monoxide-induced exacerbation of exertional angina, based on time-to-onset of angina symptoms and ST-segment depression and duration of angina symptoms, were observed at a blood COHb level of 4.1% (Sheps et al. 1987). Clinical evaluations in angina patients with ventricular arrhythmias have yielded mixed results on the effects of carbon monoxide-induced ventricular arrhythmias (Dahms et al. 1993; Hinderliter et al. 1989; Sheps et al. 1990). At a blood COHb level of 5.9%, but not 4.0%, single and complex premature ventricular depolarizations were significantly increased by 36 and 6.3%, respectively (Sheps et al. 1990). However, at this same blood COHb level, carbon monoxide exposure did not increase ventricular arrhythmias in other studies (Dahms et al. 1993; Hinderliter et al. 1989). Reasons for conflicting results in these studies are not readily apparent, although numbers of patients evaluated were small and severity of underlying coronary artery disease may have varied in each study.

*Studies of Healthy Subjects.* Several controlled clinical studies in small numbers of healthy subjects have evaluated effects of acute-duration exposure to carbon monoxide on exercise performance, including cardiovascular function. In general, study protocols were similar to those used in clinical studies on patients with coronary artery disease, although typical carbon monoxide exposures were for shorter

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Adams et al. 1988	30 nonsmokers (22 men, 8 women; mean age: 58 years; age range 36–75 years) diagnosed with coronary artery disease with exertional angina pectoris	Randomized, cross-over, double-blind; exercise challenge tests (bicycle) administered immediately following each exposure.	Patients administered one of the following treatments (via environmental chamber) on 2 test days: 1-hour exposure to room air (measured COHb prior to exercise challenge: $1.6 \pm 0.1\%^a$ ) or CO (measured COHb prior to exercise challenge: $5.9 \pm 0.1\%^a$ ; mean CO concentration: 100–200 ppm). COHb measured by CO-Ox.	Duration of exercise, time-to-onset of exercise-induced angina symptoms and ischemic ST-segment changes (EKG), radionuclide ventriculograph (ventricular ejection fraction), systolic blood pressure. Results for CO exposure compared to room air exposure.	<p><u>Mean duration of exercise</u>: Decreased by 6.5% (<math>p &lt; 0.05</math>) following CO exposure, compared to air exposure.</p> <p><u>Time-to-onset of angina symptoms</u>: Using actuarial analysis, subjects more likely to experience angina symptoms earlier in exercise following CO vs. air exposure (<math>p &lt; 0.05</math>).</p> <p><u>Time-to-onset of ischemic ST-segment changes</u>: No effect of exposure.</p> <p><u>Systolic blood pressure</u>: No effect of exposure.</p> <p><u>Ventricular ejection fraction</u>: Left ventricular ejection fraction at submaximal exercise was decreased following CO exposure, compared to air exposure (<math>p &lt; 0.05</math>).</p> <p>NOAEL: Not established LOAEL: COHb 5.9%</p>

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Allred et al. 1989, 1991	63 male nonsmokers (41–75 years) diagnosed with coronary artery disease with exertional angina pectoris and ischemic ST-segment changes on EKG	Randomized, cross-over, double-blind; exercise challenge tests (treadmill) administered prior to and immediately following each exposure.	Patients administered one of the following treatments (via environmental chamber) on 3 test days: 1-hour exposure to room air (measured COHb prior to exercise challenge: $0.70 \pm 0.02\%^a$ ), low CO (measured COHb prior to exercise challenge: $2.38 \pm 0.05\%^a$ ; mean CO concentration: 117 ppm, range 42–202 ppm), or high CO (measured COHb: $4.66 \pm 0.09\%^a$ ; mean CO concentration: 253 ppm, range 143–357 ppm). COHb measured by GC.	Time-to-onset of exercise-induced angina symptoms and ischemic ST-segment changes (EKG), total exercise duration, heart rate-systolic blood pressure double product. Effects determined based comparison (percent change) for pre- and post-exposure exercise challenge outcomes on each test day. Results for each CO exposure compared to room air exposure.	<p><u>Time-to-onset of angina symptoms:</u> Decreased by 4.2% (<math>p=0.027</math>) for low CO exposure and 7.1% (<math>p=0.002</math>) for high CO exposure, compared to room air exposure.</p> <p><u>Time-to-onset of ischemic ST-segment changes:</u> Decreased by 5.1% (<math>p=0.01</math>) for low CO exposure and 12.1% (<math>p \leq 0.0001</math>) for high CO exposure, compared to room air exposure.</p> <p><u>Mean duration of exercise:</u> Decreased by 1.7% (<math>p=0.29</math>) for low CO exposure and 6.2% (<math>p \leq 0.0001</math>) for high CO exposure, compared to room air exposure.</p> <p><u>Heart rate-systolic blood pressure double product:</u> No effect of CO exposure.</p> <p>NOAEL: Not established LOAEL: COHb 2.38%</p>

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Anderson et al. 1973	10 men (5 smokers, 5 nonsmokers; mean age 49.9 years) with diagnosed coronary artery disease with exertional angina pectoris	Randomized, cross-over, double-blind; following COHb elevation to target levels, patients were administered exercise challenge test (treadmill) under maintenance CO exposure.	Patients administered one of the following treatments (via environmental chamber) on 3 test days: 4-hour exposure to room air (measured COHb, prior to exercise challenge: 1.3%), low CO (measured COHb: 2.9%; CO concentration: 50 ppm), or high CO (measured COHb: 4.5%; CO concentration: 100 ppm). COHb measured by spectrometry.	Time-to-onset and duration of angina symptoms.	<p><u>Time-to-onset of angina symptoms:</u> Decreased by 4.2% (p=0.027) for low CO exposure and 7.1% (p=0.002) for high CO exposure, compared to room air exposure.</p> <p><u>Duration of angina symptoms:</u> No effect for low CO exposure; duration increased (p&lt;0.01) for high CO exposure, compared to room air exposure.</p> <p>NOAEL: Not established LOAEL: COHb 2.9%</p>
Dahms et al. 1993	33 nonsmokers (28 men, 5 women; mean age: 58 years; age range 36–75 years) diagnosed with coronary artery disease with chronic ventricular arrhythmias	Randomized, cross-over, double-blind; following COHb elevation to target levels, patients were administered exercise challenge test (treadmill) under maintenance CO exposure.	Patients administered one of the following treatments (via environmental chamber) on 3 test days: 1-hour exposure to room air (measured COHb by CO-Ox, prior to exercise challenge: 0.7), low CO (measured COHb: 3.2%; mean CO concentration: 159 ppm), or high CO (measured COHb: 5.1%; mean CO concentration: 292 ppm). Maintenance exposure on the CO exposure days was to 19 and 31 ppm for the low and high COHb exposures, respectively. COHb measured by CO-Ox.	Frequency of ventricular arrhythmias evaluated during exposure, during exercise (under maintenance exposure), 1–6 hours after cessation of exposure, 7–16 hours after cessation of exposure. Primary outcome was frequency of exercise-induced ventricular arrhythmias. Results for CO exposure compared to room air exposure.	<p>No effect of CO, compared to room air, on frequency of ventricular arrhythmia under any conditions assessed.</p> <p>NOAEL: COHb 5.9% LOAEL: Not established</p>

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Hinderliter et al. 1989	10 nonsmokers (7 males, 3 females; mean age 61 years) diagnosed with coronary artery disease with angina and low baseline ventricular ectopy	Randomized, cross-over, double-blind; exercise challenge tests (bicycle) administered immediately following each exposure	Patients administered one of the following treatments (via environmental chamber) on 3 test days: 1-hour exposure to room air (measured COHb following exposure: $1.4 \pm 0.5\%^b$ ), low CO (measured COHb following exposure: $4.0 \pm 0.4\%^b$ ; CO concentration: 100 ppm), or high CO (measured COHb following exposure: $5.8 \pm 0.4\%^b$ ; CO concentration: 200 ppm). COHb measured by CO-Ox.	Ventricular arrhythmia. Results for CO exposure compared to room air exposure.	No effect of CO, compared to room air, on frequency of ventricular arrhythmia.  NOAEL: COHb 5.9% LOAEL: Not established

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Kleinman et al, 1989	24 male nonsmokers (mean age: 59 years; age range: 49–66 years) diagnosed with coronary artery disease with stable angina pectoris	Randomized, cross-over, double-blind; exercise challenge tests (bicycle) administered immediately following each exposure	Patients administered one of the following treatments (via environmental chamber) on 2 test days: 1-hour exposure to room air (measured COHb following exposure: $1.4 \pm 0.1\%$ ) or CO (measured COHb following exposure: $3.0 \pm 0.1\%$ ; mean CO concentration: 100ppm). COHb measured by CO-Ox.	Time-to-onset and duration of angina symptoms, time-to-onset of ischemic ST-segment changes (EKG), heart rate, systolic blood pressure, oxygen uptake at angina. Results for CO exposure compared to room air exposure.	<p><u>Time-to-onset of angina</u>: Decreased by 5.89% (<math>p &lt; 0.046</math>) following CO exposure, compared to room air exposure.</p> <p><u>Duration of angina symptoms</u>: Increased by 8.33% (not statistically significant) following CO exposure, compared to room air exposure.</p> <p><u>Time-to-onset of ischemic ST-segment changes</u>: Decreased by 19.05% (<math>p &lt; 0.044</math>) following CO exposure, compared to room air exposure.</p> <p><u>Oxygen uptake at angina</u>: Decreased by 2.15% (<math>p &lt; 0.04</math>) following CO exposure, compared to room air exposure.</p> <p><u>Systolic blood pressure</u>: No effect of exposure</p> <p><u>Heart rate</u>: No effect of exposure.</p> <p>NOAEL: Not established LOAEL: COHb 3.0%</p>

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Kleinman et al. 1998; Leaf and Kleinman 1996b	17 male nonsmokers (mean age: 66 years; age range: 58–83 years) living at or near sea level, diagnosed with coronary artery disease with exertional angina pectoris	Randomized, cross-over, double-blind; exercise challenge tests (bicycle) administered immediately following each exposure at sea levels and under conditions simulating high altitude.	Patients administered one of the following treatments (via environmental chamber) on 4 test days: 2-hour exposure to room air at sea level (measured COHb following exposure: $0.6 \pm 0.3\%^b$ ), room air at artificial high elevation (measured COHb following exposure: $0.6 \pm 0.3\%^b$ ), CO at sea level (measured COHb: $3.9 \pm 0.5\%^b$ ; mean CO concentration: 100 ppm), or CO at artificial high elevation (measured COHb: $4.0 \pm 0.6\%^b$ ; mean CO concentration: 100 ppm), COHb measured by CO-Ox.	Time-to-onset and duration of angina symptoms, ST-segment depression and pre-ventricular contractions (EKG), peak (end-exercise) heart rate, blood pressure, oxygen uptake, and carbon dioxide output. Results for CO exposure compared to room air exposure.	<p><u>Time-to-onset of angina</u>: Decreased by 9.1% (<math>p &lt; 0.05</math>) following CO exposure at sea level and by 7.6% (<math>p &lt; 0.05</math>) following CO exposure at high altitude, compared to matched room air exposure.</p> <p><u>Duration of angina symptoms</u>: No effect of CO exposure.</p> <p><u>ST-segment changes</u>: No effect of CO exposure.</p> <p><u>Pre-ventricular contractions</u>: No effect of CO exposure</p> <p><u>Peak heart rate</u>: Increased (<math>p &lt; 0.05</math>) for CO at high altitude compared to CO at sea level and to room air at high altitude.</p> <p><u>Peak blood pressure</u>: Increased (<math>p &lt; 0.05</math>) for CO at high altitude compared to room air at CO at sea level.</p> <p><u>Oxygen uptake and carbon dioxide output</u>: Decreased (<math>p &lt; 0.05</math>) by CO exposure at sea level and high altitude.</p> <p>NOAEL: Not established LOAEL: COHb 3.9–4.0%</p>

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Sheps et al. 1987	30 nonsmokers (age range 38–75 years) with diagnosed coronary artery disease with exertional angina pectoris	Randomized, cross-over, double-blind; exercise challenge tests (bicycle) administered immediately following each exposure.	Patients administered one of the following treatments (via environmental chamber) on 2 test days: 1-hour exposure to room air (measured COHb following exposure: $1.5 \pm 0.05\%$ <sup>b</sup> ) or CO (measured COHb following exposure: $4.1 \pm 0.05\%$ <sup>b</sup> ; CO concentration: 100 ppm). COHb measured by CO-Ox.	Time-to-onset and duration of angina symptoms, ST-segment depression, cardiovascular function (heart rate, blood pressure, rate x pressure product, peak exercise workload, left ventricular ejection fraction), exercise duration. Results for CO exposure compared to room air exposure.	No statistically significant CO-induced changes for any outcomes, except for a small (0.5% decrease; $p=0.049$ ) in maximal ejection fraction. Study authors did not consider this change to be clinically significant.  NOAEL: COHb 4.1% LOAEL: Not established

**Table 3-4. Effects of Acute-duration Exposure to Carbon Monoxide on Cardiovascular Function in Patients with Coronary Artery Disease**

Reference	Study subjects	Design	Exposure	Outcomes Assessed	Results
Sheps et al. 1990	41 nonsmokers (36 men, 5 women; mean age: 62.8 years; age range: 47–77 years) with diagnosed coronary artery disease	Randomized, cross-over, double-blind; exercise challenge tests (bicycle) administered immediately following each exposure.	Patients administered one of the following treatments (via environmental chamber) on 3 test days: 1-hour exposure to room air (measured COHb following exposure: $1.46 \pm 0.07\%$ <sup>a</sup> ), low CO (measured COHb following exposure: $4.01 \pm 0.06\%$ <sup>a</sup> ; CO concentration: 100 ppm), or high CO (measured COHb following exposure: $5.91 \pm 0.08\%$ <sup>a</sup> ; CO concentration: 200 ppm). COHb measured by CO-Ox.	Ventricular arrhythmia (EKG) and ventricular ejection fraction (radionuclide ventriculography), blood pressure, heart rate, exercise duration. Results for CO exposure compared to room air exposure.	For COHb 4.01%: No effects on any parameter evaluated.  For COHb 5.91%: During exercise, statistically significant changes in single premature ventricular depolarizations (36% increase; $p=0.03$ ), complex premature ventricular depolarizations (6.3% increase; $p=0.02$ ), heart rate (3.2% increase; $p=0.01$ ),  NOAEL: COHb 4.01% LOAEL: COHb 5.91%

<sup>a</sup>mean±standard error (SE)<sup>b</sup>mean±standard deviation (SD)<sup>c</sup>means; report does not indicate if SD or SE

CO = carbon monoxide; COHb = carboxyhemoglobin; CO-Ox = CO oximetry; EKG = electrocardiogram; GC = gas chromatography; LOAEL = lowest-observed-adverse-effect level; NOAEL = no-observed-adverse-effect level

## 3. HEALTH EFFECTS

durations (a few minutes to  $\leq 1$  hour) to higher carbon monoxide concentrations (50–3,000 ppm), and produced slightly higher blood COHb levels (2.4–20.5%). However, one study exposed subjects continuously for 7 days (Davies and Smith 1980). Additional experimental details for each study are summarized in Table 3-5.

Results of several controlled clinical studies in healthy subjects show that carbon monoxide exposures that resulted in blood COHb levels between 3.35 and 20.5% decreased exercise performance, including decreased exercise duration (Adir et al. 1999); decreased maximal exercise effort or maximal aerobic capacity (Adir et al. 1999; Horvath et al. 1975; Vogel and Gleser 1972); decreased resistance to muscle fatigue (Morse et al. 2008); and induced compensatory increases in heart rate and cardiac output (Kizakevich et al. 2000; Vogel and Gleser 1972). The above effects on exercise performance are consistent with COHb-induced decreased O<sub>2</sub> carrying capacity of the blood and normal physiological compensatory mechanisms. Evaluations on the potential for carbon monoxide exposure to produce exercise-induced cardiac arrhythmias similar to those observed in patients with coronary artery disease (i.e., ST-segment depression and premature ventricular contractions) have yielded negative results at blood COHb levels up to 5.1% (Adir et al. 1999; Horvath et al. 1975; Kizakevich et al. 2000), although the potential for carbon monoxide-induced cardiac arrhythmias during exercise challenge was not evaluated at higher blood COHb levels. In subjects (including smokers and nonsmokers) exposed continuously for 7 days and evaluated at rest, EKG showed P-wave deviations in 3/15 subjects exposed to 15 ppm carbon monoxide (COHb 2.4%) and in 6/15 subjects exposed to 50 ppm carbon monoxide (COHb 5.1%), compared to 0/14 in subjects exposed to room air (Davies and Smith 1980). In one subject, a heavy smoker, marked ST-depression also was observed. However, interpretation of these findings is limited due to lack of statistical analyses and of evaluation for potential confounding factors (e.g., smoking, medication use). Other studies evaluating effects of a single 1-hour exposure to carbon monoxide (Hausberg and Somers 1997) or repeated brief exposures to carbon monoxide (approximately once every 45 minutes to simulate smoking; Zevin et al. 2001) on cardiovascular function (blood pressure, heart rate, forearm blood flow) under resting conditions show no effects at blood COHb levels up to 8.3%.

In conclusion, results of controlled clinical studies in patients with cardiovascular disease provide compelling evidence that acute-duration exposure to carbon monoxide resulting in COHb levels of 2.4–5.8% exacerbates cardiovascular morbidity (myocardial ischemia and cardiac arrhythmias). Effects at lower blood COHb levels have not been evaluated. However, it should be noted that patients with more severe cardiovascular disease may be more sensitive than patients evaluated in these clinical studies. In

**Table 3-5. Effects of Acute-Duration Carbon Monoxide Exposure on Cardiovascular Function in Healthy Subjects**

Reference	Study subjects	Design	Exposure	Outcomes assessed	Results
Adir et al. 1999	15 healthy nonsmoking men (mean age: 26 years; age range: 18–35 years)	Randomized, cross-over, double-blind; exercise challenge tests (bicycle) administered immediately following each exposure.	Subjects administered one of the following treatments (via closed breathing circuit) on 2 test days: 3–4 minute exposure to room air (COHb: $0.59 \pm 0.08\%$ <sup>a</sup> ) or CO (measured COHb: $5.1 \pm 0.65\%$ <sup>a</sup> ; CO concentration not reported). COHb measured by CO-Ox.	Level of anaerobic respiration (plasma lactate/pyruvate ratio), cardiac arrhythmia (EKG), heart rate and blood pressure (resting and peak), cardiac perfusion (thallium heart scintigraphy), exercise duration, maximal effort (metabolic equivalent units).	<p><u>Anaerobic respiration</u>: No effect of CO exposure.</p> <p><u>Cardiac arrhythmia</u>: No effect of CO exposure.</p> <p><u>Cardiac perfusion</u>: No effect of CO exposure.</p> <p><u>Heart rate and blood pressure</u>: No effect of CO exposure.</p> <p><u>Exercise duration</u>: Decreased by 10.4% (<math>p=0.0012</math>) for CO exposure compared to air exposure.</p> <p><u>Maximal effort</u>: Decreased by 12.4% (<math>p=0.001</math>) for CO exposure compared to air exposure.</p> <p>NOAEL: Not established LOAEL: COHb 5.1%</p> <p><u>P-wave changes</u>: P-wave deviations of <math>\geq 0.1</math> mV observed in 3/15 subjects at low CO and 6/15 at high CO, compared to 0/14 for room air.</p> <p><u>Marked ST-segment depression</u>: 1/15 subject (a heavy smoker) at low CO exposure.</p> <p>NOAEL: Not established LOAEL: COHb 2.4%</p>
Davies and Smith 1980	Groups (14–15/group) of matched healthy (no information on sex reported; age range 17–27 years)	Randomized, parallel group	Continuous exposure to room air (COHb 0.5%), low CO (15 ppm CO; COHb 2.4%) or high CO (50 ppm CO; COHb 7.2%) for 7 days	EKG; no statistical analysis conducted.	<p>NOAEL: Not established LOAEL: COHb 2.4%</p>

**Table 3-5. Effects of Acute-Duration Carbon Monoxide Exposure on Cardiovascular Function in Healthy Subjects**

Reference	Study subjects	Design	Exposure	Outcomes assessed	Results
Hausberg and Somers 1997	10 healthy subjects (8 men, 2 women; mean age: 27 years); 9/10 were nonsmokers	Randomized, double-blind, vehicle (air) controlled	Subjects were assessed under vehicle control (room air), then administered CO (1,000 ppm for 30 minutes, followed by 100 ppm for 30 minutes); COHb increased from $0.2 \pm 0.1\%$ <sup>b</sup> for room air exposure to $8.3 \pm 0.5\%$ <sup>b</sup> during the first 30 minutes of exposure; COHb levels were maintained during the second 30 minutes of CO exposure. COHb measured by CO-Ox.	Forearm blood flow (plethysmography), blood pressure, heart rate, minute ventilation, muscle sympathetic nerve activity (microneurography)	No effects of CO treatment on any parameter evaluated. NOAEL: COHb 8.3% LOAEL: Not established

**Table 3-5. Effects of Acute-Duration Carbon Monoxide Exposure on Cardiovascular Function in Healthy Subjects**

Reference	Study subjects	Design	Exposure	Outcomes assessed	Results
Horvath et al. 1975	4 healthy men (mean age: 24.6 years; age range: 23–33 years); 3/4 were nonsmokers	Randomized, single-blind, cross-over; exercise challenge tests (treadmill) to exhaustion administered immediately following each exposure.	Subjects administered one of the following treatments (via environmental chamber) on 3 test days exposure duration not reported): Vehicle exposure to room air (measured COHb following exposure: $0.33 \pm 0.06\%$ <sup>b</sup> ), low CO (measured COHb following exposure: $3.35 \pm 0.14\%$ <sup>b</sup> ; CO concentration: 75 ppm), or high CO (measured COHb following exposure: $4.30 \pm 0.30\%$ <sup>b</sup> ; CO concentration: 100 ppm). Method for measuring COHb not reported.	EKG, maximal aerobic capacity (maximal oxygen consumption), work time, ventilatory volume	<u>EKG</u> : No effect of CO exposure. <u>Maximal aerobic capacity</u> : Decreased at high CO exposure ( $p < 0.05$ ). Ventilatory volume and work times: Decreased at low and high CO exposure ( $p < 0.05$ ).  NOAEL: Not established LOAEL: COHb 3.35%

**Table 3-5. Effects of Acute-Duration Carbon Monoxide Exposure on Cardiovascular Function in Healthy Subjects**

Reference	Study subjects	Design	Exposure	Outcomes assessed	Results
Kizakevich et al. 2000	16 healthy nonsmoking men (age range: 18–29 years)	Randomized, cross-over (no details on blinding reported); upper (handcrank) and lower body (treadmill) exercise challenges at submaximal effort administered following exposure.	Subjects exposed to CO (concentrations 1,000–3,000 ppm) for 4–6 minutes, followed by maintenance CO exposure (27–100 ppm) on separate test days: Room air (COHb: 1.8%); CO-1 (COHb: 5.0%); CO-2 (COHb: 9.8%); CO-3 (COHb: 14.8%); CO-4 (COHb: 19.2%). COHb measured by CO-Ox.	Cardiac rhythm, output, stroke volume, heart rate, and contractility, time-to-peak ejection time, blood pressure.	<u>Cardiac rhythm</u> : No effects of CO exposure, compared to room air exposure; no arrhythmias or ST-segment depression. <u>Compensatory changes associated with reduced oxygen carrying capacity</u> : Significant increases in heart rate at all CO exposures and cardiac contractility at COHb 9.8%.  NOAEL: Not established LOAEL: COHb 5.0%
Morse et al. 2008	12 healthy nonsmoking men (mean age: 25 years)	Cross-over; skeletal muscle fatigue challenge administered following exposure	Room air or 3,000 ppm for 6 minutes (COHb 6.2%)	Leg strength, muscle fatigue assessed during voluntary and electrically evoked muscle contractions	Resistance to muscle fatigue decreased under voluntary (13% decrease; p<0.05) and electrically evoked stimulation (12.5% decrease; p<0.05). No effect on leg strength.  NOAEL: Not established LOAEL: COHb 6.2%
Resch et al. 2005	6 healthy nonsmoking men (mean age: 26 years)	Randomized, cross-over, double-blind	Subjects administered the following treatments: Room air (COHb 1.2%), 500 ppm for 30 minutes (COHb 8.5%); 500 ppm for 30 minutes (COHb 9.4%).	Retinal and submacular choroidal blood flow	Significant increases in retinal and choroidal blood flow and retinal vessel diameter at COHb ≥8.5%. Study authors state that relationship of this finding to tissue hypoxia is unknown.  NOAEL: Not established LOAEL: COHb 8.5%

**Table 3-5. Effects of Acute-Duration Carbon Monoxide Exposure on Cardiovascular Function in Healthy Subjects**

Reference	Study subjects	Design	Exposure	Outcomes assessed	Results
Vogel and Gleser 1972	8 men, including 3 smokers (age range: 20–23 years)	Randomized, cross-over (no details on blinding reported); exercise challenges administered during exposure.	1-hour exposure to air (COHb 1.0–1.7%) or 225 ppm CO (COHb 18.6–20.5%)	Maximal oxygen uptake assessed at rest and at submaximal and maximal exercise	At rest, no effect of CO exposure. At submaximal exercise, no change in oxygen uptake due to compensatory increase in cardiac output. At maximal exercise, oxygen uptake decreased by 23% (p<0.001)  NOAEL: Not established LOAEL: COHb 18.6–20.5%
Zevin et al. 2001	12 healthy men (smokers; age range: 27–47)	Single-blind, cross-over;	Subjects exposed to 3 treatments for durations of 7 days each: Room air (COHb 0.4%), 1,200 ppm CO inhalation 20 times/day (~every 45 minutes, 20 times/day to simulate smoking (COHb 6%), or 20 cigarettes/day (COHb 5%).	Heart rate, blood pressure, catecholamine release, platelet activation, white blood cell count, C-reactive protein (inflammation marker)	<u>CO exposure:</u> No effect on serum levels of CRP or platelet factor 4, heart rate, blood pressure, or catecholamine release, compared to room air.  <u>Smoking:</u> Significant elevation in serum levels of CRP and platelet factor 4, increased heart rate, or catecholamine release, compared to room air. No effect on blood pressure.  NOAEL (CO): COHb 6% LOAEL (CO): Not established

<sup>a</sup>mean± standard deviation (SD)

<sup>b</sup>mean± standard error (SE)

<sup>c</sup>means; report does not indicate if SD or SE

CO = carbon monoxide; COHb = carboxyhemoglobin; CO-Ox = CO oximetry; EKG = electrocardiogram; GC = gas chromatography; LOAEL = lowest-observed-adverse-effect level; NOAEL = no-observed-adverse-effect level

## 3. HEALTH EFFECTS

healthy subjects, acute exposure to carbon monoxide at concentrations producing blood COHb levels of 3.35 and 20.5% resulted in compensatory cardiovascular responses and decreased exercise performance consistent with COHb-induced decreased O<sub>2</sub> carrying capacity of the blood; however, subjects did not exhibit exercise-induced adverse cardiovascular effects (e.g., evidence of myocardial ischemia or cardiac arrhythmias). Continuous exposure of healthy subjects, which included smokers, to carbon monoxide resulting in blood COHb levels of 2.4 and 5.1% produced P-wave deviations under resting conditions, although similar findings have not been reported in other studies.

***Epidemiological Studies.*** Epidemiological studies of exposure to carbon monoxide and cardiovascular outcomes fall into two major categories: (1) studies that have assessed cardiovascular morbidity in cohorts or individuals (e.g., case-crossover, case-control) in association with ambient air carbon monoxide concentrations; and (2) studies that have evaluated associations between air carbon monoxide concentrations and the incidence of hospital admissions and/or emergency room visits related to cardiovascular disease (e.g., time-series studies, case-crossover). Collectively, these studies have yielded mixed results; although, in general, the weight of evidence suggests that risks of certain specific outcomes are associated with increasing ambient carbon monoxide concentrations (hospitalizations and emergency room visits related to congestive heart failure, ischemic heart disease, myocardial infarction, and stroke). Corroborated observations of association between carbon monoxide exposure and outcomes related to ischemic heart disease are consistent with results of human clinical studies that have shown that carbon monoxide-induced hypoxia exacerbates symptoms of coronary artery disease. Therefore, it is plausible that in certain susceptible populations, such as individuals who have ongoing cardiac impairment that would make them vulnerable to ischemia, exposure to carbon monoxide could contribute to triggering ischemic episodes and cardiac sequelae. While these studies provide some evidence in support of adverse cardiovascular effects of inhalation exposures to carbon monoxide, their utility for establishing dose-response relationships for these effects are limited by several factors: (1) reliance on area monitoring for estimating exposure levels; (2) uncertainty in knowledge of temporal correspondence between monitored exposure levels and outcomes; (3) relatively strong correlations between ambient air carbon monoxide concentrations and other air quality variables that can affect respiratory function; (4) outcomes investigated have tended to be serious, life-threatening events (e.g., cardiac arrhythmia, myocardial infarction, stroke, heart failure) and reflect the “late-stage” consequences of contributing pathophysiology that might be associated with exposures to lower levels of carbon monoxide for longer durations than indicated in the epidemiological studies; and (5) relatively low carbon monoxide exposures studied. Mean values of carbon monoxide exposure concentrations have ranged from 0.3 to 4.6 ppm with the highest values <30 ppm. These values correspond to predicted steady-state COHb levels of <1% for the

## 3. HEALTH EFFECTS

mean and <5% for the highest values (predicted from the CFK model). The latter range is consistent with COHb levels observed in a study of emergency department admissions for cardiopulmonary complaints (Leikin and Vogel 1986). At these COHb levels, it is possible that only highly susceptible individuals exhibit the serious cardiovascular outcomes that have been studied. Clinical studies described previously in this section of the Toxicological Profile have found that exposures to carbon monoxide that produce COHb levels 2.4–5% can exacerbate underlying cardiovascular disease, including exercise-induced angina and cardiac arrhythmia; however, higher levels (3–20%) can be tolerated without producing such effects, even under exertion, in health individuals. For these and other reasons, epidemiologic studies have typically focused on highly susceptible populations. A typical design has been to examine associations between temporal trends in ambient air carbon monoxide concentrations and hospital admissions or emergency department visits for which the reported diagnosis was some form of cardiovascular disease or impairment.

The presentation of the epidemiology is organized by major categories of cardiovascular outcomes for which the studies were designed to evaluate, which include studies of heart rate and heart rate variability, cardiac arrhythmia, ischemic heart disease and related subcategories (e.g., myocardial infarction), heart failure, stroke, and blood pressure. Study conclusions are presented in the text, with selected supporting details presented in tabular form. Although most of the studies explored various time lags between monitored air carbon monoxide concentrations and outcomes, as well as various sample strata, for the sake of brevity, only those indicative of the strongest associations to carbon monoxide are presented in the tables. Where co-pollutant models have been explored, these results are also presented.

*Studies of Heart Rate (HR) and Heart Rate Variability (HRV).* Studies of associations between exposure to carbon monoxide and various measures of HR and HRV have yielded mixed results (Table 3-6). Carbon monoxide exposure concentrations in these studies represent relatively narrow time periods (e.g., averages over 1–5 days prior to the end point assessment). Mean values of carbon monoxide exposure concentrations ranged from 0.3 to 4.6 ppm, with the highest values <30 ppm (predicted steady-state COHb%: <5). Significant associations between increasing carbon monoxide exposure and increasing HR were found in some studies (Liao et al. 2004; Peters et al. 1999) and not in others (Gold et al. 2000). Assessments of HRV found significant associations with increasing carbon monoxide exposure in some studies (Dales 2004; Riojas-Rodriguez et al. 2006; Schwartz et al. 2005; Tarkiainen et al. 2003; Timonen et al. 2006) and no associations with HRV in other studies (Chan et al. 2005; Gold et al. 2000; Holguin et al. 2003; Liao et al. 2004; Park et al. 2005b). In general, associations for HRV end points were not robust for various measures of HRV across studies. For example, assessments of standard deviation of the

**Table 3-6. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Heart rate and heart rate variability			
Chan et al. (2005) Period: 2001–2002 Location: Taipei, Taiwan	Outcome: HRV Design: Panel Sample: 83 patients, age 43–75 years	Avg time: 1 hour Mean 1.1 ppm Range: 0.1–7.7 ppm	No statistically significant effect on SDNN, rMSSD, LFV, HFV
Dales (2004) Period: NR Location: Toronto, Canada	Outcome: HRV Design: Panel Sample: 36 subjects with preexisting CAD, age 51–88 years	Avg time: 24 hours Mean 95 <sup>th</sup> ‰: 2.40 ppm Range: 0.4–16.5 ppm	Increment: NR Regression coefficient (95% CI): SDNN (not taking $\beta$ -blockers): 0.0111 (0.002–0.020, p=0.02) No statistically significant effect on HRV among those taking $\beta$ -blockers
Gold et al. (2000) Period: 1997 Location: Boston, Massachusetts	Outcome: HR and HRV Design: Panel/Cohort Sample: n=21, age 53–87	Avg time: 24 hours Mean: 0.47 ppm Range: 0.12–0.82 ppm	Increment: 0.6 ppm % Change (lag 2 day): No significant effect of CO on HR or HRV
Holguin et al. (2003) Period: 2000 Location: Mexico City, Mexico	Outcome: HRV Design: Panel Sample: n=34 age 60–90 years	Avg time: 24 hours Mean: 3.3 ppm Range: 1.8, 4.8 ppm	Increment: 10 ppm Regression coefficients (95% CI), lag 0 day: HFV: 0.003 (-0.004–0.001) LFV: 0.001 (-0.006–0.008) LFV/HFV: 0.001 (-0.005–0.002)
Liao et al. (2004) Period: 1996–1998 Location: Various U.S. cities	Outcome: HR HRV. Design: Cohort Sample: n=6784, age 45–64 years	Avg time: 24 hours Mean 0.65 ppm SD: 0.44 ppm	Increment: 0.44 ppm Regression coefficient, lag 1 day: HR (bpm): 0.404 (p<0.05) HFV (log transformed): -0.033 LFV (log transformed): 0.006 SDNN: -0.274
Park et al. (2005b) Period: 2000–2003 Location: Boston, Massachusetts	Outcome: HRV Design: Panel/Cohort Sample Description: n=497 males, age 21–81 years	Avg time: 24 hours Mean: 0.50 ppm Range: 0.13–1.8 ppm	Increment: 0.24 ppm % Change in HRV (95% CI), 4-hour moving avg: SDNN (log10): 2.0 (-2.9–7.3) HFV (log10): 8.8 (-4.6–24.1) LFV(log10): 3.2 (-7.0–14.6) LFV:HFV(log10): -5.1 (-13.5–4.1)

**Table 3-6. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Peters et al. (1999) Study: 1984–1985 Location: Augsburg, Germany	Outcome: HR Design: Cohort Age Groups Analyzed: 25–64 years Sample: n=2,681, age 25–64 years	Avg time: 24 hours Mean: 3.9 ppm Range: 0.8–10.0 ppm	Increment: 6.6 mg/m <sup>3</sup> Mean change in HR, bpm (95% CI), lag 0 day: All: 0.97 (0.02–1.91) Male: 0.95 (-0.37–2.27) Female: 0.98 (-0.37–2.34)
Riojas-Rodriguez et al. (2006) Period: 2001–2002 Location: Mexico City, Mexico	Outcome: HRV Design: Panel Sample: n=30, age 25–76 years	Avg time: 24 hours Mean: 2.9 ppm Range: 0.1–18.0 ppm	Increment: 1 ppm Regression coefficients (95% CI, lag 5 minutes): HFV: -0.006 (-0.023 to 0.010) LFV: -0.024 (-0.041 to -0.007) VLFV: -0.034 (-0.061 to -0.007)
Schwartz et al. (2005) Period: 1999 Location: Boston, Massachusetts	Outcome: HRV Design: Panel Sample: n=28, age 61–89 years	Avg time: 24 hours Median: 0.45 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.38–0.54 ppm	Increment: 0.16 ppm % Change in HRV (95% CI), lag 1 day): SDNN: -4.2 (-0.6 to -7.7) rMSSD: -10.2 (-2.4 to -17.4) PNN50: -14.8 (-3.0 to -25.2) LFV/HFV: 6.2 (-0.6 to 13.4)
Tarkiainen et al. (2003) Period: 1997–1998 Location: Kuopio, Finland	Outcome: HRV Design: Panel Sample: n=6 males, age 55–68 years	Avg time: 24 hours Mean: 4.6 ppm Range: 0.5–27.4 ppm	Increment: NR RR (95% CI, lag 5 minutes) CO>2.7 ppm: Significant increase in rMSSD (2.4 ms, p=0.034). CO <27 ppm: No significant association between CO and NN, SDNN, or rMSSD
Timonen et al. (2006) Period: 1998–1999 Location: Amsterdam, Netherlands; Erfert, Germany; Helsinki, Finland	Outcome: HRV Design: Panel Sample: n=131, age 64–72 years	Avg time: 24 hour Mean: 0.3–0.5 ppm Range: 0.1–2.2 ppm	Increment: 1 mg/m <sup>3</sup> β coefficient (95% CI), lag 2 days SDNN: -5.96 (-10.7, -0.72)   HF: -30.7 (-59.8, -1.5) LH/HF: -10.1 (-36.9, 16.7)

**Table 3-6. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
<b>Cardiac arrhythmia</b>			
Berger et al. (2006) Period: 2000–2001 Location: Erfurt, Germany	Outcome: SVT and VT 24 hour ECG Design: Panel Sample: 57 males with CHD, age 52–76 years	Avg time: 24 hours Mean: 0.45 ppm Range: 0.10–1.68 ppm	Increment: 0.22 mg/m <sup>3</sup> RR for SVT (95% CI), 5-day avg: 1.18 (1.04–1.35)
Dockery et al. (2005) Period: 1995–2002 Location: Boston, Massachusetts	Outcome: Tachycardia Design: Panel Sample: n=203 with ICDs, age 19–90 years	Avg time: 24 hours Median: 0.8 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.53–1.02 ppm	Increment: 0.48 ppm OR for VT (95% CI), lag 2-day: All: 1.14 (0.95–1.29) AR ≤3 days prior: 1.65 (1.17–2.33)
Gold et al. (2005) Period: 1999 Location: Boston, Massachusetts	Outcome: ST-segment Design: Panel Sample: n=24, age 61–68 years	Avg time: 5 hours Median: 0.52 ppm 10 <sup>th</sup> –90 <sup>th</sup> ‰: 0.20–1.08 ppm	Significant association between CO and ST-segment depression did not persist in multiple pollutant models
Metzger et al. (2007) Period: 1993–2002 Location: Atlanta, Georgia	Outcome: CAR, VT Design: Panel Sample: n=518 with ICDs, with ≥1 VT event, age 15–88 years	Avg time: 1 hour Mean 1.7 ppm Range: 0.1–7.7 ppm	Increment: 1 ppm OR for VT (95% CI), lag 0 day: All: 0.999 (0.970–1.028)
Peters et al. (2000b) Period: 1995–1997 Location: Eastern Massachusetts	Outcome: VT or fibrillation Design: Panel Sample: n=100 with ICDs, mean age 62 years	Avg time: 24 hours Mean 0.58 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.66 ppm	Increment: 0.65 ppm OR for ≥10 defibrillated discharges (95% CI): Lag 3: 1.98 (1.05–3.72) Lag 5-day avg: 1.94 (1.01–0.75)
Rich et al. (2004) Period: 2000 Location: Vancouver, Canada	Outcome: CAR Design: Case-crossover Sample: n=34 with ≥1 ICD discharge, age 15–85 years	Avg time: 24 hours Mean: 0.553 ppm Interquartile range: 0.162 ppm	No significant effect of CO (quantitative results not reported)
Rich et al. (2005) Period: 1995–1999 Location: Boston, Massachusetts	Outcome: VAR Design: Panel/Case-crossover Sample: n=203 with implanted ICDs	Avg time: 24 hours Median: 0.78 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.52–1.03	Increment: 0.5 ppm OR: (95% CI), lag 0–2 day): 1.11 (0.88–1.40) (also not significant at other lags)

**Table 3-6. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Rich et al. (2006b) Period: 2001 and 2002 Location: St. Louis, Missouri	Outcome: VAR Design: Case-crossover Sample description: n=60 with ≥1 ICD CAR recorded	Avg time: 24 hours Median: 0.5 ppm 25 <sup>th</sup> –75 <sup>th</sup> o: 0.6 ppm	Increment: 0.2 ppm OR for VAR (95% CI), lag 0–1 day: 0.99 (0.80– 1.21)
Rich et al. (2006a) Period: 1995–1999 Location: Boston, Massachusetts	Outcome: AF Design: Panel/ case-crossover Sample: n=203 with ICDs	Avg time: 24 hours Median: 0.78 ppm 25 <sup>th</sup> –75 <sup>th</sup> o: 0.52–1.03 ppm	Increment: 0: 0.58 ppm OR for AF (95% CI): Lag 0: 0.87 (0.56–1.37) Lag 0–23 hours: 0.71 (0.39–1.28)
Sarnat et al. (2006) Period: 2000 Location: Steubenville, Ohio	Outcome: CAR Design: Panel Sample 32 nonsmoking, age 53– 90 years	Avg time: 24 hours Mean: 0.02 ppm Range 0–1.5 ppm	Increment: 0.2 ppm RR (95% CI), 5-day moving avg: SVE: 0.99 (0.76–1.29) VE: 1.05 (0.75–1.46)
Vedal et al. (2004) Period: 1997–2000 Location: Vancouver, Canada	Outcome: CAR Design: Panel Sample: n=50 with ICDs and ≥1 CAR event/4 years, age 12– 77 years	Avg time: 24 hours Mean: 0.6 ppm Range: 0.3–1.6 ppm	Increment: 0.2 ppm No significant effect for CO (results shown in plots)
<b>Cardiac arrest</b>			
Levy et al. (2001) Period: 1988–1994 Location: Seattle, Washington	Outcome: Cardiac arrest Design: Case-crossover Sample: 362 cases, age 25– 75 years	Avg time: 24 hours Mean: 1.79 ppm Range: 0.52–5.92 ppm	RR (95% CI), lag 1 day: 0.99 (0.83–1.18)
Sullivan et al. (2003) Period: 1985–1994 Location: Washington State	Outcome: Cardiac arrest Design: Case-crossover Sample: 1,542 cases, median age 69 years	Avg time: 24 hours Mean: 1.92 ppm Range: 0.52–7.21 ppm	Increment: 1.02 ppm OR (95% CI): Lag 0: 0.95 (0.85–1.05) Lag 1: 0.97 (0.87–1.08) Lag 2: 0.99 (0.89–1.11)

**Table 3-6. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
<b>Myocardial infarction</b>			
Peters et al. (2001) Period: 1995–1996 Location: Boston, Massachusetts	Outcome: Onset of MI Design: Case-crossover Sample: 772 cases	Avg time: 24 hours Mean: 1.09 ppm 5 <sup>th</sup> –95 <sup>th</sup> ‰: 0.49–1.78 ppm	Increment: 1 or 0.6 ppm OR (95% CI): 2 hours: 1.22 (0.89–1.67) per 1 ppm 24 hours: 0.98 (0.70–1.36) per 0.6 ppm
Rosenlund et al. (2006) Period: 1992–1994 Location: Stockholm, Sweden	Outcome: MI Design: Case-control Sample: 1,397 cases, 1,870 controls, age range 45–70 years	Avg time: 30 years Mean: 0.058 ppm 5 <sup>th</sup> –95 <sup>th</sup> ‰: 0.012–0.258 ppm	Increment: 300 µg/m <sup>3</sup> OR: (95% CI): All cases: 1.04 (0.89–1.21) Nonfatal: 0.98 (0.82–1.16) Fatal: 1.22 (0.98–1.52) Fatal, in-hospital: 1.16 (0.89–1.51) Fatal, out-hospital: 1.36 (1.01–1.84)
<b>Blood pressure</b>			
Ibald-Mulli et al. (2001) Period: 1984–1985 Location: Augsburg, Germany	Outcome: SBP Design: Cohort Sample: n=2,607 men and women age 25–64 years	Avg time: 24 hours Mean: 3.6 ppm Range: 1.5–7.2 ppm	Increment: 5.6 mg/m <sup>3</sup> Mean change in SBP, mmHg (95% CI): All: 0.53 (-0.66–1.72) Males: 0.68 (-0.94–2.31) Females: 0.51 (-1.31–2.19)
Zanobetti et al. (2004b) Period: 1999–2001 Location: Boston, Massachusetts	Outcome: BP Design: Cohort/Panel Sample description: n=62, age 39–90 years	Avg time: 120 hours Mean: 0.66 ppm 10 <sup>th</sup> –90 <sup>th</sup> ‰: 0.48–0.86 ppm	Increment: NR RR: No significant effect of CO on BP

AF = atrial fibrillation; AP = angina pectoris; AR = arrhythmia; Avg = average; BP = blood pressure; BPM = beats per minute; CAD = coronary artery disease; CAR = cardiac arrhythmia; CI = confidence interval; CHD = coronary heart disease; CO = carbon monoxide; ECG = electrocardiogram; HFV = high frequency (HR) variability parameter; HR = heart rate; HRV = heart rate variability; ICD = implantable cardioverter defibrillator; LFV = low frequency (HR) variability parameter; Max = maximum; MI = myocardial infarction; NR = not reported; OR = odds ratio; PNN50 = proportion of interval differences of successive normal-beat intervals greater than 50 ms in EKG; rMSSD = root mean squared differences between adjacent RR intervals; RR = relative risk; SBP = systolic blood pressure; SDNN = standard deviation normal-to-normal (NN or RR) time interval between each QRS complex in the EKG; SVE = supraventricular ectopy; SVT = supraventricular tachycardia; VAR = ventricular arrhythmia; VLFV = very low frequency (HR) variability parameter; VE = ventricular ectopy; VT = ventricular tachycardia

## 3. HEALTH EFFECTS

normal-to-normal interval (SDNN) yielded significant associations in some studies (Dales 2004; Schwartz et al. 2005; Timonen et al. 2006) and not in others (Chan et al. 2005; Gold et al. 2000; Liao et al. 2004; Park et al. 2005b; Tarkiainen et al. 2003).

Three studies are particularly notable for their relatively large size. Park et al. (2005b) assessed HRV in 497 male subjects (age range: 21–82 years) of the Boston Normative Aging Study and linked these data to local measurements of air pollutant concentrations (carbon monoxide, PM<sub>2.5</sub>, O<sub>3</sub>, NO<sub>2</sub>, SO<sub>2</sub>). The mean carbon monoxide exposure concentration was 0.50 ppm (24-hour average, range: 0.13–18 ppm). The study did not find significance between carbon monoxide exposure levels and HRV end points assessed (SDNN, low frequency [LF], high frequency [HF], LF/HF). A study conducted in Augsburg, Germany evaluated possible associations between exposures to total suspended particles (TSP), SO<sub>2</sub>, and carbon monoxide (mean: 4.5 mg/m<sup>3</sup>, range: 2.4–6.8 mg/m<sup>3</sup>) and HR in 2,681 adults (age range: 25–64 years; Peters et al. 1999). A 6.6 mg/m<sup>3</sup> (5.8 ppm) increase in 24-hour average carbon monoxide concentration was associated with a small but significant increase in HR of approximately 1 beat per minute (0.97 beats/minute, 95% CI: 0.02–1.91); however, carbon monoxide exposure was not significantly associated with HR when based on a 5-day average exposure concentration (0.7 per 3.3 mg/m<sup>3</sup>, 95% CI: -0.09–1.58). A subsample of 6,232 adults (age range: 45–64 years) from the Atherosclerosis Risk in Communities (ARIC) study (Washington County, Maryland; Forsyth County, North Carolina; and selected suburbs of Minneapolis, Minnesota and Jackson, Mississippi) examined possible associations between exposure to carbon monoxide (mean: 0.65 ppm, standard deviation [SD]: 0.44) and HR and HRV, as well as to other air pollutants (PM<sub>10</sub>, O<sub>3</sub>, NO<sub>2</sub>, SO<sub>2</sub>; Liao et al. 2004). An increase in carbon monoxide exposure concentration of 0.44 ppm was associated with a small but significant, increase in HR of 0.4 beats per minute (standard error [SE]: 0.16, 95% CI: 0.09–0.7 [calculated from SE]). Carbon monoxide exposure was not significantly associated with HRV.

*Studies of Arrhythmia.* Studies of associations between exposure to carbon monoxide and cardiac arrhythmia (e.g., fibrillation, tachycardia) have yielded mixed results. Mean values of carbon monoxide exposure concentrations (i.e., averages over 1–5 days prior to the end point assessment) ranged from 0.2 to 1.7 ppm, with the highest values <8 ppm. Significant associations between increasing carbon monoxide exposure and arrhythmia episode incidence were found in some studies, particularly when the analysis was restricted to subjects with prior recent arrhythmia (Berger et al. 2006; Dockery et al. 2005; Peters et al. 2000a), but not in other studies (Metzger et al. 2007; Rich et al. 2004, 2005; Sarnat et al. 2006; Vedal et al. 2004). One of the larger studies examined associations between exposures to carbon monoxide (median: 0.8 ppm; 95<sup>th</sup> percentile: 1.37 ppm) and other air pollutants (e.g., PM<sub>10</sub>, PM<sub>2.5</sub>, O<sub>3</sub>,

## 3. HEALTH EFFECTS

NO<sub>2</sub>, SO<sub>2</sub>) and ventricular arrhythmia episodes in 203 patients who were monitored with implanted defibrillators (ICDs) and who had prior episodes of ventricular tachycardia (Dockery et al. 2005; Rich et al. 2005). In the total sample, exposure to carbon monoxide was not significantly associated with ventricular arrhythmia; however, when restricted to patients who had experienced arrhythmias within the 3 days of the exposure assessment, increasing carbon monoxide exposure of 0.48 ppm was significantly associated ventricular arrhythmia (odds ratio: 1.65; 96% CI: 1.17–2.33). A larger study examined 512 patients (age range: 15–88 years) exposed to a mean carbon monoxide concentration of 1.7 ppm (range: 0.1–7.7 ppm) and did not find a significant association with defibrillator discharges and carbon monoxide exposure concentration (Metzger et al. 2007).

Studies that have assessed arrhythmias from electrocardiogram (ECG) measurements have also yielded mixed results (Berger et al. 2006; Sarnat et al. 2006). A study of 57 coronary heart disease patients (age range: 52–76 years) found a significant association between increasing carbon monoxide exposure and risk of supraventricular tachycardia (Berger et al. 2006). Mean carbon monoxide exposure concentration was 0.52 mg/m<sup>3</sup> (0.45 ppm; range: 0.11–1.93 mg/m<sup>3</sup>). The relative risk for a 0.27 mg/m<sup>3</sup> (0.24 ppm) increase in carbon monoxide was 1.18 (95% CI: 1.04–1.35). A significant association between carbon monoxide exposure and ventricular tachycardia was not evident. Sarnat et al. (2006) did not find a significant association between carbon monoxide exposure concentrations (mean: 0.2 ppm, range: 0.1–1.5) and arrhythmia in a sample of 32 adults (age range: 53–90 years). The study did find significant associations between increased blood pressure and other air pollutants (e.g., PM<sub>2.5</sub>, SO<sub>2</sub>, O<sub>3</sub>, back carbon).

*Studies of Ischemic Heart Disease.* Numerous studies have examined possible associations between carbon monoxide exposure and ischemic heart disease (Tables 3-6 and 3-7). Although results have been mixed, most studies, in particular those that have examined hospital admissions or emergency room visits as the health outcome, have found increased risk association with increased ambient air carbon monoxide concentrations. A case-control study of myocardial infarction (1,397 cases, 1,870 controls) conducted in Sweden, did not find a significant association with carbon monoxide exposure (Rosenlund et al. 2006). Exposure concentrations were estimated from emissions data and air dispersion modeling; the mean was 61.8 µg/m<sup>3</sup> (0.067 ppm; 5<sup>th</sup>–95<sup>th</sup> percentile range: 14–296 µg/m<sup>3</sup>; predicted steady-state COHb%: <0.3). Odds ratio for myocardial infarct for a 300 µg/m<sup>3</sup> (0.3 ppm) increase in 30-year average carbon monoxide concentration was 1.04 (95% CI: 0.89–1.21). When restricted to out-of-hospital deaths, the odds ratio was 1.36 (95% CI: 1.01–1.84). A smaller case-crossover study (772 cases) conducted in Boston, Massachusetts did not find a significant association between carbon monoxide exposure and myocardial infarction (Peters et al. 2001). The odds ratio for 1 ppm increment in carbon monoxide exposure

**Table 3-7. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Hospital Admissions and Emergency Department Visits Related to Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Ischemic heart disease			
D'Ippoliti et al. (2003) Period: 1995–1997 Location: Rome, Italy	Outcome: MI HAs Design: Case-crossover Sample: n=6,531, ages ≥18 years	Avg time: 24 hours Mean: 3.8 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 2.4–3.8 ppm	Increment: 1 mg/m <sup>3</sup> OR (95% CI), lag 0–2 days: CO alone: 1.044 (1.000–0.089) CO with TSP: not significant
Hosseinpour et al. (2005) Period: 1996–2001 Location: Tehran, Iran	Outcome: AP HAs Design: Time series Sample: n=1,826	Avg time: 24 hours Mean: 9.4 ppm Range: 1.4–50.5 ppm	Increment: 1 mg/m <sup>3</sup> (0.87 ppm) RR (95% CI), lag 1 day: CO alone: 1.0096 (1.0060–1.0132) CO with NO <sub>2</sub> , O <sub>3</sub> , and PM <sub>10</sub> : 1.0093 (1.0036–1.0151)
Lanki et al. (2006) Period: 1994–2000 Location: Helsinki, Rome, Stockholm	Outcome: MI HAs Design: Time series Sample: 26,854 admissions, age ≥35 years	Avg time: 24 hours Median: 0.4–2.3 ppm 75 <sup>th</sup> ‰ Range: 0.5–2.9 ppm	Increment: 0.2 mg/m <sup>3</sup> RR (95% CI), lag 0 day: All: 1.007 (1.001–1.012) Fatal (≤75 years): 1.027 (1.006–1.048)
Lee et al. (2003b) Period: 1997–1999 Location: Seoul, Korea	Outcome: IHD, MI, AP HAs Design: Time series Sample: ~10,000 HAs	Avg time: Daily maximum Mean: 1.8 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 1.2–2.2 ppm	Increment: 1 ppm RR (95% CI), lag 5 days: All year, all ages: 0.94 (0.91–0.98) All year, ≥64 years: 1.07 (1.01–1.13) Summer, all ages: 1.19 (1.02–1.38) Summer, ≥64 years: 1.60 (1.27–2.03) All year, ≥64 years (with PM <sub>10</sub> ): 1.04 (0.98–1.11)
Maheswaran et al. (2005b) Period: 1994–1998 Location: Sheffield, United Kingdom	Outcome: IHD HAs Design: Ecological Subjects: 11,407 admissions, age ≥ 45 years	Avg time: NR 20 <sup>th</sup> –80 <sup>th</sup> ‰: 0.34–0.40 ppm	Increment: 5 <sup>th</sup> vs. 1 <sup>st</sup> quintile RR, adjusted for sex, age, deprivation, smoking (95% CI): 0.88 (0.79–0.98)
Mann et al. (2002) Period: 1988–1995 Location: California	Outcome: IHD, MI HAs Design: Time series Sample: 54,863 HAs	Avg time: 8 hours Mean: 2.07 ppm Range: 0.30–11.8 ppm	Increment: 1 ppm % Change (95% CI), lag 0 day: with arrhythmia: 2.99 (1.80–4.99) with CHF: 3.60 (1.620–5.63) without secondary diagnosis: 1.62 (0.65–2.59)

**Table 3-7. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Hospital Admissions and Emergency Department Visits Related to Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Szyszkowicz (2007) Period: 1997–2003 Location: Montreal	Outcome: IHD EDVs Design: Time series Sample: 4,979 EDVs	Avg time: 24 hours Mean: 0.5 ppm Range: 0.1–3.1 ppm	Increment: 0.2 ppm % Change (95% CI), lag 0 day: All: 5.4 (2.3–8.5) Males: 7.5 (3.6–11.6) Females: 2.7 (-2.0– 7.6) All: ≥64 years: 4.9 (1.3–8.7) Males: ≥64 years: 7.5 (2.6–12.6) Females: ≥64 years: 0: 2.4 (-3.0 to 0.0)
von Klot et al. (2005) Period: 1992–2001 Location: 5 European cities	Outcome: MI, AP, CAR, HF, HAS Design: Prospective cohort Sample: 22,006 survivors of first MI	Avg time: 24 hours Mean: 0.37–0.87 ppm	Increment: 0.2 mg/m <sup>3</sup> RR (95% CI), lag 0 day: MI: 1.022 (0.998–0.047) AP: 1.009 (0.992–0.02) Any cardiac: 1.014 (1.001–0.026)
Stroke			
Chan et al. (2006) Period: 1997–2002 Location: Taipei, Taiwan	Outcome: Stroke EDVs Design: Time series Sample: 7,341 EDVs, age 50–100 years	Avg time: 8 hours Mean: 1.7 ppm Range: 0.6–4.4 ppm	Increment: 0.8 ppm OR (95% CI), lag 2 days: All: 1.03 (1.01, 1.06) Stroke: 1.03 (1.01, 1.05) Significant after adjustment for O <sub>3</sub> , PM <sub>2.5</sub> , PM <sub>10</sub>
Henrotin et al. (2007) Period: 1994–2004 Location: Dijon, France	Outcome: Stroke HAs Design: Case-crossover Sample: 1,707 cases, age: ≥40 years	Avg time: 24 hours Mean: 0.60 ppm Range: 0–3.5 ppm	Increment: 10 µg/m <sup>3</sup> OR (95% CI), lag 0 day: Ischemic: 0.999 (0.997–1.001) Hemorrhagic: 1.000 (0.996–1.004) Other lags not significant
Maheswaran et al. (2005b) Period: 1994–1998 Location: Sheffield, United Kingdom	Outcome Stroke HAs Design: Ecological Sample: 11,407 HAs, age ≥45 years	Avg time: NR 20 <sup>th</sup> –80 <sup>th</sup> ‰: 0.34–0.40 ppm	Increment: 5 <sup>th</sup> vs. 1 <sup>st</sup> quintile RR (95% CI) adjusted for sex, age, deprivation, smoking: 1.11 (0.99–1.25)

**Table 3-7. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Hospital Admissions and Emergency Department Visits Related to Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Tsai et al. (2003) Period: 1997–2000 Location: Kaohsiung, Taiwan	Outcome: Stroke HAs Design: Case-crossover Statistical analyses: NR Sample: 23,192 HAs	Avg time: 24 hours Mean: 0.79 ppm Range: 0.24–1.72 ppm	Increment: 0.8 ppm OR (95% CI), lag 0–2 days: >20°C: 1.21 (1.14–1.28) <20°C: 1.77 (1.31–2.39) Significant OR when adjusted for PM <sub>10</sub> , SO <sub>2</sub> , or O <sub>3</sub>
Villeneuve et al. (2006a) Period: 1992–2002 Location: Edmonton, Canada	Outcome: Stroke EDVs Design: Case-crossover Sample: 12,422 EDVs, age ≥65 years	Avg time: 24 hours Mean: 0.8 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.5–1.0 ppm	Increment: 0.5 ppm OR (95% CI), lag 0–2 days: April–September: 1.32 (1.09, 1.60) (not significant for all seasons)
Wellenius et al. (2005a) Period: NR Location: 9 U.S. cities	Outcome: Stroke EDVs Design: Time series Sample: 155,503 EDVs, age ≥65 years	Avg time: NR Median: 1.02 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.73–1.44 ppm	Increment: 0.71 ppm % Change (95% CI) lag 0 days: Ischemic: 2.83 (1.23–4.46) Hemorrhagic: -1.61 (-4.79–1.68)
<b>Heart failure</b>			
Lee et al. (2007a) Period: 1996–2004 Location: Kaohsiung City, Taiwan	Outcome: CHF HAs Design: Case-crossover Sample Description: 13,475 HAs	Avg time: 24 hours Mean: 0.76 ppm Range: 0.14–1.72 ppm	Increment: 0.31 ppm OR (lag 0–2 days): ≥25 °C: 1.19 (1.09–1.31) <25 °C: 1.39 (1.24–1.54) Significant when adjusted for PM <sub>10</sub> , NO <sub>2</sub> , SO <sub>2</sub> , or O <sub>3</sub>
Symons et al. (2006) Period: 2002 Location: Baltimore, Maryland	Outcome: CHF HAs, symptom exacerbation Design: Case-crossover Sample: 125 cases, median age 70 years	Avg time: 24 hours Mean: 0.4 ppm Range: 0.1–1.0 ppm	Increment: 0.2 ppm OR (95% CI), lag 1 day: Avg: 0.90 (0.70–1.17) Cumulative: 0.82 (0.60–1.13) Other lags not significant
Wellenius et al. (2005b) Period: 1987–1999 Location: Pennsylvania	Outcome: CHF HAs Design: Case-crossover Sample: 54,019 HAs, age ≥65 years	Avg time: 24 hours Mean: 1.03 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.68–1.23 ppm	Increment: 0.55 ppm % Change (85% CI), lag 0 day: 4.55 (3.33–5.79) Significant when adjusted for NO <sub>2</sub> , PM <sub>10</sub> , NO <sub>2</sub> , SO <sub>2</sub> , or O <sub>3</sub>

**Table 3-7. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Hospital Admissions and Emergency Department Visits Related to Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Miscellaneous cardiovascular disease			
Ballester et al. (2001) Period: 1994–1996 Location: Valencia, Spain	Outcome: HD, CBVD EDVs Design: Time series Sample: >9,000 EDVs	Avg time: 1 hour Mean: 5.4 ppm Range: 0.5–15.5 ppm	Increment: 1 mg/m <sup>3</sup> RR (95% CI): All, lag 2: 1.0077 (0.9912–1.0138) HD, lag 1: 1.0092 (0.9945–1.0242) CBV, lag 1: 0.9874 (0.9646–1.0107)
Ballester et al. (2006) Period: 1995–1999 Location: 14 Cities in Spain	Outcome: HD, CVD HAs Design: Time series Sample: >250,000 HAs	Avg time: 8 hours Mean: 1.2–2.4 ppm 10 <sup>th</sup> % Range: 0.3–1.5 ppm 90 <sup>th</sup> % Range: 1.7–3.4 ppm	Increment: 1 mg/m <sup>3</sup> % Change (95% CI), lag 0–1 day: CVD: 2.06 (0.65–3.48) HD: 4.15 (1.31–7.08)
Barnett et al. (2006) Period: 1998–2001 Location: Australia, Auckland, New Zealand	Outcome: CVD HAs Design: Case-crossover Sample: age ≥15 years	Avg time: 8 hours Mean: 0.5–2.1 ppm Range: 0.0–7.9 ppm	Increment: 0.9 ppm % Change (95% CI) lag 0–1 day: Lags examined (days): 0–1 15–64 years CVD, <65 years: 1.2 (0.3–2.1) CVD, ≥65 years: 2.2 (0.9–3.4) Significant when adjusted for NO <sub>2</sub> or PM <sub>10</sub>
Chang et al. (2005) Period: 1997–2001 Location: Taipei, Taiwan	Outcome: CVD HAs Design: Case-crossover Sample 74,509 HAs	Avg time: 24 hours Mean: 1.37 ppm Range: 0.37–3.66 ppm	Increment: 0.49 ppm OR (95% CI) lag 0–2 day: ≥20°C: 1.090 (1.064–1.118) <20°C: 0.984 (0.927–1.044) Significant when adjusted for PM <sub>10</sub> , NO <sub>2</sub> , SO <sub>2</sub> , or O <sub>3</sub>
Fung et al. (2005) Period: 1995–2000 Location: Windsor, Ontario, Canada	Outcome: CVD HAs Design: Time series Sample: 11,632 HAs	Avg time: 24 hours Mean: 1.3 ppm Range: 0.0–11.8 ppm	Increment: 1.2 ppm % Change (95% CI), lag 0–2 day: <65 years: -0.5 (-6.7–6.0) ≥65 years: 2.8 (-1.1–7.0)
Jalaludin et al. (2006) Period: 1997–2001 Location: Sydney, Australia	Outcome: CVD EDVs Design: Time series Sample: >100,000 EDVs, age ≥65 years	Avg time: 8 hours Mean 0.82 ppm Range: 0.02–4.63 ppm	Increment: 0.69 ppm % Change (95% CI), lag 0–1 day: 2.35 (1.39–3.32) Significant when adjusted for PM <sub>10</sub> , NO <sub>2</sub> , SO <sub>2</sub> , or O <sub>3</sub>

**Table 3-7. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Hospital Admissions and Emergency Department Visits Related to Cardiovascular Disease**

Study	Design features	CO exposure	Effect size
Koken et al. (2003) Period: 1993–1997 Location: Denver, Colorado	Outcome: CVD HAs Design: Time series Sample: >4000 HAs, age 65 years	Avg time: 24 hours Mean: 0.9 ppm Range: 0.3, 1.6 ppm	Increment: 0.3 ppm % Change (95% CI) lag 3 day: CHF: 10.5 (0.1–22.0) CO not significantly associated with other CVD categories.
Linn et al. (2000) Period: 1992–1995 Location: Los Angeles, California	Outcome: CVD, CBVD HAs Design: Time series Sample: >500,000 HAs, age >30 years	Avg time: 24 hours Mean 1.0–2.0 ppm Range: 0.2–5.3 ppm	Increment: 1 ppm Poisson regression coefficient (SE), lag 0 day: CVD, all: 0.032 (0.003) (p<0.05) Significant for AR, CBVD, CHF, MI, stroke p<0.05
Metzger et al. (2004) Study: 1993–2000 Location: Atlanta, Georgia	Outcome: Cardiovascular: CVD EDVs Design: Time series Sample: 4,407,535 EDVs	Avg time: 1 hour Median: 1.5 ppm 10 <sup>th</sup> –90 <sup>th</sup> ‰: 0.5–3.4 ppm	Increment: 1 ppm RR (95% CI), lag 0–2 day: All CVD: 1.017 (1.008–1.027) PVD/CBVD: 1.031 (1.010–1.052) Not significant for AR, CHF, IHD categories
Peel et al. (2007) Period: 1993–2000 Location: Atlanta, Georgia	Outcome: CVD EDVs Design: Case-crossover Sample: 4,407,535 EDVs	Avg time: 1 hour Mean: 1.8 ppm SD: 1.2 ppm	Increment: 1.2 ppm OR (95% CI), lag 0–2 day: 1.020 (1.010–10.030) ORs for specific CVD categories tended to increase with co-diagnoses of HT, AR, CHF, COPD
Yang et al. (2004a) Period: 1997–2000 Location: Kaohsiung City, Taiwan	Outcome: CVD HAs Design: Case-crossover Sample: 29,661 HAs	Avg time: 24 hours Mean: 0.79 ppm Range 0.24–1.72 ppm	Increment: 0.28 ppm OR (95% CI), lag 0–2 day: ≥25 °C: 1.264 (1.205–1.326) <25 °C: 1.448 (1.357–1.545) Significant when adjusted for PM <sub>10</sub> , NO <sub>2</sub> , SO <sub>2</sub> , or O <sub>3</sub>

AP = angina pectoris; AR = arrhythmia; Avg = average; CAR = cardiac arrhythmia; CBCD = cerebrovascular disease; CHF = congestive heart failure; CI = confidence interval; CO = carbon monoxide; COPD = chronic obstructive pulmonary disease; CVD = cardiovascular disease; EDV = emergency department visit; HA = hospital admission; HF = heart failure; HT = hypertension; IHD = ischemic heart disease; MI = myocardial infarction; NR = not reported; OR = odds ratio; PVD = peripheral vascular disease; RR = relative risk; SD = standard deviation; SE = standard error

## 3. HEALTH EFFECTS

concentration, for 2-hour period prior to infarct, was 1.22 (95% CI: 0.89–1.67); the mean carbon monoxide exposure concentration was 1.09 (5<sup>th</sup>–95<sup>th</sup> percentile range: 0.49–1.78 ppm, predicted steady-state COHb%:  $\leq 0.55$ ).

Possible associations between carbon monoxide exposure (and other air pollutants; e.g., PM, NO<sub>2</sub>, O<sub>3</sub>) and ischemic heart disease have also been explored in studies of hospital admissions and/or hospital emergency department visits (D'Ippoliti et al. 2003; Hosseinpoor et al. 2005; Lanki et al. 2006; Lee et al. 2003b; Maheswaran et al. 2005b; Mann et al. 2002; Szyszkowicz 2007; von Klot et al. 2005). Although results have been mixed, most studies have found significant associations between increasing carbon monoxide exposure concentration and ischemic heart disease-related admissions or emergency room visits (Table 3-7). These included increased odds (or relative risk) for hospital admissions for angina pectoris, myocardial infarction, and aggregated ischemic heart disease categories (D'Ippoliti et al. 2003; Hosseinpoor et al. 2005; Lanki et al. 2006; Lee et al. 2003b; Mann et al. 2002; Szyszkowicz 2007; von Klot et al. 2005). Mean air carbon monoxide concentrations in these studies ranged from approximately 0.4–2 ppm, with the upper percentiles (i.e.  $\geq 75^{\text{th}}$ ) ranged from approximately 1 to 5 ppm, although one study reported a mean of 10.8 mg/m<sup>3</sup> and maximum of 57.9 mg/m<sup>3</sup> (approximately 50 ppm; Hosseinpoor et al. 2005). Associations with carbon monoxide tended to be stronger for more elderly males than for females or other age strata (e.g., Lee et al. 2003b; Szyszkowicz 2007). Risks were higher when ischemic heart disease was accompanied by a diagnosis of arrhythmia or congestive heart failure (Mann et al. 2002; see further discussion of carbon monoxide associations with congestive heart failure, below). Lee et al. (2003b) found a significant association (1997–1999) between ischemic heart disease-related hospital admissions in Seoul, Korea, and carbon monoxide exposure concentrations among males  $\geq 64$  years of age, but not for females or other age strata. A 1 ppm increase in carbon monoxide concentration was associated with a relative risk of 1.07 (95% CI: 1.01–1.1) for hospital admission. The mean daily 1-hour maximum carbon monoxide concentration was 1.8 ppm (25<sup>th</sup>–75<sup>th</sup> percentile range: 1.2–2.2 ppm). The association with carbon monoxide exposure was not significant after adjustment for exposure to PM<sub>10</sub> in a two-pollutant model (1.04, 95% CI: 0.98–1.11). Two-pollutant models with carbon monoxide and other gaseous pollutants were not reported, although significant associations were found when NO<sub>2</sub>, O<sub>3</sub>, or SO<sub>2</sub> were considered alone. Szyszkowicz (2007) examined records of 4,979 emergency room visits in Montreal, Canada for the period 1997–2002. The percent increase in visits related to ischemic heart disease for a 0.2 ppm increase in carbon monoxide concentration was 5.4% (95% CI: 2.3–8.5) for all patients, 7.5% (2.6–12.6) for males  $\geq 64$  years, and 2.4 (-3.0–0.0) for females  $\geq 64$  years. The mean carbon monoxide exposure concentration was 0.5 ppm (range: 0.1–3.1 ppm). Two other studies examined ischemic heart disease end points in multiple-pollutant models. Hosseinpoor et al. (2005)

## 3. HEALTH EFFECTS

found that the relative risk for hospital admissions in related for angina pectoris (1,826 cases; Tehran, Iran; 1996–2001) was 1.044 (95% CI: 1.000–0.089) when air carbon monoxide concentrations were considered in a carbon monoxide model, and 1.0093 (95% CI: 1.0036–1.0151) when the model was adjusted for PM<sub>10</sub>, O<sub>3</sub>, and NO<sub>2</sub>. Air carbon monoxide concentrations examined in this study were relatively high, compared to other studies (mean 10.8 ppm; range: 1.6–58 ppm). A study conducted in Rome, Italy examined records of 6,351 hospital admissions (1995–1997) for myocardial infarction and found a significant association between increasing air carbon monoxide concentration and increasing odds of hospital admission (odds ratio: 1.004, 95% CI: 1.000–0.089), which did not persist when adjusted for total suspended solids (D'Ippoliti et al. 2003). The mean air carbon monoxide concentration was 3.8 ppm (25<sup>th</sup>–75<sup>th</sup> percentile range: 2.4–3.7 ppm).

*Studies of Stroke.* Possible associations between ambient air carbon monoxide concentrations and stroke have been examined in studies of hospital admissions and emergency room visits (Chan et al. 2006; Henrotin et al. 2007; Maheswaran et al. 2005b; Tsai et al. 2003b; Villeneuve et al. 2006a; Wellenius et al. 2005a; Table 3-7). Although results of these studies have been mixed, several studies have found significant associations between stroke admissions and/or emergency room visits, and increasing air carbon monoxide concentrations (Chan et al. 2006; Tsai et al. 2003b; Villeneuve et al. 2006a; Wellenius et al. 2005a). Studies that explored two-pollutant models found that the associations persisted after adjustment for PM<sub>10</sub>, SO<sub>2</sub>, or O<sub>3</sub> (Chan et al. 2006; Tsai et al. 2003b). Mean air carbon monoxide concentrations in these studies ranged from 0.4 to 1.7 ppm and the highest reported value was <5 ppm. The largest study was a time-series analysis of ambient air carbon monoxide concentrations and 155,503 records of stroke-related emergency room visits in nine U.S. cities (Wellenius et al. 2005a). The study found that a 0.71 ppm increase in ambient air carbon monoxide concentration was associated with a 2.83% (95% CI: 1.23–4.46) increase in daily rate of emergency room visits for ischemic stroke. Chan et al. (2006) conducted a time-series analysis of ambient air carbon monoxide concentrations in Taipei, China (1997–2002) and emergency room visits related to cerebrovascular disease, including stroke (n=7,341), and found a significant odds ratio for a 0.8 ppm increase in air carbon monoxide concentration, for all cerebrovascular disease (1.03, 95% CI: 1.01–1.06) and stroke (1.03, 95% CI: 1.01–1.05), which remained significant after adjusting for O<sub>3</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> or PM<sub>2.5</sub> (1.034, 95% CI: 1.001–1.067). A larger study conducted in Kaohsiung, China (1997–2000) examined records of 23,192 hospital visits for stroke in a case-crossover design and found significant odds ratios for a 0.8 ppm increase in carbon monoxide concentration (1.77, 95% CI: 1.31–2.39), which also persisted after adjustment for SO<sub>2</sub>, O<sub>3</sub>, or PM<sub>10</sub> (Tsai et al. 2003b).

## 3. HEALTH EFFECTS

*Studies of Other Cardiovascular Outcomes.* Two case-control (i.e., case crossover) studies conducted in Seattle, Washington have examined possible associations between ambient carbon monoxide concentrations and cardiac arrest did not find significant associations (Levy et al. 2001; Sullivan et al. 2003; Table 3-6). The mean carbon monoxide exposure concentration (both studies) was 1.7 ppm, with the highest value reported as 5.9 ppm. The study reported by Levy et al. (2001) included 362 cases out-of-hospital cardiac arrests (1988–1994); the relative risk for a 1-day lag between exposure estimate and outcome was 0.88 (95% CI: 0.83–1.18). The Sullivan et al. (2003) study examined 1,542 cases of out-of-hospital cardiac arrest cases (1985–1994); the odds ratio for a 1.02 ppm increase in carbon monoxide concentration (1 day lag) was 0.97 (95% CI: 0.87–1.08).

Possible associations between air carbon monoxide concentrations and cardiac failure (primarily, congestive heart failure) have also been examined in studies of hospital admissions (Lee et al. 2007a; Symons et al. 2006; Wellenius et al. 2005b; Table 3-6). The larger of the these studies examined 13,475 hospital admissions for congestive heart failure in Kaohsiung, China (1996–2004) and found a significant odds ratios for hospital admission associated with a 0.31 ppm increase in air carbon monoxide concentration (odds ratio: 1.39, 95% CI: 1.24–1.54). The odds ratio remained significant in two-pollutant models that adjusted for PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, or O<sub>3</sub>.

Studies of possible associations between carbon monoxide exposure and abnormal blood pressure have not found significant associations (Ibald-Mulli et al. 2001; Zanobetti et al. 2004b; Table 3-6). The Ibald-Mulli et al. (2001) study included 2,607 men and women (age range 25–64 years) in the Augsburg, Germany area; the mean carbon monoxide exposure concentration was approximately 3.6 ppm and the upper end of the range was 7.2 ppm. The estimated increment in systolic blood pressure for 4.9 ppm increase in carbon monoxide concentration was a 1.06 (95% CI: -0.17–2.29). The Zanobetti et al. (2004b) study measured blood pressure in repeated examinations of 62 subjects, and found no association between blood pressure and carbon monoxide exposure concentrations (90<sup>th</sup> percentile concentration <1.2 ppm).

Several studies have evaluated more generic cardiovascular disease outcomes, for example, hospital admissions or emergency room visits for any cardiovascular disease (Ballester et al. 2001, 2006; Barnett et al. 2006; Chang et al. 2005; Fung et al. 2005; Jalaludin et al. 2006; Koken et al. 2003; Linn et al. 2000; Metzger et al. 2004; Peel et al. 2007; Yang et al. 2004a; Table 3-7). Here again, although results have been mixed, most studies have found significant associations between increased rate of hospital admissions or emergency room visits and increased air carbon monoxide concentrations. The association

## 3. HEALTH EFFECTS

was stronger for older subjects (e.g.,  $\geq 64$  years, Barnett et al. 2006) and in several studies that examined multi-pollutant models, the association persisted after adjustment for PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, or O<sub>3</sub> (Chang et al. 2005; Jalaludin et al. 2006; Yang et al. 2004a). Mean air carbon monoxide concentrations in these studies ranged from 0.8 to 6 ppm, with the highest value <18 ppm.

***Animal Studies.*** Although clinical studies in patients with coronary artery disease provide compelling evidence of adverse cardiovascular effects of acute-duration exposure to carbon monoxide, these studies are limited with respect to exposure conditions and outcomes assessed. Studies in animals, however, have investigated adverse cardiovascular effects of carbon monoxide exposure over a much wider range of exposure conditions (e.g., exposure concentration and duration) and have evaluated additional outcome measures that are not possible to assess in humans. In general, animal studies evaluated carbon monoxide exposures resulting in higher COHb levels than those evaluated in clinical studies in patients and healthy volunteers. Results of animal studies provide evidence of adverse cardiovascular effects of carbon monoxide exposure, including compensatory alterations in hemodynamics, cardiac hypertrophy, cardiac arrhythmias, and possibly atherosclerosis. For most outcomes evaluated in animal studies, conflicting evidence has been reported. Although the reasons for different results have not been established, differences in species sensitivity, exposure regimens, and experimental protocols are possible factors. Animal studies published prior to 2000 have been reviewed in detail by EPA (1991, 2000); therefore, these reviews largely form the basis of the following discussions for animals studies published prior to 2000.

Studies investigating the effects of acute-duration carbon monoxide exposure on hemodynamics have been conducted in several animal species, including monkeys, dogs, rats, and rabbits, under exposure conditions producing blood COHb levels ranging from 6.2 to 70% (EPA 1991, 2000). Results show that brief exposure (from a few minutes to approximately 3 hours) to carbon monoxide at concentrations of 80–20,000 ppm produces alterations in hemodynamics that are consistent with COHb-induced hypoxia and responsive compensatory mechanisms (e.g., vasodilation and increased cardiovascular output), including increased coronary blood flow, decreased myocardial O<sub>2</sub> consumption, increased heart rate, and alterations in blood flow to various vascular beds (e.g., cerebral, limb muscular). Generally, carbon monoxide-induced effects on hemodynamics were observed at blood COHb levels  $\geq 7.5\%$ . With longer exposure durations, COHb-induced tissue hypoxia and subsequent compensatory increases in heart rate and cardiac workload lead to the development of cardiomyopathy. Several studies confirm carbon monoxide-induced cardiomegaly in rats and rabbits exposed to 160–11,000 ppm carbon monoxide (COHb 12–58%) for intermediate durations (14 days to ~6 months) (EPA 1991, 2000). In rats exposed

### 3. HEALTH EFFECTS

chronically to 200 ppm carbon monoxide (COHb 14.7%) for 20 hours/day for 72 weeks, left and right ventricle weights increased by 20% ( $p < 0.001$ ) and 14% ( $p < 0.001$ ), respectively (Sorhaug et al. 2006). Studies evaluating developmental cardiovascular effects are reviewed in Section 3.2.6 (Developmental Effects).

Alterations in cardiac rhythm, possibly due to disturbances in cardiac conduction, have been observed in animals exposed to carbon monoxide for acute to intermediate durations (EPA 1991, 2000). Under conditions of simulated myocardial ischemia (i.e. coronary artery ligation), acute exposure (up to 15 minutes) of dogs to up to 5,000 ppm carbon monoxide (COHb 4.9–15%) enhanced myocardial ischemia, as indicated by ST-segment alterations, and severity of myocardial injury. Results of a recent study in rats exposed to 50 ppm carbon monoxide (COHb not reported) for 1 week indicate that rats with pulmonary hypertension may be more sensitive to carbon monoxide-induced myocardial ischemia (Gautier et al. 2007). However, in other studies simulating myocardial injury in dogs exposed to 100–500 ppm carbon monoxide for 6–120 minutes (COHb 5.1–20%), cardiac arrhythmias or alterations in cardiac conduction speed were not observed. In studies evaluating intermediate-duration (6–24 weeks) exposure of dogs and monkeys, 100 ppm carbon monoxide (COHb 2.6–12%) induced cardiac arrhythmias, including R-wave depression, ST-segment elevation, increased T-wave, pre-ventricular contractions, and reduced threshold for stimulus-induced ventricular fibrillation. In contrast, no evidence of cardiac arrhythmias was observed in dogs exposed continuously to 50 ppm carbon monoxide (COHb 7.3%) for 3 months or in monkeys exposed intermittently to 500 ppm (COHb 21.6%) for 14 months.

Conflicting evidence has been reported regarding the potential for carbon monoxide exposure to induce or enhance atherosclerosis (EPA 1991, 2000). Results of most studies in animals (monkeys, baboons, rabbits, and pigs) fed normal diets were negative for inducing or enhancing atherosclerosis at carbon monoxide concentrations of 50–400 ppm (COHb ~20%) for exposure durations up to 14 months. However, atherosclerosis was increased in monkeys continuously exposed to 200 ppm carbon monoxide (COHb 20.6%) for 2 weeks (Thomsen 1974) and in rabbits and monkeys fed high cholesterol diets exposed to 100–300 ppm carbon monoxide (COHb 9–33%) for 2–7 months. Based on the weight-of-evidence, EPA (1991, 2000) concluded that the available data do not strongly support a relationship between atherogenic effects and carbon monoxide exposure.

**Gastrointestinal Effects.** Studies evaluating gastrointestinal effects of exposure to low levels of carbon monoxide (i.e., producing blood COHb levels <20%) were not identified. Although the gastrointestinal tract has not been identified as a specific target organ for carbon monoxide-induced

## 3. HEALTH EFFECTS

toxicity, exposure to carbon monoxide at levels producing hypoxia would be expected to affect any tissue, in particular those tissues with high O<sub>2</sub> utilization requirements (e.g., brain, liver, kidney, heart, small intestine).

**Hematological Effects.** Hematological effects of carbon monoxide include compensatory responses to tissue hypoxia resulting from binding of carbon monoxide to Hb (e.g., increased blood volume, Hb, hematocrit, and erythrocyte count and volume). Possible associations between ambient air carbon monoxide concentrations and hematologic biomarkers of coagulation and inflammation have been examined in epidemiological studies. Collectively, these studies provide some evidence for associations that may reflect effects of carbon monoxide in modulating these systems.

*Hypoxia and Related Compensatory Responses.* Inhalation exposure to carbon monoxide results in the formation of COHb, a stable complex between carbon monoxide and Hb. Because carbon monoxide has a much higher affinity for Hb than O<sub>2</sub> (equilibrium constant for carbon monoxide is >200 times that of O<sub>2</sub>), O<sub>2</sub> is displaced from Hb by relatively low partial pressures of carbon monoxide. Binding of carbon monoxide to Hb has two effects that contribute to impaired O<sub>2</sub> delivery to tissues: (1) in the presence to carbon monoxide, the amount of O<sub>2</sub> that can be stored on Hb for delivery to tissues decreases; and (2) binding of carbon monoxide to Hb impairs release of O<sub>2</sub> from Hb for its diffusion into tissues (see Section 3.4.3 for further discussion of effects of carbon monoxide on O<sub>2</sub> dissociation from Hb). At sufficient levels of COHb, the combined effects of impaired O<sub>2</sub> storage and transport results in tissue hypoxia, the principal mechanism of many adverse effects of carbon monoxide exposure.

To maintain O<sub>2</sub> delivery to tissues under conditions of hypoxia, compensatory hematological responses (e.g., increased erythrocyte count, hematocrit, and Hb) occur. The dose-response relationship for carbon monoxide-induced compensatory hematological responses in humans has not been established. Studies in animals show that compensatory hematological effects occur in response to the COHb-induced reduction in O<sub>2</sub> delivery to tissues. Compensatory hematological effects, including increased blood volume, Hb, hematocrit, and erythrocyte count and volume, have been observed following acute- and intermediate-duration carbon monoxide exposure (Davidson and Penney 1988; Penney 1988; Penney et al. 1974a, 1974b; WHO 1999). Continuous exposure of rats to 500 ppm carbon monoxide (COHb 40%) for 5–42 days produced duration-dependent increases in hematocrit, blood Hb, erythrocyte count, and mean cell Hb concentration; mean cell volume was increased after 42 days of exposure (Davidson and Penney 1988). The largest increases (approximately 45–55%) were observed for hematocrit, blood Hb, and erythrocyte count, with increases reaching maximum levels after 30 and 42 days of exposure. Similarly,

## 3. HEALTH EFFECTS

blood Hb concentration increased by 70% in rats exposed continuously to 500 ppm carbon monoxide (COHb approximately 12% after 42 days) for up to 42 days, with the most rapid increase occurring within the first 21 days of exposure (Penney et al. 1974b). A small, but statistically significant increase in Hb concentration (6.0%;  $p < 0.001$ ) was observed in rats exposed to 100 ppm carbon monoxide (COHb 9.25%) for 30 days (Penney et al. 1974b). In rats exposed to 200–1,300 ppm carbon monoxide (COHb not reported) for 10–17 days, dose-related increases in hematocrit and blood volume were observed (Penney et al. 1988).

***Epidemiological Studies of Blood Markers Related to Coagulation.*** Studies of carbon monoxide mechanisms of action have provided evidence that carbon monoxide may participate in the regulation of vascular thrombi formation and inflammation (see Section 3.5.2, Mechanisms of Toxicity). Possible associations between ambient air carbon monoxide concentrations and biomarkers of coagulation and inflammation have been examined in epidemiological studies (Baccarelli et al. 2007; Liao et al. 2005; Pekkanen et al. 2000; Ruckerl et al. 2006, 2007; Steinvil et al. 2008). Biomarkers that have been examined include inflammation markers, C-reactive protein (CRP), serum amyloid A (SAA), and WBC count; cell adhesion markers, E-selectin, von Willebrand factor, antigen (vWF), ICAM-1; and coagulation markers, fibrinogen, factor VII (FVII), prothrombin fragment 1+2, prothrombin time (PT), and activated partial thromboplastin time (APTT). Collectively, these studies provide some evidence for associations between ambient air carbon monoxide concentration and changes in hematological indicators of inflammation and coagulation. However, interpretation of the results of these studies, in particular, quantitative estimates of the effect magnitudes on outcomes, is limited by several factors: (1) reliance on area monitoring for estimating exposure levels; (2) uncertainty in knowledge of temporal correspondence between monitored exposure levels and outcomes; (3) relatively strong correlations between ambient air carbon monoxide concentrations and other air quality variables that can affect respiratory function; and (4) relatively low carbon monoxide exposures studied. Mean values of carbon monoxide exposure concentrations have ranged from 0.3 to 3 ppm with the highest values <12 ppm.

Major findings from the body of reported epidemiological studies hematological outcomes are summarized below. Although most of the studies explored various time lags between monitored air carbon monoxide concentrations and outcomes, as well as various sample strata, for the sake of brevity, only those indicative of the strongest associations to carbon monoxide are presented. Where co-pollutant models have been explored, these results are also presented.

## 3. HEALTH EFFECTS

A cohort study conducted in London, United Kingdom (1991–1993) examined plasma fibrinogen concentrations in adult office workers (n=7,205) and found a significant association between increasing air carbon monoxide concentrations (mean 1.4 mg/m<sup>3</sup>, maximum: 9 mg/m<sup>3</sup>) and increasing plasma fibrinogen concentrations (Pekkanen et al. 2000). For a 3-day lag between measured carbon monoxide concentration and fibrinogen assessment, the odds ratio for a plasma fibrinogen exceeding 3.19 g/L was 1.22 (p<0.02) per 1.6 mg/m<sup>3</sup> increase in air carbon monoxide concentration. Increasing air concentration of NO<sub>2</sub> was also associated with increasing plasma fibrinogen concentration (odds ratio: 1.22, p<0.01), and when both were considered in two-pollutant models, the associations were no longer significant, reflecting the relative high correlation between air carbon monoxide and NO<sub>2</sub> concentrations (r=0.81). When stratified by gender, the carbon monoxide association for a 1-day lag (3-day lag data not reported) was significant in males, but not in females (Steinvil et al. 2008). A smaller cohort study conducted in Tel-Aviv, Israel (2003–2006, n=3,659, mean age: 46 years) also found a significant association between increases in air carbon monoxide concentration and decreasing plasma fibrinogen concentration among males, but not among females. The largest effect was estimated for the 1-week air carbon monoxide average (mean: 0.8 ppm, 75<sup>th</sup> percentile: 1 ppm), for which a 0.3 ppm increase in air carbon monoxide concentration was associated with a 7.7 mg/dL (95% CI: -12.1, -3.3) decrease in plasma fibrinogen in males, with no association among females (-1.6 mg/dL, 95% CI: -7.3, 4.1). Increasing air carbon monoxide concentration was also associated with a significant decrease in WBCs among males (-158 cell/μL, 95% CI: -298, -18), but not females (-182 cell/μL, 95% CI: -281, 61). However, as in the Pekkanen et al. (2000) study, the association between air NO<sub>2</sub> and fibrinogen was stronger than that of carbon monoxide and when included in two-pollutant models, the association with carbon monoxide was no longer significant. Significant associations with air carbon monoxide concentrations were not found for CRP.

A cohort study conducted in three U.S. locations (1996–1998) examined various blood biomarkers of coagulation and inflammation (i.e., fibrinogen, factor VIII-C, von Willebrand factor, serum albumin, WBC) in subjects enrolled in an *Atherosclerosis Risk in Communities Study* (n=10,208) and found a significant association between increasing air carbon monoxide concentrations (mean 1.4 ppm, SD: 0.6) and decreasing serum albumin concentrations (Liao et al. 2005). A 0.6 ppm increase in air carbon monoxide concentration corresponded to a 0.018 g/dL (SE: 0.003, p<0.01) decrease in serum albumin concentration. Other blood biomarkers evaluated were not significantly associated with air carbon monoxide concentration; results of multi-pollutant models were not reported.

## 3. HEALTH EFFECTS

A panel study conducted in Efert, German (2000–2001) examined air carbon monoxide concentrations (mean: 0.52 mg/m<sup>3</sup>, range: 0.11–1.93) and several blood biomarkers of coagulation and inflammation in 57 nonsmoking male coronary heart disease patients (age range: 51–77 years; Ruckerl et al. 2006). Biomarkers evaluated included inflammation markers, CRP and SAA; cell adhesion markers, E-selectin, vWF, ICAM-1; and coagulation markers, fibrinogen, FVII, and prothrombin fragment 1+2. Odds ratios for increase in air carbon monoxide concentration of 0.27 mg/m<sup>3</sup> (2-day lag) were: increasing CRP, 1.5 (95% CI: 1.1–2.1); increasing ICAM-1, 1.7 (95% CI: 1.3–2.3); decreasing FVII, 2.8 (95% CI: 5.1–0.4). Significant associations were not found for other blood biomarkers, including fibrinogen. A larger cohort study did not find significant associations between ambient air carbon monoxide concentrations (mean range: 0.3–1.48 mg/m<sup>3</sup>) and plasma interleukin-6 (IL-6), CRP, or fibrinogen among myocardial infarction survivors (n=1,003, age range: 31–87 years) in six European cities (2003–2004, Ruckerl et al. 2007).

A panel study conducted in Milan, Italy (1995–2005) evaluated associations (mean range: 1.14–3.11 ppm; highest value: 11.4 ppm) between air carbon monoxide concentrations and several blood coagulation, using prothrombin time (PT) and activated partial thromboplastin time (APTT) as outcome measures in 1,218 healthy adolescents and adults (age 11–84 years; Baccarelli et al. 2007). Increasing ambient air carbon monoxide concentration was associated with a decrease in PT ( $\beta$  coefficient: -0.11; 95% CI: -0.18, -0.05). Associations were also significant for PM<sub>10</sub> and NO<sub>2</sub>; however, results from multi-pollutant models were not reported. Significant associations with carbon monoxide were not found for other blood biomarkers evaluated in the study, including: carbon monoxide had no effect on fibrinogen, functional antithrombin, functional protein C, protein C antigen, functional protein S, or free protein S.

**Musculoskeletal Effects.** Studies evaluating effects of exposure to low levels of carbon monoxide (i.e., producing blood COHb levels <20%) on bone in humans or animals were not identified. In mice continuously exposed to 2,400 ppm for 180 days, increased bone mass was observed in parietal bone, sternum, lumbar vertebrae, and ribs (but not femur), and the marrow cavities of ribs, parietal bone, and femurs were expanded (Zebro et al. 1983).

Studies evaluating effects of exposure to low levels of carbon monoxide (i.e., producing blood COHb levels <20%) on muscle tissue in humans or animals were not identified. Rhabdomyolysis and myonecrosis have been reported in cases of carbon monoxide poisoning (Florkowski et al. 1992; Herman et al. 1988; Kuska et al. 1980; Wolff 1994). Damage to muscle tissue is consistent with hypoxia due to

## 3. HEALTH EFFECTS

binding of carbon monoxide to Hb and myoglobin and with inhibition of cellular aerobic metabolism due to binding of carbon monoxide to intracellular cytochrome oxidases (Wolff 1994).

**Hepatic Effects.** Studies evaluating hepatic effects of exposure to low levels of carbon monoxide (i.e., producing blood COHb levels <20%) were not identified. Although the liver has not been identified as a specific target organ for carbon monoxide-induced toxicity, exposure to carbon monoxide at levels producing hypoxia would be expected to affect any tissue, in particular those tissues with high O<sub>2</sub> utilization requirements (e.g., brain, liver, kidney, heart, small intestine).

**Renal Effects.** Studies evaluating renal effects of exposure to low levels of carbon monoxide (i.e., producing blood COHb levels <20%) were not identified. Although the kidney has not been identified as a specific target organ for carbon monoxide-induced toxicity, exposure to carbon monoxide at levels producing hypoxia would be expected to affect any tissue, in particular those tissues with high O<sub>2</sub> utilization requirements (e.g., brain, liver, kidney, heart, small intestine). Acute renal failure secondary to rhabdomyolysis has been observed in cases of acute carbon monoxide poisoning (Florkowski et al. 1992; Kuska et al. 1980; WHO 1999; Wolff 1994).

**Endocrine Effects.** Studies evaluating endocrine effects of exposure to low levels of carbon monoxide (i.e., producing blood COHb levels <20%) were not identified. Although the endocrine system has not been identified as a specific target for carbon monoxide-induced toxicity, exposure to carbon monoxide at levels producing hypoxia would be expected to affect any tissue.

**Dermal Effects.** Studies evaluating dermal effects of exposure to low levels of carbon monoxide (i.e., producing blood COHb levels <20%) were not identified. Skin lesions have been observed in patients with severe carbon monoxide poisoning (Myers et al. 1985; Torne et al. 1991). Lesions are primarily observed in pressure areas and are characterized by blisters, subepidermal vesicles, and sweat gland necrosis.

**Ocular Effects.** No information was identified to indicate that carbon monoxide is an ocular irritant in humans or animals. Visual field deficits, temporary or permanent blindness, retinal venous congestion, retinal hemorrhage, papilledema, and optic atrophy have been associated with severe carbon monoxide poisoning in humans (Choi 2001).

### 3. HEALTH EFFECTS

#### 3.2.3 Immunological and Lymphoreticular Effects

Little information is available regarding the potential of carbon monoxide exposure to produce adverse immunological effects. Epidemiological studies blood biomarkers of immunological status are summarized in the section on Hematologic Effects. Immunological biomarkers examined in these studies have included CRP, serum amyloid A (SAA), and WBC; however, other markers that can be influenced by the immune system have also been evaluated, including cell adhesion markers such as E-selectin, vWF, ICAM-1; and coagulation markers such as fibrinogen, FVII, prothrombin fragment 1+2, PT, and APTT. Although results of these studies have been mixed, collectively, they provide some evidence for associations between ambient air carbon monoxide concentration and changes in hematological indicators of inflammation and coagulation. However, these studies do not distinguish between possible effects of carbon monoxide and/or other air pollutants on the immune system from indirect effects of carbon monoxide on blood biomarkers. Mechanistic studies have revealed evidence that carbon monoxide may participate in the regulation of immune function (see Section 3.5.2, Mechanisms of Toxicity).

Results of a study in guinea pigs investigating the effects of repeated brief bursts of carbon monoxide on systemic immune function were equivocal (Snella and Rylander 1979). Exposure of guinea pigs to 3 minute bursts of 10,000 ppm carbon monoxide 12 times/day for 4 weeks resulted in an 88% reduction in the number of splenic plaque-forming cells, compared to unexposed controls; however, the reduction did not reach statistical significance. Studies evaluating effects of gestational exposure on the developing immune system function are reviewed in Section 3.2.6 (Developmental Effects) (Giustino et al. 1993, 1994).

#### 3.2.4 Neurological Effects

The literature reporting adverse nervous system effects, specifically central nervous system effects, of acute exposure of humans to carbon monoxide is extensive. This literature includes case studies of carbon monoxide poisoning, clinical studies of neurobehavioral effects of controlled exposures in humans, and experimental studies in a variety of animal models (e.g., dogs, rodents, nonhuman primates). Much of this literature has been summarized and analyzed in recent reviews and meta-analyses (Benignus et al. 1990; EPA 1991, 2000; Kao and Nañagas 2006; Raub et al. 2000; Raub and Benignus 2002; WHO 1999). This section summarizes the major findings from these analyses. Although the literature on central nervous system effects of acute experimental and accidental exposures of humans to carbon monoxide is extensive, reports of effects of longer duration exposure of humans are not available. Effects

## 3. HEALTH EFFECTS

of gestational and perinatal exposure on the developing nervous system are discussed in Section 3.2.6 (Developmental Effects).

Acute exposure to high levels of carbon monoxide produces symptoms of central nervous system toxicity (EPA 1991, 2000; Ernst and Zibrak 1998; Raub and Benignus 2002; WHO 1999). Symptoms of central nervous system toxicity include headache, dizziness, drowsiness, weakness, nausea, vomiting, confusion, disorientation, irritability, visual disturbances, convulsions, and coma. Symptoms vary depending upon the degree of exposure; headache and dizziness are the most commonly reported symptoms (Dolan 1985; Ernst and Zibrak 1998). Neuroimaging evaluations of patients with acute carbon monoxide poisoning show lesions of the basal ganglia (primarily of the globus pallidus) and white matter (Hopkins et al. 2006; Kao and Nañagas 2006; Lo et al. 2007; Parkinson et al. 2002). Following acute-onset effects, delayed development of neuropsychiatric impairment may occur within 1–4 weeks of exposure, with symptoms including inappropriate euphoria, impaired judgment, poor concentration, memory loss, cognitive and personality changes, and psychosis. Delayed neuropsychiatric impairment has been estimated to occur in up to 68% of patients with acute carbon monoxide poisoning (Ernst and Zibrak 1998; Kao and Nañagas 2006; Raub and Benignus 2002; Raub et al. 2000; WHO 1999; Wolf et al. 2008). The relationship between initial symptom severity and the likelihood of developing of delayed neuropsychiatric impairment has not been established. A cohort analysis of 256 patients with carbon monoxide poisoning suggests that initial symptom severity does not correlate to delayed-onset effects (Chambers et al. 2008). However, based on a review of available data, Kao and Nañagas (2006) concluded that patients with more severe initial symptoms are more likely to develop delayed neuropsychiatric impairment.

Mechanisms of acute and delayed adverse nervous system effects produced by carbon monoxide have not been conclusively established. Tissue hypoxia secondary to COHb formation may be a contributing factor; particularly in association with high levels of blood COHb (>60%); however, direct cellular effects of carbon monoxide (e.g., ATP depletion, excitotoxicity, oxidative stress, immunological responses) and postischemic reperfusion injury may also contribute to neurotoxicity (see Section 3.5.2, Mechanisms of Toxicity) (Ernst and Zibrak 1998; Gorman et al. 2003; Kao and Nañagas 2006). Under conditions of hypoxia, including that induced by COHb formation, cerebrovascular vasodilation and increased cardiac output occur as compensatory mechanisms to maintain O<sub>2</sub> delivery to the brain (Gorman et al. 2003; Helfaer and Traystman 1996; Raub and Benignus 2002; WHO 1999). In healthy individuals, cardiovascular compensation effectively maintains whole brain O<sub>2</sub> delivery up to COHb levels of approximately 60% (EPA 2000). However, based on studies conducted in animals, the elevated cerebral blood flow during carbon monoxide exposure is heterogeneous among different brain regions and may

## 3. HEALTH EFFECTS

not be sufficient to satisfy O<sub>2</sub> demand in all brain regions (Okeda et al. 1987; Sinha et al. 1991). However, despite these compensatory cardiovascular actions, cerebral O<sub>2</sub> consumption declines as blood COHb levels increase, with statistically significant decreases at COHb levels of 30–50% (Raub and Benignus 2002). Results of an analysis conducted by EPA (2000) of data obtained in goats (Doblar et al. 1977) and sheep (Langston et al. 1996) show that the cerebral metabolic rate for O<sub>2</sub> (CMRO<sub>2</sub>) decreased 10% at a blood COHb of approximately 23%. These results are consistent with data obtained from humans showing no statistically significant decline in CMRO<sub>2</sub> at COHb levels up to 20% (Paulson et al. 1973).

Based on an extensive database of acute carbon monoxide poisoning, it is generally accepted that central nervous system symptoms are associated with acute exposures that produce blood COHb levels  $\geq 20\%$  (EPA 1991, 2000; Raub and Benignus 2002; Raub et al. 2000; WHO 1999). However, results of clinical studies of associations between carbon monoxide exposure and carbon monoxide-induced nervous system effects at blood COHb levels between 5 and 20% are more difficult to interpret (EPA 1991, 2000; Raub and Benignus 2002; Raub et al. 2000; WHO 1999). Numerous clinical studies have investigated the potential for carbon monoxide to induce adverse nervous system effects at COHb levels  $< 20\%$ , including visual and auditory sensory effects (decreased visual tracking, visual and auditory vigilance, visual perception), fine and sensorimotor performance, cognitive effects (altered time discrimination, learning, attention level, driving performance), and brain electrical activity (Benignus et al. 1990; EPA 1991, 2000; Raub and Benignus 2002; Raub et al. 2000). Interpretation of results from most of these clinical studies is complicated by poor study design (single-blind or unblinded designs, small number of study subjects), inadequate reporting, inconsistent results, and inability to duplicate positive findings (Benignus et al. 1990; EPA 1991, 2000; Raub and Benignus 2002; Raub et al. 2000). Therefore, although evidence has been provided for central nervous system effects in humans in association with carbon monoxide exposures that result in COHb levels  $< 20\%$ , dose-response relationships for these effects have not been firmly established. Effects of exposure to ambient environmental levels of carbon monoxide on neurological function and behavior have not been investigated in epidemiological studies.


Benignus (1994) conducted a meta-analysis of the clinical literature on carbon monoxide-induced behavioral effects, (also see Benignus et al. 1990; EPA 2000; Raub and Benignus 2002). To minimize the introduction of bias from poorly designed studies, the Benignus analysis included only studies conducted using a double-blind design. The primary goal of these analyses was to compare the dose-response relationship of carbon monoxide-induced effects in humans to other extrapolated data; specifically, the dose-response functions for COHb-induced behavioral effects in rats and for hypoxic

## 3. HEALTH EFFECTS

hypoxia (HH)-induced behavioral effects in humans. Rat COHb dose-response data were converted to human COHb equivalents using a modification of the CFK human model, adjusted to predict exposure-COHb relationships in rats (Benignus and Annau 1994). Data on behavioral effects were also adjusted to account for carbon monoxide-Hb-related hypothermia that occurs in rats (Gordon 1990), but not in humans, assuming response additively of hypothermia and hypoxia (Benignus 1994). Dose-response data for HH-induced behavioral effects were converted to COHb equivalents (COHb% that yields the equivalent arterial O<sub>2</sub> content as a given level of HH) and were corrected for hyperventilation that occurs in HH, and resulting changes in alveolar ventilation rate, based on estimates of blood CO<sub>2</sub> (i.e., hypocapnia-induced by HH). For all data sets (i.e., carbon monoxide human, carbon monoxide rat, HH human), behavioral responses were transformed to percent of baseline response. Dose-response curves were fit to the extrapolated carbon monoxide rat and HH human data; curve-fitting was not conducted for carbon monoxide human data, due to the small effect level and low COHb levels. Comparison of the extrapolated carbon monoxide rat and HH human data showed very close agreement, with nearly superimposable dose-response curves (Figure 3-1); comparison of dose-response data from carbon monoxide humans to extrapolated data from HH humans is shown in Figure 3-2. The dose-response relationship from this meta-analysis predicts a 10% decrement in behavioral outcomes in humans in association with COHb levels of approximately  $\geq 20\%$  (range: approximately 18–25%, based on 90% confidence limits) in healthy, sedentary individuals. As discussed by Benignus (1994) and others (Benignus et al. 1990; Raub and Benignus 2002; EPA 1991), the magnitude of behavioral effects that may occur at blood COHb levels  $\leq 20\%$  is expected to be small. Although most task performances are not likely to be affected, it is possible that even small effects (i.e.,  $<10\%$  decrement) could interfere with successful task performance for more difficult or demanding tasks. Therefore, potential minimal behavioral changes occurring at COHb levels  $\leq 20\%$  may be important under certain conditions or in sensitive individuals (e.g., individuals with compromised cardiovascular or central nervous system function).

Clinical studies evaluating exposure to low levels of carbon monoxide (i.e., producing blood COHb levels  $<20\%$ ) have focused on the potential for carbon monoxide to induce behavioral effects; however, very little information is available on the effects of carbon monoxide on sensory system function in humans. Results of clinical studies in evaluating the effects of low carbon monoxide exposures on alterations in sensory perception (i.e., visual and auditory) are equivocal (EPA 1991, 2000). Epidemiological studies have not assessed the potential for carbon monoxide to produce alterations in sensory system function. Studies conducted in mature rats provide evidence that acute exposure ( $<10$  hours) to higher levels of carbon monoxide (500–1,500 ppm) potentiates noise-induced hearing loss, including noise-induced


## 3. HEALTH EFFECTS

**Figure 3-1. HH Human and Carbon Monoxide Rat Curves\***

\*Behavioral decrements in rats with elevated COHb (dashed line) compared to behavioral effects in hypoxic humans (solid line) in which hypoxia has been expressed in equivalence to COHb.

Source: Raub and Benignus 2002

## 3. HEALTH EFFECTS

**Figure 3-2. HH Human and Carbon Monoxide Human\***

\*Behavioral decrements in humans (circles) with elevated COHb compared to behavioral effects in hypoxic humans (solid and dotted lines) in which hypoxia has been expressed in equivalence to COHb. Closed and open circles were the mean values for studies reporting statistically and non-significant results, respectively.

Source: Raub and Benignus 2002

### 3. HEALTH EFFECTS

elevation of compound action potential threshold (Chen and Fechter 1999) and auditory threshold shifts (Fechter et al. 1988; Young et al. 1987). In these studies, exposure to carbon monoxide alone did not produce changes in auditory function. Noise-induced hearing loss was partially reversed within 4 weeks following exposure to noise alone, but not following exposure to noise and carbon monoxide (Chen and Fechter 1999). Results are consistent with carbon monoxide-induced impairment of mechanisms that repair noise-induced damage to outer hair cells. In marked contrast to results observed in adult rats, developmental studies in animals show that gestational and early postnatal exposure to low levels of carbon monoxide (i.e., 12–25 ppm carbon monoxide) may lead to altered development of the auditory system (Lopez et al. 2003, 2008; Stockard-Sullivan et al. 2003; Webber et al. 2003). These data are discussed in Section 3.2.6.

Additional evidence for effects of carbon monoxide on depressed auditory thresholds and increased cochlear blood flow have been provided from studies in which adult rats received parenteral doses of carbon monoxide (Fechter et al. 1987b). However, in these studies, auditory thresholds assessed by measurement of the compound action potential recorded at the round window remained stable up to COHb levels of at least 30%. Transient impairments of the compound action potential threshold were observed among subjects that attained COHb levels of approximately 50%. Cochlear blood flow measured using laser Doppler flowmetry also increased following carbon monoxide exposure, reaching a maximum of 140% of control values at approximately 50% COHb. In those instances of high COHb production that did produce impairment of auditory threshold, recovery of function was observed as COHb levels declined.

#### **3.2.5 Reproductive Effects**

Fetal death in humans has been reported in cases of maternal carbon monoxide poisoning during pregnancy. Epidemiological studies have examined possible relationships between exposures to ambient air concentrations of carbon monoxide and fetal mortality (see Section 3.2.6). Studies in animals provide further evidence that maternal exposure during pregnancy can result in fetal death.

*Case Reports.* Several case reports have evaluated pregnancy outcomes following acute carbon monoxide poisoning (Brown et al. 1992; Caravati et al. 1988; Cramer 1982; Elkharrat et al. 1991; Farrow et al. 1990; Greingor et al. 2001; Hollander et al. 1987; Margulies 1986; Norman and Halton 1990; Silverman and Montano 1997). In all cases, the severity of carbon monoxide poisoning required emergency room treatment or hospitalization, suggesting that maternal COHb levels exceeded 20%.

### 3. HEALTH EFFECTS

Pregnancy outcomes varied widely, from delivery of healthy infants at term to fetal death within first few days of exposure. Pregnancy outcome is most likely related to fetal age at time of exposure and severity of maternal poisoning, although, maternal COHb levels are not a good indicator of fetal outcome (Greingor et al. 2001). Fetal death may be secondary to maternal hypoxia and/or fetal hypoxia. The available information from these cases is not adequate to define dose-response relationships for acute carbon monoxide exposure and pregnancy outcomes.

*Animal Studies.* Few animal studies have assessed the effects of carbon monoxide exposure on reproductive function. Continuous exposure of male mice to 50 ppm carbon monoxide for 2 weeks prior to mating with unexposed females or of male and female mice to 50 ppm carbon monoxide for 2 weeks prior to mating had no effect on pregnancy rate or live or dead fetuses/dam (Stupfel and Bouley 1970). Exposure of pregnant rats over the entire gestational period to up to 200 ppm carbon monoxide had no effect on number of litters, fetuses per litter, or number of live/dead pups per litter (Fechter and Annau 1977; Penney et al. 1983). In contrast, increased fetal death during the first 24 hours of the postnatal period was observed in rabbits following maternal exposure to 90 and 180 ppm carbon monoxide during pregnancy (9.9 and 35% mortality) (Astrup et al. 1972).

#### 3.2.6 Developmental Effects

Epidemiological studies have examined possible associations between exposure to ambient air carbon monoxide concentrations and various developmental outcomes, including pre-term birth, birth weight, congenital anomalies, and neonatal and infant death. Results of these studies have been mixed and, collectively, do not provide strong evidence for developmental effects in association with exposures to ambient levels of carbon monoxide. Studies conducted in animals provide evidence of adverse developmental effects of gestational and early postnatal carbon monoxide exposure, including decreased fetal weight, adverse central nervous system development, altered peripheral nervous system development, cardiac effects, altered sexual behavior, immunological effects, and hematological effects. In addition, some studies showed that developmental effects persisted beyond the postnatal period.

*Epidemiological Studies.* Interpretation of the results of epidemiological studies of developmental effects of carbon monoxide, in particular, quantitative estimates of the effect magnitudes on outcomes, is limited by several factors: (1) reliance on area monitoring for estimating exposure level that may not represent exposures that occurred during any particular period of gestation; (2) uncertainty in knowledge of temporal correspondence between monitored exposure levels and outcomes; (3) relatively strong

## 3. HEALTH EFFECTS

correlations between ambient air carbon monoxide concentrations and other air quality variables that may affect developmental outcomes; and (4) relatively low carbon monoxide exposures studied. In general, these studies have examined relatively low air carbon monoxide concentrations, typical of ambient levels (e.g., mean concentrations ranging from 0.5 to 3 ppm, with the highest reported values <10 ppm). The presentation of the epidemiological studies is organized by major categories of birth and neonatal outcomes. Study conclusions are presented in the text, with selected supporting details presented in tabular form (Table 3-8). Although most studies explored various time lags between monitored air carbon monoxide concentrations and outcomes, as well as various sample strata, for the sake of brevity, only selected representative time lags (usually those indicative of the strongest associations to carbon monoxide) are presented in the tables. Where co-pollutant models have been explored, these results are also presented.

*Studies of Pre-term Birth.* Several epidemiological studies have examined possible associations between ambient air carbon monoxide concentrations and risk of pre-term birth (Huynh et al. 2006; Jalaludin et al. 2007; Leem et al. 2006; Liu et al. 2003; Ritz et al. 2000, 2007; Wilhelm and Ritz 2005). Results of these studies have been mixed and have included estimates of significantly elevated risk or lower risks of pre-term birth (Jalaludin et al. 2007). Mean air carbon monoxide concentrations evaluated in these studies ranged from 0.8 to 2.7 ppm, with the highest values <10 ppm (Ritz et al. 2000).

A retrospective cohort study (n=97,158 births) conducted in southern California (1989–1993) found a significant association between increasing air carbon monoxide concentration (mean: 2.7 ppm, range: 0.6–9.12) and pre-term birth (Ritz et al. 2000). After adjustment for other risk factors as well as ambient air concentrations of NO<sub>2</sub>, O<sub>3</sub>, and PM<sub>10</sub>, the relative risk of pre-term birth was estimated to be 1.12 (95% CI: 1.04–1.21) per 3 ppm increase in air carbon monoxide concentration during the last 6 weeks of pregnancy. Subjects were matched to ambient carbon monoxide measurements made at monitors that were <2 miles from the residence. In a follow-up study (1994–2000) that included 106,483 births, relative risk (adjusted for NO<sub>2</sub> and O<sub>3</sub>) was estimated to be 1.10 (95% CI: 1.03–1.08) per 1 ppm increment in air carbon monoxide concentration during the last 6 weeks of pregnancy (Wilhelm and Ritz 2005). The association was not significant when PM<sub>10</sub> was included in the model along with NO<sub>2</sub> and O<sub>3</sub> (relative risk: 0.98; 95% CI: 0.83–1.18). A subsequent analysis of Southern California data included a nested case-control study consisting of a subset of 2,543 cases of pre-term birth (<37 weeks of gestation) or low birth weight (<2,500 g) and an equal number of randomly selected controls for a larger cohort of 58,316 births (Ritz et al. 2007). Detailed individual data on potential co-variables were obtained, including maternal age, race, education, marriage status, birth season, parity, active and passive smoking,

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Bell et al. (2007) Period: 1999–2002 Location: Connecticut and Massachusetts	Outcome: BW, LBW Design: Retrospective cohort Sample: n=358,504 live full term singleton births	Avg time: 24 hours Mean: 0.65 ppm SD: 0.2 ppm	Increment: 0.30 ppm $\beta$ coefficient (95% CI) BW (g), total gestation: CO: -16.2 (-19.7, -12.6) Significant negative $\beta$ persisted after adjustment for NO <sub>2</sub> , SO <sub>2</sub> , PM <sub>2.5</sub> , or PM <sub>10</sub> OR (90% CI) LBW, total gestation: CO: 1.028 (0.983–1.074)
Brauer et al. (2008) Period: 1999–2004 Location: Vancouver, Canada	Outcome: PTB, LBW, SGA Design: Retrospective cohort Sample: n=70,249 live singleton births	Avg time: 24 hours Mean: 0.55 ppm Range: 0.11–1.23 ppm	Increment: 0.1 mg/m <sup>3</sup> OR (95% CI), total gestation: SGA: 1.06 (1.03–1.08) LBW: 1.02 (0.96–1.09) PTB: 1.16 (1.01–1.33)
Chen et al. (2002) Period: 1991–1999 Location: Nevada	Outcome: BW, LBW Design: Retrospective cohort Sample: n=39,338 live singleton births	Avg time: 8 hours Mean: 0.86 ppm Range: 0.42–4.25 ppm	Increment: 1ppm $\beta$ coefficient (SE), BW (g): TM 1: -1.02 (6.68) TM 2: -0.07 (6.58) TM 3: -3.95 (6.76) Not significant when adjusted for PM <sub>10</sub> or O <sub>3</sub> ORs not significant for LBW
Gilboa et al. (2005) Period: 1997–2000 Location: Texas	Outcome: CA (heart, orofacial) Design: Case-control Sample: n=4,594 cases	Avg time: 24 hours Mean: NR 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.4–0.7 ppm	Increment: <0.4 ppm (reference), 0.4–0.5 (L), 0.5–0.7 (M), >0.7 (H) OR (95% CI), 3–8 weeks of gestation: Conotruncal defect: (L) 1.38 (0.97–1.97); (M)1.17 (0.81–1.70); (H) 1.46 (1.03–2.08) Teratology of Fallot: (L) 0.92 (0.52–1.62); (M) 1.17 (0.75–2.14); (H) 2.04 (1.16–3.29) CO not associated with other heart defects or orofacial defects

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Gouveia et al. (2004) Period: 1997 Location: San Paulo, Brazil	Outcome: BW, LBW Design: Retrospective cohort Sample: n=179,460 live full term singleton births	Avg time: 8 hours Mean: 3.7 ppm Range: 1.1–11.4	Increment: 1 ppm $\beta$ coefficient (95% CI), BW (g): TM 1: -23.1 (-41.3 to -4.9) TM 2: 3.2 (-18.2 to 24.5) TM 3: 1.9 (-18.2 to 22.0) Significant negative $\beta$ did not persist after adjustment for SO <sub>2</sub> or PM <sub>10</sub> . OR (95% CI), LBW: 4 <sup>th</sup> vs. 1 <sup>st</sup> quartile exposure: TM 1: 1.02 (0.82–1.27); TM 2: 1.07 (0.88–1.30) TM 3: 0.93 (0.76–1.12)
Ha et al. (2001) Period: 1996–1997 Location: Seoul, South Korea	Outcome: LBW Design: Retrospective cohort Sample: n=276,763 live full-term singleton births	Avg time: 8 hours Median: 1.7 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.99–1.41	Increment: 0.42 ppm RR (95% CI) TM 1: 1.08 (1.04–1.12) TM 3: 0.91 (0.87–0.96)
Ha et al. (2003) Period: 1995–1999 Location: Seoul, South Korea	Outcome: Post-neonatal mortality (1–12 months) Design: Time series Sample: n=1,045 deaths	Avg time: 24 hours Median: 1.2 ppm Range: 0.39–3.38 ppm	Increment: 0.57 ppm RR (95% CI), lag 0 days: All causes: 1.020 (0.976–1.067) Respiratory: 1.388 (1.009–1.911)
Huynh et al. (2006) Period: 1999–2000 Location: California	Outcome: PTB Design: Case-control Sample: n=10,673 cases (24–36 weeks), 32,119 controls (39–44 weeks)	Avg time: 24 hours Mean: 0.8 ppm SD: 0.2 ppm	Increment: 1 ppm OR (95% CI) Month 1 (of gestation): CO: 1.10 (0.999–1.20) CO+PM <sub>2.5</sub> : 1.03 (0.93–1.13) Last 2 weeks: CO: 1.00 (0.93–1.09) CO+PM <sub>2.5</sub> : 0.97 (0.90–1.06) Total gestation: CO: 1.06 (0.95–1.18) CO+PM <sub>2.4</sub> : 0.98 (0.87–1.10) No significant trend in OR with exposure quartile

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Hwang and Jaakkola (2008) Period: 2001–2003 Location: Taiwan	Outcome: CA (oral cleft) Design: Case-control Sample: n=653 cases	Avg time: 8 hours Mean: 0.69 ppm Range: 0.25–2.7 ppm	Increment: 0.1 ppm RR (95% CI), oral cleft Cleft lip: Month 1: 1.00 (0.96–1.04) Month 2: 1.00 (0.96–1.03) Month 3: 1.00 (0.96–1.03) Cleft lip with or without cleft palate: Month 1: 1.00 (0.97–1.04) Month 2: 1.00 (0.97–1.05) Month 3: 1.00 (0.96–1.04) No significant ORs after adjustment for NO <sub>2</sub> , O <sub>3</sub> , or PM <sub>10</sub>
Jalaludin et al. (2007) Period: 1998–2000 Location: Sydney, Australia	Outcome: PTB (<37 or <42 weeks) Design: Retrospective cohort Sample: n=721,289 full-term singleton births	Avg time: 8 hours Mean: 0.9 ppm SD: 0.68 ppm	Increment: 1 ppm RR( 95% CI) <5 km of monitoring site or city-wide Month 1, <5 km: 1.03 (0.68–1.54) Month 1, city-wide: 0.89 (0.84–0.95) TM 1, <5 km): 1.24 (0.81–1.91) TM 1, city-wide:0.77 (0.71–0.83) Last month, <5 km: 1.00 (0.86–1.15) Last month, city-wide: 0.96 (0.88–1.04) TM 3, <5 km: 1.11 (0.94–1.31) TM 3, city-wide: 0.99 (0.90–1.09)
Lee et al. (2003a) Period: 1996–1998 Location: Seoul, Korea	Outcome: LBW Design: Retrospective cohort Sample: n=388,105 live full-term singleton births	Avg time: 24 hours Mean: 1.2 ppm Range: 0.4–3.4 ppm	Increment: 0.5 ppm OR (95% CI) TM 1: 1.04 (1.01–1.07) TM 2: 1.03 (1.00–1.06) TM 3: 0.96 (0.93–0.99) Total gestation: 1.05 (1.01–1.09)

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Leem et al. (2006) Period: 2001–2002 Location: Incheon, Korea	Outcome: PTB Design: Retrospective cohort Sample: n=52,113 live singleton births	Avg time: 24 hours Mean: NR Range: 0.3–1.4 ppm	Increment: quartiles (mg/m <sup>3</sup> ): 1 <sup>st</sup> 0.47–0.63; 2 <sup>nd</sup> 0.6–0.77, 3 <sup>rd</sup> 0.78–0.90, 4 <sup>th</sup> 0.91–1.27 OR (95% CI) TM 1: 2 <sup>nd</sup> , 0.92 (0.81–1.05); 3 <sup>rd</sup> , 1.14 (1.01–1.29); 4 <sup>th</sup> , 1.26 (1.11–1.44) TM 3: 2 <sup>nd</sup> , 1.07 (0.95–1.21); 3 <sup>rd</sup> , 1.07 (0.94–1.22); 4 <sup>th</sup> , 1.16 (1.01–1.34)
Lin et al. (2004a) Period: 1998–2000 Location: Sao Paulo, Brazil	Outcome: Neonatal mortality (<28 days) Design: Time series Sample: n=6,700 deaths	Avg time: 24 hours Median: 2.83 ppm Range: 0.54–10.25 ppm	Increment: 0.57 ppm β coefficient (SE), deaths per day, lag 0 days 0.0061 (0.0110)
Lin et al. (2004b) Period: 1995–1997 Location: Taiwan	Outcome: LBW Design: Retrospective cohort Sample: n=92,288 live full-term singleton births	Avg time: 24 hours Mean: Taipei, 0.84–1.31 ppm; Kaohsiun, 5.56–10.05 ppm Range: NR	Increment: <1.1 (reference), 1.1–14.2 (L) vs. >14.2 (H) ppm OR (95% CI): TM 1: (L) 1.01 (0.89–1.16); (H) 0.90 (0.75–1.09) TM 2: (L) 1.02 (0.90–1.16); (H) 1.00 (0.82–1.22) TM 3: (L) 0.88 (0.77–1.00); (H) 0.86 (0.71–1.03) Total gestation: (L) 0.89 (0.77–1.01); (H) 0.77 (0.63–0.94)
Liu et al. (2003) Period: 1985–1998 Location: Vancouver, Canada	Outcome: PTB LBW, IUGR Design: Retrospective cohort Sample: n=386,202 live singleton births	Avg time: 24 hours Mean: 1.0 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.7–1.2 ppm	Increment: 1 ppm OR (95% CI) PTB, first month: 0.95 (0.89–1.01) PTB, last month: 1.08 (1.01–1.15) LBW, first month: 1.01 (0.93–1.09) LBW, last month: 0.96 (0.88–1.04) IUGR, first month: 1.06 (1.01–1.10) IUGR, last month: 0.98 (0.94–1.03) IUGR, TM 1: 1.05 (1.00–1.10) IUGR, TM 2: 0.97 (0.92–1.01) IUGR, TM 3: 0.97 (0.93–1.02)

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Liu et al. (2007) Period: 1995–2007 Location: Canada	Outcome: IUGR Design: Retrospective cohort Sample: n=386,202 live singleton births	Avg time: 24 hours Mean: 1.1 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.6–1.3 ppm	Increment: 1 ppm RR (95% CI) TM 1: CO:1.18 (1.14–1.23) CO+NO <sub>2</sub> +O <sub>3</sub> : 1.18 (1.12–1.24) TM 2: CO:1.14 (1.10–1.18) CO+NO <sub>2</sub> +O <sub>3</sub> : 1.15 (1.08–1.21) TM 3: CO:1.09 (1.24–1.14) CO+NO <sub>2</sub> +O <sub>3</sub> : 1.20 (1.14–1.26)
Maisonet et al. (2001) Period: 1994–1996 Location: Northeast United States	Outcome: BW Design: Retrospective cohort Sample: n=89,557 live full term singleton births	Avg time: 24 hour Mean: 1.1 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.9–1.5 ppm	Increment: 1 ppm OR (95% CI): TM 1: 1.08 (0.91–1.28) TM 2: 1.14 (0.83–1.58) TM 3: 1.31 (1.06–1.62) African-Americans: TM 1: 1.43 (1.18–1.74) TM 2: 1.27 (0.87–1.86) TM 3: 1.75 (1.50–2.04) CO had no effect on strata for whites or Hispanics
Mannes et al. (2005) Period: 1998–2000 Location: Sydney, Australia	Outcome: BW, SGA Design: Retrospective cohort Sample: n=136,056 live full term singleton births	Avg time: 8 hour Mean: 0.8 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0 –4.6 ppm	Increment: 1 ppm $\beta$ coefficient (95% CI) BW (g), TM 2, <5 km from monitoring site: CO: -29.87 (-50.98, -6.76) CO+NO <sub>2</sub> : -20.17 (-43.12, 2.78) CO+ PM <sub>10</sub> : -27.31 (-55.30, 0.68) OR (95% CI) SGA <5 km from monitoring site: TM 1: 0.99 (0.86–1.14) TM 2: 1.06 (0.90–1.25) TM 3: 1.05 (0.90–1.23) One month prior to birth: 1.10 (0.96–1.27)

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Medeiros and Gouveia (2005) Period: 1998–2000 Location: Sao Paulo, Brazil	Outcome: BW and LBW Design: Retrospective cohort Sample: n=311,735 live full-term singleton births (37–41 weeks)	Avg time: 24 hours Mean: ~2–3ppm Range: ~1–12 ppm	Increment: 1 ppm $\beta$ coefficient (95% CI) BW TM 1: -11.9 (-15.5, -8.2) TM 2 4.9 (0.5–9.3) TM 3: 12.1 (7.6–16.6) OR (85% CI) LBW (75 <sup>th</sup> % vs 25 <sup>th</sup> %): TM 1: 0.98 (0.91–1.06); TM 2: 0.97 (0.90–1.05) TM 3: 1.03 (0.96–1.11)
Parker et al. (2005) Period: 2000 Location: California	Outcome: BW, SGA Design: Retrospective cohort Sample: n=18,247 live full-term singleton births	Avg time: 24 hours Mean: 0.78 ppm 25 <sup>th</sup> –95 <sup>th</sup> ‰: 0.57–0.93 ppm	Increment (ppm): <0.57 (reference), 0.57–0.76 (L), 0.76–0.93 (M), >0.93 (H) OR (95% CI) for total gestation: CO, L: 0.93 (0.80–1.09) CO, M: 0.91 (0.78–1.06) CO, H: 0.95 (0.81–1.12) CO (L) + PM <sub>2.5</sub> : 0.90 (0.77–1.06) CO (M) + PM <sub>2.5</sub> : 0.91 (0.77–1.07) CO (H) + PM <sub>2.5</sub> : 0.82 (0.68–0.99)
Ritz and Yu (1999) Period: 1989–1993 Location: California	Outcome: LBW Design: Retrospective cohort Sample: n=125,275 live full-term singleton births	Avg time: 6–9 AM Mean: 2.59 ppm 95 <sup>th</sup> ‰: 5.5 ppm	Increment (ppm): <2.2 (reference), 2.2–<5.5 (L), >5.5 (H) OR (95% CI) CO (L): 1.04 (0.96–1.13) CO, (H): 1.22 (1.03–1.44) CO (L)+ NO <sub>2</sub> +O <sub>3</sub> _PM <sub>10</sub> : 1.10 (0.91–1.32) CO (H)+ NO <sub>2</sub> +O <sub>3</sub> _PM <sub>10</sub> : 1.38 (0.86–2.22)

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Rtiz et al. (2000) Period: 1989–1993 Location: California	Outcome: PTB Design: Retrospective cohort Sample: n=97,518 live singleton births	Avg time: 6–9 AM Mean: 2.70 ppm Range: 0.36–9,12 ppm	Increment: 3 ppm RR (95% CI) Adjusted for risk factors: Last 6 weeks: 1.06 (1.02–1.10) First month: 1.01 (0.97–1.04) Adjusted for risk factors and season of birth and conception: Last 6 weeks: 1.04 (0.99–1.10) First month: 1.04 (0.99–1.09) Adjusted for risk factors, NO <sub>2</sub> , O <sub>3</sub> , and PM <sub>10</sub> : Last 6 weeks: 1.05 (0.97–1.12) First month: 1.03 (0.96–1.10)
Ritz et al. (2002) Period: 1987–1993 Location: California	Outcome: CA (heart, orofacial) Design: Case-control Sample: n=3,549 cases	Avg time: 24 hour Mean: NR 25 <sup>th</sup> –75 <sup>th</sup> ‰: 1.4–2.46 ppm	Increment: <1.4 ppm (reference), 1.14–1.57 (L), 1.57–2.39 (M), >2.39 (H) OR (95% CI), month 2 of gestation: Ventricular septal: CO: (L) (L) 1.62 (1.05–2.48); (M) 2.09 (1.19–3.67); (H) 2.95 (1.44–6.05) CO+NO <sub>2</sub> +O <sub>3</sub> +PM <sub>10</sub> : 1.63 (1.00–2.66); (M) 1.97 (1.0–3.91); (H) 2.84 (1.15–6.99) CO not associated with conotruncal, aortic artery and valve, or pulmonary artery and valve defects
Ritz et al. (2006) Period: 1989–2000 Location: California	Outcome: Post-neonatal mortality (28 days–1 year) Design: Case-control Sample: n=13,146 cases	Avg time: 24 hour Mean: 1.63 ppm Range: 0.38–3.44 ppm	Increment: 1 ppm OR (95% CI), exposure period 2 months prior to death: All causes: 1.11 (1.06–1.16) SIDS: 1/19 (1.10–1.28) OR not significant after adjustment for NO <sub>2</sub> , O <sub>3</sub> , and PM <sub>10</sub>

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Ritz et al. (2007) Period: 2003 Location: California	Outcome: PTB Design: Nested case-control Sample: n=2,543 cases, 2,543 controls from 58,316 births in registry	Avg time: 24 hours Mean: NR Range: NR	Increment (ppm): <0.58 (reference), 0.59–0.91(L), 0.92–1.25 (M), >1.25 (H) OR (95% CI), adjusted for risk factors: TM 1: L, 1.17 (1.08–1.26); M: 1.15 (1.05–1.26); H: 1.25 (1.12–1.38) Last 6 weeks: L, 1.00 (0.93–1.08); M: 1.08 (0.98– 1.20); H: 1.03 (0.93–1.14) Total gestation: L, 0.76 (0.70–0.82); M: 0.84 (0.77–0.91); H: 1.03 (0.91–1.17)
Salam et al. (2005) Period: 1975–1987 Location: California	Outcome: BW, LBW, IUGR Design: Retrospective cohort Sample: n=3,901 births (37– 44 weeks)	Avg time: 24 hours Mean: 1.8 ppm SD: 0.9 ppm	Increment: 1.4 ppm OR (95% CI): TM 1: LBW, CO: 1.0 (0.7–1.5) IUGR, CO: 1.2 (1.0–1.4) $\beta$ coefficient (95% CI) for IUGR TM 1: CO: -21.7 (-42.3, -1.1) CO + O <sub>3</sub> : -28.6 (-50.4, -6.9)
Tsai et al. (2006a) Period: 1994–2000 Location: Taiwan	Outcome: Post-neonatal mortality (27 days–1 year) Design: Case-crossover Sample: n=206 cases	Avg time: 24 hours Mean: 0.83 ppm Range: 0.23–1.77 ppm	Increment: 0.31 ppm OR (95% CI), lag 0–2 days: All causes: 1.051 (0.304–3.630)

**Table 3-8. Selected Epidemiological Studies of Associations Between Ambient Carbon Monoxide Concentrations and Developmental Outcomes**

Study	Design features	CO exposure	Effect size
Wilhelm and Ritz (2005) Period: 1994–2000 Location: California	Outcome: LBW, PTB Design: Retrospective cohort Sample: n=106,483 births for PTB outcome, 136,134 births for LBW outcome	Avg time: 24 hours Mean: 1.2–1.4 ppm Range: 0.2–5.9 ppm	Increment: 1 ppm ( $\leq 1$ mile from monitoring station) RR (95% CI), adjusted for risk factors: PTB, TM 1: CO: 1.06 (1.00–1.12) CO+NO <sub>3</sub> +O <sub>3</sub> : 1.10 (1.01–1.20) CO+NO <sub>3</sub> +O <sub>3</sub> + PM <sub>10</sub> : 0.99 (0.83–1.18) PTB, Last 6 weeks: CO: 1.04 (0.98–1.09) CO+NO <sub>3</sub> +O <sub>3</sub> : 1.10 (1.03–1.18) CO+NO <sub>3</sub> +O <sub>3</sub> + PM <sub>10</sub> : 0.98 (0.83–1.16) LBW, TM 3: CO: 1.10 (0.98–1.23) CO+NO <sub>3</sub> +O <sub>3</sub> : 1.15 (0.98–1.35) CO+NO <sub>3</sub> +O <sub>3</sub> + PM <sub>10</sub> : 1.21 (0.85–1.74)
Woodruff et al. (2008) Period: 1999–2002 Location: U.S. counties with >250,000 residents	Outcome: Post-neonatal mortality (28 days–1 year) Design: Retrospective cohort Sample: n=6,639 deaths of 3,583,495 births	Avg time: 24 hours Mean: 0.70 ppm 25 <sup>th</sup> –75 <sup>th</sup> ‰: 0.48–0.87 ppm	Increment: 0.39 ppm OR (95% CI), exposure during age 0–2 months: All causes: 1.001 (0.95–1.07) Respiratory: CO: 1.14 (0.93–1.40) CO+O <sub>3</sub> +PM <sub>10</sub> +SO <sub>2</sub> : 1.02 (0.89–1.15) SIDS: CO: 0.88 (0.76–1.03) SIDS, CO+O <sub>3</sub> +PM <sub>10</sub> +SO <sub>2</sub> : 0.96 (0.87–1.06)
Yang et al. (2006) Period: 1994–2000 Location: Taiwan	Outcome: Post-neonatal mortality (27 days–1 year) Design: Case-crossover Sample: n=471 cases	Avg time: 24 hours Mean: 1.158 ppm Range: 0.32–4.84 ppm	Increment: 0.55 ppm OR (95% CI), lag 0–2 days: All causes: 1.038 (0.663–1.624)

BW = birth weight; CA = congenital anomalies; CO = carbon monoxide; IUGR = intrauterine growth rate (BW<10<sup>th</sup> percentile of birth cohort when adjusted for gestational age); LBW = low birth weight (<2,500 g); PTB = pre-term birth (<37 weeks); OR = odds ratio; SD = standard deviation; SE = standard error; SGA = small for gestational age (BW<10<sup>th</sup> percentile of birth cohort when adjusted for gestational age); TM = trimester

## 3. HEALTH EFFECTS

and alcohol use during pregnancy. The adjusted odds ratio in the case-control study for pre-term birth (exposure to >1.25 ppm in first trimester compared to <0.58 ppm) was 1.21 (95% CI: 0.88–1.65; adjusted for co-variables other than co-pollutants). The odds ratio in the larger cohort (n=58,316) was similar (odds ratio: 1.25, 95% CI: 1.12–1.38); adjusted for maternal age, race, education, birth season, and parity). Results of multi-pollutant models were not reported. A larger case-control study, also conducted in southern California (1999–2000), did not find a significant association between ambient air carbon monoxide concentration and pre-term birth (Huynh et al. 2006). This study examined 10,673 cases of pre-term births (<37 weeks of gestation) and 32,119 full-term controls (matched for last menstrual period date within 2 weeks of cases). The adjusted odds ratios for pre-term birth were not significant, when adjusted for maternal age, maternal race/ethnicity, maternal education, marital status, and parity (with or without inclusion of PM<sub>2.5</sub> exposure). A 1 ppm increase in ambient carbon monoxide concentration during the first month of pregnancy was associated with an odds ratio of 1.10 (95% CI: 0.99–1.20) when CO was considered alone, and 1.03 (95% CI: 0.83–1.13) after adjustment for PM<sub>10</sub>. This study matched ambient carbon monoxide measurements to subjects whose residence was <5 miles of the monitoring station, whereas the Ritz et al. (2000, 2007) and Wilhelm and Ritz (2005) studies matched subjects to <2 miles from the monitoring station.

Other large retrospective cohort studies have examined pre-term births in association with ambient air carbon monoxide concentrations. Liu et al. (2003) examined a cohort of 229,085 births in Vancouver, Canada (1985–1998). The odds ratio for pre-term birth for a 1 ppm increase in ambient air carbon monoxide concentration measured during the first month of pregnancy was 1.08 (95% CI: 1.01–1.15; adjusted for maternal age, parity, infant sex, birth weight, and season of birth), and remained elevated after adjustment for NO<sub>2</sub> and O<sub>3</sub> (1.08, 95% CI: 1.00–1.20). The mean air carbon monoxide concentration in the study was 1 ppm and the highest value was 12.8 ppm. A cohort study of 52,113 births in Incheon, Korea (2001–2002) also estimated elevated risks of pre-term birth in association with increasing ambient air carbon monoxide concentrations during the first trimester. The relative risk increased with increasing estimated exposure concentration and was 1.26 (95% CI: 1.11–1.44) for the strata 0.91–1.27 mg/m<sup>3</sup> compared to the reference, >0.47–0.63 mg/m<sup>3</sup> (adjusted for maternal age, parity, sex, season of birth, and education level of father and mother; Leem et al. 2006). Results of multi-pollutant models were not reported. The ambient air carbon monoxide concentrations ranged from approximately 0.35 to 1.4 ppm and were matched to subjects by spatial kriging (a spatial statistical method for interpolating exposure levels between sampling locations) with spatial averaging on a scale of approximately 3.5 km<sup>2</sup>. A cohort study of 123,840 births in Sydney, Australia (1998–2000) did not find a significant association between ambient air carbon monoxide concentration and pre-term birth (Jalaludin

## 3. HEALTH EFFECTS

et al. 2007). The highest odds ratio estimated was 1.24 (95% CI: 0.81–1.91) or a 1 ppm increase in 8-hour maximum carbon monoxide concentration measured during first trimester, based on residence <5 km of the monitoring station. When exposure estimates were based on city-wide monitoring, the odds ratio was significantly less than one (0.89, 95% CI: 0.84–0.95), suggesting that risk of pre-term birth decreased with a 1 ppm increase in air carbon monoxide concentration. The city-wide mean 8-hour maximum air carbon monoxide concentration was 0.9 ppm (SD: 0.68).

*Studies of Low Birth Weight.* Results of studies of low birth weight (i.e., <2,500 g or <10<sup>th</sup> percentile weight for gestation age) have been mixed with most studies finding associations between increasing ambient air carbon monoxide concentrations and decreasing birth weight (Bell et al. 2007; Gouveia et al. 2004; Ha et al. 2001; Lee et al. 2003a; Liu et al. 2003, 2007; Maisonet et al. 2001; Mannes et al. 2005; Medeiros and Gouveia 2005; Parker et al. 2005; Ritz and Yu 1999; Salam et al. 2005; Wilhelm and Ritz 2005). In some studies that found significant regression coefficients (i.e.,  $\beta$  coefficients) relating air carbon monoxide concentration to birth weight (g), magnitudes of the changes in birth weight were not sufficient to produce elevated risk of low birth weight, where low birth weight is defined categorically as a birth weight <2,500 g. Furthermore, the relative contribution of co-exposures to other air pollutants has not been completely elucidated in epidemiology studies. In many, but not all, studies showing significant associations with birth weight, these associations were not significant after adjustments for exposures to other co-pollutants (e.g., NO<sub>2</sub>, O<sub>3</sub>, PM<sub>10</sub>). All of these studies described below estimated risks after adjustment for typical co-variables that are thought to affect birth weight (e.g., infant sex, maternal age, race/ethnicity, and education, interval since previous live birth, previous low birth weight or preterm infant, level of prenatal care, birth season, parity, and gestational age). Collectively, the reported studies of birth weight outcomes provide supportive evidence for an association between ambient air carbon monoxide concentrations and low birth weight.

A series of studies conducted in California found such associations that were stronger when carbon monoxide was the only pollutant considered in the model and attenuated when adjustments were made for co-pollutants (Parker et al. 2005; Ritz and Yu 1999; Salam et al. 2005; Wilhelm and Ritz 2005). Mean air carbon monoxide concentrations in these studies ranged from 0.75 to 2.4 ppm and the highest value reported was 6.70 ppm (Wilhelm and Ritz 2005). The Ritz and Yu (1999) study found elevated risk of low birth weight in a sample of 125,573 births (1989–1993). The odds ratio (>5.5 vs. <2.2 ppm during third trimester) was 1.22 (95% CI: 1.03–1.44) when carbon monoxide alone was the only air pollutant considered and 1.38 (95% CI: 0.86–2.22) when NO<sub>2</sub>, O<sub>3</sub>, and PM<sub>10</sub> were included in the model. The odds ratios remained significant after adjustment for NO<sub>2</sub>, O<sub>3</sub>, and PM<sub>10</sub>, when the analysis was restricted

## 3. HEALTH EFFECTS

to women <20 years of age (multi-pollutant model: 5.08, 95% CI: 1.77–16.63; carbon monoxide alone: 1.54, 95% CI: 1.07–2.22). In a follow-up study (Wilhelm and Ritz 2005; 1994–2000) that included 136,134 births, the odds ratio (>1.84 vs. <0.96 ppm) was estimated to be 1.36 (95% CI: 1.04–1.76) for women who resided  $\leq$ 1 mile from an air monitoring station; however, risk was no longer significant when NO<sub>2</sub> and O<sub>3</sub> (1.29, 95% CI: 0.92–1.81), or NO<sub>2</sub>, O<sub>3</sub>, and PM<sub>10</sub> were included in the model (1.39, 95% CI: 0.77–2.49). A smaller cohort study (1975–1987) of 3,901 births found significant associations between ambient air carbon monoxide concentrations and low birth weight and intrauterine growth retardation (IUGR, <15<sup>th</sup> percentile of weight at gestational age; Salam et al. 2005). The estimated odds ratio for IUGR was 1.2 (95% CI: 1.0–1.4) for a 1.4 ppm increase in air carbon monoxide concentration in the first trimester (carbon monoxide considered alone). Although the odds ratio for low birth weight was not significant (1.0, 95% CI: 0.7, 1.5), a significant association between increasing air carbon monoxide concentration and lower birth weight was observed when birth weight was treated as a continuous variable ( $\beta$  coefficient: -21.7 g, 95% CI: -42.3, -1.1), which persisted when O<sub>3</sub> was included in the model (-28.6, 95% CI: -50.4, -6.9). A cohort study of 18,247 full-term births in California (2000) did not find a significant association between ambient air carbon monoxide concentration and risk of SGA (i.e., small for gestational age, <10<sup>th</sup> percentile of weight for gestational age; Parker et al. 2005). The SGA odds ratio (>0.97 vs. <0.57 ppm for entire pregnancy) was 0.95 (95% CI: 0.81–1.12) when carbon monoxide was considered alone and 0.82 (95% CI: 0.68–0.99) when adjusted for PM<sub>2.5</sub>. When treated as a continuous variable, a significant association between increasing air carbon monoxide concentration and lower birth weight was found when carbon monoxide was modeled alone ( $\beta$  coefficient: -20.5 g, 95% CI: -40.1, -0.8); however, the association was not significant when adjusted for PM<sub>2.5</sub> (2.6, 95% CI: -20.6, 25.8).

Other studies of low birth weight conducted in the United States have yielded mixed results. A cohort study of 89,557 full term births conducted in five northeastern cities (1994–1996) estimated the odds ratio for low birth weight to be 1.31 (95% CI: 1.06–1.62) for a 1 ppm increase in air carbon monoxide concentration during the third trimester (Maisonet et al. 2001). When stratified by race, the effect was only significant among African Americans for the first trimester (odds ratio: 1.43, 95% CI: 1.18–1.74) and third trimester (odds ratio: 1.75, 95% CI: 1.50–2.04). Results of multi-pollutant models were not reported. A cohort study of 358,504 births in Connecticut and Massachusetts found a significant association between increasing ambient air carbon monoxide concentration (for entire pregnancy) and reduced birth weight ( $\beta$  coefficient: -16.2, 95% CI: -19.7, -12.6), which persisted when adjusted for NO<sub>2</sub>, O<sub>3</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, or SO<sub>2</sub> (Bell et al. 2007). However, the magnitude of the change in birth weight

## 3. HEALTH EFFECTS

did not result a significant odds ratio (odds ratio: 1.028, 95% CI: 0.983–1.074). Mean air carbon monoxide concentrations in the above studies ranged from 0.65 to 0.95 ppm.

Two studies examined low birth weight or IUGR in Canada (Liu et al. 2003, 2007). The Liu et al. (2003) study included 229,085 births in the Vancouver area (1985–1998), and found a significant association between ambient air carbon monoxide concentration (during first month of pregnancy) and IUGR risk per 1 ppm increase in carbon monoxide (odds ratio: 1.06, 95% CI: 1.01–1.10), which persisted when adjusted for NO<sub>2</sub>, O<sub>3</sub>, or SO<sub>2</sub> (OR estimates not reported). In a larger area study (Liu et al. 2007) that included 386,202 births (1995–2000), odds ratios were significant for IUGR when ambient air carbon monoxide concentrations were considered during the first trimester (1.18, 95% CI: 1.14–1.22), second trimester (1.14, 95% CI: 1.10–1.18), or third trimester (1.19, 95% CI: 1.14–1.24). The odd ratios remained significant after adjustment for exposures to NO<sub>2</sub>, and PM<sub>2.5</sub>. The mean air carbon monoxide concentration in both studies was 1 ppm and the highest value reported was 5.6 ppm.

Two studies conducted in Seoul, South Korea (1996–1997, 1996–1998) found significantly increased risk of low birth weight in cohorts of 276,763 births (Ha et al. 2001) and 388,105 births (Lee et al. 2003a). In the Ha et al. (2001) study, the relative risk of low birth weight for a 0.42 ppm increase in ambient air carbon monoxide concentration during the first trimesters was 1.08 (95% CI: 1.04–1.12). In the Lee et al. (2003a) study, the odds ratios for a 0.5 ppm increase in carbon monoxide concentration were 1.04 (95% CI: 1.01–1.07) for the first trimester, 1.03 (1.00–1.06) for the second trimester carbon monoxide, 0.96 (0.93–0.99) for the third trimester carbon monoxide, and 1.05 (1.01–1.09) for carbon monoxide concentration during the entire pregnancy. Results of multi-pollutant models were not reported. The mean air carbon monoxide concentration was 1.2 ppm (range: 0.4–3.4 ppm).

A cohort study conducted in Sydney, Australia (1998–2000) included 138,056 births and found a significant association between increased ambient air carbon monoxide concentration and decreasing birth weight (Mannes et al. 2005). For carbon monoxide exposure during the second trimester, an increase of 1 ppm was associated with a 28.7 g reduction birth weight (95% CI: 7.76–50.98). The association did not persist when the regression models were adjusted for exposure to NO<sub>2</sub>, or to PM<sub>10</sub> in two-pollutant models. Furthermore, the magnitude of the effect did not result in a significant risk of SGA (<2 SD of mean weight for gestational age). The mean ambient carbon monoxide concentration of 0.8 ppm (range: 0–4.6).

## 3. HEALTH EFFECTS

Two cohort studies conducted in Sao Paulo, Brazil also found a significant association between increasing ambient air carbon monoxide concentration and decreasing birth weight (Gouveia et al. 2004; Medeiros and Gouveia 2005). In the Gouveia et al. (2004) study, in a cohort of 179,460 births (1997), a 1 ppm increase in ambient air carbon monoxide concentration during the first trimester was associated with a 23.1 g reduction in birth weight (95% CI: 4.9.0–41.3); however, the effect was not of sufficient magnitude to result in elevated risk of low birth weight. The Medeiros and Gouveia (2005) included 311,735 births (1998–2000) and estimated a reduction in birth weight of 11.9 g (95% CI: 8.2–15.5) per 1 ppm increase in first trimester carbon monoxide concentration. Here again, odds ratios for low birth weight were not significant. The ambient carbon monoxide concentrations in these studies ranged from approximately 1 to 11 ppm.

*Studies of Congenital Anomalies.* Possible associations between exposures to carbon monoxide and developmental anomalies have been evaluated in three case-control studies (Gilboa et al. 2005; Hwang and Jaakkola 2008; Ritz et al. 2002). Although two of the studies found significant associations between carbon monoxide and heart anomalies (i.e., ventricular septic defects (Ritz et al. 2002) and conotruncal defects (Gilboa et al. 2005), the two studies did not corroborate their findings. The Ritz et al. (2002) study did not find a significant association of conotruncal defects, and the Gilboa et al. (2005) study did not find a significant association with ventricular septal defects. The evaluation in the Hwang and Jaakkola (2008) study was limited to cleft lip and palate and found no association with carbon monoxide, consistent with the Ritz et al. (2002) study. Collectively, these studies do not provide convincing evidence for associations between exposure to carbon monoxide and congenital anomalies.

The Ritz et al. (2002) study included 3,549 cases and 10,649 controls born during the period 1987–1993 in southern California. The study included various categories of anomalies including heart and pulmonary, conotruncal, cleft lip and palate, and chromosomal defects. Of these, significant odds ratios were found for ventricular septal defects with increasing ambient air carbon monoxide concentration during the second month of pregnancy. Odds ratios increased with increasing ambient air carbon monoxide concentration during the second month of pregnancy (<1.14 ppm reference): 1.14–1.60 ppm, odds ratio 1.62 (95% CI: 1.05–2.48), 1.60–<2.47 ppm, odds ratio 2.09 (95% CI: 1.19–3.67); ≥2.47 ppm, odds ratio 2.95 (95% CI: 1.44–6.05). Significant odds ratios for ventricular septal defects persisted in multi-pollutant models.

A case-control study conducted in Texas (1996–2000) included 4,594 cases and 3,667 controls (Gilboa et al. 2005). Anomalies considered in the study included heart and pulmonary, conotruncal, and cleft lip and

## 3. HEALTH EFFECTS

palate. Of these, a significant dose trend was evident for Tetralogy of Fallot (a conotruncal anomaly of the heart), with a significant odds ratio ( $>0.7$  vs.  $<0.4$  ppm) for ambient air carbon monoxide concentration during the 3<sup>rd</sup> to 8<sup>th</sup> week of pregnancy (2.04 95% CI: 1.26–3.29). The odds ratio for any conotruncal anomaly was also significant at the highest exposure category ( $>0.7$  vs.  $<0.4$  ppm) (1.46 95% CI: 1.03–2.08). Results of multi-pollutant models were not reported.

A case-control study conducted in Taiwan China (2001–2003) included 653 cases of cleft lip and/or palate and 6,530 controls and examined associations with ambient air carbon monoxide concentrations measured during the first trimester of pregnancy (Hwang and Jaakkola 2008). Air carbon monoxide concentration was not associated with odds ratio for either anomaly in single-pollutant or multiple-pollutant models ( $O_3$ ,  $PM_{10}$ , or  $SO_2$ ).

*Studies of Neonatal and Infant Mortality.* Studies of possible associations between ambient air carbon monoxide concentrations and neonatal and infant mortality have yielded mixed results (Ha et al. 2003; Lin et al. 2004a; Ritz et al. 2006; Tsai et al. 2006a; Woodruff et al. 2008; Yang et al. 2006). Although increased risk of infant mortality in association with increasing ambient air carbon monoxide concentrations has been reported (Ha et al. 2003; Ritz et al. 2006), these outcomes have not been rigorously examined for the possible confounding by birth weight and/or gestational age, two relatively influential variables in predicting infant mortality. In one study that did examine the effect of stratification by gestational age or birth weight, the association between air carbon monoxide concentration and mortality from all causes or sudden infant death syndrome (SIDS) persisted in single-pollutant models, but not in models that adjusted for  $NO_2$ ,  $PM_{10}$ , and  $O_3$  (Ritz et al. 2006). Collectively, these studies do not provide convincing evidence for an association between ambient air carbon monoxide concentrations and neonatal or infant mortality.

A time-series analysis conducted in Sao Paulo, Brazil (1998–2000) did not find a significant association between ambient air carbon monoxide concentrations and neonatal mortality (age 1–28 days; Lin et al. 2004a). The mean air carbon monoxide concentration was 2.8 ppm (range: 0.5–10). A case-control study conducted in California (1989–2000) included 13,146 cases of infant death (age 28 days–1 year) and 151,015 controls (Ritz et al. 2006). Results varied depending on the temporal window for averaging air carbon monoxide concentration. The odds ratio for infant death for a 1 ppm increase in ambient air carbon monoxide concentration, averaged for 2 months prior to death, was 1.11 (95% CI: 1.06–1.16), with higher risk for deaths attributable to SIDS (odds ratio: 1.19; 95% CI: 1.10–1.28). However, risks were not significant after adjustment for co-pollutants ( $NO_2$ ,  $PM_{10}$ , and  $O_3$ ). For an averaging period of

## 3. HEALTH EFFECTS

2 weeks prior to death, risks were significant for all respiratory-related deaths that occurred between 28 days and 1 year (odds ratio: 1.14, 95% CI: 1.03–1.25) or from age 28 days to 3 months (odds ratio: 1.20, 95% CI: 1.02–1.40). Significant respiratory deaths persisted in multi-pollutant models. The association between air carbon monoxide concentration and SIDS mortality risk was persistent when the data were stratified by gestational age and birth weight; however, in these strata, the associations were not significant after adjustment for exposure to NO<sub>2</sub>, PM<sub>10</sub>, and O<sub>3</sub>. The mean air carbon monoxide concentration in the study was 1.6 ppm (range: 0.4–3.4).

A retrospective cohort study analyzed deaths that occurred between age 28 days and 1 year in a cohort of 3,590,134 births (6,939 deaths) across U.S. counties having a population >250,000 residents (1989–2000; Woodruff et al. 2008). Ambient air carbon monoxide concentration (median: ppm, 25<sup>th</sup>–75<sup>th</sup> percentile range: 0.7–0.9) during the first 2-months postpartum was not associated with infant death of any category, including all causes (odds ratio: 1.01, 95% CI: 0.95–1.07), respiratory (odds ratio: 1.14, 0.93–1.40), or SIDS (odds ratio: 0.88, 0.76–1.03). The latter values were estimated from single-pollutant models; however, similar results were obtained in multi-pollutant models that adjusted for O<sub>3</sub> and SO<sub>2</sub>.

A time-series study conducted in Seoul, South Korea (1995–1999) found a significant risk of respiratory-related mortality (age 1 month–1 year) in association with increasing ambient air carbon monoxide concentration measured on the day of death (odds ratio: 1.388, 95% CI: 1.009–1.911, per 0.57 ppm increase in carbon monoxide concentration), but not for deaths from all causes (Ha et al. 2003). Two case-crossover studies conducted in Taiwan, China (1994–2000) did not find a significant association between ambient air carbon monoxide concentration and infant mortality (Tsai et al. 2006a; Yang et al. 2006). The Tsai et al. (2006a) study included 206 cases (age 28 days–1 year) and the mean ambient air carbon monoxide concentration was 0.83 ppm (range: 0.23–1.77). The Yang et al. (2006) study included 471 cases (age 27 days–1 year); the mean ambient air carbon monoxide concentration was 1.58 ppm (range: 0.3–4.8).

***Animal Studies.*** Numerous studies on developmental effects of gestational and early postnatal exposure to carbon monoxide have been conducted in animals. Study details are summarized in Table 3-9. In general, most studies evaluated effects of relatively low carbon monoxide concentrations (i.e., ≤300 ppm), with exposure concentrations selected to produce maternal COHb levels typically associated with smoking; however, studies did not consistently report maternal or fetal COHb levels. Studies in animals have examined effects of carbon monoxide exposure on numerous developmental outcomes, including several outcomes that have not been assessed in epidemiological studies (e.g., auditory and immune

Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals

Reference	Species	Exposure	COHb	Results
Effects on birth weight				
Astrup et al. 1972	Rabbits	Continuous exposure to 0, 90, or 180 ppm CO throughout gestation	<u>mCOHb</u> : 8–9% in 90 ppm CO group; 16–18% in 180 ppm CO group <u>fCOHb</u> : NR	<u>Birth weight</u> : Fetal weight decreased by 11 and 20% in the 90 or 180 ppm CO groups, respectively.  NOAEL (birth weight): Not established LOAEL (birth weight): 90 ppm CO (mCOHb 8–9%)
Carmines and Rajendran 2008	Rat	2 hours/day to 600 ppm CO on GDs 6–19 (nose-only exposure)	<u>mCOHb</u> : ~30% (measured on GD 20) <u>fCOHb</u> : NR	<u>Birth weight</u> : Significantly decreased by 11%, compare to control.  NOAEL (birth weight): Not established LOAEL (birth weight): 600 ppm CO (mCOHb ~30%)
Fechter and Annau 1977	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–21	<u>mCOHb</u> : 15% <u>fCOHb</u> : NR	<u>Birth weight</u> : Decreased by 4.8%, compared to controls (not statistically significant)  <u>Pre-weanling weight</u> : Pre-weanling weight significantly decreased, compared to control, on PNDs 4–21 (decreased by 16 and 27% on PNDs 4 and 21, respectively).  NOAEL (pre-weanling weight): Not established LOAEL (pre-weanling weight): 60 ppm CO (mCOHb 15%)
Fechter and Annau 1980	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>Birth weight</u> : Decreased by 7.6%, compared to controls.  NOAEL (pre-weanling weight): Not established LOAEL (pre-weanling weight): 150 ppm CO
Fechter et al. 1987a	Rats	Continuous exposure to 0, 75, 150, or 300 ppm CO from GD 0 through PND 10	<u>mCOHb</u> : 11, 18, and 27% in the 75, 150, or 300 ppm CO, respectively <u>fCOHb</u> : NR	<u>Birth weight</u> : Fetal weight decreased by 8.9 and 13.8% in 150 and 300 ppm CO groups, respectively compared to control; by PND 21, no effects of CO treatment on pup weight.  NOAEL (birth weight): 75 ppm CO (mCOHb 11%) LOAEL (birth weight): 150 ppm CO (mCOHb 18%)

**Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals**

Reference	Species	Exposure	COHb	Results
Penney et al. 1983	Rats	Continuous exposure to 0, 157, 166, or 200 ppm CO on GDs 5–22	<u>mCOHb</u> : 24.9% (in dams exposed to 200 CO, measured over period of PNDs 1–6 days) <u>fCOHb</u> : 21.8, 24.9, and 3.10–33.5% at 157, 166, or 200 ppm CO, respectively (at birth)	<u>Birth weight</u> : Statistically significant decrease (~10%) in all CO groups, compared to control.  NOAEL (birth weight): Not established LOAEL (birth weight): 157 ppm CO (fCOHb 21.8%)
Prigge and Hochrainer 1977	Rats	Continuous exposure to 0, 60, 125, 250, or 500 ppm CO on GDs 0–21	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>Birth weight</u> : Dose-related, statistically significant decrease (7.5–36.7%) in fetal weight at ≥125 ppm, compared to control.  NOAEL (birth weight): 60 ppm CO LOAEL (birth weight): 125 ppm CO
Storm and Fechter 1985b	Rats	Continuous exposure to 0, 75, 150, or 300 ppm CO for entire gestational period	<u>mCOHb</u> : 11.5, 18.5, and 26.8% in the 75, 150, or 300 ppm CO groups, respectively <u>fCOHb</u> : NR	<u>Birth weight</u> : Fetal weight decreased by 12.5% in 300 ppm CO group, compared to control.  NOAEL (birth weight): 150 ppm CO (mCOHb 18.5%) LOAEL (birth weight): 300 ppm CO (mCOHb 27%)
Tolcos et al. 2000b	Guinea pigs	10 hours/day 0 or 200 ppm CO from GDs 23 or 25 to GD 68	<u>mCOHb</u> : 8.5% <u>fCOHb</u> : 13%	<u>Birth weight</u> : Fetal weight decreased by 9.6% in CO group, compared to control.  NOAEL (birth weight): Not established LOAEL (birth weight): 200 ppm CO (mCOHb 8.5%)
Effects on central nervous system (central control of respiratory function)				
McGregor et al. 1998	Guinea pigs	10 hours/day 0 or 200 ppm CO from GDs 23 or 25 to GD 68	<u>mCOHb</u> : 8.5% <u>fCOHb</u> : 13%	<u>CNS development (function)</u> : On PND 4, pups from CO-exposed dams exhibited abnormal respiratory responses to asphyxia and hypercapnia; responses consistent with developmental alterations in brain stem.  NOAEL (CNS function): Not established LOAEL (CNS function): 200 ppm CO (mCOHb 8.5%)

**Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals**

Reference	Species	Exposure	COHb	Results
Effects on central nervous system (behavioral)				
De Salvia et al. 1995	Rats	0, 75, or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : ~15% for 150 ppm CO (NR for 75 ppm CO) <u>fCOHb</u> : NR	<u>CNS development (behavioral effects)</u> : Significant impairment of acquisition (at 3 and 18 months) and reacquisition (at 18 months) of conditioned avoidance behavior following gestational exposure of pups to 150 ppm CO.  NOAEL (CNS behavioral): 75 ppm CO LOAEL (CNS behavioral): 150 ppm CO
Fechter and Annau 1977	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–21	<u>mCOHb</u> : 15% <u>fCOHb</u> : NR	<u>CNS development (behavioral effects)</u> : During the preweaning period, CO pups were less active and showed decreased response to L-DOPA-induced movement stimulation.  NOAEL (CNS behavioral): Not established LOAEL (CNS behavioral): 60 ppm CO (mCOHb 15%)
Fechter and Annau 1980	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>CNS development (behavioral effects)</u> : Pups of exposed dams showed impaired righting reflexes, impaired negative geotaxis, and delayed homing behavior.  NOAEL (CNS behavioral): Not established LOAEL (CNS behavioral): 150 ppm CO
Giustino et al. 1999	Rats	Continuous exposure to 75 and 150 ppm CO on GDs 0–20	<u>mCOHb</u> : 7.3 and 16.1% in 75 and 150 ppm CO groups, respectively <u>fCOHb</u> : NR	<u>CNS development (behavioral effects)</u> : Alterations in habituation and working memory in young adult male offspring at PND 40, including decreased time of exploration of novel objects (75 and 150 ppm CO) and lack of habituation after the second exposure to a previously viewed object (150 ppm CO); decreased spontaneous motor activity in open field test (75 and 150 ppm CO).  NOAEL (CNS behavioral): Not established LOAEL (CNS behavioral): 75 ppm CO

Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals

Reference	Species	Exposure	COHb	Results
Mactutus and Fechter 1985	Rats	Continuous exposure to 150 ppm CO over entire gestational period	<u>mCOHb</u> : 15.6% <u>fCOHb</u> : NR	<u>CNS development (behavioral effects)</u> : Prenatal CO exposure induced learning and memory deficits in male and female offspring, as indicated by impairments in reacquisition performance, an index of retention, on PND 31 and decrements in two-way conditioned avoidance behavior (flashing light warnings followed by mild foot shock) on PND 120.  NOAEL (CNS behavioral): Not established LOAEL (CNS behavioral): 150 ppm CO (mCOHb 15.6%)
Singh 1986	Mice	Continuous exposure to 0, 65, or 135 ppm CO on GDs 7–18	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>CNS development (behavioral effects)</u> : On PNDs 1–10, pups of exposed dams showed impaired righting reflexes (125 ppm CO), geotaxis (125 ppm CO), and aerial righting reflex (65 and 125 ppm CO)  NOAEL (CNS behavioral): Not established LOAEL (CNS behavioral): 65 ppm CO
Tattoli et al. 1999	Rats	Continuous exposure to 0, 75, or 150 ppm CO on PNDs 1–10	<u>COHb in pups</u> : NR	<u>CNS development (behavioral effects)</u> : Assessed at 3 and 18 months of age, no effects of early postnatal exposure on acquisition and reacquisition of an active avoidance task in both adult and aged rats.  NOAEL: 150 ppm CO LOAEL: Not established

**Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals**

Reference	Species	Exposure	COHb	Results
Effects on central nervous system (neurotransmitter)				
Cagiano et al. 1998	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : ~16% <u>fCOHb</u> : NR	<u>CNS development (neurotransmitter changes)</u> : On PND 80, no CO effect on basal extracellular levels of dopamine in the nucleus accumbens; however, the amphetamine-induced dopamine release was decreased in pups of CO-exposed dams.  NOAEL (CNS neurotransmitter): Not established LOAEL (CNS neurotransmitter): 150 ppm CO (mCOHb 16%)
Fechter and Annau 1977	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–21	<u>mCOHb</u> : 15% <u>fCOHb</u> : NR	<u>CNS development (neurotransmitter changes)</u> : Brain dopamine content decreased by 45 and 25% on PNDs 1 and 4, respectively.  NOAEL (CNS neurotransmitter): Not established LOAEL (CNS neurotransmitter): 60 ppm CO (mCOHb 15%)
Fechter and Annau 1980	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>CNS development (neurotransmitter changes)</u> : On PND 1, no effect of CO exposure on forebrain and hindbrain levels of dopamine or norepinephrine.  NOAEL (CNS neurotransmitter): 150 ppm CO LOAEL (CNS neurotransmitter): Not established
Fechter et al. 1987a	Rats	Continuous exposure to 0, 75, 150, or 300 ppm CO from GD 0 through PND 10	<u>mCOHb</u> : 11, 18, and 27% In the 75, 150, or 300 ppm CO, respectively <u>fCOHb</u> : NR	<u>CNS development (neurotransmitter changes)</u> : On PND 21, pups of CO-exposed dams showed elevated dopamine levels in neostriatum (150 and 300 ppm CO).  NOAEL (CNS neurotransmitter): 75 ppm CO (mCOHb 11%) LOAEL (CNS neurotransmitter): 150 ppm CO (mCOHb 18%)
Storm and Fechter 1985a	Rats	Continuous exposure to 0 or 150 ppm for entire gestational period	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>CNS development (neurotransmitter changes)</u> : Norepinephrine levels (amount and concentration) in cerebellum (but not cortex) elevated from PNDs 14–42, compared to controls.  NOAEL (CNS neurotransmitter): Not established LOAEL (CNS neurotransmitter): 150 ppm CO

Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals

Reference	Species	Exposure	COHb	Results
Storm and Fechter 1985b	Rats	Continuous exposure to 0, 75, 150, or 300 ppm CO for entire gestational period	<u>mCOHb</u> : 11.5, 18.5, and 26.8% in the 75, 150, or 300 ppm CO groups, respectively <u>fCOHb</u> : NR	<u>CNS development (neurotransmitter changes)</u> : On PND 21, weight of cerebellum decreased by 8.5 and 11.4% in 150 and 300 ppm CO groups, respectively; dose-related trend for decreased norepinephrine concentration in pons/medulla (but not neocortex, hippocampus or cerebellum); decreased serotonin concentration in pons/medulla in 150 and 300 ppm CO groups. On PND 42, weight of cerebellum decreased by 8.3% in 300 ppm CO group; dose-related trend for increased norepinephrine concentration neocortex and hippocampus (but not pons/medulla or cerebellum).  NOAEL (CNS neurotransmitter): 75 ppm CO (mCOHb 11.5%) LOAEL (CNS neurotransmitter): 150 ppm CO (mCOHb 18.5%)
Tolcos et al. 2000b	Guinea pigs	10 hours/day 0 or 200 ppm CO from GDs 23 or 25 to GD 68	<u>mCOHb</u> : 8.5% <u>fCOHb</u> : 13%	<u>CNS development (neurotransmitter changes)</u> : Effects on cholinergic and catecholaminergic pathways in the medulla, including significant decrease in tyrosine hydroxylase-immunoreactivity in the nucleus tractus solitarius, dorsal motor nucleus of the vagus, area postrema, intermediate reticular nucleus, and the ventrolateral medulla; significant increase in choline acetyltransferase-immunoreactivity in the dorsal motor nucleus of the vagus and hypoglossal nucleus, compared with controls.  NOAEL (CNS neurotransmitter): Not established LOAEL (CNS neurotransmitter): 200 ppm CO (mCOHb 8.5%)

Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals

Reference	Species	Exposure	COHb	Results
Effects on central nervous system (auditory system)				
Lopez et al. 2003	Rats	Continuous exposure to 0, 12, or 25 ppm CO on PNDs 8–22	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<p><u>CNS development (auditory system)</u>: Pups examined on PND 27, nerve terminals innervating inner hair cells were swollen, with cytoplasmic vacuolization and atrophy (25 ppm CO); in the 8<sup>th</sup> cranial nerve at the level of the internal auditory canal, fibers showed “distorted myelin” with vacuolization (25 ppm CO); in spiral ganglion neurons of the organ of corti, decreased immunoreactivity of enzymes cytochrome oxidase, NADH-TR, and calcium-mediated myosin ATPase (25 ppm CO); in organ of corti, decreased immunostaining of neurofilament and myelin basic protein (25 ppm CO).</p> <p>NOAEL (CNS auditory): 12 ppm CO LOAEL (CNS auditory): 25 ppm CO</p>
Lopez et al. 2008	Rats	10–18 hours/day to 25 ppm CO on GDs 5–20 or on GDs 5–20 plus PNDs 5–20	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<p><u>CNS development (auditory system)</u>: Similar effects in pups exposed during gestation only and during gestation and PNDs 5–20. On PND 3, no morphological deterioration, or loss of inner or outer hair cells; on PND 20, vacuolization on the afferent terminals at the basal portion of the cochlea; type I spiral ganglia neurons and afferent nerve fibers showed decreased neurofilament-immunoreactivity.</p> <p>NOAEL (CNS auditory): Not established LOAEL (CNS auditory): 25 ppm CO</p>

Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals

Reference	Species	Exposure	COHb	Results
Stockard-Sullivan 2003	Rats	22 hour/day to 12, 25, 50, or 100 ppm CO on PNDs 6–22	COHb in pups: 5–7% in 100 ppm CO group (not measured in other groups)	<p><u>CNS development (auditory system)</u>: Decreased in otoacoustic emissions at 50 and 100 ppm CO, measured at age 22–24 days; attenuation of the amplitude of the 8<sup>th</sup> nerve action potential in all CO groups, measured at age 22–24 days, with effects persisting though age 73 days in the 50 and 100 ppm CO group (not examined at later ages in other groups); no effect on auditory brain stem conduction times in any CO group, measured at ages 22, 30, or 51–52 days .</p> <p>NOAEL (CNS auditory): Not established LOAEL (CNS auditory): 12 ppm CO</p>
Webber et al. 2003	Rats	Continuous exposure to 0, 12.5, 25, or 50 ppm CO on GDs 8 through 20–22	<p><u>mCOHb</u>: NR <u>fCOHb</u>: NR</p>	<p><u>CNS development (auditory system)</u>: Examined effects of CO exposed on central inferior colliculus (an area of the mid-brain with auditory integrative functions) during period of synaptogenesis/auditory development. Immunostating of c-Fos (marker of neuronal activation in the nervous system) showed significant decrease in C-Fos immunoreactivity in the central inferior colliculus on PND 27 and PNDs 75–77 in all CO groups. Results indicate that gestational exposure to CO can produce changes in the auditory system of rats that persist into adulthood.</p> <p>NOAEL (CNS auditory): Not established LOAEL (CNS auditory): 12.5 ppm CO</p>

**Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals**

Reference	Species	Exposure	COHb	Results
<b>Effect on peripheral nervous system</b>				
Carratu et al. 1993	Rats	Continuous exposure to 0, 75, or 100 ppm CO on GDs 0–20	<u>mCOHb</u> : ~15% for 150 ppm CO (NR for 75 ppm CO) <u>fCOHb</u> : NR	<u>PNS development (electrophysiological changes)</u> : In sciatic nerves isolated on PND 40, inactivation kinetics of inactive sodium channels were significantly slowed and negative shift in sodium equilibrium potential was observed in both CO groups; on PND 270, sciatic nerves showed negative shift in sodium equilibrium potential both CO groups.  NOAEL (PNS development): Not established LOAEL (PNS development): 75 ppm CO
Carratu et al. 2000a	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : 16.02% <u>fCOHb</u> : NR	<u>PNS development (neuronal development)</u> : Altered sphingomyelin composition of sciatic nerve, but no effect on motor activity in pups at PND 90.  NOAEL (PNS development): Not established LOAEL (PNS development): 150 ppm CO
Carratu et al. 2000b	Rats	Continuous exposure to 0, 75, or 100 ppm CO on GDs 0–20	<u>mCOHb</u> : 7.34 and 16.08% in the 75 and 150 ppm CO groups, respectively <u>fCOHb</u> : NR	<u>PNS development (neuronal development)</u> : Exposure of dams produced decreased myelin sheath thickness of nerve fibers (75 and 100 ppm CO) but not axon diameter or motor activity in pups at PNDs 40 and 90.  NOAEL (PNS development): Not established LOAEL (PNS development): 75 ppm CO
<b>Effects on cardiovascular system</b>				
Penney et al. 1982	Rats	Continuous exposure to 0 or 500 ppm CO on PNDs 1–32	<u>COHb in pups</u> : 38–42%	CVS (heart weight): Relative heart weight increased on PND 14 and remained above normal through PND 104.  NOAEL (developmental heart weight): Not established LOAEL (developmental heart weight): 500 ppm CO (COHb 38–42%)

Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals

Reference	Species	Exposure	COHb	Results
Penney et al. 1983	Rats	Continuous exposure to 0, 157, 166, or 200 ppm CO on GDs 5–22	<u>mCOHb</u> : 24.9% (in dams exposed to 200 CO, measured over period of PNDs 1–6 days) <u>fCOHb</u> : 21.8, 24.9, and 3.10–33.5% at 157, 166, or 200 ppm CO, respectively (at birth)	<u>CVS (heart weight)</u> : Dose-related, statistically significant increase in relative heart weight (10–108%) in all CO groups, compared to control.  NOAEL (developmental heart weight): Not established LOAEL (developmental heart weight): 157 ppm CO (fCOHb 21.8%)
Prigge and Hochrainer 1977	Rats	Continuous exposure to 0, 60, 125, 250, or 500 ppm CO on GDs 0–21	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>CVS (heart weight)</u> : Dose-related, statistically significant increase in relative heart weight (10–52%) in all CO groups, compared to control.  NOAEL (developmental heart weight): Not established LOAEL (developmental heart weight): 60 ppm CO
Sartiani et al. 2004	Rats	Continuous exposure to 0 or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>CVS (myocardial electrophysiological changes)</u> : In ventricular cardiocytes isolated from pups on PNDs 1–60, delayed development of age-related action potential duration shortening was observed. Delay was maximum at 4 weeks of age.  NOAEL (CVS function): Not established LOAEL (CVS function): 150 ppm CO

**Table 3-9. Effects of Gestational or Perinatal Exposure on Developmental Outcomes in Animals**

Reference	Species	Exposure	COHb	Results
<b>Effects on sexual behavior</b>				
Cagiano et al. 1998	Rats	Continuous exposure to 0, 75, or 150 ppm CO on GDs 0–20	<u>mCOHb</u> : ~7.5 and ~16% in 75 and 150 ppm CO groups, respectively <u>fCOHb</u> : NR	<u>Development of sexual behavior</u> : On PND 80, in pups of dams exposed to 150 ppm CO (but not 75 ppm CO), alterations in sexual behavior, including increase in mount to intromission latency, decrease in mount to intromission frequency, and decrease in ejaculation frequency.  NOAEL (developmental sexual behavior): 75 ppm CO LOAEL (developmental sexual behavior): 150 ppm CO
<b>Effects on immunological system</b>				
Giustino et al. 1993	Rats	Continuous exposure to 75 and 150 ppm CO on GDs 0–20	<u>mCOHb</u> : ~15% in 150 ppm CO group (NR in 75 ppm CO group) <u>fCOHb</u> : NR	<u>Immunological effects</u> : On PNDs 15 and 21, decreased splenic macrophage phagocytosis of <i>Candida albicans</i> (150 ppm CO); on PND 15, decreased splenic macrophage killing (75 and 150 ppm CO); no effects observed on PND 60.  NOAEL (immunological): Not established LOAEL (immunological): 75 ppm CO
Giustino et al. 1994	Rats	Continuous exposure to 75 and 150 ppm CO on GDs 0–20	<u>mCOHb</u> : ~15% in 150 ppm CO group (NR in 75 ppm CO group) <u>fCOHb</u> : NR	<u>Immunological effects</u> : On PND 21, significant decreased in number of leukocyte common antigen cells in 150 ppm CO group; no effects observed on PND 540.  NOAEL (immunological): 75 ppm CO LOAEL (immunological): 150 ppm CO (mCOHb ~15%)
<b>Effects on hematological system</b>				
Prigge and Hochrainer 1977	Rats	Continuous exposure to 0, 60, 125, 250, or 500 ppm CO on GDs 0–21	<u>mCOHb</u> : NR <u>fCOHb</u> : NR	<u>Hematological effects</u> : Dose-related, statistically significant decrease in hematocrit (14.8–34.2%) and hemoglobin (12.5–28.4%) at ≥250 ppm, compared to control.  NOAEL (hematological): 125 ppm CO LOAEL (hematological): 250 ppm CO

CNS = central nervous system; CO = carbon monoxide; COHb = carboxyhemoglobin; CVS = cardiovascular system; fCOHb = fetal COHb; GD = gestation day; LOAEL = lowest-observed-adverse-effect level; mCOHb = maternal COHb; NOAEL = no-observed-adverse-effect level; NR = not reported; PND = postnatal day

## 3. HEALTH EFFECTS

system development). Results of animal studies show adverse developmental effects of gestational and early postnatal carbon monoxide exposure, including decreased fetal weight, adverse central nervous system development, altered peripheral nervous system development, cardiac effects, altered sexual behavior, immunological effects, and hematological effects. In addition, some studies showed that developmental effects persisted beyond the postnatal period (Carratu et al. 1993, 2000a, 2000b; De Salvia et al. 1995; Mactutus and Fechter 1985; Stockard-Sullivan et al. 2003; Webber et al. 2003), although persistence of effects was not examined in all studies. The lowest LOAEL values for developmental effects were obtained in studies evaluating effects of carbon monoxide on the developing auditory system (i.e., LOAEL 12–25 ppm carbon monoxide); however, since other developmental outcomes were not assessed at this range of low carbon monoxide concentrations, it is not possible to determine if the developing auditory system is more sensitive to carbon monoxide exposure than other systems, or to identify a biological basis for such susceptibility.

*Birth Weight.* Several studies have evaluated effects of gestational exposure to carbon monoxide on birth weight in rats (Carmines and Rajendran 2008; Fechter and Annau 1977, 1980; Fechter et al. 1980, 1987a; Penney et al. 1983; Prigge and Hochrainer 1977; Storm and Fechter 1985b), rabbits (Astrup et al. 1972), and guinea pigs (Tolcos et al. 2000b). Results consistently show that gestational exposure to carbon monoxide significantly decreased birth weight and/or pre-weanling weight, with LOAEL values ranging from 60 ppm carbon monoxide (maternal COHb 15%) for continuous exposure during gestation (Fechter and Annau 1977) to 600 ppm (maternal COHb approximately 30%) for intermittent (2 hours/day) exposure (Carmines and Rajendran 2008). Decreases in birth weight exhibited exposure concentration-dependence (Astrup et al. 1972; Fechter et al. 1987a; Prigge and Hochrainer 1977).

*Central Nervous System Development.* Several studies have observed adverse effects on the developing central nervous system, including altered control of respiratory function, behavioral effects, changes in brain neurotransmitters levels, and auditory development. Gestational exposure of guinea pigs to 200 ppm carbon monoxide produced abnormal respiratory responses to asphyxia and hypercapnia, indicative of altered brain stem development (McGregor et al. 1998). In rat or mouse pups exposed to 60–150 ppm carbon monoxide during gestation, behavioral changes included decreased motor activity and response to stimulation (Fechter and Annau 1977, 1980), impaired righting reflexes (Fechter and Annau 1980; Singh 1986), and impaired homing and memory acquisition behaviors (Giustino et al. 1999; Mactutus and Fechter 1985). Effects on acquisition and reacquisition behavior persisted up to 18 months of age in rats exposed to 150 ppm carbon monoxide during gestation (De Salvia et al. 1995; Mactutus and Fechter 1985), although similar effects were not observed in pups exposed to 150 ppm carbon monoxide

## 3. HEALTH EFFECTS

during the early postnatal period (Tattoli et al. 1999). Gestational exposure of rats or guinea pigs to 60–300 ppm carbon monoxide produced altered levels of dopamine and norepinephrine in several regions of the brain (Cagiano et al. 1998; Fechter and Annau 1977; Fechter et al. 1987a; Storm and Fechter 1985a, 1985b; Tolcos et al. 2000b). Gestational or postnatal exposure to low levels of carbon monoxide (i.e., 12–25 ppm) has been shown to have effects on the developing auditory system. Gestational and/or early postnatal exposure of artificially-reared rat pups (i.e., reared in formula to allow exposure to the pups without exposure to the dams during nursing) to 25 ppm carbon monoxide produced morphological changes in the developing auditory system, including swelling, cytoplasmic vacuolization and atrophy of nerve terminals innervating inner hair cells, “distorted myelin” with vacuolization in the 8<sup>th</sup> cranial nerve at the level of the internal auditory canal, decreased immunoreactivity of the enzymes cytochrome oxidase, NADH-TR, and calcium-mediated myosin ATPase, and decreased immunostaining of neurofilament and myelin basic protein in the organ of corti, (Lopez et al. 2003, 2008). Limited physiological and acoustic testing of pups suggested potential functional impairment in the auditory system. Exposure of artificially-reared rat pups during the early postnatal period decreased action potential amplitude of the 8<sup>th</sup> cranial nerve at  $\geq 12$  ppm carbon monoxide, although the magnitude of the effect was not dose-dependent. Otoacoustic emissions showed a small reduction in amplitude (3 dB at two of three frequencies tested and one loudness level) at  $\geq 50$  ppm carbon monoxide, with effects on action potential amplitude of the 8<sup>th</sup> nerve persisting through age 73 days (Stockard-Sullivan et al. 2003). The reduction in amplitude of otoacoustic emissions observed in this study was relatively small. Larger effects are more typically associated with hearing loss (e.g., 10–29 dB across five or more frequencies). Gestational exposure gestation days ([GDs] 8–20) of dam-reared rat pups to  $\geq 12.5$  ppm carbon monoxide significantly decreased neuronal activation of the central inferior colliculus (the area of the brain with auditory integrative function), with effects observed through postnatal days 75–77 (Webber et al. 2003). Results indicate that carbon monoxide-induced effects on brain auditory integrative neurons can persist into adulthood.

*Peripheral Nervous System Development.* Gestational exposure to carbon monoxide produced effects on peripheral nervous system development. In sciatic nerves of pups exposed during gestation, myelin sheath thickness was decreased and sphingomyelin composition was altered at 75–150 ppm carbon monoxide, with changes persisting through age 90 days (Carratu et al. 2000a, 2000b). Altered nerve transmission, indicative of slowed inactivation kinetics of inactive sodium channels, was observed in sciatic nerves from pups exposed to 75 ppm carbon monoxide during gestation, with changes persisting to age 270 days (Carratu et al. 1993).

### 3. HEALTH EFFECTS

*Cardiac Effects.* Dose-related increases in heart weight was observed in rat pups exposed to 60–157 ppm carbon monoxide during gestation (Penney et al. 1983; Prigge and Hochrainer 1977) or to 500 ppm carbon monoxide during the early postnatal period (Penney et al. 1982). The development of cardiomyopathy is consistent with a pathophysiological compensatory response to hypoxia. In ventricular cells isolated from pups exposed to 150 ppm carbon monoxide (but not 75 ppm carbon monoxide) during gestation, age-related decreases in action potential duration was delayed (Sartiani et al. 2004).

*Sexual Behavior.* In male pups exposed to 150 ppm carbon monoxide during gestation, altered development of sexual behavior was observed at age 80 days (Cagiano et al. 1998). Effects included increased mount to intromission latency, decreased mount to intromission frequency, and decreased ejaculation frequency.

*Immune System Effects.* In studies conducted by Giustino et al. (1993, 1994), gestational exposure produced reversible alterations in immune system function. Decreases in splenic macrophage phagocytosis of *Candida albicans* was observed in pups exposed to 150 ppm carbon monoxide and decreased splenic macrophage killing was observed in pups exposed to 75 and 150 ppm carbon monoxide (Giustino et al. 1993); effects did not persist at age 60 days. Rats exposed to 150 ppm carbon monoxide, but not 75 ppm carbon monoxide, showed an alteration in splenic immune cell populations, with a significant decreased in leukocyte common antigen cells; effects did not persist to age 540 days (Giustino et al. 1994).

*Hematological Effects.* Dose-related decreases in hematocrit and Hb were observed in pups exposed to 250 and 500 ppm carbon monoxide during gestation (Prigge and Hochrainer 1977). Effects are consistent with physiological compensatory responses to COHb-induced hypoxia.

#### **3.2.7 Cancer**

Epidemiological studies have examined possible associations between exposure to ambient air concentration of carbon monoxide and cancer mortality (described in Section 3.2.1, Death). A prospective study examined mortality in a cohort of 552,138 adults in 151 U.S. metropolitan areas, enrolled in the study in 1982 and followed through 1998 (Jerrett et al. 2003; Pope et al. 1995, 2002). Based on estimates of average ambient air carbon monoxide concentration during the period 1982–1998, increasing air carbon monoxide concentration was significantly associated with decreasing mortality from lung cancer. A time-series analysis conducted in New York City, New York (1985–1994) evaluated data

### 3. HEALTH EFFECTS

on cancer deaths with or without contributing respiratory diseases (De Leon et al. 2003). Estimates of cancer mortality risk were significant only in the strata with contributing respiratory disease (risk ratio was approximately 1.18, the 95% lower confidence limit >1) and did not persist when estimates were adjusted for ambient concentrations of PM<sub>10</sub> (risk ratio approximately 1.06, 95% CI: approximately 0.97–1.17).

Carbon monoxide has not been assessed for carcinogenicity using animal models.

#### 3.3 GENOTOXICITY

Few studies have evaluated the genotoxic effects of carbon monoxide exposure. Results of available studies are summarized in Table 3-10. An *in vitro* bacterial replication assay in *Escherichia coli* reported that exposure to carbon monoxide inhibited deoxyribonucleic acid (DNA) synthesis (Cairns and Denhardt 1968); however, interpretation of study results is compromised by poor reporting of experimental and analytical methods (e.g., carbon monoxide concentration, exposure conditions, and statistical analyses). Exposure of pregnant mice to carbon monoxide during gestation produced dose-related increases in micronuclei and sister chromatid exchanges in maternal bone marrow and fetal blood (Kwak et al. 1986). Pregnant mice were exposed to 0, 1,500, 2,500, or 3,500 ppm carbon monoxide for 10 minutes on GDs 5, 11, or 15 or to 0 or 500 ppm carbon monoxide for 1 hour on gestation days 0–6, 7–13, or 14–20; maternal bone marrow and fetal blood were examined for micronuclei formation and sister chromatid exchanges on GD 21. Increases in micronuclei and sister chromatid exchanges in maternal bone marrow and fetal blood were observed under all carbon monoxide exposure conditions, compared to unexposed controls. The magnitude of effect was similar in dams and fetuses. Although blood COHb levels were not measured in this study, a 1-hour exposure to 500 ppm carbon monoxide would be expected to produce a maternal blood COHb concentration of approximately 20–25% (based on CFK model prediction for rats).

#### 3.4 TOXICOKINETICS

Carbon monoxide exists in the environment as a gas (Henry's law constant >50,000 atm/mol fraction, 25 °C, see Table 4-2). As a result, humans can be exposed to carbon monoxide from breathing and/or skin contact with carbon monoxide in air. No information is available on the dermal absorption of carbon monoxide resulting from exposures to gaseous carbon monoxide. However, as is the case for other gases that are avidly absorbed from the lung (e.g., O<sub>2</sub>) during exposures to carbon monoxide in air, dermal absorption of carbon monoxide through intact skin would be expected to make a minor contribution to absorbed carbon monoxide, relative to inhalation. Although carbon monoxide can dissolve in water

## 3. HEALTH EFFECTS

**Table 3-10. Genotoxicity of Carbon Monoxide**

Species (test system)	Exposure	End point	Results	Reference
Prokaryotic organisms:				
<i>Escherichia coli</i> , TAMT	CO concentration not reported	DNA replication	+	Cairns and Denhardt 1968
Mammalian cells:				
Mice (ICR strain, pregnant females, 64/dose)	0, 1,500, 2,500, or 3,500 ppm CO for 10 minutes on GD 5, 11, or 16. Fetal blood and maternal bone marrow examined on GD 21	Sister chromatid exchange	Maternal bone marrow: + Fetal blood: +	Kwak et al. 1986
Mice (ICR strain, pregnant females, 50/dose)	0 or 500 ppm CO for 1 hour on GDs 0–6, 7–13, or 14–20. Fetal blood and maternal bone marrow examined on GD 21	Sister chromatid exchange	Maternal bone marrow: + Fetal blood: +	Kwak et al. 1986
Mice (ICR strain, pregnant females, 64/dose)	0, 1,500, 2,500, or 3,500 ppm CO for 10 minutes on GD 5, 11, or 16. Fetal blood and maternal bone marrow examined on GD 21	Chromosomal aberrations, micronucleus assay	Maternal bone marrow: + Fetal blood: +	Kwak et al. 1986
Mice (ICR strain, pregnant females, 50/dose)	0 or 500 ppm CO for 1 hour on GDs 0–6, 7–13, or 14–20. Fetal blood and maternal bone marrow examined on GD 21	Chromosomal aberrations, micronucleus assay	Maternal bone marrow: + Fetal blood: +	Kwak et al. 1986

+ = positive result; CO = carbon monoxide; GD = gestation day

## 3. HEALTH EFFECTS

(23 mL CO/L water, 20 °C), appreciable concentrations in drinking water would occur only at very high partial pressures of carbon monoxide in air, conditions in which inhalation would be the dominant absorption pathway. Since the only relevant pathway of exposure to humans is the inhalation pathway, oral and dermal exposures are not considered further in the discussion of the toxicokinetics of carbon monoxide,


### 3.4.1 Absorption

Inhaled carbon monoxide is rapidly and extensively absorbed into blood (Benignus et al. 1994; Burge and Skinner 1995; Peterson and Stewart 1970, 1975; Tikuisis et al. 1987b). Inhaled carbon monoxide is transported to lung alveoli as result of convective forces in the respiratory tract and diffusion. At the alveolar gas-blood interface, carbon monoxide dissolves into pulmonary capillary plasma and, from plasma, diffuses into erythrocytes and other tissues (Figure 3-3). Binding of carbon monoxide by erythrocyte Hb (to form COHb) contributes to maintaining relatively low concentrations of dissolved carbon monoxide in erythrocyte cytosol and a partial pressure gradient to drive carbon monoxide transfer from alveolar air to blood. Diffusion of carbon monoxide into erythrocytes and binding of carbon monoxide to Hb is sufficiently rapid that near equilibrium is achieved between carbon monoxide partial pressures in alveolar air and alveolar end capillary (arterial) blood (Chakraborty et al. 2004). Continued uptake of carbon monoxide results from systemic elimination processes that lower mixed venous blood carbon monoxide concentrations returned to alveolar capillaries (Benignus et al. 1994). The primary elimination processes include diffusive transfer of carbon monoxide to extravascular tissues and binding of carbon monoxide to intracellular heme proteins (e.g., muscle myoglobin), and oxidative metabolism of carbon monoxide to CO<sub>2</sub>. Seconds after initiating inhalation exposures to carbon monoxide, COHb is detected in arterial blood. During 2–6 minutes of exposure, arterial COHb concentrations were approximately twice venous COHb concentrations and approached unity within 3–5 minutes of cessation of exposure (Benignus et al. 1994). With continued exposure to a constant air carbon monoxide concentration, a steady-state would be achieved within 200–500 minutes (Figure 3-4; Bruce and Bruce 2003; Peterson and Stewart 1970, 1975). At that point, net absorption diminishes to a value determined, primarily, by metabolic elimination of carbon monoxide (see Section 3.4.3, Metabolism).

Major factors that control absorption of carbon monoxide include those that affect delivery of inhaled air to the alveolar region of the lung and those that affect lung carbon monoxide diffusion capacity. Lung diffusion capacity refers to the driving force for diffusion of carbon monoxide across the alveolar air-


## 3. HEALTH EFFECTS

**Figure 3-3. Summary of Carbon Monoxide Uptake, Distribution, and Elimination Pathways**



## 3. HEALTH EFFECTS

**Figure 3-4. Temporal Profile of Blood Carboxyhemoglobin (%) for Hypothetical Continuous Exposures to Air Carbon Monoxide at Concentrations of 25, 200, or 1,000 ppm**



Source: Peterson and Stewart 1975

## 3. HEALTH EFFECTS

blood interface and is reflected in the carbon monoxide partial pressure difference between inspired and exhaled air. In general, factors that increase alveolar ventilation, cardiac output, and gas diffusing capacity of the lung, while maintaining alveolar ventilation perfusion matching (i.e., proportional increases in alveolar ventilation rate and cardiac output) tend to increase absorption of inhaled carbon monoxide. Factors that may increase net carbon monoxide absorption include exercise, supine position, age (increased in infancy and childhood and declines in adults with age), altitude, increasing blood Hb concentration, and decreasing partial pressure of O<sub>2</sub> in inhaled air or blood (Castillo et al. 2006; Horvath et al. 1988; Joumard et al. 1991; McGrath et al. 1993; Peterson and Stewart 1975; Tikuisis et al. 1992).

### 3.4.2 Distribution

Inhaled carbon monoxide distributes to blood and extravascular tissues. In blood, carbon monoxide rapidly distributes into erythrocytes where it exists, primarily as a complex with Hb (COHb). The binding equilibrium constant for COHb is approximately 750  $\mu\text{M}^{-1}$  (Chakraborty et al. 2004; see further discussion in Section 3.4.3, Metabolism). At a blood Hb concentration of 150 g/L (approximately 2.3 mM), the molar concentration ratio of COHb would be approximately  $1.7 \times 10^6$  times greater than the molar concentration of dissolved carbon monoxide. The relative affinity of Hb for carbon monoxide and O<sub>2</sub> greatly favors binding of carbon monoxide to Hb over O<sub>2</sub>. The CO/O<sub>2</sub> affinity ratio (Haldane constant) is approximately 234 (Chakraborty et al. 2004). This means that a partial pressure of O<sub>2</sub> (pO<sub>2</sub>) approximately 234 times that of pCO is needed to achieve an equilibrium in which equal amounts of O<sub>2</sub> and carbon monoxide are bound to Hb (i.e., %COHb = % oxyhemoglobin [O<sub>2</sub>Hb]; Engel et al. 1969; Rodkey et al. 1969; Roughton 1970). In general, typical levels of COHb in nonsmokers are <2% (Adams et al. 1988; Allred et al. 1991; Anderson et al. 1973; Hinderliter et al. 1989; Kleinman et al. 1989, 1998; Sheps et al. 1987, 1990). Of this, COHb levels of approximately 0.2–1.0% are thought to be derived from endogenous production of carbon monoxide (Coburn et al. 1963; Delivoria-Papadopoulos et al. 1974; Longo 1977). Levels of COHb can (rarely) achieve 10% immediately following cigarette smoking (Kao and Nañagas 2006; Landaw 1973).

Although bound extensively to Hb in blood, distribution of dissolved carbon monoxide out of the vasculature can occur in response to diffusion gradients for carbon monoxide created by binding of carbon monoxide to Hb, myoglobin and other heme proteins in extravascular tissues (e.g., muscle, spleen), as well as removal of carbon monoxide from extravascular tissues by oxidative metabolism (Bruce and Bruce 2006; Bruce et al. 2008; Luomanmäki and Coburn 1969). Myoglobin, the dominant binding protein for carbon monoxide in cardiac and skeletal muscle, has a lower affinity for carbon

## 3. HEALTH EFFECTS

monoxide than Hb (COMb equilibrium constant  $\approx 20 \mu\text{M}^{-1}$ , and Haldane constant of approximately 23) and a single binding site for carbon monoxide (Gibson et al. 1986); however, its abundance in muscle can result in transfer of substantial amounts of inhaled carbon monoxide into muscle tissue. At a skeletal muscle myoglobin concentration of 4.7 g/kg muscle (approximately 0.26 mmole/kg; Moller and Sylven 1981), the molar concentration ratio of COMb would be approximately  $7.4 \times 10^3$  times greater than the molar concentration of dissolved carbon monoxide. Binding of carbon monoxide to Hb in blood and myoglobin in muscle decreases the concentrations of dissolved carbon monoxide in both tissues and limits the concentration gradient for diffusion of carbon monoxide into muscle (Bruce and Bruce 2006; Bruce et al. 2008). Exercise can increase the rate of transfer of carbon monoxide from blood to skeletal muscle (Richardson et al. 2002; Werner and Lindahl 1980). The effect of exercise is thought to be related, in part, to increased  $\text{O}_2$  consumption and lower  $\text{pO}_2$  in muscle, which favors binding of carbon monoxide to myoglobin, and increased muscle blood flow. These effects are more pronounced at low COHb concentrations ( $\leq 2\%$ ) than at higher concentrations (20% COHb; Richardson et al. 2002). Hypoxia appears to increase transfer of carbon monoxide from blood to muscle (Coburn and Mayers 1971; Richardson et al. 2002).

Binding of carbon monoxide to intracellular proteins, in particular, heme proteins, contributes to maintaining a favorable gradient for carbon monoxide diffusion from plasma into cells. Measurements of total carbon monoxide concentrations in tissues obtained from human autopsies showed the highest concentrations in blood, spleen, lung, kidney, and skeletal muscle, with detectable levels also in brain and adipose (Table 3-11, Vreman et al. 2006). Higher concentrations of carbon monoxide in blood, heart, skeletal muscle, and spleen observed in human autopsy studies reflect the abundance of the major carbon monoxide binding proteins in these tissues, Hb in erythrocytes and spleen (site of destruction and storage of erythrocytes), and myoglobin in cardiac and skeletal muscle. If the values presented in Table 3-11 are converted to tissue carbon monoxide mass (pmol), based on assumed tissue masses for a 70-kg adult (Davies and Morris 1993; ILSI 1994), carbon monoxide in blood and muscle accounted for approximately 60–70 and 20–30% of total body carbon monoxide burden, respectively. Relatively large contributions of blood and skeletal muscle to total body carbon monoxide content is consistent with the contributions that these tissues make to total body Hb and muscle myoglobin (Bruce and Bruce 2003). In a typical adult, blood contains approximately 12 mmoles of Hb with a total binding capacity of approximately 48 mmole carbon monoxide when completely saturated, and muscle contains approximately 8 mmoles of myoglobin with a total binding capacity of 8 mmole carbon monoxide when completely saturated.

## 3. HEALTH EFFECTS

**Table 3-11. Post-Mortem Tissue Distribution of Carbon Monoxide in Humans<sup>a</sup>**

Exposure	Adipose	Brain	Muscle	Heart	Kidney	Lung	Spleen	Blood
Concentration (pmol/100 g)								
Background	2	3	15	31	23	57	79	165
Fire	5	7	24	54	27	131	95	286
Fire + carbon monoxide	18	17	168	128	721	1097	2,290	3,623
Carbon monoxide asphyxiation	25	72	265	527	885	2694	3,455	5,196
Tissue/blood carbon monoxide concentration ratio								
Background	0.012	0.018	0.091	0.188	0.139	0.345	0.479	1.000
Fire	0.017	0.024	0.084	0.189	0.094	0.458	0.332	1.000
Fire + carbon monoxide	0.005	0.005	0.046	0.035	0.199	0.303	0.632	1.000
Carbon monoxide asphyxiation	0.005	0.014	0.051	0.101	0.170	0.518	0.665	1.000
Body weight (kg)	70	70	70	70	70	70	70	70
Tissue mass (percent)	200	2.0	40.0	0.5	0.4	1.4	0.3	7.4
Tissue carbon monoxide burden (nmol)								
Background	0.28	0.04	4.20	0.10	0.07	0.57	0.14	8.55
Fire	0.70	0.10	6.72	0.18	0.08	1.31	0.17	14.81
Fire + carbon monoxide	2.52	0.24	47.04	0.42	2.22	10.97	4.12	187.67
Carbon monoxide asphyxiation	3.50	1.01	74.20	1.73	2.73	26.95	6.22	269.15
Tissue carbon monoxide burden (percent of total)								
Background	2.0	0.3	30.1	0.7	0.5	4.1	1.0	61.3
Fire	2.9	0.4	27.9	0.7	0.3	5.4	0.7	61.5
Fire + carbon monoxide	1.0	0.1	18.4	0.2	0.9	4.3	1.6	73.5
Carbon monoxide asphyxiation	0.9	0.3	19.2	0.4	0.7	7.0	1.6	69.8

<sup>a</sup>Carbon monoxide concentrations based on Vreman et al. (2006). Tissue masses based on Davies and Morris (1993) and ILSI (1994).

### 3. HEALTH EFFECTS

The distribution of carbon monoxide residues observed in human autopsy studies is similar to that observed in rodents. In mice and rats that had not been intentionally exposed to air carbon monoxide (i.e., sacrificed during steady-state conditions), blood and muscle accounted for most the total body carbon monoxide burden (Vreman et al. 2005). Following 30-minute exposures to carbon monoxide (5,000 ppm), blood made a much larger contribution (approximately 90%) to carbon monoxide body burden, reflecting the longer period needed for carbon monoxide to distribute from blood to extravascular tissues (Vreman et al. 2005).

Carbon monoxide in the maternal system distributes to fetal tissues. Measurements of steady-state fetal COHb concentrations in fetal and maternal blood of nonsmoking women have found fetal COHb concentrations to be approximately 10–15% higher than maternal blood (fetal/maternal ratio=1.1–1.15; Longo 1977). Carbon monoxide binding to fetal Hb is analogously similar to maternal Hb. Both Hbs have the same binding capacity of 4 moles CO/mole Hb and both exhibit cooperative binding of carbon monoxide, whereby binding affinity increases as carbon monoxide molecules are successively added to Hb (see Section 3.4.3, Metabolism). Binding affinity of fetal Hb is approximately twice that of maternal Hb (Di Cera et al. 1989). The Haldane coefficient, reflecting the relative affinities of carbon monoxide and O<sub>2</sub> (CO/O<sub>2</sub>) for fetal Hb, appears to be approximately 20% lower than adult Hb (Di Cera et al. 1989; Engel et al. 1969; Hill et al. 1977). The higher binding affinity of fetal Hb as well as the relatively small diffusion gradients for carbon monoxide between maternal and fetal blood contribute to slower kinetics of COHb in fetal blood compared to maternal blood. This has been experimentally verified in studies conducted in pregnant sheep (Longo and Hill 1977). The half-time for approach to steady-state in the human fetus has been predicted (from modeling) to be approximately 7.5 hours in the fetus, compared to approximately 4 hours in maternal blood (Hill et al. 1977). As a result, with continuous exposure to a constant level of carbon monoxide in air, fetal blood COHb concentration would be expected to lag behind maternal levels, and reach a steady-state that is approximately 10–15% higher than maternal (Longo 1977). Similarly, following cessation of exposure, elimination kinetics of carbon monoxide from fetal blood would be expected to lag behind that of maternal blood (Hill et al. 1977).

#### **3.4.3 Metabolism**

Metabolism of carbon monoxide consists of three major processes: (1) metabolic production of carbon monoxide from endogenous and exogenous precursors; (2) binding of carbon monoxide to heme proteins (e.g., Hb, myoglobin, cytochromes); and (3) oxidative metabolism of carbon monoxide to CO<sub>2</sub>.

## 3. HEALTH EFFECTS

***Metabolic Production of Carbon Monoxide.*** Carbon monoxide is produced from the enzymatic degradation of endogenous and exogenous precursors. The major endogenous source of carbon monoxide production is the oxidative metabolism of heme by the enzyme heme oxygenase (HO). The net rate of endogenous production of carbon monoxide (i.e., total production minus metabolism) has been estimated in healthy adults to be approximately 0.42 mL CO/hour (16.4  $\mu$ mole/hour) or approximately 400  $\mu$ mole/day (Coburn et al. 1963). Of this, approximately 75–80% is thought to be derived from Hb metabolism, and the remaining portion from metabolism of other heme proteins such as myoglobin, cytochromes, peroxidases, and catalase (Berk et al. 1976). Endogenous carbon monoxide production contributes to a blood COHb concentration of approximately 0.4–0.7% (Benignus 1995; Coburn et al. 1965). Carbon monoxide is produced in all tissues that express HO; however, the major tissues contributing to carbon monoxide production are also major sites of heme metabolism, including liver, spleen, and reticuloendothelial system (Berk et al. 1974).

Numerous physiological and disease factors affect the rate of endogenous production of carbon monoxide. Carbon monoxide production increased by approximately 2-fold during the post-ovulatory (progesterone) phase of the menstrual cycle (Delivoria-Papadopoulos et al. 1974; Mercke and Lundh 1976). Carbon monoxide production also increases during pregnancy, reaching values that are 2–5 times that of the estrogen phase of the menstrual cycle at partum and returning to pre-pregnancy levels within 4 days following delivery (Longo 1977). Contributing factors to the increase in carbon monoxide production during pregnancy include the contribution of the growing fetus to carbon monoxide production, as well as the increase in maternal erythrocyte production and related changes in Hb metabolism (Longo 1977). The rate of production of carbon monoxide is accelerated in conditions that increase catabolism of Hb or other heme proteins, including hemolysis, hematomas, hemolytic anemias, thalassemia, and Gilbert's syndrome (Berk et al. 1974; Coburn et al. 1964, 1965; Hampson 2007; Meyer et al. 1998; Solanki et al. 1988). Carbon monoxide production was estimated to be 2–3 times higher in patients with hemolytic anemia compared healthy individuals (Coburn et al. 1965).

HO is the rate limiting step in heme degradation and its expression and activity is modulated by numerous factors. The HO-1 isozyme (one of three isozymes) and expression of the enzyme is induced by heme and heme derivatives, oxidative stress, hypoxia (including altitude-induced hypoxia), various metals, various cytokines, and exogenous carbon monoxide (Wu and Wang 2005). These factors would tend to promote endogenous carbon monoxide production.

## 3. HEALTH EFFECTS

In addition to endogenous carbon monoxide production from metabolism of heme, carbon monoxide is produced from oxidative metabolism of certain exogenous hydrocarbons, including dichloromethane and other dihalomethanes; and carbon tetrachloride (Agency for Toxic Substances and Disease Registry 2000, 2005; Stevens et al. 1980). The oxidative metabolism of dichloromethane (DCM) by isoform CYP2E1 of cytochrome P450 results in the production of carbon monoxide and increased blood COHb concentrations in blood (Andersen et al. 1991). Elevated blood COHb levels has been observed following DCM poisoning (Leikin et al. 1990). Other sources of exogenous carbon monoxide production include the heme oxidase catalysis of products of auto-oxidation of phenols, photo-oxidation of organic compounds, and lipid peroxidation of cell membrane lipids (Rodgers et al. 1994).

***Binding of Carbon Monoxide to Heme Proteins.*** Carbon monoxide binds to heme and competes for binding of O<sub>2</sub>. Binding of carbon monoxide to heme proteins (e.g., blood Hb, muscle myoglobin) has profound effects on the kinetics of distribution and elimination kinetics of both carbon monoxide and O<sub>2</sub>. Affinities of heme for carbon monoxide for Hb and myoglobin are sufficiently high to result in concentrations of dissolved carbon monoxide being 10<sup>3</sup>–10<sup>6</sup> times lower than the concentrations of COHb or COMb. This substantially lowers the concentration of dissolved carbon monoxide in blood and tissues and decreases the diffusion gradients for carbon monoxide movement of carbon monoxide from tissues and blood to the lungs, and substantially prolongs the elimination time for carbon monoxide than would otherwise occur in the absence of binding. Furthermore, the rates of dissociation of carbon monoxide from Hb and myoglobin are substantially slower than that of O<sub>2</sub> and sufficiently slow, relative to erythrocyte capillary transit times, to limit rates exchange of carbon monoxide between tissues and capillary blood. Binding of carbon monoxide to Hb also increases the affinity of Hb for O<sub>2</sub>, decreasing the availability of O<sub>2</sub> to tissues.

Although carbon monoxide binds to many different heme proteins (see Section 3.5.2), the affinities and abundance of Hb and myoglobin in the body results in most of the body stores of carbon monoxide being associated with these two heme proteins. Estimated kinetics and equilibrium parameters for carbon monoxide and O<sub>2</sub> binding to Hb and myoglobin are presented in Table 3-12. Each of the four heme moieties in Hb can bind O<sub>2</sub> or carbon monoxide. The overall affinity (i.e., equilibrium constant, K<sub>eq</sub>) is substantially higher for carbon monoxide (≈750 μM<sup>-1</sup>) compared to that of O<sub>2</sub> (≈3.2 μM<sup>-1</sup>). As a result, at similar partial pressures of the two gases (pCO=pO<sub>2</sub>), binding of carbon monoxide will occur at the expense of binding of O<sub>2</sub> and O<sub>2</sub> carrying capacity of blood will be compromised. The ratio of the K<sub>eq</sub> values (CO/O<sub>2</sub>) is referred to as the Haldane constant (≈234). This value gives the partial pressure ratio (pO<sub>2</sub>/pCO) needed to achieve equal amounts of bound COHb and O<sub>2</sub>Hb (i.e., %COHb = %O<sub>2</sub>Hb).

## 3. HEALTH EFFECTS

**Table 3-12. Hemoglobin and Myoglobin Binding Kinetics and Equilibrium Constants for Oxygen and Carbon Monoxide**

Ligand	$k_a$ ( $\mu\text{M}^{-1}\cdot\text{s}^{-1}$ )	$k_d$ ( $\text{s}^{-1}$ )	$K_{\text{Eq}}$ ( $\mu\text{M}^{-1}$ )	M
Hemoglobin <sup>a</sup>				
Oxygen	66	20	3.2	1
Carbon monoxide	6	0.008	750	234
Myoglobin <sup>b</sup>				
Oxygen	14	12	1.2	1
Carbon monoxide	0.51	0.019	26.8	23

<sup>a</sup>Based on Chakraborty et al. 2004

<sup>b</sup>Based on Gibson et al. 1986

$k_a$  = association rate constant;  $k_b$  = dissociation rate constant;  $K_{\text{Eq}}$  = equilibrium constant; M = Haldane constant ( $K_{\text{Eq CO}}/K_{\text{Eq O}_2}$ )

## 3. HEALTH EFFECTS


At an ambient air  $pO_2$  of approximately 160 Torr (21%  $O_2$ ), the equilibrium condition of equal %COHb = % $O_2$ Hb would be achieved at a carbon monoxide exposure concentration of approximately 900 ppm. The affinity of Hb for carbon monoxide results in an equilibrium condition in which nearly all carbon monoxide in blood is bound to Hb. For example, based on the value of  $K_{eq}$  of  $750 \mu M^{-1}$ , at a blood Hb concentration of 150 g/L ( $\approx 2.3$  mM), the equilibrium ratio of bound carbon monoxide to dissolved carbon monoxide in blood would be approximately  $1.7 \times 10^6$  (i.e.,  $> 10^6$  times as much carbon monoxide is bound to Hb than is circulating in blood as dissolved carbon monoxide). The carbon monoxide dissociation rate from Hb ( $k_d \approx 0.008 s^{-1}$ ;  $t_{1/2} \approx 87$  s) is slower than from  $O_2$  ( $k_e \approx 20 s^{-1}$ ,  $t_{1/2} \approx 0.035$  s). As a result exchange of  $O_2$  bound to Hb in blood with alveolar air and tissues is much more efficient than carbon monoxide exchange (i.e., fast vs. slow dissociation relative to erythrocyte transit times in capillaries). These two characteristics (high affinity and slow dissociation) contribute to relatively long retention times of carbon monoxide in blood following exposures to carbon monoxide.

Binding of  $O_2$  to the four heme moieties of Hb is cooperative (Alcantara et al. 2007; Roughton 1970). The associative reaction rate ( $k_a$ ) becomes faster with successive additions of  $O_2$ , increasing the affinity ( $K_{eq}$ ) of Hb for  $O_2$  (values for  $k_a$  shown in Table 3-12 are for the high affinity, R, state of fully loaded Hb). Cooperative binding of  $O_2$  to Hb serves an important physiological function. Under conditions of relatively high  $pO_2$  in arterial blood entering the lung ( $pO_2 = 100$  mmHg), cooperative binding promotes the loading of Hb with  $O_2$ ; with successive binding of  $O_2$  molecules, the associative rate and affinity increases, promoting binding of a subsequent molecule of  $O_2$ . Conversely, at lower  $pO_2$  in venous blood ( $pO_2 \approx 20$  mmHg), cooperative binding promotes unloading of  $O_2$  for utilization; dissociation of a molecule of  $O_2$  decreases the affinity of Hb for  $O_2$ , promoting the dissociation of a subsequent molecule of  $O_2$ . Cooperative binding contributes to the sigmoid shape of the  $O_2$  dissociation curve for Hb (Figure 3-5).

Binding of carbon monoxide to Hb is also cooperative. Binding of carbon monoxide increases the associative rate of binding and affinity of both carbon monoxide and  $O_2$ . As a result, the presence of carbon monoxide has several effects on the dissociation curve for  $O_2$  (Figure 3-5): (1) decreases the partial pressure of  $O_2$  in blood and tissues (i.e., by an amount equal to pCO), shifting the  $pO_2$  axis downscale at each point in the arterial-venous circuit; (2) displaces  $O_2$  from Hb, decreasing the maximum %COHb achieved in arterial blood; and (3) shifts the  $O_2$  dissociation curve downscale so that less  $O_2$  dissociates  $O_2$  from Hb (more will remain bound, % $O_2$ Hb will be higher) at the lower  $pO_2$  values in venous blood. The combined result of these changes is that less  $O_2$  in blood is available for utilization in tissues.

3. HEALTH EFFECTS

**Figure 3-5. Oxyhemoglobin Dissociation Curve for Normal Human Blood Containing 50% Carboxyhemoglobin, and for Blood of Anemic Human**



Source: EPA 2000

## 3. HEALTH EFFECTS

The binding affinities of carbon monoxide and O<sub>2</sub> to myoglobin are lower than that for Hb; however, the higher affinity for carbon monoxide (Haldane constant  $\approx 23$ ), results in preferential binding of carbon monoxide over O<sub>2</sub> and displacement of O<sub>2</sub> of myoglobin at similar partial pressures of both gasses (Table 3-12). In addition to displacing O<sub>2</sub> from myoglobin, binding of carbon monoxide to myoglobin has a pronounced effect on the distribution of carbon monoxide in the body by limiting the dissolved carbon monoxide concentration in muscle and, thereby, the rates of diffusion of carbon monoxide from muscle to blood. The affinity of myoglobin ( $K_{eq} \approx 27$ ) results in an equilibrium condition in which dissolved carbon monoxide concentrations in muscle are very low relative to COMb concentrations. For example, based on a value for  $K_{eq}$  of  $27 \mu\text{M}^{-1}$ , at a muscle myoglobin content of approximately 4.7 g/kg ( $\approx 0.28 \text{ mM}$ ), the equilibrium ratio of bound carbon monoxide to dissolved carbon monoxide in muscle would be approximately  $7.4 \times 10^3$  (approximately  $10^4$  times as much carbon monoxide is bound to myoglobin than is dissolved muscle tissue). Furthermore, the carbon monoxide dissociation rate from myoglobin ( $k_d \approx 0.019 \text{ s}^{-1}$ ;  $t_{1/2} \approx 36 \text{ s}$ ) is slower than O<sub>2</sub> ( $k_e \approx 12 \text{ s}^{-1}$ ,  $t_{1/2} \approx 0.06 \text{ s}$ ). As a result exchange of carbon monoxide bound to myoglobin in muscle with blood is much less efficient than O<sub>2</sub> exchange (i.e., slow vs. fast dissociation relative to erythrocyte transit times in capillaries). These two characteristics (high affinity and slow dissociation) contribute to relatively long retention times of carbon monoxide in muscle following exposures to carbon monoxide.

***Oxidative Metabolism of Carbon Monoxide.*** Carbon monoxide can be oxidized to CO<sub>2</sub>. The rate of oxidation has been estimated in human subjects who inhaled <sup>14</sup>CO to be approximately 0.015 mL CO/hour, which corresponds to approximately 0.6  $\mu\text{mol}$ /hour; 14  $\mu\text{mol}$ /day, and approximately 3% of endogenous carbon monoxide production (Luomanmaki and Coburn 1969). Similar measurements made in dogs yielded estimates for carbon monoxide metabolism that were approximately 8% of endogenous production (Luomanmaki and Coburn 1969). The major pathway for metabolism of carbon monoxide to CO<sub>2</sub> is oxidation by mitochondrial cytochrome oxidase (Fenn 1970; Young and Caughey 1986).

#### 3.4.4 Elimination and Excretion

Absorbed carbon monoxide is eliminated from the body by exhalation and oxidative metabolism. Oxidative metabolism of carbon monoxide has been estimated to be a relatively small fraction (<10%) of endogenous carbon monoxide production (Luomanmaki and Coburn 1969). Under most conditions, the dominant route of elimination of absorbed carbon monoxide is exhalation. Rates of elimination of carbon monoxide in humans who have been experienced inhalation exposures to carbon monoxide have been

## 3. HEALTH EFFECTS

estimated from kinetic analyses of the time course of blood %COHb (Bruce and Bruce 2006; Joumard et al. 1991; Landaw 1973; Levasseur et al. 1996; Peterson and Stewart 1970; Shimazu et al. 2000; Weaver et al. 2000). The decline in blood %COHb following cessation of an inhalation exposure to carbon monoxide exhibits at least two kinetic phases (Bruce and Bruce 2006; Shimazu et al. 2000). The fast phase is thought to reflect a combination of exhalation of carbon monoxide along with slower distribution of blood carbon monoxide to tissues that continues after cessation of exposure (Bruce and Bruce 2006). The elimination half-times for both phases appears to increase with exposure intensity and/or duration; a contributing factor to this effect may be the amount of carbon monoxide accumulated in extravascular stores. In human subjects exposed to 2% carbon monoxide for 1–3 minutes (peak COHb%=30–40), values for elimination half-lives were  $5.7 \pm 1.5$  and  $103 \pm 20.5$  minutes for the fast and slow phases, respectively (Shimazu et al. 2000). The fast phase rate increased when subjects were exposed to 500 ppm for 5–10 hours (peak %COHb=34–37):  $21.5 \pm 2.1$  and  $118 \pm 11.2$  minutes, respectively (Shimazu et al. 2000). Rates have been reported for the slow phase of elimination. In subjects whose peak %COHb ranged from 2 to 15, the half-life was 320 minutes (range: 128–409 minutes, Peterson and Stewart 1970). Bruce and Bruce (2006) measured the time to achieve one half of peak %COHb in subjects whose peak %COHb values ranged from 15 to 20 and estimated a half-life of 248 minutes (range: 206–328). Clinical studies conducted on healthy individuals exposed to ambient air carbon monoxide concentrations have found that elimination half-time increases with age, with the most pronounced increase occurring from age 2 to 20 years and is marginally longer in males compared to females (Joumard et al. 1991). The gender effect may be related to relative differences in body size and/or muscle mass. Exercise decreases carbon monoxide elimination half-time (Joumard et al. 1991).

The rate of elimination of absorbed carbon monoxide is accelerated by inhaling high concentrations of O<sub>2</sub> (Landaw 1973; Peterson and Stewart 1970; Weaver et al. 2000). Elimination half-lives of approximately 20 minutes have been observed in subjects who inhaled 100% O<sub>2</sub> following cessation of exposure to carbon monoxide. An analysis of elimination rates observed in 93 acute carbon monoxide poisoning patients who were treated with 100 O<sub>2</sub> did not find significant associations between elimination half-lives and age, gender (<40 years, ≥40 years), smoke inhalation, history of loss of consciousness, concurrent tobacco smoking, degree of initial metabolic acidosis (base excess), or initial %COHb level (Weaver et al. 2000).

## 3. HEALTH EFFECTS

**3.4.5 Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models**

Physiologically based pharmacokinetic (PBPK) models use mathematical descriptions of the uptake and disposition of chemical substances to quantitatively describe the relationships among critical biological processes (Krishnan et al. 1994). PBPK models are also called biologically based tissue dosimetry models. PBPK models are increasingly used in risk assessments, primarily to predict the concentration of potentially toxic moieties of a chemical that will be delivered to any given target tissue following various combinations of route, dose level, and test species (Clewell and Andersen 1985). Physiologically based pharmacodynamic (PBPD) models use mathematical descriptions of the dose-response function to quantitatively describe the relationship between target tissue dose and toxic end points.

PBPK/PD models refine our understanding of complex quantitative dose behaviors by helping to delineate and characterize the relationships between: (1) the external/exposure concentration and target tissue dose of the toxic moiety, and (2) the target tissue dose and observed responses (Andersen and Krishnan 1994; Andersen et al. 1987). These models are biologically and mechanistically based and can be used to extrapolate the pharmacokinetic behavior of chemical substances from high to low dose, from route to route, between species, and between subpopulations within a species. The biological basis of PBPK models results in more meaningful extrapolations than those generated with the more conventional use of uncertainty factors.

The PBPK model for a chemical substance is developed in four interconnected steps: (1) model representation, (2) model parameterization, (3) model simulation, and (4) model validation (Krishnan and Andersen 1994). In the early 1990s, validated PBPK models were developed for a number of toxicologically important chemical substances, both volatile and nonvolatile (Krishnan and Andersen 1994; Leung 1993). PBPK models for a particular substance require estimates of the chemical substance-specific physicochemical parameters, and species-specific physiological and biological parameters. The numerical estimates of these model parameters are incorporated within a set of differential and algebraic equations that describe the pharmacokinetic processes. Solving these differential and algebraic equations provides the predictions of tissue dose. Computers then provide process simulations based on these solutions.

The structure and mathematical expressions used in PBPK models significantly simplify the true complexities of biological systems. If the uptake and disposition of the chemical substance(s) are adequately described, however, this simplification is desirable because data are often unavailable for

### 3. HEALTH EFFECTS

many biological processes. A simplified scheme reduces the magnitude of cumulative uncertainty. The adequacy of the model is, therefore, of great importance, and model validation is essential to the use of PBPK models in risk assessment.

PBPK models improve the pharmacokinetic extrapolations used in risk assessments that identify the maximal (i.e., the safe) levels for human exposure to chemical substances (Andersen and Krishnan 1994). PBPK models provide a scientifically sound means to predict the target tissue dose of chemicals in humans who are exposed to environmental levels (for example, levels that might occur at hazardous waste sites) based on the results of studies where doses were higher or were administered in different species. Figure 3-6 shows a conceptualized representation of a PBPK model.

If PBPK models for carbon monoxide exist, the overall results and individual models are discussed in this section in terms of their use in risk assessment, tissue dosimetry, and dose, route, and species extrapolations.


The first physiologically-based mechanistic model of carbon monoxide kinetics was reported in Coburn et al. (1965). This model, which subsequently became known as the Coburn-Forster-Kane (CFK) model, was developed to predict relationships between inhalation exposures to carbon monoxide and blood COHb levels in humans. Numerous modifications and extensions of the original CFK model have been reported that more rigorously address nonlinearities resulting from interdependence of COHb and O<sub>2</sub>Hb levels in blood (Tikuisis et al. 1987b), simulation of arterial and venous blood compartments (Smith et al. 1994), simulation of maternal-fetal transfer of carbon monoxide (Hill et al. 1977), extrapolation to nonhuman species (Benignus and Annau 1994; Hill et al. 1977; Longo and Hill 1977), linkage of the CFK model to more complex respiratory gas exchange models (Abram et al. 2007; Benignus 1995), and use in simulating the elimination kinetics of carbon monoxide produced from exogenous precursors (Andersen et al. 1991). A multi-compartment model was also developed that simulates carbon monoxide absorption and distribution kinetics in muscle (Bruce and Bruce 2003; Bruce et al. 2008).

#### **Coburn-Forster-Kane Model**

The CFK model, as originally described in Coburn et al. (1965), is a single-compartment model of instantaneous equilibrium between carbon monoxide, O<sub>2</sub>, and blood Hb, and assumes that blood O<sub>2</sub>Hb concentration is constant (i.e., independent of COHb concentration). This linear form of the CFK model

## 3. HEALTH EFFECTS

**Figure 3-6. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance**



Note: This is a conceptual representation of a physiologically based pharmacokinetic (PBPK) model for a hypothetical chemical substance. The chemical substance is shown to be absorbed via the skin, by inhalation, or by ingestion, metabolized in the liver, and excreted in the urine or by exhalation.

Source: adapted from Krishnan and Andersen 1994

## 3. HEALTH EFFECTS

is expressed in the following differential equation which is integrated over time to solve for COHb concentration (Equation 1):

$$V_b \cdot \frac{d[COHb]_t}{dt} = V_{CO} - \frac{[COHb]_{t-1} \cdot P_c O_2}{[O_2Hb] \cdot M} \cdot \left( \frac{1}{\frac{1}{D_L CO} + \frac{1}{V_A}} \right) + \left( \frac{P_I CO}{\frac{1}{D_L CO} + \frac{1}{V_A}} \right) \quad \text{Eq. (1)}$$

where  $t$  represents time at the end of the integration time step and  $t-1$  represents the time at the beginning of the time step;  $V_b$  is blood volume in mL;  $[COHb]$  is the COHb concentration in mL carbon monoxide per mL blood (STPD);  $V_{CO}$  is the endogenous carbon monoxide production rate in mL/minute (STPD);  $[O_2Hb]$  is the oxyhemoglobin concentration in mL  $O_2$  per mL blood (STPD);  $P_c O_2$  is the average partial pressure of  $O_2$  in lung capillaries in Torr;  $M$  is the Haldane constant;  $V_A$  is the alveolar ventilation in mL/minute (STPD);  $D_L CO$  is the lung diffusing capacity of carbon monoxide in mL/minute-Torr (STPD); and  $P_I CO$  is the carbon monoxide partial pressure in inhaled air in Torr.

The linear form of the CFK model is acceptable for simulating low COHb concentrations (<6%; Smith 1990). However, at higher COHb levels, appreciable errors are introduced into the simulation by ignoring the interdependence of COHb and  $O_2Hb$  levels that result from limited binding capacity of Hb. This interdependence can be introduced into the CFK model by replacing the constant term for  $[O_2Hb]$  with the following expression relating COHb and  $O_2Hb$  (Tikuisis et al. 1987b; Equation 2):

$$[O_2Hb] = 1.38 \cdot Hb - [COHb] \quad \text{Eq. (2)}$$

where  $1.38 \cdot Hb$  represents the maximum binding capacity of Hb for  $O_2$  or carbon monoxide in mL  $O_2$ /mL blood (STPD). The resulting nonlinear form of the CFK model is expressed in Equations 2, 3, and 4:

$$\frac{d[COHb]_t}{dt} = \frac{V_{CO}}{V_b} + \frac{1}{V_b \cdot \beta} \cdot \left( P_I CO - \frac{[COHb]_{t-1} \cdot P_c O_2}{[O_2Hb]_{t-1} \cdot M} \right) \quad \text{Eq. (3)}$$

$$\beta = \frac{1}{D_L CO} + \frac{P_B - 47}{V_A} \quad \text{Eq. (4)}$$

## 3. HEALTH EFFECTS

where  $P_B$  is the barometric pressure (Torr) and the value 47 represents the partial pressure of water in water saturated air (Torr). Typical parameters values used in the CFK model are presented in Table 3-13. Parameter values for application of the nonlinear CFK model for predicting COHb levels in rats were reported by Benignus and Annau (1994) and are also shown in Table 3-13.

**Validation of the model.** Numerous studies have examined the accuracy of exposure-blood COHb relationships predicted from the CFK model (Benignus et al. 1994; Coburn et al. 1965; Hauck and Neuberger 1984; Joumard et al. 1991; Peterson and Stewart 1970, 1975; Stewart et al. 1970, 1973; Tikuisis et al. 1987a, 1987b). In general, the model performs well for predicting near equilibrium conditions; however, it performs less well in predicting the early uptake phase of carbon monoxide kinetics, and COHb levels associated with rapidly changing exposures (Benignus et al. 1994; Tikuisis et al. 1987a, 1987b). The model also underestimates the arterial blood COHb level and overestimates the venous blood COHb level, an expected outcome of the simplistic assumption of a single well-mixed blood compartment. Nevertheless, the model predicts quasi-steady-state blood COHb levels that agree closely with observations (Benignus et al. 1994; Bruce and Bruce 2003; Tikuisis et al. 1987a, 1987b).

McCartney (1990) reported a sensitivity analysis of the CFK model and found that parameter value sensitivities varied considerably in magnitude at different points in the exposure simulation (Figure 3-7). At steady-state, the model is most sensitive to the values assigned to the parameters representing the input value, the inhaled carbon monoxide concentration ( $P_iCO$ ), as well as total Hb level ( $Hb$ ), and the Haldane coefficient ( $M$ ). Increases in values for any of these parameters results in higher predicted COHb levels. However, during the accumulation phase of the simulation, the predicted COHb level is also highly sensitive to alveolar ventilation rate ( $V_A$ ) and the lung carbon monoxide diffusing capacity ( $D_LCO$ ), with increases in these parameters resulting in higher COHb levels, and to the value assigned to the blood volume ( $V_b$ ), with increases in the parameter value resulting in lower predicted COHb levels.

**Risk assessment.** The CFK model has been used in establishing the National Ambient Air Quality Standard for carbon monoxide (EPA 2000) and has been used in analyses that have supported development of Threshold Limit Values for occupational safety (Tikuisis et al. 1987b).

**Target tissues.** The CFK model predicts COHb levels in blood.

**Species extrapolation.** Although originally developed to simulate blood COHb levels in humans, the model can be used to predict COHb levels in other mammalian species if physiological and chemical-

## 3. HEALTH EFFECTS

**Table 3-13. Parameter Values for Coburn-Forster-Kane (CFK) Model**

Parameter	Definition	Unit	Value	
			Human <sup>a</sup>	Rat <sup>b</sup>
BW	Body weight	kg	70	0.250
COHb <sub>0</sub>	Blood COHb level at start of simulation	mL CO/mL blood (STPD)	0.00155	0.00155
D <sub>L</sub> CO	Lung CO diffusing capacity	mL CO/minute-Torr (STPD)	30	0.027·0.563·BW
Hb	Blood Hb concentration	g/mL	0.144	0.158
M	Haldane coefficient	unitless	218	207
O <sub>2</sub> HbMAX	Maximum blood O <sub>2</sub> Hb concentration	mL O <sub>2</sub> /mL blood (STPD)	1.38	1.38
P <sub>C</sub> O <sub>2</sub>	Average partial pressure of O <sub>2</sub> in alveolar air	Torr (STPD)	49	49
P <sub>I</sub> CO	Partial pressure of inhaled CO	Torr (STPD)	Input	Input
V <sub>A</sub>	Alveolar ventilation rate	L/hour <sup>c</sup>	(15·BW <sup>0.74</sup> )	(15·BW <sup>0.74</sup> )
V <sub>B</sub>	Blood volume	L <sup>d</sup>	0.074·BW	0.074·BW
V <sub>CO</sub>	Endogenous CO production rate	mL/minute (STPD)	0.007	0.000333·BW

<sup>a</sup>Based on Tikuisis et al. (1992)

<sup>b</sup>Based on Benignus and Annau (1994)


<sup>c</sup>Must be converted to mL/minute by multiplying 1,000/60.

<sup>d</sup>Must be converted to mL by multiplying by 1,000.

BW = body weight; CO = carbon monoxide; COHb = carboxyhemoglobin; Hb = hemoglobin; STPD = standard temperature and pressure, dry (i.e., adjusted for partial pressure of water vapor in saturated air [47 Torr STP])

## 3. HEALTH EFFECTS

**Figure 3-7. Plot of Standardized Sensitivity Coefficients for Carboxyhemoglobin (COHb) Levels Predicted from the Coburn-Forster-Kane (CFK) Model for a 100 ppm Exposure to an Individual Engaged in Vigorous Exercise (Alveolar Ventilation Rate=30 L/Minute)\***



\*Each curve represents the time trend of sensitivity coefficients for a given parameter during the simulation. Time trends for six model parameters are shown:  $D_LCO$ , lung carbon monoxide diffusion capacity;  $F_I CO$ , fractional air carbon monoxide concentration (ppm);  $M$ , Haldane coefficient;  $TH_b$ , blood hemoglobin concentration;  $V_A$ , alveolar ventilation rate; and  $V_b$ , blood volume. Sensitivity coefficients and central coefficients are for 1% upward and downward perturbation of each parameter. Coefficients  $>0$  indicate an increase in predicted COHb in association with an increase in the parameter value; coefficients that are  $<0$  indicate that COHb decreased in association with an increase in the parameter value.

Source: EPA 2009g based on McCartney (1990)

## 3. HEALTH EFFECTS

specific parameter values are available for the species of interest. The latter include the Haldane coefficient ( $M$ ), the binding capacity of Hb, and the lung carbon monoxide diffusing capacity ( $D_LCO$ ). Parameter values for the rat have been reported by Benignus and Annau (1994; Table 3-13).

**Interroute extrapolation.** The CFK model simulates inhalation exposures.

**Smith et al. (1994) Multi-compartment Model**

The nonlinear CFK model, described above, represents blood as a well-mixed compartment, an acceptable assumption for simulating near equilibrium conditions. However, during uptake and elimination phases of carbon monoxide kinetics, appreciable differences have been observed between arterial and venous blood (Benignus et al. 1994). Improved prediction of arterial COHb levels was achieved with an expansion of the CFK model to include slowly and rapidly perfused compartments, and an arm compartment (representing vascular sampling sites) that includes compartments representing the major arterial and venous vasculature (Smith et al. 1994, Figure 3-8; Table 3-14).

**Validation of the model.** The Smith et al. (1994) models was evaluated with data collected from 15 healthy adults subjects who inhaled carbon monoxide under controlled conditions (Benignus et al. 1994). Each subject (15 adult males) inhaled for approximately 5 minutes from a bag that had an initial carbon monoxide concentration of approximately 7,000 ppm. The exposures achieved end exposure blood COHb concentrations ranging from 10 to 15%. The Smith et al. (1994) model provided improved predictions of COHb levels in samples taken from radial artery and antecubital vein.

**Risk assessment.** Risk assessment applications of the Smith et al. (1994) were not identified in the available literature.


**Target tissues.** The Smith et al. (1994) model predicts COHb levels in arterial and venous blood.

**Species extrapolation.** Although originally developed to simulate blood COHb levels in humans, the model could be used to predict COHb levels in other mammalian species provided that physiological and chemical-specific parameter values are available for the species of interest

**Interroute extrapolation.** The Smith et al. (1994) model simulates inhalation exposures.

## 3. HEALTH EFFECTS

Figure 3-8. Structure of the Smith et al. (1994) Model\*



\*Diffusive exchange carbon monoxide (CO) and oxygen (O<sub>2</sub>) occurs between lung alveolar space alveolar capillary blood. Instantaneous equilibrium is assumed between gases and blood hemoglobin, governed by the Haldane coefficient. Blood carboxyhemoglobin (COHb) is apportioned to various blood subcompartments according to subcompartment blood flows and volumes.

## 3. HEALTH EFFECTS

**Table 3-14. Parameter Values for the Smith et al. (1994) Model**

Parameter	Definition	Unit	Value
<b>Chemical parameters<sup>a</sup></b>			
COHb <sub>0</sub>	Blood COHb level at start of simulation	mL CO/mL blood (STPD)	0.00155
D <sub>L</sub> CO	Lung CO diffusing capacity	mL CO/minute-Torr (STPD)	30
Hb	Blood Hb concentration	g/mL	0.144
M	Haldane coefficient	unitless	218
O <sub>2</sub> HbMAX	Maximum blood O <sub>2</sub> Hb concentration	mL O <sub>2</sub> /mL blood (STPD)	1.38
P <sub>C</sub> O <sub>2</sub>	Average partial pressure of O <sub>2</sub> in alveolar air	Torr (STPD)	49
P <sub>I</sub> CO	Partial pressure of inhaled CO	Torr (STPD)	Input
V <sub>CO</sub>	Endogenous CO production rate	mL/minute (STPD)	0.007
<b>Compartment volumes<sup>b</sup></b>			
BW	Body weight	kg	70
V <sub>B</sub>	Blood volume	L	0.074·BW
V <sub>1</sub>	Volumes of pulmonary capillaries and veins and ½ heart	L	0.1·V <sub>B</sub>
V <sub>2</sub>	Volume of rapidly perfused compartment	L	0.11·V <sub>B</sub>
V <sub>3</sub>	volume of slowly perfused compartment	L	0.652·V <sub>B</sub>
V <sub>4</sub>	Volume of major arteries in arms	L	0.0069·V <sub>B</sub>
V <sub>5</sub> , V <sub>6</sub>	Volume of small vessels in arms	L	0.0076·V <sub>B</sub>
V <sub>7</sub>	Volume of intermediate veins in arms	L	0.0025·V <sub>B</sub>
V <sub>8</sub>	Volume of large veins in arms	L	0.0361·V <sub>B</sub>
V <sub>9</sub>	Volume of ½ heart and pulmonary arteries	L	0.085·V <sub>B</sub>
<b>Compartment flows<sup>c</sup></b>			
V <sub>A</sub>	Alveolar ventilation rate	L/hour	(15·BW <sup>0.74</sup> )
QC	Cardiac output	L/hour	(15·BW <sup>0.74</sup> )
Q <sub>12</sub>	Flow to rapidly perfused tissue	L/hour	0.6·QC
Q <sub>13</sub>	Flow to slowly perfused tissue	L/hour	0.35·QC
Q <sub>14</sub>	Flow to arteries of arms	L/hour	0.05·QC
Q <sub>45</sub>	Flow to small vessels in arms	L/hour	0.005·QC

<sup>a</sup>Based on Tikuisis et al. (1992)

<sup>b</sup>Must be converted to mL/minute by multiplying 1,000/60

<sup>c</sup>Must be converted to mL by multiplying by 1,000

BW = body weight; CO = carbon monoxide; COHb = carboxyhemoglobin; Hb = hemoglobin; STPD = standard temperature and pressure, dry (i.e., adjusted for partial pressure of water vapor in saturated air (47 Torr STP))

## 3. HEALTH EFFECTS

**Hill et al. (1977) Maternal-Fetal Model**

Hill et al. (1977) developed a multi-compartment model for simulating maternal-fetal transfer of carbon monoxide. This model includes simulations of maternal, placental, and fetal compartments, as well as maternal and fetal contributions to endogenous production of carbon monoxide (Figure 3-9; Table 3-15). The model assumes instantaneous equilibrium of O<sub>2</sub> and carbon monoxide with Hb in maternal, placental, and fetal tissues, with delivery to placental and fetal tissues governed by blood flow rates.


**Validation of the model.** The Hill et al. (1977) model was evaluated with data collected in experiments conducted in pregnant sheep (Longo and Hill 1977; Longo 1977). Pregnant sheep were exposed to 30, 50, or 100 ppm carbon monoxide for 24 hours, or to 300 ppm for 4 hours and maternal and fetal arterial blood was sampled from indwelling catheters for measurement of COHb levels. These exposures achieved maternal blood COHb levels ranging from approximately 4 to 24% (300 ppm). Steady-state blood COHb levels in the fetus, achieved during the exposures to 30–100 ppm, were approximately 70% higher than maternal levels. The model predicted maternal and fetal arterial blood COHb levels that agreed with observations (e.g., were within  $\pm 1$  SD of observations). The model predicts slower accumulation and elimination of COHb in the human fetus compared to the maternal system, with steady-state values in fetal blood that are approximately 10–12% higher than maternal blood. The latter prediction agrees with observations made in cord blood (Longo 1977).

**Risk assessment.** Risk assessment applications of the Hill et al. (1977) model were not identified in the available literature. However, Hill et al. (1977) present and discuss predictions made with the model regarding effects of various factors on maternal and fetal blood COHb levels, including air carbon monoxide concentration, air O<sub>2</sub> concentration, altitude, exercise, fetal O<sub>2</sub> utilization, and fetal blood flow. Of particular relevance to risk assessment is the prediction of slower accumulation and elimination of COHb in the human fetus compared to the maternal system. Based on model predictions, the elimination half-times were approximately 4 hours for maternal and 7.5 hours for fetal. Furthermore, the model predicted a more pronounced effect of breathing 100% O<sub>2</sub> on carbon monoxide elimination from the maternal system (5-fold decrease in half-time) compared to the effect on fetal half-time (2-fold decrease).

**Target tissues.** The Hill et al. (1977) model predicts COHb levels in maternal and fetal blood.

**Species extrapolation.** The Hill et al. (1977) model has been applied to predicting maternal and fetal blood COHb levels in humans and sheep (Hill et al. 1977; Longo 1977; Longo and Hill 1977).

## 3. HEALTH EFFECTS

**Figure 3-9. Structure of the Hill et al. (1977) Maternal-Fetal Model\***

\*Average blood carboxyhemoglobin (COHb) levels in maternal and fetal tissues are calculated in the Body Gas Equilibration Model, based on inputs of  $PO_2/O_2Hb$  from lung, maternal tissue, placental tissue, and fetal placental capillary models.

$P_{I,CO}$  = inhaled carbon monoxide partial pressure;  $V_{CO,M}$  = maternal endogenous carbon monoxide production;  $V_{CO,F}$  = fetal endogenous carbon monoxide production;  $VO_2$  = oxygen consumption

## 3. HEALTH EFFECTS

**Table 3-15. Parameter Values for the Hill et al. (1977) Model**

Parameter	Definition	Unit	Value
Chemical parameters			
$D_M\text{CO}$	Lung membrane diffusing capacity	mL CO/minute-Torr	50, 60
$D_L\text{CO}$	Lung CO diffusing capacity	mL CO/minute-Torr	30, 28
$D_P\text{CO}$	Placental CO diffusing capacity	mL CO/minute-Torr	1.5, 2.7
$\text{Hb}_M$	Maternal blood Hb concentration	g/mL	0.125
$\text{Hb}_F$	Fetal blood Hb concentration	g/mL	0.155
$\text{cap}_M$	Maternal blood $\text{O}_2$ capacity	mL $\text{O}_2$ /mL	0.168
$\text{cap}_F$	Fetal blood $\text{O}_2$ capacity	mL $\text{O}_2$ /mL	0.208
$M_M$	Haldane coefficient for maternal Hb	unitless	223
$M_F$	Haldane coefficient for fetal Hb	unitless	181
$P_B$	Atmospheric pressure	Torr	760
$P_{\text{H}_2\text{O}}$	Partial pressure of water vapor (STP)	Torr	47
$(\text{PO}_2/\text{O}_2\text{Hb})_L$	Average ratio of partial pressure of $\text{O}_2$ to $\text{O}_2\text{Hb}$ concentration in lung capillaries	Torr/%	1.029 (at $P_i\text{CO}=0$ )
$(\text{PO}_2/\text{O}_2\text{Hb})_M$	Average ratio of partial pressure of $\text{O}_2$ to $\text{O}_2\text{Hb}$ concentration in maternal placental capillaries	Torr/%	0.5609 (at $P_i\text{CO}=0$ )
$(\text{PO}_2/\text{O}_2\text{Hb})_F$	Average ratio of partial pressure of $\text{O}_2$ to $\text{O}_2\text{Hb}$ concentration in fetal placental capillaries	Torr/%	0.4128 (at $P_i\text{CO}=0$ )
$P_{A\text{O}_2}$	Partial pressure of $\text{O}_2$ in alveolar air	Torr	108
$P_{A\text{CO}_2}$	Partial pressure of $\text{CO}_2$ in alveolar air	Torr	32
$P_{V\text{O}_2}$	Partial pressure of $\text{O}_2$ in mixed venous blood in pulmonary artery	Torr	38
$P_{V\text{CO}_2}$	Partial pressure of $\text{CO}_2$ in mixed venous blood in pulmonary artery	Torr	38
$P_{F\text{O}_2}$	Partial pressure of $\text{O}_2$ in umbilical artery	Torr	20
$P_{F\text{CO}_2}$	Partial pressure of $\text{CO}_2$ in umbilical artery	Torr	48
$P_{M\text{O}_2}$	Partial pressure of $\text{O}_2$ in uterine artery	Torr	95
$P_{M\text{CO}_2}$	Partial pressure of $\text{CO}_2$ in uterine artery	Torr	32
$P_{50,M}$	Partial pressure of $\text{O}_2$ at half $\text{O}_2$ saturation in maternal blood	Torr	26.4
$P_{50,F}$	Partial pressure of $\text{O}_2$ at half $\text{O}_2$ saturation in fetal blood	Torr	20.0
$\text{pH}_M$	pH of uterine artery blood	M	7.40
$\text{pH}_F$	pH of umbilical artery blood	M	7.34
$\text{pH}_M$	pH of mixed venous blood	M	7.37
$P_i\text{CO}$	Partial pressure of inhaled CO	Torr	Input
$V_{\text{CO},M}$	Maternal endogenous CO production rate	mL/minute	0.0153
$V_{\text{CO},F}$	Fetal endogenous CO production rate	mL/minute	0.0006
$\text{VO}_{2M}$	Total maternal $\text{O}_2$ consumption rate	mL/minute	271
$\text{VO}_{2F}$	Fetal $\text{O}_2$ consumption rate	mL/minute	22

## 3. HEALTH EFFECTS

**Table 3-15. Parameter Values for the Hill et al. (1977) Model**

Parameter	Definition	Unit	Value
Compartment volumes			
$vol_M$	Maternal blood volume	mL	5,000
$vol_F$	Fetal blood volume	mL	400
Compartment flows			
$V_A$	Alveolar ventilation rate	mL/minute	6,000
$Q_M$	Maternal placental blood flow	mL/minute	350
$Q_F$	Fetal placental blood flow	mL/minute	350
$Q_L$	Lung blood flow	mL/minute	5,600

CO = carbon monoxide; Hb = hemoglobin; STPD = standard temperature and pressure, dry (i.e., adjusted for partial pressure of water vapor in saturated air [47 Torr STP]).

## 3. HEALTH EFFECTS

**Interroute extrapolation.** The Hill et al. (1977) model simulates inhalation exposures.


**Bruce et al. (2008) Blood-Muscle Model**

Bruce and Bruce (2003; Bruce et al. 2008) developed a multi-compartment model that simulates kinetics of carbon monoxide, blood COHb, and muscle COMb (Figure 3-10; Table 3-16). The model includes compartments for blood, muscle, and other non-vascular tissues. In blood, carbon monoxide exists as dissolved carbon monoxide in equilibrium with COHb. Binding of carbon monoxide and O<sub>2</sub> to Hb is assumed to achieve instantaneous equilibrium of carbon monoxide with Hb, defined by the Haldane equation, with adjustment for cooperative binding of carbon monoxide and O<sub>2</sub> to Hb and myoglobin. Simulation of arterial and venous blood includes a time delay to account for the time required to achieve mixing of arterial blood leaving the lung and mixed venous blood. The muscle compartment consists of two subcompartments, between which dissolved carbon monoxide exchanges by diffusion.

Subcompartment 1 represents tissue that exchanges dissolved carbon monoxide with the arterial and venous vasculature (representing arterial-venous shunting). Subcompartment 2 represents tissue that exchanges with capillary blood. Dissolved carbon monoxide in muscle is in equilibrium with myoglobin, with the equilibrium defined by the Haldane equation adjusted for cooperative binding of O<sub>2</sub>. Carbon monoxide in the other tissue compartment is assumed to consist entirely of dissolved carbon monoxide.

**Validation of the model.** The Bruce et al. (2008) model was evaluated with data collected in experiments various experiments conducted in human volunteers (Benignus et al. 1994; Burge and Skinner 1995; Peterson and Stewart 1975; Tikuisis et al. 1987a). The evaluations are reported in Bruce and Bruce (2003, 2006) and Bruce et al. (2008). The model predicted observed arterial and venous blood COHb differences and kinetics during and following single or multiple brief exposures to carbon monoxide (Benignus et al. 1994). The model predicted continued uptake of carbon monoxide into muscle during the first 300 minutes following cessation of exposure, with subsequent return of muscle carbon monoxide to blood. (Bruce and Bruce 2006) Blood-muscle carbon monoxide exchange, in combination with elimination of carbon monoxide from blood to exhaled air, resulted in multi-phase blood elimination kinetics, with the early faster phase dominated by transfer of carbon monoxide from blood to muscle, and the later slower phase contributed by return of carbon monoxide from muscle to blood and elimination from blood to exhaled air. The multi-phase kinetics predicted from the model agreed with post-exposure elimination kinetics of carbon monoxide from blood observed in Benignus et al. (1994). The model also predicted blood COHb kinetics observed in 1- and 5-minute exposures observed in Tikuisis et al. (1987a).

## 3. HEALTH EFFECTS

**Figure 3-10. Structure of the Bruce et al. (2008) Model\***

\*The model simulates diffusive exchange of carbon monoxide (CO) and oxygen (O<sub>2</sub>) between alveolar space and capillaries and tissue capillary blood and tissues. Competitive cooperative binding of carbon monoxide and O<sub>2</sub> with hemoglobin (Hb) occurs in blood and to myoglobin (Mb) in muscle. No binding is assumed in the other tissue compartments. In muscle, diffusive exchange is assumed to occur between blood and both muscle subcompartments, as well as between muscle compartments.

Sources: Adapted from Bruce and Bruce (2003); Bruce et al. (2008)

## 3. HEALTH EFFECTS

**Table 3-16. Parameter Values for the Bruce et al. (2008) Model<sup>a</sup>**

Parameter	Definition	Unit	Value
<b>Chemical parameters</b>			
D <sub>L</sub> CO	Lung CO diffusion capacity	mL CO/minute-Torr	30
D <sub>M</sub> CO	Muscle CO diffusion capacity	mL CO/minute-Torr	1.75
D <sub>T</sub> O <sub>2</sub>	Within tissue diffusion coefficient for O <sub>2</sub>	mL/minute-Torr	0.00060
D <sub>T</sub> CO	Within tissue diffusion coefficient for O <sub>2</sub>	mL/minute-Torr	0.00045
Hb	Blood Hb concentration	g/mL	0.14
Hb <sub>F</sub>	Fetal blood Hb concentration	g/mL	0.155
M <sub>Hb</sub>	Haldane coefficient for Hb	unitless	218
M <sub>Mb</sub>	Haldane coefficient for Mb	unitless	36
n	Hill equation exponent	unitless	variable <sup>a</sup>
O <sub>2</sub> Hb <sub>Max</sub>	O <sub>2</sub> capacity of Hb	mL O <sub>2</sub> /g Hb	1.381
P <sub>B</sub>	Barometric pressure	Torr	760
P <sub>A</sub> O <sub>2</sub>	Partial pressure of O <sub>2</sub> in arterial blood	Torr	100
P <sub>M</sub> O <sub>2</sub>	Partial pressure of O <sub>2</sub> in muscle	Torr	20
P <sub>50</sub> Hb	Partial pressure of at 50% Hb saturation	Torr	variable <sup>a</sup>
P <sub>50</sub> Mb	Partial pressure of at 50% Mb saturation	Torr	2.32
MRO <sub>2</sub> B	Whole body O <sub>2</sub> consumption	mL/minute/g	0.0032
MRO <sub>2</sub> M	Muscle O <sub>2</sub> consumption	mL/minute/g	0.021–0.03
SO <sub>2</sub>	Solubility of O <sub>2</sub> in plasma	mL/Torr	3.14·10 <sup>-5</sup>
SCO	Solubility of CO in plasma	mL/Torr	2.35·10 <sup>-5</sup>
VO <sub>2</sub>	Total body O <sub>2</sub> consumption	mL/minute	225
VCO	Endogenous CO production	ml/minute	0.007
<b>Volumes</b>			
V <sub>M</sub>	Volume of muscle	L	29.1
V <sub>M1</sub> F	Volume of muscle subcompartment 1	L	0.4·V <sub>M</sub>
V <sub>M2</sub> F	Volume of muscle subcompartment 1	L	0.6·V <sub>M</sub>
V <sub>OT</sub>	Volume of other tissue	L	9.6

## 3. HEALTH EFFECTS

**Table 3-16. Parameter Values for the Bruce et al. (2008) Model<sup>a</sup>**

Parameter	Definition	Unit	Value
Flows			
Q <sub>C</sub>	Cardiac output	mL/minute	5,800
Q <sub>Mf</sub>	Blood flow fraction to muscles	unitless	0.40
Q <sub>OTf</sub>	Blood flow fraction to other tissue	unitless	0.25
V <sub>A</sub>	Alveolar ventilation rate	mL/minute	5,500

<sup>a</sup>Estimated from Hill equation:

$$O_2Hb = O_2Hb_{max} \cdot \frac{(PO_2/P_{50})^n}{1 + (PO_2/P_{50})^n}$$

Values for *n* ranged from approximately 2.34 at COHb=0% to 1.8 at COHb=100%. Values for P<sub>50</sub>Hb ranged from 25 at COHb=0% to 0 at COHb=100%

CO = carbon monoxide; COHb = carboxyhemoglobin; Hb = hemoglobin

### 3. HEALTH EFFECTS

Model predictions match the CFK model when muscle diffusion is set to zero, and the model predicts lower blood COHb levels than the CFK model, when the muscle carbon monoxide diffusion is enabled. This difference reflects the blood-muscle kinetics and storage of carbon monoxide in muscle myoglobin that is not accounted for in the CFK model.

**Risk assessment.** Risk assessment applications of the Bruce et al. (2008) model were not identified in the available literature. However, the model has been applied to predicting the outcome of breathing 100% O<sub>2</sub> on blood COHb and muscle COMb kinetics (Bruce and Bruce 2006). The model predicted a decrease in blood and muscle elimination half-times. Of particular relevance to risk assessment is the prediction of continued uptake of carbon monoxide from blood to muscle that occurs for several hours after cessation of exposure, which contributes to slowing the elimination kinetics of carbon monoxide from the body.

**Target tissues.** The Bruce et al. (2008) model predicts blood COHb and muscle COMb levels and can also be used to predict blood and muscle carbon monoxide burden.

**Species extrapolation.** The Bruce et al. (2008) model has been applied only to predicting blood COHb and muscle COMb levels in humans. The model could be applied to other mammalian species providing that physiological and chemical-specific parameter values are available for species of interest.

**Interroute extrapolation.** The Bruce et al. (2008) model simulates inhalation exposures.

## 3.5 MECHANISMS OF ACTION

### 3.5.1 Pharmacokinetic Mechanisms

Pharmacokinetic mechanisms of carbon monoxide action are related to its binding to heme proteins. Kinetics and equilibria of binding to the major heme proteins that have the greatest influence on carbon monoxide pharmacokinetics (Hb and myoglobin) are discussed in the Section 3.4, Toxicokinetics.

**Absorption.** Inhaled carbon monoxide is rapidly and extensively absorbed into blood (Benignus et al. 1994; Burge and Skinner 1995; Peterson and Stewart 1970, 1975; Tikuisis et al. 1987b). Inhaled carbon monoxide is transported to lung alveoli as result of convective forces in the respiratory tract and diffusion. At the alveolar gas-blood interface, carbon monoxide dissolves into pulmonary capillary plasma and, from plasma, diffuses into erythrocytes and other tissues. Binding of carbon monoxide by erythrocyte Hb

## 3. HEALTH EFFECTS

(to form COHb) contributes to maintaining relatively low concentrations of dissolved carbon monoxide in erythrocyte cytosol and a partial pressure gradient to drive carbon monoxide transfer from alveolar air to blood. Diffusion of carbon monoxide into erythrocytes and binding of carbon monoxide to Hb is sufficiently rapid that near equilibrium is achieved between carbon monoxide partial pressures in alveolar air and alveolar end capillary (arterial) blood (Chakraborty et al. 2004). Continued uptake of carbon monoxide results from systemic elimination processes that lower mixed venous blood carbon monoxide concentrations returned to alveolar capillaries (Benignus et al. 1994).

**Distribution.** The distribution of absorbed carbon monoxide is determined, in large part, by its affinity for binding to heme. Binding of carbon monoxide to intracellular heme proteins contributes to maintaining gradients for carbon monoxide diffusion from plasma into erythrocytes and extravascular tissues. The dominant heme proteins that influence carbon monoxide distribution are erythrocyte Hb and muscle myoglobin. The distribution of carbon monoxide in the body is largely a reflection of the distribution of Hb and myoglobin, with the largest carbon monoxide burden found in blood, heart, skeletal muscle, and spleen (Vreman et al. 2005, 2006). Relatively large contributions of blood and skeletal muscle to total body carbon monoxide content is consistent with the contributions that these tissues make to total body Hb and muscle myoglobin (Bruce and Bruce 2003). In a typical adult, blood contains approximately 12 mmoles of Hb with a total binding capacity of approximately 48 mmole carbon monoxide when completely saturated, and muscle contains approximately 8 mmoles of myoglobin with a total binding capacity of 8 mmole carbon monoxide when completely saturated.

**Metabolism.** Metabolism of carbon monoxide consists of three major processes: (1) metabolic production of carbon monoxide from endogenous and exogenous precursors; (2) binding of carbon monoxide to heme proteins; and (3) oxidative metabolism of carbon monoxide to CO<sub>2</sub>.

***Metabolic Production of Carbon Monoxide.*** Endogenous carbon monoxide is produced, primarily, from the enzymatic degradation of heme by the enzyme HO. Carbon monoxide is produced in all tissues that express HO; however, the major tissues contributing to carbon monoxide production are also major sites of heme metabolism, including liver, spleen, and reticuloendothelial system (Berk et al. 1974). Three isoforms of HO have been identified, of which, HO-1 is inducible, whereas, HO-2 and HO-3 are constitutive. In general, factors that induce HO-1 increase carbon monoxide production, whereas inhibition of HO activity or expression of HO decreases carbon monoxide production. Numerous stimuli have been shown to upregulate HO-1 expression (Wu and Wang 2005): (1) those related to heme metabolism (e.g., heme and heme derivatives, endogenous carbon monoxide); (2) stimuli that produce or

## 3. HEALTH EFFECTS

are associated with physiological or oxidative stress such as hyperoxia, heat shock factors, oxidized lipids, lipopolysaccharides, hydrogen peroxide, radiation (including ultraviolet light), endotoxin, heavy metals and arsenite, and various cytokines (e.g., interleukins, TNF- $\alpha$ ); (3) substances that are mediators of vascular resistance (NO, angiotensin II); and (4) mitogens and other factors that regulate cell growth (e.g., growth factors, phorbol ester). Expression of the constitutive isoform, HO-2, is increased by estrogen, glucocorticoids, stimuli that inhibit HO, and stimuli that down regulate NO production (e.g., inhibition of NO synthase). In human cells (e.g., grown in culture), expression of HO-1 is down regulated in hypoxia, heat shock factors, and interferon- $\gamma$ . The response to the above factors is not uniform across species. For example, hypoxia and heat shock have been shown to induce HO in rodents, whereas the dominant effect in human tissues is downregulation (Nakayama et al. 2000; Shibahara et al. 2003; Wu and Wang 2005).

Carbon monoxide can be produced from oxidative metabolism of certain exogenous hydrocarbons (e.g., DCM), by isoform CYP2E1 of cytochrome P450 (Andersen et al. 1991); and from heme oxidase catalysis of products of auto-oxidation of phenols, photo-oxidation of organic compounds, and lipid peroxidation of cell membrane lipids (Rodgers et al. 1994).

***Carbon Monoxide Binding to Heme Proteins.*** Carbon monoxide competes for O<sub>2</sub> for binding to the iron site in heme. Kinetic constants and equilibrium constants for the binding of O<sub>2</sub> and carbon monoxide to Hb and myoglobin are provided in Table 3-12.

***Oxidative Metabolism of Carbon Monoxide.*** The mechanism of oxidative metabolism of carbon monoxide to CO<sub>2</sub> is thought involve catalysis by cytochrom c oxidase (Fenn 1970; Young and Caughey 1986).

**Excretion.** Absorbed carbon monoxide is eliminated from the body by exhalation and oxidative metabolism. Oxidative metabolism of carbon monoxide has been estimated to be a relatively small fraction (<10%) of endogenous carbon monoxide production (Luomanmaki and Coburn 1969). Under most conditions, the dominant route of elimination of absorbed carbon monoxide is exhalation. The mechanism of elimination of carbon monoxide by exhalation is diffusion, with the driving forces at the alveolar interface being the partial pressure difference between carbon monoxide in alveolar air and in alveolar capillary blood. The latter is maintained at a relatively low level by binding of carbon monoxide to Hb. Diffusion is also the mechanism of release of carbon monoxide from intracellular stores to blood, with the driving forces influenced by binding of carbon monoxide to extravascular heme proteins (e.g., myoglobin) and blood Hb.

## 3. HEALTH EFFECTS

**3.5.2 Mechanisms of Toxicity**

Carbon monoxide exerts effects on cell metabolism through both hypoxic and non-hypoxic mechanisms. Both types of effects are thought to be largely (if not entirely) the result of the ability of carbon monoxide to bind to heme and alter function and/or metabolism of heme proteins. Carbon monoxide may also act through mechanisms unrelated to heme binding, although evidence for this is currently more contentious.

*Hypoxic Mechanisms.* Carbon monoxide produces tissue hypoxia by binding to and displacing O<sub>2</sub> from Hb. The formation of COHb decreases the O<sub>2</sub> carrying capacity of blood and impairs release of O<sub>2</sub> from Hb for its utilization in tissues (see Section 3.4.3, Metabolism for further discussion of kinetics of binding of carbon monoxide and O<sub>2</sub> to Hb). Through similar mechanisms, carbon monoxide decreases O<sub>2</sub> storage in muscle cells by binding to and displacing O<sub>2</sub> from myoglobin. The principle mechanisms underlying the hypoxic mechanism of carbon monoxide are: (1) higher affinity of carbon monoxide for Hb than O<sub>2</sub>, and (2) increased binding affinity for O<sub>2</sub> from COHb. The overall affinity of Hb for carbon monoxide is approximately 230 times greater than that for O<sub>2</sub> (Chakraborty et al. 2004). As a result, at a partial pressure of carbon monoxide that is approximately 230-fold lower than ambient O<sub>2</sub> concentration (21%/230≈900 ppm), O<sub>2</sub> and carbon monoxide will bind to Hb in an equimolar ratio (i.e., COHb≈50%). Binding of carbon monoxide and O<sub>2</sub> to the four heme moieties of Hb is cooperative (Alcantara et al. 2007; Perrella and Di Cera 1999; Roughton 1970). The associative reaction rate (k<sub>a</sub>) becomes faster with successive additions of carbon monoxide or O<sub>2</sub>, increasing the affinity (K<sub>a</sub>) of Hb for O<sub>2</sub> and impairing the release of O<sub>2</sub> from Hb for utilization in tissues. Overall, the affinity increase is 588-fold due to cooperative events as more carbon monoxide binds to hemoglobin; however, this is not a monotonic function with each ligation. Once the first hemoglobin chain binds a carbon monoxide molecule, the affinity increases 16.5-fold when the second carbon monoxide molecule binds, 4.6-fold at the third ligation step, and 7.7-fold in the fourth ligation step (Perrella and Di Cera 1999).

Although all tissues are vulnerable to carbon monoxide-induced hypoxic injury, those having the highest O<sub>2</sub> demand are particularly vulnerable, including the brain and heart. Carbon monoxide-induced hypoxia triggers compensatory cardiovascular responses that include increased cardiac output (heart rate and stroke volume) and dilation of cardiac and cerebral vasculature. During exercise, cardiac work and O<sub>2</sub> consumption increase, and increase the vulnerability of the heart to carbon monoxide-induced hypoxic injury. In patients who have underlying coronary artery or myocardial disease, the increased O<sub>2</sub> demand of exercise, in combination with decreased O<sub>2</sub> carrying capacity of blood and impaired release of O<sub>2</sub> from

## 3. HEALTH EFFECTS

Hb, can produce cardiac toxicity (e.g., myocardial ischemia, exertional angina, arrhythmia), at lower carbon monoxide exposure levels than experienced in healthy individuals (Adams et al. 1988; Allred et al. 1989, 1991; Anderson et al. 1973; Kleinman et al. 1989, 1998; Leaf and Kleinman 1996b).

Brain hypoxia induced by carbon monoxide can result in various symptoms of impaired central nervous system function, including headache, dizziness, drowsiness, weakness, nausea, vomiting, confusion, disorientation, irritability, visual disturbances, convulsions, and coma (Dolan 1985; Ernst and Zibrak 1998). Delayed development of neuropsychiatric and neurological impairment may occur within 1–4 weeks of exposure, with symptoms including inappropriate euphoria, impaired judgment, poor concentration, memory loss, cognitive and personality changes, psychosis, and Parkinsonism (Choi 2002; Ernst and Zibrak 1998; Kao and Nañagas 2006; Klawans et al. 1982; Raub and Benignus 2002; Raub et al. 2000; Ringel and Klawans 1972; WHO 1999; Wolf et al. 2008).

*Non-Hypoxic Mechanisms.* Most of the non-hypoxic mechanisms of action of carbon monoxide have been attributed to binding of carbon monoxide to heme in proteins other than Hb. A list of heme proteins that have been shown to undergo functional modification in response to carbon monoxide is presented in Table 3-17. Notable targets of carbon monoxide include components of several important physiological regulatory systems, including: brain and muscle O<sub>2</sub> storage and utilization (myoglobin, neuroglobin); nitric oxide NO cell signaling pathway (e.g., nitric oxide synthase [NOS], guanylyl cyclase [GC]); prostaglandin cell signaling pathway (cyclooxygenase, prostaglandin H synthase); energy metabolism and mitochondrial respiration (cytochrome c oxidase, cytochrome c, NADPH oxidase); steroid and drug metabolism (cytochrome P450 [CYP450]), cellular redox balance and reactive oxygen species (ROS) (catalase, peroxidases); and various transcription factors (e.g., neuronal PAS domain protein, NPAS2).

Endogenously produced carbon monoxide may participate in the physiological regulation of some, if not all, of these systems. This has potentially important implications for the understanding of carbon monoxide toxicology and dose-response relationships for the following reasons: (1) carbon monoxide modulation of physiological processes may underlie some aspects of the toxicity of exogenous carbon monoxide; (2) exogenous carbon monoxide may disrupt physiological regulation of those systems that are responsive to endogenous carbon monoxide (e.g., vascular resistance); and (3) exposures to exogenous carbon monoxide may affect carbon monoxide-mediated physiological responses at levels that approach those resulting from endogenous production. Whole-body endogenous carbon monoxide production rate has been estimated to be approximately 16.4 µmol/hour (0.007 mL/minute, standard temperature, pressure, density [STPD]; Coburn 1970a). Local rates of production of carbon monoxide in tissues are

## 3. HEALTH EFFECTS

**Table 3-17. Heme Proteins Modulated by Carbon Monoxide**

Name of protein	Effects or carbon monoxide or heme oxygenase up-regulation	Reference
Catalase	Inhibition	Hu and Kincaid 1992
Cyclooxygenase	Inhibition	Alcaraz et al. 2003
	Activation	Lee et al. 2001
Cytochrome c	Binding	Bhuyan and Kumar 2002
Cytochrome P450	Inhibition	Estabrook et al. 1980
Cytochrome c oxidase	Inhibition	Wohlrab and Ogunmola 1971
Guanylyl cyclase	Activation	Stone and Marletta 1994
Hemoglobin	Inhibition	Forster 1970
Neuroglobin	Inhibition	Burmester et al. 2000
Myoglobin	Inhibition	Coburn and Mayers 1971
Cytoglobin	Inhibition	Geuens et al. 2003; Sawai et al. 2003
NADPH oxidase	Inhibition	Taille et al. 2004
NO synthases	Inhibition	Thorup et al. 1999; Willis et al. 1995
Peroxidases	Binding	Carlsson et al. 2005
Prostaglandin H synthase	Binding	Lou et al. 2000
Tryptophan dioxygenase	Inhibition	Piantadosi 2002
NPAS2	Inhibition	Dioum et al. 2002

Source: Wu and Wang 2005


## 3. HEALTH EFFECTS

influenced by many tissue-specific factors, including those that regulate intracellular heme oxygenase activity (e.g. some areas of brain have relatively high HO-2 activities). A similar whole-body rate of delivery of carbon monoxide to the alveolar region of the lung would be achieved by inhaling air at a carbon monoxide concentration of approximately 1–2 ppm (i.e., assuming alveolar ventilation rate is approximately 4,000 mL/minute;  $0.007 \text{ mL/minute}/4,000 \text{ mL/minute} = 1.75 \cdot 10^{-6} \text{ mL CO/mL air, STPD} = 1.75 \text{ ppm}$ ). Such levels are only modestly above national average ambient air carbon monoxide concentrations in the United States (approximately 0.5 ppm, see Chapter 6) and well within variability observed in measurements made with personnel air monitors (Chang et al. 2000). One implication of this is that, if a dose threshold for effects of exogenous carbon monoxide exists at all, it may lie near or below ambient air carbon monoxide concentrations. Although currently available toxicology and epidemiological studies provide evidence for increasing severity of adverse carbon monoxide effects in association with increasing levels of exposure, these studies do not provide strong evidence supporting or refuting a dose threshold for adverse effects.

Of the many physiological systems that may contribute to non-hypoxic mechanisms of carbon monoxide action, the following are thought to be particularly important and are the subject of ongoing intense research interests: modulation of cell signaling pathways and generation of ROS.

***Carbon Monoxide Modulation of Cell Signaling.*** Cell signaling pathways that are thought to be modulated by carbon monoxide are depicted in Figure 3-11. These pathways contribute to the regulation of various physiological processes, including inflammatory responses, apoptosis, cell proliferation, thrombosis (i.e., platelet aggregation), smooth muscle tone (including vasodilation and relaxation of extravascular smooth muscle) and synaptic neurotransmission in the central and peripheral nervous systems (e.g., glutamate). The nitric oxide radical (NO•) plays an important signal transduction role in regulating smooth muscle relaxation, synaptic neurotransmission (e.g., glutamate), platelet function, and other physiological processes (Denninger and Marletta 1999). One of the major transduction pathways mediated by NO• is the NO•-cGMP pathway, in which NO• produced by the enzyme NOS binds to the heme moiety of soluble guanylate cyclase (sGC), which catalyzes conversion of guanosine triphosphate (GTP) to cyclic guanosine monophosphate (cGMP). The latter (cGMP) activates various downstream signaling cascades that regulate cell function (e.g., cGMP-dependent protein kinase, cGMP-regulated cation channels, and cGMP-regulated phosphodiesterase). Carbon monoxide binds to the heme moieties of both NOS and sGC; and in doing so, inhibits NOS activity and activates sGC activity (Stone and Marletta 1994; Thorup et al. 1999). The potency of carbon monoxide for activating sGC is approximately 20-fold less than that of NO•; as a result, in the presence of NO•,

## 3. HEALTH EFFECTS

**Figure 3-11. Potential Signaling Pathways Modulated by Carbon Monoxide**

cGMP = cyclic guanosine monophosphate; CO = carbon monoxide; COX = cyclooxygenase; CRF = corticotropin releasing factor; CNS = central nervous system; CYP450 = cytochrome P450; ERK = extracellular signal regulated kinase; 20-HETE = 20-hydroxyeicosatetraenoic acid; HSF = heat shock factor; HSP = heat shock protein; IL = interleukin; JNK = a stress activate protein kinase; MAPK = mitogen activate protein kinase; MIP = macrophage inflammatory protein; NANC = non-noradrenergic non-cholinergic; NO• = nitric oxide radical; NOS = nitrogen oxide synthase; PDE = phosphodiesterase; PG = prostaglandins; PK = protein kinase; sGC = soluble guanylate cyclase; TNF = tumor necrosis factor

Adapted from Ryter et al. 2006; Volti et al. 2008

## 3. HEALTH EFFECTS

carbon monoxide may modulate sGC activity downward by displacing NO• from sGC (Stone and Marletta 1994). The NO•-cGMP signaling pathway has been implicated in the mechanism by which carbon monoxide produces relaxation of smooth muscle (Ny et al. 1996) and inhibition of platelet homotypic aggregation (Brune and Ullrich 1987). Carbon monoxide increases platelet-neutrophil interactions (heterotypic aggregation) through a free-radical mechanism (Thom et al. 2006).

Carbon monoxide modulates mitogen activated protein kinase (MAPK) signaling pathways, either through its interaction with sGC (Ryter et al. 2006) or through secondary effects on ROS generation (see below) MAPK pathways are important regulators of cytokine production and release in inflammatory responses, and in regulating apoptosis and proliferation.

Carbon monoxide exposure can perturb heme protein binding by nitric oxide, induce a variety of proteins including heme oxygenase (HO), superoxide dismutase, and nitric oxide synthase, alter mitochondrial function, and modify production of ATP and ROS (Brown and Piantadosi 1992; Davutoglu et al. 2006; Kim et al. 2007; Lee et al. 2006; Maulik et al. 1996; Nakao et al. 2008; Piantadosi et al. 1997; Thom et al. 1994, 1997, 2000; Zuckerbraun et al. 2003). Carbon monoxide increases production of ROS from mitochondria and also some non-mitochondrial sources, which can activate one or more protein kinases (e.g., p38 MAPK, c-Jun N-terminal kinase, ERK), signal transducers (e.g., peroxisome proliferator-activated receptor- $\gamma$ ), and transcriptional regulators (e.g., nuclear factor- $\kappa$ B, AP-1), as well as guanylate cyclase, and lead to secondary events such as vasodilation and fibrinolysis (Arruda et al. 2004; Fujita et al. 2001; Mishra et al. 2006; Ryter and Otterbein 2004; Soares et al. 2004; Wijayanti et al. 2004).

Carbon monoxide also binds to the heme moieties and inhibits CYP450 and cyclooxygenase (COX) and, thereby, can modify physiological processes regulated by products of these enzyme systems. Inhibition of CYP450 catalyzed production of eicosanoids (e.g., 20-hydroxyeicostetraenoic acid, [20-HETE]) and COX-mediated production of prostaglandins have been postulated as a potential mechanisms by which carbon monoxide may contribute to the regulation of vascular smooth muscle tone and vasodilation (Botros et al. 2002; Volti et al. 2008).

***Carbon Monoxide Modulation of ROS Generation.*** Carbon monoxide may promote and/or modify ROS generation through several different mechanisms. Binding to the heme moiety and inhibition of mitochondrial cytochrome c oxidase disrupts mitochondrial respiration and promotes ROS generation (Alonso et al. 2003; D'Amico et al. 2006; Favory et al. 2006a; Iheagwara et al. 2007; Zuckerbraun et al. 2007). Carbon monoxide also appears to promote a more general disruption of the regulation of iron and

## 3. HEALTH EFFECTS

ferritin levels, at least in part through promotion of heme protein degradation (Cronje et al. 2004; Ghio et al. 2008; Iheagwara et al. 2007).

### 3.5.3 Animal-to-Human Extrapolations

Studies conducted in nonhuman primates, dogs, and various rodent species have demonstrated the major health effects of carbon monoxide that have been observed in humans, including COHb-related hypoxia, cardiovascular responses, and neurological effects (Benignus 1994; Benignus et al. 1990; EPA 1991, 2000; Raub and Benignus 2002). These studies also provide support for the concept of nonhypoxic modes of action of carbon monoxide that operate on a variety of physiological systems (e.g., Wu and Wang 2005; also, see Section 3.5.2 for discussion and relevant references). These systems include physiological regulatory systems that function in all mammalian species, such as brain and muscle oxygen storage and utilization; nitric oxide cell signaling pathway; prostaglandin cell signaling pathway; energy metabolism and mitochondrial respiration; steroid and drug metabolism; cellular redox balance and reactive oxygen species; and various transcription factors. Therefore, results from mechanistic studies conducted in typical laboratory animal models are likely to be directly relevant to understanding mechanisms of action of carbon monoxide in humans. Manifestations of these various modes of action may differ qualitatively and quantitatively across species. For example, rats exhibit pronounced hypothermia in response to exposures to carbon monoxide whereas at low levels of exposure (e.g., COHb <20%), this response is not evident in humans (Benignus 1994; Gordon 1990). Differences in dose-response relationships between humans and animal models can also be expected, in part, due to differences in physiological and chemical determinants of carbon monoxide uptake, elimination, and binding kinetics with major heme proteins (Benignus and Annau 1994; Longo and Hill 1977) as well as interspecies differences in physiological regulation of systems that are nonhypoxic targets of carbon monoxide.

## 3.6 TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS

Recently, attention has focused on the potential hazardous effects of certain chemicals on the endocrine system because of the ability of these chemicals to mimic or block endogenous hormones. Chemicals with this type of activity are most commonly referred to as *endocrine disruptors*. However, appropriate terminology to describe such effects remains controversial. The terminology *endocrine disruptors*, initially used by Thomas and Colborn (1992), was also used in 1996 when Congress mandated the EPA to develop a screening program for "...certain substances [which] may have an effect produced by a naturally occurring estrogen, or other such endocrine effect[s]...". To meet this mandate, EPA convened a

### 3. HEALTH EFFECTS

panel called the Endocrine Disruptors Screening and Testing Advisory Committee (EDSTAC), and in 1998, the EDSTAC completed its deliberations and made recommendations to EPA concerning *endocrine disruptors*. In 1999, the National Academy of Sciences released a report that referred to these same types of chemicals as *hormonally active agents*. The terminology *endocrine modulators* has also been used to convey the fact that effects caused by such chemicals may not necessarily be adverse. Many scientists agree that chemicals with the ability to disrupt or modulate the endocrine system are a potential threat to the health of humans, aquatic animals, and wildlife. However, others think that endocrine-active chemicals do not pose a significant health risk, particularly in view of the fact that hormone mimics exist in the natural environment. Examples of natural hormone mimics are the isoflavonoid phytoestrogens (Adlercreutz 1995; Livingston 1978; Mayr et al. 1992). These chemicals are derived from plants and are similar in structure and action to endogenous estrogen. Although the public health significance and descriptive terminology of substances capable of affecting the endocrine system remains controversial, scientists agree that these chemicals may affect the synthesis, secretion, transport, binding, action, or elimination of natural hormones in the body responsible for maintaining homeostasis, reproduction, development, and/or behavior (EPA 1997). Stated differently, such compounds may cause toxicities that are mediated through the neuroendocrine axis. As a result, these chemicals may play a role in altering, for example, metabolic, sexual, immune, and neurobehavioral function. Such chemicals are also thought to be involved in inducing breast, testicular, and prostate cancers, as well as endometriosis (Berger 1994; Giwercman et al. 1993; Hoel et al. 1992).

No studies were located regarding endocrine disruption in humans and/or animals after exposure to carbon monoxide. Several cell signaling pathways that are thought to be modulated by endogenous carbon monoxide that could play a role in the regulation of activity of the hypothalamic-pituitary-adrenal (HPA) axis, including cGMP and NO• signaling (Mancuso et al. 1997; Snyder et al. 1998; Wu and Wang 2005). Modulation of endogenous carbon monoxide levels and/or production has been shown to modify various HPA axis process including release of arginine vasopressin (AVP), gonadotropin, and corticotropin releasing hormone from the hypothalamus, release of adrenocorticotrophic hormone (ACTH) from the pituitary, and response of the adrenal gland to ACTH (Wu and Wang 2005). The toxicological significance of these various actions of carbon monoxide has not been established.

#### 3.7 CHILDREN'S SUSCEPTIBILITY

This section discusses potential health effects from exposures during the period from conception to maturity at 18 years of age in humans, when all biological systems will have fully developed. Potential

## 3. HEALTH EFFECTS

effects on offspring resulting from exposures of parental germ cells are considered, as well as any indirect effects on the fetus and neonate resulting from maternal exposure during gestation and lactation. Relevant animal and *in vitro* models are also discussed.

Children are not small adults. They differ from adults in their exposures and may differ in their susceptibility to hazardous chemicals. Children's unique physiology and behavior can influence the extent of their exposure. Exposures of children are discussed in Section 6.6, Exposures of Children.

Children sometimes differ from adults in their susceptibility to hazardous chemicals, but whether there is a difference depends on the chemical (Guzelian et al. 1992; NRC 1993). Children may be more or less susceptible than adults to health effects, and the relationship may change with developmental age (Guzelian et al. 1992; NRC 1993). Vulnerability often depends on developmental stage. There are critical periods of structural and functional development during both prenatal and postnatal life, and a particular structure or function will be most sensitive to disruption during its critical period(s). Damage may not be evident until a later stage of development. There are often differences in pharmacokinetics and metabolism between children and adults. For example, absorption may be different in neonates because of the immaturity of their gastrointestinal tract and their larger skin surface area in proportion to body weight (Morselli et al. 1980; NRC 1993); the gastrointestinal absorption of lead is greatest in infants and young children (Ziegler et al. 1978). Distribution of xenobiotics may be different; for example, infants have a larger proportion of their bodies as extracellular water, and their brains and livers are proportionately larger (Altman and Dittmer 1974; Fomon 1966; Fomon et al. 1982; Owen and Brozek 1966; Widdowson and Dickerson 1964). The infant also has an immature blood-brain barrier (Adinolfi 1985; Johanson 1980) and probably an immature blood-testis barrier (Setchell and Waites 1975). Many xenobiotic metabolizing enzymes have distinctive developmental patterns. At various stages of growth and development, levels of particular enzymes may be higher or lower than those of adults, and sometimes unique enzymes may exist at particular developmental stages (Komori et al. 1990; Leeder and Kearns 1997; NRC 1993; Vieira et al. 1996). Whether differences in xenobiotic metabolism make the child more or less susceptible also depends on whether the relevant enzymes are involved in activation of the parent compound to its toxic form or in detoxification. There may also be differences in excretion, particularly in newborns who all have a low glomerular filtration rate and have not developed efficient tubular secretion and resorption capacities (Altman and Dittmer 1974; NRC 1993; West et al. 1948). Children and adults may differ in their capacity to repair damage from chemical insults. Children also have a longer remaining lifetime in which to express damage from chemicals; this potential is particularly relevant to cancer.

## 3. HEALTH EFFECTS

Certain characteristics of the developing human may increase exposure or susceptibility, whereas others may decrease susceptibility to the same chemical. For example, although infants breathe more air per kilogram of body weight than adults breathe, this difference might be somewhat counterbalanced by their alveoli being less developed, which results in a disproportionately smaller surface area for alveolar absorption (NRC 1993).

The fetus may be particularly vulnerable to maternal carbon monoxide exposure. Carbon monoxide in the maternal system distributes to fetal tissues. Measurements of steady-state fetal COHb concentrations in fetal and maternal blood of nonsmoking women have found fetal COHb concentrations to be approximately 10–15% higher than maternal blood (fetal/maternal ratio=1.1–1.15; Longo 1977). Binding affinity of fetal Hb is approximately twice that of maternal Hb (Di Cera et al. 1989). The higher binding affinity of fetal Hb as well as the relatively small diffusion gradients for carbon monoxide between maternal and fetal blood contribute to slower kinetics of COHb in fetal blood compared to maternal blood. As a result, with continuous exposure to a constant level of carbon monoxide in air, the kinetics of fetal blood COHb concentration would be expected to lag behind maternal kinetics and to reach a steady-state that is approximately 10–15% higher than maternal (Longo 1977). Similarly, following cessation of exposure, elimination kinetics of carbon monoxide from fetal blood would be expected to lag behind that of maternal blood (Hill et al. 1977).

Epidemiological studies have examined possible associations between exposure to ambient air carbon monoxide concentrations and various developmental outcomes, including pre-term birth, birth weight, congenital anomalies, and neonatal and infant death (see Section 3.2.6, Table 3-8). Results of these studies have been mixed and, collectively, do not provide strong evidence for developmental effects, including neonatal or infant mortality in association with exposures to ambient levels of carbon monoxide (<10 ppm). However, studies conducted in animals provide evidence of adverse developmental effects of gestational and early postnatal carbon monoxide in association with maternal exposure concentrations as low as 12–25 ppm. Observed effects have included decreased fetal weight, adverse central nervous system development, altered peripheral nervous system development, cardiac effects, altered sexual behavior, immunological effects, and hematological effects (see Table 3-9). In addition, some studies showed that developmental effects persisted beyond the postnatal period (Carratu et al. 1993, 2000a, 2000b; De Salvia et al. 1995; Lopez et al. 2003, 2008; Stockard-Sullivan et al. 2003; Webber et al. 2003).

### 3. HEALTH EFFECTS

Although no studies have directly compared the sensitivity of adults and children to the effects of carbon monoxide, several epidemiological studies have examined pulmonary function, asthma morbidity, and exacerbation of asthma symptoms in children (Chen et al. 1999; Fischer et al. 2002; Mortimer et al. 2008; Yu et al. 2000). Although results have been mixed, these studies provide some evidence for increased vulnerability of asthmatic children to carbon monoxide-associated respiratory effects (see Section 3.2.2, Respiratory Effects).

#### 3.8 BIOMARKERS OF EXPOSURE AND EFFECT

Biomarkers are broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility (NAS/NRC 1989).

The National Report on Human Exposure to Environmental Chemicals provides an ongoing assessment of the exposure of the U.S. population to environmental chemicals using biomonitoring. This report is available at <http://www.cdc.gov/exposurereport/>. The biomonitoring data for carbon monoxide from this report is discussed in Section 6.5. A biomarker of exposure is a xenobiotic substance or its metabolite(s) or the product of an interaction between a xenobiotic agent and some target molecule(s) or cell(s) that is measured within a compartment of an organism (NAS/NRC 1989). The preferred biomarkers of exposure are generally the substance itself, substance-specific metabolites in readily obtainable body fluid(s), or excreta. However, several factors can confound the use and interpretation of biomarkers of exposure. The body burden of a substance may be the result of exposures from more than one source. The substance being measured may be a metabolite of another xenobiotic substance (e.g., high urinary levels of phenol can result from exposure to several different aromatic compounds). Depending on the properties of the substance (e.g., biologic half-life) and environmental conditions (e.g., duration and route of exposure), the substance and all of its metabolites may have left the body by the time samples can be taken. It may be difficult to identify individuals exposed to hazardous substances that are commonly found in body tissues and fluids (e.g., essential mineral nutrients such as copper, zinc, and selenium). Biomarkers of exposure to carbon monoxide are discussed in Section 3.8.1.

Biomarkers of effect are defined as any measurable biochemical, physiologic, or other alteration within an organism that, depending on magnitude, can be recognized as an established or potential health impairment or disease (NAS/NRC 1989). This definition encompasses biochemical or cellular signals of tissue dysfunction (e.g., increased liver enzyme activity or pathologic changes in female genital epithelial

### 3. HEALTH EFFECTS

cells), as well as physiologic signs of dysfunction such as increased blood pressure or decreased lung capacity. Note that these markers are not often substance specific. They also may not be directly adverse, but can indicate potential health impairment (e.g., DNA adducts). Biomarkers of effects caused by carbon monoxide are discussed in Section 3.8.2.

A biomarker of susceptibility is an indicator of an inherent or acquired limitation of an organism's ability to respond to the challenge of exposure to a specific xenobiotic substance. It can be an intrinsic genetic or other characteristic or a preexisting disease that results in an increase in absorbed dose, a decrease in the biologically effective dose, or a target tissue response. If biomarkers of susceptibility exist, they are discussed in Section 3.10, Populations That Are Unusually Susceptible.

#### **3.8.1 Biomarkers Used to Identify or Quantify Exposure to Carbon Monoxide**

Measurement of blood COHb is the principal biomarker for identifying exposure to carbon monoxide. Nearly all carbon monoxide in the body exists as complexes with Hb and other heme proteins. Approximately 60–70% of the body burden exists as blood COHb (Vreman et al. 2006; Bruce and Bruce 2003). Therefore, measurement of COHb provides a useful indicator of the elevated carbon monoxide body burden, although the utility for blood COHb measurements for large-scale screening for elevated exposures or poisoning is questionable (Heckerling et al. 1990). The relationship between COHb levels and exposure is more complex because of the numerous physiological factors that influence carbon monoxide uptake and elimination. The elimination half-time of absorbed carbon monoxide has been estimated to be approximately 300 minutes (Bruce and Bruce 2006; Peterson and Stewart 1970); therefore, a value for COHb measured at a given time following exposure will reflect both the exposure and elimination rate of carbon monoxide. Several pharmacokinetics models have been developed that can be used to predict blood COHb levels that would correspond to a given exposure concentration and duration, in a typical adult and in the fetus resulting from maternal exposures (see Section 3.4.5). These models can be used to reconstruct possible carbon monoxide exposure scenarios (e.g., concentrations, duration, and gap between measurement of COHb and cessation of exposure) that could result in a given measured COHb level.

#### **3.8.2 Biomarkers Used to Characterize Effects Caused by Carbon Monoxide**

Although blood COHb reflects current carbon monoxide body burden, measurement of blood COHb has not been shown to be a reliable predictor of severity of acute toxicity (Hampson and Hauff 2008). Typical levels of COHb in nonsmokers are <2% (Adams et al. 1988; Allred et al. 1991; Anderson et al.

### 3. HEALTH EFFECTS

1973; Hinderliter et al. 1989; Kleinman et al. 1989, 1998; Sheps et al. 1987, 1990). Levels of 10% or more can be observed immediately after smoking a cigarette (Kao and Nañagas 2006). In general, signs and symptoms of acute carbon monoxide poisoning can present at COHb levels ranging from 3 to 24% (Hampson and Hauff 2008; Kao and Nañagas 2006). Levels ranging from 2 to 6% have been shown to exacerbate underlying cardiovascular disease, including enhanced myocardial ischemia and increased cardiac arrhythmias (Adams et al. 1988; Allred et al. 1989, 1991; Anderson et al. 1973; Kleinman et al. 1989, 1998; Leaf and Kleinman 1996b). More severe signs of carbon monoxide poisoning are poorly correlated with blood COHb, with loss of consciousness occurring at mean levels of 24.3% (range: 2–70) and fatality at mean levels of 32.1% (range: 3.0–60; Hampson and Hauff 2008). Exposures that result in COHb levels >50% are usually fatal (Kao and Nañagas 2006).

Diagnosis and characterization of carbon monoxide toxicity includes both the measurement of blood COHb and assessment of signs. In order of increasing severity, these include: (1) headache, nausea, dilation of cutaneous vasculature, vomiting, dizziness, blurred vision; (2) confusion, syncope, chest pain, dyspnea, weakness, tachycardia, tachynea rhabdomyolysis; and (3) palpitations, cardiac dysrhythmias, hypotension, myocardial ischemia, cardiac arrest, respiratory arrest, pulmonary edema, seizures, coma (Kao and Nañagas 2006). Neuropsychological testing and neuroimaging have been considered for evaluating central nervous system effects (Tomaszewski 2006).

Cardiac enzyme markers are associated with myocardial dysfunction and an elevated risk for long-term cardiac mortality following carbon monoxide poisoning; biochemical markers for brain injury such as neuron-specific enolase and S-100 beta protein have not been found to reliably correlate with severity of poisoning or clinical outcome (Brvar et al. 2004; Davutoglu et al. 2006; Kalay et al. 2007; Rasmussen et al. 2004; Satran et al. 2005).

#### 3.9 INTERACTIONS WITH OTHER CHEMICALS

The effects of combined exposure of carbon monoxide with other chemicals (e.g., pollutants, drugs) were reviewed by the EPA (1991, 2000). In general, research has mainly focused on effects of combined chemical exposures on the cardiovascular, respiratory and central nervous systems. As discussed in Section 3.10 (Populations that Are Unusually Susceptible), in addition to interactions with other chemicals, exposure to carbon monoxide at a higher altitude may result in higher body burdens of carbon monoxide and, therefore, there may be higher vulnerability to carbon monoxide at higher altitudes than at

## 3. HEALTH EFFECTS

lower altitudes (McGrath et al. 1993), in particular, for people who have not physiologically adapted to higher altitudes (Horvath et al. 1988; Hsia 2002).

**Oxygen.** The chemical interaction of highest toxicological and clinical significance is that between carbon monoxide and O<sub>2</sub>. Carbon monoxide and O<sub>2</sub> compete for binding to heme (Chakraborty et al. 2004; Gibson et al. 1986). Binding to and displacing O<sub>2</sub> from heme proteins, including blood Hb and muscle myoglobin, has profound effects on the kinetics of distribution and elimination kinetics of both carbon monoxide and O<sub>2</sub>, and is also the principal mechanism underlying the hypoxic effects of carbon monoxide (Bruce and Bruce 2006; Bruce et al. 2008; Vreman et al. 2005, 2006). Decreasing levels of O<sub>2</sub> in inspired air facilitates carbon monoxide binding to Hb and myoglobin and exacerbates carbon monoxide-induced hypoxia. This interaction can contribute to a higher vulnerability to carbon monoxide toxicity at lower O<sub>2</sub> pressures that occur at higher altitudes (Horvath et al. 1988; Hsia 2002; McGrath et al. 1993). Increasing levels of O<sub>2</sub> in inspired air facilitates displacement of carbon monoxide from Hb and myoglobin, increases the rate of carbon monoxide elimination from the body, and increases O<sub>2</sub> availability to tissues (Landaw 1973; Peterson and Stewart 1970; Weaver et al. 2000). These actions of O<sub>2</sub> form the basis for the use of 100% O<sub>2</sub> and hyperbaric O<sub>2</sub> therapy in the treatment of carbon monoxide poisoning (Lavonas 2007; Tomaszewski 2006).

**Cyanide.** Carbon monoxide and cyanide poisonings can occur concomitantly in victims of smoke inhalation (Lundquist et al. 1989; Shusterman et al. 1996; Wetherell 1966). Experimental studies conducted in animal models provide evidence that carbon monoxide and cyanide can produce synergistic toxicity (Moore et al. 1987; Norris et al. 1986; Pitt et al. 1979).

**Drugs Affecting the Central Nervous System.** Due to effects of carbon monoxide on the central nervous system, it is anticipated that combined exposure to drugs with activity in the central nervous system, including therapeutic agents and drugs of abuse, could enhance the central nervous system toxicity of carbon monoxide (EPA 1991, 2000). However, the available data are not sufficient to determine if effects of combined exposures are additive or exhibit other types of interaction relationships (e.g., synergistic). Enhanced carbon monoxide-induced central nervous system toxicity has been reported to occur with concomitant exposures to alcohol, barbiturates, amphetamine, chlorpromazine, nicotine, diazepam, and morphine (EPA 1991).

**Drugs Affecting the Cardiovascular System.** In response to carbon monoxide-induced tissue hypoxia, alterations in hemodynamics and responsive compensatory mechanisms (e.g., vasodilation and increased

### 3. HEALTH EFFECTS

cardiovascular output) occur, including increased coronary blood flow, decreased myocardial O<sub>2</sub> consumption, increased heart rate, and alterations in blood flow to various vascular beds (e.g., cerebral, limb muscular). Thus, drugs used to treat patients with coronary artery disease (e.g., beta-blockers, calcium-channel blockers, nitrates) potentially may affect susceptibility to carbon monoxide. This is of particular concern, as patients with cardiovascular disease are more susceptible to carbon monoxide-induced cardiovascular toxicity than are healthy individuals. However, little data evaluating potential interactions with therapeutic agents used to treat cardiovascular disease are available (EPA 1991, 2000).

***Air Pollution.*** Numerous epidemiological studies have examined associations between exposure to carbon monoxide and other air pollutants (e.g., NO<sub>2</sub>, O<sub>3</sub>, particular matter, SO<sub>2</sub>). In general, these studies have found evidence for multiple air pollutant contributors to outcomes observed, including cardiovascular, respiratory, and mortality end points (see discussion of epidemiology in Section 3.2). The interactive aspects of these associations have not been fully explored or elucidated. However, evidence is provided from several studies of increased risks of adverse cardiovascular and respiratory outcomes in the context of ongoing cardiovascular disease (e.g., ischemic heart disease) and respiratory diseases (e.g., asthma). These studies suggest that it is both plausible and likely that interactions contribute to outcomes observed in populations that experience co-exposures to carbon monoxide, NO<sub>2</sub>, O<sub>3</sub>, particular matter, and SO<sub>2</sub>.

***Tobacco Smoke.*** In addition to exposures to carbon monoxide, tobacco smoking results in exposures to a variety of agents that act directly on the heart to increase cardiac O<sub>2</sub> demand (e.g., nicotine, Benowitz 1997) and that contribute to respiratory and coronary vascular disease, both of which can render people more vulnerable to carbon monoxide toxicity (see Section 3.10, Populations that are Unusually Susceptible).

#### 3.10 POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE

A susceptible population will exhibit a different or enhanced response to carbon monoxide than will most persons exposed to the same level of carbon monoxide in the environment. Reasons may include genetic makeup, age, health and nutritional status, and exposure to other toxic substances (e.g., cigarette smoke). These parameters result in reduced detoxification or excretion of carbon monoxide, or compromised function of organs affected by carbon monoxide. Populations who are at greater risk due to their unusually high exposure to carbon monoxide are discussed in Section 6.7, Populations with Potentially High Exposures.

## 3. HEALTH EFFECTS

***Cardiovascular Disease.*** Studies of the cardiovascular effects of inhalation exposures to carbon monoxide conducted in controlled human clinical studies, epidemiology studies, and various animal models (monkeys, dogs, rats, and rabbits) provide convincing evidence for adverse cardiovascular effects in association with carbon monoxide exposures that result in blood COHb levels of  $\geq 2.4\%$ , with effects occurring at the lowest levels in subjects who have compromised cardiovascular function (e.g., coronary artery disease). Therefore, exposure to carbon monoxide appears to exacerbate underlying cardiovascular disease, and people with cardiovascular disease that renders them vulnerable to myocardial ischemia (e.g., coronary artery disease) represent a population that would be unusually susceptible to carbon monoxide. Epidemiological studies of exposure to ambient concentrations of carbon monoxide and cardiovascular outcomes also provide supporting evidence for higher vulnerability of people with pre-existing heart conditions such as ischemic heart disease, people with a history of cardiac arrhythmia, and the elderly (Barnett et al. 2006; D'Ippoliti et al. 2003; Dockery et al. 2005; Hosseinpoor et al. 2005; Lanki et al. 2006; Lee et al. 2003b; Mann et al. 2002; Rich et al. 2005; Szyszkowicz 2007; von Klot et al. 2005).

***Asthma and COPD.*** Several studies have examined possible associations between ambient air carbon monoxide concentrations and pulmonary function in subjects who had ongoing lung disease (e.g., asthma, chronic obstructive lung disease) and who might be more sensitive to agents that affect pulmonary function (Rabinovitch et al. 2004; Silkoff et al. 2005; Timonen et al. 2002), as well as asthma morbidity and asthma and exacerbation of symptoms (Park et al. 2005a; Rabinovitch et al. 2004; Schildcrout et al. 2006; Silkoff et al. 2005; Slaughter et al. 2003; von Klot et al. 2002; Yu et al. 2000). Results of these studies have been mixed; however, collectively, these studies provide evidence for higher vulnerability of people who have ongoing respiratory disease, including asthma and chronic obstructive pulmonary disease. This subpopulation may also include a subpopulation of former tobacco smokers (Silkoff et al. 2005). As noted in Section 3.7, Children's Susceptibility, the fetus may be particularly vulnerable to exposures that occur during pregnancy.

***Anemia.*** Conditions that produce anemia (e.g., iron deficiency, hemolysis, hemorrhage, or reduced hematopoiesis) decrease the oxygen carrying capacity of blood. Hemolytic anemia increases heme metabolism and endogenous carbon monoxide production. These factors would be expected to render individuals more vulnerable to hypoxic effects of carbon monoxide.

***Environmental and Behavioral Factors.*** In addition to ongoing diseases, other environmental and/or behavior factors may increase vulnerability to carbon monoxide toxicity. In general, factors that increase

### 3. HEALTH EFFECTS

alveolar ventilation and cardiac output, while maintaining alveolar ventilation perfusion matching (i.e., proportional increases in alveolar ventilation rate and cardiac output) tend to increase absorption of inhaled carbon monoxide. Factors that increase carbon monoxide absorption include moderate exercise, supine position, age (increased in infancy and childhood and declines in adults with age), altitude, increasing blood Hb concentration, and decreasing partial pressure of O<sub>2</sub> in inhaled air or blood.

These factors could contribute to higher body burdens of carbon monoxide and, therefore, higher vulnerability to carbon monoxide at higher altitudes than at lower altitudes (McGrath et al. 1993), in particular, for people who have not physiologically adapted to higher altitudes (Horvath et al. 1988; Hsia 2002). Exercise increases oxygen demand, absorption of carbon monoxide, and transfer of carbon monoxide from blood to muscle, and decreases elimination of carbon monoxide (Joumard et al. 1991; Richardson et al. 2002; Werner and Lindahl 1980); these factors may contribute to higher vulnerability during strenuous exercise.

#### 3.11 METHODS FOR REDUCING TOXIC EFFECTS

This section describes clinical practice and research concerning methods for reducing toxic effects of exposure to carbon monoxide. However, because some of the treatments discussed may be experimental and unproven, this section should not be used as a guide for treatment of exposures to carbon monoxide. When specific exposures have occurred, poison control centers and medical toxicologists should be consulted for medical advice. The following texts provide specific information about treatment following exposures to carbon monoxide:

Dart RC, ed. 2004. *Medical toxicology*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins, 1208.

Lavonas EJ. 2007. Carbon monoxide poisoning. In: Haddad and Winchester's clinical management of poisoning and drug overdose. 4th ed. Philadelphia, PA: Saunders, 1297-1307

Tomaszewski C. 2006. Carbon monoxide. In: Goldfrank's toxicologic emergencies. New York, NY: McGraw-Hill, 1689-1704.

##### 3.11.1 Reducing Peak Absorption Following Exposure

Human exposure to carbon monoxide occurs by the inhalation route. General recommendations for reducing absorption of carbon monoxide following acute high-level inhalation exposure have included immediate removal of the exposed individual from the contaminated area and administering 100%

### 3. HEALTH EFFECTS

oxygen through a face mask that does not allow rebreathing of expired air or endotracheal tube for a minimum of 4 hours. For patients with life-threatening carbon monoxide exposure, hyperbaric oxygen (HBO) therapy has also been suggested, although this remains a controversial method for treatment (Dart 2004; Lavonas 2007; Tomaszewski 2002; Weaver et al. 2000; Wolf et al. 2008). Hyperbaric oxygen treatment is optimal within 6 hours of exposure, as patients treated after 6 hours have displayed a higher percent of delayed sequelae (30 vs. 19%) and mortality (30 vs. 14%) (Tomaszewski 2006).

Studies conducted in animal models have provided evidence for a potential influence of isocapnia or hypercapnia (i.e., normal or elevated CO<sub>2</sub> levels) in the effect of O<sub>2</sub> therapy in promoting the reduction in carbon monoxide body burden. Studies utilizing laboratory animals (dogs and sheep) exposed to carbon monoxide have demonstrated that the carbon monoxide washout time can be significantly reduced if the subjects receive a mixture of oxygen and carbon dioxide gas designed to maintain normal PCO<sub>2</sub> values (Fisher et al. 1999; Kreck et al. 2001). This outcome has been replicated in experimental trials using volunteers exposed to carbon monoxide concentrations producing COHb levels of 10–12%. As compared to the subjects treated with hyperoxia alone, those treated with hyperoxia and carbon dioxide that maintained isocapnia showed accelerated clearance of carbon monoxide (Rucker et al. 2002; Takeuchi et al. 2000). These studies identify a potential means to enhance the treatment of acute carbon monoxide poisoning.

#### 3.11.2 Reducing Body Burden

Absorbed carbon monoxide binds to and displaces oxygen (O<sub>2</sub>) from blood Hb and other heme proteins (e.g., muscle myoglobin). Nearly all of carbon monoxide in the body exists as complexes with Hb or other heme proteins. Binding to heme proteins limits diffusion of carbon monoxide from tissues to blood, and elimination of carbon monoxide from blood to exhaled air. At ambient oxygen pressures of approximately 0.2 atmospheres, elimination of carbon monoxide occurs with a half-time of approximately 100–300 minutes (Bruce and Bruce 2006; Peterson and Stewart 1970). However, the elimination rate can be increased substantially by administering 100% oxygen (Tomaszewski 2006; Weaver et al. 2000). As noted above, the general recommendation for reducing the body burden of carbon monoxide is administration of 100% oxygen through a face mask that does not allow rebreathing of expired air, at ambient pressure, and/or hyperbaric oxygen therapy. The longer time period required for the elimination of carbon monoxide in the fetus (Hill et al. 1977; Longo 1977) has prompted a concern about the premature cessation of oxygen therapy to a pregnant woman based upon her COHb level.

### 3. HEALTH EFFECTS

#### 3.11.3 Interfering with the Mechanism of Action for Toxic Effects

Binding of carbon monoxide to Hb impairs transfer of oxygen to tissues and can produce tissue hypoxia. Tissues that have a high oxygen utilization rate, including brain and heart, are particularly vulnerable to hypoxia. Carbon monoxide-induced hypoxia is treated by administering 100% oxygen, which increases the dissolved concentration of oxygen in blood, facilitates displacement of carbon monoxide from Hb, and increases the rate of elimination of carbon monoxide from the body. Hyperbaric oxygen therapy may further reduce the risk of brain and heart injury from hypoxia. Intravenous fluids and inotropic agents may be administered to alleviate hypotension and myocardial depression (Lavonas 2007; Tomaszewski 2006).

#### 3.12 ADEQUACY OF THE DATABASE

Section 104(I)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of carbon monoxide is available. Where adequate information is not available, ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of carbon monoxide.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

##### 3.12.1 Existing Information on Health Effects of Carbon Monoxide

The existing data on health effects of inhalation, oral, and dermal exposure of humans and animals to carbon monoxide] are summarized in Figure 3-12. The purpose of this figure is to illustrate the existing information concerning the health effects of carbon monoxide. Each dot in the figure indicates that one or more studies provide information associated with that particular effect. The dot does not necessarily imply anything about the quality of the study or studies, nor should missing information in this figure be interpreted as a “data need”. A data need, as defined in ATSDR’s *Decision Guide for Identifying Substance-Specific Data Needs Related to Toxicological Profiles* (Agency for Toxic Substances and

3. HEALTH EFFECTS

**Figure 3-12. Existing Information on Health Effects of Carbon Monoxide**

	Systemic									
	Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation	●	●		●	●	●		●		●
Oral										
Dermal										

**Human**

	Systemic									
	Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation	●	●	●	●	●	●	●	●	●	
Oral										
Dermal										

**Animal**

● Existing Studies

### 3. HEALTH EFFECTS

Disease Registry 1989), is substance-specific information necessary to conduct comprehensive public health assessments. Generally, ATSDR defines a data gap more broadly as any substance-specific information missing from the scientific literature. Human clinical studies of acute exposures have investigated cardiovascular, respiratory, and neurobehavioral end points. Human epidemiology studies have examined possible associations between ambient air carbon monoxide concentrations and various health end points, including mortality, cardiovascular, respiratory, blood biomarkers of coagulation and inflammation, and developmental (pre-term birth, birth weight, congenital anomalies, neonatal and infant mortality). Studies conducted in animals have included acute, intermediate, and chronic exposures that have largely focused on cardiovascular, respiratory, and reproductive (e.g., fetal mortality) end points. Bioassays of carcinogenicity of carbon monoxide have not been reported. Carbon monoxide has been assessed for genotoxicity in an *in vivo* mouse model of micronuclei formation and sister chromatid exchange.

#### 3.12.2 Identification of Data Needs

**Acute-Duration Exposure.** Most information on effects of acute exposures to carbon monoxide derive from clinical case studies of carbon monoxide poisoning, in which blood COHb levels were >20%. Although these studies provide information on effects resulting from severe carbon monoxide-induced toxicity, they do not provide information on low dose-response relationships for carbon monoxide. Experimental clinical studies providing low dose-response information have focused on evaluating organ systems that are expected to be vulnerable to or contribute to hypoxia (e.g., brain, heart, respiratory tract). Effects of acute exposures to carbon monoxide on other organ systems have been studied far less thoroughly. Studies of low-dose carbon monoxide exposure in humans or animals have not identified NOAELs for sensitive organ systems and have not provided data that would enable low-dose response modeling (e.g., benchmark dose). Such studies would be beneficial for analyses in support of deriving MRLs.

Experimental clinical studies have examined cardiovascular effects in humans, including a major vulnerable subpopulation, people with ongoing heart disease; however, these studies have not identified NOAELs in cardiovascular disease patients (Adams et al. 1988; Allred et al. 1989, 1991; Anderson et al. 1973; Kleinman et al. 1989, 1998; Leaf and Kleinman 1996b). Studies investigating the effects of acute-duration carbon monoxide exposure on hemodynamics have been conducted in several animal species, including monkeys, dogs, rats, and rabbits, under exposure conditions producing blood COHb levels ranging from 6.2 to 70% (EPA 1991, 2000). These studies have shown that brief exposure (from a few

### 3. HEALTH EFFECTS

minutes to approximately 3 hours) to carbon monoxide at concentrations of 80–20,000 ppm produces alterations in hemodynamics that are consistent with COHb-induced hypoxia and responsive compensatory mechanisms (e.g., vasodilation and increased cardiovascular output), including increased coronary blood flow, decreased myocardial O<sub>2</sub> consumption, increased heart rate, and alterations in blood flow to various vascular beds (e.g., cerebral, limb muscular).

A few clinical experimental studies have evaluated adverse respiratory effects of carbon monoxide exposure in relatively small numbers of healthy subjects (Chevalier et al. 1966; Fisher et al. 1969; Koike et al. 1991; Ren et al. 2001; Vesely et al. 2004). Results of these studies indicate that brief exposure to relatively higher levels (e.g.,  $\geq 500$  ppm) of carbon monoxide may decrease ventilatory performance. No studies were identified that examined effects of carbon monoxide on respiratory function in patients with underlying respiratory diseases, who would be expected to have a higher sensitivity to hypoxic effects of carbon monoxide. Studies of dose-response relationships of respiratory function at low levels of carbon monoxide exposure (e.g., COHb < 20%) in patients with respiratory disease (e.g., asthma) would be beneficial for quantifying vulnerability of this subpopulation to carbon monoxide toxicity.

**Intermediate-Duration Exposure.** Controlled experimental studies in humans of effects of intermediate-duration exposures to carbon monoxide were not identified. Intermediate-duration studies have examined cardiovascular and respiratory effects of exposure to carbon monoxide in various animal models, including monkeys, dogs, and rodents (EPA 2000). The observation of cardiomegaly in rats exposed chronically to carbon monoxide (Sørhaug et al. 2006) suggests that prolonged exposure to carbon monoxide may produce effects on the heart that might not be observed with acute exposures. Additional intermediate-duration exposure studies in animals that provide dose-response relationships for cardiovascular effects and other end points (e.g., respiratory, neurological) would be beneficial for analyses in support of deriving MRLs.

**Chronic-Duration Exposure and Cancer.** Epidemiological studies have examined health outcomes in the context of chronic exposures, or acute variations in exposure concentrations that occur during chronic exposures. Outcomes that have been assessed include mortality, including cancer mortality, cardiovascular effects, respiratory effects, effects on blood biomarkers of coagulation and inflammation, and developmental effects (pre-term birth, birth weight, congenital anomalies, neonatal and infant mortality). A chronic study conducted in rats found cardiomegaly, which may have represented long-term adaptive changes to carbon monoxide-induced hypoxia (Sørhaug et al. 2006). This finding suggests that prolonged exposure to carbon monoxide may produce effects on the heart that might not be

## 3. HEALTH EFFECTS

evident from acute- or intermediate-duration studies. Additional chronic-duration exposure studies in animals would that provide dose-response relationships for cardiovascular effects and other end points (e.g., respiratory, neurological) would be beneficial for analyses in support of deriving MRLs.

Carbon monoxide has not been assessed for carcinogenicity using animal models. Cancer bioassays in animals would provide a more confident basis for establishing whether or not carbon monoxide has carcinogenic potential in humans.

**Genotoxicity.** In a bacterial replication assay with *Escherichia coli*, exposure to carbon monoxide-inhibited DNA synthesis (Cairns and Denhardt 1968). Gestational exposure of mice to carbon monoxide increased of micronuclei formation and sister chromatid exchange in dams (bone marrow) and fetuses (blood) (Kwak et al. 1986). Additional studies conducted on the mechanisms of genotoxicity in mammalian cells would provide a more confident basis for establishing whether or not carbon monoxide has carcinogenic potential in humans.

**Reproductive Toxicity.** Pregnancy outcomes, including fetal death in humans has been reported in cases of maternal carbon monoxide poisoning during pregnancy (Brown et al. 1992; Caravati et al. 1988; Cramer 1982; Elkharrat et al. 1991; Farrow et al. 1990; Greingor et al. 2001; Hollander et al. 1987; Margulies 1986; Norman and Halton 1990; Silverman and Montano 1997). Epidemiological studies have examined possible relationships between exposures to ambient air concentrations of carbon monoxide and fetal mortality (see Table 3-8). Few studies of reproductive effects in animals were identified, and these have been limited to assessments of effects of maternal exposure during pregnancy on fetal mortality (Astrup et al. 1972; Fechter and Annau 1977; Penney et al. 1983; Stupfel and Bouley 1970). The available data do not allow a confident determination of the relative sensitivity of reproductive systems to carbon monoxide. Greater confidence would be achieved from studies that examine more comprehensive end points of reproductive toxicity in females and males, in association with exposures prior to mating and pregnancy, and during pregnancy.

**Developmental Toxicity.** Epidemiological studies have examined developmental outcomes in the context of chronic exposures, or acute variations in exposure concentrations that occur during chronic exposures (see Table 3-8). Outcomes that have been assessed include pre-term birth, birth weight, congenital anomalies, neonatal and infant mortality. Studies conducted in animals provide evidence of adverse developmental effects of gestational and early postnatal carbon monoxide exposure, including decreased fetal weight, adverse central nervous system development, altered peripheral nervous system

## 3. HEALTH EFFECTS

development, cardiac effects, altered sexual behavior, immunological effects, and hematological effects. In addition, some studies showed that developmental effects persisted beyond the postnatal period (see Table 3-9). Two studies found functional impairments in the auditory system of rats in association with exposures to  $\geq 12$  ppm carbon monoxide during the neonatal periods or during gestation (Stockard-Sullivan et al. 2003; Webber et al. 2003). These values are the lowest LOAELs for external exposure that have been reported and, as a result, have a substantial impact on analyses conducted to derive MRLs. Additional studies to establish greater confidence in the dose-response relationship for neurodevelopmental effects on the auditory system would be beneficial for establishing MRLs for carbon monoxide based on this end point. Studies conducted to understand the mechanism for this effect would provide additional information about the relevance of these effects to humans exposed at similar levels.

**Immunotoxicity.** Epidemiological studies have examined effects on blood biomarkers of inflammation. Biomarkers examined have included been inflammation markers, CRP, SAA, and WBC; cell adhesion markers, E-selectin, vWF, ICAM-1 (Baccarelli et al. 2007; Liao et al. 2005; Pekkanen et al. 2000; Ruckerl et al. 2006, 2007; Steinvil et al. 2008). A study conducted in guinea pigs examined effects of carbon monoxide exposure on plaque forming cells in spleen (Snella and Rylander 1979). Effects of gestational exposure on the developing immune system have been studied in rats (Giustino et al. 1993, 1994). Studies conducted in cell culture and *in vivo* models of sepsis have examined effects of endogenous and exogenous carbon monoxide on inflammatory responses and cytokine and signaling pathways involved in inflammation (Ryter et al. 2006). These studies suggest a potential for carbon monoxide to exert effects on the immune system. Additional studies to establish modes of action and dose-response relationships would be beneficial for determining the sensitivity of the immune system to carbon monoxide, relative to other targets of toxicity, and for establishing MRLs.

**Neurotoxicity.** Experimental clinical studies (Benignus et al. 1994), clinical case studies (Chambers et al. 2008; Dolan 1985; Ernst and Zibrak 1998; Hopkins et al. 2006; Kao and Nañagas 2006; Lo et al. 2007; Parkinson et al. 2002; Raub and Benignus 2002), and studies conducted in animals (Benignus et al. 1994; Chambers et al. 2008; Chen and Fechter 1999; Dolan 1985; Ernst and Zibrak 1998; Fechter et al. 1988; Hopkins et al. 2006; Kao and Nañagas 2006; Lo et al. 2007; Parkinson et al. 2002; Raub and Benignus 2002; Young et al. 1987) have examined effects of acute carbon monoxide exposures on behavior cognitive function. Although these studies have provided strong evidence for impairments in association with exposures that result in blood COHb levels  $>20\%$ , dose-response relationships for lower levels remain controversial (Benignus 1994; Benignus et al. 1990; Raub and Benignus 2002). Further research on dose-response relationships of neurobehavioral and neurosensory effects at low levels of

## 3. HEALTH EFFECTS

carbon monoxide exposure (e.g., COHb<20%) utilizing double blind designs would be beneficial for analyses in support of deriving MRLs.

**Epidemiological and Human Dosimetry Studies.** Epidemiological studies have examined health outcomes in the context of chronic exposures, or acute variations in exposure concentrations that occur during chronic exposures (see Tables 3-3, 3-6, 3-7, and 3-8). Outcomes that have been assessed include mortality, including cancer mortality, cardiovascular effects, respiratory effects, blood biomarkers of coagulation and inflammation, and developmental effects (pre-term birth, birth weight, congenital anomalies, neonatal and infant mortality). Human dosimetry studies have examined relationships between carbon monoxide exposure levels and durations and blood COHb levels in resting subjects, during exercise, and at various altitudes (see Tables 3-4 and 3-5). Studies of elimination kinetics carbon monoxide in humans have explored effects of exercise, age, gender, and oxygen therapy on elimination half-time (Bruce and Bruce 2006; Joumard et al. 1991; Landaw 1973; Levasseur et al. 1996; Peterson and Stewart 1970; Shimazu et al. 2000; Weaver et al. 2000).

**Biomarkers of Exposure and Effect.**

*Exposure.* Measurement of blood COHb is the principal biomarker for identifying exposure to carbon monoxide. Several pharmacokinetics models have been developed that can be used to predict blood COHb levels that would correspond to a given exposure concentration and duration, in a typical adult and in the fetus resulting from maternal exposures. These models can be used to reconstruct possible carbon monoxide exposure scenarios (e.g., concentrations, duration, and gap between measurement of COHb and cessation of exposure) that could result in a given measured COHb level.

*Effect.* Although blood COHb reflects current carbon monoxide body burden, measurement of blood COHb has not been shown to be a reliable predictor of severity of acute toxicity (Hampson and Hauff 2008). Therefore, diagnoses and characterization of toxic effects of carbon monoxide relies on both the measurement of blood COHb and the assessment of signs and symptoms of carbon monoxide toxicity. Studies conducted to identify biomarkers of nonhypoxic mechanisms of carbon monoxide (those unrelated to blood COHb levels) would be beneficial for internal dosimetry in epidemiological and clinical experimental studies, for development of PBPK/PD models of nonhypoxic effects of carbon monoxide, and for improving diagnosis of carbon monoxide poisoning.

## 3. HEALTH EFFECTS

**Absorption, Distribution, Metabolism, and Excretion.** Pharmacokinetics of carbon monoxide has been extensively studied in humans and in animal models, including non-human primates (see Section 3.4). Studies conducted in humans have provided information on absorption, metabolism, distribution, and elimination kinetics, as well as physiological factors (e.g., age, gender) and environmental factors (e.g., altitude, exercise) that affect carbon monoxide kinetics. These studies have provided the bases for pharmacokinetic models that simulate carbon monoxide absorption and elimination kinetics, and blood COHb and muscle COMb levels in humans (see Section 3.4.5). Several studies have provided estimates of rates of whole-body endogenous production of carbon monoxide in adult males and females, during and following pregnancy (Coburn et al. 1963; Delivoria-Papadopoulos et al. 1974; Longo 1977; Mercke and Lundh 1976). However, measurement of local tissue production rates would also be valuable for studying mechanisms by which endogenously produced carbon monoxide modulates cellular metabolism and cell signaling pathways, as well as the relative contributions of exogenous carbon monoxide and endogenously produced carbon monoxide contribute to these processes.

**Comparative Toxicokinetics.** Major physiological and biochemical determinants of blood and skeletal muscle carbon monoxide kinetics that would be expected to contribute to interspecies differences in carbon monoxide toxicokinetics have been identified. This information provides a basis for extrapolating pharmacokinetics models across mammalian species, for which physiological (e.g., ventilation rates, blood Hb levels) and chemical-specific parameters are known (e.g., binding affinities and capacities of Hb and myoglobin, long carbon monoxide diffusion capacity). Models developed to simulate carbon monoxide kinetics in humans have been adapted for applications to animal models (e.g., rats; Benignus and Annau 1994) and to simulate maternal-fetal transfer of carbon monoxide during pregnancy in sheep (Longo and Hill 1977). Adaptation of these models to a wider range of species would be facilitated by studies that provide estimates of model parameters in these species (e.g., Haldane coefficient, binding coefficients of Hb and Mb, and the lung carbon monoxide diffusing capacity).

**Methods for Reducing Toxic Effects.** Efficacies of normobaric and hyperbaric oxygen therapy for hastening the elimination of carbon monoxide from the body and for treating carbon monoxide-induced hypoxia have been studied. Although hyperbaric oxygen therapy is considered to be a therapeutic option, its efficacy, over normobaric therapy, remains controversial (Wolf et al. 2008). Further research regarding benefits and limitations of hyperbaric oxygen therapy would improve the clinical management of treatment of carbon monoxide poisoning.

### 3. HEALTH EFFECTS

**Children's Susceptibility.** Data needs relating to both prenatal and childhood exposures, and developmental effects expressed either prenatally or during childhood, are discussed in detail in the Developmental Toxicity subsection above.

Child health data needs relating to exposure are discussed in Section 6.8.1, Identification of Data Needs: Exposures of Children.

**Mechanisms of Toxicity.** Although hypoxic mechanisms of action of carbon monoxide are well established (i.e., those related to formation of COHb), a better understanding is needed of nonhypoxic mechanisms, their contribution to observed adverse health effects of carbon monoxide, and related dose-response relationships.

#### 3.12.3 Ongoing Studies

Ongoing studies pertinent to information of health effects and/or mechanisms of action of carbon monoxide are listed in Table 3-18.

## 3. HEALTH EFFECTS

**Table 3-18. Ongoing Studies on the Health Effects of Carbon Monoxide**

Investigator	Affiliation	Research description	Sponsor
Abraham NG	New York Medical College	Heme oxygenase regulation of eicosanoid biosynthesis	National Institute of Diabetes and Digestive and Kidney Diseases
Bauer AJ	University of Pittsburgh at Pittsburgh Office of Research	Protective mechanisms of carbon monoxide in intestinal inflammation	National Institute of Diabetes and Digestive and Kidney Diseases
Bauer PM	University of Pittsburgh at Pittsburgh Office of Research	Role of caveolin-1 and eNOS in mediating the therapeutic effects of carbon monoxide in polycyclic aromatic hydrocarbons	National Heart, Lung, and Blood Institute
Choi AM	Brigham and Women's Hospital Research Administration	Role of heme oxygenase in hyperoxic lung injury	National Heart, Lung, and Blood Institute
Durante W	University of Missouri-Columbia	Carbon monoxide and vascular cell function	National Heart, Lung, and Blood Institute
Fallon MB	University of Texas	Mediators of pulmonary vasodilatation in liver disease	National Institute of Diabetes and Digestive and Kidney Diseases
Fowler AA	Virginia Commonwealth University	Role of hypoxia inducible factor-1 in inflammation	National Heart, Lung, and Blood Institute
Kibbe MR	Northwestern University	Nitric oxide eluting therapies for vascular surgery	National Heart, Lung, and Blood Institute
Lefer DJ	Yeshiva University	Mechanism of myocardial reperfusion injury in diabetes	National Heart, Lung, and Blood Institute
Leffler CW	University of Tennessee Health Science Center	Carbon monoxide in newborn cerebral circulation	National Heart, Lung, and Blood Institute
Leffler CQ	University of Tennessee Health Science Center	Control of neonatal circulation	National Heart, Lung, and Blood Institute
Lindahl PA	Texas A&M University System	Bioinorganic chemistry of carbon monoxide dehydrogenase	National Institute of General Medical Sciences
Machado RF	Clinical Center	Effects of inhaled carbon monoxide on human lung inflammation	Not applicable
Mapes JP	Rules-Based Medicine, Inc.	Biomarker profiles for carbon monoxide poisoning	National Institute of Environmental Health Sciences
McCurry KR	University of Pittsburgh at Pittsburgh Office of Research	Cytoprotective effect of carbon monoxide in lung ischemia/reperfusion	National Heart, Lung, and Blood Institute
Morse ED	University of Pittsburgh at Pittsburgh Office of Research	Inhibitor of differentiation-1 mediates antifibrotic effects of carbon monoxide	National Heart, Lung, and Blood Institute
Murase N	University of Pittsburgh at Pittsburgh Office of Research	Protective role of carbon monoxide in hepatic I/R injury	National Institute of Diabetes and Digestive and Kidney Diseases

## 3. HEALTH EFFECTS

**Table 3-18. Ongoing Studies on the Health Effects of Carbon Monoxide**

Investigator	Affiliation	Research description	Sponsor
O'Neill MS	University of Michigan at Ann Arbor	Air pollution, inflammation, and preterm birth: a mechanistic study in Mexico City	National Institute of Environmental Health Sciences
Parfenova H	University of Tennessee Health Science Center	Heme oxygenase and cerebral vascular injury	National Institute of Neurological Disorders and Stroke
Raman CS	University of Texas Health Science Center, Houston	Structural biology of gaseous messenger signaling	National Institute of General Medical Sciences
Roberts GP	University of Wisconsin Madison	Sensing mechanisms for carbon monoxide and other small molecules	National Institute of General Medical Sciences
Tolbert PE	Emory University	Air pollution and birth defects in Atlanta, 1968–2002	National Institute of Environmental Health Sciences
Tulis DA	North Carolina Central University	NO-independent cGMP regulation of vascular remodeling	National Heart, Lung, and Blood Institute
Vitali SH	Children's Hospital Boston	Hypoxic inflammation, pulmonary hypertension, and HO-1	National Heart, Lung, and Blood Institute
Williams MA	Swedish Medical Center, First Hill	Ambient air pollution, preeclampsia, and preterm delivery	National Institute of Environmental Health Sciences
Xuan U	University of Louisville	Delayed cardioprotection induced by NO and carbon monoxide	National Heart, Lung, and Blood Institute

Source: FEDRIP 2009

3. HEALTH EFFECTS

This page is intentionally blank.

## 4. CHEMICAL AND PHYSICAL INFORMATION

### 4.1 CHEMICAL IDENTITY

The chemical identity of carbon monoxide is shown in Table 4-1.

### 4.2 PHYSICAL AND CHEMICAL PROPERTIES

Carbon monoxide is a highly poisonous, odorless, colorless, and tasteless gas. It is very flammable in air over a wide range of concentrations (George 2001) and burns in air with a bright blue flame (O'Neil et al. 2006). It becomes a liquid at 81.62 K (-191.53 °C) and is insoluble in water above 70 °C (George 2001). The physical and chemical properties of carbon monoxide are shown in Table 4-2.

## 4. CHEMICAL AND PHYSICAL INFORMATION

**Table 4-1. Chemical Identity of Carbon Monoxide**

Characteristic	Carbon monoxide <sup>a</sup>
Synonym(s)	Carbon oxide, flue gas, monoxidized
Registered trade name(s)	
Chemical formula	CO <sup>b</sup>
Chemical structure	:C≡O:
Identification numbers:	
CAS registry	630-08-0 <sup>b</sup>
NIOSH RTECS	FG350000
EPA hazardous waste	No data
OHM/TADS	No data
DOT/UN/NA/IMDG shipping	1016c; 9202 <sup>d,e</sup>
HSDB	903
NCI	No data

<sup>a</sup>All information obtained from HSDB 2009, except where noted.

<sup>b</sup>O'Neil 2006

<sup>c</sup>Compressed

<sup>d</sup>NIOSH 2005

<sup>e</sup>Cryogenic liquid

CAS = Chemical Abstracts Service; DOT/UN/NA/IMDG = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances

## 4. CHEMICAL AND PHYSICAL INFORMATION

**Table 4-2. Physical and Chemical Properties of Carbon Monoxide**

Property	Carbon monoxide <sup>a</sup>
Molecular weight	28.01
Color	Colorless
Physical state	Gas
Melting point	-205 °C
Boiling point	-191.5 °C
Density at 25°C	1.145 g/L at 25 °C and 1 atm <sup>3b</sup>
Odor	Odorless
Odor threshold:	
Water	No data
Air	No data
Solubility:	
Water at 20 °C	2.3 mL/100 mL
Organic solvents	Appreciably soluble in ethyl acetate, chloroform, acetic acid; freely absorbed by a concentrated solution of cuprous chloride in hydrochloric acid or ammonium hydroxide; solubility in methanol and ethanol about 7 times as great as in water; soluble in benzene
Partition coefficients:	
Log K <sub>ow</sub>	Not applicable
Log K <sub>oc</sub>	Not applicable
Henry's law constant at 25 °C	57,978.5 atm/mol fraction (~1.04 atm·m <sup>3</sup> /mole) <sup>c</sup>
Autoignition temperature	605 °C <sup>d</sup>
Flashpoint	Flammable gas <sup>d</sup>
Flammability limits	Upper limit 74.2%; lower limit 12.5% <sup>e</sup>
Conversion factors	1 ppm = 1.16 mg/m <sup>3</sup> at 20 °C and 1 atm <sup>f</sup>
Explosive limits	No data

<sup>a</sup>All information obtained from HSDB 2009, except where noted.

<sup>c</sup>Yaws et al. 1999

<sup>b</sup>Lide 2008

<sup>d</sup>IPCS 2008

<sup>e</sup>George 2001

<sup>f</sup>Verschueren 2001

4. CHEMICAL AND PHYSICAL INFORMATION

This page is intentionally blank.

## 5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

### 5.1 PRODUCTION

Carbon monoxide is produced by the incomplete combustion of carbon in liquid, solid, and gaseous fuels (George 2001). Commercially, it is produced on an industrial scale by the partial oxidation of hydrocarbon gases from natural gas or by the gasification of coal and coke. Laboratory-scale production of carbon monoxide is accomplished by heating calcium carbonate with zinc dust (O'Neil et al. 2006). Carbon monoxide is also obtained from the dehydration of formic acid (Lewis 2007). The majority of carbon monoxide produced is used immediately downstream and at the plant site for chemical synthesis or steel manufacturing; consequently, quantitative production volumes are not available (George 2001).

Carbon monoxide is a co-product along with hydrogen in syn gas (synthetic gas) production, which are then separated and purified by pressure swing adsorption and/or cryogenic distillation.

Carbon monoxide is a major air pollutant (Lewis 2007). It is a byproduct of highway vehicle exhaust, of industrial processes and fuel combustion in boilers and incinerators, and of household appliances fueled with gas, oil, kerosene, or wood. The largest contribution comes from highway motor vehicles. Carbon monoxide exceeds all other atmospheric pollutants combined with the exception of CO<sub>2</sub> (George 2001).

The major facility within the United States that manufactures or processes carbon monoxide (produced in commercial quantities exceeding 5,000 pounds or \$10,000 in value annually) is Air Liquide America L.P. in Freeport, Texas (SRI 2008).

### 5.2 IMPORT/EXPORT

No data were located on the import and export of carbon monoxide.

### 5.3 USE

Carbon monoxide is used by the chemical industry for the synthesis of many compounds such as acetic anhydride, polycarbonates, acetic acid, polyketones, etc (George 2001). It finds application as a reducing agent in metallurgical operations, specifically the Mond process for the recovery of nickel, in the manufacture of metal carbonyls, and in organic synthesis, especially for the Fischer-Tropsch process for

## 5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

petroleum-related products and in the oxo reaction (O'Neil et al. 2006). Carbon monoxide is also used in the manufacture of zinc white pigments (Lewis 2007).

**5.4. DISPOSAL**

EPA considers a waste to be hazardous if it exhibits any of the following characteristics: ignitability, corrosivity, reactivity, or toxicity as defined in 40 CFR 261.21–261.24. Under the Resource Conservation and Recovery Act (RCRA) (40 USC 6901 et seq.). EPA has specifically listed many chemical wastes as hazardous. Although carbon monoxide is not specifically listed as a hazardous waste under RCRA, EPA requires employers to treat waste as hazardous if it exhibits any of the characteristics discussed above (OSHA 2000).

## 6. POTENTIAL FOR HUMAN EXPOSURE


### 6.1 OVERVIEW

Carbon monoxide has been identified in at least 12 of the 1,699 hazardous waste sites that have been proposed for inclusion on the EPA National Priorities List (NPL) (HazDat 2007). However, the number of sites evaluated for carbon monoxide is not known. The frequency of these sites can be seen in Figure 6-1.

Carbon monoxide is a colorless, odorless, and tasteless gas that is ubiquitous in the atmosphere (George 2001). It arises from both natural and anthropogenic sources. It is produced as a primary pollutant during the incomplete combustion of fossil fuels and biomass, including wildfires and controlled burns. Carbon monoxide is also produced indirectly from the photochemical oxidation of methane and other VOCs in the atmosphere. Vegetation can emit carbon monoxide directly into the atmosphere as a metabolic byproduct and the photooxidation of organic matter in surface waters (lakes, streams, rivers, oceans) and soil surfaces also results in the formation of carbon monoxide. Volcanic activity is another natural source of carbon monoxide in the atmosphere. Carbon monoxide is also produced endogenously in humans during the normal catabolism of Hb. The amount of anthropogenic carbon monoxide emissions versus that from natural sources is difficult to quantify since both sources of carbon monoxide vary over time. The EPA estimated that 95% of all carbon monoxide emissions in 2002 arose as a result of human activity with biogenic emissions accounting for only 5% of the total emissions (EPA 2008). However, a recent analysis of the carbon monoxide budget over North America concluded that during the summer months, when natural sources of carbon monoxide precursors such as biogenic VOCs and forest fires are high, natural sources of carbon monoxide in the atmosphere are far greater than anthropogenic contributions (Miller et al. 2008). The vast majority of anthropogenic carbon monoxide emissions arise from gasoline-powered automobile usage (EPA 2000). Data from the EPA National Emissions Inventory (NEI) suggest that over 50% of the total point and non-point emissions of carbon monoxide are historically associated with on-road automobile use (EPA 2008).

The background levels of carbon monoxide have changed significantly in the past several decades. Concentrations have decreased appreciably due to reduced emissions from automobiles as a consequence of advancements in automotive design. The development of catalytic converters for passenger vehicles, beginning in the 1970s, resulted in a substantial decrease in carbon monoxide emissions, despite large increases in miles traveled (George 2001). In the early 1980s, automakers equipped vehicles with more sophisticated catalytic converters and added on-board computers and O<sub>2</sub> sensors in order to help optimize

Figure 6-1. Frequency of NPL Sites with Carbon Monoxide Contamination



Derived from HazDat 2007

## 6. POTENTIAL FOR HUMAN EXPOSURE

the efficiency of the converter. The end result is that modern passenger automobiles emit about 90% less carbon monoxide over their lifetimes as compared to vehicles designed in previous decades (George 2001).

The annual average outdoor carbon monoxide concentrations are roughly 0.12 ppmv in the Northern Hemisphere and about 0.04 ppmv in the Southern Hemisphere (EPA 2000). These levels are variable throughout the course of the year, with seasonal maximum levels occurring during late winter in both hemispheres when inversion conditions (in which air pollutants are trapped near the ground beneath a layer of warm air) are more frequent. Minimum levels are generally observed during late summer. Carbon monoxide concentrations are reported to range from a minimum of about 0.03 ppmv during summer in the Southern Hemisphere to about 0.20 ppmv at high latitudes in the Northern Hemisphere during winter (EPA 2000). Urban locations with high automobile usage or a high volume of stationary emission sources, such as refineries or power plants, typically have greater atmospheric levels of carbon monoxide as compared to rural or remote sites. In metropolitan areas in the United States, as much as 95% of all carbon monoxide emissions result from on-road vehicle exhaust (EPA 2008). The majority of these on-road emissions are derived from gasoline-powered vehicles since diesel vehicles emit less carbon monoxide. There is also a diurnal pattern of atmospheric carbon monoxide concentrations in urban areas, with the highest levels occurring during hours with heavy vehicular usage (rush hours) and the lowest levels occurring at times that correlate with lower commuting activity (Campbell et al. 1995; EPA 2000). Maximum carbon monoxide levels frequently exceed 5 ppmv at these locations during the high commute hours.

Carbon monoxide levels in indoor air are greatly dependent upon the presence of combustion-based appliances and whether occupants smoke tobacco products. Unvented kerosene and gas space heaters; leaking chimneys and furnaces; back-drafting from furnaces, gas water heaters, wood stoves, and fireplaces; gas stoves, generators, and other gasoline-powered equipment; automobile exhaust from attached garages; and tobacco smoke all contribute to indoor air levels of carbon monoxide (EPA 2009a). Average levels in homes without gas stoves vary from 0.5 to 5 ppmv. Levels near properly adjusted gas stoves are often 5–15 ppmv and those near poorly adjusted stoves may be  $\geq 30$  ppmv (EPA 2009a).

The primary degradation pathway of carbon monoxide in the environment occurs through its reaction with photochemically-produced hydroxyl radicals. (It should be noted that the production of hydroxyl radicals requires UV radiation that does not penetrate windows and therefore, photooxidation is likely negligible indoors.) In addition, soils and coastal waters may also act as a sink for carbon monoxide since

## 6. POTENTIAL FOR HUMAN EXPOSURE

various forms of microorganisms are capable of utilizing carbon monoxide as an energy source (Tolli et al. 2006).

Exposure of the general population to carbon monoxide occurs through inhalation of outdoor and indoor air. Populations living in urban areas with heavy vehicular traffic or stationary sources such as petroleum refineries, gas and coal burning power plants, petrochemical plants, and coke oven plants are more likely to be exposed to higher levels of carbon monoxide from ambient outdoor air. Employees in these refineries and plants and workers who are subject to high levels of vehicular exhaust (such as taxi cab drivers and toll booth workers) may be occupationally exposed to higher levels of carbon monoxide. Members of the public who smoke or work in smoke-filled environments such as restaurants, bars, and casinos where smoking is allowed are exposed to higher levels of carbon monoxide than members of the population who do not smoke and are not frequently exposed to second-hand tobacco smoke.

### 6.2 RELEASES TO THE ENVIRONMENT

Carbon monoxide releases are not required to be reported for inclusion in the Toxics Release Inventory (TRI) (EPA 2005).

#### 6.2.1 Air

Title I of the Clean Air Act of 1970 establishes carbon monoxide as one of six criteria pollutants and sets national air quality standards for carbon monoxide and the other criteria pollutants (EPA 2000). Two databases developed by the EPA are particularly useful for monitoring carbon monoxide levels and emissions throughout the United States. EPA's National Emission Inventory (NEI) database contains detailed information about sources that emit criteria air pollutants and their precursors, and hazardous air pollutants for the 50 United States, Washington DC, Puerto Rico, and the U.S. Virgin Islands. The Air Quality System (AQS) database is EPA's repository of criteria air pollutant monitoring data since the 1970s. Table 6-1 supplies data for carbon monoxide emissions from 1970 to 2008 from the NEI database for 13 major emission categories. Detailed carbon monoxide emissions from these sources for individual years are available as zipped Microsoft Access database files that may be accessed directly from the EPA website; however, these data are subject to occasional revisions by EPA as emission estimates for a specific time period are amended. As indicated in Table 6-1, on-road vehicle use has historically accounted for the largest percentage of emissions in the United States as compared to the other emission sources; however, the quantity of carbon monoxide emitted from automobiles has been declining significantly over the past 4 decades due to the use of emission control devices and catalytic converters.

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-1. Carbon Monoxide Emissions (Thousands of Short Tons) from 1970 to 2008**

Year <sup>b</sup>	Emissions by category <sup>a</sup>													Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	
1970	237	770	3,625	3,397	3,644	2,179	620	NA	NA	7,059	163,231	11,371	7,909	204,042
1975	276	763	3,441	2,204	2,496	2,211	630	NA	NA	3,230	153,555	14,329	5,263	188,398
1980	322	750	6,230	2,151	2,246	1,723	830	NA	NA	2,300	143,827	16,685	8,344	185,408
1985	291	670	7,525	1,845	2,223	462	694	2	49	1,941	134,187	19,029	7,927	176,845
1990	363	879	4,269	1,183	2,640	333	537	5	76	1,079	110,255	21,447	11,122	154,188
1991	349	920	4,587	1,127	2,571	345	548	5	28	1,116	104,980	21,934	8,618	147,128
1992	350	955	4,849	1,112	2,496	371	544	5	17	1,138	99,705	22,419	6,934	140,895
1993	363	1,043	4,181	1,093	2,536	371	594	5	51	1,248	94,431	22,904	7,082	135,902
1994	370	1,041	4,108	1,171	2,475	338	600	5	24	1,225	89,156	23,389	9,656	133,558
1995	372	1,056	4,506	1,223	2,380	348	624	6	25	1,185	83,881	23,874	7,298	126,778
1996	408	1,188	2,741	1,053	1,599	354	561	1	70	2,904	78,606	24,358	15,016	128,859
1997	423	1,162	2,742	1,071	1,710	367	582	2	71	2,948	75,849	23,668	7,316	117,911
1998	451	1,151	2,727	1,081	1,702	366	590	2	72	3,121	73,244	23,689	7,184	115,380
1999	496	1,213	3,829	350	1,255	159	571	52	163	3,019	68,708	23,316	11,410	114,541
2000	484	1,219	3,081	361	1,295	161	592	51	169	1,849	68,061	24,178	12,964	114,465
2001	485	1,253	3,088	372	1,380	162	615	50	178	1,851	63,476	24,677	8,676	106,263
2002	657	1,267	3,550	284	987	357	490	2	118	1,594	60,596	22,662	18,493	111,057
2003	652	1,229	3,477	259	934	355	504	2	114	1,580	56,579	21,999	17,364	105,078
2004	647	1,190	3,404	233	882	353	519	2	111	1,567	52,562	21,336	16,235	99,041
2005	643	1,152	3,331	208	829	351	534	2	107	1,554	48,544	20,672	15,106	93,034
2006	661	1,173	3,343	227	869	352	522	2	110	1,564	45,318	19,793	13,981	87,915
2007	680	1,195	3,352	246	908	353	511	2	113	1,574	42,092	18,915	12,856	82,801
2008	699	1,216	3,369	265	947	355	500	2	115	1,584	38,866	18,036	11,731	77,685

<sup>a</sup>Categories

1. Fuel combustion electric utilities
2. Fuel combustion industrial utilities
3. Fuel combustion other
4. Chemical allied manufacturing
5. Metal processing
6. Petroleum and related categories
7. Other industrial processes
8. Solvent utilization
9. Storage and transport
10. Waste disposal and recycling
11. Highway vehicles
12. Off-highway
13. Miscellaneous

<sup>b</sup>Emission estimates are subject to updates for subsequent revised versions of the National Emissions Inventory Database (NEI) database. These numbers may be different than values published for previous versions of the data and may also be different than values for subsequent revisions of the data as generated by the Environmental Protection Agency.

NA = not available

Source: EPA 2009h

## 6. POTENTIAL FOR HUMAN EXPOSURE

Indoor air levels of carbon monoxide are highly dependent upon the smoking habits, the types of appliances and heating units that are used in the home or building, and whether or not the home or building has an attached garage for automobiles. Carbon monoxide levels from the use of appliances will depend on several factors such as the type of fuel used, ventilation, appliance design, fuel consumption rate, use pattern, and operating condition (EPA 1991, 2000). As reported in the EPA Air Quality Criteria Reports (1991, 2000), carbon monoxide emissions from ranges, ovens, and pilot lights using natural gas were typically greater under yellow tipping flame conditions (characteristic of an improper air-fuel ratio) to blue tipping flame conditions (characteristic of properly adjusted stoves). The average emission rates (mass of carbon monoxide emitted per energy unit produced) for top burning ovens and broilers ranged from 11.9 to 87.7  $\mu\text{g}/\text{kJ}$  under blue flame conditions as opposed to 53.5–156.6  $\mu\text{g}/\text{kJ}$  for similar appliances operating under yellow tipping conditions. In separate tests using three different gas ranges deliberately adjusted to produce blue tipping or yellow tipping flames, the average emission rate ranged from 34.3 to 70.9  $\mu\text{g}/\text{kJ}$  under blue flame operation to 108.4–196.9  $\mu\text{g}/\text{kJ}$  under yellow flame conditions (EPA 1991, 2000).

Quantitative estimates have been made regarding emissions of carbon monoxide from wood-burning stoves and fireplaces under normal operating conditions (Houck et al. 2006). Carbon monoxide emission factor data were based on 277 tests on 70 fireplace models. The mean emission factor was 64.1 g carbon monoxide emitted per kg dry wood with a standard deviation of 40.7 g/kg dry wood (Houck et al. 2006). It has been reported that emission rates of carbon monoxide for well-tuned unvented gas appliances range from 0.15 to 3.2 g/hour, while rates for poorly tuned unvented ones range from 1.5 to 22 g/hour (Dutton et al. 2001).

Carbon monoxide was detected in air samples at each of the 12 current or former NPL hazardous waste sites where it was detected in some environmental media (HazDat 2007).

### 6.2.2 Water

Direct anthropogenic releases of carbon monoxide to water are not expected; however, natural processes occur that result in carbon monoxide formation in waters. The photodegradation of dissolved organic matter is primarily responsible for producing carbon monoxide in sunlit surface waters (Tolli et al. 2006).

## 6. POTENTIAL FOR HUMAN EXPOSURE

**6.2.3 Soil**

The formation of carbon monoxide in soils appears to occur by abiotic processes, such as thermal decomposition or photodecomposition of organic matter. In general, warm and moist conditions found in most soils favor carbon monoxide uptake, whereas hot and dry conditions as found in deserts and some savannas favor the release of carbon monoxide from soil to the atmosphere (EPA 2000).

Carbon monoxide was detected in a soil gas sample at one current or former NPL hazardous waste sites where it was detected in some environmental media (HazDat 2007).

**6.3 ENVIRONMENTAL FATE****6.3.1 Transport and Partitioning**

Carbon monoxide is a gas that will partition to the atmosphere and is distributed globally by the advective action of the wind. Wind direction determines the horizontal transport of carbon monoxide and what impact emissions from one location will have upon another site (EPA 1991). Carbon monoxide levels may be high around local sources, and locations downwind from the source may also be elevated. The transport of carbon monoxide in urban locations is complex and can be influenced by the geometry of street canyons, topography around roadways, and presence of noise barriers, vegetation, and buildings, as well as local meteorological factors. A field study was designed to characterize the air quality and flow of pollutants near a highway with and without noise barriers (Baldauf et al. 2008). The site also contained an open field and a residential neighborhood with mature vegetation. Time-series carbon monoxide levels were collected in areas with no barriers, a noise barrier only, and a noise barrier and vegetation. Levels of carbon monoxide were reduced as much as 50% behind the noise barrier as compared to measurements taken near the road depending upon the directionality of the wind (Baldauf et al. 2008).

In most urban areas during the winter months, there is often enhanced stability in the atmospheric boundary layer, which attenuates the vertical mixing of emissions from the ground. This effectively traps carbon monoxide at street levels during these periods. Unreacted carbon monoxide in the troposphere is slowly transported to the mesosphere and stratosphere where it reacts with atomic oxygen generated by the photodissociation of O<sub>2</sub> (Fabian et al. 1979; Pressman and Warneck 1970).

The surface waters of the world's oceans are saturated with carbon monoxide with respect to the atmosphere and are therefore a net source of atmospheric carbon monoxide. Using a mean atmospheric

## 6. POTENTIAL FOR HUMAN EXPOSURE

carbon monoxide concentration over the ocean surface of 0.13 ppmv and a surface water concentration of  $6 \times 10^{-8} \text{ cm}^3$  carbon monoxide per  $\text{cm}^3$  water, the total annual flux of carbon monoxide to the atmosphere from seawater was estimated at approximately  $4.3 \times 10^{13} \text{ g/year}$  (43 Tg) (Liss and Slater 1974). Other reported estimates based on computed carbon monoxide concentrations in surface waters were reported to range from 13 to 165 Tg/year (Ohta 1997). Differences in microbial activity in surface water account for the variations in surface water levels and the flux into the atmosphere (Conrad and Seiler 1988).

### 6.3.2 Transformation and Degradation

#### 6.3.2.1 Air

Carbon monoxide is generally stable under environmental conditions. Reactions with molecular oxygen ( $\text{O}_2$ ) or water vapor are very slow at ambient temperatures and pressure. Carbon monoxide reacts with ground-state triplet oxygen atoms,  $\text{O}(^3\text{P})$ , produced by the atmospheric photodegradation of nitrogen dioxide and ozone, or atomic oxygen, formed by the photodissociation of molecular  $\text{O}_2$  in the stratosphere, to form  $\text{CO}_2$  (NRC 1977). However, the primary degradation pathway of carbon monoxide in the troposphere is through its reaction with photochemically-produced hydroxyl radicals. This results in the formation of  $\text{CO}_2$  and atomic hydrogen, which rapidly reacts with  $\text{O}_2$  to form peroxy radicals (EPA 2000). The second-order rate constant for the gas-phase reaction of carbon monoxide with hydroxyl radicals at atmospheric pressure has been measured as  $2.4 \times 10^{-13} \text{ cm}^3/\text{molecule-second}$  (EPA 1991). This corresponds to an estimated tropospheric half-life of approximately 22–67 days, assuming a hydroxyl radical concentration of  $5.0 \times 10^5$ – $1.5 \times 10^6$  hydroxyl radicals per  $\text{cm}^3$  air. No estimates of the half-life of carbon monoxide in indoor air were located. Hydroxyl radicals are formed photochemically by sunlight; therefore, their levels in indoor air will be negligible.

#### 6.3.2.2 Water

Although oceans and other water bodies are considered a net source of carbon monoxide in the environment, evidence suggests that various microorganisms may degrade carbon monoxide in the water column (Conrad and Seiler 1988). Taxonomically diverse microorganisms isolated from surface waters collected off the New England coast readily oxidized carbon monoxide (Tolli et al. 2006). Microbial oxidation rate constants for carbon monoxide in coastal waters were reported to range from 0.01 to  $0.11 \text{ hours}^{-1}$ , corresponding to half-lives on the order of several hours (Tolli et al. 2006). These rates were reported to be approximately an order of magnitude greater than in oligotrophic environments, suggesting the presence of an active carbon monoxide oxidizing microbial community near shore.

## 6. POTENTIAL FOR HUMAN EXPOSURE

**6.3.2.3 Sediment and Soil**

Soils may act as a source or a sink for carbon monoxide depending on soil moisture, intensity of sunlight reaching the soil surface, and soil temperature (Conrad and Seiler 1985). In general, warm and moist conditions found in most soils favor uptake, whereas hot and dry conditions as found in deserts and some savannas favor the release of carbon monoxide (King 1999a). Global estimates of carbon monoxide consumption by soil microorganisms ranges from 15 to 640 Tg ( $1.5 \times 10^{13}$ – $6.4 \times 10^{14}$  g) per year (King 1999a). Carbon monoxide oxidation to CO<sub>2</sub> has been demonstrated for both aerobic and anaerobic microorganisms.

**6.4 LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT**

Reliable evaluation of the potential for human exposure to carbon monoxide depends in part on the reliability of supporting analytical data from environmental samples and biological specimens.

Concentrations of carbon monoxide in unpolluted atmospheres and in pristine surface waters are often so low as to be near the limits of current analytical methods. In reviewing data on carbon monoxide levels monitored or estimated in the environment, it should also be noted that the amount of chemical identified analytically is not necessarily equivalent to the amount that is bioavailable. The analytical methods available for monitoring carbon monoxide in a variety of environmental media are detailed in Chapter 7.

**6.4.1 Air**

The concentration of carbon monoxide in air can be represented using various concentration units. Air monitoring data for carbon monoxide is usually expressed or reported as ppmv or parts per billion by volume (ppbv). A concentration of 1 ppmv implies that for every million molecules of gas in the measured volume, one of them is a carbon monoxide molecule. In order to express these concentrations in mass units, the following conversion factors may be used: 1.00 ppmv=1.16 mg/m<sup>3</sup> and 1.00 ppbv=1.16 µg/m<sup>3</sup> at 20 °C.

Annual average outdoor carbon monoxide concentrations are about 0.12 ppmv in the Northern Hemisphere and about 0.04 ppmv in the Southern Hemisphere (EPA 2000). In general, the concentration of carbon monoxide decreases with altitude in the Northern Hemisphere, but this vertical gradient may be reversed in the Southern Hemisphere due to the transport of carbon monoxide from the Northern to the Southern Hemisphere (EPA 1991). Annual 24-hour average carbon monoxide concentrations obtained at

## 6. POTENTIAL FOR HUMAN EXPOSURE

monitoring locations at rural sites in the United States are typically about 0.2 ppmv, compared with an annual 24-hour average of 1.2 ppmv across all monitoring sites (EPA 2000). Carbon monoxide levels (1- and 8-hour maximum levels) from various cities in the United States for 2008 are provided in Table 6-2. These data are derived from measurements submitted to the EPA AQS database, which collects data from EPA, state, local, and tribal air pollution control agencies. As indicated by the data presented in the table, only one measurement exceeded the EPA 8-hour carbon monoxide level of 9 ppm and no measurements exceeded the 1-hour average level of 35 ppm at the 365 different monitoring sites. Using this nationwide network of monitoring sites, EPA estimated that there has been a 76% decrease in the ambient levels of carbon monoxide in the United States from 1980 to 2007.

Urban areas with heavy vehicular traffic congestion tend to have high levels of carbon monoxide in ambient air. These levels follow a predictable diurnal pattern, which reaches maximum concentrations at times of heavy commuting and then decreases during periods of low vehicular traffic. The apex of these profiles corresponds to the morning and evening rush hour commutes when vehicle density is at its highest. For many locations, the morning peak yields higher carbon monoxide levels than the evening peak because the height of the mixed layer is much lower during the morning, thereby inhibiting vertical mixing that helps dissipate the carbon monoxide. In the late afternoon and into early evening, increased atmospheric turbulence resulting from solar activity raises the height of the mixed layer, resulting in generally lower carbon monoxide concentrations compared with those of the morning (EPA 2000). A study conducted by the Toronto (Canada) Public Health Department examined the level of carbon monoxide at four sites in Toronto as a function of traffic density (Campbell et al. 1995). The four areas of the study were categorized as low, medium, or high in terms of traffic density. Mean hourly carbon monoxide levels ranged from 0.63 ppmv at a low traffic density site (average daily vehicle count of 2,000 vehicles) to 1.54 ppmv at a site in which the vehicular count was about 45,000 per day (Campbell et al. 1995). Chen et al. (2008) measured carbon monoxide levels at 10 roadside pollution monitoring sites in the city of Leicester, England. A diurnal carbon monoxide concentration pattern was observed in this study; however, the carbon monoxide levels during the evening rush hour period generally exceeded the levels than the morning commute. Maximum carbon monoxide levels at these sites ranged from 0.8 to 3.1 ppmv during the morning commute, while maximum levels were 1.0–4.0 ppmv during the evening rush hour (Chen et al. 2008).

High levels of carbon monoxide inside vehicles and other transportation sources that use gasoline powered engines are frequently observed. Duci et al. (2003) studied the concentration of carbon monoxide in vehicles along heavy traffic routes in the Athens, Greece. The mean carbon monoxide

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
2,169	8.1	6.7	0	3.6	3.5	0	Fairbanks	AK
2,176	6.1	6	0	4.1	3.8	0	Anchorage	AK
1,988	5.1	4.5	0	2.5	2.3	0	Not provided	AK
2,176	5.6	5.5	0	3.1	2.8	0	Anchorage	AK
2,173	5.7	4.5	0	3.3	2.2	0	Fairbanks	AK
2,173	8.3	8	0	4.8	4.7	0	Anchorage	AK
7,500	19.6	15.9	0	10.7	8.1	1	Birmingham	AL
7,563	8	6.1	0	2.3	2.2	0	Fairfield	AL
8,012	3.4	3	0	2.4	2.3	0	Birmingham	AL
8,750	6.8	2.5	0	1.8	1.5	0	North Little Rock	AR
5,049	1.5	1.4	0	1	1	0	Surprise	AZ
4,584	0.7	0.7	0	0.5	0.5	0	Not provided	AZ
7,947	3.1	3.1	0	2.5	2.4	0	Phoenix	AZ
8,720	2.2	1.8	0	1.3	1	0	Tucson	AZ
8,736	2.6	2.1	0	1.1	1.1	0	Tucson	AZ
8,696	2.9	2.5	0	1.4	1.3	0	Tucson	AZ
4,369	2.5	2.5	0	1.9	1.5	0	Tucson	AZ
8,157	1.5	1.3	0	1	0.9	0	Tucson	AZ
4,315	2	1.8	0	1.3	1.2	0	Tucson	AZ
5,027	2.4	2.3	0	1.8	1.4	0	Tempe	AZ
5,043	1.8	1.7	0	1.4	1.4	0	Chandler	AZ
5,057	3.7	3.2	0	2.2	2	0	Phoenix	AZ
8,654	3	3	0	2.7	2.4	0	Phoenix	AZ
4,933	2	2	0	1.5	1.4	0	Scottsdale	AZ
5,060	2.1	2	0	1.6	1.5	0	Glendale	AZ
5,038	2.1	2	0	1.3	1.3	0	Phoenix	AZ
5,012	1.7	1.7	0	1.4	1.3	0	Mesa	AZ
8,575	4.7	4.5	0	3.1	3	0	Phoenix	AZ
8,307	3.9	3.6	0	2.8	2.8	0	Phoenix	AZ
8,397	3.6	3.5	0	2.6	2.2	0	Phoenix	AZ
6,905	2.1	1.9	0	1.1	1	0	Lompoc	CA
6,893	1.4	1.2	0	0.6	0.6	0	Goleta	CA
6,097	1.6	1.3	0	0.7	0.6	0	Vandenberg Air Force Base	CA
6,177	3.3	3	0	2.1	1.9	0	San Jose	CA
7,046	7.6	3.7	0	1.3	1.1	0	Davenport	CA

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
6,230	2.7	2.5	0	1.9	1.7	0	Vallejo	CA
6,251	0.9	0.9	0	0.7	0.6	0	Benicia	CA
6,244	3.5	1.9	0	1.3	1.1	0	Santa Rosa	CA
5,550	2.8	2.3	0	1.8	1.7	0	Modesto	CA
6,035	1.9	1.9	0	1.2	1.2	0	Turlock	CA
6,270	2.2	2.2	0	1.4	1.4	0	Livermore	CA
6,227	2.9	2.4	0	1.6	1.4	0	Oakland	CA
6,202	1.9	1.8	0	1.3	1.3	0	Fremont (Centerville)	CA
6,175	2.1	2	0	1.5	1.5	0	Berkeley	CA
5,576	3.1	3	0	2.4	2.1	0	Chico	CA
6,240	1.5	1.5	0	1	1	0	Concord	CA
6,274	1	0.9	0	0.8	0.8	0	Bethel Island	CA
6,225	2.5	2.1	0	1.3	0.9	0	San Pablo	CA
6,266	2.8	2.2	0	1.4	1.3	0	Pittsburg	CA
5,430	2.6	2.6	0	1.8	1.7	0	Fresno	CA
5,563	3.1	2.8	0	2.3	2.1	0	Fresno	CA
5,519	1.6	1.5	0	1	1	0	Fresno	CA
4,785	2.3	2.2	0	1.5	1.3	0	Clovis	CA
5,768	1.9	1.9	0	1.6	1.4	0	Eureka	CA
5,543	8.2	7	0	5.3	4.1	0	Calexico	CA
5,262	7.3	6.8	0	3.2	3.1	0	Calexico	CA
6,270	2.5	2.3	0	1.7	1.7	0	El Centro	CA
4,139	3.5	2.7	0	2.2	2.1	0	Bakersfield	CA
5,554	2.3	2.1	0	1.4	1.3	0	Azusa	CA
5,552	2.7	2.3	0	1.8	1.7	0	West Los Angeles	CA
5,555	3	2.8	0	2.5	2	0	Burbank	CA
5,560	2.9	2.8	0	2	1.9	0	Los Angeles	CA
5,594	3.4	3.3	0	2.3	2.1	0	Reseda	CA
5,579	5.8	5.7	0	4.3	4.1	0	Lynwood	CA
5,281	2.9	2.5	0	1.9	1.8	0	Pico Rivera	CA
5,557	2.6	2.6	0	1.7	1.7	0	Pomona	CA
5,550	2.4	2.4	0	2	1.9	0	Pasadena	CA
5,567	3.3	3	0	2.5	2.4	0	Long Beach	CA
5,315	3.1	2.6	0	1.9	1.8	0	Los Angeles	CA
5,495	1.6	1.5	0	0.9	0.8	0	Santa Clarita	CA
5,538	2.2	1.7	0	0.9	0.8	0	Lancaster	CA

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
6,208	1.7	1.7	0	1	1	0	San Rafael	CA
7,224	4.5	3.9	0	3.4	3.3	0	Ukiah	CA
3,282	1.5	1.4	0	1	0.9	0	Willits	CA
7,177	2.2	1.6	0	0.9	0.8	0	Salinas	CA
6,255	3.2	3.1	0	1.8	1.6	0	Napa	CA
5,560	3.3	3	0	2.4	2.1	0	Anaheim	CA
5,571	3	2.8	0	2	2	0	Costa Mesa	CA
5,524	1.5	1.5	0	1.1	1	0	Mission Viejo	CA
5,567	4.7	4.6	0	2.5	2.4	0	La Habra	CA
5,605	4.7	4.5	0	1.9	1.9	0	Riverside	CA
5,574	1	0.9	0	0.5	0.5	0	Palm Springs	CA
5,546	2.7	2.5	0	1.9	1.7	0	Rubidoux (West Riverside)	CA
5,564	1.1	1.1	0	0.8	0.8	0	Lake Elsinore	CA
6,390	2.3	2.3	0	1.8	1.8	0	North Highlands	CA
7,292	2.9	2.7	0	2.5	1.8	0	Sacramento	CA
8,036	3.1	3.1	0	2.8	2.4	0	Sacramento	CA
4,812	2.7	2.5	0	1.8	1.7	0	Sacramento	CA
5,519	1.4	1.3	0	1.2	1.1	0	Barstow	CA
5,587	1.4	1.4	0	0.9	0.9	0	Victorville	CA
5,491	2.1	1.9	0	1.4	1.2	0	Upland	CA
1,388	1.3	1.1	0	0.8	0.8	0	Fontana	CA
5,556	2.2	2.1	0	1.4	1.4	0	San Bernardino	CA
6,212	2	2	0	1.5	1.5	0	Chula Vista	CA
706	2.4	2.2	0	1.8	1.5	0	San Diego	CA
6,006	4.6	4	0	2.8	2.5	0	Escondido	CA
6,130	3.1	3.1	0	2.5	2.5	0	San Diego	CA
5,119	4.3	3.9	0	2.7	2.5	0	Otay Mesa	CA
6,247	2.1	1.9	0	1.5	1.4	0	San Francisco	CA
5,546	2.6	2.5	0	1.6	1.6	0	Stockton	CA
6,214	4.3	3.8	0	1.9	1.8	0	Redwood City	CA
4,811	5.2	3.4	0	1.2	1.2	0	Santa Barbara	CA
5,462	1.5	1.5	0	0.8	0.8	0	Santa Maria	CA
6,182	3.5	2.5	0	1.7	0.9	0	Capitan	CA
6,190	3.1	3.1	0	2.4	1.7	0	Welby	CO

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
6,505	3.5	3.3	0	2.7	2.4	0	Longmont	CO
6,507	3.4	3.2	0	2.3	2.2	0	Greeley	CO
6,481	2.2	2.1	0	1.4	1.3	0	Grand Junction	CO
6,487	5.1	4.6	0	3	2.2	0	Fort Collins	CO
4,249	0.8	0.8	0	0.7	0.7	0	Not provided	CO
6,420	4	3.5	0	2.6	2.3	0	Colorado Springs	CO
6,203	7.1	7	0	3.1	2.3	0	Denver	CO
6,488	4.8	4.5	0	2.4	1.9	0	Denver	CO
6,499	6	5.9	0	3.7	3.1	0	Hartford	CT
6,319	1.6	1.6	0	1.4	1.1	0	Westport	CT
6,305	3.3	3.2	0	2.2	2	0	Bridgeport	CT
6,491	1.2	1.1	0	0.9	0.9	0	Thomaston	CT
6,511	1.8	1.6	0	1.3	1.2	0	East Hartford	CT
3,314	2.2	2.1	0	1.9	1.6	0	New Haven	CT
6,525	6	4	0	2.6	1.8	0	Washington	DC
6,513	3	2.7	0	2.6	2.1	0	Washington	DC
7,030	1.4	1	0	0.9	0.8	0	Not provided	DE
7,171	2	2	0	1.3	1.1	0	Wilmington	DE
6,500	2.1	2	0	1.6	1.6	0	Fort Lauderdale	FL
6,513	1.6	1.5	0	1	0.9	0	Clearwater	FL
6,499	2.4	2.1	0	1.6	1.5	0	Hollywood	FL
6,269	8.7	6.2	0	3.3	2.8	0	Jacksonville	FL
6,318	2.5	2.1	0	1.4	1.2	0	Jacksonville	FL
6,250	2.3	2.2	0	1.1	1.1	0	Jacksonville	FL
6,551	2.6	2.5	0	2	1.8	0	Tampa	FL
6,511	1.2	1	0	0.5	0.4	0	Plant City	FL
6,506	2.2	2	0	1.2	1.1	0	Miami	FL
6,515	2.4	2	0	1.7	1.4	0	Not provided	FL
6,329	3.9	3.9	0	2.4	2.1	0	Miami	FL
6,550	2.8	2.6	0	2.1	1.6	0	Miami	FL
6,449	1.1	1	0	1	1	0	Winter Park	FL
2,167	1.5	1.4	0	1.1	1.1	0	Palm Beach	FL
6,280	1.5	1.4	0	1.2	1.1	0	Pompano Beach	FL
5,747	2.8	2.1	0	1.8	1.8	0	Decatur	GA
6,435	0.5	0.5	0	0.5	0.5	0	Not provided	GA

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
5,918	2.2	2.1	0	1.4	1.2	0	Atlanta	GA
6,554	2.1	1.7	0	1	1	0	Honolulu	HI
5,268	1.7	1.3	0	0.7	0.7	0	Ewa Beach	HI
2,050	1.5	1.1	0	0.7	0.6	0	Cedar Rapids	IA
5,017	1.6	1.5	0	1.3	1.2	0	Cedar Rapids	IA
7,396	1.9	1.6	0	1	1	0	Des Moines	IA
8,199	1.2	1.2	0	0.7	0.7	0	Davenport	IA
4,609	4.1	3.6	0	1.6	1.5	0	Boise	ID
7,797	2.9	2.7	0	2.3	2.2	0	East Saint Louis	IL
7,977	2.5	2.2	0	1.5	1.3	0	Springfield	IL
7,965	5.6	3	0	1.9	1.7	0	Rockford	IL
7,917	3.6	3.2	0	2.9	2.1	0	Peoria	IL
7,960	2.5	2.5	0	2.3	2.2	0	Maywood	IL
8,007	1.9	1.3	0	1.2	0.9	0	Northbrook	IL
7,926	2.4	2.3	0	1.7	1.5	0	Cicero	IL
7,783	2.3	2	0	1.8	1.5	0	Schiller Park	IL
7,982	7.3	4.4	0	2	1.5	0	Chicago	IL
6,380	1.9	1.7	0	1.2	1.2	0	Indianapolis (Remainder)	IN
5,972	2	1.8	0	1.5	1.1	0	Evansville	IN
6,538	9.1	7.1	0	3.3	3	0	East Chicago	IN
3,984	0.8	0.8	0	0.7	0.7	0	Pittsboro	IN
6,445	3.9	3.8	0	2.9	2.3	0	Fort Wayne	IN
8,710	3.6	3.5	0	3.2	2.1	0	Indianapolis	IN
6,520	2.8	2.7	0	2	1.2	0	Wichita	KS
6,394	2.7	2.4	0	1.6	1.5	0	Wichita	KS
6,389	2.2	2.2	0	1.8	1.4	0	Kansas City	KS
7,113	6.9	3.4	0	2.8	2.2	0	Louisville	KY
6,937	3.1	3.1	0	2.7	2.1	0	Louisville	KY
7,723	2.9	2.5	0	1.9	1.8	0	Baton Rouge	LA
5,769	0.9	0.7	0	0.4	0.4	0	Lynn	MA
6,586	3.4	3.4	0	3	2.5	0	Springfield	MA
6,835	3.7	3.2	0	2.6	2.1	0	Lowell	MA
6,830	1.7	1.6	0	1.3	1	0	Boston	MA
6,875	1.5	1.5	0	1.1	0.9	0	Boston	MA
6,831	2.8	2.7	0	1.7	1.3	0	Worcester	MA
6,022	2.6	2.4	0	1.6	1.5	0	Essex	MD

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
6,407	0.4	0.4	0	0.3	0.3	0	Not provided	MD
6,104	2.4	2.2	0	2.1	2	0	Baltimore	MD
6,439	1.4	1	0	0.9	0.8	0	Beltsville	MD
1,211	0.6	0.6	0	0.4	0.4	0	Presque Isle	ME
5,816	1.6	1.4	0	1.3	1	0	Portland	ME
6,351	0.4	0.4	0	0.4	0.3	0	Not provided	ME
4,569	2	1.6	0	1.2	1.1	0	Allen Park	MI
4,218	2	1.8	0	1.1	1	0	Grand Rapids	MI
7,278	2	2	0	0.8	0.7	0	Minneapolis	MN
7,262	0.5	0.5	0	0.3	0.3	0	Inver Grove Heights	MN
5,712	0.6	0.6	0	0.4	0.3	0	Rosemount	MN
2,169	1.8	1.5	0	1.2	1.2	0	Fridley	MN
7,022	3.2	3.1	0	2.4	2.4	0	St. Paul	MN
6,975	1.8	1.7	0	1.2	1.2	0	St. Cloud	MN
7,122	4.1	2.2	0	1.8	1.5	0	Duluth	MN
6,533	1.1	1	0	0.7	0.7	0	Sunset Hills	MO
6,536	1.9	1.8	0	1.2	1	0	Springfield	MO
6,528	3.3	2.9	0	2.2	1.6	0	St. Louis	MO
6,469	2.4	2.1	0	1.8	1.4	0	St. Louis	MO
2,083	3.4	3.4	0	2.9	2.7	0	Missoula	MT
2,104	2	1.9	0	0.7	0.7	0	Not provided	MT
6,916	6.7	5.7	0	2.4	2.2	0	West Yellowstone	MT
1,542	6.1	4.2	0	1.6	1.1	0	Not provided	MT
7,130	4.8	3.7	0	2.3	2	0	Billings	MT
7,244	7.7	3	0	1.8	1.5	0	Great Falls	MT
5,578	7.2	3.4	0	2.4	1.9	0	Kalispell	MT
2,819	1.5	1.5	0	1.3	1.1	0	Not provided	NC
7,811	1.9	1.8	0	1.5	1.4	0	Charlotte	NC
3,664	3.5	3.1	0	2.2	2.2	0	Raleigh	NC
7,081	4	2.8	0	2.4	1.8	0	Raleigh	NC
7,384	1.6	1.5	0	0.9	0.7	0	Rockwell	NC
2,044	2.2	2.2	0	1.7	1.7	0	Fayetteville	NC
7,927	2.3	2.1	0	1.9	1.6	0	Charlotte	NC
1,985	1.9	1.9	0	1.6	1.5	0	Greensboro	NC
6,531	2.7	2.5	0	2.3	1.9	0	Winston-Salem	NC
2,754	1.9	1.9	0	1.5	1.2	0	Durham	NC

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
7,930	6.5	1.7	0	1.2	0.7	0	Not provided	ND
6,365	3.1	2.9	0	2.5	2	0	Omaha	NE
7,252	6.5	4.5	0	1.8	1.8	0	Lincoln	NE
6,464	9.4	6	0	4.4	3.5	0	Manchester	NH
6,285	1.5	1.4	0	1.1	1.1	0	Fort Lee	NJ
2,188	1.8	1.6	0	1.1	1	0	Elizabeth	NJ
6,544	2.3	1.9	0	1.5	1.3	0	Morristown	NJ
6,508	3.4	3.4	0	2	1.6	0	Freehold	NJ
6,373	2.2	2.1	0	1.2	1.2	0	Camden	NJ
6,522	1	0.9	0	0.7	0.6	0	Not provided	NJ
6,498	2.1	2.1	0	1.7	1.5	0	Jersey City	NJ
6,318	2	1.6	0	1.2	0.9	0	Perth Amboy	NJ
6,484	2.2	2.2	0	1.8	1.6	0	Hackensack	NJ
6,326	3.7	2.7	0	1.8	1.4	0	Burlington	NJ
3,065	3.6	2.8	0	1.8	1.5	0	Albuquerque	NM
6,140	2.6	2.2	0	1.7	1.1	0	Albuquerque	NM
3,130	2.6	2.5	0	1.8	1.7	0	Albuquerque	NM
5,792	6.5	3	0	2.4	2.3	0	Albuquerque	NM
3,057	4.9	3.7	0	1.6	1.3	0	Albuquerque	NM
806	2.4	2.4	0	1.9	1.8	0	South Valley	NM
6,377	4.2	3.9	0	2.1	2	0	Sparks	NV
6,463	2.2	1.9	0	1.5	1.5	0	Lemmon Valley-Golden Valley	NV
4,243	2.9	2.7	0	2.2	1.7	0	Carson City	NV
5,033	3.5	2.9	0	2.3	2.1	0	Las Vegas	NV
6,217	5.1	4.7	0	4.2	3.7	0	Las Vegas	NV
6,275	4.1	3.9	0	3.1	2.8	0	Las Vegas	NV
6,190	3.2	2.9	0	2.1	2.1	0	Las Vegas	NV
6,271	4.2	3.6	0	2.5	2.4	0	Las Vegas	NV
6,178	5.8	5.3	0	2.6	2.4	0	Stateline	NV
6,476	2.1	2.1	0	1.5	1.5	0	Reno	NV
6,515	1.8	1.7	0	1.3	1.3	0	Reno	NV
6,400	3	2.9	0	2.3	2.2	0	Reno	NV
6,512	1.8	1.7	0	0.8	0.8	0	Reno	NV
6,466	1.8	1.8	0	1.2	0.9	0	Holtsville	NY
7,922	3	2.7	0	1.8	1.8	0	Schenectady	NY
7,445	8.6	2.3	0	1.7	1.6	0	New York	NY

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
7,956	2	1.9	0	1.7	1.4	0	Syracuse	NY
7,957	1.4	1.3	0	1.2	1	0	Niagara Falls	NY
7,769	1.5	1.2	0	1	0.9	0	Albany	NY
4,245	1.6	1.5	0	1.2	1.2	0	New York	NY
7,429	1.3	1.2	0	1.1	1	0	Rochester	NY
5,757	1.3	1	0	0.7	0.7	0	Tonawanda	NY
7,452	1.6	1.6	0	1.1	1	0	Buffalo	NY
7,962	2.3	2.1	0	1.8	1.7	0	New York	NY
3,419	1.7	1.6	0	1.3	1.2	0	New York	NY
6,967	4.8	4.8	0	2.1	1.5	0	Cleveland	OH
7,023	7.8	6.3	0	3.3	2.4	0	Cleveland	OH
6,948	2	1.5	0	0.9	0.8	0	Akron	OH
7,003	1.7	1.6	0	1.3	1.1	0	Akron	OH
6,158	2.9	2.9	0	2.6	2.5	0	Canton	OH
8,739	2.3	2.3	0	1.6	1.5	0	Dayton	OH
8,720	1.5	1.4	0	1.1	1	0	Dayton	OH
7,259	2.6	2.6	0	2.6	2.3	0	Mentor	OH
6,358	5.9	5.1	0	3.6	2.7	0	Cincinnati	OH
7,183	2.7	2.3	0	1.6	1.4	0	Columbus	OH
7,232	2	1.8	0	1.5	1.3	0	Cleveland	OH
7,252	2	2	0	1.2	1.2	0	Cleveland	OH
7,752	2.5	2.1	0	1.1	1.1	0	Oklahoma City	OK
1,382	0.4	0.4	0	0.3	0.3	0	Cherry Tree	OK
3,185	1.7	0.5	0	0.4	0.3	0	Park Hill	OK
4,625	1.7	0.3	0	0.5	0.3	0	Not provided	OK
2,278	1.7	1.6	0	1.4	1.3	0	Tulsa	OK
7,991	2.1	1.9	0	1.3	1.2	0	Tulsa	OK
5,933	2.7	2.6	0	2.3	2.2	0	Portland	OR
6,541	2.7	2.7	0	1.8	1.6	0	Portland	OR
6,437	3.4	3.4	0	2.6	2.2	0	Medford	OR
6,530	2.2	2.1	0	1.7	1.6	0	Eugene	OR
5,023	0.8	0.8	0	0.5	0.4	0	Not provided	PA
7,048	1.9	1.8	0	1.6	1.5	0	Pittsburgh	PA
7,243	2	2	0	1.3	1.2	0	York	PA
7,211	1.4	1	0	0.8	0.5	0	Greensburg	PA
6,517	1.6	1.6	0	1.3	1.1	0	Pittsburgh	PA
6,523	2.3	2.1	0	1.7	1.4	0	Pittsburgh	PA
7,054	2.6	2	0	1.6	1.3	0	Beaver Falls	PA

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
7,197	1.5	1.3	0	0.9	0.9	0	Not provided	PA
7,206	1.4	1.2	0	0.8	0.7	0	Altoona	PA
7,009	4.3	2.9	0	2	1.9	0	Bristol	PA
7,215	1.7	1.4	0	1.2	1.1	0	Johnstown	PA
7,174	1.4	1.4	0	1.1	1.1	0	Harrisburg	PA
7,152	2.8	1.6	0	1	1	0	Erie	PA
4,958	0.5	0.5	0	0.3	0.3	0	Not provided	PA
7,212	1.8	1.4	0	1.2	1	0	Scranton	PA
6,694	2.6	1.9	0	1.6	1.4	0	Lancaster	PA
7,162	1.2	1.2	0	0.7	0.7	0	New Castle	PA
7,200	2.6	2.4	0	1.9	1.5	0	Wilkes-Barre	PA
7,306	1.1	1.1	0	0.8	0.8	0	Norristown	PA
7,308	2.1	2	0	1.6	1.6	0	Freemansburg	PA
6,160	3.7	2.7	0	1.7	1.3	0	Philadelphia	PA
1,764	1.3	1.1	0	0.7	0.7	0	Philadelphia	PA
7,094	1.3	1.2	0	1.2	1.1	0	Charleroi	PA
3,795	4.2	2.8	0	2.3	2.3	0	Bayamon	PR
5,854	3.8	3.7	0	2.4	2.3	0	San Juan	PR
6,421	3.9	3.3	0	1.7	1.4	0	San Juan	PR
6,420	1.5	1.4	0	1	0.9	0	East Providence	RI
8,714	2.3	2	0	1.4	1.3	0	Greenville	SC
8,561	0.9	0.8	0	0.5	0.5	0	Not provided	SC
8,747	3.2	3.2	0	2.4	1.9	0	Nashville	TN
6,248	1.7	1.5	0	1	1	0	Kingsport	TN
8,514	3.2	2.5	0	1.5	1.4	0	Memphis	TN
6,824	1.6	1.5	0	0.8	0.7	0	Brownsville	TX
5,201	1.7	1.7	0	1.4	1.3	0	Dallas	TX
7,170	7	6.1	0	4.9	3.2	0	El Paso	TX
6,895	2.9	2.7	0	1.5	1.5	0	El Paso	TX
7,198	5	4.8	0	3	1.8	0	El Paso	TX
6,712	7.3	6	0	4.2	2.4	0	El Paso	TX
7,202	6.7	6.6	0	4.8	3.9	0	El Paso	TX
6,985	4.3	4	0	3	2.8	0	El Paso	TX
7,040	3.5	3.4	0	1.7	1.7	0	Laredo	TX
7,176	2.5	2.1	0	1.4	1.3	0	El Paso	TX
6,875	2.9	2.7	0	1.7	1.5	0	Not provided	TX
6,310	3.8	3.7	0	2.7	2.3	0	Houston	TX

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
6,258	2.1	1.9	0	1.6	1.3	0	Houston	TX
6,322	8.9	8.1	0	5.9	5.2	0	Houston	TX
5,662	1.4	1.4	0	1	0.8	0	Houston	TX
6,863	1.7	1.5	0	0.9	0.6	0	Deer Park	TX
5,594	1.8	1.7	0	1	0.7	0	Nederland	TX
7,105	0.8	0.3	0	0.3	0.3	0	Waco	TX
7,080	1.7	1.7	0	1.1	1	0	Fort Worth	TX
7,205	1.8	1.4	0	1	0.9	0	Arlington	TX
7,241	0.7	0.6	0	0.4	0.4	0	Austin	TX
6,942	3.2	2.9	0	1.4	1.3	0	Laredo	TX
6,154	0.9	0.9	0	0.5	0.5	0	Not provided	TX
6,309	2.5	2.4	0	1.9	1.4	0	San Antonio	TX
7,199	7.8	4.7	0	2.7	2.6	0	San Antonio	TX
7,235	3.2	2.9	0	1.6	1.6	0	Socorro	TX
6,469	3	2.9	0	2.2	2.1	0	Not provided	UT
6,297	4.1	3.6	0	2.4	2.3	0	Salt Lake City	UT
5,776	27.9	24.2	0	8.8	6.4	0	Ogden	UT
6,333	5	3.9	0	1.6	1.5	0	Provo	UT
5,744	3.3	3.3	0	2	2	0	Ogden	UT
2,125	3.7	3.3	0	1.8	1.8	0	West Valley	UT
5,996	1.1	1	0	0.9	0.8	0	Annandale	VA
6,479	1.3	1.2	0	1	0.8	0	Richmond	VA
6,469	1.7	1.6	0	1.2	1.1	0	Not provided	VA
7,175	1.4	1.4	0	1.2	1	0	Not provided	VA
7,133	1.9	1.9	0	1.9	1.8	0	Franconia	VA
6,489	2	2	0	1.5	1.4	0	Roanoke	VA
7,097	2	2	0	1.5	1.5	0	Mclean	VA
6,365	1.9	1.9	0	1.2	1.2	0	Alexandria	VA
6,439	4.2	4.1	0	1.6	1.3	0	Hampton	VA
6,487	4.7	4.5	0	2.2	1.9	0	Norfolk	VA
6,218	3.1	2.7	0	2.1	1.8	0	Rutland	VT
6,194	1.9	1.8	0	1.2	1.1	0	Burlington	VT
7,232	3.4	3.1	0	2.3	1.9	0	Bellevue	WA
5,468	0.3	0.3	0	0.3	0.3	0	Not provided	WA
6,885	1.4	1.2	0	0.9	0.9	0	Seattle	WA
7,140	4.3	4.1	0	2.4	2.4	0	Spokane	WA
7,674	0.5	0.5	0	0.4	0.4	0	Not provided	WI
6,541	3.9	3.5	0	2.3	2.2	0	Weirton	WV

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-2. 1-Hour and 8-Hour Maximum Carbon Monoxide Levels at Monitoring Stations Throughout the United States**

Number of measurements	1-Hour carbon monoxide level			8-Hour carbon monoxide level			City	State <sup>a</sup>
	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average	First maximum (ppmv)	Second maximum (ppmv)	Number exceeding average		
6,532	2.7	2.7	0	1.8	1.7	0	Weirton	WV
6,449	4.5	4.3	0	2.9	2.3	0	Weirton	WV
6,984	0.9	0.8	0	0.6	0.4	0	Not provided	WY
6,361	0.9	0.9	0	0.7	0.7	0	Not provided	WY

<sup>a</sup>Post office state abbreviations are used.

Source: EPA 2009i

## 6. POTENTIAL FOR HUMAN EXPOSURE

exposure level for trips greater than 30 minutes in duration were 21.4 ppmv for private car versus 10.4, 9.6, 4, and 11.5 ppmv for bus, trolley, electric train, and pedestrians, respectively, during the winter season (Duci et al. 2003). Besides the mode of transportation, route travelled, monitoring period, and the season of year had significant influences on the measured carbon monoxide concentrations. In-vehicle carbon monoxide levels were monitored in a passenger vehicle driven along a heavily traveled route of a commercial/residential area of Beirut, Lebanon, under several ventilation modes (Abi Esper et al. 2007a). Trips were conducted during morning rush hours in spring and summer months. The highest mean carbon monoxide levels were observed within vehicles driven with the windows and vents closed (37.4 ppmv) and windows closed and air conditioning on recirculation ventilation settings (30.8 ppmv) (Abi Esper et al. 2007a). Opening a window or setting the air conditioning unit to fresh air intake generally resulted in a reduction of carbon monoxide levels within the vehicle.

Indoor air levels of carbon monoxide are greatly dependent upon the presence of indoor combustion sources such as wood burning stoves, fireplaces, gas space heaters, and gas stoves, as well as the operating condition and usage patterns of these appliances. Dutton et al. (2001) studied carbon monoxide levels in two residences located in Boulder, Colorado using unvented natural gas fireplaces. Time-averaged carbon monoxide levels ranged from 1.5 to 78 ppmv at one residence and 1.6–30 ppmv at the other residence while the fireplace operated uninterrupted; however, levels exceeding 100 ppm were observed at one of the residences on several occasions. Background carbon monoxide levels were 0–7 ppmv in the homes when the fireplaces were not operating. Attached garages may also be a significant source of carbon monoxide to the indoor air of residences. The net increase of carbon monoxide levels in 16 homes with attached garages ranged from <1 to 30 ppmv following the cold start of an automobile enclosed within the attached garage (Graham et al. 2004).

#### 6.4.2 Water

The levels of carbon monoxide in the surface waters of the world's oceans are supersaturated with respect to the partial pressure of carbon monoxide in the atmosphere and are subject to diurnal cycling. Levels have been reported to vary from  $2 \times 10^{-8}$  to  $1.3 \times 10^{-7}$  cm<sup>3</sup> carbon monoxide per cm<sup>3</sup> of water (20–130 nL/L) based on measurements in the Pacific Ocean and from  $2 \times 10^{-8}$  to  $1 \times 10^{-7}$  cm<sup>3</sup> carbon monoxide per cm<sup>3</sup> of water (20–100 nL/L) in the North Atlantic (Liss and Slater 1974). Carbon monoxide concentrations in seawater measured in the upwelling region of the equatorial Pacific Ocean were reported to range from 42 to 173 nL/L on two consecutive sampling dates and reached a maximum value of 246 nL/L on subsequent sampling at a later date (Ohta 1997). These levels showed a marked diurnal variation with a

## 6. POTENTIAL FOR HUMAN EXPOSURE

maximum and minimum occurring early in the afternoon and morning, respectively. The concentrations in the water column decreased as a function of depth from the water surface to approximately 10 nL/L at a depth of 70 m. Tolli et al. (2006) reported a carbon monoxide level of 12 nanomoles/L at a coastal sampling location in Vineyard Sound, Massachusetts.

### 6.4.3 Sediment and Soil

*In-situ* soil-gas carbon monoxide levels were studied in a pine forest soil and cultivated soil as a function of depth from the surface (King 1999b). In the pine forest soil, the carbon monoxide concentrations remained elevated (250–320 ppbv) and greater than the atmospheric levels through the upper surface soil (O-horizon, 0–2 cm depth), but declined to values less than the atmospheric level in the lower depths of the soil (A-horizon 2–10 cm depth). In the cultivated soil, carbon monoxide levels decreased rapidly from approximately 250 ppbv in the upper 1 cm of the O-horizon to approximately 50 ppbv at a depth of 3 cm, which was roughly 5-fold less than ambient atmospheric levels.

### 6.4.4 Other Environmental Media

Carbon monoxide is released from tobacco smoke; however, the amount released is a function of the type of tobacco product (e.g., cigarette, cigar) and the degree to which tobacco is actively smoked. Moir et al. (2008) examined the emissions of carbon monoxide and other compounds under two smoking conditions from marijuana and tobacco cigarettes using a mechanical rotary smoking machine. The standard conditions employed a puff volume of 35 mL, puff duration of 2 seconds, and a puff interval of 60 seconds. Conditions more consistent of marijuana smoking employed a puff volume of 70 mL, puff duration of 2 seconds, and a 30-second interval between puffs. These conditions were referred to as extreme conditions. Under standard conditions, the average±standard deviation emission of carbon monoxide in mainstream smoke (n=20) was 20.8±1.9 mg carbon monoxide per tobacco cigarette and 13.4±1.6 mg carbon monoxide per marijuana cigarette. Under extreme smoking conditions, the levels were 41.5±4.0 mg carbon monoxide per tobacco cigarette and 35.3±2.9 mg carbon monoxide per marijuana cigarette. The carbon monoxide emissions in side stream smoke were 61.7±2.0 and 54.0±3.7 mg carbon monoxide per tobacco cigarette and marijuana cigarette, respectively, using the standard conditions. Under extreme smoking conditions, the levels were 61.6±2.9 and 50.6±3.9 mg carbon monoxide per cigarette for normal tobacco and marijuana cigarettes, respectively.

Emissions from plants and biomass burning are reported to contribute approximately  $1 \times 10^{14}$  g/year to the global environment (Khalil and Rasmussen 1990). The direct emission of carbon monoxide from living

## 6. POTENTIAL FOR HUMAN EXPOSURE

plants increases as a function of solar irradiance. The source of the carbon monoxide likely occurs as a result of direct photooxidation of the plant material followed by transport to the stomata; however, the exact mechanism is not fully understood (Sanderson 2002).

**6.5 GENERAL POPULATION AND OCCUPATIONAL EXPOSURE**

The general population is exposed to carbon monoxide through the inhalation of indoor and outdoor air. Since carbon monoxide is ubiquitous in the environment, all humans are exposed to some level of carbon monoxide. The CDC estimated that during 2004–2006, an estimated average of 20,636 emergency room visits occurred for nonfatal, unintentional, non-fire-related carbon monoxide exposures annually (CDC 2008). Approximately 73% of these exposures occurred in residences, 13% occurred in workplace settings, and the rest were in other or unknown settings. The greatest number of incidences (8,538 incidences representing approximately 41% of the cases) occurred during the winter months of December through February. Nearly 70% of the incidences were diagnosed as carbon monoxide poisoning. Table 6-3 displays the estimated number of exposure cases categorized by age and gender.

Blood COHb levels were measured on a cross-sectional national probability sample of persons representative of the civilian population in the United States aged 3–74 years in the second National Health and Nutrition Examination Survey (NHANES II) conducted from February 1976 to February 1980 (Radford and Drizd 1982). The statistical analysis of COHb levels in blood of the population based on smoking status and age is provided in Table 6-4. Four principle inhalation exposure routes were examined (outdoor air, indoor air, occupational exposure, and smoking). Of these four exposure routes, it was concluded that COHb levels in the population were most influenced by smoking status. Close examination of the data also indicated that COHb levels in the population tended to be greater during the winter months as opposed to the summer months, presumably due to the amount of time spent indoors where carbon monoxide levels are assumed to be higher than outdoor air.

Carbon monoxide exposure to European populations residing in five cities have been investigated through the EXPOLIS research project conducted from 1996 to 1998 and the results of this study have been summarized in several publications (Bruinen de Bruin et al. 2004a, 2004b; Hanninen et al. 2004). The geometric mean 48-hour exposure levels of nonsmoking subjects were 1.68, 0.82, 0.45, 2.17, and 1.50 mg/m<sup>3</sup> (1.45, 0.71, 0.39, 1.87, and 1.29 ppmv) in Athens, Basle, Helsinki, Milan, and Prague, respectively (Hanninen et al. 2004). Bruinen de Bruin et al. (2004a) used data for 50 office workers residing in Milan, Italy over a 1-year period to assess the contribution of local sources to exposure and

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-3. Average Annual Estimated Non Fatal Carbon Monoxide Exposure Cases in the United States Emergency Room Visits (2004–2006)**

Age (years)	Number of cases <sup>a</sup>	Percentage of the total
0–4	2,344	11.4
5–9	1,407	6.8
10–14	1,577	7.6
15–24	3,341	16.2
25–34	4,183	20.3
35–44	2,775	13.5
45–54	2,229	10.8
55–64	1,444	7.0
≥65	1,328	6.4
Gender		
Male	9,770	47.3
Female	10,866	52.7

<sup>a</sup>Age data were unavailable for eight cases.

Source: CDC 2008

## 6. POTENTIAL FOR HUMAN EXPOSURE

microenvironment concentrations. This study examined the time that the subjects spent in 11 different microenvironments and three exposure influencing activities: gas cooking, smoking, and commuting. The results of this study indicated that exposures from indoor environments contributed approximately 82% of the total carbon monoxide exposures because this is where the study population spent over 90% of their time; however, approximately 16% of the total exposure to carbon monoxide occurred from commuting activities even though these activities only accounted for about 7.5% of the populations time.

Occupations such as toll both workers, gas station attendants, and taxi drivers are potentially exposed to high levels of carbon monoxide due to emissions from automobile exhaust. Carbon monoxide levels were monitored continuously during both morning (9 AM–12 PM) and afternoon (2–4 PM) hours at seven gas stations located in New York, New Jersey, and Connecticut (API 1994). Mean (arithmetic) concentrations of carbon monoxide at the pumping islands ranged from 1.15 to 5.44 ppmv with a maximum one-minute level of 144 ppmv observed at one station. Firefighters are potentially exposed to carbon monoxide levels near or exceeding recommended occupational exposure limits. A study conducted from 1986 to 1989 in northern California was performed to estimate the level of carbon monoxide that wildland firefighters are exposed to during various job tasks associated with that profession (Materna et al. 1992). Following a series of prescribed burns, the instantaneous fireline carbon monoxide levels ranged from 3 to 80 ppmv; however, firefighters that tended gasoline powered pumping engines were exposed to levels as high as 300 ppmv. The mean time-weighted average (TWA) exposure level obtained from 46 personal measurements was 14.4 ppmv, but one employee had a TWA exposure of 38 ppmv and five firefighters had TWAs above 25 ppmv. Industrial or in-home use of methylene chloride paint strippers in poorly ventilated areas can lead to high levels of carbon monoxide in blood since carbon monoxide is a metabolic byproduct of methylene chloride.

### 6.6 EXPOSURES OF CHILDREN

This section focuses on exposures from conception to maturity at 18 years in humans. Differences from adults in susceptibility to hazardous substances are discussed in Section 3.7, Children's Susceptibility.

Children are not small adults. A child's exposure may differ from an adult's exposure in many ways. Children drink more fluids, eat more food, breathe more air per kilogram of body weight, and have a larger skin surface in proportion to their body volume. A child's diet often differs from that of adults. The developing human's source of nutrition changes with age: from placental nourishment to breast milk or formula to the diet of older children who eat more of certain types of foods than adults. A child's

## 6. POTENTIAL FOR HUMAN EXPOSURE

behavior and lifestyle also influence exposure. Children crawl on the floor, put things in their mouths, sometimes eat inappropriate things (such as dirt or paint chips), and spend more time outdoors. Children also are closer to the ground, and they do not use the judgment of adults to avoid hazards (NRC 1993).

As with adults, children are exposed to carbon monoxide through the inhalation of indoor and outdoor air. The NHANES II survey (see Table 6-4) includes COHb levels of children. Several activities may influence the levels of carbon monoxide that children or infants are exposed to. Time spent indoors versus outdoors, riding in automobiles, and exposure to second-hand tobacco smoke influence the total exposure that children receive to carbon monoxide. COHb levels in children were significantly greater when measured during the winter months of November–March as opposed to levels obtained in May–September. The average COHb level for all children in the NHANES study aged 3–11 years was 0.87% for samples collected during the winter months and 0.58% during the summer months. The location where the children in the population resided also influenced the COHb levels observed in the study. Children residing in central cities with populations over 1 million people tended to have the highest mean COHb levels (1.01% winter months and 0.77% summer months), while children living in rural areas had mean COHb levels of 0.79% in the winter months and 0.49% during the summer months (Radford and Drizd 1982).

Pregnant females who smoke potentially expose their unborn fetus to carbon monoxide, and the health consequences of this activity have been discussed in Chapter 3. Carbon monoxide crosses the placenta and can accumulate in the fetus to a greater extent than the mother.

### 6.7 POPULATIONS WITH POTENTIALLY HIGH EXPOSURES

Populations that are exposed to high levels of vehicular traffic are expected to have greater exposure to carbon monoxide as compared to individuals in low traffic density areas. As discussed in Section 6.5, certain occupations such as toll workers, gas station attendants, firefighters, and other professions exposed to combustion sources may have high levels of exposure.

Members of the general population that smoke are exposed to higher levels of carbon monoxide than nonsmoking members of the general population. Data from the NHANES II study indicates that mean COHb levels in the blood of smokers are approximately 4 times greater than levels for members of the nonsmoking population (Radford and Drizd 1982). More recent data support these conclusions. COHb levels were measured as part of pulmonary function testing in 100 subjects in one pulmonary laboratory

## 6. POTENTIAL FOR HUMAN EXPOSURE

**Table 6-4. Carboxyhemoglobin Levels in the U.S. Population Based Upon Smoking Status**

Age (years)	N <sup>2a</sup>	N <sup>3b</sup>	Mean COHb percent	Standard deviation	Standard error	50 <sup>th</sup> percentile	75 <sup>th</sup> percentile	90 <sup>th</sup> percentile	95 <sup>th</sup> percentile
All smoking status									
3-74	9,365	195,877	1.93	2.236	0.037	0.91	2.38	5.49	6.83
3-11	2,055	30,066	0.73	0.502	0.019	0.67	0.87	1.12	1.42
12-74	7,310	165,812	2.14	2.358	0.044	1.01	3.17	5.79	7.05
Never smoked									
3-74	5,459	106,042	0.83	0.671	0.021	0.72	0.97	1.33	1.65
3-11	2,055	30,066	0.73	0.502	0.019	0.67	0.87	1.12	1.42
12-74	3,404	75,976	0.87	0.726	0.025	0.74	1.01	1.38	1.77
Ex-smokers									
12-74	1,366	28,655	0.97	0.999	0.031	0.77	1.04	1.58	2.08
Current smokers									
12-74	2,533	61,015	4.30	2.533	0.072	4.15	5.89	7.56	8.68

<sup>a</sup>N<sup>2</sup> = unweighted population size.

<sup>b</sup>N<sup>3</sup> = population estimate in thousands.

Source: Radford and Drizd 1982

## 6. POTENTIAL FOR HUMAN EXPOSURE

located in a Virginia hospital outpatient setting (Mahoney et al. 2007). COHb levels for the entire group averaged 1.9% with a range of 1–8%. The average COHb level for nonsmokers (n=85) was 1.6%, while the level for smokers (n=15) was 3.5%. COHb levels were measured for smokers versus nonsmokers at 4 commercial establishments where cigarette smoking was not prohibited (Light et al. 2007). In addition, COHb levels were obtained in a control group of 50 college students and professors in a well-ventilated, nonsmoking environment. The average COHb concentration of 33 smokers from the commercial establishments was 5.04%, while the average value for the 27 nonsmokers was 2.49%. COHb levels ranged from 1 to 6% in the nonsmokers and from 1 to 14% in the smoking group. The COHb levels in the control group were 1%.

## 6.8 ADEQUACY OF THE DATABASE

Section 104(i)(5)(A) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of carbon monoxide is available. Where adequate information is not available, ATSDR, in conjunction with NTP, is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of carbon monoxide.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

### 6.8.1 Identification of Data Needs

**Physical and Chemical Properties.** Information is available on the physical and chemical properties of carbon monoxide (George 2001; Lide 2008; O’Neil et al. 2006; Verschueren 2001). These data are captured in Chapter 4. No data needs are identified.

**Production, Import/Export, Use, Release, and Disposal.** According to the Emergency Planning and Community Right-to-Know Act of 1986, 42 U.S.C. Section 11023, industries are required to submit substance release and off-site transfer information to the EPA. The TRI, which contains this information for 2007, became available in February of 2009. This database is updated yearly and should

## 6. POTENTIAL FOR HUMAN EXPOSURE

provide a list of industrial production facilities and emissions. Adequate information on the production and use of carbon monoxide was located (George 2001); no information on the import/export of carbon monoxide was found but these volumes are assumed to be low. Since carbon monoxide is not required to be reported under the TRI, the production and emissions from U.S. industrial facilities is not reported to the EPA. The EPA continuously updates the NEI database, which contains detailed information about sources that emit carbon monoxide. No data needs are identified.

**Environmental Fate.** The environmental fate of carbon monoxide is well understood. When released to the atmosphere, carbon monoxide eventually reacts with photochemically produced hydroxyl radicals and is oxidized to CO<sub>2</sub> (EPA 2000). Microorganisms have also been shown to oxidize carbon monoxide to CO<sub>2</sub> (Tolli et al. 2006). Data on the half-life of carbon monoxide in indoor environments would be desirable.

**Bioavailability from Environmental Media.** Carbon monoxide is a gas and is not considered bioavailable from environmental media other than air. No data needs are identified.

**Food Chain Bioaccumulation.** Carbon monoxide is a gas and does not bioaccumulate in the food chain. No data needs are identified.

**Exposure Levels in Environmental Media.** Reliable monitoring data for the levels of carbon monoxide in contaminated media at hazardous waste sites are needed so that the information obtained on levels of carbon monoxide in the environment can be used in combination with the known body burden of carbon monoxide to assess the potential risk of adverse health effects in populations living in the vicinity of hazardous waste sites.

The EPA continuously monitors carbon monoxide levels throughout the United States. The EPA's NEI database contains detailed information about sources that emit criteria air pollutants including carbon monoxide for the 50 United States, Washington DC, Puerto Rico, and the U.S. Virgin Islands. The AQS database is EPA's repository of criteria air pollutant monitoring data since the 1970s. These databases are updated regularly.

**Exposure Levels in Humans.** All humans are exposed to carbon monoxide through the inhalation of ambient air. Indoor air levels of carbon monoxide are highly variable and depend upon the type, condition, and venting procedures of appliances. The NHANES II survey, conducted from February 1976

## 6. POTENTIAL FOR HUMAN EXPOSURE

to February 1980, provided an analysis of COHb levels in blood of the population based on smoking status, age, race, and other factors (Radford and Drizd 1982). These data are not current; therefore, a data need exists to update these data with new monitoring results.

This information is necessary for assessing the need to conduct health studies on these populations.

**Exposures of Children.** Children are exposed to carbon monoxide by the same pathway as adults (inhalation of air). While the levels of COHb in children's blood from the NHANES II study is comprehensive, these surveys are over 2 decades old. Therefore, a data need exists to update this study with new monitoring results.

Child health data needs relating to susceptibility are discussed in Section 3.12.2, Identification of Data Needs: Children's Susceptibility.

**Exposure Registries.** No exposure registries for carbon monoxide were located. This substance is not currently one of the compounds for which a sub-registry has been established in the National Exposure Registry. The substance will be considered in the future when chemical selection is made for sub-registries to be established. The information that is amassed in the National Exposure Registry facilitates the epidemiological research needed to assess adverse health outcomes that may be related to exposure to this substance.

### 6.8.2 Ongoing Studies

The Federal Research in Progress (FEDRIP 2009) database provides additional information obtainable from a few ongoing studies that may fill in some of the data needs identified in Section 6.8.1. A study being conducted by Ohio State University (Ross Kauffman, principle investigator), seeks to study tobacco use in two Ohio prisons. One objective of this study is to examine the influence of an indoor tobacco ban on smoking behaviors among low-security prisoners. A second objective is to provide prison administrators with the information needed to develop successful policies by gathering information on prisoner attitudes toward tobacco control policies and cessation programs in correctional settings. The study is also intended to lay the groundwork for future studies of tobacco use in prison facilities. It is anticipated that this study will address COHb levels in the prison facilities.

6. POTENTIAL FOR HUMAN EXPOSURE

This page is intentionally blank.

## 7. ANALYTICAL METHODS

The purpose of this chapter is to describe the analytical methods that are available for detecting, measuring, and/or monitoring carbon monoxide, its metabolites, and other biomarkers of exposure and effect to carbon monoxide. The intent is not to provide an exhaustive list of analytical methods. Rather, the intention is to identify well-established methods that are used as the standard methods of analysis. Many of the analytical methods used for environmental samples are the methods approved by federal agencies and organizations such as EPA and the National Institute for Occupational Safety and Health (NIOSH). Other methods presented in this chapter are those that are approved by groups such as the Association of Official Analytical Chemists (AOAC) and the American Public Health Association (APHA). Additionally, analytical methods are included that modify previously used methods to obtain lower detection limits and/or to improve accuracy and precision.

### 7.1 BIOLOGICAL MATERIALS

Carbon monoxide forms a strong coordination bond with the iron atom complex of Hb forming COHb, which is a unique biomarker for carbon monoxide exposure for humans. COHb absorbs radiation in the blue wavelength region of the visible spectrum known as the Soret region (390–440 nm) and COHb levels in blood can be analyzed spectrophotometrically (Rodkey et al. 1979). Carbon monoxide-oximeters are specialized medical instruments that are employed in clinical and hospital settings to rapidly determine COHb, O<sub>2</sub>Hb, and methemoglobin (MetHb) levels in blood by taking the difference between the light absorption at each of these Hb derivative's and the total light absorption for all Hb derivatives that are present in the blood sample (Mahoney et al. 1993). This instrument is particularly useful for measuring high levels of COHb. Fetal Hb and bilirubin have the potential to interfere with spectrophotometric determination of COHb if present in sufficient quantities. Methylene blue cardiac dye and sulfhemoglobin can also adversely affect these measurements. In addition, some carbon monoxide-oximeters have been shown to overestimate low concentrations of COHb as compared to gas chromatography (GC) methods (Mahoney et al. 1993).

GC with reduction gas detectors is a selective and sensitive method to measure the amount of carbon monoxide bound to Hb in the blood and is considered a highly reliable method for measuring exposure to carbon monoxide, especially at low levels. The addition of potassium ferricyanide to Hb-bound carbon monoxide oxidizes Hb to MetHb and releases carbon monoxide, which is injected in the gas chromatograph (Vreman et al. 1984). Carbon monoxide eluted from the GC column reacts with mercuric

## 7. ANALYTICAL METHODS

oxide (HgO) to form gaseous mercury vapor; the mercury released and measured spectrophotometrically at 254 nm is proportional to the amount of carbon monoxide gas released. The detection limits for this method were reported as 0.05 nL of carbon monoxide per microliter of blood, corresponding to a saturation of approximately 0.005% COHb in blood (Vreman et al. 1984).

Carbon monoxide levels in humans can be estimated by the analysis of exhaled breath using GC and electrochemical devices. In many measurement techniques, the subject performs an inhalation-breathhold maneuver and exhales through a mouthpiece into the instrument inlet. The alveolar air is retained for analysis, and the reading in ppm of carbon monoxide can be converted to COHb using a calibration curve (EPA 2000). A rapid and sensitive technique for the analysis of carbon monoxide in a single exhaled breath sample has been described (Fritsch et al. 2007). This analysis requires the subject to exhale into a mouthpiece that is attached to a Nafion® dryer and cooling trap to remove water, CO<sub>2</sub>, and other possible interfering species. Detection is by high resolution infrared cavity ring-down laser spectroscopy in a multipath absorption cell. Amperometric electrochemical sensors have also been used for the analysis of carbon monoxide in exhaled breath and ambient air (Vreman et al. 1994). These sensors typically consist of a series of electrodes (working, reference, and counter electrodes) immersed in an electrolyte solution. Carbon monoxide diffuses through a porous teflon membrane and is oxidized at the working electrode. The transfer of electrons that accompanies the redox reaction yields the output signal of the sensor. A reference electrode measures the potential of the electrolyte. The current flowing through the working electrode is compensated by the third electrode, a counterelectrode through which an equal and opposite current flows (Vreman et al. 1994).

The analytical methods used for the detection of carbon monoxide in biological samples are summarized in Table 7-1.

## 7.2 ENVIRONMENTAL SAMPLES

Analytical methods for the detection of carbon monoxide in environmental samples have been thoroughly discussed in several review articles (EPA 1991, 2000; Novelli 1999). Table 7-2 provides a summary of many of these analytical methods that are commonly employed in environmental monitoring studies of carbon monoxide. Analytical methods used to measure carbon monoxide in environmental samples require the use of high purity standard reference gas in order to frequently calibrate the instrument and ensure consistent measurements between different laboratories.

## 7. ANALYTICAL METHODS

**Table 7-1. Analytical Methods for Determining Carbon Monoxide in Biological Samples**

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Blood	Collect sample and transfer to a 5 mL heparinized syringe and shake for 1 minute before analysis.	Spectro photometric determination using CO-oximeters	<5% COHb		Mahoney et al. 1993
Blood	Blood was placed into 2 mL CO-free GC vial containing 100 g/L $K_3Fe(CN)_6$ in 0.1 mol/L potassium phosphate buffer, pH 6.0, containing 10 g of saponin/L	GC with mercury reduction gas detector	0.005% COHb		Mahoney et al. 1993
Blood	Obtain blood by heel stick (infant), or fingerstick (adult) and venipuncture. Mix with $K_3Fe(CN)_6$ containing solution to liberate carbon monoxide to a headspace.	GC with mercury reduction gas detector	0.005% COHb		Vreman et al. 1984
Breath	Breath samples were collected with a reverse syringe pump and analyzed within 30 minutes or kept on ice until analysis within 3 hours.	Electrochemical (EC) sensor or GC with mercury reduction gas detector	EC: 0.1 $\mu$ L carbon monoxide/L air GC: 50% of the noise level of 0.01 $\mu$ L carbon monoxide/L air		Vreman et al. 1994

COHb = carboxyhemoglobin; GC = gas chromatography;  $K_3Fe(CN)_6$  = potassium ferricyanide

## 7. ANALYTICAL METHODS

**Table 7-2. Analytical Methods for Determining Carbon Monoxide in Environmental Samples**

Sample matrix	Preparation method <sup>a</sup>	Analytical method	Sample detection limit <sup>b</sup>	Percent recovery	Reference
Air	Atmospheric constituents are separated using two-column isothermic chromatography in a temperature-controlled oven, followed by analysis using FID, HgO/UV absorbance, or ECD.	GC-FID GC-HgO/UV GC-ECD	25–50 ppbv GC-FID 10 ppb GC-ECD 1–2 ppb GC-HgO/UV	NR	Novelli 1999
Air	Air samples are passed through reference and measurement optical cells. An optical filter is used to transmit only a small band width of infrared radiation.	NDIR-GFC	22–70 ppbv	NR	Novelli 1999
Air	Air samples are introduced into commercial instruments over a wide spectral bandwidth or the use of optical filters can be employed to narrow the spectral wavelength.	FTIR	1 ppbv	NR	Novelli 1999
Air	Tune laser to 4.7 $\mu\text{m}$ and measure absorption by carbon monoxide gas.	TDLS	<1 ppbv	NR	Novelli 1999
Air	Measure in fluorescence chambers operated at low pressure to avoid O <sub>2</sub> absorption.	RF	5 ppbv	NR	Novelli 1999
Workplace air	Each sample is collected by drawing a known volume of air into a five-layer aluminized gas sampling bag. A portion of the gas sample is introduced into a gas sampling loop, injected into a gas chromatograph, and analyzed using a discharge ionization detector.	GC-DID	0.12 ppmv; (0.400 ppmv quantitative)		OSHA 1991 (Method 210)

## 7. ANALYTICAL METHODS

**Table 7-2. Analytical Methods for Determining Carbon Monoxide in Environmental Samples**

Sample matrix	Preparation method <sup>a</sup>	Analytical method	Sample detection limit <sup>b</sup>	Percent recovery	Reference
Sea water	Water samples are siphoned into a glass stopcock and a portion of the water was replaced with carbon monoxide free synthetic air to create a headspace. The flask was shaken vigorously by hand for 1 minute and kept standing at room temperature for 5 minutes in the dark.	GC-FID	6 nL of carbon monoxide/L of water		Ohta 1997
Soil gas	Samples were collected using a 1-cm <sup>3</sup> syringe fitted with a 50-mm-long 26-gauge side port needle. A 1-cm <sup>3</sup> sample was obtained (sampling time, about 60 seconds); the syringe was then raised, and its contents were rapidly analyzed in the field by using GC	GC-HgO/UV	<5 ppbv	NR	King 1999b

<sup>a</sup>Hydrogen rich carrier gas is used to reduce carbon monoxide to methane prior to analysis by FID and N<sub>2</sub>O is added to a nitrogen rich carrier gas for analysis by ECD. Helium gas is used as the carrier gas for analysis by DID.

<sup>b</sup>Measurements of atmospheric carbon monoxide levels are relative to high purity reference gas used to calibrate the instruments.

DID = discharge ionization detector; ECD = electron capture detector; FID = flame ionization detector; GC = gas chromatography; FTIR = Fourier transform spectroscopy; GFC = gas filter correlation; HgO = mercuric oxide; nondispersive infrared; RF = resonance fluorescence; NR = not reported; TDLS = tunable diode laser spectroscopy; UV = ultraviolet

## 7. ANALYTICAL METHODS

Most techniques used for the measurement of carbon monoxide in the environment employ spectroscopic methods or GC. Non-dispersive infrared (NDIR) spectroscopy and GC using flame ionization detection (FID) are the two most common methods used to analyze carbon monoxide in environmental samples. Commercial instruments are capable of detection limits in the ppb range. For analysis by NDIR spectroscopy, air is pumped into sample cells and the absorption is measured at a carbon monoxide vibrational band at  $2,174\text{ cm}^{-1}$  ( $4.7\text{ }\mu\text{m}$ ). An optical filter is employed before the sample to remove light of other frequencies. Most NDIR spectrometers use gas filter correlation (GFC) methodology to improve sensitivity and reduce interferences by other gasses. In this method, the sample is simultaneously passed through both a reference cell containing a high concentration of carbon monoxide and a sample cell containing nitrogen gas. Gas species other than carbon monoxide will attenuate radiation equally in both cells; however, when carbon monoxide is present, the amount of light passing through the reference cell is unchanged, but that passing through the sample cell is absorbed at the characteristic carbon monoxide frequency. This difference in absorption between the light passing through the cells is linearly related to carbon monoxide concentrations in the air sample (EPA 2000).

In GC analysis of carbon monoxide, air samples are pumped through a silica gel or alumina pre-column followed by an analytical column composed of a molecular sieve. The pre-column captures  $\text{CO}_2$ , non-methane hydrocarbons, and water and the molecular sieve effectively separates carbon monoxide from hydrogen gas and methane. Since the sensitivity for carbon monoxide by FID is low, a hydrogen-rich carrier gas is used to reduce carbon monoxide to methane using heated nickel as a catalyst prior to FID analysis. Minimum detection limits of 0.05 ppmv or better have been achieved (Novelli 1999). With the development of the discharge ionization detector (DID) for use with GC analysis, it is possible to measure carbon monoxide concentrations directly at low levels in occupational settings (OSHA 1991). Helium is generally used as the sample carrier gas and as the ionized species. Helium passes through a chamber where a glow discharge is generated and high-energy photons are produced. The high-energy photons pass through an aperture to a second chamber and ionize the carbon monoxide in the sample stream. The resulting ions produce an electrical current, which is measured with a standard electrometer (OSHA 1991). The greater the concentration of carbon monoxide in the sample, the more ions will be produced, resulting in a greater current. OSHA method 210 discusses the GC-DID analysis for carbon monoxide and reports a detection limit of 0.12 ppmv and quantification limit of 0.40 ppmv.

As discussed in Novelli (1999), electron capture detection (ECD) can also be used in conjunction with GC to measure atmospheric levels of carbon monoxide. Carbon monoxide does not elicit a strong response from the ECD because it does not easily capture electrons; however, the addition of  $\text{N}_2\text{O}$  to a

## 7. ANALYTICAL METHODS

N<sub>2</sub> carrier gas improves sensitivity. In addition, GC/mass spectrometry, GC with mercury reduction gas detection, Fourier transform infrared (FTIR) spectroscopy, tunable diode laser spectroscopy (TDLS), and resonance fluorescence spectroscopy have also been used to measure carbon monoxide levels in air (Novelli 1999).

Commercially available electrochemical carbon monoxide detectors for home use can be purchased at most hardware or home improvement stores. These detectors use electrodes that reside in an electrolyte solution such as dilute sulfuric acid. Carbon monoxide is oxidized at one electrode to CO<sub>2</sub> and O<sub>2</sub> is consumed at the other electrode (Vreman et al. 1994). Measurement of the current in the cell provides a measure of the concentration of carbon monoxide in the atmosphere. Metal oxide semiconductor detectors such as tin oxide sensors operate on the principle that the resistance of a tin oxide layer decreases as carbon monoxide adsorbs to the surface of the sensor and is oxidized to CO<sub>2</sub> (Dobos and Zimmer 1985). When the conductance exceeds a preset threshold level, an 85 dB alarm is sounded (the same loudness as a smoke alarm). Biomimetic detectors use a gel-coated disc containing synthetic Hb that turns black when carbon monoxide is present at high levels. Reduction in the amount of transmitted light through the disc alerts the sensor to possible carbon monoxide contamination and sets off the alarm. Early carbon monoxide detectors were designed to provide an audible warning at relatively low levels of exposure and were often difficult to reset (Leikin 1996). Furthermore, the general public's knowledge regarding the use and placement of these alarms in the residence was limited. Recent changes to the design and standards of these alarms have made them more user friendly to the general population and helped to eliminate low-level activations, which can be caused by a variety of common occurrences. The use and effectiveness of these home-based alarms, including guideline standards from Underwriters Laboratory, have been reviewed by Rhee and Leikin (2007).

Ohta (1997) discussed a GC-FID method for the detection of carbon monoxide in seawater. Water samples were collected in a glass stopcock and a portion of the water was replaced with carbon monoxide free synthetic air to create a headspace. The flask was shaken vigorously by hand for 1 minute and kept standing at room temperature for 5 minutes in the absence of sunlight in order to complete the equilibration between aqueous and gaseous phases. A small portion of air (1–2 mL) was injected into the GC for analysis. Carbon monoxide concentrations were corrected based on equilibrium constants of carbon monoxide between the aqueous and gaseous phases in the flask for headspace preparation. The concentrations were further corrected for water vapor contained in the headspace gas at concentrations of 3.8–4.1% in volume at 26–27 °C. The detection limit was reported as 6 nanoliters of carbon monoxide per liter of water.

## 7. ANALYTICAL METHODS

### 7.3 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of carbon monoxide is available. Where adequate information is not available, ATSDR, in conjunction with NTP, is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of carbon monoxide.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

#### 7.3.1 Identification of Data Needs

##### **Methods for Determining Biomarkers of Exposure and Effect.**

*Exposure.* COHb is a unique biomarker of exposure for humans to carbon monoxide and accurate, sensitive methods exist for its measurement (Mahoney et al. 1993). GC methods are also available for measuring Hb-bound carbon monoxide (Vreman et al. 1984).

*Effect.* No specific biomarkers of effect for carbon monoxide were identified. Effects and symptoms of carbon monoxide poisoning are generally related to tissue hypoxia and are not specific for carbon monoxide exposure.

##### **Methods for Determining Parent Compounds and Degradation Products in Environmental**

**Media.** Sensitive analytical methods are available for determining the levels of carbon monoxide in air samples. Thorough reviews on this subject are available from the EPA (1991, 2000) and Novelli (1999). Analytical methods are also available for the determination of carbon monoxide in water samples (Ohta 1997).

## 7. ANALYTICAL METHODS

**7.3.2 Ongoing Studies**

The Federal Research in Progress (FEDRIP 2009) database provides additional information obtainable from a few ongoing studies that may fill in some of the data needs identified. Platypus Technologies LLC located in Madison, Wisconsin is developing a liquid crystal-based sensor for real-time detection of ozone, carbon monoxide, and CO<sub>2</sub>. Kestrel Labs Inc. located in Boulder, Colorado is developing a new carbon monoxide-oximeter that can use laser light sources for accurate photo-plethysmographic measurements of COHb.

7. ANALYTICAL METHODS

This page is intentionally blank.

## 8. REGULATIONS, ADVISORIES, AND GUIDELINES

MRLs are substance specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors and other responders to identify contaminants and potential health effects that may be of concern at hazardous waste sites.

MRLs were not derived for carbon monoxide, as discussed in Section 2.3.

EPA (IRIS 2009) has not established an oral reference dose (RfD) or an inhalation reference concentration (RfC) for carbon monoxide.

The International Agency for Research on Cancer (IARC), the National Toxicology Program (NTP), and EPA have not classified carbon monoxide for human carcinogenicity (IARC 2009; IRIS 2009; NTP 2005).

OSHA has required employers of workers who are occupationally exposed to carbon monoxide to institute engineering controls and work practices to reduce and maintain employee exposure at or below permissible exposure limits (PELs) (OSHA 2009). The employer must use engineering and work practice controls to reduce exposures to not exceed  $55 \text{ mg/m}^3$  (50 ppmv) for carbon monoxide at any time (OSHA 2009).

EPA has designated carbon monoxide as a hazardous air pollutant (HAP) under the Clean Air Act (CAA) (EPA 2009c). Additionally, under the National Ambient Air Quality Standards (NAAQS), EPA is required to set limits to protect public health, including the health of "sensitive" populations such as asthmatics, children, and the elderly. Carbon monoxide is required to not to exceed levels of 10 and  $40 \text{ mg/m}^3$  (9 and 35 ppmv) for 8- and 1-hour averaging times, respectively, and not to be exceeded more than once per year (EPA 2009d).

The international and national regulations, advisories, and guidelines regarding carbon monoxide in air, water, and other media are summarized in Table 8-1.

## 8. REGULATIONS AND ADVISORIES

**Table 8-1. Regulations, Advisories, and Guidelines Applicable to Carbon Monoxide**

Agency	Description	Information	Reference
<u>INTERNATIONAL</u>			
Guidelines:			
IARC	Carcinogenicity classification	No data	IARC 2009
WHO	Air quality guidelines		WHO 2000
	TWA based on effects other than cancer or odor/annoyance using an averaging time of:		
	15 minutes	100 mg/m <sup>3</sup> (87 ppm)	
	30 minutes	60 mg/m <sup>3</sup> (52 ppm)	
	1 hour	30 mg/m <sup>3</sup> (26 ppm)	
	8 hours	10 mg/m <sup>3</sup> (9 ppm)	
	Drinking water quality guidelines	No data	WHO 2006
<u>NATIONAL</u>			
Regulations and Guidelines:			
a. Air			
ACGIH	TLV (8-hour TWA)	29 mg/m <sup>3</sup> (25 ppm)	ACGIH 2008
	TLV-basis (critical effect)	Carboxyhemoglobinemia	
AIHA	ERPG-1 <sup>a</sup>	229 mg/m <sup>3</sup> (200 ppm)	AIHA 2008
	ERPG-2 <sup>a</sup>	401 mg/m <sup>3</sup> (350 ppm)	
	ERPG-3 <sup>a</sup>	573 mg/m <sup>3</sup> (500 ppm)	
EPA	AEGL-1 <sup>b</sup>	Not recommended due to insufficient data	EPA 2009b
	AEGL-2 <sup>b</sup>		
	10 minutes	481 mg/m <sup>3</sup> (420 ppm)	
	30 minutes	172 mg/m <sup>3</sup> (150 ppm)	
	60 minutes	95 mg/m <sup>3</sup> (83 ppm)	
	4 hours	38 mg/m <sup>3</sup> (33 ppm)	
	8 hours	31 mg/m <sup>3</sup> (27 ppm)	
	AEGL-3 <sup>b</sup>		
	10 minutes	1,948 mg/m <sup>3</sup> (1,700 ppm)	
	30 minutes	687 mg/m <sup>3</sup> (600 ppm)	
	60 minutes	378 mg/m <sup>3</sup> (330 ppm)	
	4 hours	172 mg/m <sup>3</sup> (150 ppm)	
	8 hours	149 mg/m <sup>3</sup> (130 ppm)	
	Hazardous air pollutant	No	EPA 2009c 42 USC 7412

## 8. REGULATIONS AND ADVISORIES

**Table 8-1. Regulations, Advisories, and Guidelines Applicable to Carbon Monoxide**

Agency	Description	Information	Reference
<u>NATIONAL</u> ( <i>cont.</i> )			
	National Ambient Air Quality Standards		EPA 2009d
	8-hour averaging time <sup>c</sup>	10 mg/m <sup>3</sup> (9 ppm)	
	1-hour averaging time <sup>c</sup>	40 mg/m <sup>3</sup> (35 ppm)	
NIOSH	REL (10-hour TWA)	40 mg/m <sup>3</sup> (35 ppm)	NIOSH 2005
	Ceiling	229 mg/m <sup>3</sup> (200 ppm)	
	IDLH	1,375 mg/m <sup>3</sup> (1,200 ppm)	
	Target organs	Cardiovascular system, lungs, blood, and central nervous system	
OSHA	PEL (8-hour TWA) for general industry	55 mg/m <sup>3</sup> (50 ppm)	OSHA 2009 29 CFR 1910.1000, Table Z-1
b. Water			
EPA	Drinking water standards and health advisories	No	EPA 2006a
	National primary drinking water standards	No	EPA 2003
	National recommended water quality criteria	No	EPA 2006b
c. Food			
FDA	EAFUS <sup>d</sup>	No	FDA 2008
d. Other			
ACGIH	Carcinogenicity classification	No	ACGIH 2008
	Biological exposure indices (end of shift)		
	Carboxyhemoglobin in blood	3.5% hemoglobin	
	Carbon monoxide in end-exhaled air	23 mg/m <sup>3</sup> (20 ppm)	
EPA	Carcinogenicity classification	No	IRIS 2009
	RfC	No	
	RfD	No	
	Superfund, emergency planning, and community right-to-know		

## 8. REGULATIONS AND ADVISORIES

**Table 8-1. Regulations, Advisories, and Guidelines Applicable to Carbon Monoxide**

Agency	Description	Information	Reference
<u>NATIONAL</u> ( <i>cont.</i> )			
	Designated CERCLA hazardous substance	No	EPA 2009e 40 CFR 302.4
	Effective date of toxic chemical release reporting	No	EPA 2009f 40 CFR 372.65
NTP	Carcinogenicity classification	No	NTP 2005

<sup>a</sup>ERPG-1 is the maximum airborne concentration below which nearly all individuals could be exposed for up to 1 hour without experiencing other than mild, transient health effects; ERPG-2 is the maximum airborne concentration below which nearly all individuals could be exposed for up to 1 hour without experiencing irreversible or other serious adverse effects; and ERPG-3 is the maximum airborne concentration below which nearly all individuals could be exposed for up to 1 hour without life-threatening health effects (AIHA 2008).

<sup>b</sup>AEGL-1 is the airborne concentration of a substance above which it is predicted that the general population, including susceptible individuals, could experience notable discomfort, irritation, or certain asymptomatic nonsensory effects, however, the effects are not disabling and are transient and reversible upon cessation of exposure; AEGL-2 is the airborne concentration of a substance above which it is predicted that the general population, including susceptible individuals, could experience irreversible or other serious, long-lasting adverse health effects or an impaired ability to escape; and AEGL-3 is the airborne concentration of a substance above which it is predicted that the general population, including susceptible individuals, could experience life-threatening health effects or death (EPA 2009b).

<sup>c</sup>Not to be exceeded more than once per year.

<sup>d</sup>The EAFUS list of substances contains ingredients added directly to food that FDA has either approved as food additives or listed or affirmed as GRAS.

ACGIH = American Conference of Governmental Industrial Hygienists; AEGL = acute exposure guideline levels; AIHA = American Industrial Hygiene Association; CERCLA = Comprehensive Environmental Response, Compensation, and Liability Act; CFR = Code of Federal Regulations; EAFUS = Everything Added to Food in the United States; EPA = Environmental Protection Agency; ERPG = emergency response planning guidelines; FDA = Food and Drug Administration; GRAS = Generally Recognized As Safe; IARC = International Agency for Research on Cancer; IDLH = immediately dangerous to life or health; IRIS = Integrated Risk Information System; NIOSH = National Institute for Occupational Safety and Health; NTP = National Toxicology Program; OSHA = Occupational Safety and Health Administration; PEL = permissible exposure limit; REL = recommended exposure limit; RfC = inhalation reference concentration; RfD = oral reference dose; TLV = threshold limit values; TWA = time-weighted average; USC = United States Code; WHO = World Health Organization

## 9. REFERENCES

Abbatiello ER, Mohrmann K. 1979. Effects on the offspring of chronic low exposure carbon monoxide during mice pregnancy. *Clin Toxicol* 14(4):401-406.

Aberg AM. 2008. Carbon monoxide in biological systems: An experimental and clinical study. *Acta Anaesthesiol Scand* 52(5):716-717.

Aberg AM, Abrahamsson P, Johansson G, et al. 2008. Does carbon monoxide treatment alter cytokine levels after endotoxin infusion in pigs? A randomized controlled study. *J Inflamm (Lond)* 5:13.

\*Abi Esper L, El-Fadel M, Nuwayhid I, et al. 2007a. The effect of different ventilation modes on in-vehicle carbon monoxide exposure. *Atmos Environ* 41:3644-3657.

Abi Esper L, El-Fadel M, Shihadeh A. 2007b. Comparison of trip average in-vehicle and exterior CO determinations by continuous and grab sampling using an electrochemical sensing method. *Atmos Environ* 41:6087-6094.

\*Abram SR, Hodnett BL, Summers RL, et al. 2007. Quantitative circulatory physiology: An integrative mathematical model of human physiology for medical education. *Adv Physiol Educ* 31:202-210.

Acevedo CH, Ahmed A. 1998. Hemeoxygenase-1 inhibits human myometrial contractility via carbon monoxide and is upregulated by progesterone during pregnancy. *J Clin Invest* 101(5):949-955.

\*ACGIH. 2008. Carbon monoxide. In: *Threshold limit values for chemical substances and physical agents and biological exposure indices*. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.

Achouh PE, Simonet S, Fabiani JN, et al. 2008. Carbon monoxide induces relaxation of human internal thoracic and radial arterial grafts. *Interact Cardiovasc Thorac Surg* 7(6):959-962.

+\*Adams KF, Koch G, Chatterjee B, et al. 1988. Acute elevation of blood carboxyhemoglobin to 6% impairs exercise performance and aggravates symptoms in patients with ischemic heart disease. *J Am Coll Cardiol* 12(4):900-909.

\*Adinolfi M. 1985. The development of the human blood-CSF-brain barrier. *Dev Med Child Neurol* 27(4):532-537.

+\*Adir Y, Merdler A, Haim SB, et al. 1999. Effects of exposure to low concentrations of carbon monoxide on exercise performance and myocardial perfusion in young healthy men. *Occup Environ Med* 56:535-538.

\*Adlercreutz H. 1995. Phytoestrogens: Epidemiology and a possible role in cancer protection. *Environmental Health Perspectives Supplement* 103(7):103-112.

---

\* Cited in text

+ Cited in supplemental document

## 9. REFERENCES

- \*Agency for Toxic Substances and Disease Registry. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Fed Regist 54(174):37618-37634.
- Agency for Toxic Substances and Disease Registry. 1990. Biomarkers of organ damage or dysfunction for the renal, hepatobiliary, and immune systems. Subcommittee on Biomarkers of Organ Damage and Dysfunction. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- \*Agency for Toxic Substances and Disease Registry. 2000. Toxicological profile for methylene chloride. Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/toxprofiles/tp14.pdf>. June 11, 2009.
- \*Agency for Toxic Substances and Disease Registry. 2005. Toxicological profile for carbon tetrachloride. Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/toxprofiles/tp30.pdf>. June 11, 2009.
- Aguilaniu B, Maitre J, Glenet S, et al. 2008. European reference equations for CO and NO lung transfer. Eur Respir J 31(5):1091-1097.
- Ahmed A, Rahman M, Zhang X, et al. 2000. Induction of placental heme oxygenase-1 is protective against TNFalpha-induced cytotoxicity and promotes vessel relaxation. Mol Med 6(5):391-409.
- \*AIHA. 2008. Emergency Response Planning Guidelines (ERPG). Fairfax, VA: American Industrial Hygiene Association. <http://www.aiha.org/1documents/Committees/ERP-erpglevels.pdf>. May 19, 2009.
- \*Alcantara RE, Xu C, Spiro TG, et al. 2007. A quantum-chemical picture of hemoglobin affinity. Proc Natl Acad Sci USA 104(47):18451-18455.
- \*Alcaraz MJ, Fernandez P, Guillen MI. 2003. Anti-inflammatory actions of the heme oxygenase-1 pathway. Curr Pharm Des 9:2541-2551.
- Alderman BW, Baron AE, Savitz DA. 1987. Maternal exposure to neighborhood carbon monoxide and risk of low infant birth weight. Public Health Report 102(4):410-414.
- Alexander PG, Chau L, Tuan RS. 2007. Role of nitric oxide in chick embryonic organogenesis and dysmorphogenesis. Birth Defects Res A Clin Mol Teratol 79:581-594.
- Alexandreaanu IC, Lawson DM. 2002. Effects of chronic administration of a heme oxygenase substrate or inhibitor on progression of the estrous cycle, pregnancy and lactation of Sprague-Dawley rats. Life Sci 72:153-162.
- Alexandreaanu IC, Lawson DM. 2003. Heme oxygenase in the rat anterior pituitary: Immunohistochemical localization and possible role in gonadotropin and prolactin secretion. Exp Biol Med (Maywood) 228(1):64-69.
- +\*Allred EN, Bleecker ER, Chaitman BR, et al. 1989. Short-term effects of carbon monoxide exposure on the exercise performance of subjects with coronary artery disease. N Engl J Med 321(21):1426-1432.
- +\*Allred EN, Bleecker ER, Chaitman BR, et al. 1991. Effects of carbon monoxide on myocardial ischemia. Environ Health Perspect 91:89-132.

## 9. REFERENCES

- \*Alonso J, Cardellach F, Lopez S, et al. 2003. Carbon monoxide specifically inhibits cytochrome c oxidase of human mitochondrial respiratory chain. *Pharmacol Toxicol* 93:142-146.
- Althaus M, Fronius M, Buchackert Y, et al. 2009. Carbon monoxide rapidly impairs alveolar fluid clearance by inhibiting epithelial sodium channels. *Am J Respir Cell Mol Biol*
- \*Altman PL, Dittmer DS. 1974. Total body water: Man. In: Altman PL, Dittmer DS, eds. *Biological handbooks: Biology data book*. Vol. III. 2nd ed. Bethesda, MD: Federation of American Societies of Experimental Biology, 1987-2008, 2041.
- \*Andersen ME, Krishnan K. 1994. Relating *in vitro* to *in vivo* exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. *Animal test alternatives: Refinement, reduction, and replacement*. New York, NY: Marcel Dekker Inc., 9-25.
- \*Andersen ME, Clewell HJ, Gargas ML, et al. 1987. Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol Appl Pharmacol* 87(2):185-205.
- \*Andersen ME, Clewell HJ, Gargas ML, et al. 1991. Physiologically based pharmacokinetic modeling with dichloromethane, its metabolite, carbon monoxide, and blood carboxyhemoglobin in rats and humans. *Toxicol Appl Pharmacol* 108:14-27.
- +Andersen ZJ, Wahlin P, Raaschou-Nielsen O, et al. 2007. Ambient particle source apportionment and daily hospital admissions among children and elderly in Copenhagen. *J Expo Sci Environ Epidemiol* 17:625-636.
- Anderson ME, Bogdan GM. 2007. Environments, indoor air quality, and children. *Pediatr Clin North Am* 54(2):295-307.
- +\*Anderson EW, Andelman RJ, Strauch JM, et al. 1973. Effect of low-level carbon monoxide exposure on onset and duration of angina pectoris. *Ann Intern Med* 79:46-50.
- +Anderson HR, Bremner SA, Atkinson RW, et al. 2001. Particulate matter and daily mortality and hospital admissions in the west midlands conurbation of the United Kingdom: Associations with fine and coarse particles, black smoke and sulphate. *Occup Environ Med* 58:504-510.
- Andersson JA, Uddman R, Cardell LO. 2000. Carbon monoxide is endogenously produced in the human nose and paranasal sinuses. *J Allergy Clin Immunol* 105(2 Pt 1):269-273.
- Andresen JJ, Shafi NI, Durante W, et al. 2006. Effects of carbon monoxide and heme oxygenase inhibitors in cerebral vessels of rats and mice. *Am J Physiol Heart Circ Physiol* 291:H223-H230.
- Antonelli T, Tomasini MC, Tatoli M, et al. 2006. Prenatal exposure to the cannabinoid receptor agonist WIN 55,212-2 and carbon monoxide reduces extracellular glutamate levels in primary rat cerebral cortex cell cultures. *Neurochem Int* 49:568-576.
- \*API. 1994. A study to characterize air concentrations of methyl tertiary butyl ether (MTBE) at service stations in the northeast. American Petroleum Institute. API publication number 4619.
- Appleton SD, Marks GS, Nakatsu K, et al. 2002. Heme oxygenase activity in placenta: Direct dependence on oxygen availability. *Am J Physiol Heart Circ Physiol* 282:H2055-H2059.

## 9. REFERENCES

- Arranz B, Blennow K, Eriksson A, et al. 1997. Serotonergic, noradrenergic, and dopaminergic measures in suicide brains. *Biol Psychiatry* 41:1000-1009.
- \*Arruda MA, Rossi AG, de Freitas MS, et al. 2004. Heme inhibits human neutrophil apoptosis: Involvement of phosphoinositide 3-kinase, MAPK, and NF- $\kappa$ B. *J Immunol* 173:2023-2030.
- +\*Astrup P, Olsen HM, Trolle D, et al. 1972. Effect of moderate carbon-monoxide exposure on fetal development. *Lancet* 2(7789):1220-1222.
- +Atkinson RW, Anderson HR, Strachan DP, et al. 1999. Short-term associations between outdoor air pollution and visits to accident and emergency departments in London for respiratory complaints. *Eur Respir J* 13:257-265.
- Ator NA, Merigan WHJ, McIntire RW. 1976. The effects of brief exposures to carbon monoxide on temporally differentiated responding. *Environ Res* 12:81-91.
- Ayres SM, Giannelli SJ, Mueller H, et al. 1973. Myocardial and systemic vascular response to low concentrations of carboxyhemoglobin. *Ann Clin Lab Sci* 3(6):440-447.
- +\*Baccarelli A, Zanobetti A, Martinelli I, et al. 2007. Effects of exposure to air pollution on blood coagulation. *J Thromb Haemost* 5(2):252-260.
- Bainbridge SA, Smith GN. 2005. HO in pregnancy. *Free Radic Biol Med* 38(8):979-988.
- Bainbridge SA, Belkacemi L, Dickinson M, et al. 2006. Carbon monoxide inhibits hypoxia/reoxygenation-induced apoptosis and secondary necrosis in syncytiotrophoblast. *Am J Pathol* 169(3):774-783.
- Bainbridge SA, Farley AE, McLaughlin BE, et al. 2002. Carbon monoxide decreases perfusion pressure in isolated human placenta. *Placenta* 23:563-569.
- Bakwin PS, Tans PP. 1994. Carbon monoxide budget in the Northern Hemisphere. *Geophys Res Lett* 21(6):433-436.
- \*Baldauf R, Thoma E, Khlystov A, et al. 2008. Impacts of noise barriers on near-road air quality. *Atmos Environ* 42:7502-7507.
- +Ballester F, Rodriguez P, Iniguez C, et al. 2006. Air pollution and cardiovascular admissions association in Spain: Results within the EMECAS project. *J Epidemiol Comm Health* 60:328-336.
- +\*Ballester F, Tenias JM, Perez-Hoyos S. 2001. Air pollution and emergency hospital admissions for cardiovascular diseases in Valencia, Spain. *J Epidemiol Community Health* 55:57-65.
- Bamberger AM, Koglin M, Kempfert J, et al. 2001. Expression and tissue localization of soluble guanylyl cyclase in the human placenta using novel antibodies directed against the  $\alpha_2$  subunit. *J Clin Endocrinol Metab* 86(2):909-912.
- Barber A, Robson SC, Lyall F. 1999. Hemoxygenase and nitric oxide synthase do not maintain human uterine quiescence during pregnancy. *Am J Pathol* 155:831-840.

## 9. REFERENCES

- Barber A, Robson SC, Myatt L, et al. 2001. Heme oxygenase expression in human placenta and placental bed: Reduced expression of placenta endothelial HO-2 in preeclampsia and fetal growth restriction. *FASEB J* 15(7):1158-1168.
- Barie PS, Wu W, Hariri RJ, et al. 1994. Alterations of pulmonary gas exchange after superimposed carbon monoxide poisoning in acute lung injury. *Surgery* 115(6):678-686.
- Barlow SM, Sullivan FM. 1982. Carbon monoxide. In: *Reproductive hazards of industrial chemicals. An evaluation of animal and human data.* New York, NY: Academic Press, Inc., 178-199.
- \*Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. *Regul Toxicol Pharmacol* 8(4):471-486.
- +\*Barnett AG, Williams GM, Schwartz J, et al. 2006. The effects of air pollution on hospitalizations for cardiovascular disease in elderly people in Australian and New Zealand cities. *Environ Health Perspect* 114(7):1018-1023.
- Bassan M. 1990. Sudden cardiac death [Letter to the editor]. *N Engl J Med* 322:272.
- Bates TS, Kelly KC, Johnson JE, et al. 1995. Regional and seasonal variations in the flux of oceanic carbon monoxide to the atmosphere. *J Geophys Res* 100(D11):23093-23101.
- Bathoorn E, Slebos DJ, Postma DS, et al. 2007. Anti-inflammatory effects of inhaled carbon monoxide in patients with COPD: A pilot study. *Eur Respir J* 30:1131-1137.
- Baum M, Schiff E, Krieser D, et al. 2000. End-tidal carbon monoxide measurements in women with pregnancy-induced hypertension and preeclampsia. *Am J Obstet Gynecol* 183:900-903.
- Beard RR, Wertheim GA. 1967. Behavioral impairment associated with small doses of carbon monoxide. *Am J Public Health Nations Health* 57(11):2012-2022.
- +Bedeschi E, Campari C, Candela S, et al. 2007. Urban air pollution and respiratory emergency visits at pediatric unit, Reggio Emilia, Italy. *J Toxicol Environ Health A* 70:261-265.
- +\*Bell ML, Ebisu K, Belanger K. 2007. Ambient air pollution and low birth weight in Connecticut and Massachusetts. *Environ Health Perspect* 115(7):1118-1124.
- +Bell ML, Levy JK, Lin Z. 2008. The effect of sandstorms and air pollution on cause-specific hospital admissions in Taipei, Taiwan. *Occup Environ Med* 65(2):104-111.
- +Bellini P, Baccini M, Biggeri A, et al. 2007. The meta-analysis of the Italian studies on short-term effects of air pollution (MISA): Old and new issues on the interpretation of the statistical evidences. *Environmetrics* 18:219-229.
- \*Benignus VA. 1994. Behavioral effects of carbon monoxide: Meta analyses and extrapolations. *J Appl Physiol* 76(3):1310-1316
- \*Benignus VA. 1995. A model to predict carboxyhemoglobin and pulmonary parameters after exposure to O<sub>2</sub>, CO<sub>2</sub>, and CO. *Aviat Space Environ Med* 66(4):369-374.

## 9. REFERENCES

- \*Benignus VA, Annau Z. 1994. Carboxyhemoglobin formation due to carbon monoxide exposure in rats. *Toxicol Appl Pharmacol* 128(1):151-157.
- Benignus VA, Coleman T, Eklund CR, et al. 2006. A general physiological and toxicokinetic (GPAT) model for simulating complex toluene exposure scenarios in humans. *Toxicol Mech Methods* 16:27-36.
- \*Benignus VA, Hazucha MJ, Smith MV, et al. 1994. Prediction of carboxyhemoglobin formation due to transient exposure to carbon monoxide. *J Appl Physiol* 76:1739-1745.
- Benignus VA, Muller KE, Barton CN, et al. 1987. Effect of low level carbon monoxide on compensatory tracking and event monitoring. *Neurotoxicol Teratol* 9(3):227-234.
- \*Benignus VA, Muller KE, Malott CM. 1990. Dose-effects functions for carboxyhemoglobin and behavior. *Neurotoxicol Teratol* 12(2):111-118.
- \*Benowitz NL. 1997. The role of nicotine in smoking-related cardiovascular disease. *Prev Med* 26:412-417.
- \*Berger GS, ed. 1994. Epidemiology of endometriosis. In: *Endometriosis: Modern surgical management of endometriosis*. New York, NY: Springer-Verlag, 3-7.
- +\*Berger A, Zareba W, Schneider A, et al. 2006. Runs of ventricular and supraventricular tachycardia triggered by air pollution in patients with coronary heart disease. *JOEM* 48(11):1149-1158.
- Bergeron M, Ferriero DM, Sharp FR. 1998. Developmental expression of heme oxygenase-1 (HSP32) in rat brain: An immunocytochemical study. *Dev Brain Res* 105:181-194.
- \*Berk PD, Blaschke TF, Scharschmidt BF, et al. 1976. A new approach to quantitation of the various sources of bilirubin in man. *J Lab Clin Med* 87(5):767-780.
- \*Berk PD, Rodkey FL, Blaschke TF, et al. 1974. Comparison of plasma bilirubin turnover and carbon monoxide production in man. *J Lab Clin Med* 83(1):29-37.
- Bernard TE, Duker J. 1981. Modeling carbon monoxide uptake during work. *Am Ind Hyg Assoc J* 42(5):361-364.
- Bessoudo R, Gray J. 1978. Carbon monoxide poisoning and nonoliguric acute renal failure. *Can Med Assoc J* 119(1):41-44.
- \*Bhuyan AK, Kumar R. 2002. Kinetic barriers to the folding of horse cytochrome c in the reduced state. *Biochemistry* 41:12821-12834.
- +Biggeri A, Baccini M, Bellini P, et al. 2005. Meta-analysis of the Italian studies of short-term effects of air pollution (MISA), 1990-1999. *Int J Occup Environ Health* 11(1):107-122.
- Borrego C, Tchepel O, Costa AM, et al. 2006. Traffic-related particulate air pollution exposure in urban areas. *Atmos Environ* 40:7205-7214.
- \*Botros FT, Laniado-Schwartzman M, Abraham NG. 2002. Regulation of cyclooxygenase- and cytochrome p450-derived eicosanoids by heme oxygenase in the rat kidney. *Hypertension* 39:639-644.

## 9. REFERENCES

- +Botter DA, Jorgensen B, Peres AA. 2002. A longitudinal study of mortality and air pollution for Sao Paulo, Brazil. *J Expo Anal Environ Epidemiol* 12(5):335-343.
- Boyes WK, Griffin VT. 1995. Acute effects of inhaled trichloroethylene (TCE) on rat visual function depend on exposure concentration and not duration. *Neurotoxicology* 16(3):543.
- +Braga AL, Saldiva PH, Pereira LA, et al. 2001. Health effects of air pollution exposure on children and adolescents in Sao Paulo, Brazil. *Pediatr Pulmonol* 31(2):106-113.
- +\*Brauer M, Lencar C, Tamburic L, et al. 2008. A cohort study of traffic-related air pollution impacts on birth outcomes. *Environ Health Perspect* 116(5):680-686.
- +Bremner SA, Anderson HR, Atkinson RW, et al. 1999. Short-term associations between outdoor air pollution and mortality in London 1992-4. *Occup Environ Med* 56(4):237-244.
- Brian JE, Heistad DD, Faraci FM. 1994. Effect of carbon monoxide on rabbit cerebral arteries. *Stroke* 25(3):639-643.
- \*Brown SD, Piantadosi CA. 1992. Recovery of energy metabolism in rat brain after carbon monoxide hypoxia. *J Clin Invest* 89:666-672.
- \*Brown DB, Mueller GL, Golich FC. 1992. Hyperbaric oxygen treatment for carbon monoxide poisoning in pregnancy: A case report. *Aviat Space Environ Med* 63:1011-1014.
- \*Bruce EN, Bruce MC. 2003. A multicompartiment model of carboxyhemoglobin and carboxymyoglobin responses to inhalation of carbon monoxide. *J Appl Physiol* 95:1235-1247.
- \*Bruce MC, Bruce EN. 2006. Analysis of factors that influence rates of carbon monoxide uptake, distribution, and washout from blood and extravascular tissues using a multicompartiment model. *J Appl Physiol* 100:1171-1180.
- \*Bruce EN, Bruce MC, Erupaka K. 2008. Prediction of the rate of uptake of carbon monoxide from blood by extravascular tissues. *Respir Physiol Neurobiol* 161(2):142-159.
- \*Bruinen De Bruin Y, Carrer P, Jantunen M, et al. 2004a. Personal carbon monoxide exposure levels: Contribution of local sources to exposures and microenvironment concentrations in Milan. *J Expo Anal Environ Epidemiol* 14:312-322.
- \*Bruinen De Bruin Y, Hanninen O, Carrer P, et al. 2004b. Simulation of working population exposures to carbon monoxide using EXPOLIS-Milan microenvironment concentration and time-activity data. *J Expo Anal Environ Epidemiol* 14:154-163.
- \*Brune B, Ullrich V. 1987. Inhibition of platelet aggregation by carbon monoxide is mediated by activation of guanylate cyclase. *Mol Pharmacol* 32(3):497-504.
- \*Brvar M, Mozina H, Osredkar J, et al. 2004. S100B protein in carbon monoxide poisoning: A pilot study. *Resuscitation* 61:357-360.
- \*Burge CM, Skinner SL. 1995. Determination of hemoglobin mass and blood volume with CO: Evaluation and application of a method. *J Appl Physiol* 79(2):623-631.

## 9. REFERENCES

- \*Burmester T, Weich B, Reinhardt S, et al. 2000. A vertebrate globin expressed in the brain. *Nature* 407(6803):520-523.
- +Burnett RT, Brook J, Dann T, et al. 2000. Association between particulate- and gas-phase components of urban air pollution and daily mortality in eight Canadian cities. *Inhal Toxicol* 12(Suppl 4):15-39.
- +Burnett RT, Smith-Doiron M, Stieb D, et al. 1999. Effects of particulate and gaseous air pollution on cardiorespiratory hospitalizations. *Arch Environ Health* 54(2):130-139.
- +\*Burnett RT, Smith-Doiron M, Stieb D, et al. 2001. Association between ozone and hospitalization for acute respiratory diseases in children less than 2 years of age. *Am J Epidemiol* 153(5):444-452.
- +\*Burnett RT, Stieb D, Brook JR, et al. 2004. Associations between short-term changes in nitrogen dioxide and mortality in Canadian cities. *Arch Environ Health* 59(5):228-236.
- +\*Cagiano R, Ancona D, Cassano T, et al. 1998. Effects of prenatal exposure to low concentrations of carbon monoxide on sexual behaviour and mesolimbic dopaminergic function in rat offspring. *Br J Pharmacol* 125:909-915.
- \*Cairns J, Denhardt DT. 1968. Effect of cyanide and carbon monoxide on the replication of bacterial DNA *in vivo*. *J Mol Biol* 36:335-342.
- +Cakmak S, Dales RE, Blanco Vidal C. 2007. Air pollution and mortality in Chile: Susceptibility among the elderly. *Environ Health Perspect* 115(4):524-527.
- +\*Cakmak S, Dales RE, Judek S. 2006. Do gender, education, and income modify the effect of air pollution gases on cardiac disease? *JOEM* 48(1):89-94.
- \*Campbell ME, Benson BA, Muir MA. 1995. Urban air quality and human health: A Toronto perspective. *Can J Public Health* 86(5):351-357.
- \*Caravati EM, Adams CJ, Joyce SM, et al. 1988. Fetal toxicity associated with maternal carbon monoxide poisoning. *Ann Emerg Med* 17(7):714-717.
- Cardell L. 1988. Bronchodilatation *in vivo* by carbon monoxide, a cyclic GMP related messenger. *Br J Pharmacol* 124:1065-1068.
- \*Carlsson GH, Nicholls P, Svistunenko D, et al. 2005. Complexes of horseradish peroxidase with formate, acetate and carbon monoxide. *Biochemistry* 44:635-642.
- +\*Carmines EL, Rajendran N. 2008. Evidence for carbon monoxide as the major factor contributing to lower fetal weights in rats exposed to cigarette smoke. *Toxicol Sci* 102(2):383-391.
- Carratu MR, Cagiano R, De Salvia MA, et al. 1995. Developmental neurotoxicity of carbon monoxide. *Arch Toxicol Suppl* 17:295-301.
- +\*Carratu MR, Cagiano R, Desantis S, et al. 2000a. Prenatal exposure to low levels of carbon monoxide alters sciatic nerve myelination in rat offspring. *Life Sci* 67:1759-1772.

## 9. REFERENCES

- +\*Carratu MR, Cagiano R, Tattoli M, et al. 2000b. Prenatal exposure model simulating CO inhalation in human cigarette smokers: Sphingomyelin alterations in the rat sciatic nerve. *Toxicol Lett* 117(1-2):101-106.
- +\*Carratu MR, Renna G, Giustino A, et al. 1993. Changes in peripheral nervous system activity produced in rats by prenatal exposure to carbon monoxide. *Arch Toxicol* 67:297-301.
- Carraway MS, Ghio AJ, Carter JD, et al. 2000. Expression of heme oxygenase-1 in the lung in chronic hypoxia. *Am J Physiol Lung Cell Mol Physiol* 278:L806-L812.
- \*Carraway MS, Ghio AJ, Suliman HB, et al. 2002. Carbon monoxide promotes hypoxic pulmonary vascular remodeling. *Am J Physiol Lung Cell Mol Physiol* 282:L693-L702.
- \*Castillo A, Llapur CJ, Martinez T, et al. 2006. Measurement of single breath-hold carbon monoxide diffusing capacity in healthy infants and toddlers. *Pediatr Pulmonol* 41(6):544-550.
- Cavallin-Stahl E, Mercke C, Lundh B. 1976a. Carbon monoxide production in patients with breast carcinoma. *Br J Haematol* 32(2):177-182.
- Cavallin-Stahl E, Mercke C, Lundh B. 1976b. Erythropoiesis and carbon monoxide production in Hodgkin's disease. *Br J Haematol* 32(2):167-175.
- \*CDC. 2007. Carbon monoxide-related deaths-United States, 1999-2004. *Morb Mortal Wkly Rep* 56(50):1309-1312. <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5650a1.htm>. August 5, 2009.
- \*CDC. 2008. Nonfatal, unintentional, non fire-related carbon monoxide exposures United States, 2004-2006. *JAMA* 300(20):2362-2363.
- Cella M, Farina MG, Keller Sarmiento MI, et al. 2006. Heme oxygenase-carbon monoxide (HO-CO) system in rat uterus: Effect of sexual steroids and prostaglandins. *J Steroid Biochem Mol Biol* 99:59-66.
- Cepinskas G, Katada K, Bihari A, et al. 2008. Carbon monoxide liberated from carbon monoxide-releasing molecule CORM-2 attenuates inflammation in the liver of septic mice. *Am J Physiol Gastrointest Liver Physiol* 294(1):G184-191.
- \*Chakraborty S, Balakotalah V, Bidani A. 2004. Diffusing capacity reexamined: Relative roles of diffusion and chemical reaction in red cell uptake of O<sub>2</sub>, CO, CO<sub>2</sub>, and NO. *J Appl Physiol* 97:2284-2302.
- \*Chambers CA, Hopkins RO, Weaver LK, et al. 2008. Cognitive and affective outcomes of more severe compared to less severe carbon monoxide poisoning. *Brain Inj* 22(5):387-395.
- +\*Chan C, Chuang K, Chien L, et al. 2006. Urban air pollution and emergency admissions for cerebrovascular diseases in Taipei, Taiwan. *Eur Heart J* 27:1238-1244.
- +\*Chan C, Chuang K, Su T, et al. 2005. Association between nitrogen dioxide and heart rate variability in susceptible population. *Eur J Cardiovasc Prev Rehabil* 12:580-586.
- +\*Chang C, Tsai S, Ho S, et al. 2005. Air pollution and hospital admissions for cardiovascular disease in Taipei, Taiwan. *Environ Res* 98:114-119.

## 9. REFERENCES

- \*Chang LT, Koutrakis P, Catalano PJ, et al. 2000. Hourly personal exposures to fine particles and gaseous pollutants—results from Baltimore, Maryland. *J Air Waste Manage Assoc* 50(7):1223-1235.
- ChemIDPlus. 2009. Carbon monoxide. ChemIDplus. Bethesda, MD: U.S. National Library of Medicine. <http://sis.nlm.nih.gov/chemical.html>. June 9, 2009.
- \*Chen GD, Fechter LD. 1999. Potentiation of octave-band noise induced auditory impairment by carbon monoxide. *Hear Res* 132(1-2):149-159.
- \*Chen H, Namedo A, Bell M. 2008. Classification of road traffic and roadside pollution concentrations for assessment of personal exposure. *Environ Modell Software* 23:282-287.
- +Chen L, Jennison BL, Yang W, et al. 2000. Elementary school absenteeism and air pollution. *Inhal Toxicol* 12:997-1016.
- +\*Chen L, Yang W, Jennison BL, et al. 2002. Air pollution and birth weight in Northern Nevada, 1991-1999. *Inhal Toxicol* 14(2):141-157.
- +\*Chen PC, Lai YM, Chan CC, et al. 1999. Short-term effect of ozone on the pulmonary function of children in primary school. *Environ Health Perspect* 107(11):921-925.
- Chen S, Weller MA, Penney DG. 1982. A study of free lung cells from young rats chronically exposed to carbon monoxide from birth. *Scan Electron Microsc (Pt 2)*:859-867.
- +Cheng M, Tsai S, Wu T, et al. 2007. Air pollution and hospital admissions for pneumonia in a tropical city: Kaohsiung, Taiwan. *J Toxicol Environ Health A* 70:2021-2026.
- \*Chevalier RB, Krumholz RA, Ross JC. 1966. Reaction of nonsmokers to carbon monoxide inhalation. Cardiopulmonary responses at rest and during exercise. *JAMA* 198(10):1061-1064.
- +Cho B, Choi J, Yum Y. 2000. Air pollution and hospital admissions for respiratory disease in certain area of Korea. *J Occup Health* 42:185-191.
- +Chock DP, Winkler SL, Chen C. 2000. A study of the association between daily mortality and ambient air pollutant concentrations in Pittsburgh, Pennsylvania. *J Air Waste Manage Assoc* 50:1481-1500.
- \*Choi IS. 2001. Carbon monoxide poisoning: Systemic manifestations and complications. *J Korean Med Sci* 16(3):253-261.
- \*Choi IS. 2002. Parkinsonism after carbon monoxide poisoning. *Eur Neurol* 48:30-33.
- Chung Y, Huang S, Glabe A, et al. 2006. Implication of CO in activation on myoglobin function. *Am J Physiol Cell Physiol* 290:C1616-C1624.
- +Cifuentes LA, Vega J, Kopfer K, et al. 2000. Effect of the fine fraction of particulate matter versus the coarse mass and other pollutants on daily mortality in Santiago, Chile. *J Air Waste Manage Assoc* 50:1287-1298.
- \*Clewell HJ, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.

## 9. REFERENCES

- Coburn RF. 1967. Endogenous carbon monoxide production and body CO stores. *Acta Med Scand Suppl* 472:269-282.
- \*Coburn RF. 1970a. Enhancement by phenobarbital and diphenylhydantoin of carbon monoxide production in normal man. *N Engl J Med* 283(10):512-515.
- Coburn RF. 1970b. The carbon monoxide body stores. *Ann N Y Acad Sci* 174:11-22.
- \*Coburn RF, Mayers LB. 1971. Myoglobin oxygen tension determined from measurements of carboxymyoglobin in skeletal muscle. *Am J Physiol* 220:66-74.
- \*Coburn RF, Blakemore WS, Forster RE. 1963. Endogenous carbon monoxide production in man. *J Clin Invest* 42(7):1172-1178.
- \*Coburn RF, Forster RE, Kane PB. 1965. Considerations of the physiological variables that determine the blood carboxyhemoglobin concentration in man. *J Clin Invest* 44(11):1899-1910.
- \*Coburn RF, Williams WJ, Forster RE. 1964. Effect of erythrocyte destruction on carbon monoxide production in man. *J Clin Invest* 43(6):1098-1103.
- Coburn RF, Williams WJ, Kahn SB. 1966. Endogenous carbon monoxide production in patients with hemolytic anemia. *J Clin Invest* 45(4):460-468.
- Cocconi F, Breen CA, Lees JG, et al. 1988. Further evidence implicating a cytochrome P-450-mediated reaction in the contractile tension of the lamb ductus arteriosus. *Circ Res* 62:471-477.
- +Conceicao GM, Miraglia SGEK, Kishi HS, et al. 2001. Air pollution and child mortality: A time-series study in Sao Paulo, Brazil. *Environ Health Perspect* 109(Suppl 3):347-350.
- Conrad R, Seiler W. 1985. Influence of temperature, moisture, and organic carbon on the flux of H<sub>2</sub> and CO between soil and atmosphere: Field studies in subtropical regions. *J Geophys Res* 90(D3):5699-5709.
- \*Conrad R, Seiler W. 1988. Influence of the surface microlayer on the flux of nonconservative trace gases (CO, H<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O) across the ocean-atmosphere interface. *J Atmos Chem* 6:83-94.
- \*Cramer CR. 1982. Fetal death due to accidental maternal carbon monoxide poisoning. *J Toxicol Clin Toxicol* 19(3):297-301.
- \*Cronje FJ, Carraway MS, Freiburger JJ, et al. 2004. Carbon monoxide actuates O<sub>2</sub>-limited heme degradation in the rat brain. *Free Radic Biol Med* 37(11):1802-1812.
- Croxford B, Leonardi GS, Kreis I. 2008. Self-reported neurological symptoms in relation to CO emissions due to problem gas appliance installations in London: A cross-sectional survey. *Environ Health* 7:34.
- Crutzen PJ, Delany AC, Greenberg J, et al. 1985. Tropospheric chemical composition measurements in Brazil during the dry season. *J Atmos Chem* 2:233-256.
- Cudmore M, Ahmad S, Al-Ani B, et al. 2007. Negative regulation of soluble Flt-1 and soluble endoglin release by heme oxygenase-1. *Circulation* 115:1789-1797.

## 9. REFERENCES

- +\*Dahms TE, Younis LT, Wiens RD, et al. 1993. Effects of carbon monoxide exposure in patients with documented cardiac arrhythmias. *J Am Coll Cardiol* 21(2):442-450.
- +\*Dales R. 2004. Ambient carbon monoxide may influence heart rate variability in subjects with coronary artery disease. *JOEM* 46(12):1217-1221.
- \*D'Amico G, Lam F, Hagen T, et al. 2006. Inhibition of cellular respiration by endogenously produced carbon monoxide. *J Cell Sci* 119:2291-2298.
- Dani C, Giannini L, Bertini G, et al. 2007. Changes of nitric oxide, carbon monoxide and oxidative stress in term infants at birth. *Free Radic Res* 41(12):1358-1363.
- \*Dart RC, ed. 2004. *Medical toxicology*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins, 1208.
- \*Davidson SB, Penney DG. 1988. Time course of blood volume change with carbon monoxide inhalation and its contribution to the overall cardiovascular response. *Arch Toxicol* 61:306-313.
- \*Davies B, Morris T. 1993. Physiological parameters in laboratory animals and humans. *Pharm Res* 10(7):1093-1095.
- +\*Davies DM, Smith DJ. 1980. Electrocardiographic changes in healthy men during continuous low-level carbon monoxide exposure. *Environ Res* 21(1):197-206.
- Davies RF, Topping DL, Turner DM. 1976. The effect of intermittent carbon monoxide exposure on experimental atherosclerosis in the rabbit. *Atherosclerosis* 24:527-536.
- \*Davutoglu V, Gunay N, Kocoglu H, et al. 2006. Serum levels of NT-ProBNP as an early cardiac marker of carbon monoxide poisoning. *Inhal Toxicol* 18:155-158.
- +de Hartog JJ, Hoek G, Peters A, et al. 2003. Effects of fine and ultrafine particles on cardiorespiratory symptoms in elderly subjects with coronary heart disease. *Am J Epidemiol* 157(7):613-623.
- +\*De Leon SF, Thurston GD, Ito K. 2003. Contribution of respiratory disease to nonrespiratory mortality associations with air pollution. *Am J Respir Crit Care Med* 167:1117-1123.
- +Delfino RJ, Gong H, Jr., Linn WS, et al. 2003. Asthma symptoms in Hispanic children and daily ambient exposures to toxic and criteria air pollutants. *Environ Health Perspect* 111:647-656.
- \*Delivoria-Papadopoulos M, Coburn RF, Forster RE. 1974. Cyclic variation of rate of carbon monoxide production in normal women. *J Appl Physiol* 36(1):49-51.
- \*Denninger JW, Marletta MA. 1999. Guanylate cyclase and the NO/cGMP signaling pathway. *Biochim Biophys Acta* 1411(2-3):334-350.
- Denschlag D, Marculescu R, Unfried G, et al. 2004. The size of a microsatellite polymorphism of the haem oxygenase 1 gene is associated with idiopathic recurrent miscarriage. *Mol Hum Reprod* 10(3):211-214.

## 9. REFERENCES

- +\*De Salvia MA, Cagiano R, Carratu MR, et al. 1995. Irreversible impairment of active avoidance behavior in rats prenatally exposed to mild concentrations of carbon monoxide. *Psychopharmacology* 122:66-71.
- Dewilde S, Kiger L, Burmester T, et al. 2001. Biochemical characterization and ligand binding properties of neuroglobin, a novel member of the globin family. *J Biol Chem* 276(42):38949-38955.
- \*Di Cera E, Doyle ML, Morgan MS, et al. 1989. Carbon monoxide and oxygen binding to human hemoglobin FO. *Biochemistry* 28(6):2631-2638.
- Di Giovanni V, Cagiano R, De Salvia MA, et al. 1993. Neurobehavioral changes produced in rats by prenatal exposure to carbon monoxide. *Brain Res* 616(1-2):126-131.
- \*Dioum EM, Rutter J, Tuckerman JR, et al. 2002. NPAS2: A gas-responsive transcription factor. *Science* 298:2385-2387.
- +\*D'Ippoliti D, Forastiere F, Ancoma C, et al. 2003. Air pollution and myocardial infarction in Rome. *Epidemiology* 14(5):528-535.
- \*Doblar DD, Santiago TV, Edelman NH. 1977. Correlation between ventilatory and cerebrovascular responses to inhalation of CO. *J Appl Physiol* 43(3):455-462.
- \*Dobos K, Zimmer G. 1985. Performance of carbon monoxide-sensitive MOSFET's with metal-oxide semiconductor gates. *IEEE Trans Electron Devices* ED-32(7):1165-1169.
- +\*Dockery DW, Luttmann-Gibson H, Rich DQ, et al. 2005. Association of air pollution with increased incidence of ventricular tachyarrhythmias recorded by implanted cardioverter defibrillators. *Environ Health Perspect* 113(6):670-674.
- \*Dolan MC. 1985. Carbon monoxide poisoning. *Can Med Assoc J* 133:392-399.
- +\*Dominici F, McDermott A, Daniels M, et al. 2003b. Mortality among residents of 90 cities. Revised analyses of time-series studies of air pollution and health. Special report. Health Effects Institute. <http://pubs.healtheffects.org/getfile.php?u=21>. October 13, 2009.
- \*Dominici F, Sheppard L, Clyde M. 2003a. Health effects of air pollution: A statistical review. *Int Stat Rev* 71:243-276.
- Doolittle DJ, Rahn CA, Lee CK. 1991. The effect of exposure to nicotine, carbon monoxide, cigarette smoke or cigarette smoke condensate on the mutagenicity of rat urine. *Mutat Res* 260(1):9-18.
- Drinkwater BL, Raven PB, Horvath SM, et al. 1974. Air pollution, exercise, and heat stress. *Arch Environ Health* 28(4):177-181.
- Dubuis E, Gautier M, Melin A, et al. 2002. Chronic carbon monoxide enhanced IbTx-sensitive currents in rat resistance pulmonary artery smooth muscle cells. *Am J Physiol Cell Physiol* 283:L120-L129.
- Dubuis E, Gautier M, Melin A, et al. 2003. Chronic carbon monoxide exposure of hypoxic rats increases *in vitro* sensitivity of pulmonary artery smooth muscle. *Can J Physiol Pharmacol* 81:711-719.

## 9. REFERENCES

- Dubuis E, Potier M, Wang R, et al. 2005. Continuous inhalation of carbon monoxide attenuates hypoxic pulmonary hypertension development presumably through activation of BKCa channels. *Cardiovasc Dis* 65:751-761.
- \*Duci A, Chaloulakou A, Spyrellis N. 2003. Exposure to carbon monoxide in the Athens urban area during commuting. *Sci Total Environ* 309(1-3):47-58.
- Duenas-Laita A, Mena-Martin FJ, Roquelai-Ruiz P, et al. 2008. Ischemic colitis associated with acute carbon monoxide poisoning. *Clin Toxicol (Phila)* 46(8):780-781.
- \*Dutton SJ, Hannign MP, Miller SL. 2001. Indoor pollutant levels from the use of unvented natural gas fireplaces in Boulder, Colorado. *J Air Waste Manage Assoc* 51:1654-1661.
- Ekblom B, Huot R, Stein EM, et al. 1975. Effect of changes in arterial oxygen content on circulation and physical performance. *J Appl Physiol* 39(1):71-75.
- \*Elkharrat E, Raphael JC, Korach JM, et al. 1991. Acute carbon monoxide intoxication and hyperbaric oxygen in pregnancy. *Intensive Care Med* 17(5):289-292.
- Ely EW, Moorehead B, Haponik EF. 1995. Warehouse workers' headache: Emergency evaluation and management of 30 patients with carbon monoxide poisoning. *Am J Med* 98(2):145-155.
- \*Engel RR, Rodkey FL, O'Neal JD, et al. 1969. Relative affinity of human fetal hemoglobin for carbon monoxide and oxygen. *Blood* 33(1):37-45.
- EPA. 1990. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment.
- \*EPA. 1991. Air quality criteria for carbon monoxide. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA600/8-90/045F.
- \*EPA. 1997. Special report on environmental endocrine disruption: An effects assessment and analysis. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum.
- EPA. 1998. Automated Form R for Windows: User's guide (RY97). Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics.
- \*EPA. 2000. Air quality criteria for carbon monoxide. Washington, DC: U.S. Environmental Protection Agency. EPA600P66001F. <http://www.epa.gov/NCEA/pdfs/coaqcd.pdf>. June 9, 2009.
- \*EPA. 2003. National primary drinking water regulations. Washington, DC: Office of Ground Water and Drinking Water, U.S. Environmental Protection Agency. <http://www.epa.gov/safewater/contaminants/index.html>. May 19, 2009.
- \*EPA. 2005. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the emergency planning and community right-to-know act (Title III of the Superfund Amendments and Reauthorization Act of 1986). U.S. Environmental Protection Agency, Office of Environmental Information.

## 9. REFERENCES

- \*EPA. 2006a. Drinking water standards and health advisories. Washington, DC: U.S. Environmental Protection Agency, Office of Water. EPA822R04005. <http://epa.gov/waterscience/criteria/drinking/>. May 19, 2009.
- \*EPA. 2006b. National recommended water quality criteria. Washington, DC: U.S. Environmental Protection Agency, Office of Water, Office of Science and Technology. <http://www.epa.gov/waterscience/criteria/wqcriteria.html>. May 11, 2009.
- \*EPA. 2008. EPA's 2008 report on the environment. Washington, DC: U.S. Environmental Protection Agency. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=190806>. July 16, 2009.
- \*EPA. 2009a. An introduction to indoor air quality. Carbon monoxide. U.S. Environmental Protection Agency. <http://www.epa.gov/iaq/co.html>. May 8, 2009.
- \*EPA. 2009b. Acute exposure guideline levels (AEGs). Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/oppt/aegl/>. May 19, 2009.
- \*EPA. 2009c. Hazardous air pollutants. Clean Air Act. U.S. Environmental Protection Agency. United States Code. 42 USC 7412. <http://www.epa.gov/ttn/atw/orig189.html>. May 19, 2009.
- \*EPA. 2009d. National Ambient Air Quality Standards (NAAQS). Washington, DC: U.S. Environmental Protection Agency, Office of Air and Radiation. <http://www.epa.gov/air/criteria.html>. May 19, 2009.
- \*EPA. 2009e. Superfund, emergency planning, and community right-to-know programs. Designation, reportable quantities, and notifications. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.4. <http://www.epa.gov/lawsregs/search/40cfr.html>. May 20, 2009.
- \*EPA. 2009f. Superfund, emergency planning, and community right-to-know programs. Toxic chemical release reporting. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 372.65. <http://www.epa.gov/lawsregs/search/40cfr.html>. May 11, 2009.
- \*EPA. 2009g. Integrated science assessment for carbon monoxide-first external review draft. Research Triangle Park, NC: U.S. Environmental Protection Agency. EPA600R09019.
- \*EPA. 2009h. 1970-2008 Average annual emissions, all criteria pollutants in MS Excel, June 2009. National Emissions Inventory (NEI) air pollutant emissions trends data. U.S. Environmental Protection Agency. <http://www.epa.gov/ttn/chieftrends/index.html>. June 10, 2009.
- \*EPA. 2009i. Carbon monoxide. Query AQS Data. U.S. Environmental Protection Agency. [http://www.epa.gov/aqspubl1/annual\\_summary.html](http://www.epa.gov/aqspubl1/annual_summary.html). June 11, 2009.
- ERG. 2000. 2000 Emergency Response Guidebook. Transport Canada. U.S. Department of Transportation. Secretariat of Transport and Communications of Mexico. <http://transit-safety.volpe.dot.gov/training/Archived/EPSSeminarReg/CD/documents/EmerPrep/erg2000.pdf>. June 9, 2009.
- \*Ernst A, Zibrak JD. 1998. Carbon monoxide poisoning. *N Engl J Med* 339(22):1603-1608.
- Esber LA, El-Fadel M, Nuwayhid I, et al. 2007. The effect of different ventilation modes on in-vehicle carbon monoxide exposure. *Atmos Environ* 41:3644-3657.

## 9. REFERENCES

- Esber LA, El-Fadel M, Shihadeh A. 2007. Comparison of trip average in-vehicle and exterior CO determinations by continuous and grab sampling using an electrochemical sensing method. *Atmos Environ* 41:6087-6094.
- \*Estabrook RW, Franklin MR, Hildebrandt AG. 1980. Factors influencing the inhibitory reactions. In: *Biological effects of carbon monoxide*. New York, NY: New York Academy of Sciences, 218.
- +Estrella B, Estrella R, Oviedo J, et al. 2005. Acute respiratory diseases and carboxyhemoglobin status in school children of Quito, Ecuador. *Environ Health Perspect* 113(5):607-611.
- Etzel RA. 2001. Indoor air pollutants in homes and schools. *Pediatr Clin North Am* 48(5):1153-1165.
- \*Fabian P, Borchers R, Weiler KH, et al. 1979. Simultaneously measured vertical profiles of H<sub>2</sub>, CH<sub>4</sub>, CO, N<sub>2</sub>O, CFCl<sub>3</sub>, and CF<sub>2</sub>Cl<sub>2</sub> in the mid-latitude stratosphere and troposphere. *J Geophys Res* 84(C6):3149-3154.
- +Fairley D. 1999. Daily mortality and air pollution in Santa Clara County, California: 1989-1996. *Environ Health Perspect* 107(8):637-641.
- +Farhat SCL, Paulo RLP, Shimoda TM, et al. 2005. Effect of air pollution on pediatric respiratory emergency room visits and hospital admissions. *Braz J Med Biol Res* 38:227-235.
- \*Farrow JR, Davis GJ, Roy TM, et al. 1990. Fetal death due to nonlethal maternal carbon monoxide poisoning. *J Forensic Sci* 35(6):1448-1452.
- Farrugia G, Irons WA, Rae JL, et al. 1993. Activation of whole cell currents in isolated human jejunal circular smooth muscle cells by carbon monoxide. *Am J Physiol* 264(6 Pt 1):G1184-1189.
- \*Favory R, Lancel S, Marechal X, et al. 2006a. Cardiovascular protective role for activated protein C during endotoxemia in rats. *Intensive Care Med* 32:899-905.
- Favory R, Lancel S, Tissier S, et al. 2006b. Myocardial dysfunction and potential cardiac hypoxia in rats induced by carbon monoxide inhalation. *Am J Respir Crit Care Med* 174(3):320-325.
- \*FDA. 2008. Everything added to food in the United States (EAFUS). Washington, DC: U.S. Food and Drug Administration. <http://vm.cfsan.fda.gov/~dms/eafus.html>. May 19, 2009.
- +\*Fechter LD, Annau Z. 1977. Toxicity of mild prenatal carbon monoxide exposure. *Science* 197(4304):680-682.
- +\*Fechter LD, Annau Z. 1980. Prenatal carbon monoxide exposure alters behavioral development. *Neurobehav Toxicol* 2(1):7-11.
- Fechter LD, Chen GD, Rao D. 2002. Chemical asphyxiants and noise. *Noise Health* 4(14):49-61.
- +\*Fechter LD, Karpa MD, Proctor B, et al. 1987a. Disruption of neostriatal development in rats following perinatal exposure to mild, but chronic carbon monoxide. *Neurotoxicol Teratol* 9(4):277-281.
- Fechter LD, Liu Y, Pearce TA. 1997. Cochlear protection from carbon monoxide exposure by free radical blockers in the guinea pig. *Toxicol Appl Pharmacol* 142(1):47-55.

## 9. REFERENCES

- \*Fechter LD, Thakur M, Miller B, et al. 1980. Effects of prenatal carbon monoxide exposure on cardiac development. *Toxicol Appl Pharmacol* 56(3):370-375.
- \*Fechter LD, Thorne PR, Nuttall AL. 1987b. Effects of carbon monoxide on cochlear electrophysiology and blood flow. *Hear Res* 27:37-45.
- \*Fechter LD, Young JS, Carlisle L. 1988. Potentiation of noise induced threshold shifts and hair cell loss by carbon monoxide. *Hear Res* 34:39-48.
- \*FEDRIP. 2009. Carbon monoxide. Federal Research in Progress database. Springfield, VA: National Technical Information Service.
- \*Fein A, Grossman RF, Jones JG, et al. 1980. Carbon monoxide effect on alveolar epithelial permeability. *Chest* 78:726-731.
- \*Fenn WO. 1970. The burning of CO in tissues. *Ann N Y Acad Sci* 174(1):64-71.
- +Fischer P, Hoek G, Brunekreef B, et al. 2003. Air pollution and mortality in the Netherlands: Are the elderly more at risk? *Eur Respir J* 21:34S-38S.
- +\*Fischer PH, Steerenberg PA, Snelder JD, et al. 2002. Association between exhaled nitric oxide, ambient air pollution and respiratory health in school children. *Int Arch Occup Environ Health* 75:348-353.
- \*Fisher AB, Hyde RW, Baue AE, et al. 1969. Effect of carbon monoxide on function and structure of the lung. *J Appl Physiol* 26(1):4-12.
- \*Fisher JA, Rucker J, Sommer LZ, et al. 1999. Isocapnic hyperpnea accelerates carbon monoxide elimination. *Am J Respir Crit Care Med* 159:1289-1292.
- \*Florkowski CM, Rossi ML, Carey MP, et al. 1992. Rhabdomyolysis and acute renal failure following carbon monoxide poisoning: Two case reports with muscle histopathology and enzyme activities. *J Toxicol Clin Toxicol* 30(3):443-454.
- \*Fomon SJ. 1966. Body composition of the infant: Part 1: The male reference infant. In: Faulkner F, ed. *Human development*. Philadelphia, PA: WB Saunders, 239-246.
- \*Fomon SJ, Haschke F, Ziegler EE, et al. 1982. Body composition of reference children from birth to age 10 years. *Am J Clin Nutr* 35(Suppl 5):1169-1175.
- +Forastiere F, Stafoggia M, Picciotto S, et al. 2005. A case-crossover analysis of out-of-hospital coronary deaths and air pollution in Rome, Italy. *Am J Respir Crit Care Med* 172:1549-1555.
- +Forastiere F, Stafoggia M, Tasco C, et al. 2007. Socioeconomic status, particulate air pollution, and daily mortality: Differential exposure or differential susceptibility. *Am J Ind Med* 50:208-216.
- \*Forster RE. 1970. Carbon monoxide and the partial pressure of oxygen in tissue. *Ann N Y Acad Sci* 174:233-241.

## 9. REFERENCES

- Fozo MS, Penney DG. 1993. Dibromomethane and carbon monoxide in the rat: Comparison of the cardiovascular and metabolic effects. *J Appl Toxicol* 13(2):147-151.
- \*Fritsch T, Hering P, Murtz M. 2007. Infrared laser spectroscopy for online recording of exhaled carbon monoxide - a progress report. *J Breath Res* 1(014002):R1-R7.
- \*Fujita T, Toda K, Karimova A, et al. 2001. Paradoxical rescue from ischemic lung injury by inhaled carbon monoxide driven by derepression of fibrinolysis. *Nat Med* 7(5):598-604.
- +Fung KY, Khan S, Krewski D, et al. 2006. Association between air pollution and multiple respiratory hospitalizations among the elderly in Vancouver, Canada. *Inhal Toxicol* 18(13):1005-1011.
- +\*Fung KY, Luginaah I, Gorey KM, et al. 2005. Air pollution and daily hospital admissions for cardiovascular diseases in Windsor, Ontario. *Can J Public Health* 96(1):29-33.
- +Fusco D, Forastiere F, Michelozzi P, et al. 2001. Air pollution and hospital admissions for respiratory conditions in Rome, Italy. *Eur Respir J* 17:1143-1150.
- Gandini C, Castoldi AF, Candura SM, et al. 2001. Carbon monoxide cardiotoxicity. *J Toxicol Clin Toxicol* 39(1):35-44.
- Garafolo MC, Seidler FJ, Auman JT, et al. 2002. Beta-adrenergic modulation of muscarinic cholinergic receptor expression and function in developing heart. *Am J Physiol Regul Integr Comp Physiol* 282:R1356-R1363.
- Garvey DJ, Longo LD. 1978. Chronic low level maternal carbon monoxide exposure and fetal growth and development. *Biol Reprod* 19(1):8-14.
- \*Gautier M, Antier D, Bonnet P, et al. 2007. Continuous inhalation of carbon monoxide induces right ventricle ischemia and dysfunction in rats with hypoxic pulmonary hypertension. *Am J Physiol Heart Circ Physiol* 293:H1046-H1052
- Gaworski CL, Carmines EL, Faqi AS, et al. 2004. *In utero* and lactation exposure of rats to 1R4F reference cigarette mainstream smoke: Effect on prenatal and postnatal development. *Toxicol Sci* 79:157-169.
- \*George C. 2001. Carbon monoxide. In: Kirk-Othmer encyclopedia of chemical technology. John Wiley & Sons, Inc., <http://mrw.interscience.wiley.com/emrw/9780471238966/kirk/article/carbpier.a02/current/pdf>. May 7, 2009.
- \*Geuens E, Brouns I, Flamez D, et al. 2003. A globin in the nucleus! *J Biol Chem* 278:30417-30420.
- \*Ghio AJ, Stonehuerner JG, Dailey LA, et al. 2008. Carbon monoxide reversibly alters iron homeostasis and respiratory epithelial cell function. *Am J Respir Cell Mol Biol* 38(6):715-723.
- \*Gibson QH, Olson JS, McKinnie RE, et al. 1986. A kinetic description of ligand binding to sperm whale myoglobin. *J Biol Chem* 261(22):10228-10239.
- +\*Gilboa SM, Mendola P, Olshan AF, et al. 2005. Relation between ambient air quality and selected birth defects, seven county study, Texas, 1997-2000. *Am J Epidemiol* 162(3):238-252.

## 9. REFERENCES

- Ginsberg MD, Myers RE. 1974. Experimental carbon monoxide encephalopathy in the primate. I. Physiologic and metabolic aspects. *Arch Neurol* 30(3):202-208.
- Ginsberg MD, Myers RE. 1976. Fetal brain injury after maternal carbon monoxide intoxication. *Neurology* 26(1):15-23.
- +\*Giustino A, Cagiano R, Carratu MR, et al. 1993. Immunological changes produced in rats by prenatal exposure to carbon monoxide. *Pharmacol Toxicol* 73(5):274-278.
- +\*Giustino A, Cagiano R, Carratu MR, et al. 1999. Prenatal exposure to low concentrations of carbon monoxide alters habituation and non-spatial working memory in rat offspring. *Brain Res* 844:201-205.
- +\*Giustino A, Carratu MR, Brigiani GS, et al. 1994. Changes in the frequency of splenic immunocompetent cells in rats exposed to carbon monoxide during gestation. *Immunopharmacol Immunotoxicol* 16(2):281-292.
- \*Giwerzman A, Carlsen E, Keiding N, et al. 1993. Evidence for increasing incidence of abnormalities of the human testis: A review. *Environmental Health Perspectives Supplement* 101(2):65-71.
- Glabe A, Chung Y, Xu D, et al. 1998. Carbon monoxide inhibition of regulatory pathways in myocardium. *Am J Physiol Heart Circ Physiol* 274:2143-2151.
- Gliner JA, Raven PB, Horvath SM, et al. 1975. Man's physiologic response to long-term work during thermal and pollutant stress. *J Appl Physiol* 39(4):628-632.
- Goda N, Suzuki K, Naito M, et al. 1998. Distribution of heme oxygenase isoforms in rat liver. Topographic basis for carbon monoxide-mediated microvascular relaxation. *J Clin Invest* 101(3):604-612.
- +\*Gold DR, Litonjua A, Schwartz J, et al. 2000. Ambient pollution and heart rate variability. *Circulation* 101:1267-1273.
- +\*Gold DR, Litonjua AA, Zanobetti A, et al. 2005. Air pollution and ST-segment depression in elderly subjects. *Environ Health Perspect* 113(7):883-887.
- +Goldberg MS, Burnett RT, Bailar JC, et al. 2001. Identification of persons with cardiorespiratory conditions who are at risk of dying from the acute effects of ambient air particles. *Environ Health Perspect* 109(Suppl 4):487-494.
- +Goldberg MS, Burnett RT, Valois MF, et al. 2003. Associations between ambient air pollution and daily mortality among persons with congestive heart failure. *Environ Res* 91:8-20.
- +Goldberg MS, Burnett RT, Yale JF, et al. 2006. Associations between ambient air pollution and daily mortality among persons with diabetes and cardiovascular disease. *Environ Res* 100:255-267.
- \*Gordon CJ. 1990. Thermal biology of the laboratory rat. *Physiol Rev* 47:963-991.
- Gorman D, Huang YL. 2008. Heme oxygenase and nitric oxide synthetase blockade and brain blood flow in sheep exposed to carbon monoxide. *Neurosci Lett* 444(2):203-207.

## 9. REFERENCES

- \*Gorman D, Drewry A, Huang YL, et al. 2003. The clinical toxicology of carbon monoxide. *Toxicology* 187(1):25-38.
- +Goss CH, Newsom SA, Schildcrout JS, et al. 2004. Effect of ambient air pollution on pulmonary exacerbations and lung function in cystic fibrosis. *Am J Respir Crit Care Med* 169:816-821.
- Gothert M, Lutz F, Malorny G. 1970. Carbon monoxide partial pressure in tissue of different animals. *Environ Res* 3(4):303-309.
- +Gouveia N, Fletcher T. 2000. Time series analysis of air pollution and mortality: Effects by cause, age and socioeconomic status. *J Epidemiol Community Health* 54:750-755.
- +\*Gouveia N, Bremner SA, Novaes HMD. 2004. Associations between ambient air pollution and birth weight in Sao Paulo, Brazil. *J Epidemiol Community Health* 58:11-17.
- \*Graham LA, Noseworthy L, Fugler D, et al. 2004. Contribution of vehicle emissions from an attached garage to residential indoor air pollution levels. *J Air Waste Manage Assoc* 54:563-584.
- \*Greingor JL, Tosi JM, Ruhlmann S, et al. 2001. Acute carbon monoxide intoxication during pregnancy. One case report and review of the literature. *Emerg Med J* 18(5):399-401.
- Groll-Knapp E, Haider M, Kienzl K, et al. 1988. Changes in discrimination learning and brain activity (ERP's) due to combined exposure to NO and CO in rats. *Toxicology* 49(2-3):441-447.
- Grover TR, Rairigh RL, Zenge JP, et al. 2000. Inhaled carbon monoxide does not cause pulmonary vasodilation in the late-gestation fetal lamb. *Am J Physiol Lung Cell Mol Physiol* 278:L779-L784.
- +Guo YL, Lin YC, Sung FC, et al. 1999. Climate, traffic-related air pollutants, and asthma prevalence in middle-school children in Taiwan. *Environ Health Perspect* 107(12):1001-1006.
- Guyatt AR, Holmes MA, Cumming G. 1981. Can carbon monoxide be absorbed from the upper respiratory tract in man? *Eur J Respir Dis* 62(6):383-390.
- \*Guzelian PS, Henry CJ, Olin SS. 1992. Similarities and differences between children and adults: Implications for risk assessment. Washington, DC: International Life Sciences and Press Institute Press, 1-283.
- +Gwynn RC, Burnett RT, Thurston GD. 2000. A time-series analysis of acidic particulate matter and daily mortality and morbidity in the Buffalo, New York, region. *Environ Health Perspect* 108(2):125-133.
- +\*Ha EH, Hong YC, Lee BE, et al. 2001. Is air pollution a risk factor for low birth weight in Seoul? *Epidemiology* 12:643-648.
- +\*Ha EH, Lee JT, Kim H, et al. 2003. Infant susceptibility of mortality to air pollution in Seoul, South Korea. *Pediatrics* 111:284-290.
- +Hajat S, Anderson HR, Atkinson RW, et al. 2002. Effects of air pollution on general practitioner consultations for upper respiratory diseases in London. *Occup Environ Med* 59:294-299.

## 9. REFERENCES

- +Hajat S, Haines A, Goubet SA, et al. 1999. Association of air pollution with daily GP consultations for asthma and other lower respiratory conditions in London. *Thorax* 54(7):597-605.
- \*Hampson NB. 2007. Carboxyhemoglobin elevation due to hemolytic anemia. *J Emerg Med* 33(1):17-19.
- \*Hampson NB, Hauff NM. 2008. Carboxyhemoglobin levels in carbon monoxide poisoning: Do they correlate with the clinical picture? *Am J Emerg Med* 26(6):665-669.
- \*Hampson NB, Zmaeff JL. 2001. Outcome of patients experiencing cardiac arrest with carbon monoxide poisoning treated with hyperbaric oxygen. *Ann Emerg Med* 38:36-41.
- \*Hampson NB, Rudd RA, Hauff NM. 2009. Increased long-term mortality among survivors of acute carbon monoxide poisoning. *Crit Care Med* 37(6):1941-1947.
- Hanada A, Sander M, Gonzalez-Alonso J. 2003. Human skeletal muscle sympathetic nerve activity, heart rate and limb haemodynamics with reduced blood oxygenation and exercise. *J Physiol* 551(2):635-647.
- \*Hanninen OO, Alm S, Katsouyanni K, et al. 2004. The EXPOLIS study: Implications for exposure research and environmental policy in Europe. *J Expo Anal Environ Epidemiol* 14:440-456.
- +Hapcioglu B, Issever H, Kocyigit E, et al. 2006. The effect of air pollution and meteorological parameters on chronic obstructive pulmonary disease at an Istanbul hospital. *Indoor Built Environ* 15(2):147-153.
- Hara S, Mukai T, Kurosaki K, et al. 2002. Modification of the striatal dopaminergic neuron system by carbon monoxide exposure in free-moving rats, as determined by *in vivo* brain microdialysis. *Arch Toxicol* 76:596-605.
- Harada T, Koi H, Kubota T, et al. 2004. Haem oxygenase augments porcine granulosa cell apoptosis *in vitro*. *J Endocrinol* 181:191-205.
- Harbin TJ, Benignus VA, Muller KE, et al. 1988. The effects of low-level carbon monoxide exposure upon evoked cortical potentials in young and elderly men. *Neurotoxicol Teratol* 10(2):93-100.
- Harley RA, Marr LC, Lehner JK, et al. 2005. Changes in motor vehicle emissions on diurnal to decadal time scales and effects on atmospheric composition. *Environ Sci Technol* 39:5356-5362.
- \*Hauck H, Neuberger M. 1984. Carbon monoxide uptake and the resulting carboxyhemoglobin in man. *Eur J Appl Physiol Occup Physiol* 53(2):186-190.
- +\*Hausberg M, Somers VK. 1997. Neural circulatory responses to carbon monoxide in healthy humans. *Hypertension* 29(5):1114-1118.
- \*HazDat. 2007. Carbon Monoxide. HazDat Database: ATSDR's Hazardous Substance Release and Health Effects Database. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- \*Heckerling PS, Leikin JB, Maturen A, et al. 1990. Screening hospital admissions from the emergency department for occult carbon monoxide poisoning. *Am J Emerg Med* 8(4):301-304.

## 9. REFERENCES

- Hegewald MJ, Crapo RO. 2007. Socioeconomic status and lung function. *Chest* 132(5):1608-1614.
- +HEI. 2000. Daily mortality and fine and ultrafine particles in Erfurt, Germany. Part I. Role of particle numbers and particle mass. Health Effects Institute. <http://pubs.healtheffects.org/getfile.php?u=189>. September 24, 2009.
- \*HEI. 2005. The national morbidity, mortality, and air pollution study. Part IV. Hierarchical bivariate time-series models-a combined analysis of PM<sub>10</sub> effects on hospitalization and mortality. Number 94. Health Effects Institute. <http://pubs.healtheffects.org/getfile.php?u=218>. August 10, 2009.
- \*Helfaer MA, Traystman RJ. 1996. Cerebrovascular effects of carbon monoxide. In: Penny DG, ed. *Carbon monoxide poisoning*. New York, NY: CRC Press, 69-86.
- Hendler I, Baum M, Kreiser D, et al. 2004. End-tidal breath carbon monoxide measurements are lower in pregnant women with uterine contractions. *J Perinatol* 24:275-278.
- +\*Henrotin JB, Besancenot JP, Bejot Y, et al. 2007. Short-term effects of ozone air pollution on ischaemic stroke occurrence: A case-crossover analysis from a 10-year population-based study in Dijon, France. *Occup Environ Med* 64:439-445.
- \*Herman GD, Shapiro AB, Leikin J. 1988. Myonecrosis in carbon monoxide poisoning. *Vet Hum Toxicol* 30(1):28-30.
- Hermans RH, McGivern RF, Chen W, et al. 1993. Altered adult sexual behavior in the male rat following chronic prenatal hypoxia. *Neurotoxicol Teratol* 15(6):353-363.
- \*Hill EP, Hill JR, Power GG, et al. 1977. Carbon monoxide exchanges between the human fetus and mother: A mathematical model. *Am J Physiol* 232(3):H311-H323.
- +\*Hinderliter AL, Adams KF, Price CJ, et al. 1989. Effects of low-level carbon monoxide exposure on resting and exercise-induced ventricular arrhythmias in patients with coronary artery disease and no baseline ectopy. *Arch Environ Health* 44(2):89-93.
- +Hinwood AL, De Klerk N, Rodriguez C, et al. 2006. The relationship between changes in daily air pollution and hospitalizations in Perth, Australia 1992-1998: A case-crossover study. *Int J Environ Health Res* 16(1):27-46.
- +Hirsch T, Weiland SK, von Mutius E, et al. 1999. Inner city air pollution and respiratory health and atopy in children. *Eur Respir J* 14:669-677.
- +Hoek G, Brunekreef B, Fischer P, et al. 2001. The association between air pollution and heart failure, arrhythmia, embolism, thrombosis, and other cardiovascular causes of death in a time series study. *Epidemiology* 12:355-357.
- +Hoek G, Brunekreef B, Verhoeff A, et al. 2000. Daily mortality and air pollution in the Netherlands. *J Air Waste Manage Assoc* 50:1380-1389.
- \*Hoel DG, Davis DL, Miller AB, et al. 1992. Trends in cancer mortality in 15 industrialized countries, 1969-1986. *J Natl Cancer Inst* 84(5):313-320.

## 9. REFERENCES

- +\*Holguin F, Tellez-Rojo MM, Hernandez M, et al. 2003. Air pollution and heart rate variability among the elderly in Mexico City. *Epidemiology* 14:521-527.
- Holland RA. 1970. Reaction rates of carbon monoxide and hemoglobin. *Ann N Y Acad Sci* 174(1):154-171.
- \*Hollander DI, Nagey DA, Welch R, et al. 1987. Hyperbaric oxygen therapy for the treatment of acute carbon monoxide poisoning in pregnancy. *J Reprod Med* 32:615-617.
- \*Honda Y, Nitta H, Ono M. 2003. Low level carbon monoxide and mortality of persons aged 65 or older in Tokyo, Japan 1976-1990. *J Health Sci* 49:454-458
- +\*Hong YC, Lee JT, Kim H, et al. 2002a. Effects of air pollutants on acute stroke mortality. *Environ Health Perspect* 110(2):187-191.
- +\*Hong YC, Lee JT, Kim H, et al. 2002b. Air pollution: A new risk factor in ischemic stroke mortality. *Stroke* 33:2165-2169.
- +\*Hong YC, Leem JH, Ha EH. 1999a. Air pollution and daily mortality in Inchon, Korea. *J Korean Med Sci* 14:239-244.
- +\*Hong YC, Leem JH, Ha EH, et al. 1999b. PM<sub>10</sub> exposure, gaseous pollutants, and daily mortality in Inchon, South Korea. *Environ Health Perspect* 107(11):873-878.
- \*Hopkins RO, Fearing MA, Weaver LK, et al. 2006. Basal ganglia lesions following carbon monoxide poisoning. *Brain Inj* 20(3):273-281.
- Horvath I, Donnelly LE, Kiss A, et al. 1998a. Raised levels of exhaled carbon monoxide are associated with an increased expression of heme oxygenase-1 in airway macrophages in asthma: A new marker of oxidative stress. *Thorax* 53(8):668-672.
- Horvath I, Loukides S, Wodehouse T, et al. 1998b. Increased levels of exhaled carbon monoxide in bronchiectasis: A new marker of oxidative stress. *Thorax* 53:867-870.
- \*Horvath SM, Bedi JF, Wagner JA, et al. 1988. Maximal aerobic capacity at several ambient concentrations of CO at several altitudes. *J Appl Physiol* 65(6):2696-2708.
- Horvath SM, Dahms TE, O'Hanlon JF. 1971. Carbon monoxide and human vigilance. A deleterious effect of present urban concentrations. *Arch Environ Health* 23(5):343-347.
- +\*Horvath SM, Raven PB, Dahms TE, et al. 1975. Maximal aerobic capacity at different levels of carboxyhemoglobin. *J Appl Physiol* 38(2):300-303.
- +\*Hosseinpour AR, Forouzanfar MH, Yunesian M, et al. 2005. Air pollution and hospitalization due to angina pectoris in Tehran, Iran: A time-series study. *Environ Res* 99:126-131.
- \*Houck JE, Crouch J, Huntley RH. 2006. Review of wood heater and fireplace emissions factors. In: 10th International Emission Inventory Conference—"One Atmosphere, One Inventory, Many Challenges" <http://www.epa.gov/ttn/chief/conference/ei10/pm/houck.pdf>. May 8, 2009.

## 9. REFERENCES

- \*HSDB. 2009. Carbon monoxide. Hazardous Substances Data Bank. National Library of Medicine. <http://toxnet.nlm.nih.gov>. June 9, 2009
- \*Hsia CC. 2002. Recruitment of lung diffusing capacity: Update of concept and application. *Chest* 122(5):1774-1783.
- \*Hu S, Kincaid JR. 1992. Resonance Raman studies of the carbonmonoxy form of catalase. Evidence for and effects of phenolate ligation. *FEBS Lett* 314(3):293-296.
- \*Hugod C. 1980. The effect of carbon monoxide exposure on morphology of lungs and pulmonary arteries in rabbits. A light- and electron-microscopic study. *Arch Toxicol* 43:273-281.
- Hutter CDD, Blair ME. 1996. Carbon monoxide - does fetal exposure cause sudden infant death syndrome? *Med Hypotheses* 46:1-4.
- +\*Huynh M, Woodruff TJ, Parker JD, et al. 2006. Relationships between air pollution and preterm birth in California. *Paediatr Perinat Epidemiol* 20:454-461.
- +\*Hwang BF, Jaakkola JJ. 2008. Ozone and other air pollutants and the risk of oral clefts. *Environ Health Perspect* 116(10):1411-1415.
- +Hwang BF, Jaakkola JJ, Lee YL, et al. 2006. Relation between air pollution and allergic rhinitis in Taiwanese schoolchildren. *Respir Res* 7(1):23.
- +\*Hwang BF, Lee YL, Lin YC, et al. 2005. Traffic related air pollution as a determinant of asthma among Taiwanese school children. *Thorax* 60:467-476.
- +Hwang JS, Chan CC. 2002. Effects of air pollution on daily clinic visits for lower respiratory tract illness. *Am J Epidemiol* 155(1):1-10.
- \*IARC. 2009. Agents Reviewed by the IARC Monographs. Volumes 1-99. Lyon, France: International Agency for Research on Cancer. <http://monographs.iarc.fr/ENG/Classification/index.php>. May 19, 2009.
- +\*Ibald-Mulli A, Stieber J, Wichmann HE, et al. 2001. Effects of air pollution on blood pressure: A population-based approach. *Am J Public Health* 91:571-577.
- \*Iheagwara KN, Thom SR, Deutschman CS, et al. 2007. Myocardial cytochrome oxidase activity is decreased following carbon monoxide exposure. *Biochim Biophys Acta* 1772(9):1112-1116.
- \*ILSI. 1994. Physiological parameter values for PBPK models. International Life Science Institute under agreement with U.S. Environmental Protection Agency. Office of Health and Environmental Assessment, Washington, DC.
- Imai T, Morita T, Shindo T, et al. 2001. Vascular smooth muscle cell-directed overexpression of heme oxygenase-1 elevates blood pressure through attenuation of nitric oxide-induced vasodilation in mice. *Circ Res* 89:55-62.
- Ingi T, Ronnett GV. 1995. Direct demonstration of a physiological role for carbon monoxide in olfactory receptor neurons. *J Neurosci* 15(12):8214-8222.

## 9. REFERENCES

- Ingi T, Cheng J, Ronnett GV. 1996. Carbon monoxide: An endogenous modulator of the nitric oxide-cyclic GMP signaling system. *Neuron* 16(4):835-842.
- Inman RE, Ingersoll RB, Levy EA. 1971. Soil: A natural sink for carbon monoxide. *Science* 172:1229-1231.
- IPCS. 2008. Carbon oxide. International chemical safety cards. International Programme on Chemical Safety. Commission of the European Communities. <http://www.cdc.gov/niosh/ipcsneng/neng0023.html>. June 9, 2009.
- \*IRIS. 2009. Integrated Risk Information System. Washington, DC. U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/index.html>. May 11, 2009.
- Ischiropoulos H, Beers MF, Ohnishi ST, et al. 1996. Nitric oxide production and perivascular tyrosine nitration in brain after carbon monoxide poisoning in the rat. *J Clin Invest* 97:2260-2267.
- Ishimaru H, Katoh A, Suzuki H, et al. 1992. Effects of N-methyl-D-aspartate receptor antagonists on carbon monoxide-induced brain damage in mice. *J Pharmacol Exp Ther* 261(1):349-352.
- Ishimaru H, Nabeshima T, Katoh A, et al. 1991. Effects of successive carbon monoxide exposures on delayed neuronal death in mice under the maintenance of normal body temperature. *Biochem Biophys Res Commun* 179(2):836-840.
- +Ito K, Thurston GD, Silverman RA. 2007. Characterization of PM2.5, gaseous pollutants, and meteorological interactions in the context of time-series health effects models. *J Expo Sci Environ Epidemiol* 17 (Suppl 2): S45-S60.
- +\*Jalaludin B, Morgan G, Lincoln D, et al. 2006. Associations between ambient air pollution and daily emergency department attendances for cardiovascular disease in the elderly (65+ years). *J Expo Sci Environ Epidemiol* 16:225-237.
- +\*Jalaludin B, Mannes T, Morgan G, et al. 2007. Impact of ambient air pollution on gestational age is modified by season in Sydney, Australia. *Environ Health* 6(1):16-24.
- +\*Jerrett M, Burnett RT, Willis A, et al. 2003. Spatial analysis of the air pollution-mortality relationship in the context of ecologic confounders. *J Toxicol Environ Health Part A* 66:1735-1777.
- Jiang G, Gao J, Xu Y, et al. 2009. Structural and functional improvement of injured brain after severe acute carbon monoxide poisoning by stem cell-based therapy in rats. *Crit Care Med* 37(4):1416-1422.
- \*Johanson CE. 1980. Permeability and vascularity of the developing brain: Cerebellum vs. cerebral cortex. *Brain Res* 190(1):3-16.
- Johnson FK, Johnson RA. 2003. Carbon monoxide promotes endothelium-dependent constriction of isolated gracilis muscle arterioles. *Am J Physiol Regul Integr Comp Physiol* 285:R536-R541.
- Johnson FK, Durante W, Peyton KJ, et al. 2004. Heme oxygenase-mediated endothelial dysfunction in DOCA-salt, but not in spontaneously hypertensive, rat arterioles. *Am J Physiol Heart Circ Physiol* 286:H1681-H1687.

## 9. REFERENCES

- Johnson FK, Johnson RA, Durante W, et al. 2006. Metabolic syndrome increases endogenous carbon monoxide production to promote hypertension and endothelial dysfunction in obese Zucker rats. *Am J Physiol Regul Integr Comp Physiol* 290:R601-R608.
- \*Joumard R, Chiron M, Vidon R, et al. 1991. Mathematical models of the uptake of carbon monoxide on hemoglobin at low carbon monoxide levels. *Environ Health Perspect* 41:277-289.
- \*Kalay N, Ozdogru I, Cetinkaya Y, et al. 2007. Cardiovascular effects of carbon monoxide poisoning. *Am J Cardiol* 99:322-324.
- Kanten WE, Penney DG, Francisco K, et al. 1983. Hemodynamic responses to acute carboxy-hemoglobinemia in the rat. *Am J Physiol* 244(3):H320-H327.
- Kao LW, Nañagas KA. 2004. Carbon monoxide poisoning. *Emerg Med Clin North Am* 22(4):985-1018.
- Kao LW, Nañagas KA. 2005. Carbon monoxide poisoning. *Med Clin North Am* 89(6):1161-1194.
- \*Kao LW, Nañagas KA. 2006. Toxicity associated with carbon monoxide. *Clin Lab Med* 26(1):99-125.
- +Karr C, Lumley T, Schreuder A, et al. 2007. Effects of subchronic and chronic exposure to ambient air pollutants on infant bronchiolitis. *Am J Epidemiol* 165(5):553-560.
- +\*Karr C, Lumley T, Shepherd K, et al. 2006. A case-crossover study of wintertime ambient air pollution and infant bronchiolitis. *Environ Health Perspect* 114(2):277-281.
- Katoue MG, Khan I, Oriowo MA. 2005. Increased expression and activity of heme oxygenase-2 in pregnant rat aorta is not involved in attenuated vasopressin-induced contraction. *Naunyn Schmiedebergs Arch Pharmacol* 372:220-227.
- Katoue MG, Khan I, Oriowo MA. 2006. Pregnancy-induced modulation of calcium mobilization and down-regulation of Rho-kinase expression contribute to attenuated vasopressin-induced contraction of the rat aorta. *Vascul Pharmacol* 44:170-176.
- Katsumata Y, Aoki M, Oya M, et al. 1980. Liver damage in rats during acute carbon monoxide poisoning. *Forensic Sci Int* 16(2):119-123.
- Kealey GP. 2009. Carbon monoxide toxicity. *J Burn Care Res* 30(1):146-147.
- +Keatinge WR, Donaldson GC. 2001. Mortality related to cold and air pollution in London after allowance for effects of associated weather patterns. *Environ Res* 86(3):209-216.
- +Kettunen J, Lanki T, Tiittanen P, et al. 2007. Associations of fine and ultrafine particulate air pollution with stroke mortality in an area of low air pollution levels. *Stroke* 38:918-922.
- \*Khalil MAK, Rasmussen RA. 1990. The global cycle of carbon monoxide: Trends and mass balance. *Chemosphere* 20(1-2):227-242.
- Khan AA, Wang Y, Sun Y, et al. 2006. Neuroglobin-overexpressing transgenic mice are resistant to cerebral and myocardial ischemia. *Proc Natl Acad Sci USA* 103(47):17944-17948.

## 9. REFERENCES

- Khosla S, Laddu A, Ehrenpreis S, et al. 1994. Cardiovascular effects of nicotine: Relation to deleterious effects of cigarette smoking. *Am Heart J* 127:1669-1672.
- Kim HP, Wang X, Nakao A, et al. 2005. Caveolin-1 expression by means of p38 $\beta$  mitogen-activated protein kinase mediates the antiproliferative effect of carbon monoxide. *Proc Natl Acad Sci USA* 102(32):11319-11324.
- Kim HS, Loughran PA, Rao J, et al. 2008. Carbon monoxide activates NF- $\kappa$ B via ROS generation and Akt pathways to protect against cell death of hepatocytes. *Am J Physiol Gastrointest Liver Physiol* 295:G146-G152.
- +\*Kim KM, Pae HO, Zheng M, et al. 2007. Carbon monoxide induces heme oxygenase-1 via activation of protein kinase R-like endoplasmic reticulum kinase and inhibits endothelial cell apoptosis triggered by endoplasmic reticulum stress. *Circ Res* 101:919-927.
- Kimmel EC, Carpenter RL, Reboulet JE, et al. 1999. A physiological model for predicting carboxyhemoglobin formation from exposure to carbon monoxide in rats. *J Appl Physiol* 86(6):1977-1983.
- \*King GM. 1999a. Characteristics and significance of atmospheric carbon monoxide consumption by soils. *Chemosphere Global Change Sci.* 1:53-63.
- \*King GM. 1999b. Attributes of atmospheric carbon monoxide oxidation by Maine forest soils. *Appl Environ Microbiol* 65(12):5257-5264.
- Kinney HC, Filiano JJ, Sleeper LA, et al. 1995. Decreased muscarinic receptor binding in the arcuate nucleus in sudden infant death syndrome. *Science* 269(5229):1446-1450.
- Kinobe RT, Vlahakis JZ, Soong JM, et al. 2006. Heme oxygenase activity in fetal and adult sheep is not altered by acclimatization to high altitude hypoxia. *Can J Physiol Pharmacol* 84:893-901.
- +\*Kizakevich PN, McCartney ML, Hazucha MJ, et al. 2000. Noninvasive ambulatory assessment of cardiac function in healthy men exposed to carbon monoxide during upper and lower body exercise. *Eur J Appl Physiol* 83:7-16.
- Kjeldsen K, Astrup P, Wanstrup J. 1972. Ultrastructural intimal changes in the rabbit aorta after a moderate carbon monoxide exposure. *Atherosclerosis* 16(1):67-82.
- Klasner AE, Smith SR, Thompson MW, et al. 1998. Carbon monoxide mass exposure in a pediatric population. *Acad Emerg Med* 5:992-996.
- \*Klawans HL, Stein RW, Tanner CM, et al. 1982. A pure parkinsonian syndrome following acute carbon monoxide intoxication. *Arch Neurol* 39(5):302-304.
- Klein JP, Forster HV, Stewart RD, et al. 1980. Hemoglobin affinity for oxygen during short-term exhaustive exercise. *J Appl Physiol* 48(2):236-242.
- +\*Kleinman MT, Davidson DM, Vandagriff RB, et al. 1989. Effects of short-term exposure to carbon monoxide in subjects with coronary artery disease. *Arch Environ Health* 44(6):361-369.

## 9. REFERENCES

- +\*Kleinman MT, Leaf DA, Kelly E, et al. 1998. Urban angina in the mountains: Effects of carbon monoxide and mild hypoxemia on subjects with chronic stable angina. *Arch Environ Health* 53(6):388-397.
- +Klemm RJ, Lipfert FW, Wyzga RE, et al. 2004. Daily mortality and air pollution in Atlanta: Two years of data from ARIES. *Inhal Toxicol* 16 (Suppl 1):131-141.
- Klimisch HJ, Chevalier HJ, Harke HP, et al. 1975. Uptake of carbon monoxide in blood of miniature pigs and other mammals. *Toxicology* 3(3):301-310.
- Knuckles TL, Lund AK, Lucas SN, et al. 2008. Diesel exhaust exposure enhances vasoconstriction via uncoupling of eNOS. *Toxicol Appl Pharmacol* 230:346-351.
- Koehler RC, Jones MD, Traystman RJ. 1982. Cerebral circulatory response to carbon monoxide and hypoxic hypoxia in the lamb. *Am J Physiol* 243(1):H27-H32.
- \*Koike A, Wasserman K, Armon Y, et al. 1991. The work-rate-dependent effect of carbon monoxide on ventilatory control during exercise. *Respir Physiol* 85(2):169-183.
- +\*Koken PJ, Piver WT, Ye F, et al. 2003. Temperature, air pollution, and hospitalization for cardiovascular diseases among elderly people in Denver. *Environ Health Perspect* 111(10):1312-1317.
- \*Komori M, Nishio K, Kitada M, et al. 1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. *Biochemistry* 29(18):4430-4433.
- +Kontos AS, Fassols SD, Deli MF. 1999. Short-term effects of air pollution on childhood respiratory illness in Piraeus, Greece, 1987-1992: Nonparametric stochastic dynamic analysis. *Environ Res* 81:275-296.
- Korres S, Riga M, Balatsouras D, et al. 2007. Influence of smoking on developing cochlea does smoking during pregnancy affect the amplitudes of transient evoked otoacoustic emissions in newborns? *Int J Pediatr Otorhinolaryngol* 71:781-786.
- Koskela RS. 1994. Cardiovascular diseases among foundry workers exposed to carbon monoxide. *Scand J Work Environ Health* 20(4):286-293.
- \*Kreck TC, Shade ED, Lamm W, et al. 2001. Isocapnic hyperventilation increases carbon monoxide elimination and oxygen delivery. *Am J Respir Crit Care Med* 163:458-462.
- Kreiser D, Baum M, Seidman DS, et al. 2004. End tidal carbon monoxide levels are lower in women with gestation hypertension and pre-eclampsia. *J Perinatol* 24:213-217.
- \*Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes AW, ed. *Principles and methods of toxicology*. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.
- \*Krishnan K, Anderson ME, Clewell HJ, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, ed. *Toxicology of chemical mixtures. Case studies, mechanisms, and novel approaches*. San Diego, CA: Academic Press, 399-437.

## 9. REFERENCES

- Kuller LH, Radford EP, Swift D, et al. 1975. Carbon monoxide and heart attacks. *Arch Environ Health* 30:477-482.
- \*Kuska J, Kokot F, Wnuk R. 1980. Acute renal failure after exposure to carbon monoxide. *Mater Med Pol* 12:236-238.
- \*Kwak HM, Yang YH, Lee MS. 1986. Cytogenetic effects on mouse fetus of acute and chronic transplacental *in vivo* exposure to carbon monoxide: Induction of micronuclei and sister chromatid exchanges. *Yonsei Med J* 27(3):205-212.
- +Kwon HJ, Cho SH, Nyberg F, et al. 2001. Effects of ambient air pollution on daily mortality in a cohort of patients with congestive heart failure. *Epidemiology* 12:413-419.
- +\*Lagorio S, Forastiere F, Pistelli R, et al. 2006. Air pollution and lung function among susceptible adult subjects: A panel study. *Environ Health* 5:11-22.
- Lamberto C, Nunes H, Le Toumelin P, et al. 2004. Membrane and capillary blood components of diffusion capacity of the lung for carbon monoxide in pulmonary sarcoidosis: Relation to exercise gas exchange. *Chest* 125(6):2061-2068.
- \*Landaw SA. 1973. The effects of cigarette smoking on total body burden and excretion rates of carbon monoxide. *J Occup Med* 15(3):231-235.
- \*Landaw SA, Callahan EW, Schmid R. 1970. Catabolism of heme *in vivo*: Comparison of the simultaneous production of bilirubin and carbon monoxide. *J Clin Invest* 49:914-925.
- \*Langston P, Gorman D, Runciman W, et al. 1996. The effect of carbon monoxide on oxygen metabolism in the brains of awake sheep. *Toxicology* 114:223-232.
- +\*Lanki T, Pekkanen J, Aalto P, et al. 2006. Associations of traffic related air pollutants with hospitalisation for first acute myocardial infarction: The HEAPSS study. *Occup Environ Med* 63:844-851.
- Lash GE, Mclaughlin BE, MacDonald-Goodfellow SK, et al. 2003. Relationship between tissue damage and heme oxygenase expression in chorionic villi of term human placenta. *Am J Physiol Heart Circ Physiol* 284:H160-H167.
- Laupeze B, Amiot L, Sparfel L, et al. 2002. Polycyclic aromatic hydrocarbons affect functional differentiation and maturation of human monocyte-derived dendritic cells. *J Immunol* 168:2652-2658.
- \*Lavonas EJ. 2007. Carbon monoxide poisoning. In: Haddad and Winchester's clinical management of poisoning and drug overdose. 4th ed. Philadelphia, PA: Saunders, 1297-1307.
- +Leaf DA, Kleinman MT. 1996a. Acute exposure to carbon monoxide does not affect plasma lipids, lipoproteins, and apolipoproteins. *Angiology* 47(4):337-341.
- \*Leaf DA, Kleinman MT. 1996b. Urban ectopy in the mountains: Carbon monoxide exposure at high altitude. *Arch Environ Health* 51(4):283-290.
- +\*Lee BE, Ha EH, Park HS, et al. 2003a. Exposure to air pollution during different gestational phases contributes to risks of low birth weight. *Hum Reprod* 18(3):638-643.

## 9. REFERENCES

- +\*Lee BS, Heo JH, Kim YM, et al. 2006. Carbon monoxide mediates heme oxygenase-1 induction via Nrf2 activation in hepatoma cells. *Biochem Biophys Res Commun* 343:965-972.
- \*Lee HU, Lee HJ, Park HY, et al. 2001. Effects of heme oxygenase system on the cyclooxygenase in the primary cultured hypothalamic cells. *Arch Pharm Res* 24:607-612.
- +\*Lee IM, Tsai SS, Ho CK, et al. 2007a. Air pollution and hospital admissions for congestive heart failure in a tropical city: Kaohsiung, Taiwan. *Inhal Toxicol* 19(10):899-904.
- +Lee IM, Tsai SS, Chang CC, et al. 2007b. Air pollution and hospital admissions for chronic obstructive pulmonary disease in a tropical city: Kaohsiung, Taiwan. *Inhal Toxicol* 19(5):393-398.
- +\*Lee JT, Kim H, Cho YS, et al. 2003b. Air pollution and hospital admissions for ischemic heart diseases among individuals 64+ years of age residing in Seoul, Korea. *Arch Environ Health* 58(10):617-623.
- +Lee JT, Kim H, Song H, et al. 2002. Air pollution and asthma among children in Seoul, Korea. *Epidemiology* 13(4):481-484.
- +Lee YL, Shaw CK, Su HJ, et al. 2003c. Climate, traffic-related air pollutants and allergic rhinitis prevalence in middle-school children in Taiwan. *Eur Respir J* 21(6):964-970.
- +Lee JT, Son JY, Cho YS. 2007c. A comparison of mortality related to urban air particles between periods with Asian dust days and without Asian dust days in Seoul, Korea, 2000-2004. *Environ Res* 105:409-413.
- \*Leeder JS, Kearns GL. 1997. Pharmacogenetics in pediatrics: Implications for practice. *Pediatr Clin North Am* 44(1):55-77.
- +\*Leem JH, Kaplan BM, Shim YK, et al. 2006. Exposures to air pollutants during pregnancy and preterm delivery. *Environ Health Perspect* 114(6):905-910.
- \*Leikin JB. 1996. Carbon monoxide detectors and emergency physicians. *J Emerg Med* 14(1):90-94.
- \*Leikin JB, Vogel S. 1986. Carbon monoxide levels in cardiac patients in an urban emergency department. *Am J Emerg Med* 4:126-128.
- \*Leikin JB, Wills BK. 2009. Long-term psychiatric consequences from carbon monoxide exposure: Progression of endogenous cause or toxicant effect? *Crit Care Med* 37(6):2116-2118.
- \*Leikin JB, Kaufman D, Lipscomb JW, et al. 1990. Methylene chloride: Report of five exposures and two deaths. *Am J Emerg Med* 8:534-537.
- \*Leung H. 1993. Physiologically-based pharmacokinetic modelling. In: Ballantyne B, Marrs T, Turner P, eds. *General and applied toxicology*. Vol. 1. New York, NY: Stockton Press, 153-164.
- \*Levasseur L, Galliot-Guilley M, Richter F, et al. 1996. Effects of mode of inhalation of carbon monoxide and of normobaric oxygen administration on carbon monoxide elimination from the blood. *Hum Exp Toxicol* 15(11):898-903.

## 9. REFERENCES

- +\*Levy D, Sheppard L, Checkoway H, et al. 2001. A case-crossover analysis of particulate matter air pollution and out-of-hospital primary cardiac arrest. *Epidemiology* 12(2):193-199.
- \*Lewis RJ. 2007. *Hawley's condensed chemical dictionary*. 15th ed. Hoboken, NJ: John Wiley & Sons, Inc., 235.
- Li XY, Zhang C, Wang SF, et al. 2008. Maternally administered lipopolysaccharide (LPS) upregulates the expression of heme oxygenase-1 in fetal liver: The role of reactive oxygen species. *Toxicol Lett* 176(3):169-177.
- Li Volti G, Vanella L, Gazzolo D, et al. 2008. Carbon monoxide: Vasoconstrictor or vasodilator? That's the question. *Am J Physiol Renal Physiol* 295(4):F901-F903.
- +\*Liao D, Duan Y, Whitsel EA, et al. 2004. Association of higher levels of ambient criteria pollutants with impaired cardiac autonomic control: A population-based study. *Am J Epidemiol* 159(8):768-777.
- +\*Liao D, Heiss G, Chinchilli VM, et al. 2005. Association of criteria pollutants with plasma hemostatic/inflammatory markers: A population-based study. *J Expo Anal Environ Epidemiol* 15(4):319-328.
- \*Lide DR, ed. 2008. *CRC handbook of chemistry and physics*. 88th ed. New York, NY: CRC Press, 4-56.
- \*Light A, Grass C, Pursley D, et al. 2007. Carboxyhemoglobin levels in smokers vs. non-smokers in a smoking environment. In: 2007 Open forum abstracts. American Association for Respiratory Care's International Respiratory Congress. Daedalus Enterprises.  
[http://www.rcjournal.com/abstracts/2007/?id=aarc07\\_111](http://www.rcjournal.com/abstracts/2007/?id=aarc07_111). May 8, 2009.
- +Lin CA, Martins MA, Farhat SC, et al. 1999. Air pollution and respiratory illness of children in Sao Paulo, Brazil. *Paediatr Perinat Epidemiol* 13(4):475-488.
- +\*Lin CA, Pereira LA, Nishioka DC, et al. 2004a. Air pollution and neonatal deaths in Sao Paulo, Brazil. *Braz J Med Biol Res* 37(5):765-770.
- +\*Lin CM, Li CY, Yang GY, et al. 2004b. Association between maternal exposure to elevated ambient sulfur dioxide during pregnancy and term low birth weight. *Environ Res* 96(1):41-50.
- +\*Lin M, Chen Y, Burnett RT, et al. 2003. Effect of short-term exposure to gaseous pollution on asthma hospitalisation in children: A bi-directional case-crossover analysis. *J Epidemiol Community Health* 57:50-55.
- +Lin M, Chen Y, Villeneuve PJ, et al. 2004c. Gaseous air pollutants and asthma hospitalization of children with low household income in Vancouver, British Columbia, Canada. *Am J Epidemiol* 159(3):294-303.
- +\*Lin M, Stieb DM, Chen Y. 2005. Coarse particulate matter and hospitalization for respiratory infections in children younger than 15 years in Toronto: A case-crossover analysis. *Pediatrics* 116(2):e235-e240.
- \*Linderholm H, Lundstrom P. 1969. Endogenous carbon monoxide production and blood loss at delivery. *Acta Obstet Gynecol Scand* 48:362-370.

## 9. REFERENCES

- +\*Linn WS, Szlachcic Y, Gong H, et al. 2000. Air pollution and daily hospital admissions in metropolitan Los Angeles. *Environ Health Perspect* 108(5):427-434.
- +\*Lipfert FW, Morris SC. 2002. Temporal and spatial relations between age specific mortality and ambient air quality in the United States: Regression results for counties, 1960-97. *Occup Environ Med* 59(3):156-174.
- +\*Lipfert FW, Baty JD, Miller JP, et al. 2006b. PM<sub>2.5</sub> constituents and related air quality variables as predictors of survival in a cohort of U.S. military veterans. *Inhal Toxicol* 18(9):645-657.
- +\*Lipfert FW, Morris SC, Wyzga RE. 2000. Daily mortality in the Philadelphia metropolitan area and size-classified particulate matter. *J Air Waste Manag Assoc* 50(8):1501-1513.
- +\*Lipfert FW, Wyzga RE, Baty JD, et al. 2006a. Traffic density as a surrogate measure of environmental exposures in studies of air pollution health effects: Long-term mortality in a cohort of US veterans. *Atmos Environ* 40:154-169.
- +Lippmann M, Ito K, Nadas A, et al. 2000. Association of particulate matter components with daily mortality and morbidity in urban populations. *Res Rep Health Eff Inst*:5-72, discussion 73-82.
- \*Liss PS, Slater PG. 1974. Flux of gases across the air-sea interface. *Nature* 247:181-184.
- +\*Liu S, Krewski D, Shi Y, et al. 2003. Association between gaseous ambient air pollutants and adverse pregnancy outcomes in Vancouver, Canada. *Environ Health Perspect* 111(14):1773-1778.
- +\*Liu S, Krewski D, Shi Y, et al. 2007. Association between maternal exposure to ambient air pollutants during pregnancy and fetal growth restriction. *J Expo Sci Environ Epidemiol* 17(5):426-432.
- Liu Y, Fechter LD. 1995. MK-801 protects against carbon monoxide-induced hearing loss. *Toxicol Appl Pharmacol* 132:196-202.
- \*Livingston AL. 1978. Forage plant estrogens. *J Toxicol Environ Health* 4(2-3):301-324.
- \*Lo CP, Chen SY, Lee KW, et al. 2007. Brain injury after acute carbon monoxide poisoning: Early and late complications. *AJR Am J Roentgenol* 189(4):W205-W211.
- Loennechen JP, Beisvag V, Arbo I, et al. 1999. Chronic carbon monoxide exposure *in vivo* induces myocardial endothelin-1 expression and hypertrophy in rat. *Pharmacol Toxicol* 85(4):192-197.
- Longo LD. 1976. Carbon monoxide: Effects on oxygenation of the fetus *in utero*. *Science* 194(4264):523-525.
- \*Longo LD. 1977. The biological effects of carbon monoxide on the pregnant woman, fetus, and newborn infant. *Am J Obstet Gynecol* 129(1):69-103.
- \*Longo LD, Hill EP. 1977. Carbon monoxide uptake and elimination in fetal and maternal sheep. *Am J Physiol Heart Circ Physiol* 232:H324-H330.
- Longo M, Jain V, Vedernikov YP, et al. 1999. Effect of nitric oxide and carbon monoxide on uterine contractility during human and rat pregnancy. *Am J Obstet Gynecol* 181:981-988.

## 9. REFERENCES

- +\*Lopez I, Acuna D, Webber DS, et al. 2003. Mild carbon monoxide exposure diminishes selectively the integrity of the cochlea of the developing rat. *J Neurosci Res* 74(5):666-675.
- +\*Lopez IA, Acuna D, Beltran-Parrazal L, et al. 2008. Oxidative stress and the deleterious consequences to the rat cochlea after prenatal chronic mild exposure to carbon monoxide in air. *Neuroscience* 151(3):854-867.
- \*Lou BS, Snyder JK, Marshall P, et al. 2000. Resonance Raman studies indicate a unique heme active site in prostaglandin H synthase. *Biochemistry* 39:12424-12434.
- Ludbrook GL, Helps SC, Gorman DF, et al. 1992. The relative effects of hypoxic hypoxia and carbon monoxide on brain function in rabbits. *Toxicology* 75(1):71-80.
- +Luginaah IN, Fung KY, Gorey KM, et al. 2005. Association of ambient air pollution with respiratory hospitalization in a government-designated "area of concern": The case of Windsor, Ontario. *Environ Health Perspect* 113(3):290-296.
- Lund AK, Knuckles TL, Obot Akata C, et al. 2007. Gasoline exhaust emissions induce vascular remodeling pathways involved in atherosclerosis. *Toxicol Sci* 95(2):485-494.
- \*Lundquist P, Lennart R, Sorbo B. 1989. The role of hydrogen cyanide and carbon monoxide in fire casualties: A prospective study. *Forensic Sci Int* 43:9-14.
- \*Luomanmäki K, Coburn RF. 1969. Effects of metabolism and distribution of carbon monoxide on blood and body stores. *Am J Physiol* 217(2):354-363.
- Lyll F, Myatt L. 2002. The role of the placenta in pre-eclampsia--a workshop report. *Placenta* 23(Suppl A):S142-S145.
- Lyll F, Barber A, Myatt L, et al. 2000. Hemeoxygenase expression in human placenta and placental bed implies a role in regulation of trophoblast invasion and placental function. *FASEB J* 14(1):208-219.
- MacMillan V. 1975. Regional cerebral blood flow of the rat in acute carbon monoxide intoxication. *Can J Physiol Pharmacol* 53(4):644-650.
- Mactutus CF, Fechter LD. 1984. Prenatal exposure to carbon monoxide: Learning and memory deficits. *Science* 223(4634):409-411.
- +\*Mactutus CF, Fechter LD. 1985. Moderate prenatal carbon monoxide exposure produces persistent, and apparently permanent, memory deficits in rats. *Teratology* 31(1):1-12.
- +Maheswaran R, Haining RP, Brindley P, et al. 2005a. Outdoor air pollution, mortality, and hospital admissions from coronary heart disease in Sheffield, UK: A small-area level ecological study. *Eur Heart J* 26(23):2543-2549.
- +\*Maheswaran R, Haining RP, Brindley P, et al. 2005b. Outdoor air pollution and stroke in Sheffield, United Kingdom: A small-area level geographical study. *Stroke* 36(2):239-243.

## 9. REFERENCES

- \*Mahoney AM, Stimpson CL, Scott KL, et al. 2007. Noninvasive measurement of carboxyhemoglobin levels for adjustment of diffusion capacity measured during pulmonary function testing. *Respir Care* 52(12):1741-1743.
- \*Mahoney JJ, Vreman HJ, Stevenson DK, et al. 1993. Measurement of carboxyhemoglobin and total hemoglobin by five specialized spectrophotometers (CO-oximeters) in comparison with reference methods. *Clin Chem* 39(8):1693-1700.
- +\*Maisonet M, Bush TJ, Correa A, et al. 2001. Relation between ambient air pollution and low birth weight in the Northeastern United States. *Environ Health Perspect* 109(Suppl 3):351-356.
- \*Mancuso C, Preziosi P, Grossman AB, et al. 1997. The role of carbon monoxide in the regulation of neuroendocrine function. *Neuroimmunomodulation* 4:225-229.
- Manier G, Moinard J, Techoueyres P, et al. 1991. Pulmonary diffusion limitation after prolonged strenuous exercise. *Respir Physiol* 83(2):143-153.
- +\*Mann JK, Tager IB, Lurmann F, et al. 2002. Air pollution and hospital admissions for ischemic heart disease in persons with congestive heart failure or arrhythmia. *Environ Health Perspect* 110(12):1247-1252.
- +\*Mannes T, Jalaludin B, Morgan G, et al. 2005. Impact of ambient air pollution on birth weight in Sydney, Australia. *Occup Environ Med* 62(8):524-530.
- +Mar TF, Norris GA, Koenig JQ, et al. 2000. Associations between air pollution and mortality in Phoenix, 1995-1997. *Environ Health Perspect* 108(4):347-353.
- \*Margulies JL. 1986. Acute carbon monoxide poisoning during pregnancy. *Am J Emerg Med* 4:516-519.
- Marshall MD, Kales SN, Christiani DC, et al. 1995. Are reference intervals for carboxyhemoglobin appropriate? A survey of Boston area laboratories. *Clin Chem* 41(10):1434-1438.
- Marthan R, Castaing Y, Manier G, et al. 1985. Gas exchange alterations in patients with chronic obstructive lung disease. *Chest* 87(4):470-475.
- +Martins LC, Latorre Mdo R, Saldiva PH, et al. 2002. Air pollution and emergency room visits due to chronic lower respiratory diseases in the elderly: An ecological time-series study in Sao Paulo, Brazil. *J Occup Environ Med* 44(7):622-627.
- +Masjedi MR, Jamaati HR, Dokouhaki P, et al. 2003. The effects of air pollution on acute respiratory conditions. *Respirology* 8(2):213-230.
- \*Materna BL, Jones JR, Sutton PM, et al. 1992. Occupational exposures in California wildland fire fighting. *Am Ind Hyg Assoc J* 53(1):69-76.
- \*Maulik N, Engelman DT, Watanabe M, et al. 1996. Nitric oxide - a retrograde messenger for carbon monoxide signaling in ischemic heart. *Mol Cell Biochem* 157:75-86.
- Mayr FB, Spiel A, Leitner J, et al. 2005. Effects of carbon monoxide inhalation during experimental endotoxemia in humans. *Am J Respir Crit Care Med* 171(4):354-360.

## 9. REFERENCES

- \*Mayr U, Butsch A, Schneider S. 1992. Validation of two *in vitro* test systems for estrogenic activities with zearalenone, phytoestrogens and cereal extracts. *Toxicology* 74(2-3):135-149.
- \*McCartney ML. 1990. Sensitivity analysis applied to Coburn-Forster-Kane models of carboxyhemoglobin formation. *Am Ind Hyg Assoc J* 51(3):169-177.
- +McGowan JA, Hider RN, Chacko E, et al. 2002. Particulate air pollution and hospital admissions in Christchurch, New Zealand. *Aust N Z J Public Health* 26:23-29.
- McGrath JJ. 1992. Effects of altitude on endogenous carboxyhemoglobin levels. *J Toxicol Environ Health* 35:127-133.
- \*McGrath JJ, Schreck RM, Lee PS. 1993. Carboxyhemoglobin levels in humans: Effects of altitude. *Inhal Toxicol* 5:241-249.
- +\*McGregor HP, Westcott K, Walker DW. 1998. The effect of prenatal exposure to carbon monoxide on breathing and growth of the newborn guinea pig. *Pediatr Res* 43(1):126-131.
- McLaughlin BE, Hutchinson JM, Graham CH, et al. 2000. Heme oxygenase activity in term human placenta. *Placenta* 21(8):870-873.
- McLaughlin BE, Lash GE, Graham CH, et al. 2001. Endogenous carbon monoxide formation by chorionic villi of term human placenta. *Placenta* 22(10):886-888.
- McLaughlin BE, Lash GE, Smith GN, et al. 2003. Heme oxygenase expression in selected regions of term human placenta. *Exp Biol Med (Maywood)* 228(5):564-567.
- McLean M, Bowman M, Clifton V, et al. 2000. Expression of the heme oxygenase-carbon monoxide signalling system in human placenta. *J Clin Endocrinol Metab* 85(6):2345-2349.
- +\*Medeiros A, Gouveia N. 2005. Relationship between low birthweight and air pollution in the city of Sao Paulo, Brazil. *Rev Saude Publica* 39(6):965-972.
- Meilin S, Rogatsky GG, Thom SR, et al. 1996. Effects of carbon monoxide on the brain may be mediated by nitric oxide. *J Appl Physiol* 81(3):1078-1083.
- Melin A, Obert P, Rebocho M, et al. 2002. Cardiac morphology and function following long-term exposure to carbon monoxide at high altitude in rats. *J Toxicol Environ Health A* 65(23):1981-1998.
- +Meng YY, Wilhelm M, Rull RP, et al. 2007. Traffic and outdoor air pollution levels near residences and poorly controlled asthma in adults. *Ann Allergy Asthma Immunol* 98(5):455-463.
- \*Mercke C, Lundh B. 1976. Erythrocyte filterability and heme catabolism during the menstrual cycle. *Ann Intern Med* 85:322-324.
- Mercke C, Cavallin-Stahl E, Lundh B. 1975a. Diurnal variation in endogenous production of carbon monoxide. Effect of caloric restriction. *Acta Med Scand* 198(3):161-164.
- Mercke C, Cavallin-Stahl E, Lundh B. 1975b. Carbon monoxide production and reticulocyte count in normal women. Effect of contraceptive drugs and smoking. *Acta Med Scand* 198(3):155-160.

## 9. REFERENCES

- Mereu G, Cammalleri M, Fa M, et al. 2000. Prenatal exposure to a low concentration of carbon monoxide disrupts hippocampal long-term potentiation in rat offspring. *J Pharmacol Exp Ther* 294(2):728-734.
- Merigan WH, McIntire RW. 1976. Effects of carbon monoxide on responding under a progressive ratio schedule in rats. *Physiol Behav* 16(4):407-412.
- +\*Metzger KB, Klein M, Flanders WD, et al. 2007. Ambient air pollution and cardiac arrhythmias in patients with implantable defibrillators. *Epidemiology* 18(5):585-592.
- \*+Metzger KB, Tolbert PE, Klein M, et al. 2004. Ambient air pollution and cardiovascular emergency department visits. *Epidemiology* 15(1):46-56.
- \*Meyer J, Prien T, Van Aken H, et al. 1998. Arterio-venous carboxyhemoglobin difference suggests carbon monoxide production by human lungs. *Biochem Biophys Res Commun* 244(1):230-232.
- Middendorff R, Kumm M, Davidoff MS, et al. 2000. Generation of cyclic guanosine monophosphate by heme oxygenases in the human testis—a regulatory role for carbon monoxide in Sertoli cells? *Biol Reprod* 63(2):651-657.
- +Migliaretti G, Dalmaso P, Gregori D. 2007. Air pollution effects on the respiratory health of the resident adult population in Turin, Italy. *Int J Environ Health Res* 17(5):369-379.
- +\*Miller KA, Siscovick DS, Sheppard L, et al. 2007. Long-term exposure to air pollution and incidence of cardiovascular events in women. *N Engl J Med* 356(5):447-458.
- \*Miller SM, Matross DM, Miller ADB, et al. 2008. Sources of carbon monoxide and formaldehyde in North America determined from high-resolution atmospheric data. *Atmos Chem Phys* 8:7673-7696.
- \*Mishra S, Fujita T, Lama VN, et al. 2006. Carbon monoxide rescues ischemic lungs by interrupting MAPK-driven expression of early growth response 1 gene and its downstream target genes. *Proc Natl Acad Sci* 103(13):5191-5196.
- Mitchell DS, Packham SC, Fitzgerald WE. 1978. Effects of ethanol and carbon monoxide on two measures of behavioral incapacitation of rats. *Proc West Pharmacol Soc* 21:427-431.
- Moessinger AC, Shen CT, Kiu GT, et al. 1990. Maternal exposure to carbon monoxide inhibits fetal lung cell multiplication [Abstract]. *Pediatr Res* 27(4 Pt 2):48A.
- \*Moir D, Rickert WS, Levasseur G, et al. 2008. A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. *Chem Res Toxicol* 21(2):494-502.
- \*Moller P, Sylven C. 1981. Myoglobin in human skeletal muscle. *Scand J Clin Lab Invest* 41(5):479-482.
- Montgomery MR, Rubin RJ. 1971. The effect of carbon monoxide inhalation on *in vivo* drug metabolism in the rat. *J Pharmacol Exp Ther* 179(3):465-473.

## 9. REFERENCES

- +\*Moolgavkar SH. 2000a. Air pollution and daily mortality in three U.S. counties. *Environ Health Perspect* 108(8):777-784.
- +Moolgavkar SH. 2000b. Air pollution and hospital admissions for chronic obstructive pulmonary disease in three metropolitan areas in the United States. *Inhal Toxicol* 12(Suppl 4):75-90.
- +Moolgavkar SH. 2003. Air pollution and daily mortality in two U.S. counties: Season-specific analyses and exposure-response relationships. *Inhal Toxicol* 15(9):877-907.
- \*Moore SJ, Norris JC, Walsh DA, et al. 1987. Antidotal use of methemoglobin forming cyanide antagonists in concurrent carbon monoxide/cyanide intoxication. *J Pharmacol Exp Ther* 242(1):70-73.
- Morita T, Mitsialis SA, Koike H, et al. 1997. Carbon monoxide controls the proliferation of hypoxic vascular smooth muscle cells. *J Biol Chem* 272(52):32804-32809.
- Morita T, Perrella MA, Lee ME, et al. 1995. Smooth muscle cell-derived carbon monoxide is a regulator of vascular cGMP. *Proc Natl Acad Sci USA* 92(5):1475-1479.
- Morris RD, Naumova EN. 1998. Carbon monoxide and hospital admissions for congestive heart failure: Evidence of an increased effect at low temperatures. *Environ Health Perspect* 106(10):649-653.
- Morris RD, Naumova EN, Munasinghe RL. 1995. Ambient air pollution and hospitalization for congestive heart failure among elderly people in seven large US cities. *Am J Public Health* 85(10):1361-1365.
- +\*Morse CI, Pritchard LJ, Wust RC, et al. 2008. Carbon monoxide inhalation reduces skeletal muscle fatigue resistance. *Acta Physiol (Oxf)* 192(3):397-401.
- \*Morselli PL, Franco-Morselli R, Bossi L. 1980. Clinical pharmacokinetics in newborns and infants: Age-related differences and therapeutic implications. *Clin Pharmacokinet* 5(6):485-527.
- +\*Mortimer K, Neugebauer R, Lurmann F, et al. 2008. Air pollution and pulmonary function in asthmatic children: Effects of prenatal and lifetime exposures. *Epidemiology* 19(4):550-557.
- \*Myers RA, Snyder SK, Majerus TC. 1985. Cutaneous blisters and carbon monoxide poisoning. *Ann Emerg Med* 14(6):603-606.
- Nabeshima T, Katoh A, Ishimaru H, et al. 1991. Carbon monoxide-induced delayed amnesia, delayed neuronal death and change in acetylcholine concentration in mice. *J Pharmacol Exp Ther* 256(1):378-384.
- Nabeshima T, Yoshida S, Morinaka H, et al. 1990. MK-801 ameliorates delayed amnesia, but potentiates acute amnesia induced by CO. *Neurosci Lett* 108(3):321-327.
- Naik JS, Walker BR. 2003. Heme oxygenase-mediated vasodilation involves vascular smooth muscle cell hyperpolarization. *Am J Physiol Heart Circ Physiol* 285:H220-H228.
- \*Nakao A, Kaczorowski DJ, Sugimoto R, et al. 2008. Application of heme oxygenase-1, carbon monoxide and biliverdin for the prevention of intestinal ischemia/reperfusion injury. *J Clin Biochem Nutr* 42(2):78-88.

## 9. REFERENCES

- \*Nakayama M, Takahashi K, Kitamuro T, et al. 2000. Repression of heme oxygenase-1 by hypoxia in vascular endothelial cells. *Biochem Biophys Res Commun* 271:665-671.
- \*NAS/NRC. 1989. Report of the oversight committee. In: *Biologic markers in reproductive toxicology*. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press, 15-35.
- Ndisang JF, Tabien HE, Wang R. 2004. Carbon monoxide and hypertension. *J Hypertens* 22(6):1057-1074.
- Neggers YH, Singh J. 2006. Zinc supplementation to protein-deficient diet in CO-exposed mice decreased fetal mortality and malformation. *Biol Trace Elem Res* 114:269-279.
- +Neidell MJ. 2004. Air pollution, health, and socio-economic status: The effect of outdoor air quality on childhood asthma. *J Health Econ* 23(6):1209-1236.
- Newby D, Cousins F, Myatt L, et al. 2005. Heme oxygenase expression in cultured human trophoblast cells during *in vitro* differentiation: Effects of hypoxia. *Placenta* 26(2-3):201-209.
- Newby MB, Roberts RJ, Bhatnagar RK. 1978. Carbon monoxide- and hypoxia-induced effects on catecholamines in the mature and developing rat brain. *J Pharmacol Exp Ther* 206(1):61-68.
- \*Niden AH. 1971. The effects of low levels of carbon monoxide on the fine structure of the terminal airways. *Am Rev Respir Dis* 103:898.
- \*NIOSH. 2005. Carbon monoxide. In: *NIOSH pocket guide to chemical hazards*. Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention. <http://www.cdc.gov/niosh/npg/>. May 19, 2009.
- \*Norman CA, Halton DM. 1990. Is carbon monoxide a workplace teratogen? A review and evaluation of the literature. *Ann Occup Hyg* 34(4):335-347.
- \*Norris JC, Moore SJ, Hume AS. 1986. Synergistic lethality induced by the combination of carbon monoxide and cyanide. *Toxicology* 40:121-129.
- +Norris G, YoungPong SN, Koenig JQ, et al. 1999. An association between fine particles and asthma emergency department visits for children in Seattle. *Environ Health Perspect* 107(6):489-493.
- \*Novelli PC. 1999. CO in the atmosphere: Measurement techniques and related issues. *Chemosphere Global Change Sci* 1:115-126.
- \*NRC. 1977. *Medical and biologic effects of environmental pollutants. Carbon monoxide*. Washington, DC: National Academy of Sciences, PB274965.
- \*NRC. 1993. *Pesticides in the diets of infants and children*. Washington, DC: National Research Council. National Academy Press.
- \*NTP. 2005. *Report on carcinogens, 11th edition*. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. <http://ntp-server.niehs.nih.gov/ntp/roc/toc11.html>. May 11, 2009.

## 9. REFERENCES

- \*Ny L, Alm P, Ekstrom P, et al. 1996. Localization and activity of haem oxygenase and functional effects of carbon monoxide in the feline lower oesophageal sphincter. *Br J Pharmacol* 118:392-399.
- Odrich MJ, Graham CH, Kimura KA, et al. 1998. Heme oxygenase and nitric oxide synthase in the placenta of the guinea-pig during gestation. *Placenta* 19(7):509-516.
- \*Ohta K. 1997. Diurnal variations of carbon monoxide concentration in the equatorial pacific upwelling region. *J Oceanogr* 53:173-178.
- Okeda R, Funata N, Takano T, et al. 1981. The pathogenesis of carbon monoxide encephalopathy in the acute phase—physiological and morphological correlation. *Acta Neuropathol (Berl)* 54(1):1-10.
- \*Okeda R, Matsuo T, Kuroiwa T, et al. 1987. Regional cerebral blood flow of acute carbon monoxide poisoning in cats. *Acta Neuropathol* 72:389-393.
- Olson KR, Donald JA. 2009. Nervous control of circulation—The role of gasotransmitters, NO, CO, and H<sub>2</sub>S. *Acta Histochem* 111(3):244-256.
- Omaye ST. 2002. Metabolic modulation of carbon monoxide toxicity. *Toxicology* 180(2):139-150.
- \*O'Neil MJ, Heckelman PE, Koch CB, et al. 2006. The Merck index. An encyclopedia of chemicals, drugs, and biologicals. 14th ed. Whitehouse Station, NJ: Merck & Co., Inc., 294.
- \*OSHA. 1991. Sampling and analytical methods: Carbon monoxide in workplace atmosphere, ID-210. Washington, DC: Occupational Safety & Health Administration. <http://www.osha.gov/dts/sltc/methods/inorganic/id210/id210.html>. May 11, 2009.
- \*OSHA. 2000. Occupational safety and health guideline for carbon monoxide. Washington, DC: Occupational Safety & Health Administration. <http://www.osha.gov/SLTC/healthguidelines/carbonmonoxide/recognition.html>. May 11, 2009.
- \*OSHA. 2009. Toxic and Hazardous Substances. Occupational safety and health standards. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1910.1000, Table Z-1. <http://www.osha.gov/comp-links.html>. May 19, 2009.
- +Ostro BD, Hurley S, Lipsett MJ. 1999. Air pollution and daily mortality in the Coachella Valley, California: A study of PM<sub>10</sub> dominated by coarse particles. *Environ Res* 81(3):231-238.
- OTA. 1990. Neurotoxicity: Identifying and controlling poisons of the nervous system. Washington, DC: Office of Technology Assessment.
- Ott WR, Mage DT. 1978. Interpreting urban carbon monoxide concentrations by means of a computerized blood COHb model. *J Air Pollut Control Assoc* 28(9):911-916.
- \*Owen GM, Brozek J. 1966. Influence of age, sex and nutrition on body composition during childhood and adolescence. In: Falkner F, ed. Human development. Philadelphia, PA: WB Saunders, 222-238.
- Ozawa N, Goda N, Makino N, et al. 2002. Leydig cell-derived heme oxygenase-1 regulates apoptosis of premeiotic germ cells in response to stress. *J Clin Invest* 109(4):457-467.

## 9. REFERENCES

- Pannen BH, Bauer M. 1998. Differential regulation of hepatic arterial and portal venous vascular resistance by nitric oxide and carbon monoxide in rats. *Life Sci* 62(22):2025-2033.
- +Park H, Lee B, Ha EH, et al. 2002. Association of air pollution with school absenteeism due to illness. *Arch Pediatr Adolesc Med* 156(12):1235-1239.
- +\*Park JW, Lim YH, Kyung SY, et al. 2005a. Effects of ambient particulate matter on peak expiratory flow rates and respiratory symptoms of asthmatics during Asian dust periods in Korea. *Respirology* 10(4):470-476.
- +\*Park SK, O'Neill MS, Vokonas PS, et al. 2005b. Effects of air pollution on heart rate variability: The VA normative aging study. *Environ Health Perspect* 113(3):304-309.
- +\*Parker JD, Woodruff TJ, Basu R, et al. 2005. Air pollution and birth weight among term infants in California. *Pediatrics* 115(1):121-128.
- \*Parkinson RB, Hopkins RO, Cleavinger HB, et al. 2002. White matter hyperintensities and neuropsychological outcome following carbon monoxide poisoning. *Neurology* 58(10):1525-1532.
- Parrish DD. 2006. Critical evaluation of US on-road vehicle emission inventories. *Atmos Environ* 40:2288-2300.
- Parving H, Ohlsson K, Buchardt Hansen HJ, et al. 1972. Effect of carbon monoxide exposure on capillary permeability to albumin and 2-macroglobulin. *Scand J Clin Lab Invest* 29:381-388.
- Patel AP, John Moody A, Handy RD, et al. 2003. Carbon monoxide exposure in rat heart: Glutathione depletion is prevented by antioxidants. *Biochem Biophys Res Commun* 302(2):392-396.
- \*Paulson OB, Parving HH, Olesen J, et al. 1973. Influence of carbon monoxide and of hemodilution on cerebral blood flow and blood gases in man. *J Appl Physiol* 35(1):111-116.
- +\*Peel JL, Metzger KB, Klein M, et al. 2007. Ambient air pollution and cardiovascular emergency department visits in potentially sensitive groups. *Am J Epidemiol* 165(6):625-633.
- +\*Peel JL, Tolbert PE, Klein M, et al. 2005. Ambient air pollution and respiratory emergency department visits. *Epidemiology* 16(2):164-174.
- +\*Pekkanen J, Brunner EJ, Anderson HR, et al. 2000. Daily concentrations of air pollution and plasma fibrinogen in London. *Occup Environ Med* 57(12):818-822.
- \*Penney DG. 1988. Hemodynamic response to carbon monoxide. *Environ Health Perspect* 77:121-130.
- Penney DG, Caldwell TM. 1984. Growth of the young heart: Influence of nutrition and stress. *Growth* 48(4):483-498.
- Penney DG, Weeks TA. 1979. Age dependence of cardiac growth in the normal and carbon monoxide-exposed rat. *Dev Biol* 71(1):153-162.
- \*Penney D, Benjamin M, Dunham E. 1974a. Effect of carbon monoxide on cardiac weight as compared with altitude effects. *J Appl Physiol* 37(1):80-84.

## 9. REFERENCES

- \*Penney D, Dunham E, Benjamin M. 1974b. Chronic carbon monoxide exposure: Time course of hemoglobin, heart weight and lactate dehydrogenase isozyme changes. *Toxicol Appl Pharmacol* 28:493-497.
- Penney DG, Barthel BG, Skoney JA. 1984. Cardiac compliance and dimensions in carbon monoxide-induced cardiomegaly. *Cardiovasc Res* 18(5):270-276.
- +\*Penney DG, Baylerian MS, Thill JE, et al. 1982. Postnatal carbon monoxide exposure: Immediate and lasting effects in the rat. *Am J Physiol* 243(2):H328-339.
- +\*Penney DG, Baylerian MS, Thill JE, et al. 1983. Cardiac response of the fetal rat to carbon monoxide exposure. *Am J Physiol* 244(2):H289-H297.
- Penney DG, Caldwell-Ayre TM, Gargulinski RB, et al. 1987. Effect of litter size (i.e. nutrition) on carbon monoxide-induced persistent heart changes. *Growth* 51(3):321-333.
- \*Penney DG, Davidson SB, Gargulinski RB, et al. 1988. Heart and lung hypertrophy, changes in blood volume, hematocrit and plasma renin activity in rats chronically exposed to increasing carbon. *J Appl Toxicol* 8(3):171-178.
- Penney DG, Tucker A, Bambach GA. 1992. Heart and lung alterations in neonatal rats exposed to CO or high altitude. *J Appl Physiol* 73(5):1713-1719.
- +Penttinen P, Tiittanen P, Pekkanen J. 2004. Mortality and air pollution in metropolitan Helsinki, 1988-1996. *Scand J Work Environ Health* 30 (Suppl 2):19-27.
- +\*Penttinen P, Timonen KL, Tiittanen P, et al. 2001. Ultrafine particles in urban air and respiratory health among adult asthmatics. *Eur Respir J* 17(3):428-435.
- Perez H, Ruiz S. 1995. Medullary responses to chemoreceptor activation are inhibited by locus coeruleus and nucleus raphe magnus. *Neuroreport* 6:1245-1248.
- \*Perrella M, Di Cera E. 1999. CO ligation intermediates and the mechanism of hemoglobin cooperativity. *J Biol Chem* 274:2605-2608.
- Pesola GR, Huggins G, Sherpa TY. 2006. Abnormal predicted diffusion capacities in healthy Asians: An inequality with a solution. *Respiration* 73(6):799-807.
- Pesola GR, Sunmonu Y, Huggins G, et al. 2004. Measured diffusion capacity versus prediction equation estimates in blacks without lung disease. *Respiration* 71(5):484-492.
- +\*Peters A, Dockery DW, Muller JE, et al. 2001. Increased particulate air pollution and the triggering of myocardial infarction. *Circulation* 103(23):2810-2815.
- +\*Peters A, Liu E, Verrier RL, et al. 2000b. Air pollution and incidence of cardiac arrhythmia. *Epidemiology* 11:11-17.
- +\*Peters A, Perz S, Doring A, et al. 1999. Increases in heart rate during an air pollution episode. *Am J Epidemiol* 150(10):1094-1098.

## 9. REFERENCES

- +\*Peters A, Skorkovsky J, Kotesovec F, et al. 2000a. Associations between mortality and air pollution in central Europe. *Environ Health Perspect* 108(4):283-287.
- Petersen LC. 1977. The effect of inhibitors on the oxygen kinetics of cytochrome c oxidase. *Biochim Biophys Acta* 460:299-307.
- \*Peterson JE, Stewart RD. 1970. Absorption and elimination of carbon monoxide by inactive young men. *Arch Environ Health* 21(2):165-171.
- \*Peterson JE, Stewart RD. 1975. Predicting the carboxyhemoglobin levels resulting from carbon monoxide exposures. *J Appl Physiol* 39(4):633-638.
- \*Piantadosi CA. 2002. Biological chemistry of carbon monoxide. *Antioxid Redox Signal* 4:259-270.
- Piantadosi CA. 2008. Carbon monoxide, reactive oxygen signaling, and oxidative stress. *Free Radic Biol Med* 45(5):562-569.
- Piantadosi CA, Carraway MS, Suliman HB. 2006. Carbon monoxide, oxidative stress, and mitochondrial permeability pore transition. *Free Radic Biol Med* 40(8):1332-1339.
- Piantadosi CA, Tatro L, Zhang J. 1995. Hydroxyl radical production in the brain after CO hypoxia in rats. *Free Radic Biol Med* 18(3):603-609.
- \*Piantadosi CA, Zhang J, Demchenko IT. 1997a. Production of hydroxyl radical in the hippocampus after CO hypoxia or hypoxic hypoxia in the rat. *Free Radic Biol Med* 22(4):725-732.
- \*Piantadosi CA, Zhang J, Levin ED, et al. 1997b. Apoptosis and delayed neuronal damage after carbon monoxide poisoning in the rat. *Exp Neurol* 147(1):103-114.
- Pimay F, Dujardin J, Deroanne R, et al. 1971. Muscular exercise during intoxication by carbon monoxide. *J Appl Physiol* 31(4):573-575.
- \*Pitt BR, Radford EP, Gurtner GH, et al. 1979. Interaction of carbon monoxide and cyanide on cerebral circulation and metabolism. *Arch Environ Health* 34:354-359.
- Plummer AL. 2008. The carbon monoxide diffusing capacity: Clinical implications, coding, and documentation. *Chest* 134(3):663-667.
- +\*Pope CA, Burnett RT, Thun MJ, et al. 2002. Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA* 287(9):1132-1141.
- \*Pope CA, Thun MJ, Namboodiri MM, et al. 1995. Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. *Am J Respir Crit Care Med* 151:669-674.
- \*Pressman J, Warneck P. 1970. The stratosphere as a chemical sink for carbon monoxide. *J Atmos Sci* 27:155-163.
- Preziosi TJ, Lindenberg R, Levy D, et al. 1970. An experimental investigation in animals of the functional and morphologic effects of single and repeated exposures to high and low concentrations of carbon monoxide. *Ann N Y Acad Sci* 174(1):369-384.

## 9. REFERENCES

- +\*Prigge E, Hochrainer D. 1977. Effects of carbon monoxide inhalation on erythropoiesis and cardiac hypertrophy in fetal rats. *Toxicol Appl Pharmacol* 42(1):225-228.
- Putz VR. 1979. The effects of carbon monoxide on dual-task performance. *Hum Factors* 21(1):13-24.
- +\*Rabinovitch N, Zhang L, Murphy JR, et al. 2004. Effects of wintertime ambient air pollutants on asthma exacerbations in urban minority children with moderate to severe disease. *J Allergy Clin Immunol* 114(5):1131-1137.
- \*Radford EP, Drizd TA. 1982. Blood carbon monoxide levels in persons 3-74 years of age: United States, 1976-80a. *Adv Data* 76:1-24.
- +Rainham DG, Smoyer-Tomic KE. 2003. The role of air pollution in the relationship between a heat stress index and human mortality in Toronto. *Environ Res* 93(1):9-19.
- Ramsey JM. 1973. Effects of single exposures of carbon monoxide on sensory and psychomotor response. *Am Ind Hyg Assoc J* 34(5):212-216.
- +Ranzi A, Gambini M, Spattini A, et al. 2004. Air pollution and respiratory status in asthmatic children: Hints for a locally based preventive strategy. AIRE study. *Eur J Epidemiol* 19(6):567-576.
- \*Rasmussen LS, Poulsen MG, Christiansen M, et al. 2004. Biochemical markers for brain damage after carbon monoxide poisoning. *Acta Anaesthesiol Scand* 48:469-473.
- \*Raub JA, Benignus VA. 2002. Carbon monoxide and the nervous system. *Neurosci Biobehav Rev* 26(8):925-940.
- \*Raub JA, Mathieu-Nolf M, Hampson NB, et al. 2000. Carbon monoxide poisoning--a public health perspective. *Toxicology* 145(1):1-14.
- Raven PB, Drinkwater BL, Ruhling RO, et al. 1974. Effect of carbon monoxide and peroxyacetyl nitrate on man's maximal aerobic capacity. *J Appl Physiol* 36(3):288-293.
- Reeves RB, Park HK. 1992. CO uptake kinetics of red cells and CO diffusing capacity. *Respir Physiol* 88(1-2):1-21.
- Remy S, Blancou P, Tesson L, et al. 2009. Carbon monoxide inhibits TLR-induced dendritic cell immunogenicity. *J Immunol* 182(4):1877-1884.
- \*Ren X, Dorrington KL, Robbins PA. 2001. Respiratory control in humans after 8 h of lowered arterial PO<sub>2</sub>, hemodilution, or carboxyhemoglobinemia. *J Appl Physiol* 90(4):1189-1195.
- +\*Resch H, Zawinka C, Weigert G, et al. 2005. Inhaled carbon monoxide increases retinal and choroidal blood flow in healthy humans. *Invest Ophthalmol Vis Sci* 46(11):4275-4280.
- \*Rhee JW, Leikin JB. 2007. Carbon monoxide detectors as preventive medicine. In: Penny DG, ed. *Carbon monoxide poisoning*. Boca Raton, FL: CRC Press, 305-311.
- +\*Rich DQ, Kim MH, Turner JR, et al. 2006a. Association of ventricular arrhythmias detected by implantable cardioverter defibrillator and ambient air pollutants in the St Louis, Missouri metropolitan area. *Occup Environ Med* 63(9):591-596.

## 9. REFERENCES

- +\*Rich DQ, Mittleman MA, Link MS, et al. 2006b. Increased risk of paroxysmal atrial fibrillation episodes associated with acute increases in ambient air pollution. *Environ Health Perspect* 114(1):120-123.
- +\*Rich DQ, Schwartz J, Mittleman MA, et al. 2005. Association of short-term ambient air pollution concentrations and ventricular arrhythmias. *Am J Epidemiol* 161(12):1123-1132.
- +\*Rich KE, Petkau J, Vedal S, et al. 2004. A case-crossover analysis of particulate air pollution and cardiac arrhythmia in patients with implantable cardioverter defibrillators. *Inhal Toxicol* 16(6-7):363-372.
- \*Richardson RS, Noyszewski EA, Saltin B, et al. 2002. Effect of mild carboxy-hemoglobin on exercising skeletal muscle: Intravascular and intracellular evidence. *Am J Physiol Regul Integr Comp Physiol* 283:R1131-R1139.
- \*Ringel SP, Klawans HL. 1972. Carbon monoxide-induced Parkinsonism. *J Neurol Sci* 16(3):245-251.
- +\*Riojas-Rodriguez H, Escamilla-Cejudo JA, Gonzalez-Hermosillo JA, et al. 2006. Personal PM2.5 and CO exposures and heart rate variability in subjects with known ischemic heart disease in Mexico City. *J Expo Sci Environ Epidemiol* 16(2):131-137.
- +\*Ritz B, Yu F. 1999. The effect of ambient carbon monoxide on low birth weight among children born in southern California between 1989 and 1993. *Environ Health Perspect* 107(1):17-25.
- +\*Ritz B, Wilhelm M, Zhao Y. 2006. Air pollution and infant death in southern California, 1989-2000. *Pediatrics* 118(2):493-502.
- +\*Ritz B, Wilhelm M, Hoggatt KJ, et al. 2007. Ambient air pollution and preterm birth in the environment and pregnancy outcomes study at the University of California, Los Angeles. *Am J Epidemiol* 166(9):1045-1052.
- +\*Ritz B, Yu F, Chapa G, et al. 2000. Effect of air pollution on preterm birth among children born in Southern California between 1989 and 1993. *Epidemiology* 11(5):502-511.
- +\*Ritz B, Yu F, Fruin S, et al. 2002. Ambient air pollution and risk of birth defects in Southern California. *Am J Epidemiol* 155(1):17-25.
- \*Rodgers PA, Vreman HJ, Dennery PA, et al. 1994. Sources of carbon monoxide (CO) in biological systems and applications of CO detection technologies. *Semin Perinatol* 18(1):2-10.
- \*Rodkey FL, O'Neal JD, Collison HA. 1969. Oxygen and carbon monoxide equilibria of human adult hemoglobin at atmospheric and elevated pressure. *Blood* 33(1):57-65.
- \*Rodkey FL, Hill TA, Pitts LL, et al. 1979. Spectrophotometric measurement of carboxyhemoglobin and methemoglobin in blood. *Clin Chem* 25(8):1388-1393.
- +\*Rodriguez C, Tonkin R, Heyworth J, et al. 2007. The relationship between outdoor air quality and respiratory symptoms in young children. *Int J Environ Health Res* 17(5):351-360.

## 9. REFERENCES

+Roemer WH, van Wijnen JH. 2001. Daily mortality and air pollution along busy streets in Amsterdam, 1987-1998. *Epidemiology* 12(6):649-653.

+\*Rosenlund M, Berglind N, Pershagen G, et al. 2006. Long-term exposure to urban air pollution and myocardial infarction. *Epidemiology* 17(4):383-390.

\*Roughton FJ. 1970. The equilibrium of carbon monoxide with human hemoglobin in whole blood. *Ann N Y Acad Sci* 174(1):177-188.

Roughton FJ, Forster RE. 1957. Relative importance of diffusion and chemical reaction rates in determining rate of exchange of gases in the human lung, with special reference to true diffusing capacity of pulmonary membrane and volume of blood in the lung capillaries. *J Appl Physiol* 11(2):290-302.

RTECS. 2008. Carbon monoxide. Registry of Toxic Effects on Chemical Substances. National Institute of Occupational Safety and Health. MDL Information Systems, Inc.

\*Rucker J, Tesler J, Fedorko L, et al. 2002. Normocapnia improves cerebral oxygen delivery during conventional oxygen therapy in carbon monoxide-exposed research subjects. *Ann Emerg Med* 40(6):611-618.

+\*Rückerl R, Greven S, Ljungman P, et al. 2007. Air pollution and inflammation (interleukin-6, C-reactive protein, fibrinogen) in myocardial infarction survivors. *Environ Health Perspect* 115(7):1072-1080.

+\*Rückerl R, Ibald-Mulli A, Koenig W, et al. 2006. Air pollution and markers of inflammation and coagulation in patients with coronary heart disease. *Am J Respir Crit Care Med* 173:432-441.

\*Ryter SW, Otterbein LE. 2004. Carbon monoxide in biology and medicine. *BioEssays* 26(3):270-280.

\*Ryter SW, Alam J, Choi AM. 2006. Heme oxygenase-1/carbon monoxide: From basic science to therapeutic applications. *Physiol Rev* 86(2):583-650.

+\*Salam MT, Millstein J, Li YF, et al. 2005. Birth outcomes and prenatal exposure to ozone, carbon monoxide, and particulate matter results from the Children's Health Study. *Environ Health Perspect* 113(11):1638-1644.

Saltzman BE, Fox SH. 1986. Biological significance of fluctuating concentrations of carbon monoxide. *Environ Sci Technol* 20(9):916-923.

+\*Samet JM, Dominici F, Zeger SL, et al. 2000. The national morbidity, mortality and air pollution study. Part 1. Methods and methodologic issues. *Res Rep Health Eff Inst*:5-14, discussion 75-84.

+\*Samoli E, Touloumi G, Schwartz J, et al. 2007. Short-term effects of carbon monoxide on mortality: An analysis within the APHEA project. *Environ Health Perspect* 115(11):1578-1583.

\*Sanderson MG. 2002. Emission of carbon monoxide by vegetation and soils. Hadley Centre technical note 36, Met Office.

Sari I, Zengin S, Ozer O, et al. 2008. Chronic carbon monoxide exposure is associated with increased electrocardiographic P-wave and QT dispersion. *Circulation* 118(12):E265-E266.

## 9. REFERENCES

- +\*Sarnat SE, Suh HH, Coull BA, et al. 2006. Ambient particulate air pollution and cardiac arrhythmia in a panel of older adults in Steubenville, Ohio. *Occup Environ Med* 63:700-706.
- +\*Sartiani L, Cerbai E, Lonardo G, et al. 2004. Prenatal exposure to carbon monoxide affects postnatal cellular electrophysiological maturation of the rat heart: A potential substrate for arrhythmogenesis in infancy. *Circulation* 109:419-423.
- \*Satran D, Henry CR, Adkinson C, et al. 2005. Cardiovascular manifestations of moderate to severe carbon monoxide poisoning. *J Am Coll Cardiol* 45:1513-1516.
- \*Sawai H, Kawada N, Yoshizato K, et al. 2003. Characterization of the heme environmental structure of cytoglobin, a fourth globin in humans. *Biochemistry* 42:5133-5142.
- +\*Schilderout JS, Sheppard L, Lumley T, et al. 2006. Ambient air pollution and asthma exacerbations in children: An eight-city analysis. *Am J Epidemiol* 164(6):505-517.
- Schoenfisch WH, Hoop KA, Struelens BS. 1980. Carbon monoxide absorption through the oral and nasal mucosae of cynomolgus monkeys. *Arch Environ Health* 35(3):152-154.
- +\*Schwartz J, Litonjua A, Suh H, et al. 2005. Traffic related pollution and heart rate variability in a panel of elderly subjects. *Thorax* 60(6):455-461.
- +Schwartz J, Norris G, Larson T, et al. 1999. Episodes of high coarse particle concentrations are not associated with increased mortality. *Environ Health Perspect* 107(5):339-342.
- Schwetz BA, Smith FA, Staples RE. 1978. Teratogenic potential of ethanol in mice, rats and rabbits. *Teratology* 18:385-392.
- Scott JR, Cukiernik MA, Ott MC, et al. 2009. Low-dose inhaled carbon monoxide attenuates the remote intestinal inflammatory response elicited by hindlimb ischemia-reperfusion. *Am J Physiol Gastrointest Liver Physiol* 296(1):G9-G14.
- Selvakumar S, Sharan M, Singh MP. 1992. Mathematical model for the exchange of gases in the lungs with special reference to carbon monoxide. *Med Biol Eng Comput* 30(5):525-532.
- Selvakumar S, Sharan M, Singh MP. 1993. A mathematical model for the elimination of carbon monoxide in humans. *J Theor Biol* 162:321-336.
- \*Setchell BP, Waites GMH. 1975. The blood testis barrier. In: Creep RO, Astwood EB, Geiger SR, eds. *Handbook of physiology: Endocrinology V*. Washington, DC: American Physiological Society, 143-172.
- Sethi JM. 2005. Carbon monoxide. *Crit Care Med* 33(12 Suppl):S496-S497.
- Shaoqing Y, Ruxin Z, Yinjian C, et al. 2008. Possible contribution of endogenous carbon monoxide to the development of allergic rhinitis in guinea pigs. *J Inflamm (Lond)* 5:23.
- Sharan M, Selvakumar S, Singh MP. 1990. Mathematical model for the computation of alveolar partial pressure of carbon monoxide. *Int J Biomed Comput* 26(3):135-147.

## 9. REFERENCES

- Sharma VS, Schmidt MR, Ranney HM. 1976. Dissociation of CO from carboxyhemoglobin. *J Biol Chem* 251(14):4267-4272.
- +Sharovsky R, Cesar LAM, Ramires JAF. 2004. Temperature, air pollution, and mortality from myocardial infarction in Sao Paulo, Brazil. *Braz J Med Biol Res* 37:1651-1657.
- Shephard RJ. 1984. Athletic performance and urban air pollution. *Can Med Assoc J* 131(2):105-109.
- +Sheppard L, Levy D, Norris G, et al. 1999. Effects of ambient air pollution on nonelderly asthma hospital admissions in Seattle, Washington, 1987-1994. *Epidemiology* 10:23-30.
- +\*Sheps DS, Adams KF, Bromberg PA, et al. 1987. Lack of effect of low levels of carboxyhemoglobin on cardiovascular function in patients with ischemic heart disease. *Arch Environ Health* 42(2):108-116.
- +\*Sheps DS, Herbst MC, Hinderliter AL, et al. 1990. Production of arrhythmias by elevated carboxyhemoglobin in patients with coronary artery disease. *Ann Intern Med* 113:343-351.
- \*Shibahara S, Nakayama M, Kitamuro T, et al. 2003. Repression of heme oxygenase-1 expression as a defense strategy in humans. *Exp Biol Med* 228:472-473.
- Shimazu T. 2001. Half-life of blood carboxyhemoglobin. *Chest* 119(2):661-663.
- \*Shimazu T, Ikeuchi H, Sugimoto H, et al. 2000. Half-life of blood carboxyhemoglobin after short-term and long-term exposure to carbon monoxide. *J Trauma* 49(1):126-131.
- \*Shusterman D, Alexeeff G, Hargis C, et al. 1996. Predictors of carbon monoxide and hydrogen cyanide exposure in smoke inhalation patients. *J Toxicol Clin Toxicol* 34(1):61-71.
- +\*Silkoff PE, Zhang L, Dutton S, et al. 2005. Winter air pollution and disease parameters in advanced chronic obstructive pulmonary disease patients residing in Denver, Colorado. *J Allergy Clin Immunol* 115:337-344.
- \*Silverman RK, Montano J. 1997. Hyperbaric oxygen treatment during pregnancy in acute carbon monoxide poisoning. A case report. *J Reprod Med* 42:309-311.
- +\*Singh J. 1986. Early behavioral alterations in mice following prenatal carbon monoxide exposure. *Neurotoxicology* 7:475-481.
- Singh J. 2006. Interaction of maternal protein and carbon monoxide on pup mortality in mice: Implications for global infant mortality. *Birth Defects Res B Dev Reprod Toxicol* 77:216-226.
- Singh J, Kaur H. 2008. Interaction of protein, zinc and carbon monoxide on daily food intake during gestation and fetal weight in mice. *Birth Defects Res A Clin Mol Teratol* 82(5):361.
- Singh J, Moore-Cheatum L. 1993. Gestational protein deficiency enhances fetotoxicity of carbon monoxide. *Ann N Y Acad Sci* 678:366-368.
- Singh J, Aggison L, Moore-Cheatum L. 1993. Teratogenicity and developmental toxicity of carbon monoxide in protein-deficient mice. *Teratology* 48(2):149-159.

## 9. REFERENCES

- Singh J, Smith CB, Moore-Cheatum L. 1992. Additivity of protein deficiency and carbon monoxide on placental carboxyhemoglobin in mice. *Am J Obstet Gynecol* 167(3):843-846.
- Singh MP, Sharan M, Selvakumar S. 1991. A mathematical model for the computation of carboxyhaemoglobin in human blood as a function of exposure time. *Philos Trans R Soc Lond B Biol Sci* 334:135-147.
- +Singh V, Khandelwal R, Gupta AB. 2003. Effect of air pollution on peak expiratory flow rate variability. *J Asthma* 40(1):81-86.
- \*Sinha AK, Klein J, Schultze P, et al. 1991. Cerebral regional capillary perfusion and blood flow after carbon monoxide exposure. *J Appl Physiol* 71(4):1196-1200.
- Sitdikova GF, Islamov RR, Mukhamedyarov MA, et al. 2007. Modulation of neurotransmitter release by carbon monoxide at the frog neuro-muscular junction. *Curr Drug Metab* 8:177-184.
- +\*Slaughter JC, Kim E, Sheppard L, et al. 2005. Association between particulate matter and emergency room visits, hospital admissions and mortality in Spokane, Washington. *J Expo Anal Environ Epidemiol* 15(2):153-159.
- +\*Slaughter JC, Lumley T, Sheppard L, et al. 2003. Effects of ambient air pollution on symptom severity and medication use in children with asthma. *Ann Allergy Asthma Immunol* 91(4):346-353.
- \*Smith MV. 1990. Comparing solutions to the linear and nonlinear CFK equations for predicting COHb formation. *Math Biosci* 99:251-263.
- Smith NJ. 1977. End-expired air technic for determining occupational carbon monoxide exposure. *J Occup Med* 19(11):766-769.
- Smith MD, Merigan WH, McIntire RW. 1976. Effects of carbon monoxide on fixed-consecutive-number performance in rats. *Pharmacol Biochem Behav* 5(3):257-262.
- \*Smith MV, Hazucha MJ, Benignus VA, et al. 1994. Effect of regional circulation patterns on observed HbCO levels. *J Appl Physiol* 77(4):1659-1665.
- \*Snella MC, Rylander R. 1979. Alteration in local and systemic immune capacity after exposure to bursts of CO. *Environ Res* 20(1):74-79.
- \*Snyder SH, Jaffrey SR, Zakhary R. 1998. Nitric oxide and carbon monoxide: Parallel roles as neural messengers. *Brain Res Rev* 26:167-175.
- \*Soares MP, Seldon MP, Gregoire IP, et al. 2004. Heme oxygenase-1 modulates the expression of adhesion molecules associated with endothelial cell activation. *J Immunol* 172:3553-3563.
- Sokal JA, Majka J, Palus J. 1984. The content of carbon monoxide in the tissues of rats intoxicated with carbon monoxide in various conditions of acute exposure. *Arch Toxicol* 56(2):106-108.
- \*Solanki DL, McCurdy PR, Cuttitta FF, et al. 1988. Hemolysis in sickle cell disease as measured by endogenous carbon monoxide production. A preliminary report. *Am J Clin Pathol* 89(2):221-225.

## 9. REFERENCES

- Song R, Mahidhara RS, Liu F, et al. 2002. Carbon monoxide inhibits human airway smooth muscle cell proliferation via mitogen-activated protein kinase pathway. *Am J Respir Cell Mol Biol* 27(5):603-610.
- Song SY, Okeda R, Funata N, et al. 1983. An experimental study of the pathogenesis of the selective lesion of the globus pallidus in acute carbon monoxide poisoning in cats. With special reference to the chronologic change in the cerebral local blood flow. *Acta Neuropathol (Berl)* 61(3-4):232-238.
- \*Sorhaug S, Steinshamn S, Nilsen OG, et al. 2006. Chronic inhalation of carbon monoxide: Effects on the respiratory and cardiovascular system at doses corresponding to tobacco smoking. *Toxicology* 228(2-3):280-290.
- \*SRI. 2008. 2008 Directory of chemical producers. Menlo Park, CA: SRI Consulting, 482.
- +Steerenberg PA, Nierkens S, Fischer PH, et al. 2001. Traffic-related air pollution affects peak expiratory flow, exhaled nitric oxide, and inflammatory nasal markers. *Arch Environ Health* 56(2):167-174.
- +\*Steinvil A, Kordova-Biezuner L, Shapira I, et al. 2008. Short-term exposure to air pollution and inflammation-sensitive biomarkers. *Environ Res* 106(1):51-61.
- \*Stevens JL, Ratnayake JH, Anders MW. 1980. Metabolism of dihalomethanes to carbon monoxide. *Toxicol Appl Pharmacol* 55:484-489.
- Stevenson DK, Vreman HJ. 1997. Carbon monoxide and bilirubin production in neonates. *Pediatrics* 100(2 Pt 1):252-254.
- Stevenson DK, Ostrander CE, Johnson JD. 1979. Effect of erythrocyte destruction on the pulmonary excretion rate of carbon monoxide in adult male Wistar rats. *J Lab Clin Med* 94(4):649-654.
- \*Stewart RD, Peterson JE, Baretta ED, et al. 1970. Experimental human exposure to carbon monoxide. *Arch Environ Health* 21(2):154-164.
- \*Stewart RD, Peterson JE, Fisher TN, et al. 1973. Experimental human exposure to high concentrations of carbon monoxide. *Arch Environ Health* 26(1):1-7.
- \*Stieb DM, Judek S, Burnett RT. 2002. Meta-analysis of time-series studies of air pollution and mortality: Effects of gases and particles and the influence of cause of death, age, and season. *J Air Waste Manag Assoc* 52:470-484.
- +Stieb DM, Judek S, Burnett RT. 2003. Meta-analysis of time-series studies of air pollution and mortality: Update in relation to the use of generalized additive models. *J Air Waste Manag Assoc* 53(3):258-261.
- +\*Stieb DM, Evans GJ, Sabaliauskas K, et al. 2007. A scripted activity study of the impact of protective advice on personal exposure to ultra-fine and fine particulate matter and volatile organic compounds. *J Expo Sci Environ Epidemiol* 18:495-502.
- +\*Stockard-Sullivan JE, Korsak RA, Webber DS, et al. 2003. Mild carbon monoxide exposure and auditory function in the developing rat. *J Neurosci Res* 74(5):644-654.

## 9. REFERENCES

- +Stolzel M, Breitner S, Cyrus J, et al. 2006. Daily mortality and particulate matter in different size classes in Erfurt, Germany. *J Expo Sci Environ Epidemiol* 17:458-467.
- \*Stone JR, Marletta MA. 1994. Soluble guanylate cyclase from bovine lung: Activation with nitric oxide and carbon monoxide and spectral characterization of the ferrous and ferric states. *Biochemistry* 33(18):5636-5640.
- +\*Storm JE, Fechter LD. 1985a. Alteration in the postnatal ontogeny of cerebellar norepinephrine content following chronic prenatal carbon monoxide. *J Neurochem* 45(3):965-969.
- +\*Storm JE, Fechter LD. 1985b. Prenatal carbon monoxide exposure differentially affects postnatal weight and monoamine concentration of rat brain regions. *Toxicol Appl Pharmacol* 81(1):139-146.
- \*Stupfel M, Bouley G. 1970. Physiological and biochemical effects on rats and mice exposed to small concentrations of carbon monoxide for long periods. *Ann N Y Acad Sci* 174(1):342-368.
- Sugi K, Theissen JL, Traber LD, et al. 1990. Impact of carbon monoxide on cardiopulmonary dysfunction after smoke inhalation injury. *Circ Res* 66(1):69-75.
- Suliman HB, Carraway MS, Tatro LG, et al. 2007. A new activating role for CO in cardiac mitochondrial biogenesis. *J Cell Sci* 120(Pt 2):299-308.
- +\*Sullivan J, Ishikawa N, Sheppard L, et al. 2003. Exposure to ambient fine particulate matter and primary cardiac arrest among persons with and without clinically recognized heart disease. *Am J Epidemiol* 157(6):501-509.
- +Sun HL, Chou MC, Lue KH. 2006. The relationship of air pollution to ED visits for asthma differ between children and adults. *Am J Emerg Med* 24:709-713.
- Sun Y, Jin K, Mao XO, et al. 2001. Neuroglobin is up-regulated by and protects neurons from hypoxic-ischemic injury. *Proc Natl Acad Sci USA* 98(26):15306-15311.
- Suner S, Jay G. 2008. Carbon monoxide has direct toxicity on the myocardium distinct from effects of hypoxia in an ex vivo rat heart model. *Acad Emerg Med* 15(1):59-65.
- +Sunyer J, Basagana X. 2001. Particles, and not gases, are associated with the risk of death in patients with chronic obstructive pulmonary disease. *Int J Epidemiol* 30(5):1138-1140.
- +Sunyer J, Basagana X, Belmonte J, et al. 2002. Effect of nitrogen dioxide and ozone on the risk of dying in patients with severe asthma. *Thorax* 57(8):687-693.
- +\*Symons JM, Wang L, Guallar E, et al. 2006. A case-crossover study of fine particulate matter air pollution and onset of congestive heart failure symptom exacerbation leading to hospitalization. *Am J Epidemiol* 164(5):421-433.
- +\*Szyszkowicz M. 2007. Air pollution and emergency department visits for ischemic heart disease in Montreal, Canada. *Int J Occup Med Environ Health* 20(2):167-174.
- \*Taille C, El-Benna J, Lanone S, et al. 2004. Induction of heme oxygenase-1 inhibits NAD(P)H oxidase activity by down-regulating cytochrome b558 expression via the reduction of heme availability. *J Biol Chem* 279:28681-28688.

## 9. REFERENCES

- \*Takeuchi A, Vesely A, Rucker J, et al. 2000. A simple "new" method to accelerate clearance of carbon monoxide. *Am J Respir Crit Care Med* 161:1816-1819.
- +\*Tarkiainen TH, Timonen KL, Vanninen EJ, et al. 2003. Effect of acute carbon monoxide exposure on heart rate variability in patients with coronary artery disease. *Clin Physiol Funct Imaging* 23(2):98-102.
- +\*Tattoli M, Carratu MR, Cassano T, et al. 1999. Effects of early postnatal exposure to low concentrations of carbon monoxide on cognitive functions in rats. *Pharmacol Res* 40(3):271-274.
- Telfer JF, Itoh H, Thomson AJ, et al. 2001. Activity and expression of soluble and particulate guanylate cyclases in myometrium from nonpregnant and pregnant women: Down-regulation of soluble guanylate cyclase at term. *J Clin Endocrinol Metab* 86(12):5934-5943.
- +\*Tenías JM, Ballester F, Pérez-Hoyos S, et al. 2002. Air pollution and hospital emergency room admissions for chronic obstructive pulmonary disease in Valencia, Spain. *Arch Environ Health* 57(1):41-47.
- ten Tusscher GW, de Weerd J, Roos CM, et al. 2001. Decreased lung function associated with perinatal exposure to Dutch background levels of dioxins. *Acta Paediatr* 90(11):1292-1298.
- Teran FJ, Johnson RA, Stevenson BK, et al. 2005. Heme oxygenase-derived carbon monoxide promotes arteriolar endothelial dysfunction and contributes to salt-induced hypertension in Dahl salt-sensitive rats. *Am J Physiol Regul Integr Comp Physiol* 288(3):R615-R622.
- Thom SR. 1990. Carbon monoxide-mediated brain lipid peroxidation in the rat. *J Appl Physiol* 68(3):997-1003.
- Thom SR. 1992. Dehydrogenase conversion to oxidase and lipid peroxidation in brain after carbon monoxide poisoning. *J Appl Physiol* 73(4):1584-1589.
- Thom SR. 1993. Leukocytes in carbon monoxide-mediated brain oxidative injury. *Toxicol Appl Pharmacol* 123(2):234-247.
- \*Thom SR, Bhopale VM, Han ST, et al. 2006. Intravascular neutrophil activation due to carbon monoxide poisoning. *Am J Respir Crit Care Med* 174(11):1239-1248.
- Thom SR, Fisher D, Manevich Y. 2001. Roles for platelet-activating factor and \*NO-derived oxidants causing neutrophil adherence after CO poisoning. *Am J Physiol Heart Circ Physiol* 281(2):H923-H930.
- Thom SR, Fisher D, Xu YA, et al. 1999. Role of nitric oxide-derived oxidants in vascular injury from carbon monoxide in the rat. *Am J Physiol* 276(3 Pt 2):H984-H992.
- \*Thom SR, Fisher D, Xu YA, et al. 2000. Adaptive responses and apoptosis in endothelial cells exposed to carbon monoxide. *Proc Natl Acad Sci USA* 97(3):1305-1310.
- \*Thom SR, Ohnishi ST, Ischiropoulos H. 1994. Nitric oxide released by platelets inhibits neutrophil B2 integrin function following acute carbon monoxide poisoning. *Toxicol Appl Pharmacol* 128(1):105-110.
- \*Thom SR, Xu YA, Ischiropoulos H. 1997. Vascular endothelial cells generate peroxynitrite in response to carbon monoxide exposure. *Chem Res Toxicol* 10(9):1023-1031.

## 9. REFERENCES

- \*Thomas K, Colborn T. 1992. Organochlorine endocrine disruptors in human tissue. In: Colborn T, Clement C, eds. Chemically induced alterations in sexual and functional development: The wildlife/human connection. Princeton, NJ: Princeton Scientific Publishing, 365-394.
- +\*Thompson AJ, Shields MD, Patterson CC. 2001. Acute asthma exacerbations and air pollutants in children living in Belfast, Northern Ireland. *Arch Environ Health* 56:234-241.
- \*Thomsen HK. 1974. Carbon monoxide-induced atherosclerosis in primates. An electron-microscopic study on the coronary arteries of *Macaca irus* monkeys. *Atherosclerosis* 20(2):233-240.
- \*Thorup C, Jones CL, Gross SS, et al. 1999. Carbon monoxide induces vasodilation and nitric oxide release but suppresses endothelial NOS. *Am J Physiol* 277(6 Pt 2):F882-F889.
- \*Tikuisis P, Buick F, Kane DM. 1987a. Percent carboxyhemoglobin in resting humans exposed repeatedly to 1,500 and 7,500 ppm CO. *J Appl Physiol* 63(2):820-827.
- \*Tikuisis P, Kane DM, McLellan TM, et al. 1992. Rate of formation of carboxyhemoglobin in exercising humans exposed to carbon monoxide. *J Appl Physiol* 72(4):1311-1319.
- \*Tikuisis P, Madill HD, Gill BJ, et al. 1987b. A critical analysis of the use of the CFK equation in predicting COHb formation. *Am Ind Hyg Assoc J* 48(3):208-213.
- +\*Timonen KL, Pekkanen J, Tiittanen P, et al. 2002. Effects of air pollution on changes in lung function induced by exercise in children with chronic respiratory symptoms. *Occup Environ Med* 59(2):129-134.
- +\*Timonen KL, Vanninen E, de Hartog J, et al. 2006. Effects of ultrafine and fine particulate and gaseous air pollution on cardiac autonomic control in subjects with coronary artery disease: The ULTRA study. *J Expo Sci Environ Epidemiol* 16(4):332-341.
- +\*Tolbert PE, Klein M, Peel JL, et al. 2007. Multipollutant modeling issues in a study of ambient air quality and emergency department visits in Atlanta. *J Expo Sci Environ Epidemiol* 17(Suppl 2):S29-S35.
- +Tolcos M, Mallard C, McGregor H, et al. 2000a. Exposure to prenatal carbon monoxide and postnatal hyperthermia: Short and long-term effects on neurochemicals and neuroglia in the developing brain. *Exp Neurol* 162:235-246.
- +\*Tolcos M, McGregor H, Walker D, et al. 2000b. Chronic prenatal exposure to carbon monoxide results in a reduction in tyrosine hydroxylase-immunoreactivity and an increase in choline acetyltransferase-immunoreactivity in the fetal medulla: Implications for Sudden Infant Death Syndrome. *J Neuropathol Exp Neurol* 59(3):218-228.
- \*Tolli JD, Sievert SM, Taylor CD. 2006. Unexpected diversity of bacteria capable of carbon monoxide oxidation in a coastal marine environment, and contribution of the roseobacter-associated clade to total CO oxidation. *Appl Environ Microbiol* 72(3):1966-1973.
- \*Tomaszewski C. 2006. Carbon monoxide. In: Goldfrank's toxicologic emergencies. 8th ed. New York, NY: McGraw-Hill, 1689-1704.
- \*Torne R, Soyer HP, Leb G, et al. 1991. Skin lesions in carbon monoxide intoxication. *Dermatologica* 183:212-215.

## 9. REFERENCES

- +Trapasso LM, Keith L. 1999. Relationships between selected meteorological/pollution parameters and hospital admissions for asthma in South Central Kentucky. *J Ky Acad Sci* 60:73-77.
- +\*Tsai S, Chen C, Hsieh H, et al. 2006a. Air pollution and postneonatal mortality in a tropical city: Kaohsiung, Taiwan. *Inhal Toxicol* 18:185-189.
- +Tsai SS, Cheng MH, Chiu HF, et al. 2006b. Air pollution and hospital admissions for asthma in a tropical city: Kaohsiung, Taiwan. *Inhal Toxicol* 18(8):549-554.
- +\*Tsai SS, Goggins WB, Chiu HF, et al. 2003b. Evidence for an association between air pollution and daily stroke admissions in Kaohsiung, Taiwan. *Stroke* 34(11):2612-2616.
- +\*Tsai SS, Huang CH, Goggins, WB, et al. 2003a. Relationship between air pollution and daily mortality in a tropical city: Kaohsiung, Taiwan. *J Toxicol Environ Sci Health A* 66:1341-1349.
- Tucciarone M, Dileo PA, Castro ER, et al. 2009. Myocardial infarction secondary to carbon monoxide poisoning: An uncommon presentation of a common condition. Case report and review of the literature. *Am J Ther* [Apr 4; Epub ahead of print]
- Tyuma I, Ueda Y, Imaizumi K, et al. 1981. Prediction of the carbonmonoxyhemoglobin levels during and after carbon monoxide exposures in various animal species. *Jpn J Physiol* 31(2):131-143.
- Utz J, Ullrich V. 1991. Carbon monoxide relaxes ileal smooth muscle through activation of guanylate cyclase. *Biochem Pharmacol* 41(8):1195-1201.
- Vallone B, Nienhaus K, Matthes A, et al. 2004. The structure of carbonmonoxy neuroglobin reveals a heme-sliding mechanism for control of ligand affinity. *Proc Natl Acad Sci USA* 101(50):17351-17356.
- +Vedal S, Brauer M, White R, et al. 2003. Air pollution and daily mortality in a city with low levels of pollution. *Environ Health Perspect* 111(1):45-51.
- +\*Vedal S, Rich K, Brauer M, et al. 2004. Air pollution and cardiac arrhythmias in patients with implantable cardioverter defibrillators. *Inhal Toxicol* 16:353-362.
- Verma A, Hirsch DJ, Glatt CE, et al. 1993. Carbon monoxide: A putative neural messenger. *Science* 259:381-384.
- \*Verschueren K. 2001. Handbook of environmental data on organic chemicals. 4th ed. New York, NY: John Wiley & Sons, Inc., 23-24.
- \*Vesely AE, Somogyi RB, Sasano H, et al. 2004. The effects of carbon monoxide on respiratory chemoreflexes in humans. *Environ Res* 94:227-233.
- \*Vieira I, Sonnier M, Cresteil T. 1996. Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 238(2):476-483.
- +Vigotti MA, Chiaverini F, Bagiola P, et al. 2007. Urban air pollution and emergency visits for respiratory complaints in Pisa, Italy. *J Toxicol Environ Health A* 70:266-269.

## 9. REFERENCES

- Villamor E, Perez-Vizcaino F, Cogolludo AL, et al. 2000. Relaxant effects of carbon monoxide compared with nitric oxide in pulmonary and systemic vessels of newborn piglets. *Pediatr Res* 48(4):546-553.
- +Villeneuve PJ, Burnett RT, Shi Y, et al. 2003. A time-series study of air pollution, socioeconomic status, and mortality in Vancouver, Canada. *J Expo Anal Environ Epidemiol* 13(6):427-435.
- +\*Villeneuve PJ, Chen L, Stieb D, et al. 2006a. Associations between outdoor air pollution and emergency department visits for stroke in Edmonton, Canada. *Eur J Epidemiol* 21:689-700.
- +Villeneuve PJ, Doiron M, Stieb D, et al. 2006b. Is outdoor air pollution associated with physician visits for allergy rhinitis among the elderly in Toronto, Canada? *Allergy* 61:750-758.
- +\*Vogel JA, Gleser MA. 1972. Effect of carbon monoxide on oxygen transport during exercise. *J Appl Physiol* 32:234-239.
- Vogel JA, Gleser MA, Wheeler RC, et al. 1972. Carbon monoxide and physical work capacity. *Arch Environ Health* 24(3):198-203.
- \*Volti GL, Vanella L, Gazzolo D, et al. 2008. Carbon monoxide: Vasoconstrictor or vasodilator? That's the question. *American Journal of physiology. Renal Physiology* 295:901-903.
- +\*von Klot S, Peters A, Aalto P, et al. 2005. Ambient air pollution is associated with increased risk of hospital cardiac readmissions of myocardial infarction survivors in five European cities. *Circ J* 112:3073-3079.
- +\*von Klot S, Wolke G, Tuch T, et al. 2002. Increased asthma medication use in association with ambient fine and ultrafine particles. *Eur Respir J* 20:691-702.
- Vreman HJ, Stevenson DK. 1994. Carboxyhemoglobin determined in neonatal blood with a CO-oximeter unaffected by fetal oxyhemoglobin. *Clin Chem* 40(8):1522-1527.
- Vreman HJ, Baxter LM, Stone RT, et al. 1996. Evaluation of a fully automated end-tidal carbon monoxide instrument for breath analysis. *Clin Chem* 42(1):50-56.
- \*Vreman HJ, Kwong LK, Stevenson DK. 1984. Carbon monoxide in blood: An improved microliter blood-sample collection system, with rapid analysis by gas chromatography. *Clin Chem* 30(8):1382-1386.
- Vreman HJ, Mahoney JJ, Stevenson DK. 1993. Electrochemical measurement of carbon monoxide in breath: Interference by hydrogen. *Atmos Environ* 27A(14):2193-2198.
- Vreman HJ, Mahoney JJ, Stevenson DK. 1995. Carbon monoxide and carboxyhemoglobin. *Adv Pediatr* 42:303-334.
- \*Vreman HJ, Stevenson DK, Oh W, et al. 1994. Semiportable electrochemical instrument for determining carbon monoxide in breath. *Clin Chem* 40(10):1927-1933.
- \*Vreman HJ, Wong RJ, Kadotani T, et al. 2005. Determination of carbon monoxide (CO) in rodent tissue: Effect of heme administration and environmental CO exposure. *Anal Biochem* 341:280-289.

## 9. REFERENCES

- Vreman HJ, Wong RJ, Kim EC, et al. 2000. Haem oxygenase activity in human umbilical cord and rat vascular tissues. *Placenta* 21:337-344.
- \*Vreman HJ, Wong RJ, Stevenson DK, et al. 2006. Concentration of carbon monoxide (CO) in postmortem human tissues: Effect of environmental CO exposure. *J Forensic Sci* 51(5):1182-1190.
- Wagner JA, Horvath SM, Dahms TE. 1975. Carbon monoxide elimination. *Respir Physiol* 23(1):41-47.
- Wallace LA. 1983. Carbon monoxide in air and breath of employees in an underground office. *J Air Pollut Control Assoc* 33(7):678-682.
- Wallace L, Thomas J, Mage D, et al. 1988. Comparison of breath CO, CO exposure, and Coburn model predictions in the U. S. EPA Washington-Denver (CO) study. *Atmos Environ* 22(10):2183-2193.
- Wang R. 1998. Resurgence of carbon monoxide: An endogenous gaseous vasorelaxing factor. *Can J Physiol Pharmacol* 76:1-15.
- \*Wang R, Wu L. 1997. The chemical modification of K<sub>Ca</sub> channels by carbon monoxide in vascular smooth muscle cells. *J Biol Chem* 272:8222-8226.
- \*Wang R, Wang Z, Wu L. 1997a. Carbon monoxide-induced vasorelaxation and the underlying mechanisms. *Br J Pharmacol* 121(5):927-934.
- \*Wang R, Wu L, Wang Z. 1997b. The direct effect of carbon monoxide on K<sub>Ca</sub> channels in vascular smooth muscle cells. *Pflugers Arch* 434:285-291.
- +Wang T, Ko Y, Chao Y, et al. 1999. Association between indoor and outdoor air pollution and adolescent asthma from 1995 to 1996 in Taiwan. *Environ Res* 81:239-247.
- \*Weaver LK, Howe S, Hopkins R, et al. 2000. Carboxyhemoglobin half-life in carbon monoxide-poisoned patients treated with 100% oxygen at atmospheric pressure. *Chest* 117(3):801-808.
- Weaver LK, Valentine KJ, Hopkins RO. 2007. Carbon monoxide poisoning. *Am J Respir Crit Care Med* 176:491-497.
- +\*Webber DS, Korsak RA, Sininger LK, et al. 2003. Mild carbon monoxide exposure impairs the developing auditory system of the rat. *J Neurosci Res* 74:655-665.
- Webber DS, Lopez I, Korsak RA, et al. 2005. Limiting iron availability confers neuroprotection from chronic mild carbon monoxide exposure in the developing auditory system of the rat. *J Neurosci Res* 80:620-633.
- Weigel PH. 1974. Inhibition of *Escherichia-coli* DNA replication by cyanide and carbon monoxide [Abstract]. *Fed Proc* 33(5 Part 2):1352.
- +\*Wellenius GA, Bateson TF, Mittleman MA, et al. 2005a. Particulate air pollution and the rate of hospitalization for congestive heart failure among medicare beneficiaries in Pittsburgh, Pennsylvania. *Am J Epidemiol* 161(11):1030-1036.
- +\*Wellenius GA, Schwartz J, Mittleman MA. 2005b. Air pollution and hospital admissions for ischemic and hemorrhagic stroke among medicare beneficiaries. *Stroke* 36:2549-2553.

## 9. REFERENCES

- Wellenius GA, Schwartz J, Mittleman MA. 2006. Particulate air pollution and hospital admissions for congestive heart failure in seven United States cities. *Am J Cardiol* 97(33):404-408.
- \*Werner B, Lindahl J. 1980. Endogenous carbon monoxide production after bicycle exercise in healthy subjects and in patients with hereditary spherocytosis. *Scand J Clin Lab Invest* 40:314-324.
- \*West JR, Smith HW, Chasis H. 1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. *J Pediatr* 32:10-18.
- \*Wetherell HR. 1966. The occurrence of cyanide in the blood of fire victims. *J. Forensic Sci* 11:167-173.
- +Wheeler A, Zanobetti A, Gold DR, et al. 2006. The relationship between ambient air pollution and heart rate variability differs for individuals with heart and pulmonary disease. *Environ Health Perspect* 114(4):560-566.
- WHO. 1998. Effects of carbon monoxide on the micronucleus test. In: *Occupational health in the chemical industry: Papers presented at the 22. ICOH Congress, Sydney, Australia, 27 Sept-2 Oct 1987*. Copenhagen: World Health Organization, 156-166.
- \*WHO. 1999. Environmental health criteria 213. Carbon monoxide. World Health Organization. <http://www.inchem.org/documents/ehc/ehc/ehc213.htm>. June 5, 2009.
- \*WHO. 2000. Air quality guidelines. 2nd edition. Geneva, Switzerland: World Health Organization. [http://www.euro.who.int/air/activities/20050223\\_4](http://www.euro.who.int/air/activities/20050223_4). May 11, 2009.
- \*WHO. 2006. Guidelines for drinking-water quality. 3rd edition. Geneva, Switzerland: World Health Organization. [http://www.who.int/water\\_sanitation\\_health/dwq/gdwq3/en/](http://www.who.int/water_sanitation_health/dwq/gdwq3/en/). May 11, 2009.
- \*Widdowson EM, Dickerson JWT. 1964. Chemical composition of the body. In: Comar CL, Bronner F, eds. *Mineral metabolism: An advance treatise. Volume II: The elements Part A*. New York, NY: Academic Press, 1-247.
- \*Wijayanti N, Huber S, Samoylenko A, et al. 2004. Role of NF- $\kappa$ B and p38 MAP kinase signaling pathways in the lipopolysaccharide-dependent activation of heme oxygenase-1 gene expression. *Antioxid Redox Signal* 6:802-810.
- +\*Wilhelm M, Ritz B. 2005. Local variations in CO and particulate air pollution and adverse birth outcomes in Los Angeles County, California, USA. *Environ Health Perspect* 113(9):1212-1221.
- \*Willis D, Tomlinson A, Frederick R, et al. 1995. Modulation of heme oxygenase activity in rat brain and spleen by inhibitors and donors of nitric oxide. *Biochem Biophys Res Commun* 214:1152-1156.
- Wittenberg BA, Wittenberg JB, Caldwell PRB. 1975. Role of myoglobin in the oxygen supply to red skeletal muscle. *J Biol Chem* 250(23):9038-9043.
- \*Wohlrab H, Ogunmola BG. 1971. Carbon monoxide binding studies of cytochrome  $\alpha_3$  hemes in intact rat liver mitochondria. *Biochemistry* 10:1103-1106.

## 9. REFERENCES

- \*Wolf SJ, Lavonas EJ, Sloan EP, et al. 2008. Clinical policy: Critical issues in the management of adult patients presenting to the emergency department with acute carbon monoxide poisoning. *Ann Emerg Med* 51(2):138-152.
- \*Wolff E. 1994. Carbon monoxide poisoning with severe myonecrosis and acute renal failure. *Am J Emerg Med* 12(3):347-349.
- +\*Woodruff TJ, Darrow LA, Parker JD. 2008. Air pollution and postneonatal infant mortality in the United States, 1999-2002. *Environ Health Perspect* 116(1):110-115.
- Woody RC, Brewster MA. 1990. Telencephalic dysgenesis associated with presumptive maternal carbon monoxide intoxication in the first trimester of pregnancy. *J Toxicol Clin Toxicol* 28(4):467-475.
- \*Wu L, Wang R. 2005. Carbon monoxide: Endogenous production, physiological functions, and pharmacological applications. *Pharmacol Rev* 57(4):585-630.
- +Xirasagar S, Lin H, Liu T. 2006. Seasonality in pediatric asthma admissions: The role of climate and environmental factors. *Eur J Pediatr* 165:747-752.
- +\*Yang C, Chang C, Chuang H, et al. 2004a. Relationship between air pollution and daily mortality in a subtropical city: Taipei, Taiwan. *Environ Int* 30:519-523.
- +Yang CY, Chen CC, Chen CY, et al. 2007. Air pollution and hospital admissions for asthma in a subtropical city: Taipei, Taiwan. *J Toxicol Environ Health A* 70(2):111-117.
- \*+Yang CY, Chen YS, Yang CH, et al. 2004b. Relationship between ambient air pollution and hospital admissions for cardiovascular diseases in Kaohsiung, Taiwan. *J Toxicol Environ Health A* 67(6):483-493.
- +\*Yang CY, Hsieh HJ, Tsai SS, et al. 2006. Correlation between air pollution and postneonatal mortality in a subtropical city: Taipei, Taiwan. *J Toxicol Environ Health Part A* 69:2033-2040.
- +\*Yang Q, Chen Y, Krewski D, et al. 2005. Effect of short-term exposure to low levels of gaseous pollutants on chronic obstructive pulmonary disease hospitalizations. *Environ Res* 99:99-105.
- +\*Yang Q, Chen Y, Shi Y, et al. 2003. Association between ozone and respiratory admissions among children and the elderly in Vancouver, Canada. *Inhal Toxicol* 15:1297-1308.
- Yang W, Jennison BL, Omaye ST. 1998. Cardiovascular disease hospitalization and ambient levels of carbon monoxide. *J Toxicol Environ Health A* 55(3):185-196.
- \*Yaws CL, Hopper JR, Wang X, et al. 1999. Calculating solubility and Henry's law constants for gases in water. *Chem Eng (NY)* 106(6):102-105.
- Yoshiki N, Kubota T, Aso T. 2001. Identification of heme oxygenase in human endometrium. *J Clin Endocrinol Metab* 86(10):5033-5038.
- \*Young LJ, Caughey WS. 1986. Oxygenation of carbon monoxide by bovine heart cytochrome c oxidase. *Biochemistry* 25(1):152-161.

## 9. REFERENCES

- \*Young JS, Upchurch MB, Kaufman MJ, et al. 1987. Carbon monoxide exposure potentiates high-frequency auditory threshold shifts induced by noise. *Hear Res* 26:37-43.
- +\*Yu O, Sheppard L, Lumley T, et al. 2000. Effects of ambient air pollution on symptoms of asthma in Seattle-area children enrolled in the CAMP study. *Environ Health Perspect* 108(12):1209-1214.
- +\*Zanobetti A, Schwartz J. 2006. Air pollution and emergency admissions in Boston, MA. *J Epidemiol Community Health* 60(10):890-895.
- \*Zanobetti A, Canner MJ, Stone PH, et al. 2004b. Ambient pollution and blood pressure in cardiac rehabilitation patients. *Circulation* 110:2184-2189.
- +\*Zanobetti A, Wellenius GA, Zeka A, et al. 2004a. The effect of particulate air pollution on the risk of myocardial infarction in elderly: A multi-city case-crossover analysis. *Epidemiology* 15(4):S21.
- \*Zebro T, Wright EA, Littleton RJ, et al. 1983. Bone changes in mice after prolonged continuous exposure to a high concentration of carbon monoxide. *Exp Pathol* 24:51-67.
- Zenclussen ML, Anegon I, Bertoja AZ, et al. 2006. Over-expression of heme oxygenase-1 by adenoviral gene transfer improves pregnancy outcome in a murine model of abortion. *J Reprod Immunol* 69:35-52.
- +\*Zevin S, Saunders S, Gourlay SG, et al. 2001. Cardiovascular effects of carbon monoxide and cigarette smoking. *J Am Coll Cardiol* 38(6):1633-1638.
- Zhang C, Li X, Zhao L, et al. 2007. Lipopolysaccharide (LPS) up-regulates the expression of haem oxygenase-1 in mouse placenta. *Placenta* 28:951-957.
- Zhang X, Shan P, Alam J, et al. 2005. Carbon monoxide differentially modulates STAT1 and STAT3 and inhibits apoptosis via a phosphatidylinositol 3-kinase/Akt and p38 kinase-dependent STAT3 pathway during anoxia-reoxygenation injury. *J Biol Chem* 280(10):8714-8721.
- Zhao H, Wong RJ, Doyle TC, et al. 2008. Regulation of maternal and fetal hemodynamics by heme oxygenase in mice. *Biol Reprod* 78(4):744-751.
- Zhuo M, Small SA, Kandel ER, et al. 1993. Nitric oxide and carbon monoxide produce activity-dependent long-term synaptic enhancement in hippocampus. *Science* 260:1946-1950.
- \*Ziegler EE, Edwards BB, Jensen RL, et al. 1978. Absorption and retention of lead by infants. *Pediatr Res* 12(1):29-34.
- \*Zuckerbraun BS, Billiar TR, Otterbein SL, et al. 2003. Carbon monoxide protects against liver failure through nitric oxide-induced heme oxygenase 1. *J Exp Med* 198:1707-1716.
- \*Zuckerbraun BS, Chin BY, Bilban M, et al. 2007. Carbon monoxide signals via inhibition of cytochrome c oxidase and generation of mitochondrial reactive oxygen species. *FASEB J* 21(4):1099-1106.

## 10. GLOSSARY

**Absorption**—The taking up of liquids by solids, or of gases by solids or liquids.

**Acute Exposure**—Exposure to a chemical for a duration of 14 days or less, as specified in the Toxicological Profiles.

**Adsorption**—The adhesion in an extremely thin layer of molecules (as of gases, solutes, or liquids) to the surfaces of solid bodies or liquids with which they are in contact.

**Adsorption Coefficient ( $K_{oc}$ )**—The ratio of the amount of a chemical adsorbed per unit weight of organic carbon in the soil or sediment to the concentration of the chemical in solution at equilibrium.

**Adsorption Ratio ( $K_d$ )**—The amount of a chemical adsorbed by sediment or soil (i.e., the solid phase) divided by the amount of chemical in the solution phase, which is in equilibrium with the solid phase, at a fixed solid/solution ratio. It is generally expressed in micrograms of chemical sorbed per gram of soil or sediment.

**Benchmark Dose (BMD)**—Usually defined as the lower confidence limit on the dose that produces a specified magnitude of changes in a specified adverse response. For example, a  $BMD_{10}$  would be the dose at the 95% lower confidence limit on a 10% response, and the benchmark response (BMR) would be 10%. The BMD is determined by modeling the dose response curve in the region of the dose response relationship where biologically observable data are feasible.

**Benchmark Dose Model**—A statistical dose-response model applied to either experimental toxicological or epidemiological data to calculate a BMD.

**Bioconcentration Factor (BCF)**—The quotient of the concentration of a chemical in aquatic organisms at a specific time or during a discrete time period of exposure divided by the concentration in the surrounding water at the same time or during the same period.

**Biomarkers**—Broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility.

**Cancer Effect Level (CEL)**—The lowest dose of chemical in a study, or group of studies, that produces significant increases in the incidence of cancer (or tumors) between the exposed population and its appropriate control.

**Carcinogen**—A chemical capable of inducing cancer.

**Case-Control Study**—A type of epidemiological study that examines the relationship between a particular outcome (disease or condition) and a variety of potential causative agents (such as toxic chemicals). In a case-controlled study, a group of people with a specified and well-defined outcome is identified and compared to a similar group of people without outcome.

**Case Report**—Describes a single individual with a particular disease or exposure. These may suggest some potential topics for scientific research, but are not actual research studies.

**Case Series**—Describes the experience of a small number of individuals with the same disease or exposure. These may suggest potential topics for scientific research, but are not actual research studies.

## 10. GLOSSARY

**Ceiling Value**—A concentration of a substance that should not be exceeded, even instantaneously.

**Chronic Exposure**—Exposure to a chemical for 365 days or more, as specified in the Toxicological Profiles.

**Cohort Study**—A type of epidemiological study of a specific group or groups of people who have had a common insult (e.g., exposure to an agent suspected of causing disease or a common disease) and are followed forward from exposure to outcome. At least one exposed group is compared to one unexposed group.

**Cross-sectional Study**—A type of epidemiological study of a group or groups of people that examines the relationship between exposure and outcome to a chemical or to chemicals at one point in time.

**Data Needs**—Substance-specific informational needs that if met would reduce the uncertainties of human health assessment.

**Developmental Toxicity**—The occurrence of adverse effects on the developing organism that may result from exposure to a chemical prior to conception (either parent), during prenatal development, or postnatally to the time of sexual maturation. Adverse developmental effects may be detected at any point in the life span of the organism.

**Dose-Response Relationship**—The quantitative relationship between the amount of exposure to a toxicant and the incidence of the adverse effects.

**Embryotoxicity and Fetotoxicity**—Any toxic effect on the conceptus as a result of prenatal exposure to a chemical; the distinguishing feature between the two terms is the stage of development during which the insult occurs. The terms, as used here, include malformations and variations, altered growth, and *in utero* death.

**Environmental Protection Agency (EPA) Health Advisory**—An estimate of acceptable drinking water levels for a chemical substance based on health effects information. A health advisory is not a legally enforceable federal standard, but serves as technical guidance to assist federal, state, and local officials.

**Epidemiology**—Refers to the investigation of factors that determine the frequency and distribution of disease or other health-related conditions within a defined human population during a specified period.

**Genotoxicity**—A specific adverse effect on the genome of living cells that, upon the duplication of affected cells, can be expressed as a mutagenic, clastogenic, or carcinogenic event because of specific alteration of the molecular structure of the genome.

**Half-life**—A measure of rate for the time required to eliminate one half of a quantity of a chemical from the body or environmental media.

**Immediately Dangerous to Life or Health (IDLH)**—The maximum environmental concentration of a contaminant from which one could escape within 30 minutes without any escape-impairing symptoms or irreversible health effects.

**Immunologic Toxicity**—The occurrence of adverse effects on the immune system that may result from exposure to environmental agents such as chemicals.

## 10. GLOSSARY

**Immunological Effects**—Functional changes in the immune response.

**Incidence**—The ratio of individuals in a population who develop a specified condition to the total number of individuals in that population who could have developed that condition in a specified time period.

**Intermediate Exposure**—Exposure to a chemical for a duration of 15–364 days, as specified in the Toxicological Profiles.

**In Vitro**—Isolated from the living organism and artificially maintained, as in a test tube.

**In Vivo**—Occurring within the living organism.

**Lethal Concentration<sub>(LO)</sub> (LC<sub>LO</sub>)**—The lowest concentration of a chemical in air that has been reported to have caused death in humans or animals.

**Lethal Concentration<sub>(50)</sub> (LC<sub>50</sub>)**—A calculated concentration of a chemical in air to which exposure for a specific length of time is expected to cause death in 50% of a defined experimental animal population.

**Lethal Dose<sub>(LO)</sub> (LD<sub>LO</sub>)**—The lowest dose of a chemical introduced by a route other than inhalation that has been reported to have caused death in humans or animals.

**Lethal Dose<sub>(50)</sub> (LD<sub>50</sub>)**—The dose of a chemical that has been calculated to cause death in 50% of a defined experimental animal population.

**Lethal Time<sub>(50)</sub> (LT<sub>50</sub>)**—A calculated period of time within which a specific concentration of a chemical is expected to cause death in 50% of a defined experimental animal population.

**Lowest-Observed-Adverse-Effect Level (LOAEL)**—The lowest exposure level of chemical in a study, or group of studies, that produces statistically or biologically significant increases in frequency or severity of adverse effects between the exposed population and its appropriate control.

**Lymphoreticular Effects**—Represent morphological effects involving lymphatic tissues such as the lymph nodes, spleen, and thymus.

**Malformations**—Permanent structural changes that may adversely affect survival, development, or function.

**Minimal Risk Level (MRL)**—An estimate of daily human exposure to a hazardous substance that is likely to be without an appreciable risk of adverse noncancer health effects over a specified route and duration of exposure.

**Modifying Factor (MF)**—A value (greater than zero) that is applied to the derivation of a Minimal Risk Level (MRL) to reflect additional concerns about the database that are not covered by the uncertainty factors. The default value for a MF is 1.

**Morbidity**—State of being diseased; morbidity rate is the incidence or prevalence of disease in a specific population.

**Mortality**—Death; mortality rate is a measure of the number of deaths in a population during a specified interval of time.

## 10. GLOSSARY

**Mutagen**—A substance that causes mutations. A mutation is a change in the DNA sequence of a cell's DNA. Mutations can lead to birth defects, miscarriages, or cancer.

**Necropsy**—The gross examination of the organs and tissues of a dead body to determine the cause of death or pathological conditions.

**Neurotoxicity**—The occurrence of adverse effects on the nervous system following exposure to a chemical.

**No-Observed-Adverse-Effect Level (NOAEL)**—The dose of a chemical at which there were no statistically or biologically significant increases in frequency or severity of adverse effects seen between the exposed population and its appropriate control. Effects may be produced at this dose, but they are not considered to be adverse.

**Octanol-Water Partition Coefficient ( $K_{ow}$ )**—The equilibrium ratio of the concentrations of a chemical in *n*-octanol and water, in dilute solution.

**Odds Ratio (OR)**—A means of measuring the association between an exposure (such as toxic substances and a disease or condition) that represents the best estimate of relative risk (risk as a ratio of the incidence among subjects exposed to a particular risk factor divided by the incidence among subjects who were not exposed to the risk factor). An OR of greater than 1 is considered to indicate greater risk of disease in the exposed group compared to the unexposed group.

**Organophosphate or Organophosphorus Compound**—A phosphorus-containing organic compound and especially a pesticide that acts by inhibiting cholinesterase.

**Permissible Exposure Limit (PEL)**—An Occupational Safety and Health Administration (OSHA) allowable exposure level in workplace air averaged over an 8-hour shift of a 40-hour workweek.

**Pesticide**—General classification of chemicals specifically developed and produced for use in the control of agricultural and public health pests.

**Pharmacokinetics**—The dynamic behavior of a material in the body, used to predict the fate (disposition) of an exogenous substance in an organism. Utilizing computational techniques, it provides the means of studying the absorption, distribution, metabolism, and excretion of chemicals by the body.

**Pharmacokinetic Model**—A set of equations that can be used to describe the time course of a parent chemical or metabolite in an animal system. There are two types of pharmacokinetic models: data-based and physiologically-based. A data-based model divides the animal system into a series of compartments, which, in general, do not represent real, identifiable anatomic regions of the body, whereas the physiologically-based model compartments represent real anatomic regions of the body.

**Physiologically Based Pharmacodynamic (PBPD) Model**—A type of physiologically based dose-response model that quantitatively describes the relationship between target tissue dose and toxic end points. These models advance the importance of physiologically based models in that they clearly describe the biological effect (response) produced by the system following exposure to an exogenous substance.

**Physiologically Based Pharmacokinetic (PBPK) Model**—Comprised of a series of compartments representing organs or tissue groups with realistic weights and blood flows. These models require a

## 10. GLOSSARY

variety of physiological information: tissue volumes, blood flow rates to tissues, cardiac output, alveolar ventilation rates, and possibly membrane permeabilities. The models also utilize biochemical information, such as air/blood partition coefficients, and metabolic parameters. PBPK models are also called biologically based tissue dosimetry models.

**Prevalence**—The number of cases of a disease or condition in a population at one point in time.

**Prospective Study**—A type of cohort study in which the pertinent observations are made on events occurring after the start of the study. A group is followed over time.

**$q_1^*$** —The upper-bound estimate of the low-dose slope of the dose-response curve as determined by the multistage procedure. The  $q_1^*$  can be used to calculate an estimate of carcinogenic potency, the incremental excess cancer risk per unit of exposure (usually  $\mu\text{g/L}$  for water,  $\text{mg/kg/day}$  for food, and  $\mu\text{g/m}^3$  for air).

**Recommended Exposure Limit (REL)**—A National Institute for Occupational Safety and Health (NIOSH) time-weighted average (TWA) concentration for up to a 10-hour workday during a 40-hour workweek.

**Reference Concentration (RfC)**—An estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious noncancer health effects during a lifetime. The inhalation reference concentration is for continuous inhalation exposures and is appropriately expressed in units of  $\text{mg/m}^3$  or ppm.

**Reference Dose (RfD)**—An estimate (with uncertainty spanning perhaps an order of magnitude) of the daily exposure of the human population to a potential hazard that is likely to be without risk of deleterious effects during a lifetime. The RfD is operationally derived from the no-observed-adverse-effect level (NOAEL, from animal and human studies) by a consistent application of uncertainty factors that reflect various types of data used to estimate RfDs and an additional modifying factor, which is based on a professional judgment of the entire database on the chemical. The RfDs are not applicable to nonthreshold effects such as cancer.

**Reportable Quantity (RQ)**—The quantity of a hazardous substance that is considered reportable under the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA). Reportable quantities are (1) 1 pound or greater or (2) for selected substances, an amount established by regulation either under CERCLA or under Section 311 of the Clean Water Act. Quantities are measured over a 24-hour period.

**Reproductive Toxicity**—The occurrence of adverse effects on the reproductive system that may result from exposure to a chemical. The toxicity may be directed to the reproductive organs and/or the related endocrine system. The manifestation of such toxicity may be noted as alterations in sexual behavior, fertility, pregnancy outcomes, or modifications in other functions that are dependent on the integrity of this system.

**Retrospective Study**—A type of cohort study based on a group of persons known to have been exposed at some time in the past. Data are collected from routinely recorded events, up to the time the study is undertaken. Retrospective studies are limited to causal factors that can be ascertained from existing records and/or examining survivors of the cohort.

**Risk**—The possibility or chance that some adverse effect will result from a given exposure to a chemical.

## 10. GLOSSARY

**Risk Factor**—An aspect of personal behavior or lifestyle, an environmental exposure, or an inborn or inherited characteristic that is associated with an increased occurrence of disease or other health-related event or condition.

**Risk Ratio**—The ratio of the risk among persons with specific risk factors compared to the risk among persons without risk factors. A risk ratio greater than 1 indicates greater risk of disease in the exposed group compared to the unexposed group.

**Short-Term Exposure Limit (STEL)**—The American Conference of Governmental Industrial Hygienists (ACGIH) maximum concentration to which workers can be exposed for up to 15 minutes continually. No more than four excursions are allowed per day, and there must be at least 60 minutes between exposure periods. The daily Threshold Limit Value-Time Weighted Average (TLV-TWA) may not be exceeded.

**Standardized Mortality Ratio (SMR)**—A ratio of the observed number of deaths and the expected number of deaths in a specific standard population.

**Target Organ Toxicity**—This term covers a broad range of adverse effects on target organs or physiological systems (e.g., renal, cardiovascular) extending from those arising through a single limited exposure to those assumed over a lifetime of exposure to a chemical.

**Teratogen**—A chemical that causes structural defects that affect the development of an organism.

**Threshold Limit Value (TLV)**—An American Conference of Governmental Industrial Hygienists (ACGIH) concentration of a substance to which most workers can be exposed without adverse effect. The TLV may be expressed as a Time Weighted Average (TWA), as a Short-Term Exposure Limit (STEL), or as a ceiling limit (CL).

**Time-Weighted Average (TWA)**—An allowable exposure concentration averaged over a normal 8-hour workday or 40-hour workweek.

**Toxic Dose<sub>(50)</sub> (TD<sub>50</sub>)**—A calculated dose of a chemical, introduced by a route other than inhalation, which is expected to cause a specific toxic effect in 50% of a defined experimental animal population.

**Toxicokinetic**—The absorption, distribution, and elimination of toxic compounds in the living organism.

**Uncertainty Factor (UF)**—A factor used in operationally deriving the Minimal Risk Level (MRL) or Reference Dose (RfD) or Reference Concentration (RfC) from experimental data. UFs are intended to account for (1) the variation in sensitivity among the members of the human population, (2) the uncertainty in extrapolating animal data to the case of human, (3) the uncertainty in extrapolating from data obtained in a study that is of less than lifetime exposure, and (4) the uncertainty in using lowest-observed-adverse-effect level (LOAEL) data rather than no-observed-adverse-effect level (NOAEL) data. A default for each individual UF is 10; if complete certainty in data exists, a value of 1 can be used; however, a reduced UF of 3 may be used on a case-by-case basis, 3 being the approximate logarithmic average of 10 and 1.

**Xenobiotic**—Any chemical that is foreign to the biological system.

## APPENDIX A. ATSDR MINIMAL RISK LEVELS AND WORKSHEETS

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) [42 U.S.C. 9601 et seq.], as amended by the Superfund Amendments and Reauthorization Act (SARA) [Pub. L. 99–499], requires that the Agency for Toxic Substances and Disease Registry (ATSDR) develop jointly with the U.S. Environmental Protection Agency (EPA), in order of priority, a list of hazardous substances most commonly found at facilities on the CERCLA National Priorities List (NPL); prepare toxicological profiles for each substance included on the priority list of hazardous substances; and assure the initiation of a research program to fill identified data needs associated with the substances.

The toxicological profiles include an examination, summary, and interpretation of available toxicological information and epidemiologic evaluations of a hazardous substance. During the development of toxicological profiles, Minimal Risk Levels (MRLs) are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration for a given route of exposure. An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse noncancer health effects over a specified duration of exposure. MRLs are based on noncancer health effects only and are not based on a consideration of cancer effects. These substance-specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors to identify contaminants and potential health effects that may be of concern at hazardous waste sites. It is important to note that MRLs are not intended to define clean-up or action levels.

MRLs are derived for hazardous substances using the no-observed-adverse-effect level/uncertainty factor approach. They are below levels that might cause adverse health effects in the people most sensitive to such chemical-induced effects. MRLs are derived for acute (1–14 days), intermediate (15–364 days), and chronic (365 days and longer) durations and for the oral and inhalation routes of exposure. Currently, MRLs for the dermal route of exposure are not derived because ATSDR has not yet identified a method suitable for this route of exposure. MRLs are generally based on the most sensitive chemical-induced end point considered to be of relevance to humans. Serious health effects (such as irreparable damage to the liver or kidneys, or birth defects) are not used as a basis for establishing MRLs. Exposure to a level above the MRL does not mean that adverse health effects will occur.

MRLs are intended only to serve as a screening tool to help public health professionals decide where to look more closely. They may also be viewed as a mechanism to identify those hazardous waste sites that

## APPENDIX A

are not expected to cause adverse health effects. Most MRLs contain a degree of uncertainty because of the lack of precise toxicological information on the people who might be most sensitive (e.g., infants, elderly, nutritionally or immunologically compromised) to the effects of hazardous substances. ATSDR uses a conservative (i.e., protective) approach to address this uncertainty consistent with the public health principle of prevention. Although human data are preferred, MRLs often must be based on animal studies because relevant human studies are lacking. In the absence of evidence to the contrary, ATSDR assumes that humans are more sensitive to the effects of hazardous substance than animals and that certain persons may be particularly sensitive. Thus, the resulting MRL may be as much as 100-fold below levels that have been shown to be nontoxic in laboratory animals.

Proposed MRLs undergo a rigorous review process: Health Effects/MRL Workgroup reviews within the Division of Toxicology and Environmental Medicine, expert panel peer reviews, and agency-wide MRL Workgroup reviews, with participation from other federal agencies and comments from the public. They are subject to change as new information becomes available concomitant with updating the toxicological profiles. Thus, MRLs in the most recent toxicological profiles supersede previously published levels. For additional information regarding MRLs, please contact the Division of Toxicology and Environmental Medicine, Agency for Toxic Substances and Disease Registry, 1600 Clifton Road NE, Mailstop F-62, Atlanta, Georgia 30333.

MRLs were not derived for carbon monoxide, as discussed in Section 2.3.

## APPENDIX B. USER'S GUIDE

### Chapter 1

#### Public Health Statement

This chapter of the profile is a health effects summary written in non-technical language. Its intended audience is the general public, especially people living in the vicinity of a hazardous waste site or chemical release. If the Public Health Statement were removed from the rest of the document, it would still communicate to the lay public essential information about the chemical.

The major headings in the Public Health Statement are useful to find specific topics of concern. The topics are written in a question and answer format. The answer to each question includes a sentence that will direct the reader to chapters in the profile that will provide more information on the given topic.

### Chapter 2

#### Relevance to Public Health

This chapter provides a health effects summary based on evaluations of existing toxicologic, epidemiologic, and toxicokinetic information. This summary is designed to present interpretive, weight-of-evidence discussions for human health end points by addressing the following questions:

1. What effects are known to occur in humans?
2. What effects observed in animals are likely to be of concern to humans?
3. What exposure conditions are likely to be of concern to humans, especially around hazardous waste sites?

The chapter covers end points in the same order that they appear within the Discussion of Health Effects by Route of Exposure section, by route (inhalation, oral, and dermal) and within route by effect. Human data are presented first, then animal data. Both are organized by duration (acute, intermediate, chronic). *In vitro* data and data from parenteral routes (intramuscular, intravenous, subcutaneous, etc.) are also considered in this chapter.

The carcinogenic potential of the profiled substance is qualitatively evaluated, when appropriate, using existing toxicokinetic, genotoxic, and carcinogenic data. ATSDR does not currently assess cancer potency or perform cancer risk assessments. Minimal Risk Levels (MRLs) for noncancer end points (if derived) and the end points from which they were derived are indicated and discussed.

Limitations to existing scientific literature that prevent a satisfactory evaluation of the relevance to public health are identified in the Chapter 3 Data Needs section.

#### Interpretation of Minimal Risk Levels

Where sufficient toxicologic information is available, ATSDR has derived MRLs for inhalation and oral routes of entry at each duration of exposure (acute, intermediate, and chronic). These MRLs are not meant to support regulatory action, but to acquaint health professionals with exposure levels at which adverse health effects are not expected to occur in humans.

## APPENDIX B

MRLs should help physicians and public health officials determine the safety of a community living near a chemical emission, given the concentration of a contaminant in air or the estimated daily dose in water. MRLs are based largely on toxicological studies in animals and on reports of human occupational exposure.

MRL users should be familiar with the toxicologic information on which the number is based. Chapter 2, "Relevance to Public Health," contains basic information known about the substance. Other sections such as Chapter 3 Section 3.9, "Interactions with Other Substances," and Section 3.10, "Populations that are Unusually Susceptible" provide important supplemental information.

MRL users should also understand the MRL derivation methodology. MRLs are derived using a modified version of the risk assessment methodology that the Environmental Protection Agency (EPA) provides (Barnes and Dourson 1988) to determine reference doses (RfDs) for lifetime exposure.

To derive an MRL, ATSDR generally selects the most sensitive end point which, in its best judgement, represents the most sensitive human health effect for a given exposure route and duration. ATSDR cannot make this judgement or derive an MRL unless information (quantitative or qualitative) is available for all potential systemic, neurological, and developmental effects. If this information and reliable quantitative data on the chosen end point are available, ATSDR derives an MRL using the most sensitive species (when information from multiple species is available) with the highest no-observed-adverse-effect level (NOAEL) that does not exceed any adverse effect levels. When a NOAEL is not available, a lowest-observed-adverse-effect level (LOAEL) can be used to derive an MRL, and an uncertainty factor (UF) of 10 must be employed. Additional uncertainty factors of 10 must be used both for human variability to protect sensitive subpopulations (people who are most susceptible to the health effects caused by the substance) and for interspecies variability (extrapolation from animals to humans). In deriving an MRL, these individual uncertainty factors are multiplied together. The product is then divided into the inhalation concentration or oral dosage selected from the study. Uncertainty factors used in developing a substance-specific MRL are provided in the footnotes of the levels of significant exposure (LSE) tables.

## **Chapter 3**

### **Health Effects**

#### **Tables and Figures for Levels of Significant Exposure (LSE)**

Tables and figures are used to summarize health effects and illustrate graphically levels of exposure associated with those effects. These levels cover health effects observed at increasing dose concentrations and durations, differences in response by species, MRLs to humans for noncancer end points, and EPA's estimated range associated with an upper-bound individual lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. Use the LSE tables and figures for a quick review of the health effects and to locate data for a specific exposure scenario. The LSE tables and figures should always be used in conjunction with the text. All entries in these tables and figures represent studies that provide reliable, quantitative estimates of NOAELs, LOAELs, or Cancer Effect Levels (CELs).

The legends presented below demonstrate the application of these tables and figures. Representative examples of LSE Table 3-1 and Figure 3-1 are shown. The numbers in the left column of the legends correspond to the numbers in the example table and figure.

## APPENDIX B

**LEGEND****See Sample LSE Table 3-1 (page B-6)**

- (1) Route of Exposure. One of the first considerations when reviewing the toxicity of a substance using these tables and figures should be the relevant and appropriate route of exposure. Typically when sufficient data exist, three LSE tables and two LSE figures are presented in the document. The three LSE tables present data on the three principal routes of exposure, i.e., inhalation, oral, and dermal (LSE Tables 3-1, 3-2, and 3-3, respectively). LSE figures are limited to the inhalation (LSE Figure 3-1) and oral (LSE Figure 3-2) routes. Not all substances will have data on each route of exposure and will not, therefore, have all five of the tables and figures.
- (2) Exposure Period. Three exposure periods—acute (less than 15 days), intermediate (15–364 days), and chronic (365 days or more)—are presented within each relevant route of exposure. In this example, an inhalation study of intermediate exposure duration is reported. For quick reference to health effects occurring from a known length of exposure, locate the applicable exposure period within the LSE table and figure.
- (3) Health Effect. The major categories of health effects included in LSE tables and figures are death, systemic, immunological, neurological, developmental, reproductive, and cancer. NOAELs and LOAELs can be reported in the tables and figures for all effects but cancer. Systemic effects are further defined in the "System" column of the LSE table (see key number 18).
- (4) Key to Figure. Each key number in the LSE table links study information to one or more data points using the same key number in the corresponding LSE figure. In this example, the study represented by key number 18 has been used to derive a NOAEL and a Less Serious LOAEL (also see the two "18r" data points in sample Figure 3-1).
- (5) Species. The test species, whether animal or human, are identified in this column. Chapter 2, "Relevance to Public Health," covers the relevance of animal data to human toxicity and Section 3.4, "Toxicokinetics," contains any available information on comparative toxicokinetics. Although NOAELs and LOAELs are species specific, the levels are extrapolated to equivalent human doses to derive an MRL.
- (6) Exposure Frequency/Duration. The duration of the study and the weekly and daily exposure regimens are provided in this column. This permits comparison of NOAELs and LOAELs from different studies. In this case (key number 18), rats were exposed to "Chemical x" via inhalation for 6 hours/day, 5 days/week, for 13 weeks. For a more complete review of the dosing regimen, refer to the appropriate sections of the text or the original reference paper (i.e., Nitschke et al. 1981).
- (7) System. This column further defines the systemic effects. These systems include respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, renal, and dermal/ocular. "Other" refers to any systemic effect (e.g., a decrease in body weight) not covered in these systems. In the example of key number 18, one systemic effect (respiratory) was investigated.
- (8) NOAEL. A NOAEL is the highest exposure level at which no harmful effects were seen in the organ system studied. Key number 18 reports a NOAEL of 3 ppm for the respiratory system, which was used to derive an intermediate exposure, inhalation MRL of 0.005 ppm (see footnote "b").

## APPENDIX B

- (9) LOAEL. A LOAEL is the lowest dose used in the study that caused a harmful health effect. LOAELs have been classified into "Less Serious" and "Serious" effects. These distinctions help readers identify the levels of exposure at which adverse health effects first appear and the gradation of effects with increasing dose. A brief description of the specific end point used to quantify the adverse effect accompanies the LOAEL. The respiratory effect reported in key number 18 (hyperplasia) is a Less Serious LOAEL of 10 ppm. MRLs are not derived from Serious LOAELs.
- (10) Reference. The complete reference citation is given in Chapter 9 of the profile.
- (11) CEL. A CEL is the lowest exposure level associated with the onset of carcinogenesis in experimental or epidemiologic studies. CELs are always considered serious effects. The LSE tables and figures do not contain NOAELs for cancer, but the text may report doses not causing measurable cancer increases.
- (12) Footnotes. Explanations of abbreviations or reference notes for data in the LSE tables are found in the footnotes. Footnote "b" indicates that the NOAEL of 3 ppm in key number 18 was used to derive an MRL of 0.005 ppm.

**LEGEND****See Sample Figure 3-1 (page B-7)**

LSE figures graphically illustrate the data presented in the corresponding LSE tables. Figures help the reader quickly compare health effects according to exposure concentrations for particular exposure periods.

- (13) Exposure Period. The same exposure periods appear as in the LSE table. In this example, health effects observed within the acute and intermediate exposure periods are illustrated.
- (14) Health Effect. These are the categories of health effects for which reliable quantitative data exists. The same health effects appear in the LSE table.
- (15) Levels of Exposure. Concentrations or doses for each health effect in the LSE tables are graphically displayed in the LSE figures. Exposure concentration or dose is measured on the log scale "y" axis. Inhalation exposure is reported in mg/m<sup>3</sup> or ppm and oral exposure is reported in mg/kg/day.
- (16) NOAEL. In this example, the open circle designated 18r identifies a NOAEL critical end point in the rat upon which an intermediate inhalation exposure MRL is based. The key number 18 corresponds to the entry in the LSE table. The dashed descending arrow indicates the extrapolation from the exposure level of 3 ppm (see entry 18 in the table) to the MRL of 0.005 ppm (see footnote "b" in the LSE table).
- (17) CEL. Key number 38m is one of three studies for which CELs were derived. The diamond symbol refers to a CEL for the test species-mouse. The number 38 corresponds to the entry in the LSE table.

## APPENDIX B

- (18) Estimated Upper-Bound Human Cancer Risk Levels. This is the range associated with the upper-bound for lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. These risk levels are derived from the EPA's Human Health Assessment Group's upper-bound estimates of the slope of the cancer dose response curve at low dose levels ( $q_1^*$ ).
- (19) Key to LSE Figure. The Key explains the abbreviations and symbols used in the figure.

## SAMPLE

1 →

Table 3-1. Levels of Significant Exposure to [Chemical x] – Inhalation


Key to figure <sup>a</sup>	Species	Exposure frequency/ duration	System	NOAEL (ppm)	LOAEL (effect)		Reference
					Less serious (ppm)	Serious (ppm)	
<b>INTERMEDIATE EXPOSURE</b>							
	5	6	7	8	9		10
3 →	Systemic	↓	↓	↓	↓	↓	↓
4 →	18	Rat	13 wk 5 d/wk 6 hr/d	Resp	3 <sup>b</sup>	10 (hyperplasia)	Nitschke et al. 1981
<b>CHRONIC EXPOSURE</b>							
	Cancer					11	
					↓		
	38	Rat	18 mo 5 d/wk 7 hr/d			20	(CEL, multiple organs) Wong et al. 1982
	39	Rat	89–104 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, nasal tumors) NTP 1982
	40	Mouse	79–103 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, hemangiosarcomas) NTP 1982

12 →

<sup>a</sup> The number corresponds to entries in Figure 3-1.<sup>b</sup> Used to derive an intermediate inhalation Minimal Risk Level (MRL) of  $5 \times 10^{-3}$  ppm; dose adjusted for intermittent exposure and divided by an uncertainty factor of 100 (10 for extrapolation from animal to humans, 10 for human variability).

**SAMPLE**

**Figure 3-1. Levels of Significant Exposure to [Chemical X] - Inhalation**



\*\*\*DRAFT FOR PUBLIC COMMENT\*\*\*

APPENDIX B

This page is intentionally blank.

## APPENDIX C. ACRONYMS, ABBREVIATIONS, AND SYMBOLS

ACGIH	American Conference of Governmental Industrial Hygienists
ACOEM	American College of Occupational and Environmental Medicine
ADI	acceptable daily intake
ADME	absorption, distribution, metabolism, and excretion
AED	atomic emission detection
AFID	alkali flame ionization detector
AFOSH	Air Force Office of Safety and Health
ALT	alanine aminotransferase
AML	acute myeloid leukemia
AOAC	Association of Official Analytical Chemists
AOEC	Association of Occupational and Environmental Clinics
AP	alkaline phosphatase
APHA	American Public Health Association
AST	aspartate aminotransferase
atm	atmosphere
ATSDR	Agency for Toxic Substances and Disease Registry
AWQC	Ambient Water Quality Criteria
BAT	best available technology
BCF	bioconcentration factor
BEI	Biological Exposure Index
BMD/C	benchmark dose or benchmark concentration
BMD <sub>x</sub>	dose that produces a X% change in response rate of an adverse effect
BMDL <sub>x</sub>	95% lower confidence limit on the BMD <sub>x</sub>
BMDS	Benchmark Dose Software
BMR	benchmark response
BSC	Board of Scientific Counselors
C	centigrade
CAA	Clean Air Act
CAG	Cancer Assessment Group of the U.S. Environmental Protection Agency
CAS	Chemical Abstract Services
CDC	Centers for Disease Control and Prevention
CEL	cancer effect level
CELDS	Computer-Environmental Legislative Data System
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CFR	Code of Federal Regulations
Ci	curie
CI	confidence interval
CL	ceiling limit value
CLP	Contract Laboratory Program
cm	centimeter
CML	chronic myeloid leukemia
CPSC	Consumer Products Safety Commission
CWA	Clean Water Act
DHEW	Department of Health, Education, and Welfare
DHHS	Department of Health and Human Services
DNA	deoxyribonucleic acid
DOD	Department of Defense
DOE	Department of Energy
DOL	Department of Labor

## APPENDIX C

DOT	Department of Transportation
DOT/UN/ NA/IMDG	Department of Transportation/United Nations/ North America/Intergovernmental Maritime Dangerous Goods Code
DWEL	drinking water exposure level
ECD	electron capture detection
ECG/EKG	electrocardiogram
EEG	electroencephalogram
EEGL	Emergency Exposure Guidance Level
EPA	Environmental Protection Agency
F	Fahrenheit
F <sub>1</sub>	first-filial generation
FAO	Food and Agricultural Organization of the United Nations
FDA	Food and Drug Administration
FEMA	Federal Emergency Management Agency
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FPD	flame photometric detection
fpm	feet per minute
FR	Federal Register
FSH	follicle stimulating hormone
g	gram
GC	gas chromatography
gd	gestational day
GLC	gas liquid chromatography
GPC	gel permeation chromatography
HPLC	high-performance liquid chromatography
HRGC	high resolution gas chromatography
HSDB	Hazardous Substance Data Bank
IARC	International Agency for Research on Cancer
IDLH	immediately dangerous to life and health
ILO	International Labor Organization
IRIS	Integrated Risk Information System
K <sub>d</sub>	adsorption ratio
kg	kilogram
kkg	metric ton
K <sub>oc</sub>	organic carbon partition coefficient
K <sub>ow</sub>	octanol-water partition coefficient
L	liter
LC	liquid chromatography
LC <sub>50</sub>	lethal concentration, 50% kill
LC <sub>Lo</sub>	lethal concentration, low
LD <sub>50</sub>	lethal dose, 50% kill
LD <sub>Lo</sub>	lethal dose, low
LDH	lactic dehydrogenase
LH	lutinizing hormone
LOAEL	lowest-observed-adverse-effect level
LSE	Levels of Significant Exposure
LT <sub>50</sub>	lethal time, 50% kill
m	meter
MA	<i>trans,trans</i> -muconic acid
MAL	maximum allowable level
mCi	millicurie

## APPENDIX C

MCL	maximum contaminant level
MCLG	maximum contaminant level goal
MF	modifying factor
MFO	mixed function oxidase
mg	milligram
mL	milliliter
mm	millimeter
mmHg	millimeters of mercury
mmol	millimole
mppcf	millions of particles per cubic foot
MRL	Minimal Risk Level
MS	mass spectrometry
NAAQS	National Ambient Air Quality Standard
NAS	National Academy of Science
NATICH	National Air Toxics Information Clearinghouse
NATO	North Atlantic Treaty Organization
NCE	normochromatic erythrocytes
NCEH	National Center for Environmental Health
NCI	National Cancer Institute
ND	not detected
NFPA	National Fire Protection Association
ng	nanogram
NHANES	National Health and Nutrition Examination Survey
NIEHS	National Institute of Environmental Health Sciences
NIOSH	National Institute for Occupational Safety and Health
NIOSHTIC	NIOSH's Computerized Information Retrieval System
NLM	National Library of Medicine
nm	nanometer
nmol	nanomole
NOAEL	no-observed-adverse-effect level
NOES	National Occupational Exposure Survey
NOHS	National Occupational Hazard Survey
NPD	nitrogen phosphorus detection
NPDES	National Pollutant Discharge Elimination System
NPL	National Priorities List
NR	not reported
NRC	National Research Council
NS	not specified
NSPS	New Source Performance Standards
NTIS	National Technical Information Service
NTP	National Toxicology Program
ODW	Office of Drinking Water, EPA
OERR	Office of Emergency and Remedial Response, EPA
OHM/TADS	Oil and Hazardous Materials/Technical Assistance Data System
OPP	Office of Pesticide Programs, EPA
OPPT	Office of Pollution Prevention and Toxics, EPA
OPPTS	Office of Prevention, Pesticides and Toxic Substances, EPA
OR	odds ratio
OSHA	Occupational Safety and Health Administration
OSW	Office of Solid Waste, EPA
OTS	Office of Toxic Substances

## APPENDIX C

OW	Office of Water
OWRS	Office of Water Regulations and Standards, EPA
PAH	polycyclic aromatic hydrocarbon
PBPD	physiologically based pharmacodynamic
PBPK	physiologically based pharmacokinetic
PCE	polychromatic erythrocytes
PEL	permissible exposure limit
pg	picogram
PHS	Public Health Service
PID	photo ionization detector
pmol	picomole
PMR	proportionate mortality ratio
ppb	parts per billion
ppm	parts per million
ppt	parts per trillion
PSNS	pretreatment standards for new sources
RBC	red blood cell
REL	recommended exposure level/limit
RfC	reference concentration
RfD	reference dose
RNA	ribonucleic acid
RQ	reportable quantity
RTECS	Registry of Toxic Effects of Chemical Substances
SARA	Superfund Amendments and Reauthorization Act
SCE	sister chromatid exchange
SGOT	serum glutamic oxaloacetic transaminase
SGPT	serum glutamic pyruvic transaminase
SIC	standard industrial classification
SIM	selected ion monitoring
SMCL	secondary maximum contaminant level
SMR	standardized mortality ratio
SNARL	suggested no adverse response level
SPEGL	Short-Term Public Emergency Guidance Level
STEL	short term exposure limit
STORET	Storage and Retrieval
TD <sub>50</sub>	toxic dose, 50% specific toxic effect
TLV	threshold limit value
TOC	total organic carbon
TPQ	threshold planning quantity
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TWA	time-weighted average
UF	uncertainty factor
U.S.	United States
USDA	United States Department of Agriculture
USGS	United States Geological Survey
VOC	volatile organic compound
WBC	white blood cell
WHO	World Health Organization

## APPENDIX C

>	greater than
≥	greater than or equal to
=	equal to
<	less than
≤	less than or equal to
%	percent
α	alpha
β	beta
γ	gamma
δ	delta
μm	micrometer
μg	microgram
q <sub>1</sub> *	cancer slope factor
-	negative
+	positive
(+)	weakly positive result
(-)	weakly negative result

APPENDIX C

This page is intentionally blank.

## APPENDIX D. INDEX

absorbed dose.....	176
adipose tissue.....	10
adrenal gland.....	172
adsorption.....	199
aerobic.....	59, 88, 209
ambient air.....	10, 12, 13, 14, 16, 21, 26, 28, 29, 30, 31, 32, 33, 34, 35, 36, 44, 48, 65, 73, 80, 84, 85, 87, 89, 95, 96, 97, 107, 108, 109, 110, 111, 112, 113, 127, 140, 143, 168, 174, 180, 185, 187, 210, 230, 234
anaerobic.....	209
anemia.....	137, 180
biomarker.....	11, 12, 16, 17, 84, 85, 86, 87, 89, 175, 176, 185, 186, 188, 189, 233, 240
blood cell count.....	17
breast milk.....	226
cancer.....	32, 127, 173, 186, 189, 244
carboxyhemoglobin (see COHb).....	7, 149, 152, 153, 155, 161, 235
carcinogenic.....	187
carcinogenicity.....	128, 185, 187, 243
cardiac arrhythmia.....	12, 13, 19, 27, 30, 48, 49, 50, 59, 65, 66, 72, 82, 83, 177, 180
cardiovascular.....	8, 11, 12, 13, 14, 20, 27, 29, 30, 32, 33, 48, 49, 50, 59, 65, 66, 81, 82, 90, 92, 165, 171, 177, 179, 180, 185, 186, 189
cardiovascular effects.....	13, 20, 48, 49, 65, 82, 83, 180, 185, 186, 189
CFK model.....	11, 14, 15, 16, 20, 26, 30, 31, 66, 128, 145, 147, 148, 151, 162
clearance.....	182
cognitive function.....	188
COHb (see carboxyhemoglobin).....	7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 20, 21, 25, 26, 27, 30, 34, 47, 48, 49, 50, 59, 65, 66, 73, 79, 82, 83, 84, 87, 88, 90, 91, 92, 93, 94, 95, 113, 125, 127, 128, 130, 133, 134, 136, 137, 138, 140, 143, 145, 147, 148, 149, 150, 151, 152, 153, 154, 155, 158, 161, 162, 163, 165, 171, 174, 176, 177, 180, 182, 185, 186, 188, 189, 190, 191, 224, 227, 228, 229, 231, 233, 234, 235, 240, 241
COMb.....	10, 134, 138, 142, 158, 162, 190
death.....	12, 14, 19, 30, 32, 95, 96, 112, 113, 174, 187, 246
deoxyribonucleic acid (see DNA).....	128
dermal effects.....	88
developmental effects.....	14, 20, 83, 89, 90, 96, 113, 125, 174, 186, 187, 189, 191
DNA (see deoxyribonucleic acid).....	128, 129, 176, 187
dopamine.....	126
elimination half-time.....	10, 143, 154, 162, 176, 189
elimination rate.....	143, 176, 182
endocrine.....	88, 171, 172
endocrine effects.....	88
endogenous production.....	9, 10, 20, 27, 28, 133, 137, 142, 154, 166, 190
fetal tissue.....	10, 136, 154, 155, 174
fetus.....	12, 27, 136, 137, 154, 173, 174, 176, 180, 182, 189, 227
gastrointestinal effects.....	83
general population.....	8, 204, 224, 227, 239, 246
genotoxic.....	128

## APPENDIX D

genotoxicity.....	185, 187
growth retardation.....	109
Haldane coefficient.....	136, 148, 149, 150, 151, 152, 153, 156, 160, 190
Haldane constant.....	133, 134, 138, 139, 142, 147
half-life.....	143, 175, 208, 230
hematological effects.....	14, 16, 84, 96, 125, 127, 174, 188
hepatic effects.....	88
Hepatic Effects.....	88
hydroxyl radical.....	203, 208, 230
hypoxic mechanisms.....	9, 11, 17, 20, 21, 27, 28, 165, 166, 168, 191
immune system.....	14, 17, 20, 89, 127, 188
immunological.....	14, 89, 90, 96, 125, 174, 188
immunological effects.....	14, 89, 96, 125, 174, 188
$K_{ow}$ .....	197
menstrual.....	9, 107, 137
micronuclei.....	128, 185, 187
musculoskeletal effects.....	87
neonatal.....	14, 96, 97, 112, 174, 185, 186, 187, 189, 192
neurobehavioral.....	89, 172, 185, 188
neurodevelopmental.....	15, 188
neurological effects.....	89, 171
nonhypoxic mechanisms.....	21, 189, 191
norepinephrine.....	126
nuclear.....	170
ocular effects.....	88
odds ratio.....	21, 35, 44, 45, 46, 73, 80, 81, 86, 107, 108, 109, 110, 111, 112, 113
pharmacodynamic.....	144
pharmacokinetic.....	144, 145, 146, 162, 173, 176, 189, 190
placenta.....	227
poisoning.....	8, 11, 12, 15, 16, 21, 26, 27, 30, 31, 33, 87, 88, 89, 90, 91, 95, 138, 143, 176, 177, 178, 181, 182, 185, 187, 189, 190, 192, 224, 240
rate constant.....	139, 208
renal effects.....	88
reproductive effects.....	95, 187
respiratory effects.....	16, 33, 34, 35, 47, 175, 186, 189
retention.....	140, 142
solubility.....	197
systemic effects.....	33
toxicokinetic.....	21, 11, 130, 190
weanling.....	125

