

# Maternity Practices in Infant Nutrition and Care in Florida —2011 mPINC Survey



More information is at [www.cdc.gov/mpinc](http://www.cdc.gov/mpinc)

This report provides data from the 2011 mPINC survey for Florida. It describes specific opportunities to improve mother-baby care at hospitals and birth centers in Florida in order to more successfully meet national quality of care standards for perinatal care.

## Breastfeeding is a National Priority

Breastfeeding is associated with decreased risk for infant morbidity and mortality as well as maternal morbidity,<sup>1</sup> and provides optimal infant nutrition. *Healthy People 2020*<sup>2</sup> establishes breastfeeding initiation, continuation, and exclusivity as national priorities.

## Changes in Maternity Care Practices Improve Breastfeeding Rates

Maternity practices in hospitals and birth centers can influence breastfeeding behaviors during a period critical to successful establishment of lactation.<sup>3</sup> Abundant literature, including a Cochrane review, document that institutional changes in maternity care practices to make them more supportive of breastfeeding increase initiation and continuation of breastfeeding.<sup>4</sup>

## Breastfeeding Support in Florida Facilities

### Strengths

	<p><b>Documentation of Mothers' Feeding Decisions</b> Staff at all (100%) facilities in Florida consistently ask about and record mothers' infant feeding decisions.</p>	Standard documentation of infant feeding decisions is important to adequately support maternal choice.
	<p><b>Provision of Breastfeeding Advice and Counseling</b> Staff at 91% of facilities in Florida provide breastfeeding advice and instructions to patients who are breastfeeding, or intend to breastfeed.</p>	The American Academy of Pediatrics (AAP) recommends pediatricians provide patients with complete, current information on the benefits and methods of breastfeeding to ensure that the feeding decision is a fully informed one. Patient education is important in order to establish breastfeeding.

### Needed Improvements

	<p><b>Appropriate Use of Breastfeeding Supplements</b> Only 18% of facilities in Florida adhere to standard clinical practice guidelines against routine supplementation with formula, glucose water, or water.</p>	The American Academy of Pediatrics (AAP) and the American College of Obstetricians and Gynecologists (ACOG) Guidelines for Perinatal Care recommend against routine supplementation because supplementation with formula and/or water makes infants more likely to receive formula at home and stop breastfeeding prematurely.
	<p><b>Inclusion of Model Breastfeeding Policy Elements</b> Only 17% of facilities in Florida have comprehensive breastfeeding policies including all model breastfeeding policy components recommended by the Academy of Breastfeeding Medicine (ABM).</p>	The ABM model breastfeeding policy elements are the result of extensive research on best practices to improve breastfeeding outcomes. Facility policies determine the nature of care that is available to patients. Facilities with comprehensive policies consistently have the highest rates of exclusive breastfeeding, regardless of patient population characteristics such as ethnicity, income, and payer status.
	<p><b>Protection of Patients from Formula Marketing</b> Only 43% of facilities in Florida adhere to clinical and public health recommendations against distributing formula company discharge packs.</p>	Distribution of discharge packs contributes to premature breastfeeding discontinuation. The ACOG, AAP, American Public Health Association (APHA), and the federal Government Accountability Office (GAO) all identify this practice as inappropriate in medical environments and recommend against it.
	<p><b>Initiation of Mother and Infant Skin-to-Skin Care</b> Only 53% of facilities in Florida initiate skin-to-skin care for at least 30 minutes upon delivery of the newborn.</p>	Upon delivery, the newborn should be placed skin-to-skin with the mother and allowed uninterrupted time to initiate and establish breastfeeding in order to improve infant health outcomes and reduce the risk of impairment of the neonatal immune system from unnecessary non-breast milk feeds.

Every two years, CDC administers the national Maternity Practices in Infant Nutrition and Care (mPINC) survey to all hospitals and birth centers in the U.S. that provide maternity care.

Data from this survey can be used to establish evidence-based, breastfeeding-supportive maternity practices as standards of care in hospitals and birth centers across the US. Improved care will help meet *Healthy People 2020* breastfeeding objectives and will help improve maternal and child health nationwide.



# Florida Summary —2011 mPINC Survey

**Survey Method** At each facility, the person who is the most knowledgeable about the facility's maternity practices related to healthy newborn feeding and care completes the CDC mPINC survey.

**Response Rate** 67% of the 133 eligible facilities in Florida responded to the 2011 mPINC Survey. Each participating facility received its facility-specific mPINC benchmarking report in October 2012.

**Florida's Composite Quality Practice Score\*** **69**  
(out of 100)

**Florida's Composite Rank†** **26**  
(out of 53)

mPINC Dimension of Care	FL Quality Practice Subscore*	Ideal Response to mPINC Survey Question	Percent of FL Facilities with Ideal Response	FL Item Rank†
<b>Labor and Delivery Care</b>	<b>67</b>	Initial skin-to-skin contact is ≥30 min w/in 1 hour (vaginal births)	53	26
		Initial skin-to-skin contact is ≥30 min w/in 2 hours (cesarean births)	37	31
		Initial breastfeeding opportunity is w/in 1 hour (vaginal births)	53	29
		Initial breastfeeding opportunity is w/in 2 hours (cesarean births)	33	49
		Routine procedures are performed skin-to-skin	27	25
<b>Feeding of Breastfed Infants</b>	<b>78</b>	Initial feeding is breast milk (vaginal births)	60	49
		Initial feeding is breast milk (cesarean births)	47	50
		Supplemental feedings to breastfeeding infants are rare	18	38
		Water and glucose water are not used	86	19
<b>Breastfeeding Assistance</b>	<b>84</b>	Infant feeding decision is documented in the patient chart	100	---
		Staff provide breastfeeding advice & instructions to patients	91	---
		Staff teach breastfeeding cues to patients	83	31
		Staff teach patients not to limit suckling time	49	24
		Staff directly observe & assess breastfeeding	90	---
		Staff use a standard feeding assessment tool	70	25
		Staff rarely provide pacifiers to breastfeeding infants	40	24
<b>Contact Between Mother and Infant</b>	<b>79</b>	Mother-infant pairs are not separated for postpartum transition	64	27
		Mother-infant pairs room-in at night	82	15
		Mother-infant pairs are not separated during the hospital stay	53	15
		Infant procedures, assessment, and care are in the patient room	9	8
		Non-rooming-in infants are brought to mothers at night for feeding	77	39
<b>Facility Discharge Care</b>	<b>49</b>	Staff provide appropriate discharge planning (referrals & other multi-modal support)	33	17
		Discharge packs containing infant formula samples and marketing products are not given to breastfeeding patients	43	28
<b>Staff Training</b>	<b>58</b>	New staff receive appropriate breastfeeding education	12	19
		Current staff receive appropriate breastfeeding education	13	36
		Staff received breastfeeding education in the past year	52	21
		Assessment of staff competency in breastfeeding management & support is at least annual	67	9
<b>Structural &amp; Organizational Aspects of Care Delivery</b>	<b>72</b>	Breastfeeding policy includes all 10 model policy elements	17	25
		Breastfeeding policy is effectively communicated	78	28
		Facility documents infant feeding rates in patient population	73	20
		Facility provides breastfeeding support to employees	81	12
		Facility does not receive infant formula free of charge	16	21
		Breastfeeding is included in prenatal patient education	92	---
		Facility has a designated staff member responsible for coordination of lactation care	71	26

\* Quality Practice scores range from 0 to 100 for each question, dimension of care, facility, and state. The highest, best possible score for each is 100. Each facility and state's "Composite Quality Practice Score" is made up of subscores for practices in each of 7 dimensions of care.

† Ranks range from 1 to 52, with 1 being the highest rank. In case of a tie, both are given the same rank.

- State ranks are not shown for survey questions with 90% or more facilities reporting ideal responses.

## References

- Ip S, Chung M, Raman G, et al. Breastfeeding and maternal and infant health outcomes in developed countries. Rockville, MD: US Dept of Health and Human Services, Agency for Healthcare Research and Quality; 2007.
- US Dept of Health and Human Services. Healthy People 2020 Summary of Objectives: Maternal, Infant, and Child Health. Available at <http://www.healthypeople.gov/2020/topics/objectives/2020/pdfs/MaternalChildHealth.pdf>
- DiGirolamo AM, Grummer-Strawn LM, Fein S. Effect of maternity care practices on breastfeeding. Pediatrics 2008;122, Supp 2:S43-9.
- Fairbank L, O'Meara S, Renfrew MJ, Woolridge M, Snowden AJ, Lister-Sharp D. A systematic review to evaluate the effectiveness of interventions to promote the initiation of breastfeeding. Health Technology Assessment 2000;4:1-171.

## Improvement is Needed in Maternity Care Practices and Policies in Florida.

Many opportunities exist to protect, promote, and support breastfeeding mothers and infants in Florida.

### Potential opportunities:

- Examine Florida regulations for maternity facilities and evaluate their evidence base.
- Sponsor a Florida-wide summit of key decision-making staff at maternity facilities to highlight the importance of evidence-based practices for breastfeeding.
- Encourage and support hospital staff across Florida to be trained in providing care that supports mothers to breastfeed.
- Establish links among maternity facilities and community breastfeeding support networks in Florida.
- Implement evidence-based practices in medical care settings across Florida that support mothers' efforts to breastfeed.
- Integrate maternity care into related hospital-wide Quality Improvement efforts across Florida.
- Promote utilization of the Joint Commission's Perinatal Care Core Measure Set including exclusive breast milk feeding at hospital discharge in Florida hospital data collection systems.

### Questions about the mPINC survey?

Information about the mPINC survey, benchmark reports, scoring methods, and complete references are available at: [www.cdc.gov/mpinc](http://www.cdc.gov/mpinc)

For more information:  
Division of Nutrition, Physical Activity, and Obesity  
Centers for Disease Control and Prevention  
Atlanta, GA USA

February 2013