

Prevention Status Report | 2013

Advancing evidence-based policy and practice

PSR | 2013



Minnesota



For more information, please visit www.cdc.gov/stltpublichealth/psr

Centers for Disease Control and Prevention

Office for State, Tribal, Local and Territorial Support

Prevention Status Report | 2013

Minnesota

PSR | 2013

Overview	1
Summary	2
Excessive Alcohol Use	4
Food Safety	8
Healthcare-Associated Infections	11
Heart Disease and Stroke	13
HIV	16
Motor Vehicle Injuries	19
Nutrition, Physical Activity, and Obesity	22
Prescription Drug Overdose	27
Teen Pregnancy	30
Tobacco Use	33

Suggested Citation:

Centers for Disease Control and Prevention.
Prevention Status Reports 2013—Minnesota.
Atlanta, GA: US Department of Health and Human
Services; 2014.

Prevention Status Report | 2013

Minnesota

The Prevention Status Reports (PSRs) highlight—for all 50 states and the District of Columbia—the status of public health policies and practices designed to prevent or reduce 10 important health problems or concerns:

Excessive alcohol use
Food safety
Healthcare-associated infections
Heart disease and stroke
HIV

Motor vehicle injuries
Nutrition, physical activity, and obesity
Prescription drug overdose
Teen pregnancy
Tobacco use

PSR Framework

The PSRs follow a simple framework:

- Describe the public health **problem** using public health data
- Identify potential **solutions** to the problem drawn from research and expert recommendations
- Report the **status** of those solutions for each state and the District of Columbia

Criteria for Selection of Policies and Practices

The policies and practices included in the PSRs were selected because they

- Can be monitored using state-level data that are readily available for most states and the District of Columbia
- Meet one or more of the following criteria:
 - Supported by systematic review(s) of scientific evidence of effectiveness (e.g., *The Guide to Community Preventive Services*)
 - Explicitly cited in a national strategy or national action plan (e.g., *Healthy People 2020*)
 - Recommended by a recognized expert body, panel, organization, study, or report with an evidence-based focus (e.g., Institute of Medicine)

Ratings

The PSRs use a simple, three-level rating scale to provide a practical assessment of the status of policies and practices in each state and the District of Columbia:

- A **green** rating indicates that the policy or practice is established in accordance with supporting evidence and/or expert recommendations.
- A **yellow** rating indicates that the policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.
- A **red** rating indicates that the policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

It is important to note that the ratings reflect the *status of policies and practices* and do not reflect the *status of efforts* by state health departments, other state agencies, or other organizations to establish or strengthen those policies and practices. Strategies for improving public health vary by individual state needs, resources, and public health priorities.

Prevention Status Reports—Summary for Minnesota | 2013

The Prevention Status Reports (PSRs) highlight—for all 50 states and the District of Columbia—the status of public health policies and practices designed to prevent or reduce 10 important health problems or concerns. Below is a summary of Minnesota's PSR ratings for 2013.

PSR Policies and Practices by Topic	2013 PSR Rating
Excessive Alcohol Use	
State beer tax	Red
State distilled spirits tax	Yellow
State wine tax	Red
Commercial host (dram shop) liability law	Green
Local authority to regulate alcohol outlet density	Green
Food Safety	
Speed of pulsed-field gel electrophoresis (PFGE) testing of reported <i>E. coli</i> O157 cases	Green
Completeness of PFGE testing of reported <i>Salmonella</i> cases	Green
Healthcare-Associated Infections (HAIs)	
State health department participation in statewide HAI prevention efforts	Green
Heart Disease and Stroke	
Implementation of electronic health records	Yellow
Pharmacist collaborative drug therapy management policy	Green
HIV	
State Medicaid reimbursement for routine HIV screening	Green
State HIV testing laws	Green
Reporting of CD4 and viral load data to state HIV surveillance program	Green
Motor Vehicle Injuries	
Seat belt law	Green
Child passenger restraint law	Yellow
Graduated driver licensing system	Red
Ignition interlock law	Green
Nutrition, Physical Activity, and Obesity	
Secondary schools not selling less nutritious foods and beverages	Red
State nutrition standards policy for foods and beverages sold or provided by state government agencies	Red
Inclusion of nutrition and physical activity standards in state regulations of licensed childcare facilities	Red
State physical education time requirement for high school students	Red
Average birth facility score for breastfeeding support	Yellow
Prescription Drug Overdose	
State pain clinic law	Red
Prescription drug monitoring programs following selected best practices	Yellow
Teen Pregnancy	
Expansion of state Medicaid family planning eligibility	Yellow
Tobacco Use	
State cigarette excise tax	Yellow
Comprehensive state smoke-free policy	Green
Funding for tobacco control	Red



PSR Rating System*

Green	The policy or practice is established in accordance with supporting evidence and/or expert recommendations.
Yellow	The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.
Red	The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

*The rating systems for the Excessive Alcohol Use (<http://www.cdc.gov/stltpublichealth/psr/alcohol/>) and Nutrition, Physical Activity, and Obesity (<http://www.cdc.gov/stltpublichealth/psr/npao/>) reports varied slightly. For details, please visit their respective pages on the PSR website. A more detailed explanation of the PSR rating system is available at <http://www.cdc.gov/stltpublichealth/psr/>.

More Information

For more information about public health activities in Minnesota, visit the Minnesota Department of Health website (<http://www.health.state.mn.us/>). For additional resources and to view reports for other states, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/>).

Prevention Status Report | 2013

Excessive Alcohol Use

Minnesota

Public Health Problem



Excessive alcohol use is responsible for about 88,000 deaths and 2.5 million years of potential life lost in the United States each year (1). Binge drinking (five or more drinks per occasion for men or four or more drinks per occasion for women) is responsible for more than half the deaths and two-thirds of the years of potential life lost resulting from excessive alcohol use (2).

Excessive drinking results in 1,257 deaths and 32,829 years of potential life lost each year in Minnesota (1).




In Minnesota, 22.1% of adults reported binge drinking in 2011 (3). Data were not available for binge drinking among high school students (4).




Excessive alcohol use cost the United States \$223.5 billion, or \$1.90 per drink consumed, in 2006 as a result of lost workplace productivity, healthcare expenses, and crime (5). In Minnesota, excessive alcohol use cost \$3.5 billion, or \$1.65 per drink (6).

Binge drinking among adults



Source: Behavioral Risk Factor Surveillance System (3)


Binge drinking among high school students



Source: Youth Risk Behavior Surveillance System (4)


Note: Minnesota data were not available for one or more years from the source used for this chart. Similar data may be available from another national or state source.

**Binge drinking intensity among adults
(in number of drinks per occasion)**



Source: Behavioral Risk Factor Surveillance System (3)

**Alcohol consumption per person aged ≥14
(in gallons)**



Source: Alcohol Epidemiologic Data System (7)

Prevention Status Report | 2013

Excessive Alcohol Use

Minnesota

Policy and Practice Solutions

This report focuses on policies and practices recommended by the Community Preventive Services Task Force on the basis of scientific studies supporting their effectiveness in reducing excessive alcohol consumption and related harms (8). These policies and practices include 1) increasing alcohol excise taxes (e.g., state taxes on beer, distilled spirits, and wine); 2) having commercial host (dram shop) liability laws; and 3) regulating alcohol outlet density (8–10). Other strategies supported by scientific evidence include avoiding further privatization of retail alcohol sales and providing adults (including pregnant women) with screening and brief intervention for excessive alcohol use (11,12). For information about why certain alcohol-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/alcohol/>).

Status of Policy and Practice Solutions in Minnesota

State beer tax

As of January 1, 2012, Minnesota's excise tax per gallon of beer was \$0.15 (13).

Task Force on Community Preventive Services recommendation: Increase alcohol excise taxes. Studies show that a 10% increase in the price of beer would likely reduce beer consumption by approximately 5% (8).



Rating	State beer tax
Green	≥\$1.00 per gallon
Yellow	\$0.50–\$0.99 per gallon
Red	\$0.00–\$0.49 per gallon

State distilled spirits tax

As of January 1, 2012, Minnesota's excise tax per gallon of distilled spirits was \$5.03 (14).

Task Force on Community Preventive Services recommendation: Increase alcohol excise taxes. Studies show that a 10% increase in the price of distilled spirits would likely reduce distilled spirits consumption by approximately 8% (8).



Rating	State distilled spirits tax
Green	≥\$8.00 per gallon
Yellow	\$4.00–\$7.99 per gallon
Red	\$0.00–\$3.99 per gallon

State wine tax

As of January 1, 2012, Minnesota's excise tax per gallon of wine was \$0.30 (15).

Task Force on Community Preventive Services recommendation: Increase alcohol excise taxes. Studies show that a 10% increase in the price of wine would likely reduce wine consumption by approximately 6% (8).



Rating	State wine tax
Green	≥\$2.00 per gallon
Yellow	\$1.00–\$1.99 per gallon
Red	\$0.00–\$0.99 per gallon

Commercial host (dram shop) liability laws

As of January 1, 2011, Minnesota had commercial host liability with no major limitations (16,17).

Task Force on Community Preventive Services recommendation: Presence of commercial host (dram shop) liability for sale or service to either underage patrons or intoxicated adults. Evidence shows these laws are associated with a reduction in alcohol-related harms, including a median 6.4% reduction in deaths from motor vehicle crashes (9).



Rating	State had
Green	Commercial host liability with no major limitations
Yellow	Commercial host liability with major limitations
Red	No commercial host liability

Prevention Status Report | 2013

Excessive Alcohol Use

Minnesota

Local authority to regulate alcohol outlet density

As of January 1, 2012, Minnesota had exclusive local alcohol retail licensing (18).

Task Force on Community Preventive Services recommendation: Use regulatory authority (e.g., through licensing and zoning) to limit alcohol outlet density. Evidence shows greater alcohol outlet density is associated with excessive drinking and related harms, including injuries and violence (10). Local control allows communities to better address density problems (18).



Rating	State had
Green	Exclusive local or joint state/local alcohol retail licensing
Yellow	Exclusive state alcohol retail licensing with local zoning authority or other mixed policies
Red	Exclusive state alcohol retail licensing

Simplified Rating System

A more detailed explanation of the rating system for excessive alcohol use is available at <http://www.cdc.gov/stltpublichealth/psr/alcohol/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations. Higher tax levels are rated green.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations. Intermediate tax levels are rated yellow.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations. Lower tax levels are rated red.

Indicator Definitions

State beer tax: The excise tax rate, in dollars per gallon, imposed by the state on beer containing 5% alcohol by volume. State beer excise tax does not include any additional taxes, such as those based on price rather than volume (e.g., ad valorem or sales taxes) that states may have implemented at the wholesale or retail level. State beer taxes ranged from \$0.02 to \$1.07 across states for which excise tax data were available.

State distilled spirits tax: The excise tax rate, in dollars per gallon, imposed by the state on distilled spirits containing 40% alcohol by volume. State distilled spirits excise tax does not include any additional taxes, such as those based on price rather than volume (e.g., ad valorem or sales taxes) that states may have implemented at the wholesale or retail level. State distilled spirits taxes ranged from \$1.50 to \$14.25 across states for which excise tax data were available. For states with different tax rates for distilled spirits sold off-sale (e.g., at liquor stores) and on-sale (e.g., at restaurants), the off-sale tax rate has been reported.

State wine tax: The excise tax rate, in dollars per gallon, imposed by the state on wine containing 12% alcohol by volume. State wine excise tax does not include any additional taxes, such as those based on price rather than volume (e.g., ad valorem or sales taxes) that states may have implemented at the wholesale or retail level. State wine taxes ranged from \$0.11 to \$2.50 across states for which excise tax data were available.

Commercial host (dram shop) liability laws: Laws that hold alcohol retailers liable for alcohol-attributable harms (e.g., injuries or deaths resulting from alcohol-related motor vehicle crashes) caused by patrons who were illegally sold or served alcohol because they were either intoxicated or under the minimum legal drinking age of 21 years at the time of sale or service. State commercial host liability laws are considered to have major limitations if they 1) cover underage patrons or intoxicated adults but not both, 2) require increased evidence for finding liability, 3) set limitations on damage awards, or 4) set restrictions on who may be sued.

Local authority to regulate alcohol outlet density: The extent to which a local government can implement zoning (land use) or licensing controls over the number of alcohol retailers (e.g., bars, restaurants, liquor stores) in its geographic area.

Prevention Status Report | 2013

Excessive Alcohol Use

Minnesota

References

1. CDC. Alcohol and Public Health: Alcohol-Related Disease Impact (ARDI) [database]. Accessed Dec 13, 2013.
2. CDC. Alcohol-attributable deaths and years of potential life lost, United States, 2001. *MMWR* 2004;53:866–70.
3. CDC. Behavioral Risk Factor Surveillance System (BRFSS) [database]. Accessed Dec 6, 2012.
4. CDC. Youth Online: High School Youth Risk Behavior Surveillance (YRBS) [database]. Accessed Feb 27, 2013.
5. Bouchery EE, Harwood HJ, Sacks JJ, et al. Economic costs of excessive alcohol consumption in the United States, 2006. *American Journal of Preventive Medicine* 2011;41:516–24; and correction, *American Journal of Preventive Medicine* 2013;44:198.
6. Sacks JJ, Roeber J, Bouchery EE, et al. State costs of excessive alcohol consumption, 2006. *American Journal of Preventive Medicine* 2013;45:474–85.
7. National Institute on Alcohol Abuse and Alcoholism. Alcohol Epidemiologic Data System. Apparent Per Capita Alcohol Consumption: National, State, and Regional Trends, 1977–2010. Bethesda, MD: National Institutes of Health; 2010.
8. Community Preventive Services Task Force. Preventing excessive alcohol consumption: increasing alcohol taxes. In: *Guide to Community Preventive Services*. Updated Jun 2007.
9. Community Preventive Services Task Force. Preventing excessive alcohol consumption: dram shop liability. In: *Guide to Community Preventive Services*. Updated Mar 2010.
10. Community Preventive Services Task Force. Preventing excessive alcohol consumption: regulation of alcohol outlet density. In: *Guide to Community Preventive Services*. Updated Feb 2007.
11. Community Preventive Services Task Force. Preventing excessive alcohol consumption: privatization of alcohol retail sales. In: *Guide to Community Preventive Services*. Updated Apr 2011.
12. US Preventive Services Task Force. Screening and Behavioral Counseling Interventions in Primary Care to Reduce Alcohol Misuse [website]. Updated Oct 2012.
13. National Institute on Alcohol Abuse and Alcoholism. Alcohol beverages taxes: beer. Alcohol Policy Information System [database]. Accessed Dec 6, 2012.
14. National Institute on Alcohol Abuse and Alcoholism. Alcohol beverages taxes: distilled spirits. Alcohol Policy Information System [database]. Accessed Dec 6, 2012.
15. National Institute on Alcohol Abuse and Alcoholism. Alcohol beverages taxes: wine. Alcohol Policy Information System [database]. Accessed Dec 6, 2012.
16. Substance Abuse and Mental Health Services Administration. Report to Congress on the Prevention and Reduction of Underage Drinking. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2011.
17. Mosher JF, Cohen EN, Jernigan DH. Commercial host (dram shop) liability: current status and trends. *American Journal of Preventive Medicine* 2013;45:347–53.
18. Mosher JF, Treffers R. State pre-emption, local control, and alcohol retail outlet density regulation. *American Journal of Preventive Medicine* 2013;44:399–405.

Prevention Status Report | 2013

Food Safety

Minnesota

Public Health Problem



Diseases spread by a wide variety of contaminated foods continue to challenge the public health system. Bacteria, viruses, parasites, and chemicals can cause foodborne diseases, which can vary from mild to fatal (1). Robust surveillance for these diseases is essential for detecting outbreaks. It also provides critical information to food regulatory agencies and the food industry so that appropriate control and preventive measures can be implemented (2).



CDC estimates that each year, roughly 1 in 6 Americans (or 48 million people) gets sick, 128,000 are hospitalized, and 3,000 die due to foodborne diseases (3). Risk for infection and severity varies at different ages and stages of health (4).



Foodborne illness is costly. According to a 2012 study, 14 pathogens alone are estimated to cost \$14.1 billion in the United States per year. This includes medical costs (doctor visits and hospitalizations), loss due to premature death, and time lost from work (5).

Policy and Practice Solutions

This report focuses on select practices recommended by the Council to Improve Foodborne Outbreak Response on the basis of scientific evidence supporting their effectiveness in improving foodborne disease surveillance and detection activities (2). These practices include 1) increasing the speed of DNA fingerprinting using pulsed-field gel electrophoresis (PFGE) testing for all reported cases of Shiga toxin-producing *Escherichia coli* (*E. coli*) O157 and 2) increasing the completeness of PFGE testing of *Salmonella*. PFGE is a technique used to distinguish between strains of organisms at the DNA level. For information about why certain food safety-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/foodsafety/>).

Prevention Status Report | 2013

Food Safety

Minnesota

Status of Policy and Practice Solutions in Minnesota

Speed of pulsed-field gel electrophoresis (PFGE) testing of reported *E. coli* O157 cases

In 2011, Minnesota tested 92.2% of *E. coli* O157 cases within 4 days (6).

CDC target: Testing of 90% of annual reported *E. coli* O157 cases within four days. The CDC Public Health Emergency Preparedness Cooperative Agreement established this and other national performance targets for food safety and provides federal funding to states and the District of Columbia. Performing DNA fingerprinting as quickly as possible for all Shiga toxin-producing *E. coli* improves detection of outbreaks. Rapid outbreak detection can help prevent additional cases and identify control and prevention measures for food regulatory agencies and the food industry (2).



Rating	Percentage of annual reported cases tested within four days:
Green	≥90.0%
Yellow	60.0%–89.9%
Red	<60.0%

Completeness of PFGE testing of reported *Salmonella* cases

In 2011, Minnesota tested 100% of reported *Salmonella* cases (6,7).

Research and experts in the field agree that performing DNA fingerprinting of all *Salmonella* cases would improve detection of outbreaks (2).



Rating	Percentage of annual reported cases tested by PFGE:
Green	≥90.0%
Yellow	60.0%–89.9%
Red	<60.0%

Simplified Rating System

A more detailed explanation of the rating system for food safety is available at <http://www.cdc.gov/stltpublichealth/psr/foodsafety/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Prevention Status Report | 2013

Food Safety

Minnesota

Indicator Definitions

Speed of PFGE testing of reported *E. coli* O157 cases: The annual proportion of *E. coli* O157 PFGE patterns reported to CDC (i.e., uploaded into PulseNet, the CDC-coordinated national molecular subtyping network for foodborne disease surveillance) within four working days of receiving the isolate in the state or District of Columbia public health PFGE lab.

Completeness of PFGE testing of reported *Salmonella* cases: The annual proportion of *Salmonella* cases reported to CDC's National Notifiable Diseases Surveillance System with PFGE patterns uploaded into PulseNet.

References

1. Scallan E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States—major pathogens. *Emerging Infectious Diseases* 2011;17:7–15.
2. Council to Improve Foodborne Outbreak Response. Guidelines for Foodborne Disease Outbreak Response. Atlanta, GA: Council of State and Territorial Epidemiologists; 2009.
3. Scallan E, Griffin P, Angulo F, et al. Foodborne illness acquired in the United States—unspecified agents. *Emerging Infectious Diseases* 2011;17:16–22.
4. Lund BM, O'Brien SJ. The occurrence and prevention of foodborne disease in vulnerable people. *Foodborne Pathogens and Disease* 2011;8:961–73.
5. Hoffmann S, Batz M, Morris JG. Annual cost of illness and quality-adjusted life year losses in the United States due to 14 foodborne pathogens. *Journal of Food Protection* 2012;75:1292–1302.
6. CDC. PulseNet [database]. Accessed Dec 19, 2012.
7. CDC. Final 2011 reports of nationally notifiable infectious diseases. *MMWR* 2012;60(32):624–37.

Prevention Status Report | 2013


Healthcare-Associated Infections

Minnesota

Public Health Problem

- ⚠️ HAIs occur in all settings where patients receive medical care, including hospital and nonhospital settings, and are associated with increased illness and death. CDC estimates that each year in the United States, 1 in 20 hospital patients gets an HAI (1).
- 👤 More than one million HAIs occur across all US healthcare settings combined. For example, *Clostridium difficile* infections kill 14,000 people in the United States each year (2).
- 💰 HAIs result in an estimated \$30 billion in excess healthcare costs nationally each year (3).

Central line-associated bloodstream infection—standardized infection ratio



Source: National and State Healthcare-Associated Infections Standardized Infection Ratio Report (4)

Note: Minnesota data were not available for one or more years from the source used for this chart. Similar data may be available from another national or state source.

What is a standardized infection ratio (SIR)?

The SIR is a summary measure used to track HAIs over time. It adjusts for the fact that each healthcare facility treats different types of patients. The SIR compares the number of infections reported to the National Healthcare Safety Network in 2011 to the number of infections that would be predicted based on national, historical baseline data:

$$\text{SIR} = \frac{\text{Observed \# of HAIs}}{\text{Predicted \# of HAIs}}$$

Policy and Practice Solutions

CDC recommends strategies for surveillance, prevention, and control of HAIs and antimicrobial resistance wherever health care is provided, including hospitals as well as ambulatory and long-term care facilities. CDC works closely with states and the District of Columbia on strategies to implement these recommendations. This collaborative effort among CDC, state and district health departments, and facilities will improve healthcare quality across the nation, working toward meeting the standards and targets set forth in the Department of Health and Human Service's *National Action Plan to Prevent Healthcare-Associated Infections* (5).

This report focuses on state health departments leading and participating in statewide HAI prevention efforts, a practice that helps improve existing prevention strategies by investing in both new and ongoing HAI prevention efforts and prioritizing HAIs as a serious public health concern. State health departments are encouraged to also engage in other practices that will provide actionable HAI data and lead to expanded HAI prevention. These include 1) state health departments validating data sent to CDC's National Healthcare Safety Network (NHSN), ideally including data on central line-associated bloodstream infections (CLABSIs); catheter-associated urinary tract infections (CAUTIs); and surgical site infections; and 2) working with CDC and other partners using NHSN data to target facilities and units most in need of consultation to prevent HAIs and antimicrobial resistance. For information about why certain HAI-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/hai/>).

Prevention Status Report | 2013

Healthcare-Associated Infections

Minnesota

Status of Policy and Practice Solutions in Minnesota

State health department participation in statewide HAI prevention efforts

In 2013, Minnesota led or participated in broad prevention collaboratives to prevent multiple HAIs in acute care facilities, including CLABSIs, CAUTIs, and *C. difficile*, as well as HAIs in long-term care facilities (6).

Implementing HAI prevention strategies and tracking the impact of those strategies have led to improvements in clinical practice and medical procedures, development of evidence-based infection control guidance, and prevention successes (7).



Rating	State health department
Green	Led or participated in a broad prevention collaborative addressing at least one HAI
Yellow	N/A
Red	Did not participate in a broad prevention collaborative addressing HAIs

Simplified Rating System

A more detailed explanation of the rating system for HAIs is available at <http://www.cdc.gov/stltpublichealth/psr/hai/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Indicator Definitions

Participation in statewide HAI prevention efforts: State health department participation in or leadership of broad prevention collaboratives addressing one or more of the following types of HAIs: central line-associated bloodstream infections, surgical site infections, catheter-associated urinary tract infections, ventilator-associated pneumonia, methicillin-resistant *Staphylococcus aureus*, and *C. difficile*.

References

1. CDC. Healthcare-Associated Infections: The Burden [website]. Updated Dec 13, 2010.
2. CDC. Vital Signs—Making Health Care Safer: Stopping *C. difficile* Infections. Atlanta, GA: US Department of Health and Human Services; 2012.
3. Scott RD 2nd. The Direct Medical Costs of Healthcare-Associated Infections in U.S. Hospitals and the Benefits of Prevention. Atlanta, GA: US Department of Health and Human Services; 2009.
4. CDC. National and State Healthcare-Associated Infections Standardized Infection Ratio Report. January–December 2011. Atlanta, GA: US Department of Health and Human Services; 2013.
5. US Department of Health and Human Services. National Action Plan to Prevent Healthcare-Associated Infections: Roadmap to Elimination [website]. Updated Apr 2012.
6. CDC. State-Based HAI Prevention [website]. Updated May 10, 2013.
7. CDC. Preventing Healthcare-Associated Infections [website]. Updated Apr 17, 2012.

Prevention Status Report | 2013

Heart Disease and Stroke

Minnesota

Public Health Problem



Cardiovascular disease—including heart disease, stroke, and other vascular diseases—is the leading cause of death in the United States. Each year, nearly 800,000 people die from cardiovascular disease, accounting for one in every three deaths (1).




An estimated 67 million American adults have high blood pressure and 71 million American adults have high levels of low-density lipoprotein (LDL) cholesterol. These are two leading risk factors for heart disease and stroke (2,3).




About one of every six healthcare dollars in the United States is spent on treating cardiovascular disease. Annual US cardiovascular disease costs exceed \$192.1 billion in direct medical expenses and \$312.6 billion when indirect expenses are included (4).

**Coronary heart disease death rate
(age-adjusted rate per 100,000 population)**




Source: National Vital Statistics System—Mortality (5)
Healthy People 2020 target: 100.8/100,000
(dotted blue line) (6)

**Stroke death rate (age-adjusted rate per
100,000 population)**




Source: National Vital Statistics System—Mortality (5)
Healthy People 2020 target: 33.8/100,000
(dotted blue line) (6)

**Prevalence of self-reported
hypertension (age-adjusted)**



Source: Behavioral Risk Factor Surveillance System (BRFSS) (7)
Note: These rates were adjusted using the direct method and the 2000 standard US population (8).

**Prevalence of self-reported high
cholesterol (age-adjusted)**



Source: Behavioral Risk Factor Surveillance System (BRFSS) (7)
Note: These rates were adjusted using the direct method and the 2000 standard US population (8).

Prevention Status Report | 2013

Heart Disease and Stroke

Minnesota

Policy and Practice Solutions

This report focuses on policies and practices recommended by the Community Preventive Services Task Force, the US Surgeon General, and the Institute of Medicine on the basis of scientific studies supporting the policies' effectiveness in the management of heart disease and stroke risks (9–12). These policies and practices include 1) implementing electronic health records and 2) developing state policies that address collaborative drug therapy management, such as the use of pharmacists to facilitate collaborative practice agreements (10). Other strategies supported by scientific evidence and practice include promoting team-based care, establishing state-level policies for patient-centered medical homes, establishing stroke systems of care, and reducing sodium consumption at the community level. For information about why certain heart disease and stroke-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/heartandstroke/>).

Status of Policy and Practice Solutions in Minnesota

Implementation of electronic health records

As of December 2012, 25.8% of office-based physicians in Minnesota met criteria for meaningful use of electronic health records (12).

Research shows that electronic health records, when used with specific goals in mind (i.e., "meaningfully"), allow physicians, nurses, pharmacists, and other healthcare providers to proactively monitor and protect the health of their patients by tracking heart disease and stroke risk factors (13–15).

Note: This indicator reflects the percentage of physicians using electronic health records that can support 13 capabilities needed to meet Stage 1 Core Set objectives to demonstrate meaningful use. Other data from the federal Office of the National Coordinator for Health Information Technology reflect the percentage of physicians using a basic system, which has seven capabilities (16).

Rating	Percentage of office-based physicians meeting meaningful use criteria:
Green	31.0%–45.0%
Yellow	16.0%–30.9%
Red	0.0%–15.9%

Pharmacist collaborative drug therapy management (CDTM) policy

As of December 31, 2012, Minnesota had a statewide pharmacist CDTM policy for all health conditions (17).

State policies such as CDTM laws, which authorize pharmacists to enter into collaborative practice agreements with prescribing providers, can increase medication adherence rates and improve health outcomes (e.g., lower blood pressure and LDL cholesterol, reduced hemoglobin A1c, fewer adverse drug events) (10).

Rating	CDTM policy
Green	Authorized pharmacists to collaborate for all health conditions
Yellow	Authorized pharmacists to collaborate but did not cover chronic diseases, or collaboration was limited to specified hospital, medical, or clinical practice settings
Red	Did not exist

Simplified Rating System

A more detailed explanation of the rating system for heart disease and stroke indicators is available at <http://www.cdc.gov/stltpublichealth/psr/heartandstroke/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Prevention Status Report | 2013

Heart Disease and Stroke

Minnesota

Indicator Definitions

Implementation of electronic health records: An electronic health record is a real-time, digital, patient-centered record that replaces paper charts. "Meaningful use" of electronic health records means meeting criteria that focus on such aspects as engaging patients in their own care, sharing information among healthcare organizations, and providing support for decisions on national high-priority conditions. It is hoped that if healthcare providers meet these criteria, "meaningful use" will lead to 1) creation of tools that measure healthcare quality to improve clinical and population health, 2) increased transparency and efficiency, 3) individuals empowered to access clinical information, and 4) more robust research data on health systems (18). Electronic health records should include clinical decision supports, such as alerts for elevated blood pressure and cholesterol levels based on laboratory results, to support guidelines-based clinical decision making.

Pharmacist collaborative drug therapy management policy: A state legislative, regulatory, or other written policy that authorizes qualified pharmacists working within the context of a defined protocol to perform patient assessments; order drug therapy-related laboratory tests; administer drugs; and select, initiate, monitor, continue, and adjust drug regimens (19).

References

1. Kochanek KD, Xu JQ, Murphy SL, et al. Deaths: final data for 2009. National Vital Statistics Report 2011;60(3).
2. CDC. Vital signs: awareness and treatment of uncontrolled hypertension among adults—United States, 2003–2010. MMWR 2012;61(35):703–9.
3. CDC. Vital signs: prevalence, treatment, and control of high levels of low-density lipoprotein cholesterol. United States, 1999–2002 and 2005–2008. MMWR 2011;60(4):109–14.
4. Fryar CD, Chen T, Li X. Prevalence of Uncontrolled Risk Factors for Cardiovascular Disease: United States, 1999–2010. NCHS Data Brief, No. 103. Hyattsville, MD: US Department of Health and Human Services; 2012.
5. CDC. Compressed Mortality File 1999–2010. CDC WONDER [database]. Accessed Jan 2013.
6. US Department of Health and Human Services. Heart disease and stroke. In: Healthy People 2020. Rockville, MD: US Department of Health and Human Services; Updated Sep 6, 2012.
7. CDC. Behavioral Risk Factor Surveillance System [database]. Accessed Jun 25, 2013.
8. Klein RJ, Schoenborn CA. Age adjustment using the 2000 projected U.S. population. Healthy People Statistical Notes, No. 20. Hyattsville, MD: National Center for Health Statistics; 2001.
9. Community Preventive Services Task Force. Cardiovascular disease prevention and control: team-based care to improve blood pressure control. In: Guide to Community Preventive Services. Updated Apr 2012.
10. Giberson S, Yoder S, Lee MP. Improving Patient and Health System Outcomes through Advanced Pharmacy Practice. A Report to the U.S. Surgeon General. Rockville, MD: US Public Health Service; 2011.
11. Institute of Medicine. Primary Care and Public Health: Exploring Integration to Improve Population Health. Washington, DC: National Academies Press; 2012.
12. CDC. National Ambulatory Medical Care Survey. National Electronic Health Records Survey 2012. Unpublished data.
13. Kinn JW, Marek JC, O'Toole MF, et al. Effectiveness of the electronic medical record in improving the management of hypertension. Journal of Clinical Hypertension 2002;4(6):415–9.
14. Ross SE, Moore LA, Earnest MA, et al. Providing a web-based online medical record with electronic communication capabilities to patients with congestive heart failure: randomized trial. Journal of Medical Internet Research 2004;6:e12.
15. Rossi RA, Every NR. A computerized intervention to decrease the use of calcium channel blockers in hypertension. Journal of General Internal Medicine 1997;12:672–8.
16. US Department of Health and Human Services. Office of the National Coordinator for Health IT, Health IT Dashboard. Accessed Dec 2013.
17. CDC. Chronic Disease State Policy Tracking System [database]. Accessed Dec 7, 2012.
18. US Department of Health and Human Services. EHR Incentives & Certification: How to Attain Meaningful Use [website]. Accessed Dec 7, 2012.
19. American College of Clinical Pharmacy. ACCP position statement: collaborative drug therapy management by pharmacists—2003. Pharmacotherapy 2003;23(9):1210–25.

Prevention Status Report | 2013

HIV

Minnesota

Public Health Problem



CDC estimates that more than 1.1 million people in the United States are living with HIV, and 15.8% (about one in six) are not aware they are infected (1). In 2010, the White House released the first National HIV/AIDS Strategy for the United States to increase the nation's sense of urgency and to improve HIV prevention and care (2).




In 2011, 302 people in Minnesota were newly diagnosed with HIV infection (1). Twenty-five percent of these people were diagnosed late in the disease and therefore were at increased risk for disease progression, death, and transmission of HIV to others. In 2010, more than 21,000 people with HIV were estimated to have died in the United States. Of these, CDC estimates that 99 were from Minnesota (1).




The lifetime cost of medical care for a person with an early HIV diagnosis is about \$400,000 (3). This means that lifetime medical costs for the 302 Minnesota residents newly diagnosed with HIV in 2011 could exceed \$120 million.

Estimated annual prevalence rate of persons living with diagnosed HIV aged 13 years and older (per 100,000 population)




Source: National HIV Surveillance System (4)
Note: The y-axis for this graph varies by state.

Estimated annual rate of new HIV diagnoses among persons aged 13 years and older (per 100,000 population)




Source: National HIV Surveillance System (4)
Note: The y-axis for this graph varies by state.

Percentage of persons newly diagnosed with HIV who have late stage HIV



Source: National HIV Surveillance System (1)
Healthy People 2020 Target: 20.8% by 2015 (dotted blue line) (5)

Estimated annual death rate among persons diagnosed with HIV (per 1,000 people living with HIV)



Source: National HIV Surveillance System (1)

Prevention Status Report | 2013

HIV

Minnesota

Policy and Practice Solutions

This report highlights policies that reflect recent scientific advances in HIV prevention and medical care. These advances create new opportunities for reducing new HIV infections and HIV-related illness and death. These policies are important state-level tools that further the goals of the 2010 National HIV/AIDS Strategy (2), including 1) facilitating state Medicaid reimbursement for HIV screening (7), 2) making state HIV testing laws compatible with the 2006 CDC HIV testing recommendations (6,10), and 3) reporting all CD4 lymphocyte and HIV viral load data to the state HIV surveillance program (7). For information about how and why certain HIV-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/hiv/>).

Status of Policy and Practice Solutions in Minnesota

State Medicaid reimbursement for routine HIV screening

As of January 1, 2013, Medicaid reimbursed for routine HIV screening of adults aged 15 to 65 regardless of risk in Minnesota (7).

CDC and the US Preventive Services Task Force recommend that adolescents, adults, and pregnant women be screened for HIV, regardless of risk (6,8). All state and District of Columbia Medicaid programs cover medically necessary HIV testing (7).

Reimbursement for routine screening, meaning broad, population-based HIV screening, in contrast with "medically necessary" testing and screening targeted at those at higher risk, increases the availability of this important preventive service for low-income populations (6,9).



Rating	State Medicaid plan
Green	Reimbursed for routine HIV screening
Yellow	N/A
Red	Did not reimburse for routine HIV screening

State HIV testing laws

As of July 2013, Minnesota's HIV testing laws were consistent with CDC's 2006 HIV testing recommendations (10).

CDC recommends that all people aged 13–64 years be tested for HIV (6). HIV testing enables individuals with HIV to become aware of their health status and to access medical care and treatment. Studies show that individuals diagnosed with HIV are less likely to transmit HIV to others (2). State and District of Columbia laws can facilitate access to HIV testing.



Rating	State HIV testing laws compared to CDC's HIV testing recommendations were
Green	Consistent with consent and counseling parameters
Yellow	N/A
Red	Inconsistent with consent or counseling parameters

Reporting of CD4 and viral load data to state HIV surveillance program

As of July 2013, Minnesota required reporting of all CD4 and viral load results (including undetectable results) for surveillance purposes (10).

CD4 and HIV viral load data are critical to the medical care and health of people living with HIV. These data are also used to monitor progress toward achieving the goals of the National HIV/AIDS Strategy and to ensure that people living with HIV are linked to HIV medical care and retained in care (2).



Rating	State law, regulation, or directive
Green	Required reporting of all CD4 and HIV viral load data
Yellow	Required reporting of some but not all CD4 and HIV viral load data
Red	Did not require reporting of any CD4 and HIV viral load data

Prevention Status Report | 2013

HIV

Minnesota

Simplified Rating System

A more detailed explanation of the rating system for HIV is available at <http://www.cdc.gov/stltpublichealth/psr/hiv/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Indicator Definitions

State Medicaid reimbursement for routine HIV screening: Medicaid reimbursement of healthcare providers for costs associated with routine HIV screening regardless of risk. Data reflect the most recent survey examining coverage as of January 2013.

State HIV testing laws: State laws governing HIV testing. Laws may or may not be consistent with key parameters of consent and counseling outlined in CDC's 2006 HIV testing recommendations (4). The consent parameters include opt-out (rather than opt-in) testing, inclusion of HIV testing consent as part of general medical consent forms (rather than HIV-specific consent forms), and permission to give consent orally. The counseling parameter includes not requiring prevention counseling prior to testing.

Reporting of CD4 and viral load data to HIV surveillance program: Existence of state statutes, regulations or directives that address the reporting of all CD4 values and all HIV viral load results (detectable and undetectable) to the state HIV surveillance program. HIV viral load and CD4 data among people with HIV infection are useful as indicators of program effectiveness. Viral load measures the amount of virus in a person's blood. CD4 results provide a measure of a person's immune function and are used for determining the stage of HIV infection. Among people with HIV, CD4 results are often used to monitor disease progression and to time clinical care, and both HIV viral load and CD4 results are used to assess response to treatment.

References

1. CDC. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data—United States and 6 U.S. dependent areas—2011. HIV Surveillance Supplemental Report 2013;18(No. 5).
2. White House Office of National AIDS Policy. National HIV/AIDS Strategy for the United States; 2010.
3. Farnham PG, Gopalappa C, Sansom SL, et al. Updates on lifetime costs of care and quality of life estimates for HIV-infected persons in the United States: late versus early diagnosis and entry into care. *Journal of Acquired Immune Deficiency Syndrome*. 2013; 64: 183–189.
4. CDC. NCHHSTP Atlas [website]. Updated Jul 2013.
5. Department of Health and Human Services. HIV. In: *Healthy People 2020*. Rockville, MD: US Department of Health and Human Services. Updated Aug 28, 2013.
6. CDC. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR* 2006;55(RR-14):1–17.
7. Kaiser Family Foundation. State Medicaid Coverage of Routine HIV Screening; 2013.
8. Moyer VA, on behalf of the US Preventive Services Task Force. Screening for HIV: US Preventive Services Task Force recommendation statement. *Annals of Internal Medicine* 2013; Apr 30.
9. Kates J. Medicaid and HIV: A National Analysis. Menlo Park, CA: Kaiser Family Foundation; 2011.
10. CDC. HIV and the Law [website]. Updated Sep 2013.

Prevention Status Report | 2013

Motor Vehicle Injuries

Minnesota

Public Health Problem



Motor vehicle crashes are a leading cause of death in the United States for people aged 30 years or younger (1).




In 2011, motor vehicle crashes killed more than 32,000 people in the United States and injured more than 2.6 million (1,2).




In 2005 alone, motor vehicle crashes cost Americans \$99 billion in medical care, rehabilitation, and lost wages (3).

**Motor vehicle-related death rate
(per 100,000 population)**




Source: National Highway Traffic Safety Administration (4)
Healthy People 2020 Target: 12.4/100,000 (dotted blue line) (5)

**Motor vehicle-related death rate
among drivers aged 15–20 years (per
100,000 population)**




Source: National Highway Traffic Safety Administration (6)

Observed seat belt use



Source: National Highway Traffic Safety Administration (7)
Healthy People 2020 Target: 92.4% (dotted blue line) (5)

**Percentage of crash-related deaths that
involved alcohol-impaired drivers**



Source: National Highway Traffic Safety Administration (8)

Policy and Practice Solutions

This report focuses on policies recommended by the Community Preventive Services Task Force and the National Highway Traffic Safety Administration on the basis of scientific studies supporting the policies' effectiveness in preventing or reducing crash-related injuries and deaths. These policies include 1) implementing primary seat belt laws, 2) improving laws mandating the use of appropriate child passenger restraints (e.g., car seats and booster seats) to cover children through at least age 8 years, 3) using comprehensive graduated driver licensing systems, and 4) requiring the use of ignition interlock devices for all convicted driving-while-intoxicated (DWI) offenders (9–16). For information about why certain motor vehicle injury-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/motorvehicle/>).

Prevention Status Report | 2013

Motor Vehicle Injuries

Minnesota

Status of Policy and Practice Solutions in Minnesota

Seat belt law

As of August 1, 2013, Minnesota had a primary enforcement seat belt law for all seating positions (17).

Task Force on Community Preventive Services recommendation: Primary enforcement seat belt laws are recommended on the basis of strong evidence that they are substantially more effective than secondary enforcement laws at reducing motor vehicle-related injuries and deaths (10,11). Rates of seat belt use are an average of 9–14 percentage points higher in primary enforcement states than in secondary states (10,11,18,19).



Rating	State had
Green	A primary enforcement seat belt law covering all seating positions
Yellow	A primary enforcement seat belt law covering only the front seats
Red	A secondary enforcement seat belt law or no law

Child passenger restraint law

As of August 1, 2013, Minnesota required that all motor vehicle passengers aged 7 years or younger be in a car seat or booster seat (17).

Evidence shows that laws mandating the use of car seats and booster seats increase their use (12). Increasing the required age for car seat or booster seat use is an effective way to keep children protected. For example, among states that increased the required age to 7 or 8 years, car seat and booster seat use tripled (13).



Rating	State law covered
Green	Children through age 8 years
Yellow	Children through age 6 or 7 years only
Red	Children aged 5 years or younger only

Graduated driver licensing (GDL) system

As of August 1, 2013, Minnesota fulfilled the recommended passenger limit restriction but not the recommended nighttime driving restriction (20).

Research indicates that more comprehensive GDL systems prevent more crashes and save more lives compared with less comprehensive GDL systems. Based on this evidence, the following five components are recommended for more comprehensive GDL systems: 1) minimum age of 16 years for a learner's permit, 2) mandatory holding period of at least six months for a learner's permit, 3) restrictions against nighttime driving between 10:00 pm and 5:00 am (or longer), 4) limit of zero or one for the number of young passengers without adult supervision, and 5) minimum age of 18 years for full licensure (9,14,15).



Rating	State policy
Green	Required all five of the GDL components
Yellow	Required both nighttime driving and young passenger limits but not all five components
Red	Lacked either the nighttime driving or young passenger limits, or both

Ignition interlock law

As of August 1, 2013, Minnesota required ignition interlocks for all convicted DWI offenders (21).

Task Force on Community Preventive Services recommendation: Use of ignition interlocks is recommended for all people convicted of alcohol-impaired driving on the basis of strong evidence of interlocks' effectiveness in reducing re-arrest rates while the interlocks are installed (16).



Rating	State had
Green	A law requiring ignition interlocks for all convicted DWI offenders (i.e., offenders with blood alcohol concentrations [BAC] ≥ 0.08 g/dL, which includes both first-time and repeat offenders)
Yellow	A law requiring ignition interlocks for convicted repeat DWI offenders or first-time offenders with a particularly high BAC (e.g., BAC ≥ 0.15 g/dL)
Red	No law requiring ignition interlocks for any convicted DWI offenders

Prevention Status Report | 2013

Motor Vehicle Injuries

Minnesota

Simplified Rating System

A more detailed explanation of the rating system for motor vehicle injuries is available at <http://www.cdc.gov/stltpublichealth/psr/motorvehicle/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Indicator Definitions

Seat belt law: A primary enforcement seat belt law allows police to stop a vehicle solely because a driver or passenger is not wearing a seat belt. A secondary enforcement seat belt law requires police to have another reason for stopping a vehicle before citing a driver or passenger for not buckling up. The most comprehensive policies are primary seat belt laws that cover all occupants, regardless of where they are sitting. Some states have primary laws that cover only the front seat occupants.

Child passenger restraint law: A law that requires child passengers to travel in appropriate child passenger restraints, such as car seats or booster seats, until adult seat belts fit them properly. All 50 states and the District of Columbia have some form of child passenger restraint laws; however, the ages covered vary.

Graduated driver licensing (GDL) system: Policy that helps new drivers gain experience under low-risk conditions by granting driving privileges in stages. As teens move through GDL stages, they are given additional privileges, such as driving unsupervised or with a passenger.

Ignition interlock law: A law that mandates the use of ignition interlocks for drivers convicted of DWI. An ignition interlock is a device that analyzes a driver's breath and prevents the vehicle from starting if alcohol is detected.

References

1. CDC. WISQARS (Web-based Injury Statistics Query and Reporting System) [database]. Accessed Jun 5, 2013.
2. National Highway Traffic Safety Administration. Traffic Safety Facts, 2011 Data: Overview. Washington, DC: US Department of Transportation; 2013.
3. Naumann RB, Dellinger AM, Zaloshnja E, et al. Incidence and total lifetime costs of motor vehicle-related fatal and nonfatal injury by road user type, United States, 2005. *Traffic Injury Prevention* 2010;11:353–60.
4. National Highway Traffic Safety Administration. Fatality Analysis Reporting System [database]. US Department of Transportation, Washington, DC. Accessed Dec 7, 2012.
5. US Department of Health and Human Services. Injury and violence prevention. In: *Healthy People 2020*. Rockville, MD: US Department of Health and Human Services. Updated Oct 30, 2012.
6. National Highway Traffic Safety Administration. Traffic Safety Facts, 2010 Data: Young Drivers. Washington, DC: US Department of Transportation; 2012.
7. National Highway Traffic Safety Administration. Traffic Safety Facts, Crash, Stats: Seat Belt Use in 2011—Use Rates in the States and Territories. Washington, DC: US Department of Transportation; 2012.
8. National Highway Traffic Safety Administration. Traffic Safety Facts, 2010 Data: State Alcohol-Impaired Driving Estimates. Washington, DC: US Department of Transportation; 2012.
9. Goodwin A, Kirley B, Sandt L, et al. Countermeasures That Work: A Highway Safety Countermeasures Guide for State Highway Safety Offices. 7th edition. (Report No. DOT HS 811 727). Washington, DC: National Highway Traffic Safety Administration; 2013.
10. Shults RA, Nichols JL, Dinh-Zarr TB, et al. Effectiveness of primary enforcement safety belt laws and enhanced enforcement of safety belt laws: a summary of the Guide to Community Preventive Services systematic reviews. *Journal of Safety Research* 2004;35(2):189–96.
11. Shults RA, Elder RW, Sleet DA, et al. Primary enforcement of seat belt laws are effective even in the face of rising belt use rates. *Accident Analysis and Prevention* 2004;36(3):491–3.
12. Zaza S, Sleet DA, Thompson R, et al. Reviews of evidence regarding interventions to increase the use of child safety seats. *American Journal of Preventive Medicine* 2001;21(4S):31–47.
13. Eichelberger AH, Chouinard AO, Jermakian JS. Effects of booster seat laws on injury risk among children in crashes. *Traffic Injury Prevention* 2012;13:631–9.
14. Baker SP, Chen LH, Li G. National evaluation of graduated driver licensing programs. Washington, DC: US Department of Transportation; 2006.
15. Williams AF, Tefft BC, Grabowski JG. Graduated driver licensing research, 2010–present. *Journal of Safety Research* 2012;43(3):195–203.
16. Elder RW, Voas R, Beirness D, et al. Effectiveness of ignition interlocks for preventing alcohol-impaired driving and alcohol-related crashes. *American Journal of Preventive Medicine* 2011;40(3):362–76.
17. Insurance Institute for Highway Safety/Highway Loss Data Institute. Safety Belt and Child Restraint Laws. Arlington, VA: Insurance Institute for Highway Safety/Highway Loss Data Institute; 2013.
18. Beck LF, West BA. Vital signs: nonfatal, motor vehicle-occupant injuries (2009) and seat belt use (2008) among adults—United States. *MMWR* 2011;59:1681–6.
19. Shults RA, Beck LF. Self-reported seatbelt use, United States, 2002–2010: does prevalence vary by state and type of seatbelt law? *Journal of Safety Research* 2012;43(5–6):417–20.
20. Insurance Institute for Highway Safety/Highway Loss Data Institute. Young Driver Licensing Systems in the U.S. Arlington, VA: Insurance Institute for Highway Safety/Highway Loss Data Institute; 2013.
21. Insurance Institute for Highway Safety/Highway Loss Data Institute. DUI/DWI Laws. Arlington, VA: Insurance Institute for Highway Safety/Highway Loss Data Institute; 2013.

Prevention Status Report | 2013

Nutrition, Physical Activity, and Obesity

Minnesota

Public Health Problem



Poor diet and physical inactivity contribute to many serious and costly health conditions, including obesity, heart disease, diabetes, some cancers, unhealthy cholesterol levels, and high blood pressure (1,2).

Obesity is associated with increased blood pressure; unhealthy cholesterol levels; chronic diseases such as heart disease, diabetes, some cancers, and osteoarthritis; complications of pregnancy; and premature death (3).

Children who are not breastfed are at greater risk for various health problems, including childhood infections and obesity (4).




During 2009-2010, based on data from the National Health and Nutrition Examination Survey, approximately 17% of children and adolescents and 36% of adults were obese (5).




US medical costs associated with adult obesity were approximately \$147 billion in 2008 (6).

Percentage of adults who were obese



Source: Behavioral Risk Factor Surveillance System (7)


Percentage of high school students who were obese



Source: Youth Risk Behavior Surveillance System (8)

Note: Minnesota data were not available for one or more years from the source used for this chart. Similar data may be available from another national or state source.


Percentage of high school students who drank a can, bottle, or glass of soda or pop at least one time per day (excludes diet soda or pop)



Source: Youth Risk Behavior Surveillance System (8)

Note: Minnesota data were not available for one or more years from the source used for this chart. Similar data may be available from another national or state source.

Percentage of high school students who attended physical education classes on one or more days in an average week when they were in school



Source: Youth Risk Behavior Surveillance System (8)


Note: Minnesota data were not available for one or more years from the source used for this chart. Similar data may be available from another national or state source.

Prevention Status Report | 2013

Nutrition, Physical Activity, and Obesity

Minnesota

Percentage of low-income children aged 2 to <5 years who were obese



Source: Pediatric Nutrition Surveillance System (9)

Policy and Practice Solutions

This report focuses on policies and practices recommended by the Institute of Medicine, Community Preventive Services Task Force, US Surgeon General, CDC, and other expert bodies. The recommendations are based on expert judgment or evidence from scientific studies that the policies and practices can improve diet, increase breastfeeding, increase physical activity, or reduce obesity (10–17). These policies and practices include 1) implementing nutrition standards to limit the availability of less nutritious foods and beverages in schools, 2) implementing nutrition standards for foods and beverages in government facilities, 3) including nutrition and physical activity standards in state regulations of licensed childcare facilities, 4) establishing physical education time requirements in high schools, and 5) promoting evidence-based practices that support breastfeeding in hospitals and birth centers.

Additional strategies to prevent obesity and promote healthy eating, physical activity, and breastfeeding have been supported by scientific evidence or expert judgment (11–15,17). For information about why certain indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/npao/>).

Status of Policy and Practice Solutions in Minnesota

Secondary schools not selling less nutritious foods and beverages

In 2012, 31.8% of secondary schools in Minnesota did not sell the following items in vending machines or at school stores, canteens, or snack bars: candy, baked goods that are not low in fat, salty snacks that are not low in fat, soda pop, or fruit drinks that are not 100% juice (18).

In addition to providing school meals, many schools offer foods and beverages in other venues, such as school stores, canteens, snack bars, vending machines, and classrooms. The Institute of Medicine recommends nutrition standards for such foods and beverages (10), and CDC recommends that schools limit the availability of less nutritious foods and beverages and ensure that “only nutritious and appealing foods and beverages are provided in all food venues in schools . . .” (15).

Rating	Percentage of secondary schools that did not sell less nutritious foods and beverages in selected venues:
Green	≥ 66.6%
Yellow	50.0%–66.5%
Red	< 50.0%

Prevention Status Report | 2013

Nutrition, Physical Activity, and Obesity

Minnesota

State nutrition standards policy for foods and beverages sold or provided by state government agencies

In 2012, Minnesota did not have a nutrition standards policy for foods and beverages sold or provided by state government agencies (19).

The Institute of Medicine recommends that government agencies implement “strong nutrition standards for all foods and beverages sold or provided through the government” and ensure “that healthy options are available in all places frequented by the public” to reduce the availability of less healthful foods and beverages and increase the availability of more healthful options (11). For purposes of this report, strong policies are those that meet the following criteria: 1) apply to at least 90% of government agencies in the state executive branch; 2) cover all food purchased, contracted, distributed, or sold by government agencies in the state executive branch; 3) provide quantifiable standards for foods or nutrients (e.g., set a maximum for the amount of sodium a food item can include); and 4) set minimal standards that limit sodium content, fat content, and the availability of high-calorie, low-nutrient foods and beverages.



Rating	State nutrition standards policy
Green	Met all criteria
Yellow	Met some but not all criteria
Red	Did not exist

Inclusion of nutrition and physical activity standards in state regulations of licensed childcare facilities

In 2012, Minnesota state regulations for licensed childcare facilities included 10.6% of the 47 components of standards for infant feeding, nutrition, physical activity, and screen time (20).

The Institute of Medicine has recommended including specific requirements related to physical activity, sedentary activity, and child feeding in childcare regulations (12). The American Academy of Pediatrics, American Public Health Association, and National Resource Center for Health and Safety in Child Care and Early Education have identified 47 components that childcare regulatory agencies and childcare providers should include in standards for infant feeding, nutrition, physical activity, and screen time in licensed childcare settings (16).



Rating	Percentage of components included in state regulations:
Green	≥80.0%
Yellow	70.0%–79.9%
Red	<70.0%

State physical education time requirement for high school students

In 2012, Minnesota did not have a physical education time requirement for high school students (21).

The Community Preventive Services Task Force recommends the implementation of quality physical education programs that increase the length of, or activity levels in, school-based physical education classes (13). This recommendation is based on strong evidence of such programs’ effectiveness in improving physical activity levels and physical fitness among school-aged children and adolescents (13). CDC and the National Association for Sport and Physical Education recommend that high school students receive at least 225 minutes of physical education per week (15,17). States and the District of Columbia can help increase physical activity among high school students by setting minimum requirements for time spent in physical education.



Rating	State had
Green	A mandate for minutes per week that high school students must participate in physical education
Yellow	N/A
Red	No mandate for minutes per week that high school students must participate in physical education

Prevention Status Report | 2013

Nutrition, Physical Activity, and Obesity

Minnesota

Average birth facility score for breastfeeding support

In 2011, Minnesota had a birth facility score of 72 out of a possible 100 (22).

The US Surgeon General recommends that maternity care practices throughout the United States fully support breastfeeding (14). A review of evidence by the Cochrane Collaboration found that institutional changes in maternity care practices effectively increased breastfeeding initiation and duration rates (23). CDC's National Survey of Maternity Practices in Infant Nutrition and Care assesses and scores the extent to which hospitals and birth centers implement multiple evidence-based strategies that support breastfeeding (22).



Rating	State average birth facility score for breastfeeding support:
Green	≥80.0%
Yellow	70.0%–79.9%
Red	<70.0%

Simplified Rating System

A more detailed explanation of the rating system for nutrition, physical activity, and obesity is available at <http://www.cdc.gov/stltpublichealth/psr/npao/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations or is widely implemented.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations or is not as widely implemented as at the green rating level.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations or is not widely implemented.

Indicator Definitions

Secondary schools not selling less nutritious foods and beverages: Percentage of middle schools and high schools that did not allow students to purchase less nutritious foods and beverages from vending machines, school stores, canteens, and snack bars. For a school to be identified as not selling less nutritious foods and beverages, the school principal had to respond "no" to each item when asked whether students could purchase the following five items: 1) chocolate candy; 2) other kinds of candy; 3) salty snacks that are high in fat, such as regular potato chips; 4) cookies, crackers, cakes, pastries, or other baked goods that are high in fat; and 5) soda pop or fruit drinks that are not 100% juice. Data were provided for 45 states and the District of Columbia and represented only those states that participated in the survey and had an overall school response rate of at least 70% (18).

State nutrition standards policy for foods and beverages sold or provided by state government agencies: The presence of statewide nutrition standards for select foods or nutrients that cover foods and beverages purchased, contracted, distributed, or sold by government agencies in the state executive branch. Information was obtained using a search of the *Westlaw* database (19). State policies captured are statutes, regulations, and administrative guidance. Data were updated November 2012. The search results did not indicate whether a policy was implemented, only whether it existed.

Inclusion of nutrition and physical activity standards in state regulations of licensed childcare facilities: Inclusion of 47 recommended components of standards in regulations for infant feeding, nutrition, physical activity, and screen time in childcare settings (16). State regulations were considered to have included a component if the regulation fully met the requirements of the component across all childcare entities licensed by the state.

State physical education time requirement for high school students: A state mandate for minimum number of minutes per week that high school students must participate in physical education (21).

Average birth facility score for breastfeeding support: The state birth facility score for breastfeeding represents the average score across participating birth facilities in a state. Each participating birth facility, based on its response to a self-administered survey, was scored on multiple evidence-based practices that support breastfeeding across seven categories: 1) labor and delivery, 2) breastfeeding assistance, 3) mother-newborn contact, 4) newborn feeding practices, 5) breastfeeding support after discharge, 6) nurse/birth attendant breastfeeding training and education, and 7) structural and organizational factors related to breastfeeding (22). The total score can range from 0 to 100, with a higher score representing more support. The national average score across all states was 70.

Prevention Status Report | 2013

Nutrition, Physical Activity, and Obesity

Minnesota

References

1. US Department of Agriculture and US Department of Health and Human Services. Dietary Guidelines for Americans, 2010. 7th edition. Washington, DC: US Government Printing Office; 2010.
2. US Department of Health and Human Services. 2008 Physical Activity Guidelines for Americans. Washington, DC: US Department of Health and Human Services; 2008.
3. National Heart, Lung, and Blood Institute. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Bethesda, MD: National Institutes of Health; 1998.
4. Ip S, Chung M, Raman G, et al. Breastfeeding and Maternal and Infant Health Outcomes in Developed Countries. Evidence Report/Technology Assessment No. 153. AHRQ Publication No. 07-E007. Rockville, MD: Agency for Healthcare Research and Quality; 2007.
5. Ogden CL, Carroll MD, Kit BC, et al. Prevalence of obesity in the United States, 2009–2010. NCHS Data Brief 2012; 82:1–8.
6. Finkelstein EA, Trogdon JG, Cohen JW, et al. Annual medical spending attributable to obesity: payer-and service-specific estimates. Health Affairs (Millwood) 2009;28(5):w822–31.
7. CDC. Behavioral Risk Factor Surveillance System [database]. Accessed Aug 9, 2013.
8. CDC. Youth Risk Behavior Surveillance System [database]. Accessed Jun 13, 2013.
9. CDC. Pediatric Nutrition Surveillance System. Accessed Aug 9, 2013.
10. Institute of Medicine. Nutrition Standards for Foods in Schools: Leading the Way Toward Healthier Youth. Washington, DC: National Academies Press, 2007.
11. Institute of Medicine. Accelerating Progress in Obesity Prevention: Solving the Weight of the Nation. Washington, DC: National Academies Press; 2012.
12. Institute of Medicine. Early Childhood Obesity Prevention Policies. Washington, DC: National Academies Press; 2011.
13. Task Force on Community Preventive Services. Recommendations to increase physical activity in communities. American Journal of Preventive Medicine 2002;22(4S):67–72.
14. Office of the Surgeon General. The Surgeon General's Call to Action to Support Breastfeeding. Washington, DC: US Department of Health and Human Services; 2011.
15. CDC. School health guidelines to promote healthy eating and physical activity. MMWR 2011;60(RR–5).
16. American Academy of Pediatrics, American Public Health Association, National Resource Center for Health and Safety in Child Care and Early Education. Caring for Our Children: National Health and Safety Performance Standards; Guidelines for Early Care and Education Programs. 3rd edition. Elk Grove Village, IL: American Academy of Pediatrics; Washington, DC: American Public Health Association; 2011.
17. National Association for Sport and Physical Education. Physical Education is Critical to Educating the Whole Child. Reston, VA: National Association for Sport and Physical Education; 2011.
18. CDC. School Health Profiles 2012. Unpublished data.
19. CDC. Public Health Law Program. Unpublished analysis. November 2012.
20. National Resource Center for Health and Safety in Child Care and Early Education. Achieving a State of Healthy Weight: 2012. Aurora, CO: University of Colorado Denver; 2013.
21. National Association for Sport and Physical Education, American Heart Association. 2012 Shape of the Nation Report: Status of Physical Education in the USA. Reston, VA: American Alliance for Health, Physical Education, Recreation, and Dance; 2012.
22. CDC. National Survey of Maternity Practices in Infant Nutrition and Care (mPINC). Atlanta, GA: US Department of Health and Human Services; 2011.
23. Fairbank L, O'Meara S, Renfrew MJ, et al. A systematic review to evaluate the effectiveness of interventions to promote the initiation of breastfeeding. Health Technology Assessment 2000;4(25):1–171.

Prevention Status Report | 2013

Prescription Drug Overdose

Minnesota

Public Health Problem



Opioid pain relievers—also called prescription painkillers—such as oxycodone, hydrocodone, fentanyl, and hydromorphone are responsible for three-fourths of all prescription drug overdose deaths and caused more than 16,600 deaths in the United States in 2010 (1). Nationally, deaths involving opioids have more than quadrupled since 1999 (1). The drug overdose mortality rate is age adjusted and includes all drugs and all intents.




The sharp rise in opioid overdose deaths closely parallels an equally sharp increase in the prescribing of these drugs. Opioid pain reliever sales in the United States quadrupled from 1999 to 2010 (2). Similarly, the substance abuse treatment admission rate for opioid abuse in 2010 was seven times higher than in 1999 (3).

The severity of the epidemic varies widely across US states and regions. For example, the state with the highest drug overdose death rate has a rate more than eight times that of the state with the lowest rate. Minnesota's overdose death rate for 2010 (7.4 per 100,000 population) is below the national rate (12.4 per 100,000 population) (1).



In addition to the human costs, the epidemic of prescription drug overdose imposes a major financial toll. Nonmedical use of opioid pain relievers—use without a prescription or simply for the feeling or experience the drug causes—costs US insurance companies up to \$72.5 billion annually in healthcare expenditures (4). The epidemic also imposes substantial costs on state Medicaid programs. A 2009 Government Accountability Office report found that in 2006–2007, roughly 65,000 Medicaid beneficiaries in five states incurred over \$60 million in drug costs related to "doctor shopping" for controlled substance prescriptions (i.e., patients obtaining controlled substances from multiple healthcare practitioners without prescribers' knowledge of other prescriptions) (5).


Drug overdose death rate (age-adjusted per 100,000 population)



Source: National Vital Statistics System (1)


Note: These rates were adjusted using the direct method and the 2000 standard US population (6).

Prevalence of nonmedical use of prescription pain relievers in the past year among persons aged ≥12 years



Source: National Survey on Drug Use and Health (7)

Kilograms of morphine equivalents of opioid pain relievers sold (per 10,000 population)



Source: Automation of Reports and Consolidated Orders System (8)

Prevention Status Report | 2013

Prescription Drug Overdose

Minnesota

Policy and Practice Solutions

The United States is in the early stages of addressing the prescription drug overdose epidemic. CDC and other agencies are working to identify and evaluate interventions to reduce overdose deaths. This report focuses on policies and practices supported by emerging evidence, expert consensus, and/or extensive review of the primary drivers of the epidemic, including 1) implementing state pain clinic laws and 2) implementing prescription drug monitoring programs that follow best practices. For information about why certain prescription drug overdose-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/prescriptiondrug/>).

Status of Policy and Practice Solutions in Minnesota

State pain clinic law

As of July 2013, Minnesota had no pain clinic law (9).

Pain clinic laws hold promise for stopping the most egregious overprescribing practices (10). A pain clinic law is rated green in the PSR if the law requires state oversight and contains other requirements concerning ownership and operation of pain management clinics, facilities, or practice locations.



Rating	State had
Green	A pain clinic law meeting selected criteria
Yellow	N/A
Red	No pain clinic law

Prescription drug monitoring programs (PDMPs) following selected best practices

As of July 2013, Minnesota had an active PDMP that followed one or two selected best practices (11).

Prescription drug monitoring programs show early signs of changing providers' prescribing practices and can yield valuable information for healthcare providers and regulatory agencies. The selected best practices for PDMPs are 1) providing prescribers and dispensers access to PDMPs, 2) interoperability with the PDMP of at least one other state or the District of Columbia, and 3) proactively reporting findings to law enforcement and regulatory agencies (12).



Rating	State PDMP
Green	Followed all three selected best practices
Yellow	Followed one or two of the selected best practices
Red	Did not follow any of the selected best practices, was authorized but was not yet operating, or did not exist

Simplified Rating System

A more detailed explanation of the rating system for prescription drug overdose is available at <http://www.cdc.gov/stltpublichealth/psr/prescriptiondrug/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Prevention Status Report | 2013

Prescription Drug Overdose

Minnesota

Indicator Definitions

State pain clinic law: A law that requires state oversight of pain management clinics or describes specific registration, licensure, or ownership requirements for pain management clinics.

PDMP following selected best practices: A state prescription drug monitoring program that tracks the prescribing and dispensing of controlled substances and that follows selected best practices articulated by the Brandeis University PDMP Center of Excellence. These best practices include 1) providing prescribers and dispensers access to PDMPs, 2) interoperability with a PDMP of at least one other state or the District of Columbia, and 3) proactively reporting findings to law enforcement and regulatory agencies (12).

References


1. CDC. National Vital Statistics System [database]. Accessed Dec 10, 2012.
2. CDC. Vital Signs: overdoses of prescription opioid pain relievers—United States, 1999–2008. MMWR 2011;60:1487–92.
3. Substance Abuse and Mental Health Services Administration. Treatment Episode Data Set (TEDS): 2000–2010. National Admissions to Substance Abuse Treatment Services. DASIS Series S-61, HHS Publication No. (SMA) 12-4701. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2012.
4. Coalition Against Insurance Fraud. Prescription for Peril: How Insurance Fraud Finances Theft and Abuse of Addictive Prescription Drugs. Washington, DC: Coalition Against Insurance Fraud; 2007.
5. General Accounting Office. Medicaid: Fraud and Abuse Related to Controlled Substances Identified in Selected States. Washington, DC: General Accounting Office; 2009.
6. Klein RJ, Schoenborn CA. Age adjustment using the 2000 projected U.S. population. Healthy People Statistical Notes, No. 20. Hyattsville, MD: National Center for Health Statistics; 2001.
7. Substance Abuse and Mental Health Services Administration. National Survey on Drug Use and Health [database], 2009–2010, 2007–2008, and 2005–2007. Accessed Dec 10, 2012.
8. Drug Enforcement Administration. Automation of Reports and Consolidated Orders System (ARCOS) [database]. Accessed Dec 10, 2012.
9. CDC. Public Health Law Program. Unpublished data; July 2013.
10. Office of National Drug Control Policy. Epidemic: Responding to America's Prescription Drug Crisis. Washington, DC: Executive Office of the President of the United States; 2011.
11. PDMP Center of Excellence, Brandeis University. Unpublished data; July 2013.
12. PDMP Center of Excellence. Prescription Drug Monitoring Programs: An Assessment of the Evidence for Best Practices. Waltham, MA: Brandeis University; 2012.


Prevention Status Report | 2013

Teen Pregnancy


Minnesota

Public Health Problem


 Each year in the United States, about 750,000 women under age 20 become pregnant (1). In 2011 in Minnesota, 3,464 teens aged 15–19 years gave birth (2).

 In 2011, young women of color—particularly Hispanic and African-American females aged 15–19 years—were disproportionately likely to give birth, with national birth rates of 49.6 and 47.3 per 1,000 population, respectively (3).

Teen mothers are more likely to experience negative social outcomes, including lower rates of school completion and reduced earnings, than teens who do not have children. The children of teenaged mothers are more likely to achieve less in school, experience abuse or neglect, have more health problems, be incarcerated at some time during adolescence, and give birth as a teenager (4).


 The annual costs of teen childbearing in 2008 were \$10.9 billion in the United States and \$173 million in Minnesota (5).

**Birth rate among females aged 15–19 years
(per 1,000 population)**



Source: National Vital Statistics System—Births (6)


**Proportion of high school students
who ever had sexual intercourse**



Source: Youth Risk Behavior Surveillance System (7)

Note: Minnesota data were not available for one or more years from the source used for this graph. Similar data may be available from another national or state source.


**Proportion of currently sexually active
female high school students who used birth
control pills, any injectable birth control, any
birth control ring or implant, or intrauterine
device before last sexual intercourse**



Source: Youth Risk Behavior Surveillance System (7)

Note: Minnesota data were not available for one or more years from the source used for this chart. Similar data may be available from another national or state source.

**Proportion of currently sexually active high
school students who used a condom during
last sexual intercourse**



Source: Youth Risk Behavior Surveillance System (7)

Note: Minnesota data were not available for one or more years from the source used for this graph. Similar data may be available from another national or state source.

Prevention Status Report | 2013

Teen Pregnancy

Minnesota

Policy and Practice Solutions

This report focuses on expanding eligibility for Medicaid family planning services to the income eligibility level for pregnancy-related services and to include women younger than age 18 years, either by amending the Medicaid waiver or by converting to the State Plan Amendment available through the Centers for Medicare and Medicaid Services, or by expanding the full Medicaid program (8–12). This policy is consistent with the US Department of Health and Human Services' National Prevention Strategy recommendations to expand access to contraceptive services and with a *Healthy People 2020* objective to "increase the number of states that set the income eligibility level for Medicaid-covered family planning services to at least the same level used to determine eligibility for Medicaid-covered, pregnancy-related care" (13,14).

Other strategies supported by scientific evidence include providing comprehensive sexual health education for adolescents, using positive youth development approaches, and improving parent-child communication and parental monitoring of youth behavior (15–17). For information about why Medicaid family planning expansion was selected as an indicator, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/teenpregnancy/>).

Status of Policy and Practice Solutions in Minnesota

Expansion of state Medicaid family planning eligibility

As of August 2013, Minnesota had expanded Medicaid coverage of family planning services to include teens and adults with incomes up to 200% of the federal poverty level (FPL). The expansion did not include all teens and adults with incomes up to 275% of the FPL, the state's income level for pregnancy-related Medicaid coverage (18,19).

Healthy People 2020 target: Increase the number of states that set the income eligibility level for Medicaid coverage of family planning services to at least the same level used to determine eligibility for Medicaid coverage of pregnancy-related care (14,18,19).



Rating	State Medicaid family planning eligibility
Green	Was income-based, met the income eligibility level for pregnancy-related care, and covered all women, including teens
Yellow	Was limited, was not income-based, did not meet the eligibility level for pregnancy-related services, and/or excluded some teens
Red	Had not been expanded

Simplified Rating System

A more detailed explanation of the rating system for teen pregnancy is available at <http://www.cdc.gov/stltpublichealth/psr/teenpregnancy/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Prevention Status Report | 2013

Teen Pregnancy

Minnesota

Indicator Definitions

Expansion of state Medicaid family planning eligibility (waiver or state plan amendment): State expansion of eligibility for Medicaid coverage of family planning services to include teens under age 18 and to be set at the eligibility level for pregnancy care (this level varies by state and the District of Columbia). This expansion is achieved by 1) securing approval (officially known as a “waiver” of federal policy) from the Centers for Medicare and Medicaid Services, 2) amending the state Medicaid plan with a State Plan Amendment (i.e., a permanent change to the state’s Medicaid program), or 3) expanding the full state Medicaid program.

References

1. Ventura SJ, Curtin SC, Abma JC, et al. Estimated pregnancy rates and rates of pregnancy outcomes for the United States, 1990–2008. *National Vital Statistics Reports*: 2012;60(7).
2. CDC. National Vital Statistics System [database]. Accessed Aug 5, 2013.
3. Martin JA, Hamilton BE, Ventura SJ, et al. Births: final data for 2011. *National Vital Statistics Report* 2013;62(1).
4. Hoffman S, Maynard R, eds. *Kids Having Kids: Economic Costs and Social Consequences of Teen Pregnancy*. Washington, DC: The Urban Institute Press; 2008.
5. National Campaign to Prevent Teen and Unplanned Pregnancy. *Counting It Up: The Public Costs of Teen Childbearing*. Washington, DC: National Campaign to Prevent Teen and Unplanned Pregnancy; Jun 2011.
6. CDC. VitalStats [database]. Accessed Aug 5, 2013.
7. CDC. Youth Risk Behavior Surveillance System [database]. Accessed Aug 5, 2013.
8. Foster DG, Biggs MA, Rostovtseva D, et al. Estimating the fertility effect of expansions of publicly funded family planning services in California. *Women’s Health Issues* 2011;21:418–24.
9. Yang Z, Gaydos LM. Reasons for and challenges of recent increases in teen birth rates: a study of family planning service policies and demographic changes at the state level. *Journal of Adolescent Health* 2010;46:517–24.
10. Kearney MS, Levine PB. Subsidized contraception, fertility, and sexual behavior. *The Review of Economics and Statistics* 2009;91(1):137–51.
11. Lindrooth RC, McCullough JS. The effect of Medicaid family planning expansions on unplanned births. *Women’s Health Issues* 2007;17:66–74.
12. Edwards J, Bronstein J, Adams K. *Evaluation of Medicaid Family Planning Demonstrations*. The CNA Corporation, CMS Contract No 752-2-415921:22; Nov 2003.
13. US Department of Health and Human Services. *National Prevention Strategy: America’s Plan for Better Health and Wellness*. Rockville, MD: US Department of Health and Human Services; 2011.
14. US Department of Health and Human Services. *Family Planning*. In: *Healthy People 2020*. Updated Sep 6, 2012.
15. US Department of Health and Human Services. *Teen Pregnancy Prevention: Evidence-Based Programs* [database]. Accessed Dec 10, 2012.
16. Community Preventive Services Task Force. *Prevention of HIV/AIDS, other STIs and pregnancy: interventions for adolescents*. In: *Guide to Community Preventive Services*. Updated Nov 30, 2010.
17. Oringanje C, Meremikwu MM, Eko H, et al. Interventions for preventing unintended pregnancies among adolescents. *Cochrane Database of Systematic Reviews* 2009;4:CD005215.
18. Guttmacher Institute. *State Policies in Brief (as of August 2013): Medicaid Family Planning Eligibility Expansions*. New York, NY: Guttmacher Institute; 2013.
19. Kaiser Family Foundation. *Adult Income Eligibility Limits at Application as a Percent of the FPL by Coverage Authority Parents (Table 4)*. In: *Performing Under Pressure: Annual Findings of a 50-State Survey of Eligibility, Enrollment, Renewal, and Cost-Sharing Policies in Medicaid and CHIP, 2011–2012*. Washington, DC: Kaiser Family Foundation; 2012.

Prevention Status Report | 2013

Tobacco Use

Minnesota

Public Health Problem



Tobacco use is the leading cause of preventable death in Minnesota and the United States overall. Smoking harms nearly every organ in the body and causes cancer, heart disease, stroke, respiratory illness, and many other health problems (1).




During 2007–08, in the United States, 37% of adult nonsmokers and 54% of children aged 3–11 years were exposed to secondhand smoke (2).




Smoking and exposure to secondhand smoke result in \$96 billion in medical expenditures and \$97 billion in lost productivity annually in the United States. In Minnesota, smoking causes \$1.93 billion in personal healthcare expenditures and \$1.28 billion in lost productivity annually (3).

Proportion of adults who smoke cigarettes



Sources: Behavioral Risk Factor Surveillance System (4), National Health Interview Survey (5)
Healthy People 2020 target: 12.0% (dotted blue line) (6)

Proportion of high school students who smoke cigarettes



Source: Youth Risk Behavior Surveillance System (7)
Healthy People 2020 target: 16.0% (dotted blue line) (6)

Note: Minnesota data were not available for one or more years from the source used for this graph. Similar data may be available from another national or state source.

Policy and Practice Solutions

This report focuses on policies and practices recommended by the Institute of Medicine, World Health Organization, Community Preventive Services Task Force, US Surgeon General, and Centers for Disease Control and Prevention on the basis of scientific studies supporting the policies' effectiveness in preventing or reducing tobacco use (8–11,13,14). These policies and practices include 1) increasing state cigarette excise taxes, 2) establishing statewide smoke-free policies, and 3) sustaining tobacco control program funding. Other strategies also supported by scientific evidence include hard-hitting media campaigns and systemic changes to increase access to and use of cessation services. For information about why certain tobacco-related indicators were selected, and for links to additional data and resources, visit the CDC website (<http://www.cdc.gov/stltpublichealth/psr/tobacco/>).

Prevention Status Report | 2013

Tobacco Use

Minnesota

Status of Policy and Practice Solutions in Minnesota

State cigarette excise tax

As of June 30, 2013, Minnesota's cigarette excise tax was \$1.23 per pack, compared with the highest state tax of \$4.35 (range = \$0.17–\$4.35) (15).

Healthy People 2020 target: An increased excise tax in all states and the District of Columbia by \$1.50 per pack by the year 2020 (6). This increase would generate millions of dollars in revenue annually, prevent more children from starting to smoke, help smokers quit, save lives, and save millions in long-term healthcare costs (16,17).



Rating	State excise tax was
Green	\$2.00 per pack or above
Yellow	\$1.00–\$1.99 per pack
Red	Less than \$1.00 per pack

Comprehensive state smoke-free policy

As of June 30, 2013, Minnesota had a statewide smoke-free policy covering workplaces, restaurants, and bars (15).

Healthy People 2020 target: A statewide ban on smoking in public places and worksites in all states and the District of Columbia (6). Studies have shown that smoke-free policies reduce secondhand smoke exposure, help smokers quit, and reduce heart attack and asthma hospitalizations (10,11,17–21).



Rating	State smoke-free policy covered
Green	Workplaces, restaurants, and bars
Yellow	Two of the three locations
Red	One or none of the locations

Funding for tobacco control

As of fiscal year 2010, Minnesota allocated 36.8% of the CDC-recommended funding for tobacco control (\$21.5 million of \$58.4 million) (22).

CDC recommendation: Tobacco control funding at 100% of CDC's recommended annual investment in all states and the District of Columbia (14). States that have made larger investments in comprehensive tobacco control programs have seen cigarette sales drop more than twice as much as sales in the United States as a whole, and smoking prevalence among adults and youth has declined faster as spending for tobacco control programs has increased (14,23,24).



Rating	Funding level was at
Green	100% or more of CDC recommendation
Yellow	50.0%–99.9% of CDC recommendation
Red	Less than 50% of CDC recommendation

Simplified Rating System

A more detailed explanation of the rating system for tobacco use is available at <http://www.cdc.gov/stltpublichealth/psr/tobacco/>.

Green

The policy or practice is established in accordance with supporting evidence and/or expert recommendations.

Yellow

The policy or practice is established in partial accordance with supporting evidence and/or expert recommendations.

Red

The policy or practice is either absent or not established in accordance with supporting evidence and/or expert recommendations.

Prevention Status Report | 2013

Tobacco Use

Minnesota

Indicator Definitions

State cigarette excise tax: The amount of state excise tax, in dollars, on a pack of 20 cigarettes.

Comprehensive state smoke-free policy: A state law that prohibits smoking in all indoor areas of private workplaces, restaurants, and bars, with no exceptions (25).

Funding for tobacco control: The amount of funding allocated for state tobacco control activities, including state and federal dollars. Note: Data provided for fiscal year 2010 funding do not include nongovernmental funding sources or federal funds from the American Recovery and Reinvestment Act Prevention Wellness Initiative announced in March 2010. Additionally, the amount allocated per fiscal year does not always match the amount spent during the year.

References

1. US Surgeon General. How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General. Rockville, MD: US Department of Health and Human Services; 2010.
2. CDC. Vital signs: nonsmokers' exposure to secondhand smoke—United States, 1999–2008. MMWR 2010;59(35).
3. Smoking—Attributable Mortality, Morbidity, and Economic Costs (SAMMEC) [database]. Accessed Dec 10, 2012.
4. CDC. Behavioral Risk Factor Surveillance System [database]. Accessed Jun 13, 2013.
5. Schiller JS, Lucas JW, Peregoy JA. Summary health statistics for U.S. adults: National Health Interview Survey, 2011. Vital Health Statistics 2012;10(256).
6. US Department of Health and Human Services. Tobacco use across the life stages. In: Healthy People 2020. Rockville, MD: US Department of Health and Human Services; Updated Nov 20, 2012.
7. CDC. Youth Risk Behavior Surveillance System [database]. Accessed Jun 13, 2013.
8. Institute of Medicine. Ending the Tobacco Problem: A Blueprint for the Nation. Washington, DC: National Academies Press; 2007.
9. World Health Organization. WHO Report on the Global Tobacco Epidemic, 2008—The MPOWER Package. Geneva, Switzerland: World Health Organization; 2008.
10. The Task Force on Community Preventive Services. The Guide to Community Preventive Services: What Works to Promote Health? New York, NY: Oxford University Press; 2005.
11. CDC. The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services; 2006.
12. CDC. Preventing Tobacco Use among Youth and Young Adults: A Report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services; 2012.
13. CDC. Reducing Tobacco Use: A Report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services; 2000.
14. CDC. Best Practices for Comprehensive Tobacco Control Programs—2007. Atlanta, GA: US Department of Health and Human Services; 2007.
15. CDC. State Tobacco Activities Tracking & Evaluation (STATE) System [database]. Accessed Dec 10, 2012.
16. Congressional Budget Office. Raising the Excise Tax on Cigarettes: Effects on Health and the Federal Budget. Washington, DC: Congressional Budget Office; 2012.
17. Hopkins DP, Razi S, Leeks KD, et al. Smoke-free policies to reduce tobacco use: a systematic review. American Journal of Preventive Medicine 2010;38(2S):275–89.
18. Hahn EJ. Smokefree legislation: a review of health and economic outcomes research. American Journal of Preventive Medicine 2010;39(6 Suppl 1):S66–S76.
19. Institute of Medicine. Secondhand Smoke Exposure and Cardiovascular Effects: Making Sense of the Evidence. Washington, DC: National Academies Press; 2010.
20. Millett C, Lee JT, Lavery AA, et al. Hospital admissions for childhood asthma after smoke-free legislation in England. Pediatrics 2013;131(2):e495–e501.
21. Herman PM, Walsh ME. Hospital admissions for acute myocardial infarction, angina, stroke, and asthma after implementation of Arizona's comprehensive statewide smoking ban. American Journal of Public Health 2011;101:491–6.
22. CDC. State tobacco revenues compared with tobacco control appropriations—United States, 1998–2010. MMWR 2012;61:370–4.
23. Farrelly MC, Pechacek TP, Chaloupka FJ. The impact of tobacco control program expenditures on aggregate cigarette sales: 1981–2000. Journal of Health Economics 2003;22(5):843–59.
24. Tauras JA, Chaloupka FJ, Farrelly MC, et al. State tobacco control spending and youth smoking. American Journal of Public Health 2005;95:338–44.
25. CDC. State smoke-free laws for worksites, restaurants, and bars—United States, 2000–2010. MMWR 2011;60:472–5.