



DEPARTMENT OF HEALTH & HUMAN SERVICES

Public Health Service

Centers for Disease Control

June 1, 1987

From: B.R. Miller

Subject: Attached "Arthropod-Borne Virus Information Exchange"

Your copy of the most recent "Arthropod-Borne Virus Information Exchange" is attached.

The next deadline for submission of contributions is March 1, 1988.

Please address all communications to the undersigned.

Barry R. Miller, Ph.D.
Division of Vector-Borne Viral Diseases
Center for Infectious Disease
Centers for Disease Control
Post Office Box 2087
Fort Collins, Colorado 80522, U.S.A.

attachment

TABLE OF CONTENTS

	Page
Guide for contributors.....	i
The American Committee on Arthropod-Borne Viruses:	
Subcommittee on Information Exchange.....	1
Arbovirus Reports from:	
WHO Collaborating Centre for Reference and Research, Dept. of Medical Microbiology, University of Malaya, Kuala Lumpur, Malaysia.....	117
Department of Virology, College of Medicine, University of Ibadan, Ibadan, Nigeria.....	118
Departamento De Patologia Tropical, Centro de Investigaciones Re- gionales "Dr. Hideyo Noguchi", Universidad Autonoma de Yucatan, Mexico, and San Juan Laboratories, Division of Vector-Borne Viral Diseases, Centers for Infectious Diseases, Centers for Disease Control, San Juan, Puerto Rico.....	120
Instituto de Medicina Tropical "Pedro Kouri" La Habana, Cuba.....	122
Pasteur Institute, Bangui, Central African Republic.....	133
NERC Institute of Virology, Mansfield Road, Oxford, UK.....	135
State of New Jersey, Dept. of Health, Virology Program, Trenton, New Jersey.....	137
WHO Collaborating Center for Arbovirus Reference and Research, Institute of Virology, Bratislava, Czechoslovakia.....	139
Arbovirus Unit, National Institute for Virology, Sandringham 2131, Republic of South Africa.....	144
Virus Laboratory, Faculty of Medicine, BP 815, 29285 Brest Cedex, France.....	149
California Department of Health Services, Viral and Rickettsial Disease Laboratory, Berkeley, California, USA.....	152
Department of Epidemiology, School of Public Health, University of Sao Paulo, Brazil.....	153
Special Pathogens Unit, National Institute for Virology, Sandringham, 2131, Republic of South Africa.....	154
Medical Research Council of New Zealand's Viral Research Unit, University of Otago, Dunedin, New Zealand.....	158
Microbiology Section, CSIRO, Division of Tropical Animal Science, Long Pocket Laboratories, Indooroopilly, Brisbane, Australia... Arbovirus Laboratory, Institut Pasteur, 97306 Cayenne Cedex, French Guiana.....	159 168

Important Notice: This exchange is issued for the sole purpose of timely exchange of information among investigators of arthropod-borne viruses. It contains reports, summaries, observations, and comments submitted voluntarily by qualified agencies and investigators. The appearance of any information, data, opinions, or views in this exchange does not constitute formal publication. Any reference to or quotation of any part of this exchange must be authorized directly by the person or agency which submitted the text.

GUIDE FOR CONTRIBUTORS

The Arthropod-Borne Virus Information Exchange is issued for the purpose of timely exchange of information among investigators of arthropod-borne viruses. It contains reports, summaries, observations, and comments submitted voluntarily by qualified investigators. The appearance of any information, data, opinions, or views in this exchange does not constitute formal publication. Any reference to or quotation of any part of the "Information Exchange" must be authorized directly by person or agency submitting the article. The editor of the "Information Exchange" cannot authorize references and quotations.

Deadlines for contributions are March 1 and September 1.

1. Heading

The heading should be typed with capital letters, including name of laboratory and address. For example:

REPORT FROM THE BIOLOGICAL PRODUCTS PRODUCTION BRANCH, CENTER FOR INFECTIOUS DISEASES, CENTERS FOR DISEASE CONTROL, ATLANTA, GA 30333

2. Body of Report

The text of the report should be as brief as possible to convey the intended message and should make reference to tables and figures included in the report. The text should be single spaced with double spacing between paragraphs.

3. Authors' Names

The names of authors should be in parentheses following the text.

4. Tables and Figures

Tables and figures should be numbered and titled if appropriate. Tables and figures should not be submitted without some description or explanation.

5. Size of Pages

Since there are specific space limitations, the typed material on each page should not exceed 7-1/8" x 9-1/4". The same dimensions apply to tables and figures. If tables and figures are larger than these dimensions, they have to be reduced before being printed. The block shown on this page represents the maximum space available for each page of your report.

Reports should be typed only on one side of each page since they have to be photographed for reproduction. Each page should be numbered. Only the original typed report should be submitted.



The AMERICAN COMMITTEE ON ARTHROPOD-BORNE VIRUSES

1986 ANNUAL REPORT ON THE CATALOGUE OF ARTHROPOD-BORNE
AND SELECTED VERTEBRATE VIRUSES OF THE WORLD*

By
THE SUBCOMMITTEE ON ARTHROPOD-BORNE VIRUS
INFORMATION EXCHANGE

SUBCOMMITTEE ON INFORMATION EXCHANGE

I. Objectives:

The objectives of the Catalogue are to register data concerning occurrence and characteristics of newly recognized arthropod-borne viruses and other viruses of vertebrates of demonstrated or potential zoonotic importance and to disseminate this information at quarterly intervals to participating scientists in all parts of the world; to collect, reproduce, collate, and distribute current information regarding registered viruses from published materials, laboratory reports and personal communications; and to prepare and distribute an annual summary of data extracted from catalogued virus registrations.

II. Materials and Methods:

Viruses are registered and information supplied on a voluntary basis, usually by scientists responsible for their isolation and identification. New registration cards, information concerning registered viruses and pertinent abstracts of published literature are distributed at quarterly intervals to participating laboratories. Abstracts of published articles dealing with catalogued viruses are reproduced by special arrangements with the editors of Biological Abstracts, Abstracts on Hygiene, and the Tropical Disease Bulletin.

Distribution of Catalogue Materials: At the start of 1986, 181 mailings of Catalogue material were being made. During the year, two addresses were dropped and nine new participants were added to the mailing list. At the end of the year, 188 mailings of Catalogue material were being made, including 59 within the U.S.A. and 129 to foreign addresses. Distribution by continent was: Africa 19, Asia 25, Australasia 9, Europe 45, North America 72, and South America 18.

Abstracts and Current Information: A total of 422 abstracts or references were coded by subject matter and distributed to participants during 1986. Of this total, 208 were obtained from Kelbry Enterprises and 214 from Abstracts on Hygiene and the Tropical Diseases Bulletin. A total of 16,099 references or units of information have been issued since the start of the program.

*The Catalogue is supported by the Centers for Disease Control, Atlanta, Georgia.

Note: This report is not a publication and should not be used as a reference source in published bibliographies.

Registration of new viruses. Two viruses were accepted for registration during the period January 1986 to December 1986. As of December 1985, the Catalogue contained 504 viruses. With the acceptance of 2 virus registrations, the total number of registered viruses is 506 as of December 1986. The 2 viruses registered between January 1986 and December 1986 are listed below.

<u>Virus Name</u>	<u>Recommended Abbreviation</u>	<u>Country</u>	<u>Source</u>	<u>Antigenic Group</u>
Corfou	CFU	Greece	<u>Phlebotomus</u> sp.	PHL
Omo	OMO	Ethiopia	Rodent	QYB

Omo virus was isolated in 1971, while Corfou virus was isolated in 1981. Both viruses have been evaluated as Possible Arbovirus by SEAS.

Neither virus has been implicated in human infections.

Antigenic grouping. There have been a number of changes during the past year in the antigenic classification of registered viruses. Two new serogroups have been formed. One serogroup includes Umatilla and Llano Seco viruses and an unregistered arbovirus named Netivot virus (1). The latter virus was isolated from mosquitoes collected in Israel. An antigenic relationship between Umatilla and Llano Seco virus was observed previously when Llano Seco virus was being characterized (see Llano Seco registration card). However, the two viruses were not grouped because there was some uncertainty concerning the antigenic relationship of Llano Seco virus to other established arboviruses. The formation of this serogroup is provisional until it is resolved that these three viruses are distinct by cross-neutralization tests. Nevertheless, all three viruses show biological differences and differ in the migration of their ds RNA segments in polyacrylamide gels (1).

A newly described virus, causing disease in humans, has been isolated from bats (Rousettus aegyptiacus) collected in Uganda and has been shown to be antigenically related to Yogue virus (2). Two virus isolates were obtained from bats and four additional isolates from humans infected in the laboratory. This new virus has been named Kasokero virus and, together with Yogue virus, they now comprise the Yogue serogroup.

There are now 65 serogroups represented among viruses registered in the Catalogue, excluding viruses placed in the Bunyamwera Supergroup but unassigned (SBU).

Zingilamo virus was shown to possess a hemagglutinin and subsequently was shown to be related to Semliki Forest virus (3,4). Provisionally, Zingilamo virus has been placed in serogroup A pending the determination of its precise relationship to Semliki Forest virus.

Keterah and Issyk-Kul viruses have been demonstrated to be identical by complement-fixation (CF) and neutralization tests (NT) (5).

Taxonomic status of registered viruses. Reported changes in the taxonomic classification of registered arboviruses are of a provisional nature, and in some instances, new taxonomic placements are based on very slight evidence.

The taxonomic status of seven registered viruses has changed as a result of observations made during 1986. Boteke, Gossas, Nkolbisson, and Rochambeau viruses have been found to possess rhabdovirus morphology (6). Studies have been undertaken in order to determine if these viruses are antigenically related to other known viruses. Boteke virus originally was considered to be related to Zingilamo virus. This is no longer the case. Zingilamo virus recently was shown to possess alphavirus morphology and to be related to Semliki Forest virus (3). In addition, the morphology of Malakal virus was found to resemble that of viruses of the family Togaviridae (7). Aside from its known relationship to Puchong virus, an antigenic relationship to other known viruses has not been found. Finally, Wanowrie virus has been shown to morphologically resemble viruses of the family Bunyaviridae (6). In addition, a hemagglutinin has been elaborated for Wanowrie virus, and the characteristics of this hemagglutinin have been described (8).

Both Issyk-Kul and Keterah viruses have been placed in the family Bunyaviridae taxon. Electron microscopic studies have shown that Issyk-Kul virus resembles viruses of the family Bunyaviridae (9). Since CF and NT studies have shown that both Issyk-Kul and Keterah are antigenically identical (5), they should be considered as a single virus.

SYNOPSIS OF INFORMATION IN THE CATALOGUE

This synopsis has been compiled primarily to provide a short review of the viruses included in the Catalogue. The following tabulations are designed to draw together groups of viruses showing certain common characteristics including taxonomic status, serological relationships, and, where appropriate, principal arthropod vector. Isolations from arthropod and animal hosts, continental distribution, involvement in human disease, and arthropod-borne status are indicated.

The recommended levels of laboratory practice and containment and the basis for assignment to these levels are shown. Most of this information was published previously by SALS (10). Several registered viruses listed in Tables 5.1 through 13.2.2 have not been rated by SALS. Appendices I and II, following Table 18.1, provide a description of recommended levels and an explanation of symbols used to define basis.

Other tables summarize the taxonomic status of registered viruses; the antigenic groups comprising a given taxon to which registered viruses have been assigned; the numbers of registered viruses assigned to presently recognized antigenic groups; chronology and areas of isolations of registered viruses; continental distribution by groups; numbers of viruses recovered from naturally infected arthropods and vertebrates; association with human disease; and evaluation of arthropod-borne status of members in various serogroups.

Appendices I and II are followed by a vector index and a host index. The vector index mostly shows registered viruses isolated from individual arthropod species collected in nature. The host index shows registered viruses isolated from vertebrate hosts collected in nature. Both Linnean taxonomic designations and common names of hosts are used depending upon information available from original sources. These indices were compiled primarily from information on virus registration cards. Other sources of information were employed as well, although these listings should not be considered as exhaustive. Because of the large number of virus names involved, official abbreviations for registered viruses were used. Please refer to Table 1.1 for the corresponding virus name.

1. Alphabetical and taxonomic listing of registered viruses. Table 1.1 presents a listing of the 506 viruses registered in the Catalogue as of December 1985. An official or provisional taxonomic classification is shown for each registered virus. If taxonomic status is not indicated, the registered virus is presently unclassified. Also, for each virus a recommended abbreviation is given, formulated according to the guidelines established by the ACAV (11). All too often, abbreviations of the author's choosing are employed in publications and do not conform to the recommended abbreviations. The use of unofficial abbreviations is confusing, is contrary to established guidelines, and erodes a portion of the effort of the Arbovirus Information Exchange program. All arbovirologists who plan to employ abbreviations in print should make every effort to use the recommended abbreviations.

Antigenic groups to which viruses have been assigned also are shown in Table 1.1. If no antigenic group is given, the virus is ungrouped and indicates that it has not been demonstrated to be related serologically to any other virus.

2. Antigenic groups of registered viruses. The originally described antigenic groups of arboviruses were designated by letters, A, B, and C; but in present practice, the first discovered virus of a newly recognized serogroup lends its name to the antigenic cluster. Before a virus can be assigned to any antigenic group, it must be shown to be serologically related to, but clearly distinguishable from a previously isolated virus.

Table 2.1 lists the serogroups comprising the various taxa to which registered viruses have been assigned. Sixty-five antigenic groups have been designated for viruses registered in the Catalogue, including the previously established rabies serogroup (12). The rabies serogroup is represented in the Catalogue because two members of that serogroup were registered in the Catalogue. Lagos bat virus was registered in 1961 and was described in the first published edition of the Catalogue (13). Kotonkan virus was registered more recently (in 1982); and had greater potential of being arthropod-borne since it was isolated from Culicoides insects. There are several instances in which only a single virus is shown in an antigenic group. That is so because one or more antigenic relatives of that virus are known but have not yet been registered.

The Bunyavirus genus comprises the Bunyamwera Supergroup to which several additional serogroups have been added. The most recent additions are the Anopheles B and Turlock serogroups (14). The Bunyamwera Supergroup originally was formulated to reflect low-level but reproducible intergroup relationships, usually by CF and/or HI reactions (15). A large number of these viruses were subsequently found to be identical morphologically and morphogenetically. The Bunyamwera Supergroup designation was thus replaced by the Bunyavirus genus in the Family Bunyaviridae (16,17). In a somewhat analogous situation, the Nairovirus genus was constructed to include six distinct serogroups which share low-level intergroup relationships among themselves (18,19). Registered viruses belonging in the Bunyavirus genus constitute slightly more than 25% of all registered viruses.

3. Initial isolations by decade and country of origin. Table 3.1 lists the initial isolation of registered viruses by the decade of discovery and according to the continent or zoogeographic region and country in which each was discovered. Because of the large number of virus names involved, abbreviations are employed. These abbreviations and the associated complete names of the respective viruses may be found in Table 1.1.

4. Initial isolation of viruses by continent, country, and chronological period. Periods or locations which show large numbers of virus isolations undoubtedly reflect the net effect of a number of contributing factors such as the change in emphasis of field programs from a search for viruses causing specific diseases to a systematic search for viruses, new or known, in their

natural ecological niche in a given geographical area; refinements in isolation and identification techniques; improved communication between arbovirus laboratories; more rapid dissemination of new information; or the presence in a given area of an arbovirus laboratory with highly active and effective field programs.

Tables 5.1 through 18.1 list registered viruses by serogroup; by recorded isolations from arthropod vectors and vertebrates; and by geographic distribution based on virus isolation. Data also are presented regarding human disease in nature or by laboratory infection, evaluation of arbovirus status, and proved or provisional taxonomic status. These tables also show the recommended laboratory practice and containment level assigned to each registered virus, and the basis for assignment to a level. Where possible, sets of viruses also were grouped according to their actual or suspected principal arthropod vector.

5. Alphaviruses. Alphaviruses clearly are mosquito associated, although a few have been isolated from other arthropods (Table 5.1). In addition, during 1985, Middelburg virus was isolated from Amblyomma variegatum ticks. About 50% of the alphaviruses are associated with avian hosts, whereas some, particularly those of the Venezuelan equine encephalitis complex, are associated with rodents. Sindbis virus has been recovered from the organs of insectivorous bats collected in Zimbabwe. Cabassou, chikungunya, eastern equine encephalitis, Highlands J, and Venezuelan equine encephalitis viruses represent the other alphaviruses which have been isolated from bats.

Eleven alphaviruses have been isolated from humans while 12 have been implicated in human disease either by infections acquired in nature or in the laboratory. Both Bebaru and Ross River viruses have been associated with clinically inapparent laboratory-acquired infections, although Ross River virus infections in nature have resulted in human disease. At least eight of these 13 alphaviruses have been responsible for epidemics: chikungunya, eastern equine encephalitis, Mayaro, o'nyong-nyong, Ross River, Sindbis, Venezuelan equine encephalitis, and western equine encephalitis. All of the 13 alphaviruses either are rated as Arbovirus (11 viruses), Probable Arbovirus (one virus) or Possible Arbovirus (1 virus).

Previously, Zingilamo virus was shown to morphologically resemble alphaviruses and subsequently was shown to be related to Semliki Forest (SF) virus by CF, HI, and NT (3,4). Upon reexamination, Boteke virus, which was registered as an antigenic relative of Zingilamo virus (see Boteke virus registration card), was found to be unrelated to SF virus both by CF and NT, but still showed a CF relationship to Zingilamo virus (4). However, Boteke virus subsequently was shown to possess rhabdovirus morphology (6) and provisionally is being listed as an ungrouped rhabdovirus separate from Zingilamo virus. Two questions must be answered in order to effect a final placement of these viruses. Firstly, is Zingilamo virus distinct from SF virus? Present evidence would indicate that both viruses differ in CF but that Zingilamo virus appears to be a subtype of SF virus by NT (4). Secondly, it must be established unequivocally whether Boteke virus is related or unrelated to Zingilamo virus.

6. Flaviviruses. Of the 66 registered flaviviruses, 47% have been placed in the mosquito-borne category (Table 6.1), 26% are considered to be tick-borne (Table 6.2), and 27% are categorized as not being associated with a proven arthropod vector (Table 6.3). Only St. Louis encephalitis, West Nile and yellow fever viruses in the mosquito-borne category (Table 6.1) and Powassan virus in the tick-borne category (Table 6.2) have been isolated from both mosquitoes and ticks. Israel turkey meningoencephalitis (IT) virus provisionally has been placed in the mosquito-associated category. Previously, it had been listed in the "no arthropod vector demonstrated" category (Table 6.3). Isolations of IT virus have been reported from Culicoides, mosquitoes (species not specified) and engorged Culex pipiens mosquitoes. Furthermore, experimentally infected Cx. molestus and Aedes aegypti mosquitoes have transmitted the virus by bite to suckling mice.

Twenty-seven of the 31 registered flaviviruses which are mosquito-borne (Table 6.1) are rated as Probable Arbovirus or Arbovirus. The tick-borne flaviviruses (Table 6.2) contain four registered viruses, Absettarov, Hanzalova, Hypr, and Kumlinge, which are very closely related or indistinguishable by conventional serological techniques, though they are purported to be clearly differentiated on the basis of clinical, epidemiological, and ecological markers from Russian spring-summer encephalitis virus and other members of that complex.

Twenty-nine (44%) registered flaviviruses have been isolated from humans; 19 of 31 (61%) mosquito-borne flaviviruses and nine of 17 (53%) tick-borne flaviviruses have been implicated in human disease. By contrast, only five of 18 (28%) flaviviruses not associated with a vector have been implicated in human disease. Modoc virus, listed in Table 6.3, was implicated in a clinically inapparent laboratory infection. Thus, a total of 33 flaviviruses have been associated with disease in humans.

With the exception of Koutango virus, none of the registered flaviviruses placed in the "no arthropod vector demonstrated" category are rated above Possible Arbovirus by SEAS. Seven members are rated as Probably Not or Not Arbovirus. Most of the flaviviruses listed in Table 6.3 have been isolated from rodents or bats. Cacipacore virus has been isolated from a wild bird and Aroa virus from a sentinel hamster. Only Dakar bat and Negishi viruses have been isolated from humans; Negishi virus has been recovered only from that source. While only 2 flaviviruses of this category have been isolated from humans, five have been implicated in human disease. These include Apoi, Dakar Bat, Koutango, Negishi, and Rio Bravo viruses.

Powassan virus was recovered in cell culture prepared from kidney tissue removed from an apparently healthy male spotted skunk (Spilogale putorius) captured in California (20). This is the first reported isolation of Powassan virus west of the Rocky Mountains.

7. Antigenically grouped and ungrouped viruses of families Togaviridae and Flaviviridae. Malakal and Puchong viruses, currently comprising the Malakal serogroup, have been placed here as possible members of the Togaviridae

family. Malakal virus was examined by electron microscopy (EM) and found to resemble viruses of the family Togaviridae (7). Thus far, Malakal virus has not shown an antigenic relationship to any of the other established or provisional togaviruses. Puchong virus has not been examined by EM and has been placed with Malakal virus because of their previously demonstrated antigenic relationship (see Malakal and Puchong virus registration cards).

Trinititi virus was recovered in Trinidad from Trichoprosopon species mosquitoes. It was rated as Probable Arbovirus by SEAS (Table 7.1). This virus was shown to possess a ribonucleic acid (RNA) genome and morphologically it resembled viruses of the family Togaviridae (21).

Simian hemorrhagic fever virus has produced severe disease in rhesus monkeys imported from India. Other monkey species developed disease following contact with the recently imported sick rhesus monkeys. Simian hemorrhagic fever virus has been classified as Not Arbovirus by SEAS. This virus has been shown to resemble the flaviviruses morphologically and structurally, although an antigenic relationship has not been demonstrated.

8. Family Bunyaviridae.

8.1 Bunyaviruses. Sixteen antigenic sets of viruses plus Kaeng Khoi virus (SBU) comprise the bunyaviruses. A total of 123 registered viruses have been placed within the Bunyavirus genus.

8.1.1 Anopheles A and Anopheles B serogroup viruses. Members of the Anopheles A serogroup have been isolated either from anopheline or both culicine and anopheline mosquitoes. Of the five members of this serogroup, only Tacaiuma virus has been isolated from and reported to cause a febrile illness in humans. In addition, this virus has been isolated from a sentinel monkey. Members of this serogroup and of the ANB serogroup appear to be geographically localized.

Viruses of the Anopheles B serogroup have been isolated only from mosquitoes collected in South America. Neither virus has been associated with infections in humans.

8.1.2 Bunyamwera serogroup viruses. All members of the Bunyamwera serogroup have been isolated from culicine or anopheline mosquitoes or both. In addition, Lokern and Main Drain viruses have been isolated from Culicoides species. Anhembi, Germiston, Kairi, Macaua, Northway, Tensaw, and Shokwe viruses have been recovered from rodents, and Lokern, Main Drain, and Tensaw viruses from lagomorphs. Northway virus also was isolated from sentinel rabbits, Kairi virus from a monkey, Macaua virus from a bird, and Tensaw virus from a fox.

Bunyamwera, Germiston, Ilesha, Shokwe, and Wyeomyia viruses have been isolated from humans. These viruses plus Calovo and Tensaw viruses have been associated with human disease, either through infections acquired in nature or in the laboratory, or both. Furthermore, Maguari virus has been isolated from

horses with encephalitis, Cache Valley from a caribou that died, a sick sheep, a cow and from an asymptomatic horse, and Main Drain virus has been isolated from brain tissue of a horse that died of encephalitis.

Fifteen of the 22 viruses registered in the Bunyamwera serogroup have been rated as Arbovirus or Probable Arbovirus. None are rated below Possible Arbovirus.

Members have been found most frequently in North America (8 viruses), South America (8 viruses) and Africa (5 viruses). Thus far, only one virus has been recovered in Asia, two in Europe, and none in Australasia.

8.1.3 Bwamba serogroup and serogroup C viruses. The group C viruses have been closely associated with mosquito vectors and small animals, particularly rodents. Eight group C viruses have been isolated from rodents, and three of these eight additionally have been isolated from marsupials. Two other viruses have been isolated from marsupials but not rodents. Ten of the twelve viruses have been isolated from humans. Only Gumbo Limbo and Vines viruses have not been isolated from humans and, with the exception of those two viruses, all members have been associated with cases of human febrile illness. In addition, Apeu, Caraparu, Marituba, Murutucu, Oriboca and Ossa viruses have been reported to infect humans as a result of laboratory mishaps. Ten of these viruses have been classified as Arboviruses and two as Probable Arboviruses.

Both Bwamba and Pongola viruses (Bwamba serogroup) are mosquito-borne, and both viruses have been isolated from humans. Bwamba virus has been reported to produce a febrile illness in humans as a result of infections acquired in nature. During 1985, it was reported that Pongola virus was isolated in 1982 from acute-phase serum of a young adult human presenting with symptoms of at least fever and headache (22). Thus far, these two viruses have been found in Africa only. Pongola virus has been rated as Arbovirus while Bwamba virus has been rated as Probable Arbovirus.

8.1.4 California and Capim serogroup viruses. All California serogroup viruses are associated with mosquito vectors and four members have been recovered from rodents (Table 8.1.4). La Crosse, Guaroa, and Tahyna viruses have been isolated from humans and, along with California encephalitis, Jamestown Canyon, snowshoe hare and Inkoo viruses, have been associated with disease as a result of infections acquired in nature. In addition, Keystone virus has been implicated in an inapparent infection in a laboratory worker. Antibody to trivittatus virus has been demonstrated in humans although the virus has not been associated with the production of disease in humans. California group infections in humans have been documented serologically in China. Only Inkoo and Tahyna viruses have been isolated on continents other than North and South America. On the basis of virus isolation, the geographic distribution of Tahyna now includes Asia as well as Africa and Europe. Ten of the California serogroup viruses have been rated as Arbovirus, one other as Probable Arbovirus, and the remaining two as Possible Arbovirus.

Viruses of the Capim serogroup are associated with mosquito vectors, and four of the members have been isolated from rodents. None of these eight viruses have been associated with disease in humans. Capim serogroup members have been recovered only in North and South America. Six of the eight Capim serogroup viruses have been rated as Arbovirus (four viruses) or Probable Arbovirus (two viruses).

8.1.5 Gamboa, Guama and Koongol serogroup viruses. In addition to Gamboa virus, the serogroup contains Pueblo Viejo and San Juan viruses (Table 8.1.5). All virus members have been isolated exclusively from Aedeomyia squamipennis mosquitoes. The viruses appear to have a limited geographic distribution, and they have not been implicated in human infections.

Guama serogroup viruses have been found only in the western hemisphere. Catu and Guama viruses have been isolated from humans and have been associated with disease in humans. Nine of the 12 Guama group viruses have been rated as Arbovirus or Probable Arbovirus. Viruses of this serogroup clearly are mosquito-borne and most appear to be associated with rodents. Ten viruses have been isolated from sentinel animals, primarily mice.

Both Koongol group viruses were isolated in Australia and very little is known about them. These two viruses were rated as Probable Arbovirus.

8.1.6 Minatitlan, Olifantsvlei and Patois serogroup viruses. The Minatitlan serogroup listed in Table 8.1.6 now contains two registered members (Minatitlan and Palestina viruses). Several isolations of Palestina virus have been made from Culex sp. mosquitoes collected in Ecuador, and from sentinel hamsters. Minatitlan virus was isolated from a sentinel hamster exposed near Minatitlan, Mexico.

The Olifantsvlei group consists of three members, and all three were isolated in Africa from mosquitoes. Little information on the properties of these viruses is extant.

Viruses of the Patois group now have been isolated in North and South America, and most appear to be associated with mosquito vectors and some with rodent hosts. Babahoyo, Patois, Shark River, and Zegla viruses also were isolated from sentinel hamsters.

None of the viruses from these three serogroups have been isolated from humans, nor have they been associated with disease.

8.1.7 Simbu serogroup viruses. Table 8.1.7 shows that essentially equal numbers of Simbu serogroup viruses have been isolated from Culicoides flies and from mosquitoes. None have been recovered from rodents. Eight Simbu serogroup viruses have been isolated from livestock. These include Sabo, Sango, Shamonda and Shuni viruses (Nigeria); Douglas and Peaton viruses (Australia); Akabane virus (Japan, Kenya and Australia); and Sathuperi virus (India and Africa). In addition, four viruses have been isolated from birds,

and Manzanilla virus has been isolated from a monkey. Oropouche and Shuni viruses are the only members that have been isolated from humans. Oropouche virus has caused frequent large outbreaks of disease in humans in Brazil.

Simbu serogroup viruses have wide distributions. Approximately 50% have been found in Africa or Africa and Asia, while others have been isolated in Asia; Asia and Australasia; Asia, Australasia and Africa; and North or South America. Only eight of the 21 members of this serogroup have been rated as Probable Arbovirus or Arbovirus. The remainder have been rated as Possible Arbovirus.

8.1.8 Tete and Turlock serogroups and unassigned (SBU) viruses. Refer to Table 8.1.8. All Tete serogroup viruses have been recovered from birds; only two of them (Bahig and Matruh viruses) have been recovered from an arthropod vector (ixodid ticks). None of these viruses have been associated with human infections. Only Bahig virus is rated above Possible Arbovirus.

All viruses of the Turlock serogroup are associated with mosquito vectors. In addition, Turlock and Umbre viruses appear to be associated with birds. Turlock virus has been found in both North and South America. All the other members have been found in single continents (Africa, Asia, and Europe).

Only Kaeng Khoi virus remains as a serologically unassigned bunyavirus. Kaeng Khoi virus was isolated from bats, sentinel mice and rats, and cimicid bugs.

8.2 Phleboviruses: Phlebotomus fever serogroup viruses. At present, the Phlebotomus fever serogroup consists of 36 members, and the entire serogroup comprises the Phlebovirus genus within the family Bunyaviridae (Table 8.2). Sicilian sandfly fever virus is the type virus for this genus.

Most of the members are associated with phlebotomine flies; only Arumowot, Chagres, Icoaraci, Itaporanga, Rift Valley fever and Zinga viruses have been isolated from mosquitoes. Nine of the phleboviruses have been isolated from humans or have been implicated in the production of disease in humans.

Gabek Forest virus has not been recovered from arthropods but has been isolated from a variety of rodents and a hedgehog collected in various areas of Africa. Gabek Forest virus has been rated as Probable Arbovirus.

Rift Valley fever (RVF) virus causes serious and extensive disease in domestic animals such as sheep and cattle, and may cause disease in veterinary personnel, field and laboratory workers, and persons who handle infected animals. Serological studies indicated that Zinga virus is closely related or identical to Rift Valley fever virus. Consequently Zinga virus has been placed in the Phlebotomus fever serogroup although it may be another strain of RVF virus. Previously it was listed as an antigenically ungrouped virus.

Corfou virus is a recently registered phlebotavirus and was isolated from Phlebotomus flies in 1982. By CF, Corfou virus is most closely related to SFS virus, although they are readily distinguished by cross-neutralization tests. There is very little information available concerning Corfou virus.

8.3 Nairoviruses. Members of the six antigenic groups shown in Tables 8.3.1 and 8.3.2 constitute the Nairovirus genus in the family Bunyaviridae (16). CHF-Congo virus was designated the type virus for this genus. Furthermore, reproducible intergroup antigenic relationships have been demonstrated for the six sets of viruses (18). Only members of the CHF-Congo and Nairobi sheep disease (NSD) serogroups have been associated with disease in humans.

8.3.1 CHF-Congo, Dera Ghazi Khan, and Hughes serogroups. Both Congo and Crimean hemorrhagic fever viruses are registered in the Catalogue. It must be reiterated that the agent of Crimean hemorrhagic fever (CHF) is antigenically indistinguishable from Congo virus. CHF virus has been implicated in more than two thousand cases of human disease in the USSR. Congo virus also has been associated with the production of disease in humans, either as a result of infections acquired in nature or in the laboratory. Thus far, Hazara virus has not been known to be involved in infections of humans, and little is known of this antigenic relative of CHF-Congo virus. All members of this serogroup appear to be associated with ixodid ticks although CHF virus was isolated from both ixodid and argasid ticks.

Members of the Dera Ghazi Khan (DGK) serogroup have not been isolated from vertebrate hosts, or from arthropod vectors other than ticks. Most of the viruses appear to be associated with argasid ticks. These viruses have been found in Africa, Asia and Australasia.

Only Hughes virus of the Hughes serogroup has been isolated from birds. It has been found in both North and South America while Soldado virus has been isolated in Africa, Europe and South America. All Hughes serogroup members have been associated with argasid ticks.

8.3.2 Nairobi Sheep Disease, Qalyub and Sakhalin serogroups. Nairobi sheep disease virus is an important cause of veterinary disease, while both Dugbe and Ganjam viruses have been isolated repeatedly from ticks removed from domestic animals. Dugbe and Ganjam viruses have caused febrile illnesses in humans. In the case of NSD virus, one infection in a person resulted in a febrile illness, while three others resulted in serologic conversions only. Thus, all three viruses have been isolated from humans and have been associated with laboratory infections. Pending further clarification of antigenic relationships, SIRACA considers Ganjam virus to be a variety of NSD virus.

All three Qalyub group viruses were found only in Africa, and both Bandia and Qalyub viruses have been isolated from ticks. In addition, Bandia and Omo viruses have been isolated from rodents. Omo virus was registered during the first half of 1986, although it had been isolated in 1971. Only a single isolate of Omo virus has been obtained from a rodent, and very little is known about this virus.

Except for Avalon virus, members of the Sakhalin serogroup were isolated only from ixodid ticks. Avalon virus also was recovered from a bird. Sakhalin serogroup viruses are distributed in Asia (Paramushir, Sakhalin), Australasia (Taggert), Europe (Clo Mor), and North America (Avalon). Antigenic studies have indicated that Avalon and Paramushir viruses are strains of the same virus.

8.4 Uukuviruses: Uukuniemi serogroup viruses. Except for Uukuniemi virus, all members of the Uukuniemi serogroup have been isolated only from ticks (Table 8.4). Uukuniemi virus also has been recovered from both rodents and birds. Two of the viruses in this serogroup were found in Asia while three others were discovered in Europe. The sixth member, Precarious Point virus, was found on Australasia. HI antibodies to Uukuniemi virus have been detected in the sera of humans residing in Europe. Grand Arbaud virus has been evaluated as Arbovirus and Uukuniemi as Probable Arbovirus. The rest of the members have been evaluated as Possible Arbovirus.

8.5 Hantaviruses and bunyavirus-like viruses.

8.5.1 Hantaan, Bhanja, Kaisodi and Upolu serogroups. At present, the Hantavirus genus is only a proposed taxon. If approved, this genus will contain the four registered viruses listed in the Hantaan serogroup shown in Table 8.5.1. In addition to Hantaan and Prospect Hill viruses, the serogroup contains Seoul and Puumala viruses. Seoul virus is the prototype virus for a group of Hantaan-related viruses isolated from rats, while Puumala virus is the etiologic agent of Nephropathia Epidemica. All four viruses have been isolated from rodents, while Hantaan virus has also been isolated from humans. Hantaan virus is the etiologic agent of hemorrhagic fever with renal syndrome (HFRS) or Korean hemorrhagic fever (KHF), and either is responsible for or is antigenically closely related to the agent(s) responsible for clinically similar diseases in the USSR, Japan, Manchuria, and Eastern and Northern Europe. More than 10,000 cases have occurred in Korea since the disease was first recognized in that country in 1951. Only Prospect Hill virus has not been shown to produce disease in humans. However, neutralizing antibodies to Prospect Hill virus were detected in the sera of four American mammalogists.

Bhanja virus is the sole registered virus member of the Bhanja serogroup. Kismayo virus, an unregistered member, has been demonstrated to share an antigenic relationship with Bhanja virus (24). Bhanja virus has been isolated from humans and has been implicated in a laboratory-acquired human infection.

Two of the Kaisodi serogroup viruses were isolated from ticks collected in Asia while the third was isolated in North America. None of these viruses have been found to infect humans. Unpublished studies suggest that the RNA species and polypeptides of Silverwater virus resemble those of uukuviruses. Kaisodi and Silverwater viruses had been evaluated as Probable Arbovirus while Lanjan virus had been rated as Possible Arbovirus.

The Upolu serogroup consists of Upolu and Aransas Bay viruses. Both viruses were isolated only from argasid ticks. Neither virus has been associated with infections in humans. One virus has been found in Australia (UPO), and the other in North America (AB).

8.5.2 Bakau, Mapputta, Matariya, Nyando, Resistencia, and Yogue serogroups. All the viruses listed in Table 8.5.2 are members of minor serogroups, and provisionally are classified taxonomically as bunyavirus-like members of the family Bunyaviridae. Most viruses in these minor serogroups have been primarily associated with mosquito vectors. Viruses of the Matariya and Yogue serogroups have not been recovered from mosquitoes.

Bakau serogroup viruses have been recovered only in Asia. Bakau virus has been isolated from mosquitoes, ticks and rodents.

Thus far, all four viruses of the Mapputta group have been found only in Australia. Maprik virus was rated as a Probable Arbovirus while the other three virus members were classified as Possible Arbovirus.

All three Matariya group viruses have been recovered from birds collected in Africa. Nothing is known concerning their possible vector association.

Nyando virus has been isolated from humans and from mosquitoes collected in Africa. The Nyando virus human infection resulted in a febrile illness.

The Resistencia antigenic group consists of three virus members isolated in Argentina from culicine mosquitoes. Electron microscopic investigations conducted with Barranqueras virus have shown that it resembles typical bunyaviruses morphologically and morphogenetically (see Barranqueras virus registration card).

Both Yogue virus and its unregistered relative, Kasokero virus, were isolated from the same species of bats. During the course of characterizing Kasokero virus, it was determined that Kasokero virus was related to Yogue virus and that Kasokero virus readily infected and caused disease in laboratory workers (2). Furthermore, Yogue virus was examined by electron microscopy and found to resemble viruses of the family Bunyaviridae.

8.5.3 Antigenically ungrouped bunyavirus-like viruses. The viruses shown in Table 8.5.3 have been subdivided according to their vector association. Those viruses in the upper part of the table have been listed together as mosquito-borne viruses. Those in the middle section have been associated with tick vectors, while the last two viruses have not been associated with any vector.

Tataguine, Tamdy and Bangui viruses have been isolated from humans and have been implicated in human disease.

Issyk-Kul and Keterah viruses have been shown to be identical or closely related by complement-fixation and neutralization tests (5). Furthermore, electron microscopic studies have indicated that Issyk-Kul virus resembles viruses of the family Bunyaviridae (9). Accordingly, both viruses provisionally are being listed with other bunyavirus-like viruses which are antigenically ungrouped (Table 8.5.3). Inasmuch as Issyk-Kul and Keterah viruses appear to be identical, one of them should be withdrawn as a registered virus. Issyk-Kul virus has been isolated on more than 20 occasions from the blood of humans infected in nature. The infections were classified as febrile illnesses. Issyk-Kul virus also has been implicated in a laboratory infection.

Wanowrie virus recently was added to the listing shown in Table 8.5.3. Electron microscopic examination has shown that Wanowrie virus is a bunyavirus (6). In addition, it recently was demonstrated that this virus possesses hemagglutinating activity (8). Equally as important, Wanowrie virus has caused human disease and appears to be rather widely distributed (see virus registration card). The virus has been isolated in India, Sri Lanka, Iran, and Egypt.

9. Orbiviruses, family Reoviridae: Colorado tick fever and Kemerovo serogroups. While the viruses listed in Table 9.1 are tick-borne agents, they differ taxonomically from those in Tables 8.3.1-8.5.3 in that they have been classified as orbiviruses in the family Reoviridae. The orbiviruses are relatively resistant to lipid solvents, are inactivated at acid pH, and possess multiple segments of a double-stranded RNA genome. It is likely that members of the genus Orbivirus, and that the criteria used to define this genus, will be reevaluated in the near future.

Only Colorado tick fever (CTF) virus of the CTF serogroup and Kemerovo (KEM) and Lipovnik viruses of the KEM serogroup have produced disease in humans or have been isolated from humans.

Members of the KEM serogroup are widely distributed with at least one virus being found in each of the listed continents. KEM and Chenuda viruses have been found in both Africa and Asia while Wad Medani virus has been discovered in Africa, Asia, and North America. Even though all members of this serogroup have been isolated from ticks, only three viruses were rated above Possible Arbovirus. All three were rated as Probable Arbovirus.

9.2 Other antigenic groups of orbiviruses. Several of the viruses in these serogroups cause significant diseases in large animals (Tables 9.2.1, 9.2.2). Bluetongue (BLU) virus causes disease in both wild and domestic ruminants; African horsesickness (AHS) virus in mules, donkeys and horses; epizootic hemorrhagic disease (EHD) virus in deer; and Ibaraki virus in cattle. Both BLU and AHS viruses have wide geographic distributions. The first isolation of BLU virus from a tick (Amblyomma variegatum) has been described (see BLU virus registration card).

Changuinola virus is the only member of these serogroups that has been isolated from humans; it has been reported to produce disease in humans. Of

the present twelve serogroup members, only Irituia, Jari, and Monte Dourado viruses have not been isolated from an arthropod. All others, including Changuinola virus, appear to be associated with phlebotomine insects. Registered viruses of the Changuinola serogroup appear to have a limited distribution. Eleven members were recovered only in South America, while Changuinola virus was isolated in Central America.

Between 1960 and 1980, a total of 178 Changuinola serogroup viruses were isolated in Brazil, Colombia, and Panama. In a detailed study, 24 of those viruses were selected as representative specimens and their antigenic, biological, and chemical properties were examined. Twelve of the viruses were distinct by neutralization tests and polyacrylamide gel electrophoresis (PAGE) (25). This study clearly showed that "a great many more Changuinola serotypes may exist" (25).

The three viruses of the Corriparta serogroup appear to be associated with mosquitoes. In addition, Corriparta virus was recovered from wild birds. All three viruses are widely separated in their distribution.

Thus far, Ibaraki and EHD viruses have not been associated with any known vector. The EHD virus has been found in Africa and North America, while Ibaraki virus has been recovered only in Asia.

Virus members of the Corriparta, Eubenangee, Palyam, and Umatilla serogroups appear to be primarily mosquito-associated, while members of the Wallal and Warrego (WAR) serogroups appear to be associated with Culicoides flies. Vector associations appear to be less clear for Eubenangee (EUB) virus of the EUB serogroup, and for WAR virus of the WAR serogroup.

Llano Seco virus is antigenically related to Umatilla virus, and both are related to the presently unregistered Netivot virus, an orbivirus isolated from mosquitoes collected in Israel (1). Thus, these two registered viruses plus Netivot virus now comprise the Umatilla serogroup.

9.3 Antigenically ungrouped Orbiviruses. The viruses in Table 9.3 are serologically ungrouped, though they have been clustered together on the basis of their association with mosquito or tick vectors.

Of the ungrouped orbiviruses associated with mosquito vectors, two viruses have been found in Africa (LEB, ORU), two in Australasia (JAP, PR) and one in South America (IERI).

Orungo virus has caused human disease as a result of laboratory-acquired infections and those acquired in nature. Lebombo virus, or a closely related virus, has been isolated from human plasma, although it has not been associated with the production of disease in humans thus far.

Chobar Gorge virus provisionally has been placed in the Orbivirus genus as a result of information originally present on the registration card but which previously had been overlooked.

If a virus has not been associated with a vector and has been isolated from bats collected in Nigeria and Cameroun.

10. Family Rhabdoviridae. Members of the serogroups listed in Tables 10.1 and 10.2 or the antigenically ungrouped viruses listed in Table 10.3 possess a "bullet-shaped" morphology and are classified as members of the family Rhabdoviridae.

The rabies serogroup listed in Table 10.1 consists of kotonkan virus and Lagos bat virus. Kotonkan virus was isolated from Culicoides species collected in Nigeria. It was rated as Probable Arbovirus by SEAS. Lagos bat virus has been isolated only from bats and SEAS has evaluated it as Not Arbovirus.

All three viruses of the Sawgrass serogroup were isolated from ticks collected in North America. The viruses of the Timbo serogroup were isolated from lizards, and none of these viruses have been isolated from arthropods.

VSV group members have been recovered from phlebotomine flies, mosquitoes, ticks, Culicoides, mites, Simulium flies and a variety of other arthropods including house flies, face flies, Chloropidae, and Anthomyidae. Piry and VS-Alagoas viruses have not been recovered from arthropods. Of the serogroups listed in this and Table 10.2, only members of the VSV serogroup and Le Dantec virus have been shown to infect humans. In the VSV serogroup, Chandipura, Piry, VS-Indiana and VS-New Jersey viruses have been isolated from man. These viruses, plus VS-Alagoas virus, have been found to produce disease in humans during infections acquired in nature or in the laboratory. Both VS-Indiana and VS-New Jersey viruses readily infect livestock, while Cocal virus has been recovered from a horse and VS-Alagoas virus from a mule.

Table 10.2 contains the newly formed bovine ephemeral fever (BEF) serogroup, the Hart Park serogroup viruses, a Kwatta serogroup virus, the recently formed Le Dantec serogroup, and an expanded Mossuril serogroup consisting of eight members.

The bovine ephemeral fever (BEF) serogroup consists of three registered Australian viruses (AR, BRM, KIM) and BEF virus. Only BEF and KIM viruses have been isolated from vectors. KIM virus has been isolated from Culicoides sp. and culicine mosquitoes, while BEF virus has been isolated from Culicoides sp., anopheline mosquitoes and a mixed pool of mosquito species. All four viruses have been isolated from cattle. Thus far, only BEF virus has been recovered outside of Australia.

All of the Hart Park serogroup members are associated with a mosquito vector and two of the viruses (Hart Park and Flanders) have been isolated from birds. None of these viruses have been associated with disease in humans. Thus far their distribution includes only North and South America.

Kwatta virus was isolated only once from mosquitoes collected in Surinam. An antigenic relative of Kwatta virus remains unregistered. This unregistered virus was recovered from a bird collected in Brazil.

The Le Dantec serogroup consists of Le Dantec and Keuraliba viruses. Prior to the discovery of an antigenic relationship between these two rhabdoviruses, Keuraliba virus was listed as a member of the VSV serogroup. However, this relationship was not reproducible and Keuraliba virus was withdrawn from the VSV serogroup when it was demonstrated to be related to Le Dantec virus. Neither virus has been isolated from an arthropod. Le Dantec virus has been isolated from humans and Keuraliba virus was isolated from rodents.

Three of the members of the Mossuril serogroup have not been isolated from arthropods. These include Cuiaba, Kern Canyon, and Marco viruses. Kern Canyon virus has been rated as Probably Not Arbovirus by SEAS. Previous studies have demonstrated that Kern Canyon virus could be propagated in an Aedes dorsalis cell culture line.

All the viruses listed in Table 10.3 are antigenically ungrouped rhabdoviruses. The first nine viruses shown in the table have been associated with mosquito vectors. Inhangapi and Sripur viruses have been associated with phlebotomine flies and Tibrogargan virus with Culicoides flies. The latter five viruses have been isolated from vertebrates only.

Only Aruac and Almpiwar viruses have been rated as Probable Arbovirus. The rest have been rated as Possible Arbovirus or Probably Not Arbovirus.

None of the viruses listed in this table have been isolated from humans or have been implicated in human disease.

Recently, Boteke, Gossas, Nkolbisson, and Rochambeau viruses were shown to possess rhabdovirus morphology (6). Their antigenic relationship to other rhabdoviruses is being actively investigated.

11. Arenaviruses: Tacaribe serogroup viruses. Tacaribe group viruses in Table 11.1 are serologically related to lymphocytic choriomeningitis virus, and they are classified taxonomically in the Arenavirus genus. They are primarily rodent viruses, and there is little or no evidence that suggests that they are associated with an arthropod vector in nature. SEAS has judged most members to be Not Arbovirus or Probably Not Arbovirus.

Ippy and Toure viruses represented provisional additions to this serogroup. Previously, Ippy virus was found to be related to Lassa virus. Its antigenic relationship to other members of the Tacaribe serogroup has yet to be determined. Characteristically, Ippy virus has been isolated from Mastomys rodents and from rodents of other species. Toure virus has been deleted from this listing. Definitive studies have shown that it is not antigenically related to members of the Tacaribe serogroup (26).

Three members of this group have been shown to cause severe, often fatal, human disease. These include Junin (Argentine hemorrhagic fever), Machupo (Bolivian hemorrhagic fever), and Lassa (Lassa disease). In addition to causing clinically frank laboratory-acquired infections, Junin virus also has

been reported to cause subclinical laboratory-acquired infections. A subclinical seroconversion to Tacaribe virus has been documented in a laboratory worker handling large quantities of Tacaribe virus. In addition, Pichinde virus has produced subclinical infections in laboratory workers. Finally, Flexal virus has produced a febrile illness in a laboratory worker following a laboratory accident. Flexal virus was recovered from rodents trapped in Brazil.

12. Thogoto serogroup viruses and antigenically ungrouped viruses of various taxa. See Table 12.1. Thogoto virus has been isolated from humans and has been involved in human disease. An unregistered antigenic relative of Thogoto virus has been isolated in Sicily (SiAr 126). In fact, there now are five other isolates from Portugal, Iran, and various areas of Africa. Their precise antigenic relationship to the registered Thogoto virus has not been determined. Molecular analysis of a Thogoto serogroup virus has indicated that its virion RNA species and structural polypeptides resemble those of members of the family Orthomyxoviridae (27).

Formerly, the Bunyaviridae study group of the ICTV had classified Dhori virus as a member of the then newly defined Nairovirus genus. Subsequently, molecular studies indicated that Dhori virus possessed seven virion polypeptides and seven single-stranded RNA segments, comparable to those of viruses of the family Orthomyxoviridae (27).

Tettnang virus was shown to cross-react in CF tests with mouse hepatitis virus (MHV). Subsequently, three isolates of Tettnang virus were compared to prototype strains of MHV by neutralization tests (28). The relationship of Tettnang virus to MHV was confirmed; however, the precise relationship of the Tettnang virus isolates to MHV strains remained unclear because of the past passage history of the Tettnang isolates. Further, whether the Tettnang isolates were, in fact, arthropod-borne remains unlikely but unanswered. Bocas virus formerly was included in the CAL serogroup until it was demonstrated that it was identical to or closely related to mouse hepatitis virus. Although both Tettnang and Bocas viruses are closely related to or identical to mouse hepatitis virus, they were not compared to each other.

Nodamura virus was isolated from wild-caught mosquitoes in Japan, and it has been demonstrated to produce disease in moths and honey bees. It also has been shown that it replicates in mosquitoes and is experimentally transmitted by mosquitoes. Nodamura virus is the type species for the Nodavirus genus within the family Nodaviridae.

Cotia virus, a poxvirus, has been reported to produce disease in humans. However, very little information is available concerning Oubangui and Salanga viruses.

13. Taxonomically unclassified viruses.

13.1 Minor antigenic groups. Nyamanini virus and the unregistered Midway virus (29) now constitute the Nyamanini serogroup. Nyamanini virus was isolated from argasid ticks and birds. It has not been associated with human disease.

Quaranfil virus has been isolated from both humans and birds and has been associated with human disease as the result of natural infections. Preliminary molecular studies conducted with Quaranfil virus indicated that this virus may resemble viruses of the family Orthomyxoviridae. At this point, further verification is required. Little is known concerning the behavior of Johnston Atoll virus in nature.

Both Marburg and Ebola viruses have caused human disease as a result of infections acquired in nature and have been associated with laboratory-acquired infections. Ebola virus was found to possess a single-stranded RNA that was noninfectious upon extraction. Recent evidence indicates that there might be different serotypes of Ebola virus (30). Marburg and Ebola viruses have been isolated from humans only.

The two viruses of the Tanjong Rabok serogroup have been isolated in Malaysia but neither has been associated with a vector. Telok Forest virus was isolated from a wild monkey and Tanjong Rabok virus from a sentinel monkey.

13.2 Antigenically ungrouped viruses. The serologically ungrouped viruses in the upper part of Table 13.2.1 have been associated with mosquito vectors, and all remain taxonomically unclassified. Gomoka and Para viruses also have been recovered from sources other than mosquitoes. Gomoka virus was isolated twice from birds collected in the Central African Republic and Para virus was isolated from sentinel mice.

With one exception, viruses shown in the lower portion of Table 13.2.1 have been associated with tick vectors or both tick and mosquito vectors. Ngaingan virus has been associated with Culicoides flies.

Only Termeil virus was rated above Possible Arbovirus and was rated as Probable Arbovirus.

None of the viruses listed in Table 13.2.2 have been isolated from an arthropod vector, and none has been rated higher than Possible Arbovirus. More than half were isolated from birds. Four other viruses have been recovered from rodents, two from bats, and two others from other vertebrates. Twelve of these viruses were recovered in Africa and Asia. The remaining five viruses were found in South America.

Table 14.1 gives continental distribution of viruses in different antigenic groups on the basis of virus isolation. Most of the registered viruses are very limited in their distribution. Approximately 86% have been isolated on a single continent only, while 23 (4.5%) have been found on three or more continents. The largest number of viruses have been isolated in South America and Africa, probably reflecting research emphasis as well as biological limitations.

Table 15.1 shows the number of viruses, according to antigenic group, isolated from various classes of arthropods. About 49% have been recovered from mosquitoes, 22% from ticks, and 18% from all other classes. One hundred and nine registered viruses (22%) have never been recovered from any arthropod vector. The largest number of viruses which have been isolated from any arthropod, have been recovered from a single class only (352 of 397, 88.7%).

Table 16.1 presents a similar type of analysis in terms of virus isolations from various classes of vertebrates. Humans and rodents have provided the largest number of virus isolations. Most of the viruses isolated from vertebrates have been recovered from a single class only (200 of 285, 70.2%).

Table 17.1 lists the viruses by antigenic group that cause disease in humans. Slightly less than 25% of all registered viruses have been associated with human disease, either as a result of infections acquired in nature, from laboratory accidents, or both. Members of serogroups A and B and those in the Bunyamwera Supergroup constitute 42.7% of all registered viruses but about 62% of the instances in which registered viruses are associated with disease production in humans.

An analysis of the SEAS ratings for all registered viruses is presented in Table 18.1. It shows that 266 registrations (52.6%) are rated as Possible Arbovirus. Clearly, additional data are required if we are to have a more precise rating of the arthropod-borne status of these viruses. Sufficient data are available for about 47% of all registered viruses so that 41% are rated Probable Arbovirus or Arbovirus, while 6% are rated Probably Not Arbovirus or Not Arbovirus.

REFERENCES

1. Tesh, R.B., et al. 1986. Biological and antigenic characterization of Netinot virus, an unusual new arbovirus recovered from mosquitoes in Israel. *Am. J. Trop. Med. Hyg.* 35:418-428.
2. Kalunda, M., et al. 1986. Kasokero virus: A new human pathogen from bats (Rousettus aegyptiacus) in Uganda. *Am. J. Trop. Med. Hyg.* 35:387-392.
3. Calisher, C.H., et al. Personal communication. 1986.
4. Digoutte, J.P., et al. 1985. Centre Collaborateur OMS De Reference Et De Recherche Pour Les Arbovirus. p. 14.
5. Main, A.J. and Karabatsos, N. Personal communication. 1986.
6. Bowen, R.A., et al. Personal communication. 1986.
7. Cropp, C.B., et al. Personal communication. 1986.
8. Addy, P.A.K., et al. 1985. Preparation of hitherto undetected haemagglutinin and precipitin antigens of Wanowrie (G-700) virus. *Ann. Inst. Pasteur* 136E:47-54.
9. Lvov, D.K., et al. 1984. Issyk-Kul fever - A new bunyavirus infection in Middle Asia. *Abst. 6th International Conference of Virology, Sendai, Japan.* p. 37.
10. The Subcommittee on Arbovirus Laboratory Safety of The American Committee on Arthropod-borne Viruses. 1980. Laboratory safety for arboviruses and certain other viruses of vertebrates. *Am. J. Trop. Med. Hyg.* 29:1359-1381.
11. American Committee on Arthropod-borne Viruses. 1969. Arbovirus names. *Am. J. Trop. Med. Hyg.* 18:731-734.
12. Shope, R.E. Rabies virus antigenic relationships. In: *The Natural History of Rabies.* Baer, G.M., ed., Academic Press, New York. 1973. pp. 141-152.
13. Taylor, R.M., Comp. 1967. *Catalogue of Arthropod-borne Viruses of the World. A Collection of Data on Registered Arthropod-borne Animal Viruses.* U.S. Public Health Service Publication No. 1760. U.S. Government Printing Office, Washington, D.C.
14. Klimas, R.A., et al. 1981. Radioimmune assays and molecular studies that place Anopheles B and Turlock serogroup viruses in the Bunyavirus genus (Bunyaviridae). *Am. J. Trop. Med. Hyg.* 3:876-887.

15. Report of WHO Scientific Group. 1967. Arboviruses and Human Disease. WHO Technical Report Series No. 369. p. 9.
16. Murphy, F.A., Harrison, A.K., and Whitfield, S.G. 1973. Bunyaviridae: Morphologic and morphogenetic similarities of Bunyamwera serologic supergroup viruses and several other arthropod-borne viruses. *Intervirology* 1:297-316
17. Porterfield, J.S., et al. 1973/74. Bunyaviruses and Bunyaviridae. *Intervirology* 2:270-272.
18. Casals, J. and Tignor, G.H. 1980. The Nairovirus genus: Serological relationships. *Intervirology* 14:144-147.
19. Bishop, D.H.L., et al. 1980. Bunyaviridae. *Intervirology* 14:125-143.
20. Johnson, H.N. 1987. Isolation of Powassan virus from a spotted skunk in California. *J. Wildl. Dis.* 23:152-153.
21. El Mekki, A.A., et al. 1981. Characterization of some ungrouped viruses. *Trans. Roy. Soc. Trop. Med. Hyg.* 75:799-806.
22. Kalunda, M., et al. 1985. Isolation of Chikungunya and Pongola viruses from patients in Uganda. *Trans. Roy. Soc. Trop. Med. Hyg.* 79:567.
23. Schmaljohn, C.S., et al. 1985. Antigenic and genetic properties place viruses linked to hemorrhagic fever with renal syndrome into a newly-defined genus of Bunyaviridae. *Science* 227:1041-1044.
24. Butenko, A., et al. 1979. Kismayo virus - A representative of the Bhanja antigenic group. *Vop. Virusol.* 24(6):661-665.
25. Travassos da Rosa, A.P.A., et al. 1984. Characterization of the Changuinola serogroup viruses. (Reoviridae: Orbivirus). *Intervirology* 21:38-49.
26. McCormick, J.B. Personal communication. 1986.
27. Clerx, J.P.M., Fuller, F. and Bishop, D.H.L. 1983. Tick-borne viruses structurally similar to orthomyxoviruses. *Virology* 127:205-219.
28. Smith, A.L., et al. 1983. Antigenic characterization of Tettang virus: Complications caused by passage of the virus in mice from a colony enzootically infected with mouse hepatitis virus. *Am. J. Trop. Med. Hyg.* 32:1172-1176.
29. Takahashi, M., et al. 1982. Isolation and characterization of Midway virus: A new tick-borne virus related to Nyamanini. *J. Med. Virol.* 10:181-193.

30. Richman, D.D., et al. 1983. Antigenic analysis of strains of Ebola virus: Identification of 2 Ebola virus serotypes. *J. Infect. Dis.* 147:268-271.

Table 1.1

ALPHABETICAL AND TAXONOMIC LISTING OF 506 VIRUSES REGISTERED
AS OF DECEMBER 1986 WITH RECOMMENDED ABBREVIATIONS
AND ANTIGENIC GROUPINGS

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
ABRAS	ABR	Bunyaviridae	<u>Bunyavirus</u>	PAT
ABSETTAROV	ABS	Flaviviridae	<u>Flavivirus</u>	B
ABU HAMMAD	AH	Bunyaviridae	<u>Nairovirus</u>	DGK
ACADO	ACD	Reoviridae	<u>Orbivirus</u>	COR
ACARA	ACA	Bunyaviridae	<u>Bunyavirus</u>	CAP
ADELAIDE RIVER	AR	Rhabdoviridae		BEF
AFRICAN HORSESICKNESS	AHS	Reoviridae	<u>Orbivirus</u>	AHS
AFRICAN SWINE FEVER	ASF	Iridoviridae		
AGUACATE	AGU	Bunyaviridae	<u>Phlebovirus</u>	PHL
AGUA PRETA	AP	Herpesviridae		
AINO	AINO	Bunyaviridae	<u>Bunyavirus</u>	SIM
AKABANE	AKA	Bunyaviridae	<u>Bunyavirus</u>	SIM
ALENQUER	ALE	Bunyaviridae	<u>Phlebovirus</u>	PHL
ALFUJ	ALF	Flaviviridae	<u>Flavivirus</u>	B
ALMEIRIM	AMR	Reoviridae	<u>Orbivirus</u>	CGL
ALMPIWAR	ALM	Rhabdoviridae		
ALTAMIRA	ALT	Reoviridae	<u>Orbivirus</u>	CGL
AMAPARI	AMA	Arenaviridae	<u>Arenavirus</u>	TCR
ANANINDEUA	ANU	Bunyaviridae	<u>Bunyavirus</u>	GMA
ANHANGA	ANH	Bunyaviridae	<u>Phlebovirus</u>	PHL

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
ANHEMBI	AMB	Bunyaviridae	<u>Bunyavirus</u>	BUN
ANOPHELES A	ANA	Bunyaviridae	<u>Bunyavirus</u>	ANA
ANOPHELES B	ANB	Bunyaviridae	<u>Bunyavirus</u>	ANB
ANTEQUERA	ANT	Bunyaviridae	<u>Bunyavirus-like</u>	RTA
APEU	APEU	Bunyaviridae	<u>Bunyavirus</u>	C
APOI	APOI	Flaviviridae	<u>Flavivirus</u>	B
ARAGUARI	ARA			
ARANSAS BAY	AB	Bunyaviridae	<u>Bunyavirus-like</u>	UPO
ARBIA	ARB	Bunyaviridae	<u>Phlebovirus</u>	PHL
ARIDE	ARI			
ARKONAM	ARK			
AROA	AROA	Flaviviridae	<u>Flavivirus</u>	B
ARUAC	ARU	Rhabdoviridae		
ARUMOWOT	AMT	Bunyaviridae	<u>Phlebovirus</u>	PHL
AURA	AURA	Togaviridae	<u>Alphavirus</u>	A
AVALON	AVA	Bunyaviridae	<u>Nairovirus</u>	SAK
BABAHOYO	BAB	Bunyaviridae	<u>Bunyavirus</u>	PAT
BAGAZA	BAG	Flaviviridae	<u>Flavivirus</u>	B
BAHIG	BAH	Bunyaviridae	<u>Bunyavirus</u>	TETE
BAKAU	BAK	Bunyaviridae	<u>Bunyavirus-like</u>	BAK
BAKU	BAKU	Reoviridae	<u>Orbivirus</u>	KEM
BANDIA	BDA	Bunyaviridae	<u>Nairovirus</u>	QYB
BANGORAN	BGN	Rhabdoviridae		MOS
BANGUI	BGI	Bunyaviridae	<u>Bunyavirus-like</u>	

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
BANZI	BAN	Flaviviridae	<u>Flavivirus</u>	B
BARMAH FOREST	BF	Togaviridae	<u>Alphavirus</u>	A
BARRANQUERAS	BQS	Bunyaviridae	<u>Bunyavirus-like</u>	RTA
BARUP	BAR	Rhabdoviridae		MOS
BATAI	BAT	Bunyaviridae	<u>Bunyavirus</u>	BUN
BATAMA	BMA	Bunyaviridae	<u>Bunyavirus</u>	TETE
BATKEN	BKN			
BAULINE	BAU	Reoviridae	<u>Orbivirus</u>	KEM
BEBARU	BEB	Togaviridae	<u>Alphavirus</u>	A
BELEM	BLM			
BELMONT	BEL	Bunyaviridae	<u>Bunyavirus-like</u>	
BENEVIDES	BVS	Bunyaviridae	<u>Bunyavirus</u>	CAP
BENFICA	BEN	Bunyaviridae	<u>Bunyavirus</u>	CAP
BERRIMAH	BRM	Rhabdoviridae		BEF
BERTIOGA	BER	Bunyaviridae	<u>Bunyavirus</u>	GMA
BHANJA	BHA	Bunyaviridae	<u>Bunyavirus-like</u>	BHA
BIMBO	BBO			
BIMITI	BIM	Bunyaviridae	<u>Bunyavirus</u>	GMA
BIRAO	BIR	Bunyaviridae	<u>Bunyavirus</u>	BUN
BLUETONGUE	BLU	Reoviridae	<u>Orbivirus</u>	BLU
BOBAYA	BOB	Bunyaviridae	<u>Bunyavirus-like</u>	
BOBIA	BIA	Bunyaviridae	<u>Bunyavirus</u>	OLI
BOCAS	BOC	Coronaviridae	<u>Coronavirus</u>	
BORACEIA	BOR	Bunyaviridae	<u>Bunyavirus</u>	ANB

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
BOTAMBI	BOT	Bunyaviridae	<u>Bunyavirus</u>	OLI
BOTEKE	BTK	Rhabdoviridae		
BOUBOUI	BOU	Flaviviridae	<u>Flavivirus</u>	B
BOVINE EPHEMERAL FEVER	BEF	Rhabdoviridae		BEF
BUENAVENTURA	BUE	Bunyaviridae	<u>Phlebovirus</u>	PHL
BUJARU	BUJ	Bunyaviridae	<u>Phlebovirus</u>	PHL
BUNYAMWERA	BUN	Bunyaviridae	<u>Bunyavirus</u>	BUN
BUNYIP CREEK	BC	Reoviridae	<u>Orbivirus</u>	PAL
BURG EL ARAB	BEA	Bunyaviridae	<u>Bunyavirus-like</u>	MTY
BUSHBUSH	BSB	Bunyaviridae	<u>Bunyavirus</u>	CAP
BUSSUQUARA	BSQ	Flaviviridae	<u>Flavivirus</u>	B
BUTTONWILLOW	BUT	Bunyaviridae	<u>Bunyavirus</u>	SIM
BWAMBA	BWA	Bunyaviridae	<u>Bunyavirus</u>	BWA
CABASSOU	CAB	Togaviridae	<u>Alphavirus</u>	A
CACAO	CAC	Bunyaviridae	<u>Phlebovirus</u>	PHL
CACHE VALLEY	CV	Bunyaviridae	<u>Bunyavirus</u>	BUN
CACIPACORE	CPC	Flaviviridae	<u>Flavivirus</u>	B
CAIMITO	CAI	Bunyaviridae	<u>Phlebovirus</u>	PHL
CALIFORNIA ENC.	CE	Bunyaviridae	<u>Bunyavirus</u>	CAL
CALOVO	CVO	Bunyaviridae	<u>Bunyavirus</u>	BUN
CANANEIA	CNA	Bunyaviridae	<u>Bunyavirus</u>	GMA
CANDIRU	CDU	Bunyaviridae	<u>Phlebovirus</u>	PHL
CANINDE	CAN	Reoviridae	<u>Orbivirus</u>	CGL
CAPE WRATH	CW	Reoviridae	<u>Orbivirus</u>	KEM

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
CAPIM	CAP	Bunyaviridae	<u>Bunyavirus</u>	CAP
CARAPARU	CAR	Bunyaviridae	<u>Bunyavirus</u>	C
CAREY ISLAND	CI	Flaviviridae	<u>Flavivirus</u>	B
CATU	CATU	Bunyaviridae	<u>Bunyavirus</u>	GMA
CHACO	CHO	Rhabdoviridae		TIM
CHAGRES	CHG	Bunyaviridae	<u>Phlebovirus</u>	PHL
CHANDIPURA	CHP	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
CHANGUINOLA	CGL	Reoviridae	<u>Orbivirus</u>	CGL
CHARLEVILLE	CHV	Rhabdoviridae		MOS
CHENUDA	CNU	Reoviridae	<u>Orbivirus</u>	KEM
CHIKUNGUNYA	CHIK	Togaviridae	<u>Alphavirus</u>	A
CHILIBRE	CHI	Bunyaviridae	<u>Phlebovirus</u>	PHL
CHIM	CHIM			
CHOBAR GORGE	CG	Reoviridae	<u>Orbivirus</u>	
CLO MOR	CM	Bunyaviridae	<u>Nairovirus</u>	SAK
COCAL	COC	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
COLORADO TICK FEVER	CTF	Reoviridae	<u>Orbivirus</u>	CTF
CONGO	CON	Bunyaviridae	<u>Nairovirus</u>	CHF-CON
CONNECTICUT	CNT	Rhabdoviridae		SAW
CORFOU	CFU	Bunyaviridae	<u>Phlebovirus</u>	PHL
CORRIPARTA	COR	Reoviridae	<u>Orbivirus</u>	COR
COTIA	COT	Poxviridae		
COWBONE RIDGE	CR	Flaviviridae	<u>Flavivirus</u>	B
CRIMEAN HEM. FEVER	CHF	Bunyaviridae	<u>Nairovirus</u>	CHF-CON

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
CSIRO VILLAGE	CVG	Reoviridae	<u>Orbivirus</u>	PAL
CUIABA	CUI	Rhabdoviridae		MOS
D'AGUILAR	DAG	Reoviridae	<u>Orbivirus</u>	PAL
DAKAR BAT	DB	Flaviviridae	<u>Flavivirus</u>	B
DENGUE-1	DEN-1	Flaviviridae	<u>Flavivirus</u>	B
DENGUE-2	DEN-2	Flaviviridae	<u>Flavivirus</u>	B
DENGUE-3	DEN-3	Flaviviridae	<u>Flavivirus</u>	B
DENGUE-4	DEN-4	Flaviviridae	<u>Flavivirus</u>	B
DERA GHAZI KHAN	DGK	Bunyaviridae	<u>Nairovirus</u>	DGK
DHORI	DHO	Orthomyxoviridae		
DOUGLAS	DOU	Bunyaviridae	<u>Bunyavirus</u>	SIM
DUGBE	DUG	Bunyaviridae	<u>Nairovirus</u>	NSD
EAST. EQUINE ENC.	EEE	Togaviridae	<u>Alphavirus</u>	A
EBOLA	EBO			MBG
EDGE HILL	EH	Flaviviridae	<u>Flavivirus</u>	B
ENSEADA	ENS	Bunyaviridae	<u>Bunyavirus-like</u>	
ENTEBBE BAT	ENT	Flaviviridae	<u>Flavivirus</u>	B
EP. HEM. DIS.	EHD	Reoviridae	<u>Orbivirus</u>	EHD
ESTERO REAL	ER			
EUBENANGEE	EUB	Reoviridae	<u>Orbivirus</u>	EUB
EVERGLADES	EVE	Togaviridae	<u>Alphavirus</u>	A
EYACH	EYA	Reoviridae	<u>Orbivirus</u>	CTF
FLANDERS	FLA	Rhabdoviridae		HP
FLEXAL	FLE	Arenaviridae	<u>Arenavirus</u>	TCR

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
FORT MORGAN	FM	Togaviridae	<u>Alphavirus</u>	A
FRIJOLES	FRI	Bunyaviridae	<u>Phlebovirus</u>	PHL
GABEK FOREST	GF	Bunyaviridae	<u>Phlebovirus</u>	PHL
GADGETS GULLY	GGY	Flaviviridae	<u>Flavivirus</u>	B
GAMBOA	GAM	Bunyaviridae	<u>Bunyavirus</u>	GAM
GAN GAN	GG	Bunyaviridae	<u>Bunyavirus-like</u>	MAP
GANJAM	GAN	Bunyaviridae	<u>Nairovirus</u>	NSD
GARBA	GAR	Bunyaviridae	<u>Bunyavirus-like</u>	MTY
GERMISTON	GER	Bunyaviridae	<u>Bunyavirus</u>	BUN
GETAH	GET	Togaviridae	<u>Alphavirus</u>	A
GOMOKA	GOM			
GORDIL	GOR	Bunyaviridae	<u>Phlebovirus</u>	PHL
GOSSAS	GOS	Rhabdoviridae		
GRAND ARBAUD	GA	Bunyaviridae	<u>Uukuvirus</u>	UUK
GRAY LODGE	GLO	Rhabdoviridae		
GREAT ISLAND	GI	Reoviridae	<u>Orbivirus</u>	KEM
GUAJARA	GJA	Bunyaviridae	<u>Bunyavirus</u>	CAP
GUAMA	GMA	Bunyaviridae	<u>Bunyavirus</u>	GMA
GUARATUBA	GTB	Bunyaviridae	<u>Bunyavirus</u>	GMA
GUAROA	GRO	Bunyaviridae	<u>Bunyavirus</u>	CAL
GUMBO LIMBO	GL	Bunyaviridae	<u>Bunyavirus</u>	C
GURUPI	GUR	Reoviridae	<u>Orbivirus</u>	CGL
HANTAAN	HTN	Bunyaviridae	<u>Hantavirus*</u>	HTN

*Proposed genus designation.

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
HANZALOVA	HAN	Flaviviridae	<u>Flavivirus</u>	B
HART PARK	HP	Rhabdoviridae		HP
HAZARA	HAZ	Bunyaviridae	<u>Nairovirus</u>	CHF-CON
HIGHLANDS J	HJ	Togaviridae	<u>Alphavirus</u>	A
HUACHO	HUA	Reoviridae	<u>Orbivirus</u>	KEM
HUGHES	HUG	Bunyaviridae	<u>Nairovirus</u>	HUG
HYPR	HYPR	Flaviviridae	<u>Flavivirus</u>	B
IACO	IACO	Bunyaviridae	<u>Bunyavirus</u>	BUN
IBARAKI	IBA	Reoviridae	<u>Orbivirus</u>	EHD
ICOARACI	ICO	Bunyaviridae	<u>Phlebovirus</u>	PHL
IERI	IERI	Reoviridae	<u>Orbivirus</u>	
IFE	IFE	Reoviridae	<u>Orbivirus</u>	
ILESHA	ILE	Bunyaviridae	<u>Bunyavirus</u>	BUN
ILHEUS	ILH	Flaviviridae	<u>Flavivirus</u>	B
INGWAVUMA	ING	Bunyaviridae	<u>Bunyavirus</u>	SIM
INHANGAPI	INH	Rhabdoviridae		
ININI	INI	Bunyaviridae	<u>Bunyavirus</u>	SIM
INKOO	INK	Bunyaviridae	<u>Bunyavirus</u>	CAL
IPPY	IPPY	Arenaviridae	<u>Arenavirus</u>	TCR
IRITUIA	IRI	Reoviridae	<u>Orbivirus</u>	CGL
ISFAHAN	ISF	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
ISRAEL TURKEY MEN.	IT	Flaviviridae	<u>Flavivirus</u>	B
ISSYK-KUL*	IK	Bunyaviridae	<u>Bunyavirus-like</u>	

*Identical to Keterah virus by CF and NT.

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
ITAITUBA	ITA	Bunyaviridae	<u>Phlebovirus</u>	PHL
ITAPORANGA	ITP	Bunyaviridae	<u>Phlebovirus</u>	PHL
ITAQUI	ITQ	Bunyaviridae	<u>Bunyavirus</u>	C
ITIMIRIM	ITI	Bunyaviridae	<u>Bunyavirus</u>	GMA
ITUPIRANGA	ITU			
JACAREACANGA	JAC	Reoviridae	<u>Orbivirus</u>	COR
JAMANXI	JAM	Reoviridae	<u>Orbivirus</u>	CGL
JAMESTOWN CANYON	JC	Bunyaviridae	<u>Bunyavirus</u>	CAL
JAPANAUT	JAP	Reoviridae	<u>Orbivirus</u>	
JAPANESE ENC.	JBE	Flaviviridae	<u>Flavivirus</u>	B
JARI	JARI	Reoviridae	<u>Orbivirus</u>	CGL
JERRY SLOUGH	JS	Bunyaviridae	<u>Bunyavirus</u>	CAL
JOHNSTON ATOLL	JA			QRF
JOINJAKAKA	JOI	Rhabdoviridae		
JUAN DIAZ	JD	Bunyaviridae	<u>Bunyavirus</u>	CAP
JUGRA	JUG	Flaviviridae	<u>Flavivirus</u>	B
JUNIN	JUN	Arenaviridae	<u>Arenavirus</u>	TCR
JURONA	JUR	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
JUTIAPA	JUT	Flaviviridae	<u>Flavivirus</u>	B
KADAM	KAD	Flaviviridae	<u>Flavivirus</u>	B
KAENG KHOI	KK	Bunyaviridae	<u>Bunyavirus</u>	SBU
KAIKALUR	KAI	Bunyaviridae	<u>Bunyavirus</u>	SIM
KAIRI	KRI	Bunyaviridae	<u>Bunyavirus</u>	BUN
KAISODI	KSO	Bunyaviridae	<u>Bunyavirus-like</u>	KSO

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
KAMESE	KAM	Rhabdoviridae		MOS
KAMMAVANPETTAI	KMP			
KANNAMANGALAM	KAN			
KAO SHUAN	KS	Bunyaviridae	<u>Nairovirus</u>	DGK
KARIMABAD	KAR	Bunyaviridae	<u>Phlebovirus</u>	PHL
KARSHI	KSI	Flaviviridae	<u>Flavivirus</u>	B
KASBA	KAS	Reoviridae	<u>Orbivirus</u>	PAL
KEMEROVO	KEM	Reoviridae	<u>Orbivirus</u>	KEM
KERN CANYON	KC	Rhabdoviridae		MOS
KETAPANG	KET	Bunyaviridae	<u>Bunyavirus-like</u>	BAK
KETERAH*	KTR	Bunyaviridae	<u>Bunyavirus-like</u>	
KEURALIBA	KEU	Rhabdoviridae		LD
KEYSTONE	KEY	Bunyaviridae	<u>Bunyavirus</u>	CAL
KHASAN	KHA	Bunyaviridae	<u>Nairovirus</u>	CHF-CON
KIMBERLEY	KIM	Rhabdoviridae		BEF
KLAMATH	KLA	Rhabdoviridae		
KOKOBERA	KOK	Flaviviridae	<u>Flavivirus</u>	B
KOLONGO	KOL	Rhabdoviridae		
KOONGOL	KOO	Bunyaviridae	<u>Bunyavirus</u>	KOO
KOTONKAN	KOT	Rhabdoviridae	<u>Lyssavirus</u>	RABIES
KOUTANGO	KOU	Flaviviridae	<u>Flavivirus</u>	B
KOWANYAMA	KOW	Bunyaviridae	<u>Bunyavirus-like</u>	
KUMLINGE	KUM	Flaviviridae	<u>Flavivirus</u>	B

*Identical to Issyk-Kul virus by CF and NT.

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
KUNJIN	KUN	Flaviviridae	<u>Flavivirus</u>	B
KUNUNURRA	KNA	Rhabdoviridae		
KWATTA	KWA	Rhabdoviridae		KWA
KYASANUR FOR. DIS.	KFD	Flaviviridae	<u>Flavivirus</u>	B
KYZYLAGACH	KYZ	Togaviridae	<u>Alphavirus</u>	A
LA CROSSE	LAC	Bunyaviridae	<u>Bunyavirus</u>	CAL
LAGOS BAT	LB	Rhabdoviridae	<u>Lyssavirus</u>	RABIES
LA JOYA	LJ	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
LAKE CLARENDON	LC			
LANDJIA	LJA			
LANGAT	LGT	Flaviviridae	<u>Flavivirus</u>	B
LANJAN	LJN	Bunyaviridae	<u>Bunyavirus-like</u>	KSO
LAS MALOYAS	LM	Bunyaviridae	<u>Bunyavirus</u>	ANA
LASSA	LAS	Arenaviridae	<u>Arenavirus</u>	TCR
LATINO	LAT	Arenaviridae	<u>Arenavirus</u>	TCR
LEBOMBO	LEB	Reoviridae	<u>Orbivirus</u>	
LE DANTEC	LD	Rhabdoviridae		LD
LEDNICE	LED	Bunyaviridae	<u>Bunyavirus</u>	TUR
LIPOVNIK	LIP	Reoviridae	<u>Orbivirus</u>	KEM
LLANO SECO	LLS	Reoviridae	<u>Orbivirus</u>	UMA
LOKERN	LOK	Bunyaviridae	<u>Bunyavirus</u>	BUN
LONE STAR	LS	Bunyaviridae	<u>Bunyavirus-like</u>	
LOUPING ILL	LI	Flaviviridae	<u>Flavivirus</u>	B
LUKUNI	LUK	Bunyaviridae	<u>Bunyavirus</u>	ANA

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
MACAUA	MCA	Bunyaviridae	<u>Bunyavirus</u>	BUN
MACHUPO	MAC	Arenaviridae	<u>Arenavirus</u>	TCR
MADRID	MAD	Bunyaviridae	<u>Bunyavirus</u>	C
MAGUARI	MAG	Bunyaviridae	<u>Bunyavirus</u>	BUN
MAHOGANY HAMMOCK	MH	Bunyaviridae	<u>Bunyavirus</u>	GMA
MAIN DRAIN	MD	Bunyaviridae	<u>Bunyavirus</u>	BUN
MALAKAL	MAL	Togaviridae		MAL
MANAWA	MWA	Bunyaviridae	<u>Uukuvirus</u>	UUK
MANZANILLA	MAN	Bunyaviridae	<u>Bunyavirus</u>	SIM
MAPPUTTA	MAP	Bunyaviridae	<u>Bunyavirus</u> -like	MAP
MAPRIK	MPK	Bunyaviridae	<u>Bunyavirus</u> -like	MAP
MAPUERA	MPR			
MARBURG	MBG			MBG
MARCO	MCO	Rhabdoviridae		MOS
MARITUBA	MTB	Bunyaviridae	<u>Bunyavirus</u>	C
MARRAKAI	MAR	Reoviridae	<u>Orbivirus</u>	PAL
MATARIYA	MTY	Bunyaviridae	<u>Bunyavirus</u> -like	MTY
MATRUH	MTR	Bunyaviridae	<u>Bunyavirus</u>	TETE
MATUCARE	MAT			
MAYARO	MAY	Togaviridae	<u>Alphavirus</u>	A
MEABAN	MEA	Flaviviridae	<u>Flavivirus</u>	B
MELAO	MEL	Bunyaviridae	<u>Bunyavirus</u>	CAL
MERMET	MER	Bunyaviridae	<u>Bunyavirus</u>	SIM
MIDDELBURG	MID	Togaviridae	<u>Alphavirus</u>	A

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
MINATITLAN	MNT	Bunyaviridae	<u>Bunyavirus</u>	MNT
MINNAL	MIN			
MIRIM	MIR	Bunyaviridae	<u>Bunyavirus</u>	GMA
MITCHELL RIVER	MR	Reoviridae	<u>Orbivirus</u>	WAR
MODOC	MOD	Flaviviridae	<u>Flavivirus</u>	B
MOJU	MOJU	Bunyaviridae	<u>Bunyavirus</u>	GMA
MOJUI DOS CAMPOS	MDC			
MONO LAKE	ML	Reoviridae	<u>Orbivirus</u>	KEM
MONT. MYOTIS LEUK.	MML	Flaviviridae	<u>Flavivirus</u>	B
MONTE DOURADO	MDO	Reoviridae	<u>Orbivirus</u>	CGL
MORICHE	MOR	Bunyaviridae	<u>Bunyavirus</u>	CAP
MOSQUEIRO	MQO	Rhabdoviridae		HP
MOSSURIL	MOS	Rhabdoviridae		MOS
MOUNT ELGON BAT	MEB	Rhabdoviridae		
M'POKO	MPO	Bunyaviridae	<u>Bunyavirus</u>	TUR
MUCAMBO	MUC	Togaviridae	<u>Alphavirus</u>	A
MUNGUBA	MUN	Bunyaviridae	<u>Phlebovirus</u>	PHL
MURRAY VALLEY ENC.	MVE	Flaviviridae	<u>Flavivirus</u>	B
MURUTUCU	MUR	Bunyaviridae	<u>Bunyavirus</u>	C
MYKINES	MYK	Reoviridae	<u>Orbivirus</u>	KEM
NAIROBI SHEEP DIS.	NSD	Bunyaviridae	<u>Nairovirus</u>	NSD
NARANJAL	NJL	Flaviviridae	<u>Flavivirus</u>	B
NARIVA	NAR	Paramyxoviridae	<u>Paramyxovirus</u>	
NAVARRO	NAV	Rhabdoviridae		

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
NDUMU	NDU	Togaviridae	<u>Alphavirus</u>	A
NEGISHI	NEG	Flaviviridae	<u>Flavivirus</u>	B
NEPUYO	NEP	Bunyaviridae	<u>Bunyavirus</u>	C
NEW MINTO	NM	Rhabdoviridae		SAW
NGAINGAN	NGA			
NIQUE	NIQ	Bunyaviridae	<u>Phlebovirus</u>	PHL
NKOLBISSON	NKO	Rhabdoviridae		
NODAMURA	NOD	Nodaviridae	<u>Nodavirus</u>	
NOLA	NOLA	Bunyaviridae	<u>Bunyavirus</u>	SIM
NORTHWAY	NOR	Bunyaviridae	<u>Bunyavirus</u>	BUN
NTAYA	NTA	Flaviviridae	<u>Flavivirus</u>	B
NUGGET	NUG	Reoviridae	<u>Orbivirus</u>	KEM
NYAMANINI	NYM			NYM
NYANDO	NDO	Bunyaviridae	<u>Bunyavirus-like</u>	NDO
OKHOTSKIY	OKH	Reoviridae	<u>Orbivirus</u>	KEM
OKOLA	OKO			
OLIFANTSVLEI	OLI	Bunyaviridae	<u>Bunyavirus</u>	OLI
OMO	OMO	Bunyaviridae	<u>Nairovirus</u>	QYB
OMSK HEM. FEVER	OMSK	Flaviviridae	<u>Flavivirus</u>	B
O'NYONG-NYONG	ONN	Togaviridae	<u>Alphavirus</u>	A
ORIBOCA	ORI	Bunyaviridae	<u>Bunyavirus</u>	C
ORIXIMINA	ORX	Bunyaviridae	<u>Phlebovirus</u>	PHL
OROPOUCHE	ORO	Bunyaviridae	<u>Bunyavirus</u>	SIM
ORUNGO	ORU	Reoviridae	<u>Orbivirus</u>	

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
PONTEVES	PTV	Bunyaviridae	<u>Uukuvirus</u>	UUK
POWASSAN	POW	Flaviviridae	<u>Flavivirus</u>	B
PRECARIOUS POINT	PP	Bunyaviridae	<u>Uukuvirus</u>	UUK
PRETORIA	PRE	Bunyaviridae	<u>Nairovirus</u>	DGK
PROSPECT HILL	PH	Bunyaviridae	<u>Hantavirus*</u>	HTN
PUCHONG	PUC	Togaviridae		MAL
PUEBLO VIEJO	PV	Bunyaviridae	<u>Bunyavirus</u>	GAM
PUNTA SALINAS	PS	Bunyaviridae	<u>Nairovirus</u>	HUG
PUNTA TORO	PT	Bunyaviridae	<u>Phlebovirus</u>	PHL
PURUS	PUR	Reoviridae	<u>Orbivirus</u>	CGL
PUUMALA	PUU	Bunyaviridae	<u>Hantavirus*</u>	HTN
QALYUB	QYB	Bunyaviridae	<u>Nairovirus</u>	QYB
QUARANFIL	QRF			QRF
RAZDAN	RAZ	Bunyaviridae	<u>Bunyavirus-like</u>	
RESISTENCIA	RTA	Bunyaviridae	<u>Bunyavirus-like</u>	RTA
RESTAN	RES	Bunyaviridae	<u>Bunyavirus</u>	C
RIFT VALLEY FEVER	RVF	Bunyaviridae	<u>Phlebovirus</u>	PHL
RIO BRAVO	RB	Flaviviridae	<u>Flavivirus</u>	B
RIO GRANDE	RG	Bunyaviridae	<u>Phlebovirus</u>	PHL
ROCHAMBEAU	RBU	Rhabdoviridae		
ROCIO	ROC	Flaviviridae	<u>Flavivirus</u>	B
ROSS RIVER	RR	Togaviridae	<u>Alphavirus</u>	A
ROYAL FARM	RF	Flaviviridae	<u>Flavivirus</u>	B

*Proposed genus designation.

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
OSSA	OSSA	Bunyaviridae	<u>Bunyavirus</u>	C
OUANGO	OUA			
OUBANGUI	OUB	Poxviridae		
OUREM	OUR	Reoviridae	<u>Orbivirus</u>	CGL
PACORA	PCA	Bunyaviridae	<u>Bunyavirus-like</u>	
PACUI	PAC	Bunyaviridae	<u>Phlebovirus</u>	PHL
PAHAYOKEE	PAH	Bunyaviridae	<u>Bunyavirus</u>	PAT
PALESTINA	PLS	Bunyaviridae	<u>Bunyavirus</u>	MNT
PALYAM	PAL	Reoviridae	<u>Orbivirus</u>	PAL
PARA	PARA			
PARAMUSHIR	PMR	Bunyaviridae	<u>Nairovirus</u>	SAK
PARANA	PAR	Arenaviridae	<u>Arenavirus</u>	TCR
PAROO RIVER	PR	Reoviridae	<u>Orbivirus</u>	
PATA	PATA	Reoviridae	<u>Orbivirus</u>	EUB
PATHUM THANI	PTH	Bunyaviridae	<u>Nairovirus</u>	DGK
PATOIS	PAT	Bunyaviridae	<u>Bunyavirus</u>	PAT
PEATON	PEA	Bunyaviridae	<u>Bunyavirus</u>	SIM
PHNOM-PENH BAT	PPB	Flaviviridae	<u>Flavivirus</u>	B
PICHINDE	PIC	Arenaviridae	<u>Arenavirus</u>	TCR
PICOLA	PIA			
PIRY	PIRY	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
PIXUNA	PIX	Togaviridae	<u>Alphavirus</u>	A
PLAYAS	PLA	Bunyaviridae	<u>Bunyavirus</u>	BUN
PONGOLA	PGA	Bunyaviridae	<u>Bunyavirus</u>	BWA

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
RUSS. SPR. SUM. ENC.	RSSE	Flaviviridae	<u>Flavivirus</u>	B
SABO	SABO	Bunyaviridae	<u>Bunyavirus</u>	SIM
SABOYA	SAB	Flaviviridae	<u>Flavivirus</u>	B
SAGIYAMA	SAG	Togaviridae	<u>Alphavirus</u>	A
SAINT-FLORES	SAF	Bunyaviridae	<u>Phlebovirus</u>	PHL
SAKHALIN	SAK	Bunyaviridae	<u>Nairovirus</u>	SAK
SAKPA	SPA			
SALANGA	SGA	Poxviridae		
SALEHABAD	SAL	Bunyaviridae	<u>Phlebovirus</u>	PHL
SAL VIEJA	SV	Flaviviridae	<u>Flavivirus</u>	B
SAN ANGELO	SA	Bunyaviridae	<u>Bunyavirus</u>	CAL
SANDFLY F. (NAPLES)	SFN	Bunyaviridae	<u>Phlebovirus</u>	PHL
SANDFLY F. (SICILIAN)	SFS	Bunyaviridae	<u>Phlebovirus</u>	PHL
SANDJIMBA	SJA			
SANGO	SAN	Bunyaviridae	<u>Bunyavirus</u>	SIM
SAN JUAN	SJ	Bunyaviridae	<u>Bunyavirus</u>	GAM
SAN PERLITA	SP	Flaviviridae	<u>Flavivirus</u>	B
SANTAREM	STM			
SANTA ROSA	SAR	Bunyaviridae	<u>Bunyavirus</u>	BUN
SARACA	SRA	Reoviridae	<u>Orbivirus</u>	CGL
SATHUPERI	SAT	Bunyaviridae	<u>Bunyavirus</u>	SIM
SAUMAREZ REEF	SRE	Flaviviridae	<u>Flavivirus</u>	B
SAWGRASS	SAW	Rhabdoviridae		SAW
SEBOKELE	SEB			

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
SELETAR	SEL	Reoviridae	<u>Orbivirus</u>	KEM
SEMBALAM	SEM			
SEMLIKI FOREST	SF	Togaviridae	<u>Alphavirus</u>	A
SENA MADUREIRA	SM	Rhabdoviridae		TIM
SEOUL	SEO	Bunyaviridae	<u>Hantavirus*</u>	HTN
SEPIK	SEP	Flaviviridae	<u>Flavivirus</u>	B
SERRA DO NAVIO	SDN	Bunyaviridae	<u>Bunyavirus</u>	CAL
SHAMONDA	SHA	Bunyaviridae	<u>Bunyavirus</u>	SIM
SHARK RIVER	SR	Bunyaviridae	<u>Bunyavirus</u>	PAT
SHOKWE	SHO	Bunyaviridae	<u>Bunyavirus</u>	BUN
SHUNI	SHU	Bunyaviridae	<u>Bunyavirus</u>	SIM
SILVERWATER	SIL	Bunyaviridae	<u>Bunyavirus</u> -like	KSO
SIMBU	SIM	Bunyaviridae	<u>Bunyavirus</u>	SIM
SIMIAN HEM. FEVER	SHF	Flaviviridae		
SINDBIS	SIN	Togaviridae	<u>Alphavirus</u>	A
SIXGUN CITY	SC	Reoviridae	<u>Orbivirus</u>	KEM
SLOVAKIA	SLO			
SNOWSHOE HARE	SSH	Bunyaviridae	<u>Bunyavirus</u>	CAL
SOKULUK	SOK	Flaviviridae	<u>Flavivirus</u>	B
SOLDADO	SOL	Bunyaviridae	<u>Nairovirus</u>	HUG
SOROROCA	SOR	Bunyaviridae	<u>Bunyavirus</u>	BUN
SPONDWENI	SPO	Flaviviridae	<u>Flavivirus</u>	B
SRIPUR	SRI	Rhabdoviridae		

*Proposed genus designation.

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
ST. LOUIS ENC.	SLE	Flaviviridae	<u>Flavivirus</u>	B
STRATFORD	STR	Flaviviridae	<u>Flavivirus</u>	B
SUNDAY CANYON	SCA	Bunyaviridae	<u>Bunyavirus-like</u>	
TACAIUMA	TCM	Bunyaviridae	<u>Bunyavirus</u>	ANA
TACARIBE	TCR	Arenaviridae	<u>Arenavirus</u>	TCR
TAGGERT	TAG	Bunyaviridae	<u>Nairovirus</u>	SAK
TAHYNA	TAH	Bunyaviridae	<u>Bunyavirus</u>	CAL
TAMDY	TDY	Bunyaviridae	<u>Bunyavirus-like</u>	
TAMIAMI	TAM	Arenaviridae	<u>Arenavirus</u>	TCR
TANGA	TAN			
TANJONG RABOK	TR			TR
TATAGUINE	TAT	Bunyaviridae	<u>Bunyavirus-like</u>	
TEHRAN	TEH	Bunyaviridae	<u>Phlebovirus</u>	PHL
TELOK FOREST	TF			TR
TEMBE	TME			
TEMBUSU	TMU	Flaviviridae	<u>Flavivirus</u>	B
TENSAW	TEN	Bunyaviridae	<u>Bunyavirus</u>	BUN
TERMEIL	TER			
TETE	TETE	Bunyaviridae	<u>Bunyavirus</u>	TETE
TETTANG	TET	Coronaviridae		
THIMIRI	THI	Bunyaviridae	<u>Bunyavirus</u>	SIM
THOGOTO	THO	Orthomyxoviridae		THO
THOTTAPALAYAM	TPM			
TIBROGARGAN	TIB	Rhabdoviridae		

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
TILLIGERRY	TIL	Reoviridae	<u>Orbivirus</u>	EUB
TIMBO	TIM	Rhabdoviridae		TIM
TIMBOTEUA	TBT	Bunyaviridae	<u>Bunyavirus</u>	GMA
TINAROO	TIN	Bunyaviridae	<u>Bunyavirus</u>	SIM
TINDHOLMUR	TDM	Reoviridae	<u>Orbivirus</u>	KEM
TLACOTALPAN	TLA	Bunyaviridae	<u>Bunyavirus</u>	BUN
TONATE	TON	Togaviridae	<u>Alphavirus</u>	A
TOSCANA	TOS	Bunyaviridae	<u>Phlebovirus</u>	PHL
TOURE	TOU			
TRIBEC	TRB	Reoviridae	<u>Orbivirus</u>	KEM
TRINITI	TNT	Togaviridae		
TRIVITTATUS	TVT	Bunyaviridae	<u>Bunyavirus</u>	CAL
TRUBANAMAN	TRU	Bunyaviridae	<u>Bunyavirus-like</u>	MAP
TSURUSE	TSU	Bunyaviridae	<u>Bunyavirus</u>	TETE
TURLOCK	TUR	Bunyaviridae	<u>Bunyavirus</u>	TUR
TURUNA	TUA	Bunyaviridae	<u>Phlebovirus</u>	PHL
TYULENIY	TYU	Flaviviridae	<u>Flavivirus</u>	B
UGANDA S	UGS	Flaviviridae	<u>Flavivirus</u>	B
UMATILLA	UMA	Reoviridae	<u>Orbivirus</u>	UMA
UMBRE	UMB	Bunyaviridae	<u>Bunyavirus</u>	TUR
UNA	UNA	Togaviridae	<u>Alphavirus</u>	A
UPOLU	UPO	Bunyaviridae	<u>Bunyavirus-like</u>	UPO
URUCURI	URU	Bunyaviridae	<u>Phlebovirus</u>	PHL
USUTU	USU	Flaviviridae	<u>Flavivirus</u>	B

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
UTINGA	UTI	Bunyaviridae	<u>Bunyavirus</u>	SIM
UUKUNIEMI	UUK	Bunyaviridae	<u>Uukuvirus</u>	UUK
VELLORE	VEL	Reoviridae	<u>Orbivirus</u>	PAL
VEN. EQUINE ENC.	VEE	Togaviridae	<u>Alphavirus</u>	A
VENKATAPURAM	VKT			
VINCES	VIN	Bunyaviridae	<u>Bunyavirus</u>	C
VIRGIN RIVER	VR	Bunyaviridae	<u>Bunyavirus</u>	ANA
VS-ALAGOAS	VSA	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
VS-INDIANA	VSI	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
VS-NEW JERSEY	VSNJ	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
WAD MEDANI	WM	Reoviridae	<u>Orbivirus</u>	KEM
WALLAL	WAL	Reoviridae	<u>Orbivirus</u>	WAL
WANOWRIE	WAN	Bunyaviridae	<u>Bunyavirus-like</u>	
WARREGO	WAR	Reoviridae	<u>Orbivirus</u>	WAR
WESSELSBRON	WSL	Flaviviridae	<u>Flavivirus</u>	B
WEST. EQUINE ENC.	WEE	Togaviridae	<u>Alphavirus</u>	A
WEST NILE	WN	Flaviviridae	<u>Flavivirus</u>	B
WHATAROA	WHA	Togaviridae	<u>Alphavirus</u>	A
WITWATERSRAND	WIT	Bunyaviridae	<u>Bunyavirus-like</u>	
WONGAL	WON	Bunyaviridae	<u>Bunyavirus</u>	KOO
WONGORR	WGR			
WYEOMYIA	WYO	Bunyaviridae	<u>Bunyavirus</u>	BUN
XIBUREMA	XIB	Rhabdoviridae		
YACAABA	YAC			

NAME	ABBR.	TAXONOMIC STATUS		ANTI-GENIC GROUP
		FAMILY	GENUS	
YAQUINA HEAD	YH	Reoviridae	<u>Orbivirus</u>	KEM
YATA	YATA	Rhabdoviridae		
YELLOW FEVER	YF	Flaviviridae	<u>Flavivirus</u>	B
YOGUE	YOG	Bunyaviridae	<u>Bunyavirus-like</u>	YOG
YUG BOGDANOVAC	YB	Rhabdoviridae	<u>Vesiculovirus</u>	VSV
ZALIV TERPENIYA	ZT	Bunyaviridae	<u>Uukuvirus</u>	UUK
ZEGLA	ZEG	Bunyaviridae	<u>Bunyavirus</u>	PAT
ZIKA	ZIKA	Flaviviridae	<u>Flavivirus</u>	B
ZINGA	ZGA	Bunyaviridae	<u>Phlebovirus</u>	PHL
ZINGILAMO	ZGO	Togaviridae	<u>Alphavirus</u>	A
ZIRQA	ZIR	Bunyaviridae	<u>Nairovirus</u>	HUG

Table 2.1 Antigenic Groups of 506 Viruses Registered in Catalogue

Virus Family and Genus	Antigenic Group	Abbreviation	No. Registered Viruses in Group	Percent
<u>ARENAVIRIDAE</u>				
<u>Arenavirus</u>	Tacaribe	TCR	11	2.2
<u>BUNYAVIRIDAE</u>				
<u>Bunyavirus</u> (Bunyamwera Supergroup)	Anopheles A	ANA	5	24.3
	Anopheles B	ANB	2	
	Bunyamwera	BUN	22	
	Bwamba	BWA	2	
	C	C	12	
	California	CAL	13	
	Capim	CAP	8	
	Gamboia	GAM	3	
	Guama	GMA	12	
	Koongol	KOO	2	
	Minatitlan	MNT	2	
	Olifantsvlei	OLI	3	
	Patois	PAT	6	
	Simbu	SIM	21	
	Tete	TETE	5	
	Turlock	TUR	4	
	Unassigned	SBU	1	
<u>Nairovirus</u>			24	4.7
	CHF-Congo	CHF-CON	4	
	Dera Ghazi Khan	DGK	5	
	Hughes	HUG	4	
	Nairobi sheep disease	NSD	3	
	Qalyub	QYB	3	
	Sakhalin	SAK	5	
<u>Phlebovirus</u>	Phlebotomus fever	PHL	36	7.1

Table 2.1 (Continued)

Virus Family and Genus	Antigenic Group	Abbreviation	No. Registered Viruses in Group	Percent
RUNYAVIRIDAE				
<u>Uukuvirus</u>	Uukuniemi	UUK	6	1.2
<u>Hantavirus*</u>	Hantaan	HTN	4	0.8
"Bunyavirus-like" (Unassigned, probable or possible members)	Bakau	BAK	2	0.4
	Bhanja	BHA	1	0.2
	Kaisodi	KSO	3	0.6
	Mapputta	MAP	4	0.8
	Matariya	MTY	3	0.6
	Nyando	NDO	1	0.2
	Resistencia	RTA	3	0.6
	Upolu	UPO	2	0.4
	Yogue	YOG	1	0.2
	Ungrouped		15	3.0
REOVIRIDAE				
<u>Orbivirus</u>	African horsesickness	AHS	1	0.2
	Bluetongue	BLU	1	0.2
	Changuinola	CGL	12	2.4
	Colorado tick fever	CTF	2	0.4
	Corriparta	COR	3	0.6
	Epizootic hemorrhagic dis.	EHD	2	0.4
	Eubenangee	EUB	3	0.6
	Kemerovo	KEM	18	3.6
	Palyam	PAL	7	1.4
	Umatilla	UMA	2	0.4
	Wallal	WAL	1	0.2
	Warrego	WAR	2	0.4
	Ungrouped		7	1.4

*Proposed genus designation.

Table 2.1 (Continued)

Virus Family and Genus	Antigenic Group	Abbreviation	No. Registered Viruses in Group	Percent
RHABDOVIRIDAE				
<u>Vesiculovirus</u>	Vesicular stomatitis	VSV	10	2.0
<u>Lyssavirus</u>	Rabies		2	0.4
Unassigned or possible members	Bovine ephemeral fever	BEF	4	0.8
	Hart Park	HP	3	0.6
	Kwatta	KWA	1	0.2
	Le Dantec	LD	2	0.4
	Mossuril	MOS	8	1.6
	Sawgrass	SAW	3	0.6
	Timbo	TIM	3	0.6
	Ungrouped		18	3.6
TOGAVIRIDAE				
<u>Alphavirus</u>	A		27	5.3
Possible members	Malakal		2	0.4
	Ungrouped		1	0.2
FLAVIVIRIDAE				
<u>Flavivirus</u>	B		66	13.0
Possible members	Ungrouped		1	0.2
CORONAVIRIDAE				
<u>Coronavirus</u>	Ungrouped		2	0.4
HERPESVIRIDAE				
	Ungrouped		1	0.2
AFRICAN SWINE FEVER*				
	Ungrouped		1	0.2

*Removed from family Iridoviridae. A new official family name has not been designated.

Table 2.1 (Continued)

Virus Family and Genus	Antigenic Group	Abbreviation	No. Registered Viruses in Group	Percent
NODAVIRIDAE <u>Nodavirus</u>	Ungrouped		1	0.2
ORTHOMYXOVIRIDAE	Thogoto	THO	1	0.2
	Ungrouped		1	0.2
PARAMYXOVIRIDAE	Ungrouped		1	0.2
POXVIRIDAE	Ungrouped		3	0.6
UNCLASSIFIED	Marburg	MBG	2	0.4
	Nyamanini	NYM	1	0.2
	Tanjong Rabok	TR	2	0.4
	Quaranfil	QRF	2	0.4
	Ungrouped		<u>37</u>	7.3
TOTAL			506	

Table 3.1 Initial Isolations of Viruses by Decade and Country of Origin

Decade	Continent	Country	Virus
1900-09	Africa	S. Africa	BLU
1910-19	Africa	Kenya	ASF, NSD
1920-29	Africa	Nigeria	YF
	Europe	Scotland	LI
	N. America	U.S.A.	VSI
1930-39	Africa	Kenya	RVF
		S. Africa	AHS
		Uganda	BWA, WN
	Asia	Japan	JE
		U.S.S.R.	RSSE
	N. America	U.S.A.	EEE, SLE, WEE
	S. America	Venezuela	VEE
1940-49	Africa	Uganda	BUN, NTA, SF, UGS, ZIKA
	Asia	Japan	NEG
		U.S.S.R.	OMSK
	Australasia	Hawaii	DEN-1
		New Guinea	DEN-2
	Europe	Czechoslovakia	HAN
		Italy	SFN, SFS
	N. America	U.S.A.	CE, CTF, TVT
	S. America	Brazil	ILH
	Colombia	ANA, ANB, WYO	
	1950-59	Africa	Egypt
Nigeria			ILE, LB
S. Africa			BAN, GER, ING, LEB, MID, MOS, NDU, NYM, PGA, SIM, SPO, TETE, USU, WIT, WSL
Asia		Sudan	WM
		Uganda	CHIK, CON, ENT, NDO, ONN, ORU
		India	ARK, BHA, GAN, KAS, KSO, KFD, MIN, PAL, SAT, VKT, UMB, WAN
		Israel	IT
		Japan	AKA, APOI, IBA, NOD, SAG, TSU
		Malaya	BAK, BAT, BEB, GET, KET, LGT, TMU
		Australasia	Australia
Europe		Philippines	DEN-3, DEN-4
		Czechoslovakia	HYPR, TAH
		Finland	KUM
N. America		U.S.S.R.	ABS
		Canada	POW
		Panama	BOC, LJ, PCA
S. America		U.S.A.	CV, EHD, HP, MML, MOD, RB, SA, SSH, TUR, VSNJ
		Argentina	JUN
		Brazil	APEU, AURA, BSQ, CAP, CAR, CATU, GJA, GMA, ITQ, MAG, MIR, MOJU, MTB, MUC, MUR, ORI, TCM, UNA
		Colombia	GRO, NAV
		Trinidad	ARU, BIM, BSB, IERI, KRI, LUK, MAN, MAY, MEL, NEP, ORO, TCR, TNT

Table 3.1 (Continued)

Decade	Continent	Country	Virus	
1960-69	Africa	Cameroun	NKO,OKO	
		Cent. Afr. Rep.	BAG,BGN,BIA,BIR,BOT,BOU,BTK,MPO, PATA,YATA,ZGA	
		Egypt	ACD,AMT,BAH,BEA,MTR,MTY,RF	
		Kenya	THO	
		Nigeria	DUG,KOT,LAS,SABO,SAN,SHA,SHU	
		Senegal	BDA,DB,GOS,KEU,KOU,LD,SAB,TAT,TOU,YOG	
		South Africa	OLI,SHO	
		Sudan	GF,MAL	
		Uganda	KAD,KAM,MEB,TAN	
		Asia	Cambodia	PPB
			India	BAR,CHP,DHO,KAN,KMP,SEM,THI,TPM,VEL
			Iran	KAR,SAL,TEH
			Japan	AINO
			Malaysia	JUG,KTR,LJN,PUC,TR
	Pakistan (West)		DGK,HAZ,MWA	
	Persian Gulf		ZIR	
	Singapore		SEL	
	Thailand		KK	
	U.S.S.R.		CHF,KYZ,OKH,SAK,TYU,ZT	
	Australasia	Australia	ALF,ALM,BEF,BEL,CHV,COR,DAG,EH,EUB, JAP,JOI,KOK,KOO,KOW,KUN,MAP,MPK,MR, RR,SEP,STR,TRU,UPO,WAR,WON	
		New Zealand	WHA	
		Pacific Island	JA	
		Europe	Czechoslovakia	CVO,KEM,LED,LIP,TRB
			Finland	INK,UUK
	France		GA,PTV	
	N. America	West Germany	MBG	
		Canada	SIL	
		Guatemala	JUT	
		Mexico	MNT,TLA	
		Panama	AGU,CHG,CHI,CGL,FRI,GAM,JD,LAT,MAD, MAT,OSSA,PAR,PAT,PT,ZEG	
		U.S.A.	BUT,CR,EVE,FLA,GL,HJ,HUG,JC,JS,KC, KEY,KLA,LAC,LOK,LS,MER,MD,MH,ML,PAH, SAW,SC,SHF,SR,TAM,TEN,UMA	
	S. America	Bolivia	MAC	
		Brazil	ACA,AMA,AMB,ANH,ANU,AP,ARA,BEN,BER, BLM,BOR,BUJ,BVS,CAN,CDU,CHO,COT,GTB, GUR,ICO,INH,IRI,ITP,JUR,MCO,OUR,PAC, PIRY,PIX,SDN,SOR,TBT,TIM,TME,URU,UTI, VSA	
		Colombia	BUE,PIC	
		French Guiana	CAB	

Table 3.1 (Continued)

Decade	Continent	Country	Virus	
1960-69	S. America	Peru	HUA, PS	
		Surinam	KWA	
		Trinidad	COC, MOR, NAR, RES, SOL	
1970-79	Africa	Cent. Afr. Rep.	BBO, BGI, BMA, BOB, GAR, GOM, GOR, IPPY, KOL, LJA, NOLA, OUA, OUB, SAF, SEB, SGA, SJA, SPA, ZGO	
		Egypt	AH, KS, PTH	
		Ethiopia	OMO	
		Nigeria	IFE	
		Seychelles	ARI	
		S. Africa	PRE	
		Zaire	EBO	
		Asia	India	CG, KAI, SRI
			Iran	ISF
			Korea	HTN
			Malaysia	CI, TF
	U.S.S.R.		BKN, CHIM, IK, KHA, KSI, PMR, RAZ, SOK, TDY	
	Australasia	Australia	BC, BF, CVG, DOU, GGY, GG, KNA, MAR, NGA, NUG, PEA, PIA, PR, PP, SRE, TAG, TER, TIB, TIL, TIN, WAL, WGR, YAC	
		Europe	Czechoslovakia	SLO
			Denmark	MYK, TDM
			Germany	EYA, TET
			Italy	TOS
			Scotland	CM, CW
			U.S.S.R.	BAKU
			Yugoslavia	YB
	N. America	Canada	AVA, BAU, GI	
		Mexico	SAR	
		Panama	CAC, CAI, NIO	
		U.S.A.	AB, CNT, FM, GLO, LLS, NM, NOR, RG, SCA, SP, SV, VR, YH	
		S. America	Brazil	ALE, ALT, CNA, CPC, CUI, ENS, FLE, IACO, ITA, ITI, ITU, JAC, JAM, MCA, MDC, MPR, MQO, PARA, ROC, SM, STM, TUA
	Ecuador		ABR, BAB, NJL, PLA, PLS, PV, SJ, VIN	
	French Guiana		INI, RBU, TON	
	Venezuela		AROA	
				Korea
	1980-86	Asia	Korea	SEO
		Australasia	Australia	AR, BRM, KIM, LC
			Europe	Finland
		France	MEA	
		Greece	CFU	
		Italy	ARB	
N. America		Cuba	ER	
		USA	PH	
S. America		Argentina	ANT, BQS, LM, RTA	
		Brazil	AMR, JARI, MDO, MUN, ORX, PUR, SRA, XIB	

Table 4.1 Initial Isolation of 506 Registered Viruses by Continent or Region, Country, and Chronological Period

Continent	Country or Area	Before 1930	1930-39	1940-49	1950-59	1960-69	1970-79	1980-86	Totals
AFRICA	Cameroon					2			2
	Cent. Afr. Rep.					11	19		30
	Egypt				5	7	3		15
	Ethiopia						1		1
	Kenya	2	1			1			4
	Nigeria	1			2	7	1		11
	Senegal					10			10
	Seychelles						1		1
	S. Africa	1	1		15	2	1		20
	Sudan					2			2
	Uganda		2	5	6	4			17
Zaire							1	1	
	Totals	4	4	5	28	46	27	0	114
ASIA	Cambodia					1			1
	India				12	9	3		24
	Iran					3	1		4
	Israel				1				1
	Japan		1	1	6	1			9
	Korea						1	1	2
	Malaysia				7	5	2		14
	W. Pakistan					3			3
	Persian Gulf					1			1
	Singapore					1			1
	Thailand					1			1
U.S.S.R. (East)		1	1		6	9		17	
	Totals	0	2	2	26	31	16	1	78
AUSTRAL-ASIA and PACIFIC ISLANDS	Australia				1	25	23	4	53
	Hawaii			1					1
	Johnston Island					1			1
	New Guinea			1					1
	New Zealand					1			1
	Philippines				2				2
	Totals	0	0	2	3	27	23	4	59
EUROPE	Czechoslovakia			1	2	5	1		9
	Denmark						2		2
	Finland				1	2		1	4
	France					2		1	3
	Greece							1	1
	Italy			2			1	1	4
	Scotland	1					2		3
	U.S.S.R. (West)				1		1		2
	West Germany					1	2		3
	Yugoslavia						1		1
	Totals	1	0	3	4	10	10	4	32

Table 4.1 (Continued)

Continent	Country or Area	Before 1930	1930-39	1940-49	1950-59	1960-69	1970-79	1980-86	Totals
NORTH AMERICA	Canada				1	1	3		5
	Cuba							1	1
	Guatemala					1			1
	Mexico					2	1		3
	Panama				3	15	3		21
	U.S.A.	1	3	3	10	27	13	1	58
	Totals	1	3	3	14	46	20	2	89
SOUTH AMERICA	Argentina				1			4	5
	Bolivia					1			1
	Brazil			1	18	37	22	8	86
	Colombia			3	2	2			7
	Ecuador						8		8
	French Guiana					1	3		4
	Peru					2			2
	Surinam					1			1
	Trinidad				13	5			18
	Venezuela		1					1	2
	Totals	0	1	4	34	49	34	12	134
	GRAND TOTALS	6	10	19	109	209	130	23	506

Table 5.1 Alphaviruses, Family Togaviridae

VIRUS	ISOLATED FROM											ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS				
	ARTHROPODS						VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis			
	Mosq.	Ticks		Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats													Marsupials	Other	Sentinels
		Culicine	Anophelinae																							
Aura	+																		2	S	22	Alphavirus				
Barmah Forest	+																		2	A7	22	"				
Bebaru	+																		2	S	22	"				
Cabassou	+																		2	S	22	"				
Chikungunya	+																		3*	IE	21	"				
Eastern equine enc.	+																		3*	S	20	"				
Everglades	+																		2V	S	20	"				
Fort Morgan	+																		3*	S	20	"				
Getah	+																		2	S	20	"				
Highlands J	+																		3	A1	20	"				
Kyzylgach	+																		2	S	20	"				
Mayaro	+																		3	IE	22	"				
Middleburg	+																		3	S	20	"				
Mucambo	+																		3	S	20	"				
Ndumu	+																		3	A1	21	"				
O'nyong-nyong	+																		3	S	20	"				
Pixuna	+																		2	S	22	"				
Ross River	+																		2	S	20	"				
Sagiyama	+																		3	A1	21	"				
Semliki Forest	+																		3	A2	20	"				
Sindbis	+																		2	S	20	"				
Tonate	+																		3*	IE	21	"				
Una	+																		2	S	21	"				
Ven. equine enc.	+																		3*	V	20	"				
Western equine enc.	+																		2V	S	20	"				
Whataroa	+																		2	S	20	"				
Zingilamo																			2	S	22	"				

* Work with these viruses at containment level 3 requires HEPA filtration of all exhaust air prior to discharge to the outside.

α Inapparent infection(s) only.

** 20 = Arbovirus
21 = Probable Arbovirus
22 = Possible Arbovirus
23 = Probably not Arbovirus
24 = Not Arbovirus

V = Vaccination with demonstration of antibody development; without such vaccination, the next higher containment level is recommended.

Table 6.1 Mosquito-Borne Flaviviruses, Family Flaviviridae

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS					
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis				
	Mosq.	Ticks		Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds													Bats	Marsupials	Other	Sentinels
		Culicine	Anophele																							
Alfuy	+								+									2	S	20	Flavivirus					
Bagaza	+																	2	S	20	"					
Banzi	+					+							+					2	S	20	"					
Boubout	+	+					+										+	2	S	20	"					
Bussuquara	+					+		+					+			+		2	S	20	"					
Dengue 1	+					+		+					+			+		2	S	20	"					
Dengue 2	+					+	+						+			+		2	S	20	"					
Dengue 3	+					+							+			+		2	S	20	"					
Dengue 4	+					+							+			+		2	S	20	"					
Edge Hill	+	+											+			+		2	S	20	"					
Ilheus	+					+			+				+			+		2	S	20	"					
Israel turkey men.	+				+				+	+			+			+		3	S	21	"					
Japanese enc.	+	+				+			+	+			+			+		2	S	20	"					
Jugra	+									+						+		2	S	20	"					
Kokobera	+														+			2	S	20	"					
Kunjin	+					+			+			+				+		2	S	20	"					
Murray Valley enc.	+					+			+			+				+		3	S	20	"					
Naranjal	+											+				+		3	IE	21	"					
Ntaya	+								+				+			+		2	S	21	"					
Rocio	+					+			+						+		3*	S	20	"						
Sepik	+														+		3	IE	21	"						
St. Louis enc.	+	+				+			+	+					+		3	S	20	"						
Spondweni	+					+							+			+		3	S	22	"					
Stratford	+																	2	S	20	"					
Tembusu	+	+											+			+		2	S	21	"					
Uganda S	+									+			+					2	S	20	"					
Usutu	+					+			+	+			+			+		2	S	22	"					
Wesselsbron	+	+				+			+			+			+		3*	S	20	"						
West Nile	+	+	+	+		+			+	+		+			+		3	S	20	"						
Yellow fever	+	+				+			+	+		+			+		3* ^v	S	20	"						
Zika	+					+	+					+				+		2	S	20	"					

* See footnote Table 5.1

V: See footnote Table 5.1

** See footnote Table 5.1

X: Arboviruses restricted by U.S. Department of Agriculture regulations or policy.

^a Inapparent infections only.

Table 6.2 Tick-Borne Flaviviruses, Family Flaviviridae

VIRUS	ISOLATED FROM											ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS						VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosp. Culicine	Anopheline	Ticks Ixodid	Argasid	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds												
Absettarov			+				+											+	+	4	A4	20	Flavivirus
Gadgets Gully			+																	4	A4	22	"
Hanzalova			+																	4	S	20	"
Hypr			+						+	+	+								+	4	S	20	"
Kadam			+																	2	S	21	"
Karshi				+																2	S	22	"
Kumlinge			+				+		+	+									+	4	A4	20	"
Kyasanur Forest dis.			+	+			+		+	+	+								+	4	S	20	"
Langat			+						+	+										2	S	20	"
Louping ill			+						+	+									+	3*	S	20	"
Meaban				+																4	S	22	"
Omsk hem. fever			+						+	+									+	3	S	20	"
Powassan		+	+						+	+									+	4	S	20	"
Royal Farm				+					+	+									+	2	S	22	"
Russ. spr. sum. enc.			+							+										4	S	20	"
Saumarez Reef				+						+										3	S	22	"
Tyuleniy			+																+	4	IE	22	"
																				2	S	21	"

* See footnote Table 5.1

** See footnote Table 5.1

X See footnote Table 6.1

Table 8.1.1 Bunyaviruses, Family Bunyaviridae:
Bunyamvera Supergroup, Anopheles A and Anopheles B Serogroup Viruses

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosq.	Ticks	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats												
<u>ANOPHELES A GR.</u>																						
Anopheles A																						
Las Maloyas																						
Lukuni																						
Tacaiuma																						
Virgin River																						
<u>ANOPHELES B GR.</u>																						
Anopheles B																						
Boraceia																						

** See footnote Table 5.1

Table 8.1.5 Bunyaviruses, Family Bunyaviridae:
Bunyamwera Supergroup, Gambia, Guama and Koongol Serogroup Viruses

VIRUS	ISOLATED FROM				ISOLATED IN						HUMAN DISEASE		SALS RATING		SEAS RATING**		TAXONOMIC STATUS		
	ARTHROPODS				VERTEBRATES						Lab Infection	Natural Infection	Level	Basis	21	22			
	Mosq.	Ticks	Ixodid		Anopheline	Other	Humans	Other Primates	Rodents	Birds								Bats	Marsupials
			Argasid																
GAMBIA GR. Gambia Pueblo Viejo San Juan	+	+	+																
	+																		
	+																		
GUAMA GR. Ananindeua Bertioga Bimiti Cananea Catu Guama Guaratuba Itimirim Mahogany Hammock Mirim Moju Timboteua	+																		
	+																		
	+																		
	+																		
	+																		
	+																		
	+																		
	+																		
	+																		
	+																		
KOONGOL GR. Koongol Wonga	+																		
	+																		
	+																		
	+																		

** See footnote Table 5.1

Table 8.1.6 Bunyaviruses, Family Bunyaviridae:
Bunyamwera Supergroup, Minatitlan, Olifantsvlei and Patois Serogroup Viruses

VIRUS	ISOLATED FROM						ISOLATED IN						HUMAN DISEASE	SALS RATING		SEAS RATING**	TAXONOMIC STATUS
	ARTHROPODS			VERTEBRATES			Africa	Asia	Australasia	Europe	North America	South America		Level	Basis		
	Mosq.	Ticks	Other	Humans	Other Primates	Rodents							Birds			Bats	Marsupials
							Culicine	Anopheline	Ixodid	Argasid							
<u>MINATITLAN GR.</u> Minatitlan Palestina													2	S		22	Bunyavirus
												+	3	1E		21	"
	+										+						
<u>OLIFANTSVLEI GR.</u> Bobia Botambi Olifantsvlei																	Bunyavirus
													3	1E		22	"
	+												2	S		22	"
													2	S		22	"
<u>PATOIS GR.</u> Abrax Bahahoyo Pahayokey Patois Shark River Zegla																	Bunyavirus
																	"
																	"
																	"
	+																"
																	"
																	"
																	"
																	"

** See footnote Table 5.1

Table 8.1.7 Bunyaviruses, Family Bunyviridae:
Bunyamvera Supergroup, Simbu Serogroup Viruses

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosq.	Ticks	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats												
Culicine	Anopheline	Ixodid	Arasid																			
Aino	+																	3	S	22	Bunyavirus	
Akabane	+																	3	S	21	"	
Buttonwillow																		2	S	20	"	
Douglas																		3	IE	21	"	
Ingwavuma	+																	2	S	20	"	
Inini																		3	IE	22	"	
Kaikalur	+																	2	S	22	"	
Manzanilla																		2	S	22	"	
Mermet																		2	S	22	"	
Nola	+																	2	S	20	"	
Oropouche	+																	3	S	21	"	
Peaton																		3	A1	21	"	
Sabo																		2	S	22	"	
Sango	+																	2	S	22	"	
Sathuperi	+																	2	S	22	"	
Shamonda																		2	S	22	"	
Shuni	+																	2	S	22	"	
Simbu	+																	2	S	21	"	
Thimiri																		2	S	22	"	
Tinaroo																		3	IE	22	"	
Utinga																		3	IE	22	"	

* See footnote Table 5.1

** See footnote Table 5.1

Table 8.1.8 Bunyaviruses, Family Bunyaviridae:
Bunyamwera Supergroup, Tete and Turlock Serogroups and Unassigned Viruses

VIRUS	ISOLATED FROM						ISOLATED IN						HUMAN DISEASE	SEALS RATING**		TAXONOMIC STATUS	
	ARTHROPODS			VERTEBRATES			Africa	Asia	Australasia	Europe	North America	South America		Level	Basis		
	Mosq.	Ticks	Other	Humans	Other Primates	Rodents							Birds			Bats	Marsupials
<u>TETE GR.</u>																	
Bahig																	21
Batama																	22
Matruh																	22
Tete																	22
Tsuruse																	22
<u>TURLOCK GR.</u>																	
Lednice																	21
H'Poko																	22
Turlock																	20
Umbre																	21
<u>UNASSIGNED - "SBU"</u>																	
Kaeng Khoi																	22

** See footnote Table 5.1

Table 8.2 Phleboviruses, Family Bunyaviridae:
Phlebotomus Fever Serogroup Viruses

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosq. Culicine	Anopheles	Ixodid	Argasid	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents												
Aguacate				+															2	S	21	Phlebovirus
Alenquer							+												3	IE	22	"
Anhanga																			2	S	22	"
Arbia				+															2	S	21	"
Arumowot	+								+	+									2	S	22	"
Buenaventura				+															3	IE	22	"
Bujaru									+										2	S	22	"
Cacao				+															2	S	21	"
Caimito				+															2	S	22	"
Candiru							+												2	S	22	"
Chagres	+			+			+												2	S	21	"
Chilibre				+															2	S	21	"
Corfou				+															2	S	22	"
Frijoles				+																	22	"
Gabek Forest									+										2	S	21	"
Gordil									+	+									3	IE	22	"
Icoaraci	+	+		+					+	+									2	S	21	"
Itaituba									+	+									3	IE	22	"
Itaporanga	+									+									2	S	20	"
Karimabad				+															2	S	21	"
Munguba				+															3	IE	22	"
Nique				+															2	S	22	"
Oriximina				+															3	IE	22	"
Pacuí				+					+										2	S	21	"
Punto Toro				+															2	S	21	"
Rift Valley fever	+							+											3*	VXS	20	"

* See footnote Table 5.1

** See footnote Table 5.1

V: See footnote Table 5.1

X: See footnote Table 6.1

Table 8.3.1 Nairoviruses, Family Bunyaviridae:
CHF-Congo, Dera Ghazi Khan and Hughes Serogroups

VIRUS	ISOLATED FROM						ISOLATED IN			HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS			
	ARTHROPODS			VERTEBRATES			Africa	Asia	Australasia	Europe	North America	South America	Natural Infection			Lab Infection	Level	Basis
	Mosq.	Ticks	Other	Humans	Other Primates	Rodents												
<u>CHF-CONGO GR.</u> Congo Crimean hem. fever Hazara Khasan				+														
				+														
				+														
				+														
<u>DERA GHAZI KHAN GR.</u> Abu Hamad Dera Ghazi Khan Kao Shuan Pathum Thani Pretoria																		
<u>HUGHES GR.</u> Hughes Punta Salinas Soldado Zirqa																		

** See footnote Table 5.1

Table 8.3.2 Nairoviruses, Family Bunyaviridae:
Nairobi Sheep Disease, Qalyub and Sakhalin Serogroups

VIRUS	ISOLATED FROM											ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS				
	ARTHROPODS						VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis			
	Mosq.	Ticks		Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats													Marsupials	Other	Sentinels
		Culicine	Anophelinae																							
<u>NAIROBI SHEEP DIS.</u> Dugbe Ganjam Nairobi sheep dis.	+					+		+	+				+					+	+	3 X X	S	21 22 20	Nairovirus " "			
<u>QALYUB GR.</u> Randia Omo Qalyub									+	+										2 2	S S	22 22 20	Nairovirus " "			
<u>SAKHALIN GR.</u> Avalon Clo Mor Paramushir Sakhalin Taggert										+										2 2 3 2 2	S S IE S S	21 22 22 21 22	Nairovirus " " " "			

** See footnote Table 5.1
X: See footnote Table 6.1

Table 8.5.1 Hantaviruses and Bunyavirus-like Viruses, Family Bunyaviridae:
Hantaan and Other Antigenic Groups

VIRUS	ISOLATED FROM											ISOLATED IN					HUMAN DISEASE	SALS RATING		SEAS RATING**	TAXONOMIC STATUS		
	ARTHIPODS					VERTEBRATES						Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection			Level	Basis
	Mosq. Culicine	Ticks Ixodid Argasid	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats	Marsupials												
HANTAAN GR. Hantaan Prospect Hill Puumala Seoul						+		+	+	+								+	+	3 [†]	S	22 23 23 23	Hantavirus* " " "
BHANJA GR. Bhanja		+				+		+			+								+	3	S	21	Bunyavirus-like
KAISODI GR. Kaisodi Lanjan Silverwater			+						+											2 2 2	S S S	21 22 21	Bunyavirus-like " "
UPOLU GR. Aransas Bay Upolu																				3 2	IE S	22 22	Bunyavirus-like "

* Hantavirus: Proposed genus designation.

** See footnote Table 5.1

† If virus is handled in very high concentrations or in animals, then level 4.

Table 8.5.2 Bunyavirus-Like Viruses, Family Bunyaviridae:
Other Antigenic Groups

VIRUS	ISOLATED FROM						ISOLATED IN						HUMAN DISEASE		SALS RATING		SEAS RATING**		TAXONOMIC STATUS
	ARTHROPODS			VERTEBRATES			Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level	Basis	22	21	
	Mosq.	Ticks	Culicine	Anopheline	Ixodid	Argasid													
<u>BAKAU GR.</u> Bakau Ketapang	+																2	2	Bunyavirus-like
<u>MAPPITTA GR.</u> Gan Gan Mapputta Maprik Trubanan	+																2	2	Bunyavirus-like
<u>MATARIYA GR.</u> Burg el Arab Garba Matariya																	2	2	Bunyavirus-like
<u>NYANDIO GR.</u> Nyando	+																2	2	Bunyavirus-like
<u>RESISTENCIA GR.</u> Antequera Barranqueras Resistencia																			Bunyavirus-like
<u>YOGUE GR.</u> Yogue																	2	2	Bunyavirus-like

** See footnote Table 5.1

Table 9.1 Orbiviruses, Family Reoviridae:
Colorado Tick Fever and Kemerovo Serogroups

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosq.	Ticks		Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds												
<u>COL. TICK FEV. GR.</u> Col. tick fever Eyach			+				+		+								+	+	2 2	S S	20 22	Orbivirus "
<u>KEMEROVO GR.</u> Baku																			2	S	22	Orbivirus
Bauline			+																2	S	22	"
Cape Wrath			+																2	S	22	"
Chenuda			+																2	S	22	"
Great Island			+																2	S	22	"
Huacho																			2	S	22	"
Kemerovo			+				+		+										2	S	21	"
Lipovnik			+																2	S	22	"
Mono Lake																			2	S	22	"
Mykines			+																	S	22	"
Nugget			+																2	S	22	"
Okhotskiy			+																2	S	22	"
Seletar			+																2	S	22	"
Sixgun City																			2	S	22	"
Tindhölmur			+																2	S	22	"
Tribec			+						+										2	S	21	"
Waa Meani			+																2	S	21	"
Yaquina Head			+																2	S	22	"

** See footnote Table 5.1

Table 9.2.1 Orbiviruses, Family Reoviridae:
Bluetongue Group and Other Antigenic Groups

VIRUS	ISOLATED FROM										ISOLATED IN		HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS					
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America			Natural Infection	Lab Infection	Level	Basis	
	Mosq.	Ticks	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats													Marsupials
<u>AFR. HORSE SICKNESS</u> Afr. horsesickness				+															X			20	Orbivirus
<u>BLUETONGUE GR.</u> Bluetongue				+															2	S		20	Orbivirus
<u>CHANGUINOLA GR.</u> Almeirim Altamira Caninde Changuinola Gurupi Irituia Jamanxi Jari Monte Dourado Ourem Purus Saraca						+													3	IE		22	Orbivirus
																			3	IE		22	"
																			2	S		21	"
																			2	S	+	22	"
																			3	IE		22	"
																			3	IE		22	"
																			3	IE		22	"
																			3	IE		22	"
<u>CORRIPARTA GR.</u> Acado Corriparta Jacareacanga																			2	S		22	Orbivirus
																			2	S		21	"
																			3	IE		22	"
<u>EIHU GR.</u> Epizootic hem.dis. Ibaraki																			2	S		21	Orbivirus
																			3	IE		22	"

** See footnote Table 5.1
X See footnote Table 6.1

Table 9.3 Orbiviruses, Family Reoviridae:
Antigenically Ungrouped Viruses

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS					
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis				
	Mosq.	Ticks		Phlebotomine	Cuticoides	Other	Humans	Other Primates	Rodents	Birds													Bats	Marsupials	Other	Sentinels
		Anopheline	Ixodid																							
Culicine																										
Ieri	+																	2	S	22	Orbivirus					
Japanaut	+																	2	S	21	"					
Lebombo	+				+					+								2	S	21	"					
Orungo	+	+				+	+										+	3	S	21	"					
Paroo River	+																	3	IE	22	"					
Chobar Gorge							+											2	S	22	Orbivirus					
Ife										+									3	IE	22	Orbivirus				

** See footnote Table 5.1

Table 10.1 Lyssaviruses and Vesiculoviruses, Family Rhabdoviridae; Family Rhabdoviridae:
Vesicular Stomatitis and Other Antigenic Groups

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SCAS RATING**	TAXONOMIC STATUS					
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis				
	Mosq.	Ticks		Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds													Bats	Marsupials	Other	Sentinels
		Culicine	Anophele																							
<u>RABIES SEROGROUP</u>																										
Kotonkan				+														2	S	21	Lyssavirus					
Lagos bat										+								2	S	24	"					
<u>SAWGRASS GR.</u>																										
Connecticut																		3	IE	22	Rhabdoviridae					
New Minto																		3	IE	22	"					
Sawgrass																		2	S	22	"					
<u>TIMBO GR.</u>																										
Chuco																		2	S	22	Rhabdoviridae					
Sena Madureira																		3	IE	22	"					
Timbo																		2	S	22	"					
<u>VES. STOMATITIS GR.</u>																										
Chandipura				+		+												2	S	20	Vesiculovirus					
Cocal	+							+										3	A1	20	"					
Isfahan	+			+														2	S	21	"					
Jurona	+																	2	S	22	"					
La Joya	+																	2	S	22	"					
Piry											+							3	S	22	"					
YS-Alagoas																		3	S	22	"					
YS-Indiana	+			+														2	A3	20	"					
YS-New Jersey	+					+												2	A3	22	"					
Yug Bogdanovac				+		+												3	IE	22	"					

** See footnote Table 5.1

Table 10.2 Family Rhabdoviridae:
Bovine Ephemeral Fever and Other Antigenic Groups

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE	SALS RATING		SEAS RATING**	TAXONOMIC STATUS					
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection			Level	Basis			
	Mosq. Culicine	Anopheleine Ixodid	Ticks Argasid	Phlebotomine	Culicoides Other	Humans Other Primates	Rodents	Birds	Bats	Marsupials Other													Sentinals		
<u>BOVINE EPHEMERAL FEVER GR.</u> Auelaiide River Berrimah Bovine ephem. fev. Kimberley	+				+														X		22 22 22 22	Rhabdovirida " " "			
<u>HART PARK GR.</u> Flanders Hart Park Hosqueiro	+															+					2 2 3	S S IE	22 21 22	Rhabdovirida " "	
<u>KWATTA GR.</u> Kwatta	+																				2	S	22	Rhabdovirida	
<u>LE DANTEC GR.</u> Keuraliba Le Dantec																					2 2	S S	22 22	Rhabdovirida "	
<u>HOSSURIL GR.</u> Bangoran Barur Charleville Cuiaba Kanese Kern Canyon Marco Hossuril	+																					2 2 2 2 2 2 2 2	S S S S S S S S	22 22 22 22 22 23 22 22	Rhabdovirida " " " " " " " "

** See footnote Table 5.1
X See footnote Table 6.1

Table 10.3 Family Rhabdoviridae:
Antigenically Ungrouped Viruses

VIRUS	ISOLATED FROM				ISOLATED IN					HUMAN DISEASE	SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS		VERTEBRATES		Africa	Asia	Australasia	Europe	North America		South America	Level			Basis
	Mosq.	Ticks	Other	Humans						Other Primates			Rodents	Birds	
Aruac	+	+													Rhabdoviridae
Boteke	+	+													"
Gray Lodge	+	+													"
Joinjakaka	+	+													"
Kununurra	+	+													"
Nkolbisson	+	+													"
Rochambeau	+	+													"
Xiburema	+	+													"
Yata	+	+													"
Inhangapi															Rhabdoviridae
Sripur															"
Tibrogargan															"
Almpiwar															Rhabdoviridae
Gossas															"
Klamath															"
Mount Elgon bat															"
Navarro															"

** See footnote Table 5.1

Table 11.1 Arenaviruses, Family Arenaviridae:
Tacaribe (LCM) Serogroup Viruses

VIRUS	ISOLATED FROM				ISOLATED IN						HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS
	ARTHROPODS		VERTEBRATES		Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level	Basis		
	Mosc.	Ticks	Other	Humans	Other Primates	Rodents	Birds	Bats	Marsupials	Other	Sentinels					
Amapari																
Flexal																
Junin																
Ippy																
Lassa																
Latino																
Machupo																
Parana																
Pichinde																
Tacaribe																
Tamiami																

** See footnote Table 5.1

Table 12.1 Families Orthomyxoviridae, Coronaviridae, Herpesviridae, Iridoviridae, Nodaviridae, Paramyxoviridae and Poxviridae: Thogoto Serogroup and Antigenically Ungrouped Viruses

VIRUS	ISOLATED FROM											ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS						VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosq.	Ticks	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats	Marsupials												
THOGOTO GR. Thogoto		+				+							+					+		2	S	21	Orthomyxoviridae
Dhori			+											+						3	S	22	Orthomyxoviridae
Bocas Tettang	+		+						+											2	S	22	Coronavirus "
Agua Preta									+									+		3	IE	22	Herpesviridae
Afr. swine fever				+							+				+	§	α			X		20	Iridoviridae
Nodamura	+													+						3	IE	23	Nodavirus
Nariva							+											+		3	IE	23	Paramyxoviridae
Cotia Oubangui Salanga	+	+		+		+						+						+	+	2 3 3	S IE IE	24 22 22	Poxviridae " "

** See footnote Table 5.1
X See footnote Table 6.1
§ Cuba
α Brazil

Table 13.1 Taxonomically Unclassified Viruses:
Quaranfil, Marburg and Other Antigenic Groups of Viruses

VIRUS	ISOLATED FROM											ISOLATED IN					HUMAN DISEASE		SALS RATING		SEAS RATING**	TAXONOMIC STATUS	
	ARTHROPODS					VERTEBRATES						Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosq. Culicine	Ticks Anopheline Ixodid	Argasid	Phlebotomine	Culicoides Other	Humans Other Primates	Rodents	Birds	Bats	Marsupials	Other Sentinels												
<u>NYAMANINI GR.</u> Nyamanini			+										+						2	S	21	Unclassified	
<u>QUARANFIL GR.</u> Johnston Atoll Quaranfil				+									+		+			+	2 2	S S	20 20	Unclassified "	
<u>MARBURG GR.</u> Ebola Marburg													+			+		+	4 4	S S	23 23	Unclassified "	
<u>TANJONG RABOK GR.</u> Tanjong Rabok Telok Forest														+					2 3	S IE	22 22	Unclassified "	

** See footnote Table 5.1

Table 13.2.1 Taxonomically Unclassified Viruses:
Antigenically Ungrouped Mosquito-, Tick-, or Culicoides-Associated Viruses

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE	SALS RATING		SEAS RATING**	TAXONOMIC STATUS		
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America		South America	Natural Infection			Lab Infection	Level
	Mosq.	Ticks		Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds						Bats			Marsupials	Other		
Arkonam	+	+																				
Gomoka	+	+																	2	S	22	"
Itupiranga	+	+						+											2	S	22	"
Minnal	+	+																	2	S	22	"
Okola	+	+																	2	S	22	"
Para	+	+											+						2	S	22	"
Picola	+	+																	3	IE	22	"
Tanga																			2	S	22	"
Tembe		+																	2	S	22	"
Termeil	+																		3	IE	21	"
Venkatapuram	+																		2	S	22	"
Wongorr	+																		2	S	22	"
Yacaaba	+																		3	IE	22	"
Aride																			2	S	22	Unclassified
Batken	+																		3	IE	22	"
Chim		+																	3	IE	22	"
Estero Real																			3	IE	22	"
Lake Clarendon																					22	"
Matucare																			2	S	22	"
Ngaingan																			2	S	22	"
Slovakia																			3	IE	24	"

** See footnote Table 5.1

Table 13.2.2 Taxonomically Unclassified Viruses:
Antigenically Ungrouped Viruses - No Arthropod Vector Known

VIRUS	ISOLATED FROM										ISOLATED IN					HUMAN DISEASE		SALS RATING		SENS RATING**	TAXONOMIC STATUS	
	ARTHROPODS					VERTEBRATES					Africa	Asia	Australasia	Europe	North America	South America	Natural Infection	Lab Infection	Level			Basis
	Mosq.	Ticks	Phlebotomine	Culicoides	Other	Humans	Other Primates	Rodents	Birds	Bats												
Culicine	Anopheleine	Ixodid	Argasid																			
Araguari																		3	IE	22	Unclassified	
Belem																		3	IE	22		
Bimbo																		3	IE	22		
Kannavanpettai																		2	S	22		
Kannamangalam																		2	S	22		
Kolongo																		2	S	22		
Landjia																		2	S	22		
Mapura																		3	IE	23		
Hojui dos Campos																		3	IE	22		
Ouango																		3	IE	22		
Sakpa																		3	IE	22		
Sandjimba																		2	S	22		
Santarem																		3	IE	22		
Sebokele																		2	S	22		
Sebalam																		2	S	22		
Thottapalayam																		2	S	22		
Toure																		2	S	22		

** See footnote Table 5.1

Table 14.1 Continental Distribution of Grouped and Ungrouped Viruses

Antigenic Group	Total in Group	Africa	Asia	Aus-tral- asia	Eur- ope	North Amer- ica	South Amer- ica	No. of Conti- nents involved					
								1	2	3	4	5	6
A	27	7	8	7	2	7	10	18	7	0	2	0	0
AHS	1	1	1	0	1	0	0	0	0	1	0	0	0
B	66	19	23	14	9	15	12	50	11	2	1	2	0
BAK	2	0	2	0	0	0	0	2	0	0	0	0	0
BEF	4	1	1	4	0	0	0	3	0	1	0	0	0
BHA	1	1	1	0	1	0	0	0	0	1	0	0	0
BLU	1	1	1	1	1	1	0	0	0	0	0	1	0
Bunyamwera Supergroup	ANA	5	0	0	0	0	1	4	5	0	0	0	0
	ANB	2	0	0	0	0	0	2	2	0	0	0	0
	BUN	22	5	1	0	2	8	8	20	2	0	0	0
	BWA	2	2	0	0	0	0	0	2	0	0	0	0
	C	12	0	0	0	0	5	9	10	2	0	0	0
	CAL	13	1	1	0	2	9	3	11	1	1	0	0
	CAP	8	0	0	0	0	3	7	6	2	0	0	0
	GAM	3	0	0	0	0	1	2	3	0	0	0	0
	GMA	12	0	0	0	0	2	11	11	1	0	0	0
	KOO	2	0	0	2	0	0	0	2	0	0	0	0
	MNT	2	0	0	0	0	1	1	2	0	0	0	0
	OLI	3	3	0	0	0	0	0	3	0	0	0	0
	PAT	6	0	0	0	0	4	2	6	0	0	0	0
	SIM	21	10	6	5	0	2	4	16	4	1	0	0
	TETE	5	4	1	0	2	0	0	3	2	0	0	0
TUR	4	1	1	0	1	1	1	3	1	0	0	0	
SBU	1	0	1	0	0	0	0	1	0	0	0	0	
CGL	12	0	0	0	0	1	11	12	0	0	0	0	
CTF	2	0	0	0	1	1	0	2	0	0	0	0	
COR	3	1	0	1	0	0	1	3	0	0	0	0	
EHD	2	1	1	0	0	1	0	1	1	0	0	0	
EUB	3	1	0	2	0	0	0	3	0	0	0	0	
HTN	4	0	2	0	1	2	1	3	0	1	0	0	
HP	3	0	0	0	0	2	1	3	0	0	0	0	
KSO	3	0	2	0	0	1	0	3	0	0	0	0	
KEM	18	3	5	1	6	6	1	15	2	1	0	0	
KWA	1	0	0	0	0	0	1	1	0	0	0	0	
LD	2	2	0	0	0	0	0	2	0	0	0	0	
MAL	2	1	1	0	0	0	0	2	0	0	0	0	
MAP	4	0	0	4	0	0	0	4	0	0	0	0	
MBG	2	2	0	0	1	0	0	1	1	0	0	0	
MOS	8	4	1	1	0	1	2	7	1	0	0	0	

Table 14.1 (Continued) Continental Distribution of Grouped and Ungrouped Viruses

Antigenic Group	Total in Group	Africa	Asia	Aus-tral- asia	Eur- ope	North Amer- ica	South Amer- ica	No. of Conti- nents involved						
								1	2	3	4	5	6	
MTY	3	3	0	0	0	0	0	3	0	0	0	0	0	0
Nairo- viruses	CHF-CON	4	2	4	0	2	0	2	0	2	0	0	0	0
	DGK	5	2	4	1	0	0	3	2	0	0	0	0	0
	HUG	4	1	1	0	1	1	3	2	1	1	0	0	0
	NSD	3	2	1	0	0	0	0	3	0	0	0	0	0
	QYB	3	3	0	0	0	0	0	3	0	0	0	0	0
	SAK	5	0	2	1	1	1	0	5	0	0	0	0	0
NDO	1	1	0	0	0	0	0	1	0	0	0	0	0	
NYM	1	1	0	0	0	0	0	1	0	0	0	0	0	
PAL	7	0	3	4	0	0	0	7	0	0	0	0	0	
PHL	36	8	5	0	5	10	14	32	2	2	0	0	0	
QRF	2	1	0	1	0	0	0	2	0	0	0	0	0	
RABIES	2	2	0	0	0	0	0	2	0	0	0	0	0	
RTA	3	0	0	0	0	0	3	3	0	0	0	0	0	
SAW	3	0	0	0	0	3	0	3	0	0	0	0	0	
TCR	11	2	0	0	0	1	8	11	0	0	0	0	0	
THO	1	1	1	0	1	0	0	0	0	1	0	0	0	
TIM	3	0	0	0	0	0	3	3	0	0	0	0	0	
TR	2	0	2	0	0	0	0	2	0	0	0	0	0	
UMA	2	0	0	0	0	2	0	2	0	0	0	0	0	
UPO	2	0	0	1	0	1	0	2	0	0	0	0	0	
UUK	6	0	2	1	3	0	0	6	0	0	0	0	0	
VSV	10	1	2	0	1	3	6	7	3	0	0	0	0	
WAL	1	0	0	1	0	0	0	1	0	0	0	0	0	
WAR	2	0	0	2	0	0	0	2	0	0	0	0	0	
YOG	1	1	0	0	0	0	0	1	0	0	0	0	0	
Ungrouped	89	29	18	14	4	9	22	85	2	1	1	0	0	
TOTAL	506	131	104	68	48	106	153	435	48	16	4	3	0	

Table 15.1 Number of Viruses Isolated from Classes of Wild-Caught Arthropods

Antigenic Group	Total in Group	Isolated From:					No. of Classes Involved		
		Mosq.	Ticks	Phlebotomine Flies	Culicoides	Other	1	2	3
A	27	25	3	0	1	5	20	4	2
AHS	1	0	0	0	1	0	1	0	0
B	66	32	20	0	1	3	41	7	1
BAK	2	2	1	0	0	0	1	1	0
BEF	4	2	0	0	2	0	0	2	0
BHA	1	0	1	0	0	0	1	0	0
BLU	1	0	1	0	1	0	0	1	0
Bunyamvera Serogroup	ANA	5	5	0	0	0	5	0	0
	ANB	2	2	0	0	0	2	0	0
	BUN	22	22	0	0	2	20	2	0
	BWA	2	2	0	0	0	2	0	0
	C	12	12	0	0	0	12	0	0
	CAL	13	13	0	0	0	12	1	0
	CAP	8	7	0	0	0	7	0	0
	GAM	3	3	0	0	0	3	0	0
	GMA	12	9	0	1	0	8	1	0
	KOO	2	2	0	0	0	2	0	0
	MNT	2	1	0	0	0	1	0	0
	OLI	3	3	0	0	0	3	0	0
	PAT	6	6	0	0	0	6	0	0
	SIM	21	10	0	0	11	11	5	0
	TETE	5	0	2	0	0	2	0	0
TUR	4	4	0	0	0	4	0	0	
SBU	1	0	0	0	0	1	0	0	
CGL	12	1	0	8	0	0	9	0	0
CTF	2	0	2	0	0	0	2	0	0
COR	3	3	0	0	0	0	3	0	0
EHD	2	0	0	0	0	0	0	0	0
EUB	3	3	0	0	1	0	2	1	0
HTN	4	0	0	0	0	0	0	0	0
HP	3	3	0	0	0	0	3	0	0
KSO	3	0	3	0	0	0	3	0	0
KEM	18	0	18	0	0	0	18	0	0
KWA	1	1	0	0	0	0	1	0	0
LD	2	0	0	0	0	0	0	0	0
MAL	2	2	0	0	0	0	2	0	0
MAP	4	4	0	0	0	0	4	0	0
MBG	2	0	0	0	0	0	0	0	0
MOS	8	4	1	1	0	0	4	1	0

Table 15.1 (Continued) Number of Viruses Isolated from Wild-Caught Arthropods

Antigenic Group	Total in Group	Isolated From:					No. of Classes Involved			
		Mosq.	Ticks	Phleboto- mine Flies	Culi- coides	Other	1	2	3	
MTY	3	0	0	0	0	0	0	0		
Nairo- viruses	CHF-CON	4	0	4	0	1	0	3	1	0
	DGK	5	0	5	0	0	0	5	0	0
	HUG	4	0	4	0	0	0	4	0	0
	NSD	3	2	3	0	2	0	0	2	1
	QYB	3	0	2	0	0	0	2	0	0
	SAK	5	0	5	0	0	0	5	0	0
NDO	1	1	0	0	0	0	1	0	0	
NYM	1	0	1	0	0	0	1	0	0	
PAL	7	3	0	0	4	0	7	0	0	
PHL	36	6	0	22	0	0	24	2	0	
QRF	2	0	2	0	0	0	2	0	0	
RABIES	2	0	0	0	1	0	1	0	0	
RTA	3	3	0	0	0	0	3	0	0	
SAW	3	0	3	0	0	0	3	0	0	
TCR	11	1	1	0	0	3	3	1	0	
THO	1	0	1	0	0	0	1	0	0	
TIM	3	0	0	0	0	0	0	0	0	
TR	2	0	0	0	0	0	0	0	0	
UMA	2	2	0	0	0	0	2	0	0	
UPO	2	0	2	0	0	0	2	0	0	
UUK	6	0	6	0	0	0	6	0	0	
VSV	10	6	1	4	1	2	4	2	2	
WAL	1	0	0	0	1	0	1	0	0	
WAR	2	1	0	0	2	0	1	1	0	
YOG	1	0	0	0	0	0	0	0	0	
Ungrouped	89	41	18	3	2	1	54	5	0	
TOTAL	506	249	110	39	34	16	351	40	6	

Table 16.1 Number of Virus Isolated from Classes of Naturally-Infected Vertebrates

Anti-genic Group	Total in Group	Humans	Other Primates	Rodents	Birds	Bats	Marsu-pials	Live-stock	All Others	No. of Classes Involved					
										1	2	3	4	5	6
A	27	11	2	6	13	4	6	6	3	8	3	4	3	1	1
AHS	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0
B	66	29	5	19	17	13	1	6	7	28	6	8	4	2	1
BAK	2	0	1	0	0	0	0	0	0	1	0	0	0	0	0
BEF	4	0	0	0	0	0	0	4	0	4	0	0	0	0	0
BHA	1	1	0	1	0	0	0	1	1	0	0	1	0	0	0
BLU	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0
ANA	5	1	1	0	0	0	0	0	0	0	1	0	0	0	0
ANB	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
BUN	22	5	1	7	1	0	0	2	4	7	5	0	0	0	0
BWA	2	2	0	0	0	0	0	0	0	2	0	0	0	0	0
C	12	10	0	8	0	1	5	0	1	2	5	3	1	0	0
CAL	13	3	0	4	0	0	0	0	1	4	2	0	0	0	0
CAP	8	0	0	4	0	0	1	0	0	3	1	0	0	0	0
GAM	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
GMA	12	2	0	8	2	2	4	0	0	5	1	1	2	0	0
KOO	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MNT	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
OLI	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
PAT	6	0	0	3	0	0	0	0	0	3	0	0	0	0	0
SIM	21	2	1	0	4	0	0	8	4	13	3	0	0	0	0
TETE	5	0	0	0	5	0	0	0	0	5	0	0	0	0	0
TUR	4	0	0	0	2	0	0	0	1	1	1	0	0	0	0
SBU	1	0	0	0	0	1	0	0	0	1	0	0	0	0	0
CGL	12	1	0	1	0	0	0	0	2	4	0	0	0	0	0
CTF	2	1	0	1	0	0	0	0	1	0	0	1	0	0	0
COR	3	0	0	0	1	0	0	0	0	1	0	0	0	0	0
EHD	2	0	0	0	0	0	0	1	1	2	0	0	0	0	0
EUB	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HTN	4	1	0	4	0	0	0	0	0	3	1	0	0	0	0
HP	3	0	0	0	2	0	0	0	0	2	0	0	0	0	0
KSO	3	0	1	0	1	0	0	0	1	3	0	0	0	0	0
KEM	18	1	0	1	1	0	0	1	0	0	2	0	0	0	0
KWA	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
LD	2	1	0	1	0	0	0	0	0	2	0	0	0	0	0
MAL	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MAP	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MBG	2	2	0	0	0	0	0	0	0	2	0	0	0	0	0
MOS	8	0	0	1	2	1	0	0	3	7	0	0	0	0	0

Table 16.1 (Continued) Number of Viruses Isolated from Naturally-Infected Vertebrates

Anti- genic Group	Total in Group	Humans	Other Primates	Rodents	Birds	Bats	Marsu- pials	Live- stock	All Others	No. of Classes Involved					
										1	2	3	4	5	6
MTY	3	0	0	0	3	0	0	0	0	3	0	0	0	0	0
Nairo- viruses CHF-CON	4	2	0	0	0	0	0	1	1	1	1	0	0	0	0
DGK	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HUG	4	0	0	0	1	0	0	0	0	1	0	0	0	0	0
NSD	3	3	0	1	1	0	0	2	1	1	1	0	1	0	0
QYB	3	0	0	2	0	0	0	0	0	2	0	0	0	0	0
SAK	5	0	0	0	1	0	0	0	0	1	0	0	0	0	0
NDO	1	1	0	0	0	0	0	0	0	1	0	0	0	0	0
NYM	1	0	0	0	1	0	0	0	0	1	0	0	0	0	0
PAL	7	0	0	0	0	0	0	4	0	4	0	0	0	0	0
PHL	36	9	0	9	3	0	2	1	3	15	4	1	0	0	0
QRF	2	1	0	0	1	0	0	0	0	0	1	0	0	0	0
RABIES	2	0	0	0	0	1	0	0	0	1	0	0	0	0	0
RTA	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
SAW	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TCR	11	3	0	10	0	1	0	0	1	8	2	1	0	0	0
THO	1	1	0	0	0	0	0	1	0	0	1	0	0	0	0
TIM	3	0	0	0	0	0	0	0	3	3	0	0	0	0	0
TR	2	0	1	0	0	0	0	0	0	1	0	0	0	0	0
UMA	2	0	0	0	1	0	0	0	0	1	0	0	0	0	0
UPO	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
UUK	6	0	0	1	1	0	0	0	0	0	1	0	0	0	0
VSV	10	4	0	1	0	0	1	3	2	1	5	0	0	0	0
WAL	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
WAR	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
YOG	1	0	0	0	0	1	0	0	0	1	0	0	0	0	0
Ungrouped	89	8	2	8	12	10	1	1	3	39	2	0	0	0	0
TOTAL	506	105	15	101	76	35	21	44	44	200	49	19	12	3	2

Table 17.1 Number of Viruses Associated with Naturally- or Laboratory-Acquired Disease in Humans

Antigenic Group	Total in Group	In Nature	Lab. Infection	Either or Both	
				Number	Percent
Group A	27	11	7	12	45.2
Afr. horsesickness	1	0	0	0	
Group B	66	31	26	33	50.0
Bakau	2	0	0	0	
Bhanja	1	0	1	1	100.0
Bluetongue	1	0	1	1	100.0
Bovine ephem. fever	4	0	0	0	
Anopheles A	5	1	0	2	100.0
Anopheles B	2	0	0	0	
Bunyamwera	22	6	2	7	31.8
Bwamba	2	2	0	2	100.0
C	12	10	6	10	83.3
California	13	7	0	7	53.8
Capim	8	0	0	0	
Gamboa	3	0	0	0	
Guama	12	2	1	2	16.7
Koongol	2	0	0	0	
Minatitlan	2	0	0	0	
Olifantsvlei	3	0	0	0	
Patos	6	0	0	0	
Simbu	21	2	1	2	9.5
Tete	5	0	0	0	
Turlock	4	0	0	0	
SBU	1	0	0	0	
Changuinola	12	1	0	1	8.3
Colorado tick fever	2	1	1	1	50.0
Corriparta	3	0	0	0	
Epizoot. hem. dis.	2	0	0	0	
Eubenangee	3	0	0	0	
Hantaan	4	3	3	3	75.0
Hart Park	3	0	0	0	
Kaisodi	3	0	0	0	
Kemerovo	18	2	2	2	11.1
Kwatta	1	0	0	0	
Le Dantec	2	1	0	1	50.0
Malakal	2	0	0	0	
Mapputta	4	0	0	0	
Marburg	2	2	2	2	100.0
Matariya	3	0	0	0	

Table 17.1 (Continued) Number of Viruses Associated with Naturally- or Laboratory-Acquired Disease in Humans

Antigenic Group	Total in Group	In Nature	Lab. Infection	Either or Both	
				Number	Percent
Mossuril	8	0	0	0	
Nairo- viruses	CHF-Congo	4	2	2	50.0
	Dera Ghazi Khan	5	0	0	
	Hughes	4	0	0	
	Nairobi sheep dis.	3	3	3	100.0
	Qalyub	2	0	0	
	Sakhalin	5	0	0	
Nyando	1	1	0	1	100.0
Nyamanini	1	0	0	0	
Palyam	7	0	0	0	
Phlebotomus fever	36	9	1	9	25.0
Quaranfil	2	1	0	1	50.0
Rabies	2	0	0	0	
Resistencia	3	0	0	0	
Sawgrass	3	0	0	0	
Tacaribe	11	3	6	6	54.5
Tanjong Rabok	2	0	0	0	
Thogoto	1	1	0	1	100.0
Timbo	3	0	0	0	
Umatilla	2	0	0	0	
Upolu	2	0	0	0	
Uukuniemi	6	0	0	0	
Vesicular stom.	10	4	3	5	50.0
Wallal	1	0	0	0	
Warrego	2	0	0	0	
Yogue	1	0	0	0	
Ungrouped	89	7	2	7	7.8
TOTAL	506	113	70	123	24.3

Table 18.1 Evaluation of Arthropod-Borne Status of 506 Registered Viruses (SEAS)

Anti- genic Group	Total in Group	Arbo- virus	Prob- ably Arbo- virus	Pos- sible Arbo- virus	Prob- ably not Arbo- virus	Not Arbo- virus	Arbo or Probably Arbo		Not or Probably Not Arbo	
							No.	%	No.	%
A	27	16	5	6	0	0	21	80.8	0	
AHS	1	1	0	0	0	0	1	100.0	0	
B	66	34	8	17	2	5	42	63.6	7	10.6
BAK	2	0	1	1	0	0	1	50.0	0	
BEF	4	0	0	4	0	0	0		0	
BHA	1	0	1	0	0	0	1	100.0	0	
BLU	1	1	0	0	0	0	1	100.0	0	
Bunyamwera Serogroup	ANA	5	0	2	3	0	2	40.0	0	
	ANB	2	0	0	2	0	0		0	
	BUN	22	8	7	7	0	0	15	68.2	0
	BWA	2	1	1	0	0	0	2	100.0	0
	C	12	10	2	0	0	0	12	100.0	0
	CAL	13	10	1	2	0	0	11	84.6	0
	CAP	8	4	2	2	0	0	6	75.0	0
	GAM	3	0	1	2	0	0	1	33.3	0
	GMA	12	5	4	3	0	0	9	75.0	0
	KOO	2	0	2	0	0	0	2	100.0	0
	MNT	2	0	1	1	0	0	1	50.0	0
	OLI	3	0	0	3	0	0	0		0
	PAT	6	1	2	3	0	0	3	50.0	0
	SIM	21	3	5	13	0	0	8	38.1	0
	TETE	5	0	1	4	0	0	1	20.0	0
TUR	4	1	2	1	0	0	3	75.0	0	
SBU	1	0	0	1	0	0	0		0	
CGL	12	0	1	11	0	0	1	8.3	0	
CTF	2	1	0	1	0	0	1	50.0	0	
COR	3	0	1	2	0	0	1	33.3	0	
EHD	2	0	1	1	0	0	1	50.0	0	
EUB	3	0	0	3	0	0	0		0	
HTN	4	0	0	1	3	0	0		3	75.0
HP	3	0	1	2	0	0	1	33.3	0	
KSO	3	0	2	1	0	0	2	66.7	0	
KEM	18	0	3	15	0	0	3	16.7	0	
KWA	1	0	0	1	0	0	0		0	
LD	2	0	0	2	0	0	0		0	
MAL	2	0	0	2	0	0	0		0	
MAP	4	0	1	3	0	0	1	25.0	0	
MBG	2	0	0	0	2	0	0		2	100.0
MOS	8	0	0	7	1	0	0		1	12.5

Table 18.1 (Continued) Evaluation of Arthropod-Borne Status of 506 Registered Viruses (SEAS)

Anti- genic Group	Total in Group	Arbo- virus	Prob- ably Arbo- virus	Pos- sible Arbo- virus	Prob- ably not Arbo- virus	Not Arbo- virus	Arbo or Probably Arbo		Not or Probably Not Arbo		
							No.	%	No.	%	
MTY	3	0	0	3	0	0	0		0		
Nairo- viruses	CHF-CON	4	2	0	2	0	0	2	50.0	0	
	DGK	5	0	0	5	0	0	0		0	
	HUG	4	1	1	2	0	0	2	50.0	0	
	NSD	3	1	1	1	0	0	2	66.7	0	
	OYB	3	1	0	2	0	0	1	33.3	0	
	SAK	5	0	2	3	0	0	2	20.0	0	
NDO	1	0	1	0	0	0	1	100.0	0		
NYM	1	0	1	0	0	0	1	100.0	0		
PAL	7	0	2	5	0	0	2	28.6	0		
PHL	36	4	13	19	0	0	17	47.2	0		
QRF	2	2	0	0	0	0	2	100.0	0		
RABIES	2	0	1	0	0	1	1	50.0	1	50.0	
RTA	3	0	0	3	0	0	0		0		
SAW	3	0	0	3	0	0	0		0		
TCR	11	0	0	1	1	9	0		10	90.9	
THO	1	0	1	0	0	0	1	100.0	0		
TIM	3	0	0	3	0	0	0		0		
TR	2	0	0	2	0	0	0		0		
UMA	2	1	1	0	0	0	2	100.0	0		
UPO	2	0	0	2	0	0	0		0		
UUK	6	2	0	4	0	0	2	33.3	0		
VSV	10	3	1	6	0	0	4	40.0	0		
WAL	1	0	0	1	0	0	0		0		
WAR	2	0	0	2	0	0	0		0		
YOG	1	0	0	1	0	0	0		0		
Ungrouped	89	3	9	69	5	3	12	13.4	8	9.0	
TOTAL	506	116	92	266	14	18	208	41.1	32	6.3	

APPENDIX I

Summary Description of Recommended Practice and Containment Levels for Arboviruses and Certain Other Viruses of Vertebrates^a (11).

Level	Laboratory Practices	Primary Containment	Secondary Containment
1	Standard microbiological practices are required.	None. Open bench.	None required.
2	Care required to limit aerosols and contamination. Limited access. ^c	Class I or II BSC ^b required for aerosol producing procedures.	Designed to facilitate cleaning and disinfection.
3	All virus materials contained. Special lab gowns required.	Class I or II BSC or equivalent required for all manipulations of infectious materials.	Restricted access, ^d air lock facility, controlled unidirectional air flow. Exhaust air discharged away from building. Work with certain viruses indicated by an * requires HEPA filtration of exhaust air.
4	Rigorous containment of all virus manipulations. Change of clothing and shower required.	Class I or II BSC adequate for work with infectious materials if all laboratory personnel are immune or insusceptible. Otherwise, Class III or one-piece positive pressure suits are required.	Facility equivalent to separate building. Includes shower facilities, heat-treated biowaste, HEPA filtration of all exhaust air, double-door autoclaves.

^aThere are also SALS recommendations concerning vector and vertebrate studies.

^bBSC = Biological Safety Cabinets.

^cAccess limited to persons with knowledge of the biohazard potential.

^dAccess restricted to persons with programmatic or support requirements for entry.

APPENDIX II

Explanation of Symbols Used to Define Basis for Assignment of Viruses to Levels of Practice and Containment (11).

- S = Results of SALS surveys and information from the Catalogue.
- IE = Insufficient experience with virus; i.e., experience factor from SALS surveys was less than 500 in laboratory facilities with low biocontainment.
- A = Additional criteria 1, 2, 3, 4, etc.
1. Disease in sheep, cattle or horses.
 2. Fatal human laboratory infection, 1978, probably aerosol (14). This is recognized to be a unique incident in a long history of work with SFV under minimal biocontainment conditions. However, since the virulence characteristics of the strain responsible in this case require further study and the prevalence of subclinical infections in laboratories working with SFV remains unknown, the committee recommends Level 3 until further information is available warranting reconsideration at a lower level.
 3. Extensive laboratory experience and mild nature of aerosol laboratory infections justifies Level 2.
 4. Placed in Level 4 based on the close antigenic relationship with a known Level 4 agent, Russian spring-summer encephalitis, plus insufficient laboratory experience.
 5. Level 2 arenaviruses are not known to cause serious acute disease in man and are not acutely pathogenic for laboratory animals, including primates. Survey experience is sufficient to conclude that laboratory aerosol infection does not occur in the course of routine work with cell cultures and animals not subject to chronic infection. In view of a reported high frequency of laboratory aerosol infection that occurred in workers manipulating high concentrations of Pichinde virus, it is strongly recommended that work with high concentrations of Level 2 arenaviruses be done at Level 3.
 6. Level assigned to prototype or wild-type virus. A lower level may be recommended for laboratory strains or geographic variants of the virus with well-defined reduced virulence characteristics, as mentioned in the text.

VECTOR INDEX

- Aedeomyia catasticta*: ALF, COR, KNA
Aedeomyia squamipennis: GAM, PV, SJ
Aedes abnormalis: BOU, MOS, NDU, MID, PGA, SF, SPO, WSL
Aedes abserratus: JC
Aedes aegypti: CHIK, DEN-1, DEN-2, DEN-3, DEN-4, DUG, ORU, USU, VEE, WN, WSL, YF, ZIKA
Aedes africanus: BOU, BUN, CHIK, DEN-2, MID, ORU, SAB, WSL, WN, YF, ZIKA
Aedes albocephalus: MID, WN
Aedes albopictus: DEN-2
Aedes albothorax: WN
Aedes and *Anopheles*: GET
Aedes and *Culiseta*, mixed pool: INK
Aedes and *Psorophora*, mixed pool: ILH
Aedes angustivittatus: SAR, VEE
Aedes arborealis: APEU
Aedes argenteopunctatus: BUN, GOM, MID, PGA, SF, SHO, WSL
Aedes argenteopunctatus and *Ae mutilis*: NKO
Aedes argyrothorax: WYO
Aedes atlanticus: EEE, LAC, TEN, TVT
*Aedes atlanticus/tormentor**: EVE, KEY, LAC, TEN, TVT
Aedes aurifer: KEY, SSH
Aedes bancroftianus: BF
Aedes butleri: BEB
Aedes caballus: MID, RVF, WSL
Aedes campestris: CV, WEE
Aedes camptorhynchus: TER
Aedes canadensis: CV, EEE, JC, KEY, LAC, SSH
Aedes cantans: TET, WN
Aedes cantator: JC
Aedes capensis: BUN
Aedes caspius: IK, ISF, TAH
Aedes caspius caspius and *Culex hortensis*: BKN
Aedes cataphylla: SSH
Aedes cinereus: CV, SSH
Aedes circumluteolus: BUN, GER, ING, LEB, MID, NDU, PGA, RVF, SHO, SIM, SPO, WSL, WN
Aedes communis: CV, JC, LAC, SSH, TAH, TVT
Aedes communis/punctor: INK
Aedes cumminsii: DEN-2, MID, NKO, PGA, RVF, SHO, SIM, SPO, WSL
Aedes dalzieli: BUN, CHIK, MID, NDU, NDO, PGA, SHO, SIM, WSL, ZIKA, ZGA
Aedes dentatus: MID, ORU, PGA, SF, SHO, WSL, YF
Aedes diantaeus: TAH
Aedes domesticus: NKO, WSL
Aedes dorsalis: CV, CE, LAC, MD, TVT, WEE
*Aedes eidsvoldensis** and *Ae pseudonormanensis*: GG
Aedes excrucians: NOR, SSH
Aedes fitchii: SSH
Aedes flavescens: CV, WEE
Aedes flavicollis: ZIKA
Aedes flavifrons: RR
Aedes fowleri: PGA, SIM
Aedes fryeri/fowleri: SPO
Aedes fulvus: MAG, SDN, WYO
Aedes fulvus pallens: EEE
Aedes funereus: MPK
Aedes furcifer: BOU, BWA, CHIK, DEN-2, ORU, RVF, YF, ZIKA
Aedes furcifer/taylori: BOU, BUN, CHIK, DEN-2, ORU, YF, ZIKA
Aedes hexodontus: NOR, SSH
Aedes infirmatus: KEY, TEN, TVT
Aedes intrudens: SSH
Aedes irritans: CHIK
Aedes jamoti: SF, ZIKA
Aedes lineatopennis: MID, RVF, TMU, WAL, WSL, WGR
Aedes lineatopennis/Ae albothorax pool: MID
Aedes luteocephalus: BOU, CHIK, DEN-2, NKO, PGA, WSL, YF, ZIKA

* Currently considered to be a subspecies of *Aedes theobaldi*.

Aedes marshallii: MID
Aedes mediolineatus: WSL
Aedes melanimon: CE, JC, SLE, WEE
Aedes melanimon-dorsalis: CE
Aedes metallicus: YF
Aedes minutus: NDU, WSL
Aedes mitchellae: EEE, TEN
Aedes neoafricanus: CHIK, YF,
 ZIKA
Aedes nigripes: SSH
Aedes nigromaculis: CV, CE, EEE,
 MD, WEE
*Aedes nocturnus**: BAT
Aedes normanensis: BF, EH, GG,
 KOK, MVE, RR, SIN, WGR
Aedes opok: BOU, BUN, CHIK,
 DEN-2, MID, ORU, YF, WSL, ZIKA
Aedes palpalis: MID, PATA, SF,
 SIM, ZGA
Aedes pembaensis: BUN, TAH
Aedes polynesiensis: DEN-1, RR
Aedes procax: BF
Aedes punctor: BAT, JC, NOR, SSH
Aedes scapularis: CV, CAR, ILH,
 KRI, LUK, MAG, MAN, MAY, MEL,
 SLE, VEE, WYO
Aedes septemstriatus: APEU, WYO
Aedes serratus: AURA, COT, GTB,
 ILH, ITU, MAG, MEL, MIR, ORO,
 SLE, UNA, VEE, WYO
Aedes sexlineatus: WYO
Aedes simpsoni: YF
Aedes simulans: MID, WSL
Aedes sollicitans: CV, JC, KEY,
 LAC, VEE
Aedes species: BUN, CHIK, GMA,
 ICO, JC, KEY, KOK, MAG, MUC,
 NOR, ORI, SIN,** SSH, TVT,
 UGS, UNA, VSI, WSL, WEE, ZIKA
Aedes (Adm) species: MID, WSL
Aedes (Can) species: JUG
Aedes (Dic) species: ORU
Aedes (Neo) species: WSL
Aedes (Och) species: UNA, WYO
Aedes (Ver) species: BF
Aedes spencerii: CV, WEE
Aedes sticticus: EEE, TAH
Aedes stimulans: CV, JC, SSH
Aedes stokesi: NKO
Aedes taeniorhynchus: CV, EEE,
 EVE, GL, KRI, KEY, NOR, ORI,
 PLA, TEN, TLA, TVT, VEE, WYO
Aedes tarsalis: MID, PATA, PGA,
 SHO, WSL, ZIKA
Aedes taylori: DEN-2, YF, ZIKA
Aedes thelcter: JC, VEE
Aedes theobaldi: GG, SIN
Aedes togoi: POW
Aedes tremulus: KUN
Aedes triseriatus: CV, JC, KEY,
 LAC, SSH, TVT
Aedes trivittatus: BOC, CV, JC,
 LAC, SSH, TVT
Aedes vexans: BAT, CV, CE, EEE,
 GET, JC, KEY, LAC, MD, SAG,
 SLE, SF, SSH, TAH, TVT, WEE
Aedes vexans nipponii: AKA, GET
Aedes vigilax: BF, DEN-1, DEN-4,
 EH, GG, KOK, RR, SIN, STR,
 TER, YAC
Aedes vittatus: BUN, CHIK, MID,
 PGA, SAB, SF, SIM, SIN, WSL,
 YF, ZIKA
Aedes vittiger: SIN
Alectorobius capensis†: SOL
Alectorobius sonrai§: CHIK, BDA,
 KOU
Alveonassus lahorensis@: CON, CHF
Amblyomma americanum: LS
Amblyomma cajennense: WM
Amblyomma cohaerens: DUG
Amblyomma loculosum: ARI
Amblyomma pomposum: DUG

* Currently considered to be a subspecies of *Ae vexans*.

**Virus of Karelian fever. About 80% of mosquitoes were *Ae communis*.

† Considered by most to be *Ornithodoros (Alectorobius) capensis*.

§ Considered by most to be *Ornithodoros (Alectorobius) sonrai*.

@ Considered by most to be *Ornithodoros (Alveonassus) lahorensis*.

Amblyomma species: DUG
Amblyomma variegatum: BHA, BLU, CON, DUG, KAS, MID, RVF, THO, WN, YF
Anopheles albimanus: TLA
Anopheles albitarsis: LM
Anopheles amictus: EH, GET, KOW, MAP, RR
Anopheles annularis: JBE
Anopheles annulipes: KOW, MAP, TIL, TRU
Anopheles aquasalis: VEE
Anopheles bancroftii: BEF, KOO
Anopheles barbirostris: JBE
Anopheles boliviensis: ANA, ANB
Anopheles bradleyi-crucians: CV
Anopheles braziliensis: TON
Anopheles brohieri: SHO
Anopheles coustani: CHIK, PGA, WN
Anopheles crucians: CV, EEE, EVE, KEY, LAC, SR, SLE, TEN, TVT, VEE
Anopheles cruzii: BOR, GTB, ICO, TCM
Anopheles farauti: EUB, KOO
Anopheles flavicosta: MID
Anopheles franciscanus: MD
Anopheles freeborni: CV, MD, VR, WEE
Anopheles funestus: AKA, BWA, CHIK, NDO, ONN, ORU, PGA, SF, TAN, TAT, WSL
Anopheles gambiae: BWA, CHIK, ILE, MID, NDO, ONN, ORU, TAT, ZIKA
Anopheles gambiae and *Anopheles pharoensis*: WSL
Anopheles grabhamii: CV
Anopheles hyrcanus: ARK, IK, JBE, POW, TAH
Anopheles maculipennis: BAT, CVO, WN
Anopheles mediopunctatus: TON
Anopheles meraukensis: EH, MAP, WAR
Anopheles neivai: ANA, ANB, GRO, YF
Anopheles neomaculipalpus: VEE
Anopheles nili: BGI, PGA, TAT
Anopheles nimbus: CATU, LUK, PIX, TME, WYO
Anopheles paludis: BOU, GOM
Anopheles pharoensis: BIR, SIN
Anopheles philippinensis: TMU
Anopheles pseudopunctipennis
pseudopunctipennis: CV, SA, VEE
Anopheles punctimacula: VEE
Anopheles punctipennis: CV, JC, LAC, SSH, TEN, VEE
Anopheles punctipennis and *Anopheles quadrimaculatus*: CV
Anopheles quadrimaculatus: CV, JC, SLE, TEN
Anopheles species: CV, MAG, MPK, TLA, UNA, WN, WEE, WYO
Anopheles squamosus: BIR
Anopheles subpictus: ARK, BAT, WN
Anopheles tessellatus: BAT
Anopheles vagus: JBE
Anopheles walkeri: CV
Anophelines: GRO
Anthomyidae: VSNJ
Argas africolumbae: PRE
Argas abdussalami: BAK, MWA
Argas arboreus: NYM, PS, QRF
Argas cooleyi: ML, SC, SCA
Argas hermanni: AH, CNU, GA, QRF, RF, UUK, WN
Argas peringueyi: CNU
Argas persicus: CHF, SLO
Argas pusillus: KTR
Argas reflexus: CNU, GA, NYM, PTV, QRF, UUK, WN
Argas robertsi: KS, LC, PTH
Argas vespertilionis: IK
Argas vulgaris: QRF
Argas walkerae: NYM
Armigeres species: SEP
Bdellonyssus: WEE
Bdellonyssus bursa: SIN
Boophilus annulatus: BHA, CON, DUG, THO
Boophilus calcaratus: CHF

Boophilus decoloratus: BHA, CON, DUG, THO
Boophilus decoloratus, Rhipicephalus appendiculatus, R. evertsi, R. simus: THO
Boophilus geigy: CON, DUG
Boophilus microplus: SEL
Boophilus microplus and Hyalomma a. anatolicum: WM
Chloropidae: VSNJ
Chrysops cincticornis: JC
Chrysops obsoletus: KEY
Cimex insuetus: KK
Coquillettidia: ILH, UNA
Coquillettidia albicosta: RBU, TON
Coquillettidia arribalzagai: UNA, WYO
Coquillettidia aurites: TAT, USU
Coquillettidia crassipes: WON
Coquillettidia fraseri: SIM
Coquillettidia fuscopennata: SIN, YF
Coquillettidia linealis: RR, TRU
Coquillettidia linealis, Culex molestus and Cx cylindricus: BF
Coquillettidia maculipennis: BTK
Coquillettidia metallica: MID, WN
Coquillettidia perturbans: CV, EEE, JC, TEN, TVT
Coquillettidia richiardii: CVO
Coquillettidia venezuelensis: ANU, BSQ, CAB, COT, ITP, MAY, MOJU, MUR, ORO, SLE, TON
Coquillettidia xanthogaster: DEN-4
Culex accelerans: BSB, NEP
Culex adamesi: ABR, NJL
*Culex aikenii**: APEU, MTB, MUR, VEE
Culex albinensis: SLE
Culex albiventris: MPO
Culex amazonensis: ACA, MOR, WYO
Culex annulirostris: BF, BEL, COR, DEN-4, EH, EUB, GG, KIM, KOK, KOO, KOW, KUN, MVE, PR, PIA, RR, SIN, TER, TRU, UMB, WAR, WON, WGR
Culex annulioris: KAM, MID
Culex antennatus: AMT, PGA, SIN, WN
Culex antennatus and Cx univittatus neavi: ACD
Culex australicus: KUN, MVE, SIN
Culex Belem complex No. 19: ANU
Culex Belem species No. 11: BEN
Culex (Mel) Belem species No. 1: BSQ, CAP
Culex (Mel) Belem species No. 17: ACA
Culex (Mel) Belem species No. 27: ANU
Culex bitaeniorhynchus: BAT, GET, MVE, SIN, UMB
Culex caudelli: ITP, SLE
Culex cinereus: MPO
Culex cinereus and Cx albiventris: MID
Culex corniger: VEE
Culex coronator: SLE
Culex crybda: BSQ
Culex (Cux) species: BUN, SLE, VEE
Culex decens: BAG, KAM, MOS, MPO, WN
Culex declarator: SLE, TUR
Culex delpontei: ANT, BQS, RTA, VEE
Culex dunni: EEE, LJ, PCA, WEE
Culex eastor: ITP
Culex edwardsi: COR, SIN
Culex elevator: BOC
Culex epanatasis: ENS
Culex erraticus: SLE, WEE
Culex fuscocephala: JBE
Culex gelidus: BAT, GET, JBE, TMU
Culex gelidus, Cx pseudovishnui and Cx tritaeniorhynchus: TMU

* *Culex (Mel) aikenii* is a synonym of *Cx ocosa*.

Culex guiarti: BAG, BOT, ING,
MID, NTA, OUB, WN
Culex ingrami and *Cx guiarti*:
BAG
Culex iolambdis: PAT
Culex (Lop) species: BAK, BEB,
KET
Culex Malaysia species No. 1: BAK
Culex Malaysia species No. 3: BAK
Culex (Mel) species: ANU, AURA,
BSQ, EVE, GL, JAC, KEY, MH,
MTB, MIR, MOJU, NJL, PAH, SR,
VEE
Culex mixed: See *Culex species*
Culex modestus: KYZ, LED, WN
*Culex molestus**: WN
Culex nakuruensis: BAN
Culex neavei: SPO
Culex nebulosus: MID, MPO, NTA
Culex nigripalpus: CAB, EEE,
EVE, HP, KEY, PAH, SR, SLE,
TEN, TVT, VEE, VSNJ, WYO
Culex ocosa: BAB, PARA, VEE, WEE
*Culex opisthopus***: PAT
Culex orostiensis: COR
Culex panocossa: EEE
Culex paracrybda: ABR, PLS
Culex paracrybda and *Cx*
ocossa: BAB
Culex perfuscus: BAG, BGN, GOM,
KAM, MPO, MOS, NOLA, ORU, USU,
WSL, WN
Culex pervigilans: WHA
Culex pipiens: CV, FLA, HP, IT,
JBE, LAC, OLI, SF, SIN, SLE,
TAH, TVT, TUR, UMA, WEE
Culex peus: SLE, TUR, WEE
Culex pipiens quinquefasciatus†:
BAN, EEE
Culex pipiens and *Cx pseudo-*
vishnui: AINO
Culex poicilipes: OLI
Culex portesi: ANU, BIM, CAB,
CAR, CATU, COC, COT, GJA, GMA,
ITQ, MAG, MTB, MOJU, MQO, MUC,
MUR, ORI, RES, SLE, TON, TUR
Culex pruina: BAG, KAM, MOS, SIN,
WN
Culex pseudosinensis: UMB
Culex pseudovishnui: KUN, SIN,
TMU, UMB
Culex pseudovishnui and *Culex*
India species No. 3: VEL
Culex pullus: ALF, SIN
Culex quinquefasciatus: CHIK,
EEE, KUN, MTB, MUR, MVE, ORO,
RR, SLE, SIN, TUR, VEE, WAN,
WEE, WN
Culex quinque-salinarius: EEE
Culex restuans: EEE, HP, SLE,
WEE
Culex rubinotus: AMT, BAN, GER,
UGS, WIT
Culex sacchettae: BER, EEE
Culex salinarius: EEE, FLA, SLE,
TEN
Culex sitiens: MOS, § TMU
Culex species: ACA, ANU, APEU,
ARU, BAG, BVS, BIM, BUN, BSB,
BSQ, CAB, CAP, CAR, CATU,
CHIK, COT, GER, GG, GJA, GMA,
ICO, ILH, ITP, ITQ, KWA, MAY,
MOS, MPO, MUC, MUR, NEP, NTA,
ORI, PAT, SLE, SIN, UNA, USU,
WEE, WIT
Culex spissipes: BVS, BIM, KRI,
SLE, TON
Culex squamosus: KUN, SIN
Culex taeniopus: ANU, BIM, BSQ,
CAP, EEE, ENS, MAG, MIR, OSSA,
PAT, SLE
Culex tarsalis: CV, CE, FLA,
GLO, HP, LLS, LOK, MD, SLE,
TUR, UMA, VEE, WEE

* *Culex molestus* is considered to be a biotype of *Cx pipiens*.

***Culex opisthopus* is a synonym for *Cx taeniopus*.

† Elevated to a full species (*Culex quinquefasciatus*).

§ Recovered from a pool identified as *Cx sitiens* but may also have contained *Cx thalassius* and *Cx tritaeniorhynchus*.

- Culex telesilla*: BAG, MID, MOS, NKO, WSL
Culex territans: EEE, FLA, SLE
Culex thalassius: BAG, ILE, SIN
Culex theileri: RVF, SHU
Culex theileri and *Cx rubinotus*: GER
Culex thriambus: PAT
Culex tigripes: BIA, KAM, MOS, SIN
Culex tritaeniorhynchus: AINO, AKA, ARK, DEN-3, GET, JBE, KAI, NOD, SAG, SIN, TMU, WN
Culex univittatus: BAG, ING, MOS, MPO, SIN, SPO, USU, WSL, WN
*Culex virgultus**: SLE, TUR
Culex vishnui: GAN, GET, ING, JBE, KAS, MIN, PAL, SAT, TMU, UMB, VKT, WN
Culex vomerifer: ANU, BSQ, CAR, GMA, ITQ, MAD, MOJU, OSSA, VIN
Culex weschei: BOU, MOS, MPO, WN
Culex zeteki: TON
Culex zombaensis: BUN, PGA
Culicines: BEF, JAP, JOI, MPK, SIN, TAH
Culicoides actoni: WAR
Culicoides austropalpalis: KNA
Culicoides brevitarsis: AINO, AKA, BC, BEF, BLU, CVG, DAG, DOU, EHD, KIM, NGA, PEA, TIB, TIN, WAL
Culicoides bundyensis: BEL
Culicoides dycei: WAL, WAR
Culicoides fulvus: BLU
Culicoides histris: SAT, THI
Culicoides imicola (= *C. pallidipennis*): BLU, SABO, SHA
Culicoides Kenya species No. 23: BLU
Culicoides kingi, *C. nivosus*, *C. bedfordi*, *C. pallidipennis*, *C. cornatus*, mixed pool: BEF
Culicoides marksii: BEL, BF, EUB, WAL, WAR
Culicoides milnei: BLU
Culicoides obsoletus: BLU
Culicoides oxystoma: BC, DAG
Culicoides oxystoma and *C. peregrinus*: MAR
Culicoides pallidothorax: WGR
Culicoides paraensis: ANU, ORO
Culicoides species: AHS, AINO, AKA, BLU, BUT, CON, CVG, DAG, DUG, EEE, EHD, IT, KOT, LOK, MAR, MR, NGA, RVF, SABO, SAN, SAT, SHA, SHU, WAL, WAR
Culicoides schultzei: BEF, BC, DAG, EHD
Culicoides schultzei and *C. peregrinus*: MAR
Culicoides tororensis: NSD
Culicoides variipennis: BLU, BUT, LOK, MD
Culicoides wadai: AKA
Culiseta: SIN,** WEE
Culiseta alaskaensis: NOR
Culiseta annulata: TAH
Culiseta impatiens: SSH
Culiseta inornata: CV, CE, JC, JS, MD, NOR, SLE, SSH, TVT, TUR, WEE
Culiseta melanura: EEE, FLA, HP, HJ
Culiseta morsitans: EEE, SIN
Culiseta tonnoiri: WHA
Deinocerites pseudes: SLE, VEE
Dermacentor albipictus: CTF
Dermacentor andersoni: CE, CTF, POW
Dermacentor auratus: KFD, LJN
Dermacentor marginatus: BHA, CHF, DHO, HYPR, OMSK, RAZ, RSSE
Dermacentor occidentalis: CTF
Dermacentor parumapertus: CTF
Dermacentor reticulatus (= *D. pictus*): HYPR, OMSK, RSSE, TET
Dermacentor silvarum: OMSK, RSSE

* Nomen Dubium.

**Virus of Ockelbo disease.

Dermacentor variabilis: SAW, SLE
Dermanyssus: WEE
Dermanyssus americanus: SLE
Dermanyssus gallinae: EEE, SLE
Echinolaelaps echidninus: JUN
Eomenocanthus stramineus: EEE
Eretmapodites chrysogaster: MID,
 NKO, OKO, RVF, SIM
Eretmapodites grahami: SF
Eretmapodites leucopous: NKO
Eretmapodites quinquevittatus:
 PGA
Eretmapodites semisimplicipes:
 OKO
Eretmapodites silvestris: SPO
Eretmapodites species: NKO, SPO
Eubrachilaelaps rotundus: JUN
Eusimulium johannseni: EEE
Ficalbia species: KOO, SEP
Gigantolaelaps inca: PIC
Gigantolaelaps species: COC, MAY,
 PIC, SLE
Haemagogus capricornii: YF
Haemagogus equinus: YF
Haemagogus janthinomys: MAY, YF
Haemagogus leucocelaenus: WYO, YF
Haemagogus lucifer: YF
Haemagogus mesodentatus: YF
Haemagogus species: ILH, MAY,
 MUC, TCM, YF
Haemagogus spegazzinii: JUR
Haemaphysalis bispinosa: KFD
Haemaphysalis concina: OMSK,
 RSSE, TDY
Haemaphysalis cuspidata: KFD
Haemaphysalis intermedia: BAR,
 BHA, GAN
Haemaphysalis inermis: HYPR
Haemaphysalis japonica douglasi:
 RSSE
Haemaphysalis kyasanurensis: KFD
Haemaphysalis leporispalustris:
 NM, SAW, SIL
Haemaphysalis longicornis: KHA,
 RSSE
Haemaphysalis minuta: KFD
Haemaphysalis neumani: POW
Haemaphysalis papuana: LGT
Haemaphysalis papuana kinneari:
 KFD
Haemaphysalis punctata: BHA, CHF,
 HYPR, TRB
Haemaphysalis semermis: LJN
Haemaphysalis species: LJN
Haemaphysalis spinigera: KSO, KFD
Haemaphysalis ticks: BHA
Haemaphysalis turturis: KSO, KFD
Haemaphysalis wellingtoni: GAN,
 KFD
Hyalomma anatolicum anatolicum:
 CHF, KEM, THO, WM
Hyalomma asiaticum asiaticum:
 CNU, CHF, ISF, TDY, WM, WAN
Hyalomma detritum: CHF
Hyalomma dromedarii: BHA, DGK,
 DHO, KAD
Hyalomma excavatum: CON
Hyalomma impeltatum: CON, DUG,
 WAN
Hyalomma impressum: CON
Hyalomma marginatum: BAH, CHF,
 DHO, MWA, MTR, SIN, TDY, WAN,
 WN
Hyalomma marginatum isaaci: WM
Hyalomma marginatum rufipes: BAH,
 BHA, CON, DUG
Hyalomma marginatum turanicum:
 CHF
Hyalomma nitidum: CON, DUG, THO
*Hyalomma plumbeum plumbeum**: BKN,
 CHF, DHO, TDY, WN
Hyalomma rufipes: CON
Hyalomma species: CHF, CON, WM,
 WN
Hyalomma truncatum: BHA, CON,
 DUG, THO, WM
Hyalomitra lasiophthalma: JC
Hyalomitra nuda: JC
Ixodes apronophorus: OMSK
Ixodes cookei: POW
Ixodes dentatus: CNT
Ixodes eudyptidis: GGY, SRE

* *Hyalomma plumbeum* = *Hyalomma marginatum* of Hoogstraal.

Ixodes granulatus: LGT, LJN
Ixodes lividus: RSSE
Ixodes marxi: POW
Ixodes persulcatus: ABS, KEM, LGT, OMSK, RSSE
Ixodes petauristae: KFD
Ixodes putus: OKH, PMR, SAK, TYU, ZT
Ixodes redikorzevi: HAZ
Ixodes ricinus: ABS, CHF, EYA, HAN, HYPR, KUM, LIP, LI, RSSE, TET, TRB, UUK
Ixodes ricinus and *I. persulcatus*: UUK
Ixodes signatus: GI, PMR
Ixodes spinipalpus: POW
Ixodes uriae (= *I. putus*): AVA, BAU, CW, CL, GGY, GI, MYK, NUG, PP, SAK, TAG, TDM, TYU, UUK, YM
Ixodes species: KFD
Ixodes tropicalis: PIC
Ixodes vespertilionis: IK
Ixodidae: DUG
Laelapidae: AMA
Laelapid ("Gamasoid") mites: RSSE
Lasiohelea taiwana: JBE
Limatus durhami: CAR, WYO
Limatus flavisetosus: GJA, WYO
Limatus pseudomethysticus: CAB, COT
Limatus species: GMA, WYO
Lutzomyia: GMA
Lutzomyia flaviscutellata: ICO, INH, PAC
Lutzomyia panamensis: NIQ
Lutzomyia species: AGU, ALT, BUE, CAC, CAN, CHG, CHI, COT, FRI, GUR, JAM, ORX, OUR, PT, SRA, TUA, VSI
Lutzomyia trapidoi: AGU, CAC, CHG, PT, VSI
Lutzomyia umbratilis: AMR, MUN
Lutzomyia ylephilator: AGU, CAI, CHG, PT
Mansonia africana: BAN, BUN, CHIK, LEB, MID, PGA, SHO, SPO, USU, WSL, ZGA
Mansonia africana/*Ma uniformis*: PGA
Mansonia dyari: VEE
Mansonia indubitans: SLE, VSNJ
Mansonia (*Man*) species: MQO
Mansonia papuensis: MPK
Mansonia pseudotitillans: SLE, TON
Mansonia septempunctata: SEP, SIN
Mansonia species: CHIK, GMA, MOJU, MUC, ORI, VEE
Mansonia titillans: BSQ, CAB, TLA, TON, VEE
Mansonia uniformis: BUN, MAL, MID, NDU, OLI, PUC, RR, SAN, SPO, WSL, YATA, ZIKA, ZGA
Mansonia uniformis, *Anopheles gambiae*, and *Culex antennatus*: BAR
Mimomyia flavens: SEP
Mimomyia flavens and *Mi modesta*: MPK
Mosquitoes: IT, MAP, MID, NEP, TAT, ZIKA
Mosquitoes, mixed species: BEF, EUB, MAG, MAP, MEL, NTA, SEP, TCR, TAH
Mosquitoes, other: BIM
Musca autumnalis: VSNJ
Musca domestica: VSNJ
Oeciacus vicarius: FM
Ornithodoros adult: KFD
Ornithodoros amblus: HUA, PS
Ornithodoros boliviensis: MAT
Ornithodoros capensis: AB, BAKU, HUG, JA, QRF, SRE, SOL, UPO
Ornithodoros capensis and/or *O. denmarki*: SOL
Ornithodoros denmarki: HUG
Ornithodoros erraticus: ASF, BDA, QYB
Ornithodoros maritimus: MEA, SOL
Ornithodoros moubata porcinus: ASF
Ornithodoros musebecki: ZIR
Ornithodoros papillipes: CHIM, KSI
Ornithodoros sonrai: CHIK

Ornithodoros species: BDA, CG,
 HUG, KFD
Ornithodoros tadaridae: ER
Ornithodoros tartakovskyi: CHIM
Ornithonyssus sylviarum: SLE
Otobius lagophilus: CTF
 Phlebotomines: CGL
 Phlebotomines species: ORX
Phlebotomus papatasi: ISF, SFN,
 TEH
Phlebotomus perfiliewi: ARB, SFN,
 YB
Phlebotomus perniciosus: ARB, TOS
Phlebotomus species: CHP, CHV,
 KAR, SAL, SFN, SFS
Phoniomyia pilicauda: AMB, BOR
Phoniomyia species: ARU
Psorophora albipes: IERI, ILH,
 ITU, PUR, UNA, WYO
Psorophora ciliata: VEE
Psorophora cilipes: VEE
Psorophora cingulata: WYO
Psorophora columbiae: CV, JC, SA,
 TEN, WEE
Psorophora confinnis: VEE
Psorophora cyanescens: VEE
Psorophora discolor: JC, VEE
Psorophora ferox: ARU, CV, COT,
 IERI, ILH, ITU, KRI, MEL, MIR,
 ORI, ROC, SLE, UNA, WYO
Psorophora ferox and *Ps albipes*:
 UNA
Psorophora howardii: LAC
Psorophora lutzii: UNA
Psorophora signipennis: CE, LOK,
 MD, SA, TVT, WEE
Psorophora species: CV, GMA, ILH,
 MAG, MAY, VEE, WEE, WYO
Rhipicephalus appendiculatus: NSD
Rhipicephalus bursa: CON, CHF,
 THO
Rhipicephalus evertsi: THO, WM
Rhipicephalus guilhoni: WM
Rhipicephalus lunulatus: DUG, WN
Rhipicephalus muhsamae: DUG, KOU,
 WSL, WN
Rhipicephalus pravus: KAD
Rhipicephalus pulchellus: BAR,
 DUG
Rhipicephalus pumilio: CHF
Rhipicephalus rossicus: CHF
Rhipicephalus sanguineus: CHF,
 CON, THO, WM
Rhipicephalus sanguineus
turanicus: MWA
Rhipicephalus species: KFD, LJN,
 MWA, THO
Rhipicephalus sulcatus: DUG
Rhipicephalus turanicus: CHIM,
 CHF, MWA, TDY
Sabethes belisarioi: SLE
Sabethes chloropterus: ARU, CHG,
 SLE, YF
Sabethes intermedius: XIB
Sabethes soperi: MCA
Sabethes species: ILH, SLE
 Sabethines: MUR
 Sabethini species: ICO, KRI, MAY,
 MOJU, MUC, ORI, SOR, WYO
Scipio aulacodi: LEB
Sergentomyia species: SRI
 Simuliidae: VSNJ
Simulium bivittatum: LOK
Simulium black flies: VSNJ
Simulium malyscheri: SSH
Simulium meridionale: EEE
Stricticimex parvus: KK
Tabanus agrestis: IK
 Ticks: BHA
Triatoma: WEE
Trichoprosopon digitatum: PIX,
 WYO
Trichoprosopon leucopus: WYO
Trichoprosopon longipes: SLE, WYO
Trichoprosopon pallidiventer: AMB
Trichoprosopon species: BSQ,
 GMA, ILH, SLE, TNT, WYO
Trichoprosopon theobaldi: ARU
Uranotaenia species: JUG
Wyeomyia aporonoma: KRI, WYO
Wyeomyia complosa: WYO
Wyeomyia melanocephala: TON, WYO
Wyeomyia occulta: CAB, TON, WYO
Wyeomyia pseudopecten: TON
Wyeomyia species: ARU, CAR, IACO,
 ILH, KRI, MCA, MAG, MQO, MUC,
 SLE, UNA, WYO
Wyeomyia ypsipola: KRI

HOST INDEX

- Ablepharus boutonii virgatus*:
 ALM, KOW
Acomys cahirinus (= *A. albigena*):
 GF
Acrocephalus schoenobaenus: SJA
Aepeomys (= *Thomasomys*) *fuscatus*:
 PIC
Aethomys medicatus: SGA
Agelaius phoeniceus phoeniceus:
 MER
Agelaius tricolor: HP
Akodon arenicola: JUN
Akodon azarae: JUN
Akodon species: KRI, SLE
Alcedo atthis: IK
Alouatta seniculus insularis: MAN
Alouatta, sentinel: BSQ, GMA
Amevia ameiva: BOC
Amevia ameiva ameiva: CHO, MCO,
 MAY, SM, TIM
Andropadus virens: GOM, MOS, USU
 Antelope: BOU, RVF
Antichromus minutus: WN
Apodemus agrarius coreae: HTN
Apodemus argenteus hokkaidi: APOI
Apodemus flavicollis: HYPR, TET,
 UUK
Apodemus sylvaticus: HYPR, LI
Ardea cinerea: SEM
Ardea novaehollandiae: MVE
Ardeola grayii: ING, THI
Artibeus jamaicensis: NEP
Artibeus jamaicensis trinitatus:
 TCR
Artibeus lituratus: NEP
Artibeus lituratus palmarum: TCR
Arvicanthus niloticus: AMT, GF,
 UGS, WN, WIT
Arvicanthus species: BDA, GF,
 IPPY, SAB
Arvicola species: TBE
Auripasser: CHIK
Bandicota indica: HTN/SEO*
 Bats: CAB, GMA, MDC, RB, SLE, WN
 Blue jay: See *Cyanocitta cristata*
 Birds, wild: ANU, EEE, HJ, ICO,
 ILH, ING, JBE, MOS, ROC, SF,
 SIN, TDY, TETE, TON, TUR, UMB,
 UUK, WEE, WN
 Boar, wild: CVO
 Bobcat: EVE
Bolomys (= *Akodon*) *obscurus*: JUN
Bos taurus: See Cattle
 Bovine: See Cattle
Bradypus tridactylus: MUR, ORO,
 UTI
Bubulcus ibis: NYM, QRF
 Buffalo: RVF
 Buffalo, water: MAR
Bufo marinus: CUI
Bycanistes sharpii: ZGO
Callithrix species: MAY
Calomys callosus: LAT, MAC
Calomys laucha: JUN
Calomys musculinus: JUN
Caluromys species: CAP
Caluromys p. philander: ANU, APEU
 Calves - See Cattle
 Calves, fetal - See Cattle
 Camel: THO, WSL, WN
 Cardinal: See *Cardinalis*
 cardinalis
Cardinalis cardinalis (= *Richmondia*
cardinalis): MER,
 SLE
Carduelis cannabina: MTR**
 Caribou: CV
Carollia subrufa: AP
Carpodacus mexicanus: HP, TUR
 Cat bird: SLE
Cathartes aura: NAV
Catharus ustulatus swainsoni (= *Hylocichla*
ustulata swainsoni):
 MER

* A single isolation of either a Hantaan or Seoul virus from *Bandicota indica*.

**Most of these isolates have not been serotyped; they may represent Bahig or Matruh viruses or other Tete group viruses.

Cattle: AINO, AKA, AR, BHA, BLU,
 BEF, BC, BRM, CV, CON, CHF,
 CVG, DAG, DOU, DUG, EHD, IBA,
 KIM, LI, MAR, NDU, PEA, RVF,
 SABO, SAN, SAT, SHA, SHU, THO,
 TIN, VSI, VSNJ, WSL
 Cattle, sentinel: BC, CVG
Cavia pamparum: JUN
Cebus, sentinel: CAR, CATU, GMA,
 ILH, ITQ, MTB, MIR, MOJU, MUC,
 MUR, ORI
Cebus apella, sentinel: APEU,
 GMA, MTB, TCM
Centropus phasianus: ALF
Cercopithecus aethiops: CHIK, YF,
 ZIKA
Cercopithecus nictitans: BOU
Chaerephon pumila [= *Tadarida*
 (*Chaerephon*) *limbata*]: ENT
Charadrius melanops: COR
 Chickens, sentinel: ANU, MVE,
 TMU, TUR, WEE, WN
 Chimney swift: SLE
Chloris chloris: BAH*
Choloepus brasiliensis: ANH
Choloepus didactylus: JARI
Cisticola chiniana: TETE
Clethrionomys glareolus: KEM, LI,
 PUU, TBE, TET, TRB
Clethrionomys rufocanus: PUU
Clethrionomys rutilus: KEM, KLA,
 NOR
Clethrionomys species: POW
Clytospiza monteiri: ING
Coendou species: GMA
Colius passer macrourus: MOS
Columbigallina: MAY
Coracopsis vasa: WN
Corvus corone sardonius: SIN
Corvus splendens: KAN
Corythornis cristata: GAR
 Cotton rat: See *Sigmodon hispidus*
Coturnix coturnix: MTR*
 Cows: See Cattle
Cricetomys gambianus: BDA, DUG,
 GF, UGS
Crocidura species: AMT, BLU
Cyanocitta cristata: HJ, MER, SLE
Cyanopica cyanus: TSU
Cynopterus brachyotis: CI, JUG,
 PPB
Cynopterus brachyotis angulatus:
 PPB
Dasypus novemcinctus: MDO
 Deer: EHD, LI, WEE
 Deer, sentinel: JC
 Deer, red: See Deer
 Deer, white-tailed: See Deer
Desmodillus auricularis: WSL
Dicrostonyx rubricatus: SSH
Didelphis m. marsupialis: ANU,
 CAB, CATU, ITA, MTB, MOJU,
 MUR, ORI
 Dog: AHS, EEE, TEN, WN
 Donkey: See Horse
 Dove, mourning: SLE
 Dove, ruddy ground: SLE
 Dove, white tipped: SLE
Dryoscopus gambensis: DUG
 Egret: WN
 Egret, nestling: NYM
Eidolon helvum: IFE, LB
Emberiza citrinella: KUM
Eonycteris spelaea: PPB
Epomophorus species: YF
Epomophorus wahlbergi: LB
Eptesicus fuscus: RB
Eptesicus (= *Vespertilio*)
serotinus: IK
Erethizon dorsatum: CTF
Erinaceus albiventris (= *Atelerix*
spiculus; *A. albiventris*):
 BHA, CHP, CON, GF, SF
Erinaceus concolor (= *E.*
roumanicus): HYPR, RSSE
Erithacus luscini (= *Luscinia*
luscini): BAH,* MTR*

* Most of these isolates have not been serotyped; they may represent Bahig or Matruh viruses or other Tete group viruses.

Erithacus megarrhynchus (= *Luscinia megarrhynchus*): BAH*
Erythrocebus patas: CHIK, DEN-2, YF, ZIKA
Estrilda melopoda: ING, WN
Euplectes afra: BMA, BBO, KOL
Euplectes orix: ING, TETE
 Flicker: SLE
Formicarius analis: CPC
 Fowl, domestic: MVE
 Fox: POW, SLE
Fringilla coelebs: BAH*
Fringilla montifringilla: BAH*
 Frogs: OMSK, WEE
Galago: WN
Galago senegalensis: CHIK, GF
Gehyra australis: CHV
 Goats: ABS, CON, CHF, NSD, SABO
 Goats, sentinel: TRB
 Goose: SLE
Gracula religiosa: SIN
Grallina cyanoleuca: RR
 Grouse: LI
Halcyon senegalensis: SF
 Hamster, sentinel: AROA, BAB, BAN, GER, GMA, GTB, ITQ, MAD, MNT, MIR, NJL, NEP, OSSA, PAT, PLS, SR, VEE, VIN, WIT, ZEG
 Hares: See Rabbits
 Heron, white-faced: MVE
Heteromys anomalus: BIM, CAR, COC, GMA, MUC, VEE
 Hipposideridae: SIN**
Hipposideros caffer: ZGA
Hipposideros terasensis: JBE
Hirundo rustica: IK
 Horse: AHS, CV, COC, EEE, GET, JBE, KUN, LI, MAG, MD, MID, RR, UNA, VEE, VSA, VSI, VSNJ, WEE, WN

House finch: See *Carpodacus mexicanus*
 House sparrow: See *Passer domesticus*
 Humans: ABS, ALE, APEU, BGI, BAN, BHA, BUN, BSQ, BWA, CDU, CAR, CATU, CHG, CHP, CGL, CHIK, CTF, CON, COT, CHF, DB, DEN-1, DEN-2, DEN-3, DEN-4, DHO, DUG, EEE, EBO, EVE, GAN, GER, GMA, GRO, HTN, HAN, HYPR, ILE, ILH, IK, ITQ, JBE, JUN, KEM, KOU, KUM, KUN, KFD, LAC, LAS, LEB, LD, LI, MAC, MAD, MBG, MTB, MAY, MID, MUC, MVE, MUR, NSD, NEG, NEP, NDO, OMSK, ONN, ORI, ORO, ORU, OSSA, PGA, PIRY, POW, PT, QRF, RES, RVF, ROC, RR, RSSE, SFN, SFS, SF, SHO, SHU, SIN, SLE, SPO, TCM, TAH, TDY, TAN, TAT, TET, THO, TOS, (UGS), † UMB, USU, VEE, VSI, VSNJ, WAN, WSL, WEE, WN, WYO, YF, ZIKA, ZGA
Hylochoerus meinertzhageni: ASF
Hylomyscus species: SEB
Hylophylax naevioides: BLM
Hylophylax poecilonota: MCA
Hyphanturgus brachypterus: ING
Hyphanturgus nigricollis: BMA
Hyphanturgus ocularius: ING
Jynx torquilla: IK
Kentropyx calcaratus: CHO
Lanius collurio: MTR*
Larus argentatus: AVA
Lavia frons: RVF
Lemniscomys barbarus: GF
Lemniscomys species: GF, KOU

* Most of these isolates have not been serotyped; they may represent Bahig or Matruh viruses or other Tete group viruses.

**Isolated from a pool of organs from bats of above family plus bats of Rhinolophidae family.

† () = questionable.

Lemniscomys striatus: AMT, GOR,
 IPPY, KOU
Lepus americanus: SIL, SSH
Lepus californicus: BUT, CTF,
 LOK, MD, TUR
Lepus europaeus: JUN
Lepus timidus: KUM
 Lizard: OMSK
Lophuromys flavopunctatus: WIT
Macaca fascicularis: BAK
Macaca mulatta: SHF
Macaca nemestrina: LJN, TR, TF
Macaca radiata: KFD
Macroglossus lagochilus: CI
Macropus (= Wallabia) agilis: RR
 Mammals, small: AMT
 Manakin, black and white: SLE
 Manakin, flame-headed: SLE
Marmosa: EEE
Marmosa cinerea: APEU
Marmosa mitis: MUC
Marmosa murina: ITQ
Marmosa species: MUR
Marmota monax: POW
 Marsupials: GMA, MUC
Meleagris gallopavo: IT
Metachirus nudicaudatus: CAB, ITQ
Microeca fascians: RR
Micropteropus pusillus: LB
Microtus agrestis: KUM
Microtus montanus: KLA
Microtus oeconomus: KLA
Microtus pennsylvanicus: PH
*Microtus gregalis (= M.
 stenocranium)*: OMSK
 Migratory finch: MTR
Miniopterus s. fuliginosus: JBE
 Mockingbird: SLE
Molossus obscurus: CATU
 Monkey: DEN-2, YF
 Monkey, sentinel: ANU, DEN-1,
 DEN-2, KRI, ORU, VEE, ZIKA
Motacilla alba: IK, SIN
Motacilla cinerea: IK
 Mouse, deer: CTF
 Mouse, sentinel: ACA, ANU, APEU,
 BVS, BEN, BER, BIM, BSB, BSQ,
 CAB, CNA, CAP, CAR, CATU,
 CHIK, COC, COT, GF, GJA, GMA,
 GTB, HJ, ICO, ITP, ITQ, JD,
 MAD, MAG, MTB, MIR, MOJU, MUC,
 MUR, NEP, ONN, ORI, PARA, PAT,
 ROC, SF, SIN, TBT, TON, TUR,
 UGS, UNA, VEE, WN, WYO, ZEG
 Mule: See Horse
Mus musculus: JUN, SAB, TBE
Mus species: LAS, WEE
Muscicapa striata: BAH*
Myotis blythii (= M. oxygnathus):
 IK
Myotis lucifugus: BOC, MML
Myotis yumanensis: KC
Neacomys guianae: AMA, ICO
Nectarinia pulchella: GAR
Nectomys: NEP
Nectomys squamipes: ACA, CAR,
 CATU, GMA, ITQ, MOJU, MUC, MUR
Nectomys squamipes amazonicus:
 BEN
Neotoma micropus: RG
Numida meleagris: ING
Nyctalus noctula: IK
Nycticorax nycticorax: SEM
Odocoileus virginianus: See Deer
Ondatra zibethicus: OMSK
 Opossum: EVE
Oriolus flavocinctus: KUN
Oriolus oriolus: BAH*
Oryzomys: EEE
Oryzomys albigularis: PIC
Oryzomys bicolor: FLE
Oryzomys (= Oecomys) bicolor:
 KRI, MOJU
Oryzomys buccinatus: PAR
Oryzomys capito: AMA, CAR, CATU,
 FLE, ITQ, MUC, ORI

* Most of these isolates have not been serotyped; they may represent Bahig or Matruh viruses or other Tete group viruses.

Oryzomys capito goeldii: PAC
Oryzomys capito velutinus: PAC
Oryzomys flavescens: JUN
Oryzomys goeldii: AMA
Oryzomys laticeps: CAR, CATU, MUC
Oryzomys palustris: TAM
Oryzomys, sentinel: CAR, GMA, MUC
Oryzomys species: BIM, GMA, IRI,
 ITI, MOJU, MUR, STM
Otomys irroratus: BLU
 Ovine: See Sheep
Papio papio: BOU, CHIK
Passer domesticus: FM, HP, SLE,
 TUR, UMA
Passer hispaniolensis: IK
Peromyscus gossypinus: TEN
Peromyscus leucopus: SV
Peromyscus maniculatus: CTF, MOD
Peromyscus species: EEE, POW, WEE
Petrochelidon pyrrhonota: FM
Phacochoerus aethiopicus: ASF
 Pheasant: EEE
Philander opossum: ARA, PIRY
Phoenicurus phoenicurus: BAH,*
 IK, KEM, MTR*
Phylloscopus collybitus: BAH*
Phylloscopus trochilus: BAH,*
 MTR*
 Pigeon: QRF, SLE
 Pig: CNU, GET, ING, JBE, VSI,
 VSNJ, WEE
Pipistrellus (= *Vespertilio*)
pipistrellus: SOK, IK
Pipra erythrocephala: MUC
Pitmys subterraneus: TRB
Platyrrhinus coronatus: MCA
Plesiositagra cucullata: ING,
 KOL, TETE
 Ploceide: ING
Ploceus cucullatus: TETE
Ploceus melanocephalus: BMA, OUA
 Porcupine: CTF
Potamochoerus species: ASF
Praomys (= *Mastomys*)
erythroleucus: SGA
Praomys (= *Mastomys*) *femelle*: CON
Praomys (= *Mastomys*) *natalensis*:
 DUG, GF, LAS, UGS
Praomys (= *Mastomys*) species:
 BDA, BLU, IPPY, KEU, KOU, SAB,
 SGA, SPA, SEB
Presbytis entellus: KFD
 Primates: YF
Proechimys iheringi: AMB
Proechimys guyannensis: BIM, BSQ,
 CAP, CAR, CATU, GJA, GMA, ITQ,
 MOJU, MUC, MUR, ORI, URU
Proechimys oris (= *Proechimys g.*
oris): BUJ, ICO, PIX
Proechimys longicaudatus: URU
Proechimys semispinosus: MAD,
 OSSA
Proechimys species: EEE, MCA,
 NEP, TBT
Progne subis: MER
Psarocolius decumanus: TON
Pteroglossus aracari: INI
Puffinus pacificus chlorohynchus:
 UPO
Pycnonotus barbatus: ING
Pyrglana leucoptera: BLM
Quelea erythroptera: SF
Quelea quelea: ING
 Rabbits (and Hares): CON, CHF,
 LI, TEN
 Rabbit, sentinel: KEY, NOR, SSH,
 TAH, TVT
 Rat: KK
 Rat, laboratory: SEO
 Rat, sentinel: MUC
 Rat, Sprague-Dawley: SEO
 Rat, Wistar: SEO
Rattus blanfordi: KFD
Rattus nitidus: SEO
Rattus norvegicus: SEO, VEE
Rattus rattus: LAS, SEO, VEE

* Most of these isolates have not been serotyped; they may represent Bahig or Matruh viruses or other Tete group viruses.

Rattus rattus wroughtoni: BAR, KFD
Rattus species: SEO
 Rhinolophidae: SIN**
Rhinolophus c. cornutus: JBE
Rhinolophus ferromequinum: IK
Rhinolophus hildebrandti eloquens: MEB
Rhinolophus rouxi: KFD
Riparia paludicola: LJA
 Robin: SLE
 Robin, pale vented: SLE
 Rodents: EEE, GER, ICO, RVF, RSSE, SHO, WEE
Rousettus aegyptiacus: YOG
Rousettus leschenaulti: WN
 Ruminants, domestic: BLU
 Ruminants, wild: BLU
 Saimiri: KRI
Saxicola rubetra: MTR,* UGS
Sciurus carolinensis: LAC
Sciurus carolinensis, sentinel: LAC
Sciurus griseus: WEE
Sciurus vulgaris: KUM
Scotophilus nigrita nigrita: DB
Scotophilus species: CHIK, DB
Scotophilus temmencki: KTR
Seiurus aurocapillus: FLA
Serinus canaria: MTR*
Serinus mozambicus: TETE
 Sheep: BHA, BLU, CV, GAN, LI, NSD, RVF, SABO, SHU, WSL
 Siberian polecat: TDY
Sigmodon hispidus: CR, EVE, GL, JUT, KEY, MH, PAT, SP, SR, TAM, TEN, TVT, VEE, WEE, ZEG
Sigmodon species: PAT, TAM, ZEG
 Snake: WEE
Sorex araneus: HYPR, LI
Spermophilus (= Citellus) beecheyi: WEE
Spermophilus (= Citellus) columbianus: CTF
Spermophilus (= Citellus) lateralis: CTF
Spermophilus (= Citellus) richardsoni: WEE
Spilogale putorius (= spotted skunk): POW
 Squirrel: CTF
 Starling, immature: FLA
Sterna fuscata: HUG
Sturnira lilium: MPR
Sturnus pagodarum: KMP
Suncus murinus: HTN/SEO, KFD, TPM
Sus scrofa: ASF
Syconycteris crassa: JAP
Sylvia atricapilla: BAH*
Sylvia borin: BAH,* MTY, MTR*
Sylvia communis: BAH,* MTR,* THI
Sylvia curruca: BAH,* BEA, MTY, MTR,* THI
Sylvia hortensis: BAH*
Sylvia nisoria: BAH*
Sylvia ruppelli: BAH*
Sylvilagus aquaticus: TEN
Sylvilagus auduboni: BUT, LOK
Tadarida brasiliensis mexicana: RB, SLE, VEE
Tadarida condylura wonderi: DB
Tadarida plicata: KK
Tadarida species: DB, GOS
Talpa europaea: HYPR
Tamias striatus: LAC
Tamiasciurus hudsonicus: POW
 Tanager, silver beak: SLE
Taphozous theobaldi: KK
Tatera valida (= T. kempi): AMT, BDA, GF, KEU, KOU, SAB, TOU
Tatera species: GF, GOR, GOS, SAF
Taterillus gracilis: GF
Taterillus nigeriae: GF

* Most of these isolates have not been serotyped; they may represent Bahig or Matruh viruses or other Tete group viruses.

**Isolated from a pool of organs from bats of above family plus bats of Hipposideridae family.

Taterillus species: BDA, GF,
KTR, KEU, KOU
Tchagra australis: WN
Thamnomys macmillani: AMT
Thamnophilus aethiops: ITP
Thryonomys swinderianus: LAT, LEB
Tolmomyias poliocephalus: CAB
Trichosurus vulpecula: WHA
Trionyx spinifer: CV
Tropidurus torquatus hispidis:
MAY
Turdus libonyanus: AMT, BGN, BOB,
MPO, USU
Turdus merula: KUM, UUK
Turdus nudigenis: CAB
Turdus philomelos: KUM
Urocyon cinereoargenteus: TEN
Vespertilio murinus: IK
Xanthocephalus xanthocephalus: HP
Xanthomyias virescens: GTB
Xerus erythropus: BDA, BHA
Zoothera citrina: KSO
Zygodontomys brevicauda: BIM,
BOC, CAR, GMA, MUC, NAR, PAC
Zygodontomys species: BIM, CAR,
CATU, GMA, PAC, VEE

REPORT FROM THE WHO COLLABORATING CENTRE FOR REFERNECE AND RESEARCH
(Dengue and Dengue Haemorrhagic Fever)

Department of Medical Microbiology, University of Malaya, Kuala Lumpur, Malaysia.

A moderately large outbreak of dengue took place in Malaysia in 1986. Out of 1399 cases reported to the Ministry of Health throughout the country, 310 were classified clinically as dengue haemorrhagic fever. There were 9 deaths from among the DHF cases, giving a case fatality rate of 2.9%. The outbreak was predicted when the Centre noticed an increase of dengue cases seen at the University Hospital in December, 1985. The peak reporting period was June-July (Table 1). Four states in the west coast of Peninsular Malaysia and one in the eastern state reported the highest number of cases.

In the University Hospital, 624 patients were tested serologically for dengue infections. Many of the samples were single specimens, making interpretation of results difficult. 81 (13%) cases were found to be positive serologically, 46 samples displaying a secondary type infection based on WHO criteria. 43 of the 81 positive cases were in males and 38 females. The majority of cases were found in those under 30 years of age. There were 47 cases among the Chinese, 24 in Malays, 9 Indians and 1 others. Thirty-two viruses were isolated in mosquito larvae from clinical specimens. Twenty-five were Den-3, four Den-1, 2 Den-4 and only 1 Den-2.

When it was evident that we have an imminent outbreak of dengue in the country, the Centre liased closely with the Ministry of Health and regular dialogue was established with the Vector Borne Diseases Control Programme. Increased surveillance was initiated by approaching general practitioners and private cinics, seeking their cooperation to submit specimens for investigation. This was particularly so in the Jinjang-Kepong area where we have an ongoing dengue project. The Ministry mounted an intensive programme of vector control. As a result of all these efforts, the number of cases reported in 1986 were much less than the previous large outbreak in 1982 when 3126 cases were reported with 36 deaths. Detailed analyses of this outbreak are in progress.

Dr. S.K. Lam

Table 1 Dengue Activity in Malaysia - 1986

<u>Month</u>	<u>DF Cases</u>	<u>DHF Cases</u>	<u>Total DF & DHF</u>	<u>Deaths</u>
January	23	9	32	1
February	30	18	48	2
March	73	32	105	3
April	107	23	130	-
May	148	36	184	-
June	196	60	256	1
July	157	43	200	-
August	113	21	134	-
September	97	15	112	1
October	47	29	76	-
November	29	8	37	1
December	69	16	85	-
TOTAL	1089	310	1399	9

COLLEGE OF MEDICINE, UNIVERSITY OF IBADAN, IBADAN, NIGERIA.

Pathogenicity of Potiskum Virus in Laboratory
Animals and Tissue Culture

The growth of Potiskum virus (Ib-AN 10069), a hitherto unknown flavivirus isolated from the liver of a giant rat (Cricetomys gambianus) collected at Fika, Bornu State of Nigeria was studied in various laboratory animal hosts and tissue culture. A ninth suckling mouse passage of the virus was used in all tests.

Mice - Suckling mice infected with 100 suckling mouse intracerebral (SMIC) LD₅₀ of Potiskum virus were susceptible to infection by intracerebral (i.c.), intraperitoneal (i.p.) and subcutaneous routes. The average survival times were 3.99, 4.67 and 6.0 respectively. All infected animals developed viraemia and this was detected from day 1 post infection (p.i.) until the time of death. Peak viraemia of 4.0 logs and 4.5 logs occurred on day 5 in i.c. and i.p. inoculated animals respectively.

Weanling (WM) and adult mice infected with Potiskum virus by i.c. and i.p. routes also developed viraemia on day 2 p.i. and lasting 3 days. Mortality rate in WM inoculated by i.c. route was 100% while 33% of those infected by i.p. route died. Peak viraemia of 3.2 logs and 2.6 logs occurred on day 5 in WM infected by i.c. and i.p. routes respectively. In adult mice, peak viraemia of 2.5 logs occurred in i.p. inoculated mice but those infected by s.c. route did not circulate the virus.

Rabbits - Adult rabbits were infected with 100 SMIC LD₅₀ of Potiskum virus. No viraemia was detected but all infected rabbits developed neutralising (N) and complement fixing (CF) antibody to Potiskum virus. Sera collected on day 28 p.i. had neutralising antibody titres of between 1:16 - 1:64 and CF antibody titres of 1:32 - 1:128.

Chicks - Four to 5 day old chicks were also susceptible to fatal Potiskum virus infection by subcutaneous (S.C.) and oral routes. Chicks were infected with 200 SMIC LD₅₀ of Potiskum virus. All chicks developed viraemia on day 1 p.i. which lasted until the time of death on day 4 p.i.

Tissue Culture

BHK-21 - Baby hamster kidney cells were infected with Potiskum virus using 100 SMIC LD₅₀/2ml of virus suspension. The virus replicated to high titres in this cell line. Highest virus titre (6×10^4 pfu/ml) was obtained on day 5 p.i. Cytopathic effect (CPE) commenced on day 3 p.i. and progressed until day 5 p.i.

Potiskum virus also produces plaques in BHK-21 under carboxyl-methyl cellulose (CMC) overlay medium. Plaques which appeared on day 1/4 p.i. were small and distinct with a diameter of about 1.5mm.

P388D₁ Cells - P388D₁ cells (a macrophage-like cell line) were also infected with Potiskum virus at two multiplicities of infection (MOI) 0.04 and 0.0004. Infectivity titres were determined by plaque assay. In cell cultures infected with high MOI, virus replication was detected on day 1. Virus titre increased from 2×10^3 pfu/ml to 8×10^3 pfu/ml. on day 3 p.i. Cultures infected with low MOI (0.0004) had late virus growth; virus replication was detected on day 2 p.i. Peak virus titres (4×10^3 pfu/ml.) was detected on day 3. Low grade CPE was detected on days 3 and 4 p.i. in cultures infected at both MOI.

Potiskum virus also produces plaques in P388D₁ cells under CMC overlay medium. Plaques appeared on day 3 p.i. and were similar in morphology to those of BHK-21 cells.

(A. H. Fagbami; S. A. Omilabu)

Clinical, serologic and virologic studies were carried out on 538 patients with a clinical diagnosis of dengue fever during epidemic in Yucatán, Mexico, 1984. Dengue virus infection was confirmed in 200 patients by serology (HI, CF or MAC ELISA) or by virus isolation. It was not possible to confirm the etiology of 207 patients on whom only a single serum sample was obtained. The rest were negative.

Of the 200 confirmed dengue patients, 189 (94.5%) were classified as secondary infections and 9 (4.5%) as primary infections using HI criteria. Two could not be classified. Virus isolation was attempted from patients from whom serum samples were collected in the first five days of illness. Dengue 4 (DEN-4) virus was isolated from 34 patients and dengue 1 (DEN-1) virus from one patient.

The signs and symptoms observed most frequently (in more than the 50% of the confirmed patients were: fever, headache, myalgias and retro-orbital pain. Rash was observed in 31.5% of the patients.

Ten cases of hemorrhagic disease were reported, four with a fatal outcome. Eight were confirmed as dengue, including the four fatal cases. The latter were all female, 8, 9, 19 and 37 years old, and from rural areas with low socio-economic conditions. DEN-4 virus was isolated from one fatal case and from three of the other cases with hemorrhagic disease. The other five cases of hemorrhagic disease were confirmed by IgM capture ELISA and/or by HI and CF tests. They included 3 males (5, 11 and 38 years old) and 2 females (19 and 43 years old). Hemorrhagic manifestations of confirmed cases are shown in Table 1.

Many of the outbreaks caused by DEN-4 virus in the American Region have been associated with relatively mild illness. The experience in tyhe Yucatán, however, demonstrates that not all DEN-4 infections are mild and reinforces the fact that all four dengue serotypes can cause severe and fatal disease.

(María Alba Loroño Pino; José Arturo Farfán Ale; D. J. Gubler; G. Kuno; G. E. Sather and J. E. Zavala Velázquez).

Table 1

Hemorrhagic Manifestations Associated with Confirmed Dengue 4
Infection, Mérida, Mexico, 1984

Hemorrhagic Manifestation	Fatal Cases	Nonfatal Cases	Total
Petechiae	4/4 (100)*	3/4 (75)	7/8 (87)
Purpura/Ecchymosis	3/4 (75)	1/4 (25)	4/8 (50)
Epistaxis	1/4 (25)	2/4 (50)	3/8 (37)
Gum Bleeding	1/4 (25)	2/4 (50)	3/8 (37)
Hematemesis	4/4 (100)	0/4 (0)	4/8 (50)
Melena	2/4 (50)	0/4 (0)	2/8 (25)
Hematuria	1/4 (25)	1/4 (25)	2/8 (25)
Thrombocytopenia	3/3 (100)	0/1 (0)	3/4 (75)

*No. positive/No. examined (% positive).

RETROSPECTIVE SEROEPIDEMIOLOGICAL
SURVEY TO DENGUE VIRUS IN
"EL CERRO" MUNICIPALITY HAVANA
C U B A.

INSTITUTO DE MEDICINA TROPICAL "PEDRO KOURI" LA HABANA, CUBA.

The well known history of Dengue in Cuba, with two epidemics reported, the first, of Classical Dengue, in 1977 (DEN-1) and the second (DEN-2) in 1981 of Dengue Hemorrhagic Fever Dengue Shock Syndrome (DHF/DSS), the lack of evidence of a wide circulation of these viruses prior to 1977 and the complete control of the outbreak since October 10 1981, allows the performance of retrospective studies on the basis of a very well defined epidemiological situation. For these reasons we consider that the retrospective seroepidemiological study carried on in an urban municipality of Havana city (El Cerro) will give us relevant information about the infection rates, related with sex, age and race; will also help to establish the proportions between the number of Classical Dengue and DHF/DSS cases, as well as the relation, both in children and adults, between the severe picture of the disease and the primary or secondary type of serological response. We will be able to validate some data obtained in different previous studies, and to go deeper in the epidemiological analysis of the disease.

To this end, we selected a representative sample of 1944 persons from whom a blood sample, in filter paper, and a filled questionnaire was obtained. In each sample we determined the presence of neutralizing antibodies to Dengue 1 and D2 viruses. In those individuals that reported a severe clinical picture, the clinical charts were reviewed at the hospitals.

THE MAIN RESULTS OBTAINED WERE:

- During the Dengue 1 epidemic (1977) 868 persons (45%) were infected and 652 (33.5%) during the Dengue 2 epidemic in 1981. This indicates the wide viral circulation occurred in both epidemics, which is a very important factor for the development of DHF/DSS outbreaks.
- The population at risk of infection was approximately the same in both epidemics. In table 1 we can see that 50% of those persons infected in 1977 had a second infection in 1981, while in those non infected during the first epidemic only a 20% of infection rate could be proved in 1981. When we made the same kind of analysis but considering the houses in which some inhabitant was infected, similar results were obtained (table 2). We consider that these facts depend on the presence of the mosquitoes in the houses, and that the infection within the house, at least in this municipality, was of utmost importance.
- In both epidemics, the infection rates in female were significantly higher (table 3), when we related this rates with age, this difference was only evident in adults. In table 4 we show the relation between infection with Dengue 2 virus and employment. The infection rates were similar in those who worked outside the house with independence to sex; additionally most of non infected adults (72%) were employees (worked outside the house). This again point out the importance of the house as the infecting site. The most the time a person stays in the house, the higher the probability of been bited by th vector. We consider that the rate of infection was not dependant of sex but of the time of permanence in the house.
- In previous studies of our group, the white race was identified as a risk factor for DHF/DSS. In this study we demonstrated that in both epidemics whites and blacks were equally infected (table 5).

The ratio between clinical and subclinical diseases was 1 to 4 in whites and 1 to 8 in blacks, this may indicate a higher frequency of asymptomatic disease in black people and could be related to a natural resistance of this race, to the disease. It must be emphasize that none of the DHF/DSS cases found in this study was black.

- DHF/DSS was found in four persons (table 6), with secondary type of serological response. This finding supports Halstead's hypothesis of the sequential infection as a risk factor for DHF/DSS.

- When we analyzed the ratio of severe cases in relation to age we found that one out of every 34.5 children and one out of every 183.5 adults had DHF/DSS. This indicates the higher frequency of the severe disease in childhood. The probability of a children to suffer DHF/DSS was 0.03 and that of adults 0.005 (six times lower).

Dra. María G. Guzmán

Prof. Gustavo Kouri

Dr. José Bravo

Lic. Maritza Soler

Lic. Luis Morier

Lic. Susana Vázquez

TABLE 1: POPULATION AT RISK IN BOTH EPIDEMICS
 Susceptible individuals in 1977: 1944

1977 Epidemic	868/1944 (45%) infected	1076/1944 (55%) non infected
RISK OF	SECOND INFECTION	FIRST INFECTION
1981 Epidemic	436/868 (50%) 1X2	216/1076 (20%) 1X5

TABLE 2: HOUSES AT RISK OF INFECTION IN BOTH EPIDEMICS
 Total number of houses: 706

1977 Epidemic	444/706 (63%) some person infected	262/706 (37%) none person infected
RISK OF	some person SECOND INFECTION	some person FIRST INFECTION
1981 Epidemic	302/444 (68%) 1X1.4	94/262 (36%) 1X2.7

TABLE 3: INFECTION AND SEX

	1977 Epidemic infection rate	1981 Epidemic infection rate
FEMALE	526/1122(47%)	410/1122(36.5%)
MALE	344/822(42%)	242/822(29%)

TABLE 4: INFECTION AND EMPLOYMENT

	Employees	Non employees
FEMALE	136/451(30%) infected with Dengue 2.	213/408(52%) non infected
MALE	138/473(29%) in fected with Dengue 2	62/115(54%) non infected

TABLE 5: INFECTION AND RACE

	1977 Epidemic	1981 Epidemic
WHITE	468/1076(43%)	382/1076(35.5%)
BLACK	166/342(48.5%)	115/342(34%)

TABLE 6: SEVERE CASES

CASE	TYPE OF Infection	SEX	AGE	RACE	C. D. *
1	second	M	25	white	-
2	second.	M	10	white	-
3	second.	F	9	Mulatto	Asthma
4	second.	F	17	Mullato	Asthma

* Chronic Disease

SOME CLINICAL ASPECTS OF FATAL
CASES OCCURED IN DHF/DSS CUBAN
EPIDEMIC

INSTITUTO DE MEDICINA TROPICAL "PEDRO KOURI". LA HABANA, CUBA.

During the DHF/DSS epidemic occurred in Cuba in 1981, 158 fatal cases were reported (101 children and 57 adults). The analysis of these fatal cases is of utmost importance since this is the first epidemic reported in America.

In 98 clinical charts (76 children and 26 adults), the evolution of the disease from the first symptoms to death, was analyzed. Fever and/or vomiting were the most frequently reported chief complainment both in children and adults, followed by the hemorrhagic manifestations. The 3 to 5 years age group was the most affected, being 4 the modal age. As compared with the Cuban population we could not find sex predominance ($p > 0.05$).

In relation to race, 80% were whites, 11% mulattos and 8% blacks were reported, this distribution shows a significant predominance of the white race as compared with the distribution of the Cuban population ($p < 0.05$).

23% of children and 13% of adults had personal antecedents of Bronchial Asthma; there was a family history of Diabetes Mellitus in 24% of the fatal cases in children, and personal history in 4% of adults. Sickle cell anemia had also a significant higher prevalence as compared with the prevalence in the Cuban population. The white race, as well as the chronic diseases mentioned above were considered as individual risk factors for DHF/DSS, and were identified for the first time during the Cuban outbreak.

Hematemesis was the most frequently reported mayor hemorrhagic manifestation, both in children and adults (Table 1). As there was an official policy of early admittance to the hospital, most of the mayor hemorrhages were recorded during the evolution in the hospital, in contrast with the minor hemorrhagic symptoms (purpura) that were more frequently seen before admittance. In general, 99% of children and 65% of adults had some hemorrhagic manifestation. Gastrointestinal bleeding was very frequently observed.

Fever, nausea, vomiting and abdominal pain were the most relevant sings and symptoms in children and adults. Hepatomegaly was reported in 78% of children and 35% of adults and with the abdominal pain were considered sings of bad clinical prognosis.

Shock was observed in 82% of children and in 100% of adults. In adults fatal cases, the evolution of the disease was almost always toward death, when shock was present.

In table 2 we can see the frequency of hemoconcentration and thrombocytopenia, both in children and adults, these two signs were of utmost importance for the diagnosis of the disease.

Dr. José Bravo

Prof. Gustavo Kourí

Dra. María G. Guzmán

TABLE 1 : MAYOR HEMORRHAGIC SYMPTOMS IN FATAL CASES.

<u>SYMPTOMS</u>	<u>CHILDREN</u>	<u>ADULTS</u>
Hematemesis	43/72 (60)	9/26 (35)*
Melena	14/72 (20)	1/26 (4)
Enterorrhagia	6/72 (8)	2/26 (8)
Metrorrhagia**	1/39 (3)	4/16 (25)

* cases with the symptom / total (%)

** female cases.

TABLE 2 : HEMOCONCENTRATION AND THROMBOCYTOPENIA IN FATAL CASES.

	<u>CHILDREN</u>	<u>ADULTS</u> *
Hemoconcentration	69/72 (96)	22/24 (92)
Thrombocytopenia	70/72 (97)	17/24 (71)

* Two cases not reported.

THE LAPSE BETWEEN THE TWO INFECTIONS WITH DENGUE VIRUS AS A RISK FACTOR FOR FHD/SSD.

INSTITUTO DE MEDICINA TROPICAL "PEDRO KOURI". LA HABANA, CUBA.

Homologous neutralizing antibodies developed after an infection with Dengue viruses lasts all life span, on the contrary, the heterologous neutralizing antibodies, that are also produced disappears in a very short time.

Halstead reported the presence of neutralizing and enhancing antibodies in immune individuals to one of the four dengue serotypes. When these individuals are infected by a different serotype (secondary infection) DHF/DSS may appear if the immunoenhancement phenomenon takes place.

In order to know the influence of time interval between the primary and secondary infections upon the enhancing capacity of the antibodies, we selected 18 persons infected by DEN 1 virus in 1977/78. Two blood specimens were taken in filter paper in 1983, 1984 and 1985 from each person. One of the samples, from each year, was eluted and tested in growing dilutions with DEN 2 (cuban strain) in order to determine the Neutralizing Index. This index decreased with time.


To determine the enhancing capacity of anti DEN 1 antibodies towards DEN 2 virus, the other filter paper from each individual by year eluted and serial dilutions, from 1/30 to 1/1222880 were made. Appropriate amounts of these dilutions, together with the DEN 2 cuban strain (low passage) in a multiplicity of 0.01 were inoculated in the P388D1 murine macrophage cell line. The viral concentration was determined in order to calculate the enhancing titre. Furthermore, by using the Detre and White's formula, the significance of the enhancement level obtained was determined. A decrease in the enhancing capacity of anti DEN 1 antibodies when

tested with DEN 2 virus during the studied lapse was observed. In graph 1 we can see the number of individuals that exhibited, by year, a significant enhancement.

A high percentage of tested sera showed a significant enhancement level five years after the first infection, which means that at that moment they had the possibility to develop the severe clinical picture of the disease. The percentage diminished from 91.3% in 1983 (5 years) to 75% in 1984 (six years) and finally to 33% in 1985, seven years from the first infection. The results obtained indicate that those persons previously infected by a certain serotype of Dengue virus (in our case DEN 1) had the highest risk to develop DHF/DSS if they are infected with a different serotype within the next five years. This risk decreases with time but still exist at least until 7 years after the primary infection.

Lic. Luis Morier
Dra. Maria G. Guzmán
Prof. Gustavo Kourí
Lic Maritza Soler
Lic. Maria R. Alemán

GRAF. I. - PER CENT OF CASES WITH SIGNIFICATIVE ENHANCEMENT
BY YEAR



«O»

FIRST HUMAN YELLOW FEVER VIRUS ISOLATION IN C.A.R.

Two strains of Yellow Fever (HB1504, HB1782) have been isolated from two fatal human cases in December 1985 and February 1986 in the Central African Republic (CAR).

The first case occurred in a 36 year-old male who presented an hemorrhagic syndrome and died a few hours after his hospital admission. He used to live in Berberati in the damaged wet tropical forest environment.

The second case, a 25 years old male, originated from near Bangui (Kapou, 20 Km south of the capital), in a similar ecosystem of a Congo-Crimean phytogeographical domaine. The clinical picture was a severe hepatonephritis syndrome. The patient died 2 days after his hospital admission.

Both cases happened at the end of the rainy season beginning of the dry season.

We performed an environment investigation in Kapou area. At the same time a Yellow Fever immunization campaign was performed by the Public health department. 65 young children were bled for an antibodies survey, 60 % of the 0-9 age group showed antibodies against Yellow Fever. A previous survey in 1978 showed that 16 % or less were positive by HIA against Yellow Fever in the age group 0-9 year. However the last immunization campaign was performed in 1982. 62 mosquito monospecific pools caught in Kapou area and 37 tick pools were used for virus isolation in suckling mice after intrathoracic (Aedes aegypti) and intracoelomic (Amblyomma variegatum) inoculation for virus enhancement. No virus were isolated. 19.5% of the mosquitoes collected on the field belong to Mansonia genus, 78.3% are distributed among 11 mosquito groups not known as potential vectors of Yellow Fever Virus.

Since 1955 no fatal human case of Yellow Fever has been reported in CAR, furthermore these human strains are the first isolated in CAR.

Up to now, Yellow Fever was considered as enzootic in

CAR and seroconversion in human population was observed without any epidemic manifestation nor severe clinical picture.

These virus isolates show active circulation of the Yellow Fever virus in CAR. Moreover another human case of Yellow Fever has been identified by specific IgM immunocapture assay without virus isolation; the patient came to the hospital for severe hepatitis with a quick evolution into renal failure and deep coma, death occurring at day 6 following the onset (A.J.Georges, unpublished). It is usual to isolate Yellow Fever Virus in CAR at the end of the rainy season, beginning the dry season. Concerning Kapou, no virus isolation were made from the environment survey probably because of the delay due to technical difficulties after notification of the case. The high prevalence of Mansonia, potential vector of Yellow Fever, can incriminate it in such observation; but in the absence of mosquito virus isolation nothing can be included for sure.

Nevertheless it is unusual to have Yellow Fever human cases in such an ecosystem. These can be explained by the fact that the "emergence zone", previously described seems to be displaced closer from the rain forest. The reason is anthropic deforestation transforming the moist forest into a forest-savannah mosaic favoring Yellow Fever Virus passage from its selvatic cycle to the human environment.

(A.J.GEORGES, M.GERMAIN, M.C.MADELON, G.GRESENGUET, J.M.DIEMER, J.L.LESBORDES and J.P.GONZALEZ)

VIRUS OF BEDBUGS

The bedbug, Cimex lectularius (Cimicidae:Hemiptera), is a cosmopolitan ectoparasite which is an obligatory haematophagous bug. It is associated with overcrowded insanitary habitation of man. Until recently there had been no evidence that Cimex lectularius harbours any virus. While examining the development of a bat trypanosome Trypanosoma (Megatrypanum) incertum in the bedbug we found large numbers of cytoplasmic arrays of virus-like particles in the epithelial cells of the ventriculus of this insect (Eley et al, 1987). The virus particles which were present in both adult and immature stages purified on 10-50% (w/v) sucrose gradients. The virions had a diameter of 56nm and appeared to be composed of an outer and inner shell. Nucleic acids contained within the virions was found to consist of 11 segments of dsRNA with a size range of 2.75 to 0.15 x 10. These particles, while having the morphology of reovirus are obviously somewhat smaller and they contain one segment too many to be members of the reovirus genus. The virions are possibly "misplaced" rotaviruses, although current dogma states that no arthropod vectors are involved in the transmission of rotaviruses. We were unable to cultivate the purified virions in suckling mice brain and a variety of insect and mammalian cell lines. Hence it is impossible to say if the "extra" dsRNA is the result of a mixed infection as the virus cannot as yet be plaque purified.

While this virus could be restricted to bedbugs the site of infection could make it transmissible during blood meals. The surface antigen of hepatitis B virus was detected in the bedbugs Cimex hemipterus from bedding in West African villages (Wills et al, 1977).

Further investigations of the aetiology of this virus is rendered more significant by a recent report (Lyons et al 1986) that Human Immunodeficiency Virus (HIV) can survive for 1 h in Cimex lectularius after feeding on blood-virus mixtures; these authors suggested that mechanical transmission by bedbugs was a possible route of transmission although Piot and Schofield (1986) have argued that the possibility of vector borne HIV infection is remote.

References

Eley S M, Gardner R A, Molyneux D H and Moore N F (1987). A reovirus from bedbugs, Cimex lectularius. J.gen.Virol., 68, 195-199.

Lyons S F, Jupp P G and Schaub B D (1986). Survival of HIV in the common bedbug. Lancet. II, 45.

Piot P and Schofield C S (1986). No evidence for Arthropod transmission of AIDS. Parasitology Today, 2, 294.

Wills W, Larowze S, London W T, Millman I, Werner B G, Ogston W, Pourtaghva M, Diallo S and Blumberg B S (1977). Hepatitis B virus in bedbugs Cimex hemipterus from Senegal. Lancet (North American Ed) 2:217-219.

1 2 2 1
S M Eley, R A Gardner, D H Molyneux and N F Moore

1
Current Address, Chemical Defence Establishment, Porton Down, Salisbury, Wilts
SP4 OJQ

2
Department of Biological Sciences, University of Salford, Salford M5 4WT

VIRUS OF BEDBUGS

The bedbug, Cimex lectularius (Cimicidae:Hemiptera), is a cosmopolitan ectoparasite which is an obligatory haematophagous bug. It is associated with overcrowded insanitary habitation of man. Until recently there had been no evidence that Cimex lectularius harbours any virus. While examining the development of a bat trypanosome Trypanosoma (Megatrypanum) incertum in the bedbug we found large numbers of cytoplasmic arrays of virus-like particles in the epithelial cells of the ventriculus of this insect (Eley et al, 1987). The virus particles which were present in both adult and immature stages purified on 10-50% (w/v) sucrose gradients. The virions had a diameter of 56nm and appeared to be composed of an outer and inner shell. Nucleic acids contained within the virions was found to consist of 11 segments of dsRNA with a size range of 2.75 to 0.15 x 10. These particles, while having the morphology of reovirus are obviously somewhat smaller and they contain one segment too many to be members of the reovirus genus. The virions are possibly "misplaced" rotaviruses, although current dogma states that no arthropod vectors are involved in the transmission of rotaviruses. We were unable to cultivate the purified virions in suckling mice brain and a variety of insect and mammalian cell lines. Hence it is impossible to say if the "extra" dsRNA is the result of a mixed infection as the virus cannot as yet be plaque purified.

While this virus could be restricted to bedbugs the site of infection could make it transmissible during blood meals. The surface antigen of hepatitis B virus was detected in the bedbugs Cimex hemipterus from bedding in West African villages (Wills et al, 1977).

Further investigations of the aetiology of this virus is rendered more significant by a recent report (Lyons et al 1986) that Human Immunodeficiency Virus (HIV) can survive for 1 h in Cimex lectularius after feeding on blood-virus mixtures; these authors suggested that mechanical transmission by bedbugs was a possible route of transmission although Piot and Schofield (1986) have argued that the possibility of vector borne HIV infection is remote.

References

Eley S M, Gardner R A, Molyneux D H and Moore N F (1987). A reovirus from bedbugs, Cimex lectularius. J.gen.Virol., 68, 195-199.

Lyons S F, Jupp P G and Schaub B D (1986). Survival of HIV in the common bedbug. Lancet. II, 45.

Piot P and Schofield C S (1986). No evidence for Arthropod transmission of AIDS. Parasitology Today, 2, 294.

Wills W, Larowze S, London W T, Millman I, Werner B G, Ogston W, Pourtaghva M, Diallo S and Blumberg B S (1977). Hepatitis B virus in bedbugs Cimex hemipterus from Senegal. Lancet (North American Ed) 2:217-219.

1 2 2 1
S M Eley, R A Gardner, D H Molyneux and N F Moore

1
Current Address, Chemical Defence Establishment, Porton Down, Salisbury, Wilts SP4 OJQ

2
Department of Biological Sciences, University of Salford, Salford M5 4WT

REPORT FROM THE VIROLOGY PROGRAM
STATE OF NEW JERSEY DEPARTMENT OF HEALTH
TRENTON, NEW JERSEY

Arbovirus Surveillance in New Jersey, 1986

During the 1986 surveillance period from June into October, 3812 mosquito pools containing up to 100 mosquitoes each were tested for viruses in day old chicks. There were twenty-two (22) mosquito pools positive for Eastern encephalitis (EE) and Western encephalitis (WE) was isolated from thirty-five (35).

Table 1 summarizes the collection area totals, species of mosquito and time of collection for the EE isolates. Activity began with the mid August collections and continued into October. All twenty-two (22) isolates were from pools containing Culiseta melanura mosquitoes at four (4) sites.

WE mosquito activity is summarized in Table 2. The July collections yielded the first isolates with continued observation of WE activity into October. There were thirty-five (35) isolates from Culiseta melanura at four (4) sites.

An EE isolate was also made in September from a horse in Atlantic County.

Sentinel chicken flocks of six (6) cockerals were placed at five (5) sites. The flocks were bled weekly and St. Louis encephalitis hemagglutination inhibition tests were conducted. There were no conversions observed in the 320 sera tested.

(Shahiedy I. Shahied, Bernard F. Taylor, Wayne Pizzuti)

Table 1
 1986

AREA COLLECTED	MOSQUITO SPECIES	EE MOSQUITO POOL ISOLATES FOR WEEK ENDING										AREA TOTALS		
		8/15	8/22	8/29	9/5	9/12	9/19	9/26	10/3	10/10	10/17			
Bass River	Cs.melanura		1											1
Dennisville	Cs.melanura		1	1		1				1				4
Green Bank	Cs.melanura	3	2	5	2		1			1	1	1		16
Ocean City	Cs.melanura										1			1
WEEKLY TOTALS		3	4	6	2	1	1	0	3	1	1			22

Table 2
 1986

AREA COLLECTED	MOSQUITO SPECIES	WE MOSQUITO POOL ISOLATES FOR WEEK ENDING														AREA TOTALS
		7/4	7/11	7/18	7/25	8/1	8/8	8/15	8/22	8/29	9/5	9/12	9/19	9/26	10/3	
Bass River	Cs.melanura							1	1	1		1	1			5
Centerton	Cs.melanura							1	1				2	1	1	6
Dennisville	Cs.melanura	1			3	4	1	5	2	1				1	1	19
Green Bank	Cs.melanura							1	1	1			1	1		5
WEEKLY TOTALS		1	0	0	3	4	1	8	5	3	0	1	4	1	3	35

The detection of the antigen and antibodies to the eastern
subtype of hemorrhagic fever with renal syndrome /HFRS/
in small rodents in Slovakia

M. Grešíková¹, O. Kožuch¹, M. Sekeyová¹, E. A. Tkačenko²,
G. V. Rezapkin²

1 Institute of Virology, Slovak Academy of Sciences,
817 03 Bratislava, Czechoslovakia;

2 Institute of Poliomyelitis and Viral Encephalitis,
Moscow, ZSSR

In 1984 we detected the antigen of hemorrhagic fever
with renal syndrome /HFRS/ in the lungs of small rodents
in Eastern Slovakia. It was found by the complement-fixation
reaction that the antigens were closely related to the wes-
tern subtype of HFRS. Antibodies to HFRS were detected in
sera of *Clethrionomys glareolus*, *Apodemus agrarius*, *Pitymys*
subterraneus collected in Eastern Slovakia and in the sera
of *Clethrionomys glareolus*, *Apodemus sylvaticus*, *Microtus*
arvalis and *Microtus economus* collected, in western Slovakia.
In the present study we used the ELISA technique for the
detection of HFRS antigen which is less time consuming than
immunofluorescence procedure.

During 1985-1986 years small rodents were captured alive
in Eastern and Western Slovakia. In the laboratory, the ani-
mals were autopsied and the lung specimens were collected;

148 specimens were collected in Western Slovakia and 47 specimens in Eastern Slovakia.

For the detection of HFRS antigen from the lungs of small rodents, enzyme-immunoassay /ELISA/ tests were used to select positive specimens. Out of 195 samples of small mammals of following species: *Apodemus flavicollis* /51 samples/, *A. sylvaticus* /48 samples/, *Clethrionomys glareolus* /74 samples/, *Microtus arvalis* /16 samples/, *Apodemus agrarius* /6 samples/, four antigens were detected by the ELISA method /Table 1/. The HFRS antigen was detected in the lungs of *Apodemus agrarius*, collected in Eastern Slovakia and *Microtus arvalis*, collected in Western Slovakia. The HFRS antigen detected in the lungs of *A. agrarius* was closely related to the eastern subtype of HFRS; the antigens detected in the lungs of *M. arvalis* were closely related to the western subtype of HFRS /Table 2/. The results of serological survey on 47 sera of small rodents collected in Eastern Slovakia with the antigen of HFRS indicate the existence of natural focus of eastern subtype of HFRS in Eastern Slovakia. Antibodies were found in the following free-living rodents: *Apodemus flavicollis* and *A. agrarius*. A total 5 out of 47 small rodents trapped in investigated localities of Eastern Slovakia had antibodies to HFRS antigen /Table 3/. The higher titers of antibodies were detected with the eastern subtype of HFRS antigen.

It is of interest that in *Apodemus agrarius* No 317 and No 320, the viral antigen of HFRS was detected in the lungs simultancously with antibodies.

Table 1

Detection of viral antigen of HFRS in the lung suspension of small rodents by enzyme-immunoassay /ELISA/

Species	No of examined species	No of positive by ELISA
Apodemus flavicollis	51	0
Apodemus agrarius	6	2 ⁺
Apodemus sylvaticus	48	0
Clethrionomys glareolus	74	0
Microtus arvalis	16	2 ⁺⁺

+ = The eastern subtype of HFRS

++ = The western subtype of HFRS

Table 2

Comparison of antigens of hemorrhagic fever with renal syndrome /HFRS/ with eastern and western type of antiserum

Strain No./isolated from	Western type of HFRS antiserum	Eastern type of HFRS antiserum
317/ Apodemus agrarius	0	+
320/ Apodemus agrarius	0	+
141/ Microtus † arvalis	+	0

† One antigen detected in *M. arvalis* was not sterile, therefore it was not used in the experiments

Table 3

Comparison of antibody titres to HFRS in small rodents by IF tests with the eastern and western subtypes of HFRS antigen

IF antibodies in species No	IF titres with Eastern type of HFRS antigen	IF titres with Western type of HFRS antigen
Apodemus flavicollis No 291	64	16
A. flavicollis No 294	128	16
A. flavicollis No 313	32	< 16
Apodemus agrarius No 317	64	< 16
A. agrarius No 320	128	< 16

HFRS = Hemorrhagic fever with renal syndrome

IF = Immunofluorescence

THE HUT TAMPAN, ORNITHODOROS MOUBATA MURRAY (ARACHNIDA : ARGASIDAE) AS VECTOR
OF HEPATITIS B VIRUS IN NAMIBIA.

Two earlier studies showed that the human population of the territories of Northern Namibia bordering on Angola had a high positivity rate for hepatitis B virus (HBV) (Bersohn *et al*, 1974, Botha *et al*, 1984). In 1985 further observations were reported in regard to one of these territories, the Kavango region, situated in the north eastern corner of the country. These showed that only 1.9% of the subjects tested were negative for all the HBV markers used indicating that infection had been present at some stage in 98.1% of the sample (Joubert *et al*, 1985¹). Since none of the usual transmission methods for HBV seemed to explain this high positivity rate in Kavango, the hut tapan tick, Ornithodoros moubata Murray, came under consideration as a possible vector. This argasid tick was found to be very prevalent occurring in about 80% of the huts of the local people where it also acts as a vector of Borrelia duttoni, the cause of relapsing fever (Joubert *et al*, 1985²). Samples of tapan ticks were collected in huts and infection rates of 150.4 and 26.4 per 1000 ticks were demonstrated for the HBsAg and HBeAg respectively in these ticks (Joubert *et al*, 1985²). These high infection rates suggested the tapan probably acted as a mechanical vector of HBV in a similar manner to that in which the bedbug, Cimex lectularius, probably acts in the northern Transvaal (Jupp *et al*, 1983). It was therefore decided to assess the vector competence of O. moubata in the laboratory to see whether such experiments would provide further evidence incriminating it as a vector of HBV.

Wild-caught and colonized tapan ticks (Ornithodoros moubata) were fed on Hepatitis B virus (HBV) - positive blood-meals in a series of four experiments. HBsAg persisted in nymphal and adult ticks for up to 779 days, while HBeAg persisted in mature nymphs up to 13 days, in adult males up to 11 days and in adult females up to 16 days. HBsAg was transmitted trans-stadially through two moults during the life cycle but transovarial transmission did not occur. The surface antigen was transmitted by two out of 15 single ticks into 0,4 ml aliquots of HBV-negative blood, although 6 groups of ticks failed to transmit into 5,5 ml aliquots of blood: this antigen was not transmitted to hamsters. HBsAg was detected in samples of the ticks' coxal and rectal fluid secretions always at the infecting feed and usually at the second feed. HBeAg was only detected in one of 2 samples of coxal fluid collected at the infecting feed. The results as a whole indicate that no biological multiplication of virus occurs in O. moubata but that mechanical transmission from ticks to man could occur by: (i) contamination of a person when crushing infected ticks; (ii) infection by bite; (iii) contamination with coxal fluid, especially by scratching bites. This is thought to take place among the Kavango tribe in their village huts in north eastern Namibia where infestations of infected O. moubata occur.

This work was done in co-operation with J.J. Joubert, C. Swanevelder and O.W. Prozesky of the Institute for Pathology, Pretoria University.

References

- Bersohn, I., MacNab, G.M., Pyzikowska, J. and Kew, M.C. (1974). The prevalence of Hepatitis B (Australia) antigen in Southern Africa. South African Medical Journal 48: 941-944.
- Botha, J.F., Dusheiko, G.M., Ritchie, M.J.J., Mouton, H.W.K. and Kew, M.C. (1984). Hepatitis B. Virus carrier state in black children in Ovamboland: role of perinatal and horizontal infection. The Lancet 1: 1210-1212.
- Joubert, J.J., Prozesky, O.W., Lourens, J.G.H., Van Straten, A.M.S., Theron, J.W., Swanevelder, C., Meenehan, G.M. and Van der Merwe, C.A. (1985)¹. Prevalence of hepatitis virus and some arbovirus infections in Kavango. South African Medical Journal 67: 500-502.
- Joubert, J.J., Jupp, P.G., Prozesky, O.W. and Lourens, J.G.H. (1985)². Infection of wild populations of the tsetse fly Ornithodoros moubata Murray, with Hepatitis B virus in the Kavango, Northern Namibia. South African Journal of Science 81: 167-168.
- Jupp, P.G., McElligott, S.E. and Lecatsas, G. (1983). The mechanical transmission of Hepatitis B virus by the common bedbug (Cimex lectularius L) in South Africa. South African Medical Journal 63: 77-81.

ARBOVIRUS ANTIBODY SURVEY ON SERA FROM POTENTIAL DENGUE REGIONS

Dengue has not been confirmed in South Africa since 1926-7, but outbreaks in East Africa, the most recent being in Mozambique in 1984, have caused concern that the virus could be re-introduced. A survey was carried out on human serum samples collected from Natal, Kwazulu and eastern Transvaal. These sera were tested against various arbovirus antigens including dengue to provide up to date information on arthropod-borne virus activity in these regions.

A total of 2233 sera were collected from 5 main sources: employees at Cato Ridge Abattoir in Durban, Natal hospitals, Kruger National Park, inpatients and outpatients at 4 Kwazulu hospitals and routine specimens submitted to the laboratory. The sera were screened using the fluorescent antibody technique on multispot slides. Positives were tested by the haemagglutination-inhibition (HAI) method with the following results:-

Source	No.	HAI Positive ($\geq 1:20$)					
		SIN	CHIK	WSL	WN	DEN	RVF
Cato Ridge	604	2	0	1	18	3	8
Natal hospitals	866	13	0	3	29	0	0
Kruger Nat. Park	481	16	9	12	18	0	3
Routine specimens	107	10	2	8	14	6	0

Source	No.	HAI Positive								
		SIN	CHIK	WSL	WN	DEN	RVF	YF	ING	GERM
Hospital Patients Kwazulu	178	2	2	0	7	0	5	5	2	19

Any dengue HAI positives were tested by the more specific complement fixation (CF) test.

The results of CF tests on dengue HAI positive sera were as follows: one from Cato Ridge was positive for dengue, this person resided in Durban at the time of the 1926-27 epidemic. Four of the six routine sera were suspicious but because of the cross reactivity with other flaviviruses could not be confirmed as dengue, one was confirmed by isolation of dengue type 1 virus and one was negative. All of these routine sera were from patients who had recently returned from India where dengue is endemic.

The conclusion from this survey was that there had been relatively low arbovirus activity in the areas tested and the population would probably be susceptible to dengue virus infection. Dengue virus was introduced into Durban by an individual returning from an endemic region, fortunately it does not appear that the virus has become established.

PRESENCE OF ARBOVIRUS ANTIBODIES IN DOGS

Dog sera from the eastern Free State and from Pretoria North were tested for the presence of arbovirus antibodies. A selection of negative sera were submitted for virus isolation.

Haemagglutination inhibition results on dog sera.

Number with HI titre \geq 1:20

No. tested	West Nile	Wesselsbron	Sindbis	RVF
377	174	57	46	0

All of the dogs with antibodies to Wesselsbron virus had higher titres to West Nile virus, so these are unlikely to be as a result of Wesselsbron virus infection. Seventy-nine per cent of the WN HI positive sera and 28 per cent of the SIN HI positive sera were positive by the neutralisation test. Because of the high incidence of WN antibodies in dogs and the particularly high titres in some cases, virus isolation was attempted on 110 WN HI negative sera. West Nile virus was isolated from the serum of one of these dogs.

MONOCLONAL ANTIBODIES AGAINST CRIMEAN-CONGO HAEMORRHAGIC FEVER

Seven monoclonal antibodies were prepared against a South African strain of Crimean-Congo haemorrhagic fever (CCHF) virus and were found to be directed against viral nucleocapsid protein. Five of the monoclonal antibodies reacted to high titre in indirect immunofluorescence (IF) tests and enzyme-linked immunosorbent assays with 22 strains of CCHF virus and failed to cross-react with the closest antigenic relative of CCHF, Hazara virus, or with 4 other nairoviruses which need to be distinguished from CCHF virus in Africa. These antibodies, used in the IF technique, readily detected antigens induced by all strains of CCHF virus included in the study in cell culture monolayers and mouse brain tissue, which represent the systems commonly used for isolation of CCHF virus. The IF technique with monoclonal antibodies constitutes a rapid and specific means of identifying newly isolated strains of CCHF virus.

ANTIGENIC RELATIONSHIP OF WEST NILE STRAINS

The antigenic relationship of several South African WN isolates to each other and to the two main WN antigenic groups was assessed. The method used was the calculation of titre ratios or 'R' values from heterologous and homologous neutralisation titres, using the formula $R = 100 \sqrt{r_1 \times r_2}$, R is expressed as a percentage.

Titre ratio or 'R' value between the South African, Indian and Egyptian WN strains expressed as a percentage.

West Nile Virus Strains

	EG101	G2266	H442	735	730	739	778	
EG101	100							
G2266	27	100						
West Nile Virus Strains	H442	16	25	100				
	735	32	7	50	100			
	730	50	10	11	46	100		
	739	33	27	21	30	42	100	
	778	30	21	23	38	65	56	100

None of the local isolates in this group were significantly closely related to either EG101 or G2266. AR20735 and AR20730 were much closer to the Egyptian strain than to the Indian. Contrary to the findings of other

workers, H442 was shown to be antigenically distinct from the Egyptian strain with an 'R' value of only 16 per cent. This is probably due to the NT test being more specific than the modified haemagglutination inhibition test used in that study.

(N.K. Blackburn, P.G. Jupp, T.G. Besselaar, G.M. Meenehan and A.J. Cornel)

ARBOVIRUSES AND SMALL WILD MAMMALS IN WESTERN TURKEY: A SEROSURVEY.

Turkey is located in the oriental mediterranean basin, Palearctic region. This country is concerned by biannual palearctic-ethiopian migrations of birds and the local climatic conditions are a priori favourable to the circulation of some arboviruses. However, relatively few arbovirus studies were so far performed here (Radda, 1971 - Tesh et al, 1976 - Serter, 1980).

During entomological surveys carried out in 1981, in the Western part of the country, 53 wild small mammals (rodents, insectivora) were caught alive and their sera collected. They were studied for antibody against 17 different arboviruses belonging in 8 different genera.

The results gave good evidence of the possible activity of a Phlebovirus in the surveyed area.

MATERIAL AND METHODS

Small wild mammals were trapped in May 1981 in the vicinity of Gölcük, near Odemis, Vilayet Izmir, about 70 km in the south-east of Izmir town (map). The mean altitude was about 1,100 meters.

Sera were obtained on blotting paper strips from 33 Apodemus sylvaticus, 6 Mus musculus, 16 Pitymys majori, 2 Microtus epiroticus and 2 Crocidura suaveolens.

Sera were studied by hemagglutination inhibition (HI) and complement fixation (CF) tests using micromethods. We used 12 antigens in HI tests: Sindbis (SIN), Chikungunya (CHIK), West Nile (WN), Dengue type 2 (DEN2), Wesselsbron (WSL), European tick-borne encephalitis (TBE), Meaban (MEA), Bhanja (BHA), Tahyna (TAH), Sandfly fever Sicilian (SFS) and Arumowot (AMT). In CF, five antigens were tested: Qalyub (QYB), Quarantfil (QRF), Tribec (TRB), Essaouira (unregistered, Chenuda complex, Orbivirus), and Soldado. Essaouira and Soldado have been isolated previously in Morocco from Ornithodoros (A.) maritimus ticks infesting herring gull'nests (Larus cachinnans) on Essaouira island (Chastel et al, 1981, 1982).

RESULTS

By HI tests, 3 sera (or 5.7%), all from field mice (A. sylvaticus), were found positive for the Arumowot (AMT) antigen only. Two sera reacted with a titer of 1:20 and the third was a titer of 1:40. No cross reaction was observed with SFS virus, another phlebovirus. All other HI and CF tests were negative (Table 1).

COMMENTS

Previous serosurveys were performed in human beings by Tesh et al (1976) on Antalya area and by Serter (1980) on the Turkish coasts of Egean sea (map).

Tesh et al demonstrated neutralizing (NT) activity to Sandfly fever Naples virus (62% of sera) and to SFS virus (22%).

Serter used another phleboviruses as antigens and found antibody to Salehabad virus (0.46% of sera), a phlebovirus giving cross serological reaction with AMT virus. Serter also found antibody to SIN, WN, Dengue type 1, TBE, SFS and TAH viruses.

Radda (1971) examined sera from domestic animals and wild rodents from Ankara area in Anatolia, and from Iskenderun and Antakya areas near the northern border of Syria. He found antibody to flaviviruses (WN, Murray Valley encephalitis, DEN2) in sheep, cattle and 'field mice' (species not recorded).


Our results gave evidence of the possible circulation of a phlebovirus, AMT or a virus very close to it, in the Gölcük area, Vilayet Izmir.

AMT virus is a widely transmitted phlebovirus circulating in Africa among mosquitoes (Culex antennatus, C. rubinotus) and rodents or insectivora: Tatera kempii (Gerbillidae), Arvicanthis niloticus, Ithamnomys macmillami, Mastomys sp. (Muridae) and Crocidura sp. (Soricidae).

New entomological and virological studies are needed in order to confirm the actual circulation of this ubiquitous virus in Western Turkey.

REFERENCE

- Le Lay-Roguès G., Hardy-Le Maux E., Ohran V., Beaucournu J.C. and Chastel C. Arbovirus et petits mammifères sauvages en Turquie: note brève. Bulletin de la Société de Pathologie Exotique, Paris, 1986, 79, 595-601.



Studied species	Number of positive reactions /Number of sera studied
Rodents:	
Muridae	
<i>Apodemus sylvaticus</i>	3/33
<i>Mus musculus</i>	0/6
Arvicolidae	
<i>Pitymys majori</i>	0/11
<i>Microtus epiroticus</i>	0/2
Insectivora:	
Soricidae	
<i>Crocidura suaveolens</i>	0/1
TOTAL	3/53 (5.7%)

Table 1. Animal species studied and positive reactions to Arumowot virus Western Turkey (1981).

Viral and Rickettsial Disease Laboratory, California Dept. of Health Services

Following an unprecedented occurrence of epidemic St. Louis encephalitis in urban and suburban areas of southern California during 1984, when 26 cases (1 fatal) were confirmed, expanded surveillance and control measures were taken in this region as well as in the traditional endemic areas of the Sacramento, San Joaquin, and Imperial Valleys. Data from 1984 and the early part of 1985 were given in previous issues of this Exchange. Fortunately, 1985 and 1986 seasons were not so severe as in 1984. During 1985, only 3 human cases of SLE were detected: a 17 year old boy from Riverside County; a 31 year old man from San Bernardino County, with exposure most probably along the Colorado River near Needles; and a 61 year old woman from the North Hollywood area of Los Angeles County. No equine or human cases of WEE were found. From 4,417 mosquito pools tested, 30 SLE, 28 WEE, 18 CEV group, 48 Turlock, 122 Hart Park, and 2 Main Drain virus isolates were recovered. All isolates were from Culex tarsalis, except for 15 CEV group from Aedes melanimon, 1 Turlock and 2 Hart Park from C. pipiens complex, 1 Turlock from C. erythrothorax, 2 Main Drain from Aedes taeniorhynchus, and 1 SLE from C. peus. Of 55 sentinel chicken flocks sampled monthly in the state, seroconversions for WEE and SLE occurred only in 6 flocks in southern California sites.

The program during 1986 was carried out as in previous years, but was expanded further. As usual, it represented collaborative efforts by local mosquito abatement agencies; local health departments; the California State Department of Health Services (Viral and Rickettsial Disease Laboratory, Vector Surveillance and Control Branch, Infectious Disease Branch, and Veterinary Public Health Unit); the Arbovirus Research Unit, Department of Biomedical and Environmental Health Sciences, School of Public Health, University of California, Berkeley; The California Department of Food and Agriculture; many private physicians and veterinarians; and the Microbiology Reference Laboratory, Cypress, Ca.

Only 3 human SLE cases were detected: (1) a 59 year old man from Covina, Los Angeles Co., with onset July 3, who was a truck driver with routes over a wide area of southern and central California and parts of Nevada, so source of mosquito exposure could not be decided; (2) a 66 year old man from Norwalk, Los Angeles Co., onset July 13, with mosquito exposure at his home; and (3) a 37 year old man from El Monte, Los Angeles Co., onset September 20, source of mosquito exposure most likely at his home site. Diagnoses were confirmed in these cases by CF, IFA, and IFA-IgM serologic tests. Two cases of WEE were found: (1) an infant girl, born May 12, 1986, who became ill August 7. The home environment near Sacramento was the likely site of exposure; and (2) a 37 year old woman from Yolo Co., with aseptic meningitis August 18, and likely exposure either at her home site or on a trip to Sierra Co. 8-10 days earlier. The CF, IFA, and IFA-IgM tests were helpful diagnostically in these cases also. Only 1 WEE case in an equine was confirmed: a 2 year old unvaccinated mare from Shasta Co., with onset September 1.

From 4,536 mosquito pools tested in cell culture systems and/or suckling mice there were 305 virus isolates: 51 SLE, 35 WEE, 47 CEV group, 133 Turlock, 38 HP, and 1 unidentified virus. All WEE and SLE isolates were from C. tarsalis, except 1 WEE from C. erythrothorax, 1 WEE from C. peus, 2 SLE from A. dorsalis, 1 SLE from C. peus, and 5 SLE from C. pipiens complex. There were 59 sentinel chicken flocks bled monthly during the season. Seroconversions for WEE were seen in June (Westside District) and August (Butte and Turlock Cos.), then fairly high seroconversion rates in September and October in northern and central counties. As usual, WEE activity was also seen in southern California (Coachella Valley, Imperial Co., and the Colorado River) beginning in July. Seroconversion for SLE was seen in Imperial Co., the Coachella Valley, Orange Co., Los Angeles Co., Kern Co., a Westside flock, and a Sacramento/Yolo Co. flock. Details of this summary are published each year in the Proceedings and Papers of the annual Conference of the California Mosquito and Vector Control Association, Inc.

R.W.Emmons, M.D.,Ph.D.

Host feeding patterns and domiciliarity development of Ribeira Valley mosquitoes, S. Paulo, Brazil.

Blood-meals identification, from mosquitoes collected in 5 different Ribeira Valley environments, were made through precipitin tests. Sources of 1444 blood-meals were identified showing that Aedes fed chiefly on mammals, while Culex (Culex) showed feeding patterns directed to avian hosts. Ae. scapularis presented preference for cattle and horses. On the whole, Culex (Melanoconion) showed large eclectic pattern including amphibian, avian, mammals and reptilian. Anthropophily was observed for several species, mainly Ae. scapularis, Cx. sacchettae and Cx. ribeirensis who showed high human blood rates in the indoor collections. Endophily studies were made through collections searching resting mosquitoes at the early morning, from all domiciliary and peridomiciliary environments. The use of diversity index and similarity quotient allows to reach some insight about resources and its distribution. So these endophily studies gave data that suggests some degree of endophilic evolution by Ae. scapularis, Cx. ribeirensis and Cx. sacchettae. This behaviour is submitted to the influence of extrinsic factors as cattle density and human behaviour for the environment management.

(Forattini, O.P.; Gomes, A.C.; Natal, D.; Kakitani, I and Marucci, D.).

CRIMEAN-CONGO HEMORRHAGIC FEVER IN SOUTHERN AFRICA.

1 Human disease and distribution of the virus.

From February 1981 to December 1986 42 cases of Crimean-Congo hemorrhagic fever (CCHF) were diagnosed in 29 outbreaks in various locations in South Africa. A further 2 cases which were imported, arose in Zaire and Tanzania while 3 cases were diagnosed in Namibia. 13 patients were infected by tickbite, 16 had contact with livestock tissues, 8 gained nosocomial infection and of the remaining 10 it is merely known that they resided in or visited a rural environment.

The clinical features of infection conformed to the classical descriptions of CCHF in the Soviet Union. Following an incubation period of usually less than a week, patients suffered sudden onset of fever, nausea, prostrating headache and myalgia. Early leukopenia, thrombocytopenia and elevated serum transaminases were followed on days 3 to 5 of illness by a petechial rash and epistaxis, hematemesis and melena. The occurrence of 13/47 deaths indicates that the African disease is no less severe than that which occurs in Eurasia.

Antibody to CCHF virus was detected by reversed passive hemagglutination-inhibition (RPHI) in 2,460/8,667 (28%) cattle sera (140/180 herds) tested in South Africa, as well as in 347/763 (45%) cattle sera (32/34 herds) in Zimbabwe. It is inferred that the virus will yet be found to be widespread in all countries in Africa and Eurasia which lie within the limits of world distribution of ticks of the genus Hyalomma.

.2 Antibody to CCHF virus in wild mammals.

Sera from 3,772 wild mammals of 87 species and from 1,978 domestic dogs collected in South Africa and Zimbabwe between 1964 and 1985 were tested for antibody to CCHF virus by RPHI and/or by indirect immunofluorescence (IF). Antibody was found to be highly prevalent in large mammals in the Orders Artiodactyla and Perissodactyla such as giraffe, Giraffa camelopardalis (3/3 positive), rhinoceros, Ceratotherium simium and Diceros bicornis (7/13), eland, Taurotragus oryx (59/127), buffalo, Syncerus caffer (56/287), kudu, Tragelaphus strepsiceros (17/78), and zebra, Equus burchelli (16/93). In small mammals antibody was found in the sera of 40/293 hares, 22/1,305 rodents and 1/74 wild carnivores but not in 522 primates, 176 insectivores or 129 hyrax. Antibody was also found in the sera of 118/1,978 domestic dogs. The fact that some of the sera had been in storage for over 20 years, indicates that CCHF virus became disseminated in southern Africa a long time ago. The species of wild mammal in which antibody was distributed (with highest antibody prevalence in hares and large herbivores) reflects the feeding

preference of immature and adult ticks of the genus Hyalomma, suggesting that Hyalomma sp. are the principal CCHF vectors in the wild. The results of the study implicate several mammalian species as possible amplifying hosts, but whether they are capable of acting in this regard awaits further study.

3 CCHF infection in 13 mammal and 7 tick species.

Thirteen species of domestic and wild mammals were infected with CCHF virus in order to determine the potential role of each in the virus transmission cycle. Low-titered viremia (maximum 4.0 log₁₀ LD₅₀/ml) followed by development of antibody occurred in sheep, calves, scrub hares (Lepus saxatilis), and 5 rodent species [the white-tailed rat (Myodomys albicaudatus), the bushveld gerbil (Tatera leucogaster), the red veld rat (Aethomys chrysophilus), the striped mouse (Rhabdomys pumilio) and the Cape ground squirrel (Xerus inauris)]. Antibody response only was detected in hedgehogs and a further 4 rodent species [the South African hedgehog (Erinaceus frontalis), the highveld gerbil (Tatera brantsii), the Namaqua gerbil (Desmodillus auricularis) and both species of multimammate mouse Mastomys coucha 2N=36 and M.natalensis 2N=32]].

Ticks failed to become infected when fed on viremic sheep and rodents, but the virus was detected transiently in a few of the Hyalomma truncatum, Amblyomma hebraeum, Rhipicephalus evertsi evertsi, Rh.appendiculatus and Boophilus decoloratus ticks which fed on calves and hares. However, transovarial transmission of virus did not occur.

The virus failed to replicate in 3 argasid tick species (Argas walkerae, Ornithodoros savignyi and O.porcinus porcinus) inoculated intracoelomically, but multiplied to high titer (maximum 6.3 log₁₀LD₅₀/ml) in 4 ixodid tick species (H.truncatum, H.marginatum rufipes, Rh.evertsi mimeticus and A.hebraeum) and persisted in the ticks for up to 205 days. Three of the ixodid tick species (H.truncatum, H.m.rufipes and Rh.e. mimeticus) transmitted CCHF infection to sheep by bite, but transovarial transmission of virus to the progeny of infected females was not detected.

4 CCHF infection in birds.

In November 1984, a case of CCHF occurred in a worker who became ill after slaughtering ostriches (Struthio camelus) on a farm near Oudtshoorn in the Cape province of South Africa. The diagnosis was confirmed by isolation of CCHF virus from the patient's serum and by demonstration of a specific antibody response. It was suspected that infection was acquired either by contact with ostrich blood or by inadvertently crushing infected Hyalomma ticks while skinning ostriches. Reversed passive hemagglutination-inhibition antibody to CCHF virus was detected in the sera of 22/92 ostriches from farms in Oudtshoorn district, including 6/9 from the farm where the patient worked, but not in the sera of 460 birds of 37 other species. In

pathogenicity studies domestic chickens proved refractory to CCHF infection, but viremia of low intensity (maximum titre $2.5 \log_{10}$ mouse ic LD₅₀) followed by a transient antibody response occurred in blue-helmeted guinea fowl (Numidia meleagris). These results constitute the first direct evidence that some bird species are susceptible to CCHF virus infection.

5 Isolation and titration of CCHF virus.

Fluorescent focus assay and plaque assay in CER cells were compared with mouse inoculation for isolation and titration of CCHF virus. Fluorescent focus assay and plaque assay were of similar sensitivity but both produced 10 to 100-fold lower titers than mouse inoculation. In specimens tested from 26 CCHF patients, virus was isolated from 20 in mice and from only 11 in cell culture. Although cell cultures were less sensitive for isolation of virus from clinical specimens they produced diagnostic results much more rapidly.

6 Antigen detection in diagnosis of CCHF fever.

Enzyme-linked immunosorbant assay (ELISA) and reversed passive hemagglutination (RPHA) tests were evaluated for rapid detection of CCHF virus antigens. Both RPHA and ELISA detected CCHF antigen in the brains of infant mice 2 to 3 days after infection; several days before the animals sickened and died. Antigen was also detected after 1 to 2 days in infected cell culture extracts and after 2 to 4 days in culture supernatant fluids. Both tests detected CCHF antigen at threshold values of approximately 10^3 tissue culture infective doses per ml of infectivity and were more sensitive than complement fixation, immunodiffusion or IF. In a comparative study on specimens from CCHF patients, virus was isolated from 38/49 sera and from 23/28 patients. Antigen was detected in 20/49 sera (15/28 patients) by RPHA and in 29/49 sera (18/28 patients) by ELISA. Antigenemia was detected more frequently in fatal cases (9/11) than in non-fatal cases (9/17). The results suggest that RPHA and ELISA may be of use in rapid diagnosis of CCHF infection, particularly in severe cases where the danger of nosocomial spread is greatest.

ARENAVIRUSES IN SOUTHERN AFRICAN RODENTS.

Sera of 4023 rodents collected over a period of 16 years in South Africa and Zimbabwe for unrelated purposes, were tested by IF for antibody to African arenaviruses using cells infected with Lassa fever and Mopeia viruses as antigen. Antibody was found in 15/28 species; most frequently in Aethomys namaquensis [20/136 (14.7%) positive; titers 1:8 - 2048], Mastomys natalensis (2N=32)[311/637 (48.8%) positive; titers 1:32 - 16384], Otomys irroratus [46/213 (21.6%) positive; titers 1:8 - 8192] and Otomys unisulcatus [33/164 (20.1%) positive; titers 1:8 - 2048]. In contrast, only 6/620 (1.0%) Mastomys coucha (2N=36) had antibody. 832 serum and organ specimens from 713 individuals of 4 species were tested in cell cultures and 17 arenavirus isolations were obtained: 10 from

M.natalensis, 5 from O.unisulcatus and 1 each from O.irroratus and A.namaquensis. Preliminary characterisation, including tests with monoclonal antibodies, suggest that the M.natalensis viruses correspond to Mopeia virus while the other isolations appear to comprise two new arenaviruses.

The medical significance of the arenaviruses in southern Africa has yet to be determined.

(R Swanepoel, A J Shepherd, P A Leman, T Whistler, D E Gill, L Searle and J C Abbott)

DENGUE TYPE 3 ON NIUE ISLAND:

Niue Island is situated some 1300 km north east of Auckland, New Zealand, and has a population of approximately 2,500 persons living in 13 separate villages. In January 1986, the Department of Health, Niue, became aware of cases resembling dengue fever occurring in Alofi, the main business centre for the island. The island had experienced major epidemics of dengue type 2 in 1972 (90% attack rate, 12 deaths) and dengue type 4 in 1980 (attack rate 33%, 4 deaths). Of 157 paired sera from suspected cases received from the Health Department, Niue, in this Unit, 137 had flavivirus HI antibody in one or both specimens. Twenty-four of these had relatively low antibody levels (1:10-1:40) and did not show diagnostic rises between acute and convalescent specimens. Of the remainder, 20 showed high antibody levels in both, but not diagnostic rises, while the remaining 93 all showed four-fold or greater increases in antibody level. Forty-nine of these had increases of equal to or greater than 64-fold and 19 had rises of 512-fold or greater when dengue was used as the HA antigen.

Acute phase sera were inoculated into cultures of *Aedes albopictus* cells (C6/36 clone) and three identical haemagglutinating agents were isolated from three different patients of 69 tested. These agents agglutinated goose erythrocytes at pH 6.0 and this agglutination was inhibited by dengue-specific rabbit antiserum. Positive identification of these viruses as dengue type 3 was accomplished by fluorescent antibody staining of infected C6/36 cells with type-specific dengue monoclonal antibodies (kindly supplied by the Centers for Disease Control, Atlanta) and by biotin-streptavidin enzyme-linked immunosorbent assay. Reisolation of virus from the original specimens was positive in each case.

Dr C.F. Tukuitonga, Acting Director of Health, Niue, reported that retrospective studies indicated that the epidemic began in November 1985 and that some 544 cases were reported (attack rate 21%) with two patients showing DSS. No deaths were recorded. Both sexes were equally affected. The epidemic ended in the first week of April, 1986.

(T Maguire)

REPORT FROM MICROBIOLOGY SECTION, CSIRO, DIVISION OF TROPICAL ANIMAL SCIENCE, LONG POCKET LABORATORIES, INDOORROOPILLY, BRISBANE, AUSTRALIA.

Isolation of Arboviruses from Cattle and Insects
at Sentinel sites in Queensland, Australia, 1979-1985.

Isolations from Cattle

Arboviral infections of cattle were monitored in and near 2 sentinel herds, 1 located in the north of the state at Kairi (145°33'E, 17°13'S) 60 km W of Cairns and 1 in the southeast at Peachester (152°53'E, 26°51'S) 70 km N of Brisbane. The sentinel herds consisted of 10-20 heifers introduced into the group during Jul-Aug, when 8-9 months old. Animals were bled monthly or more frequently until the following July-August when a fresh group was substituted. Isolations are shown in Tables 1 and 2.

TABLE 1. Isolation of Arboviruses from Sentinel Cattle - Kairi, Queensland, Australia, 1979-1982.

Summer	No. of Isolates	Virus	Period of Collection
1979-1980	31	Bunyip Creek	December-March
	4	CSIRO Village	January-February
	4	D'aguilar	January-March
	1	Peaton	January
1980-1981	2	Bunyip Creek	February-March
	5	CSIRO Village	February-March
	4	Akabane	January-February
	2	BEF	December, February
	3	BLU-1	March
	3	EHD-7	March
1981-1982	2	Bunyip Creek	February-March
	1	CSIRO Village	March
	3	D'aguilar	January-February
	1	Douglas	December
	3	EHD-5	January, March
	1	EHD-7	April

The BLU group viruses were only isolated through Aedes albopictus C6/36 cells (AA). The Simbu group viruses (Akabane, Peaton and Douglas), the Palyam group viruses (Bunyip Creek, CSIRO Village and D'aguilar) and the BEF group virus, Kimberley, were only isolated in BHK21 cells. The Simbu group virus, Aino was isolated in suckling mice, (first record from cattle). The EHD serotype 6 and serotype 7 viruses (EHD serotype numbers proposed by C.H. Campbell and T.D. St.George) and 11/13 EHD serotype 5 viruses were isolated through AA cells. The remaining two EHD-5 viruses were isolated in BHK21 cells. BEF virus was isolated in all 3 systems, 18 isolations through AA cells, 2 isolations in BHK21 cells and 1 isolation in suckling mice. Serology confirmed that cattle were infected with all viruses.

TABLE 2. Isolation of Arboviruses from Sentinel Cattle - Peachester, Queensland, Australia, 1979-1985.

Summer	No. of Isolates	Virus	Period of Collection
1979-1980	4	Bunyip Creek	February, April
	12	CSIRO Village	January-February
	1	D'aguilar	April
	1	Aino	December
	4	Akabane	December
	2	BEF	February
1980-1981	11	Bunyip Creek	December-April
	18	D'aguilar	November-January
	2	Akabane	December, March
	7	EHD-5	January-February
	4	EHD-6	February-April
1981-1982	2	Bunyip Creek	March
	119	CSIRO Village	December-February
	2	D'aguilar	March
	2	Akabane	November, January
	11	BEF	January
	1	Kimberley	January
	1	EHD-5	March
1982-1983	1	Bunyip Creek	April
	1	CSIRO Village	December
1983-1984	1	CSIRO Village	March
	3	BEF	January
1984-1985	17	Bunyip Creek	January
	8	Akabane	December
	2	EHD-5	April

Isolations from Biting Midges and Mosquitoes

During an outbreak of bovine ephemeral fever at Peachester, in Jan/Feb 1984, biting midges and mosquitoes were collected over a 3 week period by means of truck traps, light traps and cow baited collections. A total of 14338 midges of 4 species was processed in 156 pools, and 9030 mosquitoes of 27 species were processed in 232 pools. Each pool was processed in parallel in BHK21 tissue cultures, and by intrathoracic inoculation of *Ae. aegypti* mosquitoes which were then held alive for a period of 10 days at 20°C then assayed in BHKs. All pools inoculated into BHKs were passaged twice before being discarded as negative. Of the 388 pools, 90 were also processed in *Ae albopictus* tissue cultures then passaged twice in BHKs.

TABLE 3. Isolation of Arboviruses from Insects - Peachester, Queensland, Australia, Jan/Feb 1984.

Species	Virus Group	Virus	Isolation system			
			BHK only	BHK and <u>Ae aeg.</u>	<u>Ae aeg.</u> only	AA only
<u>C. brevitarsis</u> (11025 insects/ 109 pools)	Simbu	Akabane	5	5**	-	-
	Palyam	D'Aguiar	8	-	-	-
		Bunyip Ck	4	-	-	-
		CSIRO Village	3	1	-	-
		Orbi	EHD5	-	-	4
	Rhabdo	BEF	-	-	1	-
		Tibrogargan	-	-	3	-
	Undetermined		1	-	-	-
<u>C. wadai</u> (2811/38)	Simbu	Akabane	1	-	-	-
<u>Cx edwardsi</u> (1829/37)	Alpha	Sindbis	-	1*	-	-
	Corriparta		-	1*	-	-
	Maputta Gp		-	1*	-	1
	Rhabdo	Oakvale	1*	6**	2	-
	Undetermined		-	-	1*	-
<u>Cx bitaeniorhynchus</u> (362/9)	Alpha	Sindbis	-	1*	-	-
	Undetermined		-	-	-	1
<u>Cx orbostiensis</u> (3898/78)	Corriparta		-	-	-	1
	Undetermined		-	-	1	-
<u>Cx annulirostris</u> (928/24)	Undetermined		-	-	-	1
<u>Ae (Ver) No. 52</u> (920/23)	Barmah Forest		1*	-	-	-
	Undetermined		-	-	2*	-
<u>An bancroftii</u> (92/5)	Rhabdo	BEF	-	-	1	-

* Each asterisk denotes a parallel isolation in AA tissue cultures.

Table 3 shows the viruses which were isolated and the insect species from which they were isolated. The different isolation systems produced different virus groups from Culicoides brevitarsis. Palyam group viruses were isolated in BHKs but not in Ae aegypti. Orbiviruses (EHD) and rhabdoviruses (BEF, Tibrogargan) were isolated after passage through live Ae aegypti but not by direct inoculation into BHKs. The isolation of BEF virus is the first isolation of this virus from biting midges in Australia. The isolation of Akabane virus from C. wadai is the first isolation of a virus from that species.

The mosquito, Cx edwardsi produced 9 isolations of a new rhabdovirus tentatively called Oak Vale virus (BEF serogroup). No antibodies to this virus were

found in the cattle, nor was *Cx edwardsi* abundant in cow bait collections. BEF was isolated from *An bancroftii* which had been collected feeding on a febrile cow and held for 10 days before processing. This was the second isolation of BEF from this species.

On 2 occasions pools of *C. brevitarsis* yielded D'Aguiar virus in BHKs and Tibrogargan virus after processing through *Ae aegypti* mosquitoes. The pool of *C. brevitarsis* which produced BEF through *Ae aegypti* also produced D'Aguiar in BHKs.

(T.D. St. George, H.A. Standfast, M.J. Muller, H. Zakrzewski, D.H. Cybinski, D.S. Gibson)

Vector/Host Index for Australian Arboviruses

This table has been developed to update the virus/host lists in Arbovirus Information Exchange, June 1986.

Virus	Isolated from Species	Reference
Ross River	<i>Aedes vigilax</i>	44
"	<i>Gallina cyanoleuca</i>	37
"	<i>Microeca fascians</i>	37
"	<i>Anopheles amictus</i>	8
"	<i>Aedes normanensis</i>	8
"	<i>Mansonia uniformis</i>	36
"	Horse	47
"	<i>Coquilletidia linealis</i>	45
"	<i>Macropus agilis</i>	25
"	<i>Culex annulirostris</i>	25
"	<i>Aedes flavifrons</i>	56
"	Human	46
Sindbis	<i>Culex squamosus</i>	8
"	<i>Aedes vittiger</i>	8
"	<i>Aedes theobaldi</i>	8
"	<i>Culex pullus</i>	8
"	<i>Culex edwardsi</i>	57
"	<i>Culex australicus</i>	43
"	<i>Aedes normanensis</i>	24
"	<i>Aedes vigilax</i>	24
"	<i>Mansonia septempunctata</i>	24
"	<i>Culex annulirostris</i>	24
"	<i>Culex bitaeniorhynchus</i>	57
Getah	<i>Anopheles amictus</i>	24
"	<i>Culex bitaeniorhynchus</i>	24
Barmah Forest	<i>Culex annulirostris</i>	43
"	<i>Aedes procax</i>	28
"	<i>Aedes vigilax</i>	55
"	<i>Aedes (Verrallina) spp.</i>	57
"	Culicine mosquitoes	63
"	<i>Aedes normanensis</i>	8
Saumarez Reef	<i>Ornithodoros capensis</i>	33
"	<i>Ixodes eudyptidis</i>	33

Virus	Isolated from Species	Reference
Kokobera	<i>Culex annulirostris</i>	24
"	<i>Aedes vigilax</i>	25
"	<i>Aedes normanensis</i>	8
"	<i>Culex annulirostris</i>	24
"	Oriolus flavocinctus	37
"	<i>Culex squamosus</i>	38
"	<i>Aedes tremulus</i>	14
Kunjin	<i>Culex australicus</i>	39
"	<i>Culex quinquefasciatus</i>	8
Edge Hill	<i>Aedes normanensis</i>	8
"	<i>Anopheles meraukensis</i>	24
"	<i>Anopheles amictus</i>	8
"	<i>Culex annulirostris</i>	24
"	<i>Aedes vigilax</i>	24
Stratford	<i>Aedes vigilax</i>	24
Alfuy	<i>Culex pullus</i>	8
"	<i>Aedeomyia catasticta</i>	38
Gadgets Gully	<i>Ixodes eudyptidis</i>	51
Murray Valley Enceph.	<i>Culex annulirostris</i>	24
"	<i>Culex australicus</i>	43
"	<i>Culex bitaeniorhynchus</i>	24
"	Ardea novaehollandiae	43
"	<i>Culex quinquefasciatus</i>	14
"	<i>Aedes normanensis</i>	24
"	Human	52
"	Domestic fowl	53
Dengue	Human	64
Koongol	<i>Culex annulirostris</i>	24
"	<i>Anopheles bancroftii</i>	24
"	<i>Anopheles farauti</i>	24
Wongal	<i>Culex annulirostris</i>	24
"	<i>Coquilletidia crassipes</i>	35
Akabane	<i>Culicoides brevitarsis</i>	1
"	Cattle	26,57
"	<i>Culicoides wadai</i>	57
Aino	<i>Culicoides brevitarsis</i>	1
"	Cattle	2,57
Peaton	<i>Culicoides brevitarsis</i>	4
"	Cattle	4,57
Douglas	Cattle	5,57
"	<i>Culicoides brevitarsis</i>	6
Tinaroo	<i>Culicoides brevitarsis</i>	5
"	Cattle	6
(Facey's Paddock)	<i>Culex annulirostris</i>	8
"	<i>Aedes normanensis</i>	8
"	<i>Culicoides</i> spp.	9
"	<i>Culicoides austropalpalis</i>	28
Thimiri	<i>Culicoides histrio</i>	7

Virus	Isolated from Species	Reference
Kao Shaun	<i>Argas robertsi</i>	41
Taggart	<i>Ixodes uriae</i>	48
Precarious Point	<i>Ixodes uriae</i>	51
Trubanaman	<i>Culex annulirostris</i>	36
"	<i>Anopheles annulipes</i>	38
"	<i>Coquilletidia linealis</i>	63
Kowanyama	<i>Anopheles annulipes</i>	38
"	<i>Anopheles amictus</i>	38
"	<i>Culex annulirostris</i>	43
Belmont	<i>Culex annulirostris</i>	1
"	<i>Culicoides marksii</i>	9
"	<i>Culicoides bundyensis</i>	28
Mapputta	<i>Anopheles meraukensis</i>	24
"	<i>Anopheles amictus</i>	9
"	<i>Anopheles annulipes</i>	36
"	Mosquitoes (mixed pool)	34
Gan Gan	<i>Aedes normanensis</i>	8
"	<i>Aedes theobaldi</i>	8
"	<i>Culex annulirostris</i>	8
"	<i>Aedes vigilax</i>	39
"	<i>Culex</i> spp.	55
Upolu	<i>Ornithodoros capensis</i>	49
"	<i>Puffinus pacificus chlororhyncus</i>	40
Nugget	<i>Ixodes uriae</i>	48
(Mudjinbarry)	<i>Culicoides marksii</i>	32
Wallal	<i>Culicoides dycei</i>	34
"	<i>Culicoides marksii</i>	34
Warrego	<i>Anopheles meraukensis</i>	8
"	<i>Culicoides dycei</i>	34
"	<i>Culicoides marksii</i>	34
"	<i>Culex annulirostris</i>	9
"	<i>Culicoides actoni</i>	28
Mitchell River	<i>Culicoides</i> spp.	34
Eubanagee	<i>Anopheles farauti</i>	9
"	<i>Culex annulirostris</i>	9
"	Mosquitoes (mixed pool)	35
Tilligerry	<i>Anopheles annulipes</i>	39
Ibaraki (EHD 2)	Cattle	29,60
EHD 5, 6, 7, 8	Cattle	29,60,57
EHD 5, 6	<i>Culicoides brevitarsis</i>	29,60
BLU 20	<i>Culicoides</i> spp.	22
BLU 1	<i>Culicoides fulvus</i>	20
BLU 1	<i>Culicoides brevitarsis</i>	21

Virus	Isolated from Species	Reference
BLU 1,21, 23	Cattle	19,23,57
Corriparta	<i>Culex annulirostris</i>	24
"	Charadrius melanops	37
"	<i>Culex orbostiensis</i>	57
"	<i>Aedeomyia catasticta</i>	14
"	<i>Culex edwardsi</i>	57
CSIRO Village	<i>Culicoides</i> spp.	11
"	<i>Culicoides brevitarsis</i>	11
"	Cattle	11,57
Marrakai	<i>Culicoides</i> spp.	9
"	Cattle	27
"	Water buffalo	27
Bunyip Creek	<i>Culicoides brevitarsis</i>	11
"	<i>Culicoides oxystoma</i>	9
"	Cattle	11,57
D'Aguilar	<i>Culicoides brevitarsis</i>	1
"	Cattle	10,57
Paroo River	<i>Culex annulirostris</i>	43
Berrimah	Cattle	17
Adelaide River	Cattle	18
Bovine Ephemeral Fever	Cattle	12,57
"	<i>Anopheles bancroftii</i>	13
"	<i>Culicine</i> mosquitoes	13
"	<i>Culicoides brevitarsis</i>	59
Kimberley	<i>Culex annulirostris</i>	14
"	Cattle	15,57
"	<i>Culicoides brevitarsis</i>	16
(Oakvale)	<i>Culex</i> spp.	57
"	<i>Culex edwardsi</i>	57
Almpiwar	Ablepharus boutonii virgatus	42
Kununurra	<i>Aedeomyia catasticta</i>	14
"	<i>Culicoides austropalpalis</i>	28
Charleville	<i>Phlebotomus</i> spp.	34
"	Gehyra australis	34
Tibrogargan	<i>Culicoides brevitarsis</i>	61
(Coastal Plains)	Cattle	62
Parry's Creek	<i>Culex annulirostris</i>	14
Lake Clarendon	<i>Argas robertsi</i>	58
Wongorr	<i>Aedes lineatopennis</i>	34
"	<i>Culex annulirostris</i>	8
"	<i>Culicoides pallidothorax</i>	9
"	<i>Aedes normanensis</i>	14
Ngaingan	<i>Culicoides</i> (C) spp.	34,54
(Beatrice Hill)	<i>Culicoides peregrinus</i>	9
Yacaaba	<i>Aedes vigilax</i>	39
Picola	<i>Culex annulirostris</i>	43

Virus	Isolated from Species	Reference
(Leanyer)	Anopheles meraukensis	50
(Parkers Farm)	Culex annulirostris	8
"	Culicoides marksii	9
Umbre [Little Sussex]	Culex annulirostris	8
Termeil	Aedes camptorhynchus	39
"	Aedes vigilax	39
"	Culex annulirostris	39
(Humpty Doo)	Lasiohelea spp.	9
"	Culicoides marksii	9
Johnston Atoll	Ornithodoros capensis	49

(Unregistered viruses are in parenthesis.) **Bold** = Vertebrate

Listed in Vector Index in "Arthropod-Borne Virus Information Exchange"
June 1986. Reference unknown.

D'Aguilar virus	Culicoides oxystoma
Ross River "	Culex quinquefasciatus
Wallal "	Aedes lineatopennis
" "	Culicoides brevitarsis

REFERENCES

1. DOHERTY, et al., (1972). Aust Vet J **48**: 81.
2. CYBINSKI, et al., (1983). Aust Vet J **60**: 283.
3. ST.GEORGE, et al., (1978). Aust Vet J **54**: 558.
4. ST.GEORGE, et al., (1980). Aust J Biol Sci **33**: 235.
5. ST.GEORGE, et al., (1979). Aust J Exp Biol Med Sci **57**: 581.
6. CYBINSKI, (1984). Aust J Biol Sci **37**: 91.
7. STANDFAST, et al., (1982). J Med Ent **19**: 212.
8. DOHERTY, et al., (1979). Aust J Exp Biol Med Sci **57**: 509.
9. STANDFAST, et al., (1984). Aust J Biol Sci **37**: 351.
10. ST.GEORGE, et al., (1976). Aust Vet J **52**: 598.
11. CYBINSKI, et al., (1982). Aust J Biol Sci **35**: 343.
12. DOHERTY, et al., (1969). Aust J Sci **31**: 365.
13. STANDFAST, et al., (1976). Aust Vet J **52**: 242.
14. LEIHNE, et al., (1981). Aust J Exp Biol Med Sci **59**: 347.
15. CYBINSKI, et al., (1983). Vet Microbiol **8**: 221.
16. ZAKRZEWSKI, et al., (1984). Aust J Exp Biol Med Sci **62**: 779.
17. GARD, et al., (1983). Aust Vet J **60**: 89.
18. GARD, et al., (1984). Aust Vet J **61**: 333.
19. ST.GEORGE, et al., (1980). Aust Vet J **56**: 562.
20. STANDFAST, et al., (1979). Proc 2nd Sym Arbo Res Aust **0**: 20
21. ST.GEORGE, et al., (1984). Aust Vet J **61**: 95.
22. ST.GEORGE, et al., (1978). Aust Vet J **54**: 153.
23. GARD, et al., (1985) Aust Vet J **62**: 203.
24. DOHERTY, et al., (1963). Aust J Exp Biol Med Sci **41**: 17.

25. DOHERTY, et al., (1971). Trans R Soc Trop Med Hyg **65**: 504.
26. ST.GEORGE, et al., (1977). Aust Vet J **53**: 249.
27. GARD, (pers. comm.).
28. STANDFAST, (pers. comm.).
29. ST. GEORGE, et al., (1983). Aust Vet J **60**: 216.
30. CYBINSKI, et al., (1980). Vet Microbiol **5**: 301.
31. ST.GEORGE, et al., (1984). Aust J Biol Sci **37**: 85.
32. DOHERTY, et al., (1978). Aust J Biol Sci **31**: 97.
33. ST.GEORGE, et al., (1977). Aust J Exp Biol Med Sci **55**: 493.
34. DOHERTY, et al., (1973). Trans R Soc Trop Med Hyg **67**: 536.
35. DOHERTY, et al., (1968). Trans R Soc Trop Med Hyg **62**: 862.
36. WOODROOFE, (1976). Proc 1st Sym Arbo Res Aust **0**: 67.
37. WHITEHEAD, et al., (1968). Trans R Soc Trop Med Hyg **62**: 439.
38. DOHERTY, et al., (1968). Trans R Soc Trop Med Hyg **62**: 430.
39. MARSHALL, et al., (1980). Aust J Exp Biol Med Sci **58**: 91.
40. DOHERTY, (1977). Aust J Exp Biol Med Sci **55**: 103.
41. HOOGSTRAAL, et al., (1975). J Med Ent **11**: 513.
42. GRAF, et al., (1967). Rep Qld Inst Med Res **22**: 4.
43. MARSHALL, et al., (1982). Aust J Exp Biol Med Sci **60**: 457.
44. DOHERTY, et al., (1963). Aust J Sci **26**: 183.
45. GARD, et al., (1973). Am J Trop Med Hyg **22**: 551.
46. DOHERTY, et al., (1972). Med J Aust **0**: 1083.
47. PASCOE, et al., (1978). Aust Vet J **54**: 600.
48. DOHERTY, et al., (1975). Am J Trop Med Hyg **24**: 521.
49. DOHERTY, et al., (1969). Aust J Sci **31**: 363.
50. DOHERTY, et al., (1977). Aust J Exp Biol Med Sci **55**: 485.
51. ST.GEORGE, et al., (1985). Am J Trop Med Hyg **34**: 406.
52. FRENCH, (1952). Med J Aust **1**: 100.
53. CAMPBELL, et al., (1975). Aust Vet J **51**: 1.
54. KAY, et al., (1978). J Aust Ent Soc **17**: 145.
55. VALE, et al., (1986). Aust J Exp Biol Med Sci **64**: 307.
56. MC MANUS, et al., (1986). Proc 4th Sym Arbo Res Aust (in press).
57. ST.GEORGE, et al., (1987). Arbo Info Exchange (this issue).
58. ST.GEORGE, et al., (1984). Aust J Biol Sci **37**: 85.
59. MULLER, et al., (1986). Proc 4th Sym Arbo Res Aust (in press).
60. CAMPBELL, et al., (1980). Aust Vet J **63**: 233.
61. CYBINSKI, et al., (1980). Vet Microbiol **5**: 301.
62. CYBINSKI, et al., (1986). Aust J Biol Sci **39**: 225.
63. VALE et al., (1985). Commun Disease Intell Bull **85(7)**: 809.
64. KAY et al., (1984). Med J Aust **140**: 264.

S.S. DAVIS

CSIRO, Division of Animal Tropical Science,
Long Pocket Laboratories, Private Bag No. 3, Indooroopilly, Queensland 4

I. ARBOVIRUS SURVEILLANCE in French Guiana, Martinique and Guadeloupe during 1986.

A) VIRUS ISOLATION.

During this year 42 specimens, all from French Guiana, were inoculated on A.P.61 cells.

8 were found positive : 19%.

Most of the positive results occurred during the second quarter (April, May, June) i.e. long rainy season.

All the strains have been identified to dengue virus type 2 using fluorescent monoclonal antibodies tests.

It is of interest to notice that among the first symptoms of dengue disease, pulmonary symptoms were often reported, as it was already mentioned in dengue surveillance summary (D.J.GUBLER).

No case of D.H.F./D.D.S were recorded.

B) SEROLOGICAL STUDIES.

During this year, 1191 serum specimens were studied :

998 from French Guiana
135 from Martinique
68 from Guadeloupe.

The results are reported in Table I.

The serums were tested by C.F. and H.I. Tests.

These results include only the C.F. Tests.

The antigen battery of Alphavirus includes : Mucambo, Pixuna, Cabassou and Tonate.

The antigen battery of Flavivirus : Yellow Fever, S.L.E, DEN 2, DEN 3, Ilheus.

1) IN FRENCH GUIANA.

- Most of the significant results for the Flavivirus were also found during the second quarter.

Considering the two percentage of isolation and the serological results of presumptive recent infection, which are similar \approx 20%, even if several Flavivirus have been isolated in French Guiana in the past (Dengue, S.L.E, Ilheus), it seems that these results indicate a dengue 2 virus circulation only.

- For the Alphavirus, there is no evidence of recent human infection, but we can see that the main activity of these viruses occurs during the short rainy season (December, January).

2) IN MARTINIQUE AND GUADELOUPE.

The results seem to be similar, but the number of samples is not significant.

It is interesting to observe that antibodies for Alphavirus were detected in one case in Martinique.

I. VIROLOGICAL RESULTS OF AN OUTBREAK OF DHF/DDS LIKE SYNDROMS IN SURINAME IN SEPTEMBER 1984.

The results can be separated in two rubrics.

A) ONE CASE OF YELLOW FEVER.

The patient died of hepatonephritis with hemorrhagic symptoms.

From the blood sample, a yellow fever strain was isolated on A.P.61 cells incubated at 33°C.

This strain was confirmed to be a wild South American strain of Yellow Fever by the Institut Pasteur Dakar using S.D.S polyacrylamide gel electrophoresis and Fingerprinting (Vincent DEUBEL : personal communication).

This result was reconfirmed by N.KARABATSOS at C.D.C. Fort Collins using a neutralization Test.

B) TWO CASES OF MAYARO VIRUS FEVER.

Characterised by fever, headache, retro-orbital pain and myalgia.

The isolations were positive on suckling mice.

The identification was performed at C.D.C. Fort Collins.

After passage in vero cells, a high titered hemagglutinin (1/4000 at p.H.6,1) was obtained.

The antigen was inhibited by Mayaro hyper immune mouse ascitic fluid only.

During the isolation process in Cayenne a Laboratory contamination by accidental inoculation of the Mayaro containing material was observed.

After an incubation period of 5 days, the onset was brutal with fever ($38^{\circ}5$, $39^{\circ}C$) asthenia, headache, arthralgia, myalgia, retro-orbital pain, photophobia during 3 days.

Rash and fever disappear on day 3.

L. Niel, Y. Robin, Institut Pasteur).

MONTHS	J	F	M	A	M	J	J	A	S	O	N	D	TOTAL YEAR
FRENCH GUIANA													
N° OF SAMPLES	76	72	62	84	84	146	111	72	52	79	43	107	988
FLAVIVIRUS ≥64	17	15	13	27	19	37	24	18	4	13	8	7	202
<64	19	12	22	25	26	37	41	25	17	27	12	12	275
ALPHA VIRUS <64	13	3	3	2	4	4	2	1	1	6	6	11	56
MARTINIQUE													
N° OF SAMPLES	1	1	3	16	1	18	4	2	1	10	4	74	135
FLAVIVIRUS ≥64			1	5		3				2		1	12
<64	1	1	2	8	1	7	4	1	1	3	2	22	53
ALPHA VIRUS <64			0	0		0		0		0	0	1	1
GUADELOUPE													
N° OF SAMPLES	0	0	8	5	0	28	7	8	1	6	0	5	68
FLAVIVIRUS ≥64			2			15	1						18
<64			3	3		7	2	5		1		2	23
ALPHA VIRUS <64			0	0		0	0	0		0		0	0