

Web Table 1. Provision of preventive counseling to travelers attending Boston Area Travel Medicine Clinics before trip departure (March 2008 to July 2010)

Parameter	Clinic A N = 3,269 % (n/N)	Clinic B N = 2,512 % (n/N)	Clinic C N = 1,403 % (n/N)	Clinic D N = 6,668 % (n/N)	Clinic E N=1,588 % (n/N)	All subjects N=15,440 % (n/N)
Animal exposure – rabies prevention	NA ^a	92.7% (1,962/2,116)	55.9% (780/1,395)	96.8% (6,447/6,657)	78.1% (1,200/1,537)	88.8% (10,389/11,705)
Diarrhea prevention and management	99.9% (3,266/3,268)	99.0% (2,095/2,116)	NA	98.3% (6,545/6,657)	96.9% (1,491/1,539)	98.7% (13,397/13,580)
Malaria risk and prevention	97.6% (3,190/3,268)	92.6% (1,960/2,116)	NA	74.1% (4,934/6,656)	95.3% (1,467/1,539)	85.1% (11,551/13,579)
Prevention of sexually transmitted infections	NA	54.4% (1,151/2,116)	12.8% (179/1,396)	84.1% (5,595/6,655)	68.2% (1,048/1,537)	68.1% (7,973/11,704)
Medical evacuation insurance	NA	NA	NA	98.4% (6,550/6,657)	52.3% (789/1,510)	71.4% (7,339/8,167)
Vector-borne diseases and avoidance (excluding malaria)	NA	98.0% (2,073/2,116)	94.9% (1,325/1,396)	97.5% (6,492/6,656)	95.2% (1,417/1,488)	97.0% (11,307/11,656)

a. NA = not available