Center for Disease Control

Foodborne and Waterborne Disease Outbreaks

U.S. Department of Health, Education, and Welfare Public Health Service

Annual Summary 1976 Issued October 1977



PREFACE

This report summarizes information received from state and local health departments, the Food and Drug Administration, the U.S. Department of Agriculture, and other pertinent sources. The information is preliminary and is intended primarily for use by those with responsibility for disease control activities. Anyone desiring to quote this report should contact the Enteric Diseases Branch for confirmation and further interpretation.

Contributions to the report are most welcome. Please address them to:

Center for Disease Control ATTN: Bureau of Epidemiology Bacterial Diseases Division Enteric Diseases Branch Atlanta, Georgia 30333

SUGGESTED CITATION

Center for Disease Control: Foodborne and Waterborne Disease Outbreaks Annual Summary 1976. Issued October 1977

Center for Disease Control......William H. Foege, M.D., Director Bureau of Epidemiology...... Director Bacterial Diseases DivisionJohn V. Bennett, M.D., Director Eugene J. Gangarosa, M.D., Deputy Director Enteric Diseases Branch..... Roger A. Feldman, M.D., Chief Foodborne and Waterborne Disease Outbreaks Statistical Services Branch.....Stanley M. Martin, M.S., Chief Robert A. Pollard, M.A. Epidemiologic Investigations Laboratory Branch..... Chief Enteric Diseases Laboratory Section.....Joy G. Wells, M.S., Chief Editorial and Graphic Services.....Frances H. Porcher, M.A., Chief Charlotte C. Turner, Writer-Editor Bureau of Laboratories Director Bacteriology Division...... Director Enterobacteriology Branch.....V. R. Dowell, Jr., Ph.D., Chief Clinical Bacteriology Branch.....P. B. Smith, Ph.D., Chief Bacterial Immunology Branch.....John C. Feeley, Ph.D., Chief

TABLE OF CONTENTS

		Page
I.	INTRODUCTION	1
II.	FOODBORNE DISEASE OUTBREAKS	2
	A. Definition of Outbreak	2
	B. Source of Data '	2
	C. Interpretation of Data	3
	D. Analysis of Data	4
	E. Investigation of a Foodborne Outbreak, (Summary Form)	17
	F. Line Listing of Foodborne Disease Outbreaks, 1976	20
	G. Guidelines for Confirmation of Foodborne Outbreak	47
	H. Selected Foodborne Outbreak Articles, 1976, Taken from Morbidity and Mortality Weekly Report	52
III.	WATERBORNE DISEASE OUTBREAKS	59
	A. Definition of Outbreak	, 59
	B. Source of Data	59
	C. Interpretation of Data	59
	D. Analysis of Data	59
	E. Investigation of a Waterborne Outbreak, (Summary Form)	65
	F. Line Listing of Waterborne Disease Outbreaks, 1976	68
	G. Selected Waterborne Outbreak Articles, 1976, Taken from Morbidity and Mortality Weekly Report	71
IV.	OUTBREAKS ON CRUISE SHIPS AND AIRCRAFT	75
v.	REFERENCES	77
VI.	LISTING OF ARTICLES ON FOODBORNE AND WATERBORNE DISEASE OUTBREAKS, 1976, TAKEN FROM MORBIDITY AND MORTALITY WEEKLY REPORT	81

a state of the sta

. . •

J

ACKNOWLEDGEMENTS: We acknowledge Robert E. Black, M.D., former Foodborne and Waterborne Disease Outbreaks Surveillance Officer, for his assistance in collecting and evaluating foodborne and waterborne disease outbreak data.

We acknowledge the important role played by Richard Swanson, B.S., and Dan Brand, B.S., of the Field Investigations Branch, Food and Drug Administration, and John Prucha, D.V.M., L. W. Schnurrenberger, D.V.M., and James Lloyd, D.V.M., of the Animal and Plant Health Inspection Service, U.S. Department of Agriculture, in alerting the Enteric Diseases Branch to foodborne disease outbreaks which came to their attention.

We acknowledge the invaluable assistance of Gunther F. Craun, M.S., Water Quality Division, Health Effects Research Laboratory, Environmental Protection Agency, in collecting and reviewing waterborne disease outbreak data. We also acknowledge Kim A. Keele, A.B., Statistical Assistant, and Carol A. Shaw, Secretary, for their assistance in preparing this report.

•

.

. .

• •

٠

• •

. .

.

•

•

.

. .

.

I. INTRODUCTION

The reporting of foodborne and waterborne diseases in the United States began about 50 years ago when state and territorial health officers, concerned about the high morbidity and mortality caused by typhoid fever and infantile diarrhea, recommended that cases of enteric fever be investigated and reported. The purpose was to obtain information about the role of food, milk, and water in outbreaks of intestinal illness as the basis for sound public health action. Beginning in 1923, the United States Public Health Service published summaries of outbreaks of gastrointestinal illness attributed to milk. In 1938, it added summaries of outbreaks caused by all foods. These early surveillance efforts led to the enactment of important public health measures which had a profound influence in decreasing the incidence of enteric diseases, particularly those transmitted by milk and water.

From 1951 through 1960 the National Office of Vital Statistics reviewed reports of outbreaks of foodborne illness and published summaries of them annually in <u>Public</u> <u>Health Reports</u>. In 1961 the Center for Disease Control (CDC), then the Communicable Disease Center, assumed responsibility for publishing reports on foodborne illness. For the period 1961-66 CDC discontinued publication of annual reviews, but reported pertinent statistics and detailed individual investigations in the Morbidity and Mortality Weekly Report (MMWR).

In 1966 the present system of surveillance of foodborne and waterborne diseases began with the incorporation of all reports of enteric disease outbreaks attributed to microbial or chemical contamination of food or water into an annual summary. Since 1966 the quality of investigative reports has improved primarily as a result of more active participation by state and federal agencies in the investigation of foodborne and waterborne outbreaks. In this report data from foodborne and waterborne disease outbreaks reported to CDC in 1976 are summarized.

Foodborne and waterborne disease surveillance has traditionally served 3 objectives:

1. Disease Control: Early identification and removal of contaminated products from the commercial market, correction of faulty food preparation practices in food service establishments and in the home, and identification and appropriate treatment of human carriers of foodborne pathogens are the fundamental control measures resulting from surveillance of foodborne disease. Identification of contaminated water sources and adequate purification of these sources are the primary control measures in the surveillance of waterborne disease outbreaks. Rapid reporting and thorough investigation of outbreaks are important for prevention of subsequent outbreaks.

2. <u>Knowledge of Disease Causation</u>: The responsible pathogen has not been identified in 30% to 60% of foodborne disease outbreaks reported to CDC in each of the last 5 years. In many of these outbreaks pathogens known to cause foodborne illness may not have been identified because of late or incomplete laboratory investigation. In others the responsible pathogen may have escaped detection even when a thorough laboratory investigation was carried out because the pathogen is not yet appreciated as a cause of foodborne disease or because it cannot yet be identified by available laboratory techniques. These pathogens might be identified and suitable measures to control diseases caused by them might be instituted as a result of thorough clinical, epidemiologic, and laboratory investigations. Pathogens suspected of being, but not yet determined to be etiologic agents in foodborne disease include Group D <u>Streptococcus</u>, <u>Citrobacter</u>, <u>Enterobacter</u>, <u>Klebsiella</u>, <u>Pseudomonas</u>, and the presumably viral agents of acute infectious non-bacterial gastroenteritis. Other parahaemolyticus are known causes of foodborne illness, but the extent and importance of their role have not as yet been determined. The etiologic agent(s) responsible for the majority of waterborne outbreaks also awaits identification. In waterborne disease, as in foodborne disease, the roles of a variety of viral and bacterial agents, e.g., Y. enterocolitica and parasitic agents, e.g., Giardia lamblia, remain to be clarified.

3. Administrative Guidance: The collection of data from outbreak investigations permits assessment of trends in etiologic agents and food vehicles and focuses on common errors in food and water handling. By compiling the data in an annual summary, it is hoped that local and state health departments and others involved in the implementation of food and water protection programs will be kept informed of the factors involved in food and waterborne disease outbreaks. Comprehensive surveillance should result in a clearer appreciation of priorities in food and water protection, institution of better training programs, and more rational utilization of available resources.

II. FOODBORNE DISEASE OUTBREAKS

A. Definition of Outbreak

For the purpose of this report a foodborne disease outbreak is defined as an incident in which 1) 2 or more persons experience a similar illness, usually gastrointestinal, after ingestion of a common food, and 2) epidemiologic analysis implicates the food as the source of the illness. There are a few exceptions; 1 case of botulism or chemical poisoning constitutes an outbreak.

- In this report outbreaks have been divided into 2 categories:
- 1. <u>Laboratory confirmed</u>--Outbreaks in which laboratory evidence of a specific etiologic agent is obtained and specified criteria are met (see Section G).
- Undetermined etiology--Outbreaks in which epidemiologic evidence implicates

 a food source, but adequate laboratory confirmation is not obtained. These
 outbreaks are subdivided into 4 subgroups by incubation period of the illness es--less than 1 hour (probable chemical), 1 to 7 hours (probable <u>Staphylococcus</u>),
 8 to 14 hours (probable <u>Clostridium perfringens</u>), and greater than 14 hours
 (other infectious agents).

B. Source of Data

The general public and local, state, and federal agencies which have responsibility for public health and food protection participate in foodborne disease surveillance. Consumers, physicians, hospital personnel, and persons involved with food service or processing report complaints of illness to health departments or regulatory agencies. Local health department personnel (epidemiologists, sanitarians, public health nurses, etc.) carry out most epidemiologic investigations of these reports and make their findings available to state health departments. State agencies concerned with food safety frequently participate in the initial investigation of the outbreak and offer laboratory support. Occasionally, on special request, CDC participates in an investigation, particularly if the outbreak is large or involves products that move in interstate commerce. State or other officials eventually summarize the findings of the investigation on the standard CDC reporting form (see Section E) and send to CDC.

The 2 federal regulatory agencies which have major responsibilities for food protection, the Food and Drug Administration (FDA) and Department of Agriculture (USDA) report episodes of foodborne illness to CDC and to state and local health authorities. CDC and state and local health authorities, in turn, report to FDA or USDA any foodborne disease outbreaks which might involve commercial products. The U.S. Armed Forces also report outbreaks directly to CDC.

By special arrangement pharmaceutical companies immediately report all requests for botulinal antitoxin to CDC. This is sometimes, the first communication of a botulism outbreak to public health authorities, although physicians are urged to promptly report all suspect botulism cases. In botulism outbreaks CDC works closely with physicians, state and local health authorities, and FDA or USDA representatives to provide diagnostic and therapeutic consultation and to rapidly identify the responsible food or foods.

For 1976 other sources of foodborne disease data were the Morbidity and Mortality Weekly Report, the Salmonella Surveillance Activity, and the Trichinosis Surveillance Activity.

C. Interpretation of Data

The limitations on the quantity and quality of data in this report must be appreciated in order to avoid misinterpretation. The number of outbreaks of foodborne disease reported by this surveillance system clearly represents a minute fraction of the total number that occur. The likelihood of an outbreak coming to the attention of health authorities varies considerably from one locale to another depending largely upon consumer awareness and physician interest.

Interstate outbreaks, large intrastate outbreaks, and outbreaks of serious illness such as botulism or mushroom poisoning with species containing amanita toxin are more likely to come to the attention of health authorities, including CDC. The quality of the investigation conducted by state or local health department varies considerably according to the department's interest in foodborne disease outbreaks and its investigative and laboratory capabilities. The likelihood that the findings of the investigation will be reported depends upon a state's commitment to foodborne disease surveillance.

Just as this report should not be the basis of firm conclusions about the absolute incidence of foodborne disease, it should not be used to draw conclusions about the relative incidence of foodborne disease of various etiologies (Table 2). For example, foodborne diseases characterized by short incubation periods such as most outbreaks of chemical etiology or outbreaks caused by Staphylococcus are more likely to be recognized as common-source foodborne disease outbreaks than those diseases with longer incubation periods. The common source aspect of a foodborne outbreak of hepatitis A which typically has an incubation period of several weeks would be particularly likely to escape detection. Outbreaks of serious disease such as botulism or mushroom poisoning with species of mushrooms containing amanita toxin are probably more likely to be reported than less serious illnesses but, because of their rarity, they may be less likely to be recognized and diagnosed. Outbreaks of C. perfringens are recognized readily but confirmed with difficulty because of problems involved in the transport and culturing of anaerobic specimens. Outbreaks of B. cereus, E. coli, V. parahaemolyticus, and Y. enterocolitica are probably less likely to be confirmed because these organisms are less often considered clinically, epidemiologically, and in the laboratory.

The number of reported outbreaks of some etiologies may depend upon the interest of a particular health department or individual. For example, the great increase in the number of reported outbreaks of ciguatera in 1974 probably reflected greater interest in the surveillance of this disease in the states in which they occurred. If a microbiologist becomes interested in looking for <u>C</u>. perfringens, he is likely to confirm more outbreaks of this etiology.

While the relative proportions of reported outbreaks attributed to most etiologies fluctuate minimally from year to year, it is worth noting that a few outbreaks involving very large numbers of persons may vastly alter the relative proportions of cases attributed to various etiologies (Tables 2 and 3).

Information on the number of deaths associated with outbreaks was unreported in 30% of the outbreaks. In many of the others complete information was lacking. Particularly when death is not immediate, foodborne disease may not be appreciated as contributing to the demise of an elderly or debilitated person unable to withstand otherwise minor physical stresses. These limitations on the data must be understood in interpreting Table 4.

In outbreaks of unknown etiology, the accuracy of reported information is always suspect. In these outbreaks when the epidemiology incriminating a particular food item was very weak, the food was listed as unknown in this report (Table 6). Information on the place of acquisition in these outbreaks was judged reliable and recorded (Table 7). However, information on the place where food was mishandled in these outbreaks was generally judged unreliable; in many of them the place of mishandling was listed as unknown (Table 8). Only in outbreaks in which a specific etiology was highly suspected, although unconfirmed in the laboratory, and in which the information on mishandling was consistent with the suspected etiology was a known place of mishandling designated.

The implications of a food-processing establishment mishandling food are great both to the public health and the establishment concerned. Consequently the outbreaks attributed to mishandling at these establishments are thoroughly investigated and reported data carefully scrutinized. For these reasons data obtained in these investigations is considered highly reliable (Tables 8 and 9).

Much is known about contributing factors in foodborne disease. Thus in most outbreaks of botulism and trichinosis, the food is usually inadequately cooked. In most of the outbreaks of bacterial etiology other than botulism and in outbreaks of scombroid (in which bacterial growth is responsible for toxin production), the food is usually stored at improper holding temperatures. In outbreaks of ciguatera, puffer fish poisoning, mushroom poisoning, and paralytic and neurotoxic shellfish poisoning, the food is obtained from an unsafe source, almost by definition. The investigators of foodborne disease outbreaks are usually aware of these contributing factors and consequently seek and find the appropriate factors. Sometimes, however, investigators report factors which are not known to be contributing to outbreaks of the type of etiology confirmed. In such cases the factors are considered in light of the evidence presented; if they are totally unsubstantiated, they are rejected. These considerations must be borne in the mind in interpreting Table 10.

D. Analysis of Data

In 1976 there were 438 outbreaks of foodborne disease involving 12,463 cases, a decrease of 12% in the number of outbreaks reported to the CDC Foodborne Disease Surveillance Activity (Figure 1). An etiology was confirmed in 30% (132) of the outbreaks--similar to the percentage of confirmed outbreaks in 1975 (38%) and in 1974 (44%).

Of the 438 outbreaks, state, local or territorial health departments reported 408 (93%). Outbreaks were reported from 43 states, New York City, District of Columbia, Puerto Rico, Guam, and the Virgin Islands (Figure 2 and Table 1). No outbreaks were reported from 7 states or the Canal Zone. Three outbreaks involved more than 1 state. The 5 state health departments reporting the most outbreaks were Washington (48); Pennsylvania (41), California (26); New York (19); and Hawaii (17). The large number of outbreaks reported from these states undoubtedly reflects the interest of the respective state health departments in foodborne disease surveillance. The 119 outbreaks reported from New York City in 1976 and the 120 reported in 1975 represents a 60-fold increase from 1974, probably reflecting increased reporting.

Of the 132 outbreaks with confirmed etiology, the etiology was bacterial in 92 (70%), chemical in 28 (21%), parasitic in 9 (7%), and viral in 3 (2%) (Table 2). While outbreaks with known bacterial etiology accounted for only 70% of the outbreaks, they accounted for 91% of the cases. The majority of cases of bacterial etiology were caused by <u>Salmonella</u> (33%) and <u>Staphylococcus</u> (26%). The 23 outbreaks and 40 cases of botulism were both the most reported since 1935. The first foodborne outbreak of <u>Yersinia enterocolitica</u> (286 cases) and picornavirus (enterovirus) Echo, type 4 (80 cases) were documented in 1976.

No outbreaks (2 or more cases) of foodborne brucellosis were reported in 1976. However, 24 single cases of brucellosis were attributed to the ingestion of unpasteurized dairy products. Six cases were traced to milk produced in the United States, and 18 were attributed to foreign dairy products. The foreign dairy products included cows' and goats' milk and cheese.

In 1976 there were 10 deaths associated with foodborne outbreaks (Table 4). Most deaths (5) were due to eating food containing the toxin of <u>C</u>. <u>botulinum</u>, a case fatality ratio of 12.5% (5/40). The other 5 deaths (<u>Salmonella 3</u>, <u>Shigella 1</u>, and unknown 1) associated with foodborne outbreaks occurred in elderly persons. Four of the 5 were residents of nursing homes.

Table 5 lists the outbreaks of undetermined etiology by median incubation periods. If one assumes that most outbreaks in which the median incubation period was less than 1 hour were of chemical etiology, that those in which median incubation period was 1-7 hours were of staphylococcal etiology, and that those in which the median incubation period was 8-14 hours were caused by <u>C</u>. perfringens then these agents were responsible for substantially more outbreaks than suggested in Table 2.

The vehicles of transmission were identified in 256 (58%) of the outbreaks (Table 6); multiple vehicles were involved in 30 (6.8%). Of the 226 outbreaks in which a single vehicle was identified, meats or poultry were incriminated in 68 (30%), fish or shellfish in 26 (12%), dairy products in 10 (4%), fruits and vegetables in 6 (3%), salads including chicken, turkey, potato, and egg in 10 (4%), oriental food 19 (8%), mushrooms 1 (1%), and other foods in 64 (28%). Of the meat vehicles beef and ham were most frequently incriminated.

Outbreaks of <u>C</u>. <u>botulinum</u> frequently involved home preserved vegetables and fish (Alaska). <u>C</u>. <u>perfringens</u> outbreaks usually involved beef, and <u>Staphylococcus</u> outbreaks most often involved meat. <u>Salmonella</u> outbreaks were caused by many different vehicles, including meat, such as precooked roast beef, poultry, dairy products, and salads. The outbreaks of heavy metal poisoning all involved nondairy beverages. Ciguatera outbreaks involved mainly coral reef fish (grouper). All the outbreaks of paralytic shellfish poisoning were associated with the consumption of clams. <u>T</u>. <u>spiralis</u> outbreaks involved pork or sausage.

In three-fourths of the outbreaks, the food was eaten at home (24%), in a restaurant (41%), or in a school (5%) (Table 7). Of the 23 outbreaks of botulism, the food was eaten at home in 12 (52%), in a restaurant 1 (4%), and was unknown in 10 (43%). Chemical outbreaks occurred frequently in the home and in food service establishments. Outbreaks caused by parasites usually occurred at home, but hepatitis outbreaks occurred at food service establishments.

The place where the mishandling of the food responsible for an outbreak occurred was specified in 376 outbreaks (Table 8). Of these, food service establishments were specified as responsible for the mishandling of food in 78%, homes in 18%, and food processing establishments in 4%. Food service establishments are locations where food is prepared for public consumption, i.e., restaurants, cafeterias, caterers, hospitals, industrial plants, etc. Food processing establishments are locations where a food is prepared for market. The distribution of places held responsible for mishandling of food in 1976 paralleled that of the 2 previous years. As in 1975 and 1974, where a place of food mishandling was specified, the majority of outbreaks caused by C. perfringens, Salmonella, and Staphylococcus were attributed to mishandling of food in the food service establishments. In ciguatoxin fish poisoning, since there is no practical way to distinguish fish containing ciguatoxin from fish without toxin, and the presence of the toxin is not influenced substantially by the way the fish is handled or cooked, a place of food mishandling was not specified in outbreaks of ciguatera poisoning. In most reported outbreaks of trichinosis, the food handling error occurred in the home while in reported outbreaks of hepatitis, it occurred away from home.

Of the 15 outbreaks attributed to mishandling of food in food processing establishments, 8 were due to bacteria, 5 to chemicals, and 2 were unknown (Table 9).

In 242 (55%) of the 438 outbreaks, including 88 (66%) of the 132 confirmed outbreaks, a contributing factor was reported and accepted in processing data (Table 10). The data reflected patterns of disease causation seen in previous years. In reported outbreaks of botulism and trichinosis, the most frequent error was inadequate cooking of the food. Improper holding temperatures most frequently contributed to reported outbreaks of <u>C</u>. <u>perfringens</u>, <u>Salmonella</u>, and <u>Staphylococcus</u> intoxication. Storage of beverages in metal containers or in contact with tubing of a type which allowed metallic ions to dissolve in the beverage was the most important contributing factor in the outbreaks of heavy metal poisonings. In outbreaks of ciguatera, paralytic shellfish poisoning and mushroom poisoning, the food was unsafe to begin with. In the outbreaks of chemical poisoning caused by miscellaneous chemicals, the food was obtained from an unsafe source. In the 2 outbreaks of hepatitis a person suspected of having active hepatitis was involved in foodhandling.

The date of onset of an outbreak was designated as the date of onset of the first case (Table 11). Generally, outbreaks were distributed more or less equally throughout the year. Outbreaks caused by <u>Salmonella</u> and <u>Staphylococcus</u> tended to occur more frequently in the summer months probably because the warm temperatures allow bacteria to grow in unrefrigerated foods. Outbreaks of paralytic shellfish poisoning tended to occur in the spring and summer months when the growth of toxic dinoflagellates is most abundant.









ľ

K

Foodborne Dis	ease Outbreaks,	by	Location,	1974-1976
---------------	-----------------	----	-----------	-----------

State	<u>1974</u>	<u>1975</u>	<u>1976</u>	State	<u>1974</u>	<u>1975</u>	<u>1976</u>
Alabama	4	1	1	Missouri	5	8	2
Alaska	5	4	11	Montana	0	3	2
Arizona	5	2	2	Nebraska	5	3	4
Arkansas	4	2	0	Nevada	1	4	3
California	32	41	26	New Hampshire	6	2	1
Colorado	6	1	8	New Jersey	10	12	8
Connecticut	4	9	9	New Mexico	0	1	1
Delaware	0	1	1	New York City	2	120	119
District of Columbia	2	0	2	New York State	22	8	19
Florida	15	30	4	North Carolina	4	0	1
Georgia	11	17	7	North Dakota	0	0	2
Hawaii	27	15	17	Ohio	20	0	9
Idaho	3	0	5	Oklahoma	3	3	3
Illinois	15	12	5	Oregon	8	7	5
Indiana	3	4	0	Pennsylvania	86	21	41
Iowa	4	1	1	Puerto Rico	1	0	1
Kansas	1	0	1	Rhode Island	2	2	2
Kentucky	1	8	2	South Carolina	7	9	2
Louisiana	5	15	3	South Dakota	5	1	0
Maine	0	0	2	Tennessee	6	17	8
Maryland	3	2	0	Texas	5	3	5
Massachusetts	1	8	10	Utah	7	3	1
Michigan	7	5	5	Vermont	2	0	0
Minnesota	14	25	7	Virginia	3	4	5
Mississippi	2	1	2	Washington	49	44	48
Other				West Virginia	6	0	0
Virgin Islands	0	0	1	Wisconsin	8	13	9
Guam and Trust				Wyoming	0	1	0
Territories	4	2	2	Multiple	5	2	3*,**,***
Canal Zone	0	0	0				

*Oklahoma, Texas, Utah **Connecticut, Delaware, Massachusetts, New Jersey, Pennsylvania ***California, Colorado

1974	total	456
1975	total	497
1976	total	438

.

	Outb	reaks	Cases				
BACTERIAL	#	%	#	%			
B. <u>cereus</u> <u>C. botulinum</u> <u>C. perfringens</u> <u>Salmonella</u> <u>Shigella</u> <u>Staphylococcus</u> <u>Y. enterocolítica</u> <u>f</u> Total	$\begin{array}{c} 2\\ 23\\ 6\\ 28\\ 6\\ 26\\ \underline{1}\\ 92\end{array}$	1.5 17.4 4.5 21.2 4.5 19.7 <u>0.8</u> 69.6	63 40 509 1169 273 930 <u>286</u> 3,270	1.8 1.1 14.2 32.7 7.6 26.0 <u>8.0</u> 91.4			
CHEMICAL							
Heavy metal Ciguatoxin Scombrotoxin Paralytic shellfish poison Monosodium glutamate Mushroom poison Other chemicals Total	6 2 4 2 1 7 28	4.54.51.53.01.50.85.321.1	55 19 5 11 7 1 <u>59</u> 157	$ \begin{array}{r} 1.5 \\ 0.5 \\ 0.1 \\ 0.3 \\ 0.2 \\ 0.0 \\ \underline{1.6} \\ 4.2 \\ \end{array} $			
PARASITIC							
<u>T. spiralis</u> <u>E. histolytica</u> Total	8 _ <u>1</u> _ 9	6.1 <u>0.8</u> 6.9	27 <u>9</u> 36	$0.8 \\ 0.3 \\ 1.1$			
VIRAL							
Hepatitis A Echo, type 4 Total	2 1 3	$\frac{1.5}{0.8}$ 2.3	37 <u>80</u> 117	1.0 2.2 3.2			
CONFIRMED TOTAL	132	99.9	3,580	99.9			

Confirmed Foodborne Disease Outbreaks and Cases by Etiology, 1976

	No. of	Outbreaks (No. of	Cases)
	1974	1975	1976
BACTERIAL			
A. hinshawii	0(0)	1(15) [·]	0(0)
B. cereus	1(11)	3(45)	2(63)
C. botulinum	21(32)	14(19)	23(40)
C perfringens	15(863)	16(419)	6(509)
Salmonella	35(5 499)	38(1 573)	28(1,169)
Shicollo	3(212)	3(413)	6(273)
Shigella	J(212)		26(020)
Staphylococcus	43(1,303)	45(2,275)	20(930)
Group A <u>Streptococcus</u>	1(325)	0(0)	0(0)
<u>V. cholerae</u>	1(6)	0(0)	0(0)
V. parahaemolyticus	0(0)	2(222)	0(0)
Suspect Group D	2(38)	1(50)	0(0)
Streptococcus			
Y. enterocolitica	0(0)	0(0)	1(286)
Total	122(8,551)	123(5(031)	92(3,270)
Iotai	122(0,001)	123(3,031)	52(5,270)
CHEMICAL			
	((00))	((50)	((55)
Heavy metals	4(28)	4(50)	6(55)
Ciguatoxin	26(148)	19(70)	6(19)
Puffer fish tetrodotoxin	1(2)	0(0)	0(0)
Scombrotoxin	10(26)	6(16)	2(5)
Monosodium glutamate	2(4)	, 3(9)	2(7)
Mushroom poison	6(9)	5(5)	1(1)
Paralytic shellfish noison	1(4)	0(0)	4(11)
Nourotovia shallfish poison	1(1)	0(0)	0(0)
Missellerseus sheriesle	4(10)	6(38)	7(59)
Miscellaneous chemicals	$\frac{10(13)}{57(2/1)}$	(30)	- 7 (35)
Total	57(241)	43(188)	20(157)
_ _			
PARASITIC			
T. spiralis	14(58)	20(193)	8(27)
T. gondii	1(4)	0(0)	0(0)
Anisakidae	1(1)	1(1)	0(0)
D. latum	0(0)	1(1)	0(0)
E biatolution	0(0)		1(9)
E. <u>HISCOLYLICA</u>	$\frac{0(0)}{16(62)}$	22(105)	$\frac{1(3)}{0(36)}$
lotal	T0(02)	22(193)	9(30)
VIRAL			
Hepatitis A	6(282)	3(173)	2(37)
Echo, type 4	0(0)	0(0)	1(80)
Total	6(282)	3(173)	3(117)
10 La1		- ·	
CONFIRMED TOTAL	201(9.137)	191(5,587)	132(3.580)
CONT TIRTIN TOTIM		~~~~~ , ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	,,,

Confirmed Foodborne Disease Outbreaks and Cases, 1974-1976

Deaths Associated with Foodborne Outbreaks, 1974-1976

	<u>1974</u>	<u>1975</u>	<u>1976</u>
C. botulinum	7	Ź	5
C. perfringens	1	1	0
Salmonella	1	2	3
Shigella	0	0	1
Staphylococcus	0	0	0
V. cholerae	1	0	0
T. spiralis	0	1	0
Hepatitis A	1	0	0
Mushroom poison	0	2	0
Organic chemicals	2	0	0
Unknown	_1	. 2	_1
Total	14	10	. 10

Table 5

Foodborne Disease Outbreaks of Unknown Etiology, by Incubation Period, 1976

Incubation Period	Number of Outbreaks	Percent of Total Outbreaks
<1 hour	24	5.5
1-7 hours	132	30.1
8-14 hours	73	16.7
>15 hours	60	13.7
Unknown		3.8
Total	306	69.8

BACTERIAL	Beef	Lamb	Ham	Pork	Sausage	Chicken	Turkey	Other Meat	Shellfish	Mahi-Mahi	Other Fish	MIIK	Cheese	Ice Cream	Other Dairy Products	Baked Foods	Fruits and Vegetables	Potato Salad	Poultry, Fish, Egg Salad	Other Salads	Mushrooms	Chinese Food	Mexican Food	Non-Dairy Beverages	Multiple Vehicles	Other Foods	Unknown	Total
 B. cereus C. botulinum C. perfringens Salmonella Shigella Staphylococcus Y. enterocolitica 	- 2 4 - 2				- - - 1	1 1 - 2	- 1 1 - 1	- - 1 2			- 2 - - - - - -					- - - 2		- - 1 - 1 -	- - - - -				- - 1 - -		- 1 5 - 3 -	1 10 2 4 4 5 -	- 10 - 9 1 - -	2 23 6 28 6 26 1
CHEMICAL						•	-		÷	-																		
Heavy metal Ciguatoxin Scombrotoxin Paralytic shellfish poison Monosodium glutamate Mushroom poison Other chemicals	- - - 1				· · · · · ·				- - 4 -		- 6 1 - -					- - - 1	- - - 1		-		- - - - 1,			5		1 - - 2 - 2		6 2 4 2 1 7
PARASITIC						•									•		•		- •	-								
<u>T. spiralis</u> <u>E. histolytica</u>	-	-	-	3 -	3 -	1	-	- -	- -	- - -	-	-	-		-	- -	<u>-</u>	-	- -	ī		-	-	-	1 -	1 -	-	8 1
VIRAL																												
Hepatitis A Echo, type 4	-		-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	-		-	-	-	-	-	-1	2 -	2 1
CONFIRMED TOTAL	9	-	3	3	4	4	3	4	4	1	9	1	2	1	-	3	3	2	-	4	1	-	1	5	10	33	22	132
UNKNOWN	19	-	5	-	1	1	10	2	4	-	8	2	1	1	2	9	3	2	-	2	-	19	4	-	20	31	160	306
TOTAL	28	-	8	3	5	5	13	6	8	1	17	3	3	2	2	12	6	4	-	6	1	19	5	5	30	64	182	438

Foodborne Disease Outbreaks, by Vehicle of Transmission, and Specific Etiology, 1976

-

; in

Table 6

٠

BACTERIAL	Home	Restaurant	School	Picnic	Church	Camp	Other or Unknown	Total
 <u>B. cereus</u> <u>botulinum</u> <u>perfringens</u> <u>Salmonella</u> <u>Shigella</u> <u>Staphylococcus</u> <u>Y. enterocolítica</u> 	12 2 3 $-$ 9 $-$ 1 26	1 1 7 1 5 <u>-</u> 16	- 1 2 - 2 1 6	- - 1 1 2 - 4	- - 1 - - - - 1	- - 1 - - - 1	$ \begin{array}{r} 1 \\ 10 \\ 2 \\ 13 \\ 4 \\ 8 \\ - \\ \overline{38} \end{array} $	2 23 6 28 6 26 <u>1</u> 92
CHEMI CAL								
Heavy metal Ciguatoxin Scombrotoxin Paralytic shellfish poise Monosodium glutamate Mushroom poison Other chemicals Total	$ \begin{array}{c} 1\\ 3\\ 2\\ 5\\ -\\ 1\\ 5\\ 1\\ 1 \end{array} $	3 2 - 2 - 1 8			1 - - - - 1		1 - 4 - 1 7	6 2 4 2 1 7 28
PARASITIC								
<u>T. spiralis</u> <u>E. histolytica</u> Total	6 1 6				- - -		$\frac{2}{1}{3}$	8 _1 _9
VIRAL								
Hepatitis A Echo, type 4 Tota:	1 1			 			1 1	2 1 3
CONFIRMED TOTAL	45	24	6	5	2	1	49	132
UNKNOWN	62	156	18	6	4	1	59	306
Total 1976 Total 1975 Total 1974	107 137 187	180 196 128	24 29 23	11 12 16	6 16 18	2 5 6	108 102 78	438 497 456

Foodborne Disease Outbreaks, by Place Where Food Was Eaten, and Specific Etiology, 1976

Food Food Processing Service Unknown-Not Establishments Establishments Homes Unspecified Applicable Total BACTERIAL <u>B. cereus</u> <u>C. botulinum</u> 2 2 --12 11 23 _ 6 C. perfringens --.__ _ 6 3 18 4 3 Salmonella 28 _ Shigella 5 0 -1 _ 6 4 13 Staphylococcus 7 2 26 Y. enterocolitica 1 1 8(8.7) 44(47.8) 24(26.1) 16(17.4) 92 Total (%) CHEMICAL . Heavy metal 3 2 1 6 5 Ciguatoxin _ _ 1 6 Scombrotoxin 2 ----2 ----3 Paralytic 1 4 shellfish poison Monosodium 2 2 glutamate Mushroom poison 1 1 Other chemicals 7 5 5(17.9) Total (%) 5(17.9) 5(17.9) 5(17.9) 8(28.6) 28 PARASITIC 5 3 T. spiralis E. histolytica <u>1</u> 1(11.1) Total (%) 5(55.6) 3(33.3) VIRAL Hepatitis A 2 2 Echo, type 4 1 3(100.0) Total (%) CONFIRMED TOTAL (%) 13(9.8) 53(40.1) 34(25.8) 24(18.2) 8(6.1) 132 ÛNKNOWN 2 241 33 29 306 1 Total 1976 15 294 67 53 9 438 Total 1975 13 201 61 222 0 497 Total 1974 16 90 77 273 0 456

Foodborne Disease Outbreaks, by Place Where Food Was Mishandled, and Specific Etiology, 1976

.

Foodborne Disease Outbreaks Caused by Mishandling of Food in Food-Processing Establishments 1976

	Etic	1003	7		Vehicle	Number of
		106)			Venicie	Lases
Salmon	<u>ella h</u>	neide	elberg		Cheese	339
<u>S.</u> bov:	is-mor	bifi	icans		Precooked roast beef	21
S. infa	antis				Nutrient supplement	4
Staphy.	lococc	us E	Interotoxín	D	Greek spaghetti	20
Staphy.	lococc	us			Beef ravioli	4
Staphy:	lococc	us F	Enterotoxin	D	Custard-filled donuts	2
Staphy.	lococc	us			Pepperoni, sausage	3
Y. ent	erocol	itic	a		Chocolate milk	286
Sodium	nitra	te			Table salt	2
Propy1	parab	en			Cake icing	9
Histam	ine				Cheese	38
Niacin					Hamburger	2
Niacin					Cubed steak	3
Unknow	n				Tuna extender	508
Unknow	n				Milk	42
Total	1976	15	outbreaks	1,283	cases	
	1975	13	outbreaks	123	cases	
	1974	16	outbreaks	1,704	cases	

•

.

Foodborne Disease Outbreaks, by Contributing Factors, and Etiology, 1976

Etiology	Number of Reported Outbreaks	Number of Outbreaks In Which Factors Reported	Improper Holding Tempera- tures	Inade- quate <u>Cooking</u>	Contami- nated Equip- ment	Food From Unsafe Source	Poor Per- sonal Hygiene	<u>Other</u>
BACTERIAL								
<u>B. cereus</u> <u>C. botulinum</u> <u>C. perfringens</u> <u>Salmonella</u> <u>Shigella</u> <u>Staphylococcus</u> <u>Y. entero-</u> <u>colitica</u> Total	2 23 6 28 6 26 <u>1</u> 92	2 13 6 16 3 22 	2 3 5 10 1 20 	11 2 4 2 - - 19	- 1 8 - 3 - 12	- - 3 - 1 - 4	1 8 2 7 18	- 1 4 2 - 7
CHEMICAL								
Heavy metal Ciguatoxin Scombrotoxin Paralytic shellfish	6 6 2 4	6 2 1 3	1 - -	 	3 - -	- 1 3	- - - -	2 1 1 -
poison Monosodium glu-	2	ŋ	-	-	-	-	-	-
Mushroom poison Other chemicals Total	1 7 28	1 <u>5</u> 18	- - 1		- - - 3	$\frac{1}{2}$	- - -	
PARASITIC								
<u>T. spiralis</u> <u>E. bistolytica</u> Total	8 _ <u>1</u> _9	7 7	- 	7 				
VIRAL								
Hepatitis A Echo, type 4 Total	$\frac{2}{\frac{1}{3}}$	1 1		- 			1 1	·
CONFIRMED TOTAL	132	88	42	26	15	11	19	14
UNKNOWN	306	154	118	17	39	6	34	30
Total 1976 Total 1975 Total 1974	438 497 456	242 277 219	160 214 131	43 87 45	54 62 31	17 14 50	53 93 41	44 14 9

.

Foodborne Disease Outbreaks, by Month of Occurrence, and Specific Etiology, 1976

		Jan	Feb	Mar	Apr	May	Jun	<u>Ju1</u>	Aug	Sep	<u>Oct</u>	Nov	Dec	<u>Total</u>
BACTERIAL														
B. <u>cereus</u> C. <u>botulinum</u> C. <u>perfringens</u> <u>Salmonella</u> <u>Shigella</u> <u>Staphylococcus</u> <u>Y. enterocolitica</u> To	otal	- 2 1 - 1 - 5	- - - 2. - 3	- 4 1 - 1 - 6	- 2 - 1 3 	1 - - 3 - 3 - 7	$ \frac{1}{-} \\ \frac{-}{6} \\ \frac{1}{2} \\ \frac{-}{10} $	- 2 1 5 1 4 - 13	- 1 3 1 4 - 9	- 6 - 5 - 5 1 17	- 1 - 2 - - 3	- 3 - 2 1 1 - 7	- 2 - 1 1 † - - - 4	2 23 6 28 6 26 <u>1</u> 92
CHEMICAL														
Heavy metal Ciguatoxin Scombrotoxin Paralytic shellfish	1	- 1 -	1 - - -	- - -	2 - - -	- 2 - 1	1 1 1 1	2 - - 2		- - - -	- 1 1 -	- 1 - -	- - - -	6 6 2 4
poison Monosodium glutamat Mushroom poison Other chemicals To	ce otal	$\frac{1}{2}$	- - - 1	1 - - 1	- - 1 3	- 1 _1 5	- - 2 6	- - 1 5	- - -		1 - - 3	- - - 1		2 1 6* 27
PARASITIC														
<u>T. spiralis</u> <u>E. histolytica</u> Te	otal		1 	$\frac{3}{-3}$		$\frac{1}{1}$	1 1		1 1			2 2		8 _ <u>1</u> 9
VIRAL														
Hepatitis A Echo, type 4 To	otal					$\frac{1}{\frac{1}{1}}$		$\frac{1}{1}$	-	1 1				2 1 3
CONFIRMED TOTAL		7	5	10	11	14	17	19	10	18	6	10	4	131
UNKNOWN		33	14	18	37	31	24	22	33	14	23	38	19	306
Total 1976 Total 1975 Total 1974		40 39 33	19 39 21	28 35 37	48 41 33	45 66 44	41 41 42	41 48 41	43 36 43	32 33 43	29 42 39	48 31 46	23 40 29	437 491** 451

*Month of occurrence not known in 1 chemical outbreaks (Niacin) **Month of occurrence not known in 6 outbreaks of unknown etiology

⁺Delayed entry, occurred December 1975

This report is authorized by law (Public Health Service Act, 42 USC 241) and is also recommended by the Conference of State and Territorial Epidemiologists. While your response is voluntary, your cooperation is necessary for the understanding and control of the diseases.

FORM APPROVED OMB NO. 68-R557

E. INVESTIGATION OF A FOODBORNE OUTBREAK

1. Where did the outbreak occur?		2. Date of outbreak: (Date of onset 1st case)
State(1,2) City c	r Town County	(3-8)
3. Indicate actual (a) or estimated (e) numbers: Persons exposed	4. History of Exposed Persons : No, histories obtained(18-20)	5. Incubation period (hours): Shortest
Persons ill (12-14)	No. persons with symptoms121-23 Nausea(24-26) Diarrhea(33-35) Vomition (27-29) Fever (36-38)	
Hospitalized(15-16) Fatal cases(17)	Cramps(30-32) Other, specify(39)	Shortest(49-51) Longest(52-54) Approx. for majority(55-57)

7. Food-specific attack rates: (58)

Food Items Served	Number of persons who ATE specified food		Number who did NOT eat specified food					
	135	Not III	Total	Percent ill	ш	Not III	Total	Percent III
· · · · · · · · · · · · · · · · · · ·								
					<u>.</u>			
								•
		 		 		' (+
					,			
			· · · · ·					
						· ·		
8. Vehicle responsible (food item incriminated by epidemic	logical evide	ence): (59,	60)			L		l

9. Manner in which incriminated food was marketed: (Check all applicable)	10. Place of Preparation of	11. Place where eaten: (66)
	Contaminated Item: (65)	
(a) Food Industry (61) (c) Not wrapped	Restaurant 1	Restaurant 1
Raw	Delicatessen	Delicatessen 2
Processed	Cafeteria	Cafeteria
Home Produced CannedVacuum Sealed	Private Home	Private Home 2 4
Raw	Caterer	Picnic
Processed	Institution:	Institution:
106 ×	School	School
(b) Vending Machine $\Box 1^{(62)}$ (d) Boom Temperature $\Box 1^{(64)}$	Church	Church 7
Befrigerated	Camp	Camp 8
Frozen	Other, specify 🗍 9	Other, specify 🗍 9
Heated	·	
If a commercial product, indicate brand name and lot number		
· · · ·		

DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE PUBLIC HEALTH SERVICE CENTER FOR DISEASE CONTROL BUREAU OF EPIDEMIOLOGY ATLANTA, GEORGIA 30333

(Over)

Specify by "X" outbreak) or ch	s sxamir ' whethe eck-up	ned: (67) Ir food ex (prepared) kamined was <u>original</u> (eaten at time of d in similar manner but not involved in	13. Environmant	st specimens rinder	exemined: (68) Findinge C. perfringens, Hobbs Type 10
outbreak)	1	Check	Findings			
Item	Orig.	up	Ouslitative Quantitative			
Example: beer	×		C, perfringens, Hobbs type 10 2X10 ⁶ /gm			
<u></u>						
				14. Specimons fro	m patients e	examined (stool, vomitus, stc.): (69)
	 			Item	No. Persons	Findings
				Example: stool	11	C. perfringens, Hobbs Type 10
					······································	
15. Specimens from	food hi	andlers (s	tool, lesions, etc.): (70)	16. Factors contr	ibuting to av	tbreak (chack all applicable) Yes No
Item		<u> </u>	Findings	1. Improper stor	age or holdin	ng temperature
17. Etiology: (77,)	78)			5 Pasr personal 6. Other, specing	0 1991 1993 hygiene of 1 	i i 2 (76
Pathogen Chemical Other				Suspected	· · · · · · · · · · · · · ·	
to conterninatio	n of foc	od, water,	; epidemic curve; etc. (Attach additiona	(page if necessary)		
	ency: ()	80)				
Name of reporting ag						(
Name of reporting ag	:				Date o	
Name of reporting ag Investigating official: NOTE: Epidemic and ment to the O To improve national	: Center f surveille	atory Ass or Diseas ance, plea	istance for the investigation of a foodbo e Control, Atlanta, Georgia 30333. ise sand a copy of this report to: Center for Disease Control Attn: Enteric Diseases Branch,	orne outbreak is availa Bacterial Diseases (Date o ble upon req Division	vest by the State Health Depart-
Name of reporting ag Investigating official: NOTE: Epidemic and ment to the (To improve national Submitted copies sho	: d Labor: Center f surveille surveille	atory Ass or Diseas ance, plea	istance for the investigation of a foodbo e Control, Atlanta, Georgia 30333. Ise send a copy of this report to: Center for Disease Control Attn: Enteric Diseases Branch, Bureau of Epidemiology Atlanta, Georgia 30333 uch information as possible, but the cor	orne outbreak is availa Bacterial Diseases (npletion of every item	Date o ble upon req Division n is not requi	red.

×.

and the second second

F. LINE LISTING OF FOODBORNE DISEASE OUTBREAKS

F. LINE LISTING OF FOODBORNE DISEASE OUTBREAKS, 1976

		Number	Date	I	ab Data			Location Where
Etiology	State	OI Cases	or Onset	Patient	Vehicle	handler	Vehicle	And Eaten
BACTERIAL								
BACILLUS CEREUS								
B. cereus	New York	8	6-18		+		Pork fried rice	(B) restaurant
B. cereus	Wisconsin	55	5-6		+		Chicken stew	(B) cafeteria
CLOSTRIDIUM BOTULINUM								
<u>C. botulinum</u> , type E	Alaska	1	1-4	+	+		White fish	(C) home
<u>C. botulinum</u> , type E	Alaska	8	7–7	+	+		Fishheads, fish eggs	(C) home
<u>C. botulinum</u> , type A	Alaska	3	9-8		+		Fish eggs	(C) home
<u>C. botulinum</u> , type B	Alaska	2	12-23	+	+		Salted salmon	(C) home
<u>C. botulinum</u> , type A	California	1	3-27	+			Unknown	(D) unknown
<u>C. botulinum</u> , type A	California	1	9-2	+			Unknown	(D) unknown
<u>C. botulinum</u> , type A	California	1	12-20	+	+		Beef pot pie	(C) home
<u>C. botulinum</u> , type B	Connecticut	1	10-14	+			Unknown	(D) unknown
<u>C. botulinum</u> , type B	Michigan	1	3-1	+	+		Vegetable & sausage mixture	(C) home
<u>C. botulinum</u> , type unknown	Minnesota	1	4-22				Unknown	(D) unknown
<u>C. botulinum</u> , type unknown	Montana	1	3-16				Unknown	(D) unknown

20

`\

<u>C. botulinum</u> , type unknown	New Mexico	1	7-4	
<u>C. botulinum</u> , type D	North Carolina	1	11-29	+
<u>C. botulinum</u> , type B	Ohio	1	9-12	+
<u>C. botulinum</u> , type A	Ohio	1	9-16	+
<u>C. botulinum</u> , type unknown	Oregon ·	1	9-12	+
<u>C. botulinum</u> , type B	Tennessee	1	11-27	+
<u>C. botulinum</u> , type B	Virginia	1	2-7	+
<u>C. botulinum</u> , type A	Washington	2	1-3	+
<u>C. botulinum</u> , type A	Washington	1	3-7	+
<u>C. botulinum</u> , type unknown	Washington	1	9–17	
<u>C. botulinum</u> , type A	Wisconsin	1	11-10	+
<u>C. botulinum</u> , type unknown	Oklahoma, Utah, Texas	7	4-14	
CLOSTRIDIUM PERFRINGE	ENS			
<u>C. perfringens</u> , Hobbs type 1	3 Illinois	4	8-22	+
C. perfringens	New Jersey	125	4–3	
<u>C. perfringens</u> , CDC type, PS 16	New York	217	7-11	+
C. perfringens	Utah	8	4-26	+

¥

. ديد ^يد

*(A)--Food processing establishment; (B)--Food service establishment;

		Green beans	(C) home
		Tomato juice	(C) home
+		Corn	(C) home
		Carrots	(C) home
		Unknown	(D) unknown
		Unknown	(D) unknown
		Unknown	(D) unknown
+		Eels	(C) home
		Unknown	(D) unknown
		Asparagus	(C) home
		Unknown	(D) unknown
		Cherry peppers	(D) restaurant
+		Prime rib	(B) restaurant
Ŧ		Roast beef, gravy	(B) school
+		Turkey	(B) camp cafeteria
+		Beef & gravy	(B) home
	(-)		

8

; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of <u>Onset</u>
<u>C. perfringens</u>	Wisconsin	5	1-25
<u>C. perfringens</u> , Hobbs types 3, 4, & PS 52	Wisconsin	150	3-7
SALMONELLA		•	
<u>S</u> ., group È, species unknown	California	73	5-22
S. heidelberg	California	7	5-?
S. typhimurium	California	12	11-8
,			
S. <u>heidelberg</u>	Colorado	339	7-23
<u>S. typhimurium</u> , <u>S. give</u>	Connecticut	ʻ 3	11-15
<u>S. typhi</u>	, Florida	8	6-15
S. <u>heidelberg</u>	Maine	78	7-4
S. typhimurium	Maine	24	9-22
<u>S</u> . <u>newport</u>	Massachusetts	9	6-?
S. muenchen	Massachusetts	35	6-?
S. typhimurium	Massachusetts	48	10-5

*			4 1	•		
Lab Data			Location Where			
Patient	Vehicle	Food . handler	Vehicle	Food Mishandled* and Eaten		
	+		Gravy	(B) home		
+			Bread & gravy	(B) cafeteria		
			, , , , , , , , , , , , , , , , , , ,			
+	+	+	Turkey, dress- ing, squash	(B) banquet room		
+		+	Mexican food	(B) restaurant		
+			Unknown	(B) social hall - mobile home park .		
+	+		Cheese	(A) restaurant		
+		+	Unknown	(B) restaurant		
+		+	Peruvian cheese & potato dish	(C) dress factory		
+			Bread stuffing, gravy, corn, Apple Betty, & topping	(B) resort inn		
+			Unknown	(B) school .		
+	+		Roast beef	(D) home		
+		+	Unknown	(D) wedding reception		
+		÷	Unknown	(B) nursing home		

<u>s</u> .	typhimurium	Michigan	7	7-5	+
<u>s</u> .	london	Minnesota	37	6-4	+
<u>s</u> .	thompson	Missouri	15	9-7	+
<u>s</u> .	typhimurium	New Hampshire	44	7-5	+
<u>s</u> .	san-diego	New Jersey	2	7-8	+
<u>s</u> .	typhimurium	New Jersey	18	9–19	+
<u>s</u> .	saint-paul	New York	54	8-26	+
<u>s</u> .	copenhagen	Oklahoma	29	5-27	+
<u>s</u> . <u>s</u>	heidelberg, schwarzengrund	Pennsylvania	17	6-21	+
<u>s</u> . <u>s</u>	<u>saint-paul</u> , . typhimurium	Pennsylvania	42	9–28	+
<u>s</u> .	blockley	Washington	58	6-12	+
<u>s</u> .	typhimurium	Washington	27	8-8	·, +
<u>s</u> .	heidelberg	Washington	24	10-16	+
<u>s</u> .	typhimurium	Wisconsin	119	1-15	+
<u>s</u> .	<u>enteritidis</u>	New York City	15	9-24	+
<u>s</u> .	bovis morbificans	Connecticut, Delaware, Massachusetts, New Jersey, Pennsylvania	21	8-?	+

¥

*(A)--Food processing establishment; (B)--Food service establishment;

+		Ice cream	(C)	home
+	+	Prime rib, roast beef, ham	(B)	restaurant
+		Tuna & macaroni salads	(B)	hospital
	+ ·	Unknown	(D)	nursing home, jail
+	+ .	Corned beef	(B)	home
+	+	Chicken casserole	(C)	church
	+	Turkey	(B)	camp
+		Salad dressing	(B)	restaurant
		Unknown	(B)	mental institution
	+	Potato salad	(B)	restaurant
		Unknown	(B)	nursing home
		Potato salad, macaroni salad	(C)	picnic
		Unknown	(B)	cafeteria
	+	Roast beef	(B)	school
	+	Japanese food	(B)	restaurant
		Precooked roast beef	(A)	restaurant & deli

Etiology	State	Number of Cases	Date of Onset	Fatient
<u>S. infantis</u>	California, Colorado	4	12-?	+
SHIGELLA				
S. flexneri 3A **	California	46	12-22-7	'5 +
<u>S</u> ., species unknown	Colorado	12	4-26	+
<u>S. sonnei</u>	Massachusetts	5	8-13	+
<u>S. boydii</u>	Texas	176	11-5	+
<u>S. flexneri 2</u>	Washington	13	6-27	+
<u>S. sonnei</u>	Washington	21	7-31	+
STAPHYLOCOCCUS				
S. aureus	California	2	2-3	
<u>S. aureus</u> , entero- toxin A, B, & D	California	4	6-15	
<u>S. aureus</u> , 85, enterotoxin D	Colorado	20	7-20	+
S. aureus	Colorado	-5	9-6	
<u>S. aureus</u> , entero- toxin A	Georgia	. 13	3-31	
S. aureus	Hawaii	9	4-28	
S. aureus	Hawaii	3	11-4	
S. aureus	Idaho	14	8-17	

Vehicle +	Food-` <u>handler</u>	Vehicle Nutrient supple- ment	Location Where Food Mishandled* And Eaten (A) hospital
		Unknown	(B) hotel
	+	Fruit compote	(B) sorority house
		Chopped liver	(B) restaurant
	+	Spaghetti	(B) military mess hall
	+	Polynesian food	(B) youth center
		Tossed salad	(C) picnic
+		Hamburger	(B) restaurant
+		Macaroni salad	(B) restaurant
+	+	Greek spaghetti	(A) office
+		Ham	(C) home
+	•	Turkey, dress- ing, gravy	(C) civic center
+		Chicken	(D) home
+		Cooked laulau	(C) home
+		Macaroni & cheese with hot dogs	(B) day-care center

S. aureus, 83A	Illinois	324	2-9	+
<u>S. aureus</u> , 6/47/53/- 54/75/+	Iowa	13	9–28	
S. aureus	Louisiana	20	5-14	
S. aureus	Louisiana	15	8-10	
S. aureus	Nevada	3	7-15	
S. aureus	New York	4	7-?	
<u>S. aureus</u> , entero- toxin D	New York	2	9-6	
S. aureus	North Dakota	35	5-31	
S. aureus	Ohio	8	4-20	
<u>S. aureus</u> , 6/42E/47/- 53/54/75/83A/84/81/+	Pennsylvania	3	1-16	+
S. aureus	Pennsylvania	3	4-23	
S. aureus	Pennsylvania	50	6~30	
S. aureus	Pennsylvania	12	8-10	
S. aureus	Tennessee	3	7–30	4
<u>S. aureus</u> , entero- toxin D	Tennessee	22	9-11	
S. aureus	Washington	3	8-4	

•

*(A)--Food processing establishment; (B)--Food service establishment **Delayed entry--occurred 12-22-75

. К.,

25

÷ 1

+	+	Turkey salad	(B)	school
+	+	Liver paté	(B)	restaurant
+		Prime rib	(B)	convention center
+		Ham, beans	(B)	fire sta- tion
+		Shrimp salad	(B)	restaurant
+		Beef ravioli	(A)	home
+		Custard-filled donuts	(A)	home
+		Ham	(C)	picnic
+		Ham	(C)	highway rest stop
+	+	Coconut custard pie	(B)	home
+		Potato salad	(C)	picnic
+		Turkey salad	(B)	drug-alcohol rehabilita- tion center
+		Chicken	(D)	meeting hall
+		Barbecue	(B)	restaurant
+		Baked beans	(B)	home
+		Pepperoni, sausage	(A)	home

; (C)--Home; (D)--Unknown; (E)--Not applicable
	Etiology S. aureus, 1136	State Washington	Number of <u>Cases</u> 336	Date of <u>Onset</u> 9-24
	<u>S. aureus</u> , entero- toxin A & D	Wisconsin	4	5-12
	YERSINIA ENTEROCOLITICA Y. enterocolitica CHEMICAL	New York	286	9-18
	Copper	Michigan	11	4-18
26	Copper	New York	14	6-15
	Cadmium	South Carolina	5	4-25
	Copper	New York City	2	2-25
	Copper	New York City	20	7-21
	Copper	New York City	3	7-29
	Ciguatoxin	Florida	2	5-?
	Ciguatoxin	Florida	1	6-21
	Ciguatoxin	Hawali	4	10-23
	Ciguatoxin	Hawa ii	_ 2	11-18
	Ciguatoxin	Virgin Island	7	1-?
	Ciguatoxin	Puerto Rico	3	5-18

La	b Data			Loca	ation Where
Pationt	Vobialo	Food-	Vobfele	Foo	d Mishandled*
racienc	venicie	nandler	venicie	Ana	Eaten
	+	+	Club salad (macaroni)	(B)	school
	+		Polish sausage	(C)	home
+	+		Chocolate milk	(A)	school
	+		Cherry flavored soft drink	(C)	church
	+		Apple cider	(D)	school picnic
	+		Apple pancakes	(C)	home
	+		Cola	(B)	restaurant
	+		Fruit punch	(B)	restaurant
	+		Soda	(B)	restaurant
			Grouper	(E)	home
			Grouper	(E)	restaurant
			Jack Fish	(E)	unknown
			Surgeon Fish	(E)	home
			Amberjack	(E)	restaurant
			Red Snapper	(E)	home

Scombrotoxin	California	1	6-1
Scombrotoxin	Colorado ·	4	10-1
Paralytic shellfish poisoning	Alaska	4	5–20
Paralytic shellfish poisoning	Alaska	2	6-30
Paralytic shellfish poisoning	Alaska	1	7-14
Paralytic shellfish poisoning	Alaska	4	7–18
Monosodium glutamate	Hawaii	3	3-6
Monosodium glutamate	Hawaii	4	10-4
Mushroom poison	. Washington	1	5-21
Sodium nitrate	California	2	6-3
Phytolacca americana (poke weed)	Oklahoma	4	5–4
Propyl paraben	Virginia	9 -	6-7
Burdock root	Washington	1	7-26
Histamine	Washington	38	1-?
Niacin	.New York City	2	4-19
Niacin .	New York City	3	?-?

*(A)--Food processing establishment; (B)--Food service establishment;

÷	Tuna	(D) restaurant
+	Mahi-Mahi	(D) restaurant
	Clams	(E)• unknown
	Clams	(E) unknown
+	Clams	(E) ship
+	Clams	(E) ship
+	Scallop soup	(B) restaurant
÷	Won Ton Mien	(B) restaurant
+	Amanita panther- ina	(C) home
+	Table salt	(A) home
+	Green salad	(C) home
+	Cake icing	(A) business office
	Burdock root	(C) home
+	Cheese	(A) restaurant
+	Hamburger	(A) home
+	Cubed steak	(A) home
(C)Home;	(D)Unknown; (E)Not a	pplicable ·

	Etiology	State	Number of <u>Cases</u>	Date of <u>Onset</u>
	PARASITIC			
	TRICHINELLA SPIRALIS			
	<u>T</u> . <u>spiralis</u>	Alaska	4	3-?
	<u>T</u> . <u>spiralis</u>	California	5	6-21
	<u>T</u> . <u>spiralis</u>	California	5	8-?
	<u>T. spiralis</u>	Massachusetts	5	11-?
	<u>T</u> . <u>spiralis</u>	Pennsylvania	2	2-?
	<u>T. spiralis</u>	Pennsylvania	2	3-?
28	<u>T. spiralis</u>	Pennsylvania	2	3-?
	<u>T. spiralis</u>	Pennsylvania	2	11-5
	ENTAMOEBA HISTOLYTICA			
	E. <u>histolytica</u>	Colorado	9	5-11
	VIRAL			
	Hepatitis A	Georgia	26	9-3
	Hepatitis A	Pennsylvania	11	5-3
	Echo, type 4	Pennsylvania	80	7–5
	UNKNOWN			
		Alabama	226	10-7

ł

I	ab Data			Location Where
Patient	Vehicle	Food- handler	Vehicle	Food Mishandled* And Eaten
				•
+			Walrus meat	(C) home
+			Pork	(D) workshop
+			Pork	(C) home
+			Pork	(D) unknown
+	+		Pork sausage, pork steak	(C) home
+	+		Pork sausage	(C) home
+	+		Pork sausage	(D) home
+	+		Pork sausage	(C) home
+			Salad	(B) fraternity house
+		+	Unknown	(B) commune
+			Unknown	(B) restaurant
+			Cole slaw	(B) picnic
			Potatoes, meat	(B) school

Ale elee	150	5-10	Meathalls	(B)	restaurant
Alaska	190	J-10	neucouris	(2)	2000000000
Alaska	39	11-17	Unknown	(B)	school
Arizona	12	1–12	Unknown	(B)	drug reha- bilitation institution
Arizona	110	10-26	Hot turkey sand- wich, mashed potatoes & gravy	(B)	school
California	19	3-6	Unknown	(B)	restaurant
California	28	5-15	Unknown	(C)	home
California	18	6-13	Unknown	(B)	dining hall
California	23	6-20	Unknown	(C)	home
California	3	7-1	Unknown	(C)	home
California	2	8-14	Unknown	(C)	home
California	508	8-26	Tuna extender	(A)	unknown
California	37	8-31	Unknown	(B)	cafeteria
California	6	10-6	Unknown	(B)	home
California	3	11-2	Unknown	(B)	restaurant
California	8	11-4	Beef	(D)	restaurant
	Ŭ				
California	29	12-4	Unknown	(B)	restaurant

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable



Etiology	State	Number of Cases	Date of <u>Onset</u>
(UNKNOWN)	Colorado	· 2	7-14
	Colorado	70	12-3
	Connecticut	5	10-17
	Connecticut	8	10-29
	Connecticut	100	10-30
	Connecticut	125	11-11
	Connecticut	4	11-29
30	Connecticut	90	11-30
	Connecticut	3	12-9
	Delaware	79	11-8
	District of Columbia	25	11-25
	District of Columbia	21	12-3
	Florida	3	5-?
	Georgia	3	2-6
	Georgia	2	8-5
	Georgia	2	8–23

L	ab Data	 		Location Where Food Mishandled*
Patient	Vehicle	handler	Vehicle	And Eaten
			Unknown	(B) restaurant
			Unknown	(D) country club
			Unknown	(D) unknown
			Macaroni & cheese	(B) cafeteria
			Unknown	(B) wedding banquet
•	*		Turkey	(B) school
			Oriental food	(B) restaurant
			Unknown	(D) women's group luncheon
			Pizza	(B) home
			Spaghetti	(B) school
			Turkéy	(B) women's club
			Mixed vegetables	(B) school
			Snapper	(E) home
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(C) home

Georgia	350	9-16	Beef stew	(B)	correction- al insti- tution
Georgia	11	12-5	Unknown	(B)	country club
Hawaii	3	2-24	Mullet	(D)	home
Hawaii	3	3-28	Noodles	(D)	home
Hawaii	36	4-9	Oyster	(B)	restaurant
Hawaii	6	4-16	Únknown	(B)	home
Hawaii	9	4-18	Raw clams	(B)	home
Hawaii	92	7-3	Roast duck with plum sauce, oyster rolls	(B)	restaurant
Hawaii	11	8-13	Unknown	(B)	restaurant
Hawaii	150	8-21	Unknown	(B)	restaurant
Hawaii	4	11-16	White fish	(D)	home
Hawaii	20	11-18	Sausage	(B)	restaurant
Hawaii	2	11-18	Baked fish	(D)	home
Idaho	590	6-19	Potato salad	(B)	local park
Idaho	12	8-15	Unknown	(B)	restaurant
Idaho	25	10-8	Unknown	(B)	restaurant
Idaho	4	11-29	Turkey	(C)	home
Illinois	35	1-3	Unknown	(B)	wedding party dinner

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

۰.

Etiology	State	Number of Cases	Date of <u>Onset</u>
(UNKNOWN)	Illinois	9	1-7
	Illinois	27	4-21
	Kansas	3	11-4
	Kentucky	7	8-28
	Kentucky	38	10-4
	Louisiana	33	8–20
32	Massachusetts	14	5-18
	Massachusetts	38	9–?
	Massachusetts	43	10-24
	Massachusetts	61	10-?
	Massachusetts	117	11-?
_	Michigan 🕚	·5	.6-3
	Michigan	66	7-?
	Minnesota	4	4-5
	Minnesota	3	4-6

t

Patient	Lab Data <u>Vehicle</u>	Food- handler	Vehicle	Loc Foo <u>And</u>	ation Where d Mishandled* Eaten
			Ham	(C)	home
			Unknown	(B)	restaurant
			Tuna salad	(B)	truck
			Chicken with gravy, dressing	(B)	cafeteria
			Unknown	(D)	club
			Tossed salad, crab meat cocktail, cornish hen	(B)	hospital dining room
			Ham	(B)	school
			Unknown	(B)	nursing home
			Roast beef & gravy	(B)	school
			Unknown	(B)	mental health facility
			Unknown	(B)	restaurant
			Beef .	(B)	restaurant
			Roast beef	(B)	nursing home
	-		Unknown	(B)	home
			Unknown S	(C)	home

Minnesota	7	4-18	Unknown	(C) home
Minnesota	16	4-28	Unknown	(B) school
Minnesota	7	10-15	Unknown	(D) restaurant
Mississippi	50	4-24	Unknown	(B) cafeteria
Mississippi	58	12-21	Turkey, dressing	(B) plant
Missouri	30	4–10	Tuna salad	(B) fraternity house
Montana	15	12-22	Unknown	(B) restaurant
Nebraska	3	6-23	Beets	(B) restaurant
Nebraska	97	11-9	Unknown	(B) hotel
Nebraska	2	11-21	Unknown	(B) restaurant
Nebraska	6	11-28	Unknown	(B) restaurant
Nevada	4	4-28	Unknown	(B) home
Nevada	10	12-5	Chinese food	(B) home
New Jersey	58	1-3	. Unknown	(B) nursing home
New Jersey	125	1-13	Turkey, gravy	(B) school
New Jersey	26	4-24	Fried rice	(B) restaurant
New Jersey	55	6-17	Chicken, vegetables	(B) motel
New Jersey	240	10-3	Unknown	(B) fire house
New York	- 22	1-11	Hamburger with sauce	(B) restaurant

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

۲

Etiology	State	Number of Cases	Date of <u>Onset</u>
(UNKNOWN)	New York	30	2-1
	New York	3	4-13
	New York	34	5-12
	New York	29	5-12
	New York	63	5-12
	New York	17	5-16
	New York	10	7-13
	New York	13	7-15
	New York	124	10-11
	New York	34	11-1
	New York	114	11-13
	North Dakota	125	1-31
	Ohio	34	3–13
	Ohio	2	3-21
	Ohio	40	7-13

.

La	b Data	Food-		Location Where Food Mishandled*	
Patient	<u>Vehicle</u>	handler	<u>Vehicle</u>	And Eaten	
			Salad	(B) restaurant	
			Fudge cake	(B) home	
			Unknown	(B) restaurant	
			Unknown	(B) restaurant	
			Unknown	(B) restaurant	
			Unknown	(B) home	
			Unknown	(D) picnic	
			Unknown	(B) nursing home- hospital	
			Hamburger, roast beef, turkey, veal	(B) school	
			Unknown	(B) wedding reception	
			Tuna salad, carrot salad, pickles	(B) restaurant	
			Oriental food	(C) church	
			Unknown	(B) hotel	
			Unknown	(B) restaurant	
			Unknown	(B) motel restaurant	

ŝ

Ohio '	24	8-13		Beef & macaroni casserole	(B)	summer camp -
Ohio	105	11-16		Gravy	(B)	restaurant
Ohio .	244	12-11		Cherries jubilee	(B)	hotel ban- quet room
Oklahoma	9	7-27	-	Unknown	(B)	officer's club
Oregon	2	3-6		Unknown	(C)	home
Oregon	9	6-6		Turkey .	(B)	school
Oregon	8	7-19		Unknown	(B)	picnic
Oregon	13	12-31		Unknown	(B)	restaurant
Pennsylvania	2	1-19		Unknown	(D)	home
Pennsylvania	4	1-31		Chinese food	(B)	restaurant
Pennsylvania	41	2-4		Potato salad, macaroni salad	(B)	restaurant
Pennsylvania	23	2-8		Unknown	(C)	camp
Pennsylvania	2	2-18		Unknown	(B)	home
Pennsylvania	5	3-7		Hero sandwich	(B)	restaurant
Pennsylvania	42	3-11		Milk	(A)	school
Pennsylvania	2	3-22		Unknown	(B)	movie theater
Pennsylvania	2	3-30		Unknown	(B)	restaurant
Pennsylvania	30	4-3		Unknown	(B)	restaurant

S.

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset
	·	• <u></u>	
(UNKNOWN)	Pennsylvania	2	4-16
	Pennsylvania	30	4-25
	Pennsylvania	3	5-3
	Pennsylvania	3	5-9
	Pennsylvania	100	5-22
	Pennsylvania	24	5-31
	Pennsylvania	29	6-18
	Pennsylvania	4	7-3
	Pennsylvania	10	7-21
	Pennsylvania	150	7-25
	Pennsylvania	12	7-27
	Pennsylvania	2	7-?
	Pennsylvania	4	8-20
	Pennsylvania	5	8-?
	Pennsylvania	63	9-6
	Pennsylvania	48	10-30
	Pennsylvania	50	11-7
	Pennsylvania	23	11-20

La	b Data			Location Where
Patient	Vehicle	Food- handler	Vehicle	Food Mishandled* And Eaten
		<u>,</u>	The last ensure	
			Unknown	(B) restaurant
			Unknown	(B) country club
			Unknown	(B) home
			Unknown	(C) home
			Unknown	(D) wedding reception
			Meatballs	(C) block party
			Unknown	(B) country club
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Cheese topping on pizza	(B) pizza parlor
			Unknown	(D) home
			Cake	(C) home
			Unknown	(B) school
			Submarine sandwich	(B) school & factory
			Unknown	(B) restaurant
			Unknown	(B) restaurant

Pennsylvania	25	12-13	Unknown	(D)	lodge
Rhode Island	8	6-25	American goulash	(C)	automobile
Rhode Island	45	6–27	Turkey dressing	(D)	church
South Carolina	2	2–20	Unknown	(B)	home
Tennessee	100	1-8	Unknown	(B)	hotel
Tennessee	2	4–20	Unknown	(B)	restaurant
Tennessee	2	7–27	Tuna salad	(C)	home
Tennessee	2	10-5	Barbecue	(B)	home
Tennessee	120	10-19	Tuna, peas, carrots	(B)	school
Texas	29	5–8	Potato salad	(C)	picnic
Texas	950	5–11	Beef tacos	(B)	school
Texas	30	5-22	Shrimp, raw crab	(B)	restaurant
Texas	150	11-2	Corn	(B)	prison
Virginia	14	4-19	Unknown	(C)	home
Virginia	12	· 7–15	Ham	(B)	military mess hall
Virginia	4	8–26	Turkey sandwich	(B)	correctional road camp
Washington	4	1-4	Unknown	(C)	home
Washington	4	1-8	Unknown	(B)	restaurant

.

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of <u>Cases</u>	Date of Onset
(UNKNOWN)	Washington .	2	1-9
•			
	Washington	5	2-22
	Washington	9	⁻ 2−29
	Washington	11	3-26
	Washington	5	3–28
	Washington	5	4-4
	Washington	2	4-4
	Washington.	3	4-11
	Washington	2	4-13
	Washington	47	5-1
	Washington	4	5-14
	Washington	2	5–17
	Washington	7	5-23
	Washington	3	6-1
	Washington	2	6-10
	Washington	4	6-21

Lab Data				Location Where
Patient	<u>Vehicle</u>	Food <u>handler</u>	Vehicle	Food Mishandled* And Eaten
			Omelet, pickled córn, cauli- flower	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Turkey á la King	(B) restaurant
		·	Ground beef	(B) restaurant
			Turkey sandwich with gravy	(B) restaurant
			Roast beef	(B) restaurant
			Ground beef	(B) restaurant
			Unknown	(D) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restauranț
			Chicken, deviled eggs	(D) picnic
			Fried fish	(D) restaurant
			Beef enchilada	(B) restaurant
			Chinese food	(B) restaurant
				4

.



Washington	11	6-29	Sirloin tips in gravy	(B) restaurant
Washington	5	7-8	Pizza	(B) restaurant
Washington	3	7–28	Relano with ground beef & refried beans	(B) restaurant
Washington	3	8–5	Chinese food	(B) restaurant
Washington	8	8-6	Barbecue chicken, potato salad	(B) restaurant
Washington	8	8-9	Chinese food	(B) restaurant
Washington	3	8-13	Unknown	(B) home
Washington	8	8–15	Unknown	(B) restaurant
Washington	17	8–15	Roast beef	(C) picnic
Washington	5	9–5	Kippered salmon	(D) home
Washington	3	9-11	Mexican food	(B) restaurant
Washington	4	9–13	Chicken, potato salad, deviled eggs	(C) picnic
Washington	4	9-16	Ground beef	(B) restaurant
Washington	11	9-?	Hollandaise sauce	(B) restaurant
Washington	7	10-23	Pizza	(B) restaurant
Washington	7	11-21	Taco meat	(B) restaurant

.

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

.

Etiology	State	Number of Cases	Date of <u>Onset</u>
(UNKNOWN)	Washington	6	12-18
	Wisconsin	2	6-26
	Wisconsin	50	8-26
	Wisconsin	26	11-26
	Guam .	10	12-18
	Guam	18	12-27
	New York City	4	. 1-1
	New York City	4	1-2
	New York City	2	1-4
	New York City	4	1-6
	New York City	2	1-10
	New York City	2	1-12
	New York City	. 2	1–12
	New York City	2	1-16
	New York City	4	1-18
	New York City	2	1–18
	New York City	4	1-18
	New York City	2	1-24
	New York City	2	1-26

La	b Data	Food-		Location Where Food Mishandled*
Patient	Vehicle	handler	Vehicle	And Eaten
			Turkey with gravy	(B) restaurant •
			Root beer	(B) automobile
			Roast beef	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(C) home
			Unknown	(D) job site
			Unknown	(C) home
			Shrimp	(B) restaurant
			Unknown	(B) restaurant
			Chinese food	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Corned beef	(B) restaurant
			Bran muffin	(B) restaurant
			Pizza	(B) restaurant
			Unknown	(B) reataurant
			Unknown	(B) restaurant
			Hamburger	(B) restaurant

L.

.

New	York (City	2	1-26	Chicken salad	(B)	restaurant
New	York (City	2	1-26	Únknown	(B)	restaurant
New	York	City	2	1-?	Unknown	(B)	restaurant _.
New	York	City	4	1-?	Chinese food	(B)	restaurant
New	York	City	2	1-?	Chinese food	(B)	restaurant
New	York	City	2	1-?	Unknown	(B)	restaurant
New	York	City	2	1-?	Ravioli	(C)	home
New	York	City	3	2-2	Unknown	(B)	restaurant
New	York	City	2	2-14	Apple	(D)	home
New	York	City	3	2-24	Unknown	(B)	restaurant
New	York	City	2	2-25	Western omelet	(B)	restaurant
New	York	City	3	2–26	Ham & roast beef sandwich	(B)	restaurant
New	York	City	5	3-1	Chinese food	(B)	restaurant
New	York	City	3	3-1	Gefilte fish	(C)	home
New	York	City	2	3-6	Unknown	(B)	restaurant
New	York	City	2	3-22	Unknown	(B)	restaurant
New	York	City	2	3-22	Pizza	(C)	home
New	York	City	2	3–24	Tuna salad	(B)	restaurant
New	York	City	2	3-27	Unknown	(B)	restaurant
New	York	City	2	4-1	Unknown	(B)	restaurant

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

×

1

ъ.

	Etiology	State	Number of Cases	Date of <u>Onset</u>
(UNKI	NOWN)	New York City	3	4-1
		New York City	3	4-2
		New York City	2	4-5
		New York City	2	4-6
		New York City	3	4-7
		New York City	2	4-9
		New York City	2	4-15
		New York City	2	4-18
49		New York City	2	4–18
		New York City	4	4–20
		New York City	3	4-20
		New York City	2	4-21
		New York City	3	4-23
		New York City	3	4-30
		New York City	2	5-1
		New York City	4	5-1
		New York City	2	5-5
		New York City	3	5-7

Patient Vehicle handler Vehicle And Eaten Unknown (D) home	.ea*
Unknown (D) home	it
	it
Unknown (B) restaurar	
Chinese food (B) restaurar	it
Hamburger (B) restaurar	t
Pepper steak (B) work plac	e
Chinese food (B) restaurar	ıt
Pot roast with (C) home tomato sauce	
Duck (B) restaurar	ıt
Chopped liver (B) restauran	t
Unknown (B) restauran	ıt
Unknown (B) restauran	ıt
Unknown (B) restauran	it
Unknown (B) restauran	it
Chinese food (B) home	
Unknown (B) home	
Unknown (D) home	
Unknown (B) restauran	it
Unknown (B) restauran	it

						•
New	York	City	2	5-8	Unknown	(C) home
New	York	City	10	5-9	Roast beef	(C) home
New	York	City	- 2	5-16	Cake	(B) restaurant
New	York	City	7 -	5-21	Beef fried rice	(B) restaurant
New	York	City	4	5–25	Chicken	(B) restaurant
New	York	City	2	5-31	Unknown	(B) restaurant
New	York	City	2	5-?	Unknown	(B) restaurant
New	York	City	2	5-?	Unknown	(B) restaurant
New	York	City	2	6-2	Unknown	(B) restaurant
New	York	City	2	6-2	Unknown	(B) home
New	York	City	11	6–5	Unknown	(D) home
New	York	City	2	6-6	Unknown	(B) restaurant
New	York	City	2	6-7	Chinese food	(B) restaurant
New	York	City	35	6–20	Ham	(D) church
New	York	City	3	6-21	Ice cream	(D) street
New	York	City	2	6-22	Unknown	(B) restaurant
New	York	City	2	6-?	Unknown	(B) restaurant
New	York	City	3	7-6	Unknown	(B) restaurant
New	York	City	5	7-10	Unknown	(B) restaurant

÷

*(A)--Food processing establishment; (B)--Food service establishment; (B)--Home; (D)--Unknown; (E)--Not applicable

	Etiology	State	Number of <u>Cases</u>	Date of <u>Onset</u>
	(UNKNOWN)	New York City	6	7-13
		New York City	2	7–30
		New York City	3	8-2
		New York City	20	8-5
		New York City	19	8-6
		New York City	4	8-7
		New York City	2	8-7
		New York City	6	8-8
4		New York City	10	8-13
4		New York City	3	8-16
		New York City	3	8-18
		New York City	3 .	8-19
		New York City	5	8-22
		New York City	2	8-29
		New York City	2	9-2
		New York City	4	9-2
		New York City	3	9-16
		New York City	2	9-20
		New York City	6	9-24
		New York City	2	9-?

I Patient	ab Data Vehicle	Food- handler	Vehicle	Location Where Food Mishandled* And Eaten
			Egg Foo Yong	(B) home
			Unknown	(B) restaurant
			Ham	(B) restaurant
			Unknown	(D) church
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(B) restaurant
			Unknown	(C) home
			Corned beef	(B) restaurant
			Unknown	(B) work place
			Unknown	(B) automobile
			Hamburger	(B) restaurant
			Unknown	(B) restaurant
			Chinese food	(B) restaurant
			Chinese food	(B) home

New	York	City	2	10-12	Hamburger	(C)	home
New	York	City	2	10-13	Mexican food	(B)	restaurant
New	York	City	2	10-14	Unknown	(B)	restaurant
New	York	City	2	10-23	Souvlaki	(B)	restaurant
New	York	City	17	10-24	Unknown	(B)	unknown
New	York	City	3	10-31	Roast beef sandwich	(B)	adult home
New	York	City	3	11-5	Unknown	(B)	restaurant
New	York	City	4	11-10	Chinese food	(B)	home
New	York	City	9	11-12	Chicken noodle soup	(B)	job site
New	York	City	3	11-16	Unknown	(B)	restaurant
New	York	City	2	11-16	Unknown	(B)	restaurant
New	York	City	351	11-17	Turkey	(B)	school
New	York	City	5	11-18	Chinese food	(B)	home
New	York	City	35	11-20	Ice cream	(B)	nursing home
New	York	City	2	11-24	Chinese food	(B)	home
New	York	City	4	11-25	Unknown	(B)	restaurant
New	York	City	3	11-26	Clams	(B)	restaurant
New	York	City	14	11-26	Unknown	(C)	home

.

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

		Number	Date	I	ab Data			Location Where
Etiology	State	of <u>Cases</u>	of <u>Onset</u>	Patient	<u>Vehicle</u>	Food- handler	Vehicle	Food Mishandled* <u>And Eaten</u>
(UNKNOWN)	New York City	4	11-29			•	Unknown	(B) restaurant
	New York City	3	12-8				Unknown	(B) work place
	New York City	6	12-21				Chocolate milk	(D) school
	New York City	2	12-27				Unknown	(B) home
	New York City	3	12-30				Unknown	(B) restaurant

.

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

• • • .

۹

L. L.

BAG	CTER1AL	Clinical Syndrome	Laboratory and/or Epidemiologic Criteria
1.	<u>Bacillus</u> <u>cereus</u>	a) incubation period 2-16 hrs. b) gastrointestinal syndrome	 a) isolation of ≥10⁵ organisms per gram in epidemiologically incriminated food OR b) isolation of organism from stools of ill person
2.	<u>Brucella</u>	 a) incubation period several days to several months b) clinical syndrome compatible with brucellosis 	a) Four fold increase in titer <u>OR</u> b) positive blood culture
3.	<u>Clostridium</u> <u>botulinum</u>	a) incubation 2 hours - 8 days usually 12-48 hours b) clinical syndrome compatible with botulism (see CDC Botulism Manual)	 a) detection of botulinal toxin in human sera, feces, or food <u>OR</u> b) isolation of <u>C</u>. <u>botulinum</u> organism from epidemiologi- cally incriminated food or stools <u>OR</u> c) food epidemiologically incriminated
4.	<u>Clostridium</u> perfringens	 a) incubation period 9-15 hrs. b) lower intestinal syndrome majority of cases with diarrhea but little vomiting or fever 	 a) organisms of same sero- type in epidemiologically incriminated food and stool of ill individuals OR b) isolation of orgamisms with same serotype in stool of most ill individuals and not in stool of controls OR c) ≥10⁵ organisms per gram in epidemiologically incrim- inated food provided specimen properly handled
5.	<u>Escherichia coli</u>	a) incubation period 6-36 hrs. b) gastrointestinal syndrome majority of cases with diarrhea	 a) demonstration of organisms of same serotype in epidemiologically incriminated food and stool of ill individuals and not in stool of controls OR b) isolation from stool of most ill individuals, organisms of the same serotype

G. Guidelines for Confirmation of Foodborne Disease Outbreak

i

	·	Clinical Syndrome	Laboratory and/or Epidemiologic Criteria which have been shown to be enterotoxigenic or invasive by special laboratory techniques
6.	' <u>Salmonella</u>	a) incubation period 6-48 hrs. b) gastrointestinal syndrome majority of cases with diarrhea	 a) isolation of <u>Salmonella</u> organism from epidemiologi-cally implicated food b) isolation of <u>Salmonella</u> organism from stools of ill individuals
7.	<u>Shigella</u>	a) incubation period 12-50 hrs. b) gastrointestinal syndrome majority of cases with diarrhea	 a) isolation of <u>Shigella</u> organism from epidemiologically implicated food <u>OR</u> b) isolation of <u>Shigella</u> organism from stools of ill individuals
8.	<u>Staphylococcus</u> <u>aureus</u>	 a) incubation period 30 min 8 hrs. (usually 2-4 hrs.) b) gastrointestinal syndrome	a) detection of enterotoxin in epidemiologically impli- cated food b) organisms with same phage type in stools or vomitus of ill individuals and, when possible, implicated food and/or skin or nose of food handler c) isolation $\frac{OR}{of} \ge 10^5$ organisms per gram in epidemiologically impli- cated food
9.	Streptococcus Group A	a) incubation period 1-4 days b) febrile URI syndrome	 a) isolation of organisms with same M and T type from implicated food OR b) isolation of organisms with same M and T type from throats of ill individuals
10.	<u>Vibrio</u> cholerae	a) incubation period 1-3 days b) gastrointestinal syndrome majority of cases with diarrhea and without fever	 a) isolation of <u>V</u>. <u>cholerae</u> from epidemiologically incriminated food <u>OR</u> b) isolation of organisms from stools or vomitus of ill individuals

ĕ

*

Laboratory and/or Clinical Syndrome Epidemiologic Criteria OR c) significant rise in vibriocidal, bacterial agglutinating, or antitoxin antibodies in acute and early convalescent sera, or significant fall in vibriocidal antibodies in early and late convalescent sera in persons not recently immunized a) isolation of $\geq 10^5$ organisms 11. Vibrio a) incubation period 15-24 hrs. parahaemolyticus from epidemiologically implib) gastrointestinal syndrome--cated food (usually seafood) majority of cases with diarrhea OR b) isolation of Kanagawapositive organisms of same serotype from stool of ill individuals Others 12. clinical data appraised in laboratory data appraised in individual circumstances individual circumstances CHEMICAL Heavy metals a) incubation period 5 min. to 1. demonstration of high 8 hrs. (usually less than 1 hr.) concentraction of metallic Antimony ion in epidemiolologically Cadmium b) clinical syndrome compatiincriminated food or Copper ble with heavy metal poisonbeverage Iron ing--usually gastrointestinal Tin syndrome and often metallic Zinc, etc taste Ichthyosarcotoxin 2. Ciguatoxin a) incubation period 1-36 hrs. a) demonstration of cigua-(usually 2-8 hrs.) toxin in epidemiologically incriminated fish b) clinical syndrome compatible OR with ciguatera--usually initial b) ciguatera-associated gastrointestinal symptoms fish epidemiologically followed by dry mouth, paresincriminated thesia of lips, tongue, throat or extremities. A sensation of looseness and pain in the teeth and a pardoxical temperature sensation are characteristic

ÿ.

	Clinical Syndrome	Laboratory and/or Epidemiologic Criteria
Puffer fish (tetro- dotoxin)	 a) incubation period 10 min. to 3 hrs. (usually 10-45 min.) b) clinical syndrome compatible with puffer fish poisoning paresthesia of lips, tongue, face or extremities often followed by numbness, loss of proprioception or a "float- ing" sensation 	 a) demonstration of tetrodo- toxin in fish OR b) puffer fish epidemiologic- ally incriminated
Scombrotoxin	 a) incubation period 1 min. to 3 hrs. (usually less than 1 hr.) b) clinical syndrome compatible with scombroid fish poisoning often including flushing, head- ache, dizziness, burning of mouth and throat, upper and lower gastrointestinal symp- toms, urticaria and generalized pruritus 	 a) demonstration of ele- vated histamine levels in epidemiologically incri- minated fish <u>OR</u> b) fish of order Scombro- dei or fish associated with scombroid poisoning (e.g., mahi-mahi) epidemiologically incriminated
3. Monosodium glutamate	 a) incubation period 3 min. to 2 hrs. (usually less than 1 hr.) b) clinical syndrome compatible with monosodium glutamate intoxicationoften including burning sensations in chest, neck, abdomen or extremities, sensations of lightness and pressure over face, or a heavy feeling in the chest 	history of large amounts (usually ≥1.5 grams) of MSG having been added to epidemiologically incriminated food
4. Mushroom poison		
Group containing ibotenic acid and muscimol	 a) incubation period 1-12 hrs. (usually less than 4 hrs.) b) clinical syndrome compatible with mushroom poisoning by this groupoften including confu- sion, delirium, visual disturbances 	a) demonstration of toxic chemical in epidemiologi- cally incriminated mushrooms OR b) epidemiologically incriminated mushrooms identified as a toxic type
Group containing amatoxins and phallotoxins, or gyromitrin	 a) incubation period 5-18 hrs. b) characteristic clinical syndrome compatible with mushroom poisoning by this groupupper and lower gastro-intestinal symptoms followed by hepatic and/or renal failure 	 a) demonstration of toxic chemical in epidemiologi-cally incriminated mushrooms OR b) epidemiologically incriminated mushrooms identified as a toxic type

.

ALC: NO. OF ALC: NO.
		Clinical Syndrome	Laboratory and/or Epidemiologic Criteria
Groups containing muscarine, psilocybin and psilocin, gastro- intestinal irritants, disulfiram-like compounds		 a) characteristic incubation period b) clinical syndrome compati- ble with mushroom poisoning by these groups 	 a) demonstration of toxic chemical in epidemiologically incriminated mushrooms <u>OR</u> b) epidemiologically incriminated mushroom identified as toxic type
5.	Paralytic and neurotoxic shellfish poison	 a) incubation period 30 min. to 3 hrs. b) clinical syndrome compatible with paralytic shellfish poisoningoften including paresthesia of lips, mouth or face and often upper and lower gastrointestinal symptoms 	 a) detection of toxin in epidemiologically incriminated mollusks OR b) detection of large numbers of shellfish poisoning-associated species of dinoflagellates in water from which epidemiologically incriminated mollusks gathered
6.	Other chemicals	clinical data appraised in individual circumstances	laboratory data appraised in individual circumstances
PAF	ASITIC AND VIRAL		
1.	<u>Trichinella</u> <u>spiralis</u>	 a) incubation period 3-30 days b) clinical syndrome compatible with trichinosisoften including fever, high eosinophil count, orbital edema, myalgia 	 a) muscle biopsy from ill individual <u>OR</u> b) serological tests <u>OR</u> c) demonstration of larvae in incriminated food
2.	Hepatitis A	 a) incubation period 10-45 days b) clinical syndrome compati- ble with hepatitisusually including jaundice, GI symp- toms, dark urine 	liver function tests compatible with hepatitis in affected persons who consumed the epidemiolog- ically incriminated food
3.	Others	clinical evidence appraised in individual circumstances	laboratory evidence appraised in individual circumstances

H. <u>Selected Foodborne Outbreak Articles</u>, 1976, <u>Taken from Morbidity and Mortality</u> Weekly Report

Staphylococcal Food Poisoning--Colorado (MMWR 26(4):22, 1977)

The Tri-County District Health Department (Denver) notified the Colorado Department of Health July 28, 1976, of a staphylococcal food poisoning outbreak among employees of a Denver business related to a food vending machine. Eleven workers had become violently ill in the mid- and late afternoon with nausea and vomiting. A few had diarrhea, but none had chills or fever. One patient was hospitalized. There was a statistically significant association between purchase of Greek spaghetti and illness (p<.001). Persons who ate other foods from the vending machines as well as some additional purchasers of Greek spaghetti did not become sick.

Because the catering firm supplying the hot food vending machines also served 15 other businesses at 29 locations in the greater Denver area, further inquiries were made to identify other affected persons. Three sick employees from 2 other businesses were found. They had typical symptoms of staphylococcal food poisoning. All 3 had also eaten Greek spaghetti from hot food vending machines on July 28. One of the 3 also was hospitalized.

Because the vending machine company routinely prepared hot food items the day before sale, a questionnaire was given to food preparation employees about illness and food consumption at work on July 27. Two out of 10 evening employees reported illnesses characterized by abrupt onset of nausea and vomiting at 10:30 PM and lasting a few hours. They had eaten Greek spaghetti that evening at 6:00 PM. Six of 8 other evening employees also ate Greek spaghetti but did not become ill with typical symptomatology. On the regular day shift, only 3 of 15 individuals had eaten Greek spaghetti; none of them had become ill.

Coagulase-positive <u>Staphylococcus</u> aureus, phage type 85, was cultured from the stool of the 2 hospitalized patients, the hands of 1 of the 5 foodhandlers preparing Greek spaghetti, the Greek spaghetti from 2 vending machines, and from the same brand of raw meat used in the Greek spaghetti.

S. <u>aureus</u> counts on Greek spaghetti recovered on July 28 at a temperature of 60 F from a vending machine at a Denver business were greater than a million organisms per gram. Greek spaghetti taken the same day from other vending machines at 160 F was culture-negative. Other hot food items prepared on the same day taken from several machines on July 28 were also found to be contaminated with <u>S</u>. <u>aureus</u>, phage type 85, in smaller numbers. This same organism was also isolated from chili made the following day. Enterotoxin studies by the Food and Drug Administration laboratories found type D enterotoxin in samples of the implicated spaghetti taken the day of the outbreak from the machine at 160 F and in samples taken from another unheated machine with a temperature of 60 F. However, all samples of <u>Staphylococcus</u> type 85 were found to produce enterotoxin type A. Type D enterotoxin was also recovered from the chili samples taken the day of outbreak at the food preparation area of the vending machine company.

The infected foodhandler had several blisters and a bandage on his hand when the cultures were taken. He handled the raw meat before cooking and, later in the day, helped portion cooked meat onto the spaghetti. In this food operation, bare hands were routinely used for preparing cooked meat. An evaluation by Tri-County District Health Department sanitarians revealed that the central commissary refrigerator was inadequate to cool foods quickly. Many hot food items were found to be at a temperature favorable for growth of \underline{S} . aureus as long as 8 hours.

Note: This investigation shows that documentation of a small number of cases may well reveal an extensive problem from a commercial product requiring public health control measures. The widespread finding of <u>S</u>. <u>aureus</u>, phage type 85, in this outbreak indicates that there was extensive cross-contamination because of poor foodhandling practices. The presence of type D enterotoxin suggests that the original toxinproducing organism was no longer recoverable at the time of the outbreak because other staphylococcal strains had overgrown the unidentified toxin-producing strain.

Salmonella bovis-morbificans in Precooked Roasts of Beef (MMWR 25(42):333, 1976)

An outbreak of salmonellosis in New Jersey, Pennsylvania, and Connecticut during August 1976 has been epidemiologically traced to precooked, packaged, ready-to-eat beef served in several delicatessens and sandwich shops.

Clinical findings in affected cases consisted of diarrhea, cramps, chills, and fever. Six of the 21 patients were hospitalized. Most cases occurred August 7-19, and most had eaten at delicatessens and sandwich shops before their illnesses.

The outbreak was first recognized when inquiries were received from the New Jersey and Pennsylvania departments of health on successive days concerning an increase in the number of isolates of <u>Salmonella bovis-morbificans</u>. Review of the national surveillance data revealed a 3-fold increase in the isolations of this serotype over a corresponding period in 1975. Excluding a 19-case March outbreak at a home for the retarded in Philadelphia, the increase occurred during the months of June, July, but predominantly August, and involved the states of New Jersey, Pennsylvania, Connecticut, and Massachusetts. The number of isolates included 11 from New Jersey, 12 from Pennsylvania, 12 from Connecticut, and 8 from Massachusetts, with 6, 5, 7, and 4 of these, respectively, occurring during the month of August. A total of 21 ill persons were interviewed in New Jersey, Pennsylvania, Delaware (which also had an isolate during August), and Connecticut.

Initial questioning of cases revealed prominent consumption of roast beef. A case-control study comparing precooked roast beef consumption among cases in New Jersey, Pennsylvania, Connecticut, and Delaware, and among age-sex matched neighbor-hood controls demonstrated a statistically significant association with the consumption of roast beef (p=.000008). The New Jersey and Pennsylvania cases had eaten roast beef at several different delicatessens which served precooked, packaged roast beef from Company A. This company's brand was significantly associated with illness (p=.00005).

Roast beef consumed by 4 of 7 cases in Connecticut was obtained from a single grocery chain delicatessen; it had received its precooked, packaged roast beef from its commissary in Boston, Massachusetts (Company B). One of the remaining 3 Connecticut cases had consumed turkey which had been purchased from Company B.

In early October, after this information was obtained, the U.S. Department of Agriculture inspected and obtained cultures from the environment and meat products at the processing plants and reviewed the cooking procedures at these companies. Inspection of Plant A revealed that the beef there was cooked to an internal temperature of 135 F. Review of the beef processing in Company B's plant revealed that beef was cooked to an internal temperature of 115 F, then removed and the temperature allowed to rise to an internal temperature of 125 F.

No <u>Salmonella</u> organisms were obtained from the meat products. Sources of raw meat in both plants were imported beef.

Editorial Note: <u>Salmonella bovis-morbificans</u> is a rare serotype; an annual average of less than 25 isolates from humans has been reported in the years 1968-1975 with the exception of 1974 when there were 60 isolates. The majority of the 1974 isolates were from a Philadelphia outbreak in which epidemiologic investigation failed to incriminate a common vehicle.

The organism S. <u>bovis-morbificans</u> has been isolated on occasion from imported beef (<u>1</u>). It has been associated with outbreaks due to beef products reported in Australia and England (<u>2</u>).

A common source of both Company A's and Company B's products contamination is suggested by the single rare serotype that caused this outbreak. In a previously reported outbreak (3) 2 different companies' products were similarly contaminated with a different rare serotype, <u>S. saint-paul</u>. The occurrence of these 2 large outbreaks caused by the precooked roasts of beef of 4 companies within the period of about 1 year implies that additional control measures that focus on control of cross-contamination and higher cooking temperatures may be needed to insure the safety of this product.

References

 Proceedings of the National Conference on Salmonellosis. March 11-13, 1964
 Jellard CH, Jolly H, and Brown RN. An outbreak of <u>S</u>. <u>bovis-morbificans</u> infection in a children's ward. Lancet 21, Feb 1959, p 390
 MMWR 25(5):34, 1976

Salmonellosis Associated with Homemade Ice Cream--Michigan (MMWR 26(12):94, 1977)

An outbreak of intestinal illness in which homemade ice cream was the presumed vehicle of infection occurred July 5, 1976, in 3 neighboring counties in Michigan.

The outbreak involved 7 individuals, ranging in ages from 2 to 63 years, who became ill with symptoms of nausea, vomiting, diarrhea, abdominal cramps, chills, fever, and myalgia. The onset of symptoms began 12 to 24 hours after consumption of the suspect food. Four persons were hospitalized, and 2 others sought medical treatment. The mean duration of hospitalization was 5 days; no deaths occurred. Stool cultures from 3 ill individuals were positive for Salmonella typhimurium.

The 7 ill individuals were members of a family who assembled on July 3 for an Independence Day celebration. The food item common to all ill individuals was homemade ice cream; 2 persons at the gathering ate only this item. Two other family members who did not eat the ice cream remained well.

The ice cream had been prepared by mixing raw fresh eggs obtained from the family farm, evaporated milk, pasteurized milk, sugar, and vanilla flavoring. The mixture (custard) was not cooked, but was frozen immediately after it was mixed. When samples of the ice cream and the ingredients used to prepare it were cultured, <u>S. typhimurium</u> was isolated both from the ice cream and from a pool of 2 remaining unbroken eggs, but not from other ingredients.

Eggs used in this preparation had been collected from feces-encrusted metal nests. The eggs may have been in the nests for as long as 7 to 10 hours before being collected. The eggs had been washed just after collection and then stored under refrigeration for approximately 24 hours before being mixed with the ice cream custard. All eggs in the batch were reported to have had intact shells; however, 2 eggs from the same batch that were submitted for culture were cracked when received at the laboratory. One week after the outbreak, 12 eggs from the same farm were cultured for <u>Salmonella</u> organisms. Cultures of the outer shell and internal contents failed to grow the organism.

Editorial Note: <u>Salmonella</u> organisms that are pathogenic to man may be present in the intestinal tract of domestic egg-laying hens. Such organisms may contaminate eggs by penetration through the egg shell and, more rarely, by direct ovarian transmission (1,2). Cracks (checks) in the egg shell may permit <u>Salmonella</u> organisms to enter the egg rapidly; however, less frequently penetration through pores in the intact shell may occur (1,2).

As in this outbreak, ungraded eggs rather than commercially graded eggs are the usual cause of such outbreaks. Eleven outbreaks of <u>Salmonella</u> <u>enteritis</u> due to homemade ice cream have been reported to CDC from 1973 to 1975; the source was farm or home-produced chicken eggs in 6 (55%), duck eggs (presumably farm or home-produced) in 2 (18%), and unknown in 3 (27%).

References

1. Williams JE, Dillard LH, Hall GO: The penetration patterns of <u>Salmonella</u> <u>typhimurium</u> through the outer structures of chicken eggs. Avian Dis 7:445-466, 1968 2. Adler HE: Salmonella in eggs - an appraisal. Food Technology 19:191-192, 1965

Outbreak of Shigellosis--Fort Bliss, Texas (MMWR 26(13):107, 1977)

An outbreak of foodborne shigellosis occurred on November 5, 1976, in a tactical unit conducting field training exercises at Fort Bliss, Texas. Of 850 soldiers at risk, 176 became ill with diarrheal disease; 53 were hospitalized.

The onset of the majority of the cases (92) was between 1 a.m. and noon on November 5; an additional 34 cases occurred during the second half of the day. Excluding the suspected index case, the range of onset was from November 4 to 9 (Figure 3).

Fig. 3 GASTROINTESTINAL DISEASE BY TIME OF ONSET, FORT BLISS, TEXAS, NOVEMBER 1976



The disease was characterized by rapid onset with fever up to 105 F, abdominal cramps, profuse diarrhea (bloody in several cases), and frequent vomiting. Many of the more serious cases with high fevers complained of severe myalgia with backache. The mean duration of the disease was 4 days, with a range from 1 to 8 days. The longest period of hospitalization was 5 days; however, most of the hospitalized cases were discharged within 48 hours. All of the cases recovered without sequelae. Stool cultures were positive for <u>Shigella boydii</u>, serotype 2, in 29 individuals.

The distribution of times of onset and the nature of the illness typified a foodborne infection originating from a common source. Although the unit was operating under field conditions, there was a common mess hall where the majority of the personnel ate their meals. A limited number of meals were prepared separately and delivered to troops at various outlying areas; most of these meals were distributed at noon.

Interviews with a large sampling of soldiers concerning food ingestion on November 3 and 4 revealed a statistically significant association between eating spaghetti at the evening meal on November 3 and subsequent diarrheal disease (Table 12).

The mean incubation period calculated from the time of ingestion of the spaghetti at the evening meal of November 3 was 50.5 hours. The spaghetti was not available for culturing. However, water, milk, and

several other foods that were available failed to demonstrate any contamination with enteric pathogens.

Of the 26 foodhandlers working in the mess hall at the time of the outbreak, 12 were symptomatic with diarrheal disease. Nine of the symptomatic and 1 of the asymptomatic foodhandlers had positive stool cultures for <u>S</u>. <u>boydii</u>, serotype 2. One of the foodhandlers responsible for preparing the spaghetti reported having had diarrheal disease at the time he prepared the spaghetti. This foodhandler had spent the preceding weekend (October 30-31) in Jaurez, Mexico; 2 days later he had onset of illness.

Table 12

Attack Rate by History of Consumption of Spaghetti, Fort Bliss, November 1976

	<u>111</u>	Not Ill	Total
Ate spaghetti	84	64	148
Did not eat spaghetti	1	12	13
Total	85	76	161
p=.001			

The meat sauce had been prepared the morning of the outbreak, while the spaghetti was prepared in the afternoon several hours before being served. The spaghetti and sauce were reportedly reheated before serving. However, field mess facilities, including those for handwashing, were limited, and there is some question whether the reheating was performed as prescribed.

The following control measures were taken:

1. All foodhandlers associated with the outbreak were removed from the mess line and rectal swabs taken. The foodhandlers were not allowed to work at that job until they had consecutive negative cultures taken at least 24 hours apart. Cultures were not taken until at least 48 hours after discontinuance of antimicrobials. (Symptomatic foodhandlers were placed on 2 gms ampicillin daily for 7 days.)

2. Meticulous attention to food preparation procedures, especially handwashing for mess personnel, which included brushing of fingers and nails, was instituted. All food service personnel were continuously monitored for signs or symptoms of disease, and proper foodhandling techniques were emphasized.

3. All persons who were ill or had a positive culture were instructed in proper sanitary practices by a community health nurse. Special attention was given to soldiers with families to insure that secondary cases did not occur in family units. All family contacts were instructed to report any occurrence of diarrheal disease.

Editorial Note: This outbreak was unusual in 2 respects: Shigellae are an uncommon cause of foodborne illness, and <u>Shigella boydii</u> is an unusual causes of <u>Shigella</u> dysentary in the United States. In 1975, only 2% of all <u>Shigella</u> isolates reported to the <u>Shigella</u> Surveillance System at CDC were <u>S. boydii</u>. Most of these infections probably were acquired during foreign travel. Although it is difficult to incriminate conclusively the individual who apparently contaminated the spaghetti sauce, it is possible that he acquired <u>S</u>. boydii infection during his trip to Mexico.

★U.S. GOVERNMENT PRINTING OFFICE: 1978-740-265/ 7082 REGION NO. 4

. **

Yersinia enterocolitica Outbreak--New York (MMWR 26(7):53, 1977)

An outbreak of intestinal illness occurred in September 1976, among school children in Oneida County, New York. <u>Yersinia enterocolitica</u>, serotype 8, was isolated from ill children and from chocolate milk that had been epidemiologically incriminated as the vehicle of transmission.

The illness, characterized by abdominal pain, fever, and in some, diarrhea, affected 218 children attending 5 Oneida County schools. Thirty-three school children were hospitalized for suspected appendicitis; 13 had appendectomies (Figure 4). In each case at surgery the appendix was normal or only slightly inflamed. Mesenteric adenopathy and inflammation of the terminal ileum were frequently observed. The removed appendices were microscopically normal or had lymphoid hyperplasia.

Fig. 4 CHILDREN FROM 5 SCHOOLS HOSPITALIZED WITH ABDOMINAL PAIN AND FEVER, BY DATE OF ONSET, SEPTEMBER-OCTOBER, 1976



Three of the 5 schools and the central food service for all 5 schools were located in 1 village (Village A). At first, the water was suspected as the source of contamination, since in August, 1 month before the outbreak, Village A residents had been instructed to boil drinking water because of deficiencies in the treatment of village water. However, a door-to-door survey, conducted in Village A and in a nearby control village with a different water supply, demonstrated that illness - defined as abdominal pain and fever since September 1 - was not associated with consumption of Village A water. The survey did demonstrate that the illness occurred predominantly in school-age children in Village A and that their illness was associated with eating lunch at school.

あったかったいましたましたが、ころうたいのである

A case-control study was performed in which a case was defined as a child from 1 of the 5 schools who had been hospitalized for suspected appendicitis during September and October. Controls were matched by age, sex, and school classroom with the cases. Of 10 possible exposures, including consumption of school water, food, and white and chocolate milk, only drinking chocolate milk at school was significantly associated with illness. Twenty-six (81%) of 32 ill children drank chocolate milk compared with 19 (59%) of 32 control children (p<.05, McNemar Test). A survey of high school students also demonstrated the association of illness with consumption of chocolate milk.

Thirty-two ill school children were found to be infected with \underline{Y} . <u>enterocolitica</u>; 27 of the isolates have been serotyped and all are serotype 8. One well child was infected with \underline{Y} . <u>enterocolitica</u>, serotype 5. \underline{Y} . <u>enterocolitica</u>, serotype 8, was isolated from 1 of 4 unopened 8-ounce cartons of chocolate milk taken from 1 of the school cafeterias during the investigation.

A local dairy was the exclusive producer of chocolate milk for the area schools. The dairy also supplied chocolate milk to 1 small grocery. In the dairy plant, is chocolate syrup was manually added to a large open vat of pasteurized milk. This chocolate milk was not re-pasteurized before being placed in cardboard, half-pint cartons. Milk was distributed to the schools in an unrefrigerated truck. Several cultures of canned chocolate syrup were negative. No dairy employees were culturepositive. The dairy voluntarily ceased production of chocolate milk when informed in late October of the evidence associating its chocolate milk with illness.

Editorial Note: This is the first outbreak of illness from <u>Yersinia enterocolitica</u> in which foodborne transmission has been documented. In 2 previous outbreaks among school children the source and mode of spread of the infection were not established (<u>1</u>). Milk was suspected as the vehicle in a recent outbreak of <u>Y</u>. <u>enterocolitica</u> in Canada (2).

The predominant symptoms in this outbreak - abdominal pain and fever - can only closely simulate appendicitis, but actually represent mesenteric adenitis and in some cases terminal ileitis. In the Scandanavian countries, where yersiniosis has been more extensively studied, infection with Y. <u>enterocolitica</u> can be demonstrated by stool/appendix culture in 3-5% of patients with symptoms of appendicitis (3,4). Other clinical syndromes, including abscesses, acute diarrhea, erythema nodosum, and arthritis have also been reported in association with <u>Yersinia</u> infection.

References

1. Asakawa Y, Akahane S, Kagata N, et al: Two community outbreaks of human infection with Yersinia enterocolitica. J Hyg (Camb) 71:517-723, 1973

2. Laboratory Centre for Disease Control: Canada Diseases Weekly Report 2:41-44, 73-74, 1976

3. Nilehn B, Sjöström B: Studies on <u>Yersinia</u> enterocolitica. Acta Pathol Microbiol Scand 71:612-628, 1967

4. Jepsen OB, Korner B, Lauritsen KB: <u>Yersinia</u> enterocolitica infection in patients with acute surgical abdominal disease. Scand J Infect Dis 8:189-194, 1976

III. WATERBORNE DISEASE OUTBREAKS, 1976

In 1976, 35 waterborne disease outbreaks were reported to the Center for Disease Control for the United States, an increase of 46% over 1975.

A. Definition of Outbreak

A waterborne disease outbreak is defined in this report as an incident in which 1) 2 or more persons experience similar illness after consumption of water, and 2) epidemiologic evidence implicates the water as the source of illness.

There is 1 exception; 1 case of chemical poisoning constitutes an outbreak if the water is demonstrated to be contaminated by the chemical. In most of the reported outbreaks, the implicated water source was demonstrated to be contaminated; only outbreaks associated with water used for drinking are included. Β.

Sources of Data

Waterborne disease outbreaks are reported to CDC by state health departments. A standard reporting form that was pretested in 8 states is now being used (see Section E). In addition, the Health Effect's Research Laboratory, Environmental Protection Agency (EPA), contacts all state water supply agencies to obtain information about waterborne disease outbreaks and these data are included in this report. Personnel from CDC and EPA work together in the investigation and evaluation of waterborne disease outbreaks. When requested by a state health department, CDC and EPA can offer epidemiologic assistance, provide expertise in the engineering and environmental aspects of water purification, and as indicated, provide large volume water sampling for isolation of viruses, parasites (Giardia), and specific bacterial pathogens. Data obtained on outbreaks are reviewed and summarized by representatives from CDC and EPA. A line listing of reported waterborne disease outbreaks in 1976 is included (see Section F).

In this report, municipal systems are defined as public or investor-owned water supplies that serve large or small communities, subdivisions and trailer parks of at least 15 service connections or 25 year-round residents. Semipublic water systems are present systems in institutions, industries, camps, parks, hotels, service stations, etc., which have their own water system available for use by the general public. Individual water systems, generally wells and springs, are those used by single or several residences or by persons traveling outside of populated areas (e.g. backpackers).

с. Interpretation of Data

Data included in this summary of waterborne disease outbreaks have limitations similar to those outlined in the foodborne disease summary and must be interpreted with caution since they represent only a small part of a larger public health problem. These data are helpful in revealing the various etiologies of waterborne diseases, the seasonal occurrence of outbreaks, and the deficiencies in water systems that most frequently result in outbreaks. As in the past the pathogen(s) responsible for many outbreaks in 1976 remains unknown. It is hoped that advances in laboratory techniques and standardization of reporting of waterborne disease outbreaks will augment our knowledge of waterborne pathogens and the factors responsible for waterborne disease outbreaks.

Analysis of Data **D**.

In 1976, 35 waterborne disease outbreaks, an increase of 46% from 1975 (24 outbreaks) and 5,068 cases, a decrease of 53% from 1975 (10,879 cases), were reported to CDC (Table 1). Increased reporting by certain states probably accounts for the increased number of outbreaks in 1976. Of 35 outbreaks, Pennsylvania reported 14 (40%), affecting 424 individuals (median of 21 per outbreak); 12 involved semipublic water and 2 individual water systems.

Figure 1 shows the geographic distributions of outbreaks by state. Sixteen states and Puerto Rico reported at least 1 outbreak. Figure 2 depicts the trend in reported waterborne disease outbreaks in the period 1938-1976.

Table 2 shows the number of outbreaks and cases by etiology and type of water system. Of 35 outbreaks 26 (74%) were designated as "acute gastrointestinal illness." This category includes outbreaks characterized by upper and/or lower gastrointestinal

symptomatology for which no specific etiologic agent was identified. In previous years these outbreaks were grouped under the category "sewage poisoning." There were 9 (26%) outbreaks of known etiology: chemical (3), <u>Giardia lamblia</u> (3), <u>Shigella</u> (2), and <u>Salmonella</u> (1). In the 3 largest outbreaks an etiologic agent was found; <u>Shigella sonnei</u> in Puerto Rico (2,150 cases), <u>Salmonella typhimurium</u> in New York (750 cases) and Giardia lamblia in Washington (600 cases).

The 3 chemical outbreaks reported were due to lead (2.2 mg per liter in water samples), chlordane (a pesticide - 1,200 mg per liter in water samples) and polychlorinated biphenyls (pcb's - 900 mg per liter in water samples). In the 33 nonchemical outbreaks, microbiologic water sample results were reported in 28. Evidence of fecal contamination (total or fecal coliforms) or pathogens were found in water samples collected during 27 of the outbreaks. Salmonella typhimurium was isolated from water in the New York outbreak and Giardia cysts were isolated from water in outbreaks in Colorado, Vermont, and Washington. In outbreaks where pathogens were isolated from the water supply, coliforms were reported in only 1, an outbreak of giardiasis involving the use of untreated surface water where 23 coliforms per 100 ml (MPN) were found. The other outbreaks of giardiasis involved surface water sources that were disinfected, and it is possible that chlorination was sufficient to destroy indicator organisms such as colliforms but not Giardia cysts. The outbreak of salmonellosis was caused by a cross-connection, and it is not known if timely water sampling for coliforms was conducted in conjunction with the sampling for pathogens. It is important that an attempt be made to isolate pathogens from the water supply during an outbreak to help establish the etiology, but it is equally important to also document the presence of coliforms and document their relative importance as indicator organisms for use in routine surveillance of water supplies.

Most outbreaks involved semipublic (66%) and municipal (26%) water systems, and fewer involved individual (8%) systems (Table 3). This distribution is almost identical to 1975. Outbreaks attributed to water from municipal systems affected an average of 418 persons compared with 55 persons in outbreaks involving semipublic systems and 15 persons in outbreaks associated with individual water systems. Deficiencies in treatment (inadequately or untreated water) accounted for 29 (83%) of the outbreaks. Untreated water (surface or ground) accounted for 18 of the 29 outbreaks.

Of the 23 outbreaks associated with semipublic water supply systems, 17 (74%) involved visitors to areas used mostly for recreational purposes. Of these 17, 13 occurred in the summer months May through September (Table 4). Comments

The 46% increase in the number of outbreaks reported in 1976 is probably due to more complete reporting. Diligent investigation, such as was done in outbreaks reported from Pennsylvania, can uncover relatively small waterborne outbreaks that often originate from semipublic water systems. It is hoped that similar investigation and reporting will be done by other states so that major deficiencies commonly affecting semipublic water systems, especially in recreational areas, can be better understood and ultimately corrected.

As in recent years outbreaks originating from semipublic water systems in recreational areas contributed significantly to the total number of waterborne outbreaks reported in 1976. Water systems used on a seasonal basis or those that do not usually have an overwhelming demand placed on them by large numbers of visitors are showing the strains of such pressure. Water supply systems in such areas, especially national, state, and local parks, must be routinely reappraised and monitored and corrections made to insure safe water under increased demands. The large outbreak (more than 1,000 cases) that occurred in 1975 in Crater Lake National Park underscores the actual and potential problems that can occur in recreational areas.

Coliform organism identification is used as an indication of fecal contamination of water supplies and is widely employed in routine surveillance programs. Negative results have usually been interpretated as providing assurance that the water is free of enteric pathogens. This interpretation must be reevaluated in light of data available from waterborne outbreaks of giardiasis. In the 2 outbreaks of giardiasis where disinfection was provided, <u>Giardia</u> cysts were found in the water supply in the absence of coliforms. Although adequate disinfection data are not currently available, it is felt that <u>Giardia</u> cysts are as resistent to chlorination as cysts of <u>Entamoeba histolytica</u>, and high concentrations of chlorine and long contact times would, therefore, be required for cyst inactivation. Almost all of the outbreaks of giardiasis documented in the U.S. since 1965 have occurred as the result of drinking untreated surface water or surface water whose only treatment was disinfection. Disinfection practices normally employed in these systems would not provide for high concentrations of chlorine or long contact times, and it's likely that <u>Giardia</u> cysts could survive the treatment whereas coliforms would not. The coliform test in these situations would not provide assurance that an outbreak of giardiasis would be prevented.

The giardiasis outbreak in Washington is the first documented waterborne outbreak of giardiasis involving a filtered water supply. Treatment for the surface water source consisted of a mixed-media pressure filter and disinfection; no sedimentation was employed prior to filtration. In the outbreak, failure of the chlorination equipment occurred, and a number of deficiencies were noted in the installation and operation of the pressure filters, including ineffective pretreatment or conditioning of filters with appropriate chemicals. Water filtration theory indicates that organisms the size of Giardia cysts should be removed by conventional sand filters; however, effective pretreatment of the water prior to filtration must be accomplished. Conventional treatment of surface water generally includes coagulation/flocculation and settling prior to filtration or if the settling process is not used the addition of appropriate chemicals for conditioning of the filter media. Pressure filters are generally utilized for iron and manganese removal and for a number of reasons are generally not considered effective for microbiological treatment. The data to date would indicate that well operated conventional treatment plants employing coagulation/ flocculation, settling, and filtration are successful in preventing outbreaks of this disease.



Fig. 2 AVERAGE ANNUAL NUMBER WATERBORNE DISEASE OUTBREAKS, 1938-1976



*NUMBER CASES FOR 1976 ONLY

Table 1

Waterborne Disease Outbreaks, 1972--1976

	1972	<u> 1973*</u>	1974*	1975	<u>1976</u>	Total
Outbreaks	29	26	25	24	35	139
Cases	1,638	1,774	8,356	10,879	5,068	27,715
*Revised to	otals					

Table 2

Waterborne Disease Outbreaks, by Etiology and Type of Water System, 1976

	MUNICI	PAL	SEMIPUBLIC		INDIVIDUAL		TOTAL	
	Outbreaks	Cases	Outbreaks	Cases	Outbreaks	Cases	Outbreaks	Cases
Acute gastro-								
intestinal								
illness	4	229	21	1,216	1	24	26	1,469
Chemical ·								
poisoning	1	13	0	0	2	22	3	35
Giardiasis	1	600	2	39	0	0	3	639
Shigellosis	2	2,175	0	0	0	0	2	2,175
Salmonellosis	l	750	0	0	0	0	1	750
Enterotoxi-								
genic E. coli	0	0	0	0	0	0	0	0
Hepatitis	0	0	0	0	0	_0	_0	0
TOTAL	9	3,767	23	1,255	3	46	35	5,068

Table 3

Waterborne Disease Outbreaks, by Type of System, and Cause of System Deficiency, 1976

•	MUNICI	PAL	SEMIPU	BLIC	INDIVID	UAL	TOTAL	·
	Outbreaks	Cases	Outbreaks	Cases	Outbreaks	Cases	Outbreaks	Cases
Untreated surface water	1	25	2	39	1	24	4	- 88
Untreated ground water	2	77	11	790	1	20	14	887
Treatment deficiencies	3	2,900	8	362	0	0	11	3,262
Deficiencies in distribu-					_	'		0.05
tion system Miscellaneous	2 1	763 2	1 1	60 4	1 0	2	<u> </u>	825
TOTAL	9	3,767	23	1,255	3	46	35	5,068

Table 4

Waterborne	Disease Out	breaks	Involving	Semipublic	: Water	Supplies,
	by Month	, and H	Population	Affected,	1976	

	Number of	Usual	
Month	<u>Outbreaks</u>	Population*	Visitors**
Tomorem			
January	_	-	-
February	1	1	-
March		-	-
April	3	2	1
May	. 3	1	2
June	7	2	5
July	5	-	5
August	1	-	1
September	-	-	-
October	1	-	1
November	1	-	1
December	1	-	1
TOTAL	23	6	17

*Outbreaks affecting individuals using the water supply on regular basis

**Outbreaks affecting individuals not using the water supply on a regular basis

Table 5

Waterborne Disease Outbreaks, by Month of Occurrence, 1976

	Number of		Number of
Month	Outbreaks	Month	Outbreaks
January	0	July	7
February	2	August	2
March	2	September	0
April	4	October	3
May	5	November	2
June	7	December	1

TOTAL 35

DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE PUBLIC HEALTH SERVICE CENTER FOR DISEASE CONTROL BUREAU OF EPIDEMIOLOGY E. INVESTIGATION OF A WATERBORNE OUTBREAK

教育を知 1

「「「「「「「「」」」」 1

.

FORM APPROVED OMB NO. 68-R557

.

ł.

occur?						2.	Date of ou	tbreak: (Date o	of onset of 1st case)	
(1-2)	City or To	วพก		_ County					(3-8)	
timated	4. Hist	4. History of exposed persons:					5. Incubation period (hours):			
	No.	histories ob	tained		(18-20)	Sno	rtest		est (43-45)	
	No.	persons wit	h symptoms		(21-23)		Wedian	(40		
(15-16)	Nau	sea	(24-26) Dia	rrhea	(33-35)	6. Dur	ation of ill	Iness (hours):	•	
(17)	Von Crar Oth	niting mps er, specify =	(27-29) Fev (30-32) (39)	er	(36-38)	Sho	ortest Mediar	(49-51) Long 1(5)	est (52-54) 5-57)	
, attack rates [no of water consum	umber ill/i ed, anecd NU	number exp otal informa	osed) for perso ition) * (58) PERSONS WHO	ons who did or di	d not eat	or drin	k specific BER WHC	food items or w	rater,	
	DR	DRANK SPECIFIED FOOD OR WATER					SPECIFIED FOOD OR WATER			
	ILL	NOT ILL	TOTAL	PERCENT	11	L	NOT ILL	TOTAL	PERCENT ILL	
	occur? (1-2)	occur? (1-2) City or Tc timated 4. Hist (9-11) No. (12-14) No. (15-16) Nau (17) Von Crar Oth of water consumed, anecd D ILL	occur? (1-2) City or Town	occur? (1-2) City or Town	occur? (1-2) City or TownCounty timated 4. History of exposed persons: (9-11) No. histories obtained (12-14) No. persons with symptoms (17) Nausea(24-26) (17) Vomiting(27-29) (17) Vomiting(30-32) Other, specify (39)	occur? (1-2) City or TownCounty	occur? 2. (1-2) City or Town County	occur? 2. Date of or (1-2) City or Town County County timated 4. History of exposed persons: 5. Incubation persons (12-14) No. histories obtained (18-20) (12-14) No. persons with symptoms (21-23) (17) Nausea (24-26) Diarrhea (33-35) (17) Vomiting (27-29) Fever (36-38) (17) Cramps (30-32) (36-38) (17) Other, specify (39)	occur? 2. Date of outbreak: (Date of outbre	

3. Water supply characteristi	cs (/	A) Type of wa	ter sup	oply**	(61)	
			bal or c	ommu	nity sug	oply (Name)
		🗍 Individu	ual hou	sehold	supply	
		Semi-pu	ublic w	ater sup	pply	
		🗖 Insti	tution	, schoo	l, churc	h
•		🗋 Cam	p, recr	eationa	area	
		🛛 Othe	ər,			
		Bottled	water			
(B) Water source (check a	ll applicable):	(C) Tre	atment	t provid	led (cir	Cle treatment of each source checked in B):
(82-83)		а	ь	с	d	a. no treatment
Spring		а	b	c	d	b. disinfection only
🗌 Lake, po	nd	· a	b	с	d	c. purification plant - coagulation, settling, filtration,
🗋 River, st	ream	а	Ъ	С	ď	disinfection (CITCIE those applicable)
				_		
O. Point where contaminati	on occurrea: (t	201		_		
Raw water source		nent plant			Distri	bution system
*See HSM 4.245 (NCDC) I *Municipal or community Semipublic water systems to drinking water. These I obtain water from a muni	nvestigation of a water supplies a are Individual-t ocations include cipal water syste	Foodborne (re public or in ype water sup schools, cam em but have d	Dutbre vestor plies se ps, par levelop	ak, Iter owned erving a ks, reso ed and	m 7. I utilitie I group orts, ho mainta	es, Individual water supplies are wells or springs used by single residences, of residences or locations where the general public is likely to have access tels, industries, institutions, subdivisions, trailer parks, etc., that do not in their own water supply.
obtain water from a muni	cipal water syste	am but have d	evelop	ed and	mainta	in their own water supply.

					FINDIN	IGS	BACTERIOLOGIC TECHNIQ
ı	TEM	ORIGIN	AL CHECK UP	DATE	Quantitative	Qualitative	le.g., fermentation tube, membrane filter)
	Tap wat	ter X		6/12/74	10 fecal coliforms per 100 ml.		
Examples: -	Raw wa	ter	x	6/2/74	23 total coliforms per 100 ml.		
<u></u>	- <u></u>						
Treatment	ecords: (In	dicate method i	used to determine	chlorine residua	//):		
		esidual One efflu chloi Thre on 6	ent on 6/11/74 rine e samples from dis /12/74 no residu	trace of free stribution system ral found			
Specimens f	rom patient	s examined (sto	ol, vomitus, etc.)	(68)	14. Unusual occurren	ce of events:	
SPECIN	/IEN	NO. PERSONS	FINDIN	IGS	Example: Repair sewage	of water main 6/11 , no main disinfection	/74; pit contaminated with on. Turbid water reported
Example: St	fool	11	8 Salmonella typh	i	by con	sumers 6/12/74.	
			3 negative				
			·····	<u>. </u>			
····			······				
Overflov Seepage Flooding Use of u	v of sewage of sewage J, heavy rain Intreated was		nterruption of disi nadequate disinfec Deficiencies in othe Cross-connection	nfection stion er treatment pro		nproper construction lise of water not inter contamination of stor ontamination throut	on, location of well/spring ended for drinking orage facility Igh creviced limestone or fissured i
U Water in	dequately t	reated D(Contamination of n	nains during co	nstruction or repair		
Etiology: (6	9-70)						(71)
Chemical					Suspected	•••••	
Other					Unknown		
Remarks: Bi leading to co	iefly descrit Intamination	be aspects of th n of water; epid	e investigation not lemic curve; contro	covered above, ol measures imp	such as unusual age or se lemented; etc. (Attack ac	x distribution; unus Iditional page if nec	sual circumstances essary)
ne of reporti	ng agency:	(72)					
stigating Of	ficial:				Date of inve	stigation:	
Note: Epid to th	emic and La e Center for	aboratory assist r Disease Contro	ance for the invest ol, Atlanta, Georgi	igation of a wat a 30333.	erborne outbreak is avail	able upon request b	y the State Health Department
To improve	e national su	rveillance, plea	se send a copy of t	his report to:	Center for Disease Contr Attn: Enteric Diseases Bu Bureau of Epidemi	ol anch, Bacterial Dise ology	eases Division

.

F. LINE LISTING OF WATERBORNE DISEASE OUTBREAKS, 1976

F.	Line Listing	of	Waterborne	Disease	Outbreaks,	1976

÷

.

State	Month	Disease	Cases	Type of System.	Location of Outbreak	System Deficiency*
Alaska	August	Shigella flexneri	25	Municipal	Residence	1
Arkansas	July	Acute gastrointestinal illness	51	Semipublic	Camp	3
California	May	Acute gastrointestinal illness	60	Semipublic	Camp	4
California	May	Acute gastrointestinal illness	46	Municipal	Resort	2
California	July	Acute gastrointestinal illness	2	Municipal	Sewage Plant	5
Colorado	February	Giardiasis	12	Semipublic	Office, Residence	1
Colorado	May	Acute gastrointestinal illness	10	Semipublic	Restaurant	2
Colorado	June	Giardiasis	27	Semipublic	Camp	1
Colorado	December	Acute gastrointestinal illness	110	Semipublic	Camp	3
Connecticut	June	Acute gastrointestinal illness	300	Semipublic	Country Club	2
Florida	November	Acute gastrointestinal illness	31	Municipal	Residence	2
Idaho	June	Acute gastrointestinal illness	100	Semipublic	Camp	2
Massachusetts	July	Acute gastrointestinal illness	18	Semipublic	Park	2

•

•

.

Mississippi	October	Polychlorinated Biphenyls (PCB's)
New Jersey	October	Acute gastrointestinal illness
New York	March	Salmonella typhimurium
Oklahoma	April	Acute gastrointestinal illness
Pennsylvania	April	Acute gastrointestinal illness
Pennsylvania	April	Acute gastrointestinal illness
Pennsylvania	April	Lead poisoning
Pennsylvania	Мау	Acute gastrointestinal illness
Pennsylvania	June	Acute gastrointestinal illness
Pennsylvania	July	Acute gastrointestinal illness
Pennsylvania	July	Acute gastrointestinal illness
*(1) Untreated	surface water	(2) Untreated ground water

(1) Untreated surface water (2) Untreated deficiencies (5) Miscellaneous

I

20	Individual	Residence	2			
104	Semipublic	Restaurant	2			
750	Municipal /	Resort	4			
65	Semipublic	School	2			
4	Semipublic	Recreational area	5			
30	Semipublic:	Country Club	3			
2	Individual	Residence	4			
35	Semipublic	Country Club	3			
10	Semipublic	Recreational area	2			
26	Semipublic	School, Church	3			
34	Semipublic :	Camp	3			
5	Semipublic :	Recreational area	2			
10	Semipublic .	Restaurant	3			
150	Semipublic	Restaurant	2			
(3) Treatment deficiencies (4) Distribution system						

.

The Mar Land

State,	Month	Disease	Cases	Type of System	Location of Outbreak	System Deficiency*
Pennsylvania	July	Acute gastrointestinal illness	18	Semipublic	Restaurant	2
Pennsylvania	August .	Acute gastrointestinal illness	10	Semipublic	Restaurant	. 2
Pennsylvania	October	Acute gastrointestinal illness	24	Individual	Residence .	1
Pennsylvania	November	Acute gastrointestinal illness	66	Semipublic	Restaurant	3
Tennessee	March	Chlordane (pesticide)	13	Municipal	Residence	4
Vermont	February	Acute gastrointestinal illness	150	Municipal	Residence	3
Washington	May	Giardiasis	600	Municipal	Residence	3
Puerto Rico	July	<u>Shigella</u> sonnei	2,150	Municipal	Residence	3

 *(1) Untreated surface water (2) Untreated ground water (3) Treatment deficiencies (4) Distribution system deficiencies (5) Miscellaneous

. .

70

G. Selected Waterborne Outbreak Articles, 1976, Taken from Morbidity and Mortality Weekly Report

Waterborne Giardiasis Outbreaks--Washington, New Hampshire (MMWR 26(21):169, 1977

Two waterborne outbreaks of giardiasis have been reported to CDC in the past year. One occurred in Camas, Washington (pop. 6,000), in the spring of 1976; in this outbreak 128 people had laboratory-confirmed giardiasis. The other outbreak, still ongoing, is in Berlin, New Hampshire (pop. 15,000), where 205 people to date have developed confirmed giardiasis.

<u>Camas</u>: On May 6, 1976, the laboratory section of the Washington State Department of Social and Health Services contacted CDC to report a large number of <u>Giardia</u>-positive stools from Camas. Only 2 people from Camas had been stool-positive for <u>Giardia</u> in 1975, whereas the same laboratory had reported 32 positives in April and May of 1976. The 32 patients' residences were scattered throughout the town, and they had limited interpersonal contact, suggesting waterborne transmission. Therefore, an area of the city where half of the residents received Camas city water and the other half used private well water was chosen for a preliminary survey. Six of the 38 users of city water compared to none of 40 users of private water had an illness compatible with giardiasis (p=0.01), implicating waterborne transmission. A larger study was undertaken to define the extent and character of the outbreak.

Two mutually exclusive groups were investigated: those people who were ill and spontaneously sought medical care (hereafter called confirmed cases) and those people who were interviewed during a survey and found to be ill (clinical cases). The confirmed cases consisted of 128 people who voluntarily contacted their physicians reporting a diarrheal illness and were stool-positive for <u>Giardia</u>. Analysis of data obtained from confirmed cases and their medical records revealed that diarrhea for 10 or more days was the single statistically significant symptom. Among confirmed cases, the outbreak began during the first week in April and peaked the first week in May. The outbreak spontaneously declined on May 10, and on May 15 the city switched to well water exclusively to prevent any further exposures by surface water.

The second group consisted of the respondents to a randomized community questionnaire survey administered to 496 Camas residents and 318 residents in an adjacent control town (receiving only well water). Because diarrhea of 10 or more days was characteristic of confirmed cases it was used as the case definition to interpret the survey questionnaires. Nineteen people (4%) of Camas respondents fit the case definition for giardiasis; none did in the control town (p=0.01). Thus, at least 240 persons (clinical cases) were ill with giardiasis in Camas. The stools of 18 people - 9 well and 9 ill with any diarrheal illness - were examined; no viral or bacterial pathogens were found. Two of the ill persons (22%) and 1 of the not ill (11%) were stoolpositive for <u>G</u>. <u>lamblia</u>. Giardiasis was not associated with pet ownership, travel, or recreational activities such as swimming which involve raw water.

Camas has 2 water sources - a pair of mountain streams and a set of deep wells. Those residents living in areas receiving less than 70% surface water (more than 30% well water) reported no cases, while those receiving more than 70% surface water had an attack rate of 4.7%. <u>Giardia cysts were recovered from the raw surface water</u> entering the city's water treatment plant. Because the city chlorinated and filtered its surface water supplies in a closed pressurized system, flocculation efficiency was marginal. Sedimentation could not be used. <u>Giardia cysts were also recovered from</u> 2 reservoirs holding water which had already passed through the water treatment facility (finished water). Deep well water used by the city was not contaminated. An investigation of the watershed revealed 2 remote mountain streams in a fenced area with no evidence of human contamination. Several animals near the watershed were trapped. Trapping yielded 9 negative animals (including coyote, opossum, nutria, porcupine, and beaver) and 3 positive beavers. The beavers lived in a pond bordering a heavily used state park, but were within foraging distance of the water intakes for Camas.

Berlin: On April 19, 1977, a medical technologist at a local hospital in Berlin called CDC to report that 10 cases of giardiasis had been diagnosed in the past 9 days. By April 26, New Hampshire had reported a total of 90 cases in comparison to no cases of giardiasis reported in Berlin in the previous 5 years. Because cases were randomly distributed throughout the community, waterborne transmission was suspected.

Again, 2 groups were investigated: those people who were ill, voluntarily sought medical care, and were stool-positive (confirmed cases) and those people who were interviewed during a survey and were found to be ill (clinical cases). As of May 20, there were 205 confirmed cases. The outbreak began on April 8 and peaked on April 25. On April 22, Berlin residents were instructed to boil drinking water, and the city increased its level of chlorination. However, approximately 5 people per day continue to be diagnosed as stool-positive for G. lamblia.

A randomized community questionnaire survey was done in Berlin (692 surveyed) and in an adjacent control town (286). One hundred sixty-five people (24%) in Berlin and 31 people (11%) in the control town reported diarrheal illness. However, because analysis of confirmed cases is not yet complete, the case definition for giardiasis in this outbreak has not been established. Therefore, the percentage of diarrheal illness attributable to <u>Giardia</u> infection has not yet been determined.

Berlin uses 2 rivers for its water supply: The Amonoosuc and the Androscoggin. People receiving Amonoosuc River water and those receiving Androscoggin River water had similar attack rates of diarrheal illness (23% vs. 27%, respectively). <u>Giardia</u> cysts have been recovered from the raw water from both rivers. <u>Giardia</u> cysts were recovered from 3 sites within the distribution system, including the regional hospital.

An investigation of the watershed revealed that the Amonoosuc River is a small stream located in the White Mountain National Forest. However, access is not restricted, and an estimated 3,000 people used the area for recreational activities during October, November, and December 1976. The water is chlorinated and filtered under pressure without sedimentation or flocculation. The physical plant is 30 years old, and 3 of its filters were badly worn. The Androscoggin River receives untreated sewage effluent from a number of homes in 2 communities upstream from Berlin. Because of the known sewage contamination of the Androscoggin, a new water treatment plant was put in service on March 10, 1977. However, because of cross connections secondary to faulty construction and difficulty creating the proper weight floc, the new plant was ineffective. The town is repairing the plant.

Editorial Note: An outbreak of giardiasis in Rome, New York, in the spring of 1975 was the first laboratory-documented epidemic of waterborne giardiasis in the United States; it affected over 4,800 people (MMWR 24:(43), 1977). The outbreak the following spring in Camas, where at least 240 people developed giardiasis, again demonstrated the ability of <u>Giardia</u> organisms to cause citywide outbreaks of diarrhea. Significant morbidity was demonstrated, as the illness produced was characterized by prolonged diarrhea (> 10 days).

In Rome, the absence of filtration and optimum chlorination left the city unprotected against waterborne giardiasis. In Camas and in Berlin (on the Amonoosuc), pressure filters without sedimentation and proper flocculation failed to remove <u>Giardia</u> cysts. The Androscoggin water treatment plant in Berlin has sedimentation flocculation and rapid sand filters; however, flocculation difficulties and cross connections between unfiltered and finished water decreased the plant's effectiveness. Nevertheless, it has been shown that properly functioning sedimentation, flocculation, and filtration will remove particles the size of <u>Giardia</u> cysts from water, and thus can provide safe drinking water in distribution systems utilizing surface water (<u>1</u>).

Reference

1. Fair GM, Geyer JC: Water Supply and Waste-Water Disposal. New York, John Wiley & Son, Inc., 1956

Probable Viral Gastroenteritis--Colorado (MMWR 26(3):13, 1977)

An outbreak of probable waterborne viral gastroenteritis occurred during the week before Christmas among vacationers at a winter resort near Granby, Colorado. Over 700 persons were registered at the camp during the outbreak. Of 208 surveyed thus far, 53% reported symptoms of nausea, vomiting, or diarrhea. Secondary transmission appears to have occurred.

Most visitors left the camp on December 22 or 23 aboard charter buses with final destinations in Arkansas, Colorado, Mississippi, Missouri, Nebraska, and Texas. Explosive diarrhea and vomiting aboard the buses caused some groups to seek medical attention in hospital emergency rooms along the way. One group from Jackson, Mississippi, stopped in Dallas, Texas, where 60 members were seen in a single emergency room within several hours. A Beaumont, Texas, group stopped in a Denver, Colorado, hospital with approximately half its members ill with gastroenteritis. Six of the emergency room nurses caring for this group developed similar symptoms within 24 hours.

The only complete data gathered to date have been obtained from a questionnaire survey of camp personnel to which over 90% responded. The attack rate among them was 51%, with a sharp peak in the number of cases on December 23 (Figure 3). No significant differences were found between males and females. Meals consumed in the 3-camp dining rooms, serviced by a central kitchen, could not be implicated. The most common symptoms were vomiting (77%) and diarrhea (66%). Nausea without vomiting occurred in 14% (Table 6). There was no mortality. The secondary attack rate among family members of camp staff appeared to be greater than 25%. Numerous stool specimens were negative for common bacterial enteric pathogens.

Because of the widespread nature of the epidemic in the 2,500-acre camp, the occurrence of most cases over a 48-hour period, and the lack of correlation with food consumption, waterborne disease was considered. The camp is supplied with water by a natural spring in a meadow at low elevation. Water is pumped from the spring upward to the camp, and finally to a reservoir which is at still higher elevation. During heavy usage periods, the reservoir is capable of supplying water to 30 cabins by gravity. The pump house over the spring is located at the base of a small hill on top of which is located a private cabin with an attached septic tank, installed in 1959. Interviews with maintenance personnel revealed that on December 22 they discovered malfunctioning of the chlorinator and subsequently turned it off for several hours while making repairs.

On January 6 a survey of 100 guests at the camp revealed an incidence of gastroenteritis of 14% over the preceding 4-day period. Fluorescein dye flushed into the cabin sewage system rapidly appeared in the spring and in the camp tap water. The septic tank, covered by 2 feet of soil and set in fractured shale and decomposed granite, was subsequently unearthed, and a 3" x 4" hole was found in the leaching pipe several feet from its exit from the tank and directly above the pump house, at a distance of about 50 feet.

On the next day it was recommended that the camp's main water system (derived from the spring) be shut off and an auxiliary well chlorinated to provide potable water to the core buildings. All of the outlying cabins were closed. The septic tank was removed and daily monitoring of coliform count and chlorine residual was instituted.

The investigation is continuing to characterize the disease among visitors and to determine the extent of secondary transmission. Viral laboratory studies are also pending.

Editorial Note: Investigation of waterborne outbreaks of gastroenteritis often does not reveal an etiologic agent. From 1961 through 1972, gastroenteritis unassociated with known pathogens accounted for 45% of 49 municipal waterborne outbreaks investigated by CDC. The 1968 outbreak of gastroenteritis in Norwalk, Ohio, was theorized on epidemiologic grounds to be waterborne (1). In 1971 the causative agent, a



parvovirus, was identified by electronmicroscopy after transmission to volunteers. However, waterborne viral gastroenteritis has not been documented by recovery of virus from primary cases or from water.

Reference

記書がないと言語には、

ういたのにいたのように見たいのであるのであるのであるという

シートのようななののないのない

1. Adler JL, Zickl R: Winter vomiting disease. J Infect Dis 119:668-673, 1969

Fig. 3 PROBABLE VIRAL GASTROENTERITIS, COLORADO, DECEMBER 1976



Table 6

Clinical Symptoms in 36 Camp Staff with Gastroenteritis

Symptoms	<u>% I11</u>
Vomiting	77
Diarrhea	66
Muscle Aches	49
Headache	43
Dizziness	40
Abdominal Cramps	37
Fever	34
Chills	31
Nausea Without Vomiting	14
Bloody Diarrhea	0

IV. OUTBREAKS ON CRUISE SHIPS AND AIRCRAFT

This report summarizes data on outbreaks of gastrointestinal illness on cruise ships or aircraft that were reported to CDC in 1976.

A. <u>Definition</u> of <u>Outbreak</u>

Diarrheal illness on passenger vessels (vessels with 13 or more passengers) are reported by the Quarantine Stations to the Enteric Diseases Branch if 1) three percent or more of passengers or crew are ill; 2) one or more passengers or crew members is ill and the vessel has been in a cholera-infected area within the previous 5 days; 3) there has been a death or hospitalization aboard the vessel in a person who had a diarrheal illness.

After such an incident is reported, the need for a full investigation is determined by the severity, timing, and magnitude of the problem. The outbreaks tabulated in this report (Table 7) are the incidents that have been fully investigated by CDC. These investigations usually included, questionnaire surveys of passengers and crew, detailed evaluation of sanitation, and laboratory analysis of food, water, environment, and patient specimens. The Quarantine Division evaluated 6 additional incidents with medical log reviews and environmental inspections only.

Table 7

Outbreaks of Gastrointestinal Illness on Cruise Ships, 1976

·			Length of Cruise	Number of	Percent of		
Vessel	Date	Port	(Days)	Passengers	Passengers Ill	Etiology	Vehicle
А	June	Miami	14	745	35%	Unknown	Unknown
В	September	Miami	14	582	56%	Unknown	Water

B. Analysis of Data

In 1976 diarrhea outbreaks were investigated on 2 cruises (Table 7) and 1 aircraft (Table 8). The shipboard outbreaks occurred on the same ship during 2 separate Caribbean cruises (June and September). The investigation of the September cruise demonstrated coliform bacteria (TNTC) in the potable water system. Sewage contamination of the bunkered potable water had occurred. Potable water samples taken both before and after passing the U.V. light purifying system were found contaminated with coliform bacteria. There was no direct explanation for the coliform bacteria in the potable water system immediately after it passed through the U.V. system. Two possible explanations for this contamination were: 1) undiscovered cross connections existed in the potable water lines that by-passed the U.V. system (unlikely), or 2) coliform bacteria survived passage through the U.V. system.

Table 8

Outbreaks of Gastrointestinal Illness Aboard Aircraft, 1976

Aircraft	Date	Point of Embarkation	Point of Dis- embarkation	Number of Passengers	Percent of Passengers Ill	Etiology	Vehicle
A	6-20	Rio de Janeiro, Brazil	New York City, New York	185	15	Staphylococcal enterotoxin type D	Chocolate eclairs

The 1 reported outbreak on an aircraft took place on an American carrier enroute from Rio de Janeiro, Brazil, to New York City. Chocolate eclairs, consumed aboard the flight, were found to be contaminated with type D staphylococcal enterotoxin; they were prepared in Rio de Janeiro, and had been left unrefrigerated for 10 hours before being placed aboard the aircraft. A diversionary stop in San Juan, Puerto Rico, was necessary to discharge ill passengers.

いたのであるというで、

The marked decline in cruise vessel diarrheal outbreaks (8 in 1975) may be attributed to the cruise vessel sanitary inspection program which has been rigorously administered since 1974. All vessels with a home port in the United States receive a semiannual inspection. Vessels failing to meet the U.S. Public Health Service Standards are reinspected frequently until standards are achieved. Vessels meeting the standards have unscheduled spot inspections between semiannual inspections to insure that high sanitary standards are maintained.

V. REFERENCES

GENERAL

1. Foodborne Infections and Intoxications, Riemann H (ed). Academic Press, NY, 1969

2. Food Research Institute: Annual Report for 1974, University of Wisconsin-Madison, Wisconsin

3. Bryan FL: Emerging foodborne diseases. I. Their surveillance and epidemiology. II. Factors that contribute to outbreaks and their control. J Milk Food Technol 35:618-625, 632-638, 1972

4. Craun GF, McCabe LJ: Review of the causes of waterborne disease outbreaks. J Am Water Work Assoc 65:74-84, 1973

Ţ

BACTERIAL

Bacillus cereus

1. Goepfert JM, Spira WM, Kim HU: <u>Bacillus</u> cereus: Food poisoning organism. A review. J Milk Food Technol 35:213-227, 1972

2. Mortimer PR, McCann G: Food poisoning episodes associated with <u>Bacillus</u> <u>cereus</u> in fried rice. Lancet 1:1043-1045, 1974

Brucella

1. Spink WW: The Nature of Brucellosis. Minneapolis, Lund Press, Inc., 1956

2. Buchanan TM, et al: Brucellosis in the United States 1960-1972. Medicine 53: 403-439, 1974

3. Young EJ, Suvannaparrat U: Brucellosis outbreak attributed to ingestion of unpasteurized goat cheese. Arch Intern Med 135:240-243, 1975

Clostridium botulinum

1. Center for Disease Control: Botulism in the United States, 1899-1973. Handbook for Epidemiologists, Clinicians, and Laboratory Workers, CDC, Atlanta, June 1974, pp 7-11

2. Cherington M: Botulism. Ten-year experience. Arch Neurol 30:432-437, 1974

3. Koenig MG, Drutz DJ, Mushlin AI, et al: Type B botulism in man. Am J Med 42:208-219, 1967

4. Koenig MG, Spichard A, Cardella MA, et al: Clinical and laboratory observations of type E botulism in man. Medicine 43:517-545, 1964

5. Dowell VR Jr, McCroskey LM, Hatheway CL, et al: Coproexamination for botulinal toxin and <u>Clostridium botulinum</u> - A new procedure for laboratory diagnosis of botulism. JAMA 238:1829-1832, 1977

Clostridium perfringens

1. Bryan FL: What the sanitarian should know about <u>Clostridium</u> foodborne illness. J Milk Food Technol 32:381-389, 1969

2. Lowenstein MS: Epidemiology of <u>Clostridium perfringens</u> food poisoning. N Engl J Med 286(19):1026-1027, 1972

Escherichia coli

1. Marier R, Wells JG, Swanson RC, Callahan W, Mehlman IJ: An outbreak of enteropathogenic <u>Escherichia</u> <u>coli</u> foodborne disease traced to imported French cheese. Lancet 2:1376-1378, 1973

2. Sack RB: Human diarrheal disease caused by enterotoxigenic <u>Escherichia</u> <u>coli</u>. Ann Rev Microbiol 29:333-353, 1975

3. Rosenberg ML, Koplan JP, Wachsmuth IK: Epidemic diarrhea at Crater Lake from enterotoxigenic <u>Escherichia coli</u>. Ann Intern Med 86:714-718, 1977

Salmonella

1. Aserkoff B, Schroeder SA, Brachman PS: Salmonellosis in the United States--A five-year review. Am J Epidemiol 92:13-24, 1970

2. Bryan FL: What the sanitarian should know about salmonellae and

staphylococci in non-dairy foods. II. Salmonellae. J Milk Food Technol 31:131-140, 1968

3. Cohen ML, Blake PA: Trends in foodborne salmonellosis outbreaks. J Food Protection (in press), 1977

Shigella

1. Donadio JA, Gangarosa EJ: Foodborne shigellosis. J Infect Dis 119:666-668, 1969

Staphylococcus

1. Bryan FL: What the sanitarian should know about salmonellae and staphylococci in non-dairy foods. I. Staphylococci. J Milk Food Technol 31:110-116, 1968

2. Merson MH: The epidemiology of staphylococcal foodborne disease. Proceedings of the Staphylococci in Foods Conference, Pennsylvania State University, University Park, Pennsylvania, 1973, pp 20-37

3. Minor TE, Marth EH: <u>Staphylococcus aureus</u> and staphylococcal food poisoning. J Milk Food Technol 34:21-29, 77-83, 227-241, 1972, 35:447-476, 1973

Group A Streptococcus

1. Hill HR, Zimmerman RA, Reid GVK, Wilson E, Kitton RM: Foodborne epidemic of streptococcal pharyngitis at the United States Air Force Academy. N Engl J Med 280:917-921, 1969

2. McCormick JB, Kay D, Hayes M, Feldman RA: Epidemic streptococcal sore throat following a community picnic. JAMA 236:1039-1041, 1976

Vibrio cholerae

1. Finkelstein RA: Cholera. CRC Critical Reviews in Microbiology 2(4):553-623, 1973

2. Gangarosa EJ, Mosley WH: Epidemiology and surveillance of cholera: Cholera, edited by Barua D, Burrows W. Philadelphia, London, Toranto, WB Saunders Co., 1974, p 381

3. Blake PA, et al: Cholera in Portugal, 1974. I. Modes of transmission. II. Transmission by bottled mineral water. Am J Epidemiol 105:337-348, 1977

Vibrio parahaemolyticus

1. International Symposium on <u>Vibrio parahaemolyticus</u>, September 17-18, 1973, Fujino, Sakaguchi G, Sakazaki R, Takeday, (ed). Saikon Publishing Co., Ltd., Tokyo, Japan, 1974

2. Barker WH: <u>Vibrio parahaemolyticus</u> outbreaks in the United States. Lancet 1:551-554, 1974

3. Dadisman TA Jr, Nelson R, Molenda JR, et al: <u>Vibrio parahaemolyticus</u> gastroenteritis in Maryland. I. Clinical and epidemiologic aspects. Am J Epidemiol 96:414-426, 1972

CHEMICAL

Heavy Metal

Cadmium

1. Baker TD, Hafner WG: Cadmium poisoning from a refrigerator shelf used as an improvised barbecue grill. Public Health Rep 76:543-544, 1961

Copper

1. Hopper SH, Adams HS: Copper poisoning from vending machines. Public Health Rep 73:910-914, 1958

2. Semple AB, Parry WH, Phillips DE: Acute copper poisoning: An outbreak traced to contaminated water from a corroded geyser. Lancet 2:700-701, 1960

Tin

1. Barker WH, Runte V: Tomato juice-associated gastroenteritis. Washington and Oregon, 1969. Am J Epidemiol 96:219-226, 1972

Zinc

1. Brown MA, Thom JV, Orth GL, et al: Food poisoning involving zinc contamination. Arch Environ Health 8:657-660, 1964

Ichthyosarcotoxin

1. Hughes JM, Merson MH: Fish and shellfish poisoning. N Engl J Med 295:1117-1120, 1976

Ciguatoxin

1. Barkin RM: Ciguatera poisoning: A common-source outbreak. South Med J 67 (1):13-16, 1974

2. Halstead BW, Courville DA: Poisonous and venomous marine animals of the world. Vol 2 - Vertebrates. Washington, GPO, 1967, pp 63-330

Puffer Fish (tetrodotoxin)

1. Halstead BW, Courville DA: Poisonous and venomous marine animals of the world. Vol 2 - Vertebrates. Washington, GPO, 1967, pp 679-844

2. Torda TA, Sinclair E, Ulyatt DB: Puffer fish (tetrodotoxin) poisoning: Clinical record and suggested management. Med J Aust 1:599-602, 1973

Scombrotoxin

1. Halstead BW, Courville DA: Poisonous and venomous marine animals of the world. Vol 2 - Vertebrates. Washington, GPO, 1967, pp 639-668

2. Kimata M: The histamine problem: Fish as Food, edited by Borgstrom E, New York, Academic Press, 1961, pp 329-352

3. Merson MH, Baine WB, Gangarosa EJ, Swanson RC: Scombroid fish poisoning: Outbreak traced to commercially canned tuna fish. JAMA 228:1268-1269, 1974

Monosodium Glutamate

1. Schaumburg HH, Byck R, Gerstl R, Mashman JH: Monosodium L-glutamate: Its pharmacology and role in the Chinese restaurant syndrome. Science 163:826-828, 1969

Mushroom Poison

1. Wieland T, Wieland O: The toxic peptides of <u>Amanita</u> species. Vol 8 - Fungal Toxins: Microbial Toxins, edited by Kadis S, Ciegler A, Ajl SJ, New York and London, Academic Press, 1972, pp 249-280

2. Benedict RG: Mushroom toxins other than Amanita. Vol 8 - Fungal Toxins. In Microbial Toxins, edited by Kadis S, Ciegler A, Ajl SJ, New York and London, Academic Press, 1972, pp 281-320

3. Tyler VE: Poisonous mushrooms: Progress in Chemical Toxicology. Vol 1, edited by Stolman A, New York, Academic Press, 1963, pp 339-384

Paralytic and Neurotoxic Shellfish Poison

1. Music SI, Howell JT, Brumback CL: Red tide: Its public health implications. J Fla Med Assoc 60:27-29, 1973

2. Halstead BW, Courville DA: Poisonous and venemous marine animals. Vol 1 - Invertebrates. Washington, GPO, 1965, pp 157-240

PARASITIC

Anisakidae

1. Chitwood MD: Nematodes of medical significance found in market fish. Am J Trop Med Hyg 19:599-602, 1970

T. spiralis

1. Gould SE: Trichinosis in man and animals. Springfield, Ill, Charles C. Thomas, 1970

2. Zimmerman WJ, Steele JH, Kagan IG: Trichinosis in the U.S. population 1966-1970--prevalence and epidemiologic factors. Health Services Rep 88:606-623, 1973

G. lamblia

1. Petersen H: <u>Giardiasis</u> (lambliasis). Scand J Gastroenterol 7 (Suppl 14): 1-44, 1972

2. Schultz MG: Giardiasis. JAMA 233:1383-1384, 1975

3. Wolfe MS: Giardiasis. JAMA 233:1362-1365, 1975

T. gondii

1. Kean BH, Kimball AC, Christensen WN: An epidemic of acute toxoplasmosis. JAMA 208:1002-1004, 1969

VIRAL

Hepatitis A

1. Cliver DO: Implications of foodborne infectious hepatitis. Public Health Rep 81:159-165, 1966

2. Gravelle CR, Hornbeck CL, Maynard JE, et al: Hepatitis A: Report of a common-source outbreak with recovery of a possible etiologic agent. II. Laboratory studies. J Infect Dis 131:167-171, 1975

3. Leger RT, Boyer KM, Pattison CP, et al: Hepatitis A: Report of a commonsource outbreak with recovery of a possible etiologic agent. I. Epidemiologic studies. J Infect Dis 131:163, 1975

4. Denes AE, Smith JL, Hindman SH, et al: Foodborne hepatitis A infection: A report of two urban restaurant-associated outbreaks. Am J Epidemiol 105:156-162, 1977 VI. ARTICLES ON FOODBORNE AND WATERBORNE DISEASE OUTBREAKS, 1976, TAKEN FROM MORBIDITY AND MORTALITY WEEKLY REPORT

BACTERIAL

Clostridium botulinum

Botulism-Alaska 25(49):399
Botulism-Washington 25(19):150
Botulism associated with commercial cherry peppers-Oklahoma,
Utah, Texas 25(17):134
Follow-up on botulism associated with commercial cherry
peppers 25(18):148

Clostridium perfringens

Clostridium perfringens-Wisconsin 25(25):198

Salmonella

Outbreak of milk-borne <u>Salmonella</u> gastroenteritis-South Australia 26(15):127 Salmonellosis associated with homemade ice cream-Michigan 26(12):94 <u>Salmonella bovis-morbificans</u> in pre-cooked roasts of beef 25(42):333 Foodborne <u>Salmonella</u> infections contracted on aircraft 25(41):332 <u>Salmonella london-Minnesota 25(35):283</u> Salmonellosis caused by cheddar cheese 25(32):259 <u>Salmonella infantis</u>-California, Colorado 26(6):41 Follow-up on Salmonella infantis-U.S. 26(10):84

Shigella 3 8 1

Outbreak of shigellosis-Fort Bliss, Texas 26(13):107 Shigella flexneri type 2 foodborne outbreak-Washington 25(38)302

Staphylococcus

Staphylococcal food poisoning-Colorado 26(4):22 Outbreak of staphylococcal food poisoning aboard an aircraft 25(40):317

Yersinia

Yersinia enterocolitica outbreak-New York 26(7):53

CHEMICAL

Ciguatera

Ciguatera fish poisoning-Midway Atoll 25(27):219

Chlordane

Chlordane contamination of a municipal water system-Tennessee 25(15):117

Paralytic Shellfish Poisoning

Paralytic shellfish poisoning-Alaska 25(47):383 Paralytic shellfish poisoning-New Brunswick, Canada 25(43):347

Sodium Nitrite

Acute nitrite poisoning-California 25(35):278

WATERBORNE DISEASE

Giardiasis-California, Colorado 26(7):60 Probable viral gastroenteritis-Colorado 26(3):13 Gastrointestinal illness aboard a cruise ship 25(39):309 Diarrhea in bicyclers-Idaho and Montana 25(31):251



STATE EPIDEMIOLOGISTS AND STATE LABORATORY DIRECTORS

The State Epidemiologists are the key to all disease surveillance activities. They are responsible for collecting, interpreting, and transmitting data and epidemiologic information from their individual states. Their contributions to this report are gratefully acknowledged. In addition, valuable contributions are made by State Laboratory Directors; we are indebted to them for their valuable support.

STATE Alabama Alaska Arizona Arkansas California Colorado Connecticut Delaware District of Columbia Florida Georgia Hawaii Idaho Illinois Indiana Iowa Kansas Kentucky Louisiana Maine Maryland Massachusetts Michigan Minnesota Mississippi Missouri Montana Nebraska Nevada New Hampshire New Jersey New Mexico New York State New York City North Carolina North Dakota Ohio 0klahoma Oregon Pennsylvania Puerto Rico Rhode Island South Carolina South Dakota Tennessee Texas Utah Vermont Virginia Washington West Virginia Wisconsin Wyoming *Dual assignment

STATE EPIDEMIOLOGIST Frederick S Wolf, MD John Starr, MD Alexander Kelter, MD Paul C White, Jr, MD James Chin, MD Thomas M Vernon, Jr, MD John N Lewis, MD Ernest S Tierkel, VMD Martin E Levy, MD R Michael Yeller, MD John E McCroan, PhD Ned H Wiebenga, MD John A Mather, MD Byron J Francis, MD Richard D Telle, MD Laverne A Wintermeyer, MD Donald E Wilcox, MD Calixto Hernandez, MD Charles T Caraway, DVM William S Nersesian, MD Kathleen H Acree, MDCM Nicholas J Fiumara, MD Norman S Hayner, MD Ellen Z Fifer, MD Durward L Blakey, MD H Denny Donnell, Jr, MD Martin D Skinner, MD Paul A Stoesz, MD William M Edwards, MD Vladas Kaupas, MD Ronald Altman, MD Jonathan M Mann, MD Donald O Lyman, MD John S Marr, MD Martin P Hines, DVM Kenneth Mosser Thomas J Halpin, MD Patrick M Morgan, DVM, DrPH John A Googins, MD William E Parkin, DVM Henry Negron, MD Gerald A Faich, MD Richard L Parker, DVM James D Corning, BA, Acting Robert H Hutcheson, Jr, MD Charles R Webb, Jr, MD Taira Fukushima, MD Richard L Vogt, MD, Acting Grayson B Miller, Jr, MD Jack Allard, PhD* William L Cooke, MD H Grant Skinner, MD Herman S Parish, MD

STATE LABORATORY DIRECTOR Thomas S Hosty, PhD Frank P Pauls, DrPH Jon M Counts, DrPH Robert T Howell, DrPH John M Heslep, PhD C D McGuire, PhD John J Redys, BS Mahadeo P Verma, PhD Alston Shields, DrPH Nathan J Schneider, PhD Earl E Long, MS Albert I Oda D W Brock, DrPH M Louise Brown Josephine Van Fleet, MD W J Hausler, Jr, PhD Dwayne C Morse, DrPH B F Brown, MD Henry Bradford, PhD Howard E Lind, PhD J Mehsen Joseph, PhD George F Grady, MD George R Anderson DVM David Štickle, DrPH, Acting R H Andrews, MS Elmer Sputrier, DrPH David B Lackman, PhD Henry McConnell, DrPH Paul Fugazzotto, PhD Robert A Miliner, DrPH John J Nelson, MPH Aaron Bond, MPH David Axelrod, MD Bernard Davidow, PhD Mildred A Kerbaugh C Patton Steele, BS Charles C Croft, ScD William R Schmieding, P William Murphey, PhD Vern Pidcoe, DrPH Jose L Villamil Raymond G Lundgren, PhD Arthur F DiSalvo, MD A Richard Melton, DrPH M Sam Sudman, DrPH Charles Sweet, DrPh James O Mason, MD Dymitry Pomer, DVM Frank W Lambert, PhD Jack Allard, PhD* John W Brough, DrPH S L Inhorn, MD Donald T Lee, DrPH 9/29/77
U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE PUBLIC HEALTH SERVICE CENTER FOR DISEASÉ CONTROL ATLANTA, GEORGIA 30333

1

OFFICIAL BUŞINESS



POSTAGE AND FEES PAID U.S. DEPARTMENT OF HEW HEW 399

Ċ,

HEW Publication No. (CDC) 78-8185