





October 21, 2011

Suicidal Thoughts and Behaviors Among Adults Aged ≥18 Years — United States, 2008–2009











CONTENTS

ntroduction	2
Methods	
Results	
Discussion	
Conclusion	8
References	9

The MMWR series of publications is published by the Office of Surveillance, Epidemiology, and Laboratory Services, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

Suggested Citation: Centers for Disease Control and Prevention. [Title]. MMWR 2011;60(No. SS-#):[inclusive page numbers].

Centers for Disease Control and Prevention

Thomas R. Frieden, MD, MPH, Director
Harold W. Jaffe, MD, MA, Associate Director for Science
James W. Stephens, PhD, Director, Office of Science Quality
Stephen B. Thacker, MD, MSc, Deputy Director for Surveillance, Epidemiology, and Laboratory Services
Stephanie Zaza, MD, MPH, Director, Epidemiology and Analysis Program Office

MMWR Editorial and Production Staff

Ronald L. Moolenaar, MD, MPH, Editor, MMWR Series Christine G. Casey, MD, Deputy Editor, MMWR Series Teresa F. Rutledge, Managing Editor, MMWR Series David C. Johnson, Lead Technical Writer-Editor Jeffrey D. Sokolow, MA, Project Editor Martha F. Boyd, *Lead Visual Information Specialist*Maureen A. Leahy, Julia C. Martinroe, Stephen R. Spriggs, Terraye M. Starr *Visual Information Specialists*Quang M. Doan, MBA, Phyllis H. King *Information Technology Specialists*

MMWR Editorial Board

William L. Roper, MD, MPH, Chapel Hill, NC, Chairman

Virginia A. Caine, MD, Indianapolis, IN
Jonathan E. Fielding, MD, MPH, MBA, Los Angeles, CA
David W. Fleming, MD, Seattle, WA
William E. Halperin, MD, DrPH, MPH, Newark, NJ
King K. Holmes, MD, PhD, Seattle, WA
Deborah Holtzman, PhD, Atlanta, GA
John K. Iglehart, Bethesda, MD
Dennis G. Maki, MD, Madison, WI

Patricia Quinlisk, MD, MPH, Des Moines, IA
Patrick L. Remington, MD, MPH, Madison, WI
Barbara K. Rimer, DrPH, Chapel Hill, NC
John V. Rullan, MD, MPH, San Juan, PR
William Schaffner, MD, Nashville, TN
Anne Schuchat, MD, Atlanta, GA
Dixie E. Snider, MD, MPH, Atlanta, GA
John W. Ward, MD, Atlanta, GA

Suicidal Thoughts and Behaviors Among Adults Aged ≥18 Years — United States, 2008–2009

Alex E. Crosby, MD¹
Beth Han, MD, PhD²
LaVonne A. G. Ortega, MD¹
Sharyn E. Parks, PhD¹
Joseph Gfroerer, BA²

¹Division of Violence Prevention, National Center for Injury Prevention and Control, CDC ²Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Rockville, Maryland

Abstract

Problem/Condition: Suicidal thoughts and behaviors are important public health concerns in the United States. In 2008, a total of 36,035 persons died as a result of suicide, and approximately 666,000 persons visited hospital emergency departments for nonfatal, self-inflicted injuries. State-level data on suicide-related issues are needed to help establish program priorities and to evaluate the effectiveness of suicide prevention strategies. Public health surveillance with timely and consistent exchange of data between data collectors and prevention program implementers allows prevention program practitioners to implement effective prevention and control activities.

Reporting Period: January 1, 2008–December 31, 2009.

Description of System: The National Survey on Drug Use and Health (NSDUH) is a national- and state-level survey of a representative sample of the civilian, noninstitutionalized U.S. population aged ≥12 years. NSDUH collects data on health-risks related to the use of illicit drugs, alcohol, and tobacco; initiation of substance use; substance use disorders and treatment; health care; and mental health. This report summarizes data on responses to questions concerning suicidal thoughts and behaviors contained in the mental health section among sampled persons aged ≥18 years in all 50 states and the District of Columbia. This report analyzes data on the prevalence of suicidal thoughts, planning, and attempts by age, sex, race/ethnicity, and state from 92,264 respondents in the 2008 and 2009 NSDUH.

Results: Prevalence estimates of suicidal thoughts and behaviors varied by sociodemographic factors, region, and state. During 2008–2009, an estimated 8.3 million (annual average) adults aged ≥18 years in the United States (3.7% of the adult U.S. population) reported having suicidal thoughts in the past year. The prevalence of having suicidal thoughts ranged from 2.1% in Georgia to 6.8% in Utah. An estimated 2.2 million (annual average) adults in the United States (1.0% of the adult U.S. population) reported having made suicide plans in the past year. The prevalence of reports of suicide planning ranged from 0.1% in Georgia to 2.8% in Rhode Island. An estimated 1 million (annual average) adults in the United States (0.5% of the U.S. adult population) reported making a suicide attempt in the past year. The prevalence of reports of suicide attempts ranged from 0.1% in Delaware and Georgia to 1.5% in Rhode Island. The prevalence of suicidal thoughts, suicide planning, and suicide attempts was significantly higher among young adults aged 18–29 years than it was among adults aged ≥30 years. The prevalence of suicidal thoughts was significantly higher among females than it was among males, but there was no statistically significant difference for suicide planning or suicide attempts.

Interpretation: The findings in this report indicate that substantial variations exist at the regional and state level in the prevalence of adults who had suicidal thoughts, made plans to attempt suicide, and attempted suicide in the past year. Geographic differences in prevalence might be attributable to selective migration, sociodemographic composition of the population, or the local social environment (e.g., social relationship indicators such as divorce rates or resources for access to health care). These findings emphasize the importance of continued surveillance to collect locally relevant data on which to base prevention and control activities.

Public Health Action: A better understanding of the patterns of the precursors to suicide is crucial to planning and evaluating a broad spectrum of suicide prevention efforts. These results can be used by state health departments and federal agencies to measure

Corresponding author: Alex E. Crosby, MD, Division of Violence Prevention, National Center for Injury Prevention and Control, CDC, 4770 Buford Hwy, N.E., MS: F-63, Atlanta, Georgia, 30341. Telephone: 770-488-4410, Fax: 770-488-4222, E-mail: aec1@cdc.gov.


progress toward achieving national and state health objectives (e.g., those outlined in the National Strategy for Suicide Prevention). Continued surveillance is needed to design, implement, and evaluate public health policies and programs that can lead to a reduction in morbidity and mortality related to suicide-related

thoughts and behaviors. Possible strategies to implement could include universal strategies (e.g., public education campaigns that focus on improving recognition of suicide risk) and indicated strategies (e.g., cognitive-behavioral therapy) that address the needs of persons exhibiting certain risk factors (e.g., persons who have made suicide attempts).

Introduction

Self-directed violence encompasses a range of behaviors, ranging from nonsuicidal intentional self-harm (i.e., behaviors in which the intention is not to kill oneself, as in self-mutilation) to acts of fatal and nonfatal suicidal behavior (1). Injury from self-directed violence is a major public health issue throughout the United States and worldwide (2-5). In 2008, suicide was the 10th leading cause of death in the United States, resulting in 36,035 deaths (6). Suicide is a problem throughout the life span; it is the fourth leading cause of death among children aged 10-14 years, the third among persons aged 15–24 years, the fourth among persons aged 25–44 years, and the eighth among those aged 45-64 years (6). Suicides reflect only a portion of the number of persons affected by suicidal thoughts and behaviors (Figure 1). Substantially more persons are hospitalized as a result of nonfatal suicidal behaviors than are fatally injured, and an even greater number are either treated in ambulatory settings or not treated at all (7,8). During 2007-2008, an estimated average of 569,000 persons visited U.S. hospital emergency departments annually for self-directed violence, 70%, of whom had attempted suicide (9-11). According to at least one nationally representative report, 56.8% of persons who engage in suicidal behavior never seek

FIGURE 1. Public health burden of suicidal behavior among adults aged ≥18 years — United States, 2008



- * Source: CDC's National Vital Statistics System.
- [†] All rates per 100,000 population; population estimates provided by U.S. Census Bureau.
- § **Source:** Agency for Healthcare Research and Quality's Healthcare Cost and Utilization Project–Nationwide Inpatient Sample.
- Source: CDC's National Electronic Injury Surveillance System–All Injury Program.

health services (12). Consequently, prevalence figures based on health-care encounters substantially underestimate the societal burden of suicidal behavior. For this reason, a broad array of assessment tools is needed to gain a better understanding of the magnitude of the problem.

Multiple assessment tools are used to understand the extent of suicide thoughts and behaviors collect data on deaths and nonfatal injuries. The National Vital Statistics System (NVSS) compiles demographic and causal data on all death certificates filed in the United States. NVSS final mortality data include all fatalities in the United States, but these data provide limited information about the exact nature and circumstances of the injury (13). The National Hospital Discharge Survey (NHDS) and the Health Care and Utilization Project (HCUP) measure inpatient care and hospital utilization (13). Although they provide detailed information about the nature of injuries, treatment, and discharge disposition, both HCUP and NHDS are limited because not all states require external cause-of-injury coding, which identifies the intent of the injury, on hospital discharge records. In 2004, the State and Territorial Injury Prevention Directors' Association (now named the Safe States Alliance) reported that only 43.8% of the medical records for which an injury was the principal diagnosis had an accompanying E-code (14). The National Hospital Ambulatory Medical Care Survey (NHAMCS) and the National Electronic Injury Surveillance System-All Injury Program (NEISS-AIP) collect data about all types and external causes of nonfatal injuries and poisonings treated in either U.S. hospital emergency departments (EDs) or hospital outpatient departments. However, although these systems are nationally representative, they do not have state-level samples (15). NSDUH data complement data from these other systems. Community surveys such as NSDUH can obtain data on injuries that do not result in death or severe injury, occur outside the hospital environment, or occur in settings for which vital statistics or hospital-based data are unavailable (16,17).

Population surveys capture data that can convey a different understanding of the problem than information based on health-care system encounters. In general, persons self-report occurrence of disease to a different extent than clinicians diagnose; similarly, they report risk or protective factors on surveys to a different degree than they reveal to their health-care providers (18). Multiple factors account for these differences. Some persons do not seek or do not have access to health care, so events involving these persons never appear in health

records. In addition, persons tend to report fewer health problems to a health-care provider than they actually have (19,20). Anonymous population surveys can be perceived to be less intimidating to a respondent than a potential authority figure such as a health-care provider (21) and thus can provide data that complement other sources and that can be more comprehensive for policymaking and for making comparisons between population groups. A recent review of assessments examining primarily population-based surveys of suicidal thoughts and behavior among adults aged ≥18 years in the 12 months preceding the assessment indicated that 2.1%-10% of respondents reported having suicidal thoughts and 0.2%-2% reported making suicide attempts (22); these findings indicate that population surveys can provide researchers with a different perspective on the prevalence of suicidal thoughts and behaviors than those provided by death certificates or medical records.

Population surveys must be made relevant to the populations that they monitor. National estimates of the prevalence of suicidal thoughts and behaviors can be used to draw attention to the magnitude of the problem and are useful for establishing national health priorities. Nationally representative data also can be used to examine differences in rates among groups (e.g., sex, racial/ethnic, and age groups) and geographic regions. They are also useful in aggregating sufficient numbers of rare cases of a particular type of injury to identify patterns and mechanisms of injury. However, the national-level data alone convey only a partial picture of the public health burden of suicide-related events because there is substantial variation between areas. For example, rates of suicide have been demonstrated to vary more than threefold by geographic region (23).

The National Strategy for Suicide Prevention identified a need for data at the national, tribal, state, and local levels (24). Tribal, state, and local data can be used to help public health authorities establish local program priorities and evaluate the effectiveness of their suicide prevention strategies (13,25,26). The national public health surveillance capability requires a strong foundation of surveillance capacity at all levels of local, state, tribal and federal public health and the ability to rapidly, consistently, and securely exchange and share such data, detect events, and take appropriate public health actions. U.S. public health surveillance traditionally has been administered through state health departments. Because the implementation of prevention measures frequently involves interacting with persons, families, and community organizations (e.g., by providing education and performing training), most prevention programs are implemented at the tribal, state, and community levels (13). As problems and needs are identified, public support is garnered at the local level, and multiple organizations (including health-care providers,

schools, civic groups, athletic leagues, businesses, and public safety departments) frequently work together to address injury hazards and concerns.

The lack of state-level data on the extent of suicidal thoughts and behaviors limits the ability of program practitioners to act in a focused fashion where the problem is the most acute. A population-based survey can provide some of this needed information. This report is the first to present state-level data on suicidal thoughts, planning, and attempts among adults. Another reason it is important to gather data about nonfatal suicidal behavior is that persons who made suicide attempts are one of the highest risk groups for subsequent death from suicide (2). These state-level data can be used by state health departments and federal agencies to measure progress toward achieving national and state health objectives as part of the NSSP or state suicide prevention plans (24,27).

Methods

This report summarizes national estimates on suicidal thoughts and behaviors by age, sex, race/ethnicity, education, current employment, and county type (i.e., based on population size). In addition, this report presents corresponding state-specific estimates by age, sex, race/ethnicity. A total of 92,264 persons aged ≥18 years participated in the National Survey on Drug Use and Health (NSDUH) during 2008–2009 (28). NSDUH is the only ongoing survey that collects national- and state-level data on suicidal thoughts and behaviors among the civilian, noninstitutionalized U.S. population aged ≥18 years. NSDUH is conducted annually by the Substance Abuse and Mental Health Services Administration (SAMHSA); data sources have been described in detail previously (28).

Sample Design

In 2008 and 2009, NSDUH employed a state-based design with an independent, multistage area probability sample within each state and the District of Columbia. The total population within each state is based on population estimates provided by the U.S. Census Bureau. The eight states with the largest populations (California, Florida, Illinois, Michigan, New York, Ohio, Pennsylvania, and Texas), which together accounted for 48% of the total U.S. population, were designated as "large sample states" and had sample sizes ranging from 3,556 to 3,830 persons in 2008 and from 3,557 to 3,707 persons in 2009. For the remaining 42 states and the District of Columbia, the target sample sizes ranged from 876 to 981 persons in 2008 and from 886 to 984 persons in 2009. Youths and young adults aged 12–25 years were oversampled so that each state's sample was distributed approximately equally among three age

groups: persons aged 12–17 years, persons aged 18–25 years, and persons aged \geq 26 years. This design resulted in a sample size for the 2-year period of approximately 4,800 adults (2,740 persons aged 18–29 years and 2,060 aged \geq 30 years) in each of the eight largest states and 1,200 adults (685 persons aged 18–29 years and 515 persons aged \geq 30 years) in each of the other 42 states (29).

Questionnaire

In both 2008 and 2009, all adult respondents were asked whether they had thought seriously about trying to kill themselves at any time during the past 12 months. Those respondents who reported having had serious suicidal thoughts were then asked whether, in the past 12 months, they had made any plans to kill themselves and whether they had tried to kill themselves. Before 2008, these questions were asked only of persons who reported having had at least one major depressive episode during the past year. Survey questions are available at http://oas.samhsa.gov/nsduh/2k8MRB/2k8Q.pdf.

Data Collection and Processing

The NSDUH data collection method involved conducting in-person interviews with selected persons and incorporating procedures designed to increase respondents' cooperation and willingness to report honestly about sensitive issues. Confidentiality was stressed in all written and oral communications with potential respondents. Audiocomputer-assisted self-interviewing (ACASI) methods were used to provide respondents a private and confidential way to record answers. Interview data were compiled in a data file that consists of one record for each completed interview. Although some editing and consistency checks that are built into ACASI were performed during the interview, additional, more complex edits and consistency checks were applied after the interview was completed during data processing. Statistical imputation was used to replace missing or ambiguous values after editing for some key variables, but no imputation was done for suicide-related variables.

Weighting

The general approach to developing and calibrating analysis weights involved developing design-based weights as the product of the inverse of the selection probabilities at each selection stage. Adjustment factors then were applied to adjust for nonresponse, to poststratify to known population control totals, and to control for extreme weights when necessary. Because this report combines data for 2 years, person-level weights for estimates of annual averages were obtained by

dividing the analysis weights by two. Weighted response rates for household screening and for interviewing were 89.0% and 74.4%, respectively, for the 2008 survey and 88.8% and 75.6%, respectively, for the 2009 survey.

Statistical Analyses

National prevalence (population and percentage estimates, relative standard errors, and 95% confidence intervals [CIs]) were estimated for adults aged ≥18 years who had suicidal thoughts, made any suicide plans, or attempted suicide in the past year and were analyzed by selected sociodemographic characteristics. Chi-square tests were applied to test for statistical significance of the associations between suicidal thoughts and behaviors and sociodemographic characteristics at the bivariate level. A p value of <0.05 was considered statistically significant.

Then the prevalence was estimated (population and percentage estimates, relative standard errors, and 95% CIs) for adults aged ≥18 years who had suicidal thoughts in the past year by sociodemographic characteristics, overall for the United States, and by age, sex, race/ethnicity, region, county type, and state. Classification of race/ethnicity and geographic regions were based on U.S. Census Bureau recommendations (29). Definitions of county type are based on U.S. Department of Agriculture classifications: large metropolitan areas have a population of ≥1 million persons, small metropolitan areas have a population of <1 million persons, and nonmetropolitan areas are those areas that are outside of metropolitan statistical areas (30). Next, prevalence was assessed for adults who made any suicide plans in the past year by these characteristics. Finally, prevalence was estimated for adults who attempted suicide in the past year by these same factors. SUDAAN software (31) was used to account for the complex sample design and sampling weights.

Direct survey estimates produced for this report that were considered to be unreliable because of unacceptably large sampling errors are not reported. This occurred among certain racial/ethnic populations (e.g., American Indians/Alaska Natives). The suppression criteria are based on the relative standard error (RSE, defined as the ratio of the SE over the estimate), actual sample size, and effective sample size for each estimate. According to the suppression criteria, prevalence estimates are suppressed if any of the following occurred: 1) the prevalence estimate is <0.005% or >99.995%, 2) the relative standard error of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is ≤50%, 3) the relative standard error of the negative of the natural logarithm of (1-p) is >0.175 if the prevalence is >50%, 4) the actual sample size

is <100, or 5) the effective sample size (defined as the actual sample size divided by the design effect) is <68. The 2008 and 2009 NSDUH suppression rules have been published previously (29). The 95% CIs are asymmetric; asymmetric CIs are often more appropriate when dealing with small percentages (32). In this report, prevalence estimates are rounded to the nearest tenth, and therefore some numbers are recorded as <0.1%.


Results

Suicidal Thoughts and Behaviors

The prevalence rates of suicidal thoughts and behaviors are presented by sociodemographic characteristics (Table 1). During 2008–2009, an estimated 8.3 million (annual average) adults in the United States (3.7% of the adult U.S. population) reported having had suicidal thoughts in the past year. Adults aged 18–29 years were more likely than their older counterparts to have had suicidal thoughts, made any suicide plans, and attempted suicide in the past year. Females were more likely than males to have had suicidal thoughts in the past year but not more likely to have made suicide plans or attempted suicide. The racial/ethnic group with the highest prevalence for having suicidal thoughts and for making any suicide plans was non-Hispanic whites; the group with the lowest prevalence for suicidal thoughts and planning was non-Hispanic Asians. Non-Hispanic blacks were more likely than non-Hispanic Asians to report a suicide attempt in the past year. Adults with less than a high school education were more likely than college graduates to have had suicidal thoughts, made suicide plans, and attempted suicide in the past year. Unemployed adults were more likely than those with full-time employment to have had suicidal thoughts, made suicide plans, and attempted suicide in the past year. Adults in the Midwest and West were more likely to have had suicidal thoughts than those in the Northeast and South. Adults in the Midwest were more likely to have made suicide plans than those in the South. However, prevalence of suicide attempts did not vary by region. In addition, prevalence of past-year suicidal thoughts, plans, and attempts did not differ by the type of county in which adults resided (Table 1).

During 2008–2009, among adults aged ≥18 years, the estimated prevalence of suicidal thoughts ranged from 2.1% in Georgia to 6.8% in Utah (U.S. average: 3.7%)(Figure 2). The estimated prevalence of having made suicide plans ranged from 0.1% in Georgia to 2.8% in Rhode Island (U.S. average: 1.0%). The estimated prevalence of having made a suicide attempt ranged from 0.1% in Georgia and Delaware to 1.5% in Rhode Island (U.S. average: 0.5%) (Table 2).

FIGURE 2. Average annual prevalence of suicidal thoughts among adults aged ≥18 years, by state — National Survey on Drug Use and Health, United States, 2008–2009*



* U.S. average: 3.7.

Suicidal Thoughts

An estimated 2.9 million (annual average) persons aged 18–29 years in the United States had suicidal thoughts in the past year (5.7% of the age-group population). In this age group, prevalence ranged from 3.3% in North Carolina to 8.8% in Nevada. During 2008–2009, an estimated 5.5 million (annual average) persons aged ≥30 years in the United States had suicidal thoughts in the past year (3.1% of the population) (Table 3). In this age group, prevalence ranged from 1.6% in Georgia, Louisiana, and Mississippi to 6.5% in Rhode Island (Table 3).

During 2008–2009, an estimated 3.8 million (annual average) adult males in the United States had suicidal thoughts in the past year (3.5% of the adult male population). Among males, the prevalence ranged from 1.3% in Mississippi to 7.1% in Rhode Island. During 2008–2009, an estimated 4.6 million (annual average) female adults in the United States (3.9% of the adult female population) had suicidal thoughts in the past year (Table 4). Among adult females, prevalence ranged from 1.5% in Georgia to 9.1% in Nevada (Table 4).

During 2008–2009, an estimated 6.0 million (annual average) non-Hispanic white adults in the United States (3.9% of the non-Hispanic white population) had suicidal thoughts in the past year. Among non-Hispanic white adults, prevalence ranged from 1.7% in Hawaii to 7.5% in Utah. During that same period, an estimated 911,000 (annual average) non-Hispanic black adults in the United States (3.5% of the non-Hispanic black adult population) had suicidal thoughts in the past year. Among states with stable estimates for non-Hispanic black adults, prevalence ranged from 0.8% in Connecticut to 7.5% in Arkansas. During 2008–2009, an estimated 933,000

(annual average) Hispanic adults in the United States (3.0% of the Hispanic adult population) had suicidal thoughts in the past year. Among states with stable estimates for Hispanic adults, prevalence ranged from 0.6% in South Carolina to 8.6% in Wyoming. During that period, an estimated 208,000 (annual average) non-Hispanic Asian adults in the United States (2.1% of the adult non-Hispanic Asian population) had suicidal thoughts in the past year. Among states with stable estimates for non-Hispanic Asian adults, prevalence ranged from 1.2% in Florida to 3.8% in Michigan (Table 5).

Suicide Plans

During 2008–2009, an estimated 2.2 million (annual average) adults in the United States (1.0% of the U.S. adult population) reported making suicide plans in the past year (Table 1). An estimated 821,000 (annual average) persons aged 18–29 years in the United States (1.6% of the age group population) made suicide plans in the past year. In this age group, prevalence ranged from 0.4% in Georgia to 3.5% in Arkansas. During 2008–2009, an estimated 1.4 million (annual average) persons aged ≥30 years in the United States (0.8% of the age-group population) made suicide plans in the past year. In this age group, prevalence ranged from 0.1% in Mississippi to 2.8% in Rhode Island (Table 6).

During 2008–2009, an estimated 1.0 million (annual average) adult males in the United States (1.0% of the adult male population) made suicide plans in the past year. Among males, prevalence ranged from 0.1% in Georgia to 4.1% in Rhode Island. During 2008–2009, an estimated 1.2 million (annual average) adult females in the United States (1.0% of the female population) made suicide plans in the past year. Among females, prevalence ranged from 0.1% in Georgia to 3.1% in Nevada (Table 7).

During 2008-2009, an estimated 1.6 million (annual average) non-Hispanic white adults in the United States (1.0% of the adult non-Hispanic white population) made any suicide plans in the past year. Among non-Hispanic white adults, prevalence ranged from 0.1% in Georgia to 3.3% in Nevada. During that same period, an estimated 262,000 (annual average) non-Hispanic black adults in the United States (1.0% of the non-Hispanic black adult population) made suicide plans in the past year. Among states with stable estimates for non-Hispanic black adults, prevalence ranged from 0.2% in Delaware to 3.5% in Arkansas. During 2008-2009, an estimated 267,000 (annual average) Hispanic adults in the United States (0.9% of the adult Hispanic population) made suicide plans in the past year. Among states with stable estimates for Hispanic adults, prevalence ranged from <0.1% in District of Columbia to 5.0% in Connecticut. During that period, an estimated 37,000 (annual average) non-Hispanic Asian adults in the United States (0.4% of the non-Hispanic Asian adult population) made suicide plans in the past year. Among states with stable estimates for non-Hispanic Asian adults, prevalence ranged from 0.1% in Illinois to 0.5% in California (Table 8).

Suicide Attempts

During 2008–2009, approximately 1.0 million (annual average) adults in the United States (0.5% of the adult U.S. population) reported making a suicide attempt in the past year (Table 2; Figure 2) An estimated 477,000 (annual average) persons aged 18–29 years in the United States (1.0% of the agegroup population) attempted suicide in the past year. In this age group, prevalence ranged from 0.1% in Iowa to 2.4% in Arkansas. During 2008–2009, an estimated 581,000 (annual average) persons aged ≥30 years in the United States (0.3% of the age-group population) attempted suicide in the past year. In this age group, prevalence ranged from 0.1% in Hawaii, Kentucky, Montana, New Hampshire, and North Dakota to 1.3% in Rhode Island (Table 9).

During 2008–2009, an estimated 442,000 (annual average) adult males in the United States (0.4% of the adult male population) attempted suicide in the past year. Among males, prevalence ranged from <0.1% in Alaska, the District of Columbia, and Georgia to 2.2% in Rhode Island. During 2008–2009, an estimated 616,000 (annual average) adult females in the United States (0.5% of the adult female population) attempted suicide in the past year. Among females, prevalence ranged from <0.1% in Montana and Virginia to 1.3% in Connecticut (Table 10).

During 2008–2009, an estimated 663,000 (annual average) non-Hispanic white adults in the United States (0.4% of the adult non-Hispanic white population) attempted suicide in the past year. Among non-Hispanic white adults, prevalence ranged from <0.1% in Georgia to 1.5% in Rhode Island. During the same period, an estimated 182,000 (annual average) non-Hispanic black adults in the United States (0.7% of the adult non-Hispanic black population) attempted suicide in the past year. Among states with stable estimates for non-Hispanic black adults, prevalence ranged from 0.1% in Pennsylvania to 2.0% in California. During 2008-2009, an estimated 144,000 (annual average) Hispanic adults in the United States (0.5% of the population) attempted suicide in the past year. Among states with stable estimates for Hispanic adults, prevalence ranged from <0.1% in District of Columbia and Oregon to 3.0% in Connecticut. During that same period, an estimated 23,000 (annual average) non-Hispanic Asian adults in the United States (0.2% of the adult non-Hispanic

Asian population) attempted suicide in the past year. Among states with stable estimates for non-Hispanic Asian adults, prevalence ranged from 0.1% in California and Illinois to 0.3% in Michigan and New York (Table 11).

Discussion

This report is the first to present state-level data concerning answers to questions related to suicidal thoughts, plans, and attempts from the 2008 and 2009 NSDUH (29). The successful previous use of NSDUH data to describe other health behaviors (e.g., use of alcohol, illicit substances, and tobacco) (33–35) indicates its potential to measure progress in achieving national health objectives (36). The data could be particularly useful for state prevention practitioners in assessing and understanding the problem in their area and evaluating suicide prevention programs. The data may also be useful in generating research examining the factors that contribute to the sociodemographic differences in suicidal thoughts and behaviors (e.g., differences in the social determinants of health) (37,38).

The NSDUH results reveal significant sociodemographic and geographic variations in the estimated past-year prevalence of having suicidal thoughts, having made plans to attempt suicide, and having attempted suicide. Several theories have been advanced that attempt to explain these regional differences (37). These theories attribute the differences to selective migration (i.e., populations with risk factors for suicidal behavior might migrate to the same geographic areas), to the sociodemographic composition of the population (i.e., certain regions have a greater percentage of persons who are members of sociodemographic groups that are at greater risk for suicidal behavior), or to the local social environment (e.g., unemployment levels, divorce rates, social support, or environmental factors such as increased access to lethal means) (39,40). Such variations might point to areas of emphasis for planning and evaluation of prevention activities. In addition to describing the overall prevalence of suicide-related health problems, NSDUH data can be used to compare various aspects of suicidal thoughts and behaviors, analyze prevalence among subpopulations of adults (e.g., by sex or age group), assess trends in suicidal thoughts and behaviors over time, monitor progress toward achieving national or state health objectives, provide comparable state data, and evaluate and improve suicide-related prevention policies and programs.

Suicidal Thoughts

The prevalence of having seriously considered suicide differed by age, sex, state, and race/ethnicity. The prevalence of serious suicide consideration was higher among adults aged 18-29 years than it was for the overall U.S. population. Three recent population-based studies that measured 12-month prevalence of suicidal thought (the National Comorbidity Survey [NCS], the National Comorbidity Survey Replication [NCS-R], and the Injury Control and Risk Survey [ICARIS]) also revealed a higher prevalence of having suicidal thoughts among young adults (12,41,42). Previous studies consistently have demonstrated higher rates of having suicidal thoughts among females (12,41-44). In these analyses, prevalence was higher for females in the United States overall and in the majority of states. Similarly, racial/ethnic prevalences of suicidal thoughts differed; overall rates were highest for non-Hispanic whites. However, none of the racial/ethnic differences was statistically significant among non-Hispanic whites, non-Hispanic blacks, or Hispanics. Existing evidence on racial/ethnic differences in suicidal thoughts also has been mixed. Lifetime prevalence for having suicidal thoughts among respondents to the National Longitudinal Alcohol Epidemiologic Survey (NLAES) was lowest among non-Hispanic blacks and non-Hispanic Asians, but no significant differences were observed in NCS or the National Epidemiologic Survey on Alcohol and Related Studies (NESARC), which replaced NLAES in 2001 (41,43,44). Based on NSDUH data reporting, state-level prevalence of adults reporting having seriously considered suicide varied widely (range: 2.1%-6.8).

Suicide Plans

NCS, NCS-R, and ICARIS provided comparable national estimates but not state-level estimates on having made a plan to engage in suicidal behavior (12,41,42), and none of these was a regular, ongoing survey. The only significant sociodemographic difference in those samples was a higher prevalence of suicide planning among those aged 18–24 years and 35–44 years (41). The data provided in this report support this finding. Adults aged 18–29 years had a significantly higher prevalence of suicide planning than those aged ≥30 years. The only significant racial/ethnic difference was a lower prevalence of suicide planning among non-Hispanic Asians than among other racial/ethnic populations. However, NCS published data did not include Asians (42). Based on NSDUH data, state-level prevalence of reported suicide planning ranged from 0.1% to 2.8%.

Suicide Attempts

Comparable national-level (but not state-level) data on the 12-month prevalence of suicide attempts also are available from the NCS, NCS-R, NESARC and ICARIS samples (12,41,42). As in the NCS samples, the prevalence of suicide

attempts in the current analyses was significantly higher among those aged 18-29 years. The finding of no significant differences in previous 12-month suicide attempts by sex also supports the previous findings. The only significant racial/ ethnic difference identified in the current analyses was a lower prevalence of suicide attempts among non-Hispanic Asians. The NCS studies did not reveal any racial/ethnic differences in previous 12-month suicide attempts (41). In the ECA and NLAES samples, there were racial/ethnic differences in lifetime prevalence of suicide attempts; prevalence was lowest among non-Hispanic blacks (43,44). The ECA sample did not have an Asian population stratum for comparison. Although prevalence in the NLAES was qualitatively lowest among non-Hispanic blacks compared with other racial/ethnic groups, the estimated prevalence for non-Hispanic Asians was also low in this sample and did not differ significantly from that among non-Hispanic blacks. Based on the NSDUH data, state-level prevalence of adults reporting having made a suicide attempt varied widely (range: 0.1%-2.0%), this might in part reflect wide state-level differences in sociodemographic factors.

Limitations

The findings in this report are subject to at least three main limitations. First, because of low precision, certain estimates were suppressed after stratification by state and sociodemographic characteristics, which limited the ability to examine more age groups and certain racial/ethnic populations. However, the findings are comparable with those of previous studies examining similar types of thoughts and behaviors. Because this report focused on past-year prevalence of suicidal behaviors rather than on lifetime prevalence, it yielded lower estimates; however, the survey methodology reduced the potential for recall bias and thus provides more relevant information for current prevention and intervention (45). Previous studies have indicated that adults underreport suicidal thoughts that occurred when they were adolescents (46). Second, estimates from NSDUH are generalizable only to the civilian, noninstitutionalized population aged ≥18 years. Although this population includes the majority of the total U.S. population aged ≥18 years, it excludes active-duty military personnel, persons living in institutional group quarters (e.g., correctional facilities and residential drug treatment centers), and homeless persons not living in shelters on the survey date. In addition, suicidal thoughts and behaviors among American Indian/Alaska Native adults could not be reported because of low precision of sample estimates. Certain segments of these populations have a higher prevalence of some suicidal thoughts and behaviors than the general U.S. population, and thus this report might underestimate the actual prevalence rates of

suicidal thoughts and behaviors among adults in the United States (47,48). Finally, NSDUH data are based on self-reports, not observations of actual incidents in the case of suicide attempts, and the extent of underreporting or overreporting of behaviors cannot be determined. Similar survey questions that were administered among high school students demonstrated good test-retest reliability (49). However, no reliability or validity data were obtained on the measures of thoughts, plans, and attempts among adults. In addition, respondents tend to recall events in the recent past better than those occurring more distantly (50). This could reflect why certain groups would have lower prevalences if their event was 11 months prior to the survey compared with others whose event was only 1 month prior to the survey. Persons in the former group might not report their event because of recall bias, and so the finding might not mean that their rates were actually lower.

Conclusion

This report presents the first combined, representative state- and region-specific analyses of suicidal thoughts and behaviors among adults aged ≥18 years in the United States. The data presented in this report support other findings that the public health burden of suicidal thoughts and behaviors throughout the United States is much greater than the number of deaths (22). The findings in this report document important state-specific differences in the reported prevalence of suicidal thoughts, plans, and attempts both overall and when examined by age, sex, and race/ethnicity. For example, younger adults aged 18-29 years tended to report more suicidal thoughts, plans, and attempts in most states compared with those aged ≥30 years. The regional differences described in this report indicate some differences from state-level suicide analyses, which have consistently reported higher rates in the Rocky Mountain and western states (23). These studies have demonstrated that regional differences in demographic patterns (i.e., age, race/ethnicity, and sex) and in suicide methods do not account completely for variations in suicide. Nonfatal suicidal behavior has been demonstrated to have different patterns from fatal behavior, which might account for the differences between the state patterns in this report compared with patterns from mortality reports (40). For example, rates of fatal suicidal behavior are higher among males and middle aged or older adults whereas rates of nonfatal suicidal behavior are higher among females and adolescents and young adults (2).

In 1999, to maximize the benefits of suicide prevention efforts and resources, the U.S. Surgeon General issued the Call to Action to Prevent Suicide, which recommended using a public health approach to address suicide-related issues (51,52). The first step of such an approach is to define and assess the

problem, which includes measuring characteristics of persons who engage in the behavior. Whereas state vital statistics systems usually contain mortality data, sources of nonfatal population-level data on those who seriously consider or who make a plan to attempt suicide are limited. Furthermore, data that allow for state-specific assessments and comparisons of the prevalence of nonfatal suicidal behavior across states are lacking (26).

Although several other national surveys have conducted analyses of sociodemographic differences in suicidal thoughts or behaviors (12,41-43), they did not collect state-level data. For example, NCS (conducted during 1990-1992) and NCS-R (conducted during 2001–2003) were both nationally representative household surveys of U.S. adults (41,42). These surveys used fully structured research diagnostic interviews to assess the prevalence and correlates of psychiatric disorders and included sociodemographic factors; parental psychopathology; a 12-month and lifetime prevalence of suicidal thoughts, plans, gestures, and attempts together with their treatment; and mental disorders based on the criteria of the Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) (53). For each survey, approximately 10,000 respondents were interviewed. NESARC, conducted during 2001-2002, was a nationally representative survey focusing on the distribution and correlates of alcohol use disorders and associated conditions but included questions on suicidal thoughts and behaviors. It included 43,093 respondents (35). NSDUH uses epidemiologically based questions to assess sociodemographic characteristics and a 12-month prevalence of suicidal thoughts, plans, and attempts as well as other mental health and substance misuse-related issues. For NSDUH, approximately 92,000 respondents were interviewed in the 50 states and the District of Columbia.

Multiple individual, peer, family, community, and societal factors have been associated with suicidal behaviors. For example, levels of mental illness, substance misuse, family dysfunction, and violence victimization might be higher in communities with higher reports of suicidal behaviors (54). Similarly, suicidal behaviors tend to be higher in communities with low levels of social integration, high unemployment, lack of access to health-care services, and unstable social environments (54). The variations identified in this report might reflect differences in the prevalence of risk factors and socioeconomic composition of the study populations.

The wide variations identified between the age groups, sexes, and racial/ethnic populations in suicidal thoughts and behaviors indicate that different strategies are needed to address the needs of these diverse groups and eliminate health disparities. Several states have initiated multiple prevention strategies designed to reduce the impact of injuries from

suicidal behavior, including improving data collection efforts, public education campaigns, crisis intervention services, and family support programs (27,55). These strategies should be evaluated and, if documented to be effective, adapted for use in other states, particularly those with the highest reports of suicidal behavior.

Further studies are needed to better understand the effect of known risk and protective factors on the development and prevalence of suicidal thoughts and behaviors. These statespecific results can be of great utility in assessing a state's status on this health indicator in relation to neighboring states and within and across geographic regions. Only a small proportion of those who attempt suicide eventually die by suicide (56). However, persons with a past history of suicide attempts are at greater risk for dying by suicide than those who do not have such a history (3). Possible suicide-prevention strategies to implement could include universal strategies (e.g., public education campaigns that focus on improving recognition of suicide risk) and indicated strategies (e.g., cognitive-behavioral therapy) that address the needs of persons exhibiting certain risk factors (e.g., persons who have made suicide attempts) (57).

References

- Crosby AE, Ortega L, Melanson C. Self-directed violence surveillance: uniform definitions and recommended data elements, version 1.0. Atlanta, GA: US Department of Health and Human Services, CDC; 2011.
- DeLeo D, Bertolote J, Lester D. Self-directed violence. In: Krug EG, Dahlberg LL, Mercy JA, Zwi AB, Lozano R, eds. World report on violence and health. Geneva, Switzerland: World Health Organization; 2002:185–212.
- Goldsmith SK, Pellmar TC, Kleinman AM, Bunney WE, eds. Reducing suicide: a national imperative. Washington, DC: National Academy Press; 2002.
- 4. World Health Organization. Prevention of suicide: guidelines for the formulation and implementation of national strategies. Geneva, Switzerland: World Health Organization; 1996.
- 5. Murray CLJ, Lopez AD. Mortality by cause for eight regions of the world: global burden of disease study. Lancet 1997;349:1269–76.
- CDC. Web-Based Injury Statistics Query and Reporting System (WISQARS). Available at http://www.cdc.gov/injury/wisqars/index. html. Accessed October 12, 2011.
- 7. Rosenberg ML, Gelles RJ, Holinger PC. Violence: homicide, assault and suicide. In: Amler RW, Dull HB, eds. Closing the gap: the burden of unnecessary illness. New York, NY: Oxford University Press; 1987.
- Mościcki EK, Crosby A. Epidemiology of attempted suicide in adolescents: issues for prevention. Trends Evidence-Based Neuropsychiatry 2003;5:36–44.
- Niska R, Bhuiya F, Xu J. National Hospital Ambulatory Medical Care Survey: 2007 emergency department summary. National Health Stat Rep; no 26. Hyattsville, MD: US Department of Health and Human Services, CDC, National Center for Health Statistics; 2010.
- CDC, National Center for Health Statistics. National Hospital Ambulatory Medical Care Survey: 2008 emergency department summary tables. Available at http://www.cdc.gov/nchs/data/ahcd/nhamcs_ emergency/nhamcsed2008.pdf. Accessed October 12, 2011.
- 11. CDC. Nonfatal self-inflicted injuries treated in hospital emergency departments—United States, 2000. MMWR. 2002;51:436–8.

- Crosby AE, Cheltenham MP, Sacks JJ. Incidence of suicidal ideation and behavior in the United States, 1994. Suicide Life Threat Behav 1999;29:131–40.
- Bonnie RJ, Fulco CE, Liverman CT, eds. Reducing the burden of injury: advancing prevention and treatment. Washington, DC: National Academy Press; 1999.
- 14. State and Territorial Injury Prevention Directors' Association. Consensus recommendations for injury surveillance in state health departments. Marietta, GA: State and Territorial Injury Prevention Directors' Association; 2007.
- CDC. National estimates of nonfatal injuries treated in hospital emergency departments — United States, 2000. MMWR 2001;50;340-6.
- Holder Y, Peden M, Krug E, et al., eds. Injury surveillance guidelines. Geneva, Switzerland: World Health Organization; 2001.
- Sethi D, Habibula S, McGee K, et a., eds. Guidelines for conducting community surveys on injuries and violence. Geneva, Switzerland: World Health Organization; 2004.
- Harvard Scholl of Public Health. Proceedings of the workshop on using demographic and health survey (DHS) data for health sector reform. Available at http://www.hsph.harvard.edu/ihsg/publications/pdf/No-13. PDF. Accessed October 12, 2011.
- Harlow SD, Linet MS. Agreement between questionnaire data and medical records: the evidence for accuracy of recall. Am J Epidemiol 1989;129:233

 –48.
- Sadana R. Measuring reproductive health: review of community-based approaches to assessing morbidity. Bull World Health Organ 2000;78:640–54.
- Duncan T, Frankenberg E. The measurement and interpretation of health in social surveys. In: Murray C, Salomon J, Mathers C, Lopez A, eds. Measurement of the global burden of disease. Geneva, Switzerland: World Health Organization; 2002:387–420.
- Nock MK, Borges G, Bromet EJ, Cha CB, Kessler RC, Lee S. Suicide and suicidal behavior. Epidemiol Rev 2008;30:133–54.
- CDC. Regional variations in suicide rates—United States, 1990–1994.
 MMWR 1997;46:789–93.
- 24. US Department of Health and Human Services, Public Health Service. National strategy for suicide prevention: goals and objectives for action. Available at https://store.samhsa.gov/shin/content/SMA01-3517/SMA01-3517.pdf. Accessed October 12, 2011.
- 25. State and Territorial Injury Prevention Directors Association. Injury Surveillance Workgroup 5. Consensus recommendations for injury surveillance in state health departments. Available at http://ibis.health.state.nm.us/docs/Query/STIPDAConcensusRecommendations.pdf. Accessed October 12, 2011.
- Suicide Prevention Action Network. Strategies to improve non-fatal suicide attempt surveillance—recommendations from an expert roundtable. Washington, DC: Suicide Prevention Action Network; 2006.
- 27. Calgary Centre for Suicide Prevention. A summary of national, state and provincial strategies for the prevention of suicide. Calgary, Alberta, Canada: Calgary Centre for Suicide Prevention; 2004.
- 28. Substance Abuse and Mental Health Services Administration. Results from the 2008 National Survey on Drug Use and Health: national findings (Office of Applied Studies, NSDUH Series H-36, HHS Publication No. SMA 09-4434). Available at http://oas.samhsa.gov/nsduh/2k8nsduh/2k8Results.pdf. Accessed October 12, 2011.
- 29. Substance Abuse and Mental Health Services Administration. State estimates of substance use and mental disorders from the 2008–2009 National Surveys on Drug Use and Health, NSDUH Series H-40, HHS Publication No. (SMA) 11-4641. Available at http://store.samhsa.gov/shin/content/SMA11-4641/SMA11-4641.pdf. Accessed October 12, 2011.
- US Department of Agriculture, Economic Research Service. Measuring rurality: 2004 county typology codes. Available at http://www.ers.usda. gov/briefing/rurality/Typology. Accessed October 12, 2011.

- 31. Shah B, Barnwell B, Bieler G. SUDAAN user's manual. version 9.1 ed. Research Triangle Park, NC: Research Triangle Institute; 2005.
- 32. Substance Abuse and Mental Health Services Administration. 2005 National Survey on Drug Use and Health: statistical inference report. Available at http://oas.samhsa.gov/nsduh/2k5MRB/2k5statInference. pdf. Accessed October 12, 2011.
- CDC. CDC health disparities and inequalities report—United States, 2011. MMWR 2011;60(Suppl):109–13.
- Miller JW, Gfroerer JC, Brewer RD, Naimi TS, Mokdad A, Giles WH. Prevalence of adult binge drinking: a comparison of two national surveys. Am J Prev Med 2004;27:197–204.
- 35. Trosclair A, Dube SR. Smoking among adults reporting lifetime depression, anxiety, anxiety with depression, and major depressive episode, United States, 2005–2006. Addict Behav 2010;35:438–43.
- US Department of Health and Human Services. Tracking Healthy People 2010. Washington, DC: US Government Printing Office; 2000.
- 37. World Health Organization. Closing the gap in a generation: health equity through action on the social determinants of health. Report from the Commission on Social Determinants of Health. Geneva, Switzerland: World Health Organization; 2008.
- 38. Hicks J. The social determinants of elevated rates of suicide among Inuit youth. Indigenous Affairs. 2007; 4:30–37.
- 39. Lester D. Explaining regional differences in suicide rates. Soc Sci Med 1995;40:719–21.
- 40. Claassen CA, Carmody T, Bossarte R, Trivedi MH, Elliot S, Currier GW. Do geographic regions with higher suicide rates also have higher rates of nonfatal intentional self-harm? Suicide Life Threat Behav 2008;38:637–49.
- Kessler RC, Berglund P, Borges G, et al. Trends in suicide ideation, plans, gestures, and attempts in the United States, 1990–1992 to 2001–2003. JAMA 2005;293:2487–95.
- 42. Borges G, Angst J, Nock MK, et al. A risk index for 12-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med 2006;36:1747–57.
- Baca-Garcia E, Perez-Rodriguez MM, Keyes KM, et al. Suicidal ideation and suicide attempts in the United States: 1991–1992 and 2001–2002. Mol Psychiatry 2010;15:250–9.
- Moscicki EK, O'Carroll P, Regier DA, et al. Suicide attempts in the Epidemiologic Catchment Area Study. Yale J Biol Med 1988; 61:259–68.
- 45. Kruijshaar ME, Barendregt J, Vos T, De Graf R, Spijker J, Andrews G. Lifetime prevalence estimates of major depression: an indirect estimation method and a quantification of recall bias. Eur J Epidemiol 2005;20:103–11.
- Goldney RD, Smith S, Winefield AH, et al. Suicidal ideation: its enduring nature and associated morbidity. Acta Psychiatr Scand 1991;83:115–20.
- Mumola C, Noonan M. Deaths in custody statistical tables. Washington, DC: US Department of Justice, Office of Justice Programs, Bureau of Justice Statistics; 2008.
- 48. Centre for Suicide Prevention. Suicide among the homeless. Calgary, Alberta, Canada: Canadian Mental Health Association; 2010.
- Brener ND, Kann L, McManus T, Kinchen SA, Sundberg EC, Ross JG. Reliability of the 1999 Youth Risk Behavior Survey questionnaire. J Adolesc Health 2002;31:336

 –42.
- 50. Mościcki EK, O'Carroll P, Locke BZ, Rae DS, Roy AG, Regier DA. Suicidal ideation and attempts: the epidemiologic catchment area. In: US Department of Health and Human Services. Report of the Secretary's Task Force on Youth Suicide: vol. 4. Strategies for the prevention of youth suicide. Washington, DC: US Government Printing Office; 2008:115–28.
- 51. US Public Health Service. The Surgeon General's call to action to prevent suicide. Washington, DC: US Public Health Service; 1999.
- 52. Satcher DA. Bringing the public health approach to the problem of suicide. Suicide Life Threat Behav 1998;28:325–7.

Surveillance Summaries

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th ed. Washington, DC: American Psychiatric Association; 1994.
- 54. McLean J, Maxwell M, Platt S, Harris F, Jepson R. Risk and protective factors for suicide and suicidal behaviour: a literature review. Edinburgh, Scotland, UK: Scottish Government Social Research. 2008. Available at http://www.scotland.gov.uk/Publications/2008/11/28141444/0. Accessed October 12, 2011.
- 55. Lubell KM, Harber-Singer H, Gonzalez B. State suicide prevention planning: a CDC research brief. Atlanta, GA: CDC; 2008.
- 56. Kuo WH, Gallo JJ. Completed suicide after a suicide attempt. Am J Psychiatry 2005;162–3.
- 57. Mann JJ, Apter A, Bertolote J, et al. Suicide prevention strategies: a systematic review. JAMA 2005; 294:2064–74.

TABLE 1. Annual average estimated number* and percentage of adults aged ≥18 years who had suicidal thoughts or engaged in suicidal behaviors during the previous year, by selected demographic characteristics — National Survey on Drug Use and Health, United States, 2008–2009[†]

		Though	t		Plan			Attemp	t
Characteristic	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
Sex									
Male	3,789	3.5	(3.2-3.8)	1,045	1.0	(0.8-1.1)	442	0.4	(0.3-0.5)
Female	4,571	3.9	(3.7-4.2)	1,218	1.0	(0.9-1.2)	616	0.5	(0.4-0.6)
Age group (yrs)									
18–29	2,865	5.7	(5.4-6.0)	821	1.6	(1.5-1.8)	477	1.0	(0.8-1.1)
≥30	5,494	3.1	(2.9-3.4)	1,442	0.8	(0.7-0.9)	581	0.3	(0.3-0.4)
Race/Ethnicity									
White, non-Hispanic	6,044	3.9	(3.7-4.1)	1,616	1.0	(0.9-1.2)	663	0.4	(0.4-0.5)
Black, non-Hispanic	911	3.5	(3.1-4.0)	262	1.0	(0.8-1.3)	182	0.7	(0.5-0.9)
Hispanic [§]	933	3.0	(2.6-3.6)	267	0.9	(0.6-1.2)	144	0.5	(0.3-0.6)
Asian, non-Hispanic	208	2.1	(1.6-2.8)	37	0.4	(0.2-0.6)	23	0.2	(0.1-0.6)
Education									
Less than high school	1,373	4.0	(3.5-4.4)	429	1.2	(1.0-1.5)	247	0.7	(0.6-0.9)
High school graduate [¶]	2,823	4.0	(3.7-4.4)	730	1.0	(0.9-1.2)	393	0.6	(0.5-0.7)
Some college	2,359	4.1	(3.8-4.5)	672	1.2	(1.0-1.4)	304	0.5	(0.4-0.7)
College graduate or higher	1,805	2.8	(2.5-3.2)	433	0.7	(0.5-0.9)	113	0.2	(0.1-0.3)
Current employment									
Full-time	3,679	3.1	(2.9-3.3)	844	0.7	(0.6-0.8)	351	0.3	(0.2-0.4)
Part-time	1,380	4.5	(4.0-5.0)	405	1.3	(1.1-1.6)	200	0.6	(0.5-0.9)
Unemployed	768	6.5	(5.6-7.5)	244	2.1	(1.6-2.7)	118	1.0	(0.8-1.3)
Other**	2,532	3.9	(3.6-4.3)	770	1.2	(1.0-1.4)	389	0.6	(0.5-0.8)
County type									
Large metropolitan area ^{††}	4,353	3.6	(3.4-3.9)	1,148	1.0	(0.8-1.1)	559	0.5	(0.4-0.6)
Small metropolitan area ^{§§}	2,692	3.9	(3.6-4.3)	777	1.1	(1.0-1.3)	351	0.5	(0.4-0.6)
Nonmetropolitan area ^{¶¶}	1,315	3.5	(3.2-4.0)	338	0.9	(0.8-1.1)	148	0.4	(0.3-0.5)
Total**	8,359	3.7	(3.5–3.9)	2,263	1.0	(0.9–1.1)	1,058	0.5	(0.4–0.5)

Abbreviations: CI = confidence interval; thought = had serious thoughts of suicide; plan = made any suicide plan; attempt = attempted suicide.

^{*} In thousands.

[†] Estimates are based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

[§] Persons of Hispanic origin can be of any race

[¶] Includes persons with a general education diploma.

^{**} Includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

^{††} Area with a population of ≥1 million persons.

^{§§} Area with a population of <1 million persons.

Area that is outside of a metropolitan statistical area; includes urbanized counties with a population of ≥2,000 persons in urbanized areas, less urbanized counties with a population of ≥2,500 persons but <20,000 persons in urbanized areas, and completely rural counties with a population of <2,500 persons in urbanized areas

^{**} Totals exclude persons with missing or unknown race and ethnicity. Totals might vary due to rounding.

TABLE 2. Annual average estimated number* and percentage of adults aged ≥18 years who had serious thoughts of suicide, made any suicide plan, or attempted suicide during the previous year, by state and geographic region — National Survey on Drug Use and Health, United States, 2008–2009[†]

		Though	<u>t </u>		Plan			Attempt	i.
Region/State	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95%CI)
Geographic region§									
Northeast	1,469	3.5	(3.1-3.9)	358	0.9	(0.7-1.1)	182	0.4	(0.3-0.6)
Midwest	2,122	4.3	(4.0-4.7)	598	1.2	(1.0-1.4)	271	0.5	(0.4-0.7)
South	2,647	3.2	(2.9-3.5)	734	0.9	(0.8-1.1)	353	0.4	(0.3-0.5)
West	2,122	4.1	(3.7-4.5)	572	1.1	(0.9-1.3)	252	0.5	(0.4-0.6)
State									
Alabama	109	3.1	(2.1-4.7)	38	1.1	(0.6-2.0)	6	0.2	(0.1-0.4)
Alaska	16	3.4	(2.2-5.1)	5	1.0	(0.4-2.2)	1	0.2	(<0.1-0.8)
Arizona	199	4.2	(3.0-5.8)	43	0.9	(0.5–1.5)	29	0.6	(0.3-1.3)
Arkansas	105	5.0	(3.5-7.0)	30	1.4	(0.9-2.4)	19	0.9	(0.4-2.0)
California	912	3.4	(2.8-4.1)	244	0.9	(0.7-1.3)	118	0.4	(0.3-0.7)
Colorado	169	4.6	(3.1-6.9)	50	1.4	(0.6-3.1)	14	0.4	(0.2-0.7)
Connecticut	114	4.3	(3.2-5.8)	32	1.2	(0.6-2.3)	26	1.0	(0.5-1.9)
Delaware	15	2.3	(1.5-3.6)	3	0.5	(0.2-1.2)	1	0.1	(<0.1-0.3)
District of Columbia	21	4.5	(2.9-6.8)	4	0.9	(0.3-2.1)	2	0.4	(0.1-0.8)
Florida	520	3.7	(3.1-4.5)	175	1.3	(0.9-1.8)	73	0.5	(0.3-0.9)
Georgia	146	2.1	(1.3-3.3)	6	0.1	(<0.1-0.2)	6	0.1	(<0.1-0.3)
Hawaii	37	3.9	(2.7-5.6)	10	1.0	(0.5-2.0)	2	0.2	(0.1-0.5)
Idaho	66	6.0	(4.3-8.4)	12	1.1	(0.6-1.9)	6	0.6	(0.3-1.2)
Illinois	403	4.2	(3.6-5.0)	97	1.0	(0.7-1.4)	43	0.5	(0.3-0.7)
Indiana	215	4.6	(3.4-6.1)	64	1.4	(0.8-2.4)	17	0.4	(0.2-0.8)
lowa	91	4.0	(2.9-5.6)	23	1.0	(0.5-2.0)	7	0.3	(0.1-0.9)
Kansas	59	2.9	(2.1-4.1)	14	0.7	(0.3-1.3)	10	0.5	(0.2-1.2)
Kentucky	114	3.6	(2.4-5.4)	26	0.8	(0.5-1.5)	8	0.2	(0.1-0.6)
Louisiana	79	2.5	(1.6-3.7)	32	1.0	(0.5-2.0)	13	0.4	(0.2-0.8)
Maine	53	5.1	(3.6-7.2)	22	2.1	(1.1-4.2)	8	8.0	(0.3-2.3)
Maryland	122	2.9	(1.8-4.7)	27	0.6	(0.3-1.3)	11	0.3	(0.1-0.8)
Massachusetts	183	3.7	(2.5-5.3)	54	1.1	(0.5-2.2)	10	0.2	(0.1-0.4)
Michigan	332	4.4	(3.7-5.3)	117	1.6	(1.1-2.1)	58	0.8	(0.5-1.1)
Minnesota	168	4.3	(3.1-6.0)	59	1.5	(0.8-2.8)	29	0.8	(0.3-2.0)
Mississippi	50	2.4	(1.8-3.1)	13	0.6	(0.4-1.0)	9	0.4	(0.2-0.9)
Missouri	198	4.5	(3.1-6.5)	40	0.9	(0.4-2.1)	14	0.3	(0.1-1.1)
Montana	31	4.3	(3.1-5.7)	11	1.4	(0.9-2.3)	2	0.3	(0.1-0.7)
Nebraska	53	4.1	(2.8-6.0)	13	1.0	(0.6-1.7)	7	0.5	(0.3-0.9)
Nevada	124	6.5	(4.3-9.6)	42	2.2	(1.3-3.9)	10	0.5	(0.2-1.2)
New Hampshire	30	3.0	(2.2-4.0)	10	1.0	(0.6-1.6)	4	0.3	(0.2-0.8)
New Jersey	190	2.9	(1.9-4.5)	59	0.9	(0.3-2.3)	39	0.6	(0.2-2.1)
New Mexico	64	4.4	(2.8-6.8)	13	0.9	(0.6-1.4)	10	0.7	(0.3-1.3)
New York	539	3.6	(3.0-4.4)	90	0.6	(0.4-0.9)	56	0.4	(0.2-0.7)
North Carolina	232	3.4	(2.2-5.3)	74	1.1	(0.6-1.9)	33	0.5	(0.2-1.1)
North Dakota	13	2.7	(1.7–4.3)	4	8.0	(0.3-2.0)	2	0.3	(0.1-0.8)
Ohio	384	4.5	(3.7-5.4)	118	1.4	(0.9-2.1)	60	0.7	(0.5-1.0)
Oklahoma	79	3.0	(2.0-4.3)	24	0.9	(0.4-1.9)	11	0.4	(0.2-0.9)
Oregon	125	4.3	(3.0-6.1)	42	1.5	(0.7–3.1)	13	0.4	(0.2-1.1)
Pennsylvania	291	3.1	(2.5-3.7)	61	0.6	(0.4–1.0)	26	0.3	(0.1-0.5)
Rhode Island	50	6.2	(4.5-8.5)	22	2.8	(1.5–5.0)	12	1.5	(0.7-3.4)
South Carolina.	111	3.3	(2.0-5.6)	45	1.4	(0.5-3.8)	12	0.4	(0.1-1.0)
South Dakota	22	3.7	(2.6-5.3)	5	0.9	(0.5-1.6)	2	0.4	(0.2-0.7)
Tennessee	167	3.6	(2.4–5.3)	43	0.9	(0.5–1.8)	13	0.3	(0.1–0.7)
Texas	554	3.2	(2.6–4.0)	149	0.9	(0.6–1.3)	109	0.6	(0.4-1.0)
Utah	128	6.8	(5.0–9.3)	28	1.5	(0.8–2.8)	9	0.5	(0.2-1.0)
Vermont	19	3.9	(2.6-5.8)	8	1.5	(0.9–2.7)	2	0.5	(0.2-1.1)
Virginia	160	2.8	(1.9-4.1)	31	0.5	(0.2-1.4)	17	0.3	(0.1-1.1)
Washington	230	4.7	(3.4–6.4)	68	1.4	(0.7-2.8)	36	0.7	(0.3-2.0)
West Virginia	62	4.4	(3.1-6.2)	13	0.9	(0.4-2.1)	10	0.7	(0.3-2.0)
Wisconsin	184	4.3	(3.1–6.1)	45	1.1	(0.6-1.8)	21	0.5	(0.2-1.0)
Wyoming	20	5.1	(3.5–7.4)	6	1.4	(0.7–2.6)	2	0.5	(0.2-1.1)
Total	8,359	3.7	(3.5-3.9)	2,263	1.0	(0.9-1.1)	1,058	0.5	(0.4-0.5)

Abbreviations: CI = confidence interval; thought = had serious thoughts of suicide; plan = made any suicide plan; attempt = attempted suicide.

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

S Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, New Mexico, Oregon, Utah, Washington, and Wyoming.

TABLE 3. Annual average estimated number* and percentage of adults aged ≥18 years who had seriously considered suicide during the previous year, by state/area and age group — National Survey on Drug Use and Health, United States, 2008–2009[†]

Midwest 740 6.7 (6.2-7.3) 1.381 3.6 (3.2-4) (3.0-6.5) (3.0-6			18–29 yrs			≥30 yrs	
Michwest 514 57 (6.2-7.3) 1.381 3.6 (3.2-1 Mortheast 514 57 (5.0-6.5) 955 2.9 (2.5-5) South 875 4.9 (4.4-5.4) 1.772 2.8 (2.4-5) South 875 4.9 (4.4-5.4) 1.782 2.8 (2.4-5) South 875 4.9 (4.4-5.4) 1.782 2.8 (2.4-5) South 875 4.9 (4.4-5.4) 1.782 2.8 (2.4-5) South 875 4.9 (2.4-5.4) 1.8	Region/State	No.	%	(95% CI)	No.	%	(95% CI)
Midwest 740 6.7 (6.2-7.3) 1.381 3.6 (3.2-6) Northeast 514 5.7 (5.0-6.5) 955 2.9 (2.2-5) South 875 4.9 (4.4-5.4) 1.772 2.8 (2.4-5) South 875 4.9 (4.4-5.4) 1.782 2.8 (2.4-5) South 875 4.9 (2.4-5) 1.3 (2.4-5) 1.2 (2.4-5) 1.3	Geographic region§						
Northeast		740	6.7	(6.2-7.3)	1,381	3.6	(3.2-4.0)
South 875 49 (44-54) 1,772 2.8 (24-54) State State 35 (36.0) (54-68) 1,386 3.5 (36.2) Alabama 41 5.5 (37-82) (58.0) 10 2.7 (1.7-4.1) Alabama 41 5.5 (31-95) 10 2.7 (1.7-4.1) Alabama 37 81 (5-11.8) 68 41 (2.4-4.1) Arkansa 37 8.1 (5-11.8) 68 41 (2.4-4.1) Collorado 37 45 (2.8-71) 132 47 (2.8-1) Colorado 37 45 (2.8-71) 132 47 (2.8-1) Debware 6 47 (3.3-66) 9 1.7 (9.9-1) Florida 151 5.7 (46-71) 330 33 (2.5-1) Florida 151 5.7 (46-71) 330 33 (2.5-1) Georgia 5							(2.5-3.4)
Section Sect	South						(2.4-3.1)
Alabama 41 5.5 (3.7-8.2) 68 2.5 (1.3-Alashama 41 5.5 (3.7-8.2) 68 2.5 (1.3-Alashama 41 5.5 (3.7-8.2) 68 2.5 (1.7-Alizona 78 7.1 (4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 121 3.3 (2.1-4.7-10.5) 122 4.7 (2.6-10.5) 122							(3.0-4.0)
Alabama 41 5.5 G.3.7-8.2) 68 2.5 [1.3-4] Alaska 6 5.5 63.1-9.5) 10 2.7 (1.7-4) Artzona 78 7.1 (4.7-10.9) 121 33 (2.1-4) Arkamas 37 8.1 (5.5-11.8) 68 4.1 (2.4-4) Colorado 37 4.5 (2.8-7.1) 192 4.7 (2.6-1) Colorado 37 4.5 (2.8-7.1) 192 4.7 (2.6-1) Colorado 37 4.6 (2.8-7.1) 192 4.7 (2.6-1) Delaware 6 4.7 (3.3-6.6) 9 1.7 (0.9-2) Ebritido 151 5.7 (4.6-7.1) 370 3.3 (2.5-4) Horda 151 5.7 (4.6-10.7) 24 3.2 (1.9-4) Edorigia 5.8 3.7 (2.6-6.8) 2.9 3.8 3.0-4 Illinois 18 6.8				• • • • • • • • • • • • • • • • • • • •			**************************************
Alaska 6 6 5.5 (3.1-9.5) 10 0 2.7 (1.7-4 Artizona 78 7.1 (4.7-10.5) 121 3.3 (2.1-4 Artizona 78 7.1 (4.7-10.5) 121 3.3 (2.1-5 Artizonas 77 8.1 (5.5-11.8) 68 4.1 (2.4-5 (2.4-10.1) 132 4.7 (2.8-5 (2.4-10.1) 132 4.7 (2.4-10.1) 132 4.7 (2.8-		41	5.5	(3.7_8.2)	68	2.5	(1.3-4.6)
Arizona							
Arkansas 37 8.1 (55-11.8) 68 4.1 (2.4-California 362 5.6 (4.7-6.8) 549 2.7 (2.0-10 1.00 1.00 1.00 1.00 1.00 1.00 1.00							
Calfornia 362 5.6 (4,7-6,8) 549 2.7 (2.9-7) Colorado 37 4.5 (2.8-7.1) 132 4.7 (2.8-8) Connecticut 34 6.2 (3.6-6) 9 1.7 (0.9-9) District of Columbia 7 4.9 (3.2-7.3) 1.4 4.3 (2.6-1) Florida 151 5.7 (4.6-7.1) 370 3.3 (2.5-4) Florida 151 5.7 (4.6-7.1) 370 3.3 (2.5-4) Edwaid 13 6.9 (4.4-10.7) 2.4 3.2 (1.5-4) Idaho 18 6.8 (4.5-9.2) 4.8 5.8 (3.7-4) Illinois 124 5.6 (4.6-8.8) 2.79 3.8 (3.0-4) Illinois 124 5.6 (4.6-8.8) 2.79 3.8 (3.0-4) Illinois 124 5.6 (4.6-8.8) 2.79 3.8 (3.0-4) Kentucky 2.8							
Colorado 37 4.5 (2.8-7.1) 132 4.7 (2.8-1) Connecticut 34 6.2 (3.6-10.5) 80 3.8 (2.5-1) Delaware 6 4.7 (3.3-6.6) 9 1.7 (9.9-1) District of Columbia 7 4.9 (3.2-7.3) 1.4 4.3 (2.0-6.8) Florida 151 5.7 (4.6-7.1) 370 3.3 (2.5-6) Georgia 58 3.7 (2.0-6.8) 88 1.6 (0.8-1) Idaho 118 6.8 (4.9-9.2) 48 5.8 (3.7-5) Illinois 1124 5.6 (4.6-6.8) 279 3.8 (3.7-5) Illinois 124 5.6 (4.6-6.8) 279 3.8 (3.7-5) Illinois 124 5.6 (4.6-6.8) 2.79 3.8 (3.7-5) Illinois 13 7.9 (5.3-9.2) 142 3.9 2.7-2 Kentucky 2.8 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>(2.0–3.6)</td>							(2.0–3.6)
Connecticut 34 6.2 (3.6-10.5) 80 3.8 (2.5-10.5) Delayare 6 4.7 (3.3-6.6) 9 1.7 (0.9-10.5) Elorida 151 5.7 (4.6-7.1) 370 3.3 (2.6-11.5) Florida 151 5.7 (4.6-7.1) 370 3.3 (2.6-11.5) Georgia 58 3.7 (2.0-6.8) 88 1.6 (0.9-11.5) Hawaii 13 6.9 (4.4-10.7) 24 3.2 (1.9-11.5) Idaho 18 6.8 (4.9-9.2) 48 5.8 (3.7-2) Illinois 124 5.6 (4.6-6.8) 2.79 3.8 (3.0-11.5) Indiana 73 7.0 (53-9.2) 142 3.9 (2.5-2.5) Illinois 124 5.6 (4.6-8.8) 2.79 3.8 2.1 (1.2-2.5) Kansa 2.6 5.5 (3.9-2.7) 3.3 2.1 (1.2-2.5) (8.6							
Delaware 6 4.7 (3.3-6.6) 9 1.7 (0.9-1) District of Columbia 7 4.9 (3.2-7.3) 1.4 4.3 (2.6-1) Florida 151 5.7 (4.6-7.1) 370 3.3 (2.2-4) Georgia 58 3.7 (2.0-6.8) 88 1.6 (0.8-4) Idaho 18 6.8 (4.9-9.2) 48 5.8 (3.7-4) Illinois 124 5.6 (4.6-6.8) 279 3.8 (3.7-4) Illinois 124 5.6 (4.6-6.8) 279 3.8 (3.7-4) Illinois 13 7.0 (5.3-9.2) 142 3.9 (2.5-5) Illinois 13 7.0 (5.3-9.2) 142 3.9 (2.5-5) Kansas 26 5.5 (3.9-7.7) 33 2.1 (1.2-4 Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-4 Louisiana 13 <t< td=""><td></td><td></td><td></td><td>2012-2022 N C1000</td><td></td><td></td><td></td></t<>				2012-2022 N C1000			
District of Columbia 7							(0.9–3.3)
Florida							(2.6–7.2)
Georgia 58 3.7 (2.0-6.8) 88 1.6 (0.8-1) Idaho 13 6.9 (4.4-10.7) 24 3.2 (1.9-1) Idaho 18 6.8 (4.9-92) 48 5.8 (3.7-5) Illinois 124 5.6 (4.6-6.8) 2.79 3.8 (3.0-1) Indiana 73 7.0 (5.3-92) 142 3.9 (2.5-4) Kansas 26 5.5 (3.9-7.7) 33 2.1 (1.2-4) Kansas 26 5.5 (3.9-7.7) 33 2.1 (1.2-4) Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-4) Maine 13 6.9 <td< td=""><td></td><td></td><td></td><td>3</td><td></td><td></td><td>124 mar 1 mar 2</td></td<>				3			124 mar 1 mar 2
Hawaiii 13 6.9 (4.4-10.7) 24 3.2 (1.9-2) Idaho 118 6.8 (4.9-9.2) 48 5.8 (3.7-4) Illinois 124 5.6 (4.6-6.8) 279 3.8 (3.7-4) Illinois 124 5.6 (4.6-6.8) 279 3.8 (3.7-4) Illinois 13 7.0 (5.3-92) 142 3.9 (2.5-5) Kansas 26 5.5 (3.9-77) 33 2.1 (1.2-5) Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-5) Louisiana 40 5.5 (3.8-7.8) 40 1.6 (0.8-3-1) Maire 13 6.9 (4.5-10.3) 39 4.7 (3.0-1) Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-4 Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-2 Michigan 131 7.9							(0.8–3.3)
Idaho 18 6.8 (49-92) 48 5.8 (3.7-5) Illinois 124 5.6 (46-6.8) 279 3.8 (3.3-5) Indiana 73 7.0 (5.3-92) 142 3.9 (2.5-5) Iowa 34 7.1 (4.9-10.2) 57 3.2 (20-15) Kansas 26 5.5 (3.9-7.7) 33 2.1 (1.2-25) Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-35) Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-45) Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-45) Maine 13 6.9 (4.5-10.3) 39 4.7 (3.0-2-14) Maine 13 6.9 (4.5-10.3) 39 4.7 (3.0-2-14) Massachusetts 60 5.5 (3.6-8.9) 70 2.1 (10-44) Minsouri 60 6.2 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>							
Illinois 124 5.6 (4.6-6.8) 279 3.8 3.3—10.3 Indiana 73 7.0 (5.3–9.2) 142 3.9 2.5—5 Iowa 34 7.1 (4.9–10.2) 57 3.2 (2.0–6 Kansas 26 5.5 (3.9–7.7) 33 2.1 (1.2–1 Kentucky 28 4.0 (2.7–5.8) 86 8.5 (2.1–1 Louisiana 40 5.5 (3.8–7.8) 40 1.6 (0.8–3 Maine 13 6.9 (4.5–10.3) 39 4.7 (3.0–7 Maryland 52 5.7 (3.6–8.9) 70 0.2.1 (1.0–4 Massachusetts 60 5.5 (3.6–8.2) 123 3.2 (1.9–4 Michigan 131 7.9 (6.8–9.2) 201 1.3 (2.2–4 Mississippi 24 5.2 (3.5–7.4) 26 1.6 (0.9–2 Missouri 60 6.2							(3.7–9.0)
Indiana 73 7.0 (53-92) 142 3.9 (2.5-6) Iowa 34 7.1 (4.9-102) 57 3.2 (2.0-6) Kansas 26 5.5 (3.9-7.7) 33 2.1 (1.2-6) Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-6) Louisiana 40 5.5 (3.8-7.8) 40 1.6 (0.8-6) Maine 13 6.9 (4.5-10.3) 39 4.7 (3.0-7) Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-4) Massachusetts 60 5.5 (3.6-8.9) 70 2.1 (1.0-4) Missouth 60 5.5 (3.6-8.9) 10 3.2 (2.7-4) Missouth 60 6.2 (4.2-9.2) 137 40 (2.7-4) Missouth 60 6.2 (4.2-9.2) 137 40 (2.4-4) Nothasa 16 5.2				3 9			(3.0-4.9)
New Act							(2.5-5.9)
Kansas 26 5.5 (3.9-7.7) 33 2.1 (1.2-1) Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-1) Louisiana 40 5.5 (3.8-7.8) 40 1.6 (0.8-1) Maine 13 6.9 (4.5-10.3) 39 4.7 (3.0-2) Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-4) Massachusetts 60 5.5 (3.6-8.9) 201 3.4 (2.7-4) Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4) Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4) Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4) Missouri 60 6.2 (3.5-7.4) 26 1.6 (0.9-2) Missouri 60 6.2 (3.5-7.6) 3.8 3.7 (2.2-4) Nevalrampshire 11							(2.0-5.1)
Kentucky 28 4.0 (2.7-5.8) 86 3.5 (2.1-5.8) Louisiana 40 5.5 (3.8-7.8) 40 1.6 (0.8-3) Maine 13 6.9 (4.5-10.3) 39 4.7 (3.0-3) Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-4) Missachusetts 60 5.5 (3.6-8.2) 123 3.2 (1.9-4) Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4) Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4) Michigan 6.3 7.3 (5.1-10.5) 105 3.5 (2.2-5) Mississippi 24 5.2 (3.5-7.4) 2.6 1.6 (0.9-4) Mississippi 24 5.2 (3.5-7.4) 2.6 1.6 (0.9-4) Mississippi 60 6.2 (4.2-9.2) 137 4.0 (2.4-4) Mobrasach 1				• T. D. C.			
Louisiana 40 5.5 (3.8-7.8) 40 1.6 (0.8-1) Maine 13 6.9 (4.5-10.3) 39 4.7 (3.0-1) Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-4) Massachusetts 60 5.5 (3.6-8.2) 123 3.2 (1.9-5) Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4) Misnesota 63 7.3 (5.1-10.5) 105 3.5 (2.2-4) Missouri 60 6.2 (4.2-9.2) 137 4.0 (2.4-4) Mortana 12 7.7 (5.1-11.4) 20 3.4 (2.2-5) Nebaska 16 5.2 (3.6-7.6) 38 3.7 (2.2-6) Nevadad 36 8.8 (5.6-7.6) 38 5.9 (3.4-8.0) 19 2.3 (15-3-15) 8.8 5.9 (3.4-8.0) 19 2.3 (15-3-15) 8.8 5.9							100
Maine 13 6.9 (4.5-10.3) 39 4.7 (3.0-10.4) Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-4 Massachusetts 60 5.5 (3.6-8.2) 123 3.2 (1.9-5 Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-6 Misnissippi 24 5.2 (3.5-74) 26 1.6 (0.9-5 Mississippi 24 5.2 (3.5-74) 26 1.6 (0.9-5 Mississippi 24 5.2 (3.5-74) 26 1.6 (0.9-5 Mississippi 24 5.2 (3.6-76) 38 3.7 (2.2-6 Mortana 12 7.7 (5.1-11.4) 20 3.4 (2.2-6 Nebaska 16 5.2 (3.6-76) 38 3.7 (2.2-6 New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (158 New Mexico 19							
Maryland 52 5.7 (3.6-8.9) 70 2.1 (1.0-4 Massachusetts 60 5.5 (3.6-8.2) 123 3.2 (1.9-4 Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4 Minnesota 63 7.3 (5.1-10.5) 105 3.5 (2.2-5 Mississippi 24 5.2 (3.5-74) 26 1.6 (0.9-2 Missouri 60 6.2 (4.2-92) 137 4.0 (2.4-4 Mortana 12 7.7 (5.1-11.4) 20 3.4 (2.2-5 Mebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-6 Nevalda 36 8.8 (5.6-13.5) 88 5.9 (3.4-2 New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-2 New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-3 New Mork 195 <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<>							
Massachusetts 60 5.5 (3.6-8.2) 123 3.2 (1.9-2) Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4) Misnesota 63 7.3 (5.1-0.5) 105 3.5 (2.2-5) Mississippi 24 5.2 (3.5-7.4) 26 1.6 (0.9-2) Missouri 60 6.2 (4.2-9.2) 137 4.0 (2.4-6) Mortana 12 7.7 (5.1-14) 20 3.4 (2.2-6) Nebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-6) Newada 36 8.8 (5.6-13.5) 88 5.9 (3.4-6) New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-2) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-2) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-2) New York 195 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>							
Michigan 131 7.9 (6.8-9.2) 201 3.4 (2.7-4 Minnesota 63 7.3 (5.1-10.5) 105 3.5 (2.2-5 Mississippi 24 5.2 (3.5-7.4) 26 1.6 (0.9-5 Missouri 60 6.2 (4.2-9.2) 137 4.0 (2.4-6 Montana 112 7.7 (5.1-11.4) 20 3.4 (2.2-5 Nebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-5 Nevadad 36 8.8 (5.6-13.5) 88 5.9 (3.4-6 New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (15-2 New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-2 New Mexico 195 5.9 (4.8-7.2) 344 3.0 (2.3-3 North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6 Ohio 122	100 - C - C - C - C - C - C - C - C - C -						
Minnesota 63 7.3 (5.1-10.5) 105 3.5 (2.2-5) Mississpipi 24 5.2 (3.5-7.4) 26 1.6 (0.9-2) Missouri 60 6.2 (4.2-9.2) 137 4.0 (2.4-6) Montana 12 7.7 (5.1-11.4) 20 3.4 (2.2-6) Nebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-6) New Hadd 36 8.8 (5.6-13.5) 88 5.9 (3.4-5) New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-2) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-2) New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-2) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-6) Ohio 122 </td <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>							
Mississippi 24 5.2 (3.5-7.4) 26 1.6 (0.9-2) Missouri 60 6.2 (4.2-9.2) 137 4.0 (2.4-6) Montana 12 7.7 (5.1-11.4) 20 3.4 (2.2-6) Nebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-6) New Ada 36 8.8 (5.6-13.5) 88 5.9 (3.4-6) New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-2) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-2) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-7) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-7) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-7) New York 0.1 3.3 (4.0-9.8) 10.4 2.0 4.1 (1.0-2 North Carolin							
Missouri 60 6.2 (4.2-9.2) 137 4.0 (2.4-6) Montana 12 7.7 (5.1-11.4) 20 3.4 (2.2-5) Nebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-5) Nevada 36 8.8 (5.6-13.5) 88 5.9 (3.4-6) New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-3) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-3) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-3) New Mexico 195 5.9 (4.8-7.2) 344 3.0 (2.3-3) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-6) Oregon 36				Para de la companya del companya de la companya del companya de la			
Montana 12 7.7 (5.1-11.4) 20 3.4 (2.2-5) Nebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-6) New Ada 36 8.8 (5.6-13.5) 88 5.9 (3.4-6) New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-2) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-2) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-7) New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-8) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-6) Pennsylvania							
Nebraska 16 5.2 (3.6-7.6) 38 3.7 (2.2-6) Nevada 36 8.8 (5.6-13.5) 88 5.9 (3.4-5) New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-2) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-2) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-7) New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-3) North Dakota 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-6) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-6) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-3) Rhode Island 10							
Nevada 36 8.8 (5.6-13.5) 88 5.9 (3.4-6) New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-3) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-3) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-3) New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-3) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-4) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.0-6.1) 193 2.6 (2.0-3) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-5) South Carolina 3							
New Hampshire 11 5.6 (3.9-8.0) 19 2.3 (1.5-2) New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-2) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-2) New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-3) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-4) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-3) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-5) South Dakota				.2			151 3
New Jersey 86 6.3 (4.0-9.8) 104 2.0 (1.0-2) New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-7) New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-2) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-6) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-6) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-3) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-9) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-5) Texas 173<				•			
New Mexico 19 5.7 (3.9-8.3) 45 4.1 (2.1-2) New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-3) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-6) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-2) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-6) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-6) South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-6) Texas 17							
New York 195 5.9 (4.8-7.2) 344 3.0 (2.3-3-3) North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-6) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-6) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-2) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-6) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-6) South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4) Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-3) Utah 44 </td <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>							
North Carolina 49 3.3 (2.2-5.0) 183 3.4 (1.9-6) North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-4) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-3) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-6) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-5) South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4) Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-3) Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-9) Vermont 7							
North Dakota 4 3.4 (2.2-5.2) 9 2.5 (1.3-4) Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-5) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-5) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-6) South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4) Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5) Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-3) Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-5) Vermont 7				and the same of th			100 CO 10
Ohio 122 6.8 (5.6-8.1) 262 3.9 (3.0-5) Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-3) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-8) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-5) South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4) Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5) Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-5) Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-6) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5) Washington 69							
Oklahoma 30 5.3 (3.6-7.7) 49 2.4 (1.4-4) Oregon 36 6.0 (4.4-8.2) 88 3.9 (2.5-6) Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-3) Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-6) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-6) South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4) Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5) Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-3) Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-4) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5) Virginia 81 6.4 (4.1-9.8) 79 1.8 (0.9-3) Washington 69							200
Oregon 36 6.0 (4.4–8.2) 88 3.9 (2.5–6 Pennsylvania 98 5.0 (4.0–6.1) 193 2.6 (2.0–3 Rhode Island 10 5.1 (3.2–8.0) 40 6.5 (4.4–6 South Carolina 31 4.5 (3.1–6.4) 80 3.0 (1.5–6 South Dakota 12 8.3 (6.3–10.8) 10 2.3 (1.2–4 Tennessee 52 5.2 (3.5–7.6) 115 3.2 (1.8–5 Texas 173 4.2 (3.4–5.2) 380 2.9 (2.2–3 Utah 44 7.9 (5.5–11.4) 84 6.4 (4.3–8) Vermont 7 7.3 (5.2–10.2) 12 3.1 (1.8–5 Virginia 81 6.4 (4.1–9.8) 79 1.8 (0.9–3 West Virginia 16 5.9 (4.2–8.2) 47 4.1 (2.5–6							
Pennsylvania 98 5.0 (4.0-6.1) 193 2.6 (2.0-3 Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-9 South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-9 South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4 Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5 Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-3 Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-9) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5) Washington 69 6.2 (4.2-9.2) 161 4.2 (2.7-6) West Virginia 16 5.9 (4.2-8.2) 47 4.1 (2.5-6)							201
Rhode Island 10 5.1 (3.2-8.0) 40 6.5 (4.4-5) South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-5) South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4) Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5) Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-3) Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-9) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5) Virginia 81 6.4 (4.1-9.8) 79 1.8 (0.9-3) Washington 69 6.2 (4.2-9.2) 161 4.2 (2.7-6) West Virginia 16 5.9 (4.2-8.2) 47 4.1 (2.5-6)				0.500.00.000.00			A-1-1000 (1-1)
South Carolina 31 4.5 (3.1-6.4) 80 3.0 (1.5-5 South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4 Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5 Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-5 Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-9) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5 Virginia 81 6.4 (4.1-9.8) 79 1.8 (0.9-5 Washington 69 6.2 (4.2-9.2) 161 4.2 (2.7-6 West Virginia 16 5.9 (4.2-8.2) 47 4.1 (2.5-6)							181
South Dakota 12 8.3 (6.3-10.8) 10 2.3 (1.2-4) Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5) Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-5) Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-6) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5) Virginia 81 6.4 (4.1-9.8) 79 1.8 (0.9-2) Washington 69 6.2 (4.2-9.2) 161 4.2 (2.7-6) West Virginia 16 5.9 (4.2-8.2) 47 4.1 (2.5-6)							
Tennessee 52 5.2 (3.5-7.6) 115 3.2 (1.8-5 Texas 173 4.2 (3.4-5.2) 380 2.9 (2.2-3 Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-9) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5) Virginia 81 6.4 (4.1-9.8) 79 1.8 (0.9-3) Washington 69 6.2 (4.2-9.2) 161 4.2 (2.7-6) West Virginia 16 5.9 (4.2-8.2) 47 4.1 (2.5-6)							
Texas 173 4.2 (3.4–5.2) 380 2.9 (2.2–3 Utah 44 7.9 (5.5–11.4) 84 6.4 (4.3–5 Vermont 7 7.3 (5.2–10.2) 12 3.1 (1.8–5 Virginia 81 6.4 (4.1–9.8) 79 1.8 (0.9–3 Washington 69 6.2 (4.2–9.2) 161 4.2 (2.7–6 West Virginia 16 5.9 (4.2–8.2) 47 4.1 (2.5–6							
Utah 44 7.9 (5.5-11.4) 84 6.4 (4.3-6) Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5) Virginia 81 6.4 (4.1-9.8) 79 1.8 (0.9-3) Washington 69 6.2 (4.2-9.2) 161 4.2 (2.7-6) West Virginia 16 5.9 (4.2-8.2) 47 4.1 (2.5-6)							
Vermont 7 7.3 (5.2-10.2) 12 3.1 (1.8-5 Virginia 81 6.4 (4.1-9.8) 79 1.8 (0.9-3 Washington 69 6.2 (4.2-9.2) 161 4.2 (2.7-6 West Virginia 16 5.9 (4.2-8.2) 47 4.1 (2.5-6							
Virginia 81 6.4 (4.1–9.8) 79 1.8 (0.9–3 Washington 69 6.2 (4.2–9.2) 161 4.2 (2.7–6 West Virginia 16 5.9 (4.2–8.2) 47 4.1 (2.5–6							
Washington 69 6.2 (4.2–9.2) 161 4.2 (2.7–6 West Virginia 16 5.9 (4.2–8.2) 47 4.1 (2.5–6							(1.8–5.4)
West Virginia 16 5.9 (4.2–8.2) 47 4.1 (2.5–6							100
							(2.7–6.6)
wisconsin /6 8.4 (6.2–11.2) 108 3.2 (1.9–5				0.5400.00 10.000.00			(2.5-6.6)
				3			(1.9–5.4)
	0.0 At 1.0 0.0 0.0 0.0						(2.9–7.8)
Total 2,865 5.7 (5.4–6.0) 5,494 3.1 (2.9–3	IOTAI	2,865	5./	(5.4–6.0)	5,494	3.1	(2.9–3.4)

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded

[§] Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

TABLE 4. Annual average estimated number* and percentage of adults aged ≥18 years who had seriously considered suicide during the previous year, by state and sex — National Survey on Drug Use and Health, United States, 2008–2009[†]

		Male			Female	
Region/State	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region§						
Midwest	956	4.0	(3.5-4.5)	1,166	4.6	(4.1-5.1)
Northeast	643	3.2	(2.8–3.8)	826	3.8	(3.3-4.4)
South	1,256	3.2	(2.8–3.7)	1,391	3.2	(2.9-3.7)
West	934	3.7	(3.1–4.3)	1,188	4.5	(3.9–5.1)
State				S. W. Carriero		
Alabama	53	3.3	(1.7-6.2)	56	3.0	(1.6-5.5)
Alaska	7	3.0	(1.5–5.9)	9	3.7	(2.5–5.6)
Arizona	92	4.0	(2.3–6.7)	106	4.4	(3.0-6.4)
Arkansas	49	4.9	(2.9–8.1)	56	5.1	(3.3–7.8)
California	404	3.1	(2.3–4.1)	508	3.7	(3.0–4.7)
Colorado	99	5.5	(2.8–10.4)	71	3.8	(2.5–5.9)
Connecticut	59	4.6	(2.7–7.7)	55	4.0	(2.6–6.3)
Delaware	7	2.2	(1.1–4.2)	8	2.4	(1.4–4.2)
District of Columbia	7	3.2	(1.6–6.4)	14	5.6	(3.5–8.8)
Florida	224	3.3	(2.5–4.4)	296	4.1	(3.1–5.2)
	91	2.7	(1.5–5.0)	55	1.5	(0.9–2.6)
Georgia Hawaii	17	3.6	(2.5–5.2)	20	4.2	(2.3–7.3)
Idaho	27	5.1		39	7.0	
Illinois	172		(3.1–8.2)	231		(4.7–10.4)
Indiana	82	3.7	(2.8–5.0)	133	4.7 5.5	(3.8–5.9) (3.8–7.9)
	38	3.6	(2.1–6.2) (2.1–5.7)	53		(3.0–7.0)
lowa		3.5	,		4.6	
Kansas	20	2.0	(1.1–3.7)	39	3.7	(2.5–5.5)
Kentucky	47	3.1	(1.7–5.7)	67	4.1	(2.4–6.8)
Louisiana	42	2.8	(1.5–4.9)	37	2.2	(1.3–3.6)
Maine	27	5.4	(3.1–9.3)	26	4.9	(3.3–7.1)
Maryland	52	2.6	(1.4–4.7)	71	3.2	(1.7–6.1)
Massachusetts	72	3.0	(1.6–5.4)	112	4.3	(2.7–6.7)
Michigan	154	4.3	(3.3–5.5)	178	4.6	(3.7–5.7)
Minnesota	76	3.9	(2.4–6.5)	93	4.7	(3.1–7.1)
Mississippi	13	1.3	(0.8–2.0)	37	3.3	(2.3–4.7)
Missouri	85	4.0	(2.4–6.8)	113	4.9	(3.0–7.9)
Montana	15	4.3	(2.8–6.6)	16	4.2	(2.6–7.0)
Nebraska	26	4.0	(2.1–7.7)	28	4.1	(2.7–6.3)
Nevada	37	3.8	(2.2–6.6)	87	9.1	(5.4–15.0)
New Hampshire	18	3.6	(2.6–5.1)	12	2.4	(1.4–4.0)
New Jersey	63	2.0	(1.4–3.0)	127	3.7	(2.0-6.8)
New Mexico	22	3.1	(1.7–5.5)	43	5.7	(3.1–10.2)
New York	242	3.5	(2.6–4.6)	296	3.8	(3.0–4.9)
North Carolina	129	4.0	(2.4–6.7)	103	2.9	(1.4–5.7)
North Dakota	6	2.6	(1.4–5.1)	7	2.7	(1.6–4.7)
Ohio	184	4.5	(3.3–6.1)	200	4.5	(3.5–5.6)
Oklahoma	38	3.0	(1.7–5.2)	41	3.0	(1.8–4.9)
Oregon	65	4.6	(2.7–7.5)	60	4.0	(2.4–6.6)
Pennsylvania	126	2.8	(2.0–3.8)	166	3.3	(2.6–4.4)
Rhode Island	27	7.1	(4.1–11.8)	23	5.4	(4.0–7.2)
South Carolina	76	4.9	(2.4–9.4)	35	2.0	(1.1–3.5)
South Dakota	11	3.9	(2.6–6.0)	11	3.5	(2.2–5.6)
Tennessee	94	4.3	(2.5–7.2)	73	3.0	(1.9–4.6)
Texas	230	2.7	(2.0–3.8)	323	3.7	(2.8–4.8)
Utah	52	5.6	(3.6–8.5)	76	8.1	(5.4–11.9)
Vermont	10	4.2	(2.3–7.8)	9	3.6	(2.3–5.5)
Virginia	79	2.9	(1.6–5.2)	81	2.7	(1.5–4.6)
Washington	85	3.5	(2.1–5.7)	146	5.8	(3.8–8.6)
West Virginia	25	3.7	(2.3–5.6)	37	5.1	(2.9–8.9)
Wisconsin	102	4.9	(3.0–8.0)	82	3.8	(2.6–5.5)
Wyoming	12	6.0	(3.3–10.7)	8	4.1	(2.4-7.0)
Total	3,789	3.5	(3.2-3.8)	4,571	3.9	(3.7-4.2)

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded

[§] Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

TABLE 5. Annual average estimated number* and percentage of adults aged ≥ 18 years who had seriously considered suicide during the previous year, by state and race/ethnicity — National Survey on Drug Use and Health, United States, 2008–2009[†]

	White	e, non-Hi	spanic	Black	k, non-His	spanic		Hispanic	§	Asian	, non-Hi	panic
Region/State	No.	%	(95%CI)	No.	%	(95%CI)	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region [¶]												
Midwest	1,740	4.3	(3.9-4.7)	192	4.2	(3.2-5.5)	127	4.8	(3.1-7.3)	24	2.2	(1.2-3.8)
Northeast	1,161	3.8	(3.4–4.3)	128	3.0	(2.1–4.1)	131	3.0	(2.1–4.3)	27	1.2	(0.7–2.4)
South	1,787	3.3	(3.0–3.7)	470	3.2	(2.7–3.9)	284	2.6	(2.0–3.3)	51	2.6	(0.7-2.4) (1.3-5.1)
West	1,357	4.5	(3.9–5.1)	121	5.0	(3.2–7.6)	390	3.1	(2.4–4.0)	107	2.3	(1.5–3.4)
	1,337	4.5	(3.9-3.1)	121	5.0	(3.2-7.0)	390	3.1	(2.4-4.0)	107	2.3	(1.3-3.4)
State												
Alabama	75	3.0	(2.0-4.6)	33	3.8	(1.5-9.5)	_**	_	_	_	_	
Alaska	14	4.0	(2.7–6.1)	_	_	_		_		_	_	_
Arizona	144	4.7	(3.1–7.2)	_	_	_	38	3.1	(1.4–6.6)	_	_	_
Arkansas	79	4.8	(3.2–7.1)	22	7.5	(3.8–14.2)	_	_			_	_
California	439	3.5	(2.6–4.7)	85	5.0	(3.0-8.4)	255	2.9	(2.1–4.0)	74	2.2	(1.4-3.6)
Colorado	130	4.8	(2.8–8.0)	_	_		19	3.0	(1.1–8.4)	_	_	_
Connecticut	94	4.6	(3.2-6.6)	2	8.0	(0.3-2.5)	17	6.3	(3.0-12.7)		_	2.
Delaware	12	2.5	(1.5–3.9)	3	2.4	(0.8-6.6)	<1	0.7	(0.2-2.7)	-	_	
District of Columbia	5	2.9	(1.5–5.6)	15	6.1	(3.6-10.2)	1	1.9	(0.6–6.0)	_		
Florida	382	4.3	(3.3-5.7)	78	4.1	(2.7-6.2)	38	1.3	(0.8-2.3)	3	1.2	(0.4-3.8)
Georgia	89	2.1	(1.1-4.1)	42	2.1	(1.3-3.5)	_	_	-	-	-	-
Hawaii	4	1.7	(0.8-3.5)	_	_	_	3	4.4	(2.5-7.6)	12	3.1	(1.3-7.2)
Idaho	59	6.1	(4.3-8.5)	_	_	<u></u>		_		_	_	·
Illinois	297	4.6	(3.8-5.6)	38	2.9	(1.8-4.7)	47	3.7	(2.2-6.2)	6	1.6	(0.5-5.1)
Indiana	192	4.8	(3.6-6.5)	9	2.4	(1.2-4.7)	_	_	_	_	_	_
Iowa	70	3.4	(2.4-4.8)					· —	 		-	· ·
Kansas	50	2.9	(1.9-4.4)	-	-	_	3	1.7	(0.5-5.3)	_	_	· · ·
Kentucky	104	3.7	(2.4-5.6)	4	1.9	(0.7-4.7)	\rightarrow	_	-	-	-	-
Louisiana	52	2.5	(1.5-3.9)	26	2.7	(1.3-5.7)	_	_	_	_	_	-
Maine	51	5.1	(3.6-7.3)	1	-	-	_	_	_	,	_	7
Maryland	80	3.1	(1.7-5.9)	37	3.1	(1.4-6.8)	—		_	2	_	-
Massachusetts	141	3.5	(2.3-5.3)	_	-	_	_	_	_	_	_	_
Michigan	286	4.8	(4.0-5.8)	29	3.0	(1.9-4.9)	7	2.5	(1.1-5.6)	7	3.8	(1.5 - 9.4)
Minnesota	157	4.6	(3.2-6.5)	-	_	_	_	_	_	_	_	_
Mississippi	30	2.3	(1.6-3.2)	17	2.4	(1.4-4.0)	\rightarrow	-	-	-	-	_
Missouri	167	4.5	(3.1-6.6)		_	_			_	_	_	-
Montana	27	4.1	(2.9-5.7)	_	-	_	_	_	_	-	_	_
Nebraska	47	4.1	(2.7-6.2)	2000	-	_		s -	_		-	.—.
Nevada	88	7.4	(5.3-10.3)	_	-	_	_	-	_	-	_	-
New Hampshire	28	3.0	(2.2-4.0)	-	_	_		-	_	_	-	
New Jersey	158	3.8	(2.3-6.3)	15	1.9	(0.9-3.9)	11	1.1	(0.5-2.7)	_		-
New Mexico	24	3.7	(2.3-5.9)	_	_	_	19	3.1	(1.8-5.1)	_	_	-
New York	390	4.2	(3.4-5.3)	57	2.7	(1.5-4.8)	66	2.9	(1.6-5.3)	19	1.8	(0.8-4.2)
North Carolina	199	4.2	(2.5-6.9)	20	1.4	(0.6-3.4)		_	_	_		_
North Dakota	12	2.7	(1.7-4.4)	_	_		_		_		_	_
Ohio	301	4.1	(3.4-5.0)	49	5.3	(3.5-7.9)	_	-	_	_	_	-
Oklahoma	61	3.1	(1.9-4.9)	_	_	-	_	_	-	_	_	_
Oregon	110	4.6	(2.9-7.0)	_	_	_	3	1.2	(0.4-3.2)	_	_	_
Pennsylvania	234	3.0	(2.3–3.7)	37	4.2	(2.5-6.8)	10	2.7	(1.3–5.4)	-		
Rhode Island	47	7.1	(5.1–9.7)	_	_		_	_			_	_
South Carolina	88	3.9	(2.1-7.0)	22	2.5	(1.2-5.1)	1	0.6	(0.2-2.0)	-	_	_
South Dakota	18	3.5	(2.3–5.1)		_	_		_		_	_	.—.
Tennessee	122	3.3	(2.2-4.8)	35	4.9	(2.1-11.5)		7	_	1	_	
Texas	251	2.9	(2.1–3.9)	79	4.1	(2.6–6.5)	189	3.3	(2.5-4.4)	_	_	_
Utah	117	7.5	(5.3–10.5)	_	_		2	1.2	(0.3–4.8)	_	_	-
Vermont	18	3.9	(2.5–5.9)	_	_	_	_	_		_	_	-
Virginia	102	2.6	(1.6–3.9)	32	3.0	(1.1-8.3)	_	_	_	_	_	_
Washington	185	4.8	(3.3–6.9)	_	_	-	10	2.5	(0.9-6.9)	_	_	_
West Virginia	57	4.3	(2.9–6.4)	_	_	_	_	_		_	_	_
Wisconsin	143	3.8	(2.6–5.5)	_	_	_		_	-	_	_	
Wyoming	17	4.9	(3.1–7.6)		_	_	2	8.6	(4.3-16.3)	_	_	_
										<u> </u>		
Total	6,044	3.9	(3.7-4.1)	911	3.5	(3.1-4.0)	933	3.0	(2.6-3.6)	208	2.1	(1.6-2.8)

 $\textbf{Abbreviation:} \ \mathsf{CI} = \mathsf{confidence} \ \mathsf{interval}.$

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

[§] Persons of Hispanic origin can be of any race.

^{**}Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

^{***} Low precision; no estimate reported. According to the suppression criteria, prevalence estimates are suppressed if any of the following occurred: the prevalence estimate is < 0.005% or >99.995%; the relative standard error of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is ≤50%; the relative standard error of the negative of the natural logarithm of (1-p) is >0.175 if the prevalence is >50%; the actual sample size is <100; or the effective sample size (defined as the actual sample size divided by the design effect) is <68.

TABLE 6. Annual average estimated number* and percentage of adults aged ≥18 years who had made suicide plans during the previous year, by state and age group — National Survey on Drug Use and Health, United States, 2008–2009[†]

		18–29 yr	s		≥30yrs	
Region/State	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region [§]						
Midwest	221	2.0	(1.7-2.4)	377	1.0	(0.8-1.2)
Northeast	138	1.5	(1.2-1.9)	220	0.7	(0.5-1.0)
South	258	1.4	(1.2-1.7)	476	0.7	(0.6-0.9)
West	203	1.7	(1.4-2.0)	369	0.9	(0.7-1.2)
State						
Alabama	16	2.1	(1.2-3.5)	22	0.8	(0.3-2.3)
Alaska	2	1.8	(0.6-5.6)	3	0.7	(0.3-1.7)
Arizona	12	1.1	(0.5-2.3)	30	0.8	(0.4-1.6)
Arkansas	16	3.5	(1.7–7.2)	14	0.9	(0.3-2.2)
California	106	1.6	(1.2-2.3)	138	0.7	(0.4-1.2)
Colorado	15	1.8	(1.0–3.2)	34	1.2	(0.4-3.7)
Connecticut	9	1.6	(0.8-3.3)	23	1.1	(0.4-2.9)
Delaware	2	1.3	(0.6–2.5)	1	0.3	(<0.1-1.6)
District of Columbia	1	1.0	(0.5–2.0)	3	0.8	(0.2-2.8)
Florida	41	1.6	(1.1–2.3)	134	1.2	(0.8–1.8)
Georgia	6	0.4	(0.1–1.1)	13-1	_	
Hawaii	5	2.7	(1.2–6.1)	5	0.6	(0.2-1.8)
Idaho	5	1.7	(1.0-2.9)	7	0.9	(0.4–1.9)
Illinois	33	1.5	(1.1–2.0)	64	0.9	(0.5–1.4)
Indiana	30	2.9	(1.7–4.9)	34	0.9	(0.4–2.3)
Iowa	8	1.7	(0.7-4.0)	15	0.8	(0.4–2.0)
Kansas	6	1.2	(0.6–2.5)	8	0.5	(0.2–1.4)
Kentucky	8	1.2	(0.6–2.2)	18	0.7	(0.3–1.7)
Louisiana	12	1.7	(0.8–3.5)	20	0.8	(0.3–1.7)
Maine	4	1.9	(0.9–4.3)	18	2.2	(1.0-4.9)
Maryland	13	1.4	(0.7–2.7)	14	0.4	(0.1-1.4)
Massachusetts	26	2.4	(1.3–4.4)	28	0.7	(0.1–1.4)
Michigan	46	2.4	(2.1–3.7)	71	1.2	(0.3-2.0)
	22	2.5	10 harmon and market	37	1.2	(0.5–2.0)
Minnesota Mississippi	10	2.3	(1.3–4.8)	2	0.1	(<0.1–0.6)
The state of the s			(1.4–3.6)	21		
Missouri	19 4	1.9	(0.7–5.5)	7	0.6	(0.2–1.9)
Montana		2.5	(1.2–5.3)		1.2	(0.6–2.1)
Nebraska	6	2.0	(1.1–3.6)	7	0.7	(0.3–1.9)
Nevada	14	3.3	(1.5–7.2)	29	1.9	(0.9–4.1)
New Hampshire	3	1.5	(0.8–2.9)	7	8.0	(0.4–1.6)
New Jersey	19	1.4	(0.6–3.0)	40	0.8	(0.2–2.9)
New Mexico	7	1.9	(1.1–3.4)	7	0.6	(0.3–1.3)
New York	49	1.5	(1.0–2.2)	41	0.4	(0.2-0.8)
North Carolina	17	1.1	(0.5–2.6)	57	1.1	(0.5–2.1)
North Dakota	1	1.0	(0.4–2.5)	3	8.0	(0.3–2.2)
Ohio	34	1.9	(1.3–2.6)	84	1.2	(0.7-2.2)
Oklahoma	7	1.3	(0.5-3.0)	17	8.0	(0.3-2.1)
Oregon	11	1.9	(1.0-3.4)	31	1.3	(0.5-3.4)
Pennsylvania	20	1.0	(0.7–1.5)	40	0.5	(0.3-1.0)
Rhode Island	5	2.8	(1.3–5.8)	17	2.8	(1.3-5.9)
South Carolina	8	1.1	(0.6-2.0)	37	1.4	(0.4-4.7)
South Dakota	3	2.1	(1.3-3.4)	2	0.5	(0.1-1.6)
Tennessee	18	1.8	(0.9–3.7)	25	0.7	(0.3-1.7)
Texas	65	1.6	(1.1–2.3)	84	0.6	(0.3-1.3)
Utah	9	1.7	(0.9-3.0)	19	1.4	(0.6-3.5)
Vermont	3	2.9	(1.6-5.2)	5	1.2	(0.6-2.6)
Virginia	13	1.0	(0.6-1.9)	18	0.4	(0.1-1.7)
Washington	11	1.0	(0.5-2.2)	56	1.5	(0.6-3.6)
West Virginia	4	1.4	(0.8-2.5)	9	0.8	(0.3-2.5)
Wisconsin	15	1.7	(0.9–3.0)	29	0.9	(0.4–1.9)
Wyoming	2	2.3	(1.3–4.1)	4	1.1	(0.5–2.7)
Total	821	1.6	(1.5–1.8)	1,442	0.8	(0.7-0.9)

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

[§] Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

Mexico, Oregon, Utah, Washington, and Wyoming.

Low precision; no estimate reported. According to the suppression criteria, prevalence estimates are suppressed if any of the following occurred: the prevalence estimate is < 0.005% or > 99.995%; the relative standard error of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is > 0.175 if the prevalence is $\le 50\%$; the relative standard error of the negative of the natural logarithm of (1-p) is > 0.175 if the prevalence is > 50%; the actual sample size is < 100; or the effective sample size (defined as the actual sample size divided by the design effect) is < 68.

TABLE 7. Annual average estimated number* and percentage of adults aged ≥18 years who had made suicide plans during the previous year, by state and sex — National Survey on Drug Use and Health, United States, 2008–2009[†]

	S	Male			Female	
Region/State	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region§						
Midwest	278	1.2	(0.9-1.5)	320	1.3	(1.0-1.5)
Northeast	165	0.8	(0.6-1.1)	194	0.9	(0.6-1.2)
South	352	0.9	(0.7-1.1)	382	0.9	(0.7-1.1)
West	251	1.0	(0.7-1.4)	322	1.2	(0.9-1.6)
State						
Alabama	16	1.0	(0.4-2.5)	22	1.2	(0.5-2.8)
Alaska	2	0.8	(0.3-2.4)	3	1.1	(0.4-2.9)
Arizona	22	0.9	(0.4-2.4)	21	0.9	(0.4-1.9)
Arkansas	11	1.1	(0.4-3.1)	19	1.8	(0.9-3.5)
California	117	0.9	(0.5-1.5)	127	0.9	(0.6-1.5)
Colorado	30	1.6	(0.4-6.0)	20	1.1	(0.6-2.0)
Connecticut	14	1.1	(0.3-3.5)	18	1.3	(0.6-3.1)
Delaware	1	0.5	(0.1-2.6)	2	0.5	(0.2-1.1)
District of Columbia	3	1.2	(0.3-4.3)	1	0.6	(0.2-1.3)
Florida	65	1.0	(0.6-1.5)	110	1.5	(1.0-2.3)
Georgia	3	0.1	(<0.1-0.4)	3	0.1	(<0.1-0.4)
Hawaii	4	0.9	(0.4-2.3)	5	1.1	(0.4-2.8)
Idaho	5	0.9	(0.4-2.2)	7	1.2	(0.6-2.4)
Illinois	50	1.1	(0.6-1.9)	47	1.0	(0.7-1.4)
Indiana	25	1.1	(0.5-2.5)	39	1.6	(0.7-3.3)
lowa	5	0.5	(0.1-1.8)	17	1.5	(0.7-3.4)
Kansas	8	0.8	(0.2-2.3)	6	0.6	(0.3-1.1)
Kentucky	11	0.7	(0.3-1.9)	15	0.9	(0.4-2.0)
Louisiana	21	1.4	(0.6-3.1)	12	0.7	(0.2-2.2)
Maine	10	2.1	(0.6-6.9)	12	2.2	(1.0-4.6)
Maryland	15	0.7	(0.3-2.1)	12	0.6	(0.3-1.2)
Massachusetts	17	0.7	(0.2-2.1)	37	1.4	(0.7-3.2)
Michigan	51	1.4	(0.9-2.2)	67	1.7	(1.1-2.6)
Minnesota	21	1.1	(0.4-2.8)	38	1.9	(0.8-4.5)
Mississippi	3	0.3	(0.2-0.5)	10	0.9	(0.5-1.6)
Missouri	26	1.2	(0.4-3.4)	14	0.6	(0.2-2.0)
Montana	5	1.3	(0.6-3.0)	6	1.5	(0.6-3.6)
Nebraska	4	0.7	(0.2-1.8)	9	1.4	(0.7-2.5)
Nevada	13	1.4	(0.6-3.4)	29	3.1	(1.6-5.9)
New Hampshire	5	1.1	(0.6-1.9)	4	0.9	(0.4-2.0)
New Jersey	17	0.6	(0.1-2.1)	41	1.2	(0.4-4.0)
New Mexico	3	0.5	(0.2-1.1)	10	1.4	(0.8-2.3)
New York	48	0.7	(0.3-1.4)	43	0.6	(0.3-0.9)
North Carolina	44	1.4	(0.6-3.0)	30	0.8	(0.3-2.3)
North Dakota	1	0.6	(0.2-1.8)	3	1.1	(0.4-2.9)
Ohio	56	1.4	(0.6-2.9)	61	1.4	(0.9-2.1)
Oklahoma	12	1.0	(0.3-3.2)	12	0.9	(0.4-1.9)
Oregon	24	1.7	(0.6-5.0)	18	1.2	(0.5-3.1)
Pennsylvania	33	0.7	(0.4-1.4)	27	0.6	(0.3-0.9)
Rhode Island	16	4.1	(1.8-8.6)	7	1.6	(0.8-3.4)
South Carolina	40	2.6	(0.8-7.7)	5	0.3	(0.1-0.7)
South Dakota	4	1.4	(0.7-3.0)	1	0.3	(0.1-0.7)
Tennessee	21	1.0	(0.4-2.2)	22	0.9	(0.3-2.4)
Texas	59	0.7	(0.4–1.2)	90	1.0	(0.6-1.7)
Utah	11	1.1	(0.4-2.9)	18	1.9	(1.0-3.6)
Vermont	4	1.6	(0.7-3.6)	4	1.5	(0.7-3.0)
Virginia	24	0.9	(0.3-2.9)	8	0.3	(0.1-0.6)
Washington	11	0.5	(0.2-1.3)	56	2.2	(1.0-4.9)
West Virginia	4	0.6	(0.2-2.1)	9	1.2	(0.4-3.4)
Wisconsin	27	1.3	(0.6-2.8)	18	8.0	(0.4-1.6)
Wyoming	3	1.7	(0.7-4.2)	2	1.0	(0.5-2.1)
Total	1,045	1.0	(0.8-1.1)	1,218	1.0	(0.9-1.2)

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information

[§] Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

TABLE 8. Annual average estimated number* and percentage of adults aged ≥ 18 years who had made suicide plans during the previous year, by state and race/ethnicity — National Survey on Drug Use and Health, United States, 2008–2009[†]

	Whi	te, non-His	spanic	Blac	ck, non-His	panic		Hispanic [§]	<u> </u>	Asia	an, non-His	panic
Region/State	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region [¶]												
Midwest	481	1.2	(1.0-1.4)	45	1.0	(0.6-1.5)	51	1.9	(0.8-4.5)	6	0.6	(0.2-1.5)
Northeast	249	8.0	(0.6-1.1)	43	1.0	(0.6-1.7)	56	1.3	(0.7-2.4)	3	0.1	(<0.1-0.6)
South	484	0.9	(0.7-1.1)	138	0.9	(0.7-1.3)	71	0.7	(0.4-1.2)	10	0.5	(0.1-1.8)
West	402	1.3	(1.0–1.7)	36	1.5	(0.6-3.7)	88	0.7	(0.5-1.0)	18	0.4	(0.2-0.7)
State			,			,			,			,
Alabama	34	1.4	(0.7-2.8)	4	0.5	(0.1-1.5)	**	-		_	_	_
Alaska	4	1.3	(0.5-3.0)		_		_	-	-	_		-
Arizona	31	1.0	(0.5–1.9)	_	_	_	8	0.7	(0.2-2.3)	_	_	
Arkansas	20	1.2	(0.6–2.2)	11	3.5	(1.3-9.2)	_	_		_		_
California	128	1.0	(0.6–1.7)	33	2.0	(0.7–5.2)	62	0.7	(0.4-1.2)	15	0.5	(0.2-0.9)
Colorado	41	1.5	(0.6–3.9)	_	_	(o., o.,	6	0.9	(0.3–2.7)	_	_	(0.2 0.5)
Connecticut	18	0.9	(0.4-2.1)	1	0.3	(0.1-1.4)	14	5.0	(1.8–12.7)	_	_	_
Delaware	3	0.5	(0.2–1.6)	<1	0.2	(<0.1–1.3)	_	_	(1.0 12.7)	_	_	
District of Columbia	1	0.6	(0.2–1.0)	3	1.3	(0.4–3.9)	<1	<0.1	(<0.1–0.2)		_	_
Florida	124				1.2		9			-	=	
	124	1.4	(0.8–2.5)	22 5		(0.6–2.3)		0.3	(0.1–0.8)	_	_	_
Georgia		0.0	(<0.1-0.1)		0.3	(0.1-0.8)	,		_	_	_	
Hawaii	2	0.7	(0.2–2.5)	_	_	_	_	11	(0.2.26)	1	0.3	(0.2-0.6)
Idaho	10	1.1	(0.6–2.0)	_	_		1	1.1	(0.3–3.6)	_	_	
Illinois	66	1.0	(0.7–1.5)	9	0.7	(0.3–1.6)	17	1.4	(0.5-3.9)	<1	<0.1	(<0.1–0.9)
Indiana	61	1.5	(0.9–2.7)	_	-	_	-	_	_	_	_	_
lowa	16	0.8	(0.4–1.6)	_	-		_	-	_	_	-	ş -
Kansas	13	8.0	(0.4-1.6)	<1	0.5	(0.1-1.5)	<1	< 0.1	(<0.1–0.9)	_	_	_
Kentucky	23	8.0	(0.4-1.5)	_	-	_	_	-	·—	_	_	-
Louisiana	20	1.0	(0.4-2.2)	11	1.2	(0.3-4.1)	_	—	· ·	_	-	—,
Maine	22	2.2	(1.1-4.4)	-	-				* <u></u> *	_	_	_
Maryland	9	0.4	(0.1-1.0)	13	1.1	(0.4-3.5)	-		: :	-	-	
Massachusetts	41	1.0	(0.5-2.2)	_	_	_	_	_	_	_	_	_
Michigan	96	1.6	(1.1-2.3)	15	1.5	(0.7-3.3)	2	0.7	(0.2-2.3)	_	_	—
Minnesota	57	1.7	(0.9-3.2)			_	0.	—,			-	—2
Mississippi	8	0.6	(0.3-1.1)	5	0.7	(0.3-1.5)	_		_	_	_	_
Missouri	35	1.0	(0.4-2.4)	4	0.9	(0.3-2.8)	-	_	_	_	_	1-0
Montana	8	1.2	(0.7-2.1)	-	-	=	-	-	=	\rightarrow	-	_
Nebraska	12	1.1	(0.6-1.9)	_	_	_	-	_	_		_	—
Nevada	39	3.3	(1.8-6.0)	_		_	2	0.6	(0.1-2.6)	_	-	—,
New Hampshire	9	0.9	(0.5-1.5)	_	_	_	_	_	_	_	_	_
New Jersey	44	1.0	(0.3-3.6)	9	1.0	(0.3-3.6)	5	0.5	(0.1-2.0)	_	_	_
New Mexico	6	0.9	(0.5-1.5)	-	_	_	5	0.9	(0.4-1.8)	\rightarrow	_	_
New York	42	0.5	(0.3-0.8)	20	0.9	(0.4-2.2)	23	1.0	(0.3–3.3)	3	0.3	(0.1-1.3)
North Carolina	58	1.2	(0.6-2.6)	7	0.5	(0.1–2.2)	_	_		_	_	
North Dakota	4	0.8	(0.3-2.1)		_	_	_	_	-	_	-	_
Ohio	82	1.1	(0.8–1.6)	10	1.1	(0.4-2.8)	-	_	sa	_	_	
Oklahoma	18	0.9	(0.4-2.2)	_	_	_	_	_	_	_	_	_
Oregon	41	1.7	(0.8–3.6)	_	_	_	1	0.4	(0.1-2.2)	_	_	_
Pennsylvania	47	0.6	(0.4–1.0)	9	1.0	(0.4-2.6)	4	1.1	(0.3–3.3)	_	_	_
Rhode Island	20	3.0	(1.6–5.7)	_	_	_			(0.5 0.5)	_		_
South Carolina	38	1.7	(0.5-5.6)	6	0.7	(0.3-1.4)	<1	0.4	(0.1-1.9)		_	
South Dakota	4	0.8	(0.3–3.6)	_	- U.7	(0.3-1.4)	_	- U.4 	(0.1-1.9)		_	
Tennessee	26	0.8	(0.4–1.0)	10	1.4	(0.4–4.7)	_	_	_		_	
Texas	82	0.7	(0.5–1.7)	17	0.9	(0.4-4.7) (0.3-2.4)	43	0.7	— (0.3–1.7)	_	_	_
Utah	82 27	1.7		_	U.9 —	(0.3–2.4)	43 —	U.7 —	(0.3-1.7)		_	
	7		(0.9–3.3)							_		_
Vermont		1.6	(0.9–2.8)	16	1.5	(0.4.6.3)	_	_	_	_	_	_
Virginia	7	0.2	(0.1–0.4)	16	1.5	(0.4-6.3)		·	_	_	_	
Washington	60	1.5	(0.7–3.5)	_	_	_	_	·	_	_	_	_
West Virginia	12	0.9	(0.4–2.1)	_	-	_	2	_	-	_	_	_
Wisconsin	36	1.0	(0.5–1.8)	-	_	-	_	_	-	-	_	-
Wyoming	5	1.3	(0.6–2.7)	_	_		<1	1.3	(0.4–4.6)	_		
Total Abbreviation: CI = confidence	1,616	1.0	(0.9–1.2)	262	1.0	(0.8–1.3)	267	0.9	(0.6-1.2)	37	0.4	(0.2-0.6)

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

[§] Persons of Hispanic origin can be of any race.

Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

^{**} Low precision; no estimate reported. According to the suppression criteria, prevalence estimates are suppressed if any of the following occurred: the prevalence estimate is < 0.005% or >99.995%; the relative standard error of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is ≤50%; the relative standard error of the negative of the natural logarithm of (1-p) is >0.175 if the prevalence is >50%; the actual sample size is <100; or the effective sample size (defined as the actual sample size divided by the design effect) is <68.

TABLE 9. Annual average estimated number* and percentage of adults aged ≥18 years who had attempted suicide during the previous year, by state and age group — National Survey on Drug Use and Health, United States, 2008–2009[†]

		18–29 yr	5		≥30yrs	
Region/State	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region [§]						
Midwest	115	1.0	(0.9-1.3)	156	0.4	(0.3-0.6)
Northeast	77	0.9	(0.7-1.1)	105	0.3	(0.2-0.6)
South	160	0.9	(0.7–1.1)	193	0.3	(0.2-0.4)
West	125	1.0	(0.8-1.4)	127	0.3	(0.2-0.5)
tate						
Alabama	6	8.0	(0.3–2.0)		_	_
Alaska	<1	0.2	(0.1–0.7)	1	0.2	(<0.1–1.3
Arizona	9	0.9	(0.3–2.2)	19	0.5	(0.2–1.5)
Arkansas	11	2.4	(1.0-6.0)	8	0.5	(0.1-2.1)
California	66	1.0	(0.7–1.6)	52	0.3	(0.1–0.6)
Colorado	10	1.2	(0.6–2.4)	5	0.2	(<0.1–0.7
Connecticut	8	1.5	(0.7–3.2)	18	8.0	(0.3-2.5)
Delaware	1	0.6	(0.2–1.5)		_	
District of Columbia	1	0.5	(0.2–1.3)	1	0.3	(0.1–1.1)
Florida	23	0.9	(0.5–1.4)	50	0.4	(0.2-0.9)
Georgia	6	0.4	(0.1–1.4)	_	_	
Hawaii	1	0.6	(0.3–1.2)	1	0.1	(<0.1-0.5
Idaho	2	0.7	(0.3–1.6)	4	0.5	(0.2–1.4)
Illinois	18	0.8	(0.5–1.2)	26	0.4	(0.2-0.7)
Indiana	17	1.6	(0.8–3.4)	_	_	_
lowa	1	0.1	(<0.1–0.7)	7	0.4	(0.1-1.2)
Kansas	3	0.6	(0.2-1.9)	7	0.4	(0.1-1.6)
Kentucky	6	0.8	(0.3–1.8)	2	0.1	(<0.1–0.7
Louisiana	8	1.1	(0.5–2.4)	4	0.2	(<0.1-0.7
Maine	2	1.1	(0.3-3.3)	6	8.0	(0.2-2.8)
Maryland	5	0.6	(0.3–1.4)	6	0.2	(<0.1–1.3
Massachusetts	10	0.9	(0.4–1.8)	_	_	_
Michigan	26	1.6	(1.1–2.2)	32	0.6	(0.3-1.0)
Minnesota	5	0.6	(0.2–1.5)	24	0.8	(0.2-2.6)
Mississippi	6	1.4	(0.8-2.6)	3	0.2	(<0.1–1.1
Missouri	7	0.7	(0.2–2.3)	8	0.2	(<0.1–1.6
Montana	1	0.7	(0.3–1.7)	1	0.1	(<0.1-1.1
Nebraska	4	1.4	(0.7–2.9)	3	0.2	(<0.1–1.3
Nevada	7	1.8	(0.7-4.4)	3	0.2	(<0.1–1.3
New Hampshire	3	1.3	(0.6-2.8)	1	0.1	(<0.1-0.8
New Jersey	15	1.1	(0.5–2.3)	-	_	-
New Mexico	5	1.6	(0.8-3.2)	4	0.4	(0.1–1.2)
New York	26	0.8	(0.5–1.2)	30	0.3	(0.1-0.7)
North Carolina	11	0.7	(0.3-2.1)	22	0.4	(0.1-1.2)
North Dakota	1	1.1	(0.4-2.8)	<1	< 0.1	(<0.1-0.4
Ohio	23	1.3	(0.8-1.9)	37	0.5	(0.3-1.0)
Oklahoma	9	1.6	(0.8-3.3)	2	0.1	(<0.1-0.8
Oregon	3	0.5	(0.2-1.3)	10	0.4	(0.1-1.6)
Pennsylvania	9	0.5	(0.2-0.8)	17	0.2	(0.1-0.6)
Rhode Island	4	2.2	(0.8-5.6)	8	1.3	(0.4-4.1)
South Carolina	5	0.8	(0.3-1.7)	7	0.3	(0.1-1.3)
South Dakota	2	1.1	(0.6-2.1)	1	0.1	(<0.1-0.9
Tennessee	6	0.6	(0.2-2.2)	6	0.2	(<0.1-0.8
Texas	45	1.1	(0.7-1.8)	64	0.5	(0.2-1.0)
Utah	7	1.2	(0.6-2.5)	2	0.2	(<0.1-1.3
Vermont	1	1.4	(0.7-2.9)	1	0.3	(0.1-1.3)
Virginia	6	0.5	(0.2-1.2)	11	0.2	(<0.1-1.8
Washington	12	1.1	(0.4-3.2)	25	0.6	(0.2-2.6)
West Virginia	3	1.2	(0.5-2.6)	7	0.6	(0.2-2.3
Wisconsin	9	0.9	(0.4-2.1)	13	0.4	(0.1–1.1)
Wyoming	1	1.2	(0.5-3.2)	1	0.2	(0.1–1.1
otal	477	1.0	(0.8-1.1)	581	0.3	(0.3-0.4

 $\textbf{Abbreviation:} \ \mathsf{CI} = \mathsf{confidence} \ \mathsf{interval}.$

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

[§] Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

[¶] Low precision; no estimate reported. According to the suppression criteria, prevalence estimates are suppressed if any of the following occurred: the prevalence estimate is < 0.005% or >99.995%; the relative standard error of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is ≤50%; the relative standard error of the negative of the natural logarithm of (1-p) is >0.175 if the prevalence is >50%; the actual sample size is <100; or the effective sample size (defined as the actual sample size divided by the design effect) is <68.

TABLE 10. Annual average estimated number* and percentage of adults aged ≥18 years who had attempted suicide during the previous year, by state and sex — National Survey on Drug Use and Health, United States, 2008-2009+

		Male			Female	•
Region/State	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region§						
Midwest	114	0.5	(0.3-0.7)	157	0.6	(0.5-0.8)
Northeast	75	0.4	(0.2-0.6)	107	0.5	(0.3-0.8)
South	154	0.4	(0.3-0.5)	198	0.5	(0.3-0.6)
West	99	0.4	(0.2-0.6)	154	0.6	(0.4-0.8)
State						Ç,
Alabama	3	0.2	(0.1-0.6)	3	0.2	(0.1-0.5)
Alaska	<1	< 0.1	(<0.1–0.2)	1	0.3	(0.1–1.7)
Arizona	10	0.4	(0.1–2.1)	19	0.8	(0.3–1.8)
Arkansas	6	0.6	(0.1–2.9)	13	1.2	(0.5–3.0)
California	51	0.4	(0.2–0.8)	67	0.5	(0.3-0.9)
Colorado	5	0.3	(0.1–0.8)	10	0.5	(0.2–1.2)
Connecticut	8	0.6	(0.2–2.1)	18	1.3	(0.5–3.4)
Delaware	¶*	-	(0.2 2.1)	1	0.2	(0.1–0.6)
District of Columbia	<1	<0.1	(<0.1-0.3)	2	0.6	(0.2–1.5)
Florida	29	0.4	(0.2–1.1)	44	0.6	(0.3–1.1)
Georgia	1	0.4	(<0.1–0.1)	5	0.0	(<0.1–0.6)
Hawaii	<1	<0.1	15 N 7 M 10 M 10 M 10 M	2	0.1	1000 0000 00 00 0000
			(<0.1–0.3)			(0.1–0.9)
Idaho	2	0.4	(0.1–1.4)	4	0.8	(0.3–1.9)
Illinois	20	0.4	(0.2–1.0)	24	0.5	(0.3–0.8)
Indiana	4	0.2	(0.1–0.7)	13	0.5	(0.2–1.3)
lowa	2	0.2	(<0.1–1.5)	5	0.4	(0.1–1.5)
Kansas	6	0.6	(0.2–2.3)	3	0.3	(0.1–1.2)
Kentucky	5	0.3	(0.1–1.0)	3	0.2	(0.1–0.5)
Louisiana	8	0.5	(0.2–1.3)	5	0.3	(0.1–0.8)
Maine	5	1	(0.2-4.8)	3	0.6	(0.2-1.9)
Maryland	2	0.1	(<0.1–0.4)	9	0.4	(0.1-1.6)
Massachusetts	1	0.1	(<0.1-0.4)	8	0.3	(0.1-0.7)
Michigan	31	0.9	(0.5-1.5)	27	0.7	(0.4-1.1)
Minnesota	8	0.4	(0.1-2.3)	22	1.1	(0.3-3.5)
Mississippi	3	0.3	(0.1–0.7)	6	0.5	(0.2-1.4)
Missouri	5	0.2	(<0.1–1.1)	10	0.4	(0.1-2.1)
Montana	2	0.5	(0.2-1.5)	<1	< 0.1	(<0.1-0.2)
Nebraska	1	0.2	(0.1-0.8)	5	0.8	(0.4-1.6)
Nevada	3	0.3	(0.1-1.4)	7	0.8	(0.3-2.0)
New Hampshire	2	0.4	(0.1-1.1)	2	0.3	(0.1-1.0)
New Jersey	5	0.2	(0.1-0.5)	34	1.0	(0.2-4.2)
New Mexico	4	0.6	(0.2-1.9)	6	0.7	(0.4-1.5)
New York	28	0.4	(0.1-1.1)	28	0.4	(0.2-0.6)
North Carolina	19	0.6	(0.2-1.6)	14	0.4	(0.1-1.4)
North Dakota	<1	0.2	(0.1–0.7)	1	0.4	(0.1–1.3)
Ohio	19	0.5	(0.2-0.9)	42	0.9	(0.6-1.6)
Oklahoma	3	0.2	(0.1–0.7)	9	0.6	(0.3–1.5)
Oregon	8	0.6	(0.1–2.3)	5	0.3	(0.1–1.1)
Pennsylvania	15	0.3	(0.1–0.9)	10	0.2	(0.1–0.4)
Rhode Island	9	2.3	(0.8–6.5)	3	0.8	(0.3–2.0)
South Carolina	10	0.6	(0.2–2.1)	3	0.2	(<0.1-0.6)
South Dakota	2	0.6	(0.2–1.4)	1	0.2	(0.1–0.5)
Tennessee	8	0.4	(0.1–1.4)	5	0.2	(<0.1–0.7)
Texas	38	0.5	(0.2–0.8)	71	0.8	(0.5–1.4)
Utah Vermont	6	0.6	(0.2–1.7)	3	0.3	(0.1–0.9)
	1	0.6	(0.2–1.8)	1	0.5	(0.2–1.4)
Virginia	16	0.6	(0.1–2.3)	1	0	(<0.1-0.2)
Washington	7	0.3	(0.1–1.5)	30	1.2	(0.4–3.9)
West Virginia	4	0.6	(0.2–1.8)	6	8.0	(0.2–3.9)
Wisconsin	15	0.7	(0.3–1.9)	6	0.3	(0.1–0.6)
Wyoming	1	0.3	(0.1–1.3)	1	0.6	(0.2–1.7)
Total Abbreviation: Cl – confidence interval	442	0.4	(0.3–0.5)	616	0.5	(0.4–0.6)

In thousands.

 $^{^\}dagger$ Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

S Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

Low precision; no estimate reported. According to the suppression criteria, prevalence estimates are suppressed if any of the following occurred: $the \ prevalence \ estimate \ is < 0.005\% \ or > 99.995\%; the \ relative \ standard \ error \ of \ the \ negative \ natural \ logarithm \ of \ the \ estimated \ proportion \ p$ $(where \ p \ is \ the \ prevalence \ divided \ by \ 100) \ is \ > 0.175 \ if \ the \ prevalence \ is \ \le 50\%; the \ relative \ standard \ error \ of \ the \ negative \ of \ the \ natural \ logarithm$ of (1-p) is > 0.175 if the prevalence is > 50%; the actual sample size is < 100; or the effective sample size (defined as the actual sample size divided in the actual sample size of the actual samplby the design effect) is <68.

TABLE 11. Annual average estimated number* and percentage of adults aged ≥18 years who had attempted suicide during the previous year, by race/ethnicity, region, and state — National Survey on Drug Use and Health, United States, 2008–2009[†]

	Whi	te, non-Hi	spanic	Blac	k, non-His	panic		Hispanic	§	Asia	n, non-Hi	spanic
Region/State	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)	No.	%	(95% CI)
Geographic region [¶]												
Midwest	217	0.5	(0.4-0.7)	34	0.7	(0.5-1.2)	11	0.4	(0.2-0.7)	1	0.1	(<0.1-0.3)
Northeast	117	0.4	(0.2-0.6)	18	0.4	(0.2-0.8)	36	0.8	(0.4-1.8)	4	0.2	(<0.1-0.5)
South	177	0.3	(0.2-0.4)	94	0.6	(0.4-0.9)	46	0.4	(0.2-0.8)	14	0.7	(0.2-3.1)
West	152	0.5	(0.3-0.7)	36	1.5	(0.6-3.8)	51	0.4	(0.3-0.6)	4	0.1	(<0.1-0.2)
State									,			
	4	0.2	(0.1.05)	2	0.3	(0.1.0.0)	**					
Alabama	4	0.2	(0.1–0.5)	2	0.2	(0.1–0.8)		-	_	_	_	_
Alaska	1	0.2	(<0.1–1.3)	_	_	_	_	_	(0.0.0.2)	-		_
Arizona	17	0.6	(0.2–1.7)	_	_	3	9	0.7	(0.2-2.3)	-	-	_
Arkansas	13	0.8	(0.3–2.0)			_	2_3	97	_	_	-	
California	52	0.4	(0.2-0.8)	33	2	(0.7-5.3)	29	0.3	(0.2-0.6)	3	0.1	(<0.1–0.4)
Colorado	9	0.3	(0.1-0.8)	_	_	-	4	0.6	(0.2-2.1)	_	_	-
Connecticut	17	0.8	(0.3-2.3)	1	0.3	(<0.1-1.5)	8	3.0	(1.5-5.8)	_	_	_
Delaware	1	0.1	(<0.1-0.4)	0	0.2	(<0.1-1.4)	-	_	-	_	_	_
District of Columbia	<1	0.1	(<0.1-0.5)	1	0.6	(0.2-1.6)	<1	< 0.1	(<0.1-0.2)	_	_	_
Florida	44	0.5	(0.2-1.0)	5	0.3	(0.1-0.8)	6	0.2	(0.1-0.5)		_	_
Georgia	1	< 0.1	(<0.1-0.1)	5	0.2	(0.1–1.1)	_	_	`	_		_
Hawaii	_	_		_	_	_	_	_	_	1	0.2	(0.1-0.5)
Idaho	6	0.6	(0.3-1.4)		_	_	<1	0.5	(0.1-1.7)		_	(01. 0.0)
Illinois	33	0.5	(0.3-0.9)	5	0.4	(0.1-0.8)	5	0.4	(0.2–0.9)	<1	0.1	(<0.1-0.9)
Indiana	15	0.4	(0.2–0.8)	_		(0.1-0.6)	_	-	(0.2-0.9)	_	-	(<0.1-0.9)
			10.000			_			_	_		
lowa	7	0.3	(0.1–1.0)	_	_	((0.1 1.1)			(:0.1.0.0)	_	_	_
Kansas	9	0.5	(0.2–1.5)	<1	0.2	(<0.1–1.4)	<1	0.1	(<0.1–0.8)	_	<u></u>	_
Kentucky	5	0.2	(0.1–0.5)	2	1	(0.2-4.6)	-	()	_	_	_	_
Louisiana	7	0.3	(0.1–0.8)	5	0.5	(0.2-1.6)	-	_	_	_	<u></u>	_
Maine	8	8.0	(0.3-2.4)	_	_	_	_	_	_	_	_	_
Maryland	3	0.1	(0.1-0.3)	_		1—1	-	-	-	-	-	-
Massachusetts	7	0.2	(0.1-0.4)	_	_	_	1	0.3	(<0.1-2.1)	0	_	_
Michigan	46	0.8	(0.5-1.2)	9	1	(0.5-2.0)	1	0.5	(0.1-2.1)	<1	0.3	(<0.1-2.0)
Minnesota	29	0.8	(0.3-2.3)	_	_	-	_	_	_	-	_	_
Mississippi	5	0.4	(0.1-1.2)	3	0.4	(0.2-1.1)	_	_	_	_	_	_
Missouri	13	0.3	(0.1-1.3)	2	0.3	(0.1-1.3)	_	-	_	_	_	_
Montana	2	0.3	(0.1-0.8)	_	_	_	_	_	_	_	_	_
Nebraska	6	0.5	(0.3-1.0)	_		_	_	1	_	_	-	_
Nevada	7	0.6	(0.2–1.8)	_	_	_	3	0.6	(0.1-2.9)	_	_	_
New Hampshire	2	0.3	(0.1–0.6)	_	_	_	_	_	(011 215)	_	_	_
New Jersey	28	0.7	(0.1–3.7)	5	0.6	(0.1-3.0)	4	0.4	(0.1-1.4)	_	_	_
New Mexico	3	0.5	(0.2–1.3)	_	<u> </u>	(0.1-3.0)	5	0.7	(0.3–2.0)		_	_
					0.5	(0.2.1.2)			S	4		
New York	22	0.2	(0.1–0.5)	10		(0.2–1.3)	17	0.7	(0.2-3.4)	4	0.3	(0.1-1.1)
North Carolina	14	0.3	(0.1–0.9)	12	0.9	(0.3-2.9)	_	-	-	_	_	_
North Dakota	1	0.2	(0.1–0.7)	_	_	-	_	_	-	_		_
Ohio	44	0.6	(0.4–1.0)	14	1.5	(0.6-3.4)	2	1.2	(0.4-4.1)	_	_	_
Oklahoma	6	0.3	(0.1-0.8)	_	_	_	_	-	-	_	_	_
Oregon	13	0.5	(0.2-1.3)	_	_	9. 	<1	< 0.1	(<0.1-0.1)	_	_	_
Pennsylvania	20	0.2	(0.1-0.6)	1	0.1	(<0.1-0.3)	4	1	(0.3-3.3)	_	_	_
Rhode Island	10	1.5	(0.6-3.7)		_	_	_	_	_	_	-	
South Carolina	6	0.3	(<0.1-1.6)	5	0.6	(0.3-1.3)	<1	0.4	(0.1-1.9)	_	_	_
South Dakota	1	0.3	(0.1-0.7)		_	_	_	_		_		_
Tennessee	7	0.2	(0.1-0.6)	2	0.3	(<0.1-2.0)	_	_	_	_	_	_
Texas	49	0.6	(0.3-1.0)	22	1.2	(0.5-2.7)	25	0.4	(0.2-1.1)	-		_
Utah	9	0.6	(0.2–1.2)	_	_		_	_	-	_	-	
Vermont	2	0.5	(0.2–1.2)	_	_	_	_	-	_	_	_	
Virginia	2	0.3	(<0.1–0.2)	 15	1.4		_	_	20-27	-		22 - 122
	32					(0.3–5.7)			(<0.1-2.3)	_	_	_
Washington		0.8	(0.3–2.6)	_	_	i—	1	0.3	(<0.1-2.3)	_	_	_
West Virginia	10	0.8	(0.3–2.1)	\rightarrow	-	_	_	-	-	-	-	
Wisconsin	12	0.3	(0.1-0.9)	_	_	(0.5.5.5)	_	_	(0.0.0.1)	_	_	-
Total	663	0.4	(0.4–0.5)	182	0.7	(0.5-0.9)	144	0.5	(0.3-0.6)	23	0.2	(0.1–0.6)

^{*} In thousands.

[†] Estimates based only on responses to suicide items in the Mental Health module. Respondents with unknown suicide information were excluded.

[§] Persons of Hispanic origin can be of any race.

Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont. Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin. South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia. West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

^{**} Low precision; no estimate reported. According to the suppression criteria, prevalence estimates are suppressed if any of the following occurred: the prevalence estimate is < 0.005% or >99.995%; the relative standard error of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is ≤50%; the relative standard error of the negative of the natural logarithm of (1-p) is >0.175 if the prevalence is >50%; the actual sample size is <100; or the effective sample size (defined as the actual sample size divided by the design effect) is <68.

Surveillance Summaries

The Morbidity and Mortality Weekly Report (MMWR) Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, visit MMWR's free subscription page at http://www.cdc.gov/mmwr/mmwrsubscribe. html. Paper copy subscriptions are available through the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Address all inquiries about the MMWR Series, including material to be considered for publication, to Editor, MMWR Series, Mailstop E-90, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333 or to mmurq@cdc.gov.

All material in the MMWR Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in MMWR were current as of the date of publication.

U.S. Government Printing Office: 2012-523-043/ Region IV ISSN: 1546-0738