



Published in final edited form as:

Toxicol Lett. 2018 December 01; 298: 112–124. doi:10.1016/j.toxlet.2018.06.003.

Biological monitoring of workers exposed to engineered nanomaterials

P. Schulte^{a,*}, V. Leso^b, M. Niang^c, and I. Iavicoli^b

^aNational Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, 1090 Tusculum Avenue, MS C-14, Cincinnati, OH 45226, USA

^bDepartment of Public Health, University of Naples Federico II, Via S. Pansini 5, 80131 Naples, Italy

^cUniversity of Cincinnati, Cincinnati, OH, USA

Abstract

As the number of nanomaterial workers increase there is need to consider whether biomonitoring of exposure should be used as a routine risk management tool. Currently, no biomonitoring of nanomaterials is mandated by authoritative or regulatory agencies. However, there is a growing knowledge base to support such biomonitoring, but further research is needed as are investigations of priorities for biomonitoring. That research should be focused on validation of biomarkers of exposure and effect. Some biomarkers of effect are generally non-specific. These biomarkers need further interpretation before they should be used. Overall biomonitoring of nanomaterial workers may be important to supplement risk assessment and risk management efforts.

Keywords

Hazard identification; Exposure assessment; Risk management

1. Introduction

The utility and novelty of engineered matter at the nanoscale is driving a revolution in science and commerce (Roco, 1997; Stirling, 2018). Engineered nanomaterials (ENMs) offer the opportunity to make products lighter, stronger, more conductive, and generally “smarter” (Iavicoli et al., 2014a,b; Navya and Daima, 2016; Thiruvengadam et al., 2018). ENMs entered commerce in the early 2000s and found utility in many sectors and commercial products. Indeed, workers have been and are expected to be increasingly involved and exposed to ENMs in various applications as well as along their whole life-cycle, from research to end-life or recycling (Schulte et al., 2016).

*Corresponding author. PSchulte@cdc.gov (P. Schulte).

Publisher's Disclaimer: Disclaimer

The findings and conclusions in this report are those of the author(s) and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

Simultaneously, occupational health concerns have emerged regarding the possible impact of such nano-sized materials on the health of exposed employees. These concerns were based on previous evidence about the effects of ultrafine particulate matter derived from air pollution epidemiological studies, data extrapolated from investigations on the health consequences of diesel and welding fume exposures that may contain incidental nanoparticles (NPs), and preliminary toxicological findings obtained in vitro and in vivo models with ENMs (Iavicoli et al., 2011, 2012, 2016; Leso et al., 2017; Oberdörster and Yu, 1999; Peters et al., 2011; Stone et al., 2017).

Assessing the occupational risks derived from ENM exposure is a challenging task. This is because of difficulties in characterizing their potential hazard due to their huge physico-chemical variability, as well as in adequately assessing environmental exposure and individual response. In fact, to date, although there has been extensive research, no standardized environmental monitoring strategies to assess workplace and breathing zone exposures, nor the physico-chemical ENM features to be measured as the most effective dose metric parameters are widely agreed upon (Brouwer et al., 2009; Brouwer, 2010; Pietroiusti et al., 2018). Similarly there are no instances of regular monitoring of nano-material workers for biomarkers of effect. This is in part likely because these markers are not specific and have not been fully validated as indicators of adverse effects of specific exposure.

Nonetheless, biological monitoring may contribute to the identification of potential hazards of ENMs and to the assessment of occupational exposure to such xenobiotics; therefore, supporting a more adequate assessment and management of risks (Schulte and Hauser, 2012). In this study, the current status of biological monitoring for ENM workers will be addressed.

2. Methods

In order to review the available scientific literature on biological monitoring, a framework matrix was employed (Fig. 1). PubMed, Scopus, and Web of Science databases were searched. Our search strategy was aimed to identify issues concerning the development and use of biomarkers that may be helpful to address the different steps along (Fig. 1) the continuum of occupational safety and health actions from hazard identification to risk management (*x*-axis) (Schulte and Hauser, 2012). Additionally, the literature related to various classes of biomarkers (*y*-axis) was assessed for each category and subcategory in the continuum. The objective of the search was to identify papers that raised issues in the use of biomarkers for a cell or cells in the matrix. These issues will be discussed under the categories of biomonitoring, epidemiological research, and medical surveillance. However, as the major findings were from toxicological research on ENMs, these will be presented to establish a foundation for thinking about the utility of various types of biomarkers for other purposes in the continuum. There is a rich literature on the validation of biomarkers that can be consulted regarding their utility (Bonassi and Au, 2002; Schulte and Hauser, 2012; Schulte and Talaska, 1995).

3. Hazard identification

Since the late 1990s, there has been a growing knowledge-base about the potential adverse effects of ENMs. This seems a challenging issue due to the diverse universe of materials characterized by combinations of a large number of physico-chemical characteristics and other parameters that may include size, shape, chemical composition, solubility, crystalline structure, charge, surface characteristics, attached functionalized groups, agglomeration, and impurities. In humans, ENMs generally have not yet been reported to cause adverse health effects (Bergamaschi, 2012). This may be due to a number of factors: the precautions adopted by the scientific and commercial communities to control exposure levels, the guidance from authoritative agencies responsible for the development of nanotechnology, the small number of involved workers, and the limited time since first exposure, which is generally 15–20 years, considering the periods that such products were introduced in commerce (Fig. 2).

Considering that nanotechnology is an enabling technology diffused through many productive sectors, the mass of produced ENMs is generally small. In these complex industrial scenarios, ENM dose-response relationships may be underestimated by the confounding influence of other possible chemical co exposures. Nevertheless, hazard research in the last 15 years has shown that some types of ENMs appears to have the ability to result in adverse effects in animals. Table 1 shows an overview of the toxicological findings. Interestingly, toxicological evidence may be helpful in extrapolating information for biological monitoring.

3.1. In vitro studies

In vitro studies provide information concerning molecular effects of ENMs on cellular models, therefore suggesting possible modes of action in relation also to the intrinsic ENM physico-chemical features. This information may be important to define the hazardous properties of such nanoscale materials, a critical step in the identification of suitable biomarkers. Different ENMs, including silver (Ag)- (Müller et al., 2018; Murphy et al., 2016), titanium dioxide (TiO₂)- (Kongseng et al., 2016; Patil et al., 2016; Renwick et al., 2001), zinc oxide (ZnO)- (Patil et al., 2016), silica (SiO₂)- (Breznan et al., 2017), magnesium oxide (MgO)-NPs (Mahmoud et al., 2016), and carbon based-NMs (Chatterjee et al., 2017; Chortarea et al., 2017; Kim and Yu, 2014), demonstrated cyto-toxic, inflammatory, and oxidative stress effects, as well as the reduction of specific cellular function in vitro models. The small ENM size, resulting in a high surface area to volume ratio, may enhance ENM biological reactivity and relative toxicity (Guichard et al., 2016; Kim and Ryu, 2013; Soares et al., 2016; Uboldi et al., 2016).

Additionally, for adequate hazard identification, the chemical nature and surface chemistry should be also considered as possible influencing factors, since differently composed ENMs, i.e., metal oxide-NPs vs multi-walled-carbon nanotubes (MWCNTs) (Xia et al., 2013), Ag-NPs vs graphene oxide nano-sheets (Ivask et al., 2015), and differently functionalized NMs (Chatterjee et al., 2017; Magdolenova et al., 2015), resulted in diverse cytotoxic effects that could be also dependent on cellular type specificity (Chatterjee et al., 2017). The crystalline structure of the nano-compounds could also affect the toxicity of some ENMs, as in the case

of anatase and rutile TiO₂-NPs (Sayes et al., 2006). Shape-dependent effects have been reported for several ENMs, including Au- and TiO₂-NPs, and carbon based-ENMs (Allegri et al., 2016; Chithrani et al., 2006; Hamilton et al., 2009; Park et al., 2003; Porter et al., 2013; Öner et al., 2017). ENMs with a high aspect ratio morphology have raised particular concern in the field of respiratory toxicology, because they may induce toxicity similar to other fibrous substances, e.g., asbestos (Donaldson et al., 2011). Overall, these data may support a different behavior of ENMs according to their primary features as well as also in relation to those properties secondarily acquired when coming in contact with different culture media (Ritz et al., 2015).

3.2. In vivo studies

In vivo studies are essential to define the toxico-kinetic and dynamic behavior of ENMs and particularly to identify early adverse effects and possible target organ damage whose investigation, through biomarkers detected in accessible biological matrices, may be important for a suitable planning of biological monitoring programs. As previously mentioned, the biomolecular interactions that ENMs may experience when in contact with biological fluids, strictly dependent on ENM exposure route, can determine the formation of the so called “protein corona,” which may affect ENM dynamic profile, and therefore, biological monitoring findings (Monopoli et al., 2012).

3.2.1. Preclinical alterations—As reported in cellular models, and also in human and animal experiments, ENMs elicited the activation of oxidative stress responses that could be monitored through the reduction in plasma levels of anti-oxidant defense systems. This was detected in workers involved in the manufacture and/or application of ENMs in Taiwan, and also found after a six month follow up period (Liao et al., 2014; Liou et al., 2012, 2016). Additionally, the increased exhaled breath condensate (EBC) concentrations of lipid, nucleic, and protein oxidation markers were detected in workers exposed to Fe₂O₃- and Fe₃O₄-NPs (Pelclova et al., 2016a), MWCNTs (Lee et al., 2015), and TiO₂-NPs (Pelclova et al., 2017a, 2017b) compared to controls. In such epidemiological investigations, a dose-response correspondence was demonstrated according to the ENM environmental monitoring concentrations (Pelclova et al., 2016a, b) as well as the increase in risk levels established through a control binding nano-tool risk assessment approach (Liao et al., 2014).

Comparably, in animal models, reductions in plasma anti-oxidant defense systems and increased levels of Reactive Oxygen Species (ROS) and malondialdehyde (MDA) could be determined with SiO₂ (Du et al., 2013; Liu and Sun, 2013), Ag (Genter et al., 2012; Martins et al., 2017), Ag-NPs in coexposure with TiO₂-NPs (Martins et al., 2017), Fe₂O₃- NPs (Sundarraj et al., 2017), and multi-walled carbon nanotube (MWCNTs) exposures (Reddy et al., 2011).

Regarding inflammatory response, blood levels of fibrinogen and pro-inflammatory cytokines were increased in workers exposed to nano-sized carbon black and MWCNTs (Fatkhutdinova et al., 2016; Liou et al., 2012; Vlaanderen et al., 2017; Zhang et al., 2014) as well as to TiO₂-NPs compared to controls (Zhao et al., 2018). Conversely, when comparing serum proteins associated with inflammatory responses in ENM exposed and not-exposed

workers in research laboratories, significant increases in CD40, TNFR2, and CD62P serum concentrations were evident during the first shift of a working week and at the end of the week in exposed employees (Glass et al., 2017). These findings may suggest an anti-inflammatory and suppressive homeostatic cellular response in relation to an immune activation.

Enhanced levels of leukotrienes were detected in the EBCs of subjects occupationally exposed to TiO₂-NPs (Pelclova et al., 2016c) and MWCNTs (Lee et al., 2015). Comparably, increased concentrations of 8-isoprostane in EBC (Liou et al., 2017) and enhanced fractional exhaled nitric oxide were reported in workers exposed to metal oxide ENMs (Wu et al., 2014), also when the analysis was stratified according to the type of NM exposure (SiO₂, TiO₂, indium tin oxide-NPs), while potential fibrotic and pro-inflammatory biomarkers were detected in the sputum of MWCNT-involved workers (Fatkhutdinova et al., 2016). Conversely, in research laboratory workers handling ENMs, no significant alterations in exhaled nitric oxide could be detected (Glass et al., 2017).

Alterations in blood acute-phase proteins and cytokine concentrations were also detected in animal models through acute to sub-acute administrations of different types of metallic or metal oxide-NPs such as TiO₂- (Park et al., 2009), CeO₂- (Nalabotu et al., 2011; Srinivas et al., 2011), Fe₃O₄- (Chen et al., 2010; Park et al., 2010; Srinivas et al., 2012)-, Ag- (Holland et al., 2015, 2016), and SiO₂-NPs (Downs et al., 2012; Du et al., 2013; Lu et al., 2011) as well as with carbon-based NPs (Erdely et al., 2009). Interestingly, such preclinical alterations, whatever the biological matrix employed, may function both as indicators of early effects before clinical manifestations may occur and as indirect markers of exposure although with a low specificity for ENMs (Aragon et al., 2017; Erdely et al., 2009; Shvedova et al., 2016).

3.2.2. Pathological alterations—Pathological alterations pertain to a greater understanding of the potential for adverse effects following exposure. In highly exposed nanoscale carbon black workers, alterations in pulmonary functional parameters were observed compared to controls. This suggests possible adverse respiratory alterations (Zhang et al., 2014), although other investigations failed to detect such changes in MWCNT (Lee et al., 2015; Liao et al., 2014; Vlaanderen et al., 2017) and TiO₂-NP exposed employees (Pelclova et al., 2016c, 2017b) as well as in workers handling ENMs in research laboratories (Glass et al., 2017).

Interestingly, a significant dose-dependent increase in the pulmonary surfactant protein D serum levels, as a biomarker of lung damage, was detected in workers employed in a packaging workshop of a nano-TiO₂ manufacturing plant in eastern China (Zhao et al., 2018). In these workers, alterations in cardiovascular disease markers, i.e., VCAM-1, ICAM-1, LDL, and TC, were associated with occupational ENM exposure (Zhao et al., 2018). Haematological alterations were reported in animals after oral or intra-gastric exposure to ZnO-NPs (Park et al., 2014) and TiO₂-NPs (Vasantharaja et al., 2015) as well as after dermal exposure to hydroxyapatite-NPs (Parayanthala Valappil et al., 2014), while could not be detected after a sub-acute inhalation of CeO-NPs (Gosens et al., 2016). Increased ALT concentrations and decreased albumin levels in serum as markers of

hepatocyte injury, confirmed also by histopathological alterations, were reported in rats intratracheally instilled with CeO₂-NPs (Nalabotu et al., 2011). The hepatotoxic potential of ENMs has been also documented in animals orally treated with ZnO- (Park et al., 2014), Ag-, and Au-NPs (Shrivastava et al., 2016). Significant nephrotoxic alterations principally induced by metallic NPs have been demonstrated through the assessment of the kidney injury molecule-1 (KIM-1) urinary level (Blum et al., 2015; Iavicoli et al., 2016) as well as by the BUN and creatinine serum biomarkers following an oral exposure to Cu- (Chen et al., 2006; Lei et al., 2008; Liao and Liu, 2012; Sarkar et al., 2011), Au- (Shrivastava et al., 2016), and mesoporous silica-NPs (Li et al., 2015). Indicators for the abovementioned alterations may function as potential biomarkers helpful to define target organs of ENM toxicity and, in turn, suggest potential pathological health conditions susceptible to be aggravated by ENM exposure.

3.2.3. Genotoxicity—Due to their small particle size, large surface area, and physico-chemical characteristics, NPs exhibit unpredictable genotoxic properties. Studies on workers failed to show genotoxic effects (Liao et al., 2014; Liou et al., 2012) while a general dose-dependent increase in DNA strand breaks and micronucleus (MN) frequency was found in human peripheral blood cells treated in vitro with metal or metal oxide-NPs (Colognato et al., 2008; Di Bucchianico et al., 2013; Flower et al., 2012; Ghosh et al., 2010, 2012, 2013; Kang et al., 2008, 2011; Paino et al., 2012; Soni et al., 2017; Tavares et al., 2014), carbon-based NPs (Cveticanin et al., 2010; Öner et al., 2017; Tavares et al., 2014), and dendrimers (Ziemba et al., 2012). Concerning in vivo results, a significant increase in such parameters was detected in the peripheral blood cells of TiO₂- (Song et al., 2012; Trouiller et al., 2009) and ZnO-NP treated animals (Patil et al., 2016), as compared to controls, and a dose-dependent increase in DNA fragmentation percentage was detected in lymphocytes of SiO₂ and Fe₂O₃-NP treated rats (Jiménez-Villarreal et al., 2017). However, positive results were not always confirmed and conflicting results were reported also for other metallic or metal oxide ENMs (Balasubramanyam et al., 2009; Chen et al., 2014; Cordelli et al., 2017; Downs et al., 2012; Lindberg et al., 2012; Sadiq et al., 2012; Singh et al., 2013a, b; Song et al., 2012; Tiwari et al., 2011). This may suggest a possible influencing role of different ENM physico-chemical features, routes of exposure, and administered doses.

3.3. “Omic techniques” for hazard identification

Traditional toxicology is rapidly evolving into a system-based approach, able to capture almost all the interactions between living systems and endogenous and/or exogenous xenobiotics (Balbo et al., 2017). “Omic” techniques may reveal methods helpful to assess a wide range of biological responses induced by ENM exposures, therefore, they may be promising tools for the development of novel biomarkers of exposure and early effect (Fadeel, 2015; Schulte and Hauser, 2012). These techniques currently include genomics, transcriptomics, proteomics, and more recent approaches, such as metabolomics, and adductomics involved into gene expression and its consequences.

3.3.1. Genomics—Genomic studies may provide the opportunity to detect injury at the molecular level and the signaling pathways involved in organ damage long before the clinical symptoms occur (Andersen and Krewski, 2009; Klaper et al., 2014). Several

toxicogenomic studies have investigated the in vitro and in vivo response, principally pulmonary gene expression changes, to NMs using DNA microarray analysis (Aydın et al., 2017; Costa et al., 2018; Decan et al., 2016; Ellinger-Ziegelbauer and Pauluhn, 2009; Gao et al., 2012; Husain et al., 2013; Li et al., 2013, 2017; Pacurari et al., 2011; Snyder-Talkington et al., 2015). However, accessible and more easily applicable biological matrices, i.e., blood or peripheral blood cells, have been rarely employed to assess systemic alterations; determining these correlations in animal models is an important part of practical biomarker validation. The expression of genes involved in immune, inflammatory and oxidative stress responses was affected by Au-NP-oligonucleotide complexes in human peripheral blood mononuclear cells (Kim et al., 2012), by graphene oxide also functionalized with amino groups in T lymphocytes and mononuclear cell lines (Orecchioni et al., 2017). In in vivo experiments, blood expression changes in genes involved in inflammation, oxidative stress, growth factors, tissue remodeling, and endothelial function were obtained in circulating blood cells of animals treated with SW- or MWCNTs (Erdely et al., 2009).

3.3.2. Transcriptomics—Transcriptomics aims at quantifying changes in gene expression through the enumeration of the number of mRNA copies (Costa and Fadeel, 2016). A significant increase in the mRNA expression of KL-6, a marker for the diagnosis and monitoring of interstitial lung diseases, as well as its up-stream and down-stream genes, were detected in accessible biological matrices, i.e., whole blood and sputum samples collected from workers exposed to MWCNTs compared to controls (Fatkhutdinova et al., 2016; Shvedova et al., 2016). In vitro studies also reported mRNA expression changes in macrophages exposed to mesoporous silica-NPs and PEGylated mesoporous silica-NPs at doses that do not elicit acute cytotoxicity (Yazdimamaghani et al., 2017). Numerous blood mRNAs were significantly up- or down-regulated post-MWCNT inhalation in animals developing lung pathological changes (Dymacek et al., 2015; Snyder-Talkington et al., 2013, 2016). Therefore, changes in blood mRNA expression, may potentially serve as suitable biomarkers for ENM-induced lung pathological changes, providing less invasive measurements, compared to tissue analysis, that can be taken at shorter intervals with lower costs (Snyder-Talkington et al., 2016).

3.3.3. Proteomics—Proteomic investigation may be helpful to identify candidate protein biomarkers for the evaluation of ENM early effects (Matysiak et al., 2016). In vitro results reported differently expressed proteome profiles related to cellular viability, oxidative stress, and heat response processes in human lung cells exposed to MWCNTs (Phuyal et al., 2018) as well as in the human monocytes treated with Au-, CuO-, and CdTe-NPs (Tarasova et al., 2017). On the other hand, in vivo studies demonstrated an affected expression of proteins associated with metabolism, oxidative stress, and immune responses following TiO₂-NP inhalatory exposure (Maurer et al., 2016). More recently, the possible employment of proteomic analysis for the detection of protein carbonylation patterns as persistent, sensitive, and indirect biomarkers of ENM induced oxidative stress reaction has been explored (Driessen et al., 2015; Riebeling et al., 2016). Overall, the suitability of these proteomic changes as biological indicators of effect should be firstly confirmed by traditional biochemistry, thus supporting their effectiveness in detecting early NP toxicity.

3.3.4. Metabolomics—Metabolomics is the comprehensive analysis of all the metabolites of an organism or specified biological samples (Robertson et al., 2011). In vitro metabolomic data, demonstrated that CeO-, SiO₂- and CuO-NPs could affect the lipidome profile, and compromise the Phase II conjugational capacity in exposed hepatic cells (Kitchin et al., 2017). Differently hydroxylated or carboxylated functionalized CNTs induced alterations in amino acid metabolisms in hepatic and bronchial cells (Chatterjee et al., 2017) while an antioxidant effect was observed for chitosan-coated or ceria supported-Au-NPs in human peripheral blood cells (Palomino-Schätzlein et al., 2017). Concerning the metabolic profiling of ENM effects, as a rapid in vivo screening for nano-toxicity biomarkers, analyses performed in serum and urine of rats treated with Cu- (Lei et al., 2008) and TiO₂-NPs (Bu et al., 2010; Tang et al., 2010, 2011) provided evidence for the hepatotoxicity, nephrotoxicity and alterations in energy metabolism induced by these NPs. Other studies reported the ability of Ag- (Hadrup et al., 2012; Xie et al., 2018), Fe₂O₃- (Feng et al., 2010), SiO₂-NPs (Lu et al., 2011; Parveen et al., 2012), carbon 14-labeled C60 fullerenes (Sumner et al., 2010) and CNTs (Lin et al., 2013) to affect the liver, energy, lipid, glucose, and amino acid metabolism as assessed by the alterations found in serum and urine metabolic pathways, while amorphous SiO₂, zirconium dioxides, and barium sulphate-NPs failed to show a relevant impact on plasma metabolome patterns (Buesen et al., 2014). Although interesting, the different and not always well characterized ENMs employed, as well as the lack of standardized analytical techniques and procedures, make the results difficult to compare, and no definite conclusions to be extrapolated.

3.3.5. Adductomics—Adductomic approaches have been developed to comprehensively describe toxicological features in response to a genotoxic xenobiotic insult (Hemeryck et al., 2016; Villalta and Balbo, 2017). Specific classes of DNA adducts are those resulting from the reaction of ROS (%OH) with DNA, i.e., the 8-hydroxy-deoxy-guanosine (8-OH-dG). In field studies, 8-OH-dG concentrations were measured in easily available biological matrices collected from exposed workers, although with variable results. While no significant alterations were detected in urine, and plasma samples from employees of 14 manufacturing plants in Taiwan (Liao et al., 2014; Liou et al., 2012), increased concentrations were more recently reported in urine and white blood cells of workers exposed to metal oxide ENMs, whatever the type of ENM was considered (Liou et al., 2016, 2017), and in EBCs obtained from TiO₂-NP exposed subjects (Pelclova et al., 2016b). This DNA damage has been also investigated in vitro models exposed to metal or metal oxide ENMs, which demonstrated significant increases (Ng et al., 2017) although with a cellular type specificity (Mahmoud et al., 2016), and in Ag-NP treated animals that showed increased urinary concentrations of this biomarker (Chuang et al., 2013). The field of DNA adduct research is a highly promising area due to their potential. This is primarily related to the fact that, for instance, adducts may be interpreted as indicators of internal doses, biologically effective dose, and early effect, as well as susceptibility. Though the need exists to be validated specifically for each purpose.

3.3.6. Epigenetics—Epigenetics refers to heritable, reversible changes in gene expression occurring without alterations in DNA sequence (Shyamasundar et al., 2015; Sierra et al., 2016). DNA methylation, histone tail, and microRNA modifications are

considered useful epigenetic markers. Recent research demonstrated that CNTs (Brown et al., 2016; Chatterjee et al., 2017; Li et al., 2016; Öner et al., 2017; Tabish et al., 2017), Au- (Tabish et al., 2017), TiO₂- (Bai et al., 2015; Patil et al., 2016), SiO₂- (Gong et al., 2010, 2012; Mytych et al., 2017), ZnO- (Choudhury et al., 2017), and CuO-NPs (Lu et al., 2016) were able to induce changes of specific methylation patterns, including tumor suppressor-, inflammatory-, and DNA repair genes in vitro and in vivo models. Interestingly, a significantly lower global DNA methylation in white blood cells collected from workers exposed to nanoscale indium tin oxide in plants in Taiwan was recently reported by Liou et al. (2017), while Ghosh et al. (2017), although failing to detect significant differences in global methylation, found significant changes in genespecific DNA methylation in MWCNT exposed workers compared to controls.

Histone conformational modifications may either facilitate or depress the access of transcriptional machinery to the promoter region of some genes, leading to gene silencing or activation, respectively. As concerns ENMs, a global hypoacetylation in human breast carcinoma cells causing transcriptional repression of anti-apoptotic genes, thereby promoting cellular death, was detected after cadmium telluride quantum dots exposure (CdTe-QDs) (Choi et al., 2008). Ag-NPs were reported to affect histone post-translational modifications thus inducing a reduction in hemoglobin levels in mouse erythroleukemia cells (Qian et al., 2015). Non-coding microRNAs have been investigated in an attempt to identify fine, regulator molecules in ENM induced toxicity since they can modulate gene expression through the interaction with other epigenetic processes (Zhao et al., 2016; Peschansky and Wahlestedt, 2014). The changes in the microRNA expression profiling induced by exposure to Fe₂O₃-NPs, CdTe-QDs, Au-, Ag-NPs and MWCNTs were demonstrated to globally affect the mRNA and protein output of human treated cells, subsequently affecting many key biological patterns (Eom et al., 2014; Li et al., 2011a, b; Ng et al., 2017). Micro-RNA expression changes, implicated in inflammation and immune reactions, were found in the lungs of mice intratracheally exposed to TiO₂-NPs (Halappanavar et al., 2011). Changes in the blood levels of liver-specific miRNAs were identified in mice exposed to SiO₂-NPs (Nagano et al., 2013). However, as the functional consequences of the above mentioned epigenetic alterations were not assessed, further characterization of miRNA responsive genes and their role in adverse effects need to be defined.

4. Exposure assessment

Exposure assessments primarily involve area and breathing zone monitoring, job-exposure matrix modeling, or exposure-banding (Kauppinen et al., 2014; Moretto, 2015; NIOSH, 2009, 2017). Although, in recent years, there has been a growth of strong literature on task-based environmental monitoring (Asbach et al., 2017; Brouwer, 2010; Brouwer et al., 2012; Eastlake et al., 2016; Methner et al., 2010a,b,2012), some challenging issues, including determining the best measurement metrics, availability of appropriate instrumentation, and diverse sampling strategies, prevent an adequate assessment of environmental exposure levels. Therefore, human biomonitoring (HBM) may function as a complementary means for exposure assessment that can take into account the inter-individual variability in absorption,

metabolism, and excretion; the individual workload; and recent versus past exposure (Manno et al., 2010).

Currently there are no examples of HBM as a part of routine assessment of workers exposed to ENMs and no regulatory requirements are available for HBM. Nonetheless, there is a growing body of toxicological research that illustrates the possibility to verify some functional alterations detected in experimental settings as biomonitoring indicators for workers. For example, some ENMs can be detected in biological matrices, including blood or plasma, urine, and feces. Following metal- or metal-oxide-based ENM exposure, both via the inhalation or the intratracheal exposure (Balasubramanian et al., 2013; He et al., 2010; Semmler-Behnke et al., 2008; Sundarraj et al., 2017; Sung et al., 2009; Takenaka et al., 2001, 2006; Yu et al., 2007; Zhu et al., 2009) and via human skin application (Gulson et al., 2010, 2012), the elemental metal content is retrievable in blood, although generally in very small amounts. A positive dose-response relationship between the inhaled Ag-ENM and the Ag blood content was demonstrated in mice (Sung et al., 2009), although a differential size dependent biodistribution was reported for gold (Au)-NPs, because smaller 7-nm sized Au-NPs produced a greater metal content in blood compared to their larger 20-nm counterparts (Balasubramanian et al., 2013).

The duration of exposure can affect the interpretations of biomonitoring data, in fact, significant Ag-ENM accumulation in blood was evident after 15 days of treatment, but not following a shorter 5 day period of exposure. This suggests that the body burden of the ENMs is influenced by homeostatic processes (or the exposure could also be influencing homeostatic activities) rather than by the extent of the exposure (Iavicoli et al., 2014a,b). Detectable metal concentrations were reported also in urine samples collected from animals treated with metal or metal oxide ENMs via the respiratory or the dermal route of exposure (Balasubramanian et al., 2013; Gulson et al., 2010, 2012; Sundarraj et al., 2017; Zhu et al., 2009), but there would be no relationship described for any effect. Because of macrophage-mediated clearance mechanisms, the measurement of the elemental metal content in feces may be useful to evaluate the recent/current respiratory exposure to metal-NPs (Balasubramanian et al., 2013; Chuang et al., 2013; Li et al., 2016; Sundarraj et al., 2017e; tZahlu., 2009). However, it is rather difficult to routinely employ feces as a suitable biological matrix for occupational biomonitoring.

Exhaled breath condensate (EBC) is a promising matrix for human biological monitoring investigation, as demonstrated by the increased Ti levels in pre- and post-shift EBC samples collected from workers exposed to TiO₂-NPs in a pigment production plant compared to unexposed controls (Pelclova et al., 2015, 2016b). Interestingly the biological levels of such an indicator resulted positively correlated with the environmental concentrations in different production or research areas of the plant. However, not all ENMs are easily found in body fluids. Carbon nanotubes are one example. In fact, there is minimal transport of carbon nanotubes from the alveoli to the blood, and they tend to move out of the blood rapidly in to various organs (Erdely et al., 2009).

5. Risk assessment

Taking into account what was previously described for ENM hazard assessment and exposure evaluation, it is possible to perceive the difficulties in defining suitable risk assessment strategies. Biological monitoring data, including biomarkers of exposure and effect, can be used in risk assessment involving nanomaterial workers, but thus far, there are no examples where biomarkers have been used in risk assessments. In this scenario, an exposure-response modeling should be pursued to fully exploit the potential of possible exposure indicators. In fact, when the dose-response relationship is defined, the biomarker of exposure does not only indicate the dose actually adsorbed but provides also a reasonably accurate quantitative estimate of the occupational risks at the group and/or individual level (Iavicoli et al., 2014a). Additionally, scientific efforts should be focused at employing biological monitoring data to develop occupational exposure limits for ENM workers. To do this, the relationship between biomarker levels and adverse effects must be clearly demonstrated in a study design that includes no-effect exposure levels.

The efforts involving categorical approaches to developing occupational exposure limits (OELs) are beginning to identify biomarkers to predict toxicity. For example, the Nanosolutions project tested 31 different engineered ENMs and, out of 8 million data points, identified 11 biomarkers that support the toxicity of these xenobiotics (Nanosolutions, 2018). The response of these markers at human exposure levels must be determined. The 11 biomarkers will then be applied in untested engineered ENMs. These and other approaches that used ENMs with known hazards and the biomarkers of effect related to them may ultimately be used in dose-response modeling.

Another area that involves biomarkers that may be useful in risk assessment and OEL development for ENMs is the use of “omic” technologies that may be useful to understand the interactions and specific pathways impacted by different ENMs. Such biomarkers should be carefully verified in accessible biological matrices under low-dose, long-term conditions of exposure to define their specificity for different ENM insults, their predictive value, and therefore their realistic applicability. Investigations of the effect of functional silica ENMs on a human lung carcinoma cell line showed that it is possible to identify an “omic” analog to the NOAEL involving a no observed transcriptomic effect level or NOTEL (Pisani et al., 2015). This is the level of transcriptomic effect below which cellular responses are not seen to occur and may be much lower than the NOAEL. It could be used as a point of departure in deriving reference values. The development of pathways depicting the sequence of events between exposure to a stressor, progressing through intermediate events, and culminating in an adverse outcome is a growing concept for use in risk assessment (OECD, 2017). This concept may be increasingly used by regulatory agencies and, consequently, there may be more pressure for biomonitoring related to it.

Another relevant contribution to possible ENM-tailored risk assessment processes can be derived from epidemiologic studies. The review of the “first wave” of epidemiological studies of NM workers showed that most were too limited to indicate risk, although some found increased biomarkers of oxidative stress and inflammation in exposed workers (Liou et al., 2015). However, exposure assessment was generally qualitative, study designs were

generally cross-sectional, and various selection biases were possible. Therefore, suitable epidemiological research should be planned in order to verify environmental exposure to ENMs and their relationships with biological indicators of exposure and effect. However, no examples where biomarkers have been used in such process are currently available. Studies aimed to address the effectiveness of control technology in a nanomaterial workplace should be pursued not only as a primary way to test control effectiveness in workplaces but also as a useful mean to demonstrate the failure of prevention (Kreider and Halperin, 2011). It could be possible, in laboratory studies, to evaluate control designs and utilize a biomonitoring component of laboratory animals to supplement airborne exposure measures. Biological monitoring may be even more important, in consideration of the possible absorption of ENMs through the dermal route of exposure especially in conditions of compromised skin integrity due to pre-existing diseases or damaging coexposures (Brouwer et al., 2012; Gulson et al., 2010, 2012; Larese Filon et al., 2016; Osmond-McLeod et al., 2014).

6. Risk management

Biomonitoring is a risk management tool along with workplace exposure assessment and exposure controls. In this context, biological monitoring data could be used effectively to verify the efficacy of the exposure limit in protecting the health of the workers (Iavicoli et al., 2014a,b). At present, the utility of biomonitoring nanomaterial workers depends on the status of evidence about the hazard of particular ENMs that could be the target of biomonitoring. Biomonitoring should not be used as a substitute for controlling exposure or for taking precautionary control measures.

An additional application of biological monitoring in risk management is the medical surveillance of workers. This is the effort to assess asymptomatic workers for early indications of health problems. There are currently no mandated medical surveillance guidance for ENM workers except for ones involving baseline and period respiratory assessment (pulmonary function testing, x-ray) (NIOSH, 2013). No cellular or molecular biomonitoring is mandated. There is a sparse literature studying the effectiveness of medical surveillance for nanomaterial workers. Gulumian et al. (2016) conducted a review of structures of medical surveillance of nanomaterial workers and identified seven studies that met inclusion criteria. They concluded that there was very low quality of evidence that screening might detect adverse health effects associated with workplace exposures to ENMs. However, this study conflated epidemiological studies and medical surveillance and, as noted earlier, there has not been much time since first worker exposure and the possibility of identifying health effects in epidemiological studies in routine medical surveillance.

One further area of risk management that involves biomarkers of exposure and effect and shows great promise is to design out potential hazards in ENMs. This is known as prevention through design or “safety by design” (Geraci et al., 2015). The concept is illustrated by the identification of specific toxic endpoints that were observed after zebra fish were exposed to functionalized ENMs. This study allowed for the development of predictive models for the design of inherently safer ENMs. Similarly biomarkers found in large arrays in ENM studies may be the basis for material designs that will have reduced ENM hazards.

7. Research needs

For biological monitoring of nanomaterial workers to go forward, there is need for robust research to develop biological monitoring protocols (Iavicoli et al., 2014a,b). This should include the physico-chemical characterization of materials to which a population is exposed and well established understanding of toxico-kinetic and dynamic characteristics. This research may result in the determination of whether biological monitoring should measure direct parent compounds or possible metabolites or decomposition products. It may explain also the possible role of biomolecular corona formation in affecting ENM bio-disposition and, consequently, biomonitoring results.

There is an evident need to link biomarkers in experimental animals to biomarkers in humans. Candidate biomarkers must be validated with exposure and/or effects (Bergamaschi et al., 2015). Additionally, considering the extremely high doses frequently employed in experimental settings and the extremely low ENM retrieved fractions in biological fluids, potential biomarkers should be validated under low-doses and longer periods of treatment, through analytical techniques sensitive enough to address trace levels of biological markers. Moreover, the predictive significance of such biological monitoring alterations in terms of long-term effects, the influences exerted by ENM characteristics, modes and periods of exposure, and inter- and intra-individual variabilities should be assessed to define suitable biomarkers and accurate interpretation of their results.

Validation is a process—a sense of degree rather than an all-or-none-state (IPCS, 2001). Validation includes assessment not only of sensitivity, specificity, and predictive value, but also utility in studies of workers on a routine basis. Ultimately validation and developmental research to identify biomarkers for biomonitoring of workers will require human studies; these may be field, chamber, or epidemiologic studies. Such studies require attention to ethical legal and social issues. Of importance are issues of privacy, confidentiality, and notification of test and study results (Schulte and Smith, 2011).

8. Conclusion

There are no authoritative mandates or requirements currently available for biomonitoring of ENM workers. However, there is a growing body of research that is setting the stage for the use of bio-monitoring to supplement environment exposure assessments to achieve a suitable ENM risk assessment and management in occupational settings. Biomarkers may be particularly useful when exposure results in systemic effects. Biomonitoring requires the determination of the extent of the hazard of candidate nanomaterial and evidence that demonstrates biomarker presence in workers correlates with exposure. Biomarkers of effect may also be used to assess exposure and have similar developmental requirements as biomarkers of exposure with the additional need to account for non-specificity and homeostatic influences. If biomonitoring is to be used to assess early or potential health effects, there will be a need for prospective studies to support that use. Overall, there is a rich research base for biomonitoring. However, further work is required before biomonitoring workers can be routinely implemented.

Acknowledgements

The authors would like to thank: Drs. Kai Savolainen and Glenn Talaska for comments on earlier drafts; Amanda Keenan and Amanda Stammer for processing; and Cheryl Hamilton for editing.

References

- Allegri M, Bianchi MG, Chiu M, Varet J, Costa AL, Orтели S, Blosi M, Bussolati O, Poland CA, Bergamaschi E, 2016 Shape-related toxicity of titanium dioxide nanofibres. *PLoS One* 11, e0151365 <http://dx.doi.org/10.1371/journal.pone.0151365>. [PubMed: 26999274]
- Andersen ME, Krewski D, 2009 Toxicity testing in the 21st century: bringing the vision to life. *Toxicol. Sci* 107, 324–330. <http://dx.doi.org/10.1093/toxsci/kfn255>. [PubMed: 19074763]
- Aragon J, Navascues N, Mendoza G, Irusta S, 2017 Laser-treated electrospun fibers loaded with nano-hydroxyapatite for bone tissue engineering. *Int. J. Pharm* 525, 112–122. <http://dx.doi.org/10.1016/j.ijpharm.2017.04.022>. [PubMed: 28412451]
- Asbach C, Alexander C, Clavaguera S, Dahmann D, Dozol H, Faure B, Fierz M, Fontana L, Iavicoli I, Kaminski H, MacCalman L, Meyer-Plath A, Simonow B, van Tongeren M, Todea AM, 2017 Review of measurement techniques and methods for assessing personal exposure to airborne nanomaterials in workplaces. *Sci. Total Environ* 603–604, 793–806. <http://dx.doi.org/10.1016/j.scitotenv.2017.03.049>.
- Aydın E, Türkez H, Hacımüftüo lu F, Tatar A, Geyiko lu F, 2017 Molecular genetic and biochemical responses in human airway epithelial cell cultures exposed to titanium nanoparticles in vitro. *J. Biomed. Mater. Res. Part A* 105, 2056–2064. <http://dx.doi.org/10.1002/jbm.a.35994>.
- Bai W, Chen Y, Gao A, 2015 Cross talk between poly(ADP-ribose) polymerase I methylation and oxidative stress involved in the toxic effect of anatase titanium dioxide nanoparticles. *Int. J. Nanomed* 10, 5561–5569. <http://dx.doi.org/10.2147/IJN.S88059>.
- Balasubramanyam A, Sailaja N, Mahboob M, Rahman MF, Hussain SM, Grover P, 2009 In vivo genotoxicity assessment of aluminum oxide nanomaterials in rat peripheral blood cells using the comet assay and micronucleus test. *Mutagenesis* 24, 245–251. <http://dx.doi.org/10.1093/mutage/gep003>. [PubMed: 19237533]
- Balasubramanian SK, Poh KW, Ong CN, Kreyling WG, Ong WY, Yu LE, 2013 The effect of primary particle size on biodistribution of inhaled gold nano-agglomerates. *Biomaterials* 34, 5439–5452. <http://dx.doi.org/10.1016/j.biomaterials.2013.03.080>. [PubMed: 23639527]
- Balbo S, Turesky RJ, Villalta PW, 2017 DNA adductomics. *Chem. Res. Toxicol* 27, 356–366. <http://dx.doi.org/10.1021/tx4004352>.
- Bergamaschi E, 2012 Human biomonitoring of engineered nanoparticles: an appraisal of critical issues and potential biomarkers. *J. Nanomater* 2012, 564121 <http://dx.doi.org/10.1155/2012/564121> 12 pp.
- Bergamaschi E, Poland C, Guseva Canu I, Prina-Mello A, 2015 The role of biological monitoring in nano-safety. *Nanotoday* 10 (3), 274–277. <http://dx.doi.org/10.1016/j.nantod.2015.02.001>.
- Blum JL, Edwards JR, Prozialeck WC, Xiong JQ, Zelikoff JT, 2015 Effects of maternal exposure to cadmium oxide nanoparticles during pregnancy on maternal and offspring kidney injury markers using a murine model. *J. Toxicol. Environ. Health Part A* 78, 711–724. <http://dx.doi.org/10.1080/15287394.2015.1026622>. [PubMed: 26090557]
- Bonassi S, Au WW, 2002 Biomarker epidemiology studies for health risk prediction. *Mutat. Res* 511, 73–86. [http://dx.doi.org/10.1016/S1383-5742\(02\)00003-0](http://dx.doi.org/10.1016/S1383-5742(02)00003-0). [PubMed: 11906843]
- Breznan D, Das DD, O'Brien JS, MacKinnon-Roy C, Nimesh S, Vuong NQ, Bernatchez S, DeSilva N, Hill M, Kumarathanan P, Vincent R, 2017 Differential cytotoxic and inflammatory potency of amorphous silicon dioxide nanoparticles of similar size in multiple cell lines. *Nanotoxicology* 11, 223–235. <http://dx.doi.org/10.1080/17435390.2017.1287313>. [PubMed: 28142331]
- Brouwer D, 2010 Exposure to manufactured nanoparticles in different workplaces. *Toxicology* 269, 120–127. <http://dx.doi.org/10.1016/j.tox.2009.11.017>. [PubMed: 19941928]
- Brouwer D, van Duuren-Stuurman B, Berges M, Jankowska E, Bard D, Mark D, 2009 From workplace air measurement results towards estimates of exposure? Development of a strategy to assess

- exposure to manufactured nano-objects. *J. Nanopart. Res* 11, 1867–1881. <http://dx.doi.org/10.1007/s11051-009-9772-1>.
- Brouwer D, Berges M, Virji MA, Fransman W, Bello D, Hodson L, Gabriel S, Tielemans E, 2012 Harmonization of measurement strategies for exposure to manufactured nano-objects. Report of a workshop. *Ann. Occup. Hyg* 56, 1–9. <http://dx.doi.org/10.1093/annhyg/mer099>. [PubMed: 22156566]
- Brown TA, Lee JW, Holian A, Porter V, Fredriksen H, Kim M, Cho YH, 2016 Alterations in DNA methylation corresponding with lung inflammation and as a biomarker for disease development after MWCNT exposure. *Nanotoxicology* 10, 453–461. <http://dx.doi.org/10.3109/17435390.2015.1078852>. [PubMed: 26375518]
- Bu Q, Yan G, Deng P, Peng F, Lin H, Xu Y, Cao Z, Zhou T, Xue A, Wang Y, Cen X, Zhao YL, 2010 NMR-based metabonomic study of the sub-acute toxicity of titanium dioxide nanoparticles in rats after oral administration. *Nanotechnology* 21, 125105 <http://dx.doi.org/10.1088/0957-4484/21/12/125105>. [PubMed: 20203358]
- Buesen R, Landsiedel R, Sauer UG, Wohlleben W, Groeters S, Strauss V, Kamp H, van Ravenzwaay B, 2014 Effects of SiO₂, ZrO₂, and BaSO₄ nanomaterials with or without surface functionalization upon 28-day oral exposure to rats. *Arch. Toxicol* 88, 1881–1906. <http://dx.doi.org/10.1007/s00204-014-1337-0>. [PubMed: 25164825]
- Chatterjee N, Yang J, Yoon D, Kim S, Joo SW, Choi J, 2017 Differential crosstalk between global DNA methylation and metabolomics associated with cell type specific stress response by pristine and functionalized MWCNT. *Biomaterials* 115, 167–180. <http://dx.doi.org/10.1016/j.biomaterials.2016.11.005>. [PubMed: 27914347]
- Chen Z, Meng H, Xing G, Chen C, Zhao Y, Jia G, Wang T, Yuan H, Ye C, Zhao F, Chai Z, Zhu C, Fang X, Ma B, Wan L, 2006 Acute toxicological effects of copper nanoparticles in vivo. *Toxicol. Lett* 163, 109–120. <http://dx.doi.org/10.1016/j.toxlet.2005.10.003>. [PubMed: 16289865]
- Chen BA, Jin N, Wang J, Ding J, Gao C, Cheng J, Xia G, Gao F, Zhou Y, Chen Y, Zhou G, Li X, Zhang Y, Tang M, Wang X, 2010 The effect of magnetic nanoparticles of Fe₃O₄ on immune function in normal ICR mice. *Int. J. Nanomed* 5, 593–599. <http://dx.doi.org/10.2147/IJN.S12162>.
- Chen Z, Wang Y, Ba T, Li Y, Pu J, Chen T, Song Y, Gu Y, Qian Q, Yang J, Jia G, 2014 Genotoxic evaluation of titanium dioxide nanoparticles in vivo and in vitro. *Toxicol. Lett* 226, 314–319. <http://dx.doi.org/10.1016/j.toxlet.2014.02.020>. [PubMed: 24594277]
- Chithrani BD, Ghazani AA, Chan WC, 2006 Determining the size and shape dependence of gold nanoparticle uptake into mammalian cells. *Nano Lett* 6, 662–668. <http://dx.doi.org/10.1021/nl052396o>. [PubMed: 16608261]
- Choi AO, Brown SE, Szyf M, Maysinger D, 2008 Quantum dot-induced epigenetic and genotoxic changes in human breast cancer cells. *J. Mol. Med. (Berlin Ger.)* 86, 291–302. <http://dx.doi.org/10.1007/s00109-007-0274-2>.
- Chortarea S, Barosova H, Clift MJD, Wick P, Petri-Fink A, Rothen-Rutishauser B, 2017 Human asthmatic bronchial cells are more susceptible to subchronic repeated exposures of aerosolized carbon nanotubes at occupationally relevant doses than healthy cells. *ACS Nano* 11, 7615–7625. <http://dx.doi.org/10.1021/acsnano.7b01992>. [PubMed: 28505409]
- Choudhury SR, Ordaz J, Lo CL, Damayanti NP, Zhou F, Irudayaraj J, 2017 From the cover: zinc oxide nanoparticles-induced reactive oxygen species promotes multimodal cyto- and epigenetic toxicity. *Toxicol. Sci* 156, 261–274. <http://dx.doi.org/10.1093/toxsci/kfw252>. [PubMed: 28115643]
- Chuang HC, Hsiao TC, Wu CK, Chang HH, Lee CH, Chang CC, Cheng TJ, Taiwan CardioPulmonary Research Group (T-CPR), 2013 Allergenicity and toxicology of inhaled silver nanoparticles in allergen-provocation mice models. *Int. J. Nanomed* 8, 4495–4506. <http://dx.doi.org/10.2147/IJN.S52239>.
- Colognato R, Bonelli A, Ponti J, Farina M, Bergamaschi E, Sabbioni E, Migliore L, 2008 Comparative genotoxicity of cobalt nanoparticles and ions on human peripheral leukocytes in vitro. *Mutagenesis* 23, 377–382. <http://dx.doi.org/10.1093/mutage/gen024>. [PubMed: 18504271]
- Cordelli E, Keller J, Eleuteri P, Villani P, Ma-Hock L, Schulz M, Landsiedel R, Pacchierotti F, 2017 No genotoxicity in rat blood cells upon 3- or 6-month inhalation exposure to CeO₂ or BaSO₄ nanomaterials. *Mutagenesis* 32, 13–22. <http://dx.doi.org/10.1093/mutage/gew005>. [PubMed: 26861493]

- Costa PM, Fadeel B, 2016 Emerging systems biology approaches in nanotoxicology: towards a mechanism-based understanding of nanomaterial hazard and risk. *Toxicol. Appl. Pharmacol* 299, 101–111. <http://dx.doi.org/10.1016/j.taap.2015.12.014>. [PubMed: 26721310]
- Costa PM, Gosens I, Williams A, Farcal L, Pantano D, Brown DM, Stone V, Cassee FR, Halappanavar S, Fadeel B, 2018 Transcriptional profiling reveals gene expression changes associated with inflammation and cell proliferation following short-term inhalation exposure to copper oxide nanoparticles. *J. Appl. Toxicol* 38, 385–397. <http://dx.doi.org/10.1002/jat.3548>. [PubMed: 29094763]
- Cveticanin J, Joksic G, Leskovic A, Petrovic S, Sobot AV, Neskovic O, 2010 Using carbon nanotubes to induce micronuclei and double strand breaks of the DNA in human cells. *Nanotechnology* 21, 015102 <http://dx.doi.org/10.1088/0957-4484/21/1/015102>. [PubMed: 19946169]
- Decan N, Wu D, Williams A, Bernatchez S, Johnston M, Hill M, Halappanavar S, 2016 Characterization of in vitro genotoxic, cytotoxic and transcriptomic responses following exposures to amorphous silica of different sizes. *Mutat. Res. Genet. Toxicol. Environ. Mutagen* 796, 8–22. <http://dx.doi.org/10.1016/j.mrgentox.2015.11.011>. [PubMed: 26778505]
- Di Bucchianico S, Fabbri MR, Misra SK, Valsami-Jones E, Berhanu D, Reip P, Bergamaschi E, Migliore L, 2013 Multiple cytotoxic and genotoxic effects induced in vitro by differently shaped copper oxide nanomaterials. *Mutagenesis* 28, 287–299. <http://dx.doi.org/10.1093/mutage/get014>. [PubMed: 23462852]
- Donaldson K, Murphy F, Schinwald A, Duffin R, Poland CA, 2011 Identifying the pulmonary hazard of high aspect ratio nanoparticles to enable their safety-by-design. *Nanomedicine (Lond. U.K.)* 6, 143–156. <http://dx.doi.org/10.2217/nnm.10.139>.
- Downs TR, Crosby ME, Hu T, Kumar S, Sullivan A, Sarlo K, Reeder B, Lynch M, Wagner M, Mills T, Pfuhrer S, 2012 Silica nanoparticles administered at the maximum tolerated dose induce genotoxic effects through an inflammatory reaction while gold nanoparticles do not. *Mutat. Res* 745, 38–50. <http://dx.doi.org/10.1016/j.mrgentox.2012.03.012>. [PubMed: 22504169]
- Driessen MD, Mues S, Vennemann A, Hellack B, Bannuscher A, Vimalakanthan V, Riebeling C, Ossig R, Wiemann M, Schneidenburger J, Kuhlbusch TA, Renard B, Luch A, Haase A, 2015 Proteomic analysis of protein carbonylation: a useful tool to unravel nanoparticle toxicity mechanisms. *Part. Fibre Toxicol* 12, 36 <http://dx.doi.org/10.1186/s12989-015-0108-2>. [PubMed: 26525058]
- Driscoll KE, 1996 Role of inflammation in the development of rat lung tumors in response to chronic particle exposure. *Inhal. Toxicol* 8, 139–153.
- Du Z, Zhao D, Jing L, Cui G, Jin M, Li Y, Liu X, Liu Y, Du H, Guo C, Zhou X, Sun Z, 2013 Cardiovascular toxicity of different sizes amorphous silica nanoparticles in rats after intratracheal instillation. *Cardiovasc. Toxicol* 13, 194–207. <http://dx.doi.org/10.1007/s12012-013-9198-y>. [PubMed: 23322373]
- Duffin R, Tran C, Clouter A, Brown D, MacNee W, Stone V, Donaldson K, 2001 The importance of surface area and specific reactivity in the acute pulmonary inflammatory response to particles. *Ann. Occup. Hyg* 46, 242–245. <http://dx.doi.org/10.1093/annhyg/mef684>.
- Dymacek J, Snyder-Talkington BN, Porter DW, Mercer RR, Wolfarth MG, Castranova V, Qian Y, Guo NL, 2015 mRNA and miRNA regulatory networks reflective of multi-walled carbon nanotube-induced lung inflammatory and fibrotic pathologies in mice. *Toxicol. Sci* 144, 51–64. <http://dx.doi.org/10.1093/toxsci/kfu262>. [PubMed: 25527334]
- Eastlake AC, Beaucham C, Martinez KF, Dahm MM, Sparks C, Hodson LL, Geraci CL, 2016 Refinement of the nanoparticle emission assessment technique into the nanomaterial exposure assessment technique (NEAT 2.0). *Occup. Environ. Hyg* 13, 708–717. <http://dx.doi.org/10.1080/15459624.2016.1167278>.
- Ellinger-Ziegelbauer H, Pauluhn J, 2009 Pulmonary toxicity of multi-walled carbon nanotubes (Baytubes) relative to alpha-quartz following a single 6h inhalation exposure of rats and a 3 months post-exposure period. *Toxicology* 266, 16–29. <http://dx.doi.org/10.1016/j.tox.2009.10.007>. [PubMed: 19836432]
- Eom HJ, Chatterjee N, Lee J, Choi J, 2014 Integrated mRNA and micro RNA profiling reveals epigenetic mechanism of differential sensitivity of Jurkat T cells to AGNPs and Ag ions. *Toxicol. Lett* 229, 311–318. <http://dx.doi.org/10.1016/j.toxlet.2014.05.019>. [PubMed: 24974767]

- Erdely A, Hulderman T, Salmen R, Liston A, Zeidler-Erdely PC, Schwegler-Berry D, Castranova V, Koyama S, Kim YA, Endo M, Simeonova PP, 2009 Cross-talk between lung and systemic circulation during carbon nanotube respiratory exposure. Potential biomarkers. *Nano Lett* 9, 36–43. <http://dx.doi.org/10.1021/nl801828z>. [PubMed: 19049393]
- Fadeel B, 2015 Systems biology in nanosafety research. *Nanomedicine (Lond. U.K.)* 10, 1039–1041. <http://dx.doi.org/10.2217/nmm.15.17>.
- Fatkhutdinova LM, Khaliullin TO, Vasil'yeva OL, Zalyalov RR, Mustafin IG, Kisin ER, Birch ME, Yanamala N, Shvedova AA, 2016 Fibrosis biomarkers in workers exposed to MWCNTs. *Toxicol. Appl. Pharmacol* 299, 125–131. <http://dx.doi.org/10.1016/j.taap.2016.02.016>. [PubMed: 26902652]
- Feng J, Liu H, Zhang L, Bhakoo K, Lu L, 2010 An insight into the metabolic responses of ultra-small superparamagnetic particles of iron oxide using metabolomic analysis of biofluids. *Nanotechnology* 21, 395101 <http://dx.doi.org/10.1088/0957-4484/21/39/395101>. [PubMed: 20820093]
- Flower NA, Brabu B, Revathy M, Gopalakrishnan C, Raja SV, Murugan SS, Kumaravel TS, 2012 Characterization of synthesized silver nanoparticles and assessment of its genotoxicity potentials using the alkaline comet assay. *Mutat. Res* 742, 61–65. <http://dx.doi.org/10.1016/j.mrgentox.2011.12.003>. [PubMed: 22178963]
- Gao G, Ze Y, Li B, Zhao X, Zhang T, Sheng L, Hu R, Gui S, Sang X, Sun Q, Cheng J, Cheng Z, Wang L, Tang M, Hong F, 2012 Ovarian dysfunction and gene-expressed characteristics of female mice caused by long-term exposure to titanium dioxide nanoparticles. *J. Hazard. Mater* 243, 19–27. <http://dx.doi.org/10.1016/j.jhazmat.2012.08.049>. [PubMed: 23131501]
- Genter MB, Newman NC, Shertzer HG, Ali SF, Bolon B, 2012 Distribution and systemic effects of intranasally administered 25 nm silver nanoparticles in adult mice. *Toxicol. Pathol* 40, 1004–1013. <http://dx.doi.org/10.1177/0192623312444470>. [PubMed: 22549977]
- Geraci C, Heidel D, Sayes C, Hodson L, Schulte P, Eastlake A, Brenner S, 2015 Perspectives on the design of safer nanomaterials and manufacturing processes. *J. Nanopart. Res* 17, 366–379. <http://dx.doi.org/10.1007%2Fs11051-015-3152-9>. [PubMed: 26435688]
- Ghosh M, Bandyopadhyay M, Mukherjee A, 2010 Genotoxicity of titanium dioxide (TiO₂) nanoparticles at two trophic levels: plant and human lymphocytes. *Chemosphere* 81, 1253–1262. <http://dx.doi.org/10.1016/j.chemosphere.2010.09.022>. [PubMed: 20884039]
- Ghosh MJM, Sinha S, Chakraborty A, Mallick SK, Bandyopadhyay M, Mukherjee A, 2012 In vitro and in vivo genotoxicity of silver nanoparticles. *Mutat. Res* 749, 60–69. <http://dx.doi.org/10.1016/j.mrgentox.2012.08.007>. [PubMed: 22960309]
- Ghosh M, Chakraborty A, Mukherjee A, 2013 Cytotoxic, genotoxic and the hemolytic effect of titanium dioxide (TiO₂) nanoparticles on human erythrocyte and lymphocyte cells in vitro. *J. Appl. Toxicol* 33, 1097–1110. <http://dx.doi.org/10.1002/jat.2863>. [PubMed: 23616399]
- Ghosh M, Öner D, Poels K, Tabish AM, Vlaanderen J, Pronk A, Kuijpers E, Lan Q, Vermeulen R, Bekaert B, Hoet PH, Godderis L, 2017 Changes in DNA methylation induced by multi-walled carbon nanotube exposure in the workplace. *Nanotoxicology* 11, 1195–1210. <http://dx.doi.org/10.1080/17435390.2017.1406169>. [PubMed: 29191063]
- Glass DC, Mazhar M, Xiang S, Dean P, Simpson P, Priestly B, Plebanski M, Abramson M, Sim MR, Dennekamp M, 2017 Immunological effects among workers who handle engineered nanoparticles. *Occup. Environ. Med* 74, 868–876. <http://dx.doi.org/10.1136/oemed-2016-104111>. [PubMed: 28847906]
- Gong C, Tao G, Yang L, Liu J, Liu Q, Zhuang Z, 2010 SiO₂ nanoparticles induce global genomic hypomethylation in HaCaT cells. *Biochem. Biophys. Res. Commun* 397, 397–400. <http://dx.doi.org/10.1016/j.bbrc.2010.05.076>. [PubMed: 20501321]
- Gong C, Tao G, Yang L, Liu J, Liu Q, Zhuang Z, 2012 Methylation of PARP-1 promoter involved in the regulation of nano-SiO₂-induced decrease of PARP-1 mRNA expression. *Toxicol. Lett* 209, 264–269. <http://dx.doi.org/10.1016/j.toxlet.2012.01.007>. [PubMed: 22265868]
- Gosens I, Cassee FR, Zanella M, Manodori L, Brunelli A, Costa AL, Bokkers BG, de Jong WH, Brown D, Hristozov D, Stone V, 2016 Organ burden and pulmonary toxicity of nano-sized copper (II) oxide particles after short-term inhalation exposure. *Nanotoxicology* 10, 1084–1095. <http://dx.doi.org/10.3109/17435390.2016.1172678>. [PubMed: 27132941]

- Guichard Y, Fontana C, Chavinier E, Terzetti F, Gaté L, Binet S, Darne C, 2016 Cytotoxic and genotoxic evaluation of different synthetic amorphous silica nanomaterials in the V79 cell line. *Toxicol. Ind. Health* 32, 1639–1650. <http://dx.doi.org/10.1177/0748233715572562>. [PubMed: 25757481]
- Gulson B, McCall M, Korsch M, Gomez L, Casey P, Oytam Y, Taylor A, McCulloch M, Trotter J, Kinsley L, Greenoak G, 2010 Small amounts of zinc from zinc oxide particles in sunscreens applied outdoors are absorbed through human skin. *Toxicol. Sci* 118, 140–149. <http://dx.doi.org/10.1093/toxsci/kfq243>. [PubMed: 20705894]
- Gulson B, Wong H, Korsch M, Gomez L, Casey P, McCall M, McCulloch M, Trotter J, Stauber J, Greenoak G, 2012 Comparison of dermal absorption of zinc from different sunscreen formulations and differing UV exposure based on stable isotope tracing. *Sci. Total Environ* 420, 313–318. <http://dx.doi.org/10.1016/j.scitotenv.2011.12.046>. [PubMed: 22316633]
- Gulumian M, Verbeek J, Andraos C, Sanabria N, de Jager P, 2016 Systematic review of screening and surveillance programs to protect workers from nanomaterials. *PLoS One* 11, e0166071 <http://dx.doi.org/10.1371/journal.pone.0166071>. [PubMed: 27829014]
- Gwinn MR, Vallyathan V, 2006 Nanoparticles: health effects—pros and cons. *Environ. Health Perspect* 114, 1818–1825. <http://dx.doi.org/10.1289/ehp.8871>. [PubMed: 17185269]
- Hadrup N, Lam HR, Loeschner K, Mortensen A, Larsen EH, Frandsen H, 2012 Nanoparticulate silver increases uric acid and allantoin excretion in rats, as identified by metabolomics. *J. Appl. Toxicol* 32, 929–933. <http://dx.doi.org/10.1002/jat.2779>. [PubMed: 22610381]
- Halappanavar S, Jackson P, Williams A, Jensen KA, Hougaard KS, Vogel U, Yauk CL, Wallin H, 2011 Pulmonary response to surface-coated nanotitanium dioxide particles includes induction of acute phase response genes, inflammatory cascades, and changes in microRNAs: a toxicogenomic study. *Environ. Mol. Mutagen* 52, 425–439. <http://dx.doi.org/10.1002/em.20639>. [PubMed: 21259345]
- Hamilton RF, Wu N, Porter D, Buford M, Wolfarth M, Holian A, 2009 Particle length-dependent titanium dioxide nanomaterials toxicity and bioactivity. *Part. Fibre Toxicol* 6, 35 <http://dx.doi.org/10.1186/1743-8977-6-35>. [PubMed: 20043844]
- He X, Zhang H, Ma Y, Bai W, Zhang Z, Lu K, Ding Y, Zhao Y, Chai Z, 2010 Lung deposition and extrapulmonary translocation of nano-ceria after intratracheal instillation. *Nanotechnology* 21, 285103 <http://dx.doi.org/10.1088/0957-4484/21/28/285103>. [PubMed: 20562477]
- Hemeryck LY, Moore SA, Vanhaecke L, 2016 Mass spectrometric mapping of the DNA adductome as a means to study genotoxin exposure, metabolism, and effect. *Anal. Chem* 88, 7436–7446. <http://dx.doi.org/10.1021/acs.analchem.6b00863>. [PubMed: 27362284]
- Holland NA, Becak DP, Shannahan JH, Brown JM, Carratt SA, Winkle L, Pinkerton KE, Wang CM, Munusamy P, Baer DR, Sumner SJ, Fennell TR, Lust RM, Wingard CJ, 2015 Cardiac ischemia reperfusion injury following instillation of 20 nm citrate-capped nanosilver. *J. Nanomed. Nanotechnol* S6 <http://dx.doi.org/10.4172/2157-7439.S6-006.006>.
- Holland NA, Thompson LC, Vidanapathirana AK, Urankar RN, Lust RM, Fennell TR, Wingard CJ, 2016 Impact of pulmonary exposure to gold core silver nanoparticles of different size and capping agents on cardiovascular injury. *Part. Fibre Toxicol* 13, 1 <http://dx.doi.org/10.1186/s12989-016-0159-z>. [PubMed: 26746196]
- Husain M, Saber AT, Guo C, Jacobsen NR, Jensen KA, Yauk CL, Williams A, Vogel U, Wallin H, Halappanavar S, 2013 Pulmonary instillation of low doses of titanium dioxide nanoparticles in mice leads to particle retention and gene expression changes in the absence of inflammation. *Toxicol. Appl. Pharmacol* 269, 250–262. <http://dx.doi.org/10.1016/j.taap.2013.03.018>. [PubMed: 23557971]
- Iavicoli I, Leso V, Fontana L, Bergamaschi A, 2011 Toxicological effects of titanium dioxide nanoparticles: a review of in vitro mammalian studies. *Eur. Rev. Med. Pharmacol. Sci* 15, 481–508. <http://dx.doi.org/10.1155/2012/964381>. [PubMed: 21744743]
- Iavicoli I, Leso V, Bergamaschi A, 2012 Toxicological effects of titanium dioxide nanoparticles: a review of in vivo studies. *J. Nanomater* 36 <http://dx.doi.org/10.1155/2012/964381> Article ID 964381.
- Iavicoli I, Leso V, Manno M, Schulte PA, 2014a Biomarkers of nanomaterial exposure and effect: current status. *J. Nanopart. Res* 16, 2302 <http://dx.doi.org/10.1007/s11051-014-2302-9>.

- Iavicoli I, Leso V, Ricciardi W, Hodson LL, Hoover MD, 2014b Opportunities and challenges of nanotechnology in the green economy. *Environ. Health* 13, 78 <http://dx.doi.org/10.1186/1476-069X-13-78>. [PubMed: 25294341]
- Iavicoli I, Fontana L, Nordberg G, 2016 The effects of nanoparticles on the renal system. *Crit. Rev. Toxicol* 46, 490–560. <http://dx.doi.org/10.1080/10408444.2016.1181047>. [PubMed: 27195425]
- IPCS, 2001 Environmental Health Criteria 222. Biomarkers in Risk Assessment: Validity and Validation. International Programme on Chemical Safety (IPCS) (Accessed 30 January 2018). <http://www.inchem.org/documents/ehc/ehc/ehc222.htm>.
- Ivask A, Voelcker NH, Seabrook SA, Hor M, Kirby JK, Fenech M, Davis TP, Ke PC, 2015 DNA melting and genotoxicity induced by silver nanoparticles and graphene. *Chem. Res. Toxicol* 28, 1023–1035. <http://dx.doi.org/10.1021/acs.chemrestox.5b00052>. [PubMed: 25781053]
- Jiménez-Villarreal J, Rivas-Armendáriz DI, Arellano Pérez-Vertti RD, Olivas Calderón E, García-Garza R, Betancourt-Martínez ND, Serrano-Gallardo LB, Morán-Martínez J, 2017 Relationship between lymphocyte DNA fragmentation and dose of iron oxide (Fe₂O₃) and silicon oxide (SiO₂) nanoparticles. *Genet. Mol. Res* 16, 1–12. <http://dx.doi.org/10.4238/gmr16019206>.
- Kang SJ, Kim BM, Lee YJ, Chung HW, 2008 Titanium dioxide nanoparticles trigger p53-mediated damage response in peripheral blood lymphocytes. *Environ. Mol. Mutagen* 49, 399–405. <http://dx.doi.org/10.1002/em.20399>. [PubMed: 18418868]
- Kang SJ, Lee YJ, Kim BM, Choi YJ, Chung HW, 2011 Cytotoxicity and genotoxicity of titanium dioxide nanoparticles in UVA-irradiated normal peripheral blood lymphocytes. *Drug Chem. Toxicol* 34, 277–284. <http://dx.doi.org/10.3109/01480545.2010.546800>. [PubMed: 21649482]
- Kauppinen T, Uuksulainen S, Saalo A, Mäkinen I, Pukkala E, 2014 Use of the Finnish information system on occupational exposure (FINJEM) in epidemiologic, surveillance, and other applications. *Ann. Occup. Hyg* 58, 380–396. <http://dx.doi.org/10.1093/annhyg/met074>. [PubMed: 24401793]
- Kim S, Ryu DY, 2013 Silver nanoparticle-induced oxidative stress, genotoxicity and apoptosis in cultured cells and animal tissues. *J. Appl. Toxicol* 33, 78–89. <http://dx.doi.org/10.1002/jat.2792>. [PubMed: 22936301]
- Kim JS, Yu IJ, 2014 Single-wall carbon nanotubes (SWCNT) induce cytotoxicity and genotoxicity produced by reactive oxygen species (ROS) generation in phytohemagglutinin (PHA)-stimulated male human peripheral blood lymphocytes. *J. Toxicol. Environ. Health Part A* 77, 1141–1153. <http://dx.doi.org/10.1080/15287394.2014.917062>. [PubMed: 25119736]
- Kim EY, Schulz R, Swantek P, Kunstman K, Malim MH, Wolinsky SM, 2012 Gold nanoparticle-mediated gene delivery induces widespread changes in the expression of innate immunity genes. *Gene Ther* 19, 347–353. <http://dx.doi.org/10.1038/gt.2011.95>. [PubMed: 21697957]
- Kitchin KT, Stirdivant S, Robinette BL, Castellon BT, Liang X, 2017 Metabolomic effects of CeO₂, SiO₂ and CuO metal oxide nanomaterials on HepG2 cells. *Part. Fibre Toxicol* 14, 1–16. <http://dx.doi.org/10.1186/s12989-017-0230-4>. [PubMed: 28069023]
- Klaper R, Arndt D, Bozich J, Dominguez G, 2014 Molecular interactions of nanomaterials and organisms: defining biomarkers for toxicity and high-throughput screening using traditional and next-generation sequencing approaches. *Analyst* 139, 882–895. <http://dx.doi.org/10.1039/c3an01644g>. [PubMed: 24343342]
- Kongseng S, Yoovathaworn K, Wongprasert K, Chunhabundit R, Sukwong P, Pissuwan D, 2016 Cytotoxic and inflammatory responses of TiO₂ nanoparticles on human peripheral blood mononuclear cells. *J. Appl. Toxicol* 36, 1364–1373. <http://dx.doi.org/10.1002/jat.3342>. [PubMed: 27225715]
- Kreider T, Halperin W, 2011 Engineered nanomaterials: learning from the past, planning for the future. *J. Occup. Environ. Med* 53, S108–S112. <http://dx.doi.org/10.1097/JOM.0b013e31821b146a>. [PubMed: 21654410]
- Kreyling WG, Semmler-Behnke M, Seitz J, Scymczak W, Wenk A, Mayer P, Takenaka S, Oberdörster G, 2009 Size dependence of the translocation of inhaled iridium and carbon nanoparticle aggregates from the lung of rats to the blood and secondary target organs. *Inhal. Toxicol* 21, 55–60. <http://dx.doi.org/10.1080/08958370902942517>.
- Kuempel ED, Jaurand MC, Møller P, Morimoto Y, Kobayashi N, Pinkerton KE, Sargent LM, Vermeulen RC, Fubini B, Kane AB, 2017 Evaluating the mechanistic evidence and key data gaps

- in assessing the potential carcinogenicity of carbon nanotubes and nanofibers in humans. *Crit. Rev. Toxicol* 47, 1–58. <http://dx.doi.org/10.1080/10408444.2016.1206061>. [PubMed: 27537422]
- Larese Filon F, Crosera M, Mauro M, Baracchini E, Bovenzi M, Montini T, Fornasiero P, Adami G, 2016 Palladium nanoparticles exposure: evaluation of permeation through damaged and intact human skin. *Environ. Pollut* 214, 497–503. <http://dx.doi.org/10.1016/j.envpol.2016.04.077>. [PubMed: 27131807]
- Lee JS, Choi YC, Shin JH, Lee JH, Lee Y, Park SY, Baek JE, Park JD, Ahn K, Yu IJ, 2015 Health surveillance study of workers who manufacture multi-walled carbon nanotubes. *Nanotoxicology* 9, 802–811. <http://dx.doi.org/10.3109/17435390.2014.978404>. [PubMed: 25395166]
- Lei R, Wu C, Yang B, Ma H, Shi C, Wang Q, Wang Q, Yuan Y, Liao M, 2008 Integrated metabolomic analysis of the nano-sized copper particle-induced hepatotoxicity and nephrotoxicity in rats: a rapid in vivo screening method for nanotoxicity. *Toxicol. Appl. Pharmacol* 232, 292–301. <http://dx.doi.org/10.1016/j.taap.2008.06.026>. [PubMed: 18706438]
- Leso V, Fontana L, Mauriello MA, Iavicoli I, 2017 Occupational risk assessment of engineered nanomaterials: limits, challenges and opportunities. *Curr. Nanosci* 13, 55–78. <http://dx.doi.org/10.2174/1573413712666161017114934>.
- Li Z, Hulderman T, Salmen R, Chapman R, Leonard SS, Young SH, Shvedova A, Luster MI, Simeonova PP, 2007 Cardiovascular effects of pulmonary exposure to single-wall carbon nanotubes. *Environ. Health Perspect* 115, 377–382. [PubMed: 17431486]
- Li S, Wang H, Qi Y, Tu J, Bai Y, Tian T, Huang N, Wang Y, Xiong F, Lu Z, Xiao Z, 2011a Assessment of nanomaterial cytotoxicity with SOLiD sequencing-based microRNA expression profiling. *Biomaterials* 32, 9021–9030. <http://dx.doi.org/10.1016/j.biomaterials.2011a.08.033>. [PubMed: 21889204]
- Li S, Wang Y, Wang H, Bai Y, Liang G, Wang Y, Huang N, Xiao Z, 2011b MicroRNAs as participants in cytotoxicity of CdTe quantum dots in NIH/3T3. *Biomaterials* 32, 3807–3814. <http://dx.doi.org/10.1016/j.biomaterials.2011.01.074>. [PubMed: 21376392]
- Li B, Ze Y, Sun Q, Zhang T, Sang X, Cui Y, Wang X, Gui S, Tan D, Zhu M, Zhao X, Sheng L, Wang L, Hong F, Tang M, 2013 Molecular mechanisms of nanosized titanium dioxide-induced pulmonary injury in mice. *PLoS One* 8, e55563 <http://dx.doi.org/10.1371/journal.pone.0055563>. [PubMed: 23409001]
- Li L, Liu T, Fu C, Tan L, Meng X, Liu H, 2015 Biodistribution, excretion, and toxicity of mesoporous silica nanoparticles after oral administration depend on their shape. *Nanomedicine* 11, 1915–1924. <http://dx.doi.org/10.1016/j.nano.2015.07.004>. [PubMed: 26238077]
- Li J, Tian M, Cui L, Dwyer J, Fullwood NJ, Shen H, Martin FL, 2016 Low-dose carbon-based nanoparticle-induced effects in A549 lung cells determined by bio spectroscopy are associated with increases in genomic methylation. *Sci. Rep* 6, 20207 <http://dx.doi.org/10.1038/srep20207>. [PubMed: 26831369]
- Li Y, Yan J, Ding W, Chen Y, Pack LM, Chen T, 2017 Genotoxicity and gene expression analyses of liver and lung tissues of mice treated with titanium dioxide nanoparticles. *Mutagenesis* 32, 33–46. <http://dx.doi.org/10.1093/mutage/gew065>. [PubMed: 28011748]
- Liao M, Liu H, 2012 Gene expression profiling of nephrotoxicity from copper nanoparticles in rats after repeated oral administration. *Environ. Toxicol. Pharmacol* 34, 67–80. <http://dx.doi.org/10.1016/j.etap.2011.05.014>. [PubMed: 22465980]
- Liao HY, Chung YT, Lai CH, Wang SL, Chiang HC, Li LA, Tsou TC, Li WF, Lee HL, Wu WT, Lin MH, Hsu JH, Ho JJ, Chen CJ, Shih TS, Lin CC, Liou SH, 2014 Six-month follow-up study of health markers of nanomaterials among workers handling engineered nanomaterials. *Nanotoxicology* 8, 100–110. <http://dx.doi.org/10.3109/17435390.2013.858793>.
- Lin B, Zhang H, Lin Z, Fang Y, Tian L, Yang H, Yan J, Liu H, Zhang W, Xi Z, 2013 Studies of single-walled carbon nanotubes-induced hepatotoxicity by NMR-based metabolomics of rat blood plasma and liver extracts. *Nanoscale Res. Lett* 8, 1–11. <http://dx.doi.org/10.1186/1556-276X-8-236>. [PubMed: 23279756]
- Lindberg HK, Falck GC, Catalan J, Koivisto AJ, Suhonen S, Jarventaus H, Rossi EM, Nykasenoja H, Peltonen Y, Moreno C, Alenius H, Tuomi T, Savolainen KM, Norppa H, 2012 Genotoxicity of inhaled nanosized TiO₂ in mice. *Mutat. Res* 745, 58–64. <http://dx.doi.org/10.1016/j.mrgentox.2011.10.011>. [PubMed: 22094288]

- Liou SH, Tsou TC, Wang AL, Li LA, Chiang HC, Li WF, Lin PP, Lai CH, Lee HL, Lin MH, Hsu JH, Chen CR, Shih TS, Liao AY, Chung YT, 2012 Epidemiological study of health hazards among workers handling engineered nanomaterials. *J. Nanopart. Res* 14, 1–15. <http://dx.doi.org/10.1007/s11051-012-0878-5>. [PubMed: 22448125]
- Liou SH, Tsai CSJ, Pelcova D, Schubauer-Berigan MK, Schulte PA, 2015 Assessing the first wave of epidemiological studies of nanomaterial workers. *J. Nanopart. Res* 17, 1–19. <http://dx.doi.org/10.1007/s11051-015-3219-7>.
- Liou SH, Chen YC, Liao HY, Wang CJ, Chen JS, Lee HL, 2016 Increased levels of oxidative stress biomarkers in metal oxides nanomaterial-handling workers. *Biomarkers* 21, 600–606. <http://dx.doi.org/10.3109/1354750X.2016.1160432>. [PubMed: 27010249]
- Liou SH, Wu WT, Liao HY, Chen CY, Tsai CY, Jung WT, Lee HL, 2017 Global DNA methylation and oxidative stress biomarkers in workers exposed to metal oxide nanoparticles. *J. Hazard. Mater* 331, 329–335. <http://dx.doi.org/10.1016/j.jhazmat.2017.02.042>. [PubMed: 28273583]
- Liu X, Sun J, 2013. Time-course effects of intravenously administered silica nanoparticles on blood coagulation and endothelial function in rats. *J. Nanosci. Nanotechnol* 13, 222–228. <http://dx.doi.org/10.1166/jnn.2013.6910>. [PubMed: 23646720]
- Lu X, Tian Y, Zhao Q, Jin T, Xiao S, Fan X, 2011 Integrated metabonomics analysis of the size-response relationship of silica nanoparticles-induced toxicity in mice. *Nanotechnology* 22, 055101 <http://dx.doi.org/10.1088/0957-4484/22/5/055101>. [PubMed: 21178262]
- Lu X, Miousse IR, Pirela SV, Moore JK, Melnyk S, Koturbash I, Demokritou P, 2016 In vivo epigenetic effects induced by engineered nanomaterials: a case study of copper oxide and laser printeremitted engineered nanoparticles. *Nanotoxicology* 10, 629–639. <http://dx.doi.org/10.3109/17435390.2015.1108473>. [PubMed: 26559097]
- Magdolenova Z, Drlickova M, Henjum K, Rundén-Pran E, Tulinska J, Bilanicova D, Pojana G, Kazimirova A, Barancokova M, Kuricova M, Liskova A, Staruchova M, Ciampor F, Vavra I, Lorenzo Y, Collins A, Rinna A, Fjellsbø L, Volkovova K, Marcomini A, Amiry-Moghaddam M, Dusinska M, 2015 Coating-dependent induction of cytotoxicity and genotoxicity of iron oxide nanoparticles. *Nanotoxicology* 9, 44–56. <http://dx.doi.org/10.3109/17435390.2013.847505>. [PubMed: 24228750]
- Mahmoud A, Ezgi Ö, Merve A, Özhan G, 2016 In vitro toxicological assessment of magnesium oxide nanoparticle exposure in several mammalian cell types. *Int. J. Toxicol* 35, 429–437. <http://dx.doi.org/10.1177/1091581816648624>. [PubMed: 27177543]
- Manno M, Viau C, in collaboration with, Cocker J, Colosio C, Lowry L, Mutti A, Nordberg M, Wang S, 2010 Biomonitoring for occupational health risk assessment (BOHRA). *Toxicol. Lett* 192, 3–16. <http://dx.doi.org/10.1016/j.toxlet.2009.05.001>. [PubMed: 19446015]
- Martins ADC, Jr, Azevedo LF, de Souza Rocha CC, Carneiro MFH, Venancio VP, de Almeida MR, Antunes LMG, de Carvalho Hott R, Rodrigues JL, Ogunjimi AT, Adeyemi JA, Barbosa F, Jr, 2017 Evaluation of distribution, redox parameters, and genotoxicity in Wistar rats co-exposed to silver and titanium dioxide nanoparticles. *J. Toxicol. Environ. Health Part A* 80, 1156–1165. <http://dx.doi.org/10.1080/15287394.2017.1357376>. [PubMed: 28891756]
- Matysiak M, Kapka-Skrzypczak L, Brzóška K, Gutleb AC, Kruszewski M, 2016 Proteomic approach to nanotoxicity. *J. Proteom* 137, 35–44. <http://dx.doi.org/10.1016/j.jprot.2015.10.025>.
- Maurer MM, Donohoe GC, Maleki H, Yi J, McBride C, Nurkiewicz TR, Valentine SJ, 2016 Comparative plasma proteomic studies of pulmonary TiO₂ nanoparticle exposure in rats using liquid chromatography tandem mass spectrometry. *J. Proteom* 130, 85–93. <http://dx.doi.org/10.1016/j.jprot.2015.09.010>.
- Mercer RR, Scabilloni JF, Hubbs AF, Wang L, Battelli LA, McKinney W, Castranova V, Porter DW, 2013 Extrapulmonary transport of MWCNT following inhalation exposure. *Part. Fibre Toxicol* 10, 38 <http://dx.doi.org/10.1186/1743-8977-10-38>. [PubMed: 23927530]
- Methner M, Hodson L, Geraci C, 2010a Nanoparticle emission assessment technique (NEAT) for the identification and measurement of potential inhalation exposure to engineered nanomaterials—part. *J. Occup. Environ. Hyg* 7, 127–132. <http://dx.doi.org/10.1080/15459620903476355>. [PubMed: 20017054]
- Methner M, Hodson L, Dames A, Geraci C, 2010b Nanoparticle Emission Assessment Technique (NEAT) for the identification and measurement of potential inhalation exposure to engineered

nanomaterials-part b: results from 12 field studies. *J. Occup. Environ. Hyg* 7, 163–176. <http://dx.doi.org/10.1080/15459620903508066>. [PubMed: 20063229]

- Methner M, Beaucham C, Crawford C, Hodson L, Geraci C, 2012 Field application of the Nanoparticle Emission Assessment Technique (NEAT): task-based air monitoring during the processing of engineered nanomaterials (ENM) at four facilities. *J. Occup. Environ. Hyg* 9, 543–555. <http://dx.doi.org/10.1080/15459624.2012.699388>. [PubMed: 22816668]
- Monopoli MP, Aberg C, Salvati A, Dawson KA, 2012 Biomolecular coronas provide the biological identity of nanosized materials. *Nat. Nanotechnol* 7, 779–786. <http://dx.doi.org/10.1038/nnano.2012.207>. [PubMed: 23212421]
- Moretto A, 2015 Exposure assessment for chemical and physical agents. *Handb. Clin. Neurol* 131, 47–59. <http://dx.doi.org/10.1016/B978-0-444-62627-1.00004-4>. [PubMed: 26563782]
- Müller L, Steiner SK, Rodriguez-Lorenzo L, Petri-Fink A, Rothen-Rutishauser B, Latzin P, 2018 Exposure to silver nanoparticles affects viability and function of natural killer cells, mostly via the release of ions. *Cell Biol. Toxicol* 34, 167–176. <http://dx.doi.org/10.1007/s10565-017-9403-z>. [PubMed: 28721573]
- Murphy A, Casey A, Byrne G, Chambers G, Howe O, 2016 Silver nanoparticles induce pro-inflammatory gene expression and inflammasome activation in human monocytes. *J. Appl. Toxicol* 36, 1311–1320. <http://dx.doi.org/10.1002/jat.3315>. [PubMed: 26968431]
- Mytych J, Zebrowski J, Lewinska A, Wnuk M, 2017 Prolonged effects of silver nanoparticles on p53/p21 pathway-mediated proliferation, DNA damage response, and methylation parameters in HT22 hippocampal neuronal cells. *Mol. Neurobiol* 54, 1285–1300. <http://dx.doi.org/10.1007/s12035-016-9688-6>. [PubMed: 26843106]
- Nagano T, Higashisaka K, Kunieda A, Iwahara Y, Tanaka K, Nagano K, Abe Y, Kamada H, Tsunoda S, Nabeshi H, Yoshikawa T, Yoshioka Y, Tsutsumi Y, 2013 Liver-specific microRNAs as biomarkers of nanomaterial-induced liver damage. *Nanotechnology* 24, 405102 <http://dx.doi.org/10.1088/0957-4484/24/40/405102>. [PubMed: 24029385]
- Nalabotu SK, Kolli MB, Triest WE, Ma JY, Manne ND, Katta A, Addagarla HS, Rice KM, Blough ER, 2011 Intratracheal instillation of cerium oxide nanoparticles induces hepatic toxicity in male Sprague-Dawley rats. *Int. J. Nanomed* 6, 2327–2335. <http://dx.doi.org/10.2147/IJN.S25119>.
- Nanosolutions, 2018 <http://nanosolutionsfp7.com/nanosolutions-explained/> (Accessed 30 January 2018).
- Navya PN, Daima HK, 2016 Rational engineering of physicochemical properties of nanomaterials for biomedical applications with nanotoxicological perspectives. *Nano Conver* 3, 1 <http://dx.doi.org/10.1186/s40580-016-0064-z>. [PubMed: 28191411]
- Ng CT, Yong LQ, Hande MP, Ong CN, Yu LE, Bay BH, Baeg GH, 2017 Zinc oxide nanoparticles exhibit cytotoxicity and genotoxicity through oxidative stress responses in human lung fibroblasts and *Drosophila melanogaster*. *Int. J. Nanomed* 12, 1621–1637. <http://dx.doi.org/10.2147/IJN.S124403>.
- NIOSH, 2009 Approaches to Safe Nanotechnology: Managing the Health and Safety Concerns Associated With Engineered Nanomaterials U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Atlanta (Accessed 30 January 2018).. <https://www.cdc.gov/niosh/docs/2009-125/pdfs/2009-125.pdf>.
- NIOSH, 2013 Occupational Exposure to Carbon Nanotubes and Nanofibers U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Cincinnati, OH In: <https://www.cdc.gov/niosh/docs/2013-145/pdfs/2013-145.pdf>.
- NIOSH, 2017 The NIOSH Occupational Exposure Banding Process: Guidance for the Evaluation of Chemical Hazards. External Review Draft U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Atlanta (Accessed 31 January 2018).. http://www.cdc.gov/niosh/docket/review/docket290/pdfs/clean-cib-niosh-oebprocess-guidancefortheevaluationofchemicalhazards_3.8.17.pdf.

- Nurkiewicz TR, Porter DW, Hubbs AF, Cumpston JL, Chen BT, Frazer DG, Castranova V, 2008 Nanoparticle inhalation augments particle-dependent systemic microvascular dysfunction. Part. Fibre Toxicol 5, 1 <http://dx.doi.org/10.1186/1743-8977-5-1>. [PubMed: 18269765]
- Oberdörster G, Yu CP, 1999 Lung dosimetry—considerations for noninhalation studies. Exp. Lung Res 25, 1–6. <http://dx.doi.org/10.1080/019021499270385>. [PubMed: 10027075]
- Oberdörster G, Ferin J, Gelein R, Soderholm SC, Finkelstein J, 1992 Role of the alveolar macrophage in lung injury: studies with ultrafine particles. Environ. Health Perspect 97, 193–199. <http://dx.doi.org/10.2307/3431353>. [PubMed: 1396458]
- Oberdörster G, Maynard A, Donaldson K, Castranova V, Fitzpatrick J, Ausman K, Carter J, Karn B, Kreyling W, Lai D, Olin S, Monteiro-Riviere N, Warheit D, Yang H, I.L.S.I. Research Foundation/Risk Science Institute Nanomaterial Toxicity Screening Working Group, 2005 Principles for characterizing the potential human health effects from exposure to nanomaterials: elements of a screening strategy. Part. Fibre Toxicol 2, 8 <http://dx.doi.org/10.1186/1743-8977-2-8>. [PubMed: 16209704]
- OECD, 2017 Revised Guidance Document on Developing and Assessing Adverse Outcome Pathways. Series on Testing and Assessment, No. 184. 2013. ENM/JM/MONO, pp.6 (Accessed 30 January 2018).. [http://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=env/jm/mono\(2013\)6&doclanguage=en](http://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=env/jm/mono(2013)6&doclanguage=en).
- Öner D, Moisse M, Ghosh M, Duca RC, Poels K, Luyts K, Putzeys E, Cokic SM, Van Landuyt K, Vanoirbeek J, Lambrechts D, Godderis L, Hoet PH, 2017 Epigenetic effects of carbon nanotubes in human monocytic cells. Mutagenesis 32, 181–191. <http://dx.doi.org/10.1093/mutage/gew053>. [PubMed: 28011750]
- Orecchioni M, Bedognetti D, Newman L, Fuoco C, Spada F, Hendrickx W, Marincola FM, Sgarrella F, Rodrigues AF, Ménard-Moyon C, Cesareni G, Kostarelos K, Bianco A, Delogu LG, 2017 Single-cell mass cytometry and transcriptome profiling reveal the impact of graphene on human immune cells. Nat. Commun 8, 1109 <http://dx.doi.org/10.1038/s41467-017-01015-3>. [PubMed: 29061960]
- Osmond-McLeod MJ, Oytam Y, Kirby JK, Gomez-Fernandez L, Baxter B, McCall MJ, 2014 Dermal absorption and short-term biological impact in hairless mice from sunscreens containing zinc oxide nano- or larger particles. Nanotoxicology 8 (Suppl. 1), 72–84. <http://dx.doi.org/10.3109/17435390.2013.855832>.
- Pacurari M, Qian Y, Porter DW, Wolfarth M, Wan Y, Luo D, Ding M, Castranova V, Guo NL, 2011 Multi-walled carbon nanotube-induced gene expression in the mouse lung: association with lung pathology. Toxicol. Appl. Pharmacol 255, 18–31. <http://dx.doi.org/10.1016/j.taap.2011.05.012>. [PubMed: 21624382]
- Paino IM, Marangoni VS, de Oliveira Rde C, Antunes LM, Zucolotto V, 2012 Cyto and genotoxicity of gold nanoparticles in human hepatocellular carcinoma and peripheral blood mononuclear cells. Toxicol. Lett 215, 119–125. <http://dx.doi.org/10.1016/j.toxlet.2012.09.025>. [PubMed: 23046612]
- Palomino-Schätzlein M, García H, Gutiérrez-Carcedo P, Pineda-Lucena A, Herance JR, 2017 Assessment of gold nanoparticles on human peripheral blood cells by metabolic profiling with 1H-NMR spectroscopy, a novel translational approach on a patient-specific basis. PLoS One 12, e0182985 <http://dx.doi.org/10.1371/journal.pone.0182985>. [PubMed: 28793337]
- Parayanthala Valappil M, Santhakumar S, Arumugam S, 2014 Determination of oxidative stress related toxicity on repeated dermal exposure of hydroxyapatite nanoparticles in rats. Int. J. Biomater 2014, 476942 <http://dx.doi.org/10.1155/2014/476942>. [PubMed: 25587279]
- Park KH, Chhowalla M, Iqbal Z, Sesti F, 2003 Single-walled carbon nanotubes are a new class of ion channel blockers. J. Biol. Chem 278, 50212–50216. [PubMed: 14522977]
- Park EJ, Yoon J, Choi K, Yi J, Park K, 2009 Induction of chronic inflammation in mice treated with titanium dioxide nanoparticles by intratracheal instillation. Toxicology 260, 37–46. <http://dx.doi.org/10.1016/j.tox.2009.03.005>. [PubMed: 19464567]
- Park EJ, Kim H, Kim Y, Yi J, Choi K, Park K, 2010 Inflammatory responses may be induced by a single intratracheal instillation of iron nanoparticles in mice. Toxicology 275, 65–71. <http://dx.doi.org/10.1016/j.tox.2010.06.002>. [PubMed: 20540983]
- Park HS, Kim SJ, Lee TJ, Kim GY, Meang E, Hong JS, Kim SH, Koh SB, Hong SG, Sun YS, Kang JS, Kim YR, Kim MK, Jeong J, Lee JK, Son WC, Park JH, 2014 A 90-day study of sub-chronic oral

toxicity of 20 nm positively charged zinc oxide nanoparticles in Sprague Dawley rats. *Int. J. Nanomed* 9, 93–107. <http://dx.doi.org/10.2147/IJN.S57927>.

- Parveen A, Rizvi SH, Gupta A, Singh R, Ahmad I, Mahdi F, Mahdi AA, 2012 NMR-based metabonomics study of sub-acute hepatotoxicity induced by silica nanoparticles in rats after intranasal exposure. *Cell. Mol. Biol. (Noisy-le-grand)* 58, 196–203. <http://dx.doi.org/10.1170/T941>. [PubMed: 23273212]
- Patil NA, Gade WN, Deobagkar DD, 2016 Epigenetic modulation upon exposure of lung fibroblasts to TiO₂ and ZnO nanoparticles: alterations in DNA methylation. *Int. J. Nanomed* 11, 4509–4519.
- Pelcova D, Barosova H, Kukutschova J, Zdimal V, Navratil T, Fenclova Z, Vlckova S, Schwarz J, Zikova N, Kacer P, Komarc M, Belacek J, Zakharov S, 2015 Raman microspectroscopy of exhaled breath condensate and urine in workers exposed to fine and nano TiO₂ particles: a cross-sectional study. *J. Breath Res* 9, 036008 <http://dx.doi.org/10.1088/1752-7155/9/3/036008>. [PubMed: 26172946]
- Pelcova D, Zdimal V, Kacer P, Fenclova Z, Vlckova S, Syslova K, Navratil T, Schwarz J, Zikova N, Barosova H, Turci F, Komarc M, Pelcl T, Belacek J, Kukutschova J, Zakharov S, 2016a Oxidative stress markers are elevated in exhaled breath condensate of workers exposed to nanoparticles during iron oxide pigment production. *J. Breath Res* 10, 016004 <http://dx.doi.org/10.1088/1752-7155/10/1/016004>. [PubMed: 26828137]
- Pelcova D, Zdimal V, Fenclova Z, Vlckova S, Turci F, Corazzari I, Kacer P, Schwarz J, Zikova N, Makes O, Syslova K, Komarc M, Belacek J, Navratil T, Machajova M, Zakharov S, 2016b Markers of oxidative damage of nucleic acids and proteins among workers exposed to TiO₂ (nano) particles. *Occup. Environ. Med* 73, 110–118. <http://dx.doi.org/10.1136/oemed-2015-103161>. [PubMed: 26644454]
- Pelcova D, Zdimal V, Kacer P, Fenclova Z, Vlckova S, Komarc M, Navratil T, Schwarz J, Zikova N, Makes O, Syslova K, Belacek J, Zakharov S, 2016c Leukotrienes in exhaled breath condensate and fractional exhaled nitric oxide in workers exposed to TiO₂ nanoparticles. *J. Breath Res* 10, 036004 <http://dx.doi.org/10.1088/1752-7155/10/3/036004>. [PubMed: 27356965]
- Pelcova D, Zdimal V, Kacer P, Zikova N, Komarc M, Fenclova Z, Vlckova S, Schwarz J, Makeš O, Syslova K, Navratil T, Turci F, Corazzari I, Zakharov S, Bello D, 2017a Markers of lipid oxidative damage in the exhaled breath condensate of nano TiO₂ production workers. *Nanotoxicology* 11, 52–63. <http://dx.doi.org/10.1080/17435390.2016.1262921>. [PubMed: 27855548]
- Pelcova D, Zdimal V, Kacer P, Komarc M, Fenclova Z, Vlckova S, Zikova N, Schwarz J, Makes O, Navratil T, Zakharov S, Bello D, 2017b Markers of lipid oxidative damage among office workers exposed intermittently to air pollutants including nanoTiO₂ particles. *Rev. Environ. Health* 32, 193–200. <http://dx.doi.org/10.1515/reveh-2016-0030>. [PubMed: 27754970]
- Peschansky VJ, Wahlestedt C, 2014 Non-coding RNAs as direct and indirect modulators of epigenetic regulation. *Epigenetics* 9, 3–12. <http://dx.doi.org/10.4161/epi.27473>. [PubMed: 24739571]
- Peters A, Rückerl R, Cyrus J, 2011 Lessons from air pollution epidemiology for studies of engineered nanomaterials. *J. Occup. Environ. Med* 53, S8–S13. <http://dx.doi.org/10.1097/JOM.0b013e31821ad5c0>. [PubMed: 21654423]
- Phuyal S, Kasem M, Knittelfelder O, Sharma A, Fonseca DM, Vebraite V, Shaposhnikov S, Slupphaug G, Skaug V, Zienolddiny S, 2018 Characterization of the proteome and lipidome profiles of human lung cells after low dose and chronic exposure to multiwalled carbon nanotubes. *Nanotoxicology* 12, 138–152. <http://dx.doi.org/10.1080/17435390.2018.1425500>. [PubMed: 29350075]
- Pietrojusti A, Stockmann-Juvala H, Lucaroni F, Savolainen K, 2018 Nanomaterial exposure, toxicity, and impact on human health. *Wiley Interdiscip. Rev.: Nanomed. Nanobiotechnol* e1513 <http://dx.doi.org/10.1002/wnan.1513>.
- Pisani C, Gaillard JC, Nouvel V, Odorico M, Armengaud J, Prat O, 2015 High-throughput, quantitative assessment of the effects of low-dose silica nanoparticles on lung cells: grasping complex toxicity with a great depth of field. *BMC Genomics* 16, 315 <http://dx.doi.org/10.1186/s12864-015-1521-5>. [PubMed: 25895662]
- Porter DW, Wu N, Hubbs AF, Mercer RR, Funk K, Meng F, Li J, Wolfarth MG, Battelli L, Friend S, Andrew M, Hamilton R, Jr., Sriram K, Yang F, Castranova V, Holian A, 2013 Differential mouse

- pulmonary dose and time course responses to titanium dioxide nanospheres and nanobelts. *Toxicol. Sci* 131, 179–193. <http://dx.doi.org/10.1093/toxsci/kfs261>. [PubMed: 22956629]
- Prow TW, Grice JE, Lin LL, Faye R, Butler M, Becker W, Wurm EM, Yoong C, Robertson TA, Soyer HP, Roberts MS, 2011 Nanoparticles and microparticles for skin drug delivery. *Adv. Drug Deliv. Rev* 63, 470–491. <http://dx.doi.org/10.1016/j.addr.2011.01.012>. [PubMed: 21315122]
- Qian Y, Zhang J, Hu Q, Xu M, Chen Y, Hu G, Zhao M, Liu S, 2015 Silver nanoparticle-induced hemoglobin decrease involves alteration of histone 3 methylation status. *Biomaterials* 70, 12–22. <http://dx.doi.org/10.1016/j.biomaterials.2015.08.015>. [PubMed: 26295435]
- Reddy AR, Rao MV, Krishna DR, Himabindu V, Reddy YN, 2011 Evaluation of oxidative stress and anti-oxidant status in rat serum following exposure of carbon nanotubes. *Regul. Toxicol. Pharmacol* 59, 251–257. <http://dx.doi.org/10.1016/j.yrtph.2010.10.007>. [PubMed: 20955749]
- Renwick LC, Donaldson K, Clouter A, 2001 Impairment of alveolar macrophage phagocytosis by ultrafine particles. *Toxicol. Appl. Pharmacol* 172, 119–127. <http://dx.doi.org/10.1006/taap.2001.9128>. [PubMed: 11298498]
- Riebeling C, Wiemann M, Schnekenburger J, Kuhlbusch TA, Wohlleben W, Luch A, Haase A, 2016 A redox proteomics approach to investigate the mode of action of nanomaterials. *Toxicol. Appl. Pharmacol* 299, 24–29. <http://dx.doi.org/10.1016/j.taap.2016.01.019>. [PubMed: 26827820]
- Rittinghausen S, Hackbarth A, Creutzenberg O, Ernst H, Heinrich U, Leonhardt A, Schaudien D, 2014 The carcinogenic effect of various multi-walled carbon nanotubes (MWCNTs) after intraperitoneal injection in rats. Part. *Fibre Toxicol* 11, 59 <http://dx.doi.org/10.1186/s12989-014-0059-z>. [PubMed: 25410479]
- Ritz S, Schöttler S, Kotman N, Baier G, Musyanovych A, Kuharev J, Landfester K, Schild H, Jahn O, Tenzer S, Mailänder V, 2015 Protein corona of nanoparticles: distinct proteins regulate the cellular uptake. *Biomacromolecules* 16, 1311–1321. <http://dx.doi.org/10.1021/acs.biomac.5b00108>. [PubMed: 25794196]
- Robertson DG, Watkins PB, Reily MD, 2011 Metabolomics in toxicology: preclinical and clinical applications. *Toxicol. Sci* 120, S146–S170. <http://dx.doi.org/10.1093/toxsci/kfq358>. [PubMed: 21127352]
- Roco MC, 1997 Introduction, WTEC Workshop Report on R&D Status and Trends in Nanoparticles, Nanostructured Materials, and Nanodevices in the United States World Technology Evaluation Center (WTEC), Baltimore (Accessed 30 January 2018). <http://www.wtec.org/loyola/pdf/nanousws.pdf>.
- Sadiq R, Bhalli JA, Yan J, Woodruff RS, Pearce MG, Li Y, Mustafa T, Watanabe F, Pack LM, Biris AS, Khan QM, Chen T, 2012 Genotoxicity of TiO₂ anatase nanoparticles in B6C3F1 male mice evaluated using pig-a and flow cytometric micronucleus assays. *Mutat. Res* 745, 65–72. <http://dx.doi.org/10.1016/j.mrgentox.2012.02.002>. [PubMed: 22712079]
- Sargent LM, Porter DW, Staska LM, Hubbs AF, Lowry DT, Battelli L, Siegrist KJ, Kashon ML, Mercer RR, Bauer AK, Chen BT, Salisbury JL, Frazer D, McKinney W, Andrew M, Tsuruoka S, Endo M, Fluharty KL, Castranova V, Reynolds SH, 2014 Promotion of lung adenocarcinoma following inhalation exposure to multi-walled carbon nanotubes. Part. *Fibre Toxicol* 11 (3). <http://dx.doi.org/10.1186/1743-8977-11-3>.
- Sarkar A, Das J, Manna P, Sil PC, 2011 Nano-copper induces oxidative stress and apoptosis in kidney via both extrinsic and intrinsic pathways. *Toxicology* 290, 208–217. <http://dx.doi.org/10.1016/j.tox.2011.09.086>. [PubMed: 22000994]
- Sayes CM, Wahi R, Kurian PA, Liu Y, West JL, Ausman KD, Warheit DB, Colvin VL, 2006 Correlating nanoscale titania structure with toxicity: a cytotoxicity and inflammatory response study with human dermal fibroblasts and human lung epithelial cells. *Toxicol. Sci* 92, 174–185. <http://dx.doi.org/10.1093/toxsci/kfj197>. [PubMed: 16613837]
- Schmid O, Stoeger T, 2016 Surface area is the biologically most effective metric for acute nanoparticle toxicity of the lung. *J. Aerosol. Sci* 99, 133–143. <http://dx.doi.org/10.1016/j.jaerosci.2015.12.006>.
- Schulte PA, Hauser JE, 2012 The use of biomarkers in occupational health research, practice, and policy. *Toxicol. Lett* 213, 91–99. <http://dx.doi.org/10.1016/j.toxlet.2011.03.027>. [PubMed: 21477643]

- Schulte PA, Smith A, et al., 2011 Ethical issues in molecular epidemiologic research. In: Rothman (Ed.), *Molecular Epidemiology: Principles and Practices* International Agency for Research on Cancer, Lyon, pp. 9–21.
- Schulte PA, Talaska G, 1995 Validity criteria for the use of biological markers of exposure to chemical agents in environmental epidemiology. *Toxicology* 101, 73–88. [http://dx.doi.org/10.1016/0300-483X\(95\)03020-G](http://dx.doi.org/10.1016/0300-483X(95)03020-G). [PubMed: 7631325]
- Schulte PA, Iavicoli I, Rantanen JH, Dahmann D, Iavicoli S, Pipke R, Guseva Canu I, Boccuni F, Ricci M, Polci ML, Sabbioni E, Pietroiusti A, Mantovani E, 2016 Assessing the protection of the nanomaterial workforce. *Nanotoxicology* 10, 1013–1019. <http://dx.doi.org/10.3109/17435390.2015.1132347>. [PubMed: 26865347]
- Semmler-Behnke M, Kreyling WG, Lipka J, Fertsch S, Wenk A, Takenaka S, Schmid G, Brandau W, 2008 Biodistribution of 1.4- and 18-nm gold particles in rats. *Small* 4, 2108–2111. <http://dx.doi.org/10.1002/sml.200800922>. [PubMed: 19031432]
- Shrivastava R, Kushwaha P, Bhutia YC, Flora S, 2016 Oxidative stress following exposure to silver and gold nanoparticles in mice. *Toxicol. Ind. Health* 32, 1391–1404. <http://dx.doi.org/10.1177/0748233714562623>. [PubMed: 25548373]
- Shvedova AA, Kisin ER, Mercer R, Murray AR, Johnson VJ, Potapovich AI, Tyurina YY, Gorelik O, Arepalli S, Schwegler-Berry D, Hubbs AF, Antonini J, Evans DE, Ku BK, Ramsey D, Maynard A, Kagan VE, Castranova V, Baron P, 2005 Unusual inflammatory and fibrogenic pulmonary responses to single-walled carbon nanotubes in mice. *Am. J. Physiol. Lung Cell. Mol. Physiol* 289, L698–L708. <http://dx.doi.org/10.1152/ajplung.00084.2005>. [PubMed: 15951334]
- Shvedova AA, Yanamala N, Kisin ER, Khailullin TO, Birch ME, Fatkhutdinova LM, 2016 Integrated analysis of dysregulated ncRNA and mRNA expression profiles in humans exposed to carbon nanotubes. *PLoS One* 11, e0150628 <http://dx.doi.org/10.1371/journal.pone.0150628>. [PubMed: 26930275]
- Shyamasundar S, Ng CT, Yung LY, Dheen ST, Bay BH, 2015 Epigenetic mechanisms in nanomaterial-induced toxicity. *Epigenomics* 7, 395–411. <http://dx.doi.org/10.2217/epi.15.3>. [PubMed: 26077428]
- Sierra MI, Valdés A, Fernández AF, Torrecillas R, Fraga MF, 2016 The effect of exposure to nanoparticles and nanomaterials on the mammalian epigenome. *Int. J. Nanomed* 11, 6297–6306. <http://dx.doi.org/10.2147/IJN.S120104>.
- Singh SP, Kumari M, Kumari SI, Rahman MF, Mahboob M, Grover P, 2013a Toxicity assessment of manganese oxide micro and nanoparticles in Wistar rats after 28 days of repeated oral exposure. *J. Appl. Toxicol* 33, 1165–1179. <http://dx.doi.org/10.1002/jat.2887>. [PubMed: 23702825]
- Singh SP, Rahman MF, Murty US, Mahboob M, Grover P, 2013b Comparative study of genotoxicity and tissue distribution of nano and micron sized iron oxide in rats after acute oral treatment. *Toxicol. Appl. Pharmacol* 266, 56–66. <http://dx.doi.org/10.1016/j.taap.2012.10.016>. [PubMed: 23142030]
- Snyder-Talkington BN, Dymacek J, Porter DW, Wolfarth MG, Mercer RR, Pacurari M, Denvir J, Castranova V, Qian Y, Guo NL, 2013 System-based identification of toxicity pathways associated with multi-walled carbon nanotube-induced pathological responses. *Toxicol. Appl. Pharmacol* 272, 47–89. <http://dx.doi.org/10.1016/j.taap.2013.06.026>.
- Snyder-Talkington BN, Dong C, Zhao X, Dymacek J, Porter DW, Wolfarth MG, Castranova V, Qian Y, Guo NL, 2015 Multi-walled carbon nanotube-induced gene expression in vitro: concordance with in vivo studies. *Toxicology* 328, 66–74. <http://dx.doi.org/10.1016/j.tox.2014.12.012>. [PubMed: 25511174]
- Snyder-Talkington BN, Dong C, Sargent LM, Porter DW, Staska LM, Hubbs AF, Raese R, McKinney W, Chen BT, Battelli L, Lowry DT, Reynolds SH, Castranova V, Qian Y, Guo NL, 2016 mRNAs and miRNAs in whole blood associated with lung hyperplasia, fibrosis, and bronchiolo-alveolar adenoma and adenocarcinoma after multi-walled carbon nanotube inhalation exposure in mice. *J. Appl. Toxicol* 236, 161–174. <http://dx.doi.org/10.1002/jat.3157>.
- Soares T, Ribeiro D, Proença C, Chisté RC, Fernandes E, Freitas M, 2016 Size-dependent cytotoxicity of silver nanoparticles in human neutrophils assessed by multiple analytical approaches. *Life Sci* 145, 247–254. <http://dx.doi.org/10.1016/j.lfs.2015.12.046>. [PubMed: 26743952]

- Song MF, Li YS, Kasai H, Kawai K, 2012 Metal nanoparticle-induced micronuclei and oxidative DNA damage in mice. *J. Clin. Biochem. Nutr* 50, 211–216. <http://dx.doi.org/10.3164/jcbn.11-70>. [PubMed: 22573923]
- Soni D, Gandhi D, Tarale P, Bafana A, Pandey RA, Sivanesan S, 2017 Oxidative stress and genotoxicity of zinc oxide nanoparticles to pseudomonas species, human promyelocytic leukemic (HL-60), and blood cells. *Biol. Trace Elem. Res* 178, 218–227. <http://dx.doi.org/10.1007/s12011-016-0921-y>. [PubMed: 28058665]
- Srinivas A, Rao PJ, Selvam G, Murthy PB, Reddy PN, 2011 Acute inhalation toxicity of cerium oxide nanoparticles in rats. *Toxicol. Lett* 205, 105–115. <http://dx.doi.org/10.1016/j.toxlet.2011.05.1027>. [PubMed: 21624445]
- Srinivas A, Rao PJ, Selvam G, Goparaju A, Murthy PB, Reddy PN, 2012 Oxidative stress and inflammatory responses of rat following acute inhalation exposure to iron oxide nanoparticles. *Hum. Exp. Toxicol* 31, 1113–1131. <http://dx.doi.org/10.1177/0960327112446515>. [PubMed: 22699116]
- Stirling PA, 2018 *The Nanotechnology Revolution: A Global Bibliographic Perspective* CRC Press, Taylor Francis Group.
- Stone V, Miller MR, Clift MJD, Elder A, Mills NL, Møller P, Schins RPF, Vogel U, Kreyling WG, Alstrup, Jensen K, Kuhlbusch TAJ, Schwarze PE, Hoet P, Pietrousti A, De Vizcaya-Ruiz A, Baeza-Squiban A, Teixeira JP, Tran CL, Cassee FR, 2017 Nanomaterials versus ambient ultrafine particles: an opportunity to exchange toxicology knowledge. *Environ. Health Perspect* 125, 106002 <http://dx.doi.org/10.1289/EHP424>. [PubMed: 29017987]
- Sumner SC, Fennell TR, Snyder RW, Taylor GF, Lewin AH, 2010 Distribution of carbon-14 labeled C60 ([¹⁴C]C60) in the pregnant and in the lactating dam and the effect of C60 exposure on the biochemical profile of urine. *J. Appl. Toxicol* 30, 354–360. <http://dx.doi.org/10.1002/jat.1503>. [PubMed: 20063269]
- Sundarraj K, Raghunath A, Panneerselvam L, Perumal E, 2017 Iron oxide nanoparticles modulate heat shock proteins and organ specific markers expression in mice male accessory organs. *Toxicol. Appl. Pharmacol* 317, 12–24. <http://dx.doi.org/10.1016/j.taap.2017.01.002>. [PubMed: 28077287]
- Sung JH, Ji JH, Park JD, Yoon JU, Kim DS, Jeon KS, Song MY, Jeong J, Han BS, Han JH, Chung YH, Chang HK, Lee JH, Cho MH, Kelman BJ, Yu IJ, 2009 Subchronic inhalation toxicity of silver nanoparticles. *Toxicol. Sci* 108, 452–461. <http://dx.doi.org/10.1093/toxsci/kfn246>. [PubMed: 19033393]
- Tabish AM, Poels K, Byun HM, Luyts K, Baccarelli AA, Martens J, Kerkhofs S, Seys S, Hoet P, Godderis L, 2017 Changes in DNA methylation in mouse lungs after a single intra-tracheal administration of nanomaterials. *PLoS One* 12, e0169886 <http://dx.doi.org/10.1371/journal.pone.0169886>. [PubMed: 28081255]
- Takenaka S, Karg E, Roth C, Schulz H, Ziesenis A, Heinzmann U, Schramel P, Heyder J, 2001 Pulmonary and systemic distribution of inhaled ultrafine silver particles in rats. *Environ. Health Perspect* 109, 547–551. <http://dx.doi.org/10.2307/3454667>.
- Takenaka S, Karg E, Kreyling WG, Lentner B, Möller W, Behnke-Semmler M, Jennen L, Walch A, Michalke B, Schramel P, Heyder J, Schulz H, 2006 Distribution pattern of inhaled ultrafine gold particles in the rat lung. *Inhal. Toxicol* 18, 733–740. <http://dx.doi.org/10.1080/08958370600748281>. [PubMed: 16774862]
- Tang M, Zhang T, Xue Y, Wang S, Huang M, Yang Y, Lu M, Lei H, Kong L, Yuepu P, 2010 Dose dependent in vivo metabolic characteristics of titanium dioxide nanoparticles. *J. Nanosci. Nanotechnol* 10, 8575–8583. <http://dx.doi.org/10.1166/jnn.2010.2482>. [PubMed: 21121368]
- Tang M, Zhang T, Xue Y, Wang S, Huang M, Yang Y, Lu M, Lei H, Kong L, Wang Y, Pu Y, 2011 Metabonomic studies of biochemical changes in the serum of rats by intratracheally instilled TiO₂ nanoparticles. *J. Nanosci. Nanotechnol* 11, 3065–3074. <http://dx.doi.org/10.1166/jnn.2011.3604>. [PubMed: 21776672]
- Tarasova NK, Gallud A, Ytterberg AJ, Chernobrovkin A, Aranzaes JR, Astruc D, Antipov A, Fedutik Y, Fadeel B, Zubarev RA, 2017 Cytotoxic and pro-inflammatory effects of metal-based nanoparticles on THP-1 monocytes characterized by combined proteomics approaches. *J. Proteome Res* 16, 689–697. <http://dx.doi.org/10.1021/acs.jproteome.6b00747>. [PubMed: 27973853]

- Tavares A, Louro H, Antunes S, Quarre´ S, Simar S, De Temmerman PJ, Verleysen E, Mast J, Jensen KA, Norppa H, Nessler F, Silva MJ, 2014 Genotoxicity evaluation of nanosized titanium dioxide, synthetic amorphous silica and multi-walled carbon nanotubes in human lymphocytes. *Toxicol. In Vitro* 28, 60–69. <http://dx.doi.org/10.1016/j.tiv.2013.06.009>. [PubMed: 23811260]
- Thiruvengadam M, Rajakumar G, Chung IM, 2018 Nanotechnology: current uses and future applications in the food industry. *Biotechnology* 8, 74 <http://dx.doi.org/10.1007/s13205-018-1104-7>.
- Tiwari DK, Jin T, Behari J, 2011 Dose-dependent in vivo toxicity assessment of silver nanoparticle in Wistar rats. *Toxicol. Mech. Methods* 21, 13–24. <http://dx.doi.org/10.3109/15376516.2010.529184>. [PubMed: 21080782]
- Trouiller B, Reliene R, Westbrook A, Solaimani P, Schiestl RH, 2009 Titanium dioxide nanoparticles induce DNA damage and genetic instability in vivo in mice. *Cancer Res* 69, 8784–8789. <http://dx.doi.org/10.1158/0008-5472.CAN-09-2496>. [PubMed: 19887611]
- Uboldi C, Urbán P, Gilliland D, Bajak E, Valsami-Jones E, Ponti J, Rossi F, 2016 Role of the crystalline form of titanium dioxide nanoparticles: rutile, and not anatase, induces toxic effects in Balb/3T3 mouse fibroblasts. *Toxicol. In Vitro* 31, 137–145. <http://dx.doi.org/10.1016/j.tiv.2015.11.005>. [PubMed: 26571344]
- Vasantharaja D, Ramalingam V, Aadinaath, Reddy GG, 2015 Titanium dioxide nanoparticles induced alteration in haematological indices of adult male wistar rats. *J. Acad. Ind. Res* 3, 632–635.
- Villalta PW, Balbo S, 2017 The future of DNA adductomic analysis. *Int. J. Mol. Sci* 18, 1870 <http://dx.doi.org/10.3390/ijms18091870>.
- Vlaanderen J, Pronk A, Rothman N, Hildesheim A, Silverman D, Hosgood HD, Spaan S, Kuijpers E, Godderis L, Hoet P, Lan Q, Vermeulen R, 2017 A cross-sectional study of changes in markers of immunological effects and lung health due to exposure to multi-walled carbon nanotubes. *Nanotoxicology* 11, 395–404. <http://dx.doi.org/10.1080/17435390.2017.1308031>. [PubMed: 28301273]
- Wu WT, Liao HY, Chung YT, Li WF, Tsou TC, Li LA, Lin MH, Ho JJ, Wu TN, Liou SH, 2014 Effect of nanoparticles exposure on fractional exhaled nitric oxide (FENO) in workers exposed to nanomaterials. *Int. J. Mol. Sci* 15, 878–894. <http://dx.doi.org/10.3390/ijms15010878>. [PubMed: 24413755]
- Xia T, Hamilton RF, Bonner JC, Crandall ED, Elder A, Fazlollahi F, Girtsman TA, Kim K, Mitra S, Ntim SA, Orr G, Tagmount M, Taylor AJ, Telesca D, Tolic A, Vulpe CD, Walker AJ, Wang X, Witzmann FA, Wu N, Xie Y, Zink JI, Nel A, Holian A, 2013 Interlaboratory evaluation of in vitro cytotoxicity and inflammatory responses to engineered nanomaterials: the NIEHS nano GO consortium. *Environ. Health Perspect* 121, 683–690. <http://dx.doi.org/10.1289/ehp.1306561>. [PubMed: 23649538]
- Xie J, Dong W, Liu R, Wang Y, Li Y, 2018 Research on the hepatotoxicity mechanism of citrate-modified silver nanoparticles based on metabolomics and proteomics. *Nanotoxicology* 12, 18–31. <http://dx.doi.org/10.1080/17435390.2017.1415389>. [PubMed: 29251223]
- Yazdimaghani M, Moos PJ, Ghandehari H, 2017 Global gene expression analysis of macrophage response induced by nonporous and porous silica nanoparticles. *Nanomedicine* 14, 533–545. <http://dx.doi.org/10.1016/j.nano.2017.11.021>. [PubMed: 29203145]
- Yu LE, Yung LYL, Ong CN, Tan YL, Balasubramaniam KS, Hartono D, Shui GH, Wenk MR, Ong WY, 2007 Translocation and effects of gold nanoparticles after inhalation exposure in rats. *Nanotoxicology* 1, 235–242. <http://dx.doi.org/10.1080/17435390701763108>.
- Zhang R, Dai Y, Zhang X, Niu Y, Meng T, Li Y, Duan H, Bin P, Ye M, Jia X, Shen M, Yu S, Yang X, Gao W, Zheng Y, 2014 Reduced pulmonary function and increased pro-inflammatory cytokines in nanoscale carbon black-exposed workers. Part. *Fibre Toxicol* 11, 73 <http://dx.doi.org/10.1186/s12989-014-0073-1>. [PubMed: 25497989]
- Zhao Y, Sun H, Wang H, 2016 Long noncoding RNAs in DNA methylation: new players stepping into the old game. *Cell Biosci* 6, 45 <http://dx.doi.org/10.1186/s13578-016-0109-3>. [PubMed: 27408682]
- Zhao L, Zhu Y, Chen Z, Xu H, Zhou J, Tang S, Xu Z, Kong F, Li X, Zhang Y, Li X, Zhang J, Jia G, 2018 Cardiopulmonary effects induced by occupational exposure to titanium dioxide nanoparticles. *Nanotoxicology* 11, 1–16. <http://dx.doi.org/10.1080/17435390.2018.1425502>.

- Zhu MT, Feng WY, Wang Y, Wang B, Wang M, Ouyang H, Zhao YL, Chai ZF, 2009 Particokinetics and extrapulmonary translocation of intratracheally instilled ferric oxide nanoparticles in rats and the potential health risk assessment. *Toxicol. Sci* 107, 342–351. <http://dx.doi.org/10.1093/toxsci/kfn245>. [PubMed: 19023088]
- Ziemba B, Matuszko G, Appelhans D, Voit B, Bryszewska M, Klajnert B, 2012 Genotoxicity of poly(propylene imine) dendrimers. *Biopolymers* 97, 642–648. <http://dx.doi.org/10.1002/bip.22056>. [PubMed: 22605555]

Occupational Safety and Health Actions									
	Hazard Identification		Exposure Assessment		Risk Assessment		Risk Management		
Biomarker Types	In vivo	In vitro	Biomonitoring		Exposure-Response Modeling	Epidemiological Studies	Control Effectiveness	Medical Surveillance	Prevention through Design
			Direct	Indirect					
Pathological									
Genotoxic									
Transcriptomic									
Proteomic									
Metabolomic									
Epigenetic									

Fig. 1.
Framework for literature search strategy.




Fig. 2.
Conceptual timeline for growth of nanomaterial products.

Table 1

General Overview of Toxicology Findings for ENMs to date.

-
- Verified the correlation of particle surface area with biological effects (Duffin et al., 2001; Schmid and Stoeger, 2016)
 - Confirmed that particle size is generally an important factor in toxicity but other physico-chemical factors may play major roles (Driscoll, 1996; Oberdörster et al., 2005)
 - Confirmed that nanoparticles could reach the alveoli and could enter the interstitium and blood stream (Kreyling et al., 2009; Oberdörster et al., 1992)
 - Demonstrated that pulmonary exposure to carbon nanotubes (CNTs) causes alveolar interstitial fibrosis, which develops rapidly and is persistent. Fibrotic potency appears related to the physicochemical properties of the CNT (Shvedova et al., 2005)
 - Demonstrated that pulmonary exposure to nanoparticles can cause cardiovascular effects (alteration of heart rate and blood pressure, and microvascular dysfunction) (Li et al., 2007; Nurkiewicz et al., 2008)
 - Demonstrated that pulmonary exposure to some nanomaterials may be carcinogenic (Kuempel et al., 2017; Rittinghausen et al., 2014; Sargent et al., 2014)
 - Demonstrated that multi-walled carbon nanotubes (MWCNTs) (Mitsui-7) are a promoter of lung cancer (Kuempel et al., 2017)
 - Demonstrated that nanoparticles deposited in the lung can translocate to distal sites (Aragon et al., 2017; Erdely et al., 2009; Mercer et al., 2013) Demonstrated that insoluble nanoparticles do not appear to rapidly cross intact skin (Prow et al., 2011)
 - Demonstrated that nanoparticle penetration through intact skin is not likely but exposure to sensitizers and irritants still a concern (Gwinn and Vallyathan, 2006; Prow et al., 2011)
-