

## Longevity of Nonsmoking Men and Women

Many attempts have been made to determine factors which can explain the difference in male-female longevity. Possible explanatory factors include genetic differences, differences in health care use, behavioral differences in drinking and smoking habits, and behavioral differences in aggression and coping with stress.

The study by Miller and Gerstein (*1*) purports to show that all of the male-female differences in longevity can be attributed to higher rates of smoking among males. This contradicts a large body of evidence which indicates that differential smoking accounts for between one-third and one-half of the sex difference in longevity, for example, Waldron (*2*). Miller and Gerstein's work raises many issues which require more careful consideration.

Estimates of death rates which are needed for computing life expectancies are subject to sampling and nonsampling errors in estimating the populations at risk and the number of deaths. Miller and Gerstein used a 2 percent simple random sample to estimate the proportion of nonsmokers in their population. The possible effects of noncoverage error and of variance due to sampling were ignored. The sample frame for the population data covered only about 75 percent of the target population, but the estimates were applied to the entire population. Because some of the age-sex groups are fairly small the sampling errors may be relatively large. For example, the estimated percent of male nonsmokers was 36.8 percent for ages 85 years and over. Using the estimated population published in the tables in their article it can be calculated that an approximate 95 percent confidence limit is 15–60 percent nonsmokers for men 85 years and over. Similarly, when these highly variable population estimates are used to compute mortality rates, some ambiguous results are apparent, particularly at the older ages which are crucial for the estimation of life expectancy. Thus, given the sampling variation for the population estimates, the estimated life expectancy for men at age 85 could range from 5.4 years to 21.4 years.

There can be no question that sampling error in the population estimates has a substantial effect on the life table estimates. But another error, not due to sampling, may be even more important, that is, the determination of deaths of nonsmokers. Information on smoking status was obtained from relatives for only 63 percent of the deaths. There was no adjustment for nonresponse; decedents for which no information could be obtained (37 percent of all deaths) were essentially classified as "smokers." Thus, the number of deaths to nonsmokers was considerably underestimated, their death rates were similarly underestimated, and their life expectancies were therefore overestimated. Of crucial importance is the fact that Miller and Gerstein did not consider whether the biases in their estimates were the same for both males and females. For females, Miller and Gerstein classified about 65 percent of the population and 55 percent of the deaths as nonsmokers. For males, they classified about 30 percent of the population but only about 10 percent of the deaths as nonsmokers. Because of

differential nonresponse rates, it is likely that deaths for male nonsmokers may have been underestimated to a significantly greater extent than deaths for female nonsmokers.

Thus, we feel that the conclusion that all the male-female difference in longevity was due to differences in smoking is not warranted since the estimated differences may have resulted from the inherent variability present in their small data sets and from them ignoring the potential for differential biases in their methodological approach. Within these limitations their results are actually quite consistent with those previously presented in the literature that differential smoking accounts for between one-third and one-half of the sex difference in longevity.

Manning Feinleib, MD, DrPH  
Director  
National Center for Health Statistics  
Hyattsville, MD 20782

Joanne Luoto, MD, MPH  
Director  
Office of Smoking and Health  
Office of the Assistant Secretary for Health  
Public Health Service  
Rockville, MD 20857

## References .....

1. Miller, G. H. and Gerstein, D. R.: The life expectancy of nonsmoking men and women. *Public Health Rep* 98: 343–349, July-August 1983.
2. Waldron, I.: Why do women live longer than men? *Soc Sci Med* 10: 349–362 (1978).

## Others Would Get Similar Longevity Results If They Took Greater Care

Dr. Feinleib and Dr. Luoto present two methodological criticisms of the Miller-Gerstein report (*1*). They also state that our results both "contradict a large body of evidence" on smoking and the male-female longevity difference and are "actually quite consistent" with this same evidence. We will first discuss the methodological questions, then comment on the overall relation of our conclusions to the previous literature.

The first issue is sampling variability. Feinleib and Luoto illustrate the width of confidence intervals that result from data from small subsamples such as our smallest cohort, males age 85 and over. Vitality and other data on those aged 85 and older are notoriously unreliable in demographic sample analyses, and the observed results for this age category are often omitted or replaced by figures derived from large standard data bases. Rather than use this highly variable data in our life expectancy calculations, we obtained life expectancy figures for the 85+ cohort from the U.S. Bureau of Vital Statistics, as noted in the table in our report. The criticism based on this illustration is, therefore, inappropriate.