

PUBLIC HEALTH REPORTS

VOL. XXVII.

MAY 31, 1912.

No. 22.

STUDIES ON THE VIRUS OF TYPHUS.

By Joseph Goldberger, Passed Assistant Surgeon, and John F. Anderson, Director Hygienic Laboratory, Public Health and Marine-Hospital Service.

DURATION OF THE INFECTIVITY OF THE BLOOD.

The human inoculations by Moczutkowski (1900) and Otero (1908) strongly suggested, and those of Yersin and Vassal (1908) made it highly probable,¹ that the typhus-fever virus resides in the blood. Satisfactory proof of this, however, was not adduced until February, 1910, when Anderson and Goldberger showed that the fever induced in the monkey by an inoculation with human typhus blood, besides being transmissible by passage, conferred a definite immunity to a second inoculation.²

Since that time the experimental work of Anderson and Goldberger, of Ricketts and Wilder, of Gaviño and Girard, but more particularly that of Nicolle and Conseil, has developed a solid basis for the conclusion that the virus of typhus may be (or is) present in the blood, at least throughout the febrile period.

Nicolle and Conseil (August, 1910) were the first to try to determine whether the virus is present in the circulating blood before the onset of fever and after its defervescence, and from their experiments they concluded (January, 1911) that typhus blood is virulent, not only throughout the febrile period, but also before the onset of fever and at the beginning of convalescence (second day after defervescence).

In our recent work we repeated Nicolle and Conseil's experiments and have obtained results that cast grave doubt on the validity of Nicolle and Conseil's interpretations.

Before presenting our own work it will be desirable to examine Nicolle and Conseil's experiments somewhat in detail.

In order to determine whether typhus (monkey) blood is still infective after defervescence, Nicolle and Conseil inoculated a bonnet monkey (No. 31) with some blood from a chimpanzee (No. 3) obtained 36 to 48 hours after the return to normal of its temperature. After an incubation of 8 days the temperature of the bonnet monkey rose abruptly, remained up about 24 hours, then dropped, without

¹ There is some doubt whether Yersin and Vassal were really working with typhus.

² Nicolle's proof of the successful inoculation of his chimpanzee was the occurrence of a mild fever 24 days after inoculation and an eruption. Although the injection of 1 c. c. of the chimpanzee's blood was followed, in a bonnet monkey, by fever "with a most characteristic eruption," this was hardly convincing proof that the fever was typhus. It may be remarked in this connection that in his later work Nicolle has ceased to note the occurrence of an eruption.

subsequent hypothermia, but with slight emaciation. This is interpreted by Nicolle and Conseil as a typhus reaction, although when later subjected to an immunity test bonnet monkey 31 presented a well-marked reaction, showing that the fever of some 24 hours' duration, which Nicolle and Conseil interpret as a typhus reaction, had conferred no immunity.

With a view of testing the infectivity of the blood during the incubation of typhus, Nicolle and Conseil inoculated a bonnet monkey (No. 37) with blood from a chimpanzee (No. 4) obtained 3 days before the beginning of the fever. The result of this inoculation appeared to be absolutely negative, but when subjected to an immunity test 33 days later this animal (bonnet monkey 37) did not react. Whereupon Nicolle and Conseil conclude that the first inoculation, although followed by no symptoms, had conferred on their bonnet 37 a solid immunity.

It is evident that Nicolle and Conseil's conclusions that the blood is infective after defervescence and during the prefebrile period depend on the validity of their interpretations of the two experiments just cited.

The validity of their interpretation of the first of the above experiments is clearly involved in the question of what constitutes a typhus reaction. We defer a full discussion of this question to a later paper. At this time we wish simply to state that we are convinced that in the present state of our knowledge it is not permissible to interpret as a typhus reaction in the monkey a fever that confers no immunity.

In the second of the two experiments cited Nicolle and Conseil base their interpretation on the outcome of the immunity test; as this was negative they conclude that the animal, though it had presented no symptoms, was vaccinated by the injection with the blood obtained from chimpanzee No. 4.

It is obvious that in this case Nicolle and Conseil disregard the possibility of a natural immunity in the monkey. This possibility is not as remote as the early workers, including ourselves, believed. We have met with it several times in our recent work. Instances will be cited in other portions of this paper, although we shall defer a full discussion of this important matter to a later communication. For our present purpose it will be enough to say that in several instances in our recent experience monkeys have resisted one or two inoculations with virulent blood or blood serum and yet have reacted sharply and characteristically to the second or the third.

From the foregoing discussion it is clear that Nicolle and Conseil's conclusions are not justified by the results of their experiments when these are strictly interpreted.


Our own experiments follow:

Experiment No. 1.

At 12 m., February 29, about 32 hours before the beginning of a well-marked typhus reaction, we aspirated blood from the heart of rhesus No. 213 (chart 1) and after defibrination injected 8 c. c. of it diluted with 4 c. c. of saline solution into the peritoneal cavity of rhesus No. 216.

Result: During a period of observation of 30 days rhesus No. 216 failed to give any evidence of a reaction. At the end of this period he was subjected to an immunity test consisting of an intraperitoneal injection of 3 c. c. of defibrinated blood, obtained from monkeys Nos. 232 and 233, diluted with 2 c. c. of saline solution. To this test rhesus No. 216 reacted promptly and characteristically.

It appears, therefore, that monkey No. 216 was neither infected nor vaccinated by the inoculation with the blood of No. 213. This result would seem to justify the interpretation that the blood obtained from rhesus No. 213 about 32 hours before the beginning of fever was




Temperature curve of rhesus No. 213.

not virulent. This interpretation, though probably correct, is not permissible, in view of our experience that the normal monkey may resist one or two or more inoculations with virulent blood.

Experiment No. 2.

At 3.50 p. m., March 11, 1912—that is, at least 24 to 32 hours after the temperature of rhesus No. 213 reached normal—we again drew blood from this animal's heart and after defibrination, 9 c. c. of it diluted with 3 c. c. of saline solution was injected intraperitoneally into rhesus No. 227.

Result.—After an incubation period of about 24 days this monkey developed a mild fever of about 7 days' duration, which is sufficiently definite to justify us in considering it a typhus reaction (chart 2).



Temperature curve of rhesus No. 227. Mild febrile reaction following inoculation with blood obtained in beginning of convalescence.

The mild reaction obtained in rhesus No. 227 following inoculation with blood obtained at least 24 to 32 hours after the temperature of rhesus No. 213 reached normal justifies the interpretation that this blood was infective.

CONCLUSIONS.—From the foregoing discussion and the results of our experiments the following conclusions seem to us permissible:

1. The blood of the monkey infected with typhus may be virulent in the febrile stage, but no satisfactory evidence of that fact has as yet been adduced.

2. The blood of the monkey may still be virulent 24 to 32 hours after the return of the temperature to normal.

THE SEAT OF THE VIRUS OF TYPHUS.

The demonstration of the presence of the typhus virus in the circulating blood naturally leads one to inquire as to the element of the blood in which it resides. Nicolle and Jaeggy (April, 1910) were the first to consider this problem. In their hematologic study of typhus they describe a peculiar necrosis of the polynuclears which they consider significant as regards the seat of the unknown infectious agent, an agent which they therefore think is probably an intraleucocytic parasite.

Somewhat later Nicolle, Conor, and Conseil (October, 1910) point out certain apparent differences between the infectivity of typhus blood serum obtained after centrifugation of defibrinated blood and that obtained after clotting. These differences appear to them to support the intracellular hypothesis first suggested by Nicolle and Jaeggy.

In a later paper (January, 1911) Nicolle, Conor, and Conseil write that if the hypothesis of the intracellular character of the virus be accepted—

the discordant results of inoculation with serum obtained by centrifugation of defibrinated blood and that obtained after coagulation are easily explained. The former is always virulent because no matter how rapid and perfect the centrifugation may have been, it is bound to contain cellular débris and organisms liberated by the destruction of cells in the act of defibrination. The serum obtained after coagulation is, theoretically, deprived of all virulence for the monkey because it is freed of most of the liberated organisms and almost completely of the suspended cells by the formation and retraction of the clot.

They add, however, that—

the absence of virulence of the serum obtained after coagulation may not be absolute. When the serum does not separate well from the clot and when for that reason it contains a certain number of cells or when the amount inoculated is large the infection of the monkey may be realized.

In still a later paper (September, 1911) Nicolle, Conor, and Conseil report experiments especially designed to test their hypothesis. They consider that the results of these experiments support the theory of intraleucocytic localization.

On account of the great importance of the question involved it seems desirable to examine these experiments in detail. They are as follows:

A. COMPARATIVE VIRULENCE OF THE VARIOUS ELEMENTS OF THE BLOOD SEPARATED BY CENTRIFUGATION.

Experiment 1.—The blood used was drawn by venepuncture of patient 120 in the seventh day of a severe typhus, to which was added some sodium citrate, centrifuged,¹ etc. The manipulation of separation and lavage of the different elements of the blood took three-quarters of an hour.

¹ Just as we were about to go to press Nicolle, Conor, and Conseil's paper (Apr. 25, 1912) came to our hands, from which we cite the following additional details of this experiment: "After centrifugation (lasting 12 minutes) we at once removed several cubic centimeters of plasma from the upper portion of the tubes. The leucocyte layer was carefully removed and put in saline solution; several cubic centimeters of red corpuscles were then removed from the deepest portion of the tubes and placed in some of the same solution. These manipulations required 22 minutes. After that the white and the red globules suspended in saline solution were submitted to a second centrifugation of 10 minutes duration. This done, it required 6 minutes more to separate the cells from the liquid in which they were washed."

Seven bonnet monkeys were inoculated intraperitoneally as follows: A fresh control and two immunes each with 5 c. c. of the fresh blood; monkey A with the same dose of citrated blood; monkey B with the same dose of citrated plasma; monkey C with 2.5 c. c. of red corpuscles; monkey D with 1 c. mm. of the leucocyte layer.

The results were as follows: Fresh control (fresh blood), incubation 16 days, typhus of median severity of 12 days' duration (the temperature was over 40° during 6 days); immunes (fresh blood), negative. A (citrated blood), incubation 12 days, typhus of median severity of 9 days' duration (the temperature was over 40° during 9 days); B (citrated plasma), incubation 8 days, duration 9 days (6 days above 40°); C (red corpuscles), incubation 13 days, typhus abortive of 5 days (3 days at 40°, of which only one above); D (white corpuscles), incubation 6 days, typhus grave of 9 days (9 days above 40°).

Experiment No. 2.—Performed under the same conditions as No. 1 with the blood of patient 69 in the thirteenth day of a severe attack. Separation of the various elements of the blood took 30 minutes and was more perfect than in the first experiment. Three bonnet monkeys were inoculated intraperitoneally as follows: Monkey E with 5 c. c. of citrated plasma, F with the same dose of red corpuscles, G with about 1 c. mm. of leucocytes.

The results were as follows: E (citrated plasma), incubation 13 days, mild typhus of 7 days' duration (3 days above 40°); F (red corpuscles), negative; G (white corpuscles), incubation 7 days, severe typhus of 11 days (temperature above 40° during 9 days). Tested 67 days after the first inoculation by the intraperitoneal injection of 5 c. c. of blood from a monkey in the fifth day of an experimental attack; E and G have not reacted, F developed a severe typhus of 8 days after an incubation of 11 days.

B. NONVIRULENCE OF BLOOD SERUM FREED OF ITS CELLULAR ELEMENTS BY CENTRIFUGATION.

Serum obtained after clotting is, theoretically at least, free of cells. These remain entangled in the clot as it retracts; prolonged centrifugation frees it of such formed elements as it may contain. It is easy therefore to obtain a serum free of all white corpuscles. If our hypothesis as to the localization of the typhus virus in these cells is correct, such a liquid will be inoffensive. The experiment performed on man shows that it is so.

One of us received intravenously 1 c. c. of blood serum from a monkey in the third day of typhus seven hours after the blood was drawn; result negative. A previous experiment had shown us that the blood serum has no microbicidal action *in vitro* on the typhus virus.

C. NONVIRULENCE OF THE CEREBROSPINAL FLUID.

This liquid, destitute of all cellular elements, possesses no virulence.

"A bonnet monkey inoculated intraperitoneally with 15 c. c. of cerebrospinal fluid of patient 28 in the ninth day of a severe attack has presented no symptoms; tested 14 days later by an intraperitoneal inoculation of 5 c. c. of blood of a monkey in the fifth day, this monkey developed a fever of 7 days duration after an incubation period of 12 days."

From these experiments Nicolle, Conor, and Conseil draw the following conclusions:

1. Of the various elements of the blood separated by centrifugation the leucocytes are the most virulent; a minimal dose of these cells promptly determines in the monkey a grave infection.
2. The plasma, less active, seems to owe its virulence only to the leucocytes or leucocytic debris, of which it is difficult to free the plasma completely; the washed red cells are not virulent.
3. Centrifugated blood serum is inoffensive for man and a humor destitute of cells; the cerebrospinal fluid is equally inactive.


Before discussing the foregoing experiments and conclusions we desire to detail some pertinent experiments of our own.

Experiment No. 3.

On October 13, 1911, some defibrinated blood from rhesus No. 158, then in the first day of a typhus reaction, was centrifugated for about 30 minutes, after which the clear supernatant serum was pipetted off. With some of this serum two rhesus monkeys were inoculated. No. 141 received 3.5 c. c. intravenously; No. 142 received 0.5 c. c. subcutaneously and 4 c. c. intraperitoneally.

Result: Rhesus No. 141 developed a sharp reaction 7 days after the inoculation (chart 3); but during a period of observation of 29 days rhesus No. 142 gave no evidence of a reaction. At the end of this period No. 142 was subjected to an immunity test consisting of an intravenous injection of 2.5 c. c. of defibrinated blood of rhesus No. 163, to which he responded promptly and sharply. (Chart 4.)


CHART NO. 3



Temperature curve of rhesus No. 141, showing first three days of typhus reaction following inoculation with blood serum.

Although one of the two animals inoculated (with the larger dose) did not respond and was not vaccinated, it is clear that the blood serum of rhesus No. 158 was infective.

CHART NO. 4




Temperature curve of rhesus No. 142, showing reaction to immunity test.

Experiment No. 4.

On October 22, 1911, blood was drawn from rhesus No. 141, then in the third day of a marked attack of experimental typhus (chart 3), and allowed to clot in a tube. After 3 hours, in order to separate the serum more thoroughly from the clot, the tube was centrifugated for about 15 minutes. After drawing off the clear straw-tinted serum, a portion of it was used for the inoculation of two monkeys. No. 147 was given

4.5 c. c. and No. 148, 3 c. c., both intravenously. The time that elapsed between drawing the blood and inoculating was about 3 hours and 20 minutes.


Result: During a period of observation of 29 days rhesus No. 147 gave no evidence of a reaction. Rhesus No. 148, however, after an incubation period of about nine days, developed a severe typhus, as shown by marked fever (chart 5) and successful passage to another monkey. It should be noted that, although rhesus No. 147 and rhesus No. 148 were of substantially the same size and vigor, yet rhesus No. 147—the one



Temperature curve of rhesus No. 148, showing severe febrile reaction, following inoculation with blood serum.

that received the larger dose (larger by 50 per cent)—failed to react, and when subjected to an immunity test, consisting of a subcutaneous injection of about 0.5 c. c., and of an intravenous injection of 2 c. c., of defibrinated blood of rhesus No. 166, then in the third day of an experimental typhus, rhesus No. 147 reacted promptly and sharply (chart 6).

Although one of the two animals inoculated (with much the larger dose) did not become infected nor immunized, it is evident that the blood serum of rhesus No. 141, obtained after clotting and centrifugation, was infective.



Temperature curve of rhesus No. 147, showing reaction to immunity test.

Experiment No. 5.


On March 6, 1912, blood was aspirated from rhesus No. 210 and No. 213. The former was in the seventh and the latter in the fifth day of an experimental typhus. The blood was allowed to clot in tubes and, as in Experiment No. 4, after standing three hours it was centrifuged in order better to separate the serum from the clot. The serum was then pipetted off and mixed. Of this serum, 6 c. c. were intraperitoneally injected into rhesus No. 221.

Result: During a period of observation of 24 days following this inoculation monkey No. 221 manifested no recognizable symptoms of a reaction. He was therefore re-inoculated March 30 in order to test his immunity; for this purpose he was given an intraperitoneal injection of 3 c. c. of virulent defibrinated blood (rhesus Nos. 232 and 233). Twenty-one days later, April 20, not having in the meantime given any indications of a reaction, he was again given an intraperitoneal injection of 3 c. c. of virulent defibrinated blood (rhesus No. 225). So far this monkey has given no recognizable evidence of a reaction following this inoculation.

The results of the repeated immunity tests of rhesus No. 221 would seem to indicate that this animal was vaccinated by the first inoculation with 6 c. c. of serum; we think it more probable, however, that this is an instance of natural immunity in this monkey.

Experiment No. 6.

On October 27, 1911, some blood drawn from rhesus No. 141 (chart 3) was defibrinated and 3 c. c. of it tubed and centrifuged. After a centrifugation of about 15 minutes the serum was pipetted off and replaced by 5 c. c. of saline solution in which the corpuscles were thoroughly stirred up. The tube was then again put in the centrifuge for 15 minutes, after which it was taken out and the supernatant liquid pipetted



Temperature curve of rhesus No. 133, showing reaction following inoculation with washed corpuscles.

off and replaced by 5 c. c. of fresh saline solution. This maneuver was twice repeated. In other words, the corpuscles were washed three times in an excess of saline solution. After the third washing the corpuscles, in fresh saline solution, representing originally 3 c. c. of defibrinated blood, were injected intravenously into rhesus No. 133.

Result: After an incubation period of about 10 days this monkey developed a marked typhus reaction (chart No. 7).

Washing the blood corpuscles three times in an excess of saline solution had, therefore, not deprived them of the power to infect.

Experiment No. 7.

On February 12, 1912, we aspirated blood from three monkeys—Nos. 196, 198, and 322—that were in the fourth, fifth, and sixth days, respectively, of experimental typhus. The blood was defibrinated and mixed. The corpuscles of 12 c. c. of this mixed defibrinated blood were washed three times, as in experiment No. 6. After the third washing the supernatant liquid was drawn off and discarded; then the surface layer (“cream” or leucocyte layer) of corpuscles was pipetted off and injected intraperitoneally into rhesus No. 206. Of the corpuscles remaining, 3 c. c. were intraperitoneally injected into rhesus No. 207.


Result: Both monkeys developed well-marked reactions after an incubation period of seven days in No. 206 and eight days in No. 207. Although the reaction in neither was severe, that in No. 207 was perhaps the somewhat better marked (charts 8 and 9).

Both the "cream" or leucocyte layer and the red¹ corpuscles of washed blood were, therefore, infective.

DISCUSSION.—Nicolle, Conor, and Conseils's interpretations are evidently based on the assumption that the severity of the reaction is a direct index of the virulence of the infecting dose.


If this assumption were permissible it would be difficult to say from the character of the reaction noted by us in rhesus No. 206 and No.

CHART No. 8



Temperature curve of rhesus No. 206, showing reaction following an inoculation with leucocyte layer of washed corpuscles.

CHART No. 9



Temperature curve of rhesus No. 207, showing reaction following an inoculation with washed "red" corpuscles.

207 (charts 8 and 9) which was the more virulent, the "cream" layer given to No. 206 or the "reds" given to No. 207. As a matter of fact,

¹ Although centrifugation separates the white and the red corpuscles, this separation is never complete. A considerable number of leucocytes are always carried down by the reds from which they can not be completely removed, even by repeated washings and repeated removal of the "cream" layer. So that when we speak of the "reds" it must always be understood that a not inconsiderable number of whites are included. This is well illustrated by the following experiment, kindly made for us by our colleague, Dr. Leake: Some typhus monkey blood was citrated and then centrifuged for one hour. After this the serum and the cream layers were pipetted off and the remaining "reds" washed three times. After each washing the "cream" layer that formed was discarded. Leucocyte counts were made as follows: (1) Citrated blood before centrifugation, 7,000 per c. mm.; (2) plasma after centrifugation of 1 hour, less than 10 per c. mm.; (3) "cream" layer after centrifugation of 1 hour, 79,000 per c. mm.; (4) "reds" after centrifugation 1 hour and after removing the "cream," 14,000 per c. mm.; (5) "reds" after washing three times and each time discarding the "cream" layer, 9,000 per c. mm. After an hour's centrifugation, therefore, the plasma of citrated blood was not freed of leucocytes and the number of leucocytes among the "reds" was relatively increased.

there does not appear to be any necessary relation between dosage and severity of reaction.

Ricketts and Wilder (Feb. 5, 1910) and Anderson and Goldberger (Feb. 18, 1910), almost simultaneously record observations to this effect over two years ago. We have an extreme and a very pertinent illustration of this in the result above detailed of our experiment No. 4 in which, of two animals of equal size and vigor inoculated with typhus blood serum, the monkey that received the smaller dose presented a severe reaction while the monkey that received the larger dose (by 50 per cent) failed altogether to react although a month later he reacted promptly and sharply to an immunity test with a relatively small amount of defibrinated blood. But waiving this objection and accepting their interpretation that "of the various elements of the blood separated by centrifugation the leucocytes are the most virulent," "the plasma less active," "and the washed corpuscles not virulent," ("not virulent" in the sense, it is presumed of "low virulence," for in one of their experiments under "A" the "reds" which necessarily entangled some leucocytes, were infective, as they were also in our experiment No. 7), it does not seem to us necessarily to follow that because the degree of virulence of the various elements of the blood thus separated appears to run parallel¹ with their (incorrectly) assumed richness in leucocytes that the virus is localized in the leucocytes or has any relation to the leucocytes other than that of possessing, possibly, a specific gravity approximating, or identical with, that of these cells.

In other words, having due regard for the mechanics of centrifugation, one might reasonably expect that the distribution in the different layers of centrifugated blood of an extracellular parasite of approximately the same specific gravity as that of the leucocytes would be similar to, or identical with, that of the latter. This is well illustrated by the following experiment. Some blood was drawn into sodium citrate solution from the heart of a guinea pig infected with anthrax. A measured amount of this was plated in agar. The remaining citrated blood was then centrifugated for one hour, after which measured amounts of the different layers, plasma, "cream," and "reds," uniform with that of the whole blood, were plated. After incubating 18 hours at 37° the plates were examined. The dilutions not having been sufficient to permit counting the colonies on some of the plates, estimates of their number were made and then certain relative values given to each, with the following result: "Whole blood," 100; "plasma," 1; "cream," 10,000; "reds," 10. Here we have an enormous excess of organisms in the leucocyte layer and a great reduction in the plasma and in the "reds." The reason for the high virulence of the leucocytic layer, the lessened activity of the plasma and the "nonvirulence"² of the washed red corpuscles is therefore obvious.

¹ As a matter of fact, there is no evidence to show that it really does, for the leucocyte content of the various "elements" of the blood separated by centrifugation, is in the following order: "Cream," "reds," plasma (see note, Experiment No. 7).

² Nicolle, Cónor, and Conseil say: "Des divers éléments du sang, séparés par centrifugation, les globules blancs sont en effet les plus virulents; une dose minime de ces cellules détermine chez le singe une infection rapide et grave; le plasma, moins actif, semble ne devoir sa virulence qu'aux leucocytes ou débris leucocytaires dont il est malaisé de le débarrasser complètement; les globules rouges lavés n'ont pas de virulence."

We presume that by "n'ont pas de virulence"—have no virulence—they mean "low virulence," for in one of their experiments under "A" the "reds" were infective. A literal interpretation of this phrase would make their argument in favor of an intraleucocytic localization absurd, for the "reds" necessarily entangle a not inconsiderable number of leucocytes as they are precipitated by the centrifuge.

If now we bear in mind that the severity of the reaction does not necessarily correspond to the degree of virulence (dosage), even though it may do so under certain circumstances, then we believe we have accurately reproduced in this experiment with anthrax blood the results obtained by Nicolle, Conor, and Conseil after centrifugation of citrated typhus blood.

In view of the foregoing it is easy to conceive that typhus blood serum may by sufficient centrifugation be made noninfective even to so (supposedly) sensitive a subject as man. On account of the variable susceptibility of the monkey, very little significance can be attached to the negative result of the single inoculation with cerebrospinal fluid. It does not prove that this fluid is not infective, although it is in harmony with the result of a human experiment made by Otero in 1908. But even if it did, to have the significance that Nicolle, Conor, and Conseil seem to attach to it, it would be necessary to show, first, that the cerebrospinal fluid of typhus is quite free of leucocytes, and second, that in infections due to an extracellular parasite this fluid is invariably virulent.

CONCLUSIONS.—From the foregoing discussion we believe it permissible to conclude: (1) That the evidence adduced by Nicolle, Conor, and Conseil does not especially favor their hypothesis of an intraleucocytic localization of the virus of typhus. On the contrary, the infectivity of centrifugated blood serum, obtained after clotting, with its low leucocyte content would be in favor of a parasite free in the circulating plasma of the blood; (2) that the blood serum of virulent typhus blood is constantly infective, whether obtained from defibrinated blood or after clotting, instances of its apparent avirulence being explicable by a natural resistance of the monkey; (3) that it may perhaps be possible to deprive typhus blood serum (obtained after clotting) of its virulence by prolonged centrifugation, but that this does not necessarily indicate an intraleucocytic localization of the virus; and (4) that repeated washings of the blood corpuscles do not deprive them of their infectivity, a fact explicable by the physical phenomena involved in centrifugation.

FILTERABILITY.

VIRUS IN THE BLOOD.

The question of the filterability of the virus as it exists in typhus blood has been studied by several groups of workers. Ricketts and Wilder were the first to come forward with an answer to this question. On February 5, 1910, they reported a filtration experiment. In this a monkey inoculated with filtered (Berkefeld candle) blood serum (from defibrinated blood) failed to react. Almost simultaneously Goldberger and Anderson (Feb., 18, 1910) reported a similar experiment with a like result. In a paper published some weeks later (Apr. 23, 1910) Ricketts and Wilder report apparently the same result from a second experiment.

In October, 1910, Nicolle, Conor, and Conseil reported two series of experiments with filtered serum (from clotted blood) on monkeys and a third on man. In the first, one of the two monkeys that had received the filtered serum gave no indications of a reaction; the other (bonnet 47) presented an elevation of temperature of 0.5 degree

between the sixteenth and eighteenth day,¹ so that the result was doubtful. When later subjected to an immunity test bonnet 47 proved refractory, from which they conclude that the subcutaneous inoculation of filtered typhus serum produced in this monkey a feeble thermic reaction (doubtful), which, however, conferred an absolute immunity to a subsequent immunity test.

In the second series of their experiments neither the monkey that received the unfiltered serum nor the three monkeys that received the big doses of the filtered serum presented the slightest indications of a reaction.

In a third experiment one of the authors submitted himself to a subcutaneous injection of 0.25 c. c. of filtered typhus serum without developing any symptoms.

After reviewing the results of filtration recorded by Ricketts and Wilder and by Anderson and Goldberger, Nicolle, Conor, and Conseil (January, 1911) conclude "that the serum obtained by centrifugation of defibrinated blood is always inactive after filtration; that the serum obtained after coagulation is generally also inactive after filtration, but not constantly so; and that the only hypothesis that permits of an explanation of these facts is that under ordinary conditions the unknown microbe of typhus is present in the filtrate in numbers too small to cause infection or immunization of the inoculated animal. This microbe is therefore filterable and probably intraleucocytic." It should here be noted that this was written before the result of the immunity test of the monkey used in the second of Ricketts and Wilder's filtration experiments was published. This result appeared in a paper published by Wilder in July, 1911, and showed that as in one of Nicolle, Conor, and Conseil's experiments the monkey that had received the injection of filtered (defibrinated blood) serum was resistant to the immunity test.

In September, 1911, Nicolle, Conor, and Conseil reported still another filtration experiment. Believing that the virus is intraleucocytic they thought "that it would perhaps be possible to obtain, by the artificial disintegration of a large number of these cells, enough free organisms so that their filtered emulsion would infect the monkey." This, however, did not prove to be the case and they conclude that "in this instance, again, the number of organisms that passed the filter was undoubtedly insufficient," so that their "previous positive filtration experiment remains unique." They are evidently still ignorant of Ricketts and Wilder's "positive" (?) result.

In November, 1911, Gaviño and Girard published some studies in typhus, in which they report an absolutely negative result following inoculation of a monkey with filtered (through a Berkefeld candle) peritoneal exudate rich in erythrocytes and leucocytes, particularly polynuclears. This was obtained from the peritoneal cavity of a monkey at the height of an experimental typhus three hours after injecting some peptonized broth.

Summarizing the foregoing, we have eight attempts, so far recorded, to pass the virus of typhus through a Berkefeld filter. Of these, six were negative; in one of the other two (Wilder, 1911) the monkey that had been inoculated with the filtrate, without giving any evidence of a reaction, was later found to be resistant to an immunity

¹ In their summary January, 1911, this is stated as "0.5° to 0.8° between the fifteenth and twenty-first day."

test with virulent blood; in the other (Nicolle, Conon, and Conseil, January, 1911), one of a pair of monkeys inoculated with some of the filtrate is described as having presented a doubtful reaction, and later was found resistant to an inoculation with virulent blood.

Before discussing these results and their significance, we desire to record some new experiments of our own.

Experiment No. 8.

On October 13, 1911, some blood was drawn from rhesus No. 158, then in the first day of a severe experimental typhus. After defibrination this blood was centrifugated for about half an hour, after which the supernatant serum was pipetted off. A portion of the serum was diluted with three volumes of saline solution and then passed through a Berkefeld candle, after which the following inoculations were made: Of the unfiltered serum rhesus No. 141 was given 3.5 c. c. intravenously and rhesus No. 142, 4 c. c. intraperitoneally and 0.5 c. c. subcutaneously. Of the diluted filtered serum rhesus No. 143 was given 8.5 c. c. (representing 2.12 c. c. of the original) intravenously and 6 c. c. (representing 1.5 c. c. of the original) intraperitoneally; rhesus No. 144 was given 9.5 c. c. (representing 2.4 c. c. of the original) intravenously and 5 c. c. (representing 1.25 c. c. of the original) intraperitoneally.

Result: Rhesus No. 141 (unfiltered serum), after an incubation period of 7 days, developed a sharp reaction (chart 3), on the third day of which he was sacrificed in the act of aspirating blood for passage. Rhesus No. 142 (unfiltered serum), 143 (filtrate), 144 (filtrate), during a period of observation of 29 days gave no evidence of a reaction. At the end of this period these three monkeys were subjected to an immunity test, each receiving intravenously 2.5 c. c. of defibrinated blood of rhesus No. 163, then in the second day of a marked experimental typhus. All three animals responded promptly and sharply to this test (chart 4).

It appears, therefore, that of the two monkeys (Nos. 141 and 142) that were inoculated with virulent typhus serum obtained after a half hour's centrifugation of virulent defibrinated blood (virulent for two monkeys in a dose of 3.5 c. c. each, intravenously) one (the one receiving the larger dose) failed to react without being refractory to the immunity test and that neither of the two monkeys inoculated with the filtered serum was either infected thereby or had any resistance conferred upon it.

Experiment No. 9.


On October 22, 1911, blood was drawn from the carotid of rhesus No. 141, at this time in the third day of a marked attack of experimental typhus, and allowed to clot in a tube. At the end of 3 hours the tube was centrifugated for about 15 minutes in order to separate the serum more thoroughly from the clot. After drawing off the clear straw-tinted serum a portion was diluted with 3 volumes of saline solution and passed through a Berkefeld filter, after which the following inoculations were made: Of the unfiltered serum, rhesus No. 147 was given 4.5 c. c. and No. 148, 3 c. c., both intravenously. Of the dilute filtered serum rhesus Nos. 115*a* and 115*b* were each given 10 c. c. (representing 2.5 c. c. of undiluted serum) intravenously and 2 c. c. (representing 0.5 c. c. of undiluted serum) intraperitoneally.

Result: As already stated in a previous connection, rhesus No. 148 (3 c. c. serum) developed a severe typhus reaction after an incubation period of about 9 days (chart 5). Rhesus No. 147 (4.5 c. c. serum), No. 115*a* (3 c. c. filtered serum), No. 115*b* (3 c. c. filtered serum) during a period of observation of 29 days following the inoculation, presented no recognizable evidence of a reaction. At the end of this period of observation monkeys Nos. 147, 115*a* and 115*b* were subjected to an immunity test consisting of an inoculation of defibrinated blood of rhesus No. 166 then in the third day of a marked experimental typhus, No. 147 receiving 2 c. c. intravenously and 0.5 c. c. subcutaneously, No. 115*a* 2.5 c. c. subcutaneously, and No. 115*b* 2.5 c. c. intravenously. Rhesus Nos. 147 and 115*b* reacted promptly and sharply to this test (charts 6 and 10). No. 115*a*, however, gave no evidence of a reaction during a period of 33 days following this test. Sixty-two days after the inoculation with the filtered serum rhesus No. 115*a* was given a second immunity test consisting of an inoculation, part intravenous and part subcutaneous, of 2.5 c. c. of virulent defibrinated blood of

Experiment No. 10.

On January 20, 1912, some blood was drawn into a tube from the carotid of rhesus No. 309, which at this time was in the second day of a sharp attack of the experimental disease. This blood was allowed to clot at room temperature. At the end of two and one half hours it was centrifugated, after which the perfectly clear serum was pipetted off and a portion diluted with 2 volumes of saline solution. This diluted serum was then passed through a Berkefeld candle, after which the following inoculations were

CHART No. 12




Temperature curve of rhesus No. 195, showing what was probably a reaction (complicated by tuberculosis), following an inoculation with blood serum.

made: Rhesus No. 194 was given 18 c. c. of the filtrate (representing 6 c. c. of serum) intraperitoneally; rhesus No. 195 was given 6 c. c. of unfiltered serum, likewise intraperitoneally.

Result: Rhesus No. 195 developed what was probably a typhus reaction after an incubation period of 8 days (chart 12). Some doubt, however, is cast upon this interpretation by the fact that this monkey's normal temperature was somewhat high, due, probably, to a coexisting tubercular infection that later was found to be present.

CHART No. 13




Temperature curve of rhesus No. 194, showing reaction following second immunity test.

Rhesus No. 194 presented no indications of a reaction during a period of observation of 34 days. At the end of this time he was subjected to an immunity test consisting of an intraperitoneal injection of 6 c. c. of virulent defibrinated blood without eliciting any evidence of a reaction. This would suggest that this monkey had been made refractory by the inoculation with the filtrate, just as appeared to have happened in one instance in Ricketts and Wilder's experience (Wilder, 1911) and in one instance in

the experience of Nicolle, Conor, and Conseil (January, 1911). On subjecting this monkey to another immunity test (3 c. c. defibrinated blood intraperitoneally), however, 70 days after the inoculation with the filtered serum rhesus No. 194 responded promptly and sharply as may be seen from chart 13.


CHART NO. 14



Temperature curve of rhesus No. 223, showing reaction following immunity test.

It would appear therefore that the inoculation of rhesus No. 194 with 6 c. c. of filtered serum obtained after centrifugation of clotted typhus blood had neither infected nor vaccinated this animal.

CHART NO. 15



Temperature curve of rhesus No. 224, showing reaction following immunity test.

Experiment No. 11.

On March 6, 1912, blood was aspirated from rhesus No. 210 and No. 213. The former was in the seventh and the latter in the fifth day of an experimental typhus. The blood was allowed to clot. After 3 hours it was centrifuged in order better to separate the serum from the clot. The serum was then pipetted off and the two specimens were mixed. A portion of this serum was diluted with 3 volumes of saline

solution and then passed through a Berkefeld filter, after which the following inoculations were made:

Rhesus No. 221 received 6 c. c. of unfiltered serum, rhesus Nos. 223 and 224 each received 6 c. c. of filtered serum intraperitoneally.

Result: None of these animals presented any evidence of a reaction following this inoculation. Twenty-four days later the three were subjected to an immunity test consisting of an intraperitoneal injection of 3 c. c. of virulent defibrinated blood of rhesus Nos. 232 and 233, diluted with 2 c. c. of saline solution. Following this rhesus Nos. 223 and 224 responded promptly and sharply (charts 14 and 15), but No. 221 gave no evidence of a reaction. Rhesus No. 221 has been subjected to two more immunity tests since, but so far has presented no evidence of a reaction. (Cf. Exp. 5.)

It would appear, therefore, that the inoculation of two monkeys with filtered serum obtained after centrifugation of clotted typhus blood had produced neither infection nor immunization in either.

Summarizing our attempts to infect the rhesus monkey with filtered typhus blood serum, we find that in no instance was infection produced, and that when subjected to an immunity test 2 (rhesus 115a, experiment 2, and rhesus 194, experiment 3) of the 7 monkeys employed at first appeared to be resistant (apparently vaccinated), but later when the test was repeated both responded.

DISCUSSION.—Having presented first a summary of the results of the filtration experiments recorded in the literature and then detailed our own attempts to pass the virus of typhus through the Berkefeld filter, we may proceed with a consideration of their interpretation and significance.

The significance to be attached to the results of the filtration experiments, as a whole, evidently hinges on the interpretation to be given to the observation that some of the monkeys that had been inoculated with filtered typhus blood serum were subsequently resistant to an immunity test, although the inoculation with the filtrate had not been followed by any definite evidence of a febrile reaction. Nicolle, Conor, and Conseil (October, 1910, p. 3) interpret the resistance to the immunity test of one of their monkeys as indicating an immunization caused by "the microbe of typhus, which is very small and passes the Berkefeld filter"; whereas, Wilder suggests that a similar result in Ricketts's and Wilder's experience "may be explained by one or more of several different hypotheses. The animal may have been naturally immune to typhus. If such is the case, however, he is the first normal monkey with which we have had to deal that has shown such an absolute immunity when inoculated with over 1 c. c. of virulent blood. A second possibility is that the animal was immunized by the filtered serum." "Such immunization could have been accomplished either by microorganisms sufficiently small to pass the filter, by fragments of organisms, or by toxins." Although Wilder is evidently disinclined to attach any importance to the idea that the normal monkey may be naturally immune to typhus, and Nicolle, Conor, and Conseil do not even consider it, the results of our inoculations with virulent typhus blood serum (unfiltered), as well as the final results of the repeated immunity tests, clearly show that something in the nature of a natural immunity or transient unresponsiveness to typhus is possessed by some normal monkeys and we believe this to be the true explanation of the instance noted by Nicolle, Conor, and Conseil and that by Ricketts and Wilder. We believe that, had they repeated the immunity tests in these monkeys as we did in ours, their results would probably have shown

as did ours that these animals were only apparently (? transiently) resistant. These monkeys were therefore not immunized by the filtered typhus serum.

CONCLUSIONS.—It is permissible to conclude from the foregoing that there is no evidence to show that the virus in the blood of typhus is able to pass the Berkefeld filter; and, incidentally, that virulent typhus blood contains no toxin, or contains it in quantities too small to cause an appreciable increase in the normal resistance of the monkey when injected, even repeatedly, in the ordinary doses.

THE VIRUS IN THE LOUSE.

The fact that the organism of typhus, as it occurs in the blood, does not appear capable of passing the Berkefeld filter does not of necessity mean that it may not be capable of passing in the form in which it exists in the body of the louse. It was decided, therefore, to put this idea to the test of experiment.

Experiment No. 12.

On December 3, 1911, 82 living body lice of group No. 7—M and 83 of group No. 8—M were crushed and rubbed up in 16.5 c. c. of saline solution. Of this suspension 11 c. c. were diluted with an equal volume of saline solution and filtered through a Berkefeld filter. With 15 c. c. of the clear filtrate, representing 75 body lice, rhesus No. 311 was inoculated by an injection of about 13 c. c. intraperitoneally and about 2 c. c. subcutaneously. Control inoculations with the original (undiluted) suspension (reported in a previous communication by Goldberger and Anderson, Mar. 1, 1912) were made in two rhesus monkeys; one, No. 308, received 3.5 c. c. (representing 35 lice), and No. 309, 1.5 c. c. (representing 15 lice) subcutaneously.

Body lice of group No. 7—M had been allowed to feed daily during the 6 days immediately prior to the date of the experiment on various cases of typhus fever in the Hospital General, Mexico City. They were last applied to case No. 16—M, in the eleventh day of illness, at 10.50 to 11.20 a. m. December 2. Body lice of group No. 8—M were insects that had been allowed to feed daily during the 5 days immediately preceding the date of the experiment on various cases of typhus fever. They were last applied to case No. 16—M, in the eleventh day of illness, at 10.20 to 10.50 a. m., December 2. Throughout the period during which they were allowed to feed on patients with typhus the lice of groups Nos. 7—M and 8—M were kept at room temperature (about 14 to 24° C.).

Result: As has already been reported, the controls—rhesus Nos. 308 and 309—presented no evidence of a typhus reaction. No. 311 developed a markedly remittent temperature following the inoculation with the filtrate, and later was found to be infected with tuberculosis. This experiment has, therefore, no significance, so far as filtration is concerned.

Experiment No. 13.

The preceding experiment was repeated on December 10. At 5.30 p. m. 478 lice of group No. 9—M were crushed and rubbed up in 48 c. c. of saline solution. Of this suspension about 7.5 c. c. (representing about 73 lice) were subsequently injected in rhesus No. 320. Another portion of the suspension, after dilution with 2 volumes of saline solution, was filtered through a Berkefeld filter, and then 39 c. c. of the filtrate, representing not less than 130 lice, were injected intraperitoneally in rhesus No. 319.


The lice composing group No. 9—M were body lice that had been applied daily and been allowed to feed on various cases of typhus at the Hospital General, Mexico City, during the 6 days immediately prior to the date of the experiment. Throughout this period they were kept at room temperature (14° to 24° C.). They were last applied on December 9 between 10.15 a. m. and 12 m. to case No. 19—M, in the seventh day of illness.

Result: As already reported (Goldberger and Anderson, 1912), rhesus No. 320 presented no evidence of a typhus reaction following the inoculation with the crushed lice suspension. This animal, after resisting two successive immunity tests with defibrinated blood, was subjected to a third test February 23, 1912, or 75 days after inoculation with the lice suspension. This consisted of an intraperitoneal inocula-

tion of 6 c. c. of virulent defibrinated blood. During 3 days, beginning 14 days after this inoculation, rhesus No. 320 presented a well-defined perturbation in temperature that may have represented an abortive typhus reaction (chart 16).

Rhesus No. 319 presented no evidence of a reaction following the inoculation with the filtrate. Thirty-one days later he was subjected to an immunity test consisting of an intravenous inoculation of 3 c. c. of virulent defibrinated blood of rhesus No. 187; 21 days after this, having in the meantime presented no evidence of a typhus reaction, he was subjected to a second test consisting of an intravenous injection of 2.5 c. c. of virulent defibrinated blood of rhesus No. 115a. Not presenting any evidence of a reaction, he was subjected to a third test on February 23, 75 days after the inoculation with the filtered lice suspension. This test, as in the case of rhesus No. 320, consisted of an intraperitoneal injection of 6 c. c. of virulent defibrinated blood. During a period of observation of 33 days following this, rhesus No. 319 presented no recognizable indications of a typhus reaction.

DISCUSSION.—The immunity tests given monkeys No. 319 and No. 320 would indicate the possession of a complete insusceptibility to typhus by the former and an almost if not quite complete insusceptibility by the latter. In the early stages of our work we would have been inclined to interpret this resistance as due to an immunization induced by the original inoculation of filtered and unfiltered



Temperature curve of rhesus No. 320, showing abortive (?) reaction following third immunity test.

lice suspensions, respectively. Our more recent experience, however, having developed evidence that a healthy monkey may from time to time be encountered that is naturally resistant, a definite interpretation of the result of this single experiment is not permissible. Nevertheless, we undoubtedly have here a suggestion of a filterable phase in the body of the louse.

With the idea of obtaining a clearer conception of the nature of the virus of typhus we have studied its resistance to drying and to certain extremes of temperature.


DRYING.

Experiment No. 14.

At 12 m., October 22, 1911, 3 c. c. of virulent defibrinated blood of rhesus No. 141 (then in the third day of an experimental typhus) were put in one petri dish and 6 c. c. in another. The dishes with the lids tilted were placed over sulphuric acid in a desiccator from which the air was then exhausted. This apparatus with the blood was then placed at a temperature of from about 15° to 19° C. At the end of 25 hours the vacuum was broken and the dish containing the 3 c. c. of blood was removed, after which the air in the apparatus in which the dish with the 6 c. c. of blood remained was again exhausted and the apparatus continued as before.

The 3 c. c. of blood were found to have been dried and to have formed a scale on the bottom of the dish. The dry blood was rubbed up in saline solution and injected intraperitoneally into rhesus No. 130.


At the end of 6 days, that is, on October 28, the dish containing the 6 c. c. of blood was removed from the desiccator. The blood was found dried into a scale. This dried blood was rubbed up in saline solution and injected intraperitoneally into rhesus No. 129.



Temperature curve of rhesus No. 130, showing reaction following immunity test.

Result: Neither rhesus No. 130 nor No. 129 presented any evidence of a reaction. Thirty-eight days after the inoculation of the former and 32 of the latter, both were subjected to an immunity test consisting of an intravenous injection of 2.5 c. c. of defibrinated blood of rhesus No. 170, then in the fourth day of an attack of typhus. Both monkeys promptly responded to this test, No. 130 developing a grave reaction, and No. 129 a reaction of moderate severity (charts 17 and 18).

Monkeys Nos. 130 and 129 were therefore neither infected nor immunized by the inoculation, in the one of 3 c. c. of virulent defibrinated blood after drying 25 hours, and in the other of 6 c. c. after drying 6 days.



Temperature curve of rhesus No. 129, showing reaction following immunity test.

Experiment No. 15.

At 2 p. m., February 12, 1912, 3 c. c. of mixed virulent blood of rhesus Nos. 196, 198, and 322 was put in a petri dish, and the dish, with the cover tilted, placed over sulphuric acid in a desiccator as in experiment No. 14. After exhausting the air the apparatus was placed at 15° C. At 2 p. m., February 13—that is, after 25 hours—the dish with the blood now dried to a scale was removed from the apparatus. The blood was then rubbed up in saline solution and injected intraperitoneally into rhesus No. 208.

Result: During a period of observation of 39 days monkey No. 208 presented no evidence of a reaction. Forty-six days after the inoculation this monkey was subjected to an immunity test consisting of an intraperitoneal injection of 3 c. c. of defibrinated blood of rhesus No. 234 diluted with 2 c. c. of saline solution. To this test monkey No. 208 reacted promptly and sharply.

It would appear, therefore, that the inoculation with 3 c. c. of virulent defibrinated blood after drying for 25 hours had neither infected nor immunized this animal.

CONCLUSIONS.—The negative outcome of these two experiments would indicate that the typhus virus can not survive drying, or, more accurately, that it loses its virulence when dried for 25 hours under the conditions described. Unfortunately the nature of the problem does not permit final conclusions to be drawn from so small a number of experiments.

HEAT.

Gaviño and Girard were the first to study the resistance of the typhus virus to heat. In May, 1910, they inoculated a monkey with 10 c. c. of defibrinated typhus blood after heating at 50° for 40 minutes. The monkey so inoculated developed a well-marked febrile reaction after an incubation period of 14 days.

In a second experiment, performed three months later, they inoculated a monkey with 4 c. c. of defibrinated blood after heating at 55° for 15 minutes. This inoculation not being followed by any evidence of a reaction, Gaviño and Girard concluded that the virulence of the blood was destroyed by the heating to which they had subjected it. The immunity of this animal was not tested until later; when tested 56 days after the inoculation of the heated blood with 3 c. c. of defibrinated typhus blood the monkey gave no evidence of a reaction. Commenting on this result, Gaviño and Girard state that it would seem as if the first noninfecting injection with heated blood vaccinated against a subsequent injection with virulent blood, although they add that the objection might be raised that they had encountered in this animal one that was naturally immune.

In October, 1910, Nicolle, Conor, and Conseil reported an experiment in which the intraperitoneal injection of 4 c. c. of citrated blood (virulent for the controls) after heating at 50° C. for 15 minutes conferred no resistance to a subsequent immunity test. They concluded from this that the virus is destroyed by heating at 50° for 15 minutes. In view of the positive result obtained by Gaviño and Girard, and in the light of the fact several times referred to in other portions of this paper that the monkey is not invariably responsive to an infective inoculation, this conclusion is not permissible.

In a paper published in November, 1911, Gaviño and Girard report additional tests of the resistance of the typhus virus to heat. In one experiment they inoculated three guinea pigs intraperitoneally, each with 3 c. c. of defibrinated typhus blood heated at 55° for 15 minutes. None of these pigs presented any indications of a reaction. The immunity test, if made, is not recorded. In a second experiment they inoculated each of two monkeys with 7 c. c. of defibrinated typhus blood after heating at 55° for 15 minutes. Not presenting any evidence of a reaction, they were subjected to an immunity test one month later. To this both monkeys gave marked reactions. Consequently they conclude that typhus blood heated for 15 minutes at 55° loses its virulence and that a noninfecting injection of heated

blood does not vaccinate against a subsequent inoculation with virulent blood.

The heating experiments recorded in the literature may be summarized as follows:

At 50° for 40 minutes.—One experiment, 1 monkey; blood retained its virulence (Gaviño and Girard).

At 50° for 15 minutes.—One experiment, 1 monkey; not infected, not immunized (Nicolle, Conor, and Conseil).


At 55° for 15 minutes.—One experiment, 1 monkey; not infected, but resisted immunity test (Gaviño and Girard). One experiment, 3 guinea pigs; not infected; immunity test not recorded (Gaviño and Girard). One experiment, 2 monkeys; not infected, not immunized (Gaviño and Girard).

Before attempting to interpret these results, we desire to record some experiments of our own.

Experiment No. 16.

On October 22, 1911, we heated some virulent defibrinated blood of rhesus No. 141 in a tube at 55° for 15 minutes and with it inoculated 2 monkeys: Rhesus No. 131, received 3.5 c. c., and rhesus No. 132, 3 c. c., both intravenously.

Result: Neither of this pair of monkeys presented any evidence of a reaction following their inoculation, although 3 other monkeys, inoculated at the same time



Temperature curve of rhesus No. 131, showing reaction following immunity test.

with like doses of the unheated blood, all developed well marked typhus reactions. Twenty-nine days later rhesus Nos. 131 and 132 were subjected to an immunity test consisting in each of an intravenous injection of 2.5 c. c. of defibrinated blood of rhesus No. 166, then in the third day of a typhus reaction. Both animals reacted sharply (charts 19 and 20); rhesus No. 131, after an incubation of 14 days with a fever of 8 days' duration, and rhesus No. 132 after an incubation of 8 days with a fever of 6 days' duration.

The two animals were therefore neither infected nor immunized.

Experiment No. 17.

On February 14, 1912, we heated some virulent defibrinated blood in a tube at 60° for 5 minutes and injected 3 c. c. of it into the peritoneal cavity of rhesus No. 199.

Result: During a period of observation of 23 days monkey No. 199 gave no indications of a reaction. Tested at the end of this period by an intraperitoneal injection of 6 c. c. of virulent defibrinated blood, rhesus No. 199 developed a well marked febrile reaction.

This animal was therefore neither infected nor immunized by the injection of 3 c. c. of blood heated at 60° for 5 minutes.

Experiment No. 19.

On March 6, 1912, we heated some virulent blood in a tube at 55° for 5 minutes and injected 6 c. c. of it into the peritoneal cavity of rhesus No. 220.

Result: During a period of observation of 24 days this animal presented no recognizable indications of a reaction. Tested at the end of this time by an intraperitoneal injection of 3 c. c. of defibrinated blood of rhesus No. 234, diluted with 2 c. c. of saline solution, monkey No. 220 promptly developed a severe reaction (chart 22).

It would appear, therefore, that monkey No. 220 was neither infected nor immunized by the intraperitoneal injection of 6 c. c. of typhus blood heated at 55° for 5 minutes.


Our experiments may be summarized as follows:

At 55° for 15 minutes.—One experiment; neither infection nor immunization in 2 monkeys.

At 60° for 5 minutes.—One experiment; 1 monkey; neither infection nor immunization.

At 60° for 5 minutes and 55° for 5 minutes, combined.—One experiment; 1 monkey; neither infection nor immunization.

CHART NO. 22



Temperature curve of rhesus No. 220, showing reaction following immunity test.

At 55° for 5 minutes.—One experiment; 1 monkey; neither infection nor immunization.

Combining the summary of the experiments previously recorded with that of our own, we have the following:

50° for 40 minutes.—One experiment, 1 monkey; blood retained its virulence (Gaviño and Girard).

50° for 15 minutes.—One experiment, 1 monkey; not infected, not immunized (Nicolle, Conor, and Conseil).

55° for 15 minutes.—One experiment, 1 monkey; not infected, but resisted immunity test (Gaviño and Girard).

One experiment, 3 guinea pigs; not infected; immunity test not reported (Gaviño and Girard).

One experiment, 2 monkeys; not infected, not immunized (Gaviño and Girard).

One experiment, 2 monkeys; neither infection nor immunization (Goldberger and Anderson).

55° for 5 minutes.—One experiment, 1 monkey; neither infection nor immunization (Goldberger and Anderson).

55° for 5 minutes combined with 60° for 5 minutes.—One experiment, 1 monkey; neither infection nor immunization (Goldberger and Anderson).

60° for 5 minutes.—One experiment, 1 monkey; neither infection nor immunization (Goldberger and Anderson).

Reviewing this summary, we find one experiment by Nicolle, Conor, and Conseil, in which blood heated at 50° for 15 minutes neither infected nor immunized, and one by Gaviño and Girard in which blood heated at 55° for 15 minutes did not infect but apparently immunized, which appear to be out of harmony with the other results. If we bear in mind the possibility, to which we have elsewhere in this paper called attention, that the normal monkey may at times fail to respond to the first and sometimes to the second inoculation with virulent material, we will have (what we believe is) the explanation of this lack of harmony.

CONCLUSIONS.—The determination of the resistance of typhus virus to heat is obviously beset with the same difficulties as the determination of its resistance to drying. Our inability to cultivate the organism hardly permits of the final determination of these points. Nevertheless, when we combine the results of our own heating experiments with those previously recorded we have a body of facts that justify certain conclusions that may be summarized as follows:

1. The virus may (perhaps generally does) retain its virulence after heating at 50° for 40 minutes.
2. The virus is deprived of virulence (? killed) by heating at 55° for 15 minutes.
3. The virus is probably deprived of virulence (? killed) by heating at 60° for 5 minutes.
4. The virus is probably deprived of virulence (? killed) by heating at 55° for 5 minutes.

FREEZING.

So far as we are aware, the following attempts to determine the resistance of the typhus virus to "freezing" are the first to be recorded:

Experiment No. 20.

At 12.30 p. m., October 22, 1911, some defibrinated blood of rhesus No. 141 (chart 3), then in the third day of a typhus reaction, was put into a freezing mixture of ice and hydrochloric acid and frozen solid. At 2.30 p. m. it was removed from this mixture and placed in a tank of brine at 0° C. At 1.30 p. m., October 23, it was taken out of the brine, rapidly thawed and, after warming slightly, 3 c. c. of it were intraperitoneally injected into rhesus No. 155.

Result: After an incubation period of 8 days this monkey developed a well-marked typhus reaction.

The infectivity of the blood was, therefore, not destroyed by "freezing" for 25 hours.

Experiment No. 21.

On February 12, 1912, some virulent defibrinated blood from rhesus Nos. 196, 198, and 322 was frozen as in the preceding experiment and put in a tank of brine at 0° C. The blood was allowed to remain at this temperature for 8 days—that is, till February 20; it was then removed and allowed to thaw at room temperature. Having thawed, it was warmed slightly, and 6 c. c. of it was injected intraperitoneally into rhesus No. 210.

Result: After an incubation period of 9 days, rhesus No. 210 developed a well-defined typhus reaction of about 8 days' duration. He later resisted an immunity test consisting of an intraperitoneal injection of 3 c. c. of defibrinated blood of rhesus No. 324, diluted with 2 c. c. of saline solution.

It appears, therefore, that the infectivity of the blood was not destroyed by "freezing" for 8 days.

Experiment No. 22.

On February 23, 1912, some virulent blood obtained from rhesus Nos. 200, 203, and 204 was frozen and placed in brine at 0° C., as in the preceding experiments. It was kept in the brine at 0° C. for 15 days until March 9, when it was removed and permitted to thaw at room temperature. After warming slightly, 6 c. c. of it was injected intraperitoneally into Java monkey, No. 226.

Result: During a period of observation of 30 days this animal presented no evidence of a reaction. Thirty-two days after the inoculation monkey No. 226 was subjected to an immunity test consisting of an intraperitoneal injection of 3 c. c. of defibrinated blood of rhesus No. 216, diluted with 2 c. c. of saline solution. To this it responded promptly with a severe typhus reaction.

Virulent blood, after "freezing" for 15 days, did not therefore infect or vaccinate.

CONCLUSIONS.—The foregoing experiments are not extensive enough to define more than roughly the degree of resistance of the typhus virus to freezing. It is clear, however, that typhus blood may retain its infectivity after freezing (0° C.) for at least 8 days.

SUMMARY AND CONCLUSIONS.

1. The literature bearing on the duration of the infectivity of the blood is critically reviewed. Two experiments are reported, and the following conclusions are drawn:

(a) The blood of the monkey infected with typhus may be virulent in the prefebrile stage, but no satisfactory evidence of that fact has as yet been adduced.

(b) The blood of the monkey may still be virulent 24 to 32 hours after the return of the temperature to normal.

2. The literature bearing on the question of the localization of the virus is critically reviewed, and it is pointed out that the evidence adduced by Nicolle, Conor, and Conseil, in support of their hypothesis of an intraleucocytic localization, is not valid. New experiments are detailed, and the following conclusions are drawn:

(a) The available evidence favors the view that the typhus virus is extracellular and free in the circulating plasma.

(b) The serum of virulent typhus blood is constantly infective whether obtained from defibrinated blood or after clotting, instances of its apparent avirulence being explicable by a natural resistance of the monkey.

(c) It may perhaps be possible to deprive typhus blood serum of its virulence by prolonged centrifugation; but this does not necessarily indicate an intraleucocytic localization of the virus.

(d) Repeated (three) washings of the blood corpuscles does not deprive them of their infectivity.

3. The literature bearing on the question of the filterability of the typhus virus is critically reviewed. It is found that eight attempts have been recorded to pass the virus through the Berkefeld filter. Of these, six were clearly negative; in one of the other two, the monkey,

without giving any evidence of a reaction to the inoculation, was later found to be resistant to an immunity test; in the other, one of a pair of monkeys is described as having presented a doubtful reaction to the inoculation and later was found resistant to a single immunity test.

New filtration experiments are reported, but in no instance was infection produced; when submitted to an immunity test, two of seven monkeys at first appeared to be resistant, but later, when the test was repeated, both responded.

The conclusion is drawn that there is no evidence to show that the virus in the blood of typhus is able to pass the Berkefeld filter; and, incidentally, that virulent typhus blood contains no toxin, or that it contains it in quantities too small to cause an appreciable increase in the normal resistance of the monkey when injected, even repeatedly, in the doses ordinarily employed for inoculating the monkey.

Two attempts are recorded to filter the virus as it exists in the body of the louse; in one, the monkey inoculated with the filtrate, without giving any indications of a reaction, was subsequently found refractory to repeated immunity tests, suggesting that he had been vaccinated by the filtrate. While this seems to point to the existence of a filterable phase in the body of the louse, it can not be regarded as conclusive without further corroborative work.

4. The resistance of the virus to drying is tested in two experiments with results indicating that the virus is deprived of virulence at the end of 25 hours.

5. The literature bearing on the resistance of the virus to heat is critically reviewed. The results recorded are summarized, some new experiments detailed, and the following conclusions drawn:

(a) The virus may (perhaps generally does) retain its virulence after heating at 50° for 40 minutes.

(b) The virus is deprived of virulence (? killed) by heating at 55° for 15 minutes.

(c) The virus is probably deprived of virulence (? killed) by heating at 60° for 5 minutes.

(d) The virus is probably deprived of virulence (? killed) by heating at 55° for 5 minutes.

6. The resistance of the virus to freezing (0° C.) is tested. It is found that it may retain its infectivity, after freezing (0° C.), for at least 8 days.

REFERENCES.

- ANDERSON, JOHN F., and GOLDBERGER, JOSEPH. On the relation of Rocky Mountain spotted fever to the typhus fever of Mexico. A preliminary note. *Public Health Reports*, xxiv, Dec. 10, 1909, p. 1861.
- Same. A note on the etiology of "tabardillo," the typhus fever of Mexico. *Public Health Reports*, xxiv, Dec. 24, 1909, p. 1941.
- Same. On the infectivity of tabardillo or Mexican typhus for monkeys and studies on its mode of transmission. *Public Health Reports*, xxv, Feb. 18, 1910, p. 177.
- Same. On the etiology of tabardillo or Mexican typhus. An experimental investigation. *Jour. Med. Research*, Boston, June, 1910, p. 469-481.
- GAVIÑO, A., and GIRARD, J. El tifo experimental en los monos inferiores, nota preliminar; el tifo exantemático en los monos inferiores, segunda nota. *Publicaciones del Instituto Bacteriológico Nacional, México*, 1910.
- Same. Tercera nota sobre el tifo experimental en los monos inferiores. *Publicaciones del Instituto Bacteriológico Nacional, México*, Aug. 23, 1910.
- Same. Cuarta nota sobre el tifo experimental en los monos inferiores. *Ibid.*, Nov. 9, 1910.
- Same. Estudio experimental sobre el tifo exantemático. *Ibid.*, Nov. 12, 1911.

- GOLDBERGER, JOSEPH, and ANDERSON, JOHN F.** The transmission of typhus fever, with especial reference to transmission by the head louse (*Pediculus capitis*). Public Health Reports, xxvii, Mar. 1, 1912.
- MOCZUTKOWSKI, O. O.** Ueber die Impfbarkeit des Typhus exanthematicus. Allg. med. Cent.-Zeit., Berlin, 1900, p. 1055-1057.
- NICOLLE, CH.** Reproduction expérimentale du typhus exanthématique chez le singe. C. R. Acad. Sci., Paris, July 12, 1909, p. 157-160.
- Same.** Recherches expérimentales sur le typhus exanthématique entreprises à l'Institut Pasteur de Tunis pendant l'année 1909. Ann. de l'Inst. Pasteur, Paris, Apr., 1910, p. 243-275.
- NICOLLE, CH., and CONSEIL, E.** Reproduction expérimentale du typhus exanthématique chez le macaque par inoculation directe du virus humain. C. R. Acad. Sci., Paris, July 18, 1910, p. 258-260.
- Same.** Données expérimentales nouvelles sur le typhus exanthématique. C. R. Acad. Sci., Paris, Aug. 8, 1910.
- NICOLLE, CH., CONOR, A., and CONSEIL, E.** Sur quelques propriétés du virus exanthématique. C. R. Acad. Sci., Paris, Oct. 17, 1910, p. 685-688.
- NICOLLE, CH., CONSEIL, E., CONOR, A., and JAEGGY, E.** Recherches sur le typhus exanthématique entreprises à l'Institut Pasteur, Tunis, pendant l'année 1910. Mémoire d'ensemble. Ann. de l'Institut Pasteur, Paris, Jan., 1911, p. 1-103.
- NICOLLE, CH., CONSEIL, E., and CONOR, A.** Le typhus expérimental du cobaye. C. R. Acad. Sci., Paris, June 6, 1911.
- NICOLLE, CH., CONOR, A., and CONSEIL, E.** Sur la nature et la siege de l'agent pathogène du typhus exanthématique. C. R. Acad. Sci., Paris, Sept. 18, 1911.
- Same.** Recherches expérimentales sur le typhus exanthématique entreprises à l'Institut Pasteur de Tunis pendant l'année 1911. Ann. de l'Institut Pasteur Paris Apr. 25, 1912.
- OTERO, M.** In "Documentos relativos al concurso abierto para estudiar la etiologia y seroterapia de tabardillo." Dictamen de la Comision. Gaceta Medica de Mexico, Appendice, 1908.
- RICKETTS, HOWARD T., and WILDER, RUSSELL M.:** The typhus fever of Mexico (tabardillo). Preliminary observations. Jour. Am. Med. Assn., Chicago, Feb. 5, 1910, p. 463-467.
- Same.** The etiology of the typhus fever (tabardillo) of Mexico City. A further preliminary report. Journ. Am. Med. Assn., Apr. 23, 1910, p. 1373-1375.
- WILDER, RUSSELL M.:** The problem of transmission in typhus fever. Journ. infec. diseases, Chicago, July, 1911, p. 9-101.

SANITARY ADVICE FOR KEEPERS OF SUMMER RESORTS.

By W. C. Rucker, Assistant Surgeon General, Public Health and Marine-Hospital Service.

Until comparatively recent years if a keeper of a summer resort provided good housing and feeding facilities, suitable and proper social diversions, and a reasonable amount of rest and quiet, the keeper of the resort considered that he had done his duty well, and other things being equal was reasonably sure that he would have a fair return for his investment and effort. While all of these points are of no mean weight in the minds of persons who go to the seashore, the mountains, or the country to avoid the heat and discomforts of the city, they have become secondary, and the question which is now asked first of all is as to the healthfulness of the resort in question. Aside from the financial loss which the manager of a summer pleasure ground may sustain from an outbreak of sickness among his guests, there is a moral obligation to protect and further as far as possible their health interests.

In a general way it may be stated that disease is carried from the sick to the well, either directly by persons or indirectly by food and drink, dust, and insects. If the avenues for disease germs are closed

there need be comparatively little disease. It is thus seen that most disease is environmental, and it is therefore the duty of persons who are in charge of summer resorts to maintain the environment of their guests in as healthful a condition as possible.

The great source of disease in a summer resort is the toilet. In many localities it is of the most primitive pattern, and maintained in a condition of noisome filth which would not be tolerated under any other circumstances. Frequently it is merely a surface privy, to which the barnyard fowls and domestic animals have free access. Again there may be a pit privy, or a cesspool which drains into the well, or into the stream which furnishes the water for the establishment.

Inasmuch as typhoid fever, a disease which by preference spends the warm months at a summer resort, is caused by a germ which is peculiar to man, and which leaves the body of the sick patient in his discharges, it may be seen that any method of disposing of human excrement which does not absolutely insure that it will not be carried to well persons is faulty, not to say criminal. After the germ of typhoid fever has left the body of the sick patient in his discharges, it is carried to well persons in food or drink, or through the intermediation of flies or dirty fingers.

It is the duty of every keeper of a summer resort to provide clean, well-screened places of easement for his guests, and to see to it that the fecal material deposited in this place can not escape and find entrance into the food and drink of the guest. If there is a cesspool, or a pit, it must not leak, and it must be guarded against flies, because these insects, which are bred in filth and lead a life of sin and iniquity, carry the germs on their feet to the food stuffs in the kitchen or dining room, thus delivering the germs in a live and virulent condition to well people.

The selection of a water supply is of the utmost importance. It should not be from a stream which receives the drainage of human habitations. In this connection it may be remarked that the old theory of the self-purification of streams has been exploded. If well water is used, it should not be from a shallow surface well, nor should it be from a well which by any chance may receive the drainage of human dwellings. Other things being equal, driven or artesian wells are far preferable.

If there is any doubt as to the purity of the water, it should be sterilized, either by boiling or by the addition of calcium hypochlorite prior to using.

Mention has been made of the carriage of the germs of typhoid fever by dirty fingers. This does not necessarily mean fingers which are mechanically dirty. It has been found in recent years that a large number of persons who have had typhoid fever continue to pass the germs of that disease in their discharges for a considerable period after their recovery from the disease. These persons are called typhoid carriers. It is of course an impossibility for the manager of a summer resort to examine all of his employees to determine whether or not they may be carriers, but it is his duty, when selecting persons who are to be employed in the preparation and serving of food or cleaning the dishes and kitchen utensils, to inquire whether or not they have had typhoid fever recently, and to employ

none who have been afflicted with that disease within three or four months.

It has been shown that flies carry disease. The model summer resort has no flies. This condition of affairs may be brought about by preventing their breeding, and by screening the hostelry against them. Flies breed in manure. Therefore the barnyard or stables should be kept scrupulously clean, and the manure stored in well-screened metal-lined bins. The stable itself should also be screened. This adds not only to the comfort of the guests, but also saves the horses and cattle a great deal of energy, which would otherwise be expended in fighting flies.

One of the things which city people most desire when going to the country is milk. Milk which is produced by dirty cows in a dirty stable, and secured and handled by dirty dairy hands, is unfit for human consumption. The best advertisement that any country resort can have is a clean, airy, wholesome dairy, in which clean cows are milked by clean laborers, and in which the processes to which the milk is subjected follow a cleanliness of levitical scrupulosity.

It has been recommended that all of the buildings of a summer resort be screened. This not only excludes flies, but it also prevents the entrance of mosquitoes. Aside from the fact that these insects act as the producers of insomnia and irritating inflammations, they are also the purveyors of malaria, which they introduce into the system of the guests by biting them. While it is desirable that they be excluded, it is even more necessary that they be prevented from breeding. Most summer resort keepers breed their own mosquitoes, and a careful survey of the premises will discover small pools of water, unscreened rain-water barrels, and small collections of water in tin cans, flower pots, and other easily overlooked small receptacles, all of which may contain "wigglers," which are the immature forms of mosquitoes. If these are emptied or oiled, or have a considerable quantity of salt placed in them, they will cease to be producers. In fountains and ornamental pools, to which none of these methods may be applied, the planting of small fish will destroy the mosquito larvæ, and thus prevent their multiplication.

The careful tuberculosis patient may be quite harmless in a summer hotel, but those who are careless and expectorate promiscuously are a positive danger, and while such persons are always to be regarded with consideration and charity, it may become necessary to recommend that they spend their summer in a sanitarium, rather than in a public resort where they may endanger others.

There are two sorts of vermin which may be disease carriers, and with which the summer resort keeper is at constant war; these are the rat and the cockroach.

Rats stay where they can get food. Therefore if all food supplies intended for human beings, and all remnants of human food, be guarded against them their numbers will fall off considerably. Furthermore, if stables are maintained in a cleanly condition, and if the rats are excluded from feed rooms, they will find that they must move on or be starved to death. Those which remain may be killed off, either by poisoning with some reliable agent, such as phosphorus or

arsenic paste. Traps will take off a good many, and a good active terrier makes an excellent agent for the extermination of these pests.

Cockroaches may be gotten rid of by starvation, that is, the protection of foodstuffs against them, absolute cleanliness, or by the use of some agent which kills cockroaches but is harmless to man. Borax mixed with sugar has been recommended for this purpose.

Finally let it be stated that if the keeper of a summer resort has any doubt as to the sanitary condition of his premises, or if he is brought face to face with any sanitary problem with which he does not know how to deal, it is his duty to consult the local health officer.

UNITED STATES.

MUNICIPAL ORDINANCES, RULES, AND REGULATIONS PERTAINING TO PUBLIC HYGIENE.

[Adopted since July 1, 1911.]

CHICAGO, ILL.

COMMON DRINKING CUPS—USE OF, IN PUBLIC OR SEMIPUBLIC PLACES PROHIBITED.

SEC. 1. That no person, firm, or corporation as owner, agent, employee, or person in charge or control of any railroad or railway station located within the city of Chicago, any railroad or railway car or train operated from point to point within said city, any public or private school, any municipal or office building, any lodging or boarding house, hotel, club, theater, public hall, amusement ground, factory, office, store, or structure of any name or description whatsoever, shall keep, offer, exhibit, or display for use any common drinking cup, glass, or other receptacle.

SEC. 2. Any person, firm, or corporation violating any of the provisions of section 1 hereof shall be fined not less than \$5 nor more than \$50 for each offense.

SEC. 3. This ordinance shall take effect and be in force ninety (90) days from the date of its passage.

[Ordinance adopted May 8, 1911.]

ELYRIA, OHIO.

MEAT—SLAUGHTER, PREPARATION, CARE, AND SALE.

SECTION 1. No one shall kill and offer for food animals affected with the following-named diseases:

- (1) Anomalies of the blood (anemia, leucemia, hemoglobinemia, icterus, uremia).
- (2) Poisoning from any mineral drug or vegetable.
- (3) Purulent paresis (milk fever).
- (4) Animal parasitism, the parasite known to be directly or indirectly transmissible to man.
- (5) Putrid intoxication (sapremia).
- (6) Pyemia.
- (7) Septicemia.
- (8) Malignant edema.
- (9) Anthrax.
- (10) Foot-and-mouth disease.
- (11) Cowpox, sheep pox when animals show fever.
- (12) Rabies.
- (13) Glanders.
- (14) Tuberculosis.
- (15) Actinomycosis.
- (16) Contagious pleuropneumonia.
- (17) Hemorrhagic septicemia.
- (18) Diphtheritis.
- (19) Dysentery.
- (20) Hog cholera.
- (21) Swine plague.
- (22) Chicken cholera.
- (23) Mycotic gastro-enteritis.
- (24) Texas fever.
- (25) Tetanus.
- (26) Malignant epizootic head catarrh.

- (28) Malignant tumors.
- (29) Puerperal fever.
- (30) Emaciation.
- (31) Advanced pregnancy.
- (32) Recent parturition (at least 15 days must have elapsed).
- (33) Overheated, feverish animals.
- (34) Exhausted, fatigued animals should have 8 hours' rest before being slaughtered.
- (35) Fractures, bruises, contusions (when fever present).
- (36) Advanced stages of mange.
- (37) Advanced stages of footrot.
- (38) Suppurative or gangrenous inflammation of the udder (when fever is present).
- (39) Wounds (when fever is present).
- (40) Any disease of the heart, lungs, liver, kidney, spleen, peritoneum, pleura, or any organ which would render the meat unfit for human food.
- (41) Animals less than 30 days old. (Calves must dress at least 45 pounds.)
- (42) Animals killed or wounded by accident.

SEC. 2. Where an organ or part of a carcass is bruised or injured, the part injured must not be offered for sale; the rest of the carcass, if fit for food, will be allowed to go on the market.

SEC. 3. The body of any animal or part thereof, which is to be used for food, shall not be carted or carried through the streets or avenues unless it be covered so as to protect it from dust and dirt; and no meat, poultry, game, or fish shall be hung or exposed for sale in any street, or outside of any shop or store, or in the open windows and doorways thereof, in the city of Elyria, Ohio. At all times meat, poultry, game, or fish on the markets shall be protected from flies, dust, and dirt by a fine screen.

SEC. 4. The meat of boars and rams shall not be held or offered for sale.

SEC. 5. No meat above the size of a rabbit shall be taken to any public or private market to be sold for human food until the same shall have been fully cooled after killing, nor until the organs of the abdominal and thoracic cavity, head and feet, except of poultry and game, and except the head and feet of swine shall have been removed.

SEC. 6. No one shall sell or offer for sale in the city of Elyria any meat that is kept fresh by salicylic or boracic acid, or any other preservative.

SEC. 7. (a) Every butcher or other person occupying or using any room or building where any cattle are slaughtered or dressed for market or stores shall cause such room or building and all appurtenances to be thoroughly cleansed and all offal, blood, fat, garbage, refuse, and unwholesome or offensive matter to be removed therefrom once every 24 hours after the use thereof for any purpose herein mentioned; and the room or building mentioned must be well drained into a sewer or other place acceptable to the inspector.

(b) All meat dealers, butchers, or fishmongers must keep their stores, salesrooms, market stalls, slaughterhouses and all appurtenances thereto in a clean and sanitary condition, and provide proper drainage and ventilation for the same. Windows and doors shall be provided from May 1 to October 1, inclusive, with sound screens of mesh sufficiently fine to keep out flies and other insects.

SEC. 8. No meat dealer or butcher shall keep meats, fish, or fowls in any refrigerator or ice box unless the same shall be lined with lead or some other proper substance, so as to be water-tight nor unless the same be provided with drainage as prescribed in the plumbing code of this city.

SEC. 9. All meat brought into market must be placed within the stall or stalls of the owner of such meat, and all meat must be removed from the market at the close of each market day, unless the meat is placed in cold storage.

SEC. 10. Lard, when offered for sale, containing any substance other than the fat of swine must be labeled as such, and not sold as pure lard.

SEC. 11. No one shall carry, while bound or tied by their legs, or bound down in any manner, in any vehicle in the city of Elyria, any cattle, sheep, hogs, or calves. Such animals shall be allowed to stand freely in any vehicle when transported and while being therein.

SEC. 12. The fact of any cattle, sheep, hog, or lamb being in stockyard or slaughterhouse pen shall be considered sufficient evidence that the same is being exposed for sale; and the fact that the carcass of any cattle, hog, or lamb, or any part thereof, is found in any public or private market place, dressed and prepared as such meats usually are for market, it shall be deemed as sufficient evidence that the same is on sale, and no animal or part thereof, nor any fish, game, or poultry that has been examined and condemned by the meat inspector or his assistant shall be held, sold, or offered for sale for human food in any market place in the city of Elyria.

SEC. 13. Upon any meat, fowl, fish, or vegetables being found by any inspector of the health department in a condition which renders them, in his opinion, unwholesome and unfit for use as human food, he is empowered, authorized, and directed to immediately condemn the same and cause them to be removed to the garbage plant for destruction, and to report his action to the health officer.

SEC. 14. No person shall vend meat or other product of any cattle, sheep, or swine in the city of Elyria, Ohio, in quantities less than the quarter unless he shall have first obtained a permit from the board of health to do so.

SEC. 15. All meat permits shall be renewed annually in January, for which a charge of \$1 shall be made. Permits issued after July 1 shall be charged for at the rate of 50 cents for each permit covering the second half of the year only.

SEC. 16. All applicants for a permit shall state—

(a) The applicant's name, post-office address, and the place or places of business.

(b) If the applicant buys part or all of his meat from others, the name and addresses of all such persons.

(c) If applicant butchers his own meat, the location of the slaughterhouse, the days and the time of day the larger amount of the killing is done.

SEC. 17. The board of health may refuse to grant such permit and may revoke the same when granted if the applicant or person to whom the permit is issued does not comply with the lawful rules and regulations now in force, or that may hereafter be adopted by the board of health for the sale of meat.

SEC. 18. No butcher or other person shall bring into the city of Elyria or sell or offer for sale in the city of Elyria for human food any calf or any part of the meat thereof which at the time it was killed was less than four weeks old, or any pig or any part of the meat thereof which at the time it was killed was less than five weeks old, or any lamb or any part of the meat thereof which at the time it was killed was less than eight weeks old.

SEC. 19. All animals at the stockyards or in possession of any butcher, intended for slaughter for the city markets or stores, when condemned according to the provisions of this code by the meat inspector, must not be slaughtered except in the presence of the inspector, due notice being sent by the butcher to the health office for that purpose. All carcasses or parts of carcasses that are condemned by the meat inspector shall be rendered unfit for food by treatment with kerosene oil.

SEC. 20. All meat shipped into the city for sale or offered for sale shall be subject to inspection by the meat inspector.

SEC. 21. All meat condemned in the city by Government or State inspectors shall be destroyed under the supervision and subject to the directions of the dairy and food inspector.

SEC. 22. No meat, fish, or vegetables not being fresh, sound, wholesome, or any meat or fish that died of disease or accident shall be brought into the city or offered or held for sale as food anywhere in the city of Elyria, nor shall any such articles be kept or stored therein.

SEC. 23. Whoever violates any provisions of the above resolution, or obstructs or interferes with the execution thereof, or willfully or illegally omits to obey any provisions of said resolution shall be fined not to exceed \$100 or imprisonment for not to exceed 90 days, or both; but no person shall be imprisoned hereunder for the first offense, and the prosecution shall always be as and for a first offense, unless the affidavit upon which the prosecution is instituted contains the allegation that the offense is a second or repeated offense.

SEC. 24. This resolution shall be in force and effect from and after the earliest period allowed by law.

[Ordinance adopted July 28, 1911.]

FLINT, MICH.

FOODSTUFFS—PROTECTION OF FOODSTUFFS AND THE CONSTRUCTION AND MAINTENANCE OF PLACES WHERE FOOD IS PREPARED OR STORED.

Rule 1. No place for the care or storage of food or its manufacture shall be located upon grounds which are unfit and insanitary for the same.

Rule 2. The surfaces of floors, walls, and ceilings are to be hard and as smooth as possible that they may be readily cleaned. Floors which are often wet with decomposable material must be so laid that they drain properly and so that they may be easily washed down.

Tables, benches, dough troughs, etc., are to be freely movable on casters so that the floors underneath are easily accessible. Unnecessary woodwork and fixtures attached to the walls are to be avoided.

The tops of working tables, benches, dough troughs, and similar articles must be smooth and free from unnecessary roughness so that they may readily be cleaned. Resurfacing or redressing shall be done when found advisable to furnish suitable working surfaces.

Rule 3. The doors are to be screened by self-closing screen doors, and all outlets are to be screened. There must be sufficiently screened outlets for ventilation. Fans are to be used if needed to secure ventilation.

There must be sufficient light, preferably daylight, to render easy the use, care, and inspection of all the rooms. This does not apply to storerooms for fruit when ripening where darkness is preferred.

Rule 4. There must be sufficient provision, outside of the workrooms, for storage of food materials, refuse, fuel, unused clothing, and other necessary accessories, not adding to the cleanliness of the workrooms.

Rule 5. No water-closet will be allowed in any room where food or the material from which it is made is kept or used; nor shall there be any direct connection with such rooms. The doors from water-closets must be self-closing by springs or otherwise and be kept closed. Windows in such closets must be screened as well as any other openings, such as ventilators.

Rule 6. Sleeping rooms, if present, must be separated by complete partitions from rooms where food or food materials are used or stored.

Rule 7. The present water furnished by the city must not be used in the preparation of food that is not cooked after such use. No water, unless from a strictly sanitary source, shall be used for such purpose.

Rule 8. The walls and ceilings of workroom are to be whitewashed once every six months, or well painted every five years or oftener, and washed with soap and water every six months or oftener.

Rule 9. All premises and appliances must be kept strictly clean at all times. Racks, hooks, meat-blocks, tables, benches, the walls and doors of coolers, and all utensils are to be scraped or washed and scrubbed often enough to keep them free from accumulations of the materials which come in contact with them.

Rule 10. All food exposed for display or for any purpose outside of screened rooms must be screened. This applies more particularly to the exposure of fruits, etc., in front of stores.

Rule 11. No bottling of milk shall be allowed on the streets. Whenever milk-bottle caps are found carried on milk wagons it shall be considered evidence that such bottling is practiced.

In the interpretation of the above rules, screening, etc., shall be considered necessary only during fly time. And whenever there shall come to the board exceptions made necessary by the character of the work, such as cellars for ripening fruits, etc.; the board may, at its discretion, modify such rules.

[Regulations, Board of Health, adopted Jan. 1, 1912.]

NEW YORK, N. Y.

COMMON DRINKING CUP—USE OF IN PUBLIC PLACES PROHIBITED.

“SEC. 189. The use of a common drinking cup or receptacle for drinking water in any public place or in any public institution, hotel, theater, factory, public hall or public school, or in any railroad station or ferryhouse in the city of New York, or the furnishing of such common drinking cup or receptacle for use in any such place, is hereby prohibited.”

[Addition to sanitary code adopted Mar. 21, 1911. Effective Oct. 1, 1911.]

PLAGUE-PREVENTION WORK.

PLAGUE-INFECTED SQUIRRELS FOUND.

During the week ended April 20, 1912, positive diagnosis was made of 14 plague-infected ground squirrels found in Alameda County, Cal., as follows: April 3, 2 squirrels; April 5, 2 squirrels; April 6, 1 squirrel; April 8, 3 squirrels; April 9, 1 squirrel; April 13, 4 squirrels; and 1 squirrel for which no data relative to place and date of finding were obtainable, proven positive April 16.

DISTRIBUTION OF POISON.

In connection with the making and maintenance of a squirrel-free zone around the cities of California on San Francisco Bay, 10,430 acres of land in Alameda County were covered with poison during the week ended April 20, 1912.

RECORD OF PLAGUE INFECTION.

Places	Date of last case of human plague.	Date of last case of rat plague.	Date of last case of squirrel plague.	Total number of rodents found infected since May, 1907.
California:				
Cities—				
San Francisco.....	Jan. 30, 1908.....	Oct. 23, 1908.....	None.....	308 rats.
Oakland.....	Aug. 9, 1911.....	Dec. 1, 1908.....	do.....	126 rats.
Berkeley.....	Aug. 27, 1907.....	None.....	do.....	None.
Los Angeles.....	Aug. 11, 1908.....	do.....	Aug. 21, 1908.....	1 squirrel.
Counties—				
Alameda (exclusive of Oakland and Berkeley).	Sept. 26, 1909....	Wood rat, Oct. 17, 1909.	Apr. 16, 1912....	135 squirrels and 1 wood rat.
Contra Costa.....	July 21, 1911.....	None.....	Sept. 23, 1911.....	364 squirrels.
Fresno.....	None.....	do.....	Oct. 27, 1911.....	1 squirrel.
Merced.....	do.....	do.....	July 13, 1911.....	5 squirrels
Monterey.....	do.....	do.....	Aug. 6, 1911.....	Do.
San Benito.....	June 5, 1910.....	do.....	June 8, 1911.....	22 squirrels.
San Joaquin.....	Sept. 18, 1911.....	do.....	Aug. 28, 1911.....	18 squirrels.
San Luis Obispo.....	None.....	do.....	Jan. 29, 1910.....	1 squirrel.
Santa Clara.....	Aug. 23, 1910.....	do.....	Oct. 5, 1910.....	23 squirrels.
Santa Cruz.....	None.....	do.....	May 17, 1910.....	3 squirrels.
Stanislaus.....	do.....	do.....	June 2, 1911.....	13 squirrels.
Washington:				
City—				
Seattle.....	Oct. 30, 1907.....	Sept. 21, 1911....	None.....	25 rats.

PLAGUE-PREVENTION WORK—Continued.

RATS COLLECTED AND EXAMINED FOR PLAGUE INFECTION.

Places.	Week ended.	Total collected.	Found dead.	Examined.	Found infected.
California:					
Cities—					
Berkeley.....	Apr. 20, 1912	1 187	164
Oakland.....	do.....	2 789	22	591
San Francisco.....	do.....	1,854	2	1,421
Washington:					
City—					
Seattle.....	do.....	1,032	988

1 Identified: *Mus norvegicus*, 163; *Mus musculus*, 24.

2 Identified: *Mus norvegicus*, 611; *Mus rattus*, 2; *Mus musculus*, 176.

3 Identified: *Mus norvegicus*, 913; *Mus rattus*, 265; *Mus musculus*, 433; *Mus alexandrinus*, 243.

SQUIRRELS COLLECTED AND EXAMINED FOR PLAGUE INFECTION.

During the week ended April 20, 1912, 694 squirrels from Alameda County, Cal., were examined for plague infection. Fourteen were found to be plague infected.

CEREBROSPINAL MENINGITIS.

CASES AND DEATHS REPORTED BY CITY HEALTH AUTHORITIES FOR THE WEEK ENDED MAY 11, 1912.

City.	Cases.	Deaths.	City.	Cases.	Deaths.
Boston, Mass.....	1	New Orleans, La.....	2	4
Chicago, Ill.....	5	1	Newport, Ky.....	1
Cincinnati, Ohio.....	7	4	New York, N. Y.....	10	4
Danville, Ill.....	1	Omaha, Nebr.....	1
El Paso, Tex.....	5	4	Peoria, Ill.....	1
Haverhill, Mass.....	1	2	Philadelphia, Pa.....	1
Johnstown, Pa.....	1	Pittsburgh, Pa.....	1
Kansas City, Kans.....	6	St. Louis, Mo.....	2
Kansas City, Mo.....	6	8	San Antonio, Tex.....	1
Los Angeles, Cal.....	2	Saratoga Springs, N. Y.....	1
Nashville, Tenn.....	2	2	South Omaha, Nebr.....	1

ERYSIPELAS.

CASES AND DEATHS REPORTED BY CITY HEALTH AUTHORITIES FOR THE WEEK ENDED MAY 11, 1912.

City.	Cases.	Deaths.	City.	Cases.	Deaths.
Beaver Falls, Pa.....	1	New York, N. Y.....	29	13
Boston, Mass.....	4	Norristown, Pa.....	1
Cincinnati, Ohio.....	2	1	Oklahoma, Okla.....	1
Cleveland, Ohio.....	4	1	Philadelphia, Pa.....	9	2
Kalamazoo, Mich.....	1	Pittsburgh, Pa.....	6
Lancaster, Pa.....	1	Reading, Pa.....	1
Los Angeles, Cal.....	2	1	St. Louis, Mo.....	9	1
Melrose, Mass.....	1	San Diego, Cal.....	1
Milwaukee, Wis.....	3	South Bethlehem, Pa.....	2
Moline, Ill.....	1	Wilkes-Barre, Pa.....	1
Newark, N. J.....	1	Yonkers, N. Y.....	1

PELLAGRA.

During the week ended May 11, 1912, pellagra was reported as follows: Chattanooga, Tenn., 1 case; Richmond, Va., 1 case; Wilmington, N. C., 1 case, 2 deaths.

Vicinity of Brownsville, Tex.

Acting Asst. Surg. Fairbanks reports May 18 the occurrence of two deaths from pellagra in one family in the vicinity of Brownsville, Tex.

PNEUMONIA.

CASES AND DEATHS REPORTED BY CITY HEALTH AUTHORITIES FOR THE WEEK ENDED MAY 11, 1912.

City.	Cases.	Deaths.	City.	Cases.	Deaths.
Ann Arbor, Mich.	1		Newark, N. J.		5
Aurora, Ill.		1	New Bedford, Mass.		5
Baltimore, Md.		13	Newburyport, Mass.		1
Binghamton, N. Y.	7	2	New Orleans, La.		5
Boston, Mass.		34	New York, N. Y.		120
Brockton, Mass.		1	Niagara Falls, N. Y.		2
Brookline, Mass.		1	Norristown, Pa.		2
Cambridge, Mass.		4	Oakland, Cal.		2
Chelsea, Mass.		4	Omaha, Nebr.		7
Chicago, Ill.	29	85	Pasadena, Cal.		1
Chicopee, Mass.		1	Peoria, Ill.		3
Cincinnati, Ohio		5	Philadelphia, Pa.	24	45
Coffeyville, Kans.	1		Pittsburgh, Pa.	18	31
Danville, Ill.		1	Plainfield, N. J.		3
Dayton, Ohio.		7	Providence, R. I.		6
Denver, Colo.		7	Reading, Pa.		4
Duluth, Minn.		2	Richmond, Va.		3
Dunkirk, N. Y.	1		Rutland, Vt.		1
East Orange, N. J.		1	San Antonio, Tex.		3
El Paso, Tex.		2	Saginaw, Mich.		2
Evansville, Ind.		1	Salem, Mass.		3
Fall River, Mass.		5	San Francisco, Cal.	8	
Fort Wayne, Ind.		1	Saratoga Springs, N. Y.		1
Galesburg, Ill.	1	1	Schenectady, N. Y.	5	2
Grand Rapids, Mich.	1	2	South Bethlehem, Pa.	1	
Harrisburg, Pa.		2	South Omaha, Nebr.		3
Hartford, Conn.		6	Springfield, Ill.		1
Homestead, Pa.		1	Springfield, Mass.		2
Kalamazoo, Mich.	4	3	Superior, Wis.		2
Kansas City, Mo.	3	4	Taunton, Mass.		1
Lawrence, Mass.		7	Toledo, Ohio.		5
Los Angeles, Cal.	2	6	Washington, D. C.		6
Lowell, Mass.		9	Wilkes-Barre, Pa.		3
Lynchburg, Va.		1	Wilkinsburg, Pa.	1	
Lynn, Mass.		2	Williamsport, Pa.		3
Malden, Mass.		1	Wilmington, Del.		6
Marinette, Wis.		1	Wilmington, N. C.	1	1
Moline, Ill.		1	Yonkers, N. Y.		3
Montgomery, Ala.		2	Zanesville, Ohio.		1
Nashville, Tenn.		6			

POLIOMYELITIS.

During the week ended May 11, 1912, poliomyelitis was reported as follows: Boston, Mass., 1 case; New York, N. Y., 6 cases, with 2 deaths; South Bethlehem, Pa., 1 death.

TETANUS.

CASES AND DEATHS REPORTED BY CITY HEALTH AUTHORITIES
FOR THE WEEK ENDED MAY 11, 1912.

City.	Cases.	Deaths.
Chicago, Ill.		2
Dayton, Ohio.		1
New Orleans, La.		1
Philadelphia, Pa.	1	1

SMALLPOX IN THE UNITED STATES.

In the following table the States indicated by an asterisk are those from which reports of smallpox are received only from certain city, and in some cases county, boards of health. In these States, therefore, the recorded cases and deaths should not be taken as showing the general prevalence of the disease. In the States not marked by an asterisk the reports are received monthly from the State boards of health and include all cases reported to the State authorities.

REPORTS RECEIVED DURING WEEK ENDED MAY 31, 1912.

Places.	Date.	Cases.	Deaths.	Remarks.
Arizona:				
Counties—				
Cochise	Feb. 1-29	11		
Pima	do.	5	1	
Total for State		16	1	
Gila	Mar. 1-31	5		
Pima	do.	5		
Total for State		10		
Illinois:				
Counties—				
Bond	Apr. 1-30	2		
Boone	do.	2		
Bureau	do.	12		
Cook	do.	16		
Crawford	do.	1		
Dekalb	do.	1		
Henry	do.	12		
Jefferson	do.	1		
Kane	do.	12		
Lasalle	do.	24		
Lawrence	do.	1		
McDonough	do.	2		
McLean	do.	2		
Madison	do.	1		
Marion	do.	1		
Randolph	do.	2		
St. Clair	do.	5		
Sangamon	do.	2		
Stephenson	do.	1		
Vermilion	do.	1		
Swan Township	do.	5		
Taylor Springs	do.	3		
Total for State		109		
Indiana:				
Counties—				
Bartholomew	Apr. 1-30	30	1	
Brown	do.	1		
Cass	do.	1		
Clark	do.	15		
Daviess	do.	2		
Dearborn	do.	1		
Decatur	do.	3		

SMALLPOX IN THE UNITED STATES—Continued.

Reports Received during week ended May 31, 1912.

Places.	Date.	Cases.	Deaths.	Remarks.
Indiana—Continued.				
12 Counties—Continued.				
Delaware.....	Apr. 1-30.....	4		
Fayette.....	do.....	4		
Gibson.....	do.....	17		
Grant.....	do.....	3		
Greene.....	do.....	1		
Howard.....	do.....	2		
Johnson.....	do.....	9		
Knox.....	do.....	3		
Madison.....	do.....	4		
Marion.....	do.....	2		
Pike.....	do.....	2		
Randolph.....	do.....	1		
St. Joseph.....	do.....	9	1	
Sullivan.....	do.....	11		
Vanderburg.....	do.....	5		
Vigo.....	do.....	2		
Wabash.....	do.....	2		
Wayne.....	do.....	3		
Total for State.....		137	2	
*Missouri:				
St. Louis.....	May 12-18.....	1		
Montana:				
Counties—				
Chouteau.....	Apr. 1-30.....	1		
Dawson.....	do.....	1		
Fergus.....	do.....	2		
Hill.....	do.....	2		
Lewis and Clark.....	do.....	1		
Meagher.....	do.....	3		
Musselshell.....	do.....	8		
Silverbow.....	do.....	1		
Valley.....	do.....	10		
Total for State.....		29		
North Carolina:				
Counties—				
Anson.....	Apr. 1-30.....	36		
Beaufort.....	do.....	5		
Buncombe.....	do.....	2		
Caldwell.....	do.....	4		
Carteret.....	do.....	3		
Craven.....	do.....	14		
Cumberland.....	do.....	3		
Duplin.....	do.....	4		
Gates.....	do.....	2		
Granville.....	do.....	4		
Hertford.....	do.....	1		
Johnston.....	do.....	2		
Lincoln.....	do.....	11		
Madison.....	do.....	24		
New Hanover.....	do.....	13		
Pender.....	do.....	3		
Robeson.....	do.....	8		
Swain.....	do.....	1		
Wake.....	do.....	1		
Warren.....	do.....	4		
Total for State.....		145		
Vermont:				
Counties—				
Caledonia.....	Apr. 1-30.....	9		
Lamoille.....	do.....	1		
Orange.....	do.....	1		
Windsor.....	do.....	13		
Total for State.....		24		
Grand total for the United States.....		471	3	

FORT WORTH, TEX.

Dr. R. B. West, health officer of Tarrant County, reports, May 22, regarding the occurrence of smallpox in Fort Worth and Tarrant County as follows:

From January 19 to May 22, 1912, there were admitted to the isolation hospital at Fort Worth 142 cases of smallpox, of which 37 died. In the county outside of the city there was a total of 21 cases with 9 deaths. These cases were quarantined at their homes. Of the cases admitted to the hospital, 89 cases with 25 deaths were among white Americans, 30 cases with 5 deaths among negroes, and 23 cases with 7 deaths among Mexicans. Of the above cases, 16 with 4 deaths were in children under 10 years of age. Of all the cases coming under supervision, only 3 showed a vaccination scar, and of these 3, 2 had smallpox in so mild a type that they did not go to bed. The other case was in a man who had been partially paralyzed for three years, and was, therefore, confined to his bed, but, like the other 2 cases, this one was also mild.

The first cases appear to have come from Oklahoma, and were in the person of tramps cared for temporarily by the Salvation Army. Another focus was among Mexicans and negroes, and was not discovered until after the death of two Mexicans and a negro. Another focus was in a rooming house, where a mild case diagnosed as chickenpox occurred.

The disease occurred among people in all walks of life. There were a number of other cases which were quarantined in their homes in the city and were not sent to the isolation hospital. These are not included in the above figures.

MORBIDITY AND MORTALITY.

MORBIDITY AND MORTALITY TABLE, CITIES OF THE UNITED STATES, FOR WEEK ENDED MAY 11, 1912.

Cities.	Popula- tion United States Census 1910.	Total deaths from all causes.	Diph- theria.		Measles.		Scarlet fever.		Small- pox.		Tubercu- losis.		Ty- phoid fever.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
<i>Cities having over 500,000 inhabitants.</i>														
Baltimore, Md.	558,485	197	6	...	14	...	24	1	...	45	34	10
Boston, Mass.	670,585	194	28	2	167	...	38	79	19	9
Chicago, Ill.	2,185,283	629	106	12	302	...	218	15	5	102	88	13	3	...
Cleveland, Ohio.	560,663	171	31	3	142	1	43	5	...	32	9	4	2	...
New York, N. Y.	4,766,883	1,461	284	24	1,671	29	444	20	...	487	174	29	4	...
Philadelphia, Pa.	1,549,008	476	55	11	51	4	73	5	...	105	57	19	1	...
Pittsburg, Pa.	533,905	132	10	2	165	1	53	1	...	36	12	6	1	...
St. Louis, Mo.	687,029	200	25	...	45	...	39	1	...	55	27	6
<i>Cities having from 300,000 to 500,000 inhabitants.</i>														
Cincinnati, Ohio.	364,463	102	4	1	25	...	18	31	15	3
Detroit, Mich.	465,766	153	12	33	...	2
Los Angeles, Cal.	319,198	105	8	...	13	...	4	5	...	11	9	4	1	...
Milwaukee, Wis.	373,857	...	12	2	168	3	19	2	5	18	10	6	3	...
Newark, N. J.	347,469	105	28	1	63	1	20	32	17	2
New Orleans, La.	339,075	136	2	...	24	...	5	...	6	25	20	8
San Francisco, Cal.	416,912	120	5	2	24	...	6	...	1	23	12
Washington, D. C.	331,069	99	3	...	89	...	1	34	16	9

MORBIDITY AND MORTALITY—Continued.

Morbidity and mortality table, cities of the United States, for week ended May 11, 1912—Continued.

Cities.	Population United States Census 1910.	Total deaths from all causes.	Diphtheria.		Measles.		Scarlet fever.		Small-pox.		Tuberculosis.		Typhoid fever.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
<i>Cities having from 200,000 to 300,000 inhabitants.</i>														
Denver, Colo.	213,381	72	1	4	29	8	4				14			
Jersey City, N. J.	267,779	82	1								8			
Kansas City, Mo.	248,381	18		8	4	4					2	6		
Providence, R. I.	224,326	87	10	14	4	16						8		
<i>Cities having from 100,000 to 200,000 inhabitants.</i>														
Cambridge, Mass.	104,839	27	8		72	2	4				8	5		
Dayton, Ohio	116,577	43			55		2		1			4		
Fall River, Mass.	119,295	27	8	1	2		3			1	11	5	2	
Grand Rapids, Mich.	112,571	25	2		10		5				3	1	4	
Lowell, Mass.	106,294	40	3		40	1	3				3	4	3	
Nashville, Tenn.	110,364	55	2								7	5		
Oakland, Cal.	150,174	49	4	1	7		2				4	2	3	
Omaha, Nebr.	124,096	34	1				3				2	1		
Richmond, Va.	127,628	36	1		27		3		2		6	3	1	
Toledo, Ohio.	168,497	53	2	9	48	1	3	1			6	3	3	1
Worcester, Mass.	145,986	41	7		12	1	8				7	1	2	
<i>Cities having from 50,000 to 100,000 inhabitants.</i>														
Altoona, Pa.	52,127	8	2									2		
Bayonne, N. J.	55,545	16			11	1	6				3	2		
Brockton, Mass.	56,878	15	1		50		1				8	3	2	
Camden, N. J.	94,538		4		1		3				2	1	1	
Duluth, Minn.	78,466	28					7				4	4		
Evansville, Ind.	69,647	23			3		1				4	2	2	
Fort Wayne, Ind.	63,933	15	3		25		1						1	
Harrisburg, Pa.	64,186	15	4	1	53						4	2		
Hartford, Conn.	98,915	32	3	2	18		9				6	1		
Hoboken, N. J.	70,324	13	2		2		5				2			
Johnstown, Pa.	55,482	9			9		2		5			1		
Kansas City, Kans.	82,331		2		6						3		1	
Lawrence, Mass.	85,892	28	1		3		2	1			3	3		1
Lynn, Mass.	89,336	14			12		6	1			3	1	1	
Manchester, N. H.	70,063	16	1		17		3	1			2	2		
New Bedford, Mass.	96,652	30	2	1	2		1				6	5		
Oklahoma City, Okla.	64,205	6	2								1	1	4	
Pawtucket, R. I.	51,622											2		
Peoria, Ill.	66,950		6											
Reading, Pa.	96,071	33	5	1	2		1				1		1	
Saginaw, Mich.	50,510	13	1				5				2	2		
San Antonio, Tex.	96,614	84	2		2		4				11	1		
Schenectady, N. Y.	72,826	29	3		62	2	6				1	2	1	
South Bend, Ind.	53,684	9	1		1		1				1	1		
Springfield, Ill.	51,678	10	1					1				1		
Springfield, Mass.	88,926	23	3		9		13				7	2		
Trenton, N. J.	96,815	42	5		3		5				7	4		
Wilkes-Barre, Pa.	67,105	18			3		2				3	3	2	
Wilmington, Del.	87,411	26		1								5		
Yonkers, N. Y.	79,803	23	15	1	6		4				4	3		
<i>Cities having from 25,000 to 50,000 inhabitants.</i>														
Atlantic City, N. J.	46,150	14	3										1	
Aurora, Ill.	29,807	7			1							1		
Berkeley, Cal.	40,434	5			44		1				1	1		
Binghamton, N. Y.	48,443	22			2		1				1	2	4	1
Brookline, Mass.	27,792	11			16									
Chattanooga, Tenn.	44,604		1				1				1			
Chelsea, Mass.	32,452	14			29		1				2		1	
Chester, Pa.	38,537													
Chicopee, Mass.	25,401	8			4								1	
Danville, Ill.	27,871	10			2						1	1		
East Orange, N. J.	34,371	4	1		13		3				1			

MORBIDITY AND MORTALITY—Continued.

Morbidity and mortality table, cities of the United States, for week ended May 11, 1912—Continued.

Cities.	Population United States Census 1910.	Total deaths from all causes.	Diphtheria.		Measles.		Scarlet fever.		Smallpox.		Tuberculosis.		Typhoid fever.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
<i>Cities having less than 25,000 inhabitants—Con.</i>														
Nanticoke, Pa.....	18,569				3	1								
Newburyport, Mass.....	19,240	6									1	1		
North Adams, Mass.....	22,012	6		6								1		
Northampton, Mass.....	19,431	5	9	3							2			
Palmer, Mass.....		4												
Plainfield, N. J.....	22,050	7			7							1		
Pottstown, Pa.....		3												
Rutland, Vt.....			1		2							1		
Saratoga Springs, N. Y.....		5			2									
South Bethlehem, Pa.....	19,973	9		1			1				1	2		
Steelton, Pa.....	14,246	4	2		12				1		3			
Warren, Pa.....	11,080	1					1							
Wilkesburg, Pa.....	18,294	7					1				2			
Woburn, Mass.....	15,308	7			3		1				1			

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES OF THE UNITED STATES (Untabulated).

MASSACHUSETTS.—Week ended March 2, 1912. Population of reporting towns, 2,593,485. Total number of deaths from all causes 884, including diphtheria 9, measles 3, scarlet fever 2, tuberculosis 84, typhoid fever 4. Cases reported: Diphtheria 86, measles 721 scarlet fever 144, smallpox 11, tuberculosis 193, typhoid fever 12.

Week ended March 9, 1912. Total number of deaths from all causes 874, including diphtheria 8, measles 9, scarlet fever 3, tuberculosis 84. Cases reported: Diphtheria 81, measles 561, scarlet fever 119, smallpox 3, tuberculosis 150, typhoid fever 16.

Week ended March 16, 1912. Population of reporting towns, 2,577,270. Total number of deaths from all causes 880, including diphtheria 10, measles 4, scarlet fever 3, tuberculosis 86, typhoid fever 3. Cases reported: Diphtheria 90, measles 646, scarlet fever 164, tuberculosis 192, typhoid fever 18.

Week ended March 23, 1912. Population of reporting towns, 2,593,485. Total number of deaths from all causes 824, including diphtheria 4, measles 8, scarlet fever 1, tuberculosis 63, typhoid fever 4. Cases reported: Diphtheria 93, measles 628, scarlet fever 124, tuberculosis 144, typhoid fever 20.

Week ended March 30, 1912. Population of reporting towns, 2,569,087. Total number of deaths from all causes 911, including diphtheria 8, measles 3, scarlet fever 3, tuberculosis 87, typhoid fever 1. Cases reported: Diphtheria 105, measles 698, scarlet fever 123, tuberculosis 193, typhoid fever 17.

FOREIGN AND INSULAR.

AUSTRALIA.

Sydney—Examination of Rats.

The following information was taken from the bulletin issued by the department of public health of New South Wales: During the two weeks ended March 30, 1912, 689 rats were examined for plague infection. No plague-infected rats were found.

The last case of human plague was reported May 29, 1909.

The last plague-infected rat was found April 25, 1910.

BRITISH EAST AFRICA.

Smallpox and Vaccination at Mombasa.

Consul Weddell, at Zanzibar, reports the occurrence of 5 cases of smallpox at Mombasa during the month of March, 1912. During the same period 4,500 persons were vaccinated. The outbreak is believed to have originated in the Mecca pilgrimage.

EGYPT.

Cairo—Typhus Fever.

Consul Knabenshue reports the occurrence of 10 deaths from typhus fever at Cairo during the two weeks ended April 15.

GREAT BRITAIN AND IRELAND.

Typhus Fever.

At Belfast the American consul reports the occurrence of 1 death from typhus fever during the week ended April 6.

At Dublin Consul Adams reports a death from typhus fever during the week ended March 30.

At Glasgow Consul McCunn reports a death from typhus fever for the week ended April 19.

GREECE.

Cerebrospinal Meningitis.

Vice Consul Melissinos at Athens reports the occurrence of 93 cases of cerebrospinal meningitis in Greece during the week ended May 4. The cases occurred at 21 localities. Of the 93 reported cases 17 occurred at Athens and 7 at Piraeus, the port of Athens.

HAWAII.

Record of Plague Infection.

The last case of human plague at Honolulu occurred July 12, 1910. The last plague-infected rat was found at Aiea, 9 miles from Honolulu, April 12, 1910.

A case of human plague was reported at Kapulena, Hawaii, October 28, 1911.

At Hilo the last case of human plague occurred March 23, 1910. At Honokaa, 60 miles from Hilo, a fatal case occurred February 9, 1912; 2 fatal cases February 25, 1912; and a fatal case March 18, 1912. During the week ended March 2, 1912, 49 plague-infected rats were found. The last plague-infected rat was found April 17, 1912. At Hilo a plague-infected rat was found during the week ended June 10, 1911, and 2 plague-infected rats were reported found February 29, 1912.

Honolulu—Plague-prevention Work.

Chief Quarantine Officer Trotter reports:

Week ended April 27, 1912.

Total rats and mongooses taken	460
Rats trapped	419
Mongooses trapped	11
Rats found dead (<i>Mus norvegicus</i>)	1
Rats killed by sulphur dioxide	29
Rats examined bacteriologically	363
Classification of rats trapped:	
<i>Mus alexandrianus</i>	52
<i>Mus musculus</i>	108
<i>Mus norvegicus</i>	47
<i>Mus rattus</i>	212
Average number of traps set daily	1,720

INDIA.

Calcutta—Cholera and Plague.

Acting Asst. Surg. Allan reports: During the week ended March 30, 1912, 89 deaths from cholera and 133 from plague were reported at Calcutta; in all Bengal, 3,925 cases of plague with 3,455 deaths; in all India, 16,028 cases of plague with 13,940 deaths.

ITALY.

Catania—Typhus Fever.

The American consul reports the occurrence of 2 deaths from typhus fever at Catania during the 2 weeks ended April 26.

Examination of Emigrants.

Surg. Geddings at Naples reports:

Vessels inspected at Naples and Palermo week ended May 4, 1912.

NAPLES.

Date.	Name of ship.	Destination.	Steerage passengers inspected and passed.	Pieces of baggage inspected and passed.	Pieces of baggage disinfected.
Apr. 30	Oceania.....	New York.....	1,993	250	2,180
30	Canada.....	do.....	1,090	180	1,320
May 1	Cretic.....	Boston.....	859	120	1,050
3	Berlin.....	New York.....	1,734	220	2,150
	Total.....		5,676	770	6,700

PALERMO.

May 1	Oceania.....	New York.....			
1	Canada.....	do.....	679	600	450
2	Cretic.....	Boston.....			
4	Berlin.....	New York.....	341	250	125
	Total.....		1,020	850	575

JAPAN.**Plague in Formosa.**

Consul Reat at Tamsui reports the occurrence of 66 cases of plague with 56 deaths in the island of Formosa during the 4 weeks ended April 13, 1912.

JAVA.**Batavia—Typhus Fever.**

Consul Rairden reports the occurrence of 2 cases of typhus fever with 1 death at Batavia during the week ended April 6.

MEXICO.**Typhus Fever.**

The American consul reports the occurrence of a death from typhus fever at Matamoras during the week ended May 12.

Puerto Mexico—Yellow Fever.

Acting Assistant Surgeon Thompson reports the occurrence May 25 of a death from yellow fever at Puerto Mexico.

Yellow Fever at San Juan Bautista.

The American consul at Frontera reported May 25 that information had been received of the occurrence of four additional cases of yellow fever with 1 death at San Juan Bautista.

RUSSIA.

Typhus Fever.

Consul General Snodgrass at Moscow reports the occurrence of a death from typhus fever during the week ended April 20.

The American consul at Omsk reports 1 case of typhus fever for the week ended April 11.

TURKEY IN ASIA.

Basra—Plague on Vessel.

The steamship *Adalia* arrived April 10 at Basra with two cases of plague on board. The cases were bacteriologically verified. On April 12 a third case of plague occurred among the crew of the vessel. The original port of departure of the *Adalia* was New Castle. The vessel called at Bushire, on the Persian Gulf. At Bushire 78 cases of plague with 51 deaths were reported during the week ended April 6 and during the week ended April 13, 110 cases with 84 deaths.

VENEZUELA.

La Guaira—Yellow Fever in Vicinity.

Acting Assistant Surgeon Stewart reports: During the week ended April 30 a death from yellow fever was reported at Maiquetia, a suburb of La Guaira.

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX.

REPORTS RECEIVED DURING WEEK ENDED MAY 31, 1912.

[These tables include cases and deaths recorded in reports received by the Surgeon General, Public Health and Marine-Hospital Service, from American consuls through the Department of State and from other sources.]

CHOLERA.

Places.	Date.	Cases.	Deaths.	Remarks.
India:				
Bassein.....	Apr. 1-6.....	3	2	
Bombay.....	Apr. 14-27.....	13	9	
Calcutta.....	Mar. 24-30.....		89	
Madras.....	Apr. 14-27.....	7	5	
Moulmine.....	Apr. 1-6.....	8	7	
Indo-China:				
Saigon.....	Mar. 19-31.....	90	60	
Do.....	Apr. 1-15.....	110	69	And vicinity.

YELLOW FEVER.

Brazil:				
Manaos.....	Apr. 28-May 4.....		2	
Canal Zone:				
Culebra Island Quarantine Station.	Mar. 2-4.....	1	1	From s. s. Chile from Guayaquil.
Mexico:				
Puerto Mexico.....	May 25.....		1	
San Juan Bautista.....	do.....	4	1	
Venezuela:				
Caracas.....	Mar. 1-31.....		4	

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received during week ended May 31, 1912.

PLAGUE.

Places.	Date.	Cases.	Deaths.	Remarks.
India:				
Bombay.....	Apr. 14-27.....	339	320	
Calcutta.....	Mar. 24-31.....		133	
Karachi.....	Apr. 14-27.....	213	190	
Indo-China:				
Saigon.....	Mar. 19-31.....	11	5	
Do.....	Apr. 1-15.....	10	4	
Japan:				
Formosa.....	Apr. 1-13.....	27	23	
Java:				
Paseroean residency.....	Apr. 17-13.....	5	4	
Mauritius:				
.....	May 15-21.....	10	7	
Persia:				
Bushire.....	Apr. 1-13.....	188	135	
Mohammerah.....	Apr. 3 ¹	1		
Straits Settlements:				
Singapore.....	Apr. 7-13.....	2	2	
Turkey in Asia:				
Basra.....	Apr. 10-12 ²	3		From S. S. Adalia from Newcastle.

SMALLPOX.

British East Africa:				
Mombasa.....	Mar. 1-31.....	5		
Brazil:				
Para.....	Apr. 21-May 4.....	4	1	
Canada:				
Quebec.....	May 12-18.....	6		
Vinal Haven.....	May 17.....	7		5 miles from Eastport, Me.
Ceylon:				
Colombo.....	Apr. 7-13.....	1		
Egypt:				
Cairo.....	Apr. 9-29.....	6	1	
France:				
Marseille.....	Apr. 1-30.....		2	
Paris.....	Apr. 14-May 4.....	2		
Germany:				
.....	Apr. 21-27.....	6		Additional.
Do.....	Apr. 28-May 11.....	34		
Great Britain:				
Bradford.....	Apr. 28-May 4.....	1		
Bristol.....	May 5-11.....	1		
India:				
Bombay.....	Apr. 14-27.....	210	134	
Calcutta.....	Mar. 24-30.....		6	
Madras.....	Apr. 14-27.....	16	6	
Indo-China:				
Saigon.....	Mar. 19-31.....	5	2	
Do.....	Apr. 1-15.....	7	2	
Italy:				
Leghorn.....	May 5-11.....	3		
Naples.....	Apr. 28-May 11.....	3		
Palermo.....	Apr. 28-May 4.....	9	3	
Turin.....	May 6-12.....	1		
Java:				
Batavia.....	Apr. 7-13.....	5	2	
Mexico:				
Guadalajara.....	May 5-11.....	1	1	
Juarez.....	May 12-18.....	3		
Mazatlan.....	May 8-14.....		1	
Mexico.....	Mar. 24-Apr. 6.....	32	20	
San Luis Potosi.....	Mar. 9-16.....		2	
Tapachula.....	Apr. 28-May 4.....		1	
Portugal:				
Lisbon.....do.....	2		
Russia:				
Moscow.....	Apr. 14-20.....	3	1	
Odessa.....	Apr. 21-27.....	2		
Riga.....	Apr. 21-May 4.....	7		
St. Petersburg.....	Apr. 21-27.....	20	6	

¹ From the Veröffentlichungen des Kaiserlichen Gesundheitsamtes, May 8, 1912.² Bulletin Quarantenaire d'Egypte, May 9, 1912.

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received during week ended May 31, 1912.

SMALLPOX—Continued.

Places.	Date.	Cases.	Deaths.	Remarks.
Spain:				
Almeria.....	Apr. 1-30.....		1	
Barcelona.....	Apr. 27-May 3.....		1	
Seville.....	Apr. 24-30.....		2	
Valencia.....	Apr. 29-May 11.....	37	2	
Turkey in Asia:				
Beirut.....	Apr. 14-May 4.....	45		
Turkey in Europe:				
Constantinople.....	Apr. 29-May 12.....		17	
Venezuela:				
Caracas.....	Apr. 1-30.....	3		

REPORTS RECEIVED FROM DEC. 30, 1911, TO MAY 24, 1912.

[For reports received from July 1, 1911, to Dec. 29, 1911, see PUBLIC HEALTH REPORTS for Dec. 29, 1911. In accordance with custom, the tables of epidemic diseases are terminated semiannually and new tables begun.]

CHOLERA.

Places.	Date.	Cases.	Deaths.	Remarks.
Arabia:				
Hodeida.....	Jan. 21.....	2	1	
Ras-el-Ketib.....	Dec. 27-Jan. 1.....			Total cases, 22 deaths, 12; mainly in the military hospital.
Austria-Hungary:				
Coastland—				
Capodistria.....	Dec. 14-24.....	2	2	
Croatia and Slavonia.....				Total Oct. 22-Dec. 16: Cases, 36.
Sriem.....	Oct. 22-Dec. 16.....	36		
Hungary.....				Total Nov. 19-Dec. 23: Cases, 37. Free Dec. 28.
Backs-Bodog.....	Dec. 10-16.....	9	5	
Jasz-Nagykun-Szolnok.....	Dec. 3-23.....	11	7	
Torontal.....	Nov. 19-Dec. 16.....	17	2	
Bahrein Island.....	Nov. 27-Dec. 30.....		260	In the Persian Gulf.
Bulgaria:				
Burgas.....	Nov. 22-23.....	2	2	
Varna.....	Nov. 6.....	1		
China:				
Hongkong.....	Jan. 14-20.....	1	1	July 2-15: Cases, 5; deaths, 4, p. 1300, Vol. XXVI.
Dutch East Indies.....				
Batavia.....	Nov. 12-Dec. 23.....	21	8	Total year 1911: Cases, 3,624; deaths, 2,919, including report, p. 2092, vol. 1. Free Dec. 31. Year 1911: Deaths, 323,237.
India.....				
Bassein.....	Jan. 14-Mar. 30.....	128	104	
Bombay.....	Apr. 7-13.....	9	8	
Calcutta.....	Nov. 5-Mar. 23.....		1,001	
Madras.....	Nov. 26-Apr. 6.....	549	447	Madras Presidency Nov. 1-Dec. 31: Cases, 10,436; deaths, 6,545. Jan. 1-Feb. 29: Cases, 18,267; deaths, 11,563.
Moulmine.....	Feb. 18-Mar. 30.....	13	11	
Negapatam.....	Jan. 14-Feb. 24.....		79	
Pondicherry.....	Feb. 22-28.....	4	4	
Rangoon.....	Oct. 1-Feb. 29.....	104	86	
Indo-China:				
Salgon.....	Nov. 20-Mar. 18.....	1,544	1,082	
Italy.....				
Caltanissetta.....	Nov. 26-Dec. 31.....	9	7	
Girgenti.....	do.....	105	57	
Messina.....	Nov. 26-Dec. 2.....	3	2	
Syracuse.....	Nov. 26-Dec. 23.....	15	9	
Malta.....	Nov. 19-Dec. 10.....	6	6	Dec. 23 declared free from cholera.
Montenegro.....	Nov. 4-11.....	9	5	
Persia:				
Adaban.....	Nov. 4.....	1	1	
Kermansbah.....	Dec. 18-26.....		37	

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received from Dec. 30, 1911, to May 24, 1912.

CHOLERA—Continued.

Places.	Date.	Cases.	Deaths.	Remarks.
Philippine Islands.....				Third quarter, 1911: Manila, 1, fatal case; Provinces, 27 cases and 22 deaths. Fourth quarter, 1911: Manila, no case and no death; Provinces, 22 cases, 20 deaths.
Province— Union.....	Oct. 29-Dec 4.....	5	5	
Roumania.....				Total Sept. 9-Dec. 13: Cases, 192; deaths, 42, including report, p. 2094, vol. 1. Free Dec. 19.
Districts— Bralla.....	Sept. 11-Dec. 13.....	84	11	Including cases previously reported.
Convoluri.....	Oct. 31-Nov. 28.....	21	1	
Dolju.....	Nov. 6-Dec. 13.....	19	4	
Jalonitza.....	Oct. 31-Nov. 28.....	4		
Konstanza.....	Oct. 30-Nov. 28.....	8		
Prahova.....	Nov. 6-23.....	1	1	
Talomita.....do.....	2		
Tulcea.....	Nov. 24-Dec. 13.....	15	1	
Servia.....				Total year 1911: Cases, 95; deaths, 51, including report, p. 2096, vol. 1.
Belgrade district.....	Nov. 26-Dec. 16.....	6	4	Declared free Dec. 31.
Siam: Bangkok.....	Nov. 5-Mar. 24.....		1,244	
Straits Settlements: Singapore.....	Nov. 5-Feb. 3.....	4	4	
Tripoli: Tripoli.....	Oct. 13-Jan. 24.....			Cases, 2,000; deaths, from 1,000 to 1,200.
Tunis Regency.....				Total Nov. 25-Jan. 4: Cases, 462; deaths, 323. No cases since Jan. 10.
Beja district.....	Nov. 25-Dec. 21.....	71	20	
Bizerta district.....	Nov. 25-Dec. 5.....	9	15	
Turkey in Asia.....				Provinces in Asia and Europe, Apr. 16-Dec. 30, 1911; Deaths, 6,111, excluding Constantinople. Mainly among troops. Jan. 6-Feb. 27: Cases, 101; deaths, 126.
Acre.....	Jan. 21.....		33	In vicinity.
Adana.....	Dec. 2-Apr. 8.....	46	14	
Aleppo.....	Jan. 26-Apr. 20.....	46	29	
Amara.....	Oct. 15.....	1	1	
Basra.....	Oct. 22-28.....	14	10	
Erzeroum, vilayet.....	Sept. 11-16.....	50	28	
Erzeroum.....do.....	11	8	
Kaifa.....	Dec. 8.....			Present.
Kerbelah.....	Oct. 20-28.....	10	10	
Kharput.....	Nov. 19-Dec. 30.....	47	47	
Jiddah.....	Dec. 2-24.....	323	310	
Mekka.....	Dec. 4-24.....	905	879	Sept. 1-Dec. 24: Cases, 1,648; deaths, 1,565.
Mersina.....	Dec. 1-7.....	2	1	
Osmania.....	Dec. 1-6.....	2	4	
Sinope.....	Dec. 7.....	2	1	
Tor.....	Dec. 14-26.....	29		
Trebizond and vicinity.....	Sept. 18-23.....	64	34	
Tripoli.....	Jan. 4.....			Present.
Turkey in Europe: Constantinople.....	Oct. 24-Feb. 3.....	8	2	
Durazzo.....	Dec. 7-13.....	2		
Janina.....	Jan. 14-22.....	17	7	
Loros.....	Jan. 22.....	12	7	
Saloniki, vilayet.....	Nov. 6-19.....	4	3	In Serres.

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received from Dec. 30, 1911, to May 24, 1912.

YELLOW FEVER.

Places.	Date.	Cases.	Deaths.	Remarks.
Brazil:				
Bahia.....	Mar. 23-25.....	6	1	
Ceara.....	Jan. 1-Feb. 29.....		5	
Manaos.....	Nov. 19-Apr. 27.....		66	
Para.....	Mar. 3-9.....	2	2	Dec. 9-16: 1 fatal case.
Pernambuco.....	Jan. 1-Apr. 15.....		29	Apr. 2: Epidemic.
Rio de Janeiro.....	Mar. 17-Apr. 6.....	3	2	
Canal Zone:				
Culebra Island quarantine.....	Jan. 1-31.....	1		From a vessel from Guayaquil.
Chile:				
Tocopilla.....	Apr. 11.....	90	25	And vicinity. Apr. 20: Still present.
Ecuador:				
Bucay.....	Nov. 16-Feb. 29.....	7	2	
Duran.....	Dec. 1-Apr. 15.....	14	6	
Guayaquil.....	Nov. 16-Apr. 15.....	147	66	
Huigra.....	Feb. 1-29.....	1		
Milagro.....	Feb. 1-Apr. 15.....	16	7	
Naranjito.....	do.....	6	4	
Yaguachi.....	Feb. 1-29.....	1		
Mexico:				
Espita.....	Dec. 31-Jan. 6.....	1		
Kambul, hacienda.....	Feb. 21-27.....		7	
Maxcanu.....	Dec. 31-Jan. 6.....	1		
Merida.....	Nov. 12-Mar. 23.....	20	9	Total Aug. 1, 1911-May 2, 1912: Cases, 66; deaths, 30.
Puerto Mexico (Coatzacoalcos). Salina Cruz.....	Feb. 28..... Feb. 4-7.....		1	7 cases in the lazaretto from s. s. Ikalis from Guayaquil.
San Juan Bautista.....	May 11.....	1		
Temax.....	Dec. 31-Jan. 6.....	1		
Portuguese Guinea:				
Bolama.....	Dec. 19-25.....	1	1	In an engineer on a vessel.
Venezuela:				
Caracas.....	Nov. 16-Feb. 15.....	30		Dec. 1-30, 10 deaths, and Jan. 15-Feb. 29, 8 deaths, including previous reports.
La Guaira.....	Mar. 17-Apr. 1.....	2	2	
Maacuto.....	Mar. 16-19.....	1	1	
Maiquetia.....	Feb. 24-Apr. 30.....	4	4	A suburb of La Guaira.
Sabana Grande.....	Dec. 12.....			Epidemic.
West Indies:				
Barbados— Bridgetown.....	Apr. 27.....	1	1	From steamship Francis, from Para.
St. Vincent.....	Feb. 19.....	1		
At sea.....	Dec. 17-23.....	1	1	On a vessel en route from Manaos to Para.

PLAGUE.

Algeria:				
Philippeville.....	Oct. 19-Nov. 11.....	8	2	Including 5 cases, p. 2096, Vol XXVI.
Arabia:				
Aden.....	Mar. 5-25.....	2	1	
Azores:				
Fayal.....	Jan. 10.....			Still present.
Teceira.....	do.....			Do.
Brazil:				
Bahia.....	Sept. 1-30.....		2	
Para.....	Dec. 24-Apr. 20.....	24	15	
Pernambuco.....	Oct. 1-Feb. 29.....		9	
Rio de Janeiro.....	Nov. 12-Feb. 10.....	7	3	
British East Africa:				
Kismayu.....	Oct. 15-25.....	2		1 case pneumonic.
Chile:				
Iquique.....	Nov. 12-Apr. 6.....	24	11	
Pisagua.....	Nov. 1-30.....	8		
China:				
Amoy.....	Jan. 13.....		1	
Chaochowfu.....	Mar. 10-Apr. 13.....			Present.
Hongkong.....	Dec. 9-Apr. 13.....	178	157	

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received from Dec. 30, 1911, to May 24, 1912.

PLAGUE—Continued.

Places.	Date.	Cases.	Deaths.	Remarks.
Dutch East Indies:				
Java.....				Total Mar. 1—Dec. 30: Cases, 1,817; deaths, 1,324. Dec. 31—Mar. 2: Cases, 99; deaths, 95.
Paseroean Residency, Malang District.	Nov. 12—Apr. 6....	147	100	
Soerabaya.....	Oct. 17—27.....	2		
Ecuador:				
Duran.....	Feb. 1—29.....	1		
Guayaquil.....	Nov. 16—Feb. 29..	124	52	Dec. 16—Jan. 31: Reports not available because of revolution. Total Jan. 1—Dec. 31, 1911: Cases, 1,656; deaths, 1,041, including cases previously reported.
Egypt:				
Cairo.....	Apr. 22—25.....	2	1	
Alexandria.....	Mar. 22.....	1	1	
Provinces—				
Assiout.....	Jan. 1—Apr. 24....	62	40	Sept. 11—16: Cases, 50; deaths, 28.
Assouan.....	do.....	31	19	
Behera.....	Jan. 1—Apr. 15....	5	3	Sept. 11—16: Cases, 11; deaths, 8.
Beni Souef.....	Feb. 16—Apr. 20..	30	9	
Fayoum.....	Jan. 1—Apr. 24....	3	1	
Galloubeh.....	Jan. 1—Apr. 23....	5	3	Oct. 5—Dec. 26: Case, 1.
Garbieh.....	Jan. 1—Apr. 24....	24	8	
Carchieh.....	Apr. 21—22.....	6	4	
Girgeh.....	Mar. 28.....	1	1	
Kena.....	Jan. 1—Apr. 25....	96	73	Nov. 20—Dec. 13: Cases, 3; deaths, 3.
Menouf.....	Feb. 2—Apr. 23....	9	2	
Minieh.....	Jan. 1—Apr. 23....	26	6	Dec. 13: Case, 1.
German East Africa:				
Dar-es-Salaam.....	Nov. 13—15.....	1	1	From the interior via Bergamogo
Great Britain:				
Liverpool.....	Apr. 27—30.....	1	1	In the Royal Southern Hospital from s. s. Italian Prince, which arrived at Liverpool Apr. 19, laden with fruit and grain from Mediterranean ports.
Hawaii:				
Honakaa.....	Feb. 9—Mar. 18....	4	4	
India:				
Bombay.....	Nov. 19—Apr. 13....	851	744	
Calcutta.....	Nov. 11—Mar. 23....		329	
Karachi.....	Nov. 26—Apr. 13....	781	670	Total year 1911: Cases, 3,273; deaths, 3,046.
Madras.....	Jan. 1—6.....	1	1	
Rangoon.....	Oct. 1—Feb. 29....	166	158	
Bombay Presidency and Sind.....	Oct. 29—Mar. 23....	58,396	42,824	
Madras Presidency.....	do.....	10,129	7,869	
Bengal.....	do.....	37,491	31,764	
United Provinces.....	do.....	86,612	76,987	
Punjab.....	do.....	10,541	8,082	
Burma.....	do.....	1,428	1,302	
Eastern Bengal and Assam.	Jan. 1—Feb. 24....	2	2	
Central Provinces.....	Oct. 29—Mar. 23....	26,600	21,144	
Coorg.....	do.....	88	52	
Mysore State.....	do.....	8,631	6,688	
Hyderabad State.....	do.....	25,820	23,394	
Central India.....	do.....	9,096	7,453	
Rajputana and Ajmere Merwara.....	do.....	1,533	1,234	
Kashmir.....	Feb. 3—Mar. 23....	114	51	
North West Province.....	Oct. 29—Feb. 24....	2	2	Total for India, Oct. 29—Mar. 23: Cases, 276,483; deaths, 228,848. Total year 1911: Cases, 828,535; deaths, 691,849.
Indo-China:				
Saigon.....	Nov. 13—Mar. 4....	33	5	
Japan:				
Formosa.....	Mar. 7—30.....	52	40	
Nagasaki.....	Apr. 20.....	1		On the s. s. Tacoma Maru, from Hongkong and Shanghai.
Mauritius:				
.....	Nov. 3—Mar. 14....	92	54	
Persia:				
Bushire.....	Feb. 4—Mar. 30....	134	83	

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received from Dec. 30, 1911, to May 24, 1912.

PLAGUE—Continued.

Places.	Date.	Cases.	Deaths.	Remarks.
Peru:				
Departments—				
Callao.....	Oct. 1-21.....	1		City, in November, 1 case; in January, 3 cases with 2 deaths; Mar. 1-26, 12 cases.
Chiclayo.....	do.....	12	4	
Chosica.....	do.....	1	1	
Lambayeque.....	do.....	3		
Libertad.....	do.....	8		Apr. 10, 22 cases in the lazaretto at Trujillo. Apr. 22 still present.
Lima.....	do.....	13	6	
Philippine Islands:				
Cebu quarantine station.....	Dec. 4.....	1		On s. s. Montrose from Shanghai. Apr. 6, a fatal case of pneumonic form in a member of the crew on s. s. Zafiro, arrived Apr. 4 from Hongkong.
Manila.....				
Russian Empire:				
Astrakhan, government.....	Sept. 21-Jan. 7....	201	180	Including 73 cases and 63 deaths reported on p. 2098, Vol. I.
Siam:				
Bangkok.....	Nov. 4-Mar. 23.....		5	
South Africa:				
Durban.....				Total: Jan. 14-Apr. 19, cases 27, deaths 22.
Straits Settlements:				
Singapore.....	Nov. 5-Mar. 16....	29	25	
Turkey in Asia:				
Basra.....	Feb. 13.....	1	1	A stoker on s. s. Nicomedia from Bushire.
Jiddah.....	Jan. 13-Mar. 27....	16	9	
Venezuela:				
Caracas.....	Mar. 12-Apr. 22....	6	2	
West Indies:				
Trinidad.....	Apr. 2-May 1.....	6	4	
At sea.....	Mar. 1-11.....	1	1	On s. s. Macedonia from Bombay to Aden.
Do.....	Apr. 1.....	1	1	Pneumonic on s. s. Loongsang, en route from Hongkong to Manila.

SMALLPOX.

Algeria:				
Algiers.....	Nov. 1-30.....		1	
Oran.....	Jan. 1-31.....	2	1	
Arabia:				
Aden.....	Nov. 28-Apr. 8....	22	10	And vicinity.
Argentina:				
Buenos Aires.....	Jan. 1-31.....		2	Oct. 1-31, 6 deaths. No deaths in November or December, 1911.
Rosario.....	Oct. 1-Jan. 31.....		40	
Australia:				
Thursday Island.....	Jan. 2.....	1		From s. s. Taiyuan.
Austria-Hungary:				
Bohemia.....	Jan. 14-20.....	2		
Budapest.....	Jan. 4-10.....	25		
Galicja.....	Dec. 24-Apr. 13....	30	1	
Krain.....	Jan. 14-20.....	7		
Trieste.....	Dec. 3-Apr. 6.....	2		
Tyrol.....	Jan. 14-Mar. 9....	3		From s. s. Baron Call from Beirut.
Vienna.....	Mar. 25-30.....	1		
Brazil:				
Bahia.....	July 1-31.....		1	
Para.....	Mar. 24-Apr. 20....	6	4	Case Mar. 30 from Alagoas.
Pernambuco.....	Oct. 1-Apr. 15....		838	
Rio de Janeiro.....	Nov. 26-Mar. 6....	31	1	
Santos.....	Dec. 12-23.....		1	
Canada:				
British Columbia—				
Fernie.....	Feb. 26-Mar. 16....	5		
Nelson.....	Dec. 24-30.....	1		
Vancouver.....	Apr. 14-20.....	1		
Victoria.....	Feb. 4-10.....	1		
Manitoba—				
Winnipeg.....	Jan. 14-Apr. 20....	4		

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received from Dec. 30, 1911, to May 24, 1912.

SMALLPOX—Continued.

Places.	Date.	Cases.	Deaths.	Remarks.
Canada—Continued.				
New Brunswick— Summerstown	Apr. 12			Epidemic; 10 miles from Cornwall.
Nova Scotia— Halifax	Mar. 24-Apr. 6	2		
Ontario—				
Hamilton	Apr. 14-27	16		
Kingston	Dec. 19-23	1		
Ottawa	Dec. 10-May 11	105	1	
Sarnia	Oct. 17-Mar. 23	43		
Toronto	Jan. 6-Apr. 20	5	1	
Windsor	Feb. 4-Mar. 16	8		
Quebec—				
Montreal	Dec. 17-May 11	32		
Quebec	Dec. 10-May 11	289	2	
Yukon— Dawson	Apr. 1-6	1		
Ceylon: Colombo	Nov. 12-Feb. 10	3		And vicinity
Chile:				
Iquique	Dec. 10-Mar. 16	4	2	
La Serena	Nov. 21-30	14		
Santiago	Nov. 1-30	685	343	
Talcahuano	Nov. 26-Dec. 23	14	3	
Valparaiso	Dec. 3-Apr. 13	70		
China:				
Canton	Nov. 11-Dec. 30	40	6	
Chaochowfu	Mar. 30			Present.
Chenghai	Jan. 29-Apr. 13			Do.
Chungking	Nov. 18-Apr. 6			Do.
Dalny	Mar. 3-Apr. 6	11	2	
Hankow	Jan. 21-Feb. 17	2	1	
Hongkong	Nov. 12-Mar. 30	706	523	
Kityang	Jan. 21-Apr. 13			Do.
Nanking	Dec. 10-Apr. 20			Do.
Shanghai	Dec. 11-Feb. 18	1	6	Deaths among natives.
Swatow	Mar. 2			Present.
Cuba: Habana	Dec. 19-Jan. 19	2		Case Dec. 19 from German s. s. Frankenwald, from Spain and Canary Islands; case Jan. 19 from s. s. Mexico.
Dutch East Indies:				
Java— Batavia	Nov. 12-Apr. 6	55	15	
Egypt: Cairo	Dec. 10-Mar. 25	10	1	
Port Said	Jan. 30-Feb. 4	1		
France:				
Havre	Mar. 10-16		4	
Marseille	Jan. 1-Mar. 31		5	Nov. 1-30, 1 death.
Paris	Dec. 3-Apr. 27	123	2	
Germany				Total, Dec. 31-Apr. 27; cases, 129.
Hamburg	Jan. 21-Apr. 27	7		
Gibraltar	Feb. 27-Mar. 3	1		
Great Britain:				
Bristol	Jan. 29-Feb. 3	2		
Liverpool	Mar. 17-Apr. 17	2	1	Case Apr. 13, an American from the s. s. Arabic.
London	Jan. 14-Apr. 20	10	1	
Southampton	Mar. 3-9	1		
West Hartlepool	Feb. 18-Mar. 9	2		
India:				
Bombay	Nov. 19-Apr. 13	908	398	
Calcutta	Nov. 19-Mar. 23		31	
Karachi	Apr. 1-6	1	1	
Madras	Nov. 26-Apr. 13	195	77	
Rangoon	Oct. 1-Feb. 29	220	55	
Indo-China:				
Saigon	Nov. 13-Mar. 18	38	7	
Italy:				
Genoa	Dec. 1-Apr. 15	47	2	
Leghorn	Dec. 16-May 4	110	1	
Messina	Nov. 19-Jan. 31		6	
Naples	Dec. 3-Apr. 27	103	1	
Palermo	Nov. 26-Apr. 27	2,643	898	
Rome	Jan. 1-Mar. 31	31	3	
Turin	Jan. 15-Apr. 7	2		

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received from Dec. 30, 1911, to May 24, 1912.

SMALLPOX—Continued.

Places.	Date.	Cases.	Deaths.	Remarks.
Japan:				
Arima-Mura	Nov. 12-18	6	1	11 miles east from Kobe.
Formosa	Mar. 3-16	3		
Kanagawa, ken.	Dec. 17-23	1		
Kobe	Jan. 22-28	2	1	Jan. 20, 1 case from s. s. Suverio from Hongkong; Jan. 28, 1 case from Shingo Maru.
Nagasaki	Feb. 12-18	1		
Nogahama	Mar. 17-23	1		On s. s. Tenyo Maru from Hongkong.
Yokohama	Jan. 22	1		From s. s. Hydra from New York via Suez.
Malta	Dec. 24-Jan. 6	2	1	
Mexico:				
Aguascalientes	Dec. 18-Mar. 3		7	
Chihuahua	Nov. 20-Feb. 11	92	36	
Coahuila, State	Oct. 1-30		16	
Guadalajara	Jan. 14-Apr. 13	8	4	
Juarez	Dec. 19-May 11	17	5	
Magdalena	Dec. 23-Mar. 12	91	50	Mar. 12, 10 cases present.
Manzanillo	Feb. 18-24	1		
Mazatlan	Dec. 11-Apr. 30		13	Mar. 16, 25 cases in the lazaretto.
Mexico	Nov. 26-Mar. 23	224	101	
Monterey	Dec. 11-24		2	
Porfirio Diaz	Dec. 3-Mar. 23		35	
Salina Cruz	Feb. 11-Mar. 9	4	2	Mar. 23, present in vicinity.
San Antonio	Jan. 1-21	12	9	
San Carlos	do			Present.
Sandoval	Dec. 16			Do.
San Ignacio	Jan. 8	3		
Saric	Jan. 21-27		6	
Santa Ana	Jan. 8	4		
San Luis Potosi	Nov. 12-Mar. 9	8	5	
Tampico	Dec. 1-Apr. 10		16	
Tapachula	Nov. 1-Jan. 31		18	
Philippine Islands				Third quarter, 1911: Manila, 9 cases; no deaths. Fourth quarter, 1911: 38 cases.
Manila (on arriving vessel)				Apr. 1 to 6, present on the steamers Serantes and Sotolongo; Apr. 2-9, 8 cases among the crew of the United States Army transport Warren, from ports in China and Japan.
Portugal:				
Lisbon	Dec. 9-Apr. 20	56		
Roumania	Jan. 1-31	2,935	143	Total: Jan. 1-Feb. 29, cases, 5,847; deaths, 247.
Russia:				
Batun	Dec. 1-Mar. 31	2		
Libau	Dec. 17-Apr. 28	3		
Moscow	Nov. 19-Apr. 13	72	9	
Odessa	Nov. 26-Apr. 20	35	1	
Reval	Nov. 1-30	1		
Riga	Dec. 24-Apr. 16	62		
St. Petersburg	Nov. 19-Apr. 20	211	48	Oct. 1-Jan. 31: Deaths, 10.
Warsaw	Nov. 5-Mar. 16	414	200	
Siam:				
Bangkok	Nov. 5-Mar. 23		2,175	
Siberia:				
Omsk	Jan. 1-31	7		
South Africa:				
Durban	Jan. 21-Apr. 6	5		
Johannesburg	Jan. 7-Feb. 10	36		
Spain:				
Barcelona	Feb. 6-Apr. 1		2	
Cadiz	Nov. 1-Mar. 31		29	
Madrid	Dec. 1-Mar. 31		16	
Malaga	Nov. 1-30		45	
Seville	Dec. 1-Mar. 31		10	
Valencia	Dec. 3-Apr. 20	398	16	
Straits Settlements:				
Penang	Feb. 11-17	1		
Singapore	Nov. 19-Apr. 6	38	15	

CHOLERA, YELLOW FEVER, PLAGUE, AND SMALLPOX—Continued.

Reports Received from Dec. 30, 1911, to May 24, 1912.

SMALLPOX—Continued.

Places.	Date.	Cases.	Deaths.	Remarks.
Switzerland:				
Cantons—				
Aargau.....	Apr. 7-13.....	1		
Oberwalden.....	Jan. 14-20.....	1		
St. Gall.....	Mar. 30-Apr. 6.....	1		
Zurich.....	Dec. 3-23.....	6		
Teneriffe:				
Santa Cruz.....	Dec. 3-Apr. 13.....		54	
Turkey in Asia:				
Beirut.....	do.....	1,500	107	
Turkey in Europe:				
Constantinople.....	Dec. 4-Apr. 28.....		199	
Uruguay:				
Montevideo.....	Sept. 1-Dec. 31.....	25	4	
Venezuela:				
Caracas.....	Nov. 1-Jan. 15.....	11	2	
Zanzibar:				
Zanzibar.....	Oct. 28-Dec. 15.....	3	2	

MORTALITY.

WEEKLY MORTALITY TABLE, FOREIGN AND INSULAR CITIES.

Cities.	Week ended—	Estimated population.	Total deaths from all causes.	Deaths from—											
				Tuberculosis.	Plague.	Cholera.	Yellow fever.	Smallpox.	Typhus fever.	Typhoid fever.	Scarlet fever.	Diphtheria.	Measles.	Whooping cough.	
Aberdeen.....	Apr. 20	163,084	63											1	
Do.....	Apr. 27		64								1	1			
Aguascalientes.....	May 12	40,000	82	3					5		1	1	16		
Aix-la-Chapelle.....	Apr. 13	157,695	46	7								1			2
Do.....	Apr. 20	157,702	42	3							1				4
Amsterdam.....	May 4	566,131	148	26								2	5	1	1
Antwerp.....	Apr. 20	316,604	89	8							2		11	1	1
Do.....	Apr. 27		79	14						1	1	3	6		1
Athens.....	May 4	250,010	76	14								1			1
Barcelona.....	May 3	591,272	154	21				1		1	1	2			
Barmen.....	Apr. 20	171,000	32	4								2			1
Barranquilla.....	May 4	60,000	13	2						3					
Batavia.....	Apr. 6	217,630	8					1	1						
Beirut.....	Apr. 27	80,000	20							5					
Belfast.....	do.....	391,051	135	25								2			4
Do.....	May 4		134	24							1				8
Belgrade.....	Apr. 28	90,050	31								2			1	
Berlin.....	Apr. 20	2,067,928	596	87							10	14	10	7	7
Do.....	Apr. 27		597	92							5	11	10	7	
Birmingham.....	May 4	842,512	216								2	5	1	4	5
Bombay.....	Apr. 20	979,445	918	48	144	3		53					12		
Bremen.....	do.....	246,850	81	13							2	5	2		
Birmingham.....	Apr. 27	842,512	217								1	1	4		12
Bremen.....	do.....	246,850	80	11								2	4		3
Brunswick.....	Apr. 13-20.....	145,000		2							5	9			
Do.....	Apr. 27-May 4.....		5								10	17			
Brussels.....	May 4	739,684	229	21										2	1
Budapest.....	Apr. 13	1,000,000									5				
Do.....	Apr. 20										2	3			2
Cairo.....	Apr. 15	689,439	412	25				1	6	3		3	3	3	
Do.....	Apr. 22		433	23		1			9	4		4	3	3	
Calcutta.....	Mar. 30	890,493	598	34	133	89		6						1	
Catania.....	May 10	207,000	79	3							2		3	3	
Chemnitz.....	Apr. 20	300,019	83	6								2	1		3
Cienfuegos.....	do.....	33,578	19	1						1					

MORTALITY—Continued.

Weekly mortality table, foreign and insular cities—Continued.

Cities.	Week ended—	Estimated population.	Total deaths from all causes.	Deaths from—										
				Tuberculosis.	Plague.	Cholera.	Yellow fever.	Smallpox.	Typhus fever.	Typhoid fever.	Scarlet fever.	Diphtheria.	Measles.	Whooping cough.
Coburg	Apr. 20	24, 174	12	3								2		
do	do	527, 663	129	23										
Do	Apr. 27	528, 738	149	28										
Cologne	Apr. 6	227, 026	105	10								2		
Do	Apr. 13		116	14										
Constantinople	May 5	1, 000, 000	283	32				8		5	7			
Dresden	Apr. 20	555, 300	128	17						6	3	3		12
Dublin	do	406, 536	217	39						1	1	2	4	13
Do	Apr. 27		192	21						3	1	1	1	16
Do	May 4		208	38							4	4	15	4
Dundee	do	171, 006	55	5										1
Edinburgh	do	321, 200	94	8										3
Erfurt	Apr. 27	126, 560	32	1								1		1
Frankfort-on-Main	do	425, 000	88									3		1
Georgetown	May 4	57, 577	56	4						2				
Glasgow	May 10	785, 600	257								1		14	3
Gothenburg	May 4	170, 000	53	11								1		1
Guadalajara	May 11	119, 468	127					1			1			
Hamburg	May 4	953, 079	275	32					1		2			4
Havre	do	136, 159	55	12						1	6			
Hongkong	Apr. 6	336, 488			29				24		2			
Do	Apr. 13				36						4			
Hull	May 4	282, 987							21		4			
Karschi	Apr. 20	157, 290	162		109					1				
Leeds	May 4	445, 568	119	16						1	1	1		
Leipzig	Apr. 27	605, 755	154	26							1			
Liege	do		43	4							1			
Leith	May 4	81, 000	24	1									1	3
London	do	7, 340, 119	1, 734	4						2	4	7	28	56
Lubeck	May 4	100, 000	24											
Madras	Apr. 20	518, 660	334			4		4		1	1		3	
Do	Mar. 9													
Magdeburg	Apr. 13	285, 557	617	55						3		11	1	
Manaos	Apr. 20	52, 000	46	8				5						
Do	Apr. 27		37	1				4						
Matamoros	May 12	15, 000	10	3										
Montreal	May 18	466, 197	172	27					1		2	2		
Moscow	Apr. 20	1, 500, 000	886	93					1	1	4	13	4	5
Nantes	Apr. 21	170, 535	74	21						1	4		10	9
Do	Apr. 28		72	10										
Newcastle-on-Tyne	May 4	266, 671	73	8						1		1		
Nottingham	Apr. 24	260, 000	81	4									8	1
Ottawa	May 11	90, 000	38	1										
Palermo	May 4	340, 000	131	5				3						
Para	Apr. 27	185, 000	100	9			1			3	1			
Do	May 4	185, 000	87	12										
Penang	Apr. 13	102, 167	74	18								1		
Port Said	Apr. 22	52, 811	23										1	
Port of Spain	Apr. 27	60, 000	31	7						1		1	1	
Prague	Apr. 20	225, 817	91	15						1	1	1	4	2
Do	Apr. 27		102	27						1	4	1	4	
Rangoon	Apr. 6	293, 316	177					19						
Rotterdam	May 4	438, 774	106		8	4						2		
St. Petersburg	Apr. 27	1, 962, 400	946	156						1	2			11
San Luis Potosi	Mar. 16	82, 946	48	4						13	7	8	24	
Saigon	Mar. 25	220, 000	18		2	15				1				
Do	Mar. 31		49		2	45				1				
Do	Apr. 9		48		2	45								
Singapore	Apr. 6	303, 328	196	22										
South Shields	May 4	109, 676	26									1		
Stettin	Apr. 27	240, 000	68	2								2	1	1
Stoke-on-Trent	May 4	237, 153	70	2									1	1
Toronto	May 4-11	392, 000	244	11							4	12	1	4
Trieste	Apr. 27	235, 999	98							1				
Tripoli	Apr. 21	50, 000	39	2						3				
Do	Apr. 28		67	4						9				
Turin	May 5	430, 770	147	21									4	
Valencia	May 4	235, 000	94	6					1					
Vienna	Apr. 20	2, 081, 335	732	126							3	10	9	4

MORTALITY—FOREIGN AND INSULAR COUNTRIES AND CITIES
(Un tabulated).

FORMOSA.—Four weeks ended April 13, 1912. Population 3,341,217. The deaths include diphtheria 2, plague 56, typhoid fever 11.

FRANCE—Calais.—Month of April, 1912. Population 80,000. Total number of deaths from all causes 121, including diphtheria 1, tuberculosis 24, typhoid fever 1.

Roubaix.—Month of April, 1912. Population 122,723. Total number of deaths from all causes 153, including diphtheria 1, tuberculosis 33.

GREAT BRITAIN.—Week ended April 27, 1912.

England and Wales.—The deaths registered in 95 great towns correspond to an annual rate of 14.1 per 1,000 of the population which is estimated at 17,639,816.

Ireland.—The deaths registered in 21 principal town districts correspond to an annual rate of 20.3 per 1,000 of the population which is estimated at 1,157,014. The lowest rate was recorded at Armagh viz, 6.9, and the highest at Lurgan, viz, 30.1 per 1,000 of the population.

Scotland.—The deaths registered in 18 principal towns correspond to an annual rate of 16.2 per 1,000 of the population, which is estimated at 2,182,400. The lowest rate was recorded at Govan, viz, 8.1, and the highest at Kilmarnock, viz, 22.5 per 1,000 of the population. The total number of deaths reported from all causes was 679, including diphtheria 4, measles 32, scarlet fever 5.

ITALY—Catania.—Month of April, 1912. Population 127,000. Total number of deaths from all causes 159, including scarlet fever 11, tuberculosis 12, typhoid fever 1.

Milan.—Month of April, 1912. Population, 602,236. Total number of deaths from all causes 149, including diphtheria 1, measles 3, scarlet fever 1, tuberculosis 132, typhoid fever 6.

PORTUGUESE EAST AFRICA—Lourenço Marquez.—Month of January, 1912. Population 10,000. Total number of deaths from all causes 31, including tuberculosis 4.

Month of February, 1912. Total number of deaths from all causes 27, including tuberculosis 8.

SPAIN—Almeria.—Month of April, 1912. Population 50,910. Total number of deaths from all causes 114, including smallpox 1, tuberculosis 9, typhus fever 3.

Madrid.—Month of April, 1912. Population 584,117. Total number of deaths from all causes 1,143, including diphtheria 14, measles 48, scarlet fever 1, tuberculosis 118, typhoid fever 12.

Seville.—Month of April, 1912. Population 158,235. Total number of deaths from all causes 346, including diphtheria 7, measles 7, smallpox 2, tuberculosis 48, typhoid fever 5.

By authority of the Secretary of the Treasury:

RUPERT BLUE,
Surgeon General.

United States Public Health and Marine-Hospital Service.

ADDITIONAL COPIES of this publication
may be procured from the SUPERINTEND-
ENT OF DOCUMENTS, Government Printing
Office, Washington, D. C., at 5 cents per copy.

Subscription price, per volume - - - - \$2

