PUBLIC HEALTH REPORTS

VOL. 35

JANUARY 23, 1920

No. 4

PREVENTION OF STREAM POLLUTION BY DYE AND INTERMEDIATE WASTES.

By E. J. CASSELMAN, Assistant Chemist, United States Public Health Service.

Liquid wastes from the manufacture of dyes and intermediates are particularly objectionable when discharged into rivers, because of their properties of high color, bad taste, or strong odor, even in very low concentrations. When the water is to be used for drinking, these qualities are eradicated only by a comparatively great dilution; that is, the river must be large compared to the discharge from the chemical factory.

The investigation which is the basis of the report and recommendations here made was undertaken for the purpose of studying the methods of disposing of dye and intermediate wastes. Most of the work was performed at the plant of the Chemical Co. of America, with special attention given to the pollution of the Rahway River 1 by the wastes from that plant.

The plant is located at Springfield, N. J., between two forks of This river forms the only natural avenue of escape for the waste liquids. The nearest connection to a sewer is over a mile away, and the expense of building to the sewer is considered prohibitive. Other methods of disposing of the waste, except by treatment, such as trucking or shipping to tidewater, are also highly expensive. The cost of evaporation, the most obvious means of purification, depends upon the method used. If it were permissible to allow spontaneous evaporation of the waste by trickling over ricks, this method could no doubt be carried on very cheaply. But such evaporation would require a large installation for the amount of waste produced here, and, furthermore, the system would not be dependable during damp, quiet weather nor in the winter. A multiple-effect method of evaporation with recovery and discharge of distillate is hardly permissible, as the odorous constituents of the wastes are usually volatile enough to distill over with the water. It is assumed, then, that a method of evaporation would be devised

156563°—20——1 (167)

 $^{^{\}rm 1}$ For the legal aspect of the case see Public Health Reports, Oct. 17, 1919, p. 2303.

for burning coal to evaporate the waste in a single effect or stage. The cost of this would be about \$0.0006 per pound of water, assuming coal costs at \$7 per net ton and that 1 pound of coal evaporates 6.5 pounds of water.

Any chemical method of waste disposal, then, must be cheaper than \$0.0006 per pound of water in order to compete with the evaporation process.

In this investigation it was hoped that treatments cound be found for producing liquors that were not objectionable of themselves, so that they could be discharged into any river, regardless of its quantity of flow. The wastes that could not be treated to give such an effluent were studied with regard to pollution of the Rahway River at the plant.

Stream gaugings made at this point after an exceptionally dry period indicated that the minimum flow is about 5,000,000 gallons per day.

This work was carried out under the supervision of Prof. Earle B. Phelps, formerly of the Hygienic Laboratory, United States Public Health Service, whose suggestions were most helpful.

I. WASTES PRODUCED AND THEIR PROPERTIES.

Ortho-toluidine waste.—This waste results from the manufacture of ortho-toluidine. Ortho-toluidine is made from ortho-nitro-toluene by reduction with iron chips and sulphuric or hydrochloric acid. The product of the reaction is then steam distilled for purification after making alkaline. The distillation gives rise to two liquid wastes—a still residue and an aqueous layer in the receiver. The latter is a solution containing 0.015 per cent 1 toluidine and is not an objectionable waste. The still residue, however, is a very disagreeable waste.

The properties of this waste are a very dark color, a highly disagreeable odor, and a composition which may conceivably consist of every high boiling intermediate reduction product of nitrotoluol. In addition, there are a slight acidity and high concentration of ferrous salt.

The quantity of this waste is, roughly, four-tenths of a gallon per pound of toluidine.

Nitrobenzene waste.—In the manufacture of nitrobenzene, benzene is nitrated with mixed sulphuric and nitric acids in a nitration kettle. The products are two liquid layers, one of which is spent acid and the other crude nitrobenzene. The spent acid is not considered a waste, as it is still strong enough to use for other purposes.

The nitrobenzene is next washed with soda ash solution to remove acidity, and two layers are again formed, the nitrobenzene and the wash solution. The nitrobenzene layer is distilled in vacuum stills

and there is no waste in connection with this procedure. The wash water is not used again and is the one waste of the process.

This wash water is naturally alkaline, from the excess of soda ash. It is a high orange yellow in color, and it smells strongly of nitrobenzene. The quantity is about one-sixth of a gallon per pound of nitrobenzene.

Monoethyl aniline wastes.—In the manufacture of monoethyl aniline from aniline and ethyl alcohol, these two substances are heated together with a catalyzer under certain conditions of temperature and pressure, and the monoethyl aniline is produced as a salt. It is then thrown out of combination with soda ash and distilled with steam, from which process two wastes are produced—a still residue and an aqueous layer in the receiver. This latter is salted in order to separate more monoethyl aniline from the water.

The still residue is of a medium brown color that of itself is not very objectionable. The odor, however, is very strong and disagreeable. In composition it is alkaline and may contain some unchanged alcohol and aniline, also monoethyl aniline and diethyl aniline, as well as side products like orthoethyl aniline, paraethyl aniline, orthoethyl monoethyl aniline, paraethyl monoethyl aniline. Inorganic sodium salts are also present. The quantity of still residue is about 13 gallons per pound of monoethyl aniline.

The receiver residue is little else than a suspension of monoethyl aniline in salt water. At first it is colorless, but on standing it turns to a pinkish shade. Its principal objection is its odor, which is very strong and disagreeable. In quantity it is about 23 gallons per pound of monoethyl aniline.

Dye wastes.—The dyes manufactured at the Chemical Co.'s plant consist of several triphenyl methane dyes and several azo-dyes. The processes generally take place in one solution, from which the final step is the removal of the solid dye by filtration or otherwise. The waste liquor comprises the residual solution.

The properties of these wastes vary considerably with the dye manufactured, but in every case they are highly colored because of failure to remove all the dye from them. In addition they frequently contain odorous compounds and concentrated salt. Their volumes are very considerable, ranging from 10 to 20 gallons per pound of dye.

Lagoon liquor.—For a year or more the wastes given above, as well as some others not studied, were stored in an artificial lagoon, and comprised some four or five million gallons at the time of this study. These wastes had, of course, had ample time for aeration and the evaporation of some of their more volatile constituents. Furthermore, it had been considerably diluted with rain water. The composition of this body of wastes varied to some extent from point to point

as a result of the difference in the wastes discharged at various times. In the main, however, it was a fairly uniform mixture except near the point of discharge from the plant, where dye wastes predominated.

The color of the main body of this liquor was a rather deep red, but there was scarcely any odor. It was 0.005 normal in acid, corresponding to about 0.25 per cent sulphuric acid; and it contained some iron, zinc, and sodium salts in small quantities. The object of treating this waste was to prevent the lagoon from becoming filled up and overflowing.

II. EXPERIMENTAL WORK.

The properties and treatment of the separate wastes were first studied, and then their treatment as mixtures in the proportions in which they were likely to be when discharged from the works, was undertaken.

In making these studies the first experiments were qualitative in nature, no regard being paid to cost of treatment. Quantitative tests were made upon the best results offered from the experiments. Most of these experiments were made in 100 or 200 cc. bottles with 100 or 200 cc. samples. Some of the quantitative tests were made with 2-liter volumes. Filtrations were made with a 10 cm. Büchner funnel and filter paper.

From data gained by the quantitative tests the wastes were treated on a scale of about 40 cubic feet (300 gallons). The apparatus consisted of the following: Two 350-gallon wooden tubs fitted with gauge glasses graduated in cubic feet, one wooden tower 3 by 3 by 4 feet, filled with coke, and another one filled with cinders; two sand filter-beds, each having an area of 1 square yard, and a depth of about 2 feet of sand and gravel; one small filter press; one small pump connected to each of the tubs and piped to discharge into any desirable piece of apparatus; two barrels arranged to empty into the tubs, for the purpose of mixing reagents. (Figs. 1 and 2.)

It is assumed that any results observed on this 300-gallon scale can be duplicated on any large scale without essential change in treatment.

Ortho-toluidine Waste.

Effect of making alkaline.—When ortho-toluidine waste was made alkaline, the iron it contained was thrown down as a sludge of ferrous hydroxide. The supernatant liquid was at first without much color, but it retained the very obnoxious odor. This liquid became very deeply colored on standing. That this was due to oxidation was shown by the fact that the reaction started at the surface, and that it could be more quickly brought about by aeration. The shade was somewhat dependent upon the alkalinity, being more bluish in strong alkali, and more greenish in weak. In acid solutions it was deep red. When nitrobenzene waste instead of lime was used for the precipi-






Fig. 1.—Precipitation tubs and filter press of experimental waste disposal plant, Springfield, N. J.

Fig. 2.—Precipitation tubs and coke tower of experimental waste disposal plant, Springfield, N. J.

tation of the iron, the deep color developed immediately, presumably from oxidation by the nitrobenzene.

The amount of lime required to neutralize and just precipitate all the iron appeared to be about 3 grams per cubic foot, but this would naturally vary with each batch of waste produced. The effect of any excess of lime over this was to make the iron sludge settle more slowly, and to decrease the blue shade of the color of the solution after oxidation.

Other precipitants.—Attempts to carry down color or odor of this alkaline solution with alum, silica, zinc hydrate, and lead sulphate were unsuccessful.

It may be concluded that precipitants are of no avail in treating toluidine waste.

Aeration.—Air was blown through alkaline filtered toluidine waste for two hours, and failed to remove much of the odor or any of the color.

Bromine.—Bromine caused a precipitation of a finely divided solid, leaving a light colored supernatant liquid (after removal of the iron).

Bleaching powder.—Bleaching powder and acid caused the precipitation of a solid tar-like substance which could be filtered off.

Quantitative experiments, made after removal of the iron by lime and after filtration, showed that the best results were obtained by the use of 10 to 30 grams per liter of chlorine, depending upon the strength of the waste. Upon filtering such a solution a light brown filtrate was obtained which turned dark on being made neutral, and still darker on being made alkaline with lime. This color could be extracted by ether in acid solution, but not in alkaline solution.

It was noted that this chlorine requirement was about the same for any particular waste no matter what the condition of its previous aeration might have been.

The color of the chlorinated solution could not be removed by precipitants like ferrous hydroxide; in fact, it became worse if nitrobenzene waste instead of slaked lime had been used for neutralizing in the first place. When enough chlorine to get the best color was used, the color removal seemed to be about 90 per cent in acid solution, 80 per cent in neutral solution, and 75 per cent in alkaline solution.

The odor of this treated waste was not much improved in intensity, but it seemed much less unpleasant.

Absorbent carbons.—Boiling alkaline, filtered toluidine waste with bone black was found to remove both odor and color. The same result was accomplished when the waste was filtered through a layer of bone black one-quarter of an inch thick. Quantitative experiments showed that roughly 1.1 cubic feet could be passed through 1 pound of the bone black before it became exhausted.

A new carbon of several grades has appeared on the market under the trade name of Superfiltchar.¹ The different grades were tried out and found to be effective against toluidine waste. In quantitative tests an amount of the carbon was weighed out into a measured volume of waste (usually 100 cc.) and allowed to stand 18 to 24 hours with occasional shaking. Table I shows the effect of various amounts of these carbons.

TABLE I.—Effect of carbons on alkaline, filtered toluidine wast	TABLE	IEffect	of carbons	on	alkaline,	filtered	toluidine	waste
---	-------	---------	------------	----	-----------	----------	-----------	-------

Per cent.	Substance.	Residual odor.	Time	
1.44 .2 .3	Bone blackSuperfiltchar, oil gradedo	60	Faint	1 50
.35 .40 .7 2.0	dododododo	3 2 Nonedo.	Very faint	20 20 16 16
.46	Filtchar do	40	Slightdo	- 19
.80 .90 .30	do do do	4.6	dodo	18 18
.40 .46 .70	do do	65 52 7.1 6.4	do do Not observed	1
.90 .40 .80	dodododododo	2 11	do	ì

A Bismarck brown standard solution of 100 p. p. m. was made by dissolving 100 mg. of Bismarck brown in 1 liter of distilled water, and filtering.
 By filtration.

If it be assumed that a practically odorless liquid is wanted with a color of not more than five parts per million on the arbitrary scale selected, it is seen that this can be obtained by the use of 0.3 per cent Superfiltchar, or 0.8 per cent Filtchar with a treatment of 18 hours or so. The indication of the last part of the table is that no further improvement was obtained, however, by using a time of 18 hours instead of $2\frac{1}{2}$ hours in the case of Filtchar. One-quarter of an hour was too short a time.

Other absorbents.—Coke and cinders failed to remove any odor or color on a large scale in the treating plant. In the laboratory, clay from the river bank failed to remove any odor or color.

Large scale experiments.—Experiment 1: Used 35.5 cubic feet of toluidine waste and aerated it for one hour; then added enough lime (about 8 pounds) to make neutral and precipitate all the iron. This precipitate settled over night to less than 20 per cent of the total volume.

From Industrial Chemical Co., 200 Fifth Avenue, New York City.

Part of the supernatant liquid was pumped through the coke filter and then through the sand filter. The coke effluent was very deeply colored as was also the sand effluent. There was some slight improvement in odor.

Another part of the supernatant liquid was pumped through the cinder filter and then through the sand filter. The effluents from both cinders and sand were highly colored and unsatisfactory, although there was some improvement in odor.

Experiment 2: To 44.2 cubic feet of toluidine waste, after aeration, 8 pounds of lime were added as milk of lime. Pumped part of the supernatant liquid through the cinders first and sand filter afterwards.

To 4.75 cubic feet of this effluent was added 34 pounds of bleaching powder (10 per cent chlorine) and 1,220 cc. of 94 per cent sulphuric acid. This should have released 54 ounces of chlorine for chlorination, corresponding to 11.5 ounces per cubic foot of solution, or 11.5 grams per liter of solution.

After standing several hours, a thick foam was observed. The liquid beneath was medium brown in color, and neutral; it had a rather penetrating odor of chlorine. No improvement was obtained by filtering this liquid through cinders and sand.

This result checked the best results that could be expected from the laboratory experiments, but the effluent could hardly be considered satisfactory to discharge into the Rahway River.

Nitrobenzene Waste.

Precipitants.—The following precipitants failed to remove odor or color: Fe(OH)₂, Fe(OH)₃, Al(OH)₃, Si(OH)₄, Ca(OH)₂, CaSO₄, Zn(OH)₂, and PbSO₄.

Absorbents.—Clay from the river bank, blast-furnace slag, open-hearth slag, coke, charcoal, cinders, and bone black failed to remove odor or color.

Acid.—Acid caused a precipitation which was very finely divided. The supernatant liquid retained a yellow color and the original odor. The precipitate went into solution on warming, and fell out again on recooling.

Solvents.—Benzene and ether both failed to remove color from the nitrobenzene waste.

Bleaching powder.—Bleaching powder in acid solution gave a precipitate, which removed most of the color and odor.

Large scale experiments were not performed because of the discontinuance of the manufacture of nitrobenzene.

Monoethyl Aniline Waste. Receiver Liquor.

Bromine.—Bromine precipitated a yellow solid that turned brown on standing. The supernatant liquid was yellowish, and had no odor.

Salt.—Salt threw out a little monoethyl aniline, but the water layer retained a high color.

Precipitants.—Fe(OH)₂, Fe(OH)₃, Ca(OH)₂ were without effect on either color or odor.

Boiling.—Boiling did not improve odor in five minutes.

Filtchar.—Filtchar removed some odor, but even large quantities failed to remove very much.

Bleaching powder.—Bleaching powder in acid solution gave a dark purplish coloration. The character of the odor was changed and seemed phenolic. The color was not removed by Filtchar.

Dye.—Attempts to make an azo dye by adding the waste to diazotized arsol solution were fruitless.

Nitrous acid.—Nitrous acid caused the formation of a light colored precipitate that turned brown. The supernatant liquid was light yellow in color and had an odor that was somewhat improved, though phenolic in character.

Dilution.—A hundredfold dilution gave a liquid without odor or color.

Monoethyl Aniline Waste. Still Residue.

Bromine.—Bromine precipitated a brown solid, leaving a light yellow, odorless, supernatant liquid.

Filtchar.—Filtchar, above 1 per cent, removed all the odor and part of the color.

Bleaching powder.—Bleaching powder in acid solution caused a yellow colloidal solution to form, having no odor.

Ether.—Ether removed both color and odor.

Precipitants.—Fe(OH)₂, Al(OH)₃, Ca(OH)₂ all removed some color, but no odor.

Nitrous acid.—Nitrous acid caused a brown colloidal precipitate which could be carried down by Fe(OH)₂, leaving a light colored, odorless solution.

Dilution.—Upon dilution 100 times, both odor and color were no longer perceptible.

Large scale experiments.—The experiments 1 and 2 under "Dye wastes" show the effect of monoethyl aniline waste upon the precipitation treatment of dye wastes. In experiment 1 (page 180) the dilution was about 21 times for the combined monoethyl aniline wastes, and it is seen that the effluent had a little odor.

In experiment 2 the final dilution was 84 times, and the odor was practically all gone.

Lagoon Liquor.

Precipitants.—When lagoon liquor was made faintly alkaline with lime a slight flocculent precipitation was noted, which consisted of the hydroxides of iron and zinc. These precipitants failed to carry down any of the color. Other precipitants, including PbSO₄, Al(OH)₃ also failed to carry down any color.

It was found, however, that a large amount of ferrous iron would carry down color when precipitated. A 200 cc. sample of lagoon liquor was treated with ferrous sulphate containing 0.017 grams of iron, and made alkaline. The filtrate effluent from this sample had most of the color removed. On standing 24 hours, however, the solution took on a purplish color. It is believed that the effect of the iron was more one of reduction than one of carrying down color. The purplish tinge was so faint that it disappeared nearly altogether when the liquor was diluted to six times its volume.

The iron present in the lagoon liquor was found to be in the ferrous state when tested with ferricyanide, ferrocyanide, and thiocyanate.

Distillation.—When lagoon liquor was distilled, the first 20 per cent distillate was found to have a strong earthy odor, but very little color. The remaining distillate had neither color nor odor.

Oxidation.—Lagoon liquor decolorized permanganate without itself being decolorized. Boiling lagoon liquor with strong nitric acid and then making it alkaline seemed to remove some of the color.

Bromine.—Bromine caused the precipitation of a fine solid, and practically the decolorization of the solution.

Bleaching powder.—Bleaching powder in acid solution had practically the same effect. Table II shows the effect of varying amounts of chlorine on the color, and the effect of various conditions of chlorination.

Sample.	Grams of chlorine per liter.	Series C, alkaline lagoon liquor fil- tered. Color removal, alkaline.	Series D, acidified series C liquor. Color re- moval, acid.	Series E, lagoon liquor as found. Color removal.
1 2 3 4 5	0. 4 . 8 1. 2 1. 8 3. 0	Darker	dodo	Darker. Do. None. 75 per cent. 90 per cent.


TABLE II.—Effect of bleaching powder on lagoon liquor.

The 90 per cent color removal appears to be satisfactory, and it is seen that this requires 3 grams of chlorine per liter.

Absorbents.—In the laboratory it was found that charcoal, clay, blast furnace slag, open hearth slag failed to remove color from lagoon liquor.

Bone black, the Filtchar series, and Kelcar, were all found to be effective in removing color. In testing the efficacy of these compounds they were simply stirred up in measured proportions with the lagoon liquor, and allowed to stand overnight. The solutions were then filtered, and the color removal was determined by testing against the Bismarck brown standard (see footnote to Table I). The set of curves (Fig. 3) shows the efficacy of the various absorbents used.

From these curves it is seen that in order to obtain an effluent having an absolute color of 10 p. p. m. Bismarck brown, the require-




ment was about 0.9 or 1.0 gram per liter of either Superfiltchar, about 1.2 gram per liter of Kelcar, and about 2.3 grams per liter of Filtchar. (The special Superfiltchar is not on the market.)

It is also seen that the color removal was better in acid than in alkaline solution. When these acid-treated solutions were made alkaline the colors were about the same as if they had been treated alkaline in the first place.

¹ Produced from Pacific coast kelp by R. B. Stringfield, Los Angeles, Calif.

Attempts were made to rejuvenate the Filtchar and Superfiltchar which had been used up in treating lagoon liquor. The result of these attempts was to develop a process by which the material was heated to about 1,300° F. in air for a short time, and allowed to cool in the absence of air. If air was not excluded during the cooling, the material smouldered at low red heat and never became cooled off.

A continuous process for treating exhausted Filtchar was contemplated, by which it was slowly passed through a trough con-



veyor of the helical type, heated to the desired temperature. A small model of such a conveyor was operated and found to give good results. It was seen to be necessary, however, to break up all lumps, and to grind large particles fine, before passing through the heating chamber. The loss caused by conducting this heating in one stage in the laboratory averaged only 10 per cent. The set of curves (Fig. 4) shows the efficacy of rejuvenated Filtchar and Superfiltchar as compared with Filtchar as bought on the market.

From these graphs it is seen that the rejuvenated material is approximately in the same class as the fresh Filtchar.

One impractical feature of the continuous process is that the amount of fine dust caused is very great, and may perhaps entail large losses.

Large scale experiments.—In the 300-gallon treating plant, the best laboratory results of the chlorination and the Filtchar treating processes were tried out.

Experiment 1: Used 42.8 cubic feet of lagoon liquor; added 1.4 pound of bleaching powder (30 per cent chlorine, equivalent to 67.2 ounces chlorine); added 600 cc. of oleum.

The result of this treatment was a light brown sludge and a very light yellow, very sligh ly acid, supernatant liquid which had a slight odor of chlorine. It was considered a satisfactory effluent for discharge into the river.

Experiment 2: 29.4 cubic feet of lagoon liquor was treated with 500 cc. of slaked lime; at this point the solution was alkaline to litmus.

To 25.5 cubic feet of supernatant liquid from this treatment were added 23 ounces of Superfiltchar (oil grade) and this was allowed to stand overnight (corresponds to 0.090 per cent Superfiltchar). When the resulting suspension was filtered the effluent had a yellow color of 38 p. p. m. Bismarck brown, but little odor or taste. From the color point of view it was unsatisfactory to discharge into the river.

Experiment 3: Same as experiment 2, except that 50 ounces of Superfiltchar were used with 35 cubic feet of alkaline lagoon liquor (0.142 per cent Superfiltchar).

The result of filtering this treated waste was an effluent having only 4 p. p. m. color, no odor, and only a taste of lime. It was considered a satisfactory product to discharge.

Experiment 4: Used 40 cubic feet of lagoon liquor and 100 cc. of slaked lime. Upon settling, the supernatant liquid was neutral to litmus, alkaline to methyl orange, and acid to phenolphthalein.

To 25.7 cubic feet of this supernatant liquid were added 68 ounces of ordinary Filtchar (0.260 per cent), and the mixture stood overnight.

The effluent from filtering this mixture had a color of 8 p. p. m., practically no odor and practically no taste.

Dye Wastes: Lake-forming Dyes.

The color in the wastes from the particular dyes, "guinea green" and "formyl violet," were found to be carried down by the precipitants, ferrous hydroxide and aluminum hydroxide. The results were much better when lime instead of soda ash was used as the precipitating agent.

Quantitative experiments showed that a slight excess of lime was necessary in order that the iron should give the best results. The color of the resulting supernatant liquor depended upon the amount of iron used, but could be almost entirely removed with enough iron. Any residual color was not much affected by the use of Filtchar. The effect of dilution was to weaken the carrying down power of the same amount of iron.

Experiments were performed on a mixture of wastes in the proportion in which they were expected to be discharged from the plant, including wash water, as well as some monoethyl aniline waste, which was a relatively uncolored waste. This proportion was as follows:

Pe	r cent.
Lake-forming wastes	45. 1
Monoethyl aniline waste	
Wash water	50.0
•	100. 0

With this mixture it was found that the actual coloring matter could be carried down by 0.4 pound of iron as FeSO, per thousand pounds of waste. But this left a rather intensely colored yellowish solution. By using 1.2 pound of iron per thousand pounds of waste over 99.5 per cent of all the color could be removed. This also required 9.6 pounds of lime as milk of lime per thousand pounds. The result seemed to be partially due to the reducing action of the ferrous sulphate, because some yellow color returned slowly upon oxidation in the air. The best results were inhibited by mixture of any nonlake-forming dye waste with this, because a brownish coloring matter was formed, which could not be removed with Filtchar.

The odor of the waste when treated with copperas and lime was not very pleasant, as there were faint traces of benzaldehyde, monoethyl aniline, and other odors in it. This odor was not much improved upon treatment with Filtchar, but disappeared on dilution of the waste with water. Partial evaporation of 40 per cent of the treated waste failed to improve the odor to any extent.

The taste was very salty from the salt used in the manufacturing process. Table III shows the effect of dilution upon the odor, taste, and color of the treated waste.

TABLE III.—Effect of dilution upon the odor, taste, and color of the treated waste.

Dilution.	Oder.	Taste.	Color.	
One-eighteenth One-fortieth One sixty-seventh One one-hundredth One two-hundredth		do	4.5 2.3 1 0	

From this table it is observed that a dilution of one-fourth gives an effluent satisfactory from the color standpoint, a dilution between one-fourth and one-eighth from the odor standpoint, and about one one-hundredth from the taste standpoint.

Azo-phloxine waste could be decolorized by copperas and lime. It appeared that a large quantity of each chemical was required to give the necessary decolorization.

Evaporation.—Evaporation experiments were performed to determine whether the salt contained in the wastes could be used again. Some of the salt obtained from both the guinea green and the formyl violet was used again for salting out guinea green and formyl violet dyes, respectively. The dyes thus precipitated were tested in the local dye laboratory and found to be as clean and as good as the dye salted out by fresh salt.

Bleaching powder.—Azo-aphloxine waste could be largely decolorized with bleaching powder and acid. The resulting liquid was brownish in color and quite odorous from the chlorine and chlorinated products.

Large-scale experiments.—Experiment 1: A mixture of wastes, to simulate working proportions, was made as follows:

•	Cubic feet.
Formyl violet	6. 5
Guinea green	10. 8
Monoethyl aniline still residue	
Monoethyl aniline receiver waste	1. 20
Water	20. 8
m . •	
Total	40. 0

To this were added solutions of ferrous sulphate containing about 15 ounces of iron and a half bucket of moist slaked lime weighing about 15 pounds. After agitating, this was allowed to stand overnight.

The supernatant liquid was of a quite intense yellow color and had a moderate odor. The color removal was 80 per cent.

A quantity of 14.2 cubic feet was treated in another tank with 30 ounces of Superfiltchar and 11 ounces of rejuvenated Filtchar.

At the end of two days the odor removal was nearly complete and the color removal was still about 80 per cent. In other words, the Filtchar removed odor but not color.

Experiment 2: Wastes were mixed together as follows:

C	ubic feet.
Guinea green	. 4.8
Formyl violet	. 2.9
Monoethyl aniline still residue.	
Monoethyl aniline receiver liquor	
Water	
Total	17 1

Added ferrous sulphate containing 30 ounces of iron (1.75 ounces per cubic foot, equivalent to 1.75 pounds per thousand pounds); added, with agitation, 20 pounds of moist slaked lime (9.4 ounces per cubic foot, equivalent to 9.4 pounds per thousand pounds).

Samples taken of the supernatant liquid after standing overnight showed a color removal of 99.5 per cent, absolute color of 18 p. p. m., and a rather limy odor which, on standing, gave way to a faint one of organic compounds. The volume of the sludge was about 70 per cent of the total volume.

This waste was agitated and filtered through the sand filters. It was found that with a 3-inch covering of iron sludge and a head of about 5 inches, the rate was about 34 gallons per square yard per hour.

The effluent from the sand filter was greenish in color, but it was seen that this color was caused by suspended iron hydroxide. On oxidation by air and filtering in the laboratory, the suspension was all removed; but the filtrate was brown in color—about 80 p. p. m. Bismarck brown. On diluting this six times, the laboratory sample was seen to have a satisfactory color and practically no odor.

Nonlake-Forming Dyes.

Precipitants.—The nonlake-forming dye wastes were those from the manufacture of the "direct green" series of dyes. It was found that the following precipitants failed to carry down any color: Fe(OH)₂. Fe(OH)₃, Al(OH)₃.

When the waste was mixed with the lake-forming wastes and given the precipitation treatment, it was found that the effluent was of dark green color instead of a light yellow. This color was not removed by treatment with 1.5 per cent Filtchar.

If the waste was acidified and filtered before addition to the lakeforming wastes, and then given the precipitation treatment, the color of the filtrate was dark brown, and could be reduced to 15 p. p. m. Bismarck brown by treatment with 3 per cent Filtchar or 3 per cent Kelcar.

Acid.—Sulphuric acid caused a considerable foaming, but no color change. The liquid was then filtered and a dark slimy substance was removed. Upon treatment of this filtrate with Kelcar or Filtchar, there was little improvement in color. If, instead, it was treated with copperas and lime, and then with Filtchar, there was still little color improvement.

Reduction.—The reducing effect of ferrous sulphate was practically nil. By vigorous reduction with hydrogen from zinc and acid, most of the color was removed. Part of this returned on oxidation by the air.

III. RESULTS.

The results of this investigation show by what methods the various wastes produced in the manufacture of some dyes and intermediates can be treated. These treatments include chlorination, precipitation, and absorption.

The treatment for each waste or group of wastes will depend upon the relative cost, it being desired, of course, to achieve the best results at the lowest cost. The estimated relative costs of the various processes are given below. These include only those costs of each process that are not common to all the processes. For example, such a common cost would be the making alkaline of an acid waste.

TABLE IV.—Comparative costs of treating Toluidine waste.

Method.	Color of treated waste, p. p. m. Bismarck brown.	Odor of treated waste.	Cost per pound of waste.	Cost per 1,000 gallons.	Cost per pound of product.
Chlorination, 30 grams of chlorine to the liter (at \$0.06).	Medium brown	Rather strong	\$0.0018	\$15.00	\$0.0060
77	About 5	None	.0006	5.00 10.80	. 0020 . 0043
Filtchar, 8 pounds per 1,000 (\$0.06 per pound).	do	do	.00048	4.00	.0016
Superfiltchar, 3 pounds per 1,000 (\$0.14 per pound).	do	Faint	. 00042	3.50	.0014

The cost of the Filtchar and Superfiltchar treatments is seen to be slightly under the cost of evaporation, while all others are higher. The advantages of the Filtchar treatment over evaporation are that it requires less equipment, and is more flexible as regards the quality of the treated waste. Furthermore, there is a possibility that exhausted Filtchar can be cheaply rejuvenated. The advantage of the Filtchar treatment over the Superfiltchar treatment is that the Filtchar is likely to be a stable product on the market, whereas the Superfiltchar is more of a special product.

LAGOON LIQUOR.

The cost of treating the lagoon liquor should be cut down as low as possible because it is of such a large volume and represents no income from production. The rejuvenation of exhausted Filtchar offers the best possibilities. If it is to be considered that the loss by rejuvenation is 10 per cent each time, we can determine the life of 1 pound of Filtchar by summing the decreasing geometrical series, $1 + (0.9 \times 1) + [0.9 \times (0.9 \times 1)] + \dots$, which converges to a definite limit. The sum of this series is 10. In other words, by rejuvenation,

1 pound of Filtchar becomes the equal of 10 pounds of fresh Filtchar, theoretically. If Filtchar is \$0.06 per pound, the equivalent in rejuvenated Filtchar is \$0.006 per pound. Adding \$0.01 per pound to this for the cost of rejuvenation we arrive at a total estimate of \$0.016 per pound for material.

TARLE	V.—Comparative	costs of	treating	laaoon	limor.
TVDDD	1.—Consparative	coole uj	creating	iayoon	uguoi.

Method.	Color of treated waste, p. p. m. Bismarck brown.	Odor of treated waste.	Cost per pound of waste.	Cost per 1,000 gallons.
Evaporation. Chlorination: 3 grams of chlorine to the liter (at \$0.06 per pound). Superflitchar: 0.85 gram to the liter (pharmaceutical, at \$0.19 per pound). Superflitchar: 0.95 gram to the liter (pharmaceutical, at \$0.19 per pound). Filtchar: 1.7 grams to the liter (\$0.06 per pound). Filtchar: 2.5 grams to the liter (\$0.06 per pound). Rejuvenated Filtchar: 2.5 grams to the liter (\$0.016 per pound). Kelcar: 1.1 grams to the liter (\$0.045 per pound). Precipitation and dilution: Lime, at \$0.008 per pound, 9.6 pounds to 1,000 pounds of waste; FeSO ₄ at \$0.010 per pound Fe 1.2 pounds to 1,000 pounds of waste.	12 12 12 12 12 8 8 12	Very slight do	\$0.0006 .00018 .00016 .00013 .00010 .00015 .00004 .000060 \$\int 000077 .000012	\$5.50 1.00 1.33 1.08 .83 1.25 .332 .41

Table V indicates that the treatment of the waste by Filtchar in connection with a process for rejuvenation for the latter offers the best prospects for treatment at low cost. The treatment by precipitation and dilution is nearly as cheap in material, and is likely to be much more convenient in practice as there will not be the rejuvenation process to be handled. Furthermore, the precipitation process has the advantage of being a duplicate of the process recommended for the lake-forming dye wastes.

The treatment by Kelcar without rejuvenation is also a very cheap process, and its only disadvantages are the uncertainty of the supply and the distance it must be shipped (Los Angeles, Calif., to Springfield, N. J.).

LAKE-FORMING DYE WASTES.

The most effective treatment for lake-forming dye wastes was found to be the precipitation treatment followed by dilution.

Table VI.—Comparative costs of evaporation and precipitation methods of treating lake-forming dye wastes.

Method.	Residual color,	Cost per	Cost per
	p. p. m. Bis-	pound of	1,000
	marck brown.	waste.	gallons.
Evaporation	15	\$0.0006 .000089	\$5.00 .74

Table VI does not give a true comparison of the costs of the two methods enumerated, because salt having value would be obtained as a product of an evaporation. To carry out such a method of

evaporation it would be necessary to separate each actual waste liquor from the wash water of the tanks, filter presses, etc., and evaporate only the former, each waste separately. The wash water, on account of very low concentration of coloring matter, could be quite cheaply treated with copperas and lime.

IV. SUMMARY. .

Investigations have been carried out as to properties and disposal treatment of several wastes of the dye and intermediate industries. Laboratory studies were made of the properties of toluidine waste, nitrobenzene waste, monoethyl aniline waste, dye wastes, and lagoon liquor.

Treatments that were successful on a 300-gallon scale and that were cheaper than evaporation were found for the lagoon liquor, the monoethyl aniline wastes, and the lake-forming dye wastes; and a method of treating toluidine waste was indicated in laboratory experiments.

Specific treatments of wastes were recommended to the Chemical Company of America, based on their proposed production schedule. For the lake-forming dye wastes and monoethyl aniline wastes this treatment was as follows:

- 1. A mixture together of all the wastes and wash waters except the waste and wash water from the manufacture of the "Direct green" series of dyes.
- 2. Addition and dissolving of ferrous salt in the proportion of 1.2 pounds of iron per thousand pounds of solution. This can be done in a wooden vat.
- 3. Addition, with agitation, of slaked lime in the proportion of 8 pounds of Ca(OH), per pound of Fe (or 9.6 pounds of Ca(OH), per thousand pounds of solution). This amount is in addition to the amount necessary to neutralize the acid in the waste if any should occur.
- 4. Filtration with or without settling of the supernatant liquid. This can be done by means of a sand filter bed or any other suitable filtering medium.
- 5. Dilution of the effluent from the filter with five times its volume of water.
- 6. Daily removal of the solid sludge from the filter to suitable dumps or hollows.
- 7. The only cheap method found for disposal of the "Direct green" dye wastes is evaporation.

APPENDIX.

There is given below a systematic set of steps for experimental treatment of wastes. Realizing that evaporation is the simplest form of treatment for liquid effluents, the figures given below are

¹ The plant recommended has been constructed and is operating with entire satisfaction.

the maximum quantities of materials that can be used in order that the process be cheaper than evaporation. They are based on market prices in June, 1919.

(1) Try aeration for blowing out bad odors.

- (2) Try Filtchar up to 8 pounds per thousand of waste for removing odor and color.
- (3) Try chlorine (in both acid and alkaline solution) up to 8 pounds per thousand for improving odor and color.
- (4) Try copperas and lime for precipitating color; also alum and lime, etc.
- (5) Try boiling for removing bad odors.
- (6) Try partial evaporation for removing bad odors.
- (7) Try strong reducing agents; also strong oxidizing agents.
- (8) Try strong alkali up to 5 pounds per thousand of waste.
- (9) Investigate the manufacturing process with reference to making use of the waste again several times. For instance, in the case of the residue from the manufacture of toluidine by acid reduction of nitrotoluene, this waste might be fortified with strong acid and used again.
- (10) Try shaking out with ether up to 5 pounds per thousand of waste.
- (11) Try dilution.
- (12) Try bacterial action.

If no satisfactory improvement is obtained by any one or combination of these schemes, then the special chemical and physical properties of the objectionable constituents should be studied.

For instance, an aldehyde might be oxidized to an acid having an insoluble lime salt, or a nitro compound might be reduced to form an amine removable by Filtchar. An aromatic primary amine might be precipitated as an azo compound.

Other absorption compounds than Filtchar, such as Kelcar, Norite, coke, talc, fuller's earth, should be studied if they offer cheaper possibilities.

With special reference to dyes which are salted out, there is a good field for studying the recovery of salt from the wastes by evaporation.

It should be remembered in any case that the waste must be made alkaline after treatment if it is not already so, and should be allowed to stand in air as a test, for it will be oxidized as well as made alkaline in the river.

The removal of the dyes by electrical precipitation may be a field for study. It could not be studied at the experimental plant for lack of direct current power.

NATIONAL DENTAL ASSOCIATION COOPERATING IN VENE-REAL DISEASE CONTROL.

At the twenty-third annual session of the National Dental Association, held in New Orleans on October 20-24, the following resolution was unanimously adopted:

Whereas, it has been the accepted policy of the National Dental Association of the United States to cooperate with and support the Federal Government in its every department in all measures undertaken for the advancement of the public health and the welfare of the Nation; and

Whereas, the United States Public Health Service, under authority of Congress, and by direction of the President, is engaged in a Nation-wide campaign for the control of the venereal diseases; and

Whereas the United States Public Health Service has supplied to the convention a most unusual showing of social hygiene films at the Globe Theater, and a most comprehensive exhibit of slides, charts, and literature in the Grunewald Hotel in demonstration and explanation of the propaganda; and

Whereas the 4,500 officers, delegates, members, and guests of the convention have, through the untiring efforts of Acting Asst. Surg. Charles V. Herdliska, been made cognizant of the necessity for cooperation on the part of the dental profession of the United

States in this vital work of venereal disease control; and

Whereas the United States Public Health Service, following its usual policy of cooperation with the various State boards of health, is about to launch its campaign from Washington with the 45,000 dentists of the country to secure the cooperation of each individual dentist, through an agreement card to be sent to him for his sig-

nature: and

Whereas under medical measures the campaign already carried on by the Public Health Service, and since their transfer to the various State boards of health for completion with advertising media, druggists, physicians, medical and allied colleges have resulted in the cooperation, through signing and returning of agreement cards, of over 99 per cent of the 20,000 advertising media, of more than 60 per cent of the 48,500 druggists, of more than 50 per cent of the 132,000 physicians, and approximately all the medical and allied colleges of the country; and

Whereas conventions of advertising media, druggists, and physicians have unequivocally indorsed this propaganda of the Public Health

Service; Therefore, be it

Resolved, That the twenty-third annual convention of the National Dental Association of the United States, following the course already pursued by the advertising media, druggists, physicians, and the medical and allied colleges of the United States, hereby approves and indorses the propaganda for the control of the venereal diseases, undertaken by the United States Public Health Service, and hereby gives assurance that it will use its best endeavors not only to secure the interest and the cooperation of every member of the dental profession, but will also use its best endeavors to assist in the general plan outlined by the Public Health Service for the education of the civilian population in this vital subject.

DEATHS DURING WEEK ENDED JAN. 10, 1920.

[From the "Weekly Health Index" Jan. 13, 1020, issued by the Bureau of the Census, Department of Commerce.]

Deaths from all causes in certain large cities of the United States during the week ended Jan. 10, 1920, infant mortality (per cent), annual death rates, and comparison with corresponding week of preceding years.

	Population		Week ended Jan. 10, 1920.		Per cent of deaths under 1 year.	
City.	July 1, 1918, esti- mated.	Total deaths.	Death rate.1	annual death rate per 1,000.2	Week ended Jan. 10, 1920.	Previous year or years.2
Albany, N. Y Atlanta, Ga Baltimore, Md Boston, Mass Buffalo, N. Y Cambridge, Mass Chicago, Ill. Cincinnati, Ohio Cleveland, Ohio. Cleveland, Ohio. Columbus, Ohio. Dayton, Ohio Dayton, Ohio Denver, Colo. Fall River, Mass Grand Rapids, Mich Indianapolis, Ind Jersey City, N. J Kansas City, Mo Los Angeles, Calif Louisville, Ky Lowell, Mass. Memphis, Tenn Milwaukee, Wis. Minneapolis, Minn Nashville, Tenn Newark, N. J Newark, N. J Newark, N. J Newark, N. Y Oakland, Calif. Oomaha, Nebr Philadelphia, Pa Pittsburgh, Pa Portland, Oreg. Providence, R. I Richmond, Va Rochester, N. Y St. Louis, Mo St. Paul, Minn San Francisco, Calif. Seattle, Wash Spokane, Wash Spokane, Wash Spokane, Wash Syracuse, N. Y Toledo, Ohlo. Washington, D. C. Worcester, Mass.	810, 306 225, 296 130, 655 128, 392 135, 450 290, 389 318, 770 313, 759 109, 081 154, 759 453, 481 119, 215 383, 442 119, 215 383, 442 119, 215 382, 273 5, 215, 879 214, 206 180, 264 1, 761, 371 593, 303 160, 719 264, 856 779, 951 257, 699 478, 530	39 194 194 688 141 194 688 195 28 95 151 35 95 151 167 64 1,534 167 167 167 167 167 167 167 167 167 167	18. 1 17. 8 15. 5 19. 15. 5 18. 0 14. 0 11. 8 17. 1 10. 8 17. 1 15. 8 13. 9 17. 2 12. 5 11. 6 13. 9 14. 6 19. 6 19. 6 19. 6 19. 6 19. 6 19. 1 19. 1	C 22, 7 C 19, 6 A 20, 2 C 16, 6 A 17, 4 C 20, 2 C 14, 1 C 212, 4 C 212, 4 C 18, 8 C 19, 1 A 17, 4 C 21, 1 C 18, 8 C 19, 1 A 17, 4 C 21, 1 C 18, 8 C 19, 1 A 17, 4 C 21, 1 C 18, 8 C 19, 1 C 11, 1 C 21, 1 C 14, 1 C 21, 1 C 11, 1 C 21, 1 C 11, 1 C 11, 2 C 11, 1 C 11, 2 C 11, 1 C 11, 2 C 11, 3 C 11, 4 C 11, 5 C 11, 4 C 11, 5 C 11, 7 C 11, 8 C 1, 1 C 11, 7 C 11, 8 C 1, 1 C 1 C 1, 1 C 1 C 1, 1 C 1 C 1, 1 C	10.3 13.0 14.4 17.6 15.6 15.8 8.5 22.9 8.4 37.9 19.3 8.4 9.3 8.9 16.7 10.2 4.8 13.9 14.7 10.2 4.8 13.9 14.9 13.5 13.5 13.5 13.5 13.5 14.9 13.8 13.5 13.5 13.5 13.6 13.8 13.9 13.8 13.9 13.8 13.9 13.8 13.9 13.8 13.9 13.8 13.9 13.8 13.9 13.8 13.9 13.8 13.8 13.8 13.8 13.8 13.8 13.8 13.8	C 18.4 C 17.9 A 11.7 C 24.5 A 16.2 C 17.2 C 14.8 C 17.2 C 16.3 C 10.3 C 12.1 C 10.3 C 12.1 C 11.5 A 16.9 C 11.15 C 11.5 C 11.5 C 12.6 C 11.7 C 13.8 C 11.7 C 13.8 C

Summary of information received by telegraph from industrial insurance companies for week ended Jan. 10, 1920.

Policies in force	42, 358, 989
Number of death claims	8, 914
Death claims per 1 000 policies in force annual rate	

Annual rates per 1,000 estimated population.
 "A" indicates data for the corresponding week of the years 1913 to 1917, inclusive. "C" indicates data for the corresponding week of the year 1917.
 Population estimated as of July 1, 1919.
 Data are based on statistics of 1915, 1916, and 1917.

PREVALENCE OF DISEASE.

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring.

UNITED STATES.

CURRENT STATE SUMMARIES.

Telegraphic Reports for Week Ended Jan. 17, 1920.

These reports are preliminary, and the figures are subject to change when later returns are received by the State health officers.

ALABAMA.		CALIFORNIA—continued.	
Cerebrospinal meningitis	ases.	Lethargic encephalitis:	306.
Chicken pox.		Los Angeles	2
Diphtheria		San Francisco	
Malaria.		Santa Clara	ī
Measles		Shasta County	ī
Pneumonis, broncho.		Poliomvelitis:	-
Pneumonia, lobar	-	Les Angeles	1
Scarlet fever		Smallpox:	-
Smallpox		Alameda	4
Tuberculosis, pulmonary		Les Angeles County	6
Typhoid fever	9	Oakland	7
Venereal diseases	64	Riverside	15
Whooping cough	8	Solano County	10
• • •		Scattering	25
arkansas.		Typhoki fever	11
Cerebrospinal meningitis	3		
Chancroid	1	CONNECTICUT.	
Chicken pox	32		
Diphtheria	34	Cerebrospinal meningitis:	
Gonorrhea.	33	New London	1
Hookworm	2	Waterbury	1
Influenza	53	Influenza	14
Malaria	76	Smallpox:	_
Measles	12	Waterbury	1
Ophthalmia neonatorum	1		
Pellagra	6	DELAWARE.	
Scarlet fever	17	Chicken pox	7
Smallpox	18	Diphtheria:	
Syphilis	16	Wilmington	9
Trachoma	5	Scattering	4
Tuberculosis	49	Erysipelas:	
Typhoid fever	12	Wilmington	1
Whooping cough	18	Measles	50
CALIFORNIA.	- 1	Mumps	6
CALIFORNIA.	- 1	Pneumonia:	
Cerebrospinal meningitis:	i	Dover	3
Oakland	1	Wilmington	4
Orange	1	Scarlet fever	8
Sacramento	1	Tuberculosis	3
Influenza	264	Whooping cough	8
		, =	

FLORIDA.		ILLINOIS—continued.	
Diphtheria		Scarlet fever—Continued.	isos.
Dysentery	7	Evanston.	10
Influenza	10	Forest Park	
Malaria	18	Oak Park	
Ophthalmia neonatorum	1	Peoria	4
Pneumonia	14	Quincy	5
Scarlet fever	3	Riverside	
Smallpox	70	Woodstock	
Membeld force			
Typhoid fever	8	Scattering	80
GEORGIA.		Smallpox:	
		Benton.	10
Anchylostomiasis	60	Carriers Mills	7
Cerebrospinal meningitis	2	Eldorado	7
Chicken pox	18	Galesburg	9
Diphtheria	8		_
Gonorrhea	17	Glasgow	4
		Hamilton County—	
Influenza	27	Crook Township	6
Malaria	15	Jersey ville	13
Measles	4	Monmouth	16
Mumps	4	Rock Island	8
Paratyphoid fever	1	Warren County—	•
Pneumonia, lobar	21		
		Kelly Township	10
Poliomyelitis	2	Scattering	50
Scarlet fever	7	Syphilis	152
Septic sore throat	2	Typhoid fever	16
Smallpox	30	′	
Syphilis	17	INDIANA.	
	- 1		
Tuberculosis, pulmonary	9	Cerebrospinal meningitis:	
Typhoid fever	11	Johnson County	1
Whooping cough	9	Vigo County	1
	- 1	Chancroid	6
ILLINOIS.	- 1	Diphtheria:	•
Combine minel meninglish	ı		
Cerebrospinal meningitis:	. 1	Delaware County	6
Aurora	1	Lake County	14
Chicago	4	Maricn County	9
Chancroid	18	Scattering	20
Diphtheria:	- 1	Gonorrhea	82
Chicago 1	50	Influenza:	
		Dekalb County	10
East Alton	4		10
East St. Louis	6	- Koseiusko County	10
Peoria	4	Lake County	8
Streator	7	Tipton County	6
Scattering	55	Scattering	10
Gonorrhea. 2		Measles:	
Influenza:	~	Dearborn County	-
		•	5
Camp Grant 3		Delaware County	7
Chicago	27	Greene County	5
Forest Park	4	Hancock County	8
Freeport	7	Jay County	25
Great Lakes Naval Training Station 7	- 1	Jackson County	63
	25	Johnson County.	4
		-	_
	29	Lake County	14
	12	Marion County	35
Poplar Grove	4	Rush County	5
Rockford84	49	Wabash County	11
Waverly 16		Wayne County.	21
Zion City	5	_ *	18
	-	_	10
	30	Scarlet fever:	
Lethargic encephalitis:	-1	• • • • • • • • • • • • • • • • • • • •	10
Chicago	4	Boone County	10
Poliomyelitis:		Carroll County	10
*	2	Decatur County	5
Scarlet fever:	-	Delaware County	6
	.		-
	9	Elkhart County	10
	4	Hendricks County	8
Chicago 39	19 I	Kosciusko County	11

INDIANA—continued.	~	LOUISTANA—continued.
Scarlet fever—Continued.	Cases	Smallpex. Cases. 13
Lake County	1	1 Syphilis
Marion County	3	
Rush County		2
Tippecanoe County		7 MAINE.
Wayne County.		
Scattering		6 Chicken pox 14
Smallpox:		Conjunctivitis
Dearborn County	. 2	Diphtheria 10
Dubois County.		Gonorrhea
Fountain County		Influenza 4
Gibson County		Measles:
Jackson County		Delan 1
Lake County.		Sanford 12
Marion County		1 0
Spencer County		35
Sullivan County		Pneumonia 1
Warrick County	-	Constant formers
Scattering.		77
Syphilis		TP144
Typhoid fever:		Paris. 5
Elkhart County	. 4	Dantland as
Scattering		Goodening 91
Dearteining	•	Smallpox:
IOWA.		Bridgewater. 4
Cerebrospinal meningitis:		Lewiston 6
Dubuque		Scattering 8
Chancroid		Syphilis. 11
		Tuberculosis
Chicken pox	. 2	Typhoid fever
Diphtheria:	_	Whooping cough. 6
Des Moines		whooling confinence of
Scattering		Massachusetts.
Gonorrhea	97	
Influenza:		Cerebrospinal meningitis 6
Davenport		Chicken pox
Eagle Grove		Diphtheria 218
Fort Dodge		Dog bite
Scattering	10	German measles
Measles:		Gonorrhea
Mason City		Influenza. 54
Scattering	5	Measles. 821
Mumps	3	Mumps
Scarlet fever:		Ophthalmia neonatorum
Boone	4	Pneumonia, lobar. 145
Council Bluffs	5	Scarlet fever. 415
Des Moines	11	Septic sore throat
Humboldt County	4	Smallpox
· Onawa	5	Suppurative conjunctivitis
Scattering	40	Syphilis 103
Smallpox:		Trachoma. 3
Cedar Rapids	5	Tuberculosis, pulmonary 136
Council Bluffs	4	Tuberculosis, other forms
Davenport	30	Typhoid fever 13
Dike	5	Whooping cough
Franklin County	5	Winodang congn
Greenfield	7	Minnesota.
Scattering	21	Chancroid. 1
Syphilis	17	Gonorrhea. 87
Whooping cough	8	Smallpox:
· ·	•	Duluth 8
LOUISIANA.		Marshall County—
Cerebrospinal meningitis	5	Harrison Township 6
Chancroid	32	Otter Tail County—
Diphtheria	18	Fergus Falls
Gonorrhee	98	Steele County—
Influenza	27	Owatoms 4
	1	~ ************************************

MINNESOTA—continued.	.	NEW MEXICO—continued.	
Smallpox—Continued.	285 65.	C ₁	asos.
Swift County—		Pneumonia	. 7
	_	Scarlet fever:	
Clontarf		Taiban	6
Scattering		Van Houten	4
Syphilis	. 96	Scattering.	-
		Septic sore throat	6
MONTANA.		C	1
Diphtheria		Smallpox:	
Influenza	. 1	Tularosa	18
Measles	. 25	Scattering.	14
Scarlet fever	. 72	Syphilis	1
Smallpox		Tetanus	ī
Typhoid fever	. 4	Trachoma	
1 ypnoru iever	. 4	Tuberevlesie	1
NEBRASKA.		Tuberculosis	25
Chancroid	. 12	Typhoid fever	2
		Whooping cough	2
Chicken pox.	. 11		
Gonorrhea		NEW YORK.	
Influenza	. 1	477	
Measles:		(Exclusive of New York City.)	
Central City	. 9	Anthrax:	
Omaha		Gloversville	1
			1
Scattering		Cerebrospinal meningitis:	
Mumps	. 8	Riverhead	1
Scarlet fever:		Diphtheria	209
Callaway	6	Gonorrhea	60
Chadron		Influenza	61
Holt County		Measles	
		Pneumonia	100
Omaha		T Redifforms	182
Otoe		Scarlet fever.	239
Sutton	4	Smallpox	7
Wilber	4	Syphilis	203
Scattering		Typhoid fever	17
		trees.	
Smellner		W booning cough	262
Smallpox:		Whooping cough	262
Central City	9		262
Central City	9 7	Whooping cough NORTH CAROLINA.	262
Central City	- 1	NORTH CAROLINA.	
Central City	7	NORTH CAROLINA. Cerebrospinal meningitis	3
Central City Dorchester Elmwood Humboldt	7 4 5	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid	3 15
Central City Dorchester Elmwood Humboldt Kearney	7 4 5 6	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox.	3 15 87
Central City Dorchester Elmwood Humboldt Kearney	7 4 5 6 9	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox. Diphtheria.	3 15 87 55
Central City Dorchester Elmwood Humboldt Kearney Lincoln Omaha	7 4 5 6 9 8	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox. Diphtheria. German measles.	3 15 87 55 5
Central City. Dorchester. Elmwood. Humboldt. Kearney Lineoln. Omaha. Seward.	7 4 5 6 9 8	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chieken pox. Diphtheria. German measles. Gonorrhea.	3 15 87 55
Central City. Dorchester. Elmwood. Humboldt. Kearney Lineoln. Omaha. Seward. University Place.	7 4 5 6 9 8	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox. Diphtheria. German measles.	3 15 87 55 5
Central City. Dorchester. Elmwood. Humboldt. Kearney Lineoln. Omaha. Seward.	7 4 5 6 9 8	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox. Diphtheria. German measles. Gonorrhea. Measles.	3 15 87 55 5
Central City. Dorchester. Elmwood. Humboldt. Kearney Lineoln. Omaha. Seward. University Place.	7 4 5 6 9 8 9 7 4	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox. Diphtheria. German measles. Gonorrhea. Measles. Ophthalmia neonatorum.	3 15 87 55 5 97 79
Central City Dorchester Elmwood Humboldt Keerney Linceln Omaha. Seward. University Place York Scattering	7 4 5 6 9 8 9 7 4	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho	3 15 87 55 5 97 70 1
Central City. Dorchester. Elmwood. Humboldt. Kearney Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis.	7 4 5 6 9 8 9 7 4 39 54	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar	3 15 87 55 5 97 70 1 33 47
Central City. Dorchester. Elmwood. Humboldt. Kearney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis.	7 4 5 6 9 8 9 7 4 39 54	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis	3 15 87 55 5 97 79 1 33 47
Central City. Dorchester. Elmwood. Humboldt. Kearney Linceln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever.	7 4 5 6 9 8 9 7 4 39 54 1 2	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox. Diphtheria. German measles. German measles. Gonorrhea. Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar. Poliomyelitis. Scarlet fever.	3 15 87 55 5 97 70 1 33 47
Central City. Dorchester. Elmwood. Humboldt. Kearney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis.	7 4 5 6 9 8 9 7 4 39 54	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis	3 15 87 55 5 97 79 1 33 47
Central City Dorchester Elmwood Humboldt Kearney Lincoln Omaha Seward University Place York Scattering Syphilis Tuberculosis Typhoid fever Whooping cough	7 4 5 6 9 8 9 7 4 39 54 1 2	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chicken pox. Diphtheria. German measles. German measles. Gonorrhea. Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar. Poliomyelitis. Scarlet fever.	3 15 87 55 5 97 79 1 33 47 1 45 4
Central City. Dorchester. Elmwood. Humboldt. Kearney Linceln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever.	7 4 5 6 9 8 9 7 4 39 54 1 2	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox	3 15 87 55 5 97 70 1 33 47 1 45 4
Central City. Dorchester Elmwood Humboldt Keerney Lincoln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough	7 4 5 6 9 8 9 7 4 39 54 1 2	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis	3 15 87 55 5 79 70 1 33 47 1 45 4 61
Central City. Dorchester. Elmwood. Humboldt. Kearney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent.	7 4 5 6 9 8 9 7 4 39 54 1 2	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Trachoma	3 15 87 55 5 5 97 70 1 33 47 1 45 4 61 49
Central City. Dorchester. Elmwood. Humboldt. Kearney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent. Influenza.	7 4 5 6 9 8 9 7 4 39 54 1 2	NORTH CAROLINA. Cerebrospinal meningitis. Chancroid. Chieken pox. Diphtheria. German measles. Gonorthea. Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis. Scarlet fever. Septic sore throat. Smallpox. Syphilis. Trachoma. Tryphoid fever.	3 15 87 55 5 5 97 70 1 33 47 1 45 4 61 49 1 9
Central City. Dorchester Elmwood. Humboldt. Keerney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent.	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Trachoma	3 15 87 55 5 5 97 70 1 33 47 1 45 4 61 49 1 9
Central City. Dorchester Elmwood. Humboldt Keerney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia.	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox Syphilis Trachoma Typhoid fever Whooping cough	3 15 87 55 5 5 97 70 1 33 47 1 45 4 61 49 1 9
Central City. Dorchester Elmwood Humboldt Keerney Linceln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox. Diphtheria. German measles. German measles. Gonorrhea. Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis. Scarlet fever. Septic sore throat Smallpox. 1 Syphilis. Trachoma Typhoid fever. Whooping cough.	3 15 87 55 5 5 97 70 1 33 47 1 45 4 61 49 1 9
Central City. Dorchester Elmwood. Humboldt Keerney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia.	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorthea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox Smallpox 1 Syphilis Trachoma Typhoid fever Whooping cough 1 OHIO. Diphtheria:	3 15 87 55 5 5 97 70 1 33 47 1 45 49 1 9 07
Central City. Dorchester Elmwood Humboldt Keerney Linceln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever. Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair.	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox Smallpox 1 Syphilis Trachoma Typhoid fever Whooping cough OBIO. Diphtheria: Youn;stown	3 15 87 55 5 5 97 70 1 33 47 1 45 4 61 49 1 9
Central City. Dorchester Elmwood Humboldt Keerney Linceln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO.	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Truchoma Typhoid fever Whooping cough 1 OHIO Diphtheria: Youngstown	3 15 87 55 5 5 97 70 1 33 47 1 45 49 1 9 07
Central City. Dorchester Elmwood Humboldt Keerney Lincoln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Precumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chancroid.	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Trachoma Typhoid fever Whooping cough OHIO. Diphtheria: Youngstown. Scarlet fever:	3 15 87 55 5 5 97 70 1 33 47 1 45 49 1 9 07
Central City. Dorchester Elmwood Humboldt Keerney Lincoln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Precumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chancroid.	7 4 5 6 9 8 9 7 4 39 54 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid. Chicken pox. Diphtheria. German measles. Gonorrhea. Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever. Septic sore throat Smallpox. 1 Syphilis. Truchoma. Typhoid fever. Whooping cough. Diphtheria: Youn;stown. Scarlet fever: Akron.	3 15 87 55 5 97 79 1 33 47 1 45 4 61 49 1 9 07 28 79
Central City. Dorchester. Elmwood. Humboldt. Kearney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chancroid. Chicken pox.	7 4 5 6 9 8 9 7 4 39 54 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid. Chicken pox. Diphtheria. German measles. Gonorrhea. Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever. Septic sore throat Smallpox. 1 Syphilis. Trachoma. Typhoid fever. Whooping cough. OHIO. Diphtheria: Youn;stown. Scarlet fever: Akron. Cincinnati.	3 15 87 55 5 97 79 1 33 47 1 45 4 61 49 1 9 007 28 79 81
Central City. Dorchester Elmwood Humboldt Keerney Lincoln Omaha. Seward. University Place York Scattering Syphilis Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montelair. NEW MEXICO. Chancroid Chicken pox. Diphtheria:	7 4 5 6 9 8 9 7 4 39 54 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Trachoma Tryphoid fever Whooping cough 10 Diphtheria: Youngstown Scarlet fever: Akron Cincinnati Youngstown	3 15 87 55 5 97 79 1 33 47 1 45 4 61 49 1 9 07 28 79
Central City. Dorchester Elmwood. Humboldt Keerney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent. Jufluenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chancroid. Chicken pox. Diphtheria: Arroye Hondo.	7 4 5 6 9 8 9 7 4 39 1 1 2 7	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Trachoma Typhoid fever Whooping cough OHIO. Diphtheria: Youn;stown Scarlet fever: Akron Cincinnati Youn;stown.	3 15 87 55 5 97 779 1 333 47 1 45 4 61 49 9 07 28 79 61 333
Central City. Dorchester Elmwood Humboldt Keerney Linceln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chapcroid. Chicken pox. Diphtheria: Arroye Hondo. Scattering.	7 4 5 6 9 8 9 7 4 39 54 1 2 7 9 8 9 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Trachoma Typhoid fever Whooping cough OHIO Diphtheria: Youngstown Scarlet fever: Akron Cincinnati Youngstown Smallpox:	3 15 87 55 5 97 779 1 33 47 1 45 4 61 49 1 9 007 228 79 61 33 31 12
Central City. Dorchester Elmwood Humboldt Keerney Lincoln Omaha. Seward. University Place York Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chancroid. Chicken pox. Diphtheria: Arroye Hondo. Scattering. Gonorrhea.	7 4 5 6 9 8 9 7 4 39 54 1 2 7 7 98 219	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox. Diphtheria. German measles Gonorrhea Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox. 1 Syphilis Trachoma Typhoid fever Whooping cough OHIO. Diphtheria: Youngstown Scarlet fever: A kron Cineinnati Youn gstown Smallpox: Smallpox: Piqua. Port Clinton	3 15 87 55 5 97 779 1 333 47 1 45 4 61 49 9 07 28 79 61 333
Central City. Dorchester Elmwood. Humboldt. Keerney. Lincoln. Omaha. Seward. University Place. York. Scattering. Syphilis. Tuberculosis. Typhoid fever. Whooping cough. NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chancroid. Chicken pox. Diphtheria: Arroye Hondo. Scattering. Gonorrhea.	7 4 5 6 9 8 9 7 4 39 54 1 2 7 9 8 9 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1	NORTH CAROLINA. Cerebrospinal meningitis Chancroid Chicken pox Diphtheria German measles Gonorrhea Measles Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever Septic sore throat Smallpox 1 Syphilis Trachoma Typhoid fever Whooping cough OHIO Diphtheria: Youngstown Scarlet fever: Akron Cincinnati Youngstown Smallpox:	3 15 87 55 5 97 779 1 33 47 1 45 4 61 49 1 9 007 228 79 61 33 31 12
Central City. Dorchester Elmwood Humboldt Keerney Linceln Omaha. Seward. University Place York. Scattering Syphilis. Tuberculosis Typhoid fever Whooping cough NEW JERSEY. Chicken pox, unusually prevalent. Influenza. Measles, unusually prevalent. Pneumonia. Smallpox, reported at Mount Laurel Township and Montclair. NEW MEXICO. Chancroid. Chicken pox. Diphtheria: Arroyo Hondo. Scattering. Gonorrhea. Influenza.	7 4 5 6 9 8 9 7 4 39 54 1 2 7 7 98 219	NORTH CAROLINA. Cerebrospinal meningitis Chancroid. Chicken pox. Diphtheria. German measles. Gonorrhea. Measles. Ophthalmia neonatorum Pneumonia, broncho Pneumonia, lobar Poliomyelitis Scarlet fever. Septic sore throat Smallpox. 1 Syphilis. Truchoma. Typhoid fever. Whooping cough. Diphtheria: Youn;stown. Scarlet fever: A kron. Cincinnati. Youn;stown. Smallpox. Smallpox. Smallpox. Smallpox. Smallpox. Smallpox. Smallpox. Smallpox. Port Clinton. Steubenville.	3 15 87 55 5 97 979 1 33 47 1 45 4 61 49 1 9 07 28 79 133 12 6

VERMONT.		WISCONSIN.	
	BS65.		Cases.
Chicken pox	55	Cerebrospinal meningitus	1
Diphtheria	3	Chicken pox	39
Gonorrhea	21	Diphtheria	36
Measles	55	Measles	
Mumps	165	Scarlet fever	
Scarlet fever	8	Smallpox	. e
Syphilis	11	Tuberculosis	
Typhoid fever	3		
Whooping cough	58	Typhoid fever	39
	•	State (exclusive of Milwaukee):	
VIRGINIA.		Chicken pox	126
Smallpox:		Diphtheria	
Bedford County	6	Erysipelas	
Scattering	4	Influenza—	• -
		Clayton	. 6
WEST VIRGINIA.		Portage	
Diphtheria	16	Scattering	
Measles:		Measles.	
Wheeling	6	Poliomyelitis	
Scattering	3	Scarlet fever	
Scarlet fever	14	Smallpox	
Smallpox:	**	Tuberculosis	
Bluefield	12	Typhoid fever.	
Huntington.	5	Whooping cough	
Scattering	9	Entire State:	, 15
Typhoid fever:	"	Chancroid	
••			
Hinton	6	Gonorrhea	
Scattering	0 1	Syphilis	. 24

SUMMARY OF CASES REPORTED MONTHLY BY STATES.

Tables showing, by counties, the reported cases of cerebrospinal meningitis, malaria, pellagra, poliomyelitis, smallpox, and typhoid fever are published under the names of these diseases. (See names of these and other diseases in the table of contents.)

The following monthly State reports include only those which were received during the current week. These reports appear each week as received.

State.	Cerebro- spinal menin- gitis.	Diph- theria.	Malaria.	Measles.	Pel- lagra.	Polio- mye- litis.	Scarlet fever.	Small- pox.	Ty- phoid fever.
1919. Colorado (November) Fictida (December) Maryland (December) West Virginia (December) Wisconsin (December) Wyoming (November)	4 1 4 7 6 1	64 89 419 352 309 7	222 3	5 4 350 26 492 63	8 2 1	2 2 1 4 3	130 14 342 336 485 42	272 2 4 203 740 6	7 42 82 84 11

ANTHRAX.

East Orange, N. J.-Week Ended Jan. 3, 1920.

During the week ended January 3, 1920, one case of anthrax was reported at East Orange, N. J.

CEREBROSPINAL MENINGITIS.

State Reports for November and December, 1919.

Place.	New cases reported.	Place.	New cases reported.
Colorado (November): Denver El Paso County Total Florida (December): Pinellas County Maryland (December): Baltimore. West Virginia (December): Fayette County Kanawha County	1 4	West Virginia (December)—Con. Mercer County. Tucker County. Total Wisconsin (December): Dane County. Iron County. Manitowoc County. Miwaukee County Total Wyoming (November): Sheridan County.	7

City Reports for Week Ended Jan. 3, 1920.

. Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Akron, Ohio. Baltimore, Md Boston, Mass Chattanooga, Tenn Chicago, III. Chicopee, Mass. Cicero, III. Dallas, Tex. Detroit, Mich. East St. Louis, III. Evanston, III. Haverhill, Mass. Kansas City, Mo Lexington, Ky. Los Angeles, Calif. Lowell, Mass.	1 1 3 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Milwaukee, Wis. Minneapolis, Minn Nashville, Tenn Newark, N. J. New Bedford, Mass. New Haven, Conn New York, N. Y Oshkosh, Wis. Paterson, N. J. Peoria, III. Portland, Oreg. Providence, R. I. Sacramento, Callí. Salt lake City, Utah San Francisco, Calif. Stockton, Calif. Stockton, Calif. Stockton, Calif.	2 2 3 1 2 1 1	

DIPHTHERIA.

See Telegraphic weekly reports from States, p. 188; Monthly summaries by States, p. 192; and Weekly reports from cities, p. 202.

INFLUENZA.

Cases Reported by State Health Officers, Week Ended Jan. 17, 1920.

	Cases.	1 [*]	Cases.
Arkansas	. 53	Louisiana	. 27
California	. 264	Maine	. 4
Connecticut	. 14	Massachusetts	. 54
District of Columbia	. 126	Montana	. 1
Florida	. 10	Nebraska	. 1
Georgia	. 27	New Jersey	. 98
Idaho	270	New Mexico	. 4
Illinois		New York (exclusive of New York City)	. 61
Indiana	. 44	Wisconsin	. 67
Iowa			

LEPROSY.

Christiana, Wis.

During December, 1919, one death from leprosy was reported at Christiana, Wis. The disease was diagnosed clinically and bacteriologically June 3, 1919.

LETHARGIC ENCEPHALITIS.

Topeka, Kans.—Week Ended Jan. 3, 1920.

During the week ended January 3, 1920, one case and one death from lethargic encephalitis were reported at Topeka, Kans.

MALARIA.

Florida and Maryland Reports for December, 1919.

Place.	New cases reported.	Place.	New cases reported.
Florida: Alachua County. Bradford County. Brevard Coun y Calhoun County. Citrus County. Clay County. Columbia County De Soto County Duval County. Jacksonville Escambia County Pensacola. Gadsden County Hernando County Hillsboro County.	1 1 2 2 2 1 4 11 13 2 6 2 4 3	Florida—Continued. Okeechobee County. Orange County. Osceola County. Paim Beach County. Pasco County. Pinellas County. Polk County. Putnam County. St. Johns County. Santa Rosa County. Sumter County. Suwannee County. Taylor County. Volusia County. Wakulla County. Wakulla County.	2 4 1 1 1 4 4 4 3 3 2 2 1 10 4 6
Jackson County Jefferson County Lafayette County Leon County Leon County Liberty County Madison County Manatee County Marion County Marion County Okaloosa County	10 10 7 11 1 3 10	Total	222 2 1 3

City Reports for Week Ended Jan. 3, 1920.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Alexandria, La	47 5 2		Pontiac, MichSavannah, Ga	1	i

MEASLES.

See Telegraphic weekly reports from States, p. 188; Monthly summaries by States, p. 192; and Weekly reports from cities, p. 202.

PELLAGRA.

State Reports for December, 1919.

Place.	New cases reported.	Place.	New cases reported.
Florida: Brevard County Columbia County De Soto County Duval County Gadsden County Oscoola County Santa Rosa County Total	2 1 1 1	Maryland: Allegany County— Frostburg. Frederick County— Brunswick. Total. West Virginia: Lewis County.	1

City Reports for Week Ended Jan. 3, 1920.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Atlanta, Ga Baton Rouge, La. Charleston, S. C. Memphis, Tenn	1	1 1 2	Montgomery, Ala. Nashville, Tenn. Waco, Tex. Washington, D. C.	1	1 4 1

PLAGUE.

Rodent Cases of Plague, New Orleans, La.

During the period from January 10 to 16, inclusive, 33 cases of plague in rodents were confirmed at New Orleans, La.

No human case occurred there between December 30, 1919, and January 16, 1920.

PNEUMONIA (ALL FORMS).

City Reports for Week Ended Jan. 3, 1920.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
lliance, Ohio			Camden, N. J	3	
lton, Ill	. 0	i	Cape Girardeau, Mo Chanute, Kans. Charleston, S. C Charlotte, N. C		j
nn Arbor, Mich	. 2		Charleston, S. C		3
anniston, Alashland, Ky	- 2				1
tlanta, Ga		6	Cheyenne, Wyo	ī	1
urora, III	66	29	Chicago Heights, Ill	255	9
Saltimore, Md		2	Cheyenne, Wyo Chicago Heights, Ill Chicago, Ill Chillicothe, Offio	300	9.
aton Rouge, Lasayonne, N. Jseaumont, Tex	. 1	1			
sayonne, N. J	- 1	2	Cincinnati, Ohio	22	1.
elleville, N. J erkeley, Calif. erlin, N. H.	. 2		Clinton Moss	1	l
erkeley, Calif		2	Coloredo Springs Colo	2	
Avorly Magg	l i	11	Columbus, Ga	5	
illings, Mont		2	Coffeyville, Kans Colorado Springs, Colo. Columbus, Ga. Columbus, Ohio Cortland, N. Y		
Sirmingham, Ala		9	Council Bluffs Iowa	1	
loomfield, N. J		i	Council Bluffs, Iowa Covington, Ky Cranston, R. I.		
oston. Mass	. 21	24	Cranston, R. I	1 16	
razil, Ind ridgeport, Conn	1	. 71	Danville III		
ristol, Conn rockline, Mass uffalo, N. Y	. 1		Dayton, Ohio	2	
rookune, mass	13	13	Denver, Colo Detroit, Mich	27	1. 5
nte. Mont		4	Debith Minn	3.1	
airo, III ambridge, Mass		3	Dunkirk, N. Y	1	

PNEUMONIA (ALL FORMS)—Continued.

City Reports for Week Ended Jan. 3, 1920—Continued.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
East Chicago, Ind East Orange, N. J East St. Louis, III Elizabeth, N. J El Paso, Tex Englewood, N. J Fall River, Mass. Fond du Lac, Wis. Fort Wayne, Ind Fort Worth, Tex Freeport, III. Freeno, Calif. Jalesburg, Ilh. Jalesbu		2	Norfolk, Va	1	
East Orange, N. J.	3	1	North Adams, Mass	1	
East St. Louis, III	1	i	North Attlebore Mass	i	
Elmira, N. Y.	1	1	North Attleboro, Mass Norwalk, Conn Norwood, Ohio Oakland, Calif Oak Park, Ill	-	j
El Paso, Tex		7	Norwood, Ohio	1	1 1
Englewood, N. J		1	Oakland, Calif	2	
Fond du Lee Wis	อ	2	Ogden, Utsh	3	·
Fort Wayne, Ind		Ī	Oak Park, III. Ogden, Utah. Oklahoma City, Okla. Omaha, Nebr. Orange, N. J. Oshkosh, Wis Paducah, Ky. Pasadena, Calif. Passaic, N. J. Paterson, N. J. Peoria, III.		1
Fort Worth, Tex	7	1 7 1	Omaha, Nebr		· :
Freeport, Ill	3	1 1	Oshkosh Wis	3	
Talesburg, Ill		i	Paducah, Ky.	1	
alveston, Tex		4	Pasadena, Calif	4	3
ardner, Mass	•••••	2	Passaic, N. J	1	4
iary, ind			Peoria III	3	
Frand Rapids, Mich	5		Perth Amboy, N. J		
reen Bay, Wis		4	Petersburg, Va		2
dart ford, Conn	•••••••••••••••••••••••••••••••••••••••	6 2	Philadelphia, Pa	84	100
lighland Park, Mich	6	3	Pittsfield, Mass		2 2 2 62 1 3
loboken, N. J.		3 3	Plainfield, N. J	1	ì
lolyoke, Mass	3	5 7	Pontiac, Mich		1
Juntington W. Va		7	Paterson, N. J. Peoria, Ili Perth Amboy, N. J. Petersburg, Va. Philadelphia, Pa. Pittsburg, Kans. Pittsfield, Mass. Plainfield, N. J. Pontiac, Mich. Port Huron, Mich. Port Huron, Oreg. Portsmouth, Va. Poughkeepsie, N. Y. Pueblo, Colo. Quincy, Mass. Racine, Wis. Rahway, N. J. Redlands, Calif. Richmond, Va.	9	4
ndependence, Mo	1	1 3	Portsmouth, Va		i
onwood, Mich	1	1	Poughkeepsie, N. Y	2	
shpeming, Mich	2 2	1	Pueblo, Colo		9
shpeming, Michthaca, N. Yamestown, N. Y	1	. 1	Racine Wis		í
efferson City. Mo		1	Rahway, N. J.		1
efferson City, Moersey City, N. J	9		Redlands, Calif		2
alamazoo, Mich	3 4	3	Richmond, Va	1 2	3 2 1 1 2 7 2 8
ansas City, Kans	21	12	Richmond, Va	8	8
Cearny, N. J	1		Rock Island, Ill	1	1
Cansas City, Kans Cansas City, Mo. Cearny, N. J. Ceene, N. H	2	1	Rock Island, III Rome, Ga. Rome, N. Y Rutland, Vt Sacramento, Calif St. Joseph, Mo. St. Paul, Minn Salt Lake City, Utah San Angelo, Tex San Diego, Calif Sanford, Me.	1	• • • • • • • • • • • • • • • • • • • •
a Fayette, Ind	2	2 1 1 1 1 1	Rome, N. Y	1	•••••
awrence, Kans	i	î	Sacramento, Calif		i
iceno, N. H. & Fayette, Ind. ancaster, Ohio awrence, Kans. exington, Ky incoln, Nebr. ockport, N. Y. ogansport, Ind. ong Beach, Calif. os Angeles, Calif. ouisville, Kv		1	St. Joseph, Mo		6
incoln, Nebr		1	St. Paul, Minn		7 6
ockport, N. 1	9	2	San Angelo, Tex		ĭ
ong Beach, Calif		ī	San Bernardino, Calif		4
os Angeles, Calif	45	18	San Diego, Calif	4	3 1
		3	San Francisco Callf		7
ynchburg, Va		ĭ	Santa Barbara, Calif		1
ynn, Mass	2	1	Savannah, Ga		4
adison, Wis		1	Schenectady, N. I	3	1 3
anchester N H	3	3	Seringfield, Mass	5	
ankato, Minn		ĭ	Springfield, Ohio		2
edford, Mass	3	2	Staunton, Va	••••••	1 1
owell, Mass ynchburg, Va ynn, Mass adison, Wis adison, Wis alden, Mass anchester, N. H ankato, Minn edford, Mass emphis, Tenn ethuen, Mass iddletown, N. Y illwaukce, Wis inneapolis, Minn issoula, Mont ooble, Ala oomnouth, Ill oontgomery, Ala		6	San Bernardino, Calif. San Diego, Calif. San Diego, Calif. Sanford, Me. San Francisco, Calif. Santa Barbara, Calif. Savamah, Ga. Schenectady, N. Y. South Bend, Ind. Springfield, Mass. Springfield, Ohio. Staunton, Va. Stockton, Calif. Syracuse, N. Y. Taunton, Mass. Terre Haute, Ind. Toledo, Ohio. Topeka, Kans. Trenton, N. J. Troy, N. Y. Waco, Tex. Waltham, Mass. Washington, D. C.		6
iddletown, N. Y		i i	Taunton, Mass	i	3
ilwaukce, Wis		14	Terre Haute, Ind		3 1 7 2 5 5 5
inneapolis, Minn		19	Toledo, Ohio		7 9
obile Ala		1 3	Trenton, N. J.	7	5
onmouth, Ill		ĭ∥	Troy, N. Y	1	5
ontclair, N. Jontgomery, Ala	2 .		Waco, Tex		5 1
ontgomery, Ala			Washington, D. C.	2	
ornistown, N. Jount Vernon, N. Y	5	····i	Wausau, Wis		31 1 2 1 2
owark N. I	86	16	Webster, Mass		2
ewark, Ohio		.2	West Hoboken, N. J		1 9
ew Dedicru, Mass	4 2	10	West Orange, N. J.	·····i.	
ewburgh, N. Y	2 2 1	10 2 2 2 1	Wheeling, Va		2
w Castle, Ind	1	1	Wichita, Kans		1
ow Bedford, Mass. ow Britain, Conn. ow burgh, N. Y ow Castle, Ind ow Haven, Conn. ow Orleans, La owton, Mass. ow York, N. Y agara Falls, N. Y		10	Wilmington, Del	2	2
w Orleans, La	2	13	Waltham, Mass. Washington, D. C. Wausau, Wis. Webster, Mass. West Hoboken, N. J. West New York, N. J. West Orange, N. J. Whoeling, Va. Wichita, Kans. Wilmington, Del Winston-Salem, N. C. Yonkers, N. Y. Youngstown, Ohio Zanesville, Ohio.	9	2 1 2 2 5 1 2
w York N V	331	189	Youngstown, Ohio		ĭ
	2		Zanesville, Ohio	1	

POLIOMYELFTIS (INFANTILE PARALYSIS).

State Reports for November and December, 1919.

Plage.	New cases reported.	Place.	New cases reported.
Calorado (November): Denver. Florida (December): Escambia County Hilisbero County Total. Maryland (December): Baltimore.	1 1 2 2	West Virginia (December): Summers County Wisconsin (December): Manitowoc County Milwaukee County Rock County. Total	1 1 1 1 3

City Reports for Week Ended Jan. 3, 1920.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Hackensack, N. J. Manitowoc, Wis St. Cloud, Minn	1 1 1	1	Salt Lake City, Utah Topeka, Kans	i	1

RABIES IN ANIMALS.

Kansas City, Mo., and Newburyport, Mass.

During the week ended January 3, 1920, one case of rabies in animals was reported at Kansas City, Mo., and one at Newburyport, Mass.

SCARLET FEVER.

See Telegraphic weekly reports from States, p. 188; Monthly summaries by States, p. 192; and Weekly reports from cities, p. 202.

SMALLPOX.

Tularosa, N. Mex.

On January 16, 1920, approximately one hundred cases of smallpox were reported at Tularosa, N. Mex.

Vaccinations on Canadian Border, 1920.

Reports from Public Health Service officers on the Canadian border show that vaccinations were performed at points of entry to the United States from January 1 to January 9, inclusive, as follows:

Vaccinations.

Buffalo	493
Detroit	4, 826
Niagara Falls	499
Ogdensburg	108
Port Huron	836
Sault Ste. Marie 1	71

SMALLPOX—Continued. State Reports for November and December, 1919—Vaccination Histories.

		İ	Vaccination history of cases.			
Place.	New cases reported.	Deaths.	Vaccinated within 7 years pre- ceding attack.	Last vacci- nated more than 7 years preceding attack.		History no obtained o uncertain.
Colorado (November):						
Adams County	7		1		5]
Alamosa County	1 9		1		5	
Boulder County	ľi		l		5	l :
Denver County—						
Denver	123		14		103	
Elbert County	i				1	
Frement County	1					
Grand County	1	ļ			1	
Hueriano County— Hueriano	3			1 1	1	
Lake County	ı				i	l
La Plata County	15				15	
Larimer County	63		2		53	
Las Animas County Montrose County	4		. 1		3	
Montezuma County	1				1	
Morgan County Pueblo County	8		3		3	
Pueblo County—					_	
Pueblo Rio Blanco County—	1	• • • • • • • • • • • • • • • • • • • •				1
Rio Blanco	1				1	
Summit County	2				•	
Weld County	27				26]
m. 4.1						
Total	272		22	•••••	220	30
lorida (December):						
Escambia County	2				2	·
faryland (December): Alfegany County— Cumberland	4				4	
Visconsin (December):						
Adams County	13	• • • • • • • • • • • • • • • • • • • •		2	11	
Barron County	6 53	•••••••			5 8] 48
Buffalo County	7				7	796
Calumet County	21		2		1i	8
Chippewa County	2		••••••		1	. 1
Clark County Columbia County	5 2		•••••		5	······i
Crawford County	2			·····i	11	
Dane County	7		6	<u>-</u> .	. .	1
Dodge County	19				11	8
Douglas County Eau Claire County	12 5		2	2	7	1
Fond du Lac County	128				121	5
Grant County	6		2		4	• • • • • • • • • • • • • • • • • • •
Green Lake County	1				1 .	· · · · · · · · · · · · · · · · · · ·
Iowa County	9				9	• • • • • • • • • • • • • • • • • • • •
Jackson County	5 3				1 3	
Kenosha County	3 7					3
La Crosse County					7	
Langlade County Lincoln County		•••••	1		7	25
Manitowoc County	1 21		3		4	•••••••••
Marathon County	22				20	3 2
Marinette County	21 22 3 2 43 17				20 3 2	
Milwankee County	2	······ ·			2 .	•••••••
Monroe County	17		1	•••••••••		43
	69		8		16	48
Oconto County	المما			1	32	30
Oconto CountyOutagamie County	63	· · · · · · · · · · · •				
Portage County	13				2	11
Portage County	13		14			
Portage County	63 13 14 18 4 7		14 1		3 1	11 14 3

SMALLPOX—Continued.

State Reports for November and December, 1919—Vaccination Histories—Contd.

			Vaccination history of cases.			
Place.	New cases reported.	Deaths.	Vaccinated within 7 years pre- ceding attack.	Last vacci- nated more than 7 years preceding attack.	Never suc- cessfully vaccinated.	obtained or
Wisconsin—Continued. Taylor County. Vernon County. Waupaca County. Waushara County. Winnebago County. Wood County. Total.	1 25 22 18 11 17 740		3 1	1 1 1 4	14 8 17 3 13	1 8 12 7 7

West Virginia and Wyoming-November and December, 1919.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
West Virginia (December): Braxton County. Brooke County. Cabell County. Clay County Fayette County Hancock County. Hardy County. Harrison County. Lewis County Lewis County Lincoln County McDowell Marion County Mason County Mercer County Mercer County	2 7 2 16 12 1 20 1 1 1 14		West Virginia (Dec.)—Con. Mingo County Monroe County Ohio County Raleigh County Taylor County Upshur County Wayne County Wood County Total. Wyoming (November): Albany County Sheridan County Uinta County	1 1 9 8 2	

City Reports for Week Ended Jan. 3, 1920.

·					
P.ace.	Cases.	Deaths.	Place.	Cases.	Deaths.
Alameda, Calif	3 1 1 12 1 7 1 6 8 4 1 1 2 4 1 1 1 47 16 16 11 11 12 17 16 17 17 17 17 17 17 17 17 17 17 17 17 17		Erie, Pa. Eugene, Oreg. Everett, Wash Fargo, N. Dak. Fond du Lac, Wis. Galesburg, Ill. Green Bay, Wis. Hammond, Ind. Hoquiam, Wash. Huntington, Ind. Independence, Mo. Indianapolis, Ind. Janesville, Wis. Kalamazoo, Mich. Kansas City, Mo. Kokomo, Ind. La Crosse, Wis. La Fayette, Ind. Lawrence, Kans Lincoln, Nebr. Logansport, Ind. Los Angeles, Calif.	2 4 4 2 1 3 3 2 8 8 6 5 5 2 1 1 8 4 4 4 10 24 1 3 3 2 6	
Dubuque, Iowa	1 1 1		Lincoln, Nebr Logansport, Ind Los Angeles, Calif Madison, Wis	3 2 6 3	

SMALLPOX—Continued. City Reports for Week Ended Jan. 3, 1929—Continued.

Place.	Cases.	Deaths.	Piace.	Cases.	Deaths.
Marion, Ohio. Marshalltown, Iowa. Mason City, Iova. Milwaukee, Wis. Minneapolis, Minn. Monnouth, Ill. Montclair, N. J. Muskogee, Okla. New Orleans, La. Norfolk, Va. Norwoo'l, Ohio. Oaklan'l, Calif. Ogden, Utah. Oklahoma City, Okla.	6 7 9 21 16 3 2 7 3 1 5 30 2		Pueblo, Colo Remo, Nev Riverside, Calif. Rock Island, IM. Sacramento, Calif. St. Cloud, Minn St. Joseph, Mo. St. Louis, Mo. St. Paul, Minn Sal Jose, Calif. San Jose, Calif. Sante Barbara, Calif. Seattle, Wash. Spokane, Wash.	1 7 2 1 7 29 1 4 2 2 5	
Omha, Nebr. Oshkosh, Wis Paducah, Ky Pine Bluff, Ark Pon'iac, Mich Porland, Oreg. Portsmouth, Va.	1 1 1 2		Superior, Wis. Tacoma, Wash. Toledo, Ohio. Walla Walla, Wash. Wichita, Kans. Winston-Salem, N. C. Yakima, Wash.	4 5 1 4 1 1	

TETANUS.

Barberton, Ohio, and Oakland, Calif.

During the week ended January 3, 1920, one case of tetanus was reported at Barberton, Ohio, and one at Oakland, Calif.

TUBERCULOSIS.

See Telegraphic weekly reports from States, p. 188, and Weekly reports from cities, p. 202.

TYPHOID FEVER.

State Reports for November and December, 1919.

Place.	New cases reported.	Place.	New cases reported.
Colorado (November): Archuleta County. Denver. La Plata County. Montrose County. Prowers County. Saguache County. Total. Florida (December): Baker County. Bradford County. Columbia County De Soto County. Duval County. Jacksonville Escambia County Pensacola Flagler County. Gadsen County Hamilton County Hamilton County Tampa Jefferson County Lafayette (ounty	3 1 1 2 2 2 1 1 1 1 1 1 1	Florida (December)—Continued. Putnam Cornty. Santa Rosa County Taylor County. Walton County. Washington County. Total. Maryland (December): Allegany County— Luke. Eckhart. Cumberland. McCool. Ellerslie. Miners' Hospital Baltimore. Baltimore County— Oella. Warren. Caroline County— Federalsburg. Denton. Goldsboro.	2 1 1 1 42 2 2 1 2 1 1 1 21
Lake County	1	Cecil County— Chesapeake City	2

TYPHOID FEVER—Continued.

State Reports for November and December, 1919—Continued.

Place.	New cases reported.	Place.	New cases reported.
Maryland (December)—Continued.		West Virginia (December):	
Charles County—	1	Clay County	. 1
Malcolm	1	Fayette County	1 8
White Plains	1 2	Greenbrier County	[i
Mason Springs	2	Hancock County	l i
Rel Alton	2	Harrison County	
Bel Alton (R. D.)	î	Jackson County	1 7
Dorchester County—	•	Kanawha County	
Airey's (R. D.)		Lewis County.	
Mirey's (R. D.)		T County	1 14
Vienna East New Market	2	Logan County	1 1
East New Market	2	Marion County	1
Cambridge	2 2 2 2	Mason County	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Honga (R. D.)	1	MineralCounty	!
Frederick County-		Mingo County	
Walkersville		Monongalia County	1 4
Woodsboro	1	Monroe County	1 2
Libertytown (R. D.) Lewistown (R. D.)	1	Ohio County	3
Lewistown (R. D.)	1	Pleasants County	1 3
Howard County—		Preston County	1 2
West Elk Řidge	1	Putnam County	1 2
Montgomery County—	_	Raleigh County	1 1
Martinsburg	1	Randolph County	1 3
Prince Georges County-	- 1	Summers County	l š
Formington	1	Tucker County.	i
Farmington. Fairmont Heights	i	Tyler County	
. Queen Annes County—		Wayne County	1
Centerville (R. D.)	. 1	Wetzel County	7
Pandatawa	i	Wood County	1
Pondstown	i	Wood County	
Marydel (R. D.)		Total	84
St. Marys County— Ceder Point.	_	10(81	04
Ceder Point	1		
Mechanicsville	1	Wisconsin (December):	_
Somerset County—	_	Iron County	3
Crisfield	1	Kenosha County	1
Kingston Marion (R. D.)	1	Marinette County	1
Marion (R. D.)	1 2 1	Milwaukee County	
Marion	1	Price County	1
Pocomoke City (R. D.)	2	Racine County	
Princess Anne	1	Winnebago County	1
Marion Station	1	•	
		Total	11
Easton	1		
Washington County—	7	Wyoming (November):	
Washington County— Indian Spring.	1	Converse County.	2
Hagerstown	2	Sheridan County	ī
Hagerstown	- 1	Natrona County	2 1 5
Relighter	1	Johnson County	ī
Wornester County-	*	Washakie County	î
Worcester County— Bishopville	1	Uinta County.	î
		Omea County	-
Total	82	Total	11

City Reports for Week Ended Jan. 3, 1920.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
	0	- 000000			
			a:		
Akron, Ohio	1		Cincinnati, Ohio	1	
Alexandria, Va			Cleveland, Ohio	. 3	
Alliance, Ohio	1		Coffeyville, Kans	1	
Amesbury, Mass	1		Columbus, Ohio	1	
Ashland, Ky	1		Dallas, Tex	1	
Aurora, Ill	1		Decatur, III Durham, N. C	1	
Baltimore, Md	6	1	Durham, N. C	1	
Beaver Falls, Pa	1		El Paso, Tex		
Birmingham, Ala			Erie, Pa	1	
Boston, Mass	1		Fall River, Mass		
Braddock, Pa	4		Fargo, N. Dak	1	
Butler, Pa	1		Fort Wayne, Ind Fort Worth, Tex	2	
Camden, N. J	1		Fort Worth, Tex	1	
Centralia, Ill	3		Galveston, Tex		
Chelsea, Mass	1		Haverhill, Mass		
Chicago Heights, Ill	1	1	Highland Park, Mich	1	
Chicago, Ill	6		Independence, Mo	1 /	

TYPHOID FEVER—Continued. City Reports for Week Ended Jan. 3, 1920—Continued.

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Jamestown, N. Y. Jefferson City, Mo. Kansas City, Mo. La Fayette, Ind. Lawrence, Mass. Lima, Ohlo. Lockport, N. Y. Long Beach, Calif. Los Angeles, Calif. Louisville, Ky. I owell, Mass. Mirneapolis, Minn. Missoula, Mont. Mobile, Ata. Newark, N. J. New Bedford, Mass. New London, Conn. Newton, Mass. New Uondon, Conn. Newton, Mass.	2 2 1 3 1 1 1 2 1 1	1 1 1 1 1	Omaha, Nebr. Oshkosh, Wis. Philadelphia, Pa. Plainfield, N. J. Port Huron, Mich. Portland, Oreg. Quin'y, Mass. Racine, Wis. Reading, Pa. Riverside, Calif. St. Joseph, Mo. St. Paul, Minn. San Angelo, Tex. Sault Ste. Marie, Mich. Spokane, Wash. Springfield, Mo. Ta'oma, Wash. Trenton, N. J. Winston-Salem, N. C.	1 7 2 1 2 1 1 1 1 1 1	1 1 2 1

DIPHTHERIA, MEASLES, SCARLET FEVER, AND TUBERCULOSIS. City Reports for Week Ended Jan. 3, 1920.

•										
 .	Popula- tion as of July 1, 1917	Total deaths	Diph	theria.	Mea	sles.		rlet er.		ber- osis.
City.	(estimated by U. S. (ensus Bureau).	from all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Aberdeen, Wash. Adams, Mass. Akron, Ohio Alameda, Calif. Alexandria, Va Allentown, Pa. Alliance, Ohio Alpena, Mich Alton, ill. Altoona, Pa. Amesbury, Mass. Anaconda, Mont. Ann Arbor, Mich Anniston, Ala Ansonia, Conn Arlington, Mass. Asbury Park, N. J. Ashland, Ky. Ashtabula, Ohio. Atchison, Kans Atlanta, Ga. Atlanta City, N. J. Attleboro, Mass Aurora, Ill. Austin, Tex.	93, 604 28, 433 17, 959 65, 109	1 7 8 2 2 11 10 2 2 2 14 6 3 2 2 57 10 5 2 2 19	26 22 26 5	1	1 24 9 3 7 5 1 119 9 2 2 3 32 32 1		3 3 3 12 1 1 1 1 3 2 10	1	1 3	1 1 1 5 5
Battmore, Md Barberton, Ohio Baton Rouge, La Battle Creek, Mich Bayonne, N. J. Beaumont, Tex Belleville, N. J. Bellingham, Wash Berkeley, Calif Berlin, N. H Beverly, Mass.	35,01,02 594,02 14,187 17,544 30,159 72,201 28,851 12,797 34,362 60,427 15,892 22,128 16,123	12 5 9 12 14 7 5 4	5 11 3	3	1 2 1 1 19	1	32 1 7 3 2 2 2 3		12 1 1 1	2 27
Birmingham, Ala Bloomfield, N. J. Bloomington, Ill.	189, 716 19, 013 27, 462	57 3 8	4 1 2		3	1	2 3 1		2	7

	Popula- tion as of July 1, 1917 (estimated	Total deaths		ntheria	Me	asles.		arlet ver.		nber- losis.
City.	(estimated by U. S. Census Bureau).	from all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Bloomington, Ind	11,661	2	 		6		<u>.</u>		.	
Bluefield, W. Va	16, 123 35, 951 767, 815		·	·		-	1 2		· ·····	· ·····
Boston, Mass	767, 815	253	56	ii	232	i	86	i	35	20
Braddock, Pa	722 (94)	6			34		. 1		·	
Bridgeport, Conn	124, 724	39	14		26	2	4		i	6
Boston, Mass. Braddock, Pa Brasil, Ind. Bridgeport, Conn Bristol, Conn Brookline, Mass. Brunswick, Ga Burfalo, N. Y Burlington, Iowa Burlington, Vt. Butler, Pa. Butte, Mont Cairo, Ill. Cambridge, Mass. Camden, N. J Canton, Ill. Canton, Ill. Canton, Ill. Canton, Ill.	10, 472 124, 724 16, 318 33, 526	3 6	2		5	· ·····	3		1	. i
Brunswick, Ga	10.984	3								
Buffalo, N. Y	475, 781 25, 144	131	68	8	19		19 1	1	11	9
Burlington, Vt	25, 144 21, 802	4	2				1		i	ļī
Butler, Pa	28, 677 44, 057	21	1			·····	7		•••••	4
Cadillac, Mich	10 159	4	3	i		Į		ļ	ļ	l
Cairo, Ill	15, 995 114, 293	14 19	11		1 15		6	·····	5	5 2
Camden, N. J.	108,117		5		iĭ		4		ľ	
Canton, Ill	13, 674 62, 566	3 16	6	i	i		4		4	ļ
Cape Girardeau, Mo	11.146	11	3	<u>.</u>						i
Canton, Ohio Cape Girardeau, Mo. Carbondale, Pa. Carlisle, Pa. Centralia, Ill	19, 597 10, 795	•••••	1				····i	·····	2	
Centralia, Ill	11,838	4	<u>.</u> .							
Chambersburg, Pa	12,475 12,968		ļ		2			·····	·····	····i
Centrains, Inc. Chambersburg, Pa. Chanute, Kans. Charleston, S. C. Charleston, W. Va. Charlotte, N. C.	61,041	4 23	i							i
Charleston, W. Va	31.060	4					i		3	
Chattanooga, Tenn	40, 759 61, 575	14 13	5 4		1				3	3
Chelsea, Mass	48, 405	8	6		. 1		1		1	
Chevenne. Wvo	41,857 111,320	4	3				1		1	
Cheyenne, Wyo. Chicago Heights, Ill. Chicago, Ill.	1 11,320 22,863 2,547,201	7	2		6					
Chicopee, Mass	2, 547, 201 29, 950	715 29	132	14	175	1	277 2	9	169	57
Chicopee, Mass. Chillicothe, Ohio.	15 625	4	2							
Cicero, Ill. Cincinnati, Ohio. Cleveland, Ohio. Clinton, Iowa	20, 846 414, 248 692, 259	129	3 19	1 1	2 48		1 46		14	10
Cleveland, Ohio.	692, 259		63	2	132	1	17		21 .	
Clintop, Iowa	27, 678 1 13, 075	•••••	•••••		•••••		1 2	•••••		····i
Clinton, Iowa Clinton, Mass Coatesville, Pa Coffeyville, Kans Cohoes, N. Y Colorado Springs, Colo Columbia, S. C Columbia, G.	14,995 [.		i		i					
Cohoes N. Y	18, 331 25, 292	5 15	1	j			1	•••••	2 2	4
Colorado Springs, Colo	38, 965	9	i						28	2
Columbia, S. C	35, 165 26, 306	10	····i		•••••		1	•••••	•••••	• • • • • •
Columbus, GaColumbus, Ohio	220, 135	71	4	ī	3 1		8	1	4	6
Concord, N. H.	22, 858 13, 321	13 7	•••••	• • • • • •	1		5 2		•••••	•••••
ouncil Bluffs, Iowa	31,838	12			2		1			
Covington, Ky	59, 623 26, 773	17 6	6 2		8		3	•••••	2	6
Cortiand, N. Y. Council Bluffs, Iowa. Covington, Ky Transton, R. I. Cumberland, Md.	26, 686 129, 738	5	3		1				1	
Danville III	129, 738 32, 969	37 10	14		1		7	1	3	5
Danville, Va	20, 183 .									·····ż
Danville, Ill Danville, Va Dayenport, Iowa Dayton, Ohio	49, 618 . 128, 939 .		2 6				1		···· ₂ ·	
Decatur, III. Dedham, Mass. Denver, Colo. Des Moines, Iowa Detroit, Mich. Dover, N. H. Detroit Be	41,483	7	2		50					•••••
Denver, Colo	10, 618 268, 439	75	11		48		7			····ii
Des Moines, Iowa	104,052 [.		11	2			7		<u></u>	••••
Petroit, Mich	619, 648 13, 276	261 6	93	8	69 1	1	76	4	37	24
Du Bois, Pa.	14,994		i	::::: <u>.</u> .			3			•••••
Du Bois, Pa Dubuque, Iowa Duluth, Minn. Dunkirk, N. Y.	40,096 97,077 21,311	24	2 9	•••••	7 2		5		3	•••••
	01,011	7	•	• • • • • • •	- 1	•••••	5	•••••	-	~

Population April 15, 1910.

City.	July 1, 1917	Total deaths	1 -	htheria.	H 6	asles.	fe	arlet ver.	cul	ber- osis.
· · · · · · · · · · · · · · · · · · ·	(estimated by U. S. Čensus Bureau).	from all causes.		Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Dunmore, Pa	21, 286			<u></u>	1		1	ļ		
Dunmore, Pa. Durham. N. C. Fast Chicago, Ind.	26, 160 30, 286	11 8	1					·····	ļ	3
Fast Cleveland. (hio	13.864	1			7				l	
Fasthampton, Mass	10,656	1					4	1		
Faston, Pa East Crange. N. J	30, 854 43, 761	14	2	• • • • • • • • • • • • • • • • • • • •	5		2		1	i
Fast St. Louis. Ill	77.312	13	2		21		1		17	3
	18, 887 28, 562	7	· ·····		1 1				1	٠٠٠٠٠;
Flizabeth, N. J	88, 830		3		31		9			2 1 1
Flmira, N. Y.	38.272	14			34		11		i	.1
Fl Paso, Tex	69, 149 19, 503	40 8	1	1	16		2 1			12 1
Figlewood, N. J.	12.603	Ĭ	2							
F lyria, Chio Englewood, N. J. Frie. Pa. Fureka, Calif.	76. 592 15. 142	4	16		9		6 2 3 9		1	•••••
	29,304	5 5			2		3			
Formount W Va	40.160 16,111	5	20	1			.9		1.	•••••
Fverett, Mass. Fairmount, W. Va Fall River, Mass. Fargo, N. Dak Farrell, Pa.	129.828	29 7	10	i	8	1	2		3	3
Fargo. N. Dak	17,872 1 10,190	7	1				13			•••••
Findlay, Chio	1 14. 858		1				•••••			•••••
Fond du Lac. Wis	21,486	5								i
Fort Dedge, lows	21, 039 10, 564	1 3					••••••			•••••
Farren, ra Findlay, Chio Ford du Lac. Wis Fort Dedge. Iowa Fort Scott. Kans. Fort Smith, Ark.	29, 390	.					4			•••••
Fo t Wayn., Ind	29, 390 78, 014 109, 597 10, 259	19 39	1 5	1 1	1		8	1	7 2	1
Fosteria, Chio	10,959	2 2	3				···i'	:::::1	2	2
Framingham, Mass	14.159 1	2			1				1	•••••
Fremont, M	19, 844 10, 080	10 5							1	• • • • • •
Fremont. (hio	11,034	5 6								•••••
Fresno, Calif	36, 314 24, 629	12 7	2		1	••••• •				3 1
Galesburg, Ill	42.650	20	2	1						
Gardner, Mass.	17, 534 56, 000	5 13	1 2	i	···i	-	···i		1	1
Gary, Ind	13,915	2			i	:::::Ì.				i
Grand Rapids. Mich. Granite City. Ill. Green Ray. Wis. Greenfeld, Mass.	132 861	2 29 2	4		2		3		1 .	•••••
Green Bay. Wis.	15.810 30.017	12	5	•••••	1		i		1	1
Greenfeld, Mass	12.251	5			1		1			
Greensburg, Pa	15.881 19,594				1 2		1			••••
Hackensack, N. J	17, 412	5			2		2		2	i
dammond, Ind	27, 016 73, 276	6	4		1 .		2 5 1	• • • • • • • • • • • • • • • • • • • •	-	•••••
Harrisburg, Pa. Harrison, N. J	17 345		2				1		, i .	•••••
Hartford, Conn. Haverhill, Mass.	112,831 49,180	26 11	9 2	1 .			20		4 .	••••
	28, 981 1.				8					••••
Highland Park, Mich	33,859	12	11	1	8 .		1 .	-	;- -	••••;
Icland. Mich	78,324 12,459	3	2		9 .		3		1	1 1
Iolyoke, Mass	66, 503	32 58	1		3 .				1	• 4
Hazelouf Fak, Mich. Hoboken, N. J. Holland, Mich. Holyoke, Mass. Houston, Tex. Hudson, N. Y.	116, 878 12, 898	9 !	23	1 -			5 .	····· ·		1 1
Tuntington, Ind	10.982	4 .			i .					···· <u>·</u>
Tuntington, M. Va	47,686 21,461	9	1	-		•••••	1 .			1
ndependence, Mondianapolis, Ind	11,964 283,622	5 .			1:		2 .			· · · · ·
ndianapolis, Ind	283, 622	96 9	10	•••••	7 .		20	1	6	11
ronwood, Mich	15,695 16,710	٠	''i		:::: :		3 .	::::	2	
shpeming, Michthaca, N. Y	16,710 112,448	4.					2 .	-		• • • • •
amestown, N. Y	16,017 37,431	6 .	5		···;· ·		···· -		1 -	

¹ Population Apr. 15, 1919.

CUL	Popula- tion as of July 1, 191 (estimated	Total deaths	1 -	theria	Me	asles.		arlet ver.		iber- losis.
City.	by U. S. Census Bureau).	from all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Janesville, Wis	14,411	6	 		.		. 4			. 1
Janesville, Wis	14,411 13,712 312,557	8	20		ii		6		15	· -
Johnstown, N. Y	10 K78	1	20		1				13	
T12 1/	70, 473 33, 400 50, 408 102, 096		1		64		1	ļ. 		
Walesmanne Mich	50, 408	6 25	2 6 8 9 2	i			1 3		1 1	2
Kansas City, Kans. Kansas City, Mo Kearny, N. J Keene, N. H Knoxville, Tenn Kokomo, Ind	102,096		8	ļ <u>.</u>		<u>.</u> .	4		. 9	l
Kansas City, Mo	305, 816 24, 325 10, 725 59, 112	98	9		44	3	11		11	12
Keene, N. H.	10, 725	2	li		1				i	
Keene, N. H. Kokomo, Ind. Lackawanna, N. Y. La Crosse, Wis. La Fayette, Ind. Lancaster, Ohio. Lancaster, Pa. La Salle, Ill. Laurel, Miss. Lawrence, Kans. Lawrence, Kans. Leavenworth, Kans. Leoninster, Mass. Lecanger, Chio. Lincoln, Nebr. Little Rock, Ark. Lockport, N. Y. Locansport, Ind. Long Beach, Calif.	59,112		1	1	13		7		4	4
Lackawanna N. Y	21, 929 16, 219	9	5					·····	2 3	1
La Crosse, Wis	31,833	3 3	ļ							ļ <u>.</u>
La Fayette, Ind	21,481	4		ļ	1	ļ	3	ļ		ļ
Lancaster, Unio	16,086 51,437	9	5	·····			l····i	·····	····i	ļ
La Salle, Ill	51, 437 12, 332	3					ļ . .			
Laurel, Miss	12.313	·····i	1	1				ļ		
Lawrence, Mass	13,477 102,923 1 19,363	26	5	·····2	i		12		3	3
Leavenworth, Kans	1 19, 363	4							3	ĭ
Leominster, Mass	21.365	8	1				3 2 1 2		1	i
Lima. Ohio	41,997 37,145 46,957	21 8			21		i i		4	1
Lincoln, Nebr	46, 957	13	i				2		i	3
Little Rock, Ark	58,716	······	4				2		···· <u>·</u>	
Lockport, N. I	20,028 21,338	6 16			5		7		1	
Long Beach, Calif	29, 163	24					3			i
Long Branch, N. J	29, 163 15, 733 38, 266	1			17				ļ <u>.</u> .	
Loram, Unio	535, 485	184	34	i	10		g	•••••	79	22
Louisville, Ky	240,808	67	25	l [.]			8 12		8	5
Lockport, N Y Logansport, Ind. Long Beach, Calif Long Branch, N. J Lorain, Ohio Los Angeles, Calif Louisville, Ky Lowell, Mass.	114,366 10,566	31	5	1	2		12	i	8	2
Ludington, Mich Lynchburg, Va Lynn, Mass McKeesport, Pa Madison, Wis Mahanoy City, Pa. Malden, Mass	33, 497	2 5		•••••		•••••	1	•••••	i	
Lynn, Mass	33, 497 101, 534	24	9	3			41	1	2	i
McKeesport, Pa	48, 299 31, 315 17, 709	4			1				Ī	
Mahanov City. Pa	17, 709	2	1	•••••	2 1	•••••	3			
Malden, Mass	52, 243	11	4	1	i		5		2 1	i
Manchester, Conn	15,859	3 21	2 2				1 1		1	
Manitowoc. Wis	79, 607 13, 931 1 10, 365 1 14, 610	5	2		•••••	• • • • • • •	3			
Mankato, Minn	1 10, 365	5 3								
Malden, Mass Manchester, Conn. Manchester, N. H. Manitowoe, Wis. Mankato, Minn Marinette, Wis. Marion, Ind Marion, Ohio.	1 14,610	····i	····2		····i	•••••	2 1	• • • • • •	····i	
Marion, Ohio	19, 923 24, 129		4				il			
Marquette, Mich	12,555	2								
Marshalltown, lowa	14,519 12,984	•••••	3 4				····i	•••••	• • • • • •	•••••
Martins Ferry, Ohio	10, 135					l	il			
Mason City, Iowa	14 038	5			31					•••••
Mattoon, Ill	12,764 26,681 17,724		····i		11		1		····i	
Melrose, Mass	17, 724	4					8		i	
Memphis, Tenn	151.877 I	19	25 3	1			3		9	2
Marion, Ind Marquette, Mich Marquette, Mich Marshalltown, Iowa Martinsburg, W. Va. Martins Ferry, Ohio. Mason City, Iowa Mattoon, Ili Mediord, Mass Melrose, Mass Melrose, Mass Memphis, Tenn Mertden, Conn Methnen, Mass. Middletown, N. Y. Middletown, N. Y. Middletown, Ohio. Milwaukee, Wis Mimeapolis, Minn Mishawaka, Ind Missoula, Mont Mothe, Mas Mont Missoula, Mont Mothe, Ala	29, 431 14, 320 15, 890	•••••	3 2	•••••	•••••	•••••	11 5	••••••	1	i
Middletown, N. Y.	15.890		î				1			
fiddletown, Ohio	16,384	3			3		1		2	
finneanolis Minr	445, 008 373, 448	101 84	34 12	1	18	•••••	29	1	20 29	6
(ishawaka, Ind.	373,448 17,083 19,075	2			<u>°</u>					9
dissoula, Mont	19,075	4 .					ا.ي		1	1
Cobile, Ala	59, 201	31	5		12	•••••	2		1	•••••
fonessen Pa	23 1170 1									
fonessen, Pa fonmouth, Ill fontclair, N. J.	23,070 10,346 27,087	4					5		i	

¹ Population, Apr. 15, 1910.

	i	1	T				1		Ŧ	
ov.	Popula- tion as of July 1, 1917 (estimated	Total deaths		theria	Mea	sles.	Sc.	arlet ver.	Tu cu	iber- losis.
Cit y .	by U.S. Census	from all causes.	. s	Desths.	88.	Deaths.	88.	Deaths.	es.	Deaths.
	Buresu).		Cases.	Å	Cases.	Ã	Cases.	Ã	Cases.	Ã
Mon'gomery, Ala Morgantown, W. Va Morristown, N. J. Moun'sville, W. Va. Mount Carmel, Pa Mount Vernon, III Muscatine, Iova Muskoree, Okla Nanticoke, Pa Nashua, N. H. Nashville, Tenn. Newark, N. J. Newark, N. J. Newark, N. J. Newark, N. J. New Britain, Conn. New Britain, Conn. New Brunswick, N. J. Newburry, N. Y. New Castle, Ini. New Castle, Pa. New Haven, Conn. New Orleans, La. Newyort, R. L. Newton, Mass. New York, N. Y. Niacara Falls, N. Y. Norolk, Va.	44, 039 14, 444	27	3		ļ		2 6	ļ	ļ	ļ
Morristown, N. J.	13 410	9	1				l			
Moundsville, W. Va	11,513	2					ļ			
Mount Carmel, Pa	11,513 20,709 10,043				···-		1			ļ <u>.</u>
Mount Vernou, III	17,713	13 4	1 *		23		5			1 1
Muskoree, Okla	47 173 1		i						1	
Nanticoké, Pa	23,811 27,541 118,136 418,789	• • • • • • • • • • • • • • • • • • • •	2				1			
Nashua, N. H	27,541	4 47			2		3 6		····-	·····
Newark N. J	418, 789	116	18		106	1	19		5 41	11
Newark, Ohio	481.517 (7	1				\ 1		2	
New Berford, Mass	121,622 55,3%5	30	7 3	2	60		4		5	2
New Britam, Conn	25, 855	12		1	2		1		3	
Newburgh, N. Y.	29 803	15	2			1	l	1	2	ı i
Newburyport, Mass	15, 291	4			1			 		
New Castle, In 1	14, 144	2	•••••		6 2		5			
New Haven Conn	14, 144 41, 915 152, 275 21, 199	40	14	i	68	i	8	1	8	4
New London, Conn	21, 199	•••••	2				1			
New Orleans, La		138	16	1			7		15	15
Newton Wass	30, 585 44, 345 5, 737, 492	8 1 2	4	• • • • • • • • • • • • • • • • • • • •	1		5		1	1
New York, N. Y	5,737,492	1,401	320	38	1,246	17	129		196	127
Niagara Falls, N. Y	39, 466 91, 148	· 11	2		42		6		4	2
Norristown, Pa	31, 969	•••••	4 3		····i		1	•••••		
North Adams, Mass	31,969 1 22,019 20,006	8								
North Adams, Mass. North Adams, Mass. Northampton, Mass. North Braddock, Pa. North Braddock, Pa. North Tonavania, N. Y. Norwalk, Conn. Norwich, Conn. Norwich, Conn. Norwood, Ohio. Oaklan ¹ , Calif. Oak Park, Ill. Ogdensburg, N. Y. Ogdeng, Utah. Oil City, Pa.	20,006	13					1			
North Braddock, Pa	11, 248 15, 684	3			4	•••••	•••••	•••••	2	2
North Tona van la, N. Y	14,060	4	1				5	i		
Norwalk, Conn	14,060 27,312 21,923	6 3 5	····i		1		1			1
Norwood Ohio	21,923	3 5	2	1	1 13	•••••	1	•••••		
Oaklan , Calif	000 400	60	3		73		47			
Oak Park, Ill	27, 816 16, 845 32, 343 2°, 162 97, 588	11			2		7			1
Orden Utah	32 343	7 10	····i		•••••		2	•••••		
Oil City, Pa	20, 162		i		i					
Oklahoma City, Okla	97,588	12	6				i		1	
Omaha Nahr	16,927	5 47		•••••	····i		28			
Orange, N. J.	33,636	12	8 1				5	2	1	2
ogdem, Utan. Oil City, Pa Okiahoma City, Okla. Olean, N. Y. Omaha, Nebr. Orange, N. J. Oshkosh, Wis. Parkersburg, W. Va. Parsons, Kans. Passadema Calif	16,927 177,777 33,636 36,549	. 8	1		2 1		2		2 1	
Porkarchurg W Va	25, 178 21, 059	9	4	i	1				1	•••••
Parsons, Kans.	15,952						i			
Parsons, Kans. Passadena, Calif. Passade, N. J. Paterson, N. J. Pawtucket, R. I. Peekskill, N. Y. Pekin, III.	15, 952 49, 620				1		4		. 1	3
Passaic, N. J	74, 478 140, 512	23	10	1	1 5	•••••	3		2	. 1
Pawtucket, R. I	69,666 1	15	2				7			•••••
Peekskill, N. Y	19,034 10,973	4								1
Pekin, III	10,973 . 72,184	12	1 4		····i		1 6	i'	···· <u>·</u> -	i
Perth Amboy, N. J.	42 R4R 1	7	il				4		5	i
Petersburg, Va	25,817	9	2						2 55	
Peoria, III. Perth Amboy, N. J. Petersburg, Va. Philadelphia, Pa. Phillipsburg, N. J. Pine Bluff, Ark. Pitteburgh Pa	25,817 1,735,514 15,879 17,777	526 5	88	9	130	······	75	1	55	46 2
Pine Bluff, Ark	17,777								···i'l	
Pittsburg, Pa. Pittsburg, Kans. Pittsfield, Mass. P ainfield, N. J.			16		80		2		8	
Pittsburg, Kans	18, 340 (2]					•••••	
P ainfield, N. J.	39,678 24,330 13,111	5	···i		28		···i			
P attsburg, N. Y Plymouth, Mass	13,111	5								
Pontice Mich	14,001 18,006	13 5 5 3 6 7			20-		;-		;:	····i
P.rt Chester, N. Y.	16,727	7	2 2 3		39 1		1		1	
P.rt Chester, N. Y	16,727 118,863	9	3	. 1	95	1				

¹ Population Apr. 15, 1910.

	Popula- tion as of July 1, 1917	Total deaths	Diph	theria	Me	asles.	Ser	arlet ver.	Tu	ber- osis.
City.	(estimated by U. S. Census Bureau).	from all causes.	Сазея.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Port'and, Me	64,720 308,399	29	ļ _. .	ļ <u>.</u> .	<u>.</u>	ļ	8		ļ <u>.</u> .	1 3
Port'and, Me	308,309 11,730	69	3	1	6		10		3 3	i
Prismcuth, Va	40,003	15	3 1 1				2 1		3 1	i
P ttst wn, Pa. P tts ille, Pa.	16,987 22,717 30,786	·····	3	 				l:::::		l::::::
P ughkeepsia, N. Y	30,786 250,805	7 61	3 2 20 20 2	3	3		18			1 6
P tts il'e, Pa. P ughkeepsie, N. Y. Providence, R. I. Pueble, Colo.	56,084	1	2		ļ		1 7			ľ
Quincy, Mass Racine, Wis Rahway, N. J	30 (122	6 12	1 2	ļ	6		7		1	3
Rahway, N. J	47, 465 10, 361 111, 607	l i								
Reading, Pa. Radlands, Ca if	111,607 14,573	4	4	·····	····i	·····	1		···· <u>·</u>	·····i
Dadwing Minn	10.158	[3	 	1			
Renr., Nev Richmend, Ind Richmend, Va Ri erside, Calif Roanoke, Va R chester, N. Y R ckf rd, Ill	15.514	8 5	4		4		2	<u> </u>	1 1	1
Richmond, Va	25, 080 158, 702	50	3	1	6		12		13	6
Ri erside, Calif	46 292	9 2	1	1			···· ₂ ·		i	1
R chester, N. Y	264,714	65	20	ï	93	1.	7	i	2	5
R ck Is and, Ill		12 7	····i		····i		6		····i	4
R cky Mount, N. C	29, 452 12, 673 15, 607	6	ļ <u>-</u>				ļ		1	
R me, Ga	15,607 24,759		5		•••••		i		6 2	
Rut and, Vt	TR LEOK	5								
Sacrament, Ca if	68,984 12 013	17 5	3	<u>-</u> i	5		2		6	1
St. Joseph, Mo	68,584 12,013 84,498 768,630	45	1	1			1			5
R ck Is and, III. R cky Mount, N. C. R me, Ga R me, N. Y. Sacrament, Ca if St. Cloud, Minn. St. Jeseph, Mo. St. Lusis, Mo. St. Lusis, Mo.	768, 630	210 48	100 5	11	180		30 9		32 31	5 9 4
Salem, Mass	252, 465 49, 346 121, 623 10, 321 17, 616	12	14	1			ő			
Salt Lake City, Utah	121,673	34 22	1	•••••	• • • • •			····i	•••••	4 11
Salem, Mass Salt Lake City, Utah San Angelo, Tex San Bernardino, Calif	17, 616	22 11 33								2
San Diego, Ca if	56, 412 20, 276	33 1	1		•••••		1	•••••	5	4
Sanf rd, Me	11,217	4			11					
San Francisco, Calif	11,217 471,023 30,810	151	•••••	•••••	7	1	·····i′		····i	12 1
Santa Barbara, ('alif	10.00	8								ī
Sarat ga Springs, N. Y	13, 839 10, 210	8 3 0	•••••			•••••	•••••	•••••	•••••	
Sau t Ste. Marie, Mich	14, 130				i		2			
Savannah, Ga	14,130 69,250 103,774	49 10	2 5		••••2		5 1 7		3 1	2 1
Scrant n, Pa	149.541		4		7				2	
Seattle, Wash	366 445 1		8	•••••	90 11		11		····i	· · · · · · ·
Shenand ah, Pa	21,274 29,753 58,568				•••••		1			
Sioux Falls, S. Dak	58, 568 16, 887	4	1				2 8 3	1		
Somervil'e, Mass	88,618	23	4		20		3 2		6	•••••
South Bend, Ind	70, 967 14, 465	16 2	3							
Spokane, Wash	14, 465 157, 656 108, 668		2		1		2 5		···· <u>è</u> ·	
Sau't Ste. Marie, Mich Savannah, Ga. Schenectady, N. Y Scrant n, Pa. Seattle, Wash Shamokin, Pa. Shenand ah, Pa. Sicux City, Iowa Sicux Gity, Iowa Sioux Falls, S. Dak Somervil'e, Mass South Bend, Ind Southbridge, Mass Springfield, Mass Springfield, Mos. Springfield, Mo. Springfield, Mo.	108, 668 41, 169	31 10	4				3			1
Springfield, Ohio	52 296	10								• • • • • •
Springfield, Mo. Springfield, Ohlo. Staunt n, Va. Steelt n, Pa. Stillwater, Minn. St ett n, Calif.	11,8°3 15,759 10,198 36,209	7			:::::				2	· · • • · ·
Stillwater, Minn	1 10, 198	3			3		····i	····• ·		i
Sunbury, Pa	36, 209 16, 661	20	····2			:::::]	1			
Superior, Wis	47.167	7	1		4		4 9	···i	3	1
Tac ma. Wash	158,559 117,446	41	8 7		1 2	:::::	3 1			
Sunbury, Pa. Superior, Wis. Syracuse, N. Y. Tac ma, Wash. Taunton, Mass.	36,610	13	1				1 2	•••••	1	2 2
Terre Haute, Ind	67,361	20					2		1	2

¹ Population Apr. 15, 1910.

	Popula- tion as of July 1, 1917	Total deaths	Diph	theria.	Mea	sles.	Scar fev	let er.		ber- osis.
City.	(estimated by U. S. Census Bureau).	U.S. all causes.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Toledo, Ohio Topeka, Kans Trenton, N. J Troy, N. Y Tucson, Aris Vancouver, Wash Walla Walla, Wash Wa'tham, Mass Warren, Pa Washington, D. C Watertown, Mass Watertown, Mass Watertown, Mass West Chester, Pa West Hoboken, N. J West Hoboken, N. J West Hoboken, N. J West Orange, N. J Wheeling, W. Va Wikinsburgh, Pa Wilkinsburgh, Pa Wilkinsburgh, Pa Williamsport, Pa Williamsport, Pa Williamsport, Pa Williamsport, Pa Williamsport, Minn Winthrop, Mass Woburn, Mass Woburn, Mass Woburn, Mass Yonkers, N. Y York, Pa Yornestown, Ohio	113, 974 78, 094 17, 324 13, 805 34, 015 26, 067 31, 011 15, 083	15 148 148 15 0 2 - 4 18 18 10 11 17 25 21 18	7 65 15 2 2 19 6 6 1 3 2 2 1 1 1 3 3 9 1 1 1 1 4 4 1 5 6 6 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 1 1 1 1	203 3 7 1 10 10 10 3 2 2 2 2 4 8 8	1	24 15 2 25 3 3 3 4 4 4 1 1 4 1 1 4 1 1 5	1	3 4 1 1 7 7 4 2 3 3 1	111 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

¹ Population Apr. 15, 1910.

FOREIGN AND INSULAR.

CUBA.

Communicable Diseases-Habana.

Communicable diseases have been notified at Habana as follows:

	Dec. 11-	-20, 1919.	Remain- ing under
Disease.	Cases.	Deaths.	Dec. 20, 1919.
Broncho-pneumonia	1		1
Chicken pox	.] 		2
Diphtheria	. 2		1
Infliten &			.2
eprosy			19 174
Malaria. Measies			22
Measies			1
Scarlet fever.			3
Smallpox.	1 .		2 8
Typhoid fever	11		341

¹ From the interior, 26.

INFLUENZA.

The following information was taken from reports received during the week ended January 23, 1920:

P ace.	Date.	Cases.	Deaths.	Remarks.
Brazil: Pahia	Oct. 26-Nov. 1	2	2	
British Celumbia— Vancouver	Dec. 21-27	1		
France: Paris	Nov. 30-Dec. 6		5	
Great Britain: London	Nov. 30-Dec. 13		58	Greater London, including Outer Ring and 14 large towns.
Spain: Madrid	Nov. 1-30		36	Tring and I I large Company
Switzer and: Zurich	Nov. 30-Dec. 6	2		·

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER.

Reports Received During Week Ended Jan. 23, 1920.1

CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.
India: Calcutta	Nov. 9-15	37	35	

¹ From medical officers of the Public Health Service, American consuls and other sources.

From the interior, 1. From the interior, 15.

Reports Received During Week Ended Jan. 23, 1920—Continued.

PLAGUE.

Place.	Date.	Cases.	Deaths.	Remarks.
Chile: Antofagasta	Dec. 8-14	1		
Peru:		l		
Salaverry (Trujillo) Svria:	Dec. 1-14	5		
Beirut	Dec. 22	29		
Turkey: Constantinople	Dec. 11	ļ	·	Present.
	SMAL	LPOX.		•
Bolivia:				
La Paz	Nov. 23-Dec. 13	25	32	
Bahia Sanada:	Oct. 26-Nov. 8	736	444	**
Ontario				Dec. 28, 1919-Jan. 3, 1920; Case 272. In 34 localities.
Gloucester County Hami'ton	OctNov Jan. 4-10	3 2		ZIZ. III 34 IOCANTIES.
Ottawa	Dec. 28-Jan. 3	1		
Prescott	Jan. 4-10 Dec. 28-Jan. 3	1		
Toronto	do	182		
Saskatchewan— Moosejaw	do	1		
Thina: Canton Chungking	Nov. 23-29 Nov. 16-22			Present.
Foochow.	do			Do.
ndia: Calcutta	Nov. 9-15	14	12	
taly: Genoa	Nov. 24-30	7		
Yewfoundland: St. Johns	Dec. 27-Jan. 2	1		Outports, 4 cases. Present at
ortugal: Lisbon	Nov. 23-Dec. 6		18	other localities.
pain:				
BarcelonaBilbao	Nov. 20-Dec. 9 Nov. 21-30		12 3	
	TYPHUS	FEVE	R.	
Solivia:				
La Paz	Nov. 23-Dec. 13	5	5	
Antofagasta	Dec. 8-14	4		
SantiagoValparaiso	Nov. 24-30	500	27	Jan. 12-Sept. 30, 1919: Cases 5,153; deaths, 1,023. Outbreak in October, 1918.
	Dec. 1-13	700	18	in October, 1918.
eru: Cerro de Pasco	Dec. 7-13	1		
Barcelona	Nov. 20-26	2		
urkey: Constantinople	Nov. 30-Dec. 6	7		
<u>-</u>	YELLOW	FEVE	R.	
razil:	: 1			
Bahia	Oct. 26-Nov. 8	1	2	i

Reports Received from Dec. 27, 1919, to Jan. 16, 1920.1

CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.
China:				
Amoy	Nov. 4-17	ı	. 2	
Chosen (Korea)			-	Oct. 20-Nov. 16, 1919: Cases
Chemulpo	Oct. 1-31	6	4	3,525; deaths, 3,144. Aug. 15-
Fusan	do	34	30	Nov. 16 1010: Cocco 15 100
Provinces—	J	34	30	Nov. 16, 1919: Cases, 15, 192
Provinces— Kelti Kogen Ko'si. North Chusei North Heian North Kankyo North Keisho North Kenses South Chusei	A 15 No. 10			deaths, 9,823.
Kelvi	Aug. 15-Nov. 16	224	135	
Kogen]do	64	38	1
Ko al	do	4,015	2,770	ĺ
North Chusei	do	1	1	
North Heian	ldo	3,196	2,434	ĺ
North Kankvo	do	497	275	
North Keisho	do	63	35	l
North Zenra	do	1,326	692	
South Chusei	do	930	590	l
• South Union	do			· ·
South Heian	qo	3,031	1,858	
South Kan'tyo	ao	870	551	
South Keisno	do	318	156	
South Zenra	do	657	288	
Proces				
Saloniki	Oct. 10	1		
ndia:	000.10	•		
	Mary 0.0			
Bombay	Nov. 2-8	1	1	
Calcutta	Oct. 26-Nov. 8	23	23	
apan:				
Kobe	Nov. 24-30	2		
Taiwan				For entire island: Oct. 22-Nov
Tokyo.	Nov. 10-20	i	1	20, 1919: Cases, 645; deaths, 374
ava:	1101. 10-20		-	20, 1515. Cases, 040, deaths, 374
Fast Java				Oat 5 11 1010: One area 1 death
hilimping Inlands		• • • • • • • • •		Oct. 5-11, 1919: One case, 1 death.
hilippine Islands:	÷=	1	_	At Pasoeroean.
Manila	Nov. 2-29	12	7	
Provinces				Nov. 2-29, 1919: Cases, 1,153;
Albay	Nov. 2-29	307	212	deaths, 843.
Ambos Camarines	Nov. 2-22	39	20	
Antique	Nov. 2-29	18	10	
Batangas	do	34	24	
Pohol	Nov 0 00			
Bohol	Nov. 2-22	19	16	
Cagayan	Nov. 3-15	35	20	•
Capiz	Nov. 2-8	6 [5	
Cavite	Nov. 2-29	12	13	
Cebu	do	21	13	
Davao	Nov. 9-15	6	4	•
Ilocos Norte	Nov. 2-29	42	40	
Ilocos Sur	Nov. 2-22	18		
Theile	Nov. 2-22		15	
Iloilo	Nov. 2-29	38	22	
Isabela	Nov. 2-15	111	40	
Laguna	Nov 9_90 I	17	11	
Mindoro. Mountain Occidental Negros	do	76	28	
Mountain	Nov 2-22	6	4	
Ossidental Negros	Nov. 2 22	65	39	
Den marinen	NOV. 2-29			
1 2015 2011 2011	uo	29	19	
Rizal	do	36	14	
Sorsogon	do	161	106	
Tarlac	do	11	11	
Tavahas	do	38	28	
Tarlac Tayabas Union	Nov. 9-15	5	5	
oland:	1101. 5-10	3	9	
Commolin		i	1	December Manageber 1010
Garwolin		·		Present in November, 1919.
Kowai				Do.
Stryi		1.		Do.
iam:				
	Oct. 5-18	i	4	
traits Settlements:	001. 0-10		*	
	1			
Singapore	Oct. 5-Nov. 1	13	12	

¹ From medical officers of the Public Health Service, American consuls, and other sources. For reports received from June 28 to Dec. 26, 1919, see Public Health Reports for Dec. 26, 1919. The tables of epidemic diseases are terminated semiannually and new tables begun.

Reports Received from Dec. 27, 1919, to Jan. 16, 1929—Continued.

PLAGUE.

Place.	Date.	Cases.	Deaths.	Remarks.
British East Africa: Kisumu	Sept. 28-Nov. 1	. 6	6	
Ceylon: Colombo Ecuador:	Oct. 26-Nov. 8	. 14	14	,
Guayaquil Egypt	Nov. 1-30	1		Jan. 1-Nov. 20, 1919: Cases, 4,070;
Province—				deaths, 3,044.
AssioutGreece:	Nov. 15-20	15	8	
SalonikiIndia	Oct. 6-Nov. 30		7	Oct. 19-Nov. 8, 1919: Cases, 5,771;
Bombay Karachi	Oct. 19-Nov. 18 Nov. 9-15	3	3	deaths, 4, 382.
Madras Presidency Rangoon	Nov. 9-22 Nov. 2-8	329 5	207 3	Oct. 19-Nov. 1, 1919: Cases, 10; deaths, 7.
Java: East Java				Sept. 28-Oct. 18, 1919: Cases, 121;
Peru:				deaths, 121.
Salaverry Straits Settlements:	Nov. 23-29	.3	1	
Singapore Turkey:	Oct. 26-Nov. 2	1	1	,
Constantinople On vessel:	Nov. 14-20 Nov. 28	10		At Port Said Fount From
S. S. Kaisar-i-Hind	1404. 20	3		At Port Said, Egypt. From Bombay, Nov. 15, for London.
	QM AT	LPOX.		
	SWAI	AAFUA.		
Algeria: Departments—			•	
Algiers	Nov. 11-Dec. 11do	34 13		,
Oran	do	55		
Bolivia: La Paz	June 29-Nov. 28	<u>.</u>	174	Dec. 29-June 28, 1919: Cases, 86; deaths, 44.
Canada: British Columbia—				,
Vancouver Nova Scotia—	Nov. 30-Dec. 6	1		
Sydney Halifax	Dec. 7-13 Dec. 21-27	1 2		
Counties— Cumberland	Dec. 14-20			Present.
Inverness Pictou.	do			Do. Do.
Ontario				Nov. 30-Dec. 6, 1919: Cases, 125. In 45 localities, exclusive of
Hamilton Kingston	Dec. 14-20 Dec. 21-27	3 1		Dysart and Toronto.
Ottawa	Dec. 14-20	i		Dysair and Toronto.
Peterborough	Dec. 7-27	4		
Sault Ste. Marie	do Dec. 7-27	727		
Toronto Windsor	Dec. 14-27	2		
Quebec-				•
Montreal	Dec. 7-27	3 4		
QuebecSaskatchewan— Saskatoon.	Dec. 14-20	1		
China:	Nov. 4-10	•		Present.
Amoy Canton	Nov. 2-29			Do.
Chungking Chosen (Korea):	Nov. 2-15			Do.
FusanSeoul	Oct. 1-31do	2 9	1	
Colombia: Barranquilla		50	1	
				•

Reports Received from Dec. 27, 1919, to Jan. 16, 1920—Continued.

SMALLPOX-Continued.

Place.	Date.	Cases.	Deaths.	Remarks.
Egypt:				
Alexandria	Nov. 12-Dec. 2	. 17	9	
Cairo	Oct. 1-14	18		
Port Said	Oct. 1-7	i		
Finland:	l .	İ		
Provinces	l	i	1	
Nyland Tavastehus	. July 16-31	1	1	
Tavastehus	do.:	. 1		
Viborg	do	23	1	
Finland			.	. Oct. 15-31, 1919: Cases, 6.
Provinces—	1	l	ļ	
Nyland	Oct. 15-31	4		Helsingfors.
Tavastehus	do	1		Rural district.
Viborg		1		. Do.
Finland.			•	Nov. 1-15, 1919: Cases, 27.
Provinces—	N 1 15	١.		1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
Abo Och Borneborg	Nov. 1-15	1		Rural district.
Nyland	qo	16		Helsingfors.
St. Michael	do	5		Rural district.
St. Michael	do	1		Do.
Germany	ao	4		Do.
Germany		•••••		Oct. 5-11, 1919; Cases, 32. In ad-
		i	:	Oct. 5-11, 1919: Cases, 32. In addition to previously reported cases, Sept. 28-Oct. 4, 1919: Cases, 26.
	I	l	I	Cases, Sept. 28-Oct. 4, 1919;
Greece:	1		l	Cases, 20.
Saloniki	Nov. 10-22	9		İ
India:	NOV. 10-22	9	8	i
Bombay	Oct 19 Nov 9	10	3	
Calcutta.	Oct. 12-Nov. 8	11		
Madras	Oct. 12-Nov. 8 Oct. 26-Nov. 8 Nov. 2-13	10	8 7	
Rangoon	Oct. 19-Nov. 8	21	l ii	
Italy:	Oct. 15-110V. 8	21	1 11	
Genoa	Nov. 16-22	5		
Messina.	Nov. 10-30.	10	6	
Japan:	2.01. 10 00	10	1	
Taiwan	Nov. 1-20	10	2	Entire island.
Java:			_	Danie India.
East Java			1	Sept. 28-Oct. 18, 1919: Cases, 12
West Java				Oct. 17-29, 1919: Cases, 117.
			1	deaths, 31.
Batavia	Oct. 17-29	11	3	,
Mexico:				
Acapulco	Nov. 9-15	2		
Mexico City	Nov. 16-Dec. 6	8		
Mexico City	Dec. 14-20.		1	
Chihuahua	Dec. 21-27	2		
Newfounaland:				
St. John's	Dec. 20-26	3		Dec. 13-26, at outports, 6 cases.
				Present at 8 other localities.
Panama:				
_ Colon	Dec. 15-21	1		
Portugal:				
Lisbon	Dec. 14-20		15	
Spain:				
Barcelona	Nov. 6-12		2	
Cadiz	Oct. 1-31		5	
Val. ncia	Nov. 10-Dec. 20	33	. 8	
Vigo	Nov. 18-Dec. 13	• • • • • • • •	14	
Funis:	Dec 00 00		i	
Tunis	Dec. 23-29	1		
Furkey:	37 0.15	ا ہ		
Constantinople	Nov. 9-15	5		
Union of South Africa:	04 1 91		1	
Johannesburg	Oct. 1-31	10		
S. S. Sarcoxie	Dog 22	1	1	At Poute Delgade Azores From
D. D. Dan CUAIC	Dec. 23	- 1	•••••	At Ponta Delgada, Azores. From Rotterdam for New York.
i			i	TOURCHAMM TOURS TOUR.

Reports Received from Dec. 27, 1919, to Jan. 16, 1920—Continued. TYPHUS FEVER.

Place.	Date.	Cases.	Deaths.	Remarks.
Algeria:				
Departments-	ł	ı	1	1
Constantine	Nov. 11-30	. 2		· ·
Oran	Nov. 21-Dec. 11	3		1
Austria:	1101.21-200.11	1 .	1	1
Vienna	Sept. 7-14	5		Sept. 7-14, 1919: Cases, 7.
Rolivia	DC pt. 7-14	1 "	1	Dept. 1-11, 1010. Cusco, 1.
La Paz	June 29-Nov. 22	I	24	Dec. 29-June 28, 1919: Deaths, 5
Chile:	• unic 20-1101. 22	1	1 -3	Dec. 20 - Julie 20, 1010. Deadle, 0
Antofagasta	Nov. 17-30	10	1	Į.
Valparaiso	Nov. 9-22	275	41	
China:	1104. 5-22	213	31	
Antung	Nov. 3-9	1	ł	!
Famel.	1104.9-9			
Egypt: Alexandria	Nov. 12-18	4		
Alexandria	Oct. 1-14		1 8	
Cairo Port Said	do		, 8	_
Finland:	ao	1		•
		į	,	
Province—	T-1- 10 01			<u>.</u> .
Viborg	July 16-31	. 2		0-4 5 05 1010: 0 10 -
Germany		• • • • • • • • • • • • • • • • • • • •		Oct. 5-25, 1919: Cases, 10—civ population, 3; military, 7.
Great Britain:			l i	population, s, mineary, 1.
Glasgow	Nov. 30-Dec. 6	2		
Greece:	NOV. 30-Dec. 0	2		
Cavalla	Nov. 17-23	1		
Drama	Nov. 24-30	1		
Saloniki	Oct. 6-Nov. 23	1		'
		• • • • • • • • • • • • • • • • • • • •	41	Aug 05 Cant 14 1010: Canas 6
Hungary				Aug. 25-Sept. 14, 1919: Cases, 6.
taly: Venice	Nov. 17-23			
	Nov. 17-23	4	1	
apan:	Dec 1.7		i	
Nagasaki	Dec. 1-7	2		
dexico:	70-01-07	_		
Chihuahua	Dec. 21-27	2		
Mexico City	Nov. 16-22	46		Donound
San Luis Potosi	Dec. 14-27			Present.
Spain:	N 04 D 7	اہ	l	
Corunna	Nov. 24-Dec. 7	2	•••••	
Curkey:	37 14 00		1	
Constantinople	Nov. 14-20	3		

YELLOW FEVER.

Mexico: Merids	Dec. 7-20	3	.2	The cases were sent from Opi- chen, vicinity of Muna. One death in case from Muna. To- tal to Dec. 20: Cases, 46; deaths,
				20.