

PUBLIC HEALTH REPORTS

VOL. 41

DECEMBER 31, 1926

NO. 53

CURRENT WORLD PREVALENCE OF DISEASE

A REVIEW OF THE MONTHLY EPIDEMIOLOGICAL REPORT ISSUED OCTOBER 15, 1926,
BY THE HEALTH SECTION OF THE LEAGUE OF NATIONS' SECRETARIAT¹

Only five of the ports in the Far East, which report to the Singapore Bureau, reported either cases of cholera or deaths from this disease during the two weeks ending October 9, and in nearly all localities in eastern Asia, where cholera was prevalent during the past summer, there was marked improvement in the situation at the end of August or during September according to the data made available in the October Epidemiological Report published by the Health Section of the League of Nations. The table below gives the reports from port towns for the five weeks ending October 9.

TABLE 1.—*Cholera cases reported in the principal maritime towns of the Far East between September 5 and October 9, 1926*

Towns	Weeks ending—				
	September			October	
	11	18	25	2	9
Deaths:					
Calcutta.....	18	9	7	-----	14
Madras.....	0	1	3	0	0
Negapatam.....	1	3	0	0	0
Cases:					
Bangkok.....	7	2	3	0	2
Amoy.....	53	50	42	23	18
Shanghai.....	57	66	31	22	6
Antung.....	0	0	0	2	0
Dairen.....	3	1	1	0	0
Harbin.....	27	-----	0	0	0

At Shanghai and Amoy a marked diminution in the number of new cases of cholera was shown by the reports for the two weeks ending October 9, and the outbreaks at Harbin and Dairen apparently had come to an end. In southern China, at Hoihow, Hainan, the number of deaths from cholera declined from 410 in the four weeks ending August 1 to 116 in the five weeks ending September 5. Cholera was reported also from Swatow and other parts of the island of Hainan. At Kwang-Chow-Wan, where 483 cholera cases were reported in August, 194 cases were reported from September 1 to 20.

¹ From the Office of Statistical Investigations.

"Cholera remained prevalent in Tonkin and Annam in September but disappeared from the remainder of French Indo-China except for a few sporadic cases," says the Report. Cases reported in each Province from June to September are shown in the table below.

TABLE 2.—Cholera cases reported in French Indo-China, June–September, 1926

Month	Cam-bodia	Cochin-China	Laos	Annam	Tonkin
June.....	521	1,159	0	128	724
July.....	362	403	7	212	784
August.....	120	39	32	297	234
September.....	4	5	0	138	200

The incidence of cholera in Siam has been declining since last May when 2,660 cases were reported. During the four weeks ending September 11, 204 cases were reported as against 674 cases in the previous four weeks.

Plague.—Three cases of plague were reported at Constantinople during September and none in other Mediterranean ports during the month. Twelve cases occurred in the western desert district of Egypt at Sidi-Barrani between August 19 and September 4, but no cases were reported from any part of Egypt during the remainder of September.

At Beirut, Syria, there were 2 cases of plague on October 11 and another on October 12.

A marked decrease in plague occurred in Senegal during August when only 37 new cases were notified as against 178 in July and 192 in June, the peak month of the outbreak.

In Madagascar the number of cases showed an increase during August, and more cases were reported than in August of any previous year. "The disease is, as usual, most prevalent in the Province of Tananarive," states the Report, "but is spreading also elsewhere and especially at Majunga, where 42 cases were reported during the first half of September. Pulmonary and septicemic cases were very common, as is seen from the table below."

TABLE 3.—Plague cases reported in Madagascar, showing type of disease, July 16 to September 15, 1926

Date	Bubonic	Pulmonary	Septicemic	Total
July 16-31.....	2	5	0	7
Aug. 1-15.....	21	7	2	30
Aug. 16-31.....	42	32	38	112
Sept. 1-15.....	58	17	12	87

"In India plague was spreading during August in Lower Burma and in the Central Provinces and Berar. Outside of these two areas only a few plague cases were reported. In the ports plague cases were reported in September at Rangoon and at Bombay."

No case of plague was reported in August or September at Kwang-Chow-Wan, where 52 cases occurred in June and July. In French Indo-China there were 3 plague cases in September as against 11 in August. Siam reported 6 plague cases in August and only 1 in July.

Plague incidence in Java during the past summer reached the lowest level since 1919. The table below shows that the improvement was general throughout the island with the exception of the Province of Pekalongan, where plague was not very prevalent in 1925. In Surakarta, which was the worst infected Province, a remarkable decline is shown.

TABLE 4.—*Plague deaths reported in Java between June 21 and August 14, 1925 and 1926*

Provinces	1925	1926
Bantam, Batavia, and Preanger.....	0	0
Cheribon.....	122	42
Pekalongan.....	94	150
Semarang.....	39	1
Banjumas.....	182	32
Kedu.....	167	55
Djakakarta.....	13	36
Surakarta.....	631	38
East Java and Madura.....	8	1
Total.....	1,256	355

Twenty-one plague cases were reported in Peru in August, nearly all in the Department of Lima. Seven cases were reported at Guayaquil, Ecuador, in the same month. Argentina reported 3 cases in the first week of October, all in inland localities of the Provinces of Cordoba and Chubut.

Yellow fever.—Cases of yellow fever reported were as follows:

TABLE 5.—*Yellow fever*

Localities	Date	Cases	Deaths
Africa:			
Dahomey—			
Porto-Novo.....	Sept. 10.....	2	1
Gold Coast.....	July 1-31.....	8	3
Nigeria.....	do.....	4	3
America:			
Brazil— ¹			
Bahia.....	May 23-29.....	1	1
	June 6-19.....	4	3
	June 20-26.....	2	1
	July 4-10.....	1	-----

¹ Public Health Reports.

In the Gold Coast 17 cases had been reported from March to July, more than during either of the previous two years, when 6 and 8 cases were reported in 1924 and 1925, respectively.

Typhus and relapsing fever.—Typhus reaches its minimum seasonal incidence in Europe in the late summer, and the disease was little in evidence during August and September. A slight recrudescence

in Poland occurred in September, when 55 cases were reported during the two weeks ending September 18 as against 15 in the previous two weeks.

In Korea, where 118 cases of typhus were reported in June, the incidence declined markedly and only 37 cases were reported in July. No cases have been reported in Japan since May.

Typhus has shown a declining incidence in Chile since 1920, and its decline was accelerated during the first half of 1926, when 83 cases were reported, compared with 317 during the corresponding period of 1925.

An outbreak of relapsing fever occurred in Nigeria in June and July and 324 cases with 41 deaths were reported.

Smallpox.—"Smallpox is becoming increasingly rare on the European Continent," says the Report, but "A new increase of smallpox began early in September in England; 443 cases were reported during the four weeks ended October 2, as against 305 cases during the previous four weeks and 119 during the corresponding period last year. The great majority of cases occur, as usual, in northern England, but there have been a few cases also in London and in Middlesex. There was 1 death from smallpox at South Shields and 1 in the suburbs of London during the week ended October 2."

Dysentery.—The usual seasonal rise in the incidence of dysentery occurred in August or September in most of the central European countries. In Germany 887 cases were reported in the four weeks ending September 18 as against 565 in the previous four weeks, but the incidence was no higher than in either of the previous two years. Czechoslovakia reported 206 cases, Hungary 426 cases, and the Kingdom of the Serbs, Croats, and Slovenes 236 cases in August. In Poland 1,310 cases were reported during the four weeks ended September 18, an increase over the 849 cases in the corresponding period of 1925, but not much more than 50 per cent of the 2,303 cases reported in the corresponding weeks of 1924.

In Japan dysentery shows the same seasonal variation common in Europe, and this year the disease has been more prevalent than it was in 1925. In Java bacillary dysentery was reported to be very prevalent in the first quarter of the year, but the incidence subsided during the spring. There were a number of local outbreaks in scattered districts of the island and no general epidemic.

TABLE 6.—*Dysentery cases reported in Japan and Java by four-weekly periods, 1924, 1925, and 1926*

Four weeks ending—	Japan			Java ²		
	1924	1925	1926	1924	1925	1926
Jan. 31.....	¹ 153	¹ 139	174	1,554	347	1,153
Feb. 28.....	¹ 161	¹ 116	180	1,136	110	3,000
Mar. 28.....	¹ 192	¹ 183	202	493	40	2,733
Apr. 25.....	¹ 265	¹ 214	254	218	21	1,215
May 23.....	¹ 577	¹ 289	472	159	62	396
June 20.....	¹ 676	¹ 678	895	180	41	519
July 18.....	2,179	1,947	1,859	134	24	243
Aug. 15.....	3,021	2,953	3,210	64	12	118
Sept. 12.....	4,013	2,560	3,550	36	60	-----
Oct. 10.....	3,041	2,297	-----	33	162	-----
Nov. 7.....	1,477	1,061	-----	108	163	-----
Dec. 4.....	491	550	-----	122	251	-----
Jan. 2.....	243	302	-----	355	646	-----

¹ Bacillary dysentery only.² Data for calendar months.³ Data for period June 1-20.

Enteric fever.—The report states: "The incidence of enteric fever in European countries in August did not on the whole differ greatly from last year. The situation was more favorable than in August, 1925, in Denmark, Norway, Great Britain, and the Balkan countries. More cases were reported in Poland in August and September than during the corresponding months of 1925. A sudden and severe outbreak of enteric fever occurred in September at Hanover, in Germany, where over 2,000 cases were reported in three weeks. During the two weeks ended September 25, 111 deaths were attributed to enteric fever in the city of Hanover alone. During the week ended August 28 also, 100 cases of 'meat poisoning' were reported at Hanover. The Deutsche Medizinische Wochenschrift states that during that week numerous cases of infectious enteritis occurred at Hanover and were ascribed to the unusually high bacterial content of the drinking water. The bacilli disappeared after chlorination of the water."

In Palestine 421 cases of enteric fever were reported during June and July, compared with 147 cases during the corresponding two months of 1925.

In the United States the incidence was slightly lower than last year. During the four weeks ended September 4, 38 States reported 4,849 cases.

Influenza.—In Mauritius 910 influenza cases and 35 deaths were reported in June; the seasonal maximum is usually in July. Mild outbreaks were reported in July in Basutoland and southern Rhodesia.

New Zealand reported an outbreak of influenza which started in June and reached its maximum in July. During the 12 weeks ending September 6, 117 deaths were attributed to influenza as against 7 during the corresponding period last year.

Acute poliomyelitis.—"An unusual prevalence of poliomyelitis was reported in England and Wales, where more cases were notified in August and September than during the corresponding months of any of the previous eight years," says the Report. The highest number of cases was reported in the county of Leicester, where there were 102 cases during the eight weeks ending October 9, and in Essex, where there were 52 cases in the same period.

An extensive outbreak also occurred in Germany, where it seems to have reached its maximum in the first weeks of September. The cases were scattered throughout northern Germany, while Bavaria, Wurtemberg, and Baden were practically free from the disease.

TABLE 7.—*Cases of poliomyelitis notified in England and Wales and in Germany in 1925 and 1926*

Four weeks ending—	England and Wales		Germany	
	1925	1926	1925	1926
Jan. 31.....	26	17	17	22
Feb. 28.....	23	20	28	14
Mar. 28.....	17	14	21	18
Apr. 25.....	12	14	18	18
May 23.....	16	17	25	22
June 20.....	15	23	18	21
July 18.....	17	26	20	57
Aug. 15.....	28	98	31	180
Sept. 12.....	61	181	57	454
Oct. 10.....	57	227	53	-----
Nov. 7.....	44	-----	45	-----
Dec. 5.....	28	-----	37	-----

Poliomyelitis was much less prevalent in August in the United States than during the previous two years. The disease was also less prevalent in the Scandinavian countries.

Scarlet fever.—Scarlet-fever cases increased in Germany, the Netherlands, and especially in Poland during August and September, and in all three countries the incidence is higher than last year.

TABLE 8.—*Scarlet-fever cases reported in Poland, Germany, and the Netherlands, July 18–October 9, 1925 and 1926*

3 weeks ending—	Poland		Germany		Netherlands	
	1925	1926	1925	1926	1925	1926
Aug. 7.....	1,151	1,813	1,960	2,182	567	711
Aug. 28.....	1,200	2,388	2,167	2,812	673	704
Sept. 13.....	1,511	3,752	2,535	3,756	744	873
Oct. 9.....	1,798	-----	2,965	-----	1,040	1,211

In Poland scarlet fever was reported to be most prevalent in the populous centers and the highest incidence to be among the Jewish population. At Warsaw 14,000 children had been vaccinated against scarlet fever and only 2 cases out of 410 cases reported occurred among those previously vaccinated.

In Germany the disease is most prevalent in east Prussia, Brandenburg, Silesia, Saxony, and the Rhineland; least prevalent in Bavaria and Wurttemberg.

Diphtheria.—The incidence of diphtheria in Europe, on the whole, was slightly lower in August and the first half of September than it was last year. A slight increase over last year, however, was indicated in the reports for Poland, Hungary, Kingdom of the Serbs, Croats, and Slovenes, and Bulgaria.

In the United States about the same number of diphtheria cases were reported early in September as at the corresponding date last year.

Tuberculosis.—The mortality from tuberculosis in a number of large towns during the first half of 1926 is compared with the corresponding rates for 1925 in the following table. While the mortality from tuberculosis is usually higher in the first half year than in the second half year, and these rates, therefore, are not representative of the annual rate, they show, nevertheless, that the decline in tuberculosis mortality has continued in nearly all the towns.

TABLE 9.—*Mortality from tuberculosis in various cities during the first half year of 1925 and 1926*

Cities	Popula- tion in thousands	1925		1926		Increase or decrease
		Deaths	Rates per 100,000	Deaths	Rates per 100,000	
(a) Tuberculosis, all forms						
Europe:						Per cent
Lille.....	201	299	298	227	226	-24.2
Breslau.....	555	336	139	308	111	-20.1
Dresden.....	619	418	135	341	110	-18.5
Lyons.....	562	813	289	669	238	-17.6
Budapest.....	961	1,631	340	1,378	287	-15.6
Dublin.....	438	471	215	398	182	-15.3
Tallinn.....	127	198	312	170	268	-14.1
Berlin.....	4,014	2,559	128	2,221	111	-13.3
Munich.....	681	439	129	382	112	-13.2
Edinburgh.....	427	326	153	285	133	-13.1
Hamburg.....	1,079	703	130	611	113	-13.1
Oslo.....	258	230	178	204	158	-11.2
Glasgow.....	1,057	796	151	720	136	-9.9
Venice.....	201	234	233	211	210	-9.9
London.....	4,602	2,652	115	2,399	104	-9.6
Cologne.....	727	483	133	439	121	-9.0
Prague.....	713	690	194	642	180	-7.2
The Hague.....	398	177	80	170	85	-4.5
Rotterdam.....	552	330	120	316	115	-4.2
Stockholm.....	439	360	164	348	159	-3.0
Trieste.....	249	380	305	368	296	-3.0
Genoa.....	335	389	232	381	227	-2.6
Paris.....	2,906	4,488	309	4,373	301	-2.6
Thirty Swiss cities ¹	1,176	812	139	804	137	-1.4
Copenhagen.....	587	354	121	352	120	-0.8
Madrid.....	783	1,113	284	1,149	293	+3.1
Belfast.....	415	433	209	472	227	+8.6
Milan.....	857	685	160	747	174	+8.8
Bologna.....	221	192	174	217	196	+12.6
Amsterdam.....	718	352	98	399	111	+13.2
Cracow.....	187	230	246	285	304	+23.6

¹ In 1925, 26 cities only.

TABLE 9.—Mortality from tuberculosis in various cities during the first half year of 1925 and 1926—Continued

Cities	Popula- tion in thousands	1925		1926		Increase or decrease
		Deaths	Rates per 100,000	Deaths	Rates per 100,000	
(a) Tuberculosis, all forms—Continued						
America:						Per cent
San Francisco.....	558	323	116	303	106	-6.0
Sao Paulo.....	850	444	104	422	99	-4.8
St. Louis.....	822	312	76	244	59	-2.2
New Orleans.....	414	400	193	405	196	+1.6
Boston.....	780	424	109	433	111	+1.8
Chicago.....	2,995	1,350	90	1,398	93	+3.3
Asia: Manila.....	308	795	516	833	541	+4.8
(b) Pulmonary tuberculosis						
Europe: Sofia.....	154	366	475	444	577	+21.5
America:						
Montevideo.....	423	780	369	567	268	-27.4
New York.....	6,252	2,683	86	2,700	86	0
Asia:						
Madras ¹	527	653	307	625	294	-4.2
Singapore.....	396	615	311	663	335	+7.7
Bombay ²	1,176	516	104	590	118	+13.5

¹ Twenty-two weeks only.

Trachoma.—Information on the prevalence of trachoma is shown in the table below:

TABLE 10.—Trachoma cases reported in various countries, 1924-1926

Country	Total, 1924	1925				1926	
		First quarter	Second quarter	Third quarter	Fourth quarter	First quarter	Second quarter
Germany.....	1,764	487	757	619	914	575	684
Austria.....	341	175	255	104	293	414	172
Danzig.....	54	9	11	17	12	11	9
Estonia.....	528	168	142	76	85	91	81
France.....	173	8	29	11	6	12	9
Lithuania.....	2,375	571	531	372	644	565	146
Malta.....		89	71	123	259	107	184
Poland.....	2,954	1,012	1,057	962	1,720	1,400	2,094
Switzerland.....	13	2	12	1	1	5	4
Czechoslovakia.....	2,782	651	1,001	760	823	810	1,354
Saar Territory.....	3	4	0	1	10	4	0
U. S. R.:							
Governments and territories in							
Europe.....	362,890	139,401	166,602	149,645	105,057	78,210	
Ukraine.....	49,592	18,022	17,160	15,874	19,160	23,660	³ 16,009
Transcaucasia.....	45,982	4,474	11,326	15,603	14,579	280	
Siberia.....	48,158	10,627	10,486	12,216		1,561	
Kirghiz Republic.....	12,045		21,143			1,037	
Turkestan.....	6,648		23,181				
Waterways, railways.....	648	956	994	614	842	1,590	
Tunisia.....	102	24	1	0	0	1	0
United States.....	3,260	392	487	444	628	316	734
New Zealand.....	20	10	5	4	10	3	5

¹ Compulsorily notifiable from Apr. 1, 1926.² Month of March only.³ Data for April and May only.

SYNTHESIS AND INDICATOR PROPERTIES OF SOME NEW SULFONPHTHALEINS

By BARNETT COHEN, Chemist, Hygienic Laboratory, United States Public Health Service

Sensitiveness, brilliant color, and general stability place the simpler sulfonpht haleins in the front rank of acid-base indicators. Although a few of these compounds have been known for some time, a fuller realization and utilization of their unique properties as indicators did not appear until Lubs and Clark (1915, 1916) reported some new syntheses, and Clark and Lubs (1916) proposed their selection of indicators for the determination of hydriions. Not only was a useful set of indicators presented by these authors but, as will be shown presently, there was implicit in their data the means for extending and modifying the series almost at pleasure. Indeed, given the requisite skill in organic synthesis, it would be no great exaggeration to claim the possibility of producing a sulfonpht halein of any desired apparent dissociation constant (useful indicator range) and of almost any color characteristics, within limitations.

The writer's attention was drawn to the problem specifically by the need for a sulfonpht halein substitute for methyl red (an azo compound) in the Clark and Lubs series. Hydriion color standards containing methyl red are notoriously unstable, and the indicator becomes unreliable when used in a biologically active medium, owing, presumably, to more or less reversible reduction and to decompositions. By comparison, the sulfonpht haleins as a class are much more stable. Hence the development of a sulfonpht halein substitute for methyl red would serve two useful purposes—(1) eliminate the unreliable methyl red and (2) render the Clark and Lubs series more uniform chemically.

Analysis of the data of Clark and Lubs led to the decision that di-halogenation of a meta-methyl phenol should produce the desired substitute, and experimental test resulting in the synthesis of tetrabrom-m-cresol sulfonpht halein (brom cresol green) verified this conclusion. Incidentally, a number of other compounds were prepared; and six of them appeared of sufficient value as indicators to merit further study and introduction into the Clark and Lubs series.¹

The following report includes a description of the synthesis of the new sulfonpht haleins and of their absorption curves in the visible spectrum. The apparent dissociation constants were also determined, and data are presented on the salt and protein errors.


¹ A preliminary report on five of these compounds was made in 1923: Public Health Reports, 88, 199. Circumstances have prevented a more detailed report until now, but in the meantime the essential details for the synthesis of these compounds were made available to all inquirers.

EFFECTS OF SIMPLE SUBSTITUTIONS IN PHENOLSULFONPHTHALEIN UPON IONIZATIONS

Structurally phenolsulfonphthalein is a triphenylmethane derivative containing a sulfophenyl and two phenol radicals, the latter attached in their para positions to the methane carbon.

Rather little is known of the effects of substitution in the sulfophenyl radical upon dissociations in the compound (cf. Lubs and Acree, 1916). The dissociation in the strong sulfonic acid radical occurs in the extreme acid ranges, and theoretical considerations suggest that pronounced changes in the constant of this dissociation should be produced by substitutions in the sulfobenzoic acid nucleus. Consequently, should the need arise for the development of sulfonphthalein indicators for extremely acid regions, this type of substitution would be likely to yield compounds of the required indicator properties.

Although accurate comparative measurements of dissociation constants of the sulfonic acid in the different known sulfonphthaleins are not available, we do know that alkyl substitution in the position meta to the phenolic hydroxyl (such as is found in thymol sulfon-



phthalein and m-cresol sulfonphthalein) suppresses this dissociation greatly, enough at least to enable the employment of these compounds as indicators in the pH region between 1.0 and 3.0.

More useful and definite data are available as to the effects of substitution in the phenolic radicals upon the dissociation of the phenolic hydrion, although many gaps still remain to be filled in. It is this dissociation which determines the zones of color-change of most of the useful sulfonphthalein indicators. Such information as is available will be found in Table 1. The value of the dissociation constant K_a is expressed in terms of pK_a , which equals $\log \frac{1}{K_a}$.

The names in parentheses are the common laboratory names proposed for the unwieldy ones of the more common compounds.

TABLE 1.—*Apparent dissociation constant of the phenolic hydrion in the sulfonphthaleins*

Substituted phenol	pK_a
2-isopropyl-5-methyl phenol (thymol blue).....	8. 90
2, 3-dimethyl phenol (xylenol blue).....	8. 9?
*2, 6-dimethyl phenol.....	8. 6
*3-methyl phenol (m-cresol purple).....	8. 32

Substituted phenol	pK _a
2-methyl phenol (o-cresol red)	8. 20
phenol (phenol red)	7. 90
o-iodophenol	6. 6
*o-bromophenol (brom phenol red)	6. 16
*o-chlorophenol (chlor phenol red)	5. 98
2, 6-dibromophenol (brom phenol blue)	4. 05
*2-bromo-6-chlorophenol (brom-chlor phenol blue)	3. 98
*2, 6-dichlorophenol	4. 07
2, 6-dinitrophenol	3. 37
6-bromothymol (brom thymol blue)	7. 10
6-bromo-2, 3-xyleneol	7. 1?
6-bromo-2-methyl phenol (brom cresol purple)	6. 30
*2, 6-dichloro-3-methyl phenol (chlor cresol green)	4. 8
*2, 6-dibromo-3-methyl phenol (brom cresol green)	4. 67

The compounds marked with an asterisk were synthesized by the author, and, with the exception of the 2, 6-dichlorophenol derivative (tetrachloro-phenolsulfonphthalein) were of sufficient purity to give well-defined dissociation constants. The sample of di-iodo-phenol-sulfonphthalein (o-iodophenol derivative) was obtained from the National Aniline & Chemical Co. The data for the other compounds were obtained from Brode (1924), Clark, Cohen, and Elvove (1922), and A. Cohen (1922, 1923).

A mere inspection of this table discloses the following important facts: Alkyl groups depress the dissociation of the phenolic hydrogen and halogens increase it. Considering the effects of alkyl substitution more in detail, it will be noted that meta-substitution has a greater effect than ortho-substitution, that di-substitution has a greater effect than mono-substitution, and that a combination of ortho plus meta-substitution is more effective than di-ortho substitution. The data are not extensive enough to disclose the effect of the heavier isopropyl group as compared with the methyl.

Mono-halogenation in the ortho position increases the ionization of the phenolic hydrogen in the order, iodo < bromo < chloro. Attempts to prepare meta-halogen sulfonphthaleins have been unfruitful, but should the synthesis be accomplished, it will probably be found that the effect on ionization is rather less than that of ortho-halogenation. In terms of pK_a differences, di-halogenation has twice the effect of mono-halogenation. This mode of designating the effects on ionizations is very useful, but the reader should keep in mind that the pK_a differences are direct functions of differences between the *energies* of ionizations and not between the *magnitudes* of the dissociation constants.

Rather noteworthy is an apparent reversal in the order of effect upon ionization of the phenolic hydrion by chlorine and bromine in di-ortho halogenation on the one hand and tetra-ortho halogenation on the other. In phenolsulfonphthalein, dichlorination produces a

greater effect than dibromination; and while this effect seems to be only diminished (but not reversed) in tetra-chlor- and tetrabrom-phenolsulfonphthaleins, we find in the case of *m*-cresolsulfonphthalein that tetra-bromination has a greater effect than tetra-chlorination.

Analogous effects of approximately the same magnitude were found among the indophenols by Cohen, Gibbs, and Clark (1924a). Their data are reproduced here for purposes of comparison, because the parallelism with the sulfonphthaleins is instructive.

Indophenol system	pK _a
carvacrol indophenol.....	8.8
thymol indophenol.....	8.7
<i>m</i> -cresol indophenol.....	8.5
<i>o</i> -cresol indophenol.....	8.4
phenol indophenol.....	8.1
<i>m</i> -bromophenol indophenol.....	7.7
<i>o</i> -bromophenol indophenol.....	7.2
<i>o</i> -chlorophenol indophenol.....	7.0

The effects of alkyl substitutions in both the indophenols and the sulfonphthaleins are almost identical. The papers by Cohen, Gibbs, and Clark (1924b) and by Gibbs, Cohen, and Cannan (1925) contain additional information of possible value in predicting the effects of substitution upon ionization of the phenolic hydrion in the sulfonphthaleins and perhaps other systems.

The effect of substitution on the dissociation of the phenolic hydrogen may be visualized somewhat as follows: If a group (or groups) substituted for hydrogen in the phenol nucleus pulls electron pairs toward itself more than the dissociable hydrogen pulls electron pairs toward itself, the escaping tendency of an electron pair should be lowered at least in the immediate neighborhood. This should become evident in an increased ionization of the hydrogen. The converse of this effect should occur if the substituent group tends to repel electron pairs. If alkyl groups be considered repellant and halogen attractive the effects would be those found here.

The dissociation constants of the ionizable groups depend on three factors—(1) the nature of the groups, (2) the influence of other groups, and (3) the effect of electrostatic forces between the ionizing groups. The nature of the groups determines the general order of magnitude of each constant. The other two factors have an influence dependent upon conditions. Each substituent produces an effect upon an ionizable group dependent upon the nature of the substituent and its position. In addition, work is expended in the liberation of the dissociable hydrogen from the electrostatic attraction of the charge or charges on the rest of the molecule.

The complete formulation of all these factors appears to be hopeless at the moment, but some promising attempts in this direction are being made (cf. Simms, 1926).

The above rather incomplete summary regarding the effects of substitution on ionization in the sulfonphthaleins was only partly available at the time we decided to seek the substitute for methyl red, but enough of it was implicit in Clark and Lubs' data to point the way.

Knowing approximately the magnitude and the direction of shift in pK_a value caused by introduction of halogen or methyl groups in the phenol nuclei of the phenolsulfonphthalein molecule, it was deduced that a tetra-halogenated m-cresolsulfonphthalein should have a pK_a value close to that of methyl red. Experimental test verified our deduction and resulted in the synthesis of m-cresol purple (pK_a 8.32) and brom cresol green (pK_a 4.67). The latter was proposed as a substitute for methyl red (pK 5.0). Since then chlor cresol green (pK 4.8) has been added. The useful pH ranges of these indicators are given below.

Methyl red.....	4. 4-6. 0
Chlor cresol green.....	4. 0-5. 6
Brom cresol green.....	3. 8-5. 4

Although these ranges are not identical, they are sufficiently close for practical indicator use; for it is well known that skillful manipulation of conditions permits accurate colorimetric readings beyond the "limits" of the useful ranges. In actual practice we have found that the new indicators can function as adequate substitutes for methyl red.


EXPERIMENTAL

m-Cresolsulfonphthalein.—A preliminary report by B. Cohen (1923) was the first announcement of the synthesis of this compound. While the present paper was in preparation there appeared the paper by Orndorff and Purdy² (1926) giving a competent elucidation of the chemistry of m-cresolsulfonphthalein.

Orndorff and his associates have shown that the condensation of a phenol with the anhydride of sulfobenzoic acid takes place in two stages, there being first formed an addition product, the intermediate acid, which then reacts with a second molecule of the phenol to give the sulfonphthalein. This process also takes place in the

² Orndorff and Purdy, referring to the preliminary report by B. Cohen (1923), state that the latter gave no details as to the method of preparation or the properties of these sulfonphthaleins (m-cresolsulfonphthalein and its tetrabromo derivative), nor were any analyses given. While this statement is correct, these authors appear to have unintentionally overlooked an exchange of letters between Orndorff and Cohen in 1923, in which Cohen responded to a request for information and gave the following essential facts: (1) m-cresol sulfonphthalein was made by condensation of m-cresol with sulfobenzoic acid anhydride at a temperature not exceeding 110° for 10 hours; (2) purification was obtained by dissolving the crude dye in an alkaline medium of about pH 10 to 11, filtering and reprecipitating with acid; (3) slow crystallization from approximately normal HCl or H₂SO₄ yields well-formed crystals with a metallic luster; and (4) analysis of the brominated product had shown it to be the tetra-brom product.

synthesis of *m*-cresolsulfonphthalein which possesses the following structure:



m-Cresol.—The *m*-cresol was obtained from Eastman (*m*-cresol, "practical") and was redistilled before use. The distillate boiled between 200–201° (755 mm.).

o-Sulfobenzoic acid anhydride.—This was made according to the method of White and Acree (1919) from saccharin. It was crystallized out of benzol, and retained a strong odor of benzol. The presence of the benzol was found to be not detrimental.

Condensation of m-cresol with o-sulfobenzoic acid anhydride.—The process must be carried out at a temperature below 110° if *m*-cresolsulfonphthalein is to be obtained. This has been confirmed by Orndorff and Purdy, who find that higher temperatures favor the formation of dimethylsulfonfluoran, the anhydride of the di-ortho compound. No particular advantage was noted in the employment of condensing agents like zinc chloride so far as improvement in the yield is concerned. The yield is low, between 15 and 20 per cent at the best, and is probably due to a retardative effect exerted by the meta-methyl group.

Crystalline *o*-sulfobenzoic acid anhydride, 30.8 gm., was added to 36.2 gm. of dry, redistilled *m*-cresol which had been warmed to 110°. The mixture was stirred and held for six hours in a bath kept at a temperature of 106°. The compound formation was followed by observing the amount of color produced by a test drop in 10 per cent sodium carbonate solution and in dilute acid. The fusion was terminated when color reached a maximum. The mixture was then steam distilled to remove *m*-cresol. Solid sodium carbonate was then added carefully to the hot solution until the color became deep purple. The solution was allowed to stand overnight to cool and settle out. It was filtered, the precipitate was discarded, and to the filtrate was slowly added concentrated hydrochloric acid until a deep red color developed. This solution was evaporated on the water bath under reduced pressure. Uniform, small green crystals of the sulfonphthalein separated as evaporation progressed. The crystalline product may be washed with cold water to remove adherent acid and salt, and is sufficiently pure (over 95 per cent) for ordinary indicator purposes. The yield up to this point was 12 gm., or about 19 per cent. The residues contain a considerable amount

of coloring matter, but attempts to recover more of the crystalline m-cresolsulfonphthalein from them have not been profitable. Purification is easily effected by dissolving the crystals in hot sodium carbonate solution, filtering, acidifying, and recrystallizing as above by evaporating the solution under reduced pressure.

*Analyses.*³—The air-dried crystals contained from 1 to 3 per cent of moisture. The material dried to constant weight at 110° gave the following analyses for sulfur: Substance, 0.1500, 0.1500, 0.1500; BaSO₄, 0.0891, 0.0903, 0.0911. Calculated for C₂₁H₁₈O₅S, S, 8.39 per cent. Found, 8.16, 8.27, 8.34 per cent. The compound has no definite melting point. It darkens and contracts at 230° and carbonizes at higher temperatures.

Indicator properties.—m-Cresolsulfonphthalein is a brilliant acid-base indicator, and the common name we have suggested for it is *meta-cresol purple*. Like the other sulfonphthaleins, it exhibits two distinct sets of color changes corresponding to the dissociations of the sulfonic acid and the phenolic hydrion, respectively. Unlike most sulfonphthaleins, however, its sulfonic acid dissociation (pK_a 1.51) is sufficiently repressed to make it useful as an indicator of acidity in the pH range 1.2 to 2.8, the corresponding color change being from red to yellow. This pH range and virage are identical with those of thymol sulfonphthalein in the Clark and Lubs series.

It has been found, however, that hydrion color standards of thymol blue in the acid range tend to fade with time.⁴

Since addition of alkali does not regenerate the faded color of thymol blue, it would seem that the loss of color is due to a decomposition rather than to a mere agglomeration of the dye by the high acidity. Under strictly comparable conditions, hydrion color standards containing meta-cresol purple do not suffer the disadvantage of this slow fading. (It is important to emphasize that the color fading we are now discussing is a slow one, being a matter of days or weeks, and does not affect the ordinary use of thymol blue).⁵ We discover, therefore, in meta-cresol purple a brilliant and stable indicator of acidity in the pH region 1.2 to 2.8.

The second color change in meta-cresol purple is from yellow to purple in the pH region 7.4 to 9.0, corresponding to the dissociation (pK_a 8.32) of the phenolic hydrion. In this zone this indicator shares with the other purple indicators the disadvantage of dichromatism, which interferes with the accurate matching of colors. The spectrophotometric data and measurements of the dissociation constants, salt, and protein effects are given in a later section.

³ I am indebted to Chemist Elias Elvove and Assistant Chemist C. G. Remsburg not only for the final analyses presented in this paper but also for numerous preliminary analyses controlling the steps in purification.

⁴ This has also been noted by Dr. W. A. Taylor, of the LaMotte Chemical Co., Baltimore, who now proposes the use of meta-cresol purple as a substitute for thymol blue in the acid range.

⁵ Holmes and Snyder (1925a) found this change to be appreciable spectrophotometrically within 24 hours.

Tetrabrom-m-cresol sulfonphthalein (Brom cresol green).—The synthesis of this compound was first announced in the preliminary report of B. Cohen (1923). It is briefly described by Orndorff and Purdy (1926). A solution of 25 gm. of bromine in 150 c. c. glacial acetic acid was added slowly to a suspension of 15 gm. m-cresol sulfonphthalein in 150 c. c. glacial acetic acid. The mixture was stirred and not allowed to heat above 30°. At intervals a drop was tested in buffer of pH 7. When the blue color reached a maximum residual bromine was removed by aeration. The mixture was then poured into 300 c. c. water and solid sodium bicarbonate was added until the solution turned definitely green. This was allowed to stand overnight and then filtered. Hydrochloric acid was then added and the solution evaporated. As the acetic acid evaporated off the product separated as a dark, reddish-brown amorphous mass. This material on recrystallization from glacial acetic acid gave a light yellowish product which melted at 217–218° (corr.).

Analyses.—Several lots of the compound dried to constant weight at 110° yielded the following analyses for sulfur and bromine: Substance, 0.3565, 0.2277, 0.1910, 0.2570; BaSO₄, 0.1185, 0.0765, 0.0623, 0.0817; substance, 0.1650, 0.1810, 0.1932, 0.2228; AgBr, 0.1780, 0.1933, 0.2175, 0.2543. Calculated for C₂₁H₁₄ Br₄O₅S; S, 4.59 per cent; Br, 45.80 per cent. Found, S, 4.56, 4.62, 4.48, 4.36 per cent; Br, 45.91, 45.45, 45.05, 46.09 per cent.

Indicator properties.—Tetrabrom-m-cresolsulfonphthalein is the compound proposed by B. Cohen (1923) as a substitute for methyl red, and the common name proposed for it is *brom cresol green*. It is far more stable in solution than methyl red, and its color changes are distinct. Brom cresol green may be used in the colorimetric determination of hydron concentration in bacterial cultures to the same extent as the other sulfonphthaleins, although it should be remembered that even these rather stable indicators may be attacked by very active species. Hydron color standards containing brom cresol green remain unaltered under proper conditions for long periods. The color change associated with the ionization of the phenolic hydron is from yellow to blue (corresponding to the pH zone 3.8 to 5.4), the color at the midpoint, pK_a 4.67, being green. Owing to this moderately high dissociation, brom cresol green gives in ordinary tap water the characteristic blue color of the fully dissociated dibasic salt; and for the same reason this indicator is practically insensitive to CO₂.

The spectrophotometric data and measurements of the dissociation constants, salt, and protein effects are given in a later section.

Tetrachlor-m-cresolsulfonphthalein (Chlor cresol green).—Pure m-cresolsulfonphthalein, 8 gm., was suspended in 175 c. c. glacial acetic acid and was chlorinated by bubbling commercial tank chlorine

through the suspension. The subsequent procedure was substantially the same as in the preparation of the tetrabrom derivative. The tetrachlor product was finally recrystallized from glacial acetic acid, from which it separated out in small, brown, velvety tufts, melting at 200–201° (corr.). On analysis it was found to contain 6.1 per cent S and 27.0 per cent Cl; calculated for $C_{21}H_{14}Cl_4O_3S$, 6.17 per cent S, 27.27 per cent Cl.

Except for a determination of the pK_a by the Salm method, no very detailed examination of the compound was made, hence the data here given should be regarded as only approximate. The original purpose in preparing the compound was to determine the effect of tetrachlor substitution as compared with tetrabrom upon the dissociation of the phenolic hydrion.

The pK_a of the tetrachlor derivative was found to be 4.8, and we have seen above that in the tetrabrom compound it is 4.67. The color change in both compounds is the same, from yellow to blue, but the pH ranges are slightly different, corresponding to the differences in pK_a values. The pH range of chlor cresol green is 4.0 to 5.6, a slightly closer approach to the range of methyl red than is given by brom cresol green.

Dibrom-phenolsulfonphthalein.—In the colorimetric determination of hydrion concentration, a matter of minor importance but yet of great convenience is the color of the indicator itself, a factor which is determined by the nature of the solution as well as by the physiology and psychology of color perception. We may encounter amongst apparently normal persons a greater ease in distinguishing color gradations in the reds than in the blues, and vice versa. Another factor of still greater importance in this connection is the dichromatism especially of the purple indicators, which introduces real difficulties in the accurate matching of colors.

The elimination of such troublesome indicators is greatly to be desired if adequate substitutes can be found. In the Clark and Lubs series brom cresol purple and brom phenol blue are the chief offenders, and we have succeeded in producing an excellent substitute for the former in *brom phenol red* (dibrom-phenolsulfonphthalein) which is a clear red in solutions where brom cresol purple is either blue or red, according as the liquid layer is thin or thick.

Brom cresol purple has a pK_a value of 6.3, and from the fact that tetra-brom phenolsulfonphthalein has a pK_a of 4.05 while that of phenolsulfonphthalein is 7.90, it is to be expected that the dibrom compound should have a pK_a about midway between these two and therefore approximately that of brom cresol purple.

In addition it was expected that the color of the new compound in alkaline solution would show more of the red of phenol red and less

of the blue of brom phenol blue. This was deduced from the fact that halogenation in the sulfonphthaleins tends to introduce a blue component in the color of the unhalogenated compound. Thus, tetrabrom-phenolsulfonphthalein is blue while phenolsulfonphthalein is red, dibrom-o-cresolsulfonphthalein is purple while o-cresolsulfonphthalein is red, and dibrom-thymolsulfonphthalein is blue while thymolsulfonphthalein is purplish blue. This deduction was confirmed, but the elimination of the blue component occurred to a greater degree than was expected, for the alkaline color of dibrom-phenolsulfonphthalein exhibits only a slight suggestion of blue.

Sohon (1898) describes the synthesis and properties and gives analyses of a compound alleged to be dibrom-phenolsulfonphthalein. We find that although his analyses correspond to such a compound, the properties described are those of tetra-brom-phenolsulfonphthalein and the method of synthesis yields the tetra-brom product and nothing else. We are unable to account for the apparent discrepancy.

When phenolsulfonphthalein is brominated in glacial acetic acid (the method followed by Sohon) there results tetrabrom-phenolsulfonphthalein, and when the bromination is incomplete the result is a mixture of the unbrominated and tetra-brominated compounds. Analogous effects are produced by chlorination.

We have prepared the dibrom compound and found it to possess properties distinct from the tetra-brom. Moreover, we have confirmed its identity by brominating it and producing the tetra-brom derivative. The change in pK value of the successively brominated derivatives furnishes independent confirmatory evidence.

Synthesis.—The o-bromophenol employed came from two sources—some was prepared in this laboratory and some was purchased from Eastman Kodak Co. Thirty-four grams of o-bromophenol was heated to 140° and 18.1 grams o-sulfobenzoic acid anhydride was added and stirred in. The mixture was kept in an oil bath at 140° for about 10 hours, or until a test drop showed maximum color production. Water was then added and the mixture steam distilled to remove residual bromophenol. Solid sodium bicarbonate was then cautiously added until the solution was a deep bluish red. After standing overnight the solution was filtered. The filtrate was poured slowly into 20 per cent hydrochloric acid and the compound separated out in bright red granular masses. On standing the material assumed a crystalline form with a greenish lustre.

The product is surprisingly soluble in water. It was therefore thoroughly washed with dilute hydrochloric acid, dried over stick sodium hydroxide, and heated in the oven to remove adherent hydrochloric acid. The mother liquor contained a considerable proportion of the compound, which was recovered by evaporation and extraction

with *n*-butyl alcohol. The purified dibrom-phenolsulfonphthalein, recrystallized from glacial acetic acid, melted at 230° (corr.).

Analyses.—Substance, 0.2538, 0.1587; BaSO₄, 0.1159, 0.0713; substance, 0.1500, 0.1500; AgBr, 0.1100, 0.1104. Calculated for C₁₉ H₁₂ O₅ SBr₂, 6.26 per cent S and 31.21 per cent Br. Found, 6.27, 6.17 per cent S, and 31.21, 31.32 per cent Br.

Indicator properties.—Dibrom-phenolsulfonphthalein, *brom phenol red* for short, has the brilliant indicator properties characteristic of the sulfonphthaleins. It is soluble in water to the extent of at least 0.2 per cent, yielding a golden yellow solution. In strongly acid solution it gives an orange-red color, in intermediate zones the color is yellow, and in alkaline solution it is deep red. Its useful range for the colorimetric determination of hydrions is between pH 5.2 and 6.8, corresponding to a pK_a of 6.16. Brom phenol red is an almost perfect substitute for brom cresol purple and is free from the disturbing property of dichromatism.

Purified brom phenol red appears to be perfectly stable, but we have noted in the case of some of our crude preparations a certain amount of fading. This tendency was eliminated by purification of the material. In this connection it is interesting to note that we found a specimen, labeled brom phenol red and sent to us for examination, to have the properties of phenol red and not of brom phenol red. This has also been encountered by Rous (1925). These, may however, be merely cases of accidental mislabeling.

In this connection it is possible that the fading of acid solutions of thymol blue, which we have previously discussed, may be due to impurities in that compound.

Dichlor-phenolsulfonphthalein.—This compound was made in order to determine the effect of dichlor substitution as against dibrom substitution on the ionization of the phenolic hydrion in the corresponding sulfonphthaleins.

Synthesis.—*o*-Chlorophenol was the starting material. Both the Eastman product and material prepared in this laboratory were used. Thirty-two grams of dry *o*-chlorophenol was heated to 130° and 23 grams crystalline *o*-sulfobenzoic acid anhydride was stirred in. The mixture was heated for six hours at 130°; or until a test drop showed maximum color formation. Water was then added and the mixture steam-distilled to remove residual chlorophenol. Sodium bicarbonate was then carefully added until the color became a deep bluish red, and the solution was allowed to stand overnight before filtering. Concentrated hydrochloric acid was added to the filtrate until a precipitate formed. This was filtered off and washed with dilute hydrochloric acid. Water can not be used for washing because of the solubility of the compound. The mother liquor was evaporated and a second crop of crystals was obtained. The adherent moisture

and hydrochloric acid may be driven off with heat. The crystals are very small and are of dark green color with a reddish tinge, and when ground the material is dark red. When recrystallized from glacial acetic acid it yields a product melting at 261–262° (corr.).

Analyses.—Substance, 0.1500, 0.1500; BaSO₄, 0.0810, 0.0815; substance, 0.1500, 0.1500; AgCl, 0.1017, 0.1020. Calculated for C₁₉H₁₂O₅·SCl₂, 7.58 per cent S and 16.76 per cent Cl. Found, 7.42, 7.46 per cent S, and 16.77, 16.82 per cent Cl.

Indicator properties.—Dichlor-phenolsulfonphthalein, *chlor phenol red* for short, is very similar to brom phenol red in solubility and in indicator properties. Its alkaline color is a deep red with even less of a bluish cast than is seen in brom phenol red. The alkaline color of a commercial specimen of the corresponding di-iodo compound was found to be decidedly purplish. We see, therefore, that increasing weight of the halogen substituent tends to introduce an increasing amount of blue in the colors of the corresponding compounds.

The useful range of chlor phenol red for the colorimetric determination of hydriions is between 4.8 and 6.4, corresponding to a pK_a of 5.98. Chlor phenol red overlaps the range of brom cresol green on the one hand and of brom thymol blue on the other. Consequently, both methyl red and brom cresol purple, two objectionable compounds as above indicated, may be eliminated from the Clark and Lubs series of indicators without leaving a gap.

Dibrom-dichlor-phenolsulfonphthalein (Brom-chlor phenol blue).—By brominating dichlor-phenolsulfonphthalein or chlorinating the dibrom compound it is possible to obtain a dibrom-dichlor derivative. It was of interest to obtain this compound and compare its properties with those of the tetrabrom and tetrachlor derivatives.

Synthesis.—Dichlor-phenolsulfonphthalein was brominated in glacial acetic acid at room temperature. The bromination was terminated when a test drop showed maximum development of purple-blue color in alkaline solution. Residual bromine and hydrobromic acid were removed by aeration. Water was added and then solid sodium bicarbonate until the yellow color changed to a deep wine red. The solution was filtered after settling overnight, and to the filtrate was added concentrated hydrochloric acid. The compound separated out as a dark brown precipitate. The mother liquor was evaporated under reduced pressure, and a second crop was obtained. The material was recrystallized from benzol and glacial acetic acid, yielding a flesh-pink powder melting at 250–251° (corr.).

Analyses.—Substance, 0.2000, 0.2000; BaSO₄, 0.0773, 0.0752; substance, 0.2000; AgBr 0.1271; AgCl, 0.1013. Calculated for C₁₉H₁₀O₅SB₂Cl₂, 5.52 per cent S, 27.51 per cent Br, 12.21 per cent Cl. Found, 5.31 and 5.16 per cent S, 27.05 per cent Br, 12.53 per cent Cl.

Indicator properties.—Dibrom-dichlor-phenolsulfonphthalein, *brom-chlor phenol blue* for short, is very similar to brom phenol blue in indicator properties. It imparts a yellow color to mineral acid solutions of around 0.01N and a purplish blue to more alkaline solutions, in which is exhibited the troublesome dichromatism shared by brom phenol blue. Its useful range for the colorimetric determination of hydrons is between pH 3.0 and 4.6, corresponding to a pK_a of 3.98 for the ionization of the phenolic hydron.

Comparing the pK_a values of the tetra-brom compound (4.05) and the dibrom-dichlor compound (3.98), we note that the effect is only a relatively slight increase in ionization of the phenolic hydron when two bromine atoms are replaced by two chlorines. From this it may be inferred that the pK_a value of the tetra-chlor derivative will be shifted still further and to the same slight degree. A crude specimen of tetrachlor-phenolsulfonphthalein was prepared (but not purified or analyzed) and its pK_a value, as determined by the Salm (1906) method, was found to be about 4.0.

Xylenol-sulfonphthaleins.—Xylenol blue, made from 2, 3-dimethyl phenol has been synthesized by A. Cohen (1922). Its pK_a value is approximately 8.9, like that of thymol sulfonphthalein. It is to be expected that the compound made with 2, 5-dimethyl phenol will have approximately the same dissociation constant for the phenolic hydron. On the other hand, the compound made with 2, 6-dimethyl phenol should show a lower pK_a value because of a lesser suppression of the phenolic ionization by o-methyl substitution as compared with m-methyl substitution. By the same reasoning, the compound made with 3, 5-dimethyl phenol (symmetrical m-xylenol) should show a much higher suppression of ionization of the phenolic hydron (pK_a about 9.5).

We did not have available any 2, 5-dimethyl phenol for confirming the one aspect of our predictions, but 2, 6-dimethyl phenol⁶ and 3, 5-dimethyl phenol were available. We found that condensation of 2, 6-dimethyl phenol with o-sulfobenzoic acid anhydride yielded a sulfonphthalein similar in indicator properties to xylenol blue and having a pK_a of 8.6. The compound (2, 6-xylenol sulfonphthalein) crystallized in beautiful, reddish bronze masses melting at 253–254° (corr.). No analyses were made. The behavior of this compound confirmed our prediction.

However, numerous attempts to prepare the symmetrical xylenol derivative were unsuccessful. Pure 3, 5-dimethyl phenol was prepared by the Knoevenagel reaction from ethyl aceto-acetate according to the method described by Gattermann (1923). The *sym*-xylenol was condensed in a variety of ways with the sulfobenzoic acid anhy-

⁶ We are grateful to Dr. L. H. Marks, of the National Aniline & Chemical Co., who supplied us with a pure specimen of this compound.

dride, but no sulfonphthalein was obtained. We have already seen that the yield on condensing m-cresol with the anhydride is very low, and it would seem that the presence of two meta-methyl groups completely hinders the condensation to a sulfonphthalein by the ordinary procedure. Some other method of synthesis will have to be devised to produce this compound, and when it is accomplished we believe our prediction as to its pK_a value will be verified.

SPECTROPHOTOMETRIC MEASUREMENTS

Measurements in the visible and ultraviolet ranges of the spectrum made by Orndorff, Gibbs, Scott, and Jackson (1921) have shown that the sulfonphthaleins in neutral aqueous solution have two absorption bands. The addition of either acid or alkali results in the disappearance of one of the bands and the appearance of two new absorption bands, one on each side of the position of the band that disappears. The other band of the neutral solutions seems to be modified by the addition of acid, but with the addition of alkali it also disappears and a new band with lower frequency appears. In the case of dilute alkaline solutions the new type of absorption is not stable but reverts more or less rapidly to the two absorption bands found in the corresponding neutral solution. These changes and reversions indicate that in the neutral aqueous solutions the carbinol and hydrate forms of the sulfonphthalein are present and that on the addition of either acid or alkali a salt having a quinoid structure is formed.

The absorption curves of aqueous solutions of most of the new sulfonphthaleins were determined in the visible region with a Keuffel & Esser, Model E, direct reading color analyzer, employing tubes 10 cm. long. The wave-length scale was graduated to 5 millimicron intervals and the photometer in unit steps from zero to 100.

A stock aqueous solution was prepared containing 0.04 per cent of the indicator plus one equivalent of NaOH. The solution was diluted 1:9 with water, and this dilution was further diluted 1:19 with acid, alkali, or the required buffer (as indicated below) to produce complete color transformation and to permit viewing through a 10 cm. thickness of solution. The final concentration of indicator was 2.0 mg. per litre, except as noted in certain cases. The measurements were carried out at 30°.

The peaks of the absorption bands were found to be at the following wave lengths ($1 \mu = 10^{-6} \text{ mm} = 10 \text{ Angstrom}$).

	m μ
Meta-cresol purple (acid range).....	533
Meta-cresol purple (alkaline range).....	580
Brom cresol green.....	617
Chlor cresol green.....	612
Brom phenol red.....	574
Chlor phenol red.....	573
Brom-chlor phenol blue.....	596

We found the absorption peak of brom phenol blue to be at $593m\mu$, which is in fair agreement with the value ($592m\mu$) found by Brode (1924). For brom cresol green, Holmes and Snyder (1925b) report the peak at approximately $614m\mu$, and we find it near $617m\mu$.

The absorption curves in the visible spectrum are shown in the accompanying charts (fig. 1) and the experimental data in Table 2. The absorptions are given in terms of $-\log_{10}$ transmittancy ($-\log_{10}T$), which is identical with the product of the thickness, concentration, and the specific transmissive index, k . (cf. Gibson et al., 1922).

m-Cresol purple.—The full acid color was developed in conc. HCl, the neutral color in Clark and Lubs' phthalate buffer of pH 4.6, and the alkaline color in N/5 NaOH.

Brom cresol green, chlor cresol green, brom phenol red.—The full alkaline colors were developed in Clark and Lubs' borate buffer of pH 10. The concentration of brom phenol red was 1.6 mg. per liter.

Chlor phenol red.—The full alkaline color was developed in N/5 NaOH.

Brom-chlor phenol blue.—The full alkaline color was developed in Clark and Lubs' borate buffer of pH 9.2.

TABLE 2.—Absorption values in the visible spectrum ($-\log$ transmittancy)

Wave length in $m\mu$	m-cresol purple			Brom cresol green	Brom phenol red	Chlor phenol red	Brom-chlor phenol blue	Chlor cresol green
	Acid range in conc. HCl	Neutral range in buffer pH 4.6	Alk. range in 0.2 N NaOH					
700				0.039				
690				.061				0.000
680			0.000	.140				.041
670			.018	.252			0.000	.119
660	0.000		.022	.403			.022	.194
650	.004		.071	.620	0.000	0.000	.041	.314
640	.009		.131	.878	.013	.013	.066	.458
630	.013		.222	1.076	.027	.018	.201	.620
620	.018	0.000	.377	1.122	.051	.081	.553	.708
610	.027	.013	.585	1.094	.155	.237	1.187	.744
600	.041	.018	.854	.991	.482	.620	1.602	.713
590	.092	.022	1.046	.870	.979	1.294	1.668	.668
580	.155	.025	1.097	.739	1.372	1.648	1.448	.588
570	.319	.026	1.071	.602	1.398	1.668	1.280	.523
560	.569	.027	.959	.498	1.222	1.462	1.046	.446
550	.921	.036	.824	.415	1.018	1.252	.903	.369
540	1.181	.051	.699	.337	.854	1.046	.710	.319
530	1.301	.071	.577	.244	.678	.886	.538	.244
520	1.097	.097	.475	.174	.538	.683	.432	.194
510	.921	.137	.372	.114	.398	.530	.314	.119
500	.620	.187	.297	.076	.319	.409	.208	.092
490	.469	.222	.237	.046		.328	.174	.056
480	.347	.292	.155		.194	.229	.119	.046
470	.244	.367	.143			.174	.092	.022
460	.149	.432	.119		.097	.131	.046	
450	.125	.444	.032			.102	.000	
440		.469	.000			.027		




FIG. 1

SPECTROPHOTOMETRIC DETERMINATION OF THE APPARENT DISSOCIATION CONSTANTS

The degree of transformation of an indicator within the range of its utility depends upon the hydrion concentration of its solution; and Brode (1924) and Holmes (1924, 1925) have shown how it may be measured with relative accuracy with the aid of the spectrophotometer. All that is necessary is to determine under comparable conditions of concentration and temperature the ratio of color absorption within the useful pH range of any wave length in the absorption band (preferably at the peak) to that of the same wave length in the completely transformed compound. This ratio gives the percentage dissociation of the indicator at the particular pH of the solution measured. The apparent dissociation constant, pK_a , of the indicator can then be calculated by the familiar equation:

$$pK_a = pH - \log \frac{\alpha}{1 - \alpha}$$

where α is the degree of dissociation.

This procedure yields consistent results, and in the case of brom cresol green we have been able to confirm the value obtained by Holmes and Snyder (1925b).

Suitable dilutions of each indicator were made in acid, alkali, or Clark and Lubs' buffers, as the case required, to produce complete color transformation, and these were compared with the same quantity of indicator in buffers of known pH.

All measurements were carried out at 30°.

m-Cresol purple, acid range.—The acid form of the indicator is red, with an absorption band in the yellow, the peak lying at 533 $m\mu$. The accurate determination of the dissociation constant depends on obtaining complete dissociation of the indicator, and we found concentrated HCl necessary to produce complete acid transformation of *m-cresol purple*, just as Holmes and Snyder did for the acid range of thymol blue. The results are summarized in Table 3.

TABLE 3.—*m-Cresol purple (acid range). Absorption maximum at 533 $m\mu$*

Buffer (pH)	pH electro- metric	Conc. mg. per liter	T	—log T	α	$\log \frac{\alpha}{1 - \alpha}$	pK_a
Conc. HCl.....		1.6	7.7	1.1135	1.000		
1.2.....	1.224	1.6	18.3	.7375	.662	+0.2626	1.52
1.4.....	1.419	1.6	24.8	.6055	.544	+ .0766	1.50
1.6.....	1.609	1.6	32.5	.4881	.438	— .1082	1.50
1.8.....	1.806	3.2	17.7	.3760	.338	— .2520	1.51
2.0.....	2.000	3.2	26.1	.2917	.262	— .4498	(1.55)
Average.....							1.51

* Corrected to concentration of 1.6 mg. per liter.

It will be seen from the table that the pK_a values obtained near the middle of the dissociation curve agree fairly closely, the average value being 1.51.

m-Cresol purple, alkaline range.—As the alkalinity of the indicator solution is increased through the pH range 7.0 to 11.0 the indicator becomes progressively and completely transformed to its alkaline form, which is purple in color, with an absorption band in the yellow, the peak lying at $580m\mu$. The apparent dissociation constant was determined spectrophotometrically, the fully transformed alkaline form of the indicator being produced in N/5 NaOH. The data are summarized in Table 4. The average pK_a value found is 8.32.

TABLE 4.—*m-Cresol purple (alkaline range). Absorption maximum at $580m\mu$*

Buffer (pH)	pH electro-metric	Conc. mg. per liter	T	—log T	α	$\log \frac{\alpha}{1-\alpha}$	pK_a
8.0	7.905	3.2	29.6	¹ 0.2644	0.274	+0.4242	8.33
8.2	8.108	3.2	18.3	¹ .3688	.382	+ .2099	8.32
8.4	8.304	3.2	11.6	¹ .4678	.484	+ .0278	8.33
8.6	8.500	3.2	7.1	¹ .5744	.594	— .1655	8.33
8.8	8.700	1.6	20.3	.6925	.716	— .4024	8.30
N/5 NaOH	-----	1.6	10.8	.9666	1.000	-----	-----
Average	-----	-----	-----	-----	-----	-----	8.32

¹ Corrected to concentration of 1.6 mg. per liter.

Brom cresol green.—The peak of the absorption band of the alkaline form of this indicator lies at $617m\mu$. Clark and Lubs' borate buffer pH 9.6 was used to produce the alkaline transformation. The average pK_a value found is 4.67, which agrees with that found by Holmes and Snyder (1925b). Our experimental data are summarized in Table 5. In our preliminary note, B. Cohen (1923) this value was stated to be 5.0, as determined by the Salm method. We have found on subsequent repurification of the indicator that the apparent dissociation constant went down and remained constant at 4.67, although the bromine and sulfur analyses remained substantially unchanged (cf. Holmes and Snyder (1925b)).

TABLE 5.—*Brom cresol green. Absorption maximum at $617m\mu$*

Buffer (pH)	pH electro-metric	Conc. mg. per liter	T	—log T	α	$\log \frac{\alpha}{1-\alpha}$	pK_a
4.4	4.397	3.2	22.3	¹ 0.3259	0.343	+0.2827	4.68
4.6	4.597	1.6	37.8	.4225	.444	+ .0970	4.69
4.8	4.800	1.6	28.6	.5436	.572	+ .1254	4.67
5.0	4.998	1.6	22.3	.6517	.685	— .3383	4.66
5.2	5.193	1.6	18.3	.7375	.776	— .5389	4.65
9.6	-----	1.6	11.2	.9508	1.000	-----	-----
Average	-----	-----	-----	-----	-----	-----	4.67

¹ Corrected to concentration of 1.6 mg. per liter

Brom phenol red.—The absorption peak of the alkaline form of this indicator lies near $574m\mu$. Clark and Lubs' borate buffer pH 10.0 was used to produce the full alkaline color. The average pK_a value found is 6.16, and the data are summarized in Table 6. We have noted in some specimens of this indicator a peculiarity not observed in the other sulfonphthaleins examined. Alkaline solutions of these specimens showed a progressive decrease in absorption with time. In other repurified specimens there was no such change, and we are therefore inclined to ascribe this peculiar behavior to impurities.

TABLE 6.—*Brom phenol red.* Absorption maximum at $574m\mu$

Buffer (pH)	pH electro-metric	Conc. mg. per liter	T	—log T	α	$\log \frac{\alpha}{1-\alpha}$	pK_a
6.0	5.956	1.6	27.8	0.5560	0.385	+0.2032	6.16
6.2	6.165	1.6	18.4	.7352	.509	— .0160	6.15
6.4	6.349	1.6	13.6	.8665	.600	— .1764	6.17
6.6	6.567	1.6	9.0	1.0458	.724	— .4197	6.15
6.8	6.769	1.6	7.1	1.1487	.796	— .5906	6.18
10.0	-----	1.6	3.6	1.4437	1.000	-----	-----
Average	-----	-----	-----	-----	-----	-----	6.16

Chlor phenol red.—The absorption peak of the alkaline form of this indicator lies near $573m\mu$. Clark and Lubs' borate buffer of pH 10.0 was used to produce the full alkaline color. The average pK_a value found was 5.98, and the data are summarized in Table 7.

TABLE 7.—*Chlor phenol red.* Absorption maximum at $573m\mu$

Buffer (pH)	pH electro-metric	Conc. mg. per liter	T	—log T	α	$\log \frac{\alpha}{1-\alpha}$	pK_a
5.6	5.592	3.2	11.5	¹ 0.4697	0.296	+0.3757	5.97
5.8	5.783	1.6	25.9	.5867	.370	+ .2309	6.01
6.0	5.956	1.6	17.1	.7670	.484	+ .0279	5.98
6.2	6.165	1.6	10.8	.9666	.610	— .1939	5.97
6.4	6.349	1.6	7.6	1.1192	.706	— .3807	5.97
10.0	-----	1.6	2.6	1.5850	1.000	-----	-----
Average	-----	-----	-----	-----	-----	-----	5.98

¹ Corrected to concentration of 1.6 mg. per liter.

Brom-chlor phenol blue.—The absorption peak of the alkaline form of this indicator lies at $596m\mu$. Clark and Lubs' borate buffer of pH 9.4 was used to produce the full alkaline color. The average pK_a value found is 3.98, and the data are summarized in Table 8.

TABLE 8.—*Brom-chlor phenol blue. Absorption maximum at 596m μ*

Buffer (pH)	pH electro-metric	Conc. mg. per liter	T	—log T	α	$\log \frac{\alpha}{1-\alpha}$	pK _a
3.8	3.803	3.2	8.0	¹ 0.5485	0.386	+0.2013	4.00
4.0	3.997	1.6	18.9	.7235	.509	— .0163	3.98
4.2	4.195	1.6	12.6	.8996	.633	— .2375	3.96
4.4	4.397	1.6	9.0	1.0458	.736	— .4401	3.96
4.6	4.597	1.6	6.7	1.1739	.827	— .6783	(3.92)
9.4		1.6	3.8	1.4202	1.000		
Average							3.98

¹ Corrected to concentration of 1.6 mg. per liter.

"SALT ERRORS" OF THE NEW INDICATORS

In the absence of a satisfactory theory that will permit calculation of the salt errors of indicators, the only reliable procedure is to determine these errors by direct hydrogen electrode measurements. This has been done for those of the new sulfonphthaleins that are regarded as useful supplements to the Clark and Lubs series.

Sodium chloride was added to various Clark and Lubs buffers so as to bring the solutions to 1 molar concentration (the electrolyte in the 0.02 M buffer being calculated in terms of NaCl). For measuring the salt error of m-cresol purple, acid range, the "1 M" buffer contained 53.110 gm. NaCl, 250 c. c. M/5 KCl and 207.5 c. c. M/5 HCl per liter; for the alkaline range the "molar" buffer contained 52.268 gm. NaCl, 250 c. c. M/5 H₃BO₃, M/5 KCl and 29.5 c. c. M/5 NaOH. For brom phenol red and chlor phenol red the "molar" buffer contained 55.204 gm. NaCl, 250 c. c. M/5 KH₂PO₄ and 89.00 c. c. M/5 NaOH per liter. For brom cresol green and brom-chlor phenol blue the "molar" buffer contained 55.32 gm. NaCl, 250 c. c. M/5 KH-phthalate, and 18.5 c. c. M/5 NaOH per liter.

These "molar" solutions were then diluted to 0.5 M, 0.2 M, and 0.005 M.

Hydrogen ion measurements were taken of the various solutions, both electrometrically and colorimetrically, the basis of the colorimetric comparisons being the standard Clark and Lubs buffers. The measurements were all made at 30°. The hydrogen electrode determinations were referred to M/20 KH-phthalate (pH 3.97) as a standard. The indicator solutions were 0.04 per cent concentrations of the mono-sodium salts in water. These were prepared in the manner outlined by Clark (Determination of Hydrogen Ions, 2d ed., p. 80–81), the equivalents of N/20 NaOH per 100 mg. indicator being as follows:

TABLE 9.—Quantities of NaOH to produce mono-sodium salts of indicators

Indicator	Mol. weight	N/20 NaOH per 100 mg.
m-Cresol purple.....	382.2	c. c. 5.3
Brom cresol green.....	698.0	2.9
Brom phenol red.....	512.1	3.9
Chlor phenol red.....	423.2	4.7
Brom-chlor phenol blue.....	581.0	3.4

The differences between the electrometric and colorimetric pH values were determined and are summarized in Table 10 as corrections.

TABLE 10.—Salt effect on the new sulphonphthaleins

[The values given below are corrections to be added to the colorimetric pH determinations to bring the values to the electrometric pH of the corresponding Clark and Lubs' buffers]

Molar conc. salt	m-Cresol purple		Brom cresol green	Brom phenol red	Chlor phenol red	Brom-chlor phenol blue
	Acid range	Alkaline range				
1.0	-0.14	-0.29	-0.32	-0.26	-0.26	-0.33
0.5	-.09	-.22	-.26	-.22	-.20	-.28
0.2	-.02	-.16	-.16	-.12	-.10	-.16
0.005	+.11	+.09	+.09	+.25	+.23	+.14

Similar results were obtained in another series of experiments with brom cresol green. In this case a normal sodium citrate solution was used, containing 250 c. c. molar citric acid and 500 c. c. normal NaOH. The colorimetric comparisons were made against Clark and Lubs' buffers. The pH corrections found for the salt effect are given below.

1.0 normal.....	-0.20
0.5 normal.....	-.19
0.2 normal.....	-.09
0.1 normal.....	-.03

The salt effect at high salt concentrations appears to be least for m-cresol purple in acid ranges and greatest for brom-chlor phenol blue. As the salt concentration decreases toward 0.1 molar the effect becomes practically negligible. With still greater dilutions the sign of the salt "error" changes and becomes quite appreciable at 0.005 molar salt (cf. Kolthoff, 1925, and Lepper and Martin, 1926).

"PROTEIN" EFFECT

Protein material in solutions containing indicators exerts specific effects on the colors, effects dependent on the nature not only of the indicator but also of the protein and apparently on its previous

treatment. The only safe procedure when attempting the colorimetric determination of pH in protein solutions is to calibrate the readings of the particular indicator in the specific protein solution with the hydrogen electrode.

The following data are presented merely to show the magnitude of the effect produced by a certain peptone upon the sulfonphthalein indicators. Incidentally, the sulfonphthaleins in the Clark and Lubs series were also included so as to have a comparable set of data. All observations were made at least in duplicate and were consistent. Colorimetric readings were made to the nearest 0.05 pH against Clark and Lubs' buffers. Quadruplicate hydrogen electrode measurements had to agree within 0.5 mv. before being accepted. The determinations were made at 30°.

The peptone solution was a 5 per cent concentration of Witte's peptone, which was boiled and filtered. To aliquots of this stock peptone solution were added small quantities of conc. HCl or NaOH to bring the pH within the range of the particular indicator studied. In most of the cases a more or less distinct opalescent haze appeared after the addition of acid or alkali, but the colorimetric and electro-metric readings remained unchanged after filtration of such solutions.

The same experiment was repeated two months later. The same lot of peptone was used and apparently the same technic. The results disclosed certain divergencies which we are unable to explain. There was substantial agreement in the results for the indicators of the acid regions down to brom cresol green, but below that the two series tend to diverge. The data are summarized in Table 11.

TABLE 11.—“Protein effect” on sulfonphthalein indicators

[The values listed are the corrections to be added to colorimetric pH readings to bring them to the electro-metric]

Indicator	In 5 per cent Witte peptone		Clark and Lubs ¹
	Series 1	Series 2	
m-Cresol purple (acid).....	-0.20	-0.20	
Thymol blue (acid).....	-.19	-.20	
Brom phenol blue.....	-.28	-.43	+0.05
Brom-chlor phenol blue.....	-.28	-.43	
Brom cresol green.....	-.10	-.13	
Chlor phenol red.....	+ .09	-.07	
Brom phenol red.....	+ .11	-.10	
Brom cresol purple.....	+ .11	-.10	+ .01
Brom thymol blue.....	+ .34	+ .07	+ .10
Phenol red.....	+ .24	-.01	+ .04
Cresol red.....	+ .02	-.03	+ .03
m-Cresol purple (alk.).....	+ .03	-.02	
Thymol blue (alk.).....	+ .09	-.03	+ .04

¹ In a 1 per cent peptone-beef infusion broth.

For purposes of comparison there is included in Table 11, under the column headed Clark and Lubs, the corrections found by these authors (1917) for a 1 per cent peptone-beef infusion broth. While a strict comparison could scarcely be considered valid, it nevertheless is useful in a rough survey of the ground. Our results show the general magnitude of the effect produced on the colorimetric reading by the presence of 5 per cent peptone. Their main value lies in again emphasizing the dictum that "protein effects" have to be determined experimentally for the material under examination and that calibration is not a simple matter.

SUMMARY

The following new sulfonphthaleins have been synthesized: m-cresolsulfonphthalein, tetra-brom-m-cresolsulfonphthalein tetra-chlor-m-cresolsulfonphthalein, dibrom-phenolsulfonphthalein, dichlor-phenolsulfonphthalein, dibrom-dichlor-phenolsulfonphthalein, and 2, 6-xyleneol sulfonphthalein. The effects of substitution on dissociations in the sulfonphthaleins are discussed, and certain predictions based on an empirical formulation of the effects have been verified.

All but the last mentioned of these compounds are recommended as useful supplements to the Clark and Lubs series of acid-base indicators, or as substitutes for certain unsatisfactory members in that series. The new compounds have been studied as to their indicator properties, spectrophotometric behavior in the visible region, apparent dissociation constants, salt, and protein effects, The essential characteristics are summarized in Table 12.

TABLE 12.—*Summary of characteristics of the new sulfonphthalein indicators*

Sulfonphthalein	Common name	Absorption max. ¹	pK _a	Useful pH range	Color change
m-Cresol	m-Cresol purple	$\left\{ \begin{array}{l} 533 \\ 580 \end{array} \right.$	1.51	1.2-2.8	Red-yellow.
Tetra-brom m-cresol	Brom cresol green	617	8.32	7.4-9.0	Yellow-purple.
Tetra-chlor m-cresol	Chlor cresol green	612	4.67	3.8-5.4	Yellow-blue.
Dibrom phenol	Brom phenol red	574	4.8	4.0-5.6	Do.
Dichlor phenol	Chlor phenol red	573	6.16	5.2-6.8	Yellow-red.
Dibrom dichlor phenol	Brom-chlor phenol blue	596	5.98	4.8-6.4	Do.
			3.98	3.0-4.6	Yellow-blue.

¹ The absorption maxima are for the (alkaline) disodium salts, except in the case of m-cresol purple, acid range, where the value given refers to the absorption of the free acid.

² Acid.

³ Alk.

References

- Brode, W. R. 1924. Jour. Am. Chem. Soc., **46**, 581.
 Clark, W. M., and Lubs, H. A. 1916. Jour. Wash. Acad. Sci., **6**, 483.
 Clark, W. M., and Lubs, H. A. 1917. Jour. Bact., **2**, 1, 109, 191.
 Clark, W. M., Cohen, B., and Elvove, E. 1922. In "The Determination of Hydrogen Ions," by W. M. Clark, 2d edition, p. 93.
 Cohen, A. 1922. Biochem. Jour., **16**, 31.

- Cohen, A. 1923. *Biochem. Jour.*, **17**, 535.
Cohen, B. 1923. *Public Health Reports*, **38**, 199.
Cohen, B., Gibbs, H. D., and Clark, W. M. 1924a. *Public Health Reports*, **39**, 381.
Cohen, B., Gibbs, H. D., and Clark, W. M. 1924b. *Public Health Reports*, **39**, 804.
Gattermann, L. 1923. *Practical Methods of Organic Chemistry*, 3d edition, N. Y., p. 202.
Gibbs, H. D., Cohen, B., and Cannan, R. K. 1925. *Public Health Reports*, **40**, 649.
Gibson, K. S., et al. 1922. *Sci. Papers, U. S. Bur. Standards*, **18**, 124. (No. 440.)
Holmes, W. C. 1924. *Jour. Am. Chem. Soc.*, **46**, 627.
Holmes, W. C., and Snyder, E. F. 1925a. *Jour. Am. Chem. Soc.*, **47**, 221.
Holmes, W. C., and Snyder, E. F. 1925b. *Jour. Am. Chem. Soc.*, **47**, 226.
Kolthoff, I. M. 1925. *Rec. trav. chim.*, **44**, 275.
Lepper, E. H., and Martin, C. J. 1926. *Biochem. J.* **20**, 45.
Lubs, H. A., and Acree, S. F. 1916. *Jour. Am. Chem. Soc.*, **38**, 2772.
Lubs, H. A., and Clark, W. M. 1915. *Jour. Wash. Acad. Sci.*, **5**, 609.
Lubs, H. A., and Clark, W. M. 1916. *Jour. Wash. Acad. Sci.*, **6**, 481.
Orndorff, W. R., et al. 1921. *Phys. Rev.*, **17**, 437.
Orndorff, W. R., and Purdy, A. C. 1926. *Jour. Am. Chem. Soc.*, **48**, 2212.
Rous, P. 1925. *Jour. Exp. Med.*, **41**, 451.
Salm, E. 1906. *Z. physik. Chem.*, **57**, 471.
Simms, H. S. 1926. *Jour. Am. Chem. Soc.*, **48**, 1251.
Sohn, M. D. 1898. *Am. Chem. Jour.*, **20**, 257.
White, E. C., and Acree, S. F. 1919. *Jour. Am. Chem. Soc.*, **41**, 1190.

SMALLPOX IN MINNESOTA—1913–1925

In the PUBLIC HEALTH REPORTS of December 3, 1926, page 2789, appears the statement that 1,298 cases of smallpox with 63 deaths had occurred in Minneapolis, Minn. Dr. A. J. Chesley, State health officer of Minnesota, advises that the number of deaths was 363, not 63, which gives a case fatality in this series of cases of more than 25 per cent.

Doctor Chesley sends the following statement of vaccination histories of smallpox cases which have occurred in Minnesota:

MINNESOTA, SMALLPOX

1925—Total cases, 973; total deaths, 198

Class A.—Successfully vaccinated within seven years before attack, 15 cases, or 1.54 per cent; 1 death, or 0.51 per cent.

Class B.—Successfully vaccinated over seven years before attack, 191 cases, or 19.63 per cent; 41 deaths, or 20.71 per cent.

Classes C and D.—Class C, never successfully vaccinated, and Class D, unable to give definite history of vaccination and no scar found, 767 cases, or 78.83 per cent; 156 deaths, or 78.78 per cent.

1913-1925—Total cases, 39,250; total deaths, 613

Class A.—661 cases, or 1.68 per cent of all; 1 death, or 0.16 per cent of all deaths from smallpox.

Class B.—1,976 cases, or 5.03 per cent of all; 89 deaths, or 14.52 per cent of all deaths from smallpox.

Classes C and D.—36,613 cases, or 93.28 per cent of all; 523 deaths, or 85.32 per cent of all deaths from smallpox.

Vaccination history of 505 fatal cases in Minnesota

	1924	1925	Total
Class A.....	0	1	1
Class B.....	47	41	88
Class C.....	243	151	394
Class D.....	17	5	22
Total.....	307	198	505

PUBLIC HEALTH ENGINEERING ABSTRACTS

Lead Poisoning From Food.—Anon. *The Lancet*, No. 5375, September 4, 1926, p. 507:

"The story is told in the *Presse Médical* of an outbreak of lead poisoning which commenced at Vidin, in Bulgaria, during April, 1923, and terminated only when its origin was detected five months later. By that time there had been 314 cases, occurring in 153 families, with three deaths, while several other deaths occurred among those poisoned as a result of other disorders considered to have been brought on or at least accentuated by the lead absorbed; notably one case of cancer of the rectum in a woman aged 28 is attributed to this cause. The number of cases ranks the epidemics with others, such as have followed upon accidentally mixing of white lead with flour, from drinking plumbo-solvent water by the royal household at Versailles in the eighteenth century, and from Loch Katrine, in Glasgow, at the end of the nineteenth century, and the 350 cases at Saint-Georges-sur-Eure in 1865. The signs and symptoms appear to have been quite typical; the blue line was present in 99 per cent of cases; lead colic was frequent and affections of the nervous system, with 3 cases (1 fatal) of encephalopathy. Nephritis and wrist drop, symptoms of chronic lead poisoning, were not observed. The source of the poisoning was found to be adulterated red pepper. This pepper, prepared from capsicum fruit, is much used in Hungary and Bulgaria, and it is often adulterated with such things as maize or vetch flour, sawdust, iron filings, or brick dust, but on this occasion analysis revealed the presence of 20.5 per cent of red lead and 4.1 per cent of sand, while very small amounts of true pepper were found.

This fraudulent "red pepper" was placed on the market by only one firm, and inhabitants who bought their red pepper elsewhere or made it themselves escaped. Most of the cases followed the ingestion of a number of small doses, a teaspoonful of pepper serving two to four persons for several days; but one case followed a single dose of two teaspoonfuls of "pepper," containing about 6 grains of red lead, taken at one meal; the case recovered."

Outbreak of Paratyphoid Fever Due to Infected Ice Cream.—J. P. Kinloch. (*The Medical Officer*. 1925, v. 34, pp. 191-192.) Abstract by W. G. Savage in *Bulletin of Hygiene*, vol. 1, No. 2, February, 1926, p. 101.

"An outbreak of 23 cases, all but 2 in Aberdeen, of paratyphoid fever in August, 1925. The symptoms were all of this disease and none were of food-poisoning type. For example, in all the cases the onset was characteristically insidious; and while vomiting was not uncommon, when present it was slight. In general, constipation and not diarrhea was the rule. Rose spots were present in most cases. Although a number of the cases were severe, there were no deaths. The incubation period was about 15 days.

"Careful epidemiological inquiry showed that the one article of food consumed in common was ice cream, obtained from one particular shop. The milk used to make the ice cream was naturally the object of suspicion, but milk from the same source used elsewhere produced no disease. About 6 gallons of milk were daily converted into ice cream on the incriminated premises. The staff on these premises was investigated, but no evidence of previous illness of any member was forthcoming and the bacteriological examinations were negative. It was not possible to ascertain how the ice cream became infected, but direct or indirect contamination by a paratyphoid B 'carrier' was considered as the most likely source. Investigations could not be undertaken until three weeks after the ice cream became infective.

"The outbreak was definitely proved to be paratyphoid fever by the isolation of *B. paratyphosus* B from the feces of eight sufferers and from the urine of two cases, and by the demonstration of specific agglutinins in the blood of all the cases.

"A point of interest is that the ice cream on the day it was infective was distributed to between 120 to 360 people, while only 23 developed the disease. Probably only part of it was infected and the freezing prevented distribution of the bacilli throughout the whole mass."

Clean and Safe Milk Campaign to Stimulate Use.—S. J. Crumbine, general executive, American Child Health Association, New York City. *The Nation's Health*, vol. 8, No. 8, August, 1926, pp. 530-532. (Abstract by H. N. Old.)

Some very significant facts and figures on milk-borne epidemics of communicable diseases are tabulated and commented upon, as obtained from questionnaires covering 42 States, the District of Columbia, 3 Territories, and 3 Canadian Provinces.

During 1924 and 1925, 43 and 44 such epidemics are reported, respectively, with a total for the two years of 3,286 cases and 130 deaths, typhoid epidemics constituting 72 per cent of the total reported.

The danger from tuberculous cows is dwelt upon at length and the conclusion reached by Park and Krumwiede is stated—that 27 per cent of tubercular children under 5 and 25 per cent of those between 5 and 16 years of age are found to have infection of the bovine type.

Up to the present time 12 States have been surveyed, 179 towns covered, 3,945 supplies and 4,928 samples examined for visible dirt, bacteriological contamination, and detection of adulteration, Standard Methods being followed. Summarizing the survey results, it is stated that 77 per cent of the samples were classed as dirty, 58 per cent showing bacterial count over 100,000, with 43 per cent positive for colon bacillus, and about 14 per cent adulteration.

The objectives of the campaign are, first, stimulation of the production and distribution of an abundant, clean, and safe milk supply; second, to center the responsibility for such production and distribution on dairymen, milk dealers, and State and local dairy and health officials; and third, to promote increased consumption of milk after reasonable assurance of its safety.

The by-products of the survey thus far are said to be most encouraging, resulting in the promotion of a clean-up among the dairymen and dealers, provision of local supervision using laboratory examinations in many instances, and commitments made toward a general tightening of milk-control regulations.

While the survey shows the daily per capita milk consumption to be only 0.6 to 0.8 pint, the conclusion is reached that, in many communities, increased consumption should not be urged until the safety of the supply is assured.

The organizations cooperating actively in this work are the American Child Health Association, the Association of Dairy, Food, and Drug Officials, and the Conference of State and Provincial Health Authorities.

Bacterial Flora of the Market Oyster.—Calista Eliot. *The American Journal of Hygiene*, vol. 6, No. 6, November, 1926, p. 755.

(1) Shucked oysters and shell oysters kept at the laboratory temperatures show a sudden and maximum rise in total count from the second to the fourth day of storage; (2) the *Bacillus coli* score of oysters stored in a cool basement increased from 4 to 500,000 in 14

days; as signs of spoilage appeared, the *Bacillus coli* score decreased; (3) hydrogen ion determinations on oysters spoiling in the shell showed little change in acidity; shucked oysters, however, became markedly acid during the first few days of spoilage; later, there was a reversal of reaction and the original pH was regained and maintained; (4) in the ice box the rise in acidity lagged two or three days behind and remained at a slightly lower level than at room temperature; at ice-box temperature the maximum total count was about one-tenth of the maximum count at room temperature; (5) the bacteria of the decomposing oyster may be divided into five principal groups—(a) the colon-aerogenes group; (b) the streptococci; (c) the 'water bacteria,' including members of the green fluorescent, the yellow pigmented, and the nonpigmented groups, and vibrios; (d) the anaerobes; and (e) the incidental organisms, such as the chromogenic cocci and the aerobic spore formers; (6) in shucked oysters the souring process may be initiated by either the colon-aerogenes group or the streptococci; if the streptococci are present in large numbers, the colon-aerogenes group is inhibited by the second day; (7) after a varying period of time, 12 days or longer, the water forms multiply rapidly, there is a reversion in reaction and actual decomposition of the oyster meat begins. Certain members of the green pigmented and the yellow pigmented groups produce changes in sterilized oysters comparable to those observed in the decomposition of market oysters; there is a slimy chromogenic growth and a marked softening, and, in some instances, liquefaction of the oyster meat when these organisms are grown upon them; members of these groups are always found abundantly in spoiling oysters; other water forms which are also abundant in the spoiling oyster do not initiate decomposition processes in sterilized oysters; (8) several types of anaerobes multiply in spoiling oysters and produce large amounts of gas, but apparently bring about no putrefactive changes.

Summary of the Purpose and Principles of Aeration of Water Supplies.—C. A. Emerson, jr. Proceedings of Eighth Texas Water Works Short School, Bulletin No. 1, January 23, 1926, pp. 78–83. (Abstract by W. H. Wendler.)

The aeration of ground water is usually for the purpose of the oxidation of iron, manganese, or organic matter and for removing volatile odors and gases such as carbon dioxide and hydrogen sulphide. These constituents, when present to excess, impart color, turbidity, and sometimes taste to the water, and by deposit cause staining of plumbing fixtures and white clothing in the laundry. Carbon dioxide also dissolves iron from the interior of the mains. There have been instances in which samples of tap water showed six or more parts per million of iron in contrast to one part per million at the well.

Where iron is present it is readily changed by oxidation from the soluble ferrous form to the insoluble ferric hydrate, only one part of oxygen being required to oxidize seven parts of iron. It has been found that if the dissolved oxygen content were permitted to rise above 50 per cent saturation, the iron and manganese could not be satisfactorily removed. In some Massachusetts plants where manganese and organic matter interfered with precipitation of iron in fully aerated water, the tricklers were operated as submerged contact beds.

Aeration for removal of tastes and odors due to industrial waste pollution, particularly 'phenol' wastes from by-product coke ovens and wood distillation plants, has been of little practical value in most instances.

Sometimes surface supplies, taken from the lower levels of large reservoirs or from rivers which have been ice blocked for long periods, are somewhat deficient in oxygen, and in these instances aeration was helpful.

Connecting Safe and Unsafe Water Supplies.—Anon. *Public Works*, vol. 57, No. 8, September, 1926, pp. 281-282. (Abstract by Dana E. Kepner.)

At the conference this year of the State sanitary engineers a committee on cross-connections presented a report recommending the adoption by the conference of resolutions providing that "no physical connections should be permitted between any potable public water supplies, either through cross-connections, auxiliary intakes or by-passes, and other supplies except as follows: (1) With another potable public water supply; or (2) with a potable supply which is regularly examined as to its quality by those in charge of the potable public supply to which the connection is made." A cross-connection is defined as any physical connection whereby a potable public water-supply system is connected with another water-supply system, whether public or private, in such a manner that a flow of water into the potable supply is possible therefrom, directly through the manipulation of gate valves, because of ineffective check or back-pressure valves, or otherwise.

The results from a questionnaire sent to the various State boards of health dealing with existing regulations in this respect are given.

Garbage Collection and Disposal.—Anon. *Public Works*, vol. 57, No. 10. November, 1926, pp. 385-387. (Abstract by C. L. Pool.)

This article is the first of a series in review of a symposium on garbage collection and disposal held by the sanitary engineering division of the American Society of Civil Engineers. Six papers constituting the article were as follows: A general review of the problem, by Samuel A. Greeley; a description of practice at Lansing,

Mich., by Edward D. Rich; the same for the hog feeding for Los Angeles, Calif., by W. T. Knowlton; a description of the Beccari system at Scarsdale, N. Y., by Arthur Boniface; one of high temperature incineration at Toronto, Canada, by A. J. Burnett; and one of the Cobwell system of garbage reduction at Rochester, N. Y., by John V. Lewis.

Mr. Greeley discussed administrative and engineering problems encountered and outlined the procedure recommended to cities confronted with the problem. In connection with incineration specifications recently prepared by him the work was classified under five heads: (1) Incinerator furnaces and appurtenances; (2) incinerator building and scale; (3) chimney; (4) runway; and (5) sewers and sewage-disposal plant. A list of reduction plants in operation noted whether each was operated by the city or by a contractor. The Kansas City contract allows disposal by any satisfactory method. The contract price (1925) was \$6.45 a ton for collection and \$1 a ton for disposal.

Can collection practice at Lansing, Mich., is emphasized and constructional details of cans are given. Frequency and methods of collection are outlined and costs given include \$0.91 per capita of the population served for collection in 1924.

Los Angeles practice is discussed, with quantities and costs noted. Material rejected by the pigs is covered with gypsum to conserve the ammonia content, dried, and ground for use as fertilizer.

Examination for Entrance into the Regular Corps of the United States Public Health Service

Examinations of candidates for entrance into the Regular Corps of the United States Public Health Service will be held at the following-named places on the dates specified:

Washington, D. C.	February 7, 1927
Chicago, Ill.	February 7, 1927
New Orleans, La.	February 7, 1927
San Francisco, Calif.	February 7, 1927

Candidates must be not less than 23 nor more than 32 years of age, and they must have been graduated in medicine at some reputable medical college, and have had one year's hospital experience or two years' professional practice. They must pass satisfactorily, oral, written, and clinical tests before a board of medical officers and undergo a physical examination.

Successful candidates will be recommended for appointment by the President, with the advice and consent of the Senate.

Requests for information or permission to take this examination should be addressed to the Surgeon General, United States Public Health Service, Washington, D. C.

DEATHS DURING WEEK ENDED DECEMBER 18, 1926

Summary of information received by telegraph from industrial insurance companies for week ended December 18, 1926, and corresponding week of 1925. (From the Weekly Health Index, December 22, 1926, issued by the Bureau of the Census, Department of Commerce)

	Week ended Dec. 18, 1926	Corresponding week, 1925
Policies in force-----	65, 797, 778	62, 410, 497
Number of death claims-----	12, 674	12, 128
Death claims per 1,000 policies in force, annual rate--	10. 0	10. 1

Deaths from all causes in certain large cities of the United States during the week ended December 18, 1926, infant mortality, annual death rate, and comparison with corresponding week of 1925. (From the Weekly Health Index, December 22, 1926, issued by the Bureau of the Census, Department of Commerce)

City	Week ended Dec. 18, 1926		Annual death rate per 1,000 cor- respond- ing week, 1925	Deaths under 1 year		Infant mortality rate, week ended Dec. 18, 1926 ²
	Total deaths	Death rate ¹		Week ended Dec. 18, 1926	Cor- respond- ing week, 1925	
Total (65 cities)-----	7, 237	13. 1	13. 0	787	771	3 66
Akron-----	27			2	3	22
Albany ⁴ -----	37	16. 2	23. 0	1	3	21
Atlanta-----	64			5	10	
White-----	29			2	7	
Colored-----	35	(⁵)		3	3	
Baltimore ⁴ -----	222	14. 3	13. 3	24	15	73
White-----	162			16	11	60
Colored-----	60	(⁵)		8	4	127
Birmingham-----	55	13. 6	20. 0	10	3	
White-----	28			4	1	
Colored-----	27	(⁵)		6	2	
Boston-----	209	13. 8	15. 4	31	31	87
Bridgeport-----	33			5	6	85
Buffalo-----	127	12. 2	14. 1	12	20	50
Cambridge-----	28	12. 0	13. 5	8	6	142
Camden-----	37	14. 7	12. 1	9	4	151
Canton-----	29	13. 7	8. 8	3	3	66
Chicago ⁴ -----	696	11. 9	12. 2	64	80	56
Cincinnati-----	127	16. 1	17. 7	9	17	56
Cleveland-----	184	10. 0	10. 3	16	30	42
Columbus-----	93	17. 0	12. 1	7	3	65
Dallas-----	50	12. 8	16. 4	5	18	
White-----	39			5	16	
Colored-----	11	(⁵)		0	2	
Denver-----	90	16. 5	14. 7	15	7	
Des Moines-----	40	14. 3	8. 5	6	0	100
Detroit-----	274	11. 1	11. 5	48	45	78
Duluth-----	24	11. 1	11. 3	6	2	139
El Paso-----	26	12. 4	12. 4	6	4	
Erie-----	36			5	7	98
Fall River ⁴ -----	33	13. 1	10. 5	6	5	94
Flint-----	40	15. 3	7. 6	16	5	271
Fort Worth-----	38	12. 5	10. 1	3	6	
White-----	34			2	5	
Colored-----	4	(⁵)		1	1	
Grand Rapids-----	32	10. 7	9. 5	6	5	86
Houston-----	46			4	11	
White-----	35			4	5	
Colored-----	11	(⁵)		0	6	
Indianapolis-----	104	14. 8	15. 5	7	7	53
White-----	87			5	6	44
Colored-----	17	(⁵)		2	1	115
Jersey City-----	62	10. 2	11. 9	6	15	45
Kansas City, Kans-----	25	11. 1	11. 7	1	4	19
White-----	22			1	3	22
Colored-----	3	(⁵)		0	1	0
Kansas City, Mo-----	116	16. 1	12. 1	9	7	
Los Angeles-----	261			31	22	86

(See footnotes at end of table.)

Deaths from all causes in certain large cities of the United States during the week ended December 18, 1926, infant mortality, annual death rate, and comparison with corresponding week of 1925. (From the Weekly Health Index, December 22, 1926, issued by the Bureau of the Census, Department of Commerce)—Continued

City	Week ended Dec. 18, 1926		Annual death rate per 1,000 corresponding week, 1925	Deaths under 1 year		Infant mortality rate, week ended Dec. 18, 1926 ¹
	Total deaths	Death rate ²		Week ended Dec. 18, 1926	Corresponding week, 1925	
Louisville.....	86	14.4	13.6	4	6	34
White.....	62			3	5	29
Colored.....	24	(³)		1	1	70
Lowell.....	22			1	7	19
Lynn.....	25	12.5	13.7	1	5	26
Memphis.....	66	19.4	21.2	9	8	
White.....	29			2	5	
Colored.....	37	(³)		7	3	
Milwaukee.....	107	10.8	9.2	19	11	90
Minneapolis.....	101	12.1	14.2	6	11	33
Nashville ⁴	45	17.1	15.0	5	1	
New Bedford.....	20			2	2	35
New Haven.....	46	13.2	11.1	4	3	55
New Orleans.....	163	20.3	19.5	14	12	
White.....	90			8	7	
Colored.....	73	(³)		6	5	
New York.....	1,524	13.4	12.3	148	140	60
Bronx Borough.....	173	10.0	10.8	17	16	57
Brooklyn Borough.....	550	12.8	10.9	57	45	58
Manhattan Borough.....	630	17.5	16.1	56	62	62
Queens Borough.....	128	8.7	7.9	15	14	68
Richmond Borough.....	43	15.7	16.6	3	3	53
Newark, N. J.....	90	10.2	13.1	14	11	67
Norfolk.....	23	6.9	12.0	4	4	81
White.....	11			1	3	33
Colored.....	12	(³)		3	1	159
Oakland.....	52	10.4	11.1	6	5	70
Oklahoma City.....	26			4	4	
Omaha.....	58	14.0	15.7	6	9	64
Paterson.....	35	12.8	11.4	5	2	84
Philadelphia.....	508	13.2	14.6	46	55	61
Pittsburgh.....	176	14.4	13.4	25	18	83
Portland, Oreg.....	68			2	2	20
Providence.....	59	11.2	14.0	5	2	42
Richmond.....	53	14.6	14.9	8	3	100
White.....	35			4	0	78
Colored.....	18	(³)		4	3	139
Rochester.....	62	10.1	13.8	5	6	40
St. Louis.....	220	13.8	14.5	15	21	
St. Paul.....	58	12.2	12.1	2	4	18
Salt Lake City ⁴	28	11.0	11.5	8	3	122
San Antonio.....	40	10.2	14.7	11	10	
San Diego.....	51	24.2	20.7	3	4	64
San Francisco.....	154	14.2	12.0	6	8	36
Schenectady.....	25	14.0	8.4	7	4	201
Seattle.....	76			10	4	96
Somerville.....	29	15.1	15.3	1	3	28
Spokane.....	24	11.5	14.8	1	3	23
Springfield, Mass.....	37	13.3	10.3	5	3	77
Syracuse.....	53	14.9	13.2	3	7	38
Tacoma.....	29	14.3	10.0	1	0	24
Toledo.....	75	13.3	10.0	10	7	96
Trenton.....	34	13.2	17.8	6	8	102
Utica.....	36	18.2	15.4	3	3	68
Washington, D. C.....	120	11.9	12.6	13	16	74
White.....	74			8	9	67
Colored.....	46	(³)		5	7	91
Waterbury.....	21			1	4	24
Wilmington, Del.....	21	8.8	13.2	4	2	89
Worcester.....	53	14.3	11.5	5	3	60
Yonkers.....	35	15.7	8.3	6	1	135
Youngstown.....	38	12.0	11.1	9	4	114

¹ Annual rate per 1,000 population.

² Deaths under 1 year per 1,000 births. Cities left blank are not in registration area for births.

³ Data for 63 cities.

⁴ Deaths for week ended Dec. 17, 1926.

⁵ In the cities for which deaths are shown by color, the colored population in 1920 constituted the following percentages of the total population: Atlanta, 31; Baltimore, 15; Birmingham, 39; Dallas, 15; Fort Worth, 14; Houston, 25; Indianapolis, 11; Kansas City, Kans., 14; Louisville, 17; Memphis, 38; Nashville, 30; New Orleans, 26; Norfolk, 38; Richmond, 32; and Washington, D. C., 25.

PREVALENCE OF DISEASE

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

UNITED STATES

CURRENT WEEKLY STATE REPORTS

These reports are preliminary and the figures are subject to change when later returns are received by the State health officers

Reports for Week Ended December 25, 1926

ARIZONA		CONNECTICUT—continued	
	Cases		Cases
Chicken pox.....	2	Septic sore throat.....	1
Diphtheria.....	3	Trachoma.....	1
German measles.....	23	Tuberculosis (all forms).....	14
Paratyphoid fever.....	1	Typhoid fever.....	2
Scarlet fever.....	3	Whooping cough.....	23
Smallpox.....	1		
Tuberculosis.....	21		
Typhoid fever.....	1		
ARKANSAS		DELAWARE	
Chicken pox.....	21	Influenza.....	2
Diphtheria.....	8	Scarlet fever.....	14
Influenza.....	35	Tuberculosis.....	4
Malaria.....	3	Typhoid fever.....	2
Measles.....	1	Whooping cough.....	3
Scarlet fever.....	9		
Smallpox.....	1		
Tuberculosis.....	1		
Typhoid fever.....	3		
COLORADO		FLORIDA	
Cerebrospinal meningitis.....	1	Cerebrospinal meningitis.....	1
Chicken pox.....	15	Chicken pox.....	21
Diphtheria.....	9	Diphtheria.....	30
Measles.....	4	Influenza.....	3
Mumps.....	1	Malaria.....	4
Pneumonia.....	3	Measles.....	8
Scarlet fever.....	31	Mumps.....	16
Smallpox.....	1	Paratyphoid fever.....	1
Tuberculosis.....	97	Pneumonia.....	5
Whooping cough.....	1	Scarlet fever.....	13
		Smallpox.....	39
		Tetanus.....	1
		Tuberculosis.....	11
		Typhoid fever.....	4
		Whooping cough.....	5
CONNECTICUT		IDAHO	
Cerebrospinal meningitis.....	1	Chicken pox.....	4
Chicken pox.....	81	Diphtheria.....	3
Diphtheria.....	18	Measles.....	57
German measles.....	3	Pneumonia.....	4
Influenza.....	2	Scarlet fever:	
Measles.....	29	Nampa.....	15
Mumps.....	17	Scattering.....	27
Pneumonia (all forms).....	55	Smallpox.....	1
Polioomyelitis.....	1	Tuberculosis.....	1
Scarlet fever.....	60	Typhoid fever.....	1

ILLINOIS	Cases
Cerebrospinal meningitis:	
Cook County.....	3
Chicken pox.....	360
Diphtheria.....	110
Influenza.....	37
Lethargic encephalitis:	
Cook County.....	1
Measles.....	577
Mumps.....	82
Pneumonia.....	291
Scarlet fever.....	234
Smallpox.....	20
Tuberculosis.....	267
Typhoid fever.....	16
Whooping cough.....	148

KANSAS	Cases
Chicken pox.....	169
Diphtheria.....	19
Influenza.....	6
Measles.....	34
Mumps.....	14
Pneumonia.....	35
Scarlet fever.....	77
Septic sore throat.....	1
Smallpox:	
Seneca.....	13
Topeka.....	10
Scattering.....	6
Tuberculosis.....	62
Typhoid fever.....	1
Whooping cough.....	9

LOUISIANA	Cases
Diphtheria.....	17
Influenza.....	11
Lethargic encephalitis.....	1
Malaria.....	6
Measles.....	24
Pneumonia.....	26
Poliomyelitis.....	1
Scarlet fever.....	9
Smallpox.....	1
Tuberculosis.....	32
Typhoid fever.....	6

MAINE	Cases
Chicken pox.....	43
Diphtheria.....	1
German measles.....	1
Influenza.....	4
Measles.....	78
Mumps.....	16
Pneumonia.....	17
Scarlet fever.....	42
Tuberculosis.....	6
Typhoid fever.....	1
Vincent's angina.....	2
Whooping cough.....	33

MARYLAND ¹	Cases
Cerebrospinal meningitis.....	2
Chicken pox.....	146
Diphtheria.....	48
German measles.....	1

MARYLAND—continued	Cases
Influenza.....	42
Measles.....	27
Mumps.....	22
Paratyphoid fever.....	1
Pellagra.....	1
Pneumonia (all forms).....	77
Scabies.....	1
Scarlet fever.....	68
Septic sore throat.....	4
Tuberculosis.....	27
Typhoid fever.....	11
Typhus fever.....	1
Whooping cough.....	67

MASSACHUSETTS	Cases
Cerebrospinal meningitis.....	2
Chicken pox.....	265
Conjunctivitis (suppurative).....	4
Diphtheria.....	104
German measles.....	7
Influenza.....	14
Lethargic encephalitis.....	1
Measles.....	59
Mumps.....	140
Ophthalmia neonatorum.....	25
Pneumonia (lobar).....	72
Poliomyelitis.....	1
Scarlet fever.....	236
Septic sore throat.....	2
Tuberculosis (all forms).....	113
Typhoid fever.....	31
Whooping cough.....	115

MICHIGAN	Cases
Diphtheria.....	63
Measles.....	66
Pneumonia.....	60
Scarlet fever.....	154
Smallpox.....	19
Tuberculosis.....	13
Typhoid fever.....	1
Whooping cough.....	89

MONTANA	Cases
Chicken pox.....	11
Diphtheria.....	7
Measles.....	73
Mumps.....	9
Scarlet fever.....	103
Smallpox.....	10
Typhoid fever.....	3

NEW JERSEY	Cases
Anthrax.....	1
Chicken pox.....	145
Diphtheria.....	78
Influenza.....	11
Measles.....	21
Pneumonia.....	86
Scarlet fever.....	127
Typhoid fever.....	1
Whooping cough.....	102

¹ Week ended Friday.

NEW YORK

Cases

(Exclusive of New York City and Syracuse)

Anthrax.....	1
Cerebrospinal meningitis.....	3
Chicken pox.....	278
Diphtheria.....	76
Dysentery.....	1
German measles.....	41
Lethargic encephalitis.....	1
Measles.....	571
Mumps.....	103
Ophthalmia neonatorum.....	2
Paratyphoid fever.....	1
Pneumonia.....	200
Polioomyelitis.....	2
Scarlet fever.....	149
Smallpox.....	6
Trachoma.....	1
Typhoid fever.....	9
Vincent's angina.....	6
Whooping cough.....	154

OREGON

Cerebrospinal meningitis.....	1
Chicken pox.....	18
Diphtheria.....	12
Influenza.....	15
Malaria.....	1
Measles.....	32
Mumps.....	4
Pneumonia.....	9
Scarlet fever.....	32
Septic sore throat.....	1
Smallpox.....	17
Tuberculosis.....	23
Typhoid fever.....	1
Whooping cough.....	1

SOUTH DAKOTA

Cerebrospinal meningitis.....	1
Chicken pox.....	19
Diphtheria.....	1
Influenza.....	1
Measles.....	35

SOUTH DAKOTA—continued

Cases

Pneumonia.....	1
Scarlet fever.....	27
Smallpox.....	4
Whooping cough.....	1

UTAH

Chicken pox.....	10
Diphtheria.....	2
German measles.....	3
Measles.....	179
Mumps.....	19
Pneumonia.....	6
Scarlet fever.....	4

VERMONT

Chicken pox.....	16
Measles.....	23
Mumps.....	16
Scarlet fever.....	4
Whooping cough.....	24

WASHINGTON

Cerebrospinal meningitis.....	1
Chicken pox.....	110
Diphtheria.....	23
German measles.....	17
Measles.....	117
Mumps.....	30
Scarlet fever.....	91
Smallpox.....	38
Tuberculosis.....	4
Typhoid fever.....	4
Whooping cough.....	2

WEST VIRGINIA

Chicken pox.....	101
Diphtheria.....	33
Influenza.....	30
Measles.....	103
Scarlet fever.....	59
Smallpox.....	1
Tuberculosis.....	13
Typhoid fever.....	16
Whooping cough.....	69

Reports for Week Ended December 18, 1926

NORTH DAKOTA

Cases

Chicken pox.....	24
Diphtheria.....	7
Measles.....	361
Mumps.....	10
Pneumonia.....	5

1 Deaths.

NORTH DAKOTA—continued

Cases

Scarlet fever.....	54
Smallpox.....	1
Tuberculosis.....	4
Typhoid fever.....	1
Whooping cough.....	2

SUMMARY OF MONTHLY REPORTS FROM STATES

The following summary of monthly State reports is published weekly and covers only those States from which reports are received during the current week:

State	Cerebro-spinal meningitis	Diphtheria	Influenza	Malaria	Measles	Pellagra	Polio-myelitis	Scarlet fever	Small-pox	Typhoid fever
<i>July, 1926</i>										
Massachusetts.....	8	174	9	4	917	4	21	628	0	46
<i>October, 1926</i>										
Texas.....	0	180	57	1,392			0	81		160
<i>November, 1926</i>										
Illinois.....	12	581	80	5	1,368	0	12	1,124	25	202
Iowa.....	3	132			82		1	220	27	12
Louisiana.....	4	180	61	57	52	19	2	87	14	57
Maryland.....	3	208	71	2	89	0	1	192	0	96
Michigan.....	0	711	15	1	325		7	962	90	49
Minnesota.....	1	430	9		511		0	1,054	23	15
New York.....	18	1,178		7	2,657		41	1,213	76	206
North Dakota.....	0	26	0		423		1	226	32	3
Ohio.....	6	1,333	23	1	134		7	1,387	132	159
Wisconsin.....	11	301	140		1,837		7	606	0	32
Wyoming.....	7	5	3		92		1	88	6	7

<i>November, 1926</i>			
Actinomycosis:	Cases	Lethargic encephalitis:	Cases
Minnesota.....	1	Illinois.....	6
Anthrax:		Louisiana.....	1
New York.....	2	Maryland.....	6
Chicken pox:		Michigan.....	6
Illinois.....	1,860	Minnesota.....	4
Iowa.....	290	New York.....	14
Louisiana.....	14	North Dakota.....	1
Maryland.....	501	Ohio.....	4
Michigan.....	1,212	Wisconsin.....	1
Minnesota.....	1,121	Mumps:	
New York.....	2,619	Illinois.....	239
North Dakota.....	146	Iowa.....	21
Ohio.....	2,376	Louisiana.....	2
Wisconsin.....	1,503	Maryland.....	43
Wyoming.....	115	Michigan.....	132
Dysentery:		New York.....	911
Illinois.....	23	North Dakota.....	11
Louisiana.....	5	Ohio.....	207
Maryland.....	9	Wisconsin.....	461
Michigan.....	1	Wyoming.....	17
Minnesota.....	4	Ophthalmia neonatorum:	
New York.....	3	Illinois.....	42
Wyoming.....	1	Iowa.....	1
German measles:		Maryland.....	1
Illinois.....	30	New York.....	2
Iowa.....	1	Ohio.....	107
Maryland.....	11	Paratyphoid fever:	
New York.....	243	Illinois.....	3
North Dakota.....	21	Minnesota.....	1
Ohio.....	9	New York.....	8
Wisconsin.....	22	Ohio.....	2
Hookworm disease:		Wyoming.....	1
Louisiana.....	15	Puerperal septicemia:	
Minnesota.....	1	Illinois.....	6
Impetigo contagiosa:		New York.....	9
Maryland.....	8	Rabies in animals:	
Lead poisoning:		Maryland.....	6
Illinois.....	22		
Ohio.....	15		

Rabies in man:	Cases	Typhus fever:	Cases
Ohio.....	1	Illinois.....	1
Septic sore throat:		Maryland.....	1
Illinois.....	4	Vincent's angina:	
Maryland.....	14	Maryland.....	2
Michigan.....	21	New York.....	59
New York.....	12	Whooping cough:	
Ohio.....	5	Illinois.....	953
Wyoming.....	1	Iowa.....	25
Tetanus:		Louisiana.....	13
Illinois.....	2	Maryland.....	298
Minnesota.....	1	Michigan.....	493
New York.....	8	Minnesota.....	98
Trachoma:		New York.....	1,352
Iowa.....	1	North Dakota.....	23
Ohio.....	3	Ohio.....	938
Tularæmia:		Wisconsin.....	910
Illinois.....	1	Wyoming.....	55

RECIPROCAL NOTIFICATIONS

Notifications regarding communicable diseases sent during the month of November, 1926, to other State health departments by departments of health of certain States

Referred by—	Acti- nomy- cosis	Blasto- myco- sis	Chick- en pox	Diph- theria	Dysen- tery	Measles	Polio- myeli- tis	Small- pox	Tuber- culo- sis	Ty- phoid fever
California.....									2	
Illinois.....								2	24	6
New Jersey.....										1
New York.....				3		2	1			6
Minnesota.....	1	1	1		3				28	1

GENERAL CURRENT SUMMARY AND WEEKLY REPORTS FROM CITIES

Diphtheria.—For the week ended December 11, 1926, 39 States reported 2,143 cases of diphtheria. For the week ended December 12, 1925, the same States reported 1,679 cases of this disease. One hundred cities, situated in all parts of the country and having an aggregate population of more than 30,360,000, reported 1,169 cases of diphtheria for the week ended December 11, 1926. Last year for the corresponding week they reported 911 cases. The estimated expectancy for these cities was 1,350 cases. The estimated expectancy is based on the experience of the last nine years, excluding epidemics.

Measles.—Thirty-seven States reported 5,089 cases of measles for the week ended December 11, 1926, and 4,561 cases of this disease for the week ended December 12, 1925. One hundred cities reported 1,160 cases of measles for the week this year and 2,451 cases last year.

Poliomyelitis.—The health officers of 39 States reported 29 cases of poliomyelitis for the week ended December 11, 1926. The same States reported 41 cases for the week ended December 12, 1925.

Scarlet fever.—Scarlet fever was reported for the week as follows: Thirty-nine States—this year, 3,576 cases; last year, 3,203 cases; 100 cities—this year, 1,387 cases; last year, 1,280 cases; estimated expectancy, 1,062 cases.

Smallpox.—For the week ended December 11, 1926, 38 States reported 667 cases of smallpox. Last year for the corresponding week they reported 365 cases. One hundred cities reported smallpox for the week as follows: 1926, 65 cases; 1925, 119 cases; estimated expectancy, 69 cases. No deaths from smallpox were reported by these cities for the week this year. .

Typhoid fever.—Three hundred and eighty-four cases of typhoid fever were reported for the week ended December 11, 1926, by 39 States. For the corresponding week of 1925 the same States reported 444 cases of this disease. Ninety-nine cities reported 61 cases of typhoid fever for the week this year and 111 cases for the corresponding week last year. The estimated expectancy for these cities was 87 cases.

Influenza and pneumonia.—Deaths from influenza and pneumonia were reported for the week by 94 cities, with a population of more than 29,600,000, as follows: 1926, 830 deaths; 1925, 799 deaths.

City reports for week ended December 11, 1926

The "estimated expectancy" given for diphtheria, poliomyelitis, scarlet fever, smallpox, and typhoid fever is the result of an attempt to ascertain from previous occurrence how many cases of the disease under consideration may be expected to occur during a certain week in the absence of epidemics. It is based on reports to the Public Health Service during the past nine years. It is in most instances the median number of cases reported in the corresponding week of the preceding years. When the reports include several epidemics or when for other reasons the median is unsatisfactory, the epidemic periods are excluded and the estimated expectancy is the mean number of cases reported for the week during nonepidemic years.

If reports have not been received for the full nine years, data are used for as many years as possible, but no year earlier than 1917 is included. In obtaining the estimated expectancy the figures are smoothed when necessary to avoid abrupt deviations from the usual trend. For some of the diseases given in the table the available data were not sufficient to make it practicable to compute the estimated expectancy.

Division, State, and city	Population, July 1, 1925, estimated	Chicken pox, cases re-reported	Diphtheria		Influenza		Measles, cases re-reported	Mumps, cases re-reported	Pneumonia, deaths re-reported
			Cases, estimated expectancy	Cases re-reported	Cases re-reported	Deaths re-reported			
NEW ENGLAND									
Maine:									
Portland.....	75,333	22	2	0	0	0	0	1	4
New Hampshire:									
Concord.....	22,546	0	1	0	0	0	26	0	0
Manchester.....	83,097	0	5	0	0	0	6	0	2
Vermont:									
Barre.....	10,008	3	0	0	0	0	21	0	1
Massachusetts:									
Boston.....	779,620	145	64	37	1	2	14	52	23
Fall River.....	128,993	5	5	4	1	0	1	4	0
Springfield.....	142,065	9	5	7	2	0	4	1	0
Worcester.....	190,757	27	5	0	0	1	1	3	9
Rhode Island:									
Pawtucket.....	69,760	6	2	5	0	0	0	0	0
Providence.....	267,918	0	10	8	0	0	1	0	7
Connecticut:									
Bridgeport.....	(1)	2	11	4	2	1	1	2	4
Hartford.....	160,197	6	9	3	0	0	0	0	5
New Haven.....	178,927	26	4	1	0	0	1	1	4

1 No estimate made.

City reports for week ended December 11, 1926—Continued

Division, State, and city	Population July 1, 1925, estimated	Chick- en pox, cases re- ported	Diphtheria		Influenza		Meas- les, cases re- ported	Mumps, cases re- ported	Pneu- monia, deaths re- ported
			Cases, esti- mated expec- tancy	Cases re- ported	Cases re- ported	Deaths re- ported			
MIDDLE ATLANTIC									
New York:									
Buffalo.....	538,016	27	26	16	-----	0	3	7	15
New York.....	5,873,356	237	227	194	77	16	18	172	157
Rochester.....	316,786	5	9	7	5	2	1	1	10
Syracuse.....	182,003	20	10	0	-----	0	18	8	0
New Jersey:									
Camden.....	128,642	8	6	18	0	0	0	1	0
Newark.....	452,513	25	20	12	6	0	1	8	9
Trenton.....	132,020	6	8	2	0	0	0	0	1
Pennsylvania:									
Philadelphia.....	1,979,364	165	85	48	1	5	3	21	62
Pittsburgh.....	631,563	100	29	22	1	1	2	2	23
Reading.....	112,707	13	5	3	-----	0	0	1	3
EAST NORTH CENTRAL									
Ohio:									
Cincinnati.....	409,333	35	20	12	0	3	1	24	10
Cleveland.....	936,485	103	47	101	2	5	8	0	22
Columbus.....	279,836	18	8	13	0	1	1	0	6
Toledo.....	287,380	81	17	9	0	0	8	0	5
Indiana:									
Fort Wayne.....	97,846	9	5	6	0	0	3	0	3
Indianapolis.....	358,819	62	13	22	0	0	1	0	16
South Bend.....	80,091	4	2	5	0	0	21	0	2
Terre Haute.....	71,071	6	3	0	0	0	0	0	1
Illinois:									
Chicago.....	2,995,239	201	148	52	13	5	184	65	47
Peoria.....	81,564	12	3	1	0	0	85	10	3
Springfield.....	63,923	20	2	4	0	0	68	0	3
Michigan:									
Detroit.....	1,245,824	121	72	82	1	5	6	28	24
Flint.....	130,316	25	14	4	0	0	1	0	3
Grand Rapids.....	153,698	15	6	0	1	1	0	0	3
Wisconsin:									
Kenosha.....	50,891	34	2	1	0	0	7	2	0
Madison.....	46,385	37	1	1	0	0	16	0	0
Milwaukee.....	509,192	88	30	22	1	0	10	51	11
Racine.....	67,707	27	3	2	0	0	0	5	0
Superior.....	39,671	1	1	1	0	0	0	0	0
WEST NORTH CENTRAL									
Minnesota:									
Duluth.....	110,502	10	2	0	0	0	38	0	1
Minneapolis.....	425,435	198	26	33	0	2	0	0	8
St. Paul.....	246,001	33	21	3	0	0	7	0	12
Iowa:									
Davenport.....	52,469	2	2	0	0	-----	6	0	-----
Des Moines.....	141,441	0	6	1	0	-----	0	0	-----
Sioux City.....	76,411	17	3	1	0	-----	0	0	-----
Waterloo.....	36,771	44	1	0	0	-----	0	0	-----
Missouri:									
Kansas City.....	367,481	56	14	9	3	3	2	0	17
St. Joseph.....	78,342	3	4	1	0	0	0	0	5
St. Louis.....	821,543	49	59	44	1	2	8	5	-----
North Dakota:									
Fargo.....	26,403	4	0	0	0	0	3	0	2
Grand Forks.....	14,811	1	0	0	0	-----	29	0	-----
South Dakota:									
Aberdeen.....	15,036	25	1	0	0	-----	2	0	-----
Sioux Falls.....	30,127	1	0	0	0	-----	0	0	-----
Nebraska:									
Lincoln.....	60,941	9	2	0	0	0	2	0	2
Omaha.....	211,768	10	6	5	0	0	5	13	10
Kansas:									
Topeka.....	55,411	3	3	0	0	0	1	0	0
Wichita.....	88,367	25	8	0	0	0	0	1	1

City reports for week ended December 11, 1926—Continued

Division, State, and city	Population July 1, 1925, estimated	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases reported	Mumps, cases reported	Pneumonia, deaths reported
			Cases, estimated expectancy	Cases reported	Cases reported	Deaths reported			
SOUTH ATLANTIC									
Delaware:									
Wilmington	122,049	2	4	1	0	0	1	0	
Maryland:									
Baltimore	796,296	102	36	33	20	3	2	8	32
Cumberland	33,741	4	2	1	0	0	0	1	0
Frederick	12,035	12	1	1	0	0	0	0	0
District of Columbia:									
Washington	497,006	52	24	23	0	2	0	0	12
Virginia:									
Lynchburg	30,395	6	2	2	0	0	1	1	2
Norfolk	(1)	0	4	3	0	0	0	0	5
Richmond	186,403	3	13	17	0	2	9	1	4
Roanoke	58,208	2	4	4	0	2	0	0	2
West Virginia:									
Charleston	49,019	14	3	4	1	0	0	0	0
Huntington	63,485	0	2	1	0	0	0	0	1
Wheeling	56,208	14	3	3	0	0	0	1	2
North Carolina:									
Raleigh	30,371	6	2	3	0	0	0	0	1
Wilmington	37,061	8	1	2	0	0	0	0	1
Winston-Salem	69,031	2	2	6	0	0	0	0	4
South Carolina:									
Charleston	73,125	0	2	0	15	1	0	0	3
Columbia	41,225	3	1	1	0	0	1	0	0
Greenville	27,311	8	0	2	0	0	0	0	1
Georgia:									
Atlanta	(1)	4	5	18	22	6	7	0	5
Brunswick	16,809	4	0	0	0	0	0	0	0
Savannah	93,134	1	2	1	14	2	0	0	5
Florida:									
Miami	69,754	6		2	1	0	1	0	3
St. Petersburg	26,847		1			0			1
Tampa	94,743	1	2	2	0	0	8	0	2
EAST SOUTH CENTRAL									
Kentucky:									
Covington	58,309		3						
Louisville	305,935	14	13	8	1	2	1	0	12
Tennessee:									
Memphis	174,533	13	16	6	0	3	8	0	6
Nashville	136,220	3	5	14	0	0	0	0	5
Alabama:									
Birmingham	205,670	18	6	11	7	3	0	2	6
Mobile	65,955	0	2	3	0	0	6	0	2
Montgomery	46,481	15	1	8	0	0	0	0	0
WEST SOUTH CENTRAL									
Arkansas:									
Fort Smith	31,643	0	2	5	0		0	5	
Little Rock	74,216	4	2	1	0		1	0	2
Louisiana:									
New Orleans	414,493	1	12	13	10	6	30	0	13
Shreveport	57,857	10	1	4	0	0	1	5	4
Oklahoma:									
Oklahoma City	(1)	0	3	0	0	0	0	0	3
Texas:									
Dallas	194,450	4	12	21	1	1	0	0	2
Galveston	48,375	0	1	0	0	0	0	0	3
Houston	164,954	5	5	8	0	1	0	0	5
San Antonio	198,069	0	4	10	0	1	2	0	5
MOUNTAIN									
Montana:									
Billings	17,971	2	0	0	0	0	39	0	0
Great Falls	29,883	14	0	0	0	1	3	0	0
Helena	12,037	0	0	0	0	0	1	0	0
Missoula	12,668	2	0	1	0	0	0	3	0
Idaho:									
Boise	23,042	4	0	0	0	0	11	0	0

1 No estimate made.

City reports for week ended December 11, 1926—Continued

Division, State, and city	Population July 1, 1925, estimated	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases reported	Mumps, cases reported	Pneumonia, deaths reported
			Cases, estimated expectancy	Cases reported	Cases reported	Deaths reported			
MOUNTAIN—continued									
Colorado:									
Denver.....	280,911	15	13	15	—	3	30	0	7
Pueblo.....	43,787	6	5	0	0	0	0	0	0
New Mexico:									
Albuquerque.....	21,000	4	1	0	0	0	5	1	0
Arizona:									
Phoenix.....	38,669	0	0	0	0	0	0	0	2
Utah:									
Salt Lake City.....	130,948	25	4	10	0	0	269	1	5
Nevada:									
Reno.....	12,665	1	0	1	0	0	0	0	0
PACIFIC									
Washington:									
Seattle.....	(1)	55	7	9	0	—	9	27	—
Spokane.....	108,897	38	5	1	0	—	97	36	—
Tacoma.....	104,455	14	3	7	0	0	0	0	2
Oregon:									
Portland.....	282,383	11	10	13	0	0	6	1	10
California:									
Los Angeles.....	(1)	80	37	56	23	3	13	10	25
Sacramento.....	72,260	7	3	2	0	0	33	22	3
San Francisco.....	557,520	32	19	14	2	0	77	38	2

Division, State, and city	Scarlet fever		Smallpox			Tuber- culosis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, estimated expectancy	Cases re- ported	Cases, estimated expectancy	Cases re- ported	Deaths re- ported		Cases, estimated expectancy	Cases re- ported	Deaths re- ported		
NEW ENGLAND											
Maine:											
Portland.....	2	3	0	0	0	0	0	0	0	21	20
New Hampshire:											
Concord.....	0	4	0	0	0	0	0	0	0	6	13
Manchester.....	2	3	0	0	0	1	0	0	0	6	16
Vermont:											
Barre.....	1	0	0	0	0	1	0	0	0	1	4
Massachusetts:											
Boston.....	43	78	0	0	0	15	1	0	0	33	226
Fall River.....	2	1	0	0	0	4	1	0	0	0	26
Springfield.....	8	8	0	0	0	0	0	0	0	1	23
Worcester.....	11	16	0	0	0	4	0	0	0	4	42
Rhode Island:											
Pawtucket.....	1	1	0	0	0	0	0	0	0	2	15
Providence.....	6	5	0	0	0	1	1	0	0	5	57
Connecticut:											
Bridgeport.....	7	20	0	0	0	0	0	0	0	4	29
Hartford.....	6	6	0	0	0	1	1	0	0	1	33
New Haven.....	7	2	0	0	0	1	1	1	0	0	37
MIDDLE ATLANTIC											
New York:											
Buffalo.....	22	15	1	0	0	9	2	0	1	8	116
New York.....	153	201	0	2	0	91	18	14	3	49	1,484
Rochester.....	12	14	0	0	0	5	1	14	4	4	96
Syracuse.....	12	15	0	0	0	1	1	0	0	9	44
New Jersey:											
Camden.....	3	5	0	0	0	1	0	1	0	0	34
Newark.....	16	26	0	0	0	5	1	2	0	26	114
Trenton.....	3	1	0	0	0	5	1	0	0	5	35
Pennsylvania:											
Philadelphia.....	66	66	0	0	0	30	4	4	0	39	521
Pittsburgh.....	36	12	0	0	0	6	1	0	0	10	143
Reading.....	1	1	0	0	0	0	0	1	0	6	30

¹ No estimate made.² Pulmonary tuberculosis only.

City reports for week ended December 11, 1926—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuber- culosis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, esti- mated expect- ancy	Cases re- ported	Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		
EAST NORTH CENTRAL											
Ohio:											
Cincinnati.....	13	24	0	0	0	14	1	0	0	0	152
Cleveland.....	33	25	0	2	0	17	2	1	0	11	180
Columbus.....	12	14	0	0	0	2	0	0	0	0	82
Toledo.....	13	10	0	0	0	5	0	0	0	27	65
Indiana:											
Fort Wayne.....	2	1	0	0	0	3	1	1	0	0	22
Indianapolis.....	12	17	4	7	0	7	0	1	0	10	107
South Bend.....	3	4	0	0	0	2	0	0	0	0	12
Terre Haute.....	3	3	0	0	0	1	0	0	0	0	17
Illinois:											
Chicago.....	117	100	1	0	0	38	7	1	0	50	657
Peoria.....	6	0	0	0	0	0	0	0	0	3	27
Springfield.....	2	1	0	0	0	1	1	0	0	9	23
Michigan:											
Detroit.....	85	96	2	0	0	23	3	1	2	52	290
Flint.....	8	21	0	1	0	2	1	0	0	4	30
Grand Rapids.....	8	12	0	0	0	0	1	0	0	0	42
Wisconsin:											
Kenosha.....	1	5	1	0	0	0	0	0	0	9	5
Madison.....	2	6	0	0	0	0	0	0	0	0	5
Milwaukee.....	26	18	2	0	0	6	1	0	0	59	121
Racine.....	4	2	1	0	0	0	0	0	0	2	11
Superior.....	2	2	1	0	0	1	0	0	0	0	11
WEST NORTH CENTRAL											
Minnesota:											
Duluth.....	6	11	1	0	0	2	0	0	0	1	21
Minneapolis.....	47	79	4	0	0	1	1	0	0	2	99
St. Paul.....	20	28	10	2	0	1	1	1	0	5	55
Iowa:											
Davenport.....	1	2	1	0	0	0	0	0	0	0	1
Des Moines.....	7	1	0	0	0	0	0	0	0	0	1
Sioux City.....	3	10	1	2	0	0	0	0	0	2	2
Waterloo.....	3	1	0	0	0	0	0	0	0	2	2
Missouri:											
Kansas City.....	11	23	0	2	0	10	1	0	0	2	97
St. Joseph.....	3	2	0	0	0	2	0	0	0	0	28
St. Louis.....	34	39	0	1	0	4	2	1	0	25	237
North Dakota:											
Fargo.....	2	0	0	0	0	0	0	0	0	0	6
Grand Forks.....	0	0	1	0	0	0	0	0	0	0	0
South Dakota:											
Aberdeen.....	1	18	0	0	0	0	0	0	0	4	0
Sioux Falls.....	2	0	1	0	0	0	0	0	0	0	0
Nebraska:											
Lincoln.....	2	4	0	0	0	0	0	0	0	0	16
Omaha.....	5	11	4	0	0	1	1	0	0	0	49
Kansas:											
Topeka.....	2	2	0	12	0	0	0	0	0	2	10
Wichita.....	3	8	0	0	0	1	0	0	1	0	23
SOUTH ATLANTIC											
Delaware:											
Wilmington.....	3	18	0	0	0	1	1	0	0	1	30
Maryland:											
Baltimore.....	23	22	1	0	0	14	3	3	0	53	216
Cumberland.....	0	1	1	0	0	0	1	1	0	4	10
Frederick.....	1	1	0	0	0	1	1	0	0	5	6
District of Colum- bia:											
Washington.....	20	8	0	0	0	9	4	1	1	8	136
Virginia:											
Lynchburg.....	0	7	0	0	0	0	0	0	0	0	15
Norfolk.....	2	0	0	0	0	2	0	1	0	3	0
Richmond.....	6	6	0	0	0	3	1	1	0	0	59
Roanoke.....	1	4	0	1	0	1	1	0	0	0	20

Division, State, and city	Scarlet fever		Smallpox			Tuber- culosis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, esti- mated expect- ancy	Cases re- ported	Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		
SOUTH ATLANTIC— continued											
West Virginia:											
Charleston	1	2	1	0	0	0	0	1	0	0	11
Huntington	1	1	0	0	0	3	0	0	0	0	13
Wheeling	2	0	0	0	0	0	1	1	0	5	16
North Carolina:											
Raleigh	1	1	0	0	0	1	0	0	0	20	14
Wilmington	0	0	0	0	0	1	0	0	0	2	11
Winston-Salem	2	3	1	0	0	1	0	0	0	4	17
South Carolina:											
Charleston	1	0	0	0	0	1	0	2	0	0	27
Columbia	0	0	0	0	0	0	0	0	0	0	0
Greenville	0	1	0	0	0	0	0	0	0	4	5
Georgia:											
Atlanta	5	15	1	8	0	5	1	2	1	4	65
Brunswick	0	0	0	0	0	0	0	0	0	0	3
Savannah	1	2	1	1	0	2	1	0	0	1	26
Florida:											
Miami		1		0	0	0		1	1	1	34
St. Petersburg	1		0		0	0	0		0	15	15
Tampa	0	2	0	0	0	2	0	0	1	0	26
EAST SOUTH CENTRAL											
Kentucky:											
Covington	2		0				0				
Louisville	5	6	0	1	0	1	1	1	0	14	92
Tennessee:											
Memphis	5	11	0	1	0	2	1	5	1	32	53
Nashville	3	6	1	0	0	6	1	2	0	0	42
Alabama:											
Birmingham	4	4	1	0	0	4	2	0	0	1	54
Mobile	1	0	0	2	0	1	0	0	0	0	18
Montgomery	1	0	1	0	0	0	0	0	0	0	28
WEST SOUTH CENTRAL											
Arkansas:											
Fort Smith	1	0	0	0			0	0		0	
Little Rock	2	1	0	0		3	1	0		0	
Louisiana:											
New Orleans	6	9	0	0	0	8	1	0	0	0	119
Shreveport	1	2	1	0	0	0	1	0	0	0	16
Oklahoma:											
Oklahoma City	3	1	0	0	0	2	0	0	0	0	27
Texas:											
Dallas	4	9	0	1	0	3	1	0	0	0	43
Galveston	0	7	0	0	0	0	0	1	0	0	12
Houston	2	4	1	1	0	9	0	0	0	0	63
San Antonio	1	1	0	0	0	9	1	2	0	0	55
MOUNTAIN											
Montana:											
Billings	1	0	0	0	0	0	0	0	0	0	7
Great Falls	2	3	1	0	0	0	0	0	0	0	6
Helena	0	1	0	0	0	0	0	0	0	0	
Missoula	1	5	1	0	0	0	0	0	0	0	7
Idaho:											
Boise	1	3	1	1	0	0	0	0</			

City reports for week ended December 11, 1926—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuber- culosis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, esti- mated expect- ancy	Cases re- ported	Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		
PACIFIC											
Washington:											
Seattle	7	8	3	0			0	2		0	
Spokane	6	24	3	2			1	0		9	
Tacoma	3	6	3	12	0	1	0	1	0	0	27
Oregon:											
Portland	7	24	6	5	0	7	0	1	0	0	87
California:											
Los Angeles	20	34	4	1	0	22	2	2	0	5	257
Sacramento	2	2	2	1	0	2	0	0	0	0	21
San Francisco	10	12	1	0	0	8	1	1	0	11	157

Division, State, and city	Cerebrospinal meningitis		Lethargic encephalitis		Pellagra		Polioomyelitis (infantile paralysis)			
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases, estimated expectancy	Cases	Deaths	
NEW ENGLAND										
Massachusetts:										
Boston.....	1	0	0	0	0	0	1	0	0	0
Fall River.....	0	1	0	0	0	0	0	0	0	0
Rhode Island:										
Pawtucket.....	0	0	0	0	0	0	0	1	0	0
Providence.....	0	0	0	0	0	0	0	1	0	0
MIDDLE ATLANTIC										
New York:										
New York.....	4	0	3	2	0	1	2	1	1	1
Rochester.....	0	0	0	0	0	0	0	1	0	0
New Jersey:										
Newark.....	0	0	0	0	0	0	1	1	0	0
Pennsylvania:										
Pittsburgh ¹	0	1	0	0	0	0	0	0	0	0
EAST NORTH CENTRAL										
Ohio:										
Cleveland.....	1	0	0	0	0	0	0	1	0	0
Columbus.....	0	0	0	0	0	0	0	0	1	1
Illinois:										
Chicago.....	2	0	1	0	0	0	1	1	0	0
Michigan:										
Detroit.....	1	0	2	0	0	0	1	0	0	0
WEST NORTH CENTRAL										
Minnesota:										
Duluth.....	1	0	0	0	0	0	0	0	0	0
Missouri:										
St. Louis.....	2	1	0	0	0	0	0	0	0	0
SOUTH ATLANTIC										
Maryland:										
Baltimore.....	0	0	2	1	0	0	0	0	1	1
Virginia:										
Norfolk.....	0	0	1	0	0	0	0	0	0	0
North Carolina:										
Wilmington.....	0	0	0	0	0	1	0	0	0	0
South Carolina:										
Charleston.....	0	0	0	0	1	0	0	0	0	0
Georgia:										
Atlanta.....	0	0	0	0	1	0	0	0	0	0

¹ Rabies (human); 1 death at Pittsburgh, Pa.

City reports for week ended December 11, 1926--Continued

Division, State, and city	Cerebrospinal meningitis		Lethargic encephalitis		Pellagra		Poliomyelitis (infantile paralysis)		
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases, estimated expectancy	Cases	Deaths
EAST SOUTH CENTRAL									
Tennessee:									
Memphis.....	1	1	0	0	0	0	0	0	0
Nashville.....	0	0	0	0	0	0	0	1	0
Alabama:									
Birmingham.....	0	0	0	0	1	0	0	0	0
WEST SOUTH CENTRAL									
Louisiana:									
Shreveport.....	0	0	0	0	0	1	0	0	0
Texas:									
Galveston.....	0	0	0	0	0	1	0	0	0
MOUNTAIN									
Montana:									
Missoula.....	1	0	0	0	0	0	0	0	0
New Mexico:									
Albuquerque.....	0	0	0	0	0	0	0	1	1
PACIFIC									
Washington:									
Spokane.....	1	0	0	0	0	0	0	0	0
California:									
Los Angeles.....	0	0	0	0	0	0	0	1	0
San Francisco.....	0	0	1	0	0	0	1	1	0

The following table gives the rates per 100,000 population for 101 cities for the five-week period ended December 11, 1926, compared with those for a like period ended December 12, 1925. The population figures used in computing the rates are approximate estimates as of July 1, 1925 and 1926, respectively, authoritative figures for many of the cities not being available. The 101 cities reporting cases had an estimated aggregate population of nearly 30,000,000 in 1925 and nearly 30,500,000 in 1926. The 95 cities reporting deaths had more than 29,200,000 estimated population in 1925 and more than 29,730,000 in 1926. The number of cities included in each group and the estimated aggregate populations are shown in a separate table below.

Summary of weekly reports from cities, November 7 to December 11, 1926—Annual rates per 100,000 population, compared with rates for the corresponding period of 1925¹

DIPHTHERIA CASE RATES

	Week ended—									
	Nov. 14, 1925	Nov. 13, 1926	Nov. 21, 1925	Nov. 20, 1926	Nov. 28, 1925	Nov. 27, 1926	Dec. 5, 1925	Dec. 4, 1926	Dec. 12, 1925	Dec. 11, 1926
101 cities.....	169	229	176	230	154	212	165	² 225	159	³ 201
New England.....	122	135	139	139	101	132	120	173	103	163
Middle Atlantic.....	140	162	143	159	150	154	137	176	138	160
East North Central.....	185	264	180	292	155	257	164	267	158	223
West North Central.....	235	222	221	213	170	191	272	² 221	239	193
South Atlantic.....	236	391	271	278	207	284	207	242	192	239
East South Central.....	63	265	121	368	110	218	116	301	121	³ 275
West South Central.....	203	379	167	327	172	301	264	318	176	267
Mountain.....	240	182	305	146	129	200	231	228	166	246
Pacific.....	138	232	177	326	157	305	122	270	191	240

MEASLES CASE RATES

101 cities.....	169	105	222	135	205	133	342	² 177	427	³ 199
New England.....	903	31	1,090	47	798	57	1,526	102	1,953	165
Middle Atlantic.....	170	44	255	28	238	30	338	37	451	23
East North Central.....	84	100	97	121	118	131	243	145	293	218
West North Central.....	10	147	14	197	29	109	18	² 127	25	129
South Atlantic.....	217	24	271	54	330	23	516	49	539	54
East South Central.....	16	10	47	31	32	16	37	26	21	³ 83
West South Central.....	9	26	9	26	4	103	4	142	4	146
Mountain.....	46	1,529	28	1,948	9	2,540	9	2,840	37	3,214
Pacific.....	19	280	30	491	25	340	55	704	52	617

SCARLET FEVER CASE RATES

101 cities.....	182	207	178	213	197	215	211	² 242	223	³ 238
New England.....	237	352	201	331	206	286	216	326	187	340
Middle Atlantic.....	142	125	143	129	149	137	166	156	172	177
East North Central.....	180	185	187	202	210	202	261	239	288	236
West North Central.....	354	346	401	407	438	411	405	² 459	476	431
South Atlantic.....	161	178	115	145	134	158	119	182	152	175
East South Central.....	168	296	126	228	168	239	163	244	110	² 149
West South Central.....	114	142	88	116	132	198	106	211	141	142
Mountain.....	176	701	157	637	166	783	240	929	157	801
Pacific.....	196	280	188	337	237	251	215	267	185	232

SMALLPOX CASE RATES

101 cities.....	8	5	16	5	16	5	13	² 14	21	³ 11
New England.....	0	0	0	0	0	0	0	0	0	0
Middle Atlantic.....	0	0	0	0	0	0	0	1	0	1
East North Central.....	13	10	31	3	31	7	13	21	33	7
West North Central.....	4	10	16	4	10	30	18	² 57	18	38
South Atlantic.....	6	2	19	4	2	4	4	19	8	19
East South Central.....	32	10	11	0	11	5	11	0	5	² 22
West South Central.....	0	30	0	4	9	4	13	9	9	9
Mountain.....	18	9	18	0	9	0	0	18	102	18
Pacific.....	41	5	75	49	94	5	105	35	124	43

¹ The figures given in this table are rates per 100,000 population, annual basis, and not the number of cases reported. Populations used are estimated as of July 1, 1925 and 1926, respectively.

² Kansas City, Mo., not included.

³ Covington, Ky., not included.

Summary of weekly reports from cities, November 7 to December 11, 1926—Annual rates per 100,000 population, compared with rates for the corresponding period of 1925—Continued

TYPHOID FEVER CASE RATES

	Week ended—									
	Nov. 14, 1925	Nov. 13, 1926	Nov. 21, 1925	Nov. 20, 1926	Nov. 28, 1925	Nov. 27, 1926	Dec. 5, 1925	Dec. 4, 1926	Dec. 12, 1925	Dec. 11, 1926
101 cities.....	11	21	17	16	13	12	19	10	20	11
New England.....	2	9	31	7	17	7	22	7	22	2
Middle Atlantic.....	8	21	20	21	14	13	26	9	25	11
East North Central.....	9	10	3	5	3	4	8	6	12	3
West North Central.....	16	16	14	6	8	8	10	19	12	4
South Atlantic.....	10	36	29	23	27	19	19	17	23	24
East South Central.....	42	52	32	36	21	31	53	42	26	44
West South Central.....	57	34	31	13	31	17	40	9	31	13
Mountain.....	9	27	18	27	18	18	0	9	18	9
Pacific.....	3	30	6	30	14	22	14	16	14	16

INFLUENZA DEATH RATES

95 cities.....	11	14	8	10	9	10	11	14	13	17
New England.....	7	2	2	2	12	9	10	7	10	9
Middle Atlantic.....	14	10	6	10	8	7	10	13	12	12
East North Central.....	10	10	6	10	5	9	6	9	11	14
West North Central.....	13	13	2	6	2	2	6	12	6	15
South Atlantic.....	2	17	13	8	10	15	17	21	8	34
East South Central.....	26	26	42	31	26	42	42	42	47	44
West South Central.....	29	71	10	33	34	33	39	43	44	43
Mountain.....	0	27	18	9	9	36	18	46	18	36
Pacific.....	4	14	18	4	4	0	4	11	4	11

PNEUMONIA DEATH RATES

95 cities.....	132	106	146	123	126	126	144	123	130	129
New England.....	120	90	139	104	156	132	180	118	132	135
Middle Atlantic.....	143	114	160	135	145	138	161	150	132	139
East North Central.....	131	85	139	106	95	99	142	87	116	103
West North Central.....	81	76	101	120	81	74	54	72	84	118
South Atlantic.....	152	139	146	143	134	165	159	105	173	154
East South Central.....	163	166	221	171	179	104	131	135	184	171
West South Central.....	102	113	155	156	150	213	155	161	208	151
Mountain.....	176	155	222	109	157	146	157	209	176	109
Pacific.....	109	99	87	75	98	124	98	153	76	114

¹ Kansas City, Mo., not included.

² Covington, Ky., not included.

³ Rochester, N. Y., and Covington, Ky., not included.

⁴ Rochester, N. Y., not included.

Number of cities included in summary of weekly reports, and aggregate population of cities in each group, approximated as of July 1, 1925 and 1926, respectively

Group of cities	Number of cities reporting cases	Number of cities reporting deaths	Aggregate population of cities reporting cases		Aggregate population of cities reporting deaths	
			1925	1926	1925	1926
Total.....	101	95	29, 900, 058	30, 427, 598	29, 221, 531	29, 733, 613
New England.....	12	12	2, 176, 124	2, 206, 124	2, 176, 124	2, 206, 124
Middle Atlantic.....	10	10	10, 346, 970	10, 476, 970	10, 346, 970	10, 476, 970
East North Central.....	16	16	7, 481, 656	7, 655, 436	7, 481, 656	7, 655, 436
West North Central.....	12	10	2, 550, 024	2, 589, 131	2, 431, 253	2, 468, 448
South Atlantic.....	21	21	2, 716, 070	2, 776, 070	2, 716, 070	2, 776, 070
East South Central.....	7	7	993, 103	1, 004, 953	993, 103	1, 004, 953
West South Central.....	8	6	1, 184, 057	1, 212, 057	1, 078, 198	1, 103, 695
Mountain.....	9	9	563, 912	572, 773	563, 912	572, 773
Pacific.....	6	4	1, 888, 142	1, 934, 084	1, 434, 245	1, 469, 144

FOREIGN AND INSULAR

PLAGUE ON VESSEL

Steamship "Dacia"—*At Haifa, Syria.*—On November 17, 1926, a case of plague was reported on the steamship *Dacia* at Haifa, Syria, occurring in a seaman. The vessel came from Rumania.

BERMUDA

Leprosy—Care and treatment of patients.—Reports of leprosy in the island of Bermuda, received under date of December 10, 1926, show for September, 1925, 8 lepers present, 3 male, 5 female, and for September, 1926, 9 lepers, 3 male and 6 female; one man and one woman, white; the remaining cases, colored. The isolation hospital not being equipped for the treatment of these cases, the lepers are cared for mainly by the parishes in which they reside and where they are segregated. The treatment includes administration of chaulmestrol.

CANADA

Communicable diseases—Week ended December 4, 1926.—The Canadian Ministry of Health reports cases of certain communicable diseases in seven Provinces of Canada for the week ended December 4, 1926, as follows:

Disease	Nova Scotia	New Brunswick	Quebec	Ontario	Manitoba	Saskatch- ewan	Alberta	Total
Influenza.....	27							27
Smallpox.....				11	2	20	6	39
Typhoid fever.....	2	4	9	12	2	1	3	33

CHINA

Plague—Mongolia.—Information received under date of December 18, 1926, shows epidemic pneumonic plague present at Urga and Sanbese, Mongolia. Prophylactic measures were stated to have been put in force at Manchuria Station, on the South Manchuria Railway.

EGYPT

Plague—November 12-18, 1926.—During the week ended November 18, 1926, 1 case of plague, occurring in the district of Tantah, was reported in Egypt, making a total from January 1 to November 18, 1926, of 143 cases as compared with 137 cases reported for the corresponding period of 1925.

Alexandria—November 23, 1926.—On November 23, 1926, a case of bubonic plague was reported at Alexandria.

Gharbieh.—From November 22 to 23, 1926, 2 cases of plague with 1 death were reported at Tanta, Province of Gharbieh.

FRENCH SUDAN

Yellow fever—Segou—November 23, 1926.—Under date of November 23, 1926, a fatal case of yellow fever was reported at Segou, French Sudan, West Africa.

GREECE

Plague—Patras—November 9–13, 1926.—Three cases of plague have been reported at Patras, Greece, occurring November 9, 11, and 13, respectively.

Typhus fever.—During the month of October, 1926, 7 cases of typhus fever with 1 death were reported in Greece.

MADAGASCAR

Plague—October 1 to 15, 1926.—During the two weeks ended October 15, 1926, 121 cases of plague with 111 deaths were reported in the island of Madagascar. The occurrence was distributed according to provinces as follows: Maevatanana, cases 17, deaths 17; Majunga, cases 6, deaths 2; Moramanga, cases 18, deaths, 18; Tamatave, cases 1, deaths 1; Tananarive (town), cases 16; deaths 14; other localities, cases 63, deaths 59.

Deaths among Europeans.—Of the 14 deaths from plague reported in the town of Tananarive 3 deaths were in Europeans, making a total of 5 deaths of Europeans from plague since August, 1926.

MEXICO

Malaria—Vicinity of Vera Cruz.—Information received from Vera Cruz under date of December 8, 1926, shows malaria present at Palmar, a small locality in the vicinity of Vera Cruz, with 2 fatalities reported to November 10, 1926. A physician of the State medical service has been in charge of the situation since that date. It was stated that at the outset he treated daily from 100 to 120 cases of malaria, with a few cases of dysentery. Population of Palmar, 350, including residents of near-by ranches.

SENEGAL

Further relative to plague—November 22, 1926.—Under date of November 23, 1926, 2 new cases of bubonic plague were reported in the interior of Senegal, West Africa. The cases occurred in natives and in the district of Diourbel.

Yellow fever.—Yellow fever was reported in Senegal, November 23, 1926, as follows: Four cases with 4 deaths, 1 case occurring in the

district of Kolda (Casamance) and 3 cases in the district of Sine Saloum. Of these cases, 3 were in Syrians and 1 in a European.

UNION OF SOUTH AFRICA

Plague—Cape Province—October 31—November 6, 1926.—During the week ended November 6, 1926, a case of plague, occurring in a native on a farm in Colesberg district, was reported in the Cape Province, Union of South Africa.

Smallpox—Natal.—During the same period 7 additional cases of smallpox were reported at Durban, Natal, making a total of 49 cases with 9 deaths reported to date, occurring in Hindus or natives.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER

The reports contained in the following tables must not be considered as complete or final as regards either the lists of countries included or the figures for the particular countries for which reports are given.

Reports Received During Week Ended December 31, 1926¹

CHOLERA

Place	Date	Cases	Deaths	Remarks
China:				
Amoy.....	Oct. 31-Nov. 6....	1		
India:				
Calcutta.....	Oct. 24-30.....	18	11	
Rangoon.....	Oct. 31-Nov. 6....	1	1	
Persia:				
Teheran.....	Aug. 23-Sept. 23..	1		

PLAGUE

China:				
Mongolia—				
Sanbese.....	Dec. 18.....			Epidemic pneumonic.
Urga.....	do.....			Do.
Egypt:				Jan. 1-Nov. 18, 1926: Cases, 143.
Garbieh Province.....	Nov. 22-23.....	2	1	Corresponding period, 1925:
Tantah District.....	Nov. 12-18.....	1		Cases, 137.
City—				
Alexandria.....	Nov. 23.....	1		
Greece:				
Patras.....	Nov. 9-13.....	3		
India:				
Madras Presidency.....	Oct. 17-23.....	97	55	
Rangoon.....	Oct. 31-Nov. 6....	3	4	
Java:				
Batavia.....	do.....	5	4	Province.
Madagascar				Oct. 1-15, 1926: Cases, 121;
				deaths, 111.
Maevatanana.....	Oct. 1-15.....	17	17	Bubonic, pneumonic, septicemic.
Majunga.....	do.....	6	2	Bubonic.
Moramanga.....	do.....	18	18	Bubonic, septicemic.
Tamatave.....	do.....	1	1	Bubonic.
Tananarive (Town).....	do.....	16	14	Bubonic, pneumonic, septicemic.
				Of the deaths, 3 were in Euro-
				peans; total European deaths
				of plague from August, 1926, 5.
				Bubonic, pneumonic, septicemic.
Other localities.....	do.....	63	59	
Senegal.....	Nov. 23.....	2		
Union of South Africa:				
Cape Province—				
Colesberg District.....	Oct. 31-Nov. 6....	1		
On vessel:				
Steamship Dacia.....	Nov. 17.....	1		At Haifa, Syria. Seaman on
				Rumanian steamship.

¹ From medical officers of the Public Health Service, American consuls, and other sources.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received During Week Ended December 31, 1926—Continued

SMALLPOX

Place	Date	Cases	Deaths	Remarks
Algeria:				
Algiers.....	Nov. 1-10.....	1		
Constantine.....	Nov. 24.....	44		Eastern department.
Brazil:				
Bahia.....	Oct. 24-30.....	6	2	
British South Africa:				
Northern Rhodesia.....	Oct. 30-Nov. 5....	1		
Canada:				
Alberta.....	Nov. 28-Dec. 4....	6		
Manitoba.....	do.....	2		
Ontario.....	do.....	11		
Toronto.....	Dec. 5-11.....	3		
Saskatchewan.....	Nov. 28-Dec. 4....	20		
France:				
Paris.....	Nov. 11-20.....	4	1	
India:				
Calcutta.....	Oct. 24-30.....	4	4	
Madras.....	Nov. 7-13.....	3		
Java:				
Batavia.....	Oct. 31-Nov. 6....	5		
Surabaya.....	Oct. 17-23.....	4	1	For East Java and Madura.
Mexico:				
Ciudad Juarez.....	Dec. 7-13.....	1		
Mexico City.....	Nov. 28-Dec. 4....	2		Including municipalities in Federal District.
Torreón.....	Nov. 21-27.....		1	
Persia:				
Teheran.....	Aug. 23-Sept. 23..		4	
Poland.....				Sept. 27-Oct. 9, 1926: One case.
Portugal:				
Lisbon.....	Nov. 21-27.....	5		
Union of South Africa:				
Natal—				
Durban.....	Oct. 10-Nov. 6....	50	10	
Polela.....	Oct. 31-Nov. 6....			Outbreak. In Nkandhla District.
Transvaal—				
Johannesburg.....	Nov. 7-13.....	1		

TYPHUS FEVER

Chosen:				
Seoul.....	Oct. 25-31.....	1		
Greece.....				Oct., 1926: Cases, 7; deaths, 1.
Ireland (Irish Free State):				
Ennistymon.....	July 4-10.....	5		
Mexico:				
Mexico City.....	Nov. 28-Dec. 4....	10		Including municipalities in Federal district.
Palestine:				
Haifa.....	Nov. 9-15.....	1		
Persia:				
Teheran.....	Aug. 23-Sept. 23..		2	
Poland.....	Sept. 27-Oct. 16..	52	5	
Krakow.....	Oct. 17-23.....	31	5	
Union of South Africa:				
Cape Province—				
Alexandria District.....	Oct. 31-Nov. 6....			Outbreak. In one locality.

YELLOW FEVER

French Sudan:				
Segou.....	Nov. 23.....	1	1	
Senegal.....				Nov. 23, 1926: Cases, 4; deaths, 4. One European.
Kolda District.....	Nov. 23.....	1	1	
Sine Saloum.....	do.....	3	3	

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received From June 26 to December 31, 1926¹

CHOLERA

Place	Date	Cases	Deaths	Remarks
Ceylon.....				Apr. 18-May 29, 1926: Cases, 31; deaths, 29.
China:				
Amoy.....	Aug. 8-Nov. 6.....	275		
Antung.....	Aug. 1-31.....	500		
Canton.....	June 1-30.....	38	14	
Do.....	July 15-31.....	54	28	
Do.....	Aug. 25-31.....	30	8	
Changsha.....	Oct. 3-18.....	2		
Foochow.....	Aug. 15-Oct. 2.....		1	In foreign population.
Kulangsu.....	Sept. 12-18.....		2	
Manchuria—				
Changshun.....	Aug. 1-31.....	320		
Dairen.....	do.....	10	1	
Harbin.....	Aug. 5-Sept. 12.....	289	83	
Newchwang.....	Aug. 1-31.....	167		
Nanking.....	July 25-Oct. 2.....			Present.
Shanghai.....	Reported July 20.....	35	8	
Do.....	July 25-Oct. 23.....	43	420	Cases, foreign; deaths, native and foreign.
Swatow.....	July 11-Oct. 16.....	50	63	
Tsingtao.....	July 11-Aug. 30.....	4	4	Japanese settlements, 10 deaths; Chinese, 30 to 40 deaths daily, estimated.
Do.....	Oct. 10-30.....			Present.
Chosen:				
North Heian Province.....	Sept. 3-16.....	70	30	Deaths estimated.
Shingishu.....	Sept. 13.....	19		Including places in vicinity.
French Settlements in India.....	Mar. 7-June 26.....	31	30	
Do.....	June 27-Aug. 28.....	94	83	
India:				
Bombay.....	May 30-June 5.....	1	1	Apr. 25-June 26, 1926: Cases, 18,526; deaths, 11,531.
Do.....	July 18-Oct. 16.....	4	4	June 27-Oct. 9, 1926: Cases, 28,544; deaths, 17,966.
Calcutta.....	Apr. 4-May 29.....	478	418	
Do.....	June 13-26.....	73	69	
Do.....	June 27-Oct. 30.....	366	320	
Madras.....	May 16-June 5.....	2	1	
Do.....	Aug. 1-Sept. 25.....	7	6	
Rangoon.....	May 9-June 26.....	67	44	
Do.....	June 27-Nov. 6.....	33	31	
Indo-China:				
Saigon.....	May 2-15.....	52	48	
Do.....	May 22-June 26.....	42	32	
Do.....	June 27-Aug. 14.....	31	17	
Japan:				
Ken (Prefecture)—				To Sept. 10, 1926: Cases, 35.
Hiroshima.....	To Sept. 10.....	1		
Ilyogo.....	do.....	7		
Kagakawa.....	do.....	8		
Kanagawa.....	do.....	3		Including Yokohama.
Kochi.....	do.....	1		
Ookayama.....	do.....	7		
Osaka.....	do.....	6		
Taihoku.....	Sept. 1-10.....	2		
Wakayama.....	To Sept. 10.....	2		
Taiwan Island.....	Sept. 21-Oct. 10.....	11		
Persia:				
Teheran.....	Aug. 23-Sept. 23.....	1		
Philippine Islands:				
Manila.....	Dec. 29, 1925-Oct. 30, 1926.....	27	6	
Provinces—				
Albay.....	Apr. 18-21.....	1	1	
Davao.....	May 23-29.....	1		
Mindoro.....	Feb. 21-Mar. 6.....	3	3	
Pampanga.....	July 25-31.....	1	1	
Rizal.....	July 18-24.....	1		
Romblon.....	Dec. 14-31.....	42	43	
Do.....	Jan. 2-Mar. 27.....	41	35	
Siam:				
Bangkok.....	May 2-June 12.....	1,325	736	Apr. 1-Oct. 30, 1926: Cases, 7,706; deaths, 5,075.
Do.....	June 20-26.....	56	26	
Do.....	June 27-Oct. 30.....	99	69	
Straits Settlements:				
Singapore.....	July 4-17.....	2	1	
On vessel:				
Steamship Macedonia.....	Aug. 5.....	7		At Yokohama, Japan. Vessel sailed from Singapore July 19, 1926.

¹ From medical officers of the Public Health Service, American consuls, and other sources.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

PLAGUE

Place	Date	Cases	Deaths	Remarks
Algeria:				
Algiers.....	June 21-30.....	1	-----	Under date of July 16, 2 cases reported.
Do.....	July 1-20.....	1	-----	
Do.....	Sept. 23.....	1	-----	
Bona.....	Aug. 14.....	1	-----	
Oran.....	Sept. 21-Nov. 13.....	10	5	
Philippeville.....	Sept. 7.....	1	-----	
Sfax.....	Nov. 13.....	7	-----	
Argentina:				
Cordoba Province.....	Nov. 20.....	5	-----	
Azores:				
Fayal Island—				
Horta.....	Aug. 2-29.....	2	2	
St. Michaels Island.....	May 9-June 16.....	4	1	
Do.....	June 27-July 10.....	3	1	
Brazil:				
Paranagua.....	Oct. 8.....	-----	-----	Present.
British East Africa:				
Kenya—				
Kisumu.....	May 16-22.....	1	1	
Do.....	Aug. 17-Sept. 11.....	3	2	
Uganda.....	Mar. 1-June 30.....	732	574	
Do.....	July 1-Aug. 31.....	312	267	
Canary Islands:				
Las Palmas.....	Nov. 2.....	3	-----	Stated to be in locality removed from port.
Teneriffe.....	Aug. 2.....	2	-----	
Ceylon:				
Colombo.....	May 29-June 5.....	1	1	
Do.....	Oct. 31-Nov. 6.....	1	1	Provisional diagnosis.
Chile:				
Iquique.....	June 20-26.....	-----	1	
China:				
Amoy.....	Apr. 18-June 26.....	40	30	
Do.....	June 27-Aug. 7.....	28	-----	
Foochow.....	June 6-July 31.....	-----	-----	Several cases. Not epidemic.
Mongolia—				
Sanbese.....	Dec. 18.....	-----	-----	Epidemic, pneumonic.
Urga.....	do.....	-----	-----	Do.
Nanking.....	May 9-Oct. 23.....	-----	-----	Prevalent.
Swatow.....	July 25-31.....	14	-----	
Ecuador				
Chimborazo.....	January-June.....	9	2	January-June, 1926: Cases, 385; deaths, 154.
Guayaquil.....	May 16-June 30.....	6	-----	Rats taken, 766.
Do.....	July 1-Oct. 31.....	19	3	Rats taken, 30,914; found infected, 31.
Leon.....	January-June.....	43	19	Rats taken, 82,774; found infected, 115.
Loja.....	do.....	176	75	Localities, 2.
Tungurahua.....	do.....	83	29	Cantons, 2.
				At Ambato, Huachi, and Pica-yhua. Rats taken, 1,542.
Egypt.....				Jan. 1-Nov. 18, 1926: Cases, 143.
City—				
Alexandria.....	July 27-Nov. 23.....	7	1	
Suez.....	May 21-July 1.....	9	5	
Do.....	July 29.....	2	-----	
Provinces—				
Beheran.....	July 23-Aug. 15.....	4	1	
Beni-Suef.....	May 23-June 8.....	8	2	
Charkieh.....	July 27.....	1	1	
Gharbieh.....	June 2.....	1	1	
Do.....	Nov. 22-23.....	2	1	
Minieh.....	July 24.....	1	1	
Sidi Barrani.....	Sept. 30-Oct. 21.....	23	3	In western desert.
Tanta District.....	Oct. 22-Nov. 18.....	3	-----	
France:				
Marseille.....	July 8.....	1	1	Reported July 24.
Paris.....	Oct. 18.....	1	-----	
St. Denis.....	Reported Aug. 2.....	1	-----	Vicinity of Paris.
St. Ouen.....	Aug. 14.....	2	-----	Suburb of Paris.
Great Britain:				
Liverpool.....	Aug. 29-Sept. 4.....	2	1	
Greece:				
Athens.....	Apr. 1-May 31.....	16	4	Including Piræus.
Do.....	Aug. 1-Sept. 30.....	20	5	Do.
Patras.....	May 27-June 12.....	4	1	
Do.....	July 25-Nov. 13.....	12	5	
Zante.....	May 17.....	1	-----	

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

PLAGUE—Continued

Place	Date	Cases	Deaths	Remarks
Hawaii Territory:				
Hamakua	June 9			1 plague rodent trapped near Hamakua Mill.
Honokaa	Oct. 6	1	1	
Panauhau	July 18-24			Plague-infected rat trapped.
India				Apr. 25-June 16, 1926: Cases, 53,001; deaths, 41,576. June 27-Oct. 9, 1926: Cases, 10,024; deaths, 5,660.
Bombay	May 2-June 26	16	15	
Do	July 18-Oct. 9	13	12	
Karachi	May 23-June 26	15	13	
Do	July 11-17	1	1	
Madras Presidency	Apr. 25-June 26	162	93	
Do	July 4-Oct. 23	1,159	562	
Rangoon	May 9-June 26	20	15	
Do	June 27-Nov. 6	92	81	
Indo-China:				
Saigon	May 23-June 26	8	3	
Do	July 18-Aug. 7	2	1	
Iraq:				
Baghdad	Apr. 18-June 12	161	108	
Do	July 18-Sept. 11	4	4	
Japan:				
Yokohama	July 2-Aug. 10	9	8	
Java:				
Batavia	Apr. 24-June 19	65	65	
Do	June 26-Nov. 6	102	99	
Cheribon	Apr. 11-24	3	3	
Do	Sept. 12-18	1	1	
East Java and Madura	June 13-19	1	1	
Do	July 25-Oct. 16	1	2	
Surabaya	Aug. 22-Sept. 25	18	2	
Madagascar:				
Amboisira Province	May 1-15	4	4	Septicemic.
Antisirabi Province	June 16-30	4	4	
Itasy Province	do	17	10	
Do	Aug. 16-Sept. 30	8	8	
Maevatanana Province	Aug. 16-Oct. 15	19	19	
Majunga Province	June 16-30	10	6	
Do	Aug. 16-Oct. 15	72	68	
Mananjary Province	do	1	1	
Moramanga Province	Apr. 1-15	2	2	
Do	Sept. 1-Oct. 15	49	49	
Tamatave Province	Aug. 16-Oct. 15	21	16	
Tananarive Province				Apr. 1-June 30, 1926: Cases, 130, deaths, 120. July 1-Oct. 15, 1926: Cases, 276; deaths, 262.
Towns—				
Majunga	Aug. 1-15	14	10	
Tamatave (port)	May 16-31	1	1	
Do	July 1-Aug. 15	6	5	
Tananarive	Apr. 1-June 30	7	7	
Do	July 1-Oct. 15	48	45	
Mauritius:				
Port Louis	July 31	1	1	
Nigeria				Feb. 1-June 30, 1926: Cases, 191; deaths, 163. July 1-31, 1926: Cases, 121; deaths, 112.
Peru				May-June, 1926: Cases, 57; deaths, 16. July 1-Oct. 31, 1926: Cases, 125; deaths, 65 Present.
Departments—				
Ancash	May 1-31			
Do	July 1-Sept. 30	2		
Cajamarca	May 1-June 30	10	4	
Do	Aug. 1-Oct. 31	1		
Ica	May 1-31	1		
Do	July 1-31	1		
Junin	Sept. 1-30	21	20	
Lambayeque	Sept. 1-Oct. 31	5	2	
Libertad	May 1-31	4		
Do	Sept. 1-Oct. 31	11	2	
Lima	May 1-June 30	29	12	
Do	July 1-Oct. 31	82	40	
Piura	June 1-30	13		
Do	Oct. 1-31	2	1	
Russia				Jan. 1-Mar. 31, 1926: Cases, 37.
Senegal				Nov. 1-30, 1925: Cases, 3; deaths, 2. Mar. 1-June 30, 1926: Cases, 342; deaths, 213. Nov. 1-23, 1926: Cases, 57; deaths, 27.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

PLAGUE—Continued

Place	Date	Cases	Deaths	Remarks
Siam				Apr. 1-Oct. 30, 1926: Cases, 15; deaths, 10.
Bangkok	May 23-June 26	2	2	
Do.	July 18-24	1	1	
Straits Settlements:				
Singapore	May 2-8	1	1	
Do.	July 4-17	1	1	
Syria:				
Beirut	July 1-Aug. 10	2		
Do.	Oct. 15-20	3		
Tunisia	May 11-June 30	174		
Do.	July 1-Aug. 20	13		
Do.	Reported Nov. 27	57		
Kairouan	June 9	3		9 cases 30 miles south of Kairouan.
Turkey:				
Constantinople	Aug. 1-Sept. 25	7	4	
Union of South Africa:				
Cape Province	May 16-22	5	3	
Do.	Oct. 17-23	4	3	
Calvinia District	June 13-26	12	6	
Do.	June 27-Aug. 21	3	3	
Colesberg District	Oct. 31-Nov. 6	1		
Hanover District	Oct. 10-16	1	1	Native. On farm.
Kimberley District	Oct. 17-23	2	2	European.
Williston District	June 13-26	2		
Do.	June 27-July 3	1		
Do.	Oct. 17-30	4	3	
Orange Free State—				
Hoopstad District	Aug. 15-21	1		
Protestpan	May 9-22	3	3	
On vessel:				
Steamship Zaria	September, 1926	2	2	At Liverpool, England, from Lagos, Nigeria, West Africa; 29 plague-infected rats found on board.
Steamship Dacia	Nov. 17	1		At Haifa, Syria. Seaman on Rumanian steamship.

SMALLPOX

Algeria				July 21-Sept. 20, 1926: Cases, 230.
Algiers	May 21-June 30	14		
Do.	July 1-Nov. 10	4		
Constantine	Nov. 24	44		Eastern department.
Arabia:				
Aden	Oct. 3-9	1		Imported.
Belgium				Sept. 1-30, 1926: Cases, 2.
Antwerp	Aug. 1-7	1	1	
Bolivia:				
La Paz	May 1-June 30	14	7	
Do.	July 1-Aug. 31	16	8	
Brazil:				
Bahia	June 20-26	1		
Do.	June 27-Oct. 30	82	43	
Manaos	Apr. 1-30		5	
Para	May 16-June 26	26	25	
Do.	June 27-Oct. 30	33	27	
Pernambuco	July 11-Oct. 16	236	26	
Porto Alegre	Aug. 10-31	2		
Rio de Janeiro	May 2-June 19	132	91	
Do.	July 4-Sept. 25	2,534	1,338	
Do.	Oct. 3-Nov. 13	475	300	Jan. 1-Oct. 16, 1926: Cases, 3,601; deaths, 1,596.
Sao Paulo	June 27-Aug. 22		5	
Santos	Mar. 1-7		1	
British East Africa:				
Mombasa	July 5-11	5	4	
Tanganyika	May 1-31	252	46	
Do.	Aug. 29-Sept. 18	7		
Uganda	Mar. 1-May 31	3		
Do.	Aug. 1-31	1		
British South Africa:				
Northern Rhodesia	May 18-24	17	6	Natives.
Do.	June 8-14	5		
Do.	Sept. 11-Nov. 5	2		

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

SMALLPOX—Continued

Place	Date	Cases	Deaths	Remarks
Canada				May 30-June 26, 1926: Cases, 70. June 27-Dec. 4, 1926: Cases, 571.
Alberta				May 30-June 12, 1926: Cases, 3. June 27-Dec. 4, 1926: Cases, 92.
Calgary	Sept. 5-Nov. 27	50		
British Columbia—				
Vancouver	Aug. 16-Sept. 12	3		
Manitoba				May 30-June 26, 1926: Cases, 15. June 27-Dec. 4, 1926: Cases, 90.
Winnipeg	June 6-12	5		
Do	July 4-Dec. 11	18		
New Brunswick				Oct. 31-Nov. 6, 1926: 1 case.
Northumberland County.	Oct. 11-23	1		
Ontario				May 30-June 26, 1926: Cases, 36. June 27-Dec. 4: Cases, 223.
Fort William	July 25-Aug. 7	2		
Kingston	May 23-June 26	5		
Do	July 11-Nov. 6	3		
Kitchener	Apr. 26-May 29	3	1	
North Bay	May 2-22	5		
Do	July 25-31	2		
Orillia	Apr. 26-May 29	7		
Ottawa	July 18-24	1		
Do	Nov. 28-Dec. 4	1		
Packenham	do	10		
Peterboro	Sept. 1-30	10		
Toronto	July 18-Dec. 11	49		
Waterloo	July 18-24	6		
Saskatchewan				May 30-June 26, 1926: Cases, 16. June 27-Dec. 4: Cases, 165.
Regina	July 4-Sept. 25	3		Mar. 14-May 29, 1926: Cases, 41; deaths, 3. Sept. 12-13, 1926: Cases, 2.
Ceylon				
Colombo	Sept. 19-Oct. 16	7		
Chile				
Antofagasta	June 6-12	1		
China				
Amoy	May 1-June 26	4	8	
Do	July 4-10	1		
Antung	May 17-June 19	5		
Do	July 4-18	2		
Canton	May 1-31	4	2	
Do	Sept. 1-30	1		
Changsha	Aug. 8-14	1		
Chungking	May 2-Oct. 23			Present.
Foochow	May 2-Oct. 30			Do.
Fushun	Sept. 12-18	1		
Hongkong	May 2-June 26	19	10	
Do	June 27-July 3	1	1	
Manchuria	July 4-31	18		Railway stations.
An-shan	May 16-June 12	5		South Manchurian Railway.
Antung	May 16-June 19	5		
Changchun	May 16-June 26	6		Do.
Do	June 27-Sept. 11	2		Do.
Dairen	Apr. 26-June 20	69	16	
Do	June 28-Aug. 8	5	3	
Fushun	May 16-June 5	4		Do.
Harbin	May 14-June 30	21		Do.
Do	July 1-28	12		
Kai-yuan	May 16-June 30	10		Do.
Kungchuling	June 13-19	1		Do.
Liaoyang	May 16-June 30	4		Do.
Mukden	do	4		Do.
Penhshu	May 16-June 19	4		Do.
Do	Aug. 8-Oct. 3	3		Do.
Ssuninghai	May 16-June 30	2		Do.
Do	Aug. 1-7	1		Do.
Teshihchiao	May 16-June 30	2		Do.
Tieh-ling	Sept. 27-Oct. 3	1		
Wa-feng-tien	do	3		Do.
Do	Aug. 1-7	1		Do.
Nanking	May 8-Oct. 30			Present.
Shanghai	May 2-June 26	10	25	Cases, foreign: Deaths, popula-
Do	June 27-July 24	3	3	tion of international conces-
Do	Oct. 3-9	1		sion, foreign and native.
Swatow	May 9-Oct. 30			Sporadic.
Tientsin	June 2-26		1	Reported by British municipal-
Wanshien	May 1			ity.
				Prevalent.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

SMALLPOX—Continued

Place	Date	Cases	Deaths	Remarks
Chosen				Mar. 1-June 30, 1926: Cases, 667; deaths, 146. July 1-31, 1926: Cases, 82; deaths, 27.
Fusan	May 1-31	1		
Selshun	do	2	1	
Egypt:				
Alexandria	May 15-July 1	18	3	
Do	July 23-Oct. 28	15	7	
Cairo	Jan. 29-June 10	56	14	
Estonia				May 1-June 30, 1926: Cases, 3.
France				Mar. 1-June 30, 1926: Cases, 141. July 1-Aug. 31: Cases, 24.
Paris	Sept. 1-Nov. 20	69	19	
St. Etienne	Apr. 18-June 15	7	3	
Do	Sept. 16-30	2	1	
French Settlements in India	Mar. 7-June 26	282	282	
Do	June 27-Aug. 28	68	68	
Germany:				
Coblenz	Oct. 24-30	1		
Gold Coast	Mar. 1-June 30	671		
Do	July 1-31	20	1	
Great Britain:				
England and Wales				May 23-June 26, 1926: Cases, 933; June 27-Nov. 13, 1926: Cases, 2,415.
Birmingham	Sept. 26-Oct. 2	1		
Bradford	May 23-29	1		
Do	Aug. 29-Sept. 4	1		
Hull	Oct. 17-23	1		
London	Sept. 26-Oct. 23	4		
Newcastle-on-Tyne	June 6-12	1		
Do	July 11-Nov. 30	7		At Gateshead, several cases reported.
Nottingham	May 2-June 5	7		
Do	July 18-24	1		
Sheffield	June 13-19	1		
Do	July 4-Nov. 27	49		
South Shields	Oct. 3-9	1		
Stoke-on-Trent	Nov. 7-13	1		
Greece:				
Athens	July 1-31	71	6	Including Piræus.
Saloniki	June 1-14		3	
Guatemala:				
Guatemala City	June 1-30		2	
India:				
Bombay	May 2-June 26	220	134	Apr. 25-June 26, 1926: Cases, 54,851; deaths, 14,771. June 27-Oct. 9, 1926: Cases, 27,840; deaths, 8,445.
Do	June 27-Nov. 6	137	75	
Calcutta	Apr. 4-May 20	171	152	
Do	June 13-26	24	18	
Do	June 27-Oct. 30	53	47	
Karachi	May 6-June 26	44	18	
Do	June 27-Oct. 30	15	7	
Madras	May 16-June 26	7	4	
Do	June 27-Nov. 13	83	21	
Rangoon	May 9-June 26	10	5	
Do	July 4-Sept. 23	21	5	
Indo-China:				
Saigon	May 9-June 26	2		
Iraq:				
Baghdad	do	8	3	
Do	July 4-Sept. 11	3	1	
Basra	Apr. 18-June 22	34	25	
Do	Aug. 15-21	1		
Italy				
Catania	Aug. 9-15	2		Mar. 28-June 26, 1926: Cases, 34. June 27-Aug. 7, 1926: Cases, 12.
Rome	June 14-20	4		Entire consular district, including island of Sardinia.
Do	Aug. 30-Sept. 5	2		Do.
Jamaica				Apr. 25-June 26, 1926: Cases—201. (Reported as alastrim.)
Do				June 27-Nov. 27, 1926: Cases, 347. (Reported as alastrim.)
Japan				Apr. 11-June 26, 1926: Cases, 653. June 27-Aug. 28, 1926: Cases, 70.
Kobe	May 30-June 5	1		
Nagoya	May 16-June 22		1	
Do	July 4-10	1		
Taiwan Island	May 11-20	24		
Do	June 1-20	23		
Do	July 11-Aug. 10	2		
Tokyo	June 26-July 17	3		
Yokohama	May 2-8	2		

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

SMALLPOX—Continued

Place	Date	Cases	Deaths	Remarks
Java:				
Batavia.....	May 15-June 25...	2	-----	Province.
Do.....	July 24-Nov. 6.....	22	-----	
East Java and Madura.....	Apr. 11-July 3.....	100	6	
Do.....	July 4-Oct. 23.....	79	6	
Malang.....	Apr. 4-10.....	6	1	Interior.
Surabaya.....	May 16-22.....	14	1	
Do.....	July 18-Sept. 25.....	143	8	
Latvia.				Apr. 1-June 30, 1926: Cases, 5.
Mexico.				Feb. 1-June 30, 1926: Deaths, 1,525.
Aguascalientes.....	June 13-26.....	-----	5	
Ciudad Jaurez.....	Dec. 7-13.....	1	-----	
Guadalajara.....	June 8-14.....	-----	2	
Do.....	June 29-Sept. 27.....	-----	8	
Mexico City.....	May 16-June 5.....	3	-----	Including municipalities in Federal district.
Do.....	July 25-Dec. 4.....	9	-----	Do.
Saltillo.....	July 18-24.....	-----	1	
San Antonio de Arenales.....	Jan. 1-June 30.....	-----	-----	Present: 100 miles from Chihuahua.
San Luis Potosi.....	June 13-26.....	-----	7	
Do.....	July 4-Dec. 4.....	-----	30	
Torreón.....	May 1-June 30.....	-----	17	
Do.....	July 1-Nov. 27.....	-----	17	
Netherlands:				
Amsterdam.....	July 18-24.....	-----	9	
Nigeria.				Feb. 1-June 30, 1926: Cases, 521; deaths, 49.
Persia:				
Teheran.....	Apr. 21-Sept. 23.....	-----	18	
Peru:				
Arequipa.....	June 1-30.....	-----	1	
Do.....	Sept. 1-Oct. 31.....	-----	-----	Present.
Poland.				Mar. 28-May 1, 1926: Cases, 12; deaths, 1. June 27-Oct. 9, 1926: Cases, 417; deaths, 1.
Portugal:				
Lisbon.....	Apr. 26-June 19.....	10	3	
Do.....	July 11-Nov. 27.....	46	7	
Oporto.....	May 23-June 5.....	4	-----	
Do.....	July 11-Nov. 6.....	3	1	
Russia.				Jan. 1-Apr. 30, 1926: Cases, 2,529.
Siam.				Apr. 1-Oct. 30, 1926: Cases, 626; deaths, 251.
Bangkok.....	May 2-June 12.....	23	20	
Do.....	July 4-Oct. 30.....	87	68	
Spain.				Jan. 1-June 30, 1926: deaths, 99.
Valencia.....	Aug. 22-Oct. 23.....	3	-----	
Straits Settlements:				
Singapore.....	Apr. 25-May 1.....	1	-----	
Do.....	July 11-17.....	1	-----	
Sumatra:				
Medan.....	Aug. 22-28.....	-----	-----	1 case varioloid.
Switzerland:				
Lucerne Canton.....	June 1-30.....	1	-----	
Do.....	July 1-Sept. 30.....	3	-----	
Tripolitania.....	Apr. 1-June 30.....	12	-----	
Tunisia.				Apr. 1-June 30, 1926: Cases, 17.
Tunis.....	Sept. 11-30.....	2	-----	July 1-Sept. 30, 1926: Cases, 38.
Union of South Africa.				
Cape Province.....	June 1-30.....	8	1	
Do.....	June 20-26.....	-----	-----	Outbreaks.
Idutya district.....	Aug. 15-Oct. 30.....	-----	-----	Do.
Do.....	May 23-29.....	-----	-----	Do.
Natal.....	May 30-June 5.....	-----	-----	Do.
Durban.....	Oct. 10-Nov. 6.....	50	10	
Polela.....	Oct. 31-Nov. 6.....	-----	-----	Outbreak. In Nkandhla district.
Orange Free State.				Outbreak.
Transvaal.....	June 20-Aug. 28.....	-----	-----	June 6-12, 1926: Outbreaks in Pietersburg and Rustenburg districts.
Do.....	Aug. 29-Sept. 4.....	1	-----	Native.
Johannesburg.....	May 9-June 12.....	5	-----	
Do.....	July 11-Nov. 13.....	5	-----	
Praetoria.....	Sept. 19-25.....	1	-----	
Yugoslavia.				Apr. 15-30, 1926: Cases, 2; deaths, 1.
Zagreb.....	Aug. 9-15.....	2	-----	

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

SMALLPOX—Continued

Place	Date	Cases	Deaths	Remarks
On vessels: S. S. Karapara.....				At Zanzibar, June 7, 1926: 1 case of smallpox landed. At Durban, Union of South Africa, June 16, 1926: 1 suspect case landed.
Steamship.....	July 2.....	1		Vessel from Glasgow, Scotland, for Canada. Patient from Glasgow; removed at quarantine on outward voyage.

TYPHUS FEVER

Algeria.....				July 21-Sept. 20, 1926: Cases, 34; deaths, 1.
Algiers.....	May 21-June 30.....	7	1	
Do.....	July 21-Aug. 31.....	3		
Argentina: Rosario.....	Feb. 1, 28.....	2		
Bolivia: La Paz.....	June 1-30.....		1	
Do.....	Aug. 1-31.....	9	1	
Bulgaria.....				Mar. 1-June 30, 1926: Cases, 87; deaths, 14.
Chile: Antofagasta.....	May 23-June 26.....	4		
Do.....	June 27-July 3.....	1		
Concepcion.....	June 1-7.....		1	
Do.....	Oct. 1-31.....			Stated to be present in gaol.
Iquique.....	Aug. 8-Oct. 16.....	1	2	
Valparaiso.....	Apr. 29-May 5.....		1	
Do.....	Aug. 14-Nov. 6.....	11		
China: Antung.....	June 14-27.....	7	1	
Do.....	June 28-Oct. 31.....	45	1	
Canton.....	May 1-31.....	1		
Chungking.....	Aug. 29-Sept. 4.....			Present.
Ichang.....			1	Reported May 1, 1926. Occurring among troops.
Manchuria— Harbin.....	Oct. 14-20.....	1		
Wanshien.....				Present among troops May 1, 1926. Locality in Chungking consular district.
Chosen.....				Feb. 1-June 30, 1926: Cases, 1,005; deaths, 112. July 1-31, 1926: Cases, 37; deaths, 6.
Chemuipo.....	May 1-June 30.....	38		
Do.....	July 1-31.....	7	2	
Gensan.....	June 1-30.....	1		
Seoul.....	do.....	8	3	
Do.....	July 1-Oct. 31.....	9		
Czechoslovakia.....				Jan. 1-June 30, 1926: Cases, 156; deaths, 6.
Egypt: Alexandria.....	July 16-Aug. 19.....	3		
Do.....	Oct. 1-7.....	1	1	
Cairo.....	Jan. 29-May 13.....	89	27	
Do.....	July 23-Aug. 5.....	1		
Port Said.....	June 4-24.....	4	1	
Do.....	July 9-Oct. 7.....	5	1	
France.....	Aug. 1-31.....	5		
Great Britain: Scotland— Glasgow.....	July 30-Aug. 21.....	9	1	
Do.....	Reported Dec. 10.....	8		
Greece.....				Oct. 1-31, 1926: Cases, 7; deaths, 1.
Athens.....	Sept. 1-30.....		17	Including Piræus.
Hungary.....	May 1-June 30.....	3		
Baghdad.....	Oct. 10-16.....	1		
Ireland (Irish Free State): Cork.....	June 5.....	1		
Cork County.....	Oct. 17-23.....	1		
Ennistymon.....	July 4-10.....	5		
Kerry County— Dingle.....	June 27-July 3.....	1		

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

TYPHUS FEVER—Continued

Place	Date	Cases	Deaths	Remarks
Italy.....				Mar. 28-May 8, 1926: Cases, 3.
Palermo.....	Sept. 12-18.....	1		
Japan.....				Mar. 28-May 29, 1926: Cases, 37.
Latvia.....				May 1-June 30, 1926: Cases, 19.
				Aug. 1-31, 1926: Cases, 2.
Lithuania.....				Mar. 1-June 30, 1926: Cases, 199; deaths, 22. July 1-Aug. 31, 1926: Cases, 23.
				Feb. 1-June 30, 1926: Deaths, 189.
Mexico.....			1	
Durango.....	July 1-31.....			
Mexico City.....	May 16-June 5.....	20		Including municipalities in Federal District.
Do.....	June 13-19.....	9		Do.
Do.....	July 25-31.....	3		Do.
Do.....	Aug. 15-Dec. 4.....	99		Do.
San Luis Potosi.....	June 13-26.....			Present, city and country.
Morocco.....				Mar. 1-June 30, 1926: Cases, 426.
				July 1-Aug. 31, 1926: Cases, 20.
Norway:				
Stavanger.....	Sept. 6-12.....	1		
Palestine.....				Mar. 1-June 30, 1926: Cases, 14; deaths, 1. Aug. 1-Oct. 25, 1926: Cases, 22.
Birtuvia.....	Oct. 31-Nov. 6.....	1		
Gaza.....	July 6-12.....	1		
Haifa.....	July 13-Nov. 15.....	6		
Halalal.....	Aug. 17-23.....	1		
Jaffa district.....	June 15-28.....	5		
Do.....	Sept. 28-Nov. 8.....	4		
Jerusalem.....	Sept. 14-27.....	2		
Majdal district.....	July 13-Aug. 2.....	2		
Nazareth district.....	July 13-Nov. 8.....	7		
Petah Tokvah.....	Oct. 5-11.....	3		
Tiberias.....	Aug. 3-9.....	1		
Yavneil.....	Aug. 17-23.....	1		
Persia:				
Teheran.....	May 23-June 22.....		1	
Do.....	July 24-Sept. 23.....		5	
Peru:				
Arequipa.....	Jan. 1-31.....		2	
Lima.....	Aug. 1-31.....	1		
Poland.....				Mar. 28-June 26, 1926: Cases, 1,272; deaths, 85. June 27-Oct. 16, 1926: Cases, 346; deaths, 27.
Krakow.....	Oct. 17-23.....	31	5	
Tarnopol district.....	Oct. 10-16.....	1	1	
Rumania.....				Mar. 1-June 30, 1926: Cases, 899; deaths, 83. July 1-31, 1926: Cases, 65; deaths, 9.
				Jan. 1-Apr. 30, 1926: Cases, 18,647.
Russia.....				
Spain.....	Jan. 1-June 30.....		13	
Tunisia.....				Apr. 1-June 30, 1926: Cases, 110.
Tunis.....	June 11-30.....	3		July 1-Sept. 20, 1926: Cases, 101.
Turkey:				
Constantinople.....	June 16-22.....	1		
Union of South Africa.....				Apr. 1-May 31, 1926: Cases, 153; deaths, 19.
Do.....				July 1-31, 1926: Cases, 90; deaths, 17.
Cape Province.....				Apr. 1-June 30, 1926: Cases, 202; deaths, 24, native. July 1-Sept. 30, 1926: Cases, 82; deaths, 17.
				Outbreak in one locality.
Alexandria District.....	Oct. 31-Nov. 6.....			Outbreaks.
Clydesdale.....	Oct. 17-23.....			
Elliot District.....	Oct. 24-30.....	1		
Glengray district.....	June 27-July 3.....			Do.
Grahamstown.....	do.....	1		
Natal.....			1	
Durban.....	July 25-Sept. 18.....	11		Apr. 1-June 30, 1926: Cases, 28.
				July 1-31, 1926: Cases, 23; deaths, 2.
Orange Free State.....				Apr. 1-June 30, 1926: Cases, 24; deaths, 4. July 1-Sept. 30, 1926: Cases, 31.
				Outbreak on farm.
Brandford district.....	Oct. 10-16.....			

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

Reports Received from June 26 to December 31, 1926—Continued

TYPHUS FEVER—Continued

Place	Date	Cases	Deaths	Remarks
Union of South Africa—Con. Transvaal				Apr. 1-June 30, 1926: Cases, 10; deaths, 5. July 1-31, 1926: Cases, 2. Aug. 15-21, 1926, outbreaks.
Johannesburg	Aug. 29-Sept. 4	1		Outbreaks.
Walkkerstrom district	June 20-26			Do.
Wolmaransstad district	do.			Do.
Yugoslavia				Apr. 15-June 30, 1926: Cases, 48; deaths, 7. July 1-Oct. 31, 1926: Cases, 4; deaths, 1.
Zagreb	May 15-21	1		

YELLOW FEVER

Brazil	Reported June 26			Present in interior of Bahia, Pirapora, and Minas.
Bahia	May 9-June 26	10	7	
Do.	July 4-10	1		
French Sudan:				
Segou	Nov. 23	1	1	
Gold Coast	Apr. 1-June 30	8	4	
Nigeria	June 1-30	1	1	
Senegal	Nov. 1-23		9	