

CONTENTS

	Page
A study of endemic pellagra in some cotton-mill villages of South Carolina.	2645
Fumigation with cyanogen products—Report of experiments conducted with cyanogen products used in the fumigation of vessels for quarantine purposes at the New York quarantine station, Rosebank, Staten Island, N. Y.	2647
Court decision relating to public health.	2670
Public health engineering abstracts.	2671
Deaths during week ended September 29, 1928:	
Death claims reported by insurance companies.	2674
Deaths in certain large cities of the United States.	2674
PREVALENCE OF DISEASE	
United States:	
Current weekly State reports—	
Reports for weeks ended September 29, 1928, and October 1, 1927.	2676
Summary of monthly reports from States.	2678
Number of cases of certain communicable diseases reported for the month of July, 1928, by State health officers.	2679
Case rates per 1,000 population (annual basis) for the month of July, 1928.	2680
General current summary and weekly reports from cities.	2681
City reports for week ended September 22, 1928.	2682
Summary of weekly reports from cities, August 19 to September 22, 1928—Rates—Comparison with 1927.	2690
Foreign and insular:	
The Far East—Report for the week ended September 15, 1928.	2692
Argentina—Santiago del Estero—Further relative to plague.	2692
Canada—	
Provinces—Communicable diseases—Week ended September 22, 1928.	2692
Quebec Province—Communicable diseases—Week ended September 22, 1928.	2693
Cuba—Provinces—Communicable diseases—February 12—June 30, 1928.	2693
Czechoslovakia—Communicable diseases—July, 1928.	2693
Germany—Bavaria—Vital statistics—January 1—June 30, 1928.	2693
Greece—Dengue.	2694
Japan—Osaka—Cholera.	2694
Yugoslavia—Communicable diseases—August, 1928.	2694
Cholera, plague, smallpox, typhus fever, and yellow fever—	
Cholera.	2695
Plague.	2698
Plague rats on vessels.	2703
Smallpox.	2703
Typhus fever.	2710
Yellow fever.	2712

PUBLIC HEALTH REPORTS

VOL. 43

OCTOBER 12, 1928

NO. 41

A STUDY OF ENDEMIC PELLAGRA IN SOME COTTON-MILL VILLAGES OF SOUTH CAROLINA¹

An Abstract

By JOSEPH GOLDBERGER and G. A. WHEELER, *Surgeons*, EDGAR SYDENSTRICKER, *Statistician*, and WILFORD I. KING, *Special Consultant in Statistics*, with the cooperation of Wm. S. BEAN, JR., R. E. DYER, J. D. REICHARD, P. M. STEWART, *Surgeons*, M. C. EDMUNDS, *Assistant Surgeon*, R. E. TARBETT, *Sanitary Engineer*, DOROTHY WIEHL, *Assistant Statistician*, and JENNIE C. GODDARD, *Senior Statistical Clerk, United States Public Health Service*

As a part of the field investigations of pellagra conducted by the Public Health Service there was begun in the spring of 1916 a study of the relation of certain social, hygienic, sanitary, and economic factors to pellagra incidence in some representative South Carolina textile-mill communities, so-called cotton-mill villages, in which the disease was believed to be endemic. On a varying scale, but without interruption, this study was continued until the fall of 1921; that is, during a period of about five and a half years.

During 1916 this study was carried on in 7 villages. As it progressed it was more and more felt that the mass of data being collected would prove to be too small to afford entirely convincing indications with respect to certain important phases of the investigation. For this reason and because it seemed desirable to observe the possible fluctuations in the incidence of the disease from year to year and to study some of the factors possibly related to such fluctuations, it was arranged to continue the study, and for at least one year to carry it out on a much larger scale. Accordingly, early in January, 1917, a considerable number of additional villages were taken under observation, and by the end of February, 17 villages in addition to the 7 of 1916 were settled upon for study. These 24 villages were kept under surveillance for pellagra throughout the year 1917.

With the beginning of 1918 the scale of the investigation was reduced to about that of 1916, surveillance of 18 of the 24 villages studied during 1917 being discontinued. Of the 6 continued under observation during 1918, 2 had been among the 7 studied in 1916. At the beginning of 1919 the scale of the investigation was further reduced by discontinuing observation of all but 1 of the villages.

¹ The complete report will appear in *Hygienic Laboratory Bull.* No. 153.

This 1 village (*In.*) was 1 of the original 7 and was continued under surveillance throughout 1919, 1920, and up to October 15, 1921—or, in all, for about five and a half years.

The results of the first year's study have already been published.¹ In the present communication, much delayed by, among other reasons, the pressure of other continuing studies, we desire to record certain of the results of that phase of the subsequent study concerned with the incidence of the disease and the relation of this incidence to certain social, climatic, sanitary, economic, and dietary factors.

During 1917 in an aggregate population of 22,653 individuals, 1,147 cases of pellagra (an incidence rate of 50.6 per 1,000) were observed. Of the 4,104 households among which that population was distributed, 18.5 per cent had at least one member affected by the disease in that year.

Pellagra (in an endemic locality) is very much (two to six times) more prevalent than the experience of the physicians of the locality would seem to indicate.

The fatality rate of the endemic disease, when definitely marked cases of all grades of severity are considered, would appear not to exceed 3 per cent.

Striking peculiarities of age and sex distribution of the disease were observed.

The observations of age incidence appear to indicate, what seems not to have been recognized heretofore, that endemic pellagra is preponderatingly a disease of children of from 2 to 15 years of age.

Explanations of the peculiarities of age and of sex incidence are suggested.

The single woman, as compared with the married, widowed, or divorced, is relatively exempt from the disease. In the population group under consideration, the single woman is usually a wage earner, which may place her in a somewhat more advantageous position with respect to diet than her married or widowed sister.

The incidence of the disease was found to be markedly seasonal; 80 to 90 per cent of all cases had their "onset" within the period April to July, inclusive. One explanation suggested, in view of the proved dietary relation of the disease, is the variation in diet brought about by the seasonal modification of the food supply.

The seasonal incidence of cases distinguished by their occurrence singly or otherwise in a household, and as initial and recurrent attacks, was studied.

The disease was found to have a marked and very sharply limited season of prevalence the curve of which, with a slight lag, paralleled that of incidence.

¹Public Health Reports, Mar. 19, 1920 (Reprint No. 587), July 9, 1920 (Reprint No. 601), July 16, 1920 (Reprint No. 603), Nov. 12, 1920 (Reprint No. 621).

The study failed to disclose any consistent correlation between sanitary conditions and pellagra incidence. Such association as may at times be observed is regarded as accidental and to be explained by the intimate relation of the endemic disease to economic status, of which the sanitary condition may be an index.

The study reveals the existence of a striking inverse correlation between the incidence of the endemic disease and family income.

The continuous study of a selected village during a period of nearly six years appears to demonstrate that income shortage was a fundamental, though indirect, controlling factor in relation to the year-to-year fluctuation in the incidence of the disease. It is therefore inferred that the year-to-year fluctuations in the incidence of the endemic disease are bound up with fluctuations in economic conditions that influence the ability of a certain section of the population to procure an adequate diet.

Marked seasonal variations in the food supply of a selected village are demonstrated. A relation of this variation in food supply to the striking seasonal incidence and prevalence of the disease is suggested

FUMIGATION WITH CYANOGEN PRODUCTS

Report of Experiments Conducted with Cyanogen Products Used in the Fumigation of Vessels for Quarantine Purposes at the New York Quarantine Station, Rosebank, Staten Island, N. Y.

By C. V. AKIN, *Surgeon*, and G. C. SHERRARD, *Acting Assistant Surgeon, United States Public Health Service*

During the period February 15 to May 29, 1926, an extended series of experiments was conducted at the New York quarantine station to determine the relative efficiency of certain cyanogen products used in the fumigation of vessels for the destruction of rats. Tests of all products under consideration were made, both under control in the laboratory and under practical conditions on board ship.

For the conduct of this work an informal board of officers on duty at the station was formed, consisting of Surg. C. V. Akin, Acting Assist. Surgs. G. C. Sherrard and G. H. Guth, and Chief Pharmacist B. E. Holsendorf. All of the experimental work reported herein was done by Surg. C. V. Akin and Acting Assist. Surg. G. C. Sherrard.

The general purposes of the tests were to determine with reasonable exactness the relative merits of several cyanogen products used in ship fumigation for the destruction of rodents, from the standpoints of (1) lethal efficiency, (2) safety to fumigators and others, and (3) cost.

PRODUCTS TESTED

The following products were tested:

(1) Liquid hydrocyanic acid (96 to 98 per cent):

(a) A mixture of 80 per cent liquid HCN and 20 per cent of cyanogen chloride (CNCl), an irritating warning-giving component.

(b) Liquid hydrocyanic acid (96 to 98 per cent).

(2) Hydrocyanic acid gas generated from a mixture of sodium cyanide, mineral acid, and water.

(a) Sodium cyanide, sodium chlorate, hydrochloric acid, and water in proportions designated by the quarantine regulations of the Public Health Service.

(b) Sodium cyanide, sulphuric acid, and water in proportions designated by the quarantine regulations of the Public Health Service.

NOTE.—Formula (a) in (2) above gives a mixture of HCN and tear gas, CNCl; while formula (b) gives HCN only.

(3) Zyklon-B, a product of German manufacture, containing liquid hydrocyanic acid and, in various lots, 10, 6, and 4 per cent of chloropicrin, an irritating warning-giving gas.

(4) Calcium cyanide, a cyanogen product of American manufacture, in the form of a fine dust, one-half of the total volume by weight of which is HCN.

DESCRIPTION OF PRODUCTS USED

Liquid hydrocyanic acid.—Commercial liquid hydrocyanic acid averaging from 96 to 98 per cent HCN, with from 2 to 4 per cent of water, slightly aciduated with sulphuric acid.

Liquid hydrocyanic acid-cyanogen chloride mixture.—Commercial hydrocyanic acid to which has been added 20 per cent of liquid cyanogen chloride for lachrymatory effect.

Sodium cyanide (NaCN, 96 to 98 per cent).—Egg sodium cyanide containing approximately 52 per cent cyanogen and showing less than 2 per cent chlorides.

Sulphuric acid.—Commercial sulphuric acid 92 to 94 per cent pure (66° Baumé), free from nitric acid and metals.

Hydrochloric acid.—Commercial hydrochloric acid, 20° Baumé.

Sodium chlorate.—Sodium chlorate crystals.

Zyklon-B.—Zyklon-B is the trade name given to a chemical preparation of German manufacture composed of liquid hydrocyanic acid absorbed in a porous granulated earthy substance, named diatomite.

The mixture consists of equal, or nearly equal, parts of liquid hydrocyanic acid and diatomite (by weight), plus small quantities

(about 5 per cent) of one of two irritating gases which have a markedly lachrymatory effect, and which serve the double function of warning exposed persons and of stabilizing the HCN in the product. Additional stabilization is secured through the diatomite and a small amount of sulphuric acid.

Zyklon-B is marketed in heavy tin cans which withstand a pressure test of five atmospheres. The cans are filled with a guaranteed HCN content of 20 grams, 100 grams, 500 grams, 1,000 grams, or 1,200 grams, which makes the "dosing" of a compartment of any size an easy matter.

When the can is first opened, Zyklon-B has the appearance of dried or only slightly moist particles of sandy clay, varying from a pale reddish yellow to an orange-yellow color. The amount of appreciable moisture varies from that in the small can, the content of which is thoroughly dry, to that in the larger cans, containing from 500 to 1,200 grams HCN, wherein the material is of the consistency of moist sawdust.

In all sizes of cans the material runs freely from a comparatively small opening (1 inch to 1½ inches), as the contents of the cans are being emptied into the holds of a vessel or spread on the floor of a smaller compartment.

The large surface afforded by the carrying material, diatomite, promotes rapid evolution of the HCN gas even when the product is exposed in relatively thick layers up to three-eighths of an inch in thickness. Under all ordinary circumstances the HCN content is quickly given off, and by the end of a two-hour fumigation period the residue is practically inert. (See Public Health Reports, vol. 42, No. 50 (Dec. 16, 1927), p. 3071.) While there is no tendency for the residue to retain or take up HCN in gaseous form, it is well to remove the residue after fumigation, especially in the quarters or superstructure of a vessel, as a marked odor and some tear effect are noted for a considerable time if the residue be not removed.

The opening of the cans is easily accomplished with a special hammer having a tempered cutting head which cuts a 1½-inch opening through the top of the can with one stroke. Cone-headed peen hammers or sharpened chipping hammers are used for this purpose. Two or three such openings permit the discharge of the contents of the largest cans in 15 to 20 seconds. A small air hole should be punched in the opposite end of container to facilitate emptying its contents.

In dosing holds the contents of the requisite cans are poured from deck down through the hatches and spread over the floor of the hold with a sowing motion. In the holds the residue may be swept up and thrown overboard if desired, or it may be allowed to remain without danger. The dose for superstructure compartments is

thinly and evenly spread on protecting sheets of paper previously laid on the floor. The subsequent removal of this residue is then easy to accomplish.

Calcium cyanide.—Calcium cyanide is a cyanogen carrier of the formula $\text{Ca}(\text{CN})_2 \cdot 2\text{HCN}$, formed by the reaction of hydrocyanic acid containing a slight amount of water on calcium carbide, with the formation of calcium cyanide and liberating acetylene.

According to the manufacturers, the product contains only such impurities as are common to calcium carbide, together with the small amount of polymer which is responsible for the light tan color of the compound. It is in a very fine state of subdivision, passing freely through a 300-mesh screen.

The product is extremely sensitive to moisture, being decomposed, with the liberation of hydrocyanic-acid gas, as follows:



The usual moisture in the air quickly sets up the above reaction upon exposure, and a satisfactory liberation of HCN gas takes place when the humidity is 25 per cent or even less.

The material used for these tests was packed for shipping in 1-gallon tin buckets with friction tops. As a precaution against accidental opening, the tops were spotted in place with solder.

Each bucket contained 4 pounds of a light tan powder of the approximate consistency of the finest wheat flour. This powder is so dry and finely divided that a dust cloud is formed by the slightest agitation or draft. The odor of hydrocyanic acid gas can be noted the instant the powder is exposed to the air.

There is no appreciable change in the color or consistency of the powder after prolonged exposure, nor can the amount of HCN evolved be estimated by change in weight.

The manufacturers state that the cyanogen content ranges between 50 and 55 per cent, averaging about 53 per cent. In computing the test "doses" it was assumed that one-half of the calcium cyanide, by weight, was available HCN.

"Calcium cyanide" may be applied either in the form of a dust, through being blown into the compartment to be fumigated, or by being laid down in very thin layers. The most efficient action of the product is promoted by "dusting," as a more general distribution is secured and the HCN gas more promptly liberated. Satisfactory gas evolution is obtained, however, in layers up to one-sixteenth inch in thickness, as slightly over 95 per cent of the HCN content is given off in two hours when so distributed.

Owing to the fact that, roughly, 5 per cent of the HCN content remains in the residue at the end of a two to four hour fumigation

period, and, further, because the reaction is reversible and HCN is taken up by the residue, it is essential that as much as possible of the residue be removed at the end of fumigation. When the powder is dusted into a compartment its subsequent removal is a practical impossibility, and this method of "dosing" superstructure compartments is further contraindicated, for the dust which settles on and clings to everything exposed to it is disfiguring. This criticism would not, of course, obtain when the powder is used in the hold of a vessel. When the material is distributed in thin layers on sheets of paper, the removal of the residue is easily accomplished.

LOCATION OF EXPERIMENTAL ROOMS

Satisfactory experimental rooms were available in a vacant building on Hoffman Island, a part of the New York quarantine station. Several rooms, averaging between 1,180 and 1,185 cubic feet capacity, were selected and prepared for the tests by carefully sealing all cracks and openings, no matter how small. As the walls, ceilings, and floors of these rooms were covered with cement mortar and painted over, there was little opportunity for leakage.

In order to test the diffusion of gases, two adjoining rooms were connected by introducing a number of 2-inch metal pipes through the partition wall. These pipes could be plugged gas-tight or opened as desired, and the number and location of the pipes afforded a variety of combinations to test gas circulation.

TEST ANIMALS

A large number of white rats which had been bred on Hoffman Island were available as test animals. Adult white rats were used in all tests both on the island and on shipboard.

No direct evidence was obtained as to the relative resistance or susceptibility of these animals to HCN gas as compared with wild rats; but a considerable variation in resistance between these white rats was noted when several animals were simultaneously exposed to the same concentration of gas. For this reason two or more white rats were used in all of the more delicate tests, such as when the effects of greatly reduced doses were being studied.

PROGRAM OF PROPOSED EXPERIMENTAL WORK

- (1) Diffusion of gas from one compartment to another of equal size through relatively small orifices located at various levels.
- (2) Retention of HCN gas in residues of certain fumigants.
- (3) Reabsorption of HCN gas by residues of certain fumigants.
- (4) Fumigating with reduced dosages of cyanogen products to determine the minimum lethal rat dose and to compare the lethal efficiency of the various preparations tested.

(5) Absorption and holding of HCN gas by absorptive materials exposed to fumigation.

(6) Penetration by gaseous fumigants of porous materials used to protect test animals.

(7) Miscellaneous tests of cyanogen products to secure information as to the properties, behavior, etc., of HCN gas.

(8) Fumigation of ships without cargo with various cyanogen products controlled with test animals.

(1) DIFFUSION OF GAS WITHIN A SINGLE COMPARTMENT AND FROM ONE COMPARTMENT TO ANOTHER OF EQUAL SIZE THROUGH RELATIVELY SMALL ORIFICES

Adjoining rooms of 1,181 and 1,185 cubic feet air capacity, respectively, were made to communicate through three short sections of 2-inch iron pipe which perforated the intervening partition wall at equal intervals along the mid-perpendicular line from ceiling to floor.

An attempt was made to have these rooms thoroughly gas-tight by plastering and pasting all openings in the walls, ceilings, and floors, and by papering over windows and doors at the time of fumigation. We proposed to determine the regular or average rate of diffusion from one compartment to another through such orifices as above described and to observe to what extent the passage of gas was affected by changes in temperature, artificially operated air currents, etc.

Room temperatures could not be made to fluctuate during the actual fumigation, but by utilizing steam heat and coal-oil stoves a considerable difference in the temperature of the adjoining test rooms was secured and maintained. Briefly, four temperature combinations were tried: (1) Both rooms equally chilled (50° F. or lower); (2) both rooms equally warmed (70° F. or higher); (3) room A (in which total amount of gas for both rooms was introduced) chilled while room B (containing test animals) was kept at least 20° F. warmer; and (4) room A (gas room) warmed while room B (test animals) was kept at least 20° F. colder.

Artificially induced air currents were supplied by a 10-inch electric fan operated for one series of tests in room A (gas room) and for another series of tests in room B (test animals).

For all tests the standard dosage of 2 ounces HCN per 1,000 cubic feet of space was used. In testing diffusion, sufficient gas was introduced or generated in room A to furnish an average of 2 ounces HCN per 1,000 cubic feet for both rooms A and B.

In all of these tests four test animals were placed in open-mesh wire cages in room B, two at the ceiling and two at the floor level, in opposite corners of the room, so as to afford the maximum distance

from the gas intakes represented by the pipe orifices in the partition wall.

Twenty separate experiments were conducted, in which the several cyanogen products under consideration were used. Liquid HCN (80 per cent) CNCl (20 per cent) mixture was used as the standard of comparison.

Results.—(1) With exposures of from four to seven hours not enough gas passed from room A to room B through one, two, or three 2-inch pipes to affect the test animals when both rooms were otherwise tightly sealed.

(2) When small openings were left around the window frames and the wind blew directly against the windows, a lethal quantity of gas passed from room A to room B within from one to four hours.

(3) No variation in the temperature secured between the two rooms modified the passage of gas from room A to room B through the 2-inch pipes when both rooms were otherwise tightly sealed.

(4) No observable effect was produced by the air currents set up by a 10-inch electric fan running at full speed alternately in room A and room B.

A visual check on the experiences noted in Nos. 1 to 4 above was secured by burning double the standard amount of sulphur in room A and watching for the passage of the smoke through the pipes leading into room B. The first smoke seen came through the pipe nearest the ceiling 18 minutes after the sulphur had been ignited. No smoke was seen to pass through the middle and lower pipes. At no time during the 5-hour experiment was there more than a very faint cloud in room B, and at the end of the time none of the test animals in room B showed any effect.

(5) Diffusion of gas *within* the room into which it was introduced or generated indicated clearly that when no gross air currents were present, HCN gas, whether alone or mixed with "tear gas," showed a constant tendency to rise. Test animals exposed at the ceilings were invariably killed before the animals placed directly beneath them on the floor. This observation holds good only when the rats on the floor were at least as far from the center of gas generation as those at the ceiling level. In our experiments, liquid hydrocyanic acid (alone and mixed) was sprayed into the test rooms through a prepared vent in the door which made the gas distribution more or less central, while Zyklon-B was placed on the floor, as were the buckets for generating HCN from sodium cyanide and acid.

Conclusions.—Hydrocyanic acid gas shows little tendency to flow or diffuse from one compartment to another through small apertures when both compartments are otherwise tightly sealed. The importance of this conclusion is apparent when applied to conditions ordinarily existing in holds and other parts of ships.

It is a well-known fact that rats escape cyanide fumigation as fumigation is usually done, and the reason for this becomes clear when it is seen that a lethal quantity of the gas will not flow through small openings into tightly closed sections not primarily exposed to the gas. The practical proof of this was witnessed on shipboard when test animals, concealed in the covered bilges, closed drawers, and similarly tight compartments, were unaffected by standard amounts of cyanide during the course of a two-hour fumigation.

The movement of HCN gas from one room into the other through small openings as a result of the action of extraneous air currents was clearly demonstrated in those tests in which the communicating compartments were not sealed equally tight and were subject to the influence of natural ventilation. As the movement from room A (gas room) to room B (test-animal room) occurred only when the wind came directly at small outside openings and did not occur when the air in the gas chamber was agitated by a fan inclosed in the tightly sealed chamber, it appears that the diffusion or flow of gas through small orifices (2-inch pipes) resulted from the movement of the total volume of gas-air mixture in room A and the air in room B rather than from localized movements in either or both rooms.

Observations on the behavior of gas within tightly sealed rooms indicate clearly that the natural flow or diffusion of HCN and HCN mixtures is primarily upward, which emphasizes the importance of locating the center of gas generation at the lower rather than the upper part of large compartments. Even if this were not invariably true, it is obvious that the highest gas concentration in ship deratization should be contrived and maintained at or near those points where rats are most numerous and where rats have the greatest opportunity for escape; i. e., the lower parts of the ship. This detail will be further developed in the section relating to the generation method.

The conclusions regarding hydrocyanic gas diffusion in usual fumigation procedures may be briefly summed up by stating that no ordinary concentration and no ordinary exposure time will insure the infiltration of gas into so-called dead spaces commonly communicating with compartments under gas through such small openings as are customarily used by rats. The opening up or complete elimination of such small contiguous spaces and the competent blocking of such escape openings must therefore be considered as of equal, if not greater, importance than the gas dosage, the exposure time, or the kind of fumigant used. A vessel not thus properly prepared for fumigation will more or less nullify the potential good effects of the most careful gassing.

(2) RETENTION OF HYDROCYANIC ACID GAS IN RESIDUES OF CERTAIN FUMIGANTS

Residues remaining after fumigation by the generation method and following the use of Zyklon-B and calcium cyanide were considered from the standpoint of residual or unexpended cyanogen which might render dangerous subsequent human habitation of the fumigated compartment.

(a) *Generation method.*—All residues of sodium cyanide and acid mixtures are dangerous, as they contain varying amounts of HCN. The mere removal of containers used in the generation of HCN and the proper disposal of contents effects the necessary safeguard. The residual or unexpended HCN in such residues often comes off freely when the residue is agitated, as during careless removal, but otherwise is given off slowly.

(b) *Zyklon-B.*—A negligible quantity of HCN remains in the diatomite residue at the end of two to four hour exposure, even when the material is spread in very thin layers. By concentrating relatively large amounts of residue into small, tight containers, test animals were rarely killed by exposures varying from 4 to 12 hours. (See PUBLIC HEALTH REPORTS, vol. 42, No. 50 (December 16, 1927), p. 3071.)

(c) *Calcium cyanide.*—Tables prepared by the manufacturers of calcium cyanide show that the major portion of the HCN content of this product is given off rapidly, but that the rate of evolution is modified by the thickness of the layer in which the material is distributed. Under optimum conditions, with layers only one-sixteenth inch in thickness, 79.2 per cent of HCN is given off in 1 hour, 95.7 per cent in 2 hours, 95.9 per cent in 4 hours, 96.9 per cent in 8 hours, and 97.5 per cent in 24 hours. It will be seen that 3 or 4 per cent of available HCN (one-half of material by weight) remains after a period much greater than any afforded under practical conditions in ship fumigation. These figures serve to confirm preliminary experimental findings which showed the lethal effects of this residue before the exact figures were available, in which relatively large amounts of residue exhausted for 35 hours and concentrated in small, tight containers, killed test animals in from 30 minutes to 1 hour.

(3) REABSORPTION OF HYDROCYANIC-ACID GAS BY RESIDUES

Residues of Zyklon-B and of calcium cyanide were tested to determine whether or not they were inert in the presence of hydrocyanic acid gas. Quantities of these products sufficient to fumigate rooms of 1,000 cubic feet capacity were exposed until all HCN had been exhausted. The residues were then exposed to standard doses

of HCN (gas from the liquid HCN-CNCl mixture) for two hours. The residues were then separately concentrated into small, tight containers and test animals directly exposed. The animals exposed to the Zyklon-B residue survived all exposures, while the calcium-cyanide residue killed test animals in 30 minutes.

This result indicates that the reaction of calcium cyanide in giving off HCN in the presence of atmospheric moisture is reversible to the extent that the residue will take up HCN when the atmospheric content of HCN exceeds that of the residue.

Conclusions.—The results outlined in the last two sections above indicate clearly that, while the residues from both Zyklon-B and especially from calcium cyanide should be removed carefully following fumigation, the residue of Zyklon-B is apparently much less dangerous than that of calcium cyanide.

Of additional importance is the time saved in "clearing" the vessel of gas when the residue is removed soon after opening up following fumigation. There is an apparent persistence of tear effect in Zyklon-B residue which can be accounted for only by the fact that the irritating gases are more slowly released from the diatomite than are the hydrocyanic acid component.

(4) TESTING THE LETHAL EFFECT OF REDUCED DOSES OF CYANOGEN PRODUCTS AS THE BEST MEANS OF COMPARING EFFICIENCY

The most interesting and perhaps the most fruitful tests performed in our series of tests with cyanide products were those undertaken with reduced doses or fractional parts of the standard dose of 2 ounces per 1,000 cubic feet.

It is obvious that when a concentration of 2 ounces of HCN per 1,000 cubic feet is provided, a substantial overdose is insured. The question is not whether 2 ounces of HCN per 1,000 cubic feet will kill, but whether with the materials in the proportions used, the desired dose of cyanogen is made available in the compartment under fumigation.

In all of our tests, liquid hydrocyanic acid (96 to 98 per cent) was used as the standard of comparison. This eliminated the constant variability of the generation method and the uncertainty associated with the use of two comparatively new fumigants—Zyklon-B and calcium cyanide. Once the average working dose, the minimum lethal dose, and the threshold dose of liquid HCN were determined with reasonable certainty, the other cyanogen products were measured by this standard.

It was at once apparent that, if usual doses of the products to be tested were used, no real comparison of their performance would be afforded. Under all ordinary circumstances all test animals would be killed, the only variation being in the *time* required to kill. In

view of the known difference in the resistance of rats to HCN gas, *the time required to kill* is of uncertain value unless several other factors are considered.

Before testing the action of reduced doses of cyanide, all test rooms were carefully sealed. Accepting 2 ounces of HCN per 1,000 cubic feet as the standard, fractional parts of this dose were used. Test animals were carefully selected as to size, and two animals were used in each test, one being placed at the ceiling and one at the floor level. It is interesting to note that, in these experiments, the animals at the higher level were always affected before those on the floor, which further indicates the natural tendency for HCN gas to rise.

Fifty-five separate fumigations were done, using the liquid HCN-CNCl mixture, liquid HCN alone, calcium cyanide, HCN-CNCl mixture generated, and HCN generated. As the smallest amount of Zyklon B available represented 20 grams of HCN, less than one-third of the standard dose per 1,000 cubic feet, it could not be used in certain tests. It is significant to note, however, that in all concentrations from double the standard dose down to a one-third dose, Zyklon-B gave results exactly comparable with liquid hydrocyanic acid (96 to 98 per cent).

Allowing for the variation in resistance in test animals, the average killing time for fractional doses when using liquid hydrocyanic acid, mixed and alone, may be accepted as follows:

- One-eighth dose, or one-fourth ounce per 1,000 cubic feet, 15 to 20 minutes.
- One-tenth dose, or one-fifth ounce per 1,000 cubic feet, 20 to 25 minutes.
- One-twelfth dose, or one-sixth ounce per 1,000 cubic feet, 30 to 45 minutes.
- One-sixteenth dose, or one-eighth ounce per 1,000 cubic feet, 60 to 180 minutes.
- One-twentieth dose, or one-tenth ounce per 1,000 cubic feet, overnight.
- Animals withstood smaller doses for as long as 36 hours without ill effect.

In doses between one-twelfth and one-twentieth of the standard 2 ounces per 1,000 cubic feet, liquid hydrocyanic acid is slightly more lethal than the liquid HCN-CNCl mixture. For all practical purposes, however, there is no choice between the two preparations.

From the above experience we feel justified in concluding that much less than the present standard dose of liquid hydrocyanic acid will serve to kill rats directly exposed to its fumes, and that a concentration as low as one-tenth ounce of HCN per 1,000 cubic feet must be considered as dangerous to human beings exposed over a long period of time. It is logical, therefore, to assume that a reduction in the amount of gas customarily used to fumigate living and sleeping quarters (superstructure) would effectually advance the safety of persons subsequently occupying them without materially interfering with a satisfactory deratization. Our experiences indicate clearly that rats which are well enough protected in living quarters

to escape a dose of 1 ounce of HCN per 1,000 cubic feet will also survive a 2-hour exposure to the standard dose.

In the light of our findings with reduced doses it is obvious that the same results as those obtained with liquid hydrocyanic acid may not be expected from the generation method. This is particularly true as regards the HCN-CNCl mixture.

Making due allowance for the loss of time in generation, one is forced to the conclusion that much less than the theoretically obtainable amount of HCN is actually delivered in the course of the average fumigation. Our tests indicate that the generation method for the production of HCN is much more effective than the same procedure for the generation of the HCN-CNCl mixture. This is to be expected in view of the fact that 1 ounce, or 20 per cent less, of sodium cyanide per 1,000 cubic feet, is used in the production of the mixed gas, and in addition some of the CN liberated from the sodium cyanide is utilized in the formation of the irritating gas, CNCl. The more rapid evolution of HCN from sodium cyanide and sulphuric acid also plays a part, as maximum gas concentration is more rapidly reached. This is of importance when it is appreciated that a very brief exposure to a high concentration is more uniformly fatal than prolonged exposure to doses approaching the threshold concentration. It is desired at this point to emphasize the extreme importance of the length of exposure under circumstances where diffusion or circulation of gas is rendered difficult. This consideration is separate and distinct from the proposition of reduced dosage which, for the purposes of this discussion, is applicable only to the results to be expected when animals are exposed immediately and directly to the gas.

Based on the results of tests of reduced doses of HCN, the conclusion is reached that the relative lethal efficiency of the several products and methods under consideration warrants their listing in the following order:

- (i) Liquid hydrocyanic acid (96-98 per cent).
- (ii) Liquid hydrocyanic acid-cyanogen chloride mixture.
- (iii) Zyklon-B (10 per cent irritating gases). Equal to the liquid HCN-CNCl mixture and compares favorably with straight liquid HCN.
- (iv) Hydrocyanic acid gas generated by mixing sodium cyanide with sulphuric acid and water.
- (v) Hydrocyanic acid-cyanogen chloride mixture generated by mixing sodium cyanide, sodium chlorate, hydrochloric acid, and water.
- (vi) Calcium cyanide (50-55 per cent HCN).

It must be understood that the above arrangement is arrived at by considering lethal efficiency alone. In the proper place a final comparison will be made, in which other factors affecting the value of a fumigant will be considered.

(5) **ABSORPTION AND HOLDING OF HYDROCYANIC ACID GAS BY ABSORPTIVE MATERIALS EXPOSED TO FUMIGATION**

The absorption and subsequent holding of hydrocyanic acid (gas) by permeable materials, such as bedding, clothing, floor coverings, sacking, and baled goods exposed to fumigation, is of more than ordinary interest on two accounts: First, the presence of considerable amounts of such material lengthens the time required to clear a given compartment of gas and, consequently, increases the time required for completion of fumigation; second, the retention of minute quantities of HCN in such materials as are always found in sleeping quarters represents a distinct hazard for persons who, without assuming all necessary precautions, subsequently occupy the quarters.

Conclusions.—The conclusions drawn from our experimental work along this line are as follows:

(i) Owing to the ready solubility of HCN in water, moist materials take up more HCN than dry materials. Materials dry to the touch, however, will take up a lethal quantity of HCN, and for purposes of safety no distinction should be drawn on the basis of supposed moisture content.

(ii) Hydrocyanic acid taken up by moist or wet materials will be held longer and given off more slowly than is the case with dry materials.

(iii) The *quantity* of absorptive materials exposed is apparently of greater importance than the concentration of gas used.

(iv) When the hydrocyanic acid-cyanogen chloride mixtures are used, the presence of moist materials or of actual collections of water in the fumigated compartment or in the bilges of a vessel, gives rise to a persistence of HCN after all "tear effect" has disappeared. This is more nearly constant for liquid HCN-CNCl mixtures and for sodium cyanide-sodium chlorate-acid mixtures than Zyklon-B (10 per cent irritator content), as in the instance of the latter the character of the irritating gases makes for persistence of the "tear effect." (See PUBLIC HEALTH REPORTS, vol. 38, No. 27 (July 6, 1923), p. 1532.)

(v) When using liquid hydrocyanic acid it is of the greatest importance to avoid spraying or pouring the acid directly on bedding, permeable floor coverings, or clothing. The danger is greatly increased if the materials are moist.

(vi) Proper aeration and drying of absorptive materials exposed to cyanide fumigation in sleeping quarters is of vital importance. In two instances this can be accomplished by exposure of from one to two hours in the open air. The process is expedited by beating and shaking the materials, or exposure to the warmth of the sunlight.

(vii) In view of the customary carelessness and disregard for these precautions by crews, and on account of the inclement weather which

frequently occurs during and after fumigation, it would obviously be much safer if all bedding, floor mats, etc., were removed from sleeping compartments prior to fumigation. This is particularly true of crew's quarters, which, owing to location, are frequently poorly ventilated and damp.

(viii) The reduction of dosage for fumigation of sleeping quarters is worthy of serious consideration, as rats are killed with much less than the standard dose (2 ounces HCN per 1,000 cubic feet) of gas, and the hazard to human life diminishes with the amount of hydrocyanic acid introduced.

(6) PENETRATION OF PERMEABLE MATERIALS SERVING TO PROTECT RATS

It is our belief, based on numerous experiments, that rats which escape fumigation do so either because at the time gas is introduced they are safely ensconced in the gas-free atmosphere of a "dead" space, or, through minor structural defects, they get away from the gas into otherwise well-closed spaces not directly affected by fumigation. It is in such spaces that ship's rats naturally harbor, and it is to such places that they instinctively turn when menaced by the introduction of gas or disturbed by the preparations incident to the proposed fumigation.

To a much lesser extent do rats find protection in cargo and the dunnage customarily found in ship's holds and compartments. Both the quantity and kind of cargo must be considered, however, and the quantity and arrangement of dunnage require attention if the vessel is to be properly prepared. The mere presence or absence of cargo and dunnage does not, in the final analysis, determine the efficacy of fumigation, but rather the quantity of material and its disposition, as hydrocyanic acid gas will penetrate either bagged or loosely boxed parcels if the gas is permitted to *surround* the container.

In testing penetration various materials with various sized perforation were used to "protect" the test animals. Included in the list are wooden boxes made gas-tight except for a predetermined number of quarter-inch holes, gas-tight containers in which the animals were protected by layers of gunny sacking varying in number from 10 to 80, blankets, rolls of matting, mattresses, and heavy paper sacks. Not only did these devices serve to check the penetrating power of measured concentrations of HCN, but they permitted a rigorous comparison of the fumigations afforded by the several cyanogen preparations under consideration.

Details of penetration tests.—(i) Boxes: A number of tightly jointed wooden boxes, with gas-tight doors for admitting test animals, were used. A series of such boxes was prepared by boring from

none to four quarter-inch holes through one end. Adult white rats were placed in the boxes and exposed to fumigation with 2 ounces of HCN per 1,000 cubic feet. A 2-hour fumigation with liquid HCN-CNCl mixture, Zyklon-B, generation method, and calcium cyanide invariably killed all rats in boxes with two or more holes, whereas all rats in boxes with no holes were spared.

(ii) Gunny sacking: Test animals in open-mesh wire cages were placed in gas-tight buckets of about one-half cubic foot capacity. The tops of these buckets were covered with pads of new gunny sacking varying from 10 layers to 80 layers in thickness. These pads were so affixed that it was necessary for the gas to go through the sacking to get to the animal in the bucket. In one series of tests the pads of sacking were used *dry* and in another *wet*.

When dry sacking was used, test animals were invariably *killed* through 70 layers. Eighty layers always protected.

When *wet* sacking was used, test animals were invariably *protected* by 40 layers. Wet sacks were prepared by saturating them in water, wringing them as dry as possible, and then hanging them in the air for one hour, by which time their moisture content seemed uniform.

(iii) Paper sacks: 8, 10, and 12 pound sacks of kraft paper were used. By slipping one sack inside another, from one to four layers of sacking were secured, and after the rats had been introduced into the bags, the open ends were tightly pasted up.

When using 2 ounces of liquid HCN-CNCl mixture per 1,000 cubic feet, test animals were killed by a 2-hour exposure when sealed up in four sacks.

Rats protected by from 16 to 20 layers of blankets, and others rolled in matting or hidden in piles of loose sacking were invariably killed by the standard dose and exposure.

The use of artificial protection afforded a definite comparison of the lethal efficiency of the cyanogen products tested.

The liquid hydrocyanic acid-cyanogen chloride mixture (80-20 per cent) gave a slightly higher percentage of kills than Zyklon-B, but both are more lethal than calcium cyanide and the sodium cyanide-sodium chlorate-mineral acid mixture. It is apparent that the superiority of the liquid gas and Zyklon-B rests on the higher proportion of HCN gas evolved within the permissible fumigation time, and the additional fact that a higher gas concentration is reached more promptly. With accurately proportioned doses, calcium cyanide will furnish as much HCN per 1,000 cubic feet, but the maximum gas concentration is reached more slowly. The generation method is not only the slowest of the four, but actually much less HCN in gas form is produced, much HCN remaining in solution or unexpended.

(7) MISCELLANEOUS TESTS TO SECURE ADDITIONAL INFORMATION OF PROPERTIES AND BEHAVIOR OF CYANOGEN FUMIGANTS

ZYKLON-B

(A) *Gas leakage from small openings in containers.*—A Zyklon-B can containing 20 grams of HCN (approximately five-eighths the standard dose) was inclosed with test animals in a tightly sealed room. Five punctures were made in one end of the can with an 8-penny nail. The gas leakage from these holes was insufficient to affect test animals in three hours.

To check the potency of the contents this can was then opened in a gas-free room and the Zyklon-B spread in a thin layer on the floor. Exposed test animals were killed in 10 minutes.

(B) *Evolution of gas from Zyklon-B.*—To test the evolution of gas from Zyklon-B three sets of conditions were arranged: (1) Material spread in thin layers; (2) contents of can poured into one small compact pile; (3) can thoroughly opened, but material left in. Three rooms were used.

(1) Material spread in thin layers: Can of Zyklon-B containing 20 grams HCN was opened and contents were spread on the floor in thinnest possible layer. Exposed test animals were killed in nine minutes. Residue was then collected and placed in small gas-tight container with rat. Animal was unaffected after one hour, residue showing almost complete exhaustion of gas.

(2) Material spread in thick layer: Can of Zyklon-B containing 20 grams of HCN was opened and contents were carefully dumped so as to form pile of least circumference. Exposed test animals were killed in 23 minutes. Residue was then collected and placed in small gas-tight container with rat. Animal killed in 1-hour exposure.

(3) Can opened but contents left in: Can of Zyklon-B containing 20 grams HCN opened by cutting away the entire end. Contents of can were not disturbed. Exposed test animals were killed in 39 minutes. Contents of can were then poured into a small gas-tight container with rat. Animal was killed in two minutes.

(C) *Resistance of Zyklon-B containers to hard usage.*—Cans of Zyklon-B containing 20 grams and 100 grams of HCN and weighing (gross) approximately 200 grams and 400 grams, respectively, were dropped from a 45-foot tower to concrete pavement and thrown by a man with full force against solid brick walls without causing leakage. Test for leakage was made by placing these cans together with test animals in small gas-tight containers. The animals were unaffected after three hours' exposure.

(D) *Increase of pressure within Zyklon-B containers.*—Some pressure was exerted within in about one-half of the cans of Zyklon-B handled in all sizes from 20 to 1,200 grams HCN, as indicated by bulging

ends. A number of cans with unbulged ends were heated by direct exposure to sunlight on the steel deck of a vessel. With the increase in pressure due to temperature, the ends of these cans bulged with loud popping noises, but no leakage occurred.

The tests outlined above (A)-(D), indicate the safety with which Zyklon-B can be handled. No apprehension need be felt when transporting it through crowded streets, as, even if the transporting vehicles were wrecked, the small, strong containers would hardly be opened up; and if they were the relatively slow escape of gas into the open air would be without danger to the public.

(E) *Persistence of "tear effect" with Zyklon-B.*—Repeated observations and experiments indicate definitely that the "tear effect" produced by the irritating gases included with HCN in Zyklon-B is always effective as long as a dangerous quantity of HCN is present in the fumigated compartment. When large amounts of the material are used, as in holds, the "tear effect" persists for some time after the HCN has disappeared. This conclusion is based on combined human and rat tests wherein the test animal is lowered into the hold at the same time the officer goes below.

The claim of the manufacturers in this connection seems to be sufficiently substantiated: "The tear gas contained in Zyklon-B is intended only as a rear guard, and not as an advance guard; it being so much heavier than HCN, its rate of diffusion is much slower. Its useful function is as a warning when airing fumigated compartments, as during the period of fumigation it has time to develop fully, and remains irritant for some time after all traces of HCN have disappeared."

Our experience with the mixtures secured for test assures us that from 4 to 6 per cent of irritating gas affords ample protection. The mixture containing 10 per cent of irritating gas so prolongs the clearing of the larger compartments as to necessitate the holding of a vessel much longer than demanded by safety.

It is well here to insert a comparison with the cyanogen chloride mixture (HCN-CNCl) and to discuss the general proposition of protective gases. The writers are firmly of the belief that, when used in connection with liquid HCN, none of the irritating gases exert a dependable *pre-warning* effect. In the slower gas evolution of the generation method, time is sometimes given for escape, but this may not be relied upon. Cyanogen chloride does serve an important function during the process of active fumigation, as even the uninitiated would not get far into a compartment filled with the HCN-CNCl mixture. It is our opinion, however, that, with the present fumigating routine, the most dangerous phase of the operation is after the vessel has been opened up. The ideal warning gas is one which will deter both fumigators and other persons from

entering a gassed compartment until all traces of HCN have disappeared.

While thoroughly appreciating the theoretical integrity of the HCN-CNCl mixture, we have been forced to the conclusion that, in certain comparatively rare instances, considerable amounts of HCN did remain in moist, poorly ventilated holds after all tear effect had disappeared. This was confirmed experimentally by fumigating compartments in which were placed quantities of moist materials and subsequently testing for tear effect and retained HCN. On numerous occasions, not only was the presence of HCN easily detected by odor and taste, but test animals have been killed by direct exposure to the fumigated materials long after all traces of tear effect had disappeared. It was further observed that dry materials retained lethal quantities of HCN after the tear gas had been dissipated.

We feel that we can not too urgently stress the importance of not placing too much reliance on "tear effect." When using HCN alone, an experienced cyanide fumigator can, with the aid of test animals, declare a vessel safe for human occupancy with every assurance that no trouble will follow. On the other hand, no fumigator, regardless of his experience, is warranted in declaring a vessel safe merely because his eyes did not water when he inspected the sleeping quarters and holds. This applies particularly to the HCN-CNCl mixture; but even when using Zyklon-B the testimony of one's senses should be supplemented by exposing test animals at the danger points.

(8) SPECIAL CONSIDERATIONS OF THE USEFULNESS AND ADAPTABILITY OF VARIOUS CYANIDE PRODUCTS AND VARIOUS METHODS OF FUMIGATION

Liquid hydrocyanic acid, alone and mixed.—Considered only as a lethal agent, liquid hydrocyanic acid is the fumigant of choice. When public safety and ease of handling are likewise considered, it is not better than, if as good as, Zyklon-B. After all mechanical and other preparations are made and liquid HCN is placed on board in suitable applicators, one finds that fumigation has been simplified to the utmost degree. To gas a hold containing 120,000 cubic feet of air space one opens a valve, and when the indicator on a hand scale shows that 15 pounds of the material have entered the hold the valve is closed. Fifteen minutes will suffice to gas a cargo vessel of from 3,000 to 5,000 tons capacity. The dosing of superstructure, however, comprising some large and numerous small compartments, develops one of the major problems incident to the use of liquid HCN. As yet, no instrument has been devised which will rapidly and accurately deliver small doses of the gas; and until the cyanide fumigator is given an accurate

dosing machine, the fumigation of quarters will be a matter of guess-work.

In the light of our experiences, the tendency to overdose small compartments is constant. The danger of this practice is obvious, and in the absence of the suggested mechanical equipment can be counteracted only by deliberate underdosing. This perplexity is entirely overcome through the use of Zyklon-B with accurately measured doses of from 20 grams to 100 grams HCN always at hand.

Considerable mechanical equipment is required for handling liquid hydrocyanic acid. It is transported by truck from the manufacturer to the station using it, in large (75 pound) I. C. C. cylinders. The application of air pressure is required to pipe the liquid from these tanks to smaller (10 and 15 pound) cylinders used as applicators. These in turn have to be "pumped up" to a pressure sufficient to expel the liquid gas in the form of a fine spray. Numerous mechanical and chemical problems were met in connection with the applicators. These were successfully overcome, as regards administering large doses for hold fumigation, by the manufacturers of liquid cyanide, and the personnel of the New York quarantine station prior to the beginning of the studies dealt with in this report.

The explosive instability of liquid hydrocyanic acid has been recognized for many years. In addition to the explosion of gaseous mixtures of HCN and air, it is known that liquid hydrocyanic acid undergoes violent decomposition produced entirely by exothermic reactions occurring in the liquid in a closed container. The nature and mechanism of the exothermic polymerization and decomposition of liquid hydrocyanic acid has been carefully studied by the research chemists of one of the largest American chemical corporations, and the findings of these experts were such as to give rise to the following statement from the manufacturers: "This investigation and others made by the same company led to the conclusion that not enough is known about ways and means of stabilizing liquid hydrocyanic acid to warrant its shipment by *common carrier*. The — Chemical Company will continue indefinitely its present policy of shipping liquid hydrocyanic acid only under such conditions that its employed representatives may supervise and be entirely responsible for the product until it passes into the care of the ultimate consumer or of some equally responsible party." It is significant of the high character of the organization concerned that, at the time the above determination was expressed, the shipment of liquid hydrocyanic acid by common carrier was permitted under the regulations of the Interstate Commerce Commission.

From the standpoint of station fumigation, such findings regarding liquid HCN sharply define the responsibility accepted in storing and transporting the material.

Considered on a cost basis, liquid HCN is one of the least expensive of fumigants. The purchase price of the material is relatively low, there is no waste, the amount of cyanogen paid for is actually delivered into the compartment to be fumigated, and about one-half the personnel required for a generation method of cyaniding is needed. If the material were absolutely stable, and an easily portable and accurate device were available for delivering small doses, liquid hydrocyanic acid would be the ideal fumigant.

Zyklon-B.—All of the objections to liquid HCN are met and overcome with Zyklon-B. At the present time Zyklon-B is manufactured only in Germany, but the price per pound of HCN content is the same as that of the liquid HCN manufactured in the United States.

Compared on the basis of weight and bulk of materials required for fumigation, Zyklon-B runs a close second to liquid HCN. Prepared for fumigation, the liquid HCN applicators represent twice the weight of available HCN. The average package of Zyklon-B represents three times the weight of contained HCN. This is equivalent to saying that for a 3,000-ton ship, 75 pounds gross weight is carried for liquid cyanide fumigation and 114 pounds for Zyklon-B fumigation. The difference in weight is offset by the fact that when using Zyklon-B, empty cans and residue are thrown away so that there is no load on the return trip. As only four men are required for a liquid cyanide or Zyklon-B fumigation of a cargo vessel up to 5,000 net tons, the personnel cost is the same. All things being equal, the transportation costs of handling liquid cyanide and Zyklon-B would be about equal; but the complete safety with which Zyklon-B can be handled permits the use of much simpler and less expensive transport.

There is every assurance that Zyklon-B will be manufactured in the United States at an early date; and it is reasonable to assume that the cost of the product can be greatly reduced by local manufacture.

Generation method.—Results secured experimentally and on a large scale lead us to the conclusion that HCN generated by the barrel method is relatively much more efficient than the HCN-CNCl mixture similarly generated. So far as can be determined, this is due to the fact that the standard formula for the mixed HCN-CNCl gas produces less HCN than does HCN alone. Twenty per cent less sodium cyanide is used per 1,000 cubic feet, and the production of cyanogen chloride naturally utilizes some of the available cyanogen. In view of the fact that the "tear effect" of cyanogen chloride does not persist any longer than, if as long as, the HCN, even in proportions from 20 to 40 per cent, it is believed that its usefulness as a warning gas is thereby seriously vitiated.

We are firmly of the opinion that the generation method for the production of hydrocyanic acid gas for ship fumigation can not be justified in comparison with the liquid gas method and Zyklon-B method, when the items of cost, handling, and transportation are considered.

The excessive quantity and weight of equipment in the generation method necessitates the use of large trucks for transportation and a larger personnel than either of the other methods. Having arrived at the vessel to be fumigated, approximately four times the amount of time is required to get the ship under gas; and when the fumigation is completed, an equal length of time is required to remove barrels, buckets, etc., and to prepare them for return to the station. This loss of time not only affects the fumigating squads, which are frequently needed for other ships, but the vessels as well; and it is the loss of ship time after all that is of the most serious moment.

The least expensive item of generated HCN fumigation is that for the chemicals used. It is self-evident, however, that if such use of material is not uniformly productive of the results desired, i. e., a maximum rat kill, waste ensues which inevitably adds to cost, even though it is not immediately apparent.

An item of constant and increasing expense is that for barrels and buckets, which quickly break down under wear and tear and exposure to the diluted acid. The cost of heavy truck transportation plus depreciation adds to the steadily mounting expense, exclusive of the greater personnel required.

Calcium cyanide.—As has been previously stated, calcium cyanide compares favorably with both liquid HCN and Zyklon-B in killing efficiency when HCN content is used as the basis of comparison. It is most effective when applied in the form of a dust; but when applied in layers the rapidity of the evolution of HCN is, within limits, proportional to the thickness of the layers. For all practical purposes, calcium cyanide can not be dusted into vessels, as the residue is objectional, and unless laid down on sheets of paper or otherwise so that it can be completely removed, its use must be criticized not only on grounds of cleanliness, but because a variable proportion of its HCN content is retained for periods greatly in excess of the time permitted in routine ship fumigation.

Inasmuch as only one-half the volume of calcium cyanide, by weight, is HCN, its present price of \$1 per pound gross weight is excessively costly. We understand that reductions in cost up to 15 per cent are made on large quantities, but even so the cost of HCN content will then be about \$1.70 per pound as compared with \$0.90 to \$1 for liquid HCN and Zyklon-B.

In this connection, we wish to state that none of our experimental work substantiated the claim of the manufacturers of calcium cyanide

that in lethal efficiency, it was the equivalent of liquid cyanide "pound for pound."

If the points of objection raised were of no moment, a competent fumigation with calcium cyanide would be acceptable, but as the material has no qualities superior to the other cyanogen products tested, we do not recommend it for prior consideration at this time.

GENERAL ASPECTS OF FUMIGATION WITH SPECIAL REFERENCE TO
THE APPLICABILITY OF CYANIDE FUMIGANTS AT VARIOUS QUARANTINE STATIONS OF THE SERVICE

A sincere effort has been made to view the question of cyanide fumigation from the angle of the small and sometimes isolated quarantine station with limited personnel and equipment. While we have dealt only with cyanogen products, we have not lost sight of the fact that some such stations are not yet, and may not for a long time be, ready for this type of fumigation. We believe, however, that if ship fumigation is ever to reach the plane of a scientifically controlled procedure, some radical changes will have to be made, and these involve the selection and development of a highly lethal agent which can be handled with comparative safety under all circumstances.

If managed with due care and proper respect, hydrocyanic acid is the best of all fumigants for rodents. Careless use of it will be attended by human fatalities, but it will kill rats under conditions that the use of sulphur can not meet; and it is obvious that if fumigation does not kill rats, time and money are wasted.

It is apparent that the present-day routine fumigation does not kill all of the rats in a vessel even when cyanide is used. This is hardly a criticism of HCN as a lethal agent (as it has been shown that even one-fifteenth of the standard dose is uniformly fatal to exposed rats), but points rather to the fact that usual procedure, method of application, and other factors on board the ship operate against complete success.

It is certain that, if the concentration of gas theoretically obtained by introducing a predetermined number of pounds or ounces of HCN reaches the rats on board a vessel, the rats will be killed. It is no less certain that, if the animals survive, the expected concentration and the rat did not meet. Proper preparation of a vessel for fumigation (and this includes the fulfillment of certain structural requirements from the time the keel is laid) is absolutely essential if gaseous fumigation is ever to become a more exact rat-eradicator measure. So long as there are contiguous dead air spaces or pockets into which rats may escape, only partial results will be secured. No gas, no matter how lethal nor in what concentration used, can be expected to follow comparatively long and tortuous rat runs nor to pass through small openings into practically dead air spaces by diffusion within the two hours usually allotted for fumigation.

Successful fumigation also depends largely on proper location and distribution of the gas generation centers. To insure maximum efficiency the gas should be introduced as near as possible to rat-harboring places, and several well distributed small "shots" in a large compartment are far more efficacious than one big one. When using the generation method, containers should be placed on the floor of the lower hold and on the "between decks" and not swung from the hatch coamings as has been suggested and advocated heretofore.

To kill rats is the prime object of ship fumigation. To accomplish this purpose in the interest of the public welfare, the most thorough and painstaking measures are warranted. If, however, it could be known that there were very few or no rats on board a vessel, the fumigation of the vessel would be unjustifiable. It is believed that it is usually possible to determine by competent inspection whether or not a significant number of rats infest a vessel at a given time. It seems evident that further experimentation along this line will be productive of fruitful results as the cooperation and support of shipping interests can be counted on for the furtherance of a plan which would promise definite relief from unnecessary delay and expense. With inspection as the basis for determining the fumigation status of a vessel, only such vessels as showed evidence of rats would be fumigated, and these would be handled in a thoroughly competent manner. After adequate preparation, repeated protracted fumigations would be undertaken for the purpose of *ridding* the vessel of rats. These vessels would not be fumigated again until evidence of rats showed renewed infestation. Shipping companies could be depended on to prosecute active rat trapping, rat-proofing, and other eradivative measures as the best means of deferring fumigation. The fact must not be lost sight of that it is the presence of rats on board and not merely the plague status of ports of departure, nor the interval since the last fumigation, which defines the potential infectiousness of a vessel.

Liquid cyanide can be used only at certain of the larger quarantine stations having trained personnel and adequate mechanical facilities. Supplies of liquid gas must also be procurable from nearby depots and "service stations," for such equipment must be readily available. At stations where liquid cyanide is used, supplies of Zyklon-B should be maintained for the fumigation of superstructure compartments, where the demand for accurate small doses of HCN is constant. The combination of liquid cyanide for the holds and Zyklon-B for the sleeping quarters and deck compartments is almost ideal.

Zyklon-B will meet every requirement of the smaller stations. It is compact and easily stored. The package withstands rough handling and lasts indefinitely or until punctured. Convenient doses of HCN are provided, and dosing with the material is practically

"foolproof." Because of the granular consistency and the free-running quality of the material, highly satisfactory gas distribution is accomplished without effort. Three men can fumigate a 3,000- or 4,000-ton empty cargo vessel in less time than can six men using the generation method. When fumigation is completed, empty cans and residue are thrown overboard, so there is no return load.

For stations not yet ready to relinquish sulphur as a fumigant, a useful combination will be found in sulphur for the holds and Zyklon-B in all upper deck compartments where the destructive effects of sulphur are objectionable. Such stations should be encouraged in the use of cyanide, however, as a sulphur fumigation is time consuming and, except in the instance of unusually well-prepared vessels, does not compare with cyanide.

Hydrocyanid acid gas does not affect metals, fabrics, or foodstuffs. Its relatively high rate of solubility in water, however, indicates the advisability of pouring out all drinking water and other beverages directly exposed to the gas during fumigation, and the prompt pumping out of bilges following fumigation.

All persons directly engaged in cyanide fumigation should be equipped with an efficient anticyanide gas mask and compelled to wear it both while dosing and opening up compartments. In entering holds to test gas, the mask should be carried in such a position that it can be instantly applied. If gas is encountered in a hold, panic should be avoided. Apply the mask and *walk* to the well-ventilated area immediately beneath the hatch opening. Do not attempt to climb the hold ladders immediately, but wait until the head is clear, the heart beat steady and slow, and the knees are strong. When leaving the hold, climb slowly.

The subject of ship fumigation is one of the greatest sanitary importance. Until a more efficient method for the eradication of potentially plague-infected rats is devised and its specificity proved beyond any reasonable doubt, fumigation will stand as the procedure of choice. It will repay all who are concerned to study the subject carefully to the end that the manner of its performance and the measures used are made more efficient.

The lessons learned in the course of the experimental work covered in this report point insistently to one conclusion: FUMIGATE FEWER SHIPS BETTER.

COURT DECISION RELATING TO PUBLIC HEALTH

Occupational disease held not compensable.—(Maine Supreme Judicial Court; Dillingham's Case, 142 A. 865; decided August 20, 1928.) The Maine workmen's compensation act provided:

If an employee * * * receives a personal injury by accident, * * * he shall be paid compensation.

The question was presented to the supreme court as to whether an occupational disease was a personal injury by accident under said statute. In deciding that an occupational disease was not compensable the court used this language:

Accident has been defined, in cases under the act [workmen's compensation act], as an unusual, undesigned, unexpected, and sudden event resulting in injury. [Cases cited.] Disease, to be compensable, must be interpreted both as an "injury" and an "accident." An occupational or industrial disease is one normally peculiar to and gradually caused by the occupation in which the afflicted employee is or was regularly engaged, and to which everyone similarly working in the same industry is alike constantly exposed. * * *

Cases of occupational disease, remarked Mr. Justice Philbrook in Brodin's Case, 124 Me. 162, 126 A. 829, can not be said to have arisen from accidental causes, since they lack the element of sudden or unexpected event. Obiter dictum and not adjudication was that remark, surely. But it served well to differentiate in the case where it was made, and in the present case it is entitled to, and does, receive respect, when for the first time the point necessarily arises whether disease caused by occupation, in the restricted sense of a disease which is not merely a risk of the particular employment, but also of gradual growth, may as matter of law be ruled to be personal injury by accident.

Without examining all the decided cases in States where the workmen's compensation enactments are in similarity to our own, apparently the weight of authority is to the effect that cases of occupational or industrial poisoning can not be regarded as accidents, within the meaning of statutes which provide for money payments to workmen for injuries caused by accident arising out of and in the course of their employment. The ground fixed by the statute, says Mr. Justice Swayze, in New Jersey, is the injury by accident, not the results of an indefinite something which may not be an accident. * * * In Massachusetts, where the statute is for personal injury without reference to accident, the court has said that "personal injury by accident" is not so broad in scope as "personal injury." [Case cited.]

It is the conclusion of this court that, as disability caused by personal injury by accident arising out of and in the course of his employment is a statutory prerequisite for the payment of compensation to an injured employee, this claimant's injury, from what in a like situation some judge phrased the insensible progress of occupational disease, was not as matter of law received by accident.
* * *

PUBLIC HEALTH ENGINEERING ABSTRACTS

Streptococcus as an Indicator of Swimming-Pool Pollution. W. L. Mallman. *American Journal of Public Health and The Nation's Health*, vol. 18, No. 6, June, 1928, pp. 771-776. (Abstract by C. T. Butterfield.)

The author discusses the standards adopted by the California State Board of Health, 1919, and the 1923 report of the committee appointed by the American Public Health Association. Results, *B. coli* and total count of a two years' study of swimming pools are given. *B. coli* were absent in 1 c. c. portions and the total count was below 1,000 per c. c. No cases of disease were traced to the use of these pools during this period. Results are next presented covering a one-year period in which tests for streptococci were also included.

Mallman concludes in part that in swimming pools: (1) *B. coli* is not always a reliable indicator of pollution; (2) streptococci are constant indicators of pollution; (3) *B. coli* tend to multiply in such water; streptococci die out; (4) the

presence of streptococci indicate an unsafe condition; (5) the presence of *B. coli* does not necessarily indicate pollution but their absence is an excellent index of safety.

White Lead. Anon. *Public Health*, vol. 41, No. 5, February, 1928, pp. 132-133. (Abstract by W. L. Havens.)

The prevalence of lead poisoning in the paint industry caused by the inhalation of dust impregnated with lead has resulted in rather drastic legislation abroad. Since 1921 the White Lead Convention of Geneva has been ratified by 13 countries undertaking to prohibit the use of white lead and sulphate of lead in the internal painting of buildings. In Great Britain the lead paint act of 1926 empowers the Home Secretary to make regulations prohibiting the employment of women and young persons in painting with lead paints and providing for reports and registration of plumbism among employees.

Sewage Agitation and Chlorination Tests at Havre, Mont. Emil Sandquist and H. B. Foote. *Engineering News-Record*, vol. 100, No. 26, June 28, 1928, pp. 1001-1002. (Abstract by R. J. Faust.)

During the summer months the Milk River, on which Havre is located, has not sufficient flow adequately to oxidize the city's sewage. An agitation and chlorination plant was established at the outlet of one of three main sewer outlets with the following results:

Agitation reduced suspended solids from 0.882 per cent to 0.102 per cent, a reduction of 88 per cent; no marked difference was noted in the amount of settleable solids before and after agitation. With agitation only, there was an average reduction of hydrogen sulphide of 26.5 per cent (from 4.08 to 3 p. p. m.); agitation combined with chlorination gave an average reduction of hydrogen sulphide of 63 per cent; bacterial reductions were great (shown by an accompanying table); chlorine residual after 10 minutes was maintained between 0.5 and 1.0 p. p. m.; cost of operating plant 1 month was estimated at \$110. This plant treated about 0.122 m. g. d.

Calculating the Capacity of Sludge Digestion Tanks. Karl Imhoff. (Trans. from German by Gordon M. Fair.) *American City*, vol. 38, No. 5, May, 1928, pp. 124-125. (Abstract by J. B. Harrington.)

The required capacity of sludge digestion tanks is described in this article. Ordinarily, with a mean temperature of 59° F., 1 cubic foot per capita sludge volume is required. At 50° F., the sludge volume must be doubled. In the northern part of the United States, where the winters are more severe, and in towns of less than 5,000, the capacity of the sludge tank should be increased. The size depends upon the time interval between drawing sludge in the fall and spring. At 59° F. two months' time is necessary for complete digestion. Additional allowance must also be made for wastes likely to retard digestion.

The Function of Aeration in Water Purification. N. T. Veatch, jr. Proceedings of Tenth Texas Water Works Short School, January, 1928, pp. 172-177. Abstract by Chester Cohen.

Experiments with simple aerators have proved the following: Oxygen may be added to water to the point of saturation; carbon dioxide may be largely but not completely removed; odors and tastes due to gases of decomposition may be partially removed, but if due to products of organic decomposition the effect is negligible; odors and tastes due to coal tar, phenol, and similar wastes are little affected; hydrogen sulphide and odors due to over-chlorination may be satisfactorily removed; the oxidation of organic matter is negligible, but inorganic compounds, as iron and manganese, are largely but not completely oxidized; the bacterial content of water is often reduced; the economy and effectiveness of treatment with ordinary coagulants is usually increased; the corrosiveness of

soft or peaty waters is usually increased, whereas surface supplies are not usually affected.

Four different types of aerators are discussed; namely, the air lift, the injection, the gravity, and the fountain aerator. "Results of tests on aerators of different types, operating under different heads with different waters, vary so greatly that no general conclusion can be made as to the effect any particular type of aerator will have on a given water. * * *"

The Microscopic Life in Water. Asa C. Chandler. Proceedings of Tenth Texas Water Works Short School, January, 1928, pp. 155-159. (Abstract by Chester Cohen.)

This article presents a very interesting discussion of the part played by microscopic organisms in water. A discussion of the factors which influence the development of microscopic organisms is given, with explanations of the influence of sunlight, temperature, overturn, food supplies, and aeration. The relation between stream pollution and the abundance and character of micro-organisms is explained. The by-products of plant life, such as odor, taste, and color in water, are accounted for; the effect of the growths on pipes, sand filter beds, and oxygen content of water is discussed, together with methods for control of the organisms, either through the use of copper sulphate or chlorine. The author mentions, in concluding, that there is an important field of biological investigation which could be developed with relation to assisting nature in establishing an equilibrium in the growth of the plant life, which would tend to maintain stable conditions and prevent the sudden fluctuations and changes that take place in the water supplies.

Dosing Apparatus. Wynkoop Kiersted. Proceedings of Tenth Texas Water Works Short School, January, 1928, pp. 54-57. (Abstract by Chester Cohen.)

This article brings out the fact that local conditions modify both the design and mechanics required for the proper application of coagulants. The solution method of applying lime is probably cheaper in most cases, due to the economy possible in the purchase of lime and subsequent slaking.

"Two basic methods of preparing and distributing the coagulants by the solution method are in common use. One is the constant-volume method and the other the constant-strength method." The constant-volume method is used in dissolving and applying crystalline coagulants, as alum and iron, and the constant-strength method is used in the case of application of lime or when a gravity or some other easily regulated flow may be added to the point of application.

The various methods of mixing the coagulants with the raw water are discussed. "I believe that when there is sufficiently violent agitation of the water in the mixing chamber, little or no time is required for reaction before passing into the settling basin, but when handling a large volume of water where only a portion of the raw water passes through the mixing chamber, a little additional time will be required for mixing the coagulated portion of the raw water with the noncoagulated portion."

Mr. Kiersted sums up his paper by explaining that although solution feed would probably be more economical in the larger water-works plants, the dry-feed method finds greatest favor in the smaller plants.

Experiences with Covered and Open Reservoirs. Carl J. Lauter. *Engineering News-Record*, vol. 100, No. 25, June 21, 1928, pp. 963-964. (Abstract by R. J. Faust.)

"Three years' bacterial, microscopic, and chemical tests on an open and a closed reservoir in the city of Washington, D. C., each receiving the same supply from the slow sand filtration plant, chlorination then being intermittent, showed the following results:

"Average bacteria count, 4 for filtered water, 7.6 for the covered reservoir, an increase of only 90 per cent, as compared with 44 for the open basin, an increase

of 1,000 per cent. There was no micro-organism growth in the covered reservoir, whereas it was abundant in the open one. Albuminoid ammonia, nitrate, and nitrite tests also showed much better conditions in the covered basin at all times."

DEATHS DURING WEEK ENDED SEPTEMBER 29, 1928

Summary of information received by telegraph from industrial insurance companies for the week ended September 29, 1928, and corresponding week of 1927. (From the Weekly Health Index, October 3, 1928, issued by the Bureau of the Census, Department of Commerce)

	Week ended Sept. 29, 1928	Corresponding week, 1927
Policies in force.....	71, 769, 909	68, 810, 404
Number of death claims.....	12, 623	10, 934
Death claims per 1,000 policies in force, annual rate.....	9. 2	8. 3

Deaths from all causes in certain large cities of the United States during the week ended September 29, 1928, infant mortality, annual death rate, and comparison with corresponding week of 1927. (From the Weekly Health Index, October 3, 1928, issued by the Bureau of the Census, Department of Commerce)

City	Week ended Sept. 29, 1928		Annual death rate per 1,000 corresponding week 1927	Deaths under 1 year		Infant mortality rate, week ended Sept. 29, 1928 ¹
	Total deaths	Death rate ¹		Week ended Sept. 29, 1928	Corresponding week 1927	
Total (68 cities).....	6, 370	11. 0	10. 8	726	735	60
Akron.....	47			10	7	109
Albany ¹	34	14. 8	17. 0	4	6	82
Atlanta.....	73	15. 0	14. 2	9	5	
White.....	36		10. 4	5	4	
Colored.....	37	(¹)	23. 3	4	1	
Baltimore ¹	238	15. 0	13. 3	28	22	89
White.....	188		11. 1	21	14	84
Colored.....	50	(¹)	26. 0	7	8	110
Birmingham.....	49	11. 5	15. 1	8	9	68
White.....	23		11. 4	6	6	83
Colored.....	26	(¹)	20. 9	2	3	45
Boston.....	168	11. 0	12. 0	16	27	44
Bridgeport.....	20			1	3	18
Buffalo.....	141	13. 3	12. 9	16	22	69
Cambridge.....	18	7. 5	10. 5	2	2	36
Camden.....	24	9. 3	12. 9	4	5	64
Canton.....	21	9. 4	12. 9	2	3	48
Chicago ¹	648	10. 7	9. 9	62	80	53
Cincinnati.....	146	18. 5	13. 9	24	10	145
Cleveland.....	167	8. 6	7. 9	27	23	73
Columbus.....	61	10. 7	12. 2	8	7	75
Dallas.....	41	9. 9	9. 9	8	4	
White.....	28		9. 1	7	3	
Colored.....	13	(¹)	15. 2	1	1	
Dayton.....	37	10. 5	11. 3	6	5	99
Denver.....	74	13. 2	11. 7	15	6	
Des Moines.....	29	10. 0	10. 5	3	1	50
Detroit.....	267	10. 1	9. 2	42	36	65
Duluth.....	24	10. 7	10. 9	4	1	93
El Paso.....	24	10. 7	13. 3	3	6	
Erie.....	22			2	0	41
Fall River ¹	25	9. 7	9. 0	3	1	51
Flint.....	36	10. 5	12. 1	8	10	102
Fort Worth.....	28	8. 7	10. 5	4	4	
White.....	26		9. 4	4	4	
Colored.....	2	(¹)	18. 6	0	0	
Grand Rapids.....	31	9. 9	8. 4	4	5	60
Houston.....	70			5	5	
White.....	49			4	4	
Colored.....	21	(¹)		1	1	
Indianapolis.....	81	11. 1	13. 1	10	10	76
White.....	68		11. 7	7	8	61
Colored.....	13	(¹)	23. 3	3	2	182
Jersey City.....	47	7. 6	9. 7	4	12	30

(Footnotes at end of table)

Deaths from all causes in certain large cities of the United States during the week ended September 29, 1928, infant mortality, annual death rate, and comparison with corresponding week of 1927—Continued

City	Week ended Sept. 29, 1928		Annual death rate per 1,000 corresponding week 1927	Deaths under 1 year		Infant mortality rate, week ended Sept. 29, 1928 ¹
	Total deaths	Death rate ¹		Week ended Sept. 29, 1928	Corresponding week 1927	
Kansas City, Kans.	22	9.7	11.1	1	2	21
White	14		11.4	1	2	25
Colored	8	(²)	9.8	0	0	0
Kansas City, Mo.	82	11.0	10.8	9	9	64
Knoxville	36	14.9	11.7	8	5	174
White	19		12.8	6	5	145
Colored	11	(²)	4.3	2	0	427
Los Angeles	240			28	27	80
Louisville	104	16.5	9.4	13	8	109
White	75		7.3	12	8	114
Colored	29	(²)	21.3	1	0	69
Lowell	30	14.2	9.0	2	2	42
Lynn	18	8.0	11.4	1	0	25
Memphis	65	17.9	12.2	3	2	35
White	30		16.8	1	1	19
Colored	35	(²)	14.8	2	1	63
Milwaukee	100	9.6	10.2	11	19	49
Minneapolis	72	8.3	6.5	1	3	6
Nashville	44	16.6	15.9	2	2	31
White	28		11.6	0	0	0
Colored	16	(²)	26.8	2	2	120
New Bedford	16	7.0	7.4	2	2	43
New Haven	33	9.2	9.0	5	4	71
New Orleans	134	16.3	18.7	20	18	97
White	76		14.4	16	9	116
Colored	58	(²)	20.7	4	9	58
New York	1,221	10.6	10.0	134	112	54
Bronx Borough	160	8.8	7.7	10	7	30
Brooklyn Borough	379	8.6	8.9	50	43	50
Manhattan Borough	604	15.0	13.8	52	52	62
Queens Borough	130	8.0	7.2	19	8	76
Richmond Borough	48	16.7	11.4	3	2	54
Newark, N. J.	68	7.5	9.7	7	13	36
Oakland	44	8.4	12.7	4	6	43
Oklahoma City	20			1	3	
Omaha	53	12.4	11.7	3	3	35
Paterson	35	12.6	8.0	3	0	52
Philadelphia	387	9.8	10.4	49	43	66
Pittsburgh	133	10.7	11.4	20	32	65
Portland, Oreg.	74			2	4	21
Providence	50	9.1	10.8	6	6	52
Richmond	50	13.4	12.2	8	5	104
White	29		8.4	3	2	61
Colored	21	(²)	21.6	5	3	184
Rochester	61	9.7	10.0	5	3	41
St. Louis	188	11.6	15.2	18	21	60
St. Paul	51	10.6	11.3	3	5	29
Salt Lake City ¹	29	11.0	9.2	4	6	65
San Antonio	47	11.3	9.4	11	8	
San Diego	33	14.4	14.9	1	4	19
San Francisco	161	14.4	9.6	4	7	25
Schenectady	20	11.2	6.7	2	1	63
Somerville	9	4.6	9.2	0	3	0
Spokane	23	11.0	12.4	1	2	26
Springfield, Mass.	25	8.7	9.2	1	7	16
Syracuse	60	15.7	11.1	2	6	24
Tacoma	25	11.8	10.2	1	2	26
Toledo	57	9.5	9.7	6	2	58
Trenton	35	13.2	13.0	5	6	85
Washington, D. C.	116	11.0	13.9	13	16	74
White	69		11.9	9	11	74
Colored	47	(²)	19.9	4	5	74
Waterbury	15			4	2	116
Wilmington, Del.	20	8.1	12.0	2	2	53
Worcester	46	12.2	13.1	5	6	61
Yonkers	18	7.8	8.3	3	2	68
Youngstown	24	7.2	10.5	3	7	40

¹ Annual rate per 1,000 population.

² Deaths under 1 year per 1,000 births. Cities left blank are not in the registration area for births.

³ Deaths for week ended Friday, Sept. 28, 1928.

⁴ In the cities for which deaths are shown by color, the colored population in 1920 constituted the following percentages of the total population: Atlanta, 31; Baltimore, 15; Birmingham, 39; Dallas, 15; Fort Worth, 14; Houston, 25; Indianapolis, 11; Kansas City, Kans., 14; Knoxville, 15; Louisville, 17; Memphis, 36; Nashville, 30; New Orleans, 26; Richmond, 32; and Washington, D. C., 28.

PREVALENCE OF DISEASE

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

UNITED STATES

CURRENT WEEKLY STATE REPORTS

These reports are preliminary and the figures are subject to change when later returns are received by the State health officers

Reports for Weeks Ended September 29, 1928, and October 1, 1927

Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended September 29, 1928, and October 1, 1927

Division and State	Diphtheria		Influenza		Measles		Meningococcus meningitis	
	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927
New England States:								
Maine.....	1	3	2	-----	20	9	0	0
New Hampshire.....	1	-----	9	-----	15	-----	0	-----
Vermont.....	-----	2	-----	-----	2	-----	0	0
Massachusetts.....	34	74	8	6	50	41	1	0
Rhode Island.....	7	5	-----	-----	-----	-----	0	1
Connecticut.....	17	19	2	1	11	9	2	3
Middle Atlantic States:								
New York.....	50	168	11	13	88	59	18	7
New Jersey.....	80	105	5	2	12	8	4	1
Pennsylvania.....	163	24	-----	-----	103	96	3	2
East North Central States:								
Ohio.....	83	-----	6	-----	65	-----	8	-----
Indiana.....	46	29	23	3	10	7	0	0
Illinois.....	100	100	17	7	40	18	9	10
Michigan.....	82	79	2	2	17	17	6	2
Wisconsin.....	10	26	15	36	27	62	0	2
West North Central States:								
Minnesota.....	30	39	1	-----	7	4	0	2
Iowa.....	12	14	-----	-----	-----	3	0	1
Missouri.....	33	40	6	-----	2	3	1	1
North Dakota.....	13	11	2	-----	3	21	9	0
South Dakota.....	-----	-----	-----	-----	1	-----	0	0
Nebraska.....	10	9	-----	-----	1	1	0	0
Kansas.....	16	51	1	1	4	24	0	1
South Atlantic States:								
Delaware.....	1	2	-----	1	-----	1	0	0
Maryland ¹	25	29	4	2	13	9	0	1
District of Columbia.....	12	16	-----	-----	4	2	0	0
Virginia.....	-----	-----	-----	-----	-----	-----	-----	-----
West Virginia.....	20	19	11	-----	12	6	1	0
North Carolina.....	147	176	-----	-----	5	106	0	2
South Carolina.....	59	72	664	216	-----	97	0	0
Georgia.....	28	48	118	33	-----	8	1	0
Florida.....	-----	6	-----	2	-----	1	-----	2

¹ New York City only.

² Week ended Friday.

Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended September 29, 1928, and October 1, 1927—Continued

Division and State	Diphtheria		Influenza		Measles		Meningococcus meningitis	
	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927
East South Central States:								
Kentucky.....	27						1	0
Tennessee.....	66	26	27	6		26	2	0
Alabama.....	75	77	46	7	73	11	1	0
Mississippi.....	33	39					0	1
West South Central States:								
Arkansas.....	14	9	34	43	9	6	0	1
Louisiana.....	23	37	20	13		10	0	1
Oklahoma.....	65	68	14	6	8	13	1	1
Texas.....	26	44	19	30	6	2	0	1
Mountain States:								
Montana.....	2	5			4	2	0	1
Idaho.....		1			1	2	1	0
Wyoming.....		2			1	4	0	0
Colorado.....	12	34	2		1	4	3	1
New Mexico.....	13	2	1		6	14	0	0
Arizona.....		1				1	0	0
Utah.....	3	9	2	3		1	0	0
Pacific States:								
Washington.....	12	18	2		19	19	3	4
Oregon.....	11	6	2	25	7	2	1	4
California.....	69	88	21	12	33	25	6	1

Division and State	Poliomyelitis		Scarlet fever		Smallpox		Typhoid fever	
	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927
New England States:								
Maine.....	3	5	24	16	0	0	4	6
New Hampshire.....	2		1		6		0	0
Vermont.....	1	2	4	6	0	0	0	0
Massachusetts.....	20	79	99	129	0	0	9	19
Rhode Island.....	0	1	3	10	0	0	1	2
Connecticut.....	2	13	14	22	0	0	2	5
Middle Atlantic States:								
New York.....	63	60	91	138	0	6	78	51
New Jersey.....	1	38	30	51	0	4	20	10
Pennsylvania.....	11	35	116	149	0	0	77	65
East North Central States:								
Ohio.....	19	87	134		4		60	
Indiana.....	2	18	34	59	4	5	14	21
Illinois.....	1	50	114	129	16	5	46	46
Michigan.....	3	21	161	100	18	7	14	15
Wisconsin.....	1	19	53	41	3	7	6	17
West North Central States:								
Minnesota.....	14	15	56	61	0	1	3	9
Iowa.....	3	6	21	20	0	5	8	5
Missouri.....	0	20	52	41	7	5	37	9
North Dakota.....	4	4	29	47	0	0	3	3
South Dakota.....	0	7	9	18	8	0	1	3
Nebraska.....	2	7	24	23	79	0	4	1
Kansas.....	2	19	70	64	3	2	20	28
South Atlantic States:								
Delaware.....	0	0	4	0	0	0	4	
Maryland.....	13	3	16	16	0	0	46	24
District of Columbia.....	0	3	7	10	0	0	1	3
Virginia.....	1							
West Virginia.....	11	22	41	50	1	8	27	31
North Carolina.....	0	1	70	76	5	9	36	60
South Carolina.....	2	2	6	26	0	2	51	69
Georgia.....	0	0	11	29	0	0	28	27
Florida.....		1		7		1		3

¹ Week ended Friday.

² Figures for 1928 are exclusive of Oklahoma City and Tulsa and for 1927 are exclusive of Tulsa.

Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended September 29, 1928, and October 1, 1927—Continued

Division and State	Poliomyelitis		Scarlet fever		Smallpox		Typhoid fever	
	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927	Week ended Sept. 29, 1928	Week ended Oct. 1, 1927
East South Central States:								
Kentucky.....	0		43		0		20	
Tennessee.....	1	3	33	35	1	0	71	59
Alabama.....	4	0	28	25	1	0	47	36
Mississippi.....	0	0	23	29	1	1	14	9
West South Central States:								
Arkansas.....	0	4	22	7	0	0	30	15
Louisiana.....	0	3	10	7	3	9	36	13
Oklahoma ¹	0	7	35	21	6	7	76	86
Texas.....	1	12	10	19	0	4	22	22
Mountain States:								
Montana.....	5	0	10	9	12	6	9	6
Idaho.....	1	1	7	4	11	0	1	1
Wyoming.....	0	1	8	9	1	0	0	
Colorado.....	2	9	12	31	2	1	2	9
New Mexico.....	1	9	13	4	10	2	13	15
Arizona.....	0	1	2	1	0	0	0	0
Utah ²	0	2	9	2	1	6	1	4
Pacific States:								
Washington.....	11	16	20	17	23	12	7	16
Oregon.....	2	30	12	25	15	24	8	5
California.....	6	46	84	73	26	8	22	16

¹ Week ended Friday.

² Figures for 1928 are exclusive of Oklahoma City and Tulsa and for 1927 are exclusive of Tulsa.

SUMMARY OF MONTHLY REPORTS FROM STATES

The following summary of monthly State reports is published weekly and covers only those States from which reports are received during the current week:

State	Menin- gococ- cus menin- gitis	Diph- theria	Infl- uenza	Malaria	Measles	Pella- gra	Polio- myelitis	Scarlet fever	Small- pox	Ty- phoid fever
<i>August, 1928</i>										
California.....	15	274	44	7	67	6	31	224	48	137
District of Columbia.....	0	60			29		8	9	0	7
Florida.....	6	42	149	123	16	5	1	9	1	36
Idaho.....	3	12			4		16	23	21	22
Indiana.....	0	38	21		60		2	94	53	87
Mississippi.....	2	60	2, 203	20, 882	155	1, 514	7	31	1	289
Montana.....	6	8			29		33	11	21	12
Oregon.....	5	27	15		37		8	35	58	23
Pennsylvania.....	24	399		1	732	1	44	296	2	247
Virginia.....	2	93	571	168	220	44	23	63	9	232
Washington.....	5	28	11		47		89	34	61	33
West Virginia.....	1	37	41		46		69	84	18	139

<i>August, 1928</i>		Chicken pox—Continued.	
Actinomycosis:	Cases	Indiana.....	Cases
California.....	1	Mississippi.....	218
Anthrax:		Montana.....	8
California.....	2	Oregon.....	25
Mississippi.....	1	Pennsylvania.....	150
Oregon.....	4	Virginia.....	39
Chicken pox:		Washington.....	66
California.....	199	West Virginia.....	15
District of Columbia.....	3	Dengue:	
Florida.....	40	Florida.....	1
Idaho.....	3	Mississippi.....	487

Dysentery:	Cases	Puerperal septicemia:	Cases
California (amebic).....	7	Mississippi.....	56
California (bacillary).....	7	Pennsylvania.....	6
Florida.....	14	Rabies in animals:	
Mississippi (amebic).....	104	California.....	63
Mississippi (bacillary).....	1,135	Mississippi.....	9
Oregon.....	1	Oregon.....	4
Pennsylvania.....	1	Washington.....	1
Virginia.....	602	Rabies in man:	
German measles:		Pennsylvania.....	1
California.....	65	Rocky Mountain spotted or tick fever:	
Pennsylvania.....	16	Montana.....	1
Washington.....	14	Oregon.....	1
Hookworm disease:		Scabies:	
Florida.....	43	Oregon.....	1
Mississippi.....	394	Septic sore throat:	
Virginia.....	9	Oregon.....	6
Impetigo contagiosa:		Tetanus:	
Oregon.....	3	California.....	12
Washington.....	5	Florida.....	8
Leprosy:		Pennsylvania.....	8
California.....	6	Trachoma:	
Pennsylvania.....	1	California.....	6
Lethargic encephalitis:		Mississippi.....	5
California.....	2	Oregon.....	1
District of Columbia.....	1	Pennsylvania.....	3
Florida.....	1	Tularaemia:	
Pennsylvania.....	11	California.....	1
Virginia.....	1	Indiana.....	1
Washington.....	3	Typhus fever:	
Mumps:		California.....	11
California.....	261	Florida.....	11
Florida.....	15	Virginia.....	4
Idaho.....	3	Undulant fever:	
Indiana.....	9	California.....	2
Mississippi.....	185	Oregon.....	1
Oregon.....	14	Whooping cough:	
Pennsylvania.....	227	California.....	779
Washington.....	43	District of Columbia.....	41
Ophthalmia neonatorum:		Florida.....	28
California.....	3	Idaho.....	8
Mississippi.....	10	Indiana.....	97
Pennsylvania.....	14	Mississippi.....	569
Paratyphoid fever:		Montana.....	8
California.....	3	Oregon.....	27
Idaho.....	5	Pennsylvania.....	1,746
Oregon.....	2	Virginia.....	276
Plague:		Washington.....	65
California.....	1	West Virginia.....	72

Number of Cases of Certain Communicable Diseases Reported for the Month of July, 1928, by State Health Officers

State	Chick- en pox	Diph- theria	Measles	Mumps	Scarlet fever	Small- pox	Tuber- culosis	Ty- phoid fever	Whoop- ing cough
Maine.....	37	8	229	47	32	0	35	10	95
New Hampshire.....		9			12	0		0	
Vermont.....	47	2	177	33	14	0	11	1	126
Massachusetts.....	290	163	1,350	164	317	1	529	28	405
Rhode Island.....	12	29	634	10	41	0	41	2	21
Connecticut.....	54	51	798	110	66	0	152	4	356
New York.....	701	837	3,804	415	439	5	1,305	111	1,421
New Jersey.....	178	347	1,255		131	2	406	36	618
Pennsylvania.....	363	391	2,688	527	390	37	713	155	1,377
Ohio.....	323	153	1,400	132	228	54	770	111	1,007
Indiana.....	43	46	247	19	103	104	164	19	82
Illinois.....	551	312	379	248	405	88	1,035	86	921
Michigan.....	222	210	1,208	179	390	87	270	26	790
Wisconsin.....	624	68	90	160	244	65	129	5	549

Number of Cases of Certain Communicable Diseases Reported for the Month of July, 1928, by State Health Officers—Continued

State	Chick- en pox	Diph- theria	Meas- les	Mumps	Scarlet fever	Small- pox	Tuber- culosis	Ty- phoid fever	Whoop- ing cough
Minnesota	131	92	57	---	191	5	171	4	192
Iowa	66	18	25	53	60	64	162	11	58
Missouri	46	87	221	65	127	65	199	52	231
North Dakota	38	10	34	7	87	5	16	6	34
South Dakota	12	4	112	3	58	16	3	3	18
Nebraska	14	13	10	17	47	57	126	5	42
Kansas	54	16	69	174	107	124	147	65	333
Delaware	2	1	49	8	5	0	17	6	5
Maryland	63	48	197	74	55	6	314	61	498
District of Columbia	8	69	158	---	42	0	112	2	41
Virginia	125	43	572	---	69	13	1199	196	290
West Virginia	31	21	63	---	53	33	29	52	30
North Carolina	68	62	270	---	64	58	---	286	461
South Carolina	43	78	87	---	15	31	135	433	276
Georgia	23	16	90	24	23	7	72	246	100
Florida	2	33	95	6	10	14	96	41	45
Kentucky ¹	---	---	---	---	---	---	---	---	---
Tennessee	22	28	103	55	34	41	175	267	69
Alabama	21	37	188	26	90	26	386	246	118
Mississippi	200	43	374	341	26	2	289	343	954
Arkansas	36	11	88	10	8	19	167	138	82
Louisiana	4	32	121	2	14	6	128	134	41
Oklahoma ²	2	35	36	19	21	103	58	166	77
Texas ²	---	---	---	---	---	---	---	---	---
Montana	15	6	71	17	13	64	22	17	11
Idaho	11	10	5	---	15	32	---	2	5
Wyoming	13	8	2	5	19	3	11	5	12
Colorado ³	---	---	---	---	---	---	---	---	---
New Mexico ²	---	---	---	---	---	---	---	---	---
Arizona	---	3	54	2	4	4	50	6	3
Utah ²	---	---	---	---	---	---	---	---	---
Nevada ²	---	---	---	---	---	---	---	---	---
Washington	176	27	62	76	52	82	157	18	62
Oregon	36	32	67	23	20	119	60	14	6
California	364	229	87	234	268	69	879	60	780

Case Rates per 1,000 Population (Annual Basis) for the Month of July, 1928

State	Chick- en pox	Diph- theria	Meas- les	Mumps	Scarlet fever	Small- pox	Tuber- culosis	Ty- phoid fever	Whoop- ing cough
Maine	0.55	0.12	3.40	0.70	0.48	0	0.52	0.15	1.41
New Hampshire	---	.23	---	---	.31	0	---	0	---
Vermont	1.57	.07	5.93	1.11	.47	0	.37	.03	4.22
Massachusetts	.80	.45	3.72	.45	.87	0	1.46	.06	1.11
Rhode Island	.20	.48	10.45	1.16	.68	0	.68	.03	.35
Connecticut	.38	.36	5.65	.78	.47	0	1.08	.03	2.52
New York	.72	.86	3.89	.42	.45	0.01	1.33	.11	1.45
New Jersey	.55	1.07	3.88	---	.40	.01	1.25	.11	1.91
Pennsylvania	.43	.47	3.22	.63	.47	.04	.85	.19	1.65
Ohio	.56	.26	2.42	.23	.39	.09	1.33	.19	1.74
Indiana	.16	.17	.92	.07	.38	.39	.61	.07	.30
Illinois	.88	.50	.60	.40	.65	.14	1.65	.14	1.47
Michigan	.57	.54	3.11	.46	1.00	.22	.69	.07	2.03
Wisconsin	1.30	.27	.36	.64	.98	.26	.52	.02	2.19
Minnesota	.57	.40	.25	---	.83	.02	.74	.02	.83
Iowa	.32	.00	.12	.26	.24	.31	.79	.05	.28
Missouri	.15	.29	.74	.22	.43	.22	.67	.17	.77
North Dakota	.70	.18	.63	.13	1.60	.09	.29	.11	.63
South Dakota	.20	.07	1.89	.05	.97	.27	.05	.05	.30
Nebraska	.12	.11	.08	.14	.30	.48	1.22	.04	.35
Kansas	.35	.10	.44	1.12	.60	.80	.95	.42	2.14
Delaware	.10	.05	2.37	.39	.24	0	1.34	.29	.24
Maryland	.46	.35	1.44	.54	.40	.04	2.20	.05	3.64
District of Columbia	.17	1.48	3.38	---	.90	0	2.40	.04	.88
Virginia	.57	.19	2.62	---	.32	.06	1.64	.90	1.33
West Virginia	.21	.14	.47	---	.36	.23	.20	.36	.21
North Carolina	.27	.25	1.09	---	.26	.23	---	1.66	1.85

¹ Pulmonary.

² Reports received weekly.

³ Exclusive of Oklahoma City and Tulsa.

⁴ Reports not received at time of going to press.

⁵ Reports received annually.

**Case Rates per 1,000 Population (Annual Basis) for the Month of July, 1928—
Continued**

State	Chick- en pox	Diph- theria	Mea- sles	Mumps	Scarlet fever	Small- pox	Tuber- culosis	Ty- phoid fever	Whoop- ing cough
South Carolina.....	0.27	0.49	0.55	0.03	0.10	0.20	0.86	2.74	1.75
Georgia.....	.08	.06	.33	.09	.08	.03	.27	.91	.37
Florida.....	.02	.28	.79	.05	.08	.12	.80	.34	.38
Kentucky ¹									
Tennessee.....	.10	.13	.49	.26	.16	.19	.83	1.26	.33
Alabama.....	.10	.17	.86	.12	.09	.13	1.77	1.13	.54
Mississippi.....	1.32	.28	2.47	2.25	.17	.01	1.91	2.26	6.29
Arkansas.....	.22	.07	.53	.06	.05	.12	1.41	.84	.50
Louisiana.....	.02	.19	.73	.01	.08	.04	1.78	.81	.25
Oklahoma ²01	.19	.20	.10	.12	.57	.32	.91	.42
Texas ³									
Montana.....	.32	.13	1.53	.37	.28	1.38	.47	.37	.24
Idaho.....	.24	.22	.11		.32	.69		.04	.11
Wyoming.....	.62	.38	.10	.24	.91	.14	1.05	.24	.57
Colorado ⁴									
New Mexico ⁵									
Arizona.....		.07	1.35	.05	.10	.10	1.25	.15	.07
Utah ²									
Nevada ⁵									
Washington.....	1.31	.20	.46	.57	.39	.61	1.17	.13	.46
Oregon.....	.47	.42	.88	.37	.26	1.56	.79	.18	.08
California.....	.94	.59	.23	.61	.69	.18	2.28	.16	2.02

¹ Pulmonary.² Reports received weekly.³ Exclusive of Oklahoma City and Tulsa.⁴ Reports not received at time of going to press.⁵ Reports received annually.

**GENERAL CURRENT SUMMARY AND WEEKLY REPORTS FROM
CITIES**

The 99 cities reporting cases used in the following table are situated in all parts of the country and have an estimated aggregate population of more than 31,605,000. The estimated population of the 93 cities reporting deaths is more than 30,910,000. The estimated expectancy is based on the experience of the last nine years, excluding epidemics.

Weeks ended September 22, 1928, and September 24, 1927

	1928	1927	Estimated expectancy
<i>Cases reported</i>			
Diphtheria:			
43 States.....	1,177	1,526	
99 cities.....	476	612	713
Measles:			
42 States.....	443	593	
99 cities.....	106	162	
Poliomyelitis:			
45 States.....	305	681	
Scarlet fever:			
43 States.....	1,115	1,332	
99 cities.....	380	396	431
Smallpox:			
43 States.....	113	167	
99 cities.....	7	34	13
Typhoid fever:			
43 States.....	866	1,041	
99 cities.....	161	164	191
<i>Deaths reported</i>			
Influenza and pneumonia:			
93 cities.....	412	358	
Smallpox:			
93 cities.....	1	0	
South Bend, Ind.....	1	0	

City reports for week ended September 22, 1928

The "estimated expectancy" given for diphtheria, poliomyelitis, scarlet fever, smallpox, and typhoid fever is the result of an attempt to ascertain from previous occurrence the number of cases of the disease under consideration that may be expected to occur during a certain week in the absence of epidemics. It is based on reports to the Public Health Service during the past nine years. It is in most instances the median number of cases reported in the corresponding weeks of the preceding years. When the reports include several epidemics or when for other reasons the median is unsatisfactory, the epidemic periods are excluded and the estimated expectancy is the mean number of cases reported for the week during non-epidemic years.

If the reports have not been received for the full nine years, data are used for as many years as possible, but no year earlier than 1919 is included. In obtaining the estimated expectancy, the figures are smoothed when necessary to avoid abrupt deviation from the usual trend. For some of the diseases given in the table the available data were not sufficient to make it practicable to compute the estimated expectancy.

Division, State, and city	Population July 1, 1926, estimated	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases reported	Mumps, cases reported	Pneumonia, deaths reported
			Cases, estimated expectancy	Cases reported	Cases reported	Deaths reported			
NEW ENGLAND									
Maine:									
Portland	76,400	3	1	0	0	0	0	0	0
New Hampshire:									
Concord	122,546	0	0	0	0	0	0	0	0
Vermont:									
Barre	110,008	1	0	0	0	0	0	0	0
Massachusetts:									
Boston	787,000	3	28	11	5	1	2	0	15
Fall River	131,000	0	2	1	1	0	10	0	1
Springfield	145,000	0	2	3	0	0	0	1	2
Worcester	193,000	0	4	4	1	0	1	1	1
Rhode Island:									
Pawtucket	71,000	0	1	0	0	0	0	0	3
Providence	275,000	0	5	3	0	0	5	0	4
Connecticut:									
Bridgeport	(?)	0	5	4	0	0	3	0	3
Hartford	164,000	1	3	2	0	0	0	1	3
New Haven	182,000	0	2	1	0	0	0	0	1
MIDDLE ATLANTIC									
New York:									
Buffalo	544,000	0	13	5	0	0	0	1	5
New York	5,924,000	11	100	68	8	4	12	10	92
Rochester	321,000	3	5	1	0	0	1	1	3
Syracuse	185,000	2	3	0	0	0	0	1	5
New Jersey:									
Camden	131,000	1	3	2	0	0	0	1	2
Newark	459,000	3	8	14	0	0	1	5	5
Trenton	134,000	0	2	0	0	0	0	0	2
Pennsylvania:									
Philadelphia	2,008,000	5	41	25	0	4	18	4	21
Pittsburgh	637,000	3	17	11	0	2	0	2	16
Reading	114,000	2	2	2	0	0	0	0	0
EAST NORTH CENTRAL									
Ohio:									
Cincinnati	411,000	2	8	3	0	0	1	0	7
Cleveland	960,000	5	34	22	3	2	6	4	9
Columbus	285,000	0	4	0	1	0	0	0	3
Toledo	295,000	2	10	0	0	0	1	0	4
Indiana:									
Fort Wayne	99,900	0	2	1	0	0	0	0	1
Indianapolis	367,000	0	8	4	0	0	0	3	8
South Bend	81,700	1	1	0	0	0	0	0	2
Terre Haute	71,900	0	0	0	0	0	0	0	1
Illinois:									
Chicago	3,048,000	15	53	77	5	2	15	8	33
Springfield	64,700	0	1	0	1	1	1	1	2
Michigan:									
Detroit	1,242,044	11	47	25	2	1	2	6	14
Flint	136,000	3	7	2	0	0	1	1	2
Grand Rapids	156,000	3	3	1	0	0	1	0	2

¹ Estimated, July 1, 1925.

² No estimate made.

³ Special census.

City reports for week ended September 22, 1928—Continued

Division, State, and city	Population July 1, 1926, estimated	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases reported	Mumps, cases reported	Pneumonia, deaths reported
			Cases, estimated expectancy	Cases reported	Cases reported	Deaths reported			
EAST NORTH CENTRAL—continued									
Wisconsin:									
Kenosha.....	52,700	2	1	1	0	0	0	2	0
Milwaukee.....	517,000	5	11	5	0	0	4	3	5
Racine.....	69,400	1	1	0	0	0	0	1	1
Superior.....	139,671	0	0	0	0	0	0	0	0
WEST NORTH CENTRAL									
Minnesota:									
Duluth.....	113,000	4	1	0	0	0	0	0	4
Minneapolis.....	434,000	11	20	14	0	0	6	6	4
St. Paul.....	248,000	4	15	8	0	0	0	0	2
Iowa:									
Davenport.....	152,469	0	1	0	0	0	0	0	0
Des Moines.....	146,000	0	4	1	0	0	0	0	0
Sioux City.....	78,000	0	2	0	0	0	1	0	0
Waterloo.....	36,900	0	1	1	0	0	0	5	0
Missouri:									
Kansas City.....	375,000	5	5	1	0	0	0	2	3
St. Joseph.....	78,400	1	1	0	0	1	0	2	1
St. Louis.....	830,000	5	27	18	0	0	1	1	0
North Dakota:									
Fargo.....	126,403	0	0	0	0	0	0	0	0
Grand Forks.....	114,811	0	0	0	0	0	0	0	0
South Dakota:									
Aberdeen.....	115,036	0	0	0	0	0	0	0	0
Sioux Falls.....	130,127	0	0	0	0	0	0	0	0
Nebraska:									
Lincoln.....	62,000	1	1	0	0	0	0	1	0
Omaha.....	216,000	0	12	4	0	0	0	1	4
Kansas:									
Topeka.....	56,500	0	2	0	2	0	0	2	1
Wichita.....	92,500	1	2	1	0	0	1	0	1
SOUTH ATLANTIC									
Delaware:									
Wilmington.....	124,000	0	1	0	0	0	2	0	3
Maryland:									
Baltimore.....	808,000	2	19	8	4	0	2	2	17
Cumberland.....	133,741	1	0	0	0	0	0	0	1
Frederick.....	112,035	0	0	0	0	0	0	0	0
District of Columbia:									
Washington.....	528,000	0	9	12	1	1	3	0	9
Virginia:									
Lynchburg.....	138,493	0	2	3	0	0	0	1	0
Norfolk.....	174,000	1	2	0	1	0	0	0	4
Richmond.....	189,000	0	16	11	0	0	1	5	0
Roanoke.....	61,900	0	4	6	0	0	0	0	0
West Virginia:									
Charleston.....	50,700	0	1	1	0	1	0	0	3
Wheeling.....	156,208	0	1	0	0	0	1	3	4
North Carolina:									
Raleigh.....	130,371	0	5	0	0	0	0	0	1
Wilmington.....	37,700	1	0	0	0	0	0	0	0
Winston-Salem.....	71,800	0	3	0	0	0	0	0	0
South Carolina:									
Charleston.....	74,100	0	0	1	4	0	0	0	2
Columbia.....	41,800	0	2	1	0	0	0	0	0
Greenville.....	127,311	1							
Georgia:									
Atlanta.....	(²)	0	7	3	9	0	0	0	2
Brunswick.....	116,809	0	0						
Savannah.....	94,900	0	1	1	0	0	0	1	2
Florida:									
Miami.....	131,286	0	3	0	2	0	0	0	1
St. Petersburg.....	47,629	0	0						0
Tampa.....	102,000	0	1	1	0	0	0	0	0

¹ Estimated, July 1, 1925.² No estimate made.³ Special census.

City reports for week ended September 22, 1928—Continued

Division, State, and city	Population July 1, 1928, estimated	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases reported	Mumps, cases reported	Pneumonia, deaths reported
			Cases, estimated expectancy	Cases reported	Cases reported	Deaths reported			
EAST SOUTH CENTRAL									
Kentucky:									
Covington.....	58,500	0	1	0	0	0	0	0	1
Louisville.....	311,000	0	5	6	0	0	0	2	0
Tennessee:									
Memphis.....	177,000	2	4	5	0	1	0	0	4
Nashville.....	137,000	0	5	8	0	1	0	0	2
Alabama:									
Birmingham.....	211,000	0	6	4	1	0	1	1	2
Mobile.....	66,800	0	1	3	0	0	0	0	0
Montgomery.....	47,000	0	2	5	0	0	0	0	0
WEST SOUTH CENTRAL									
Arkansas:									
Fort Smith.....	¹ 31,643	0	1	2	0	0	0	0	0
Little Rock.....	75,900	0	1	0	0	0	0	2	0
Louisiana:									
New Orleans.....	419,000	0	7	4	2	0	0	0	0
Shreveport.....	59,500	0	1	0	0	0	0	0	0
Oklahoma:									
Tulsa.....	133,000	0	1	1	0	0	0	0	0
Texas:									
Dallas.....	203,000	1	7	10	0	0	1	0	0
Fort Worth.....	159,000	0	3	2	0	0	0	0	0
Galveston.....	49,100	0	0	0	0	0	0	0	0
Houston.....	¹ 164,954	0	3	6	0	0	0	0	1
San Antonio.....	205,000	0	2	1	0	1	0	0	2
MOUNTAIN									
Montana:									
Billings.....	¹ 17,971	0	0	0	0	0	0	0	0
Great Falls.....	¹ 29,883	0	0	0	0	0	0	0	0
Helena.....	¹ 12,037	0	0	1	0	0	0	0	0
Missoula.....	¹ 12,668	1	0	0	0	0	0	0	0
Idaho:									
Boise.....	¹ 23,042	0	0	0	0	0	0	0	0
Colorado:									
Denver.....	285,000	0	17	3	0	0	0	5	7
Pueblo.....	43,900	0	2	1	0	0	0	2	0
New Mexico:									
Albuquerque.....	¹ 21,000	0	0	0	0	0	0	0	0
Utah:									
Salt Lake City.....	133,000	8	4	2	0	0	0	5	1
Nevada:									
Reno.....	¹ 12,665	0	0	0	0	0	0	0	0
PACIFIC									
Washington:									
Seattle.....	(?)	5	4	1	0	0	2	0	0
Spokane.....	109,000	3	2	0	0	0	1	0	0
Tacoma.....	106,000	0	3	0	0	0	0	4	1
Oregon:									
Portland.....	¹ 282,383	0	6	7	1	1	2	3	1
California:									
Los Angeles.....	(?)	5	20	15	7	0	1	5	16
Sacramento.....	73,400	0	2	0	0	0	0	9	1
San Francisco.....	567,000	7	14	5	7	0	0	5	9

¹ Estimated, July 1, 1925.

² No estimate made.

City reports for week ended September 22, 1928—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuber- culosis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, esti- mated expect- ancy	Cases re- ported	Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		
NEW ENGLAND											
Maine:											
Portland.....	0	2	0	0	0	0	1	2	0	1	12
New Hampshire:											
Concord.....	0	0	0	0	0	0	0	0	0	0	8
Vermont:											
Barre.....	0	1	0	0	0	1	0	0	0	0	4
Massachusetts:											
Boston.....	20	16	0	0	0	10	4	3	0	13	190
Fall River.....	1	4	0	0	0	6	1	1	0	3	29
Springfield.....	2	3	0	0	0	5	0	0	0	0	29
Worcester.....	4	7	0	0	0	1	0	0	0	8	45
Rhode Island:											
Pawtucket.....	1	1	0	0	0	1	0	0	0	0	18
Providence.....	2	5	0	0	0	5	0	2	1	4	64
Connecticut:											
Bridgeport.....	2	2	0	0	0	0	0	0	0	3	34
Hartford.....	2	1	0	0	0	2	1	0	0	4	23
New Haven.....	2	2	0	0	0	1	2	1	0	5	21
MIDDLE ATLANTIC											
New York:											
Buffalo.....	8	7	0	0	0	11	2	2	0	13	134
New York.....	40	23	0	0	0	76	41	24	2	53	1,194
Rochester.....	3	0	0	0	0	0	1	0	0	6	48
Syracuse.....	4	1	0	0	0	1	2	0	0	16	55
New Jersey:											
Camden.....	2	0	0	0	0	0	1	0	0	1	19
Newark.....	5	1	0	0	0	8	2	0	0	28	78
Trenton.....	0	2	0	0	0	2	1	13	0	1	30
Pennsylvania:											
Philadelphia.....	27	9	0	0	0	23	13	14	2	70	438
Pittsburgh.....	13	7	1	0	0	9	3	2	0	28	144
Reading.....	0	0	0	0	0	1	1	3	0	15	12
EAST NORTH CENTRAL											
Ohio:											
Cincinnati.....	6	8	0	0	0	13	2	1	0	6	132
Cleveland.....	14	13	0	0	0	11	3	2	0	58	171
Columbus.....	4	1	0	0	0	6	0	0	0	4	71
Toledo.....	6	4	0	0	0	6	2	6	1	22	62
Indiana:											
Fort Wayne.....	1	0	0	0	0	1	1	0	0	0	26
Indianapolis.....	5	9	1	0	0	5	2	4	0	4	107
South Bend.....	2	1	0	0	1	1	0	1	0	0	14
Terre Haute.....	0	0	0	0	0	2	0	0	0	0	39
Illinois:											
Chicago.....	40	41	0	0	0	48	8	10	0	52	594
Springfield.....	1	1	0	0	0	1	1	0	0	0	26
Michigan:											
Detroit.....	33	35	1	0	0	26	6	4	0	163	250
Flint.....	7	3	0	1	0	0	1	2	0	10	39
Grand Rapids.....	4	1	0	0	0	0	0	0	0	7	33
Wisconsin:											
Kenosha.....	1	4	0	0	0	0	0	0	0	0	6
Milwaukee.....	13	19	0	1	0	6	1	1	0	61	94
Racine.....	3	0	0	0	0	1	0	0	0	18	9
Superior.....	1	3	0	0	0	0	0	0	0	0	4
WEST NORTH CENTRAL											
Minnesota:											
Duluth.....	4	4	0	0	0	2	0	0	0	2	22
Minneapolis.....	22	13	0	0	0	2	1	2	0	10	85
St. Paul.....	8	7	2	0	0	2	2	2	2	25	42

12 cases nonresident.

City reports for week ended September 22, 1928—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuberculosis, deaths reported	Typhoid fever			Whooping cough, cases reported	Deaths, all causes
	Cases, estimated expectancy	Cases reported	Cases, estimated expectancy	Cases reported	Deaths reported		Cases, estimated expectancy	Cases reported	Deaths reported		
WEST NORTH CENTRAL—continued											
Iowa:											
Davenport	0	1	0	0		0	0		0		
Des Moines	4	2	0	0		0	0				22
Sioux City	1	0	0	0		0	0				2
Waterloo	1	1	0	0		1	0		1		
Missouri:											
Kansas City	4	4	0	1	0	5	2	1	1	2	66
St. Joseph	1	0	1	0	0	1	0	1	0	1	26
St. Louis	13	7	0	0	0	6	6	7	2	19	167
North Dakota:											
Fargo	1	9	0	0	0	1	0	0	0	0	6
Grand Forks	1	0	0	0			0	0	0	0	
South Dakota:											
Aberdeen	0	0	0	0			0	0	0	3	
Sioux Falls	1	0	0	0			1	0	0	0	6
Nebraska:											
Lincoln	0	0	0	0	0	0	0	0	0	1	12
Omaha	2	1	0	1	0	1	0	1	0	2	46
Kansas:											
Topeka	1	1	0	0	0	0	0	0	0	0	14
Wichita	2	6	1	0	0	0	2	2	0	7	34
SOUTH ATLANTIC											
Delaware:											
Wilmington	1	0	0	0	0	1	1	1	0	1	25
Maryland:											
Baltimore	7	5	0	0	0	13	10	3	2	103	186
Cumberland	0	0	0	0	0	0	1	0	0	2	15
Frederick	0	0	0	0	0	0	0	1	0	0	2
District of Columbia:											
Washington	7	4	0	0	0	9	4	4	1	10	114
Virginia:											
Lynchburg	0	0	0	0	0	0	0	0	0	0	5
Norfolk	1	1	0	0	0	1	1	0	0	0	
Richmond	5	5	0	0	0	4	2	0	0	2	41
Roanoke	1	8	0	0	0	0	1	0	0	0	18
West Virginia:											
Charleston	2	2	0	0	0	1	1	0	0	0	15
Wheeling	3	2	0	0	0	0	0	1	0	0	18
North Carolina:											
Raleigh	1	1	0	0	0	3	0	0	0	0	15
Wilmington	2	0	0	0	0	1	1	0	0	0	8
Winston-Salem	2	3	0	0	0	2	1	0	0	0	17
South Carolina:											
Charleston	0	2	0	0	0	1	3	3	0	0	18
Columbia	0	0	0	0	0	0	1	2	1	0	17
Greenville	0		0				1				
Georgia:											
Atlanta	5	5	0	0	0	4	4			1	73
Brunswick	0	0	0	0	0	0	0	0	0	0	
Savannah	1	0	0	0	0	2	0	0	0	0	29
Florida:											
Miami	0	0	0	0	0	1	1	1	0	0	13
St. Petersburg	0	0	0	0	0	0	0	0	0	0	4
Tampa	0	0	0	0	0	0	0	2	0	0	15
EAST SOUTH CENTRAL											
Kentucky:											
Covington	1	1	0	0	0	1	0	1	0	0	29
Louisville	3	5	0	0	0	2	5	3	2	5	65
Tennessee:											
Memphis	2	2	0	0	0	3	4	1	3	2	64
Nashville	2	0	0	0	0	10	5	8	2	4	53
Alabama:											
Birmingham	4	4	1	0	0	5	4	6	0	0	66
Mobile	0	1	0	0	0	1	0	0	0	0	16
Montgomery	1	0	0	0	0		0	0	0	0	

City reports for week ended September 22, 1928—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuber- culosis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, estimated expectancy	Cases reported	Cases, estimated expectancy	Cases reported	Deaths reported		Cases, estimated expectancy	Cases reported	Deaths reported		
WEST SOUTH CENTRAL											
Arkansas:											
Fort Smith.....	0	1	0	0	0	1	0	0	0	0	
Little Rock.....	0	0	0	0	0	3	2	1	0	0	
Louisiana:											
New Orleans.....	2	1	0	0	0	19	4	5	0	0	147
Shreveport.....	0	0	0	0	0	0	1	0	0	0	19
Oklahoma:											
Tulsa.....	2	2	0	0	0	0	1	2	0	0	
Texas:											
Dallas.....	3	4	0	0	0	1	2	10	1	1	53
Fort Worth.....	2	1	0	0	0	2	2	1	0	0	33
Galveston.....	0	0	0	0	0	2	0	0	0	0	7
Houston.....	0	1	0	0	0	2	0	1	1	0	47
San Antonio.....	0	0	0	1	0	4	1	0	0	0	51
MOUNTAIN											
Montana:											
Billings.....	1	0	0	0	0	0	0	0	0	0	3
Great Falls.....	1	0	0	0	0	0	0	0	0	0	5
Helena.....	0	1	0	0	0	0	0	0	0	0	5
Missoula.....	1	0	1	0	0	0	0	0	0	4	2
Idaho:											
Boise.....	0	0	0	0	0	0	0	0	0	0	6
Colorado:											
Denver.....	5	1	0	0	0	5	3	2	2	5	88
Pueblo.....	1	0	0	0	0	0	2	0	1	0	10
New Mexico:											
Albuquerque.....	0	0	0	0	0	1	2	1	0	0	6
Utah:											
Salt Lake City.....	1	3	0	0	0	1	3	1	1	3	26
Nevada:											
Reno.....	0	1	0	0	0	0	0	0	0	0	6
PACIFIC											
Washington:											
Seattle.....	6	1	0	1	0	1	0	0	0	6	
Spokane.....	4	2	1	0	0	0	0	0	0	0	
Tacoma.....	1	0	1	0	0	2	0	0	0	4	23
Oregon:											
Portland.....	5	6	4	6	0	1	1	1	0	0	59
California:											
Los Angeles.....	10	9	2	0	0	19	4	5	2	47	211
Sacramento.....	1	9	1	0	0	4	1	0	0	0	31
San Francisco.....	6	9	0	1	0	11	2	2	0	11	159

Division, State, and city	Meningococcus meningitis		Lethargic encephalitis		Pellagra		Poliomyelitis (Infantile paralysis)		Deaths
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases, estimated expectancy	Cases	
NEW ENGLAND									
Vermont:									
Barre.....	0	0	0	0	0	0	0	1	0
Massachusetts:									
Boston.....	1	0	0	0	0	0	3	9	1
Fall River.....	0	0	0	0	0	0	1	1	0
Springfield.....	0	0	0	0	0	0	1	1	0
Rhode Island:									
Providence.....	2	0	1	1	0	0	1	0	0
Connecticut:									
Hartford.....	0	0	1	0	0	0	0	0	0
New Haven.....	1	0	0	0	0	0	1	0	0

City reports for week ended September 22, 1928—Continued

Division, State, and city	Meningococcus meningitis		Lethargic encephalitis		Pellagra		Poliomyelitis (infantile paralysis)		
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases, estimated expectancy	Cases	Deaths
MIDDLE ATLANTIC									
New York:									
Buffalo.....	0	0	0	0	0	0	0	3	1
New York City.....	23	10	5	1	0	0	16	31	4
Rochester.....	0	0	0	0	0	0	1	5	0
Syracuse.....	0	0	0	0	0	0	1	1	0
New Jersey:									
Newark.....	0	0	0	0	0	0	1	2	0
Trenton.....	0	0	0	1	0	0	0	0	0
Pennsylvania:									
Philadelphia.....	1	0	1	1	0	1	2	2	0
Pittsburgh.....	0	0	0	0	0	0	1	1	0
EAST NORTH CENTRAL									
Ohio:									
Cleveland.....	3	1	0	0	0	0	1	9	2
Columbus.....	0	0	0	0	0	0	0	2	0
Toledo.....	0	0	0	0	0	0	0	1	0
Indiana:									
Indianapolis.....	1	0	0	0	0	0	0	0	0
Illinois:									
Chicago.....	1	2	0	0	0	0	6	0	0
Springfield.....	0	0	0	0	0	0	0	1	0
Michigan:									
Detroit.....	6	3	0	1	0	0	2	1	1
Wisconsin:									
Milwaukee.....	1	0	0	0	0	0	1	0	0
WEST NORTH CENTRAL¹									
Minnesota:									
Duluth.....	1	0	0	0	0	0	0	0	0
Minneapolis.....	0	0	0	0	0	0	0	4	0
St. Paul.....	0	0	0	0	0	0	1	1	0
Missouri:									
Kansas City.....	0	0	0	0	0	0	0	2	0
St. Louis.....	3	0	1	0	0	0	1	1	0
SOUTH ATLANTIC²									
Delaware:									
Wilmington.....	1	0	0	0	0	0	0	0	0
Maryland:									
Baltimore.....	0	0	0	0	0	0	2	2	0
District of Columbia:									
Washington.....	0	0	0	0	0	0	0	2	0
Virginia:									
Richmond.....	0	0	0	1	0	1	1	0	0
North Carolina:									
Winston-Salem.....	0	0	0	0	1	0	0	0	0
Georgia: ³									
Atlanta.....	0	0	0	0	0	1	0	0	0
Florida:									
Tampa ³	0	0	0	0	0	1	0	0	0
EAST SOUTH CENTRAL									
Tennessee:									
Nashville.....	1	0	0	0	1	0	0	1	0
Alabama:									
Birmingham.....	0	0	0	0	1	1	0	0	0

¹ Rabies (in man): 1 death at Omaha, Nebr.

² Dengue: 12 cases at Charleston, S. C.

³ Typhus fever: 3 cases; 1 case at Savannah, Ga., 1 case at Tampa, Fla., and 1 case at Houston, Tex.

City reports for week ended September 22, 1928—Continued

Division, State, and city	Meningococcus meningitis		Lethargic encephalitis		Pellagra		Poliomyelitis (Infantile paralysis)		
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases, estimated expectancy	Cases	Deaths
WEST SOUTH CENTRAL									
Arkansas:									
Little Rock.....	0	0	0	0	0	2	0	0	0
Louisiana:									
New Orleans.....	1	0	0	0	7	1	1	0	0
Texas:									
Dallas.....	0	0	0	0	1	1	1	0	0
Fort Worth.....	0	0	0	0	0	1	0	0	0
Houston ¹	0	0	0	0	0	1	0	0	0
San Antonio.....	0	0	0	0	0	0	0	2	0
MOUNTAIN									
Colorado:									
Denver.....	2	1	0	0	0	0	0	0	0
Pueblo.....	0	1	0	0	0	0	0	2	0
Nevada:									
Reno.....	0	0	0	1	0	0	0	0	0
PACIFIC									
Washington:									
Seattle.....	0	0	0	0	0	0	1	5	0
Spokane.....	0	0	0	0	0	0	0	3	0
Tacoma.....	0	0	0	0	0	0	1	2	0
Oregon:									
Portland.....	0	1	0	0	0	0	0	0	0
California:									
Los Angeles.....	0	0	1	0	1	0	1	1	1
San Francisco.....	0	0	0	0	0	1	1	0	0

¹ Typhus fever: 3 cases; 1 case at Savannah, Ga., 1 case at Tampa, Fla., and 1 case at Houston, Tex.

The following table gives the rates per 100,000 population for 101 cities for the 5-week period ended September 22, 1928, compared with those for a like period ended September 24, 1927. The population figures used in computing the rates are approximate estimates as of July 1, 1928 and 1927, respectively, authoritative figures for many of the cities not being available. The 101 cities reporting cases had estimated aggregate populations of approximately 31,657,000 in 1928 and 31,050,000 in 1927. The 95 cities reporting deaths had nearly 30,961,000 estimated population in 1928 and nearly 30,370,000 in 1927. The number of cities included in each group and the estimated aggregate populations are shown in a separate table below.

Summary of weekly reports from cities, August 19 to September 22, 1928—Annual rates per 100,000 population compared with rates for the corresponding period of 1927¹

DIPHTHERIA CASE RATES

	Week ended—									
	Aug. 25, 1928	Aug. 27, 1927	Sept. 1, 1928	Sept. 3, 1927	Sept. 5, 1928	Sept. 10, 1927	Sept. 15, 1928	Sept. 17, 1927	Sept. 22, 1928	Sept. 24, 1927
101 cities.....	64	81	56	84	51	94	75	101	79	103
New England.....	62	86	37	88	34	93	87	53	67	91
Middle Atlantic.....	66	78	58	77	49	90	57	105	62	95
East North Central.....	67	81	61	87	51	90	67	82	92	105
West North Central.....	64	53	51	69	70	63	97	125	92	87
South Atlantic.....	79	88	67	89	47	108	107	112	86	105
East South Central.....	35	61	40	51	30	106	125	117	160	81
West South Central.....	64	95	100	161	76	149	140	136	92	203
Mountain.....	44	134	44	117	53	152	35	224	62	233
Pacific.....	41	94	20	73	49	91	49	91	54	76

MEASLES CASE RATES

101 cities.....	28	25	21	21	19	20	18	20	18	27
New England.....	85	58	90	58	55	63	39	30	48	40
Middle Atlantic.....	21	24	16	18	18	16	15	14	15	30
East North Central.....	31	13	28	11	24	15	24	18	20	18
West North Central.....	16	16	4	16	2	10	14	28	18	20
South Atlantic.....	33	31	4	18	5	14	11	14	16	36
East South Central.....	10	25	10	10	0	10	10	10	5	15
West South Central.....	0	17	0	41	4	17	0	17	4	0
Mountain.....	9	27	18	9	35	36	44	45	0	45
Pacific.....	31	52	13	42	28	34	13	44	10	52

SCARLET FEVER CASE RATES

101 cities.....	33	54	32	57	37	52	58	69	63	67
New England.....	30	81	64	60	46	53	78	102	101	123
Middle Atlantic.....	18	37	14	38	18	30	28	46	24	42
East North Central.....	44	61	32	81	44	65	88	89	91	69
West North Central.....	49	61	55	69	39	91	68	87	103	59
South Atlantic.....	32	63	30	60	49	60	54	78	68	106
East South Central.....	45	56	44	76	60	96	100	46	65	46
West South Central.....	52	58	44	58	56	45	44	41	28	50
Mountain.....	62	63	35	63	27	54	27	99	53	162
Pacific.....	33	37	31	34	59	31	64	55	77	71

SMALLPOX CASE RATES

101 cities.....	2	5	0	4	1	4	1	5	1	6
New England.....	0	0	0	0	0	0	0	0	0	0
Middle Atlantic.....	0	0	0	0	0	0	0	0	0	0
East North Central.....	5	6	1	7	1	4	2	4	1	1
West North Central.....	0	4	0	2	4	12	4	22	4	8
South Atlantic.....	0	0	0	0	0	2	0	4	0	0
East South Central.....	0	25	0	0	0	10	0	0	0	10
West South Central.....	0	0	0	0	0	4	4	4	4	0
Mountain.....	9	27	0	36	9	9	9	27	0	161
Pacific.....	0	31	5	18	8	13	3	37	5	21

¹ The figures given in this table are rates per 100,000 population, annual basis, and not the number of cases reported. Populations used are estimated as of July 1, 1928 and 1927, respectively.

² South Bend, Ind., not included.

³ Greenville, S. C., not included.

⁴ Lynchburg, Va., and Savannah, Ga., not included.

⁵ Greenville, S. C., and Brunswick, Ga., not included.

Summary of weekly reports from cities, August 19 to September 22, 1928—Annual rates per 100,000 population compared with rates for the corresponding period of 1927—Continued

TYPHOID FEVER CASE RATES

	Week ended—									
	Aug. 25, 1928	Aug. 27, 1927	Sept. 1, 1928	Sept. 3, 1927	Sept. 8, 1928	Sept. 10, 1927	Sept. 15, 1928	Sept. 17, 1927	Sept. 22, 1928	Sept. 24, 1927
101 cities.....	31	31	29	32	24	30	28	33	27	28
New England.....	16	33	23	21	16	40	14	47	21	63
Middle Atlantic.....	23	21	18	28	25	27	29	37	23	24
East North Central.....	18	11	15	15	13	7	14	16	16	10
West North Central.....	25	20	39	10	19	32	25	24	31	14
South Atlantic.....	51	68	44	71	33	58	42	31	30	45
East South Central.....	165	203	135	183	80	112	100	152	95	88
West South Central.....	52	74	72	54	28	74	28	37	68	79
Mountain.....	62	45	44	54	80	63	18	36	27	36
Pacific.....	26	21	26	8	13	8	38	16	18	13

INFLUENZA DEATH RATES

95 cities.....	4	5	3	4	3	4	5	5	4	3
New England.....	2	2	0	2	0	5	0	0	2	0
Middle Atlantic.....	3	2	3	3	2	3	4	4	5	2
East North Central.....	3	3	3	5	2	4	5	2	4	1
West North Central.....	0	2	2	4	2	0	10	4	2	2
South Atlantic.....	9	11	4	7	9	5	7	9	4	11
East South Central.....	0	16	5	5	16	11	16	0	10	11
West South Central.....	16	21	4	13	8	13	8	17	4	8
Mountain.....	0	9	18	0	18	9	0	9	0	0
Pacific.....	3	7	3	0	7	7	3	10	0	0

PNEUMONIA DEATH RATES

95 cities.....	56	46	55	56	57	62	63	60	66	59
New England.....	44	51	32	49	48	65	62	40	76	70
Middle Atlantic.....	68	55	60	72	56	66	69	60	74	69
East North Central.....	41	34	50	51	60	59	64	53	59	44
West North Central.....	35	31	31	23	22	43	43	46	41	25
South Atlantic.....	60	36	72	42	78	49	65	76	86	65
East South Central.....	84	69	105	48	78	117	37	106	47	85
West South Central.....	86	64	66	81	57	64	70	59	12	68
Mountain.....	44	36	53	54	44	90	44	99	71	54
Pacific.....	51	62	41	55	73	52	61	86	91	66

¹ South Bend, Ind., not included.

⁴ Lynchburg, Va., and Savannah, Ga., not included.

³ Greenville, S. C., not included.

⁵ Greenville, S. C., and Brunswick, Ga., not included.

Number of cities included in summary of weekly reports, and aggregate population of cities of each group, approximated as of July 1, 1928 and 1927, respectively

Group of cities	Number of cities reporting cases	Number of cities reporting deaths	Aggregate population of cities reporting cases		Aggregate population of cities reporting deaths	
			1928	1927	1928	1927
Total.....	101	95	31,657,000	31,050,300	30,960,700	30,369,500
New England.....	12	12	2,274,400	2,242,700	2,274,400	2,242,700
Middle Atlantic.....	10	10	10,732,400	10,594,700	10,732,400	10,594,700
East North Central.....	16	16	7,991,400	7,820,700	7,991,400	7,820,700
West North Central.....	12	10	2,683,500	2,634,500	2,683,500	2,518,500
South Atlantic.....	21	21	2,961,900	2,890,700	2,961,900	2,890,700
East South Central.....	7	6	1,048,300	1,028,300	1,000,100	980,700
West South Central.....	8	7	1,307,600	1,260,700	1,274,100	1,227,800
Mountain.....	9	9	591,100	581,600	591,100	581,600
Pacific.....	6	4	2,046,400	1,966,400	1,548,900	1,512,100

FOREIGN AND INSULAR

THE FAR EAST

Report for the week ended September 15, 1928.—The following report for the week ended September 15, 1928, was transmitted by the Eastern Bureau of the Health Section of the Secretariat of the League of Nations, located at Singapore, to the headquarters at Geneva.

Plague, cholera, or smallpox was reported at the following ports:

PLAGUE

Ceylon.—Colombo.
India.—Bombay, Cochin, Rangoon.
Indo-China.—Pnompenh.

CHOLERA

India.—Rangoon, Calcutta, Madras, Negapatam.
French India.—Pondicherry.
China.—Shanghai.

SMALLPOX

India.—Bombay, Madras, Negapatam.
French India.—Pondicherry.
Indo-China.—Pnompenh.
China.—Hong Kong.
Dutch East Indies.—Pontianak, Belawan Deli, Surabaya.

ARGENTINA

Santiago del Estero—Further relative to plague.—The information regarding the occurrence of pneumonic plague at Santiago del Estero, Argentina, published in PUBLIC HEALTH REPORTS for October 5, 1928, page 2622, was received from official sources. More recent information, however, dated October 1, states that up to that date only five cases had been definitely confirmed and two other suspected cases were being bacteriologically examined; and a dispatch dated October 3 states that two more cases had been confirmed, with no new cases reported.

CANADA

Provinces—Communicable diseases—Week ended September 22, 1928.—The Canadian Ministry of Health reports cases of certain communicable diseases from seven Provinces of Canada for the week ended September 22, 1928, as follows:

Disease	Nova Scotia	New Brunswick	Quebec	Ontario	Manitoba	Saskatchewan	Alberta	Total
Influenza.....	1							1
Lethargic encephalitis.....				1				1
Poliomyelitis.....			4	9	63	7	4	87
Smallpox.....			3	10				13
Typhoid fever.....	13	10	24	27	3	9	1	87

Quebec Province—Communicable diseases—Week ended September 22, 1928.—The Bureau of Health of the Province of Quebec reports cases of certain communicable diseases for the week ended September 22, 1928, as follows:

Disease	Cases	Disease	Cases
Chicken pox.....	7	Smallpox.....	3
Diphtheria.....	26	Tuberculosis.....	54
Measles.....	9	Typhoid fever.....	24
Poliomyelitis.....	4	Whooping cough.....	8
Scarlet fever.....	59		

-CUBA

Provinces—Communicable diseases—February 12–June 30, 1928.—During the period from February 12 to June 30, 1928, cases of communicable diseases were reported from the Provinces of Cuba as follows:

Disease	Pinar del Rio	Habana	Matanzas	Santa Clara	Camagui	Oriente	Total
Cerebrospinal meningitis.....						3	3
Chicken pox.....	13	287	70	28	31	174	603
Diphtheria.....	4	32	7	17	14	14	88
Malaria.....	2	49	6	9	109	831	1,006
Measles.....	19	115	18	17	1	1	171
Paratyphoid fever.....	13	13	14	47	11	25	123
Scarlet fever.....	2	51	6	7		1	67
Tetanus (infantile).....				3	3		6
Typhoid fever.....	56	283	82	230	103	105	859

CZECHOSLOVAKIA

Communicable diseases—July, 1928.—During the month of July, 1928, communicable diseases were reported in Czechoslovakia as follows:

Disease	Cases	Deaths	Disease	Cases	Deaths
Anthrax.....	9	2	Puerperal fever.....	45	15
Cerebrospinal meningitis.....	23	6	Rabies.....	1	1
Diphtheria.....	532	43	Scarlet fever.....	1,110	19
Dysentery.....	35		Trachoma.....	197	
Malaria.....	304		Typhoid fever.....	697	53
Paratyphoid fever.....	26	1			

GERMANY

Bavaria—Vital statistics—January 1–June 30, 1928.—The Bavarian Bureau of Statistics has recently issued the following preliminary figures on marriages, births, and deaths in Bavaria during the first six months of 1928:

Year	Estimated population, July 1	Marriages	Births, including stillbirths	Deaths	Infant mortality (deaths under 1 year)	Excess of births over deaths
1913.....	7,062,395	25,427	104,772	67,498	13,774	37,274
1927.....	7,485,785	28,193	81,252	54,580	10,054	26,672
1928.....	7,500,000	32,280	81,797	52,534	9,023	29,263

GREECE

*Dengue.*¹—Complete statistics of cases of dengue fever in the provinces of Greece are not yet available. The health section of the League of Nations states, however, that the southern and south-eastern part of the country is the most severely infected. The Provinces reporting the largest number of cases up to September 15 are the following: Attica, Laconia, Messenia, Samos, Phthiotis and Phocis, Cephalonia, Zante, Chios, Cyclades, and Salonica.

JAPAN

Osaka—Cholera.—A case of cholera was reported at Osaka, Japan, on October 4, 1928, in a member of the personnel of the Bacteriological Institute.

YUGOSLAVIA

Communicable diseases—August, 1928.—During the month of August, 1928, communicable diseases were reported in Yugoslavia as follows:

Disease	Cases	Deaths	Disease	Cases	Deaths
Anthrax.....	292	33	Measles.....	155	5
Cerebrospinal meningitis.....	9	6	Poliomyelitis.....	1	—
Diphtheria.....	222	32	Rabies.....	3	3
Dysentery.....	489	36	Scarlet fever.....	1,484	204
Leprosy.....	1	1	Tetanus.....	28	15
Lethargic encephalitis.....	1	—	Typhoid fever.....	478	42

¹ See PUBLIC HEALTH REPORTS, Sept. 21, 1928, p. 2497, Sept. 28, 1928, p. 2563, and Oct. 5, 1928, p. 2623.

C	Manila.....						4	1	
C	Pangasinan Province—						1		
D	Bayambang.....						1		
C	Surigao Province—								
D	Surigao.....						1		
D	Slam.....						163	9	
D	Ayudhya.....	200	205	201	202	203	139	144	
D	Bangkok.....	139	214	218	224	127	123	4	
D	Songkla.....	101	60	60	74	60	7	1	
D	Tred.....	66	36	33	35	26	4	1	
D	Straits Settlements: Singapore.....								
D	On vessel.....								
D	S. S. Hawaii Maru at Singapore from Saigon, French Indo-China.....	3		2	1				
D	S. S. Kambangan at Batavia from Jeddah via Sabang and Palembang.....	1		2	1				
C	S. S. Talres at Penang from Madras via Nagapatam.....			11					

Place	January-March, 1923		April, 1923		May, 1923		June, 1923			July, 1923			August, 1923			Sept. 1-10, 1923	
	1-10	11-31	1-10	11-31	1-10	11-31	1-10	11-20	21-31	1-10	11-20	21-31	1-10	11-20	21-31	1-10	21-31
Indo-China (French) (see also table above):																	
Annam.....	389		46	43			22	8	9	8	5	3		4	7	2	
Cambodia.....	312		196	101	26	82	26	111	111	92	25	38		19	19	15	
Cochin-China.....	1,407		933	404	143	109	77	65	33	65	55	33		13	15	1	
Laos.....																	
Tonkin.....	1		6	26	8	2	1	1	1	5	1	1		2			
Kwangchow-Wan.....	1			16			2								1		

Canary Islands:																							
Azores.....	0																						
Lanzarote Village.....	0																						
Las Palmas.....	0			1																			
Palma Island.....	0																						
Tenerife.....	0																						
Ceylon: Colombo.....	0			3																			
Plague-infected rats.....	0			2																			
China:																							
Kunoy.....	0			5																			
Hong Kong.....	0																						
Mongolia:																							
Tungshoo.....	0																						
Urga.....	0																						
Dutch East Indies:																							
Celebes—Makassar.....	0																						
Java:																							
Batavia and West Java.....	0			6																			
Plague-infected rats.....	0			737																			
East Java and Madura.....	0			137																			
Celebes.....	0			136																			
Kedoe Residency.....	0			7																			
Surabaya Residency.....	0			7																			
Ecuador (see also table below):																							
Alaua.....	0																						
Alexandria.....	0																						
Amrieh District.....	0			2																			
Assiout Province.....	0			2																			
Behaira.....	0																						
Bent-Suef.....	0																						
Girga.....	0																						
Magbagha District.....	0																						

1 See p. 2692.

2 During the 4 weeks ended Sept. 15, 1928, 160 cases of plague were reported at Tungshoo, Mongolia.

Shanghai—	5	6	7	3	11	7	2	1	1	1	1
Foreigners only.....	11	10	0	0	10	11	10				
Including natives.....	48	20	13	17	9	14	1				
Tientsin.....											
Chosen (see table below).....											
Gurosoo (elasmim).....	2	3	2	1			2				
Dominican Republic; Santo Domingo.....											
Dutch East Indies:											
Balkpapan.....											
Belawan Deli.....						11	7				
Borneo—Pontianak.....						3	1				
Java—							2				
Batavia and West Java.....	4	2	5	1	2		2				
East Java and Madura.....	2	9	2	2		2	3				
Surabaya.....							3		1	3	2
Sumatra—											
Medan.....	23	14	7	2	10	4	5	5	1	2	
Palembang.....	11	2	5	3	5		2	2	1		
Ecuador (see table below).....	1	2	52	12	1		1				
Egypt.....			33	7	1						
Belgium Province.....							1				
Caro.....	1										
France (see table below).....											
Gold Coast (see table below).....											
Great Britain:											
England and Wales.....	1,630	1,473	1,341	1,344	1,190	1,140	681	126	111	114	141
Birmingham.....	44	3	1	1		2					102
Bradford.....	24	13	19	14	17	10	2				123
Bristol.....	3	4	12	17	10	2			2		
Cardiff.....			5	3	2						
Castleford.....		4	26	69	24	18	14	6	2	1	1
Hull.....						1	24	6	2	3	4
Leeds.....	3	9	14	1		8	1	1			8
Liverpool.....											1
London.....	4	14	17	42	25	90	19	3	5	1	2
Manchester.....	25	8	14	8	5	1					3
Newcastle-on-Tyne.....	27	12	9	4	12	6	28		1		
Nottingham.....	27	11	13	17	20	3	6		3	5	2
Plymouth.....											1
Sheffield.....	8	6	1	14	4	2			6	6	1
Stoke-on-Trent.....	1	15	12	32	24	14			1	2	1
Weymouth.....								4	4	1	5
Scotland—											
Ayrbroath.....											
Dundee.....									1	3	

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

TYPHUS FEVER

[C, indicates cases; D, deaths; F, present]

Place	Week ended—											
	July, 1928			August, 1928			September, 1928					
	7	14	21	28	4	11	18	25	1	8	15	22
Algeria (see also table below):												
Algiers.....	C	1	2		1	4	3	2	1			
Oran.....	D	2	6	2	1	1	1	1				
Austria: Vienna.....	C	1										
Bulgaria.....	D	13	18	26				3	5	2		
Sofia.....	D	1	4					2				
Chile:	D	1										
Iquique.....	C											
Talcahuano.....	D											
Valparaiso.....	D	1										
China (see also table below):	D											
Manchuria—	D											
Harbin.....	C											
Kwantung.....	C			2	16							
South Manchuria Railway Zone.....	C			17	283							
Tientsin.....	C			10	5	1						
Chosen (see table below).	C											
Czechoslovakia (see table below).	C											
Egypt.....	C	9	17	2	3	11	7					
Alexandria.....	D	2	8	1	1	2	3					
Assiut Province.....	D											
Assuan Province.....	C											
Benara Province.....	D											
Cairo.....	C			29	32	43	7	2				
Dakallah.....	D			2	4	7	2					
	D			1	4	4						

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

TYPHUS FEVER—Continued
 [C, indicates cases; D, deaths; F, present]

Place	Jan- uary- March, 1928	April, 1928	May, 1928	June, 1928	July, 1928	Aug- ust, 1928	Place	Jan- uary- March, 1928	April, 1928	May, 1928	June, 1928	July, 1928	Aug- ust, 1928
	886	210	241	182	---	---		46	2	1	---	---	---
Chosen.....	C	19	32	24	---	---	D	2	1	---	---	---	---
Chemulpo.....	C	1	2	---	---	---	D	17	---	---	---	---	---
Gonsan.....	C	1	---	---	---	---	D	1	---	---	---	---	---
Seoul.....	D	10	18	4	---	---	D	1	---	---	---	---	---
Czechoslovakia.....	C	1	---	---	---	---	C	199	---	---	---	---	---
Greece: Athens.....	C	25	4	---	---	---	C	---	---	---	---	---	---
Japan.....	C	4	13	11	9	6	C	17	---	---	---	---	---
Latvia.....	C	27	4	65	2	---	C	1,476	---	---	---	---	---
Lithuania.....	C	223	66	64	32	8	C	5,167	---	---	---	---	---
	D	22	3	3	1	---	D	34	10	19	16	12	3
								3	1	3	1	3	---

YELLOW FEVER

Place	Jan. 15- Feb. 15, 1928	Feb. 15- Mar. 10, 1928	Mar. 11- Apr. 5, 1928	Apr. 6- May 2, 1928	May 3-30, 1928	Week ended—											
						July, 1928				August, 1928				September, 1928			
	7	14	21	28	4	11	18	25	1	8	15	22	29				
Belgian Congo: Matadi.....	C	1	---	2	2	---	---	---	---	---	---	---	---	---	---	---	---
Brazil:	D	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Aracaju.....	D	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Bahia.....	D	---	---	---	2	---	---	---	---	---	---	---	---	---	---	---	---
Estancia.....	D	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Perambuco.....	C	1	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Rio de Janeiro.....	C	---	---	---	2	48	10	12	10	8	6	4	4	4	4	1	8
Sao Felix.....	C	---	---	---	2	15	6	7	9	---	---	---	---	---	---	---	---
Dahomey: Grand Popo.....	C	---	---	---	2	3	---	---	---	---	---	---	---	---	---	---	---
Gold Coast.....	C	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Ivory Coast.....	C	---	---	---	2	---	---	---	---	---	---	---	---	---	---	---	---
Abidjan.....	C	---	---	---	---	1	---	---	---	---	---	---	---	---	---	---	---
Ferkessedougou.....	C	---	---	---	---	1	1	---	---	---	---	---	---	---	---	---	---