CONTENTS

| Prevalence of poliomyelitis |
|---|
| National Cancer Institute building completed. |
| Disabling morbidity among industrial workers, second quarter and first half of 1939 |
| Studies in chemotherapy. X. Colorimetric tests for aromatic hydroxylamines and for further oxidation products of aromatic amines. Their demonstration in the urine following sulfanilamide administration |
| Provisional mortality rates for the first 6 months of 1939 |
| Court decision on public health |
| Deaths during week ended September 30, 1939: |
| Deaths in a group of large cities in the United States |
| Death claims reported by insurance companies |
| PREVALENCE OF DISEASE |
| United States: |
| Current weekly State reports: |
| Reports for week ended October 7, 1939, with case rates and |
| comparison with former years |
| Summary of monthly reports from States |
| Weekly reports from cities: |
| City reports for week ended September 30, 1939 |
| Foreign reports: |
| Cuba— |
| Habana—Communicable diseases—4 weeks ended September 23, |
| Provinces-Notifiable diseases-4 weeks ended September 16, |
| 1939 |
| Italy—Communicable diseases—4 weeks ended July 16, 1939 |
| Japan—Tokyo—Encephalitis, lethargic |
| Reports of cholera, plague, smallpox, typhus fever, and yellow fever |
| received during the current week- |
| Cholera |
| Plague |
| Smallpox |
| Yellow fever |