CONTENTS

| Administrative factors involving effectiveness of communicable disease control. |
|---|
| Cerebrospinal pathology of experimental poliomyelitis in the eastern cotton rat, Sigmodon hispidus hispidus, and in the white mouse, Mus musculus |
| A note on the pathology in monkeys of the Lansing strain of poliomyelitis virus before and after passage in the cotton rat |
| A highly virulent strain of Rocky Mountain spotted fever virus isolated in the eastern United States |
| Cancer mortality in the United States according to site, by sex and age, 1938 |
| Court decision on public health |
| Deaths during week ended April 6, 1940: |
| Deaths in a group of large cities in the United States |
| Death claims reported by insurance companies |
| PREVALENCE OF DISEASE |
| United States: |
| Reports from States for week ended April 20, 1940, and comparison |
| with former years |
| Weekly reports from cities: |
| City reports for week ended April 6, 1940 |
| Foreign reports: |
| Cuba—Provinces—Notifiable diseases—4 weeks ended March 30, |
| 1940 |
| Sweden—Notifiable diseases—February 1940 |
| World distribution of cholera, plague, smallpox, typhus fever, and |
| yellow fever— |
| Cholera |
| Plague |
| Smallpox |
| Typhus fever |
| Yellow fever |