## CONTENTS

Prevalence of communicable diseases in the United States, February 23- March 22, 1941
Mobile laboratory units of the Ohio River Pollution Survey
Domestic water and dental caries. Il. A study of 2,832 white children,
aged 12-14 years, in 8 suburban Chicago communities, including Lactobacillus acidophilus studies of 1,761 children
The application of the human serum opacity reaction for evaluating the antitoxin binding power (Lb) of Clostridium perfringens (type A) toxoid
Disabling morbidity among industrial workers, final quarter of 1940, with an index of the previous publications of this series
The prevalence of disabling illness among male and female workers and housewives
Additional contributions to outside journals of personnel of the Public
Court decision on public health
Deaths during week ended March 29, 1941:
Deaths in a group of large cities in the United States
Death claims reported by insurance companies
PREVALENCE OF DISEASE
United States:
Reports from States for week ended April 5, 1941, and comparison with former years
Weekly reports from cities:
City reports for week ended March 22, 1941
Territories and possessions:
Hawaii Territory—Plague (rodent)
Foreign reports:
Canada—Provinces—Communicable diseases—Week ended March 1,
1941
Cubs—Provinces—Notifiable diseases—4 weeks ended March 1, 1941_
Jamaica—Communicable diseases—4 weeks ended March 15, 1941
Switzerland—Notifiable diseases—December 1940
Reports of cholera, plague, smallpox, typhus fever, and yellow fever
received during the current week—
Plague
Yellow fever