



HHS Public Access

Author manuscript

Ann Epidemiol. Author manuscript; available in PMC 2020 January 01.

Published in final edited form as:

Ann Epidemiol. 2019 January ; 29: 67–73.e1. doi:10.1016/j.annepidem.2018.10.006.

Associations between County-level Voter Turnout, County-level Felony Voter Disenfranchisement, and Sexually Transmitted Infections among Women in the Southern United States

Danielle F. Haley, PhD, MPH^a, Andrew Edmonds, PhD^b, Victor J. Schoenbach, PhD^c, Catalina Ramirez, MPH, MPA^d, DeMarc A. Hickson, PhD^e, Gina M. Wingood, ScD^f, Hector Bolivar, MD^g, Elizabeth Golub, PhD^h, and Adaora A. Adimora, MD, MPHⁱ

^aInstitute for Global Health and Infectious Diseases, School of Medicine, University of North Carolina at Chapel Hill 130 Mason Farm Road, Chapel Hill, NC, 27599 USA and Northeastern University, Bouvé College of Health Sciences, 360 Huntington Avenue, Boston, MA 02115, USA; d.haley@northeastern.edu

^bDepartment of Epidemiology, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, 137 East Franklin Street, Chapel Hill, North Carolina USA; aedmonds@email.unc.edu

^cDepartment of Epidemiology, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, 2104D McGavran-Greenberg Hall, Chapel Hill, North Carolina USA; vjs@unc.edu

^dInstitute for Global Health and Infectious Diseases, School of Medicine, University of North Carolina at Chapel Hill 130 Mason Farm Road, Chapel Hill, NC, 27599 USA; catalina_ramirez@med.unc.edu

^eDepartment of Epidemiology and Biostatistics, Jackson State University School of Public Health, 350 West Woodrow Wilson Drive, Room 222, Jackson, MS 39213 USA; dhickson@uhupil.org

^fDepartment of Sociomedical Sciences, Lerner Center for Public Health Promotion, Mailman School of Public Health at Columbia University, 722 West 168th Street, New York, NY, 10032 USA; gw2326@cumc.columbia.edu

^gDivision of Infectious Diseases, University of Miami Miller School of Medicine, 1611 NW 12th Ave, Miami, FL 33136 USA; hbolivar@med.miami.edu

Corresponding Author Contact Information Danielle F. Haley, Northeastern University, Bouvé College of Health Sciences, 360 Huntington Avenue, Boston, MA 02115, USA, T: 919-357-1045, d.haley@northeastern.edu.

Role of Funder. The funders were not involved in study design; in the collection, analysis and interpretation of data; in the writing of the articles; and in the decision to submit it for publication.

Conflict of Interest

The authors have no financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

^hDepartment of Epidemiology, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe Street, Baltimore, Maryland 21205 USA; egolub@jhu.edu

ⁱInstitute for Global Health and Infectious Diseases, School of Medicine, University of North Carolina at Chapel Hill and Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, 130 Mason Farm Road, Chapel Hill, NC, 27599 USA; adimora@med.unc.edu

Abstract

Purpose.—Voting may play a critical role in the allocation of social and structural resources to communities, which in turn shapes neighborhood environments, and ultimately, an individual’s sexually transmitted infection (STI) risk. We assessed relationships among county-level voter turnout and felony voter disenfranchisement, and STIs.

Methods.—This cross-sectional multilevel analysis included 666 women in Alabama, Florida, Georgia, Mississippi, and North Carolina enrolled in the Women’s Interagency HIV Study between 2013 and 2015. Having a baseline bacterial STI (chlamydia, gonorrhea, trichomoniasis, or early syphilis) was determined by laboratory testing. We used generalized estimating equations to test relationships between county-level voter turnout in the 2012 general election, county-level percentage of felony disenfranchised voters, and STI prevalence.

Results.—Eleven percent of participants had a STI. Higher voter turnout corresponded to lower STI prevalence (PR=0.84, 95% CI=0.73–0.96 per 4 percentage point higher turnout). Greater felony voter disenfranchisement corresponded to higher STI prevalence (PR =1.89, 95% CI=1.10–3.24 per 4 percentage point higher disenfranchisement).

Conclusion.—STI prevalence was inversely associated with voter turnout and positively associated with felony voter disenfranchisement. Research should assess causality and mechanisms through which civic engagement shapes sexual health. Expanding political participation, including eliminating discriminatory voting laws, could influence sexual health.

Keywords

women; sexually transmitted infections; political participation; felony voter disenfranchisement

Introduction

The United States (US) has experienced a dramatic increase in the rate of reportable sexually transmitted infections (STIs) in the past decade, despite advances in diagnosis and treatment.

¹ STIs contribute to comorbidities (e.g., pelvic inflammatory disease) and facilitate transmission of HIV.¹ The burden of STIs is especially notable in the South, which consistently reports higher rates of STIs than other regions of the US.^{1,2}

African American women are at greater risk of acquiring STIs, including HIV, than non-African American women, despite engaging in the same or lower levels of sexual risk behavior^{3,4} Features of the social and built environment are powerful determinants of racial disparities in STIs, including HIV. Specifically, living in areas with comparatively greater poverty, a shortage of men relative to women, and more social disorder (e.g., violent crime)

is associated with higher risk sexual networks (e.g., sexual partnerships that overlap in time) which facilitate the spread of STIs⁵⁻⁹ Social and political factors may differentially allocate fiscal and structural capital to communities, which in turn reduce residents' access to resources needed to support sexual health (e.g., health care services, educational opportunities), while simultaneously over-exposing residents to community phenomena (e.g., violent crime, illicit drug use) associated with greater sexual risk^{2,10-12} In the US, the legacy of segregation, coupled with ongoing discriminatory macroeconomic and social forces, has produced community environments that vary by race^{2,4,13, 14} Nationally representative samples demonstrate stark contrasts in area socioeconomic and social (e.g., ratio of men to women) contexts by race, highlighting potential pathways between social inequities and racial disparities in HIV/STIs.⁹

Political parties and the policies elected officials implement have important downstream implications for resource allocation and resulting health disparities.^{11, 12, 15} For example, the implementation of the Patient Protection and Affordable Care Act (ACA) has the potential to ameliorate inequities in health care access, including STI screening and treatment, by decreasing the number of low-income people who lack health insurance. Yet, implementation of the ACA, especially in the South, has been limited by many states' refusal to expand Medicaid.¹⁶ Because laws and policies are shaped by elected officials, voting plays a critical role in determining the allocation of social and structural resources to communities, which in turn shapes community environments and, ultimately, an individual's risk of acquiring an STI.

Non-voters are more likely to be poor, have less formal education, be in poor health, and report unmet needs for medical care.^{15, 17, 18} Living in areas with socioeconomic inequalities in voter turnout is associated with poorer self-rated health.¹⁵ Systematic restrictions on voting which disproportionately and negatively impact racial and ethnic minorities and the poor may promote health disparities by excluding substantial portions of the population from participating in the political process.^{2, 10, 15, 19} Over 6 million Americans are prohibited from voting due to felony disenfranchisement laws restricting voting rights for individuals convicted of felony offenses.^{20, 21} The length of voting prohibitions vary by state, with the period of disenfranchisement for persons convicted of a felony offense ranging from the duration of the sentence to lifetime prohibition.^{20, 21} The potential impact of felony disenfranchisement varies substantially across states and racial and ethnic groups.^{2, 10, 19, 21, 22} One in every 13 African American adults is disenfranchised, as compared to 1 in 56 non-African American adults.²¹ The South has some of the strictest (e.g., lifetime prohibition) and highest rates of felony disenfranchisement, with as many as 1 in 4 African Americans in some states prohibited from voting.^{2, 20}

The objective of this cross-sectional multilevel study was to assess the relationship between two related constructs capturing county-level political participation (i.e., voter turnout and felony voter disenfranchisement) and having a bacterial STI, among a predominantly African American cohort of women living in the South. We hypothesized that county-level voter turnout would be inversely associated with having an STI, whereas county-level voter disenfranchisement would be positively associated with having an STI.

Materials and Methods

The Women's Interagency HIV Study (WIHS) is a multisite, prospective cohort study designed to characterize the natural history, clinical, and behavioral impact of HIV among US women.^{23, 24} The WIHS includes HIV-seronegative women whose sociodemographic characteristics are similar to those of the HIV-seropositive women in the cohort, who, in turn are representative of the race/ethnicity of HIV-seropositive US women.²³⁻²⁶ Between October 2013 and September 2015, WIHS clinical research sites in five states in the South (Alabama, Florida, Georgia, Mississippi, and North Carolina) enrolled HIV-seropositive and HIV-seronegative women, aged 25–60 years. Women with HIV were antiretroviral therapy (ART)-naïve or began taking highly active antiretroviral therapy (HAART) only after December 31, 2004 and had never been on non-HAART ART (except possibly when pregnant). All had documented CD4 counts and HIV viral load before initiation of ART. HIV-seronegative women reported that either they or a sexual partner met at least one criterion associated with increased risk of HIV acquisition in the last five years (e.g., illicit drug use).

Participants were recruited by WIHS using multiple strategies, including community-based organization referrals. Institutional Review Board (IRB) approval was obtained at each institution and written informed consent was obtained prior to study procedures. Methods are described in detail elsewhere.^{23, 24} This secondary analysis, approved by the University of North Carolina IRB, was restricted to Southern site participants providing written informed consent to collect and geocode their residential addresses.

Measures

Outcome: Current bacterial sexually transmitted infection—The binary outcome, having a current bacterial STI, was defined as a laboratory-confirmed diagnosis (Appendix A) at baseline of: chlamydia, gonorrhea, trichomoniasis, or early syphilis (titer and confirmatory test results consistent with primary, secondary, or early latent [<1 year duration] infection). Participants with a STI were referred to medical providers for treatment.

County-level exposures: Voter turnout and felony voter disenfranchisement—Our county-level exposures captured two dimensions of political participation: voter turnout (the percentage of voting age residents casting a vote in the general election) and felony voter disenfranchisement (the percentage of voting age residents excluded from voting due to a felony conviction).

County-level percentage voter turnout (continuous, mean-centered) was calculated as the percentage of the voting age population (aged 18 and older, based on American Community Survey [ACS] 2010–2014 5-year estimates) casting a vote for any presidential candidate in the 2012 general election. Total votes cast per county were obtained from Dave Leip's Atlas of US Presidential Elections.²⁷

County-level percentage of felony disenfranchised voters was estimated by applying 2010 state-level felony disenfranchisement rates for African Americans, as calculated by Uggen

and colleagues,²⁰ to counties based on ACS 2010–2014 5-year estimates of voting age African American county residents. All states included in these analyses have voter disenfranchisement laws, with estimates of disenfranchised African Americans ranging from 2.8% in North Carolina to 23.3% in Florida.²⁰

Because past research suggests that area socioeconomic conditions are associated with STIs as well as voter turnout, we also evaluated whether the relationships between our county-level exposures (voter turnout, felony voter disenfranchisement) and having an STI were independent of county-level socioeconomic advantage.^{5–8, 15, 28} Measures of county socioeconomic advantage (i.e., median household income, percentage of residents 25 or older with a college degree or greater, and percentage of residents over age 16 who were employed) were created using ACS 2010–2014 5-year estimates. Because these socioeconomic measures were correlated, we used principal component analysis (PCA) with orthogonal rotation (varimax) to capture underlying constructs and avoid multicollinearity.^{5, 29–31} The PCA produced one component with eigenvalue >1 which accounted for 88% of the variability explained by these factors. Continuous, standardized component scores were extracted for each participant and included in multivariable models. County-level political participation measures were not correlated with county socioeconomic advantage (Pearson's $r < 0.60$).

Covariates—The WIHS collected all demographic and behavioral data using interviewer-administered questionnaires. We adjusted for the following participant-level demographics and behaviors in the past six months, which were binary unless otherwise noted, and included: age in years (continuous, mean-centered), non-Hispanic African American race/ethnicity, annual household income of \$18,000 or less, less than a high school education or equivalent (e.g., GED), and being HIV-seropositive (i.e., reactive serologic enzyme-linked immunosorbent assay test and a confirmed positive western blot or a detectable plasma HIV-1 ribonucleic acid). Given the geographic diversity of our sample and the relationship between voter turnout and travel distance,[>] we also controlled for population density (continuous, mean-centered), defined as persons per square mile, using ACS 2010–2014 5-year estimates.³²

Analysis

All analyses were conducted using SAS 9.4 (SAS Institute Inc., Cary, NC). We used descriptive statistics to assess distributions of variables, and to compare these distributions by HIV status and missingness. We modeled bivariable and multivariable relationships using generalized estimating equations (GEE; PROC GENMOD) with a binomial distribution and log link (to estimate prevalence ratios [PRs] and 95% confidence intervals [CIs]), and an exchangeable correlation structure. We selected PRs over odds ratios because: 1) Odds ratios overstate effect sizes relative to PRs, this overstatement occurs to a greater extent with increasing prevalence of the outcome; and 2) PRs are more easily interpretable because they represent a ratio of proportions rather than a ratio of odds.^{33–35} We selected GEEs over mixed models because our goal was to generate population-average estimates of the associations between county-level political participation and STIs.³⁶ Because of the WIHS study design, which consists predominantly of HIV-seropositive women, we tested

statistically for multiplicative interactions between county-level political participation and having an STI by HIV status (e.g., HIV status*voter turnout, HIV status*felony voter disenfranchisement), retaining terms with $p < 0.05$ in the final multivariable model.

We conducted sensitivity analyses to test the robustness of findings. First, in order to test whether relationships between our county-level exposures (voter turnout, felony voter disenfranchisement) and STI prevalence were independent of county-level socioeconomic advantage, we compared PR estimates for each county-level exposure in models with and without county-level socioeconomic advantage. We determined a priori that <10% difference in county-level PR estimates between models suggested that the relationship between each county-level exposure and having an STI was independent of county-level socioeconomic advantage. Second, in order to examine whether the contribution of felony voter disenfranchisement to having an STI was distinct from underlying racial composition of counties, we compared PR estimates for models assessing relationships between voter disenfranchisement and having an STI to PR estimates for models assessing relationships between percentage of voting age African American residents and having an STI. We determined a priori that >10% difference in PR estimates in county-level voter disenfranchisement, as compared to the percentage of voting age African American residents suggested that these exposure variables were capturing different underlying constructs. Third, parallel analyses were conducted using random effects models (PROC GLIMMIX) in order to evaluate the robustness of findings. Prevalence ratios for our county-level exposures (voter turnout, felony voter disenfranchisement) generated using random effects models, as compared to prevalence ratios generated using GEEs, were within 7%. Furthermore, the random effect for the intercept in unconditional models was near zero, suggesting low intra-county variability and further supporting the use of GEEs.

Results

Participant Characteristics

A total of 845 women enrolled at WIHS sites in the South. One hundred and seventy-nine women were excluded from these analyses because they (1) did not have geocoded address data ($n=116$; the majority of these women did not consent for geocoding [$n=65$]) or (2) were missing one or more STI laboratory test results ($n=63$). Participants excluded because they were missing geocoded address data were more likely to report annual household incomes \$18,000 (83.2% vs. 69.0%; Chi-square Fisher's Exact Test, $p < 0.05$). Participants enrolled at study sites other than Georgia were more likely (Chi-square Fisher's Exact Test, $p < 0.05$) to have missing STI data, as were participants with HIV. We controlled for income, site, and HIV status in all multivariable models in order to minimize potential bias.

The 666 participants included in these analyses lived in 61 unique counties, with a median of 2 participants per county and first [Q1] and third [Q3] quartiles of 2 and 5, respectively. Across the 666 participants in the final analytic sample, the distribution of county median household income had a median of \$50,596 (Q1: \$43,099, Q3: \$56,642). County voter turnout percentage in the 2012 general election ranged from 41.0% to 79.4% (median: 57.0%, Q1: 53.8%, Q3: 60.2%); felony disenfranchisement percentages ranged from 0.22% to 11.8% (median: 3.9%, Q1: 1.6%, Q3: 5.6%). Eleven percent of participants ($n=76$) tested

positive for at least one bacterial STI at baseline. The mean age of participants was 43.5 years (SD=9.4), 70.7% were HIV-seropositive, and 84.6% identified as non-Hispanic African American. HIV-seropositive and HIV-seronegative participants, respectively, were similar for both the outcome (having an STI: 11.7% vs. 10.7%) and exposures (voter turnout: 56.7% vs. 57.5%, felony voter disenfranchisement: 3.9% vs. 3.9%).

Relationships between voter turnout and having a bacterial STI

County-level voter turnout was associated with lower STI prevalence. In bivariable analyses (Table 2), county-level voter turnout was modestly associated with having a bacterial STI (PR=0.87, 95% CI=0.77–1.00). In the multivariable model controlling for participant- and county-level characteristics (Model A), a four-percentage point higher county-level voter turnout (e.g., from 57 to 61) corresponded to a lower STI prevalence (Model A; PR =0.84, 95% CI=0.73–0.96). County-level socioeconomic advantage was not associated with having an STI in the bivariable (PR =1.19, 95% CI=0.97–1.47) and multivariable models (Model A; RR PR =1.33, CI=0.91–1.95), providing insufficient evidence of an association between county-level socioeconomic advantage and having an STI. Prevalence ratios for county-level voter turnout in models controlling for and not controlling for county-level socioeconomic advantage were within 10% (results not shown), suggesting that relationships between county-level voter turnout and having a bacterial STI were independent of county-level socioeconomic advantage. There was no statistically significant interaction between county-level voter turnout and HIV status on the multiplicative scale ($p>0.05$).

Relationships between voter disenfranchisement and having a bacterial STI

In the multivariable model controlling for participant- and county-level characteristics (Model B), a four-percentage point higher county-level felony voter disenfranchisement (e.g., from 0 to 4 percent) corresponded to a greater STI prevalence (PR =1.89, 95% CI=1.10–3.24). The bivariable association (PR =1.18, 95% CI=0.90–1.54) was much weaker or absent, suggesting that the association was obscured by confounding. County-level socioeconomic advantage was not associated with having an STI in the bivariable (PR =1.19, 95% CI=0.97–1.47) and multivariable models (Model B; PR =1.36, CI=0.94–1.97), providing insufficient evidence of an association between county-level socioeconomic advantage and STI prevalence. Prevalence ratios for county-level voter disenfranchisement in models not controlling for county-level socioeconomic advantage were shifted towards the null (PR =1.46, CI=0.99–2.14). Given that more socioeconomically advantaged areas experience less crime and incarceration, it logically follows that adjusting for this factor would result with an effect estimate with a higher magnitude. There was no statistically significant interaction between county-level felony voter disenfranchisement and HIV status on the multiplicative scale ($p>0.05$).

The percentage of county-level voting age African American residents was weakly associated with STI prevalence in the bivariable model (PR =1.05, 95% CI=1.01–1.10), but not in the multivariable model controlling for participant- and county-level characteristics (Model C; RR=1.16, 95% CI=0.85–1.58). We compared PR estimates for models assessing relationships between felony voter disenfranchisement and having an STI (Model B; PR =1.89, 95% CI=1.103.24) to PR estimates for models assessing relationships between

percentage of voting age African American residents and having an STI (Model C; PR =1.16, 95% CI=0.85–1.58) and concluded county-level felony voter disenfranchisement and county-level percentage of African American residents captured distinct underlying constructs, with county-level felony voter disenfranchisement being the more powerful explanatory variable. There was no statistically significant interaction between the percentage of county-level voting age African American residents and HIV status on the multiplicative scale ($p>0.05$).

Discussion

In this multilevel analysis controlling for participant-level characteristics, we found that greater county-level voter turnout was associated with a lower prevalence of having a bacterial STI among women living in the Southern US. Moreover, greater county-level felony voter disenfranchisement, but not county-level percentage of African-Americans, was associated with greater prevalence STIs. These relationships did not vary by HIV status.

The ability to vote is a symbol of inclusion that affirms one's sense of collective identity and provides opportunities to shape policy.^{15,19, 22, 37} Politically active neighborhoods promote politically active citizens.²⁸ Communities with higher levels of civic engagement may be more effective in garnering human and physical resources needed to forward agendas and policies with positive community health benefits. For example, natural experiments exploring relationships between voter turnout, alcohol availability, and gonorrhea rates before and after the 1992 civil unrest in Los Angeles, California found that areas with increased voter turnout had fewer alcohol outlets (businesses such as liquor stores or convenience stores that sell alcohol for off-premise consumption) reopen following the unrest,³⁸ and subsequently, areas with lower alcohol availability also had lower neighborhood gonorrhea rates.³⁹ These findings support the possibility that voter turnout can change neighborhood characteristics in ways that improve health.

The disproportionate incarceration of people of color in the US, combined with felony voting restrictions, has resulted in the exclusion of substantial portions of African Americans from the political process.^{2, 20} Our finding that a 4-percentage point higher felony voter disenfranchisement was associated with a nearly 2-fold greater prevalence of having a bacterial STI is notable, especially since in over 50% of US states, greater than 4% of the African American adult population is disenfranchised due to a felony conviction.²¹ Racial and socioeconomic inequalities in political participation result in detrimental policies for financially vulnerable groups.^{11, 12, 15} Political participation has the potential to determine policies and policy makers, which in turn dictates resource allocation; systematic exclusion of African Americans from voting promotes inequitable distribution of assets and subsequent health disparities.^{2, 10, 15, 19} Research by Uggen and Manza demonstrates that felony voter disenfranchisement skews elections in favor of more conservative parties.^{22, 37} This electorate imbalance is further heightened when simultaneously accounting for excess mortality among African Americans.¹⁰ US states with more liberal leanings also have policies supporting positive reproductive health outcomes for women (e.g., fewer teen births, more human papillomavirus vaccination coverage).^{40, 41}

Findings are subject to limitations. Although the WIHS provides high-quality data concerning women who have or are at risk for HIV infection in the Southern US, participants agree to long-term follow-up and therefore may not be representative of the general population. Participants excluded from this analysis due to a lack of geocoded address data may have lived in qualitatively different communities. However, participants with and without geocoded address data were not different with respect to our outcome (STIs). We did not have sufficient sample size to assess relationships between county-level political participation and each STI type (e.g., chlamydia alone). However, past research demonstrates that high incidence areas of reportable STIs overlap geographically.⁴² Due to the cross-sectional nature of our study, we have not attempted to draw conclusions regarding the causality of the relationship between county-level political participation and having an STI. We assessed the relationship between two related constructs capturing county-level political participation and bacterial STI prevalence, controlling for a mixture of individual and area-level covariates. It is possible that the county-level measures of political participation do not adequately represent the individual under study (ecologic fallacy). Findings may also be subject to residual confounding. Our measures of political participation may be markers for social conditions (e.g., incarceration rates, ratio of men to women) not included in the analysis that are strong predictors of STIs.^{9, 43–45} County-level felony voter disenfranchisement data are not publicly available. Our measure of disenfranchisement represents a proxy based on best available data. The fact that the association of STI prevalence with our constructed measure of felony disenfranchisement (based on percentage of voting age African Americans) is much stronger than the corresponding association with percentage of African American voting age residents itself suggests that felony disenfranchisement is highly informative, highlighting once again that inequities in STIs are a function of social and structural inequities, and of racism rather than of race.^{2, 13, 14}

This study found an inverse association between county-level voter turnout and having an STI and a positive association between county-level felony voter disenfranchisement and having an STI. Additional research is needed to assess causality and to elucidate the mechanisms through which community-level political participation is linked to STIs among women living in the Southern US. Additional research examining the relationships between area-level civic engagement and sexual health using nationally-representative samples with geographically-linked information is warranted. Expanding opportunities for civic engagement, including voter registration campaigns and eliminating discriminatory voting laws, could provide positive benefits for individual- and community-level sexual health.

Acknowledgements

The authors thank the Women's Interagency HIV Study participants for sharing their time and experiences. The authors also thank Camden Bay of the UNC Center for AIDS Research Biostatistics Core (P30 AI50410) and Jonathan Purtle for their valuable feedback on study methods and manuscript drafts.

Sources of Support

The Women's Interagency HIV Study (WIHS) is funded primarily by the National Institute of Allergy and Infectious Diseases, with additional co-funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, the National Cancer Institute, the National Institute on Drug Abuse, and the National Institute on Mental Health. Participant data in this manuscript were collected by the UAB-MS WIHS (PIs: Michael

Saag, Mirjam-Colette Kempf, and Deborah Konkle-Parker), U01-AI-103401; Atlanta WIHS (PIs: Ighovwerha Ofotokun and Gina Wingood), U01-AI-103408; Miami WIHS (PIs: Margaret Fischl and Lisa Metsch), U01-AI-103397; UNC WIHS (PI: Adaora Adimora) U01-AI-103390; WIHS Data Management and Analysis Center (PIs: Stephen Gange and Elizabeth Golub) U01-AI-042590. Targeted supplemental funding for specific projects is also provided by the National Institute of Dental and Craniofacial Research, the National Institute on Alcohol Abuse and Alcoholism, the National Institute on Deafness and other Communication Disorders, and the NIH Office of Research on Women's Health. WIHS data collection is also supported by U11-TR000454 (Atlanta CTSA). DAH's time was supported by the Centers for Disease Control and Prevention under Cooperative Agreement U01PS003315 as part of the Minority HIV/AIDS Research Initiative. The contents of this publication are solely the responsibility of the authors and do not represent the official views of the National Institutes of Health.

Appendix A. Specimen type, test, and sensitivity and specificity for sexually transmitted infection (STI) evaluation at The Women's Interagency HIV Study sites in Alabama, Florida, Georgia, Mississippi, and North Carolina between 2013–2015

STI	Specimen Type	Test (Sensitivity, Specificity) by Site				
		Alabama	Florida	Georgia	North Carolina	Mississippi
Chlamydia	Cervical swab ¹	APTIMA Nucleic Acid Amplification Test (NAAT) (94.30, 98.00)	Becton Dickinson (BD) Probetec ET System (93.80, 99.80)	Aptima Combo 2 for CT/NG (96.60, 98.50)	Gen Probe Aptima (98.30, 96.10)	ROCHE Cobas polymerase chain reaction (PCR) (94.90, 99.40)
Gonorrhea	Cervical swab ¹	APTIMA NAAT (92.00, 99.80)	BD Probetec ET System (88.00, 99.80)	Aptima Combo 2 for CT/NG (96.60, 98.50)	Gen Probe Aptima (97.30, 99.00)	ROCHE Cobas PCR (96.60, 99.90)
Trichomoniasis	Vaginal swab ²	APTIMA NAAT (100.00, 98.10)	Wet mount ³ (N/A)	Wet mount ³ (N/A)	Wet mount ³ (N/A)	Wet mount ³ (N/A)
Early Syphilis ⁴	Serum	BD Screening rapid plasma regain (RPR) with confirmatory <i>Treponema pallidum</i> haemagglutination assay or <i>Treponema pallidum</i> particle agglutination assay (Screening: 3.00-100.00, 98.00, Confirmatory: 99.40, 100.00)	Arlington Scientific RPR Card (95.00, 98.00)	BD RPR titer, with confirmatory IgG enzyme immunoassay if reactive (N/A)	Labcorp Screening RPR with Confirmatory Quantitative RPR (Screening: 99.00, 98.40; Confirmatory: 100.00, 99.80)	BD RPR (86.00, N/A)

¹Urine was used for testing at Alabama site.

²Cervical swab was used for testing at Alabama site

³Trichomoniasis is defined as the presence of motile trichomonads on a vaginal wet mount.

⁴Titer and confirmatory testing results consistent with primary, secondary, or early latent (<1 year duration) infection.

Abbreviations and Acronyms

ACA	Affordable Care Act
ACS	American Community Survey
ART	Antiretroviral therapy
CI	Confidence interval

HAART	Highly active antiretroviral therapy
HIV	Human immunodeficiency virus
IRB	Institutional Review Board
PCA	Principal components analysis
Q	Quartile
PR	Prevalence ratio
STI	Sexually transmitted infection
US	United States
WIHS	Women's Interagency HIV Study

References

- Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2015. Atlanta; 2016.
- Adimora AA, Ramirez C, Schoenbach VJ, et al. Policies and politics that promote HIV infection in the Southern United States. *AIDS* 2014;28(10):1393–7. [PubMed: 24556871]
- Adimora AA, Schoenbach VJ. Social context, sexual networks, and racial disparities in rates of sexually transmitted infections. *J Infect Dis* 2005;191 Suppl 1:S115–22. [PubMed: 15627221]
- Hogben M, Leichliter JS. Social determinants and sexually transmitted disease disparities. *Sex Transm Dis* 2008;35(12 Suppl):S13–8. [PubMed: 18936725]
- Ford JL, Browning CR. Neighborhood social disorganization and the acquisition of trichomoniasis among young adults in the United States. *Am J Public Health* 2011;101(9):1696703.
- Ford JL, Browning CR. Neighborhoods and infectious disease risk: acquisition of chlamydia during the transition to young adulthood. *J Urban Health* 2014;91(1):136–50. [PubMed: 23494850]
- Cooper HL, Hunter-Jones J, Kelley ME, et al. The aftermath of public housing relocations: relationships between changes in local socioeconomic conditions and depressive symptoms in a cohort of adult relocaters. *J Urban Health* 2014;91(2):223–41. [PubMed: 24311024]
- Haley DF, Kramer MR, Adimora AA, et al. Relationships between neighbourhood characteristics and current STI status among HIV-infected and HIV-uninfected women living in the Southern USA: a cross-sectional multilevel analysis. *Sex Transm Infect* 2017;10.1136/sextrans-2016-052889.
- Adimora AA, Schoenbach VJ, Taylor EM, et al. Sex ratio, poverty, and concurrent partnerships among men and women in the United States: a multilevel analysis. *Ann Epidemiol* 2013;23(11):716–9. [PubMed: 24099690]
- Rodriguez JM, Geronimus AT, Bound J, et al. Black lives matter: Differential mortality and the racial composition of the U.S. electorate, 1970–2004. *Soc Sci Med* 2015;136–137:193–9.
- Navarro V, Muntaner C, Borrell C, et al. Politics and health outcomes. *Lancet* 2006;368(9540):1033–7. [PubMed: 16980120]
- Navarro V, Shi L. The political context of social inequalities and health. *Soc Sci Med* 2001;52(3):481–91. [PubMed: 11330781]
- Massey DS, Denton NA. *American apartheid : segregation and the making of the underclass*. Cambridge, Mass.: Harvard University Press; 1993.
- Massey DS, Russell Sage Foundation. *Categorically unequal : the American stratification system*. New York: Russell Sage Foundation; 2007.
- Blakely TA, Kennedy BP, Kawachi I. Socioeconomic inequality in voting participation and self-rated health. *Am J Public Health* 2001;91(1):99–104. [PubMed: 11189832]

16. Mayer M, Kenter R, Morris JC. Partisan Politics or Public-Health Need? An empirical analysis of state choice during initial implementation of the Affordable Care Act. *Politics Life Sci* 2015;34(2): 44–51.
17. Gollust SE, Rahn WM. The Bodies Politic: Chronic Health Conditions and Voter Turnout in the 2008 Election. *J Health Polit Policy Law* 2015;40(6):1115–55. [PubMed: 26447024]
18. Ziegenfuss JK, Davern M, Blewett LA. Access to health care and voting behavior in the United States. *J Health Care Poor Underserved* 2008;19(3):731–42. [PubMed: 18677067]
19. Purtle J Felon disenfranchisement in the United States: a health equity perspective. *Am J Public Health* 2013;103(4):632–7. [PubMed: 23153146]
20. Uggen C, Shannon S, Manza J. State-level estimates of felon disenfranchisement in the United States, 2010.: The Sentencing Project; 2012.
21. Uggen C, Larson R, Shannon S. 6 Million Lost Voters: State-Level Estimates of Felony Disenfranchisement, 2016. Washington, DC: The Sentencing Project; 2016.
22. Uggen C, Manza J. Democratic contraction? Political consequences of felon disenfranchisement in the United States *American Sociological Review* 2002;67(6):777–803.
23. Hessol NA, Weber KM, Holman S, et al. Retention and attendance of women enrolled in a large prospective study of HIV-1 in the United States. *J Womens Health (Larchmt)* 2009;18(10):1627–37. [PubMed: 19788344]
24. Bacon MC, von Wyl V, Alden C, et al. The Women’s Interagency HIV Study: an observational cohort brings clinical sciences to the bench. *Clin Diagn Lab Immunol* 2005;12(9):1013–9. [PubMed: 16148165]
25. Hessol NA, Anastos K, Levine AM, et al. Factors associated with incident self-reported AIDS among women enrolled in the women’s interagency HIV study (WIHS). WIHS Collaborative Study Group. *AIDS Res Hum Retroviruses* 2000;16(12):1105–11.
26. Hessol NA, Schneider M, Greenblatt RM, et al. Retention of women enrolled in a prospective study of human immunodeficiency virus infection: impact of race, unstable housing, and use of human immunodeficiency virus therapy. *Am J Epidemiol* 2001;154(6):563–73. [PubMed: 11549562]
27. Leip D Dave Leip’s Atlas of U.S. Presidential Elections (<http://uselectionatlas.org>). In.
28. Tam Cho WK, Gimpel JG, Dyck JJ. Residential Concentration, Political Socialization, and Voter Turnout. *The Journal of Politics* 2006;68(1):12.
29. O’Rourke N, Hatcher L. A step-by-step approach to using SAS for factor analysis and structural equation modeling: SAS Institute; 2013.
30. Cooper HL, Linton S, Haley DF, et al. Changes in Exposure to Neighborhood Characteristics are Associated with Sexual Network Characteristics in a Cohort of Adults Relocating from Public Housing. *AIDS Behav* 2015;19(6):1016–30. [PubMed: 25150728]
31. Dembo R, Belenko S, Childs K, et al. Individual and community risk factors and sexually transmitted diseases among arrested youths: a two level analysis. *J Behav Med* 2009;32(4):30316.
32. Dyck JJ, Gimpel JG. Distance, Turnout, and the Convenience of Voting. *Social Science Quarterly* 2005;86(3):18.
33. Viera AJ. Odds ratios and risk ratios: what’s the difference and why does it matter? *South Med J* 2008;101(7):730–4. [PubMed: 18580722]
34. Cummings P The relative merits of risk ratios and odds ratios. *Arch Pediatr Adolesc Med* 2009;163(5):438–45. [PubMed: 19414690]
35. Zhang J, Yu KF. What’s the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. *JAMA* 1998;280(19):1690–1. [PubMed: 9832001]
36. Hirai A, Kaufman J. Fixed versus random effects models for multilevel and longitudinal data In: Oakes J, Kaufman J, editors. *Methods in Social Epidemiology*. 2 ed. San Francisco, California: Jossey-Bass; 2017 p. 369–397.
37. Manza J, Uggen C. Punishment and democracy: Disenfranchisement of nonincarcerated felons in the United States. *Perspectives on Politics* 2004;2(3):491–505.

38. Scribner R, Theall KP, Ghosh-Dastidar B, et al. Determinants of social capital indicators at the neighborhood level: a longitudinal analysis of loss of off-sale alcohol outlets and voting. *J Stud Alcohol Drugs* 2007;68(6):934–43. [PubMed: 17960312]
39. Theall KP, Scribner R, Ghosh-Dastidar B, et al. Neighbourhood alcohol availability and gonorrhoea rates: impact of social capital. *Geospat Health* 2009;3(2):241–55. [PubMed: 19440966]
40. Kimball R, Wissner M. Religion, Poverty, and Politics: Their Impact on Women’s Reproductive Health Outcomes. *Public Health Nurs* 2015;32(6):598–612. [PubMed: 25879714]
41. Bernstein S, North A, Schwartz J, et al. State-Level Voting Patterns and Adolescent Vaccination Coverage in the United States, 2014. *Am J Public Health* 2016;106(10):1879–81. [PubMed: 27552268]
42. Law DC, Serre ML, Christakos G, et al. Spatial analysis and mapping of sexually transmitted diseases to optimise intervention and prevention strategies. *Sex Transm Infect* 2004;80(4):294–9. [PubMed: 15295129]
43. Green TC, Pouget ER, Harrington M, et al. Limiting options: sex ratios, incarceration rates, and sexual risk behavior among people on probation and parole. *Sex Transm Dis* 2012;39(6):424–30. [PubMed: 22592827]
44. Khan MR, Doherty IA, Schoenbach VJ, et al. Incarceration and high-risk sex partnerships among men in the United States. *J Urban Health* 2009;86(4):584–601. [PubMed: 19459050]
45. Pouget ER, Kershaw TS, Niccolai LM, et al. Associations of sex ratios and male incarceration rates with multiple opposite-sex partners: potential social determinants of HIV/STI transmission. *Public Health Rep* 2010;125 Suppl 4:70–80.

Research highlights

- County-level voter turnout was inversely associated with STIs among women.
- County-level felony voter disenfranchisement was positively associated with STIs.
- County-level voting age African American residents was not associated with STIs.
- These relationships were independent of county-level socioeconomic advantage.

Table 1.

Distributions of county and participant characteristics among 666 women enrolled in the Women's Interagency HIV Study sites in Alabama, Florida, Georgia, Mississippi, and North Carolina between 2013–2105

Characteristics of participants and counties	Overall n (%) or median (interquartile range)n=666	HIV-Seropositive n (%) or median(interquartile range) n=471	HIV-Seronegative n (%) or median (interquartile range)n=195
Outcomes			
Laboratory-confirmed sexually transmitted infection	76 (11.4)	55 (11.7)	21 (10.8)
Chlamydia	8 (1.2)	4 (0.8)	4 (2.0)
Gonorrhea	6 (0.9)	5 (1.1)	1 (0.5)
Trichomoniasis	44 (6.6)	32 (6.8)	12 (6.1)
Syphilis	21 (3.2)	17 (3.6)	4 (2.0)
County-level characteristics			
Percentage voter turnout	57.0 (53.8–60.2)	56.7 (53.8–60.2)	57.5 (53.8–60.2)
Percentage felony voter disenfranchisement	3.9 (1.6–5.6)	3.9 (1.4–6.1)	3.9 (1.7–5.1)
Percentage of African American residents 18 and older	40.5 (20.2–48.9)	40.5 (20.4–52.5)	40.5 (24.8–42.9)
Socioeconomic advantage ¹	-0.18 (-0.7–0.8)	-0.5 (-0.8–0.8)	0.4 (-0.7–0.8)
Median household income ¹	\$50,596 (\$43,099-\$56,642)	\$45,239 (\$43,099-\$56,642)	\$52,038 (\$44,409-\$56,642)
Percentage of residents >16 who are employed ¹	58.8 (55.5–60.2)	56.6 (55.2–60.2)	59.8 (55.6–61.9)
Percentage of residents >25 with a college degree or greater ¹	30.3 (26.4–48.3)	30.3 (26.4–45.6)	40.3 (27.5–48.3)
Population density (residents per square mile)	1140.4 (496.7–1836.4)	1140.4 (283.3–1836.4)	1140.4 (592.9–1836.4)
Participant-level characteristics			
Age in years	44.5 (35.8–51.5)	44.7 (36.5–51.8)	43.6 (34.1–50.4)
Non-Hispanic African American ²	554 (83.2)	388 (82.4)	166 (85.13)
Annual household income <\$18,000 ³	443 (68.7)	325 (70.8)	118 (63.4)
Less than a high school education or equivalent	202 (30.3)	147 (31.2)	55 (28.2)
HIV-seropositive	471 (70.7)	--	--

¹Statistically significantly different distribution by HIV status ($p < 0.05$) using Wilcoxon median tests.

²Race/ethnicity missing for 11 (1.6%) participants.

³Annual household income missing for 21 (3.1%) participants.

Table 2.

Bivariable and multivariable relationships between county-level voter turnout and having a current sexually transmitted infection among women enrolled in Women's Interagency HIV Study sites in Alabama, Florida, Georgia, Mississippi, and North Carolina between 2013–2105 (n=666)^{1,2}

Characteristics of counties and participants	Bivariable PR (95% CI)	Multivariable Model A aPR (95% CI) ³	Multivariable Model B aPR (95% CI) ⁴	Multivariable Model C aPR (95% CI) ⁵
County-level exposures				
Voter turnout	0.87 (0.77–1.00)	0.84 (0.73–0.96)	--	--
Felony voter disenfranchisement ⁶	1.18 (0.90–1.54)	--	1.89 (1.10–3.24)	--
Voting age African American residents ⁶	1.05 (1.01–1.10)	--	--	1.06 (0.99–1.14)
Socioeconomic advantage ⁷	1.19 (0.97–1.47)	1.33 (0.91–1.95)	1.36 (0.94–1.97)	1.16 (0.85–1.58)
Covariates				
Age ⁸	1.03 (1.00–1.05)	1.01 (0.99–1.03)	1.01 (0.99–1.04)	1.01 (0.99–1.04)
Non-Hispanic African American	1.54 (0.76–3.12)	1.31 (0.68–2.51)	1.17 (0.59–2.31)	1.20 (0.61–2.33)
Annual household income of \$18,000 or less	1.15 (0.76–1.74)	1.02 (0.59–1.75)	1.06 (0.61–1.86)	1.05 (0.60–1.82)
Less than a high school education or equivalent	1.26 (0.78–2.06)	1.14 (0.69–1.89)	1.21 (0.75–1.96)	1.22 (0.76–1.97)
HIV-seropositive	1.08 (0.70–1.68)	1.17 (0.72–1.92)	1.18 (0.72–1.93)	1.17 (0.71–1.94)
Population density ⁹	1.00 (1.00–1.01)	1.00 (0.99–1.00)	1.00 (1.00–1.01)	1.00 (0.99–1.00)
Model fit				
QIC	--	458.70	457.75	460.99
QICu	--	458.27	456.68	459.71

¹We modeled bivariable and multivariable relationships using generalized estimating equations with a binomial distribution and log link (to estimate prevalence ratios [PRs] and 95% confidence intervals [CIs]), and an exchangeable correlation structure in SAS 9.4.

²Multivariable models restricted to participants with no missing covariates (n=632).

³Model assessing multivariable relationships between county-level voter turnout, controlling for county-level socioeconomic advantage and other individual and area-level covariates.

⁴Model assessing multivariable relationships between county-level felony voter disenfranchisement, controlling for county-level socioeconomic advantage and other individual and area-level covariates.

⁵Model assessing multivariable relationships between county-level voting age African American residents, controlling for county-level socioeconomic advantage and other individual and area-level covariates.

⁶Per 4 percentage points.

⁷Per standard deviation.

⁸Per year.

⁹Per 10 residents per square mile.