Plans to Evaluate Recommendations for Booster Doses of Yellow Fever Vaccine

Joseph A. Bocchini, Jr., M.D. ACIP, Workgroup Chair

Japanese Encephalitis and Yellow Fever Vaccines Workgroup

October 24, 2013



National Center for Emerging and Zoonotic Infectious Diseases

Division of Vector-borne Diseases

Yellow fever (YF) vaccine in the U.S.

- Live-attenuated YF vaccine (YF-VAX[®]) is the only YF vaccine licensed for use in U.S.
- Vaccine recommended for people residing in or traveling to YF endemic areas
- In accordance with International Health Regulations, ACIP recommends a booster dose every 10 years
- ACIP recommendations for use of YF vaccine were updated in 2009 and published in 2010

WHO Strategic Advisory Group of Experts (SAGE) Recommendations for YF Vaccine

- In April 2013, based on systematic evidence review, SAGE concluded:
 - Single dose of YF vaccine sufficient to confer sustained immunity and lifelong protection against YF disease
 - Booster doses of YF vaccine are not needed

Countries can still require YF vaccine booster doses under International Health Regulations

JE and YF Vaccine Workgroup: Plans

- Restructure JE Vaccine Workgroup to form JE and YF Vaccines Workgroup
- Review long-term immunogenicity and vaccine failure data for YF vaccine
- Develop recommendations regarding YF vaccine booster doses in U.S. travelers and laboratory workers
- Anticipate ACIP vote on YF vaccine booster dose in 2014

JE and YF Vaccines Workgroup Members

<u>ACIP members</u> Joseph Bocchini (Chair) Lorry Rubin

Liaison representatives Cody Meissner (AAP) Robert Schechter (AIM)

<u>CDC Leads</u> Erin Staples (NCEZID/DVBD) Marc Fischer (NCEZID/DVBD) Ex Officio members Doran Fink (FDA) Jesse Geibe (DoD) Michael Holbrook (NIH) Lewis Markoff (FDA) Pat Repik (NIH)

Invited consultants Elizabeth Barnett Alan Barrett Lin Chen David Shlim