

HHS Public Access

Am J Health Promot. Author manuscript; available in PMC 2019 March 12.

Published in final edited form as:

Author manuscript

Am J Health Promot. 2013; 27(6): 410–416. doi:10.4278/ajhp.100827-QUAN-285.

Neighborhood Environment and Urban Schoolchildren's Risk for Being Overweight

Damiya Whitaker, PsyD, MA,

Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland.; National Institute on Drug Abuse, Intramural Research Program, Clinical Pharmacology and Therapeutics Research Branch, National Institutes of Health Biomedical Research Center, Baltimore, Maryland.

Adam J. Milam, PhD, MHS,

Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland.

Camelia M. Graham, PhD, MSPH,

Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland.

Michele Cooley-Strickland, PhD,

Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland.; Semel Institute/NPI, Center for Culture and Health, University of California, Los Angeles, California.; Center for Health Disparities Solutions, Johns Hopkins University, Baltimore, Maryland.

Harolyn M. Belcher, MD, MHS, and

Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland.; Department of Pediatrics, Johns Hopkins University School of Medicine, Baltimore Maryland. Kennedy Krieger Institute, Family Center, Baltimore, Maryland.

C. Debra Furr-Holden, PhD

Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland.; Center for Health Disparities Solutions, Johns Hopkins University, Baltimore, Maryland.

Abstract

Purpose.—Child and adolescent obesity is increasingly prevalent and predisposes risk for poor physical and psychosocial health. Physical and social factors in the environment, such as neighborhood disorder, may be associated with childhood obesity. This study examines the association between living in a disordered neighborhood and being overweight among a sample of urban schoolchildren.

Send reprint requests to Damiya Whitaker, PsyD, MA, Johns Hopkins University Bloomberg School of Public Health, Department of Mental Health, DIVE Studies Laboratory, 111 Market Place, Suite 850, Baltimore, MD 21202; dwhitake@jhsph.edu.

Manuscript format: research; Research purpose: test of association (odds) cross-sectional; Study design: experimental; Outcome measure: overweight/obesity; Setting: school/community; Health focus: weight control; Strategy: built environment; Target population age: youth; Target population circumstances: race/ethnicity, geographic location, income/education level

Design.—Baseline interview data, including height, weight, and hip circumference, were obtained from 313 elementary school-aged participants in a community-based epidemiologic study.

Setting.-The setting was Baltimore, Maryland, a large metropolitan city.

Subjects.—Subjects were elementary school students ages 8 to 12 years.

Measures.—To assess neighborhood characteristics, independent evaluators conducted objective environmental assessments using the Neighborhood Inventory for Environmental Typology instrument on the block faces (defined as one side of a city block between two intersections) where the children resided.

Analysis.—Logistic regression models with generalized estimating equations were used to examine the association between neighborhood disorder and children being overweight.

Results.—Neighborhood disorder showed a trend toward a statistically significant association with being overweight during childhood (odds ratio [OR], 1.03; confidence interval [CI], .99–1.07; p = .07) in the unadjusted model. Gender was significantly associated with being overweight, with female gender increasing the odds of being overweight by 50% in the sample (OR, 1.50; CI, 1.18–1.92; p < .01). After controlling for race, age, and comparative time spent on a sport, multivariable analyses revealed that gender (adjusted odds ratio [AOR], 2.42; CI, 1.63–3.59; p < .01) and neighborhood disorder (AOR, 1.09; CI, 1.03–1.15; p < .01) were associated with being overweight. Further, an examination of interactions revealed girls (AOR, 2.40; CI, 1.65–3.49; p < .01) were more likely to be overweight compared with boys (AOR, 2.20; CI, 1.57–3.11; p < .01) living in neighborhoods with the same level of neighborhood disorder.

Conclusion.—Results suggest neighborhood hazards warrant additional consideration for their potential as obesogenic elements affecting gender-based disparities in weight among urban schoolchildren. Future studies in this area should include longitudinal examinations. (Am J Health Promot 2013;27[6]:410–416.)

Keywords

Youth; Overweight; Gender; Environmental Exposure; Prevention Research

INTRODUCTION

The prevalence of obesity among children and adolescents has tripled since the 1970s.^{1–3} In the United States alone, 32% of children and adolescents ages 2 to 19 years are overweight.⁴ Although obesity has increased significantly throughout the population, sharply higher rates of obesity and being overweight occur among children and adolescents of urban lower socioeconomic status^{5,6} and minority groups.⁴ Weight increases among children have prompted nationwide programs to prevent obesity and obesity-related illnesses among children, including First Lady Michelle Obama's "Let's Move" initiative aimed at improving food quality in schools, increasing healthy food accessibility, and enhancing parents' nutrition and exercise knowledge. The significant impact that obesity has on the physical and psychosocial health of children^{7,8} calls for a change in policy. There is an increased prevalence of insulin resistance, glucose intolerance, advanced growth/early maturity,

hypertension, sleep apnea, and weight stigmatization^{9–11} associated with obesity and being overweight during childhood. Further, these children are at greater risk for ill health persisting into adulthood.¹² Current estimates confirm that obesity is one of the nation's most serious childhood health problems.^{2,3,13} As such, there is a need to examine obesity-related risk^{14–17} and protective factors (e.g., behavioral, environmental, socioeconomic, and cultural) among urban children.

Pioneering work of urban planners like Jane Jacobs observed that the design of physical spaces can create a repelling or an inviting effect.¹⁸ Using those findings, social scientists undertook empirical studies on neighborhood geography and its contribution to obesity and obesity-related disparities. For example, Franco et al.¹⁹ examined associations between neighborhood income and differential availability of healthy foods. Using a "healthy food availability index" and 159 census tracts, it was found that predominantly African-American, lower-income neighborhoods have lower access to healthy, less energy-dense food options (e.g., fruits, vegetables, low-fat foods) because of the types and locations of stores compared with predominantly white, higher-income neighborhoods.¹⁹ In a similar study by Morland et al.,²⁰ the authors provided additional evidence that making healthy dietary choices was difficult for people in lower-income communities because of the differential distribution of quality supermarkets and convenience stores by neighborhood wealth and racial segregation.

Other studies, using primarily adult samples, examined neighborhood physical and social surroundings and obesity outcomes. Research using the socioecologic model with adults noted associations between obesity and residential density, land use patterns,²¹ poor terrain/ sidewalk conditions, fewer community recreational outlets,^{22,23} proximity to fast food restaurants,^{24,25} and reduced access to supermarkets.²⁶ Further, indicators of neighborhood quality, safety, disorder, and perceptions of neighborhood conditions as stressful contributed to decreased healthy nutrition seeking and physical activity engagement^{27,28} among adults, ^{29–36} affecting obesity. Yet, there are few analyses of the association between aspects of the built environment and childhood obesity in published peer-reviewed journals. This is especially true for studies including ethnic minority youth.

Existing studies using youth samples have found correlations between the availability of safe places to play,^{37,38} carbohydrate-rich dietary intake,^{39–41} increased caloric intake, hours spent watching television,^{42,43} physical activity,^{44,45} and childhood obesity. Molnar et al.⁴⁶ showed that an environment (including social disorder and neighborhood safety) can affect children's recreational involvement/physical activity and that unsafe neighborhoods may deprive children of time spent in outdoor play and increase their susceptibility for obesity.⁴⁶ Moreover, disadvantaged African-American and Hispanic youth fare worse than their majority white counterparts regarding their neighborhood's features (e.g., abandoned buildings, poor lighting, poorly maintained sidewalks, and fewer green spaces),⁴⁷ and this pattern of risk also correlates with gender, with the highest obesity rates found among girls. ⁴⁸

Despite these early findings, results do not consistently demonstrate an association between neighborhood safety and disorder and childhood obesity.^{49–51} For example, in a study of the

effects of neighborhood deprivation on strenuous exercise, Ross⁵² found no association between neighborhood disadvantage and walking; however, high levels of walking in lowincome neighborhoods are likely due to greater reliance on public transportation.⁵³ Further, schoolchildren in such neighborhoods are more likely to walk to school.⁵⁴ The conflicting results regarding neighborhood effects on childhood obesity prompted literature reviews to identify conceptual gaps and methodologic limitations. The use of cross-sectional study designs and compositional measures derived from U.S. Census data (e.g., aggregated data used to approximate certain neighborhood characteristics) limits conclusions derived from studies using these designs.¹⁷ Additionally, the lack of theoretical foundations and understanding of interdependency among various built and social environmental variables was frequently identified as a limitation of existing studies.^{55,56}

In more recent years, a new method called the Neighborhood Inventory for Environmental Typology (NIfETy) was devised to objectively measure physical and social disorder in a neighborhood. The NIfETy evaluates both positive and negative features of the social environment and neighborhood structures, with an emphasis on those that are malleable.^{57,58} This study uses the NIfETy method to examine associations between 11 physical and social hazards and being overweight in a sample of urban, primarily African-American, elementary schoolchildren enrolled in an ongoing study.⁵⁹ This study also examined the interaction of gender and environmental determinants of being overweight, because evidence shows that these determinants affect girls differently.^{3,4}

The conceptual model for the current study (Figure) is guided by the socioecologic model of behavior.^{60,61} The socioecologic model of behavior holds that multiple contexts and elements in the environment are interrelated. Given that individuals and their behavior are nested within socioenvironmental influences, the socioecologic model of behavior in this conceptual model illustrates that neighborhood physical and social disorder hazards may restrict child residents' range of behavior (e.g., by limiting safe opportunities for physical activity), which in turn increases the risk of obesity and being overweight.

METHODS

Data Sources

Data from the baseline interviews of 313 schoolchildren enrolled in a longitudinal community-epidemiologic study called the Multiple Opportunities to Reach Excellence (MORE) Project were used for this analysis. Within the MORE Project, six schools were selected from the 55 Baltimore Community Statistical Areas; the Community Statistical Areas were ranked into three violence strata based on the number of homicides per 100,000 residents.⁵⁹ Within each stratum, the two schools with the largest enrollment of third through fifth grade students were recruited for participation. To be eligible for the MORE Project, students had to be ages 8 to 12 years, speak English, and live with an Englishspeaking parent/guardian. A total of 425 students consented to participate in this study. Most of the non-African-American portion of the sample (n = 49; 11%), self-reported their race as biracial or white. Participating schools were incentivized with honorariums, and students and parents received gift cards worth \$5 to \$25.

The MORE Project examined the impact of community violence exposure on the emotional, behavioral, and physical health of urban schoolchildren.⁵⁹ The assessment protocol included a battery of survey measures administered via computer-assisted interview.⁵⁹ Trained research assistants obtained measurements for height (inches), waist (inches), and weight (pounds) while participants were in lightweight clothing and bare feet, with their heads held in a horizontal position and their chins parallel to the floor.

Upon completion of the MORE Project interviews, independent raters made physical disorder observational assessments on the residential block faces of MORE participants, such as deteriorated landscapes (e.g., boarded abandoned buildings, broken windows, graffiti, and trash) and social disorder (e.g., loitering, drug sales, noise, and public drug/ alcohol consumption) using the NIfETy Instrument.^{57, 8} NIfETy assessments were conducted by a pair of trained field raters. The NIfETy includes seven domains: (1) physical layout, (2) dwellings, (3) adult activity, (4) youth activity, (5) physical disorder, (6) social disorder, and (7) violence, alcohol, and other indicators. The NIfETy has good total scale reliability. The internal consistency reliability (ICC) is .84, and the violence, alcohol, and other drug (VAOD) subscale ICC is .71.⁵⁷ It has acceptable reliability across raters (ICC, .67 —.79). Validity metrics are good as well; NIfETy indicators of VAOD exposure correlated strongly with local crime data.⁵⁸ More detailed information regarding the NIfETy instrument has been published.^{57,58,62,63} The NIfETy and MORE data were collected during spring and summer 2007. All data collection was approved by the Johns Hopkins Bloomberg School of Public Health Institutional Review Board.

Measures

Outcome.—Body mass index (BMI) was calculated according to the Centers for Disease Control and Prevention (CDC) BMI guidelines for children and adolescents.⁶⁴

 $BMI = [weight(1b)/height^{2}(in)] \times 703$

The resultant BMI value was plotted on a gender-specific growth chart, which reveals a percentile used to assess overweight and risk of obesity between childhood and adolescence. ⁶⁴ The CDC's BMI-for-age percentile is classified into four categories: underweight (<5th percentile), normal weight (5th to <85th percentile), overweight (85th to <95th percentile), and obese (95th percentile). In this study, youth with underweight BMI values (n = 39) were excluded. Overweight and obese values were categorized as a single group. Overweight among children is defined as at or above the 85th percentile of BMI-for-age.

Control Variables.—Demographic data, including children's self-reported race (African-American or non-African-American), gender, and age in years were included as control variables. Comparative time spent on a sport was obtained via responses to the question: "How much time do you spend on sport A." Responses were scored as "less than others" (=1), "average" (=2), and "more than average" (=3).

Independent Variables/Environmental Predictors of Being Overweight in Childhood.—A total of 21 items were considered for use from the NIfETy.^{65–67}

Exploratory factor analysis techniques identified 11 items that consistently loaded together: structures with broken windows, unboarded abandoned buildings, unmaintained property, trash in open spaces, broken bottles, graffiti, noise, people yelling, public alcohol consumption, drug paraphernalia, and discarded alcohol bottles. The presence or absence of these 11 items on each participant's residential block face (Cronbach $\alpha = .806$) was used to create a neighborhood disorder score, which ranged from 0 to 11 (mean, 5.38; SD, 2.89). To address the validity of the neighborhood disorder score, the score was regressed on self-reported neighborhood safety and the violence stratum based on the number of homicides per 100,000 residents. Neighborhood disorder among children living in the highest- and moderate-violence strata was statistically higher than children living in the lowest-violence strata (odds ratio [OR], 4.5; p < .01). Neighborhood safety was also associated with the neighborhood disorder score (OR, 1.72; p < .01).

Analysis

The analytic sample included 313 children. A total of 425 children were interviewed for the MORE Project; 406 children had valid height and weight data. Additionally, 43 children who were underweight were excluded, and 50 children were missing neighborhood data. The children with missing neighborhood and height/weight data were not statistically different from the children with complete data in terms of age, race, or gender. Logistic regression models were used to measure the association between neighborhood disorder on children's residential block face and overweight status. The data were analyzed using STATA 10.0 (Intercooled STATA 10.0, 2008; Stata-Corp, College Station, Texas). Generalized estimating equations were used to adjust for correlations among students attending the same school (resulting in six clusters). Male gender was used as the reference. Gender-specific estimates for each unit increase in odds of being overweight were obtained using the lincom command in STATA; the model also included an interaction term comparing girls to boys with the same disorder score. This postestimation technique enabled a test of gender as a modifier of the neighborhood disorder effect on being overweight among the children in the sample. All *p* values <.05 were considered statistically significant.

RESULTS

Descriptive Statistics

Table 1 presents the sample statistics and the bivariate correlates of being overweight. The sample mean age was 10 years (male age SD \pm 1.04 years; female age SD \pm 1.07 years). A total of 87% of the sample was African-American (n = 271), which is comparable to the ethnic composition of students in the Baltimore City Public School System.⁶⁸

In the model, 53% of the sample was female (n = 167) and two thirds (n = 201) of the total 313 children had BMIs above normal (BMI-for-age percentile 85%) according to the CDC guidelines.⁶⁴ Overweight status and neighborhood disorder did not vary by gender (χ^2 , p > . 10).

Logistic Regression Results

Table 2 presents both the independent and adjusted odds for being overweight. In the model of independent effects, neighborhood disorder score showed a trend toward an association with being overweight (OR, 1.03; confidence interval [CI], .99–1.07; p = .07). Girls were 1.5 times more likely to be overweight (OR, 1.50; CI, 1.18–1.92; p < .01). No statistically significant associations were found between race, age, and overweight status.

The adjusted logistic regression model included race, comparative time spent on a sport, and age, as well as an interaction term comparing girls and boys with the same disorder score. In the adjusted model, the interactions (female × disorder and female × male) were statistically significant (Table 2). Postestimation commands were used in STATA to provide point estimates and standard errors. Among girls, the odds of being overweight increased 2.4 times for each unit increase in disorder on the block (adjusted odds ratio [AOR], 2.42; CI, 1.63–3.59; p < .01). A similar but weaker effect was seen among boys, with the odds of being overweight increasing 9% for each unit increase in disorder (AOR, 1.09; 1.03–1.15; p < .01). Relative to boys, girls living on disordered blocks were two times more likely to be overweight than their male counterparts when other factors were kept constant (AOR, 2.20; 1.57–3.11; p < .01). There were no race- or age-related variations noted in the model.

Subsidiary Analyses

To assess whether socioeconomic status had an impact on study findings, the effect of free and reduced-price lunch, reported by the teachers and participants and used here as a proxy for socioeconomic status, was examined among a subpopulation of youth with available data (n = 170; 54%). Free/reduced-price lunch status was not a significant predictor of being overweight (OR, 1.08; .61–1.92; p = .79). However, when free lunch status was added as a covariate to the adjusted model, the OR for neighborhood disorder increased (OR, 1.16; p < .01), as did the ORs from the linear combinations of neighborhood disorder and gender.

The proxy for physical activity (comparative time spent on a sport) was assessed as a potential mediator for the association between neighborhood disorder and being overweight. Although the trend (OR, 1.03; p = .07) toward an association between neighborhood disorder and risk for being overweight decreased with the addition of the physical activity variable (OR, 1.03; p = .139), no significant effect was noted.

DISCUSSION

This study investigated the association between urban neighborhood disorder and the risk of being overweight in childhood. Among urban schoolchildren, gender was associated with an increased likelihood of being overweight. An interaction term comparing girls with boys living in the same level of disorder was also associated with increases in overweight status in childhood, indicating that although urban male and female youth are affected by unhealthy weight status, urban neighborhood hazards may impact school-aged girls more than boys. Lastly, neither race nor comparative time spent on a sport was significantly associated with being overweight, after adjusting for demographic characteristics. The results of this study on children are similar to those in the extant adult literature, suggesting that neighborhood

characteristics¹⁴ and physical surroundings encourage obesity, obesity-generating behaviors, ^{15,16} such as sedentary lifestyle and unhealthy eating habits,¹⁷ and poorer physical health status.^{23,27,69, 70}

Evidence from this and other studies tends to support the notion that disorder at the neighborhood level may affect obesity by keeping residents indoors. The lack of significant race effects is not surprising given the work of Powell et al.³⁷ Evidence from their work suggests that the reduced availability of physical activity settings in ethnic minority and socioeconomically disadvantaged communities is the likely culprit for higher weight and lower physical activity among this population. Similarly, the finding of an association between gender and being overweight is consistent with the literature.¹ Between 1999 and 2004, African-American and Hispanic female children and adolescents were significantly more likely to be overweight compared with their white counterparts.¹

The results of this study should be interpreted in view of its limitations. There are exogenous variables potentially related to obesity and neighborhood disorder that were not available for study, including a dietary intake measure. In addition, we did not quantitatively measure physical activity because of a lack of available metrics. Future studies should use a more robust and empirically validated measure for physical activity. Also, data on parental or familial history of obesity were not available and may have strengthened the study. Lastly, this is a cross-sectional study. Longitudinal prospective studies are needed to evaluate the association of neighborhood characteristics, gender, and weight over time.

CONCLUSION

A major criticism of prior research was the degree to which aggregated individual compositional measures limited the investigation of causal links between specific neighborhood features and health outcomes.⁷¹ The current study used an objective measure of neighborhood features, the NIfETy, which enabled a more direct exploration of environmental triggers for being overweight in childhood. Additionally, this work suggests measures for intervention. Intervention designs may benefit from the inclusion of family, school, and/or community involvement in efforts to increase active-recreational orientation for urban youth, particularly those that are gender specific, given that girls' and boys' obesity risk profiles (defined as the social, cultural, and environmental factors influencing obesity for children) differ. Further, strategies might also include increasing the development of safe neighborhood structures that support gender-specific physical activity. In addition, addressing neighborhood-based obesogenic risks in urban communities (e.g., revitalization projects to improve neighborhood quality and reduce crime) may lead to a more healthpromoting environment. Finally, the Child Nutrition Act and campaigns such as the "Let's Move" Initiative must include the importance of neighborhood factors associated with the development and remediation of obesity.

Although this study represents a first exploration, further research is warranted. Obesity and overweight status among urban, ethnic minority children is likely related to neighborhood socioeconomic status, neighborhood disorder, and the availability of safe physical activity settings. Accordingly, efforts to avert, suspend, and reverse this alarming trend may be more

effective if culturally and ecologically relevant, long-term, and gender-specific interventions were employed at the individual, family, school, and/or community levels.

Acknowledgments

This research was supported by awards from the National Institute on Alcoholism and Alcohol Abuse (R01AA015196) to principal investigator C. Debra Furr-Holden, PhD; the National Institute on Drug Abuse (NIDA; R01DA018318); the NIDA Drug Dependence Epidemiology Training Program (T32 DA007292); Kenzie Preston, PhD, the Intramural Research Program, National Institutes of Health, NIDA; the Centers for Disease Control and Prevention (1U49CE000728) to principal investigator Philip Leaf, PhD; and the Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland. The authors would like to acknowledge the MORE Project participants; the MORE Project coordinator; the NIFETy project field supervisor; and the study field rating teams.

References

- 1. Ogden C, Carroll M, Curtin L, et al. Prevalence of overweight and obesity in the United States, 1999–2004. JAMA. 2006;295:1549–1555. [PubMed: 16595758]
- Koplan J, Liverman C, Kraak V. Preventing Childhood Obesity: Health in Balance. Washington, DC: The National Academies Press; 2005.
- 3. National Institute of Environmental Health Sciences. Obesity and the environment. Available at: http://www.niehs.nih.gov/health/docs/obesity-fs.pdf.
- 4. Ogden C, Carroll M, Curtin L, et al. Prevalence of high body mass index in U.S. children and adolescents, 2007–2008. JAMA. 2010;303:242–249. [PubMed: 20071470]
- 5. Shrewsbury V, Wardle J. Socioeconomic status and adiposity in childhood: a systematic review of cross-sectional studies 1990–2005. Obesity. 2008;16:275–284. [PubMed: 18239633]
- 6. Dept of Health & Human Services. The Surgeon General's Call to Action to Prevent and Decrease Overweight and Obesity. Rockville, Md: US Dept of Health and Human Services; 2001.
- Reilly J, Houston-Callaghan K, Donaghey Z, Hammed S. Physical health consequences of child and adolescent obesity In: Crawford D, Jeffery RW, Ball K, Brug J, eds. Obesity Epidemiology – From Aetiology to Public Health. 2nd ed. New York, NY: Oxford University Press; 2010.
- Lobstein T Prevalence and trends in childhood obesity In: Crawford D, Jeffery RW, Ball K, Brug J, eds. Obesity Epidemiology – From Aetiology to Public Health. 2nd ed. New York, NY: Oxford University Press; 2010.
- 9. Weiss R, Dziura J, Burgert T, et al. Obesity and the metabolic syndrome in children and adolescents. N Engl J Med. 2004;350:2362–2374. [PubMed: 15175438]
- Ebbeling C, Pawlak D, Ludwig D. Childhood obesity: public-health crisis, common sense cure. Lancet. 2002;360:473–482. [PubMed: 12241736]
- Dietz W, Gortmaker S. Preventing obesity in children and adolescents. Annu Rev Public Health. 2001;22:337–353. [PubMed: 11274525]
- 12. Must A, Strauss R. Risks and consequences of childhood and adolescent obesity. Int J Obes Relat Metab Disord. 1999;23(suppl 2):S2–S11.
- 13. Dept of Health and Human Services. Healthy People 2010: Understanding and Improving Health. 2nd ed. Washington, DC: US Government Printing Office; 2000.
- Mair J, Mair M. Violence prevention and control through environmental modifications. Annu Rev Public Health. 2003;24:209–225. [PubMed: 12428032]
- 15. Jackson R, Kochtitzky C. Creating a Healthy Environment: The Impact of the Built Environment on Public Health. Washington, DC: Sprawl Watch Clearinghouse Monograph Series; 2001.
- Centers for Disease Control and Prevention. Guidelines for school and community programs to promote lifelong physical activity among young people. MMWR Recomm Rep. 1997;46(RR-6):1– 36.
- Papas M, Alberg A, Ewing R, et al. The built environment and obesity. Epidemiol Rev. 2007;29:129–143. [PubMed: 17533172]
- 18. Jacobs J The Death and Life of Great American Cities. New York, NY: Random House; 1993.

- Franco M, Diez-Roux A, Glass T, et al. Neighborhood characteristics and availability of healthy foods in Baltimore. Am J Prev Med. 2008;35:561–567. [PubMed: 18842389]
- Morland K, Wing S, Diez-Roux A, Poole C. Neighborhood characteristics associated with the location of food stores and food service places. Am J Prev Med. 2002;22: 23–29. [PubMed: 11777675]
- 21. Frank L, Andresen M, Schmidt T. Obesity relationships with community design, physical activity and time spent in cars. Am J Prev Med. 2004;27:87–96. [PubMed: 15261894]
- Giles-Corti B, MacIntyre S, Clarkson J, et al. Environmental and lifestyle factors associated with overweight and obesity in Perth, Australia. Am J Health Promot. 2003;18:93–102. [PubMed: 13677967]
- 23. Ewing R Can the physical environment determine physical activity levels? Exerc Sport Sci Rev. 2005;33:69–75. [PubMed: 15821427]
- 24. Inagami S, Cohen D, Brown A, Asch S. Body mass index, neighborhood fast food and restaurant concentration and car ownership. J Urban Health. 2009;86: 683–695. [PubMed: 19533365]
- 25. Li F, Harmer P, Cardinal B, et al. Obesity and the built environment: does the density of neighborhood fast-food outlets matter? Am J Health Promot. 2009;23: 203–209. [PubMed: 19149426]
- Inagami S, Cohen D, Finch B. You are where you shop: grocery store locations, weight and neighborhoods. Am J Prev Med. 2006;31:10–17. [PubMed: 16777537]
- Handy S, Boarnet M, Ewing R, Killingsworth R. How the built environment affects physical activity: views from urban planning. Am J Prev Med. 2002;23(2 suppl):64–73. [PubMed: 12133739]
- 28. Centers for Disease Control and Prevention. Physical Activity and Health: A Report of the Surgeon General. Atlanta, Ga: US Dept of Health and Human Services; 1996.
- Harrison R, Gemmell I, Heller R. The population effect of crime and neighborhood on physical activity: an analysis of 15, 461 adults. J Epidemiol Community Health. 2007;61:34–39. [PubMed: 17183012]
- Hoehner C, Brennan-Ramirez L, Elliott M, et al. Perceived and objective environmental measures and physical activity among urban adults. Am J Prev Med. 2005;28:105–116. [PubMed: 15694518]
- Suminski R, Poston W, Petosa R, et al. Features of the neighborhood environment and walking by U.S. adults. Am J Prev Med. 2005;28:149–155. [PubMed: 15710269]
- Humpel N, Owen N, Iverson D, et al. Perceived environment attributes, residential location, and walking for particular purposes. Am J Prev Med. 2004;26:119–125. [PubMed: 14751322]
- Moudon A, Lee C. Walking and bicycling: an evaluation of environmental audit instruments. Am J Health Promot. 2003;18:21–37. [PubMed: 13677960]
- 34. Sampson R The neighborhood context of well-being. Perspect Biol Med. 2003;46(suppl):S53–S64. [PubMed: 14563074]
- Centers for Disease Control and Prevention. Neighborhood safety and the prevalence of physical inactivity—selected states, 1996. MMWR Morb Mortal Wkly Rep. 1999;48:143–146. [PubMed: 10077460]
- Watts R, Rabow J. Alcohol availability and alcohol related problems in 213 California cities. Alcohol Clin Exp Res. 1983;7:47–58. [PubMed: 6342449]
- 37. Powell LM, Slater S, Chaloupka F. The relationship between community physical activity settings and race, ethnicity, and socioeconomic status. Evid Based Pract Med. 2004;1:135–144.
- Sallis J, McKenzie T, Elder J, et al. Factors parents use in selecting play spaces for young children. Arch Pediatr Adolesc Med. 1997;151:414–417. [PubMed: 9111442]
- Popkin B, Duffey K, Gordon-Larsen P. Environmental influences on food choice, physical activity and energy balance. Physiol Behav. 2005;86:603–613. [PubMed: 16246381]
- French SA, Story M, Jeffery RW. Environmental influences on eating and physical activity. Annu Rev Public Health. 2001;22:309–335. [PubMed: 11274524]
- Grundy SM. Multifactorial causation of obesity: implications for prevention. Am J Clin Nutr. 1998;67(suppl):563S–572S. [PubMed: 9497171]

- 42. Wiecha J, Peterson K, Ludwig D, et al. When children eat what they watch: impact of television viewing on dietary intake in youth. Arch Pediatr Adolesc Med. 2006;160:436–442. [PubMed: 16585491]
- Delva J, Johnston L, O'Malley P. The epidemiology of overweight and related lifestyle behaviors: racial/ethnic and socioeconomic status differences among American youth. Am J Prev Med. 2007;33(4 suppl):S178–S186. [PubMed: 17884566]
- 44. Anderson PM, Butcher KE. Childhood obesity: trends and potential causes. Future Child. 2006;16:19–45.
- 45. Goran M, Reynolds K, Lindquist C. Role of physical activity in the prevention of obesity in children. Int J Obes Relat Metab Disord. 1999;23(suppl 3):S18–S33.
- Molnar B, Gortmaker S, Bull F, Buka S. Unsafe to play? Neighborhood disorder and lack of safety predict reduced physical activity among urban children and adolescents. Am J Health Promot. 2004;18:378–386. [PubMed: 15163139]
- 47. Kumanyika S, Grier S. Targeting interventions for ethnic minority and low-income populations. Future Child. 2006;16:187–207. [PubMed: 16532664]
- 48. Hedley A, Ogden C, Johnson C, et al. Prevalence of overweight and obesity among US children, adolescents, and adults, 1999–2002. JAMA. 2004;291:2847–2850. [PubMed: 15199035]
- Powell L, Bao Y. Food prices, access to food outlets and child weight. Econ Hum Biol. 2009;7:64– 72. [PubMed: 19231301]
- 50. Sturm R, Datar A. Body mass index in elementary school children, metropolitan area food prices and food outlet density. Public Health. 2005;119:1059–1068. [PubMed: 16140349]
- Gomez J, Johnson B, Selva M, Sallis J. Violent crime and outdoor physical activity among innercity youth. Prev Med. 2004;39:876–881. [PubMed: 15475019]
- Ross CWalking, exercising, and smoking: does neighborhood matter? Soc Sci Med. 2000;51:265– 274. [PubMed: 10832573]
- 53. Yen I, Kaplan G. Poverty area residence and changes in physical activity level: evidence from the alameda county study. Am J Public Health. 1998;88:1709–1712. [PubMed: 9807543]
- 54. Rossen LM, Pollack KM, Curriero FC, et al. Neighborhood incivilities, perceived neighborhood safety, and walking to school among urban-dwelling children. J Phys Act Health. 2011;8:262–271. [PubMed: 21415453]
- 55. Davison K, Lawson C. Do attributes in the physical environment influence children's physical activity? A review of the literature. Int J Behav Nutr Phys Act. 2006;3:19. [PubMed: 16872543]
- 56. Ferreira I, van der Horst K, Wendel-Vos W, et al. Environmental correlates of physical activity in youth a review and update. Obes Rev. 2007;8:129–154. [PubMed: 17300279]
- Furr-Holden CD, Smart M, Pokorni J, et al. The NIfETy method for environmental assessment of neighborhood-level indicators of violence, alcohol, tobacco, and other drug exposure. Prev Sci. 2008;9:245–255. [PubMed: 18931911]
- 58. Furr-Holden CD, Campbell KD, Milam AJ, et al. Measuring indicators of violence, alcohol, tobacco, and other drug typology (NIfETy): an environmental assessment tool for measuring indicators of violence, alcohol, tobacco, and other drug exposures. Eval Rev. 2010;34:159–184. [PubMed: 20479211]
- Cooley-Strickland M, Quille T, Griffin R, et al. Community violence and youth: affect, behavior, substance use, and academics. Clin Child Fam Psychol Rev. 2009;12: 127–156. [PubMed: 19472053]
- 60. Wicker A An Introduction to Ecological Psychology. Pacific Grove, Calif: Brooks/Cole; 1979.
- Sallis J, Owen N. Ecological models In: Glanz K, Lewis FM, Rimer BK, eds. Health Behavior and Health Education: Theory, Research, and Practice. 2nd ed. San Francisco, Calif: Jossey-Bass; 1996:403–424.
- 62. Whitaker D, Graham C, Furr-Holden CD, et al. Neighborhood disorder and incarceration history among urban substance users. J Correct Health Care. 2011;17:309–318. [PubMed: 21969040]
- 63. Milam A, Furr-Holden CD, Whitaker D, et al. Neighborhood environment and internalizing problems in African American children. Community Ment Health J. 2012;38:39–44.

- 64. Centers for Disease Control and Prevention. CDC growth charts: United States. Available at: http://www.cdc.gov/growthcharts/. Accessed December 11, 2009
- 65. Perkins D, Brown B, Taylor R. The ecology of empowerment: predicting participation in community organizations. J Soc Issues. 1996;52:85–110. [PubMed: 15156863]
- 66. Cohen D, Farley T, Mason K. Why is poverty unhealthy? Social and physical mediators. Soc Sci Med. 2003;57:1631–1641. [PubMed: 12948572]
- 67. Sampson R, Raudenbush S, Earls F. Neighborhoods and violent crime: a multilevel study of collective efficacy. Science. 1997;227:918–923.
- 68. Baltimore City Public Schools. Baltimore city public schools: what drives us. Available at: http://www.baltimorecityschools.org/news/pdf/fastfacts2008_09.pdf. Accessed December 11, 2009
- 69. Burdette H, Whitaker R. Neighborhood playgrounds, fast food restaurants, and crime: relationships to overweight in low-income preschool children. Prev Med. 2004;38:57–63. [PubMed: 14672642]
- Ford J, Beveridge A. Bad neighborhood, fast food, sleazy businesses and drug dealers: relations between the location of licit and illicit businesses in the urban environment. J Drug Issues. 2002;34:51–76.
- Mujahid S, Diez-Rouz A, Morenoff J, et al. Neighborhood characteristics and hypertension. Epidemiology. 2008;19: 590–598. [PubMed: 18480733]

SO WHAT? Implications for Health Promotion Practitioners and Researchers

What is already known on this topic?

There is a growing body of research documenting associations between physical and social factors in the environment and increased risk of obesity.

What does this article add?

The current investigation adds to existing literature by providing preliminary evidence, via objective neighborhood assessment, that links the interaction between living in a distressed neighborhood (e.g., crime, violence, substance sales/use) and gender to the overweight status of urban youth.

What are the implications for health promotion practice and research?

The use of tools like the NIfETy instrument can expand health promotion and research efforts by providing appropriate methodology for taking the impact of ecologic settings (e.g., neighborhood-level disadvantage) into account when studying obesity in childhood.



Figure.

Conceptual Model Linking Neighborhood Physical Disorder and Childhood Obesity

Author Manuscript

Variable	Total No. (%) $(N = 313)$	Male, No. (%)	Female, No. (%)
Total	313	146 (46.6)	167 (53.4)
Normal weight	112 (35.8)	45 (30.8)	67 (40.1)
Overweight	201 (64.2)	101 (69.2)	100 (59.9)
African-American	271 (86.6)	128 (87.7)	143 (85.6)
Non-African-American	42 (13.4)		
Neighborhood disorder	I	5.35 (2.9)	5.41 (2.9)
Mean age, y (SD)	I	10.08 (1.04)	10.06 (1.07)
<avg< td=""><td>I</td><td>26 (17.8)</td><td>52 (31.1)</td></avg<>	I	26 (17.8)	52 (31.1)
AVG	I	48 (32.9)	45 (26.9)
>AVG	I	72 (49.3)	70 (41.9)

* Reference groups: male gender, African-American race, overweight; comparative time spent on sport (average [AVG]).

Author Manuscript

Whitaker et al.

Independent and Adjusted Effects of Covariates and Overweight vs. Normal Body Mass Index $(BMI)^*$

		Logistic 1	Regression	
	BMI	Overwe	ight vs. Normal	
Variable	OR (CI)	d	AOR (CI)	d
Female	1.50 (1.18–1.92)	<0.01	2.42 (1.63–3.59)	<0.01
Non-African-American	1.12 (0.68–1.85)	0.65	1.12 (0.64–1.96)	0.69
Neighborhood disorder	1.03 (0.99–1.07)	0.07	1.09 (1.03–1.15)	<0.01
Female $ imes$ disorder	I		$2.40~(1.65{-}3.49)^{\uparrow}$	<0.01
Female $ imes$ male	Ι	I	$2.20(1.57{-}3.11)^{\ddagger}$	<0.01
Age, y	1.01 (0.72–1.42)	0.935	1.00 (0.72–1.39)	0.99
AVG	0.65 (0.41–1.02)	0.06	0.69 (0.43–1.09)	0.11
>AVG	0.81 (0.50–1.30)	0.38	0.84 (0.53–1.35)	0.47
* AOR indicates adjusted o	odds ratio; OR, odds	ratio; AV	G, average.	

 ${}^{\sharp}$ djiusted odds for girls compared with boys with same neighborhood disorder score.

 \dot{f}^{t} Adjusted odds for girls being overweight.