



Published in final edited form as:

Sex Transm Dis. 2014 March ; 41(3): 151–157. doi:10.1097/OLQ.0000000000000100.

An Assessment of the *GYT: Get Yourself Tested* Campaign: An Integrated Approach to Sexually Transmitted Disease Prevention Communication

Allison L. Friedman, MS*, Kathryn A. Brookmeyer, PhD*, Rachel E. Kachur, MPH*, Jessie Ford, MS*, Matthew Hogben, PhD*, Melissa A. Habel, MPH*, Leslie M. Kantor, MPH†, Elizabeth Clark, MA†, Jamie Sabatini, MSW‡, and Mary McFarlane, PhD*

*Division of STD Prevention, Centers for Disease Control and Prevention, Atlanta, GA

†Division of Communications, New York City, NY

‡Division of Planned Parenthood Federation of America, New York City, NY

Abstract

Background: Youth in the United States bear a disproportionate burden of sexually transmitted diseases (STDs). Stigma, misconceptions, and access challenges keep many from getting tested or treated. The *GYT: Get Yourself Tested* campaign was launched in 2009 to reduce stigma and promote STD communication and testing. This evaluation sought to assess the first 2 years of campaign engagement and associations with STD testing among youth.

Methods: Campaign engagement with select GYT on-the-ground events, social media sites, and STD testing locator tools was measured through process/media tracking metrics. Sexually transmitted disease testing patterns were assessed using data from Planned Parenthood affiliates (2008–2010) and national trend data from clinics participating in national infertility prevention activities (2003–2010).

Results: On-the-ground events reached an estimated 20,000 youth in 2009 and 52,000 youth in 2010. Across 2009 to 2010, GYT's Facebook page gained 4477 fans, Twitter feed gained 1994 followers, and more than 140,000 referrals were made to the STD testing locator. From April 2008 to 2010, there was a 71% increase in STD testing and a 41% increase in chlamydia testing at reporting Planned Parenthood affiliates (representing ~118 health centers). Chlamydia case positivity rates during this period were stable at 6.6% (2008) and 7.3% (2010). Trend data indicate that testing was higher in spring 2009 and 2010 compared with other periods during those years; this pattern is commensurate with STD Awareness Month/GYT activities.

Conclusions: Data quality is limited in a manner similar to many STD prevention efforts. Within these limitations, evidence suggests that GYT reaches youth and is associated with increased STD testing.

Correspondence: Allison L. Friedman, MS, Centers for Disease Control and Prevention, 1600 Clifton Rd, NE, Mail Stop E-44, Atlanta, GA 30333. afriedman@cdc.gov.

Conflicts of interest and sources of funding: None declared.

Notes: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

There are an estimated 20 million incident sexually transmitted diseases (STD) annually, half of which occur among youth aged 15 to 24 years.¹ However, many youth do not receive the services needed to detect and treat STDs and prevent further transmission.² Numerous individual, social, and structural challenges prevent STD testing and treatment among youth. These include a culture of silence and stigma, which not only inhibits access to services but also hinders STD information seeking, open communication, and STD disclosure.^{3–6} A Centers for Disease Control and Prevention (CDC) study released in March 2008 indicated that 1 in 4 teen girls had an STD.⁷ In response, the national *GYT: Get Yourself Tested* campaign was launched in April 2009 (STD Awareness Month) to promote STD testing among sexually active youth 25 years and younger.

Campaign Background and Development

GYT is produced through a partnership between MTV, the Kaiser Family Foundation, and Planned Parenthood Federation of America (PPFA). The CDC has provided assistance to ensure scientific accuracy of GYT's health information. Each of these agencies mobilizes its media, communication, and public relations resources through teams of 2 to 6 existing staff members.³ Campaign strategies were informed by formative research^{4,8,9} and key constructs of the Health Belief Model¹⁰ and the Theory of Planned Behavior.¹¹ Formative research revealed that many youth are unaware of the asymptomatic nature and pervasiveness of STDs, that routine STD screening is advised, or that confidential, noninvasive, and youth-friendly services are available. Youth often do not perceive themselves to be at risk for STDs and fear various aspects of STD testing (e.g., uncomfortable procedures and what others will think and say).^{4,9} Although providers, family members, friends, and partners emerged as important intermediaries for reaching youth, opportunities to engage youth in discussions about STDs are often missed in clinical, school, and home settings.¹² Campaign messages were therefore developed to increase STD awareness and perceived risk, reduce fear and stigma, and promote open communication with sex partners and health care providers.

GYT uses television, Web, print, short message service (SMS), and on-the-ground efforts to reach youth with information and link them to testing. The hub of the campaign is the Web site www.GYTNOW.org (Fig. 1). Through this site, as well as an SMS locator tool, youth can access nearby, free/low-cost, and teen-friendly STD testing services (www.findSTDtest.org). National promotions and celebrity endorsements further incentivize testing. The GYT campaign runs year-round, with refreshed messaging each April. GYT toolkits containing promotional materials are provided each year to facilitate state and local efforts.

In its first 2 years, the campaign primarily drove youth to the Web site, with a limited presence on Facebook and Twitter. Sexually transmitted disease testing was promoted across platforms and through partners' on-the-ground efforts. In year 1 (2009), on-the-ground activities relied predominantly on local Planned Parenthood health centers and advocates. By year 2 (2010), the campaign had expanded its partner base to include school-based health centers, state and local health departments, community organizations, colleges and

^aWhen available, GYT products may be supported through additional industry funding, obtained through private sector partnerships.

universities, and high schools. GYT toolkit distribution increased from 1500 in year 1 to 4000 in year 2. Evidence from select local sites suggests that on-the-ground GYT efforts are associated with increases in testing at participating health centers, schools, and community organizations.^{13–15} A 2010 consumer mail survey of teens indicated that awareness of the GYT campaign had reached 18.3%; some respondents (10%–21%) reported that GYT prompted information seeking or discussions about STDs/testing with a sex partner, family member, friend, or provider.¹⁶ However, this survey assessed only a subset of GYT's intended audience and was unable to link reported campaign exposure to behavioral outcomes.

This article presents the first available national evaluation of GYT-associated youth engagement and STD testing during the campaign's first 2 years. Two major factors have inhibited a rigorous evaluation. First, GYT is produced through existing, in-kind resources from key partners and various state and local organizations. Second, national dissemination precluded randomized approaches to testing community-level effects (exposure cannot be controlled). Instead, we have tracked campaign reach and effectiveness through available data sources. We have measured changes in STD testing at select partner health centers nationwide, observed national testing patterns during that same period, and examined national trend data to assess if patterns of testing cohered with GYT's peak promotion period and STD Awareness Month activities (April). Although each data source is limited with respect to establishing effectiveness, a consistent pattern of GYT's "fingerprints" across multiple independent data sources provides some convergent validity for the campaign. Evaluation findings help to identify challenges and successes that can inform other sexual health prevention efforts.

MATERIALS AND METHODS

Three data sources were used to assess the first 2 years of campaign engagement and associations with STD testing and communication. These include event and media tracking metrics, STD patient data from PPFA health centers during the campaign's major promotion period (April), and national trend data from clinics participating in national infertility prevention activities.

Campaign Tracking (Media Metrics)

Campaign reach and engagement in 2009 and 2010 were tracked through youth participation at Planned Parenthood–affiliated events in April, GYT Facebook fans, and GYT Twitter followers. Tracking metrics also gauged the GYT pledges made during a collegiate Facebook contest during the month of April 2010, and the number of GYT badges unlocked during a Foursquare^b promotion during the month of September 2010. In addition, GYT traffic to the STD testing locator was measured.

^bFoursquare is a location-based social networking Web site for mobile device.

Patient and STD Testing Data from Planned Parenthood Affiliates

Sexually transmitted disease testing was assessed using data from Planned Parenthood, the campaign's primary health service provider. In April 2009 and 2010, 69 affiliates^c (representing approximately 850 health centers) consistently reported the number of STD tests run, the proportion of positive tests (test positivity), and patient demographic data. A subset of 9 Planned Parenthood affiliates (representing approximately 118 health centers dispersed across the United States) reported STD testing in April 2008 (precampaign), allowing for a limited pre-post comparison. These data were compared against a national-level data set from family planning settings.¹⁷

National Trend Data

To assess the extent to which GYT may affect national testing behaviors, we examined national trend data in chlamydia testing from 2003 to 2010. Data were provided quarterly to CDC by more than 6500 clinics participating in national infertility prevention activities. These data permit an examination of quarterly testing patterns before and after GYT's peak promotion period.

Data Analysis

Data analyses are descriptive for campaign media and event tracking, testing trends at Planned Parenthood affiliates, and national trend data in chlamydia testing. For the subset of 9 Planned Parenthood affiliates that reported 3 years of data, researchers calculated the change from year to year to examine affiliate-level trends from April 2008 (precampaign) through April 2010, comparing the proportion of affiliates reporting testing increases against a null hypothesis of random changes and an alternative of 5% increases^d (guided by chlamydia test numbers in family planning clinics across 2008–2010).¹⁷ To examine trends at the affiliate level on a larger scale, the percent change in STD tests performed by 69 Planned Parenthood affiliates in April 2009 and 2010 are described. Analyses used a prepost evaluation design to examine changes in STD testing before and during the period of the GYT campaign. Percent changes in testing patterns and patient demographics as well as the odds of observing a change from 2008 to 2010 against the null hypothesis of no change were used.

^cEach locally governed affiliate consists of one or more health centers within a defined geographical area and is separately incorporated.

^dFor any year-to-year comparison, the 9 affiliates present 512 combinations of increases and decreases or ties. The probability of all 9 affiliates reporting increases is $1/512 = 0.002$; the probability of reporting 8 increases and 1 decrease/tie is $9/512 = 0.018$, and so on. Because we are testing the odds of seeing a distribution of testing increases as or more extreme than the observed pattern, the composite *P* value for 8 increases and 1 decrease or tie = $0.18 + 0.002 = 0.02$. The calculations are analogous to tossing a coin and calculating the probability of the observed pattern if the expected value is 50% heads. For the "no-change" hypothesis, the 2010 observed values are compared with the 2008 baseline; for the 5% model, the 2010 observed values are compared with the 2008 baseline $\times 1.05$.

RESULTS

Campaign Tracking (Events and Media Metrics)

In year 1, Planned Parenthood health centers and advocates hosted 450 education and outreach events, reaching an estimated 20,000 youth. Year 2 on-the-ground efforts included 745 events, reaching almost 52,000 youth.

In year 1, the GYT Facebook page gained 2726 likes (fans) and GYT's Twitter feed gained 1445 followers. A total of 83,404 referrals were made to the STD testing locator from the GYT Web site or SMS code in that same year. At the end of the second year, there were a total of 4477 Facebook fans and 1994 Twitter followers. The number of referrals in year 2 dropped to 61,119, but it should be noted that the testing locator was unavailable to clients of a large mobile network carrier for a portion of this period due to contractual issues.

GYT was also promoted through a Facebook "Campus Challenge" in April 2010, in which 1365 students from 167 schools pledged to get tested during the month. A one-time promotion on Foursquare during September 2010 resulted in 4819 GYT badges unlocked.

Patient and STD Testing Data from Planned Parenthood Affiliates

Three-Year STD Patient and Testing Trends (2008–2010).—Data from the 9 Planned Parenthood affiliates from baseline year through the 2 implementation years indicate increases between 2008, 2009, and 2010. There was a 71.0% increase in patients receiving STD testing in April 2010, compared with the same period in 2008 (Table 1). The increases in testing between 2008 and 2010 were in populations deemed most vulnerable to STD infection, including African Americans, Latinos, and people living at or below 150% of the federal poverty level.¹⁸ Increases were also seen among youth younger than 25 years, GYT's target audience (Table 1). Chlamydia testing and test positivity were of particular interest to the campaign, given chlamydia's high prevalence, potential health consequences, and annual testing recommendations for young women. There was an increase in chlamydia testing of 41.2% among women and 49.9% among men between April 2008 and 2010 (Table 2). Chlamydia positivity from these tests ranged between 5.1% in 2008 and 6.6% in 2010 (Table 2).

Two-Year STD Patient and Testing Trends (2009–2010).—Table 3 presents the total number of STD tests performed by 69 reporting Planned Parenthood affiliates in April 2009 and 2010. Planned Parenthood affiliates tested a total of 106,948 patients for STDs in April 2009, compared with 120,370 patients in 2010, a 12.5% increase. The increases in testing were greater in populations most affected by STDs, including African Americans, and for women living at or below 150% of the federal poverty level, although slight decreases were observed in STD testing among white men. Increases were also seen among youth younger than 25 years. Sexually transmitted disease test positivity rates (chlamydia, Table 4) were 5.5% in 2009 and 6.5% in 2010, similar to the national median.¹⁹

Probability Analyses (2008–2010).—A post hoc probability analysis of chlamydia testing in the subset of 9 Planned Parenthood affiliates providing data between 2008 and 2010 ($n = 9$) examined the pattern of changes in numbers testing between 2008 and 2010.

Analyses indicate that increases in chlamydia testing at the affiliate level were not likely due to chance alone. Between 2008 and 2009, nearly all affiliates (8 of 9) showed an increase in chlamydia testing among women ($P = 0.02$), and all affiliates showed an increase in chlamydia testing among men ($P = 0.002$). Between 2009 and 2010, there was no marked pattern of increases; the distribution of testing increases for both sexes was close to what one would expect from random data (4 of 9 affiliates increased for women and 5 of 9 for men, both $P > 0.40$). Last, between 2008 and 2010 (baseline and implementation year 2), nearly all affiliates (7 of 9) indicated an increase in chlamydia testing among women ($P = 0.09$), and nearly all affiliates (8 of 9) indicated an increase in chlamydia testing among men ($P = 0.02$). These conclusions hold for men and women, even if the probability of the observed increases is compared with a 5% increase in testing from 2008 to 2010, although for women, the comparison drops from 7 of 9 to 6 of 9 (both $P > 0.05$).

National Trends in STD Testing

For 2009 to 2010, trend data from clinics participating in national infertility prevention activities indicate that testing is higher in spring compared with other periods during those years. Since 2003, the number of tests reported is highest in the spring of each year, but also increases across time (Fig. 2). The pattern of increases around April (which is sustained up to the fourth quarter of each year) precedes GYT; however, the trend is clearest in 2009 and 2010. For example, in 2006, the second-quarter peak is high, but the drop-off is steep and seen in the third quarter.

DISCUSSION

GYT marks the first comprehensive national campaign promoting STD awareness, testing, and communication among youth in the United States. In its first 2 years of implementation, GYT event and media tracking show a growing number of youth engaged with the campaign. A near-doubling of Planned Parenthood on-the-ground activities in year 2 reached almost 3 times as many youth as that in year 1, reflecting both increased youth engagement as well as enthusiasm and uptake of GYT from clinical partners and youth advocates. Online engagement also increased on GYT's social media platforms, without any paid advertising. Reported metrics do not capture engagement with GYT through partner and celebrity platforms (e.g., social media sites, blogs, YouTube), which are difficult to track but may extend campaign reach. For example, one of GYT's prelaunch celebrity videos on YouTube attracted more than a quarter million views and generated thousands of comments.

Although there is ample evidence from other campaigns linking consumer awareness and exposure to behavior change,^{20,21} it is more difficult to assess the value of social media engagement and whether it translates into meaningful behavior change. Research in the area of digital media interventions for the promotion of sexual health has been limited, with better evidence for knowledge changes than for behavior change.²² It is also difficult to gauge the "success" of these efforts, given the lack of other national STD-focused campaigns in the United States. Tracking numbers may be modest relative to other national efforts, perhaps because of limited investment in advertising, or youth's reluctance to associate with a stigmatized topic. Although GYT's fan base grew throughout the first 2

years, some of its social media efforts (e.g., Campus Challenge) were met with limited youth engagement. Nonetheless, thousands of individuals were engaged in GYT and may have been influenced to get tested or talk about the issue (i.e., normalizing the conversation). Encouragingly, nearly 145,000 Web and SMS referrals were made to clinics/health centers through GYT during these 2 years, which suggests that users are linking to campaign information, and intending to get tested.

Positive changes in testing behavior were observed in a subset of 9 Planned Parenthood's affiliates. The number of STD patients visiting these Planned Parenthood affiliates was higher during campaign implementation periods compared with the same period precampaign. Patient demographic data reveal that increases were most notable among GYT's intended audience (youth) and those at increased risk for STDs, including racial/ethnic minorities and low-income persons. Data from these 9 affiliates are bolstered by data from a broader network of 69 Planned Parenthood affiliates, where testing increased between April 2009 and April 2010, commensurate with greater dissemination of GYT. Importantly, distinct increases were seen among young (<25 years), low-income and minority patients. Through the first 2 years of campaign implementation, STD test positivity rates remained consistent with the national averages reported to CDC,¹⁷ indicating that at-risk populations were being reached. These testing increases exceed those observed in family planning clinics nationwide during this same period,¹⁷ as well as what would be expected from random chance alone.

Finally, national trend data show that GYT activities occurred with a marked increase in testing in spring 2009 and 2010, which is sustained through most of those years. Interpretation of this finding is open to caution as similar patterns are visible in previous years, although with less clarity. It should be noted that STD Awareness Month activities predated the campaign, albeit on a smaller scale and in a less coordinated manner. It is plausible that raising awareness of STD prevention is generally useful and that GYT is a particularly efficacious version of these activities. The overall rise in chlamydia testing across time is certainly affected by efforts other than GYT (e.g., the HEDIS measure).²³

Limitations

As noted in the "Introduction," the data sources we have used in this evaluation are limited in that we had to seek evidence of GYT reach and effects in data primarily designed for other purposes. Tracking data were limited to a small number of events, media channels, and resources, not including television and other key channels used to promote GYT. Data described in this report do not explicitly connect campaign exposure to actual STD information-seeking, talking or testing behaviors. We regard these data as a starting point for more sophisticated analyses using data from later years, which will assess the campaign's impact on youth testing and communication, as well as stigma and theoretical mediators of behavior change.

Data from Planned Parenthood sources and from clinics participating in national infertility prevention efforts may not be representative of all facilities in the United States that test young persons for chlamydia and may be biased toward serving low-income women. The screening criteria and coverage vary by facility, and some may not send data on all tests

completed. Moreover, observed increases in testing may have resulted from factors unrelated to GYT. Finally, the specific health centers reporting data may have changed over time due to changes in local programs, which limits the use of approaches such as time series analyses.

CONCLUSIONS

There is limited but promising evidence that youth-oriented marketing and normalizing messaging, coupled with links to testing services and on-the-ground promotions, can yield increases in STD testing. Evidence presented here indicates that GYT is reaching youth—prompting dialogue about STDs/testing, linking youth to STD testing resources and services—and is associated with increased STD testing at select testing sites nationwide.

Cooperation among public and private sector partners and the use of existing data to measure outcomes may inform other STD prevention evaluation efforts in situations where ability to influence data collection is limited.

Acknowledgments:

The authors gratefully acknowledge the contributions of Elizabeth Torrone at the Centers for Disease Control and Prevention; Tina Hoff, Meredith Mishel, and Sarah Levine at Kaiser Family Foundation; Jason Rzepka, Lily Williamson, and Liza Vadnai at MTV; and Vanessa Cullins at Planned Parenthood Federation of America.

REFERENCES

1. Satterwhite CL, Torrone E, Meites E, et al. Sexually transmitted infections among US women and men: prevalence and incidence estimates, 2008. *Sex Transm Dis* 2013; 40:187–193. [PubMed: 23403598]
2. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2008. Atlanta, GA: US Department of Health and Human Services, 2009.
3. Pavlin NL, Gunn JM, Parker R, et al. Implementing chlamydia screening: What do women think? A systematic review of the literature. *BMC Public Health* 2006; 6:221–232. [PubMed: 16948838]
4. Hoff T, Greene L, Davis J. National survey of adolescents and young adults: sexual health knowledge, attitudes and experiences. Menlo Park, CA: Henry J Kaiser Family Foundation 2003 Available at: <http://kff.org/hiv/aids/report/national-survey-of-adolescents-and-young-adults/>. Accessed May 10, 2013.
5. Hood J, Friedman AL. Unveiling the hidden epidemic: A review of stigma associated with sexually transmissible infections. *Sex Health* 2011; 8:159–170. [PubMed: 21592429]
6. Institute of Medicine. The Hidden Epidemic: Confronting Sexually Transmitted Diseases. In: Eng TR, Butler WT, eds. Washington, DC: National Academy Press, 1997.
7. Forhan SE, Gottlieb SL, Sternberg MR, et al. Prevalence of sexually transmitted infections among female adolescents aged 14 to 19 in the United States. *Pediatrics* 2009; 124:1505–1512. [PubMed: 19933728]
8. CDC. Summary of a Review of the Literature: Programs to Promote Chlamydia Screening. Prepared by the Academy for Educational Development. Available at: www.cdc.gov/std/HealthComm/ChlamydiaLitReview2008.pdf.
9. CDC. Knowledge, Attitudes, and Behaviors about STDs, Chlamydia and Chlamydia Testing: Final Report from Telephone and In-Person In-Depth Interviews with Females Ages 15–25. Submitted by AED 2008.
10. Rosenstock IM, Strecher VJ, Becker MH. Social learning theory and the health belief model. *Health Educ Q* 1988; 15:175–183. [PubMed: 3378902]
11. Ajzen I From intentions to actions: A theory of planned behavior In: Kuhl J, Beckman J, eds. Action-Control: From Cognition to Behavior. Heidelberg, Germany: Springer, 1985:11–39.

12. Friedman AL, Bloodgood B. “Something we’d rather not talk about”: Findings from CDC exploratory research on STD communication with sexually active girls and women. *J Womens Health* 2010; 19:1823–1831.
13. Habel M, Levine S, Eastman-Mueller H, et al. Bringing GYT to Your Campus: 2011 Campaign Results, 2012 Update, and the Logistics of Hosting a STD/HIV Testing Event on Campus. Presented at the American College Health Association Annual Meeting, May 2012, Chicago, IL.
14. Friedman A, Bozniak A, Ford J, et al. Building on lessons learned from locally tailored adaptations of the United States’ National GYT: Get Yourself Tested Campaign. Presented at the World Social Marketing Conference, Toronto, Canada, 4 2013 Available at: http://wsconference.com/downloads/wsm_presentations/tuesday/Friedman-%20GYT-%20lessons%20learned%20from%20locally%20tailored%20adaptations.pdf
15. Planned Parenthood Federation of America. Get Yourself Tested: Helping Young People Stay Healthy. 2010 <http://issuu.com/actionfund/docs/gyt10-final-final>. Accessed May 2013.
16. Friedman AL, Habel M, McFarlane M, et al. Year 3 of GYT: Lessons Learned about Sustaining Youth’s Attention While Expanding Reach and Partnerships of the Get Yourself Tested Campaign. Abstract 27632 Presented at the National Conference on Health Communication, Marketing and Media August 2011, Atlanta, GA.
17. Fowler CI, Lloyd SW, Gable J, et al. Family planning annual report: 2010 national summary. Research Triangle Park, NC: RTI International, 2011 Available at: <http://www.hhs.gov/opa/pdfs/fpar-2010-national-summary.pdf>. Accessed May 10, 2013.
18. Centers for Disease Control and Prevention. Establishing a Holistic Framework to Reduce Inequities in HIV, Viral Hepatitis, STDs, and Tuberculosis in the United States. Atlanta, GA: US Department of Health and Human Services, 2010.
19. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2011. Atlanta, GA: US Department of Health and Human Services, 2012.
20. Hornik RC. Exposure: Theory and evidence about all the ways it matters. *Soc Market Q* 2002; 8:30–37.
21. Noar SM, Palmgreen P, Chabot M, et al. A 10-year systematic review of HIV/AIDS mass communication campaigns: Have we made progress? *J Health Commun* 2009; 14:15–42. [PubMed: 19180369]
22. Guse K, Levine D, Martins S, et al. Interventions using new digital media to improve adolescent sexual health: A systematic review. *J Adolesc Health* 2012; 51:535–543. [PubMed: 23174462]
23. Burstein GR, Snyder MH, Conley D, et al. Chlamydia screening in a Health Plan before and after a national performance measure introduction. *Obstet Gynecol* 2005; 106:327–334. [PubMed: 16055583]



Figure 1.
Screen shot of the 2010 GYT Web site.

Number of Chlamydia Tests (Males and Females) Reported Quarterly, 2003-2010




Figure 2. Number of chlamydia tests (men and women) reported quarterly, 2003 to 2010.

TABLE 1.

PPFA Data—STD Clients and Demographics at 9 Planned Parenthood Affiliates

	April 2008 (pre-GYT baseline)	April 2009 (GYT Year 1)	April 2010 (GYT Year 2)	Percent Change 2008–2010
STD clients	18,257*	22,929*	31,227*	71.0%
Female	15,927	19,627	27,376	71.9%
Male	2330	3302	3851	65.3%
Demographics				
Age < 25 y	9473	11,971	14,862	56.9%
Female	8649	10,747	13,493	56.0%
Male	824	1224	1369	66.1%
Race/Ethnicity, white	7458	10,480	11,013	47.7%
Female	6545	9058	9519	45.4%
Male	913	1422	1494	63.6%
Race/Ethnicity, black	1355	2253	3162	133.3%
Female	1188	1964	2823	137.6%
Male	167	289	339	102.9%
Race/Ethnicity, Latino	3613	4535	5569	54.1%
Female	3259	3989	4969	52.5%
Male	354	546	600	69.5%
150% of the federal poverty line	6224	12,479	15,766	153.3%
Female	5532	11,381	14,439	161.0%
Male	692	1098	1327	91.8%

* No unknowns (unknown not an option).

TABLE 2.

PPFA Data--STD Testing at 9 Planned Parenthood Affiliates

Testing	April 2008 (pre-GYT baseline)	April 2009 (GYT Year 1)	April 2010 (GYT Year 2)	Percent Change 2008-2010
Chlamydia tests	16,398*	21,179*	23,151*	41.2%
Female	14,391	17,571	20,320	41.2%
Male	1887	2996	2830	49.9%
Positive chlamydia tests (test positivity)	1079* (6.6%)	1379* (6.5%)	1687* (7.3%)	56.3%
Female	735 (5.1%)	1015 (5.8%)	1335 (6.6%)	81.6%
Male	191 (10.1%)	320 (10.7%)	352 (12.4%)	84.3%
Gonorrhea tests	14,903*	23,504*	28,391*	90.5%
Female	13,007	19,819	25,247	94.1%
Male	1776	3094	3144	77.0%
Positive gonorrhea tests (test positivity)	119* (0.1%)	263* (1.1%)	268* (0.9%)	125.2%
Female	87 (0.7%)	157 (0.8%)	178 (0.7%)	104.6%
Male	31 (1.7%)	89 (2.9%)	90 (2.9%)	190.3%
HIV tests	6265*	9877*	13,327*	112.7%
Female	4661	7353	10,504	125.4%
Male	1500	2350	2801	86.7%
Positive HIV tests (test positivity)	6 (0.0%)	29* (0.0%)	18* (0.0%)	†
Female	5	0	4	†
Male	1	20	14	†

* Includes unknowns for sex.

† N is too small to reliably calculate percent change.

PPFA Data--STD Clients and Demographics at All Planned Parenthood Affiliates Reporting in 2009 and 2010

TABLE 3.

	April 2009 (GYT Year 1)	April 2010 (GYT Year 2)	Percent Change 2009–2010
Reporting Affiliates	69	69	
STD clients	106,948*	120,370*	12.5%
Female	93,026	106,024	13.9%
Male	13,922	14,346	3.0%
Demographics			
Age 25 y	65,538	89,814	37.0%
Female	58,133	80,939	39.2%
Male	7,405	8,875	19.8%
Race/Ethnicity, white	50,268	50,799	1.0%
Female	43,589	44,654	2.4%
Male	6,679	6,145	-8.0%
Race/Ethnicity, black	16,147	21,419	32.6%
Female	14,022	19,040	35.8%
Male	2,125	2,379	11.9%
Race/Ethnicity, Latino	21,424	23,830	11.2%
Female	19,150	21,117	10.3%
Male	2,274	2,713	19.3%
150% of the federal poverty line	64,988	87,795	35.1%
Female	58,133	80,939	39.2%
Male	6,855	6,856	0.0%

* No unknowns (unknown not an option).

PPFA Data--STD Testing at All Planned Parenthood Affiliates Reporting in 2009 and 2010

TABLE 4.

	April 2009 (GYT Year 1)	April 2010 (GYT Year 2)	Percent Change 2009-2010
Reporting affiliates	69	69	
Testing			
Chlamydia tests			
Female	98,872*	106,932*	8.1%
Male	98,807	111,061	12.4%
Total	13,895	22,248	60.1%
Positive chlamydia tests (test positivity)	5515* (5.5%)	7027* (6.5%)	27.4%
Female	4221 (4.2%)	5036 (5.0%)	19.3%
Male	1095 (7.9%)	1262 (5.6%)	15.2%
Gonorrhea tests			
Female	91,335*	103,734*	13.6%
Male	75,970	85,039	11.9%
Total	11,729	10,294	-12.2%
Positive gonorrhea tests (test positivity)	788* (0.9%)	956* (0.9%)	21.3%
Female	540 (0.7%)	638 (0.7%)	18.1%
Male	224 (1.9%)	210 (2.0%)	-6.2%
HIV tests			
Female	42,354*	50,454*	19.1%
Male	31,704	36,405	14.8%
Total	9315	10,366	11.3%
Positive HIV tests (test positivity)	55* (0.0%)	47* (0.0%)	-14.5%
Female	12	19	†
Male	34	27	†

* Includes unknowns for sex.

† N is too small to reliably calculate percent change.