

SUPPLEMENTARY BOX. Studies analyzed in systematic review of screening and testing of health care personnel for tuberculosis

1. Choudhary M, Ramirez L, Long R, et al. A university hospital's 10-year experience with tuberculin testing: value of the 2-step tuberculin skin test. *Am J Infect Control* 2006;34:358–61.
2. Frenzel EC, Thomas GA, Hanna HA. The importance of two-step tuberculin skin testing for newly employed healthcare workers. *Infect Control Hosp Epidemiol* 2006;27:512–4.
3. Gillenwater KA, Sapp SC, Pearce K, Siberry GK. Increase in tuberculin skin test converters among health care workers after a change from Tubersol to Aplisol. *Am J Infect Control* 2006;34:651–4.
4. Baussano I, Bugiani M, Carosso A, et al. Risk of tuberculin conversion among healthcare workers and the adoption of preventive measures. *Occup Environ Med* 2007;64:161–6.
5. Soborg B, Andersen AB, Larsen HK, et al. Detecting a low prevalence of latent tuberculosis among health care workers in Denmark detected by M. tuberculosis specific IFN-gamma whole-blood test. *Scand J Infect Dis* 2007;39:554–9.
6. Barsegian V, Mathias KD, Wrighton-Smith P, Grosse-Wilde H, Lindemann M. Prevalence of latent tuberculosis infection in German radiologists. *J Hosp Infect* 2008;69:69–76.
7. Nienhaus A, Schablon A, Bacle CL, Siano B, Diel R. Evaluation of the interferon-gamma release assay in healthcare workers. *Int Arch Occup Environ Health* 2008;81:295–300.
8. Pollock NR, Campos-Neto A, Kashino S, et al. Discordant QuantiFERON-TB Gold test results among US healthcare workers with increased risk of latent tuberculosis infection: a problem or solution? *Infect Control Hosp Epidemiol* 2008;29:878–86.
9. Casas I, Latorre I, Esteve M, et al. Evaluation of interferon-gamma release assays in the diagnosis of recent tuberculosis infection in health care workers. *PLoS ONE* 2009;4:e6686.
10. Cummings KJ, Smith TS, Shogren ES, et al. Prospective comparison of tuberculin skin test and QuantiFERON-TB Gold In-Tube assay for the detection of latent tuberculosis infection among healthcare workers in a low-incidence setting. *Infect Control Hosp Epidemiol* 2009;30:1123–6.
11. Girardi E, Angeletti C, Puro V, et al. Estimating diagnostic accuracy of tests for latent tuberculosis infection without a gold standard among healthcare workers. *Euro Surveill* 2009;14:29.
12. Miranda C, Yen-Lieberman B, Terpeluk P, Tomford JW, Gordon S. Reducing the rates of indeterminate results of the QuantiFERON-TB Gold In-Tube test during routine preemployment screening for latent tuberculosis infection among healthcare personnel. *Infect Control Hosp Epidemiol* 2009;30:296–8.
13. Tripodi D, Brunet-Courtois B, Nael V, et al. Evaluation of the tuberculin skin test and the interferon-release assay for TB screening in French healthcare workers. *J Occup Med Toxicol* 2009;4.
14. Vinton P, Mhrshahi S, Johnson P, Jenkin GA, Jolley D, Biggs BA. Comparison of QuantiFERON-TB Gold In-Tube Test and tuberculin skin test for identification of latent Mycobacterium tuberculosis infection in healthcare staff and association between positive test results and known risk factors for infection. *Infect Control Hosp Epidemiol* 2009;30:215–21.
15. Gandra S, Scott WS, Somaraju V, Wang H, Wilton S, Feigenbaum M. Questionable effectiveness

of the QuantiFERON-TB Gold Test (Cellestis) as a screening tool in healthcare workers. *Infect Control Hosp Epidemiol* 2010;31:1279–85.

16. Ringshausen FC, Nienhaus A, Schablon A, Schlosser S, Schultze-Werninghaus G, Rohde G. Predictors of persistently positive Mycobacterium-tuberculosis-specific interferon-gamma responses in the serial testing of health care workers. *BMC Infect Dis* 2010;10:220.
17. Schablon A, Harling M, Diel R, Nienhaus A. Risk of latent TB infection in individuals employed in the healthcare sector in Germany: a multicentre prevalence study. *BMC Infect Dis* 2010;10:107.
18. Ringshausen FC, Nienhaus A, Torres Costa J, et al. Within-subject variability of Mycobacterium tuberculosis-specific gamma interferon responses in German health care workers. *Clin Vaccine Immunol* 2011;18:1176–82.
19. Sherman HA, Karakis I, Heimer D, et al. Housekeeping health care workers have the highest risk for tuberculin skin test conversion. *Int J Tuberc Lung Dis* 2011;15:1050–5.
20. Fong KS, Tomford JW, Teixeira L, et al. Challenges of interferon-gamma release assay conversions in serial testing of health-care workers in a TB control program. *Chest* 2012;142:55–62.
21. Freeman JT, Marshall RJ, Newton S, et al. Screening for Mycobacterium tuberculosis infection among healthcare workers in New Zealand: prospective comparison between the tuberculin skin test and the QuantiFERON-TB Gold In-Tube assay. *N Z Med J* 2012;125:21–9.
22. Joshi M, Monson TP, Woods GL. Use of interferon-gamma release assays in a health care worker screening program: experience from a tertiary care centre in the United States. *Can Respir J* 2012;19:84–8.
23. Larcher C, Frizzera E, Pretto P, Lang M, Sonnleitner N, Huemer HP. Immunosurveillance for Mycobacterium tuberculosis of health care personnel in a third level care hospital. *Med Lav* 2012;103:26–36.
24. Moucaut A, Nienhaus A, Courtois B, et al. The effect of introducing IGRA to screen French healthcare workers for tuberculosis and potential conclusions for the work organisation. *J Occup Med Toxicol* 2013;8:12.
25. Slater ML, Welland G, Pai M, Parsonnet J, Banaei N. Challenges with QuantiFERON-TB Gold assay for large-scale, routine screening of U.S. healthcare workers. *Am J Respir Crit Care Med* 2013;188:1005–10.
26. Dorman SE, Belknap R, Graviss EA, et al. Interferon-gamma release assays and tuberculin skin testing for diagnosis of latent tuberculosis infection in healthcare workers in the United States. *Am J Respir Crit Care Med* 2014;189:77–87.
27. Joshi M, Monson TP, Joshi A, Woods GL. IFN-gamma release assay conversions and reversions. Challenges with serial testing in U.S. health care workers. *Ann Am Thorac Soc* 2014;11:296–302.
28. Nienhaus A, Garipey PK, Trouve C, Lhaumet C, Toureau J, Peters C. Tuberculosis screening at the Sainte-Anne Hospital in Paris—Results of first and second IGRA. *J Occup Med Toxicol* 2014;9:24.
29. Schablon A, Nienhaus A, Ringshausen FC, Preisser AM, Peters C. Occupational screening for

tuberculosis and the use of a borderline zone for interpretation of the IGRA in German healthcare workers.[Erratum appears in PLoS One. 2015;10:e0142541; PMID: 26540199]. PLoS ONE 2014;9:e115322.

30. Weddle G, Hamilton M, Potthoff D, Rivera D, Jackson MA. QuantiFERON-TB gold in-tube testing for tuberculosis in healthcare professionals. *Lab Med* 2014;45:207–10.
31. Durando P, Alicino C, Orsi A, et al. Latent tuberculosis infection among a large cohort of medical students at a teaching hospital in Italy. *Biomed Res Int* 2015; 2015:746895.
32. King TC, Upfal M, Gottlieb A, et al. T-SPOT.TB Interferon-gamma Release Assay Performance in Healthcare Worker Screening at Nineteen U.S. Hospitals. *Am J Respir Crit Care Med* 2015;192:367–73.
33. Lamberti M, Muoio M, Arnese A, et al. Prevalence of latent tuberculosis infection in healthcare workers at a hospital in Naples, Italy, a low-incidence country. *J Occup Med Toxicol* 2016;11:53.
34. Olivieri R, Scarnera S, Ciabattini A, et al. Using IFN-gamma release assay to confirm tuberculin skin test improves the screening of latent tuberculosis infection in Italian healthcare workers. *J Occup Med Toxicol* 2016;11:29.
35. Farah WH, Breeher LE, Newcomb RD, et al. Late boosting phenomenon in TST conversion among health care workers. *Occup Med (Oxford, England)* 2017;67:484–9.
36. Dobler CC, Farah WH, Alsawas M, et al. Tuberculin skin test conversions and occupational exposure risk in US healthcare workers. *Clin Infect Dis* 2018;66:706–11.