

# *Bacillus cereus*–Attributable Primary Cutaneous Anthrax-Like Infection in Newborn Infants, India

## Appendix

**Appendix Table.** Primer sequences used in this study\*

Target gene	Sequence (5'–3')	Amplicon size, bp	Annealing temperature, °C	Reference/GenBank accession no.
<i>cytK</i>	F: CAACCCAGTTTGCAGTTCCG R: GGCGCTAGTGCAACATTACG	601	55	AJ277962
<i>nheA</i>	F: GTTAGGATCACAATCACCGC R: ACGAATGTAATTTGAGTCGC	755	48	(1)
<i>nheB</i>	F: TTTAGTAGTGGATCTGTACGC R: TTAATGTTTCGTTAATCCTGC	743	45	(1)
<i>nheC</i>	F: TGGATTCCAAGATGTAACG R: ATTACGACTTCTGCTTGTGC	683	47	(1)
<i>entFM</i>	F: AAAGAAATTAATGGACAAACTCAA R: GTATGTAGCTGGGCCTGTACGT	596	50	(2)
<i>hblC</i>	F: GATACTCAATGTGGCAACTGC R: TTGAGACTGCTCGTCTAGTTG	740	52	(3)
<i>hblD</i>	F: ACCGGTAACACTATTCATGC R: GAGTCCATATGCTTAGATGC	829	48	(1)
<i>hblA</i>	F: AAGCAATGGAATACAATGGG R: AGAATCTAAATCATGCCACTGC	1154	48	(1)
<i>sph</i>	F: CGTGCCGATTTAATTGGGGC R: CAATGTTTTAAACATGGATGCG	558	48	(4)
<i>pi-plc</i>	F: CGCTATCAAATGGACCATGG R: GGACTATTCCATGCTGTACC	569	51	(5)
<i>cya</i>	F: GCGTTTTCTTTAGCGTTTTTC R: CTACATTTTAGCCATCACTTCT	727	48	(6)
<i>cap</i>	F: CTGAGCCATTAATCGATATG R: TCCCCTTACGTAATCTGAG	846	55	(7)
<i>pag</i>	F: TCCTAACACTAACGAAGTCG R: GAGGTAGAAGGATATACGGT	596	55	(8)
<i>lef</i>	F: TGAACCCGTAATTGTAATCCAATC R: ATCGCTCCAGTGTTGATAGTGCT	475	53	(9)

\*F, forward; R, reverse.

## References

1. Guinebretière MH, Broussolle V, Nguyen-The C. Enterotoxigenic profiles of food-poisoning and food-borne *Bacillus cereus* strains. J Clin Microbiol. 2002;40:3053–6. [PubMed](https://pubmed.ncbi.nlm.nih.gov/12011288/) <http://dx.doi.org/10.1128/JCM.40.8.3053-3056.2002>
2. Kim J-B, Kim J-M, Cho S-H, Oh H-S, Choi NJ, Oh D-H. Toxin genes profiles and toxin production ability of *Bacillus cereus* isolated from clinical and food samples. J Food Sci. 2011;76:T25–9. [PubMed](https://pubmed.ncbi.nlm.nih.gov/21111211/) <http://dx.doi.org/10.1111/j.1750-3841.2010.01958.x>

3. Chon JW, Kim JH, Lee SJ, Hyeon JY, Song KY, Park C, et al. Prevalence, phenotypic traits and molecular characterization of emetic toxin-producing *Bacillus cereus* strains isolated from human stools in Korea. J Appl Microbiol. 2012;112:1042–9. PubMed <http://dx.doi.org/10.1111/j.1365-2672.2012.05277.x>
4. Hsieh YM, Sheu SJ, Chen YL, Tsen HY. Enterotoxigenic profiles and polymerase chain reaction detection of *Bacillus cereus* group cells and *B. cereus* strains from foods and food-borne outbreaks. J Appl Microbiol. 1999;87:481–90. PubMed <http://dx.doi.org/10.1046/j.1365-2672.1999.00837.x>
5. Hansen BM, Damgaard PH, Eilenberg J, Pedersen JC. Molecular and phenotypic characterization of *Bacillus thuringiensis* isolated from leaves and insects. J Invertebr Pathol. 1998;71:106–14. PubMed <http://dx.doi.org/10.1006/jipa.1997.4712>
6. Shangkuan YH, Chang YH, Yang JF, Lin HC, Shaio MF. Molecular characterization of *Bacillus anthracis* using multiplex PCR, ERIC-PCR and RAPD. Lett Appl Microbiol. 2001;32:139–45. PubMed <http://dx.doi.org/10.1046/j.1472-765x.2001.00881.x>
7. Hutson RA, Duggleby CJ, Lowe JR, Manchee RJ, Turnbull PC. The development and assessment of DNA and oligonucleotide probes for the specific detection of *Bacillus anthracis*. J Appl Bacteriol. 1993;75:463–72. PubMed <http://dx.doi.org/10.1111/j.1365-2672.1993.tb02803.x>
8. Beyer W, Glöckner P, Otto J, Böhm R. A nested PCR method for the detection of *Bacillus anthracis* in environmental samples collected from former tannery sites. Microbiol Res. 1995;150:179–86. PubMed [http://dx.doi.org/10.1016/S0944-5013\(11\)80054-6](http://dx.doi.org/10.1016/S0944-5013(11)80054-6)
9. Kim K, Seo J, Wheeler K, Park C, Kim D, Park S, et al. Rapid genotypic detection of *Bacillus anthracis* and the *Bacillus cereus* group by multiplex real-time PCR melting curve analysis. FEMS Immunol Med Microbiol. 2005;43:301–10. PubMed <http://dx.doi.org/10.1016/j.femsim.2004.10.005>