



HHS Public Access

Author manuscript

Anesthesiology. Author manuscript; available in PMC 2018 January 01.

Published in final edited form as:

Anesthesiology. 2017 January ; 126(1): 6–8. doi:10.1097/ALN.0000000000001384.

Monkey in the middle: Translational studies of pediatric anesthetic exposure

Mark G. Baxter, Ph.D. and

Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY (M.G.B.)

Maria C. Alvarado, Ph.D.

Division of Developmental and Cognitive Neuroscience, Yerkes National Primate Research Center, Atlanta, GA (M.C.A.)

A seminal report by Jevtovic-Todorovic and colleagues¹ described neurotoxicity and long-term cognitive impairments after exposure to general anesthesia in infant (postnatal day 7) rats. This phenomenon has been observed repeatedly in multiple species and with a variety of different anesthetic drugs. It has also sparked parallel retrospective and prospective studies in humans that suggest that pediatric anesthesia might be associated with increased risk for adverse neurocognitive outcomes when exposure is repeated or prolonged^{2–9}. The recent investigation by Coleman and colleagues¹⁰ joins a growing number of studies investigating the impact of exposure to general anesthesia in infancy on neurobehavioral development in nonhuman primates. This highly translationally-relevant model makes unique contributions to understanding the phenomena and potential mechanisms of long-term cognitive and behavioral changes after general anesthesia early in life.

There are a number of important advantages to studying this question in nonhuman primate models, not the least of which is the ability to study the effects of general anesthesia in the absence of surgery. Physiological monitoring and support is much more feasible in an infant monkey than an infant rodent. Thus, long-term effects of anesthesia exposure cannot easily be attributed to hypoxia, hypercapnia, and so forth. The stage of brain development at birth of a rhesus monkey is similar to that of a 6-month old human infant^{11,12} whereas based on many neurobiological markers, postnatal day 7 rats that are at peak vulnerability to anesthetic neurotoxicity¹³ may more closely match the stage of brain development of a third-trimester human fetus. Nonhuman primates also have the capacity to perform more complex tasks and have a sophisticated social structure, like that of humans, allowing more translationally relevant tests of cognition and socioemotional behavior. Because puberty occurs between 3–5 years of age in rhesus monkeys, it is possible to examine changes in behavior over the course of development that would be challenging to study in rodent models, in which sexual maturity occurs around the age of 6 weeks.

Corresponding author's address: Mark G. Baxter, Ph.D., Department of Neuroscience, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1639, New York, NY 10029 USA, Office: +1 212 824 9303, FAX: +1 646 537 9585, mark.baxter@mssm.edu.

Neither author has any conflicts of interest to declare.

The primary outcomes in this study from Coleman and colleagues are measures of early motor reflexes and emotional behavior in a number of different situations. They employ test batteries for monkeys that were developed to be analogous to behavioral testing procedures in humans, including the Laboratory Temperament Assessment Battery (LabTAB¹⁴) and "human intruder" tests. These tests have been used extensively in studies of socioemotional development in macaque monkeys¹⁵ and therefore are well-validated and highly translational tools. Indeed, the use of these paradigms is a significant strength of studies in nonhuman primates, where tests can be given in similar formats to humans and monkeys. As another example, the ongoing Mayo Anesthesia Safety in Kids (MASK) study¹⁶ includes an operant behavioral testing battery that has been validated for developmental research in both humans and monkeys¹⁷.

Coleman and colleagues found abnormal motor reflexes at one month of age (about three weeks after anesthesia exposure) as well as heightened anxiety in the home cage social environment at twelve months of age in monkeys that were exposed three times to isoflurane (for five hours each time) between postnatal days 6 and 12. Changes in these measures were not statistically significant in monkeys exposed to isoflurane only once for five hours on postnatal day 6. Because monkeys did not undergo surgery, changes in anxiety in monkeys cannot be attributed to experience of postoperative pain. These findings suggest that negative behavioral changes in children after surgery with general anesthesia¹⁸ might result, in part, from long-term effects of multiple exposures to the anesthetic agent during the surgical procedure.

Because of the time and expense involved in a prospective study with nonhuman primates, investigators have based their anesthesia protocols on durations of exposure that are known to result in increased neuro- and glioapoptosis^{12,17,19,20}, a candidate mechanism for the long-term effects of pediatric anesthesia on behavior. Because these durations tend to be longer than those commonly encountered in the pediatric operating room, many of these studies face the criticism that the anesthesia exposures are not clinically relevant. Single anesthesia exposures in infants that have been investigated recently^{8,9} are of short duration relative to those in the studies to date with monkeys. It is worth noting that prolonged anesthesia exposure is certainly not unknown in pediatric surgery; Coleman et al.¹⁰ cite 30% of infant anesthesia exposures being longer than 3 hours at their institution, and prolonged sedation in the neonatal intensive care unit may last for weeks¹⁷. However, repeated exposures to anesthesia appear rare in these populations; for example, 7.4% of cases (44/593) in the study by Wilder et al.⁴ received 3 or more anesthetic exposures, so three long exposures to anesthesia, especially within a week, would be extremely unusual in a clinical context. Although this is a limitation of extant preclinical studies, these exposure protocols establish boundary conditions upon which future work can build, to carry out more finely-grained analyses of duration and frequency of anesthesia exposure.

Another issue in preclinical studies, especially with nonhuman primates, is that of repeated testing with a relatively small subject pool. This is a practical constraint of research with rare and expensive animals. Although this can incur statistical concerns about a large number of tests on a limited population, it is also a strength of these studies that data can be collected on the same study population longitudinally and across a number of different behavioral

domains. As multiple research groups carry out studies using similar behavioral protocols, the generality of individual findings becomes clearer, as in all other areas of biomedical research. Similarly, this study like others tests both male and female monkeys but is underpowered to detect subtle sex differences. Nevertheless, this is a more desirable state of affairs than limiting preclinical studies to a single sex to evade this criticism²¹.

Two studies in monkeys^{10,22} have independently reported increased anxiety-related behaviors in monkeys that were repeatedly exposed to general anesthesia as infants, despite using different anesthetic agents and different schedules and durations of exposure. The finding that repeated exposure is associated with adverse neurocognitive outcomes is consistent with some human epidemiological studies^{4,23}. Recent prospective studies in humans have emphasized the safety of single, relatively brief exposures to general anesthesia in the context of pediatric surgery^{8,9}; Coleman et al.¹⁰ also find limited effects of single exposures to general anesthesia in monkeys. These observations reinforce one another and suggest that concerns about repeated or prolonged exposure of children to general anesthesia remain significant. Long-term neurobehavioral changes after repeated exposure to anesthesia in infancy may not simply be a consequence of cumulative exposure to the anesthetic agent; one study in rodents found a greater decline in synaptic density in adult rats exposed to sevoflurane three times for two hours each as infants compared to those exposed once for six hours²⁴, so the mechanisms of long-term effects of repeated anesthesia exposure also merit further investigation.

The translation of these findings to the clinical setting remains subject to interpretation. It appears relatively easier to detect neurocognitive impairments in animal models after pediatric anesthesia than it does in clinical studies with humans. This may reflect uncertainty about the animal models that are used, with regard to details of anesthesia exposure, dosing, and so forth. It may also reflect the greater degree of experimental control in animal studies, where all subjects receive identical anesthetic regimens, and behavioral assessments may be more extensive and sensitive to subtle effects.

Findings that single exposures to general anesthesia for pediatric surgery are not associated with adverse neurocognitive outcomes^{8,9} are a great relief to anxious parents (and anesthesiologists). However, it does not seem safe to conclude that because single exposures to general anesthesia for pediatric surgery are apparently without substantial long-term adverse neurocognitive effects, that any pattern of pediatric anesthesia exposure will be similarly benign. This work with nonhuman primates forms a critical middle ground for future investigations of impaired neurobehavioral function after repeated and/or prolonged anesthesia exposure, and for testing potential interventions to mitigate them.

Acknowledgments

The authors' research on this topic is supported by NIH R01-HD068388.

References

1. Jevtovic-Todorovic V, Hartman RE, Izumi Y, Benshoff ND, Dikranian K, Zorumski CF, Olney JW, Wozniak DF. Early exposure to common anesthetic agents causes widespread neurodegeneration in

the developing rat brain and persistent learning deficits. *J Neurosci*. 2003; 23:876–882. [PubMed: 12574416]

2. O'Leary JD, Janus M, Duku E, Wijeyesundera DN, To T, Li P, Maynes JT, Crawford MW. A population-based study evaluating the association between surgery in early life and child development at primary school entry. *Anesthesiology*. 2016; 125:272–279. [PubMed: 27433745]
3. Graham MR, Brownell M, Chateau DG, Dragan RD, Burchill C, Fransoo RR. Neurodevelopmental Assessment in Kindergarten in Children Exposed to General Anesthesia before the Age of 4 Years: A Retrospective Matched Cohort Study. *Anesthesiology*. 2016
4. Wilder RT, Flick RP, Sprung J, Katusic SK, Barbaresi WJ, Mickelson C, Gleich SJ, Schroeder DR, Weaver AL, Warner DO. Early exposure to anesthesia and learning disabilities in a population-based birth cohort. *Anesthesiology*. 2009; 110:796–804. [PubMed: 19293700]
5. Sprung J, Flick RP, Katusic SK, Colligan RC, Barbaresi WJ, Bojani K, Welch TL, Olson MD, Hanson AC, Schroeder DR, Wilder RT, Warner DO. Attention-deficit/hyperactivity disorder after early exposure to procedures requiring general anesthesia. *Mayo Clinic Proceedings*. 2012; 87:120–129. [PubMed: 22305025]
6. Ing CH, Dimaggio CJ, Malacova E, Whitehouse AJ, Hegarty MK, Feng T, Brady JE, Ungern-Sternberg von BS, Davidson AJ, Wall MM, Wood AJJ, Li G, Sun LS. Comparative analysis of outcome measures used in examining neurodevelopmental effects of early childhood anesthesia exposure. *Anesthesiology*. 2014; 120:1319–1332. [PubMed: 24694922]
7. Ing C, DiMaggio C, Whitehouse A, Hegarty MK, Brady J, Ungern-Sternberg von BS, Davidson A, Wood AJJ, Li G, Sun LS. Long-term differences in language and cognitive function after childhood exposure to anesthesia. *Pediatrics*. 2012; 130:e476–e485. [PubMed: 22908104]
8. Sun LS, Li G, Miller TLK, Salorio C, Byrne MW, Bellinger DC, Ing C, Park R, Radcliffe J, Hays SR, Dimaggio CJ, Cooper TJ, Rauh V, Maxwell LG, Youn A, McGowan FX. Association between a single general anesthesia exposure before age 36 months and neurocognitive outcomes in later childhood. *JAMA*. 2016; 315:2312–2320. [PubMed: 27272582]
9. Davidson AJ, Disma N, de Graaff JC, Withington DE, Dorris L, Bell G, Stargatt R, Bellinger DC, Schuster T, Arnup SJ, Hardy P, Hunt RW, Takagi MJ, Giribaldi G, Hartmann PL, Salvo I, Morton NS, Ungern-Sternberg von BS, Locatelli BG, Wilton N, Lynn A, Thomas JJ, Polaner D, Bagshaw O, Szmuk P, Absalom AR, Frawley G, Berde C, Ormond GD, Marmor J, et al. Neurodevelopmental outcome at 2 years of age after general anaesthesia and awake-regional anaesthesia in infancy (GAS): an international multicentre, randomised controlled trial. *Lancet*. 2016; 387:239–250. [PubMed: 26507180]
10. Coleman K, Robertson ND, Dissen GA, Neuringer MD, Martin LD, Cuzon Carlson VC, Kroenke C, Fair D, Brambrink A. Infant exposure to anesthesia has long-term, dose-dependent consequences on motor and behavioral development in rhesus macaques. *Anesthesiology*. 2016:1–42.
11. Dobbing J, Sands J. Comparative aspects of the brain growth spurt. *Early Hum Dev*. 1979; 3:79–83. [PubMed: 118862]
12. Brambrink AM, Evers AS, Avidan MS, Farber NB, Smith DJ, Zhang X, Dissen GA, Creeley CE, Olney JW. Isoflurane-induced neuroapoptosis in the neonatal rhesus macaque brain. *Anesthesiology*. 2010; 112:834–841. [PubMed: 20234312]
13. Yon JH, Daniel-Johnson J, Carter LB, Jevtovic-Todorovic V. Anesthesia induces neuronal cell death in the developing rat brain via the intrinsic and extrinsic apoptotic pathways. *Neuroscience*. 2005; 135:815–827. [PubMed: 16154281]
14. Goldsmith, HH.; Rothbart, MK. Contemporary instruments for assessing early temperament by questionnaire and in the laboratory, *Explorations in temperament*. Angleitner, A.; Strelau, J., editors. Springer; 1991. p. 249-272.
15. Kalin NH, Shelton SE. Defensive behaviors in infant rhesus monkeys: environmental cues and neurochemical regulation. *Science*. 1989; 243:1718–1721. [PubMed: 2564702]
16. Gleich SJ, Flick R, Hu D, Zaccariello MJ, Colligan RC, Katusic SK, Schroeder DR, Hanson A, Buenvenida S, Wilder RT, Sprung J, Voigt RG, Paule MG, Chelonis JJ, Warner DO. Neurodevelopment of children exposed to anesthesia: design of the Mayo Anesthesia Safety in Kids (MASK) study. *Contemporary Clinical Trials*. 2015; 41:45–54. [PubMed: 25555440]

17. Paule MG, Li M, Allen RR, Liu F, Zou X, Hotchkiss C, Hanig JP, Patterson TA, W Slikker J, Wang C. Ketamine anesthesia during the first week of life can cause long-lasting cognitive deficits in rhesus monkeys. *Neurotoxicology and Teratology*. 2011; 33:220–230. [PubMed: 21241795]
18. Stargatt R, Davidson AJ, Huang GH, Czarnecki C, Gibson AJN, Stewart SA, Jansen K. A cohort study of the incidence and risk factors for negative behavior changes in children after general anesthesia. *Paediatr Anaesth*. 2006; 16:846–859. [PubMed: 16884468]
19. Brambrink AM, Back SA, Riddle A, Gong X, Moravec MD, Dissen GA, Creeley CE, Dikranian KT, Olney JW. Isoflurane-induced apoptosis of oligodendrocytes in the neonatal primate brain. *Ann Neurol*. 2012; 72:525–535. [PubMed: 23109147]
20. Slikker W, Zou X, Hotchkiss CE, Divine RL, Sadovova N, Twaddle NC, Doerge DR, Scallet AC, Patterson TA, Hanig JP, Paule MG, Wang C. Ketamine-induced neuronal cell death in the perinatal rhesus monkey. *Toxicol Sci*. 2007; 98:145–158. [PubMed: 17426105]
21. Clayton JA, Collins FS. Policy: NIH to balance sex in cell and animal studies. *Nature*. 2014; 509:282–283. [PubMed: 24834516]
22. Raper J, Alvarado MC, Murphy KL, Baxter MG. Multiple anesthetic exposure in infant monkeys alters emotional reactivity to an acute stressor. *Anesthesiology*. 2015; 123:1084–1092. [PubMed: 26313293]
23. Flick RP, Katusic SK, Colligan RC, Wilder RT, Voigt RG, Olson MD, Sprung J, Weaver AL, Schroeder DR, Warner DO. Cognitive and behavioral outcomes after early exposure to anesthesia and surgery. *Pediatrics*. 2011; 128:e1053–e1061. [PubMed: 21969289]
24. Amrock LG, Starner ML, Murphy KL, Baxter MG. Long-term effects of single or multiple neonatal sevoflurane exposures on rat hippocampal ultrastructure. *Anesthesiology*. 2015; 122:87–95. [PubMed: 25289484]