

2011-2012 Influenza Season Week 42 ending October 22, 2011

All data are preliminary and may change as more reports are received.

Synopsis: During week 42 (October 16-22, 2011), influenza activity remained low in the United States.

- **U.S. Virologic Surveillance:** Of the 1,479 specimens tested by U.S. World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NREVSS) collaborating laboratories and reported to CDC/Influenza Division, 12 (0.8%) were positive for influenza.
- **Pneumonia and Influenza (P&I) Mortality Surveillance:** The proportion of deaths attributed to P&I was below the epidemic threshold.
- **Influenza-associated Pediatric Mortality:** No influenza-associated pediatric deaths were reported.
- **Outpatient Illness Surveillance:** The proportion of outpatient visits for influenza-like illness (ILI) was 1.1%, which is below the national baseline of 2.4%. All 10 regions reported ILI below region-specific baseline levels. All 50 states and New York City experienced minimal ILI activity and the District of Columbia had insufficient data.
- **Geographic Spread of Influenza:** The geographic spread of influenza in the District of Columbia, Puerto Rico, the U.S. Virgin Islands, and 18 states was reported as sporadic and Guam and 32 states reported no influenza activity.

National and Regional Summary of Select Surveillance Components

HHS Surveillance Regions*	Data for current week			Data cumulative since October 2, 2011 (Week 40)				
	Out-patient ILI†	% of respiratory specimens positive for flu‡	Number of jurisdictions reporting regional or widespread activity§	A (H3)	A (2009 H1N1)	A (Subtyping not performed)	B	Pediatric Deaths
Nation	Normal	0.8%	0 of 54	16	3	17	8	0
Region 1	Normal	4.2%	0 of 6	0	0	0	3	0
Region 2	Normal	0.0%	0 of 4	0	0	0	0	0
Region 3	Normal	0.5%	0 of 6	0	0	2	0	0
Region 4	Normal	1.3%	0 of 8	0	0	12	1	0
Region 5	Normal	2.1%	0 of 6	5	1	0	1	0
Region 6	Normal	0.4%	0 of 5	2	0	1	1	0
Region 7	Normal	0.0%	0 of 4	0	0	0	0	0
Region 8	Normal	0.7%	0 of 6	3	0	2	0	0
Region 9	Normal	1.2%	0 of 5	4	2	0	1	0
Region 10	Normal	0.6%	0 of 4	2	0	0	1	0

*HHS regions (Region 1 CT, ME, MA, NH, RI, VT; Region 2: NJ, NY, Puerto Rico, U.S. Virgin Islands; Region 3: DE, DC, MD, PA, VA, WV; Region 4: AL, FL, GA, KY, MS, NC, SC, TN; Region 5: IL, IN, MI, MN, OH, WI; Region 6: AR, LA, NM, OK, TX; Region 7: IA, KS, MO, NE; Region 8: CO, MT, ND, SD, UT, WY; Region 9: AZ, CA, Guam, HI, NV; and Region 10: AK, ID, OR, WA).

† Elevated means the % of visits for ILI is at or above the national or region-specific baseline.


‡ National data are for current week; regional data are for the most recent three weeks.

§ Includes all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands.

U.S. Virologic Surveillance: WHO and NREVSS collaborating laboratories located in all 50 states and Washington D.C. report to CDC the number of respiratory specimens tested for influenza and the number positive by influenza type and subtype. The results of tests performed during the current week are summarized in the table below.

	Week 42
No. of specimens tested	1,479
No. of positive specimens (%)	12 (0.8%)
Positive specimens by type/subtype	
Influenza A	10 (83%)
A (2009 H1N1)	0 (0.0%)
A (subtyping not performed)	8 (80%)
A (H3)	2 (20%)
Influenza B	2 (17%)

Influenza Positive Tests Reported to CDC by U.S. WHO/NREVSS Collaborating Laboratories, National Summary, 2011-12 Season



Antigenic Characterization: CDC has antigenically characterized four influenza viruses [four influenza A (H3N2) viruses] collected by U.S. laboratories since October 1, 2011.

Influenza A (H3N2) [4]

- All four were characterized as A/Perth/16/2009-like, the influenza A (H3N2) component of the 2011-12 influenza vaccine for the Northern Hemisphere.

Antiviral Resistance: Testing of 2009 influenza A (H1N1), influenza A (H3N2), and influenza B virus isolates for resistance to neuraminidase inhibitors (oseltamivir and zanamivir) is performed at CDC using a functional assay. Additional 2009 influenza A (H1N1) clinical samples are tested for a single mutation in the neuraminidase of the virus known to confer oseltamivir resistance (H275Y). The data summarized below combine the results of both testing methods. These samples are routinely obtained for surveillance purposes rather than for diagnostic testing of patients suspected to be infected with antiviral resistant virus.

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among 2009 influenza A (H1N1) and A (H3N2) viruses (the adamantanes are not effective against influenza B viruses). As a result of the sustained high levels of resistance, data from adamantane resistance testing are not presented in the table below.


**Neuraminidase Inhibitor Resistance Testing Results
on Samples Collected Since October 1, 2011.**

	Oseltamivir		Zanamivir	
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)
Influenza A (H3N2)	1	0 (0.0)	1	0 (0.0)
Influenza B	1	0 (0.0)	1	0 (0.0)
Influenza A (2009 H1N1)	0	0 (0.0)	0	0 (0.0)

The majority of currently circulating influenza viruses are susceptible to the neuraminidase inhibitor antiviral medications oseltamivir and zanamivir; however, rare sporadic cases of oseltamivir resistant 2009 influenza A (H1N1) and A (H3N2) viruses have been detected worldwide. Antiviral treatment with oseltamivir or zanamivir is recommended as early as possible for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at greater risk for influenza-related complications. Additional information on recommendations for treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at <http://www.cdc.gov/flu/antivirals/index.htm>.


Pneumonia and Influenza (P&I) Mortality Surveillance: During week 42, 6.2% of all deaths reported through the 122-Cities Mortality Reporting System were due to P&I. This percentage was below the epidemic threshold of 6.7% for week 42.

Pneumonia and Influenza Mortality for 122 U.S. Cities Week ending 10/22/2011



Influenza-Associated Pediatric Mortality: No influenza-associated pediatric deaths were reported to CDC during week 42.


**Number of Influenza-Associated Pediatric Deaths by Week of Death:
2008-09 season to present**













Influenza-Associated Hospitalizations: The Influenza Hospitalization Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza related hospitalizations in children (persons younger than 18 years) and adults. The network covers more than 80 counties in the 10 Emerging Infections Program (EIP) states (CA, CO, CT, GA, MD, MN, NM, NY, OR, and TN) and four additional states (MI, OH, RI and UT). FluSurv-NET estimated hospitalization rates will be updated weekly starting later this season.

Outpatient Illness Surveillance: Nationwide during week 42, 1.1% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). This percentage is below the national baseline of 2.4%. (ILI is defined as fever (temperature of 100°F [37.8°C] or greater) and cough and/or sore throat.)

Percentage of Visits for Influenza-like Illness (ILI) Reported by the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Weekly National Summary, September 30, 2007 – October 22, 2011



On a regional level, the percentage of outpatient visits for ILI ranged from 0.6% to 1.9% during week 42. All 10 regions reported a proportion of outpatient visits for ILI below their region-specific baseline levels.



NOTE: Scales differ between regions


*Use of the regional baselines for state data is not appropriate.

ILINet Activity Indicator Map: Data collected in ILINet are used to produce a measure of ILI activity* by state. Activity levels are based on the percent of outpatient visits in a state due to ILI and are compared to the average percent of ILI visits that occur during spring and fall weeks with little or no influenza virus circulation. Activity levels range from minimal, which would correspond to ILI activity from outpatient clinics being below the average, to intense, which would correspond to ILI activity from outpatient clinics being much higher than average.

During week 42, the following ILI activity levels were experienced:

- All 50 states and New York City experienced minimal ILI activity (Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming).
- Data were insufficient to calculate an ILI activity level from the District of Columbia.

**Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet
2011-12 Influenza Season Week 42 ending Oct 22, 2011**



*This map uses the proportion of outpatient visits to health care providers for influenza-like illness to measure the ILI activity level within a state. It does not, however, measure the extent of geographic spread of flu within a state. Therefore, outbreaks occurring in a single city could cause the state to display high activity levels.

Data collected in ILINet may disproportionately represent certain populations within a state, and therefore, may not accurately depict the full picture of influenza activity for the whole state.


Data displayed in this map are based on data collected in ILINet, whereas the State and Territorial flu activity map is based on reports from state and territorial epidemiologists. The data presented in this map is preliminary and may change as more data is received.

Differences in the data presented here by CDC and independently by some state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.

Geographic Spread of Influenza as Assessed by State and Territorial Epidemiologists: The influenza activity reported by state and territorial epidemiologists indicates geographic spread of influenza viruses, but does not measure the severity of influenza activity.

During week 42, the following influenza activity was reported:

- Sporadic influenza activity was reported by the District of Columbia, Puerto Rico, the U.S. Virgin Islands, and 18 states (Alaska, Arkansas, California, Colorado, Connecticut, Florida, Hawaii, Indiana, Louisiana, Massachusetts, Minnesota, Nevada, New York, North Carolina, Ohio, Pennsylvania, Texas, and Wisconsin).
- No influenza activity was reported by Guam and 32 states (Alabama, Arizona, Delaware, Georgia, Idaho, Illinois, Iowa, Kansas, Kentucky, Maine, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, North Dakota, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, and Wyoming).



* This map indicates geographic spread & does not measure the severity of influenza activity

A description of surveillance methods is available at: <http://www.cdc.gov/flu/weekly/overview.htm>
Report prepared: October 28, 2011.

Additional National and International Influenza Surveillance Information

Distribute Project: Additional information on the Distribute syndromic surveillance project, developed and piloted by the International Society for Disease Surveillance (ISDS), now working in collaboration with CDC to enhance and support Emergency Department (ED) surveillance, is available at <http://isdistribute.org/>.

Google Flu Trends: Google Flu Trends uses aggregated Google search data in a model created in collaboration with CDC to estimate influenza activity in the United States. For more information and activity estimates from the U.S. and worldwide, see <http://www.google.org/flutrends/>.

Europe: For the most recent influenza surveillance information from Europe, please see WHO/Europe at <http://www.euroflu.org/index.php> and visit the European Centre for Disease Prevention and Control at http://ecdc.europa.eu/en/publications/surveillance_reports/influenza/Pages/weekly_influenza_surveillance_overview.aspx.

Public Health Agency of Canada: The most up to date influenza information from Canada is available at <http://www.phac-aspc.gc.ca/fluwatch/>.

World Health Organization FluNet: Additional influenza surveillance information from participating WHO member nations is available at http://www.who.int/influenza/gisrs_laboratory/flunet/en/index.html.