Albania Reproductive Health Survey



ALBANIA

Final Report

2002









REPRODUCTIVE HEALTH SURVEY ALBANIA, 2002

FINAL REPORT

Edited by: Leo Morris, Ph.D. MPH Joan Herold, Ph.D. Silva Bino, M.D. Alban Yili, M.D. Danielle Jackson, MPH

Institute of Public Health (IPH) Albania Ministry of Health Institute of Statistics (INSTAT) TIRANA, ALBANIA

Division of Reproductive Health Centers for Disease Control and Prevention (DRH/CDC) ATLANTA, GEORGIA USA

United States Agency for International Development (USAID) United Nations Population Fund, Albania (UNFPA) United Nations Children's Fund (UNICEF)

May, 2005

PREFACE

In Albania, some models of services related to reproductive health have traditionally existed; this is especially the case for mother and child care services. Meanwhile, reproductive health programs are relatively new and began to function after the fall of the old system. During this period of time, Albania has faced a range of health problems which have followed economic and especially social changes in the country.

For 50 years Albania followed a pro-natalist policy, with modern family planning methods banned; and it was almost taboo to discuss sexuality and contraception in public. Abortion was also banned before the year 1991 and half of the maternal deaths in the 80's were caused by abortion complications.

The Ministry of Health of Albania has started to adapt specific policies to cope with increasing risks, which are mostly related to changing life styles, and to meet as well the increasing demands from the population for specific services. Products of such policies are new programs of family planning and programs of sex education. Through these approaches we aim to support the Albanian women and men of reproductive age taking rational decisions for the number of children they want to have, the time they want them, child spacing, and safe sex.

Vice Minister of Health Saemira Pino Only a few years ago the Albanian parliament passed a law on reproductive health, which regulates management and functioning of all services concerning reproductive health in public and private institutions. The law guarantees the rights of every individual and every couple related to reproduction in coherence with national policies and well known international principles.

The reproductive health survey of Albania has provided baseline data for new developments in recent years in this field, developments which have affected lifestyle, legislation, policies and social services. Until this survey was in place there was no comprehensive comparative analysis of the main indicators of family planning and reproductive health, utilizing data from a nation-wide representative sample.

A main objective of this survey is to assess reproductive health status and needs, which will help us design new programs and adjust existing ones according to the needs of the population. I strongly believe that the results presented in this report will serve most managers and professionals who operate in the field of reproductive health, as a important reference in their everyday activities.

EXECUTIVE SUMMARY

ALBANIA REPRODUCTIVE HEALTH SURVEY: 2002

BACKGROUND

The Albania Reproductive Health Survey (RHS) 2002 was conducted by the Institute of Public Health, with the support of the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA) and United Nations Children's Fund (UNICEF), and with the technical assistance of the Division of Reproductive Health (DRH) of the Centers for Disease Control and Prevention (CDC) in Atlanta, Georgia, USA. It is the first national survey on reproductive health, for both female and male respondents, in Albania, and the 18th survey of this type in 13 countries of Eastern Europe and the Former Soviet Union since 1993. The results obtained from this population-based survey provide the Ministry of Health, NGO's and donor agencies the current status of the reproductive health situation in the country and make possible the comparison of results with other national and international surveys.

The Institute of Statistics (INSTAT) provided information from the 2001 census to serve as the sampling frame for the national sample and also was responsible for data processing.

A principal objective of the study was to examine the reproductive health status of the population and needs that can be used to help direct or modify program interventions. Until this survey, relatively little detailed and reliable population-based information was available about important reproductive health topics in the country. The RHS examines patterns and levels of fertility, contraceptive use and method selection, health behaviors, knowledge of HIV/AIDS, attitudes toward specific contraceptive methods, domestic violence and sexual abuse, as well as sex education and sexual behavior of young adults. These data are particularly useful in assisting policy makers and program officials in evaluating health service needs and identifying reproductive health behaviors associated with poor health outcomes.

METHODOLOGY

Results of the Albania RHS are based on inperson face-to-face interviews with 5,697 women and 1,740 men in their homes. The household-based survey was designed to collect information from a representative sample of reproductive age men (15-49 years of age) and women (15-44 years of age), regardless of marital status, who were living in Albania when the survey was conducted in late 2002. Male and female samples were selected independently. For analysis purposes, three strata were constructed for the sample design: Metropolitan Tirana, other urban areas and other rural areas.

As in other countries in Eastern Europe, the survey had a three-stage sampling design: (1) selection of census sectors with probability proportional to the number of households in the 2001 census (2) clusters of households randomly selected in each census sector chosen in the first stage of the sample and (3) random selection of one eligible respondent in each household.

CHARACTERISTICS OF THE SAMPLE

The average size of a household in Albania with at least one eligible female respondent is 5.1, ranging from 4.6 in urban areas to 5.4 in rural areas. Almost two-thirds (65%) of the female sample were currently married compared with 60% of the male sample and 36% of females reported secondary complete or post-secondary education compared with 43% of the male sample; there was a marked differential in educational attainment of urban and rural Albanians. About 80% of both samples reported their religion to be Muslim, but only 5% reported that they attend religious services at least once a month.

FERTILITY TRENDS AND LEVELS

Albania has had a 21% decline in the Total Fertility Rate (TFR) between 1993 and 2002. The survey data show that the TFR has declined from 3.3 children per woman during the three-year period from August 1993 to July 1996 to 2.8 from 1996-1999 to 2.6 from 1999-2002. Women 20-29 years of age at birth contribute 65% of the fertility rate. The TFR for women with post-secondary education is estimated to be 2.0 compared with 2.7 for women with primary school education.

Over 90% of married women (93%) have had a live birth and 11% have had four or more children. The median age of first intercourse for all women is 21.1, first marriage is 21.9 and first live birth is 23.4.

Unfortunately, the survey rate of 73 abortions per 1,000 live births for the three years prior to the survey is 64% lower than the officially reported rate of 200 per 1000 live births. Since the apparent underreporting by survey respondents is

at least 50% and may be as high as 77% (upper limit of 95% confidence interval), and since the underreporting is most likely not random but associated with the characteristics of the respondent, further analysis of the abortion data is probably unreliable and beyond the scope of this report.

MATERNAL and CHILD HEALTH

Nineteen percent of women with births in the past five years reported that they did not have prenatal care during their pregnancy. No prenatal care was highest among rural women (26%), older women (28%), women primary education (25%) and women classified as low socioeconomic status (SES) (26%). Of women with prenatal care, one-fourth (24%) reported a pregnancy complication, including risk of preterm delivery and anemia related to pregnancy as the most frequent complications reported.

Almost all women (94%) gave birth in a medical facility. Ten percent of births to rural women, women older than 34, low SES women and women with birth order 3 or greater were reported to be at home. Fifteen percent of women with no prenatal care reported a home birth. Only 19% of women reported a post-partum care visit following delivery.

Postnatal baby clinic visits were more common with 86% of women reporting a postnatal clinic visit for their baby. The overwhelming majority of babies were breastfed (93%) with a mean duration of 14.3 months. However, exclusive breastfeeding was only 2.8 months on the average and full breastfeeding 4.9 months on the average.

The infant mortality rate (IMR) calculated for the period from August 1992 to July

2002 is 26.2 per 1000 and the under five mortality rate for the same period is 31.9 per 1000. These results are similar to the results of the MICS survey carried out in 2000: infant mortality was 28 per 1000 and the child mortality rate was 33 per 1000. The survey IMR of 26.2/1000 for the period, 1992-2002, is 29% higher than the official rate reported between 1995 and 2000.

AWARENESS AND USE OF CONTRACEPTION

Nine out of 10 Albanian women have heard of at least one modern method and 87% have heard of at least one traditional method, predominantly withdrawal. However, while 81% of women have heard of the condom, only two-thirds of women have heard of oral contraception or tubal ligation. Less than 35% have heard of other modern methods. There are 13 percentage point differences for most modern methods between women living in urban areas vs. rural areas.

Males have principally heard of the condom (89%) and withdrawal (89%). Knowledge of other modern methods is very low, reaching only 33% for oral contraceptives.

Women who have heard of contraception believe that withdrawal is much more effective at preventing pregnancy than are modern methods. Males said that tubal ligation was best at preventing pregnancy followed by the condom and withdrawal.

Contraceptive prevalence is 75% for married women (8% modern methods) and 77% for married men (3% modern methods). For both genders, the predominant method is withdrawal, 67% reported by the females sample and 74% reported by the male sample. Only three groups of married women reported at least a 15% contraceptive prevalence for modern methods: High SES (18%), post-secondary education (17%) and living in metro Tirana (15%). Married men with post-secondary education (13%) and high SES (12%) were the only groups to surpass a level of 9% using modern methods.

Almost three out of four women (73%) using traditional methods thought that withdrawal was more effective at preventing pregnancy than modern methods. Among men using traditional methods, 48% thought withdrawal was more effective than modern methods and 42% thought that they were equally effective.

NEED FOR CONTRACEPTIVE SERVICES

Unmet need for contraception is a very specific estimate that measures the gap between desired fertility levels and the contraceptive practices adopted to ensure that fertility preferences are met. In addition to unmet need for any contraceptive method, the Albania survey also estimated the need for modern contraception – an indicator used in other Eastern European surveys.

It is estimated that 46% of all women and 68% of married women have unmet need for modern contraception if unintended pregnancies were to be prevented. Among married women, since only 8% report using modern contraception, only 12% of the potential "demand" for modern contraception has been met. The highest percentage of "met demand" was for women living in Metro Tirana (22%), women with a post-secondary education (26%) and women classified as high SES (27%).

Two-thirds of women (68%) expressed a desire for more information about contraception. The desire for more information was highest among non-users (75%), young adults 15-24 years of age (84%), and never married women (85%). Ninety percent of women and 77% of men said that information about contraception should be broadcast on radio or television.

REPRODUCTIVE HEALTH KNOWLEDGE AND ATTITUDES

Women said that the ideal number of children for a young family in Albania was 2.6, coincidentally equal to the total fertility rate found in the three years prior to the survey. For men, the ideal number of children was 2.4. Almost all women (96%) and 89% of men agree that both the husband and wife should decide together on how many children a couple should have.

Only 26% of women and 11% of men knew the most likely time during the menstrual cycle that a woman would get pregnant. Only 9% of unmarried women and 10% of 15-19 year olds responded correctly to this question. Also, only about one-fifth of women and men knew that the likelihood of pregnancy was lower if the mother was breastfeeding.

Almost three-fourths of women (72%) and 53% of men agreed that seeking an abortion is a woman's personal decision. Of those not agreeing, 26% of women and 45% of men said that abortion was alright under certain circumstances. Only one percent of females and males said that abortion is never acceptable.

More than 75% of women and more than 84% of men agree that the husband should help with chores if the wife works, the main job of women is housework and every individual should get married. Threequarters of women (75%) and 64% of men agree that a married woman needs her husband's permission to work.

HEALTH BEHAVIORS

Only 24% of women and 14% of men visited a health facility in the past 12 months. Only 16% of those women visiting a health facility received counseling for family planning. The main reasons expressed by women as a barrier in getting medical advice or treatment for themselves were "lack of money" (46%) and "not wanting to go alone" (41%). The two main reasons expressed by men ware "lack of money" (54%) and "did not know where to go" (40%).

Of sexually experienced women, twothirds (69%) have never had a routine gynecological exam. About one-half of women (52%) were aware of breast selfexam, but only 8% have ever practiced breast self-exam. Only 3% of sexually experienced women had ever had a pap smear for cervical cancer screening. Fully 70% of these women had never heard of screening for cervical cancer.

Almost one-half of men (46%) smoke compared with only 3% of women.

YOUNG ADULTS

More than 90 percent of women 15-44 years of age (92%) agree that age appropriate sex education topics concerning human reproduction, contraception and sexually transmitted infections should be taught in school. For men 15-49 years of age, 84% agree.

For young adult women 15-24 years of age, two-thirds (64%) have discussed sex education topics with a parent before they reached age 18, but only 15% discussed HIV/AIDS and 8% discussed contraception.

For young adult men, only 11% discussed any sex education topic with a parent before age 18, and only 9% discussed HIV/AIDS and 2% methods of contraception.

Three fourths (77%) of young adult women and 64% of young adult men said that they were taught some sex education topic in school by age 18. However, only about one-half of females and males received information about HIV/AIDS and only 30% of males and 24% of females received information about contraceptive methods.

One-third of young adult women (32%) reported sexual experience and 14%, or 42% of those with sexual experience, had premarital sex; almost all (99%) reported to be their fiancée or boy friend. Among young adult males, 29% reported having had sexual experience and 27%, or 91% of those with sexual experience, had premarital sex. Most men with premarital sexual experience reported their first partner to be a girl friend (43%), a lover (19%) or a friend (14%). Only 1% reported that their first sexual encounter was with a prostitute.

Four out of five (81%) females said that they or their partner used a contraceptive method at first intercourse, mostly withdrawal, with the following distribution: withdrawal (96%), condoms (3%) and other modern methods (1%). Eight-five percent of men reported contraceptive use at first intercourse, including withdrawal (56%), condoms (43%) and other modern methods (1%).

Eighteen percent of unmarried males and 5% of unmarried females15-24 years of age were sexually active at least once in the past three months. Only 15% of sexually active unmarried males reported using a modern method at last intercourse similar to the 11% of sexually active unmarried females.

Of all sexually experienced men, 72% report two or more lifetime partners versus only 3% of females.

KNOWLEDGE OF HIV/ AIDS TRANSMISSION AND PREVENTION

Almost all women and men of reproductive age (96%) have heard of or are aware of HIV/AIDS; However, only 56% of women and 45% of men believe that HIV can be asymptomatic, and only 17% of women and 33% of men know where HIV tests are provided.

Seventy-three percent of all women and 69% of all men identified monogamy, partner limitation and condom use as prevention measures against HIV/AIDS (UNAIDS indicator no. 1). However only 1% of both women and men had correct knowledge that HIV could be asymptomatic, and is not spread by mosquito bites or through medical treatment (UNAIDS indicator no. 2).

Seventy-nine percent of both women and men say they have no risk of contracting HIV/AIDS; 95% of the women who say that they are not at risk say they are monogamous (41%), not sexually active (32%) or they trust their partner (23%), and 87% of the males give the same three reasons plus 8% saying they use condoms. Of those that think they have some risk, the overwhelming proportion of women (91%) stated that their risk was from medical or dental treatment. Slightly over half of the men (54%) also gave this reason followed by unprotected sex with casual partners (16%).

VIOLENCE AGAINST WOMEN

The data in this report on violence against women, also known as "gender-based violence", represent the first national population-based information on the issue of violence against women in Albania. Women who have ever been married reported both lifetime intimate partner violence (IPV) and IPV during the past 12 months. During their lifetime, among these women, 30% report verbal abuse, 8% physical abuse and 3% sexual abuse; during the past 12 months, the corresponding reports are 23%, 5% and 2%, respectively. Except for sexual abuse, men report inflicting more abuse on their partners than reported by women indicating a possible reluctance of women to report IPV even in a private interview. Lifetime, 33% of men reported inflicting verbal abuse, 14% physical abuse and 1% sexual abuse. In the past 12 months the corresponding reports are 19%, 5% and <1%, respectively.

Of women reporting physical violence in the past year, less than half (46%) talked to anyone about this violence, mostly with family, a relative or a friend. Only 20% talked to the police, a health provider or a legal adviser.

All women were asked if they have been forced to have sexual intercourse against their will during their lifetime. Only 2% of women reported that they have ever experienced forced sexual intercourse, lower than the prevalence reported in other countries of this region. About 90% of these women reported that the perpetrator of forced sex was their husband or partner or ex-husband or ex-partner. It is notable that no woman reported forced sexual intercourse by a casual partner or stranger.

Ι	
II	
Х	
II	
II	
1	0
9	
	CO
5	TN
.7	ENTS

PREFACE	I
EXECUTIVE SUM	MARY III
TABLE OF CONT	ENTS IX
LIST OF TABLES	XIII
LISTOFFIGURES	XXVII
CHAPTER 1:	BACKGROUND
CHAPTER 2:	METHODOLOGY
CHAPTER 3:	CHARACTERISTICS OF THE SAMPLE
CHAPTER 4:	FERTILITY AND PREGNANCY
CHAPTER 5:	MATERNAL AND CHILD HEALTH

CHAPTER 6:	CONTRACEPTION AWARENESS AND KNOWLEDGE OF USE
CHAPTER 7:	CONTRACEPTIVE USE AND PREFERENCE
CHAPTER 8:	NEED FOR CONTRACEPTIVE SERVICES AND CONTRACEPTIVE COUNSELING
CHAPTER 9:	 OPINIONS ABOUT CONTRACEPTION AND ABORTION
CHAPTER 10:	REPRODUCTIVE HEALTH KNOWLEDGE AND ATTITUDES

CHAPTER 11:	HEALTH BEHAVIORS1(Alban Yili, Florina Serbanescu)Use of Health Care and Preventive ServicesPrevalence of routine gynecologic visits among sexuallyexperienced womenBreast Self ExamCervical Cancer ScreeningPrevalence of Selected Health ProblemsImpaired FecunditySmoking and Alcohol Consumption	73
CHAPTER 12:	SEX EDUCATION	,,
CHAPTER 13:	YOUNG ADULT SEXUAL AND CONTRACEPTIVE EXPERIENCE	31
CHAPTER 14:	KNOWLEDGE AND EXPERIENCE OF SEXUALLY TRANSMITTED INFECTIONS AND KNOWLEDGE OF HIV/AIDS TRANSMISSION AND PREVENTION	ls Is

CHAPTER 15:	VIOLENCE AGAINST WOMEN
	(Mary Goodwin, Silva Bino, Florina Serbanescu)
	Introduction
	Comparison of Violence Across Countries of Eastern Europe
	History of Witnessing or Experience Abuse During Childhood
	Women's Experience of Intimate Partner Violence
	Male Infliction of Intimate Partner Violence
	Characteristics of Physical or Sexual Violence Experienced by
	Women in the Past Year
	Women's Attitudes and Beliefs About the Effects of Violence
	Prevalence of Forced Sexual Intercourse
GLOSSARY	i
REFERENCES	V
APPENDIX A:	SAMPLING ERROR ESTIMATES xiii
APPENDIX B:	PARTICIPANT INSTITUTIONS xvii
APPENDIX C:	FEMALE QUESTIONNAIRE xxi
APPENDIX D:	MALE QUESTIONNAIRE lxxiii

List of Tables

Table 2.1 A Results of Household Visits and Interview Status of Eligible Women by Stratum (Percent Distribution) Reproductive Health Survey: Albania 2002

Table 2.1 B Results of Household Visits and Interview Status of Eligible Men by Stratum (Percent Distribution) Reproductive Health Survey: Albania 2002

Table 3.1 A Size of Households with at Least One Eligible Respondent by Stratum (Percent Distribution) Female Sample, Reproductive Health Survey: Albania 2002

Table 3.1 B Size of Households with at Least One Eligible Respondent by Stratum (Percent Distribution). Male Sample, Reproductive Health Survey: Albania 2002

Table 3.2 A Percentage of Households with Basic Household Amenities and Goods, by Stratum, for Women Aged 15–44 Years Reproductive Health Survey: Albania 2002

Table 3.2 B Percentage of Households with Basic Household Amenities and Goods, by Stratum, for Men Aged 15–49 Years Reproductive Health Survey: Albania 2002

Table3.3APercentDistributionofCharacteristics of Female Sample by ResidenceandStratumReproductiveHealthSurvey:Albania 2002

Table3.3BPercentDistributionofCharacteristicsofMaleSamplebyResidenceandStratumReproductiveHealthSurvey:Albania2002

Table 3.4 A Percent Distribution of Marital Status by Selected Characteristics, for Women Aged 15–44 Years Reproductive Health Survey: Albania 2002

Table 3.4 B Percent Distribution of Marital Status by Selected Characteristics, for Men Aged 15–49 Years Reproductive Health Survey: Albania 2002

Table 4.1 Three-Year Period Age-Specific Fertility Rates for Several Time Periods Among All Women Aged 15–44 Reproductive Health Survey: Albania 2002 Table 4.2 Three-Year Period* Age–Specific and Total Fertility Rates, Among Women Aged 15–44 Reproductive and Demographic Health Surveys (RHS and DHS) in Selected Eastern European and Former Soviet Union Countries Albania Reproductive Health Survey 2002, Final Report

Table 4.3 A Age-Specific Fertility Rates and Total Fertility Rates by Selected Characteristics Among All Women Aged 15–44 Reproductive Health Survey: Albania 2002

Table 4.3 B Age-Specific Fertility Rates and Total Fertility Rates by Selected Characteristics Among All Men Aged 15–49 Reproductive Health Survey: Albania 2002

Table 4.4 A Percent Distribution of Number of Live Births by Current Age of Respondents Among all Women and Among Married Women Aged 15–44 Reproductive Health Survey: Albania 2002

Table 4.4 B Percent Distribution of Number of Live Births by Current Age of Respondents Among All Men and Among Married Men Aged 15–49 Reproductive Health Survey: Albania 2002

Table 4.5 A Percent of Women Aged 15–44 Who Had Their First Sexual Relation, First Marriage, and First Birth Before Selected Ages, and Median Age at These Events, by Current Age Reproductive Health Survey: Albania 2002

Table 4.5 B Percent of Men Aged 15–49 Who Had Their First Sexual Relation and First Marriage Before Selected Ages and Median Age at These Events, by Current Age Reproductive Health Survey: Albania 2002

Table 4.6 Median Age at First Sexual Intercourse, First Marriage and First Birth Among Women Aged 15–44 and Men Aged 15–49 by Selected Characteristics Reproductive Health Survey: Albania 2002

Table 4.7 Sexual Activity Status by Current Marital Status for Women Aged 15–44 and Men Aged 15–49 (Percent Distribution) Reproductive Health Survey: Albania 2002 Table 4.8 Abortions * per 1000 Births Reported in Reproductive Health Survey and By Albanian Institute of Statistics (INSTAT)† Three Year Period: 1999–2001

Table 4.9 A Planning Status of the Last Pregnancy Among Women 15–44 Years of Age With at Least One Pregnancy Since January 1997, by Selected Characteristics (Percent Distribution) Reproductive Health Survey: Albania 2002

Table 4.9 B Planning Status of the Last Live Birth Among Men 15–49 Years of Age With Partner Having at Least One Live Birth Since January 1997, by Selected Characteristics (Percent Distribution) Reproductive Health Survey: Albania 2002

Table 4.10 A Fertility Preferences of Currently Married Women Aged 15–44 Years By Number of Living Children and by Age Group (Percent Distribution) Reproductive Health Survey: Albania 2002

Table 4.10 B Fertility Preferences of Currently Married Men Aged 15–49 Years By Number of Living Children and by Age Group (Percent Distribution) Reproductive Health Survey: Albania 2002

Table 4.11 Percentage of Fecund Married Women Reporting They Want no More Children by Number of Living Children and Selected Characteristics Fecund Women 15–44 Years of Age. Reproductive Health Survey: Albania 2002

Table 5.1 Prenatal Care by Pregnancy Trimester of First Visitand Number of Prenatal Visits, for Births in 1997–2002, by Selected Characteristics (Percent Distribution) Reproductive Health Survey: Albania 2002

Table 5.2 Percentage Counseled on Specific Topics During Prenatal Care Visits for Births in 1997–2002 by Selected Characteristics, Among Women with Any Prenatal Care Reproductive Health Survey: Albania 2002

Table 5.3 Percentage Experiencing Selected Procedures During Prenatal Care Visits for Births in 1997–2002 by Selected Characteristics, Among Women with Any Prenatal Care. Reproductive Health Survey: Albania 2002 Table 5.4 Percentage with Routine Measurement of Blood Pressure (BP) During Pregnancy, Reported High Blood Pressure (HBP) During Pregnancy, and Hospitalization Rate for HBP, for Births in 1997–2002, Among Women with Any Prenatal Care. Reproductive Health Survey: Albania 2002

Table 5.5 Use of Ultrasound Exams During Pregnancy and Time of First Ultrasound Exam, By Selected Characteristics, for Births in 1997– 2002, Among Women with Any Prenatal Care. Reproductive Health Survey: Albania 2002

Table 5.6 Percentage of Pregnancy Complications that Required Medical Attention, by Selected Characteristics, for Births in 1997–2002, Among Women with Any Prenatal Care. Reproductive Health Survey: Albania 2002

Table 5.7 Place of Delivery for Births in 1996–2001 by Selected Characteristics (PercentDistribution) Reproductive Health Survey:Albania 2002

Table 5.8 Average Time Between Admission and Delivery and Percent Distribution of Number of Nights Spent in a Medical Facility, for Births in 1997–2002. Reproductive Health Survey: Albania 2002

Table 5.9 Percentage of Cesarean Deliveries by Selected Characteristics, for Births in 1997– 2002, Among Women Currently Aged 15–44. Reproductive Health Survey: Albania 2002

Table 5.10 Percentage of Poor Birth Outcomes by Selected Characteristics, for Births in 1997– 2002, Among Women Currently Aged 15–44. Reproductive Health Survey: Albania 2002

Table 5.11 Percentage Who Attended Postpartum Care Visit within Specified Time Periods by Selected Characteristics, for Births in 1997– 2002, Among Women Currently Aged 15–44. Reproductive Health Survey: Albania 2002

Table 5.12 Percentage of Women with InformationReceived During Postpartum Care by SelectedCharacteristics, for Births in 1997–2002.Reproductive Health Survey: Albania 2002

Table 5.13 Time Between Delivery and First Baby Clinic Visit by Selected Characteristics, for Births in 1997–2002. Reproductive Health Survey: Albania 2002

Table 5.14 Percentage of Babies With Birth Certificates Issued and Time Between Delivery and Certificate Issued for Births in 1997–2002. Reproductive Health Survey: Albania 2002

Table 5.15 Percentage of Women ReportingPostpartumComplications, bySelectedCharacteristics, forBirths in 1997–2002.Reproductive Health Survey: Albania 2002

Table 5.16 Percent of Children Ever Breastfed and Percent Distribution of Initiation of Breastfeeding, by Selected Characteristics, for Births in 1997–2002. Reproductive Health Survey: Albania 2002

Table 5.17 Mean Duration of Breastfeeding in Months, by Type of Breastfeeding and Selected Characteristics, for Births in 1997–2002. Reproductive Health Survey: Albania, 2002

Table 5.18 Infant and Child Mortality Rates (Infant and Child Deaths per 1,000 Live Births) by Selected Characterisitcs, for Children Born August 1992–July 2002. Reproductive Health Survey: Albania 2002

Table 6.1 A Percentage of Women Aged 15–44 Who Have Heard of Specific Methods ofContraception by Selected CharacteristicsReproductive Health Survey: Albania, 2002

Table 6.1 B Percentage of Men Aged 15–49 Who Have Heard of Specific Methods of Contraception by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 6.2 A Percentage of Women Aged 15–44 Who Say They Know How Specific Methods of Contraception Are Used by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 6.2 B Percentage of Men 15–49 Years Who Say They Know How Specific Methods of Contraception are Used by Selected Characteristics. Reproductive Health Survey: Albania, 2002 Table 6.3 A Percentage of Women 15–44 Years Who Say They Know Where to Get Specific Contraceptive Methods by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 6.3 B Percentage of Men 15–49 Years Who Say They Know Where to Get Specific Contraceptive Methods by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 6.4 A Percent Distribution of Most Important Source of Information about Contraception by Specific Method Among Women Aged 15–44 Who Have Heard About Specific Methods of Contraception. Reproductive Health Survey: Albania, 2002

Table 6.4 B Percent Distribution of Most Important Source of Information about Contraception by Specific Method Among Men Aged 15–49 Who Have Heard About Specific Methods of Contraception. Reproductive Health Survey: Albania, 2002

Table 6.5 A Percentage Distribution of Women 15–44 by Their Opinion About Contraceptive Effectiveness if the Method is Used Correctly and Consistently. Reproductive Health Survey: Albania, 2002

Table 6.5 B Percentage Distribution of Men 15–49 by Their Opinion About Contraceptive Effectiveness if the Method is Used Correctly and Consistently. Reproductive Health Survey: Albania, 2002

Table 7.1 Percent of Women Currently Using Any Contraceptive Method by Marital Status Among Women Aged 15–44 Reproductive and Demographic Health Surveys (RHS and DHS) In Selected Eastern European and Former Soviet Union Countries. Albania Reproductive Health Survey 2002, Final Report

Table 7.2 A Current Use of Contraception Among Women 15–44 Years of Age, by Marital Status (Percent Distribution). Reproductive Health Survey: Albania, 2002 Table 7.2 B Current Use of Contraception Among Men 15–49 Years of Age, by Marital Status (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 7.3 A Percentage Currently Using Modern and Traditional Methods by Selected Characteristics Currently Married Women Aged 15–44 Years. Reproductive Health Survey: Albania, 2002

Table 7.3 B Percentage Currently Using Modern and Traditional Methods by Selected Characteristics Currently Married Men Aged 15– 49 Years. Reproductive Health Survey: Albania, 2002

Table 7.4 A Current Use of Specific Contraceptive Methods by Selected Characteristics Currently Married Women Aged 15–44 Years. Reproductive Health Survey: Albania, 2002

Table 7.4 B Current Use of Specific Contraceptive Methods by Selected Characteristics Currently Married Men Aged 15–49 Years. Reproductive Health Survey: Albania, 2002

Table 7.5 Source of Supply for Last Modern Contraceptive Used Among Currently Married Women Aged 15–44 Years Currently Using a Modern Method by Specific Methods (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 7.6 Percent Distribution of Type of Counseling by a Health Care Provider for Current Contraceptive Method Among Married Women Aged 15–44 Currently Using a Modern Method. Reproductive Health Survey: Albania, 2002

Table 7.7 A Percentage of Contraceptive Users Who Stated that Selected Reasons Read to Them Were Very Important or Somewhat Important When Deciding To Use a Non-Supplied Method Instead of a Modern Method, by Selected Characteristics Women Aged 15–44 Currently Using Traditional Methods. Reproductive Health Survey: Albania, 2002

Table 7.7 B Percentage of Contraceptive UsersWho Stated That Selected Reasons Read to ThemWere Very Important or Somewhat Important

When Deciding To Use a Non-Supplied Method Instead of a Modern Method, by Selected Characteristics Men Aged 15–49 Currently Using Traditional Methods. Reproductive Health Survey: Albania, 2002

Table 7.8 A Perceived Effectiveness of Traditional Methods Compared to Modern Methods by Selected Characteristics Women Aged 15–44 Currently Using Traditional Methods (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 7.8 B Perceived Effectiveness of Traditional Methods Compared to Modern Methods by Selected Characteristics Men Aged 15–49 Currently Using Traditional Methods (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 7.9 A Most Commonly Cited Reasons for Not Currently Using Contraception Among Currently Married Women Aged 15–44 Years (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 7.9 B Most Commonly Cited Reasons for Not Currently Using Contraception by Age Group Among Currently Married Men Aged 15– 49 Years (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 7.10 A Desire to Use Contraception in the Future by Number of Living Children Fecund Currently Married Women Aged 15–44 Years Who Are Not Using Contraception (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 7.10 B Desire to Use Contraception in the Future by Number of Living Children Fecund Currently Married Men Aged 15–49 Years Who Are Not Using Contraception (Percent Distribution). Reproductive Health Survey: Albania, 2002 Table 7.11 Desire to Use Contraception in the Future by Fertility Preferences Fecund Currently Married Women Aged 15–44 Years Who Are Not Using Contraception (Percent Distribution). Reproductive Health Survey: Albania, 2002

Table 8.1 Potential Demand For Family Planning (FP) Services by Age Group And Marital Status (Percent Distributions and Percentage). Reproductive Health Survey: Albania, 2002

Table 8.2 Percentage of Women Aged 15–44Years in Need of Any or Modern ContraceptiveMethodsbySelectedCharacteristics.Reproductive Health Survey: Albania, 2002

Table 8.3 Percentage of Unmet Need for Contraception Among Fecund Married Women of Reproductive Age by Future Fertility Preferences *. Reproductive Health Survey: Albania, 2002

Table 8.4 Percentage of Potential Demand for Modern Contraceptive Methods Satisfied Married Women 15–44 Years of Age At Risk of an Unintended Pregnancy By Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 8.5 Advise on Using Modern Methods by Method and Type of Counseling: Ever Users of Modern Methods. Reproductive Health Survey: Albania, 2002

Table 8.6 Counseling and Contraception Offered at the Time of Legally Performed Abortions. Reproductive Health Survey: Albania, 2002

Table 9.1 Percentage Who Want More Information about Contraception by Selected Characteristics Among Women Aged 15–44 and Men Aged 15– 49. Reproductive Health Survey: Albania, 2002

Table 9.2 A Percent Distribution of Women's Opinion on the Best Source of Contraceptive Information by Selected Characteristics Among Women 15–44 Who Want More Information About Contraception. Reproductive Health Survey: Albania, 2002

Table 9.2 B Percent Distribution of Men's Opinion on the Best Source of Contraceptive Information by Selected Characteristics Among Men Aged 15–49 Who Want More Information About Contraception. Reproductive Health Survey: Albania, 2002 Table 9.3 Percentage Who Think That Information about Contraception Should be Broadcast on Radio or Television by Selected Characteristics Among Women Aged 15–44 and Men Aged 15– 49. Reproductive Health Survey: Albania, 2002

Table 9.4 Percentage Who Agree with Selected Statements Concerning Possible Advantages and Disadvantages of Using the Pill, by Residence and Education Among Women Aged 15–44 Who Have Heard of the Pill. Reproductive Health Survey: Albania 2002

Table 9.5 Percentage Who Agree with Selected Statements Concerning Possible Advantages and Disadvantages of Using the IUD, by Residence and Education Among Women Aged 15–44 Who Have Heard of the IUD. Reproductive Health Survey: Albania 2002

Table 9.7 Percent Distribution of Opinion of Degree of Risk to a Woman's Health From Using the Pill by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania, 2002

Table 9.8 A Percent Distribution of Opinion of Degree of Risk to a Woman's Health From Using the Condom by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania, 2002

Table 9.8 B Percent Distribution of Opinion of Degree of Risk to a Woman's Health From Using the Condom by Selected Characteristics Among Men Aged 15–49. Reproductive Health Survey: Albania, 2002

Table 9.9 Percent Distribution of Opinion of Degree of Risk to a Woman's Health from Tubal Ligation by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania, 2002 Table 9.10 A Percent Distribution of Opinion of Degree of Risk to a Woman's Health From Abortion by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania, 2002

Table 9.10 B Percent Distribution of Opinion of Degree of Risk to a Woman's Health from Abortion by Selected Characteristics Among Men Aged 15– 49. Reproductive Health Survey: Albania, 2002

Table 10.1 Mean Ideal Number of Children for a Young Family in Albania by Selected Characteristics Among Women Aged 15–44 and Men Aged 15–49. Reproductive Health Survey: Albania, 2002

Table 10.2 A Percent Distribution of Women's Opinion on the Most Likely Time in the Menstrual Cycle for a Woman to Become Pregnant by Selected Characteristics Among Women 15–44. Reproductive Health Survey: Albania, 2002

Table 10.2 B Percent Distribution of Men's Opinion on the Most Likely Time In the Menstrual Cycle for a Woman to Become Pregnant by Selected Characteristics Among Men Aged 15– 49. Reproductive Health Survey: Albania, 2002

Table 10.3 A Percent Distribution of Women's Opinion on the Likelihood of Pregnancy While Breastfeeding by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania 2002

Table 10.3 B Percent Distribution of Men's Opinion on the Likelihood of Pregnancy While Breastfeeding by Selected Characteristics Among Men Aged 15–44. Reproductive Health Survey: Albania 2002

Table 10.4 A Percentage Distribution of Women's Agreement that Abortion Is a Woman's Personal Decision by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania, 2002

Table 10.4 B Percentage Distribution of Men's Agreement that Abortion Is a Woman's Personal Decision by Selected Characteristics Among Men Aged 15–49. Reproductive Health Survey: Albania, 2002 Table 10.5 Percent Distribution of Opinion Regarding Abortion Under Selected Circumstances by Selected Characteristics Among Women and Men of Reproductive Age Who Do Not Believe That Abortion Is Always a Woman's Personal Decision. Reproductive Health Survey: Albania, 2002

Table 10.6 A Percentage Who Agree with the Acceptability of Abortion Under Selected Circumstances by Selected Characteristics Among Women Aged 15–44 Who Do Not Believe That Abortion is Always a Woman's Personal Decision. Reproductive Health Survey: Albania, 2002

Table 10.6 B Percentage Who Agree with the Acceptability of Abortion Under Selected Circumstances by Selected Characteristics Among Men Aged 15–49 Who Do Not Believe That Abortion Is Always a Woman's Personal Decision. Reproductive Health Survey: Albania, 2002

Table 10.7 A Percent Distribution of Women's Opinion on What a Woman Should Do If a Pregnancy is Unwanted by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania, 2002

Table 10.7 BPercent Distribution of Men's Opinion on What a Woman Should Do If a Pregnancy is Unwanted by Selected Characteristics Among Men Aged 15–49. Reproductive Health Survey: Albania, 2002

Table 10.8 Percentage Who Agree with Statements on Gender Norms by Residence and Education Among Women Aged 15–44 and Men Aged 15– 49. Reproductive Health Survey: Albania 2002

Table 10.9 A Percentage Who Agree with Statements on Gender Norms by Selected Characteristics Among Women Aged 15–44. Reproductive Health Survey: Albania, 2002

Table 10.9 BPercentage Who Agree withStatements on Gender Norms by SelectedCharacteristics Among Men Aged 15–49.Reproductive Health Survey: Albania, 2002

Table 10.10 Percent Distribution of Opinion on Who Should Decide How Many Children A Couple Will Have by Selected Characteristics Among Women Aged 15–44 and Men Aged 15– 49. Reproductive Health Survey: Albania 2002

Table 11.1 A Percentage Visiting Any Health Facility* in the Past 12 Months, by Selected Characteristics Women Aged 15–44 Years. Reproductive Health Survey: Albania 2002

Table 11.1 B Percentage Visiting Any Health Facility* in the Past 12 Months, by Selected Characteristics, Among Men Aged 15–49 Years. Reproductive Health Survey: Albania 2002

Table 11.2 A Percentage of Women Who Visited a Health Facility in the Past 12 Months That Received Counseling For Family Planning Methods by Type of Facility and Selected Characteristics, Women Aged 15–44 Years. Reproductive Health Survey: Albania 2002

Table 11.2 B Percentage of Men Aged 15–49 Years Who Visited a Health Facility in Past 12 Months That Received information/Counseling on Selected Topics, By Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 11.3 A Percentage Agreeing that Selected Factors May Be a Major Problem Preventing WomenFromGettingMedicalAdviceorTreatment for Themselves, by Selected Characteristics, Among Women Aged 15–44 Years. Reproductive Health Survey: Albania 2002

Table 11.3 B Percentage Agreeing that Selected Factors May Be a Major Problem Preventing Men From Getting Medical Advice or Treatment for Themselves, by Selected Characteristics, Among Men Aged 15–49 Years. Reproductive Health Survey: Albania 2002

Table 11.4 Time since Last Routine Gynecologic Examination by Selected Characteristics, Among Women Aged 15–44 Years Who Have Ever Had Sexual Intercourse. Reproductive Health Survey: Albania 2002 Table 11.5 Most Important Reason That Women Have Never Had a Routine Gynecologic Exam, by Selected Characteristics, Among Women Aged 15–44 Years Who Have Ever Had Sexual Intercourse. Reproductive Health Survey: Albania 2002

Table 11.6 Percentage with Awareness of Breast Self-Examination (BSE) and Frequency of BSE, by Selected Characteristics, Among Women Aged 15–44 Years Who Have Ever Had Sexual Intercourse. Reproductive Health Survey: Albania 2002

Table 11.7 Frequency of Cervical Cancer Screening by Selected Characteristics, Among Women Aged 15–44 Years Who Have Ever Had Sexual Intercourse. Reproductive Health Survey: Albania 2002

Table 11.8 Main Reason that Women Have Never Had a Pap Smear, by Selected Characteristics, Among Women Aged 15–44 Years Who Have Ever Had Sexual Intercourse. Reproductive Health Survey: Albania 2002

Table 11.9 A Percentage of Women Aged 15– 44 Years Who Have Been Told by a Doctor That They Have Selected Health Problems, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 11.9 B Percentage of Men Aged 15–49 Years Who Have Been Told by a Doctor That They Have Selected Health Problems, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 11.10 Percentage of Currently Married Women Aged 15–44 Years Who Reported FecundityImpairment,bySelectedCharacteristics. Reproductive Health Survey: Albania 2002

Table 11.11 Percentage of Currently Married Women Aged 15–44 Years Who Reported Fecundity Impairment by Type of Problem and Selected Characteristics. Reproductive Health Survey: Albania 2002 Table 11.12 A Percentage of Women Aged 15– 44 Who Have Ever Smoked and Who Currently Smoke by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 11.12 B Percentage of Men Aged 15–49 Who Have Ever Smoked and Who Currently Smoke, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 11.13 A Percentage of Women Aged 15–44 Who Report Having a Drink Containing Alcohol in past 3 Months, and Frequency of Drinking, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 11.13 B Percentage of Men Aged 15–49 Who Report Having a Drink Containing Alcohol in Past 3 Months, and Frequency of Drinking, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.1 A Percentage of Women Aged 15–44 Who Agree Certain Sex Education Topics Be Taught in School, By Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.1 B Percentage of Men Aged 15–49 Who Agree Certain Sex Education Topics Be Taught in School, By Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.2 A Percent Distribution of Best Age to Start School-Based Courses on Human Reproduction and Contraception, as Reported by Women Aged 15–44 Who Are in Favor of Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.2 B Percent Distribution of Best Age to Start School-Based Courses on Human Reproduction and Contraception, as Reported by Men Aged 15–49 Who Are in Favor of Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.3 A Percent Distribution of Best Age to Start School-Based Courses on Sexually Transmitted Infections, as Reported by Women 15–44 Who are in Favor of Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002 Table 12.3 B Percent Distribution of Best Age to Start School-Based Courses on Sexually Transmitted Infections, as Reported by Men 15– 49 Who are in Favor of Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.4 A Percentage of Young Adult Women Aged 15–24 Who Discussed The Indicated Family Life Education Topics With a Parent Before They reached Age 18, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.4 B Percentage of Young Adult Men Aged 15–24 Who Discussed The Indicated Family Life Education Topics With a Parent Before They Reached Age 18, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.5 A Percentage of Young Adult Women Aged 15–24 Who Were Taught Indicated Family Life Education Topics in School Before They Reached Age 18, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.5 B Percentage of Young Adult Men Aged 15–24 Who Were Taught Indicated Family Life Education Topics in School Before They Reached Age 18, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.6 A Percentage of Young Adult Women Aged 15–24 Who Were Taught in School About Indicated Family Life Education Topics by Specific Ages. Reproductive Health Survey: Albania 2002

Table 12.6 B Percentage of Young Adult Men Aged 15–24 Who Were Taught in School About Indicated Family Life Education Topics by Specific Ages. Reproductive Health Survey: Albania 2002

Table 12.7 A Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Women 15–24 Who Had Received Any Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002 Table 12.7 B Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Men 15–24 Who Had Received Any Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.8 A Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Women 15–24 Who Never Received Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.8 B Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Men 15–24 Who Never Received Sex Education in School, by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 12.9 A Knowledge of Young Adult Women Aged 15–24, on Selected Reproductive Health Issues by Whether or Not Specific Topics Were Discussed With a Parent or Taught in School (Percent Distribution). Reproductive Health Survey: Albania 2002

Table 12.9 B Knowledge of Young Adult Men Aged 15–24, on Selected Reproductive Health Issues by Whether or Not Specific Topics Were Discussed With a Parent or Taught in School (Percent Distribution). Reproductive Health Survey: Albania 2002

Table 13.1 A Percent Distribution of Reported Sexual Experience by Marital Status at Time of First Sexual Experience by Selected Characteristics Among Young Women Aged 15– 24. Reproductive Health Survey: Albania, 2002

Table 13.1 B Percent Distribution of Reported Sexual Experience by Marital Status at Time of First Experience by Selected Characteristics, Among Young Men Aged 15–24. Reproductive Health Survey: Albania, 2002 Table 13.2 Percent Distribution of Reported Sexual Experience by Marital Status at Time of First Sexual Experience by Current Age Among Young Women Aged 15–24 Reproductive and Demographic Health Surveys (RHS and DHS) In Selected Eastern European and Former Soviet Union Countries. Albania Reproductive Health Survey 2002, Final Report

Table 13.3 A Percent Distribution of Relationship to Partner at First Sexual Intercourse Among Sexually Experienced Young Women Aged 15– 24. Reproductive Health Survey: Albania, 2002

Table 13.3 B Percent Distribution of Relationship to Partner at First Sexual Intercourse Among Sexually Experienced Young Men Aged 15–24. Reproductive Health Survey: Albania, 2002

Table 13.4 A Percent of Women Aged 15–24 Who Had Their First Sexual Intercourse Before Selected Ages, by Various Characteristics. Reproductive Health Survey: Albania 2002

Table 13.4 B Percent of Men Aged 15–24 Who Had Their First Sexual Intercourse Before Selected Ages, by Various Characteristics. Reproductive Health Survey: Albania 2002

Table 13.5 A Percent Distribution of Age Difference Between Partners at First Sexual Intercourse Among Sexually Experienced Young Women Aged 15–24 by Residence, Age at first Sex and Marital Status at First Sex. Reproductive Health Survey: Albania, 2002

Table 13.5 B Percent Distribution of Age Difference Between Partners at First Sexual Intercourse Among Sexually Experienced Young Men Aged 15–24 by Residence, Age at first Sex and Marital Status at First Sex. Reproductive Health Survey: Albania, 2002

Table 13.6 A Percent Who Used Contraception at First Sexual Intercourse and Percent Distribution of Method Used Among Sexually Experienced Young Women Aged 15–24. Reproductive Health Survey: Albania, 2002 Table 13.6 B Percent Who Used Contraception at First Sexual Intercourse And Percent Distribution of Method Used Among Sexual Experienced Young Men Aged 15–24. Reproductive Health Survey: Albania, 2002

Table13.7PercentDistributionofMostCommonlyCitedReasonsforNotUsingContraception at First Sexual Intercourse AmongSexuallyExperiencedYoungWomenAged15–24byMaritalStatusatFirstSexualIntercourse.ReproductiveHealthSurvey:Albania, 2002

Table 13.8 Percent Distribution of Current Sexual Activity Status by Sex, Current Marital Status and Age Group Among Young Adults Aged 15– 24. Reproductive Health Survey: Albania, 2002

Table 13.9 A Percent Who Used Contraception at Last Sexual Intercourse and Percent Distribution of Methods Used by Current Marital Status and Age Group Among Sexually Experienced Young Women Aged 15–24. Reproductive Health Survey: Albania, 2002

Table 13.9 B Percent Who Used Contraception at Last Sexual Intercourse and Percent Distribution of Methods Used by Current Marital Status and Age Group Among Sexually Experienced Young Men Aged 15–24. Reproductive Health Survey: Albania, 2002

Table 13.10 Percent Distribution of Main Reason Not Currently Using Contraception by Sex and Current Marital Status Among Sexually Active Young Adults Aged 15–24. Reproductive Health Survey: Albania, 2002

Table 13.11 A Percent Distribution of Number of Sexual Partners in Last Three Months and in Lifetime by Current Marital Status and Education Level Among Sexually Experienced Young Women Aged 15–24. Reproductive Health Survey: Albania, 2002

Table 13.11 B Percent Distribution of Number of Sexual Partners in Last Three Months And in Lifetime by Marital Status and Education Level Among Sexually Experienced Young Men Aged 15–24. Reproductive Health Survey: Albania, 2002 Table 13.12 Percent Agreeing With Statements About Condoms and Condom Use, by Sex and Condom Experience for Sexually Experienced Young Adults Aged 15–24 Years. Reproductive Health Survey: Albania, 2002

Table 13.13 A Percent Agreeing With Hypothetical Responses to a Partner Requesting Condom Use During Sex, by Selected Characteristics, Among Sexually Experienced Young Women 15–24 Years of Age. Reproductive Health Survey: Albania, 2002

Table 13.13 B Percent Agreeing With Hypothetical Responses to a Partner Requesting Condom Use During Sex, by Selected Characteristics, Among Sexually Experienced Young Men 15–24 Years of Age. Reproductive Health Survey: Albania, 2002

Table 13.14 Percent Who Have Ever Talked to a Partner About Using Condoms, by Sex and Selected Characteristics for Sexually Experienced Young Adults 15–24 Years of Age. Reproductive Health Survey: Albania, 2002

Table 14.1 A Percentage of Women Aged 15–44 Who Have Heard of Specified Sexually Transmitted Infections by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.1 B Percentage of Men Aged 15–49 WhoHave Heard of Specified Sexually TransmittedInfections by Selected Characteristics.Reproductive Health Survey: Albania, 2002

Table 14.2 A Percentage of Women Aged 15–44 With Knowledge of Symptoms Associated With STIs, Other Than HIV/AIDS, in a Woman by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.2 B Percentage of Men Aged 15–49 With Knowledge of Symptoms Associated With STIs, Other Than HIV/AIDS, in a Man by Selected Characteristics. Reproductive Health Survey: Albania, 2002 Table 14.3 A Most Important Source of Information About Sexually Transmitted Infections All Women Aged 15–44 Who Have Heard of at Least One STI by Selected Characteristics. Reproductive Health Survey: Albania, 2002 (Percent Distribution)

Table 14.3 B Most Important Source of Information on Sexually Transmitted Infections All Men Aged 15–49 Who Have Heard of at Least One STI by Selected Characteristics. Reproductive Health Survey: Albania, 2002 (Percent Distribution)

Table 14.4 A Percentage of Women of Reproductive Age Who Had Received Radio And Television Messages About HIV/AIDS and Other STIs During The Past Six Months By Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.4 B Percentage of Men of Reproductive Age Who Had Received Radio And Television Messages About HIV/AIDS and Other STIs During The Past Six Months By Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.5 A Percentage of Women Aged 15–44 Years with a Diagnosis of Specified Sexually Transmitted Infections by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.5 B Percentage Men Aged 15–49 Witha Diagnosis of Specified Sexually TransmittedInfectionsbySelectedCharacteristics.Reproductive Health Survey: Albania, 2002

Table 14.6 A Level of Awareness, Testing, Diagnosis, And Treatment For STIs Among Women Aged 15–44 Years Who Have Ever Had Sexual Intercourse by Specific Sexually Transmitted Infections. Reproductive Health Survey: Albania, 2002

Table 14.6 B Level of Awareness, Testing, Diagnosis, And Treatment For STIs Among Men Aged 15–49 Years Who Have Ever Had Sexual Intercourse by Specific Sexually Transmitted Infections. Reproductive Health Survey: Albania, 2002 Table 14.7 A Percent Distribution of Women Aged 15–44 Who Have Heard of at Least One Sexually Transmitted Infection By Self-Perceived Risk of Acquiring an STI by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.7 B Percent Distribution of Men Aged 15–49 Who Have Heard of at Least One Sexually Transmitted Infection By Self-Perceived Risk of Acquiring an STI by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.8 A Percentage of Women Aged 15–44 Who Have Heard of HIV/AIDS, Who Believe HIV/AIDS Infection Can Be Asymptomatic, and Who Know Where HIV Testing Is Provided by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.8 B Percentage of Men Aged 15–49 Who Have Heard of HIV/AIDS, Who Believe HIV/AIDS Infection Can be Asymptomatic, and Who Know Where HIV Testing is Provided by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.9 A Percentage of Women Aged 15–44 Who Do Not Know Principle Mechanisms of HIV Transmission by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.9 B Percentage of Men Aged 15–49 Who Do Not Know Principle Mechanisms of HIV Transmission by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.10 A Percentage of Women Aged 15– 44 Who Correctly Reject Misconceptions About HIV Transmission by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.10 B Percentage of Men Aged 15–49 Who Correctly Reject Misconception About HIV Transmission by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.11 A Percentage of Women Aged 15–44 Who Know HIV Infection Can Be Asymptomatic, And is Not Spread by Dental Treatment or Mosquito Bite, by Selected Characteristics. Reproductive Health Survey: Albania, 2002 Table 14.11 B Percentage of MenAged 15–49 Who Know HIV Infection Can Be Asymptomatic, And is Not Spread by Medical Treatment or Mosquito Bite, by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.12 A Percent Distribution of Women Aged 15–44 Who Mentioned Possible Means of Preventing HIV/AIDS Spontaneously And After Probing. Reproductive Health Survey: Albania, 2002

Table 14.12 B Percent Distribution of Men Aged 15–49 Who Mentioned Possible Means of Preventing HIV/AIDS Spontaneously and After Probing. Reproductive Health Survey: Albania, 2002

Table 14.13 A Percent of Women Aged 15–44 Who Know Possible Means of Preventing HIV/AIDS Transmission Spontaneously and After Probing, by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.13 B Percent of Men Aged 15–49 Who Know Possible Means of Preventing HIV/AIDS Transmission Spontaneously and After Probing, by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.14 A Percent of Women Aged 15–44 Who Believe HIV Can Be Prevented By Limiting Number of Sexual Partners, Being Monogomous, And Using Condoms, by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.14 B Percent of Men Aged 15–49 Who Believe HIV Can Be Prevented By Limiting Number of Sexual Partners, Being Monogamous, And Using Condoms, by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.15 A Percent Distribution of Women Aged 15–44 by Self-Perceived Risk of Contracting HIV/ AIDS by Selected Characteristics. Reproductive Health Survey: Albania, 2002

Table 14.15 B Percent Distribution of Men Aged 15–49 by Self-Perceived Risk of Contracting HIV/ AIDS by Selected Characteristics. Reproductive Health Survey: Albania, 2002 Table 14.16 A Opinion About The Main Risk Factor of Contracting HIV/AIDS Among Women 15–44 Who Have Heard About HIV/AIDS And Believe They Have Any Risk of Contracting HIV/ AIDS. Reproductive Health Survey: Albania, 2002

Table 14.16 B Opinion About The Main Risk Factor of Contracting HIV/AIDS Among Men 15–44 Who Have Heard About HIV/AIDS And Believe They Have Any Risk of Contracting HIV/ AIDS. Reproductive Health Survey: Albania, 2002

Table 14.17 A Opinion About The Main Factor That Protects From Contracting HIV/AIDS Among Women Aged 15–44 Who Have Heard of HIV/AIDS And Believe That They Have No Risk of Contracting HIV/AIDS, by Selected Characteristics. Reproductive Health Survey: Albania, 2002 (Percent Distribution)

Table 14.17 B Opinion About The Main Factor That Protects From Contracting HIV/AIDS Among Men Aged 15–49 Who Have Heard of HIV/AIDS And Believe That They Have No Risk of Contracting HIV/AIDS, by Selected Characteristics. Reproductive Health Survey: Albania, 2002 (Percent Distribution)

Table 15.1 A Percentage of Women Aged 15–44 Who Witnessed or Experienced Parental Abuse by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 15.1 B Percentage of Men Aged 15–49 Who Witnessed or Experienced Parental Abuse by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 15.2 A Percentage Who Reported Intimate Partner Violence (IPV) in Their Lifetime and Percentage Who Reported Intimate Partner Violence in The Last Year, by Type of Abuse and Selected Characteristics, Among Women Aged 15–44 Ever Married or in Union. Reproductive Health Survey: Albania 2002 Table 15.2 B Percentage of Men Who Reported Perpetration of Intimate Partner Violence (IPV) in Their Lifetime And Percentage Who Perpetrated Intimate Partner Violence in The Last Year by Type of Abuse and Selected Characteristics, Among Men Aged 15–49 Ever Married or in Union. Reproductive Health Survey: Albania 2002

Table 15.3 A Percentage Who Reported Lifetime Physical Violence and Recent Violence by Severity of Violent Act by Selected Characteristics, Among Women Aged 15–44 Ever Married or in Union. Reproductive Health Survey: Albania 2002

Table 15.3 B Percentage Who Reported Inflicting Lifetime Physical Violence And Recent Violence by Severity of Violent Act and Selected Characteristics, Among Men Aged 15–49 Ever Married or in Union. Reproductive Health Survey: Albania 2002

Table 15.4 Percent Distribution of Number of Incidents of Physical Abuse by Type of Abuse and Percentage Resulting in Physical Injuries Among Women Who Were Physically Abused During The Past Year Women Aged 15–44 Ever Married or in Union. Reproductive Health Survey: Albania 2002

Table 15.5 Women Who Were Physically Abused by an Intimate Partner During The Past Year Who Talked to Anyone About This Incident of Violence by Selected Characteristics. Reproductive Health Survey: Albania 2002 Table 15.6 Percentage of Women Who Were Physically Abused by an Intimate Partner During The Past Year Who Discussed the Abuse With Other Persons by Type of Person by Selected Characteristics. Reproductive Health Survey: Albania 2002

Table 15.7 Percentage Distribution of Women Who Were Ever Physically Abused by an Intimate Partner Who Did not Report the Abuse to a Health Provider, Police, or Lawyer by reasons for Not Reporting The Abuse by Residence and Socio-Economic Index. Reproductive Health Survey: Albania 2002

Table 15.8 Situations That Make Partner Violent as Reported by Women Who Were Physically Abused by an Intimate Partner During The Past Year by Residence. Reproductive Health Survey: Albania 2002

Table 15.9 Percentage of Women Aged 15–44 Ever Married or in Union Who Ever Experienced Any Form of Abuse Who Want to Know of The Tirana Hotline Number Where They Can Ask Questions About Violence Against Women. Reproductive Health Survey: Albania 2002

Table 15.10 Percentage of Women Age 15–44, Who Stated That a Man Has a Right to Hit or Beat His Wife Under The Following Situations by Residence and Education. Reproductive Health Survey: Albania 2002

List of Tables

List of Figures

Figure 4.1 Three-year period age-specific fertility rates for several time periods among all women

Figure 4.2 Median age at first sexual intercourse, first marriage, and first birth by education level

Figure 5.1 Adequacy of prenatal care utilization index, births in the 5 years prior to the survey, across Eastern European countries

Figure 5.2 Adequacy of prenatal care utilization index, births in the 5 years prior to the survey

Figure 5.3 Infant mortality rates by district in 1995 and in 2000

Figure 5.4 Distribution of neonatal and post-neonatal deaths in Albania, 1995 and 2000

Figure 6.1 Awareness and knowledge of how to use contraceptive methods among women aged 15-44 years

Figure 6.2 Awareness and knowledge of how to use contraceptive methods among men aged 15-49 years

Figure 7.1 Currently married women and contraceptive use status

Figure 7.2 Currently married women who are using modern, traditional, or no contraceptive method by educational level

Figure 7.3 Percentage of contraceptive users who stated that selected reasons read to them were very important or somewhat important when deciding to use a non-supplied method instead of a modern method Figure 8.1 Unmet need for modern contraception among fecund married women of reproductive age by future fertility preferences

Figure 8.2 Counseling and contraception offered at the time of last legally performed abortion

Figure 9.1 Percentage who want more information about contraception

Figure 9.2 Opinions on the best source of contraceptive information

Figure 11.1 Percentage of respondents visiting any health facility in the past 12 months by residence

Figure 11.2 Reproductive health behaviors among women who have ever had sex

Figure 12.1 Best age to start schoolbased courses on human reproduction, contraception, and STIs in the opinions of female respondents

Figure 12.2 Best age to start schoolbased courses on human reproduction, contraception, and STIs in the opinions of male respondents

Figure 12.3 Young adult women who discussed or were taught about various family life education topics before they reached age 18

Figure 12.4 Young adult men who discussed or were taught about various family life education topics before they reached age 18

Figure 13.1 Marital status at time of first sexual experience among young adults

Figure 13.2 Contraception used at first sexual intercourse

Figure 13 3.Percent of sexually experienced young adults who have ever talked to a partner about using condoms

Figure 14.1 Awareness of certain sexually transmitted infections among men and women 14-44

Figure 14.2 Recent exposure to mass media messages on HIV/AIDS and other STIs, all women and men of reproductive age

Figure 15.1 Lifetime and recent physical violence, CDC-assisted Reproductive Health Surveys, selected countries in Eastern Europe and NIS

Figure 15.2 Percentage of respondents who witnessed or experienced parental abuse

Figure 15.3 Questions asked using modified Conflict Tactics Scale (CTS)

Figure 15.4 Percentage of women ever married or in union who reported lifetime intimate partner violence (IPV), by marital status

Figure 15.5 Women's and men's reports of IPV in lifetime and past year

Chapter 1 BACKGROUND

lbania is a small country of about 28,748 km², situated in southeastern Europe. It borders on Greece to the south, the Former Yugoslav Republic of Macedonia to the east, the UN administered province of Kosova and Republic of Serbia and Montenegro to the north and the Adriatic and Ionian Seas to the west. There are 720 km of land borders and 362 km of coastline. The terrain is mountainous except along the central coast. About 42% of the 3.1 million people of Albania live in urban areas and approximately 20% of the population live in the capital of Tirana (INSTAT, 2002). Albanians are the majority ethnic group, representing over 95% of the population. Albania is a multi-religious country and three major religions, Muslim, Orthodox Christian, and Roman Catholic, have been important in contributing to the Albanian heritage and culture. All religious practice was outlawed and mosques and churches closed in 1967. However, private religious practice was again legalized in 1990, with a separation of religion and state functions.

There are no official data on the prevalence of religious identity among Albanians. A recent unofficial study indicated that about 72% of the country identifies themselves as Muslim, 18% as Orthodox Christian, and 10% as Roman Catholic (Neza, 2000), but religious affiliation is relative and linked mainly with inheritance from the past and not with current practice and beliefs.

Albania is administratively divided into 36 districts, 12 prefectures, 311 communes, and 64 municipalities (INSTAT, 2002). The population distribution between districts is quite different; there are districts with less than 10,000 inhabitants and there are districts with as many as 200,000 inhabitants. Seven cities with more than 50,000 inhabitants represent 62% of the

total urban population. This diversity is reflected in the average number of persons per square kilometer. There are districts with as low as 21 persons per km^2 while there are a number of districts with more than 400 persons per km^2 (INSTAT, 2001).

The modern history of the Albanian state starts at the beginning of the last century when it gained independence from the Ottoman Empire in November 1912. Between the two world wars Albania was first a parliamentary democracy, then a monarchy. After the end of World War II, the National Liberation Front (NLF) led by the Communist Party, controlled the country and a one party system headed by the Workers' Party was established. At that time Albania was part of the socialist bloc but gradually became isolated not only from the capitalist West but from almost all other Communist countries. This isolation was complete after 1978 when the country broke ties with China.

The collapse of Communism across Eastern Europe in 1990–1991 brought a number of social, economic, and political changes to the region. Albania began a transition toward a democratic government and new, market-oriented economy, which has presented formidable challenges. The current Constitution of Albania was ratified in 1998 and established the government as a parliamentary republic, with the capital in Tirana.

During the past 12 years Albania has faced continuous political and social changes, and after a period of transition, interrupted many times by social crises such as those of the years 1991–1992 and 1997 (the collapse of pyramid schemes) and the Kosovo crisis in 1999, Albania is now a country undergoing profound economic and structural reforms.

The economy of the country is changing from a central economic planning system to a free-market system; many questions related to privatization, property ownership claims, and the appropriate regulation of business still remain unresolved. The country has experienced slow but steady economic progress; however, according to poverty baseline statistics, 25.4% of the population is poor and 5% of Albanian citizens live in extreme poverty. The rates of poverty are higher in rural and remote areas of the country (UNDP, 2004). The official unemployment rate is 16%, with two-thirds of all workers employed in agriculture, mostly at the subsistence level. Remittances from citizens working abroad remain extremely important, as does foreign assistance.

Albania is a lower middle income country with a Gross National Income (GNI) per capita of US\$ 1,380. The agricultural sector accounts for 34% of Gross Domestic Product (GDP). Workers' remittances account for an additional 12% of GDP, with industry and services contributing 13% and 32%, respectively.

While economic growth in Albania has fluctuated during the last five years, it is now on a positive track for growth. The government is projecting that economic growth will increase to 6% from 2003-2005, with inflation rates of 2-4% (Ministry of Finance, 2002). The country has good potential for growth in agriculture, livestock, fisheries, forestry, tourism, mining, and light industry.

Nonetheless, the country faces considerable challenges as it remains one of the poorest countries in Europe. According to available data, as mentioned above, about one out of four Albanians is poor and lack access to basic services. A weak and deteriorating

infrastructure and related services have left up to 40% of households without access to necessities such as basic education, water, sanitation, and heating. The Government of Albania has developed a National Strategy for Social and Economic Development (NSSED) to lower the level of poverty and improve social and economic development. The main objectives of the NSSED for the three-year period, 2002-2004, are the following: (i) real GDP growth of 22-25 percent; (ii) reduction of number of people living in poverty, particularly for the worst-affected social groups and areas; (iii) tangible improvement of infrastructure and related services, e.g. supply of potable water and electricity, particularly for the impoverished populations; (iv) reduction of infant and maternal mortality rates and disease incidence; and (v) increase in the level of elementary and secondary education school enrollment (Ministry of Finance, 2002).

Reductions in infant and maternal mortality are one of the objectives of the Strategy and it follows that the health sector is responsible for this objective. Public spending on health is low compared to the average of 3.0% of GDP for lower middle income countries. As a consequence, one of the strategic priorities for public expenditures is to increase the share of GDP allocated to the health sector. The government planned to increase the share of GDP in 2005 to 4% compared with 1.85% in 2001. The percentage of total public expenditures has increased to 11% in 2004 from 7.4% in 1998.

Albania started negotiations in January 2003 with the European Union for a Stabilization and Association Agreement (SAA). Negotiations have so far focused on political issues.

The Albanian Government committed to

achieve Millennium Development Goals (MDG) following the July 2003 parliament resolution in support of MDGs, and recently prepared a MDG Progress Report as a result of the work of national consensus building groups comprising all partners and stakeholders. The Albania relevant MDG targets and indicators have been identified at the national level and regional and local levels, adopting different regional strategies consistent with local and national indicators (UNDP 2004).

An Albanian-European Partnership Action Plan document has also been prepared by the Government with clearly identified measures to improve food safety, the environment and control and surveillance of communicable diseases, including process and outcome indicators. Also. the Government, in collaboration with donor agencies and other partners, is coordinating the MDG process and strategy using the objectives of the NSSED as well as related long term development efforts towards European and regional integration through the Stabilization and Association Agreement process. Strong links have been formed between global, national and local development agendas emphasizing national ownership of the process, unification of stakeholders and common advocacy campaigns, and establishing strong mechanisms for monitoring and reporting.

During the years 1992–1993 there was a huge wave of emigration, mainly to neighboring European countries (Greece and Italy). During the period 1990– 1995, it is estimated that the number of emigrants fluctuated between 300,000 and 600,000, representing 9%–11% of the total population in 1995. Around 40% of them are estimated to be women and a new pattern of women emigrating alone without family members has been seen in recent years. Female trafficking and their exploitation as sex workers has appeared in the aftermath of the fall of old regime and in the vacuum of legislation (National Equity Committee, 2002; Lesko et al., 2003).

The social changes have also been associated with the internal migration of the population towards the big cities and particularly towards the capital city of Tirana. The internal migration during these years brought changes in the ratio of the urban/rural population. In 1989, the urban population was 36% of the total population and by the year 2001, the percentage had increased to 42% (INSTAT, 2002).

The demographic changes and the urbanization process are directly reflected in the decrease of the average household size. While in 1979 the average household size was 5.6 persons with 4.6 in the urban zones and 6.2 in the rural areas, the same indicator for 1989 was 4.7, 3.9, and 5.3 persons, respectively, and in the 2001 Census was 4.2, 3.9, and 4.5 persons, respectively (INSTAT, 2002). It appears that the tendency toward a family with two parents and two children is the new norm in Albania, as in many other countries. Internal migration has affected the lives of both women and children due to difficult access of services, unemployment and dependence on the male as the workers in the family.

Life expectancy for females in 2001 was estimated to be 77.5 compared with 72.5 for males. Women typically marry and begin families at a relatively young age; the average age at marriage for women is 23 years (INSTAT, 2002).

The provisions of the Albanian Constitution of 1946 sanctioned for the first time basic rights for women. However, Albanian women tend to follow a traditional model where women defer to the men in the family and the society has been very conservative

in preserving traditional family structures and inequalities in the family. The Albanian Parliament ratified in December 1993 the Convention on the Elimination of all Forms of Discrimination Against Women (AFPA et al, 2002). However, during the transition period, due to lack of work and high unemployment, women became more vulnerable and male or family support was necessary to survive. Many social services helping women in the past were eliminated due to lack of funding. In the northern parts of Albania the influence of the Kanun, a code of traditional law dating back several centuries, is still practiced although it was prohibited by the state after the second world war. In the aftermath of the fall of the hard-line Communist regime, the Kanun is reported to have regained strength in the north of the country (National Equity Committee, 2002). The Kanun declares a woman to be property transferable from her father to her husband.

When Albania was a Communist country, the healthcare system was centrally controlled and based on the Semashko model as in other countries of Eastern Europe and the Former Soviet Union. Health care was free to all with emphasis on infectious disease prevention and some health education programs, but an important percentage of the budget went to medical treatment. A variety of changes in the health legislation have been proposed very recently with the drafting or revision of important laws related to the financing and organization of the health care systems, and the creation of a health insurance fund and the patient rights card. Also, Albania has recently developed a new national ten-year strategy for healthcare reform. The implementation of a National Health Promotion Strategy has been discussed, consistent with the objectives of the health sector reform (under The National Strategy for Social-Economic Development) and includes the following

goals: 1) increased effectiveness and efficiency in use of resources; 2) increased access to quality health services nationwide; and 3) improvement of health indicators through specific targeted interventions. The increased effectiveness and efficient use of resources will be achieved through: (i) improvement of the planning process and needsassessmentwithimprovedmanagement and fairer distribution of resources; (ii) decentralization of management functions to local institutions, including regional health authorities, and strengthening of the role of professional organizations; (iii) reduction in corruption; (iv) the gradual establishment of information systems; and (v) support for the privatization process in providing health services and monitoring of the private sector (Ministry of Health, 2004)

The Primary Health Care Policy adopted in 1997 aims to offer accessible and financially affordable healthcare to all. Officially. health care still remains free, but while physicians are still employed by the state, many people pay for healthcare services in the form of gifts or unofficial fees for service. This unofficial expense comes out of limited household budgets. Very recently, user fees were introduced in hospitals and some primary health care facilities which are meant to limit and prevent "unofficial expenses." Even though many health care facilities were rehabilitated since 1997 there are still facilities that urgently need basic repairs. Also, due to the closed nature of the country until 1990, physicians may not have the most up-to-date information and skills. In addition to Ministry of Health operated clinics, there are now private providers of healthcare and clinics operated by international organizations.

Infant and under five mortality are officially reported as 23 deaths per 1,000 live births and 32 deaths per 1,000 live births, respectively, as of 2000, the

highest officially reported rate in Europe (INSTAT, 2002). Although these rates have both declined considerably from rates in the late 1950s and early 1960s, and the official statistics appear to show improvement in infant mortality since the transition, the rates may in fact be affected by problems such as nonregistration of births in which the infant dies shortly thereafter and the definition of early neonatal deaths and early deaths of premature infants as stillbirths. Mortality rates are estimated to be more than twice as high among children in the more rural northern areas of the country than in the more urban central and coastal areas (World Bank, 1997).

For 50 years Albania had a pronatalist population policy; modern family planning was forbidden and it was taboo to speak about sexuality and contraception in public. Reproductive health care and basic family planning services were introduced into the country in 1992 after a Decision of the Council of the Ministers that declared family planning should be seen as a basic human right from which all citizens should be able to benefit on their own free will. Eleven services are now provided with three levels of care, starting with primary healthcare services (health centers, ambulances in villages, and consulting centers for women and children in cities), maternity and pediatric hospitals at the secondary level, and the Hospital of Obstetrics and Gynecology in Tirana and the University Hospital Center "Mother Teresa" in Tirana at the tertiary level of care. Also, national and international NGOs provide family planning services in addition to advocacy on this issue. However, women often have limited access to information and services regarding reproductive health, especially in rural areas.

Abortion, prior to 1991, was also forbidden and about half of the maternal deaths during the decade of the 1980's were due to complications following illegal abortion. In April 1991, through the "order of the Minister of Health" and the "Decision of the Council of Ministers for the approval of activities of Family Planning in Albania," abortion was legalized and modern methods of contraception were introduced in the public health services. The "Law for the Voluntary Interruption of Pregnancy until 12 Weeks Gestation" passed the Parliament in May 1995. Following its legalization in 1991, abortion declined as a cause of maternal mortality from 50% in 1989 to 25% in 1993 and to 6% in 1997 (unpublished paper, Population in Europe and North America on the Eve of the Millenium: Dynamic and Policy Responses, presented at the UNFPA Regional Population Meeting, December 1998, Budapest, Hungary). During the past 12 years, the figures for maternal and infant mortality remain relatively high despite the fact that they had declined by about 50% compared with the years before 1990.

A national law on reproductive health passed the parliament in June 2001 which regulates the management, administration, supervision functioning and of all reproductive health services and activities in public and private health institutions. The law protects the reproductive rights of individuals and couples in accordance with national policies and laws as well as known and accepted international principles. According to the law, the overall goals of reproductive healthcare services are to offer good access and quality reproductive healthcare; to improve the health status of women during their reproductive years, especially during childbearing and delivery; to improve the health status of newborns, infants, and children; and to improve the health of adolescents and young adults.

In addition to the striking similarities in socioeconomic conditions inherited from the Communist era, there have also been demographic and health similarities among countries in the region, in particular a heavy reliance on abortion rather than on modern contraception as a means of preventing unintended births. Therefore, reproductive health is an issue of critical importance for the countries of this region (CDC and ORC MACRO, 2004). Also, reproductive health has been considered as one of the priorities of the national health promotion and public health strategy (MOH, 2003)

Beginning in 1993, several surveys on family planning and reproductive health attitudes and behaviors were conducted in Eastern Europe (CDC and MACRO, 2003). To this end, the Division of Reproductive Health of the U.S. Centers for Disease Control and Prevention (CDC/DRH) in Atlanta has provided technical assistance for Reproductive Health Surveys (RHS) in collaboration with local counterparts. Between 1993 and 2001, ten Reproductive Health Surveys were conducted in seven countries in Eastern Europe.

A Reproductive Health Survey (RHS) was conducted in Albania in 2002 making it the 8th country in the region to conduct this type of survey. This survey represents the first systematic effort to gather representative national data on population and reproductive health issues in Albania. Population-based surveys of women of reproductive age using nationally representative samples are an effective mechanism for collecting information on topics such as family planning, fertility, contraceptive use, knowledge about HIV/ AIDS, and other reproductive health Until recently, relatively little issues. detailed and reliable population-based information was available about the situation in the country with regard to important reproductive health topics.

The RHS, supported by USAID, UNFPA and UNICEF, examines patterns and levels of fertility, family planning, contraceptive use and method selection, health behaviors, knowledge of HIV/AIDS, as well as attitudes towards specific contraceptive methods and abortion. These issues are of particular importance in Albania, since for many years women and healthcare providers had limited access to up-todate and reliable information on these topics. The survey also provides data on key maternal and child health indicators, infant feeding, and the extent to which mothers receive medical care during pregnancy and at delivery.

A principal objective of the survey is to examine the reproductive health status and needs that can be used to help direct or modify program interventions. These data are particularly useful in assisting policy makers and health planners in evaluating health service needs, and identifying reproductive health behaviors associated with poor health outcomes. They could also play a significant role in designing programs better targeted to meet the needs of population subgroups. A key programmatic difference between policy objectives in Albania and other countries in Eastern Europe, compared with those in some developing countries, is that the emphasis is not on promoting a decline in fertility and population growth, but on bringing about improvements in women's health through increased availability and improved use of modern contraceptive methods and reduced reliance on abortion. Until now, a comprehensive comparison of key family planning and reproductive

health indicators had never been compiled in Albania.

The nationally representative data on key indicators presented in this report can be used to design or modify health interventions, identify high-risk behaviors amenable to change and highlight reproductive health areas that warrant greater attention. These data can be translated into policy and programmatic activities to improve services and findings may be combined with other existing information to contribute to a more profound understanding of reproductive health in Albania.

CHAPTER 2 METHODOLOGY

Sampling Design

The Albania RHS 2002 is based on inperson, face-to-face interviews with 5,697 women and 1,740 men in their homes. The household-based survey was designed to collect information from a representative sample of men and women of reproductive age throughout Albania. Respondents were selected from the universe of all females aged 15–44 years and all males aged 15– 49 years, regardless of marital status, who were living in Albania when the survey was conducted. Male and female samples were selected independently.

For analysis purposes, three strata were constructed for the sampling design: Metro Tirana, other urban areas and other rural areas. Metro Tirana includes 6 of the 19 communes in Tirana district: Bashkia Tirane (capital city of Tirana), Kamez, Vore, Farke, Kashar and Paskuqan. These six communes include 85% of the District population and an estimated 92% of the urban population in the District. The "Other Urban Area" stratum includes urban areas outside of Metro Tirana and the "Other Rural Area" stratum includes all rural areas outside of Metro Tirana.

As in other countries in Eastern Europe with Reproductive Health Surveys, the survey had a three-stage sampling design, which allows independent estimates for the female and male samples. The first stage of the sample design was a selection of census sectors with probability proportional to the number of households recorded in the 2001 Census. During this stage, 300 census sectors, 100 in each of the strata defined above, were selected as primary sampling units (PSUs) throughout Albania. This step was accomplished by using a systematic sample with a random start in each strata for the female sample. A 33% sub-sample (every third PSU) of the census sectors selected in the female sample constituted the first stage of the male sample. Thus, the first-stage selection included 300 sectors for the female sample and 100 sectors for the male sample.

In the second stage of sampling, clusters of households were randomly selected in each PSU that was chosen in the first stage (separate households were selected for the female and male samples). Finally, in the third stage of sampling, in each of the households in the female sample, one woman aged 15–44 years was selected at random for interview and in the male sample one man aged 15–49 years was randomly selected for interview.

Metro Tirana and Other Urban Areas were over-sampled, and rural areas were undersampled, so that more precise estimates could be made for the two mostly urban strata. Two variables are used in this report: STRATA, including metro Tirana, other urban areas and other rural areas as defined above and RESIDENCE representing urban or rural residence independent of strata. Urban residence includes the 100 PSUs in the "other urban areas" stratum and 86 of the 100 PSUs in Metro Tirana. Rural residence includes the 100 PSUs in the "other rural areas" stratum and 14 of the 100 sectors in Metro Tirana.

Some PSUs intended for both the male and female samples were not large enough to provide non–overlapping clusters. In these cases, an adjacent enumeration area in the same location was identified for the male sample, and in a few instances the male sample was drawn from a combination of both areas due to small population size.

Because only one respondent was selected from each household with women (or men) of reproductive age, all results have been weighted to compensate for the fact that some households included more than one eligible respondent. Survey results were also weighted to adjust for over-sampling of the metro Tirana stratum and other urban areas and under-sampling of rural areas. A review of the sample data compared with results from the 2001 census showed that there was differential non-response in certain age groups for both females and males and also differential non-response by marital status among females. Thus, a third weight was added to adjust for differential non-response. Response rates were lowest for unmarried women 30-44 years of age but they represent only 4% of all women of reproductive age (WRA). For women 15-19, married women were underrepresented and unmarried women overrepresented in the sample; however, married teenagers represent only 2% of all WRA. Thus, the differential non-response weight for females is not a significant adjustment. For males, teenagers 15-19 were over-represented in the sample and older working age men from 20-49 years of age slightly underrepresented. The third weight for males adjusted for this differential non-response.

Presentation of Tables

All tables in this report represent weighted results. However, the un-weighted number of cases, used for variance estimates, is shown in each table (see Appendix A on sampling errors). Thus, the survey can be used to make national and sub-national estimates because of the process to "weight" the data —that is, to determine how many women in the population were represented by each woman in the sample.

Another note concerning data presented in tables in this report relates to percent distributions: although all percent distributions are shown to add to 100.0%, they may actually add to 99.9% or 100.1% due to rounding. Also tables for females (A) and males (B) that relate to the same topic are positioned to face each other for easier comparisons by the reader. Tables labeled A (female) will always be on an evennumbered page and corresponding B tables will be on the following odd-numbered page. To maintain these comparisons, there sometimes will be blank numbered pages in the table sections of the report.

Questionnaire

The individual questionnaire included information on each respondent's education, employment, living arrangements, and other background characteristics, as well as histories of marriage, divorce and cohabitation, sexual experience, pregnancy and contraceptive use. Additional questions investigated health risk behaviors that may affect reproductive health (smoking and drinking habits), women's health screening practices, young adult sexual and contraceptive behavior, knowledge and attitudes related to HIV/AIDS, and intimate partner violence. The questionnaire was developed in English and translated into Albanian and underwent two pretests. The second pretest, in May 2002, was performed to test changes in the questionnaire made after the first pretest.

Data Collection

The interviews were performed by 25 female and 8 male interviewers specially trained in interview techniques, survey procedures, and questionnaire content before the beginning of fieldwork. Interviewer training took place in the Health Authority Training Center, a facility next to the headquarters of the Albanian National Institute of Public Health (IPH), just before data collection began and consisted of one week of classroom

training in fieldwork procedures and proper administration of the questionnaire and one week of practical training in the field with close monitoring by the trainers. At the end of the training period, five female and two male teams were selected for the fieldwork. Each team consisted of one Supervisor, four Interviewers, and a Driver. Fieldwork was managed by staff of IPH with technical assistance from CDC/DRH. The overall fieldwork implementation was supervised by two fieldwork coordinators. Fieldwork lasted from August through December 2002. Each team was assigned to visit a number of primary sampling units in all regions of the country and traveled by car throughout the country on planned itineraries. Interviews were conducted at the homes of the respondents and lasted on average about 35 minutes for both men and women. Interviews were conducted in Albanian. Completed questionnaires were first reviewed in the field by team supervisors and then were taken by the fieldwork coordinators, who also reviewed them, to the National Institute for Statistics (INSTAT) headquarters where they were reviewed again by a data quality consultant before data processing.

Response Rates

Of the 10,316 households selected in the female household sample, 5,866 (57%) included at least one eligible woman (age 15-44 years) (Table 2.1A). One-third (34%) of households did not include an eligible woman and 8% of households were unoccupied, principally in rural areas (12%). Of the identified respondents, 5,697 were successfully interviewed, yielding an individual response rate of 97% for an overall response rate of 97% for women. Virtually all respondents who were selected to participate in the sample agreed to be interviewed and were very cooperative. Less than one percent refused to be interviewed. Response rates were similar in all three strata.

The male sample totaled 3,965 households with 1,831 (46%) including at least one eligible man (age 15–49 years) (Table 2.1B). A lower percentage of households included an eligible male than did the female sample (include an eligible female) due, in part, to the emigration of males of working age. A total of 1,740 eligible males were interviewed for a 95% individual response rate, yielding an overall response rate of 94% (.95 x 99%). As with the female sample, refusals were less than one percent and response rates in the three strata were similar.

Table 2.1 A
Results of Household Visits and Interview Status of Eligible Women by Stratum
(Percent Distribution)
Reproductive Health Survey: Albania 2002

			Strata	
Households Visits	Total	Metro Tirana	Other Urban	Rural
Identified eligible women*	56.9	59.7	52.5	58.6
No eligible woman lives in household	34.0	35.5	37.0	28.8
Residents not at home	0.8	0.3	1.4	0.6
Household refusal	0.0	0.0	0.0	0.0
Unoccupied house	8.4	4.5	9.1	12.0
Total	100.0	100.0	100.0	100.0
No. of Households Visited	10,316	3,594	3,593	3,129
Eligible Women				
Completed interview	97.1	98.2	96.3	96.7
Selected respondent not home	1.5	0.7	2.3	1.7
Selected respondent refusal	0.1	0.1	0.2	0.0
Other reasons**	0.2	0.4	0.2	0.0
Total	100.0	100.0	100.0	100.0
No. of Eligible Women Identified	5,866	2,146	1,886	1,834
No. of Completed Interviews	5,697	2,108	1,816	1,773

* Includes women 15–44 years of age who had complete or incomplete interviews, who were absent or handicapped, or who refused to be interviewed.

**Includes women with a handicap preventing them to be interviewed and women having incomplete interviews.

			Strata	
Households Visits	Total	Metro Tirana	Other Urban	Other Rural
Identified eligible men*	46.2	54.3	42.2	41.2
No eligible man lives in household	42.0	34.7	45.5	46.5
Residents not at home	1.3	1.6	1.1	1.5
Household refusal	0.1	0.1	0.1	0.0
Unoccupied house	10.4	9.3	11.1	10.8
Total	100.0	100.0	100.0	100.0
No. of Households Visited	3,965	1,402	1,380	1,183
Eligible Men				
Completed interview	95.0	94.3	93.8	97.5
Selected respondent not home	4.4	4.3	6.2	2.5
Selected respondent refusal	0.2	0.5	0.0	0.0
Other reasons**	0.3	0.8	0.0	0.0
Total	100.0	100.0	100.0	100.0
No. of Eligible Men Identified	1,831	761	583	487
No. of Completed Interviews	1,740	718	547	475

Table 2.1 B Results of Household Visits and Interview Status of Eligible Men by Stratum (Percent Distribution) Reproductive Health Survey: Albania 2002

* Includes men 15–49 years of age who had complete or incomplete interview, who were absent or handicapped, or who refused to be interviewed.

**Includes men with a handicap preventing them to be interviewed and men having incompleted interviews.

CHAPTER 3 CHARACTERISTICS OF THE SAMPLE

Household characteristics

In the 2002 Albania Reproductive Health Survey, the household questionnaire included a roster for the interviewer to list all members of the household who met the criteria to be eligible for the sample. From this list, one eligible respondent was randomly selected for the female sample or one eligible man was randomly selected for the male sample. The household questionnaire also listed the total number of persons living in the household. In addition, the individual questionnaire collected data on amenities and durable consumer goods belonging to the household, which would later be used to construct a socioeconomic scale to allow assignment of a socioeconomic status indicator to women and men in the sample. In this way, we would be able to examine the association of socioeconomic status with reproductive health indicators or behaviors measured in the survey. Tables 3.1 (A & B) and 3.2 (A & B) present data from the household questionnaire.

In Tables 3.1 (A & B), the average size of Albanian households can be seen to be 5.1 persons in the female sample and 4.6 persons in the male sample. The most frequent household sizes are four- and five-person households. More than half of all the households, for the total country and for specific geographic strata in both female and male samples, have an average size of four or five persons. Households of one or two persons are relatively rare, with less than 1% of households being singleperson households and less than 4% being two-person households in both the female and male samples. However, the percentage of one- or two-person households may be understated due to the greater likelihood of the interviewer finding no one at home for the household interview visit compared to households having three or more members.

As would be expected, rural areas have a slightly larger household size than urban areas, especially in the female sample.

Household amenities and consumer goods are shown in Tables 3.2A and 3.2B. Roughly one-quarter of Albanian households in 2002 had a telephone line, and 63% had flush toilets. Less than 10% had access to 24-hour electricity. Dramatic differences in these amenities can be seen between the urban and rural strata. This is particularly so for telephone lines and flush toilets. Almost half of the households in urban areas had telephone lines, whereas less than 10% of rural households had these lines. However, in both samples, approximately 60% of respondents (females-62%, males-58%) reported having cell phones; about twothirds of households in urban areas and approximately half of households in rural areas. While more than three quarters of urban households had flush toilets. less than half of rural households reported having them. There is strong consistency between the female and male samples.

Among the durable goods possessed by households, the most frequent are TVs and refrigerators, with more than 90% of all households possessing these items. Ownership of these items appears to be geographically ubiquitous, with only minor differences between urban and rural areas. Seventy-eight percent of Albanian households have a gas or electric stove, roughly 60% have cell phones, and 21% have a working automobile. These three possessions vary markedly by strata, with their presence more common among the urban compared to the rural population. Computers and air-conditioning are rare. Less than 5% of households outside of Metro Tirana have these modern goods, and in Tirana only 12% of the households reported having them. Vegetable gardens,

on the other hand, are quite common, with 90% of rural households and 20 to 40% of urban households reporting access to a vegetable garden. Again, as with household amenities, there is strong consistency in percentages of durable goods between the female and male samples.

Characteristics of the Respondents

Tables 3.3A and 3.3B present selected sociodemographic characteristics of the samples.

Regarding the age distribution, 38% of female respondents and 33% of male respondents were young adults (15–24 years of age). The age distribution is slightly younger in rural areas for both females and males. Overall, the age distributions are similar to those found in official statistics (Instituti I Statistikes, 2002).

Sixty-five percent of women and 60% of men reported that they were currently married. Divorce and widowhood are very infrequent in Albania. Only 2% of women and less than 1% of men reported themselves in either of these categories. There is no significant urban-rural difference in marital status. More than a third of both women and men reported being childless, and among those with children, the modal number of children is two. Although the percentage of childless respondents did not significantly differ between urban and rural areas, more women and men reported three or more children in rural areas than in Tirana or other urban areas. Among women in rural areas, 28% had 3 or more children compared to 15% in Tirana and 16% in other urban areas. The corresponding percentages for men were 25% in rural areas, 13% in Tirana, and 11% in other urban areas.

Eight percent of women and 9% of men have had post-secondary education. Respondents

in urban areas were significantly more likely to have post-secondary education than those in rural areas (17% vs. 2%). Most women (54%) and men (48%) have had only primary or no schooling. In rural areas, three-quarters of the female population and two-thirds of the male population fall into this latter category.

The majority of respondents reported that they were Muslim; 80% of women and 84% of men. This is somewhat higher than the 72% reported in the census, but this difference may be affected by differential emigration patterns. Another 8% of women and 9% of men said they were Orthodox, and Catholics made up 12% of the women and 4% of the men. Ethnically, the population is almost 100% Albanian.

As for religiosity — measured by frequency of attendance at religious services, less than 50% of respondents reported that they attend services at least once a month. The one exception to this norm is Catholic men living in rural areas. Among the latter group, two-thirds (67%) report they attend church once a month or more frequently. Otherwise, only 5% of all Muslim women and men report attending religious services at least once a month, and the comparable percentages for the other religions are 34% and 25% for Orthodox women and men, respectively, and 44% for Catholic women and 43% for Catholic men.

Only 49% of men and 15% of women reported that they were working outside the home for 20 or more hours per week. These percentages rise to 60% for men and 25% for women in urban areas, and drop to 39% for men and 8% for women in rural areas. As for geographic mobility, the women report more mobility than the men. More than half of the women (53%) reported to have ever migrated, with only 22% of the men reporting the same. As would be expected, the Metro Tirana population has a much higher percentage of migrants than other urban or other rural places. Malefemale differences in migration history are greatest in the rural population. Only 3% of rural men have ever migrated, whereas 51% of rural women report having migrated.

Tables 3.4A and 3.4B show the marital status distribution controlling for various sociodemographic characteristics. There is little to no variation in marital status by residence for both women and men. The positive association between age and marriage occurs before the age of 35 for both sexes, although women appear to marry at younger ages than men. By age 20-24, half of the women are married, whereas for men this proportion is not married until age 25-29. None of the men in the sample are married at ages 15–19, while 10% of women are married in these young ages. On the other hand, at ages 40-44, 4% of women and less than 1% of men have remained never married. The median age at first marriage is 21.9 years for the women and 26.5 years for the men (see Chapter 4). An association for number of living children with marital status is observed only between no children and any children. Once there are any living children present, currently married status reaches 95% or higher. The relation of educational level with marital status appears to be linked in part with school attendance. Women and men with secondary incomplete and postsecondary educational levels are more likely to be still in school (data not shown) and, consequently, less likely to be married. For both sexes, those with completed secondary education are more likely to be married than those with primary or less. At the same time socioeconomic status has no effect for women and a very modest effect for men, with men at the lowest status level somewhat more likely to be married than those at the other two levels. Employed women and men have higher percentages married than those not working (74% vs. 64% for women and 76% vs. 45% for men), also likely reflecting current school attendance and young age among those not working.

Table 3.1 ASize of Households with at Least One Eligible Respondent by Stratum
(Percent Distribution)
Female Sample,
Reproductive Health Survey: Albania 2002

	Strata							
No. Persons per Household	Total	Tirana	Other Urban	Rural				
1	0.2	0.9	0.2	0.0				
2	2.5	3.5	3.7	1.5				
3	10.7	14.4	11.4	9.3				
4	28.6	34.8	36.8	22.3				
5	24.0	22.1	25.9	23.4				
6	16.5	12.5	12.8	19.8				
7	8.7	4.8	4.8	12.0				
8 +	8.8	7.0	4.3	11.7				
Total	100.0	100.0	100.0	100.0				
Average No. of Persons	5.1	4.7	4.6	5.4				
No. of Cases	5,788*	2,125	1,859	1,804				

* Exludes 15 households whose number of inhabitants was unknown.

Table 3.1 BSize of Households with at Least One Eligible Respondent by Stratum
(Percent Distribution)
Male Sample,
Reproductive Health Survey: Albania 2002

		Strata					
No. Persons Per Household	Total	Metro Tirana	Other Urban	Other Rural			
1	0.5	0.9	0.7	0.2			
2	3.9	5.3	4.6	2.9			
3	14.4	15.6	19.7	10.8			
4	30.8	32.1	40.3	24.6			
5	27.4	21.9	24.1	31.4			
6	12.6	13.2	8.2	15.0			
7	6.7	6.0	2.0	9.7			
8 +	3.8	4.9	0.3	5.4			
Total	100.0	100.0	100.0	100.0			
Average No. of Persons	4.6	4.6	4.1	4.9			
No. of Cases	1825*	755	583	487			

* Excludes 6 households whose number of inhabitants was unknown

Tables

Table 3.2 A Percentage of Households with Basic Household Amenities and Goods, by Stratum, for Women Aged 15–44 Years Reproductive Health Survey: Albania 2002

	_		Strata	
Household Amenities	Total	Metro Tirana	Other Urban	Other Rural
Flush Toilet	62.9	77.4	80.9	48.8
Cell Phone	61.6	69.5	66.5	56.7
Telephone Line	24.7	47.6	48.6	4.9
Electricity (24 Hours)	9.5	31.5	5.3	5.3
Vacation Home	1.5	5.6	1.1	0.5
Household Goods				
TV	96.4	97.7	98.5	94.9
Refrigerator	90.0	97.3	96.3	84.5
Gas/Electric Stove	78.1	84.0	93.7	67.9
Family Has Access to Vegetable Garden	60.7	33.9	22.9	89.0
VCR	31.8	44.6	42.4	22.3
Satellite Antenna	30.2	22.2	33.3	30.8
Auto	20.9	29.9	25.1	15.9
Computer	3.7	12.1	5.1	0.6
Air Conditioner	2.9	11.2	2.6	0.6
Percentage of Households With				
Crowded Conditions*	92.6	92.1	94.1	92.0
No. of Cases	5,697	2,108	1,816	1,773

* Total number of persons living in the household divided by total number rooms in the house (not including kitchen and bathroom) was higher than one

Table 3.2 BPercentage of Households with Basic Household Amenities and Goods,
by Stratum, for Men Aged 15–49 Years
Reproductive Health Survey: Albania 2002

			Strata	
Households Amenities	Total	Metro Tirana	Other Urban	Other Rural
Flush Toilet	62.9	83.2	76.5	46.9
Cell Phone	58.3	73.6	68.9	46.0
Telephone Line	24.0	46.8	39.9	5.6
Electricity (24 Hours)	5.3	19.2	3.3	1.1
Vacation Home	1.3	2.8	1.4	0.6
Household Goods				
TV	98.4	99.2	99.7	97.4
Refrigerator	92.7	98.4	98.7	86.9
Gas/Electric Stove	77.6	93.5	94.5	61.3
Family Has Access to Vegetable Garden	60.2	39.4	19.4	92.6
VCR	32.5	51.1	40.0	20.8
Auto	20.8	35.8	29.8	9.6
Satellite Antenna	17.3	18.7	16.1	17.5
Computer	3.9	12.7	4.1	0.4
Air Conditioner	3.2	12.2	2.2	0.3
Percentages of Households With				
Crowded Conditions*	88.5	86.9	87.6	89.6
No. of Cases	1,740	718	547	475

* Total number of persons living in the household divided by total number rooms in the house (not including kitchen and bathroom) was higher than one

	9
C	
5	

Table 3.3 A
Percent Distribution of Characteristics
of Female Sample by Residence and Stratum
Reproductive Health Survey: Albania 2002

Residence Strata								
Characteristics	Total	Urban	Rural	Metro Tirana	Other Urban	Other Rural		
Age Group								
15–19	21.0	18.5	22.9	18.3	18.8	23.0		
20–24	17.0	15.5	18.1	18.2	14.7	17.9		
25–29	15.6	15.3	15.8	16.7	14.8	15.7		
30–34	15.7	17.4	14.5	16.1	18.1	14.4		
35–39	15.6	17.9	13.9	15.7	18.5	14.0		
40–44	15.0	15.4	14.7	15.0	15.1	15.0		
Marital Status								
Currently Married	65.1	66.3	64.2	63.1	68.0	64.1		
Previously Married	2.1	2.4	1.8	2.9	2.1	1.9		
Never Married	32.8	31.3	34.0	34.0	30.0	34.0		
Living Children								
0	37.8	36.7	38.6	39.6	35.4	38.6		
1	12.8	13.8	12.0	16.3	12.6	11.9		
2	27.1	34.1	21.9	29.3	35.7	21.8		
3	14.9	11.5	17.4	10.6	12.2	17.5		
4 +	7.4	3.8	10.1	4.3	4.0	10.2		
Education Level								
Primary or Less	53.9	27.5	73.7	33.8	27.7	74.0		
Secondary Incomplete	10.2	14.3	7.2	12.0	15.1	7.1		
Secondary Complete	27.7	41.3	17.6	35.9	42.3	17.4		
Post-Secondary	8.1	16.8	1.6	18.2	14.9	1.5		
Socioeconomic Index	10.0			aa <i>i</i>				
Low	42.2	20.0	58.8	22.1	21.4	59.4		
Medium	49.5	63.0	39.5	53.9	66.6	39.0		
High	8.2	17.0	1.7	24.0	12.0	1.6		
Religion*								
Muslim	79.6	76.6	81.8	84.5	74.1	81.1		
Orthodox	8.1	12.7	4.7	7.9	14.1	4.9		
Catholic	11.5	9.2	13.2	5.7	10.7	13.6		
Other/Undeclared	0.8	1.5	0.3	2.0	1.1	0.3		
Employment	4 - 0			a- a				
Working	15.3	25.7	7.6	27.8	23.5	7.3		
Not Working	84.7	74.3	92.4	72.2	76.5	92.7		
Migration Status								
Ever Migrated	52.8	53.1	52.6	67.5	48.3	51.0		
Never Migrated	47.0	46.7	47.3	31.7	51.7	48.9		
Do Not Know	0.1	0.3	0.0	0.7	0.0	0.0		
Total	100.0	100.0	100.0	100.0	100.0	100.0		
No. of Cases	5,697	3,572	2,125	2,108	1,816	1,773		
*With regard to religious service at								
Muslim	<u>Total</u>	<u>Urban</u>	Rural	Metro Tirana	Other Urban	Other Rural		
Muslim	5.4	6.5	4.6	6.3	6.7	4.5		
Orthodox	33.7	34.9	31.2	37.3	34.2	31.3		
Catholic	43.8	36.1	47.8	44.6	34.2	47.9		

Table 3.3 BPercent Distribution of Characteristicsof Male Sample by Residence and StratumReproductive Health Survey: Albania 2002

		Residence Strata						
Characteristics	Total	Urban	Rural	Metro Tirana	Other Urban	Other Rural		
Age Group								
15–19	18.9	16.6	20.9	16.3	16.9	21.0		
20–24	14.5	13.2	15.6	14.9	13.3	15.0		
25–29	13.3	12.9	13.7	14.2	12.0	13.8		
30–34	13.6	14.6	12.8	12.4	15.8	12.8		
35–39	14.1	14.6	13.7	11.7	16.3	13.8		
40–44	14.2	15.6	12.9	16.2	14.8	13.0		
45–49	11.4	12.4	10.6	14.3	11.0	10.6		
Marital Status								
Married	60.3	59.6	61.0	59.8	59.8	60.9		
Previously Married	0.6	0.9	0.4	1.2	0.5	0.4		
Never Married	39.0	39.5	38.6	39.0	39.7	38.7		
Living Children								
0	45.5	45.5	45.5	47.4	45.0	45.1		
1	11.5	13.4	9.8	13.5	12.8	9.9		
2	24.9	30.0	20.5	26.6	31.5	20.3		
3	11.8	8.5	14.7	7.7	8.9	15.1		
4 +	6.3	2.6	9.5	4.8	1.8	9.5		
Education Level								
Primary or Less	48.4	28.4	65.7	30.0	30.5	66.1		
Secondary Incomplete	8.7	10.6	7.1	11.8	9.7	6.9		
Secondary Complete	33.9	43.6	25.5	39.6	44.4	25.4		
Post-Secondary	9.1	17.4	1.8	18.7	15.4	1.6		
Socioeconomic Index								
Low	47.2	25.1	66.4	20.9	29.2	68.2		
Medium	42.2	53.6	32.3	52.3	54.8	30.9		
High	10.6	21.3	1.2	26.8	16.0	1.0		
Religion*								
Muslim	84.0	75.9	91.1	84.0	71.6	91.4		
Orthodox	8.8	15.0	3.4	7.6	18.4	3.6		
Catholic	4.3	5.6	3.1	3.4	7.3	2.8		
Other/Undeclared	2.9	3.5	2.4	5.0	2.7	2.2		
Employment								
Working	48.9	60.3	39.1	63.7	57.1	38.3		
Not Working	51.1	39.7	60.9	36.3	42.9	61.7		
Migration Status								
Ever Migrated	21.7	38.3	7.2	58.8	29.4	2.6		
Never Migrated	78.1	61.2	92.7	39.8	70.6	97.4		
Do Not Know	0.3	0.5	0.0	1.3	0.0	0.0		
Total	100.0	100.0	100.0	100.0	100.0	100.0		
No. of Cases	1,740	1,155	585	718	547	475		
*With regard to religious service a					-			
	Total	Urban	Rural	Metro Tirana	Other Urban	Other Rural		
Muslim	5.4	3.1	7.2	4.9	2.8	6.9		
Orthodox	25.0	23.6	30.5	23.0	23.7	30.5		
Catholic	42.7	27.0	67.0	36.1	23.1	76.0		
			0110					

Chapter 3

Table 3.4 A Percent Distribution of Marital Status by Selected Characteristics, for Women Aged 15–44 Years Reproductive Health Survey: Albania 2002

		Marital Status	;		
Characteristics	Married	Previously Married	Never Married	Total	No. of Cases
Total	65.1	2.1	32.8	100.0	5,697
Strata					
Metro Tirana	63.1	2.9	34.0	100.0	2,108
Other Urban	68.0	2.1	30.0	100.0	1,816
Other Rural	64.1	1.9	34.0	100.0	1,773
Residence					
Urban	66.3	2.4	31.3	100.0	3,572
Rural	64.2	1.8	34.0	100.0	2,125
Age Group					
15–19	9.5	0.0	90.5	100.0	1,094
20–24	49.5	0.3	50.2	100.0	936
25–29	78.9	2.8	18.3	100.0	946
30–34	89.5	1.8	8.6	100.0	1,067
35–39	92.3	3.9	3.8	100.0	958
40–44	92.5	4.6	2.8	100.0	696
Living Children					
0	12.9	0.3	86.8	100.0	1,943
1	94.8	5.2	0.0	100.0	828
2	96.9	3.1	0.0	100.0	1,840
3 +	97.9	2.1	0.0	100.0	1,086
Education Level					
Primary or Less	66.8	2.2	31.0	100.0	2,519
Secondary Incomplete	35.2	1.9	62.9	100.0	653
Secondary Complete	75.1	2.2	22.7	100.0	1,830
Post-Secondary	57.5	1.0	41.5	100.0	695
Socioeconomic Index					
Low	67.3	2.1	30.6	100.0	1,940
Medium	63.0	2.2	34.8	100.0	2,985
High	66.7	1.1	32.2	100.0	772
Employment					
Working	73.6	4.2	22.2	100.0	1,118
Not Working	63.6	1.7	34.7	100.0	4,579

Table 3.4 B Percent Distribution of Marital Status by Selected Characteristics, for Men Aged 15–49 Years Reproductive Health Survey: Albania 2002

		Marital Status			
Characteristics	Married	Previously Married	Never Married	Total	No. of Cases
Total	60.3	0.6	39.0	100.0	1740
Strata					
Metro Tirana	59.8	1.2	39.0	100.0	718
Other Urban	59.8	0.5	39.7	100.0	547
Other Rural	60.9	0.4	38.7	100.0	475
Residence					
Urban	59.6	0.9	39.5	100.0	1155
Rural	61.0	0.4	38.6	100.0	585
Age Group					
15–19	0.0	0.0	100.0	100.0	401
20–24	13.3	0.4	86.3	100.0	189
25–29	60.2	1.0	38.8	100.0	218
30–34	87.3	0.4	12.3	100.0	253
35–39	94.9	0.7	4.5	100.0	255
40–44	98.7	0.7	0.6	100.0	277
45–49	97.4	1.5	1.0	100.0	147
Living Children					
0	13.7	0.5	85.8	100.0	815
1	99.2	0.8	0.0	100.0	221
2	98.8	1.2	0.0	100.0	468
3 +	100.0	0.0	0.0	100.0	236
Education Level					
Primary or Less	60.9	0.6	38.5	100.0	689
Secondary Incomplete	13.9	0.3	85.9	100.0	199
Secondary Complete	72.1	0.6	27.2	100.0	626
Post-Secondary	58.0	0.8	41.2	100.0	226
Socioeconomic Index					
Low	63.6	0.5	35.9	100.0	638
Medium	57.4	0.5	42.1	100.0	814
High	57.3	1.8	40.8	100.0	288
Employment					
Working	75.9	0.8	23.3	100.0	913
Not Working	45.4	0.4	54.2	100.0	827


CHAPTER 4 FERTILITY AND PREGNANCY

Fertility Experience

Current levels of fertility presented in Tables 4.1, 4.2, 4.3A and 4.3B were estimated with the use of five-year agespecific fertility rates (ASFR) calculated from information collected through the respondents' lifetime pregnancy histories. ASFRs are expressed per 1000 women. The total fertility rate (TFR) is computed by multiplying the age-specific fertility rates by five (the number of years in each age group) and summing them over the reproductive ages. When this number is divided by 1000, the TFR can be defined as the average number of live births a women would have during her reproductive lifetime (15-44) if she experienced the observed ASFRs of a given time period. Numerators for the ASFRs were calculated by selecting live births that occurred during the 36-month period preceding the

survey and grouping them (in five-year age groups) by the age of the mother at the time of the reported date of birth. The denominators for the rates represent the number of woman-years lived in each five-year age group during the specified three-year period.

Table 4.1 and Figure 4.1 present agespecific fertility rates calculated from the live birth history asked of every woman in the survey. These rates were calculated for three three-year periods over the last decade, 1993–1996, 1996–1999, and 1999–2002. As can be observed in the Table, the total fertility rate has declined substantially over the 10-year period, from 3.3 to 2.6 children per woman. This was due to pronounced declines in the most fertile ages of 20–24 and 25–29.



As with women in other countries of the region (Table 4.2), Albanian women initiate and complete childbearing at an early age. The highest fertility rates in Albania are among women 20-24 and 25-29 years of age, accounting for 33% and 32%, respectively, of the TFR of 2.6. Women aged 35-44 make a minimal contribution to total fertility of only 8% of the TFR. The adolescent fertility rate is very low, only 35 live births per 1000 women 15-19 years of age, representing 7% of the total fertility rate. The estimated TFR of 2.6 is a bit higher than the rate published by WHO for 2001 (2.4) and the rate of 2.3 published by the UN population Division (WHO, 2003; UN, 2003). The TFR of 2.6 is the highest in Europe and higher than the TFR in 9 of the 13 countries in Eastern Europe and the Former Soviet Union that have conducted similar Reproductive Health Surveys (CDC and MACRO, 2003).

There is no difference in the TFR in Albania by urban or rural residence, although the rate is slightly lower in Metropolitan Tirana than in the rest of the country (Table 4.3A). The TFR for women with a postsecondary education (2.0) is lower than the TFR for those with a primary (2.7) or secondary (2.5) education. Also, although the principal childbearing years for all women are 20–29 years of age, those with a post-secondary education tend to bear their children somewhat later at ages 25–34. There is also a tendency for age at childbearing to increase as SES increases.

Fertility rates for men are shown in Table 4.3B. These rates are based upon the responses of men to questions about children fathered by them. The male TFRs are universally lower than the female TFRs, due to very low fertility rates at ages 15– 24 reflecting a later age at marriage and possibly indicating a tendency for men to underreport children fathered by them. Also, because of the smaller sample size for males with a small number of births reported by men, many of the male ASFRs are considered to be unstable. Nevertheless, the male rates show the same relationship with the selected characteristics as were observed for females in Table 4.3A. The male ASFRs also differ from the female in that they reveal older ages for childbearing, reflecting age differences between partners engaging in sexual intercourse.

Cumulative fertility of Albanian women and men is shown in Tables 4.4A and B. The number of live births (also known as "children ever born") in Table 4.4A shows that 38% of Albanian women in the reproductive years had not had a live birth at the time of the survey, but only 8% of women in union were without a live birth. By age 40–44, only 1% of married women reported never having had a live birth. The median number of live births for married women was 2, and for women at the end of their reproductive years the median increased to 3 live births.

Cumulative fertility levels observed for married men are similar to those of the women. Among all men, however, lower levels of fertility are reported, with 46% stating that they were childless. It is probable that single men are less knowledgeable about the number of children they may have fathered. It is worth noting that all men at ages 45– 49 report completed fertility at similar levels to married women, most likely because most men in this age group are married.

Age at First Intercourse, Union and Birth

Tables 4.5A and B present data on age at first sexual intercourse, first union and first live birth for women and first sexual intercourse and first union for men. respectively, according to their age cohort at the time of the survey. By examining the percentages that have experienced sexual intercourse or marriage or a live birth by current age cohort, it is possible to determine whether the ages at which these events first take place are changing over time. For example, in Table 4.5A, the percentage of women who experienced sexual intercourse before age 18 increased from 10% among current 40-44 year olds to 16% among current 20-24 year olds. Similarly, the percentage of 40-44 year olds who married before age 18 is 7%. whereas 11% of 20-24 year olds married before age 18. In turn, a higher percentage of first live births before age 18 and before age 20 result from the younger ages of first intercourse and first union. Of course, these data cannot reveal the temporal relationship between marriage and intercourse - that is, which of these events preceded the other. It is also important to note that less than half of the women had had intercourse (47%) or had been married (41%) before age 22 and only a quarter had had a live birth (26%) before reaching 22 years of age. These findings are reflected in an average age 21.1 years at first intercourse, 21.9 years at first marriage, and 23.4 years at first live birth. These averages show little change across the age cohorts, suggesting little change over the last two decades in the timing of these events in a woman's life cycle.

Data for the men are shown in Table 4.5B. Here we see a more dramatic change in age at first intercourse across time. Only 5% of men currently aged 45-49 reported having had first sexual intercourse before age 18, compared to 22% of men currently aged 20-24. This substantial increase is also seen for percentages that had first intercourse by the age of 20. However, these increases in sexual experience at younger ages are not reflected in the average age at first intercourse for all ages and cohorts, which remained relatively stable over time. Also, age at first sexual intercourse for men does not correspond as closely to age at first marriage as it does for women. Thus, while the median age at first intercourse for all men is 21.5 years, the median age at first marriage for men is 26.5 years.

When observed by residence (Table 4.6), age at first intercourse for women is rather stable. The median age at first intercourse does not vary between urban and rural areas. However, on average, women in urban areas marry and have their first births a year later than those in rural areas. Educational level shows an even stronger and consistent effect on ages of these three events, resulting in a three-year difference in average age at first intercourse (20.5 vs. 23.3), a four-year difference in age at first marriage (21.0 vs. 25.1) and a fouryear difference in age at first birth (22.5 vs. 26.6), when comparing the lowest and highest educational categories. (See also Figure 4.2.)


The patterns of age at first intercourse for men are somewhat different from those of women (Table 4.6). In contrast to women, the average age at first intercourse for men is higher in rural areas and inversely affected by educational level. For men in rural areas, the median age at first intercourse is 22.2 years compared to 20.6 years for men in urban areas. Likewise,

men with primary or lower levels of education have a median age of 22.0 years at first intercourse compared to 20.3 years for men with post-secondary education. Urbanization and education have the opposite effect on age at marriage for men. The median age at first marriage increases from 25.8 years for men living in rural areas to 27.4 years for those residing in urban areas. Similarly, men with the lowest educational levels marry, on average, at 26.0 years of age compared to 28.6 years of age for men with the highest level of education. The education effects for men are shown also in Figure 4.2.

Current Sexual Activity

Current sexual activity is an important measure for determining the women who are risk of an unintended pregnancy and thus in need of contraceptive services. Table 4.7 shows that 70% of Albanian women in their reproductive years

have had sexual intercourse. This percentage comprises all married and previously married women and 8% of never married women. Seventy-five percent of currently married women had intercourse within the last month, and another 13% had last intercourse 1–3 months prior to the survey. Another 7% reported а pregnancy-related reason for no current sexual activity. Among previously married women, 76% last had intercourse one or more years ago, presumably when they were still married. Almost half (48%) of the sexually experienced never married women have had intercourse within the last month (3.9%/8.2%) and another 23% had their last intercourse 1-3months ago (1.9%/8.2%). Current sexual activity among men is higher than that of women. Seventy-six percent of all men have ever had sexual intercourse, including 38% of never married men. The percentage of men who had



intercourse in the last month was 83% of the currently married and 14% of the never married men. Among the sexually experienced never married men, 38% (14.4%/37.5%) had sex in the last month and another 29% (11.0%/37.5%) had sex 1–3 months prior to the survey.

Induced Abortion

For several decades one of the most outstanding demographic features of most of the Eastern European countries has been the high reliance on induced abortion as a means of birth prevention (David, 1992). Induced abortion was the single most important means of controlling fertility. In recent years, abortion rates and ratios in many of these countries have been among the highest in the world. Among the factors frequently cited as contributing to the reliance on abortion has been the limited availability of modern contraceptive methods, poor quality of methods available, fears about possible side effects, and easy access to and low cost of induced abortion. However, since 1990, data show that an increase in use of modern contraception has been associated with a decline in abortion in many countries of the region (C Westoff et al., 1998, 2000, 2002; CDC and MACRO, 2003).

As with the calculation of the total fertility rate, age-specific induced abortion rates are calculated by using the age of the woman at age of pregnancy termination and then summed over the ages 15–44 to produce a total abortion rate. In Table 4.8, the number of induced abortions per 1000 live births reported in the reproductive health survey for the three years prior to the survey is compared with the official data reported to the Institute of Statistics (INSTAT) for 1999–2001. The survey rate of 73 abortions per 1,000 live births is 64% lower than the official data of 200 per 1,000 live births (three-year average) reported to INSTAT. Over the last three-year period, the official ratio reported to INSTAT has declined from 241 per 1,000 live births to 172 per 1,000, a 29% decrease (Instituti I Statistikes, 2003).

Romania In and the countries of the Former Soviet Union that have conducted Reproductive Health Surveys or Demographic and Health Surveys, reporting of induced abortion by survey respondents has been close to, and in some cases, has exceeded official reporting. (CDC and MACRO, 2003). Only in the Czech Republic has there been severe under-reporting of induced abortion by respondents as appears to be the case in Albania. It is estimated that respondents in the Czech survey only reported between 45% and 50% of induced abortions they underwent (Czech Statistical Office et al., 1995).

There are three principal factors that may affect the under-reporting of induced abortions, even in a country where they are legal, by survey respondents: (1) Under-reporting of unwanted pregnancies that have a higher probability of being terminated by the voluntary interruption of the pregnancy (see next section of this chapter); (2) Under-reporting of clandestine abortions outside of the medical system; and (3) a tendency to declare induced abortion as spontaneous abortions or miscarriages (see next section of this chapter).

Since the apparent underreporting of abortion by survey respondents is at least 50% and may be as high as 77%, and since the underreporting is most likely not a random event but associated with characteristics of the respondent, data from the survey on abortion is probably unreliable. For this reason, the further analysis of abortion data is considered beyond the scope of this report.

Planning Status of the Last Pregnancy

For every pregnancy ended since January 1997, respondents were asked the planning status of their pregnancies at the time of conception. Each pregnancy was classified as either intended (wanted at the time it occurred), mistimed (occurring earlier than intended), unwanted (the respondent did not want any more children), or the respondent was unsure. Mistimed and unwanted pregnancies together constitute unintended pregnancies (Westoff, 1976).

Despite the under-reporting of induced abortions, strongly associated with unwanted pregnancies, the results in Table 4.9A are somewhat useful for examining relative levels of the planning status of the last pregnancy among the various population sub-groups. The sharp differential between pregnancies ending in induced abortion and a live birth (or a current pregnancy) is obvious. Almost two-thirds (65%) of pregnancies ending in induced abortion were reported as unwanted compared with only 3% of current pregnancies and live births. Also, 11% of pregnancies ending in stillbirth, spontaneous abortion, or an ectopic pregnancy, were reported as unwanted. although the proportion would not be expected to be significantly higher than the 3% of live births reported as unwanted. This suggests that some women who experienced an induced abortion reported their pregnancy outcome as a spontaneous abortion.

The proportion of unwanted pregnancies increases as age group and number of living children, two correlated variables, increase, reaching 20% for 35–44 year old women and 28% for women with four or more living children. Mistimed pregnancies are highest for young adults 15–24 years of age and women with no living children. No major differences are seen by residence or by education. A rough adjustment for the underreporting of abortions puts the percentage of unwanted pregnancies closer to 12% (one out every eight pregnancies) compared with the 7% shown in the table.

Among men, for whom there is no pregnancy data, Table 4.9B shows men reporting 98% of last live births as intended, with no differentials observed by residence, age, number of living children or educational level.

Future Fertility Preferences

Data on fertility preferences are needed so that a determination can be made of the appropriate forms of contraception required by couples in the society. Tables 4.10A and B present future fertility preferences of currently married women and men. Approximately two-thirds (63%) of married women want no more children (Table 4.10A). Another 12% want a child after two or more years. The desire for no more children increases with parity from 2% of married women with no living children to 92% of women who have four or more children. The desire to delay the next birth for two years or longer is highest (45%) among women with one child. Age shows the same direct relationship with wanting no more children as observed for parity, with a low of 5% among women 15-19 and a high of 90% among women 40-44 years of age. Wanting to postpone the birth of a child for two or more years is inversely related to age, with 54% of 15-19 year olds and 1% of 35-39 year olds reporting a desire to postpone the next birth by two years or longer.

In Table 4.10B the percentage of married men who want no more children (58%) is similar to that of women, with a corresponding increase with parity and

age. However, compared to women, smaller percentages of men report wanting to postpone the next birth for two or more years (6% vs. 12% for women) and a larger percentage is undecided (13% of men compared to 7% of women). The differences between men and women appear to be most pronounced at parity one. Nineteen percent of men with one living child compared to 45% of women in that category want to postpone the next birth for two or more years. Furthermore, at parity one, 19% of men are undecided about having any more children and another 11% say they want more but do not know when. This contrasts to lower percentages for women at parity one, where the corresponding percentages are 9% and 5%, respectively.

In order to better understand the relationship between number of living children and desire for no more births, Table 4.11 shows this association controlling for fecundity and selected demographic characteristics. While in the aggregate, there appears to be

no affect of residence on desire for more children, the trend by parity indicates that urban low-parity women are more likely to want no more births than rural low-parity women. Urban women with 0, 1, and 2 living children report wanting no more children at levels of 5%, 16% and 80%, respectively, compared to rural women at 0%, 5% and 67%, respectively. When the correlation between parity and age is controlled, both variables appear to have strong independent effects on desire for no more children. This is most noticeable at parities one and two. Only 3% of women 15-24 years of age with one living child state they want no more children, whereas 64% of women 35-44 years of age want no more children. Similarly, 47% of 15-24 year old women with two living children want no more births, and 90% of 35-44 year olds at the same parity want no more births. Education, while in the aggregate appears to have no effect, at parities one and two there is a direct relationship between wanting no more children and level of education.

Chapt	ter 4
-------	-------

Table 4.1

Three-Year Period Age-Specific Fertility Rates for Several Time Periods Among All Women Aged 15–44 **Reproductive Health Survey: Albania 2002**

	Albania	Albania	Albania
Age Group	1993–1996 *	1996–1999 **	1999–2002 ***
15–19	41	38	35
20–24	207	182	170
25–29	221	185	167
30–34	120	92	105
35–39	76	43	34
40–44	0	10	8
Total Fertility Rate	3.3	2.8	2.6

*Period from August 1993 - July 1996

**Period from August 1996 - July 1999

***Period from August 1999 - July 2002

Table 4.2

Three-Year Period* Age–Specific and Total Fertility Rates, Among Women Aged 15–44 Reproductive and Demographic Health Surveys (RHS and DHS) in Selected Eastern European and Former Soviet Union Countries Albania Reproductive Health Survey 2002, Final Report

	Time	Age-Specific Fertility Rates (per 1,000 women) [†]							
Region and Country	Time Period	15–19	20–24	25–29	30–34	35–39	40–44	TFR [‡] 15–44	GFR § 15–44
Eastern Europe									
Albania, 2002	1999–2001	35	170	167	105	34	8	2.6	89
Czech Rep., 1993	1990–1992	49	176	92	41	11	4	1.9	62
Moldova, 1997	1994–1996	57	158	88	40	17	6	1.8	64
Romania, 1999	1997–1999	36	100	83	29	13	2	1.3	49
Russia (three oblasts), 1999 [¶]	1996–1998	39	101	73	28	11	7	1.3	44
Ukraine, 1999	1997–1999	49	115	66	36	14	4	1.4	49
Caucasus									
Armenia, 2000	1998–2000	50	149	88	35	16	3	1.7	56
Azerbaijan, 2001	1998–2000	44	151	133	58	19	9	2.1	71
Georgia, 1999	1997–1999	64	113	92	48	21	7	1.7	61
Central Asia									
Kazakhstan, 1999	1997–1999	40	167	106	64	24	9	2.1	67
Kyrgyz Rep., 1997	1995–1997	75	246	179	113	47	13	3.4	118
Turkmenistan, 2000	1998–2000	30	184	195	105	48	14	2.9	103
Uzbekistan, 1996	1994–1996	61	266	176	114	39	9	3.3	123

* Three years prior to the interview.

[†] Age at pregnancy outcome.

[‡] TFR: Total Fertility Rate (number of births per woman).

§ GFR: General Fertility Rate (births divided by the number of women age 15–44), expressed per 1,000 women

[¶] Yekaterinburg, Perm, and Ivanovo, respectively (predominantly urban sample).

Table 4.3 AAge-Specific Fertility Rates and Total Fertility Rates by Selected CharacteristicsAmong All Women Aged 15–44Reproductive Health Survey: Albania 2002

	Age-	Specific	SFR)	Total				
Characteristics	15–19	20–24	25–29	30–34	35–39	40–44	Fertility Rate *	GFR [†]
Total	35	170	167	105	34	8	2.6	89
Strata								
Metro Tirana	37	143	170	98	30	(5)	2.4	85
Other Urban	38	184	159	100	45	(4)	2.7	92
Other Rural	33	172	171	111	27	(11)	2.6	89
Residence								
Urban	37	167	164	100	41	(4)	2.6	89
Rural	33	172	169	109	27	(11)	2.6	90
Education Level								
Primary or Less	51	187	171	105	28	(6)	2.7	96
Secondary	14	180	169	99	34	11	2.5	83
Post-Secondary	(6)	54	136	134	67	(4)	2.0	75
Socioeconomic Index								
Low	37	196	194	103	34	(9)	2.9	98
Medium	31	153	142	105	30	(8)	2.3	80
High	50	147	181	116	51	(0)	2.7	100

Note: All rates in this table are calculated based on births in the last three years (August 1999–July 2002) and ages of mothers at time of birth.

* The total fertility rate (TFR) is calculated as the sum of ASFR's for each year of age from age 15 to 44.

[†] The general fertility rate (GFR) is calculated as the number of births per 1000 women 15-44.

() Rates considered unstable due to numerators of less than 15 cases.

	Among All Men Aged 15–49 Reproductive Health Survey: Albania 2002								
		Age-Spec	ific Fert	ility Rate	(per 100	0) (ASFR	R)	Total	GFR †
Characteristics	15–19	20–24	25–29	30–34	35–39	40-44	45–49	Fertility Rate *	
Total	(1)	31	134	137	58	30	(8)	2.0	59
Strata									
Metro Tirana	(3)	36	125	125	46	(19)	(15)	1.8	52
Other Urban	(0)	(14)	109	163	52	38	(12)	2.0	62
Other Rural	(0)	38	152	122	68	29	(0)	2.1	60
Residence									
Urban	(0)	25	111	150	51	32	(15)	1.9	58
Rural	(1)	36	154	123	66	28	(0)	2.0	60
Education Level									
Primary or Less	(1)	61	142	143	70	28	(10)	2.3	65
Secondary	(1)	(59)	144	127	49	30	(9)	1.8	61
Post-Secondary	(0)	(4)	65	144	(53)	(43)	(0)	1.5	50
Socioeconomic Index									
Low	(0)	41	134	156	63	31	(3)	2.1	59
Medium	(1)	20	141	114	54	(10)	(8)	1.7	55
High	(0)	(45)	108	155	(54)	93	(40)	2.5	77

Table 4.3 BAge-Specific Fertility Rates and Total Fertility Rates by Selected CharacteristicsAmong All Men Aged 15–49Reproductive Health Survey: Albania 2002

Note: All rates in this table area calculated based on births in the last three years (August 1999–July 2002) and ages of fathers at time of birth.

* The total fertility rate (TFR) is calculated as the sum of ASFR's for each year of age from ages 15 to 49.

[†] The general fertility rate (GFR) is calculated as the number of births per 1000 men 15-49.

() Rates caonsidered unstable due to numerators of less than 15 cases.

Table 4.4 APercent Distribution of Number of Live Births by Current Age ofRespondents Among all Women and Among Married Women Aged 15–44Reproductive Health Survey: Albania 2002

	All Women									
	Age Group (Current Age)									
Number of Live Births	Total	15–19	20–24	25–29	30–34	35–39	40–44			
0	37.8	95.8	64.0	23.0	9.6	6.6	4.1			
1	12.8	3.8	25.0	28.3	10.9	5.8	4.6			
2	27.1	0.4	9.7	39.0	47.5	41.7	35.5			
3	14.9	0.0	1.3	9.5	25.4	31.6	28.2			
4 +	7.4	0.0	0.0	0.2	6.7	14.3	27.5			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0			
No. of Cases	5,697	1,094	936	946	1,067	958	696			

	Married Women									
		Age Group (Current Age)								
Number of Live Births	Total	15–19	20–24	25–29	30-34	35–39	40-44			
0	7.5	55.6	27.5	5.7	1.1	2.9	1.2			
1	18.6	40.0	50.3	34.0	11.8	5.1	3.7			
2	40.4	4.5	19.5	48.4	51.4	43.6	36.4			
3	22.4	0.0	2.6	11.7	28.3	33.4	29.9			
4 +	11.1	0.0	0.0	0.2	7.4	15.0	28.8			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0			
No. of Cases	3,965	97	502	800	1,004	906	656			

Table 4.4 BPercent Distribution of Number of Live Births by Current Age of Respondents
Among All Men and Among Married Men Aged 15–49
Reproductive Health Survey: Albania 2002

	All Men									
		Age Group (Current Age)								
Number of Live Births	Total	15–19	20–24	25–29	30–34	35–39	40–44	45–49		
0	45.5	100.0	94.9	63.4	21.4	7.0	1.6	2.7		
1	11.5	0.0	2.6	22.9	33.5	15.4	6.8	3.2		
2	24.9	0.0	1.5	12.0	36.8	51.3	45.8	38.0		
3	11.8	0.0	1.0	1.6	6.8	20.0	32.3	27.6		
4 +	6.3	0.0	0.0	0.2	1.6	6.3	13.5	28.5		
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
No. of Cases	1,740	401	189	218	253	255	277	147		

		Married Men									
				Age Gro	oup (Curr	ent Age)					
Number of Live Births	Total	15–19	20–24	25–29	30–34	35–39	40–44	45–49			
0	10.3	**	61.7	39.1	10.4	2.7	1.0	1.4			
1	18.9	**	19.2	38.0	38.3	16.2	6.2	3.3			
2	40.8	**	11.5	20.0	41.7	53.4	46.4	37.6			
3	19.6	**	7.6	2.6	7.8	21.1	32.7	28.3			
4 +	10.4	**	0.0	0.3	1.8	6.6	13.7	29.3			
Total	100.0	**	100.0	100.0	100.0	100.0	100.0	100.0			
No. of Cases	1,023	0	32	123	215	242	270	141			

**Percentages are not shown when base is less than 25 cases.

Table 4.5 A

Percent of Women Aged 15–44 Who Had Their First Sexual Relation, First Marriage, and First Birth Before Selected Ages, and Median Age at These Events, by Current Age Reproductive Health Survey: Albania 2002

	Age at First Sexual Intercourse					. Has Had	Never Had	Median	No. of
Current Age	<15	<18	<20	<22	<25	Intercourse	Intercourse	Age †	Cases
15–19	1.2	(10.8)	(13.8)	NA	NA	13.8	86.2	**	1,094
20–24	1.6	16.3	35.1	(50.2)	(55.2)	55.2	44.8	21.5	935
25–29	0.3	12.4	36.9	61.0	80.7	85.1	14.9	20.7	945
30-34	0.3	8.7	30.8	59.4	82.3	92.4	7.6	21.0	1,066
35–39	0.4	10.8	32.2	55.2	82.9	97.2	2.8	21.3	957
40–44	0.6	9.6	31.4	56.6	83.4	97.2	2.8	21.1	696
Total	0.8	11.5	29.2	47.4	63.3	69.9	30.1	21.1	5693*

	Age at First Marriage					Ever in	Never in	Median	No. of	
Current Age	<15	<18	<20	<22	<25	Union	Union	Age †	Cases	
15–19	0.6	(6.8)	(9.5)	NA	NA	9.5	90.5	**	1,094	
20–24	0.9	11.2	27.8	(43.6)	(49.8)	49.8	50.2	22.2	936	
25–29	0.1	7.9	28.0	51.8	75.6	81.7	18.3	21.8	946	
30-34	0.6	6.7	23.5	51.4	79.1	91.4	8.6	21.9	1,067	
35–39	0.4	7.7	24.6	47.6	76.9	96.2	3.8	22.2	958	
40–44	0.2	6.8	25.4	51.5	79.0	97.2	2.8	21.9	696	
Total	0.5	7.8	22.4	40.7	58.5	67.2	32.8	21.9	5,697	

Current Age		Age at	First Liv	e Birth		- Has Had Live Birth	Never Had Live Birth	Median Age⁺	No. of Cases
	<15	<18	<20	<22	<25				
15–19	0.2	(2.5)	(4.2)	NA	NA	4.2	95.8	**	1,094
20–24	0.2	3.8	15.1	(28.3)	(36.0)	36.0	64.0	23.6	936
25–29	0.0	1.6	13.7	37.0	65.8	77.0	23.0	23.3	946
30–34	0.5	2.7	9.7	31.9	69.2	90.4	9.6	23.3	1,066
35–39	0.2	2.1	11.7	31.0	65.3	93.4	6.6	23.7	955
40–44	0.0	2.6	9.8	33.2	66.5	95.9	4.1	23.5	696
Total	0.2	2.6	10.4	26.3	48.3	62.2	37.8	23.4	5693*

* Excludes 4 cases not reporting the date at first sexual intercourse and 4 cases not reporting date at first birth

** Omitted because less than 50% in that age group reported the variable of interest by the end of the interval

() Time exposed partially truncated because not all cases have exposure throughout the period of analysis

NA. Not applicable

⁺ Life table method used in calculation of median age at first intercourse, first marriage and first birth to control for truncated cases

Table 4.5 B

Percent of Men Aged 15–49 Who Had Their First Sexual Relation and First Marriage Before Selected Ages and Median Age at These Events, by Current Age Reproductive Health Survey: Albania 2002

	A	ge at Firs	t Sexual	Intercours	se	- Has Had	Never Had	Median	No. of
Current Age	<15	<18	<20	<22	<25	Intercourse	Intercourse	Age	Cases
15–19	0.6	(4.5)	(4.8)	N/A	N/A	4.8	95.2	**	401
20–24	1.5	21.5	40.9	(56.1)	(61.3)	61.3	38.7	21.5	188
25–29	2.1	15.3	38.6	56.1	80.4	93.6	6.4	20.9	209
30–34	0.4	8.3	39.2	53.8	76.1	99.4	0.6	20.9	241
35–39	0.2	8.3	26.7	48.1	72.2	99.8	0.2	21.9	235
40–44	0.0	8.3	30.8	53.8	74.5	100.0	0.0	21.3	247
45–49	0.0	5.4	22.1	51.3	75.4	100.0	0.0	21.7	132
Total	0.7	10.2	27.9	43.7	59.5	74.3	25.7	21.5	1,653*

Current Age		Age a	t First Ma	rriage		Ever in Union	Never in Union	Median Age	No. of Cases
	<15	<18	<20	<22	<25				
15–19	0.0	(0.0)	(0.0)	N/A	N/A	0.0	100.0	**	401
20–24	0.0	1.0	4.3	(7.7)	(13.7)	13.7	86.3	**	189
25–29	0.0	0.0	3.1	11.1	28.7	61.2	38.8	26.4	218
30–34	0.0	0.0	2.1	6.0	31.6	87.7	12.3	26.9	253
35–39	0.0	0.7	1.0	6.4	30.8	95.5	4.5	26.7	255
40-44	0.0	1.1	5.1	11.0	33.6	99.4	0.6	26.6	277
45–49	0.0	0.5	1.5	13.8	35.8	99.0	1.0	25.8	147
Total	0.0	0.5	2.4	7.4	23.3	61.0	39.0	26.5	1,740

* Excludes 87 cases not reporting the date at first sexual intercourse.

** Omitted because less than 50% in that age group reported the variable of interest by the end of the interval.

() Time exposed partially truncated because not all cases have exposure throughout the period analisys

NA. Not applicable

† Life table method used in calculation of median age at first intercourse, first marriage and first birth to control for truncated cases.

Table 4.6 Median Age at First Sexual Intercourse, First Marriage and First Birth Among Women Aged 15-44 and Men Aged 15-49 by Selected Characteristics **Reproductive Health Survey: Albania 2002**

		Women		Men			
Characteristics	Median Age at First Intercourse	Median Age at First Marriage	Median Age at First Birth	Median Age at First Intercourse	Median Age at First Marriage		
Total	21.1	21.9	23.4	21.5	26.5		
Strata							
Metro Tirana	21.1	22.5	24.1	20.3	27.0		
Other Urban	21.4	22.7	24.0	20.8	27.4		
Other Rural	21.0	21.5	22.9	22.3	25.9		
Residence							
Urban	21.3	22.7	24.1	20.6	27.4		
Rural	21.0	21.5	22.9	22.2	25.8		
Education Level							
Primary or Less	20.5	21.0	22.5	22.0	26.0		
Secondary Incomplete	20.7	22.1	23.5	21.5	25.2		
Secondary Complete	21.7	22.7	24.0	21.2	26.7		
Post-Secondary	23.3	25.1	26.6	20.3	28.6		

Table 4.7 Sexual Activity Status by Current Marital Status for Women Aged 15–44 and Men Aged 15–49 (Percent Distribution) Reproductive Health Survey: Albania 2002

		Women Marital Status						
Sexual Activity Status	Total	Married	Previously Married	Never Married				
Never Had Intercourse	30.1	0.0	0.0	91.8				
Ever Had Intercourse	69.9	100.0	100.0	8.2				
Within The Last Month	50.3	75.3	2.7	3.9				
1–3 Months Ago	8.9	12.5	2.7	1.9				
Over 3 Months But Within Last Year	2.6	3.2	9.9	0.8				
One Year or Longer	3.7	2.5	76.4	1.4				
Currently Pregnant	3.6	5.4	2.5	0.0				
Postpartum	0.7	1.1	0.0	0.0				
Unknown Interval	0.2	0.1	5.8	0.1				
Total	100.0	100.0	100.0	100.0				
No. of Cases	5,697	3,965	88	1,644				
			Men					

		Marital Status							
Sexual Activity Status	Total	Married	Previously Married	Never Married					
Never Had Intercourse	24.4	0.0	**	62.5					
Ever Had Intercourse	75.6	100.0	**	37.5					
Within The Last Month	55.6	82.5	**	14.4					
1–3 Months Ago	8.1	6.2	**	11.0					
Over 3 Months But Within Last Year	2.7	0.8	**	5.5					
One Year or Longer	2.9	1.4	**	4.9					
Partner Currently Pregnant	2.4	3.9	**	0.0					
Partner Postpartum	0.9	1.5	**	0.0					
Unknown Interval	3.0	3.7	**	1.8					
Total	100.0	100.0	**	100.0					
No. of Cases	1,740	1,023	14	703					

** Percentages are not shown when base is less than 25 cases.

Table 4.8 Abortions * per 1000 Births Reported in Reproductive Health Survey and By Albanian Institute of Statistics (INSTAT)[†] Three Year Period: 1999–2001

	INSTAT	RHS (CI) [‡]	Under-Reporting in RHS (CI) [‡]
Abortion/1000 Births	200	73 (46 to 100)	64% (-77% to -50%)

* Arborteve me Nderprerje (Induced Abortions)

[†] Instituti I Statistikes (2003): Http://www.Instat.gov.Al/graphics/doc/tablelat/shno1.html

[‡] 95% confidence interval

ables

Table 4.9 A

Planning Status of the Last Pregnancy Among Women 15–44 Years of Age With at Least One Pregnancy Since January 1997, by Selected Characteristics (Percent Distribution) Reproductive Health Survey: Albania 2002

	Planr	ning Status o	of Last Pregr	nancy		No. of
Characteristics	Intended	Mistimed	Unwanted	Not Sure	Total	Cases
Total	86.7	5.8	7.0	0.5	100.0	2,275
Last Pregnancy Outcome						
Current Pregnancy	86.7	9.3	3.0	1.0	100.0	222
Live Birth	91.7	4.8	3.2	0.3	100.0	1,811
Induced Abortion	22.2	11.6	64.7	1.5	100.0	149
Other Pregnancy Outcomes *	78.8	8.9	11.4	0.8	100.0	93
Strata						
Metro Tirana	83.3	7.9	8.5	0.3	100.0	844
Other Urban	85.3	7.1	7.3	0.3	100.0	730
Other Rural	88.5	4.5	6.4	0.6	100.0	701
Residence						
Urban	84.5	7.6	7.6	0.3	100.0	1,410
Rural	88.4	4.5	6.5	0.6	100.0	865
Age at The Time of The Last Pregna	ncv Outcom	ne†				
< 20	88.5	10.4	1.1	0.0	100.0	135
20–24	92.0	6.6	1.4	0.1	100.0	653
25–29	87.1	6.7	5.6	0.5	100.0	798
30–34	82.0	3.5	13.2	1.3	100.0	498
35+	78.2	1.7	20.1	0.0	100.0	191
Marital Status at Last Pregnancy						
Currently Married	86.7	5.7	7.1	0.5	100.0	2,171
Not Currently Married	86.7	8.2	5.2	0.0	100.0	104
Living Children						
0	87.4	11.3	1.3	0.0	100.0	105
1	93.7	5.0	1.0	0.2	100.0	679
2	87.6	6.7	5.6	0.0	100.0	934
3	80.1	5.9	12.1	2.0	100.0	419
4 +	71.2	0.9	27.9	0.0	100.0	138
Education Level						
Primary or Less	87.5	5.4	6.7	0.4	100.0	1,072
Secondary Incomplete	84.4	8.5	7.1	0.0	100.0	158
Secondary Complete	86.5	5.0	7.8	0.8	100.0	795
Post-Secondary	84.1	10.0	5.9	0.0	100.0	250

* Includes pregnancies resulting in stillbirth, miscarriage or ectopic pregnancy.

[†] Age of the woman at the time of pregnancy outcome.

Table 4.9 B Planning Status of the Last Live Birth Among Men 15–49 Years of Age With Partner Having at Least One Live Birth Since January 1997, by Selected Characteristics (Percent Distribution) Reproductive Health Survey: Albania 2002

	Planr	ning Status o	nancy		No. of	
Characteristics	Intended	Mistimed	Unwanted	Not Sure	Total	Cases
Total	98.4	1.0	0.1	0.5	100.0	488
Strata						
Metro Tirana	96.4	3.1	0.5	0.0	100.0	181
Other Urban	98.8	0.6	0.0	0.6	100.0	162
Other Rural	98.7	0.7	0.0	0.6	100.0	145
Residence						
Urban	98.3	1.2	0.2	0.4	100.0	312
Rural	98.5	0.9	0.0	0.6	100.0	176
Age at Time of The Last Live Birth*						
< 25	98.6	1.4	0.0	0.0	100.0	27
25–29	97.9	2.1	0.0	0.0	100.0	152
30–34	99.3	0.7	0.0	0.0	100.0	189
35+	97.4	0.0	0.4	2.2	100.0	120
Living Children						
0–1	98.9	1.1	0.0	0.0	100.0	164
2	98.0	1.6	0.0	0.4	100.0	227
3 +	98.2	0.0	0.4	1.4	100.0	97
Education Level						
Primary or Less	99.0	0.3	0.0	0.6	100.0	212
Secondary Incomplete	**	**	**	**	**	6
Secondary Complete	97.6	1.8	0.2	0.5	100.0	214
Post-Secondary	97.8	2.2	0.0	0.0	100.0	56

* Age of the man at the time of live birth outcome

 $\ensuremath{^{**}\text{Percentages}}$ are not shown when base is less than 25 cases.

Table 4.10 A Fertility Preferences of Currently Married Women Aged 15-44 Years By Number of Living Children and by Age Group (Percent Distribution) **Reproductive Health Survey: Albania 2002**

		Number of Living Children [†]						
Preference for Children	Total	0	1	2	3	4 +		
Want a Child Now	3.4	29.2	6.5	1.4	0.3	0.3		
Want a Child Within a Year	3.1	21.0	6.1	1.4	1.0	0.3		
Want a Child in 1–2 Years	5.8	14.1	17.7	3.6	1.0	0.3		
Want a Child After 2 or More Years	12.4	8.7	44.6	7.6	1.9	0.1		
Want More But Do Not Know When	2.7	3.5	4.8	2.9	1.3	0.9		
Want No (no more) Children	62.5	2.0	10.0	72.5	86.7	91.5		
Undecided	7.3	6.6	8.5	8.6	5.8	3.6		
Subfecund, Infecund	2.8	14.9	1.8	2.1	1.9	2.9		
Total	100.0	100.0	100.0	100.0	100.0	100.0		
No. of Cases	3,961*	211	783	1,864	809	294		
		Age Group						

		Age Gloup							
Preference for Children	Total	15–19	20–24	25–29	30–34	35–39	40–44		
Want a Child Now	3.4	11.3	7.8	4.5	2.8	1.7	1.2		
Want a Child Within a Year	3.1	5.1	9.6	3.7	2.9	1.1	0.5		
Want a Child in 1–2 Years	5.8	6.4	15.0	12.1	5.1	1.1	0.3		
Want a Child After 2 or More Years	12.4	53.6	38.4	23.2	5.5	1.1	0.0		
Want More But Do Not Know When	2.7	5.7	3.9	3.7	3.1	2.2	0.8		
Want No (no more) Children	62.5	5.1	15.2	41.1	68.4	84.2	89.9		
Undecided	7.3	12.9	9.5	10.6	11.1	4.8	0.9		
Subfecund, Infecund	2.8	0.0	0.6	1.1	1.1	3.8	6.5		
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
No. of Cases	3,961*	96	501	800	1,003	906	655		

* Excludes 4 women with missing information.

⁺ Women who were pregnant at the time of interview are classified as having one more child than their actual number of living children.

Table 4.10 B Fertility Preferences of Currently Married Men Aged 15–49 Years By Number of Living Children and by Age Group (Percent Distribution) Reproductive Health Survey: Albania 2002

		Number of Living Children [†]							
Preference for Children	Total	0	1	2	3	4 +			
Want a Child Now	4.1	26.2	8.2	0.7	0.0	0.0			
Want a Child Within a Year	4.3	25.6	8.6	0.5	0.8	0.4			
Want a Child in 1–2 Years	6.2	9.1	21.5	2.9	0.0	0.0			
Want a Child After 2 or More Years	5.6	3.2	18.6	4.1	0.0	0.0			
Want More But Do Not Know When	3.7	6.9	10.9	2.0	1.0	0.0			
Want No (no more) Children	58.2	4.5	9.9	74.7	85.5	76.2			
Undecided	12.9	3.8	18.8	13.4	9.4	14.1			
Subfecund, Infecund	4.9	20.7	3.5	1.7	3.4	9.3			
Total	100.0	100.0	100.0	100.0	100.0	100.0			
No. of Cases	1,022*	88	221	473	170	70			

		Age Group							
Preference for Children	Total	20–24	25–29	30–34	35–39	40–44	45–49		
Want a Child Now	4.1	15.6	14.6	5.7	2.1	0.3	0.3		
Want a Child Within a Year	4.3	19.1	12.0	6.9	1.4	1.6	0.3		
Want a Child in 1–2 Years	6.2	8.9	15.2	13.9	3.4	1.7	0.3		
Want a Child After 2 or More Years	5.6	17.8	14.6	12.7	2.7	0.2	0.0		
Want More But Do Not Know When	3.7	5.1	8.9	6.6	4.0	0.5	0.6		
Want No (no more) Children	58.2	14.0	15.3	32.0	64.7	84.0	84.2		
Undecided	12.9	19.5	11.6	19.4	18.2	7.8	6.0		
Subfecund, Infecund	4.9	0.0	7.8	2.9	3.5	4.0	8.4		
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0		
No. of Cases	1,022*	32	122	215	242	270	141		

* Excludes 1 man with missing information.

[†] Men whose partner was pregnant at the time of interview are clasified as having one more child than their actual number of living children.

Table 4.11 Percentage of Fecund Married Women Reporting They Want no More Children by Number of Living Children and Selected Characteristics Fecund Women 15–44 Years of Age Reproductive Health Survey: Albania 2002

			Number o	of Living C	hildren [†]	
Characteristics	Total	0	1	2	3	4 +
Total	64.3	2.3	10.1	74.0	88.5	94.2
No. of Cases	3,866*	183	769	1,833	795	286
Strata						
Metro Tirana	62.6	0.0	17.5	78.8	90.3	94.1
Other Urban	66.9	7.2	13.8	79.7	87.2	93.3
Other Rural	63.3	0.0	5.3	66.9	88.6	94.5
Residence						
Urban	65.8	4.8	16.1	79.8	87.9	93.8
Rural	63.1	0.0	5.2	67.2	88.8	94.4
Age Group						
15–24	13.3	2.5	2.7	46.8	69.4	**
25–34	56.3	0.0	7.7	65.4	78.1	81.2
35–44	91.7	**	63.7	90.3	95.6	96.8
Education Level						
Primary or Less	61.1	0.8	4.9	66.3	87.5	94.0
Secondary Incomplete	58.4	**	16.0	66.0	90.5	‡
Secondary Complete	71.0	8.5	15.3	81.7	90.3	94.3
Post-Secondary	63.1	0.0	20.9	85.6	85.1	‡

* Excludes 4 women with missing information.

[†] Women who were pregnant at the time of interview are clasified as having one more child than their actual number of living children.

[‡] Percentages are not shown when base is less than 25 cases.

Chapter 5 Maternal and Child health

dequate perinatal care is an essential approach in identifying and addressing risk factors that may affect the health of mothers and their babies. In Albania, perinatal care has been organized as a vertical program controlled by the Ministry of Health and for many years women have had free access. Currently, under the new health reform, it is included in all three sub systems: primary health care, secondary health care and tertiary health care.

Perinatal care consists of three components: preconception care, prenatal care and postnatal care. Preconceptional counseling and prenatal care are generally offered by primary care providers and consists of a wide range of information. Information includes risks associated with pregnancies, health risk factors that can affect the development of the fetus (e.g. tobacco and alcohol), maternal infection (e.g. rubella, toxoplasma, HIV and other sexually transmitted diseases), risks associated with maternal health conditions, and risks associated with genetic conditions.

Efforts are being made by the Ministry of Health of Albania to organize preconception counseling, especially in addressing the high prevalence of genetic conditions in some areas of the country. Nevertheless, preconception counseling is not routinely provided during health care visits in spite of the essential role the primary care provider plays in modifying women's health behaviors (many healthy behaviors must be in place before the pregnancy is recognized) and in identifying medical conditions that require special attention during pregnancy.

The use of timely and periodic prenatal care can assist in the identification and/ or prevention of perinatal morbidity and

prevention of mortality. In Albania, public prenatal care is organized within the primary health care subsystem, and in urban areas, it is offered in women's clinics (policlinics), and in both urban and rural areas the service is provided by family doctors (GP's) in their health centers. There are 95 women's clinics and 582 health centers in the urban areas. Prenatal care includes a general risk assessment, consisting of a medical examination and a series of laboratory tests, such as blood, urine, vaginal bacteriological exams, screening for sexually transmitted infections. and isoimmunization Rh. Pregnant women in Albania are entitled to use these public services free of charge. In urban areas, mainly in Tirana, there are an increasing number of private clinics that offer prenatal services, especially the use of ultrasound exams during pregnancy. Although women's clinics are now separated from well baby clinics in urban areas, postpartum care is performed jointly with infant care visits during the first year postpartum.

The Albanian Reproductive Health Survey looked at a number of factors which can have a considerable impact on the health of a woman, the health of her baby, and the outcome of her pregnancy. The instrument used for the survey covered issues such as: the use of health care services related to pregnancy; health related behaviors during pregnancy; the place of delivery; type and assistance at delivery; and postpartum behaviors. including infant feeding practices. However, the sample size allows the ability to estimate infant and child mortality indicators for only the ten year period prior to the survey.

In this chapter selected aspects of maternal and child care in Albania will be examined. Such aspects include sources of health care, utilization of maternal health care clinics, quality of care, etc. The aim is to identify subgroups with specific needs of care and to investigate maternal and child outcomes, which may be related to the availability and quality of maternity care services. All this information will be used to help direct or modify program interventions.

Prenatal care

Prenatal care is most effective when it is initiated in the early stages of pregnancy, is continued throughout gestation, and is comprehensive. For the optimal health of the mother and child, it is recommended that every pregnant woman starts seeing a health care provider for prenatal care examinations during her first trimester of pregnancy.

This section describes the use of prenatal care for all pregnancies ending in a live birth since January 1997. Women participating in the survey were asked about the total number of prenatal care visits they have had during their pregnancy (information did not include visits made just to confirm pregnancy or the use of health care services for the delivery only). Another question regarded the week or month of gestation when they had their first visit for prenatal care.

Table 5.1 displays prenatal care which is distributed by pregnancy trimester of first visit and number of prenatal visits by selected categories. Nineteen percent of all pregnancies ending in a live birth since January 1997 have not received any prenatal care by health professionals. The figure is the highest when compared with Eastern European countries and it is among the highest even when compared with the Caucasus Region and Central Asian countries (CDC and ORC MACRO, 2004).

Although there is a considerable amount of

variation between countries on how many prenatal visits a pregnant woman should make and when they should make their first visit, it is generally accepted that the first visit should take place within the first three months after conception. Among pregnant women who had a live birth since January 1997, including those who have not had any prenatal visits, only 59% received their first prenatal care during the first trimester. During the second trimester the first visit was made by 18 % of pregnant women, and the remaining 3% had their first prenatal visit only during the third trimester. Almost one-fifth of women (19%) reported no prenatal care.

While there are virtually no differences between women living in Tirana and women living in other urban areas in Albania. rural women are by far, more likely (two and a half times more) to carry their birth to term without having any prenatal visits. In addition to rural women there are two additional socio-economic factors. which increases the risk of not having any prenatal visits: level of education and socio economic index. One out of four women among those with only a primary education or less have not had any visits, compared to 7% for women who have had their university studies. The picture is similar when considering the socioeconomic index; women with a low socioeconomic index are almost two and a half times more likely to not have had any prenatal care visits.

Other characteristics that influence not having any prenatal care visits are the age of the mothers and the birth order. Table 5.1 shows a steady trend; women of older ages are more inclined to not receive any prenatal care and among those in the 35–44 age group the likelihood of not receiving prenatal care is almost twice as high as those under 20 years old. Almost 30% of women who have had three or more births made no prenatal visits.

The place of residence, education, and socioeconomic index are also three important factors which influence the early starting of prenatal care. More women living in an urban area started their prenatal care earlier compared to women living in a rural area: 71% versus 51%. Differences among various educational and socio-economic groups are even higher. Only one-fourth of women with a post-secondary education had their first visit after the first trimester or no visits while almost half of the women with a primary education do so. When analyzing the socioeconomic index, the picture is the same: 79% of women classified in the high socioeconomic index start their prenatal care during the first trimester versus only 49 % of those classified in the low index.

Prenatal care should not only start early but also continue throughout pregnancy, according to the recommended standards of periodicity. To asses the adequacy of prenatal care it is necessary to monitor both the time of the first visit and the number of prenatal care visits. In our study the relative majority of women (slightly less than half of those who had some prenatal care) have had only 1–3 visits. The average number of prenatal visits among all pregnant women was around 3 ranging from no visits at all to 27 visits.

Women living in urban areas, women with higher education, and those with a higher socioeconomic status use the prenatal care services more frequently compared to the women living in rural areas, with less education and a lower socioeconomic status.

When compared to other countries in the sub-region, the same differences between areas of residence and education of mothers are noticed, but the range of differences is more similar to the Caucasus countries than Eastern Europe. Generally, in all of these countries, urban living and better educated women use the health services more frequently for prenatal care, compared to women living in rural areas and having a lower educational status. Nevertheless, only among countries such as Georgia, Azerbaijan or Armenia are differences as high as those seen in Albania (CDC and ORC MACRO, 2004).


The baby's weight at birth does not seem to be associated with onset of prenatal care. The proportions of the two categories – under 2500 kg and 2500 kg and above– are quite similar.


Prenatal care is inadequate in some of the countries of Eastern Europe and the former Soviet Union (successor states of USSR). In recent reproductive health surveys (RHS) and demographic and health surveys (DHS) conducted in the region, the proportion of pregnant women with no prenatal care was less than 1% in Czech Republic, 1% in Moldova, 4% in Russia, 8% in Armenia, 9% in Ukraine and Georgia, between 2% and 55% in Central Asian Republics, 11% in Romania, and 30% in Azerbaijan (CDC and ORC MACRO, 2004). Late prenatal care is also common. With the exception of Czech Republic, where more than 90% of women began receiving care in the first trimester, in all other countries less than three-fourths of women entered prenatal care early. Late prenatal care was more prevalent in the Caucasus region than in other regions. In the United States, in 2000, 83% of pregnant women began prenatal care in the first trimester, while only 4% had no prenatal care or late care (third trimester) (CDC, 2002).

In Albania there are no official indicators to measure the adequacy of prenatal care. In the United States the adequacy of prenatal care is assessed by using the Adequacy of Prenatal Care Utilization Index (APNCU), also known as the Kotelchuck index (Kotelchuck, 1994). This index is based on the recommendation of the American College of Obstetricians and Gynecologists, and it is used in all similar reproductive surveys, which makes it a good comparison indicator. It combines the month when prenatal care begins with the number of visits received. Inadequate care is defined as no or late prenatal care or less than 50% of recommended visits. The three remaining levels (intermediate, adequate and adequate plus) require an early initiation of care by the fourth month of gestation. Intermediate care requires 50%–79% of the recommended number of visits; adequate care requires

80–109%; and adequate plus level requires 110% or more of the recommended number of visits.

Using the recommendations of the American College of Obstetricians and Gynecologists for number of visits, as has been used in other reproductive health surveys in Eastern Europe, the adequacy of prenatal care in Albania assessed by the Kotelchuck Index compared with four other countries is shown below. The percentage of women in each category is shown across multiple countries (figure 5.1) and for Albania (figure 5.2)





*Also known as the Kotelchuck Index, it is a measure of adequacy of prenatal care based on initiation of such care (no prenatal care automatically warrants "Inadequate" level) and the number of required visits adjusted for the length of gestation and the gestational age at first visit.

Inadequate care ranges from 13% in Moldova to 81% in Azerbaijan. For Albania, inadequate care is estimated at 70% of the women eligible for prenatal care with live births since 1997. Adequate and Adequate Plus care ranges from 6% in Azerbaijan to 68% in Moldova. The percentage of women with adequate or adequate plus care in Albania is estimated at 9%, somewhat higher than Azerbaijan and somewhat lower than Romania.

In the prenatal health care package an important component is the dissemination of health information. Especially when preconception care is missing, the first prenatal visit is a critical opportunity to screen women for behavioral risk factors such as tobacco or alcohol use, medical and genetical risks, occupational risks and to provide comprehensive counseling. Counseling should include information about maternal behaviors and exposures that may affect the health of the fetus, nutrition, rest, and early signs and symptoms of pregnancy complications. In addition, approaching the time of delivery, counseling should prepare women for what they will face when giving birth, distribute accurate information regarding labor and delivery, and advise about techniques to reduce pain and anxiety during labor. Also, counseling about breastfeeding and family planning after birth should be initiated during the prenatal period and reinforced during post partum care.

The majority of the women (37%) had most of their prenatal care visits in a district hospital or in the Tirana maternity hospital. A similar proportion (33%) used policlinics, especially in Tirana or in another city, for most of their prenatal care visits and 23% of pregnant women used health centers or health posts for their visits (data not shown). It seems that some proportion of women living in rural areas prefer to use health services located in Tirana or other urban areas instead of going to the nearest health center or health post for examinations and counseling. Only 7% of pregnant women sought prenatal care in private clinics, ranging from 13% in Tirana to 5% in rural areas.

To complete the picture of the quality of prenatal care, besides the utilization of services discussed above, the reproductive health survey included additional questions aimed at assessing information received and measurements performed during the prenatal visit; that is the adequacy of the content of prenatal care. Table 5.2 shows the percentage of pregnant women who received some information about specific educational topics during their prenatal care visit.

Only 60% of women who attended prenatal care clinics received some counseling about nutrition during pregnancy. The proportion of women who received counseling on breastfeeding and delivery is approximately the same, slightly over 58%. These three topics are, nevertheless, the most discussed during the prenatal visits among pregnant women in Albania. Other topics, like postnatal care, pregnancy complications, effects of smoking and alcohol, are remembered to have been discussed with a health professional by roughly half of the pregnant women. Contraception was less frequently mentioned as a counseling topic during prenatal care visits; less than 40% of women reported it.

Maternal characteristics that appear to be associated with lower levels of counseling for almost all the topics include rural residence, low levels of education (less than secondary education), and a low socioeconomic index. Other maternal characteristics related with poor counseling practices during prenatal care are age of the mother, under 20 years old, and having two or more previous pregnancies. The number of prenatal visits was directly related to the proportion of women receiving information during their prenatal care visits.

In addition to counseling, the first prenatal visit should include a detailed medical history of the women and her family, including information about risk factors and genetic disorders, a detailed obstetrical history, a comprehensive physical examination, measurement of blood pressure, urine tests, basic blood tests, an ultrasound, and tests for various types of infection. Monitoring of mothers weight, blood pressure, and basic blood tests are extended during the follow up visits.

The proportion of women who have received examinations such as blood tests, urine tests and measurement of blood pressure is above 80% (Table 5.3). About three-fourths (77%) of women had an ultrasound examination during their prenatal care visit, while only 62% stated that they received a tetanus immunization. Only one in four pregnant women received iron supplements as part of their prenatal care.

Residing in a rural area, a low educational level and a low socioeconomic status remains associated with lower proportions of exposure to most of the selected procedures presented in table 5.3. On the other hand, having more than three prenatal care visits increases the chances of receiving an ultrasound examination. A private clinic, as a source of prenatal care, raised the odds for having an ultrasound exam and blood and urine tests but not for the measurement of blood pressure or other selected procedures.

Routine measurement of blood pressure is an essential component of health risk assessment during prenatal visits. Table 5.4 clearly shows that the majority of women who gave birth during the 1997–2002 period had routine measurements of their blood pressure during pregnancy (83.3%). Among them, more than one in ten has been told by a doctor that they have high blood pressure. The proportion of those hospitalized exclusively for this condition was very low (only 0.5%), while more than half of the pregnant women diagnosed with high blood pressure received treatment for it.

Although blood pressure was measured more frequently among more educated, higher socioeconomic status women and those living in urban areas, high blood pressure was found at a higher prevalence among rural women and women with lower educational and socioeconomic level. Treatment and hospitalization for high blood pressure among pregnant women followed the same profile. The number of prenatal visits steadily increases the identification rate of high blood pressure. As a result, the prevalence of high blood pressure among women, having more than 10 prenatal care visits, is almost three times higher compared to women who have attended only one to three prenatal visits.

Table 5.5 shows the prevalence of ultrasound exams during pregnancies, carried to term, between 1997 and 2002. Overall, about three in four pregnant women (77%) had at least one ultrasound exam. This prevalence is lower than that observed in the Czech Republic in 1993, similar to the more recent data of the Ukraine (78%) and Moldova (75%), but much higher than the prevalence observed in Romania (47%) and some Caucasus Countries; Georgia (54%) and Azerbaijan (26%).

Ultrasound exam rates in Albania, similarly to those observed elsewhere, are higher in urban areas, among better educated and higher socioeconomic status women, and among women having their first child. Ultrasound examinations rates are positively associated with the number of prenatal visits.

Almost one in three women (30%) had their first ultrasound exam very early in their pregnancy (less than 14 weeks) while the relative majority of first exams occurred between 14 and 26 weeks of pregnancy (44%). The three characteristics mentioned before (urban residence, a high educational level and a high socioeconomic status) are associated to starting an ultrasound examination early among pregnant women. The utilization of a private clinic for prenatal care visits is another important factor that is positively associated to the starting ultrasound examinations early; around 43% of pregnant women using a private clinic as a source of prenatal care receive their first ultrasound exam within the first 14 weeks of pregnancy.

The reproductive health survey data does not allow differentiation between selected specific indications (e.g. confirmation of gestational age, assessment of fetal viability, fetal malformations, fetal growth, fetal presentation and multiple pregnancy, examination of the placenta, assessment of amniotic fluid) or for routine ultrasound screening, either during early pregnancy (less than 14 weeks) or in late pregnancy (after 27 weeks).

Almost one in three women had their first ultrasound exam very early in their pregnancy (less than 14 weeks) while the relative majority of first exams occurred between 14 and 26 weeks of pregnancy (43.8%). The three characteristics mentioned before (urban residence, a high educational level and a high socioeconomic status) are associated with having an ultrasound examination early among pregnant women. The utilization of a private clinic for prenatal care visits is another important factor that is positively associated with starting ultrasound examinations early; around 43% of pregnant women using a private clinic as a source of prenatal care receive their first ultrasound exam within the first 14 weeks of pregnancy.

Pregnancy complications

Table 5.6 presents pregnancy complications that required medical attention distributed by selected characteristics. Among all pregnancies brought to term since January 1997 and that have some prenatal care, almost one in four were reported to have some kind of pregnancy complication. The most frequently mentioned complication was the risk of preterm delivery (10%), followed by anemia related to pregnancy (7%), a weak cervix, water retention or edema, an urinary tract infection (around 6% each), followed by high blood pressure related to pregnancy, bleeding, and Rh isoimunisation. There are no significant differences among the different subgroups, based on background characteristics of the mothers

Intrapartum care

In Albania all births are recommended to occur in medical facilities where adequately trained personnel can monitor the progress of labor and delivery. The reproductive health survey data demonstrates that 85% of births in the country are occurring in a district maternity hospital or in Tirana maternity (Table 5.7). Only around 8% of deliveries take place in "birth houses"; 13% in rural areas. Births delivered outside medical facilities or at home are rare but still represent a significant 6% of total births in the country.

The proportion of births occurring outside a hospital or in birth houses become even higher in several subgroups of the population reaching 22% in rural areas, 19% among women with primary education, and almost 21% among those of low socioeconomic status. Giving birth at home is highly associated with inadequate prenatal care; women who had no prenatal care visits have a four times higher risk of delivering their baby at home, compared to those who have had some prenatal care visits. The rate of giving birth at home is more than 15% among this subgroup. Other characteristics which are likely to increase the risk of giving birth at home are age of the women (those 35–44 years have a rate of 11%) and birth order (10% for women who already have two children)

Delivery at a birth house is affected by the same factors as the delivery at home. The phenomenon of giving birth in a private clinic or hospital remains extremely rare in Albania; only among women of higher socioeconomic status does this proportion reach 2.5%.

Table 5.8 shows the time spent in a medical facility prior to delivery and the length of stay after delivery. The average time spent prior to delivery in the hospital was about 7 hours (shorter than the times observed in Romania and Georgia, but similar to that observed in Azerbaijan) (CDC and ORC MACRO, 2004). The average duration of labor generally ranges from 6 hours (for multiparous women) to 10 hours (for nulliparous women). Thus, many women, particularly those giving birth for the first time were admitted for delivery during or right after the onset of labor. The average time spent in the medical facilities prior to delivery was slightly shorter for women living in urban areas, and those of higher socioeconomic status, probably indicating better access to hospitals by women of these subgroups. The time spent in the hospital prior to delivery was longer for nulliparous women (almost 10 hours), those who deliver by C-section and those who gave birth to babies weighing less than 2500 g.

The majority of Albanian women (59%) stay in the medical premises less than three days after giving birth. Less than 20% stay in the hospital for more than four days and most of these are women who delivered by C-section. Only low birth weight babies and C-section deliveries are factors which highly increase the period of stay in the hospital after delivery. Other characteristics do not seem to be associated with period of stay in the hospital after delivery.

Table 5.9 shows the percentage of births by C-section by selected delivered characteristics. The Caesarian section (Csection) rate varies considerably among countries, from about 5% to more than 20% of all deliveries. The optimal rate is not known, but little improvement in birth outcomes has been demonstrated if the rate is higher that 7%. In addition to unequivocal obstetrical indications, a C-section is often performed in less clear situations (e.g. prolonged labor) and often if a previous Csection was performed, which is rarely an adequate indication by itself.

The reproductive health survey shows that although most births are delivered vaginally, in Albania, between 1997 and 2002, the rate of C-section deliveries was 13.4% and this figure is only slightly higher than those reported by several Eastern European countries and the Caucasus region, where similar surveys were carried out. The Csection rate in those countries ranges from 3% in Azerbaijan to 11% in Romania and 12% in Russian areas with surveys (CDC and ORC MACRO, 2004).

Women residing in Tirana were almost twice as likely to have a C-section delivery compared to women in rural areas, demonstrating that the bases of decision for a C-section may include other reasons apart from medical ones. The socioeconomic index was another factor seemingly associated with this type of delivery. Women of a high socioeconomic status are more than twice as likely to have a C-section delivery. Factors which might increase the chances of having a C-section are the age of the mother (especially over 35 years), prolonged labor, pregnancy complications, low weight of the baby at birth, and to a lesser extent, being a first time mother.

The Albania RHS also included a question directed to women who gave birth during the study period on the most important reasons for a C-section delivery. The most frequent reasons given by them included mainly clinical factors like malpresentation of fetus (35%), prolonged labor or baby started to suffer (22%), and baby too big for vaginal delivery (7%). Having had a previous C-section is also mentioned quite frequently as a reason for the actual C-section (22%). There were only 6% of women that requested a C-section delivery. Although small numbers do not allow important conclusions to be made, this last reason is more frequently found in urban areas compared to rural areas (9% versus 2%).

Poor birth outcomes

Poor birth outcomes are considered stillbirths, preterm births (live births within 37 weeks of gestation) and infants weighting less than 2500 grams at birth (low birth rate-LBW). Selected poor birth outcomes in the five years preceding the survey are shown in table 5.10.

The incidence of low birth weight (LBW) for infants born alive during the study period was 4.6%. It is only slightly higher than the rate reported by a recent survey in Albania (UNICEF and INSTAT, 2000). Nevertheless, it is lower than the rates produced by similar studies in some Eastern European countries and the former Soviet Union. There are no clear trends in low birth rate among selected background categories of women apart from age and birth order; LBW rate was respectively 5.7% and 6.7% among women younger than 20 years and first time mothers, while only 1.5% among the 35–44 age group and 2.4% among women who have had three or more births.

The preterm birth rate is slightly lower than the LBW rate (3.6%) but the birth order characteristic has the same profile as the LBW rate. Prolonged labor increases more than twofold the risk of LBW (slightly less that of prematurity birth), and this fact is consistent with other findings in other similar studies.

Postnatal care

After the birth of the child, it is important to provide appropriate postnatal care for both the health of the mother and the child, which must include counseling about breastfeeding, nutrition, and family planning. The postnatal period is a critical time that allows the health care provider to evaluate the physical and psychological health of the new mother and her infant, to detect and treat possible postpartum complications and to provide the support needed to address any specific problems related to child care.

The Albania RHS provided information on the use of postnatal care and the content of postnatal counseling. As it is clearly demonstrated by Table 5.11, the percentage of women who attended a postpartum care visit within a specified time period is only a fraction of those who have been receiving some kind of prenatal care. Less than one in five women (19%), currently aged 15-44, who delivered live birth babies during the 1997-2002 period, had at least one postnatal care visit. Furthermore, only 36% had their first postnatal visit within the first week after the delivery. About 61% of women had their first visit within the first two weeks after the delivery. Urban residency, high

level of education and high socioeconomic category were the main factors influencing a higher rate of postpartum care. Those giving birth at the Tirana Maternity Hospital, first time mothers, and those with postpartum complications were also more inclined to use the postnatal care services than other women.

Information on whether women received a postpartum examination following their most recent live births was collected by the RHS or DHS surveys in five other countries in Eastern Europe and the Caucasus Region. There was a high variation in the proportion of women across these countries who received postpartum care, raging from 74% in Moldova down to 11% in Georgia. Except in Moldova, postpartum care coverage was always higher in urban areas than in rural areas. Coverage also tended to increase with education.

During the postnatal care visits more than 70% of the women received information on immunization, child care, nutrition, breastfeeding and breast care by a health professional (Table 5.12). Although less frequent, contraception counseling was also received by slightly more than half of the women (55%). The differences, regarding this topic, between selected categories are not very sharp but proportions are always lower among women living in the rural areas, with lower education, and a lower socioeconomic category.

The Albania RHS questionnaire included questions regarding healthcare for the baby after the delivery. Table 5.13 shows the proportion of live births followed by postnatal baby care, whether that care was received during a routine health exam or sickness, and the timing when the care happened. More than 85% of babies were seen by a health professional and there are very small differences between the various categories. In only a few cases were the visits made to examine a sick baby (8%). Almost everyone used baby health care as a part of a routine health exam; 45% of these visits took place during the first week after the delivery, while slightly more than 20% was done only after the second week of the baby's life.

The proportion of women who registered their newborns was more than 92%. Only women under the age of 20 seem to have lower than a 90% rate of registration. Between the other categories, there are no significant differences. Around 20% of mothers registered their babies during the first week after the delivery. This proportion increases to almost 70% after the second week (Table 5.14).

Smoking and drinking during pregnancy

Smoking and alcohol drinking during pregnancy does not seem to be a problem in Albania; very few women smoke before becoming pregnant (1.3%) and almost half of them stop smoking when they become pregnant. Among 3.1% of women drink some alcohol during pregnancy, the majority drinks less than once a week (data not shown).

Postpartum complications

Postpartum complications reported by women who gave birth in the five years prior to the survey are shown in table 5.15. Roughly one in four women experienced at least one postpartum complication. From Tirana City, postpartum complications were reported more frequently than from other areas of the country, rural or urban. This increase in frequency might be influenced by the presence of the University Hospital in Tirana (as the likelihood of being diagnosed with a pregnancy complication increases to diagnose them). Complications during pregnancy increase two times the odds for having a postpartum complication. Higher postpartum complications rates for women under age of 20 are likely to be caused by a two times higher rate of experiencing a painful uterus among these women compared to those women of older ages.

Breastfeeding

Infant feeding practices influence the health of both the child and the mother. Breast milk is the most complete food an infant can receive during the first few months of life. Early initiation of breastfeeding – within an hour following birth-permits the newborn to benefit immediately from colostrum, which is highly nutritious and contains antibodies necessary to protect babies before they are able to produce their own appropriate immune system. Early initiation also takes advantage of the newborn's sucking reflex alertness immediately during postpartum. In addition, breastfeeding has been shown to contribute to longer birth intervals, improve maternal health by reducing postpartum bleeding, allowing an earlier return to prepregnancy weight, and reduce the risk of premenopausal breast cancer.

In early infancy, frequent breastfeeding, including night feeds, is important to ensure that the infant receives sufficient breast milk and is able to increase its weight. Current recommendations are that infants should be breastfeed 8–10 times every 24 hours, and even more frequently during the first month of life. On the other hand, frequent feeding also ensures that the mother maintains her ability to produce sufficient quantities of breast milk.

Optimal breast feeding for infants includes:

- 1. Initiation of breastfeeding within one hour of birth
- 2. Frequent on demand feeding (including night feeds)

3. Exclusive breastfeeding (defined as breast milk only and no other foods or liquids) until the infant is about 6 months of age

The Albania RHS included questions about breastfeeding patterns and duration for all children under 5 years of age. Table 5.16 shows the percentage of children ever breastfed and percent distribution of initiation of breastfeeding by selected categories. The overwhelming majority (93%) of all babies born during the 1997-2002 period were breastfed for at least a short period of time. This result is consistent with other recent surveys in the country. The rate is also similar to ever breastfeeding rates reported by the RHS or DHS studies in the East European countries and the former USSR countries. The rate for Albania is slightly higher than that of Armenia and Georgia (89%, 87%), almost equal to that of Romania (93%), and to some extent lower than those of Central Asian countries (95–97%)

The percentage of babies ever breastfed was not significantly influenced by background characteristics. Only babies having a low birth weight were slightly less likely to be breastfed than those with a birth weight of 2500 grams or more (79% versus 94%). The same was true for babies born with a cesarean section compared to those having a vaginal birth (88% versus 94%).

In the same table (5.16), in the right panel, time elapsed between delivery and initiation of breastfeeding is included. Only 15% of infants ever breastfed started to do so within the first hour of life. Most children (52%) began breastfeeding between two hours after birth and completion of the first day of life. Less than 10% started breastfeeding the second day and the remaining 23% after the second day of life. There is a significant difference among those who started breastfeeding within the first hour of life between Tirana and other parts of the country (rural or urban). Women who give birth with a C-section are visibly inclined to start breastfeeding their babies late. To a lesser extent, the same is true for those giving birth to low birth weight babies.

An infant is "exclusively" breastfed when he or she receives only breast milk and "almost exclusively" breastfed or predominantly breastfed when he receives breast milk accompanied by water or other liquids (except other types of milk). Children who are exclusively or almost exclusively breastfed are considered to be "fully" breastfed (Labbok and Krasovec, 1990). These indicators are recommended by WHO to assess the adequacy of breastfeeding practices in a population and allow for comparisons with findings from other countries. The WHO recommendations state that "all infants should be fed exclusively on breast milk from birth to 4–6 months of age" and that some breastfeeding should be maintained until the child is at least 1 year old (WHO 1991).

The indicator used to estimate the duration of breastfeeding is the mean duration of three different types of breastfeeding; exclusive breastfeeding, full breastfeeding, and any breastfeeding. The results are presented in Table 5.17. The proportion of children under 5 years old still being breastfed at the time of the survey was calculated by the single months of age (0-59 months). Those proportions were summed together to calculate the mean duration of breastfeeding. This method is known as the "current status mean" method (WHO 1991). Duration of exclusive and full breastfeeding were calculated in the same way, where babies who did not yet initiate any other liquids or food were classified as exclusively breastfed and those who were either exclusively breastfed or started to take liquids but no other food were classified as fully breastfed.


The mean duration of any breastfeeding was 14.3 months and it does not seem to be highly influenced by background characteristics included in the analysis. Obviously the mean time duration of the two other indicators was much smaller, indicating that most of these 14 months infants were being fed only partially with their mother's milk. The mean duration of exclusive breastfeeding was 2.8 months and that of full breastfeeding was almost 5 months. The differences between selected categories remain small.

Infant and child mortality

Childhood mortality consists of deaths occurring among children from birth until the age of five. It can be broken down by an age classification: perinatal mortality, neonatal mortality, postneonatal mortality, infant mortality, child mortality (not to be confused with overall childhood mortality), and under five mortality. In this section perinatal mortality is not included. The age classifications used in this report are as follows:

- Neonatal mortality: the probability of dying between birth and less than 29 days
- Post neonatal mortality: the difference between infant and neonatal mortality rates
- Infant mortality: the probability of dying between birth and the exact age of one year
- Child mortality: the probability of dying between the exact ages of one and five years
- Under five mortality: the probability of dying between birth and the exact age of five years

Infant mortality is considered one of the crucial indicators for the health and social welfare of a country, as it synthesizes the quality of health care, nutrition, education and many other aspects of a society. Infant mortality for many years has been very high in Albania, ranked among the highest in the European Region. Despite a steady decrease from 45 per 1000 in 1990 to 16 per 1000 in 2000 in official reports, it remains far from the European Union standards. Official reports by the Ministry of Health of Albania demonstrate an infant mortality rate of 24.2/1,000 in 1995 and 16.0/1,000 in 2000. The official rates for 1995 and 2000 are presented by district in figure 5.3.




The survey questionnaire included a series of questions in the pregnancy history, which can help provided detailed information on infant and child mortality. For each live birth data are gathered on the date of the birth, sex of the child, survival status, and for children who have died, the age of when the death occurred. Respondents were asked to report pregnancy outcomes (e.g stillbirths and live births) according to international definitions. Accordingly, a live birth was defined as any infant, irrespective of the duration of the pregnancy, that breathes or shows any other signs of life after separation from the mother. Survey data on mortality levels among respondents' children were used to calculate the infant mortality (deaths before the first birthday per 1000) and child mortality (deaths between 12 and 59 completed months of age per 1000) rates. Infant mortality was further categorized into neonatal (0-28 days) and post-neonatal (29 days to 11 completed months) groups. Infant and child mortality rates were calculated by means of life table analysis.

Table 5.18 shows the survey estimates of infant and other childhood mortality rates distributed by selected categories. The infant mortality rate calculated for the

period of August 1992 – July 2002 is 26.2 per 1000 and the under-5 mortality rate (children under five years), for the same period, is 31.9 per 1000. The survey IMR of 26.2 per 1000 for the 10 year period from 1992 to 2002 is 29% higher than the average official rate reported between 1995 and 2000. The MICS 2000 survey carried out two years prior to the Albania RHS has produced similar results; infant mortality was 28 per 1000 and child mortality was 33 per 1000 (UNICEF, 2000).

For this 10-year period of time, the estimated neonatal mortality rate is 11.9 per 1000 and the post-neonatal mortality result is 14.3 per 1000. In this type of survey and in vital statistics systems, underestimation of neonatal mortality tends to be greater than underestimation of child mortality at older ages. Some women do not always consider their births to be live births, especially when the death occurred in the first days of life. For this reason, the estimated 10 year neonatal and infant mortality rates should be considered minimum values. The RHS recorded 55% of infant deaths in the post-neonatal period, a similar result when compared to the 60% reported by the MOH in 2000 (see Figure 5.4).



As shown in table 5.18, all indicators included in the table, with the exception of child mortality, neonatal mortality and post neonatal mortality are higher in rural areas. Accordingly, the infant mortality rate is around 40% higher in rural areas (30%) compared to urban areas. (21%). Only child mortality among 1-4 year olds seems quite similar in both areas. Another characteristic seemingly influencing the higher of the infant mortality rates, and to some extent under 5 mortality rates as well, is a low socioeconomic index. Nevertheless, one of the most strongly associated factors with infant mortality and under 5 mortality is education; among the lower education status group (not having finished secondary school), infant mortality is 2.3 times higher than in the more educated group. The difference is statistically significant. The ratio is almost as high when comparing child mortality among the two categories. Interestingly, the two later factors (socioeconomic index and education) do not seem to influence the risk of infants dying within the first month of life, a period of life, when other factors may be more important.

Infant mortality and post-neonatal mortality is higher for birth orders 3 or higher. Infants born after their mother had already two other children have higher rates. The infant mortality rate among them is almost 34 per 1000 while among the first born it is only 22.6 per 1000. The infant mortality rate estimate from the survey is 26.2/1000with a 95% confidence interval of 21.8 to 30.6. The most strongly associated factor with infant mortality and child mortality in Albania seems to be birth spacing; children born within 24 months after their mothers had a previous birth have a three to four times higher risk of dying, either within the first year, or within the first five years of life, compared to children born after more than two years of time between the two births. Despite the small numbers, the difference is statistically significant.

		R	eprod		ercent l e Health				2002				
	Trim	Trimester of First Prenatal Visit Number of Prenatal Visits											
Characteristics	No visits	1st	2nd	3rd	Don't Know⁺	0	1–3	4–6	7–9	10+	Don't Know	Total	No. of Cases*
Total	19.1	59.3	18.2	2.9	0.4	19.1	38.3	27.0	10.5	3.6	1.4	100.0	2,551
Strata													
Metro Tirana	10.9	71.1	15.2	2.3	0.5	10.9	27.4	31.7	18.0	9.9	2.1	100.0	900
Other Urban	9.9	70.3	17.0	2.4	0.4	9.9	28.6	38.1	16.9	5.0	1.4	100.0	807
Other Rural	26.0	50.6	19.7	3.3	0.4	26.0	46.1	20.0	5.3	1.3	1.3	100.0	844
Residence													
Urban	9.0	72.0	16.3	2.4	0.4	9.0	27.2	37.0	18.2	7.0	1.7	100.0	1,522
Rural	26.0	50.6	19.6	3.3	0.4	26.0	46.0	20.1	5.3	1.3	1.2	100.0	1,029
Age Group at Birth	1												
< 20	15.1	53.9	28.0	2.8	0.2	15.1	42.6	31.2	7.1	2.5	1.4	100.0	196
20–24	18.4	62.1	16.2	2.8	0.3	18.4	41.2	24.0	12.1	3.1	1.1	100.0	867
25–34	19.2	59.4	17.9	3.0	0.5	19.2	36.9	27.8	10.5	4.1	1.5	100.0	1,350
35–44	28.2	50.4	18.8	2.6	0.0	28.2	27.3	30.7	6.5	4.4	3.0	100.0	138
Education Level													
Primary or Less	25.0	50.9	21.0	2.7	0.4	25.0	43.6	21.1	6.9	1.9	1.4	100.0	1,228
Secondary Incomplete	10.7	68.6	16.1	4.7	0.0	10.7	29.3	38.3	13.6	4.9	3.3	100.0	179
Secondary Complete	12.3	69.8	14.2	3.2	0.5	12.3	33.8	33.0	14.0	5.6	1.3	100.0	880
Post-Secondary	7.4	75.4	14.9	2.1	0.2	7.4	22.8	38.9	22.3	8.0	0.6	100.0	264
Socioeconomic Ind	dex												
Low	26.4	49.4	20.1	3.5	0.6	26.4	43.7	21.3	4.8	1.9	1.7	100.0	1,020
Medium	13.0	66.4	18.0	2.4	0.3	13.0	36.3	30.7	14.1	4.8	1.1	100.0	1,222
High	10.4	78.5	8.7	2.2	0.2	10.4	17.3	39.5	24.2	7.2	1.4	100.0	309
Birth Order													
First	12.4	64.3	19.5	3.2	0.6	12.4	39.2	29.3	13.2	4.1	1.7	100.0	963
Second	19.0	61.6	16.7	2.4	0.4	19.0	38.1	27.7	9.7	4.3	1.2	100.0	964
Third or More	28.9	49.2	18.5	3.1	0.3	28.9	37.4	22.6	7.7	2.1	1.4	100.0	624
Baby Weight at Bir	th												
< 2500 Grams	18.6	55.4	21.1	2.3	2.5	18.6	36.6	26.7	9.1	6.6	2.3	100.0	123
≥ 2500 Grams	19.1	59.9	17.8	2.9	0.3	19.1	38.2	27.1	10.6	3.5	1.4	100.0	2,414
Don't Know	**	**	**	**	**	**	**	**	**	**	**	**	14

Table 5.1 Prenatal Care by Pregnancy Trimester of First Visitand Number of Prenatal Visits, for Births in 1997–2002, by Selected Characteristics (Percent Distribution) Reproductive Health Survey: Albania 2002

* Data are missing for 5 live births in the last 5 years

 ** Percentages are not shown when base is less than 25 cases

† Women who did attend prenatal care but did not know in which trimester they began prenatal care

Table 5.2

Percentage Counseled on Specific Topics During Prenatal Care Visits for Births in 1997–2002 by Selected Characteristics, among Women with Any Prenatal Care **Reproductive Health Survey: Albania 2002**

Characteristics	Nutrition	Breast Feeding	Delivery	Postnatal Care	Pregnancy Complications	Effects of Smoking	Effects of Alcohol	Contra- ception	No. of Cases
Total	60.0	58.7	58.2	52.8	50.8	53.3	53.3	37.9	2,152
Strata									
Metro Tirana	69.9	69.9	67.5	64.5	63.6	60.0	60.8	46.4	805
Other Urban	66.7	66.2	65.5	58.8	57.4	62.0	61.2	41.2	725
Other Rural	52.6	50.5	50.8	45.4	42.7	45.8	46.1	33.2	622
Residence									
Urban	68.6	68.4	67.0	61.7	60.3	62.4	62.1	44.0	1,394
Rural	52.7	50.5	50.8	45.3	42.8	45.6	45.9	32.8	758
Age Group at Birth									
< 20	56.5	56.3	58.9	49.8	49.1	49.2	48.4	33.7	168
20–24	61.8	60.0	57.4	54.3	51.9	55.4	55.7	38.6	733
25–34	58.2	57.1	58.0	51.6	49.7	51.5	51.3	36.9	1,141
35–44	70.2	68.7	64.6	59.5	56.5	63.5	65.1	49.6	110
Education Level									
Primary or Less	51.8	50.1	50.6	44.9	43.0	45.2	45.1	31.4	943
Secondary Incomplete	61.6	62.5	60.5	57.4	53.8	56.9	57.6	40.8	163
Secondary Complete	68.2	66.7	66.1	60.0	58.8	60.5	60.6	43.8	797
Post-Secondary	79.7	79.6	75.0	72.5	67.8	74.9	74.9	54.6	249
Socioeconomic Index									
Low	50.7	49.7	49.9	45.2	42.1	43.9	44.2	29.9	785
Medium	64.3	62.6	62.4	56.2	54.5	57.9	58.1	42.7	1,080
High	81.6	81.0	76.3	71.6	73.6	74.1	72.1	50.7	287
Birth Order									
First	63.6	62.6	61.8	57.6	53.5	57.5	57.3	39.1	858
Second	61.4	60.0	59.8	54.3	52.9	54.4	54.6	41.1	818
Third or More	51.5	49.7	49.6	42.1	42.8	44.2	44.4	31.0	476
Number of Prenatal Visits									
1–3	54.3	51.5	52.7	48.0	47.1	48.3	48.1	39.2	872
4–6	63.4	65.0	63.0	55.4	52.9	57.7	57.9	35.9	767
7–9	67.8	66.0	62.8	57.6	54.0	58.7	58.7	35.1	334
10+	71.7	66.0	65.4	61.9	59.6	59.6	60.1	44.3	137
Don t Know	60.4	60.1	65.5	71.6	64.6	48.7	49.8	45.8	42
Place of Prenatal Care									
District/Tirana Maternity/									
Hospital	55.0	54.1	53.7	50.2	47.9	50.3	50.3	36.8	682
Policlinic	64.7	64.1	64.2	57.6	56.3	57.1	57.4	38.9	841
Rural/Urban Health Center	66.0	64.9	62.9	60.1	57.7	59.0	58.9	48.0	249
Health Post	62.3	58.5	52.1	44.4	42.5	51.9	52.4	29.5	195
Private Office/Clinic/ Hospital	52.1	50.1	54.6	49.9	46.1	46.6	45.0	40.9	173
At Home/Other	**	**	**	**	**	**	**	**	12

** Percentages are not shown when base is less than 25 cases

Та	b	le	5.	3
		.	Υ.	-

Percentage Experiencing Selected Procedures During Prenatal Care Visits for Births in 1997–2002 by Selected Characteristics, among Women with Any Prenatal Care Reproductive Health Survey: Albania 2002

Characteristics	Blood Sample Taken	Urine Sample Taken	Blood Pressure Measured	Had Ultrasound Exam	Tetanus Inmunization	Iron Supplements	No. of Cases
Total	85.4	85.0	83.3	76.5	61.6	25.1	2,152
Strata							
Metro Tirana	91.6	92.0	88.2	88.4	48.2	32.4	805
Other Urban	92.7	93.3	89.1	82.4	67.1	35.9	725
Other Rural	78.9	77.7	78.2	69.1	62.3	16.0	622
Residence							
Urban	93.1	93.7	89.0	84.7	60.9	36.1	1,394
Rural	79.0	77.8	78.6	69.6	62.1	15.8	758
Age Group at Birth							
< 20	90.0	91.4	80.1	74.9	55.0	21.5	168
20–24	85.5	85.3	81.6	75.2	62.8	26.0	733
25–34	83.8	83.0	84.3	77.8	61.8	24.4	1,141
35–44	91.8	91.8	90.6	75.1	62.4	31.4	110
Education Level							
Primary or Less	80.4	80.5	79.8	71.1	59.9	16.3	943
Secondary Incomplete	87.7	81.2	80.6	81.0	68.0	35.1	163
Secondary Complete	89.9	90.0	87.2	80.5	61.7	32.9	797
Post-Secondary	98.5	98.5	93.8	92.2	66.3	43.0	249
Socioeconomic Index							
Low	79.7	79.7	80.1	65.4	58.2	11.5	785
Medium	88.5	87.8	84.9	83.2	65.1	31.9	1,080
High	96.2	96.6	90.3	93.8	58.4	54.1	287
Birth Order							
First	88.3	87.9	83.9	80.3	59.9	29.0	858
Second	87.9	87.3	83.8	74.5	64.6	24.5	818
Third or More	76.6	76.7	81.5	72.7	59.9	18.9	476
Number of Prenatal Visits							
1–3	77.3	77.0	78.0	67.6	60.8	15.3	872
4–6	91.6	90.9	85.7	81.8	62.8	29.8	767
7–9	96.7	96.3	92.3	90.1	59.9	40.6	334
10+	95.9	95.9	94.0	91.0	63.8	49.4	137
Don t Know	78.5	79.6	87.2	76.8	65.3	22.3	42
Place of Prenatal Care							
District/Tirana Maternity/Hospital	87.7	88.2	82.6	79.4	61.4	23.8	682
Policlinic	88.0	88.0	86.8	78.4	59.8	26.8	841
Rural/Urban Health Center	91.0	87.4	86.1	76.5	64.6	25.1	249
Health Post	66.2	65.1	75.3	54.1	70.4	10.4	195
Private Office/Clinic/Hospital	91.4	90.6	80.4	90.7	53.3	47.8	173
At Home/Other	**	**	**	**	**	**	12

 ** Percentages are not shown when base is less than 25 cases

Table 5.4

Percentage with Routine Measurement of Blood Pressure (BP) During Pregnancy, Reported High Blood Pressure (HBP) During Pregnancy, and Hospitalization Rate for HBP, for Births in 1997–2002, among Women with Any Prenatal Care Reproductive Health Survey: Albania 2002

Characteristics	Measurement of Blood Pressure	Told High Blood Pressure	Treatment HBP	Pregnancies Hospitalized for HBP (Exclusive)*	Pregnancies Hospitalized for HBP (Not Exclusive)*	No. of Cases
Total	83.3	11.4	51.0	0.5	2.3	2,152
Strata						
Metro Tirana	88.2	13.7	36.2	0.2	2.0	805
Other Urban	89.1	9.9	50.3	0.3	2.1	725
Other Rural	78.2	11.6	57.7	0.7	2.5	622
Residence						
Urban	89.0	10.9	54.3	0.3	2.1	1,394
Rural	78.6	11.8	44.3	0.7	2.4	758
Age Group at Birth						
< 20	80.1	11.9	48.2	0.0	3.0	168
20–24	81.6	7.9	42.7	0.1	1.8	733
25–34	84.3	12.9	52.3	0.8	1.9	1,141
35–44	90.6	17.6	67.6	1.4	7.8	110
Education Level						
Primary or Less	79.8	13.5	53.1	0.5	2.8	943
Secondary Incomplete	80.6	9.9	55.7	0.3	2.4	163
Secondary Complete	87.2	9.2	52.5	0.6	1.6	797
Post-Secondary	93.8	8.7	23.4	0.2	1.2	249
Socioeconomic Index						
Low	80.1	12.5	50.0	0.8	2.4	785
Medium	84.9	10.8	53.1	0.3	2.1	1,080
High	90.3	9.7	44.7	0.0	2.5	287
Birth Order						
First	83.9	11.7	47.5	0.5	2.4	858
Second	83.8	8.7	45.4	0.4	1.7	818
Third or More	81.5	15.0	61.1	0.6	3.0	476
Number of Prenatal Visits						
1–3	78.0	8.6	40.8	0.1	1.4	872
4–6	85.7	10.7	45.9	0.7	1.8	767
7–9	92.3	17.3	61.1	0.6	4.9	334
10+	94.0	24.5	71.9	3.2	5.6	137
Don t Know	87.2	9.8	87.6	1.0	6.5	42
Place of Prenatal Care						
District/Tirana Maternity/Hospital	82.6	12.9	63.7	0.9	3.4	682
Policlinic	86.8	9.5	45.2	0.2	1.9	841
Rural/Urban Health Center	86.1	7.9	44.7	0.2	1.5	249
Health Post	75.3	10.8	53.9	0.6	0.6	195
Private Office/Clinic/Hospital	80.4	14.7	26.7	0.0	2.4	173
At Home/Other	**	**	**	**	**	12

* Exclusive - hospitalized only for high blood pressure; non-exclusive - hospitalized with high blood pressure as only one of conditions. ** Percentages are not shown when base is less than 25 cases

	Had Ultrasound Exam		<u>Time c</u>	of First Ultrasound Exam (Percent Distric				
Characteristics	%	No. of Cases	< 14 weeks	14–26 weeks	27 +	Dk/Dr	Total	No. of Cases
Total	76.5	2,152	29.6	43.8	26.0	0.6	100.0	1,736
	70.5	2,132	29.0	43.0	20.0	0.0	100.0	1,730
Strata							100.0	= 1 0
Metro Tirana	88.4	805	35.5	34.6	29.0	0.9	100.0	712
Other Urban	82.4	725	30.6	47.1	21.7	0.6	100.0	599
Other Rural	69.1	622	26.5	45.1	28.0	0.4	100.0	425
Residence								
Urban	84.7	1,394	32.7	42.3	24.3	0.7	100.0	1,204
Rural	69.6	758	26.5	45.4	27.7	0.4	100.0	532
Age Group at Birth								
< 20	74.9	168	23.5	45.8	30.4	0.3	100.0	132
20–24	75.2	733	33.1	42.9	23.5	0.4	100.0	586
25–34	77.8	1,141	28.9	43.6	26.7	0.8	100.0	928
35–44	75.1	110	24.2	48.7	27.1	0.0	100.0	90
Education Level								
Primary or Less	71.1	943	25.6	47.0	26.9	0.4	100.0	698
Secondary Incomplete	81.0	163	26.2	50.2	22.1	1.6	100.0	136
Secondary Complete	80.5	797	34.4	39.1	25.9	0.6	100.0	667
Post-Secondary	92.2	249	36.2	38.6	24.7	0.5	100.0	235
Socioeconomic Index								
Low	65.4	785	23.6	45.8	30.2	0.4	100.0	544
Medium	83.2	1,080	31.1	44.2	24.0	0.7	100.0	923
High	93.8	287	43.2	35.0	21.4	0.5	100.0	269
Birth Order								
First	80.3	858	31.2	41.5	27.1	0.2	100.0	719
Second	74.5	818	26.3	46.9	25.8	1.0	100.0	657
Third or More	72.7	476	31.6	43.7	24.1	0.7	100.0	360
Number of Prenatal Visits								
1–3	67.6	872	22.1	48.6	29.1	0.2	100.0	615
4–6	81.8	767	29.7	47.8	22.2	0.4	100.0	651
7–9	90.1	334	45.0	30.2	23.3	1.6	100.0	307
10+	91.0	137	43.5	22.9	31.9	1.8	100.0	128
Don t Know	76.8	42	33.1	31.0	34.5	1.4	100.0	35
Place of Prenatal Care								
District/Tirana Maternity/Hospital	79.4	682	34.3	43.1	22.4	0.2	100.0	567
Policlinic	78.4	841	24.4	47.0	27.7	0.9	100.0	693
Rural/Urban Health Center	76.5	249	29.0	44.7	26.4	0.0	100.0	196
Health Post	54.1	195	18.1	52.0	28.2	1.7	100.0	109
Private Office/Clinic/Hospital	90.7	173	42.7	28.9	28.1	0.3	100.0	164
At Home/Other	**	**	**	**	**	**	**	7

Table 5.5Use of Ultrasound Exams During Pregnancy and Time of First Ultrasound Exam, BySelected Characteristics, for Births in 1997–2002, among Women with Any Prenatal Care

** Percentages are not shown when base is less than 25 cases

ables

Table 5.6

Percentage of Pregnancy Complications that Required Medical Attention, by Selected Characteristics, for Births in 1997–2002, among Women with Any Prenatal Care **Reproductive Health Survey: Albania 2002**

Characteristics	Any Pregnancy Complication	Risk of Preterm Delivery	Anemia Related to Preg.	Weak Cervix	Water Retention or Edema	Urinary Tract Infection	High BP Related to Preg.	Bleeding During First 6 Mth	Bleeding at 6 Mth or More	Rh Isoimmu- nization	Other	No. of Cases
Total	24.2	9.9	7.1	6.4	6.3	6.1	5.3	4.2	1.5	1.5	1.8	2,152
Strata												
Metro Tirana	22.1	7.8	6.9	6.3	5.6	6.4	5.2	5.4	1.0	1.8	2.2	805
Other Urban	22.4	8.6	7.9	5.6	4.8	5.2	4.7	4.3	1.5	0.6	2.3	725
Other Rural	26.0	11.3	6.7	6.9	7.4	6.5	5.7	3.7	1.6	1.9	1.4	622
Residence												
Urban	22.6	8.3	7.6	5.9	5.1	5.6	4.9	4.8	1.3	1.1	2.3	1,394
Rural	25.5	11.1	6.6	6.8	7.3	6.4	5.6	3.7	1.6	1.8	1.4	758
Age Group at Bi	rth											
< 20	24.0	11.0	10.8	6.9	7.2	8.5	5.1	3.2	2.8	0.8	2.6	168
20–24	24.8	10.4	7.1	7.8	5.9	6.4	3.5	4.8	0.5	1.5	1.9	733
25–34	23.9	9.2	6.3	5.3	5.9	5.0	5.9	3.9	1.6	1.7	1.6	1,141
35–44	23.6	10.6	7.9	7.1	10.8	9.1	10.8	4.5	4.0	0.0	2.2	110
Education Level												
Primary or Less	23.9	9.6	6.9	7.4	6.1	6.2	6.1	3.8	1.7	1.8	1.2	943
Secondary Incomplete	28.0	11.3	7.4	3.5	12.4	6.1	6.1	2.9	1.6	0.3	3.5	163
Secondary Complete	23.6	10.2	7.1	5.0	6.0	6.2	4.5	4.8	1.4	1.3	1.9	797
Post-Secondary	24.8	8.6	8.5	8.2	3.7	4.7	1.9	5.8	0.5	0.9	4.8	249
Socioeconomic	Index											
Low	24.2	8.9	7.7	6.4	7.3	5.4	5.2	3.5	1.6	1.6	1.8	785
Medium	23.7	10.2	5.8	5.8	5.7	6.4	5.4	3.8	1.5	1.3	1.8	1,080
High	26.7	12.7	11.4	9.6	5.0	7.5	4.9	9.6	0.2	1.6	2.1	287
Birth Order												
First	26.6	12.6	8.3	7.4	6.4	6.9	5.3	6.0	0.8	1.8	1.5	858
Second	22.1	7.3	6.1	5.9	5.8	5.3	3.8	3.0	1.2	1.7	2.5	818
Third or More	23.0	8.8	6.4	5.5	6.9	5.7	7.5	2.8	3.0	0.6	1.6	476

Table 5.7Place of Delivery for Births in 1996–2001 by Selected Characteristics
(Percent Distribution)
Reproductive Health Survey: Albania 2002

			Place o	of Delivery			
Characteristics	District Maternity Hospital	Tirana Maternity	Birth House/ Health Center	Private Clinic/ Hospital	At Home	Total	No. of Cases
Total	70.9	14.3	8.2	0.6	6.0	100.0	2,551
Strata							
Metro Tirana	16.6	78.0	2.0	0.9	2.5	100.0	900
Other Urban	92.8	2.9	2.2	1.3	0.8	100.0	807
Other Rural	73.8	3.6	12.9	0.2	9.6	100.0	844
Residence							
Urban	70.2	25.6	2.0	1.2	1.0	100.0	1,522
Rural	71.3	6.6	12.5	0.2	9.4	100.0	1,029
Age Group at Birth							
< 20	73.5	15.3	4.5	0.0	6.8	100.0	196
20–24	72.2	14.2	8.2	0.7	4.8	100.0	867
25–34	70.2	14.3	8.7	0.7	6.1	100.0	1,350
35–44	64.9	13.8	10.1	0.3	11.0	100.0	138
Education Level							
Primary or Less	70.3	10.1	11.0	0.4	8.3	100.0	1,228
Secondary Incomplete	74.5	16.0	5.6	0.0	3.9	100.0	179
Secondary Complete	70.9	19.8	4.9	1.1	3.2	100.0	880
Post-Secondary	71.9	24.4	1.9	1.3	0.4	100.0	264
Socioeconomic Index							
Low	69.7	8.9	10.6	0.2	10.6	100.0	1,020
Medium	74.4	16.0	6.7	0.7	2.1	100.0	1,222
High	56.9	37.3	2.5	2.5	0.8	100.0	309
Birth Order							
First	73.8	17.8	5.4	0.7	2.4	100.0	963
Second	68.5	14.7	9.4	0.9	6.6	100.0	964
Third or More	69.8	8.8	10.8	0.2	10.4	100.0	624
Trimester of First Prenata	al Visit						
No Visits	65.6	6.8	12.2	0.3	15.1	100.0	399
1st	70.6	18.0	7.1	0.8	3.5	100.0	1,634
2nd	75.3	10.7	8.4	0.6	5.1	100.0	438
3rd	85.1	10.0	2.4	0.0	2.6	100.0	69
Don't Know	**	**	**	**	**	**	11

** Percentages are not shown when base is less than 25 cases

Table 5.8
Average Time Between Admission and Delivery and Percent
Distribution of Number of Nights Spent in a Medical Facility,
for Births in 1997–2002
Reproductive Health Survey: Albania 2002

	Average Time Between Admission		een De	livery		Facility charge n)		
	and Delivery					8 or		No. of
Characteristics	(Hours)	0–2	3–4	5	6–7	More	Total	Cases
Total	7.2	59.0	21.8	4.0	7.0	8.2	100.0	2,444
Strata								
Metro Tirana	6.8	60.4	19.2	6.0	6.9	7.5	100.0	876
Other Urban	6.0	56.9	22.0	4.8	8.5	7.8	100.0	800
Other Rural	8.0	59.8	22.4	3.0	6.2	8.6	100.0	768
Residence								
Urban	6.2	57.3	21.7	5.3	8.2	7.5	100.0	1,503
Rural	7.9	60.3	21.8	3.1	6.2	8.6	100.0	941
Age Group at Birth								
< 20	9.8	57.8	24.4	3.4	8.6	5.8	100.0	185
20–24	7.9	59.4	20.9	3.7	7.6	8.4	100.0	835
25–34	6.3	59.7	21.8	4.2	6.2	8.1	100.0	1,292
35–44	7.2	52.5	23.4	5.1	8.0	11.0	100.0	132
Education Level								
Primary or Less	7.7	58.8	23.1	3.2	7.3	7.6	100.0	1,150
Secondary Incomplete	5.5	59.2	20.6	7.3	4.2	8.8	100.0	174
Secondary Complete	7.0	58.9	20.2	4.7	6.8	9.4	100.0	858
Post-Secondary	5.4	61.3	19.1	4.3	8.8	6.4	100.0	262
Socioeconomic Index								
Low	7.9	59.6	21.0	3.7	7.7	8.0	100.0	939
Medium	6.7	59.4	22.6	3.5	5.8	8.7	100.0	1,199
High	5.9	53.3	21.3	8.6	10.5	6.3	100.0	306
Birth Order								
First	9.9	53.5	20.9	4.2	10.8	10.7	100.0	948
Second	5.3	60.7	24.3	4.2	4.9	6.0	100.0	919
Third or More	5.7	65.3	19.8	3.6	4.1	7.2	100.0	577
Baby Weight at Birth								
< 2500 Grams	12.6	21.6	16.2	3.7	15.8	42.7	100.0	116
≥ 2500 Grams	6.9	60.8	22.1	4.0	6.6	6.5	100.0	2,320
Dk/Dr	**	**	**	**	**	**	**	8
Type of Delivery								
Vaginal	6.3	67.2	22.7	2.8	4.0	3.2	100.0	2,075
Cesarean Section	13.4	5.9	15.8	11.8	26.4	40.1	100.0	369
Pregnancy Complications								
Yes	9.6	45.2	24.2	6.3	11.7	12.6	100.0	495
No	6.6	62.7	21.1	3.4	5.8	7.0	100.0	1,949
** Percentages are not shown when base	se is less than 25 cases							

A 1
6
Ch
Chi
Chil
Child
Child
Child
Child
Child H
Child H
hild He
Child He
hild He
hild He
hild He
hild He
hild He
hild Healt

Table 5.9
Percentage of Cesarean Deliveries by Selected
Characteristics, for Births in 1997–2002,
among Women Currently Aged 15–44
Reproductive Health Survey: Albania 2002

Characteristics	Cesarean Deliveries	No. of Cases
Total	13.4	2,444
Strata		
Metro Tirana	19.2	876
Other Urban	14.4	800
Other Rural	11.2	768
Residence		
Urban	16.0	1,503
Rural	11.4	941
Age Group at Birth		
< 20	8.6	185
20–24	11.7	835
25-34	14.0	1,292
35–44	25.3	132
Education Level		
Primary or Less	12.0	1,150
Secondary Incomplete	12.1	174
Secondary Complete	15.3	858
Post-Secondary	16.7	262
Socioeconomic Index		
Low	11.0	939
Medium	14.0	1,199
High	22.7	306
Birth Order		
First	16.4	948
Second Third or More	12.2 10.1	919 577
	10.1	577
Pregnancy Complications	10.0	
Yes	19.0	495
No	11.9	1,949
Baby Weight at Birth		
< 2500 Grams	36.0	116
≥ 2500 Grams	12.3 **	2,320
Don't know		8
Prolonged Labor		
No	4.4	2,094
Yes	21.4	65
No Labor Don't know	98.9 16.8	233 52
DOILEKIIOW	10.0	52

** Percentages are not shown when base is less than 25 cases

Table 5.10

Percentage of Poor Birth Outcomes by Selected Characteristics, for Births in 1997–2002, Among Women Currently Aged 15–44 Reproductive Health Survey: Albania 2002

Characteristics	Low Birth Weight < 2,500 grams	Preterm Birth	No. of Cases	
Total	4.6	3.6	2,551	
Strata				
Metro Tirana	5.3	4.3	900	
Other Urban	4.1	3.6	807	
Other Rural	4.7	3.4	844	
Residence				
Urban	4.4	4.0	1,522	
Rural	4.7	3.3	1,029	
Age Group at Birth				
< 20	5.7	2.5	196	
20–24	5.6	3.9	867	
25–34	4.1	3.7	1,350	
35–44	1.5	2.5	138	
Education Level				
Primary or Less	5.4	3.1	1,228	
Secondary Incomplete	3.8	1.5	179	
Secondary Complete	3.2	4.7	880	
Post-Secondary	5.0	4.6	264	
Socioeconomic Index				
Low	5.5	3.3	1,020	
Medium	3.4	3.5	1,222	
High	6.6	5.8	309	
Prenatal Care Visits				
Yes	4.6	3.7	2,152	
No	4.5	2.9	399	
Birth Order				
First	6.7	5.1	963	
Second	4.0	2.8	964	
Third or More	2.4	2.3	624	
Pregnancy Complications				
Yes	8.9	5.5	498	
No	3.5	3.1	2,053	
Prolonged Labor				
No	3.9	3.5	2,195	
Yes	9.5	5.8	70	
No Labor	9.0	3.8	233	
Don't know	10.5	0.8	53	

Table 5.11

Percentage Who Attended Postpartum Care Visit within Specified Time Periods by Selected Characteristics, for Births in 1997–2002, among Women Currently Aged 15–44 Reproductive Health Survey: Albania 2002

	Postpartum Care				Time betwen Delivery and First Postpartum Visit (Percent Distribution of Weeks)						
Characteristics	%	No. of Cases	< 1	1–2	> 2	Don't Remember	Total	No. of Cases			
Total	18.7	2,551	36.0	24.9	34.3	4.8	100.0	549			
Strata											
Metro Tirana	26.9	900	25.2	35.7	36.8	2.3	100.0	243			
Other Urban	21.9	807	33.3	27.1	36.5	3.1	100.0	179			
Other Rural	14.8	844	43.1	18.2	31.5	7.2	100.0	127			
Residence											
Urban	23.8	1,522	30.0	29.5	37.9	2.7	100.0	384			
Rural	15.1	1,029	42.5	20.0	30.5	7.1	100.0	165			
Age Group at Birth											
< 20	15.7	196	45.5	26.8	27.7	0.0	100.0	39			
20–24	18.7	867	38.9	19.3	40.6	1.2	100.0	174			
25–34	19.2	1,350	31.4	29.7	31.2	7.6	100.0	303			
35–44	18.3	138	46.4	13.8	32.2	7.5	100.0	33			
Education Level											
Primary or Less	16.0	1,228	44.2	19.0	34.0	2.8	100.0	215			
Secondary Incomplete	21.2	179	41.8	30.0	28.2	0.0	100.0	45			
Secondary Complete	20.6	880	27.7	29.3	34.1	8.8	100.0	203			
Post-Secondary	30.2	264	20.5	34.4	40.0	5.0	100.0	86			
Socioeconomic Index											
Low	13.9	1,020	39.7	27.2	28.5	4.6	100.0	158			
Medium	21.3	1,222	37.8	23.3	34.2	4.7	100.0	292			
High	31.8	309	18.9	25.1	50.3	5.7	100.0	99			
Birth Order											
First	20.5	963	34.9	22.9	36.2	6.0	100.0	230			
Second	18.6	964	36.6	25.2	35.2	3.1	100.0	205			
Third or More	16.2	624	37.1	28.1	29.5	5.2	100.0	114			
Place of Delivery											
District Maternity Hospital	17.8	1,527	40.0	22.2	34.7	3.0	100.0	294			
Tirana Maternity	30.2	744	24.8	31.9	33.5	9.7	100.0	216			
Private Clinic/Hospital	**	**	**	**	**	**	**	9			
Birth House/Health Center	9.7	154	38.6	26.3	23.7	11.4	100.0	19			
At Home	8.9	107	**	**	**	**	**	11			
Postpartum Complications											
Any Complications	26.8	676	27.5	27.7	41.9	2.9	100.0	214			
No Complications	16.0	1,875	40.7	23.4	30.1	5.9	100.0	335			

 ** Percentages are not shown when base is less than 25 cases

lables

Total 76.1 74.7 72.9 72.8 71.9 55.0 549 Strata Metro Tirana 82.6 79.1 80.2 76.4 78.0 60.1 243 Other Urban 79.7 79.6 80.2 77.5 78.6 60.6 179 Other Rural 70.4 68.9 64.0 67.5 63.9 48.3 127 Residence Urban 82.0 81.2 81.5 78.8 80.0 62.9 384 Rural 69.8 67.8 63.7 66.3 63.1 46.5 165 Age Group at Birth 20 81.1 72.6 76.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 74.1 65.8 33 Education Level Primary or Less 67.3 65.1 65.5 63.3 64.4 47.4	Characteristics	Inmunization	Child Care	Nutrition	Breastfeeding	Breast Care	Contraception	No. of Cases
Metro Tirana 82.6 79.1 80.2 76.4 78.0 60.1 243 Other Urban 79.7 79.6 80.2 77.5 78.6 60.6 179 Other Rural 70.4 68.9 64.0 67.5 63.9 48.3 127 Residence Urban 82.0 81.2 81.5 78.8 80.0 62.9 384 Rural 69.8 67.8 63.7 66.3 146.5 165 Age Group at Birth - - - - 25.4 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.6 71.4 75.1 71.4 53.6 303 35 36 303 35 36 303 35 544 47.4 215 Secondary Incomplete 81.2 81.2 81.2 78.8 78.8 69.9 45.1 252 39 36 36.4 86.9 <t< th=""><th></th><th></th><th></th><th></th><th></th><th></th><th>-</th><th></th></t<>							-	
Metro Tirana 82.6 79.1 80.2 76.4 78.0 60.1 243 Other Urban 79.7 79.6 80.2 77.5 78.6 60.6 179 Other Rural 70.4 68.9 64.0 67.5 63.9 48.3 127 Residence Urban 82.0 81.2 81.5 78.8 80.0 62.9 384 Rural 69.8 67.8 63.7 66.3 63.1 46.5 165 Age Group at Birth - - - - 25.4 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.6 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 75.2 71.4 75.7 74.1 65.8 33 Education Level - - - - - 174 255 Secondary Incomplete 81.2	Strata							
Other Urban 79.7 79.6 80.2 77.5 78.6 60.6 179 Other Rural 70.4 68.9 64.0 67.5 63.9 48.3 127 Residence 01man 82.0 81.2 81.5 78.8 80.0 62.9 384 Rural 69.8 67.8 63.7 66.3 63.1 46.5 165 Age Group at Birth 72.6 76.7 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 33 Education Level		82.6	79.1	80.2	76.4	78.0	60.1	243
Residence Urban 82.0 81.2 81.5 78.8 80.0 62.9 384 Rural 69.8 67.8 63.7 66.3 63.1 46.5 165 Age Group at Birth 52.4 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 303 Education Level 75.4 75.2 71.4 75.7 74.1 65.8 303 Secondary lncomplete 81.2 81.2 81.2 83.8 64.4 74.4 215 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 8								
Urban 82.0 81.2 81.5 78.8 80.0 62.9 384 Rural 69.8 67.8 63.7 66.3 63.1 46.5 165 Age Group at Birth 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 333 Education Level 30.4 47.4 215 Secondary Incomplete 81.2 81.2 81.2 78.8 78.8 69.9 45 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 88.7 68.2 99 45	Other Rural	70.4	68.9	64.0	67.5	63.9	48.3	127
Urban 82.0 81.2 81.5 78.8 80.0 62.9 384 Rural 69.8 67.8 63.7 66.3 63.1 46.5 165 Age Group at Birth 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 333 Education Level 30.4 47.4 215 Secondary Incomplete 81.2 81.2 81.2 78.8 78.8 69.9 45 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 88.7 68.2 99 45	Residence							
Rural 69.8 67.8 63.7 66.3 63.1 46.5 165 Age Group at Birth 72.0 81.1 72.6 76.9 68.1 70.2 52.4 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 74.1 65.8 33 Education Level 81.2 81.2 81.2 78.8 69.9 45.5 Secondary Incomplete 81.2 81.2 81.2 78.8 78.8 69.9 45.5 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Secondary Complete 84.7 85.6 86.2 88.7 86.6 82.2 89.6 292 Host 83.2 80.4 81.2 79.2 80.4 55.2		82.0	81.2	81.5	78.8	80.0	62.9	384
< 20 81.1 72.6 76.9 68.1 70.2 52.4 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 33 Education Level 71.4 72.7 74.1 65.8 33 Feducation Level 81.2 81.2 81.2 78.8 78.8 69.9 45 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 88.7 66.4 86 Socioeconomic Index 104.7 65.9 65.1 64.7 65.2 49.6 158 Medium 82.1 79.3 76.2 76.7 74.3 58.6 292 <	Rural			63.7	66.3	63.1		
< 20 81.1 72.6 76.9 68.1 70.2 52.4 39 20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 33 Education Level 71.4 72.7 74.1 65.8 33 Feducation Level 81.2 81.2 81.2 78.8 78.8 69.9 45 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 88.7 66.4 86 Socioeconomic Index 104.7 65.9 65.1 64.7 65.2 49.6 158 Medium 82.1 79.3 76.2 76.7 74.3 58.6 292 <	Age Group at Birth							
20-24 76.2 74.5 74.4 70.5 72.6 55.7 174 25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 33 Education Level </td <td>• .</td> <td>81.1</td> <td>72.6</td> <td>76.9</td> <td>68.1</td> <td>70.2</td> <td>52.4</td> <td>39</td>	• .	81.1	72.6	76.9	68.1	70.2	52.4	39
25-34 75.4 75.2 71.4 75.1 71.4 53.6 303 35-44 75.6 74.1 72.7 72.7 74.1 65.8 33 Education Level 53.6 33 Secondary Incomplete 81.2 81.2 81.2 78.8 78.8 69.9 45 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 88.7 66.4 86 Socioeconomic Index 79.3 76.2 76.7 74.3 58.6 292 High 83.2 80.4 81.2 79.2 80.4 55.2 99 Birth Order 75.3 74.8 75.7 <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<>								
Education Level Primary or Less 67.3 65.1 65.5 63.3 64.4 47.4 215 Secondary Incomplete 81.2 81.2 81.2 78.8 78.8 69.9 45 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.6 86.2 88.7 66.9 45 Socioeconomic Index U U 1 75.7 76.7 41.8 76.7 74.3 58.6 292 High 83.2 80.4 81.2 79.2 80.4 55.2 99 Birth Order U U 1 76.7 74.3 54.6 205 Second 77.2 76.7 75.3 74.8 75.7 60.1 205 Birth Order U U 114 11.7 71.7 72.5 70.5 71.0 53.4 294 Tirana Maternity 83.8 82.5 74		75.4		71.4	75.1	71.4	53.6	303
Primary or Less67.365.165.563.364.447.4215Secondary Incomplete81.281.281.278.878.869.945Secondary Complete84.782.876.781.175.759.1203Post-Secondary86.488.788.686.288.766.486Socioeconomic IndexLow64.765.965.164.765.249.6158Medium82.179.376.276.774.358.6292High83.280.481.279.280.455.299Birth OrderFirst81.979.375.176.673.354.2230Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital************9Birth House/Health Center84.898.185.296.298.166.519At Home**************11Postpartum ComplicationsAny Complications74.873.375.1	35–44	75.6	74.1	72.7	72.7	74.1	65.8	33
Secondary Incomplete 81.2 81.2 81.2 81.2 78.8 78.8 69.9 45 Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 88.7 66.4 86 Socioeconomic Index U U 1 75.7 74.3 58.6 292 High 83.2 80.4 81.2 79.2 80.4 55.2 99 Birth Order E E E E E 203 203 Second 77.2 76.7 74.3 58.6 292 99 99 9 Birth Order E E 81.9 79.3 75.1 76.6 73.3 54.2 230 230 Second 77.2 76.7 75.3 74.8 75.7 60.1 205 Third or More 64.0 63.6 65.4 62.9 63.6	Education Level							
Secondary Complete 84.7 82.8 76.7 81.1 75.7 59.1 203 Post-Secondary 86.4 88.7 88.6 86.2 88.7 66.4 86 Socioeconomic Index U U U 158 64.7 65.2 49.6 158 Medium 82.1 79.3 76.2 76.7 74.3 58.6 292 High 83.2 80.4 81.2 79.2 80.4 55.2 99 Birth Order E <td>Primary or Less</td> <td>67.3</td> <td>65.1</td> <td>65.5</td> <td>63.3</td> <td>64.4</td> <td>47.4</td> <td>215</td>	Primary or Less	67.3	65.1	65.5	63.3	64.4	47.4	215
Post-Secondary86.488.788.686.288.766.486Socioeconomic IndexLow64.765.965.164.765.249.6158Medium82.179.376.276.774.358.6292High83.280.481.279.280.455.299Birth OrderFirst81.979.375.176.673.354.2230Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital**********9Birth House/Health Center84.898.185.296.298.166.519At Home************11Postpartum Complications74.873.375.169.670.952.8214	•	81.2	81.2	81.2	78.8	78.8	69.9	45
Socioeconomic Index Low 64.7 65.9 65.1 64.7 65.2 49.6 158 Medium 82.1 79.3 76.2 76.7 74.3 58.6 292 High 83.2 80.4 81.2 79.2 80.4 55.2 99 Birth Order E E Second 77.2 76.7 74.3 54.2 230 Second 77.2 76.7 75.3 74.8 75.7 60.1 205 Second 77.2 76.7 75.3 74.8 75.7 60.1 205 Third or More 64.0 63.6 65.4 62.9 63.6 48.9 114 Place of Delivery E E E E E 9 District Maternity Hospital 74.1 71.7 72.5 70.5 71.0 53.4 294 Tirana Maternity 83.8 82.5 74.8 78.2 72.3 57.5 216 <td< td=""><td>Secondary Complete</td><td>84.7</td><td>82.8</td><td>76.7</td><td>81.1</td><td>75.7</td><td>59.1</td><td>203</td></td<>	Secondary Complete	84.7	82.8	76.7	81.1	75.7	59.1	203
Low64.765.965.164.765.249.6158Medium82.179.376.276.774.358.6292High83.280.481.279.280.455.299Birth OrderFirst81.979.375.176.673.354.2230Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital************9Birth House/Health Center84.898.185.296.298.166.519At Home************11Postpartum Complications74.873.375.169.670.952.8214	Post-Secondary	86.4	88.7	88.6	86.2	88.7	66.4	86
Medium82.179.376.276.774.358.6292High83.280.481.279.280.455.299Birth OrderFirst81.979.375.176.673.354.2230Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital************9Birth House/Health Center84.898.185.296.298.166.519At Home**************11Postpartum Complications74.873.375.169.670.952.8214	Socioeconomic Index							
High83.280.481.279.280.455.299Birth OrderFirst81.979.375.176.673.354.2230Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital**********9Birth House/Health Center84.898.185.296.298.166.519At Home************11Postpartum Complications74.873.375.169.670.952.8214	Low	64.7	65.9	65.1	64.7	65.2	49.6	158
Birth OrderFirst81.979.375.176.673.354.2230Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital**********9Birth House/Health Center84.898.185.296.298.166.519At Home************11Postpartum Complications74.873.375.169.670.952.8214	Medium	82.1	79.3	76.2	76.7	74.3	58.6	292
First81.979.375.176.673.354.2230Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital**********9Birth House/Health Center84.898.185.296.298.166.519At Home**********11Postpartum Complications74.873.375.169.670.952.8214	High	83.2	80.4	81.2	79.2	80.4	55.2	99
Second77.276.775.374.875.760.1205Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital**********9Birth House/Health Center84.898.185.296.298.166.519At Home**********11Postpartum Complications74.873.375.169.670.952.8214	Birth Order							
Third or More64.063.665.462.963.648.9114Place of DeliveryDistrict Maternity Hospital74.171.772.570.571.053.4294Tirana Maternity83.882.574.878.272.357.5216Private Clinic/Hospital**********9Birth House/Health Center84.898.185.296.298.166.519At Home************11Postpartum Complications74.873.375.169.670.952.8214	First	81.9	79.3	75.1	76.6	73.3	54.2	230
Place of Delivery District Maternity Hospital 74.1 71.7 72.5 70.5 71.0 53.4 294 Tirana Maternity 83.8 82.5 74.8 78.2 72.3 57.5 216 Private Clinic/Hospital ** ** ** ** ** 9 Birth House/Health Center 84.8 98.1 85.2 96.2 98.1 66.5 19 At Home ** ** ** ** ** ** 11 Postpartum Complications Any Complications 74.8 73.3 75.1 69.6 70.9 52.8 214	Second	77.2	76.7	75.3	74.8	75.7	60.1	205
District Maternity Hospital 74.1 71.7 72.5 70.5 71.0 53.4 294 Tirana Maternity 83.8 82.5 74.8 78.2 72.3 57.5 216 Private Clinic/Hospital ** ** ** ** ** ** 9 Birth House/Health Center 84.8 98.1 85.2 96.2 98.1 66.5 19 At Home ** ** ** ** ** ** 11 Postpartum Complications 74.8 73.3 75.1 69.6 70.9 52.8 214	Third or More	64.0	63.6	65.4	62.9	63.6	48.9	114
Tirana Maternity 83.8 82.5 74.8 78.2 72.3 57.5 216 Private Clinic/Hospital ** ** ** ** ** ** 9 Birth House/Health Center 84.8 98.1 85.2 96.2 98.1 66.5 19 At Home ** ** ** ** ** ** 11 Postpartum Complications 74.8 73.3 75.1 69.6 70.9 52.8 214	Place of Delivery							
Private Clinic/Hospital ** ** ** ** ** ** 9 Birth House/Health Center 84.8 98.1 85.2 96.2 98.1 66.5 19 At Home ** ** ** ** ** 11 Postpartum Complications 74.8 73.3 75.1 69.6 70.9 52.8 214	District Maternity Hospital	74.1	71.7	72.5	70.5	71.0	53.4	294
Birth House/Health Center 84.8 98.1 85.2 96.2 98.1 66.5 19 At Home ** ** ** ** ** ** 11 Postpartum Complications 74.8 73.3 75.1 69.6 70.9 52.8 214	Tirana Maternity	83.8	82.5	74.8	78.2	72.3	57.5	216
At Home ** ** ** ** ** 11 Postpartum Complications	Private Clinic/Hospital	**	**	**	**	**	**	9
Postpartum Complications 74.8 73.3 75.1 69.6 70.9 52.8 214	Birth House/Health Center	84.8	98.1	85.2	96.2	98.1	66.5	19
Any Complications 74.8 73.3 75.1 69.6 70.9 52.8 214	At Home	**	**	**	**	**	**	11
	Postpartum Complications	i						
No Complications 76.9 75.5 71.7 74.6 72.4 56.2 335	Any Complications	74.8	73.3	75.1	69.6	70.9	52.8	214
	No Complications	76.9	75.5	71.7	74.6	72.4	56.2	335

** Percentages are not shown when base is less than 25 cases

		tal Baby <u>c Visit</u>	y Sick or Well Time Between Delivery and First Postnatal Baby Clinic Visit Baby Clinic Visit (Percent Distribution of Me							
Characteristics	%	No. of Cases	Health Exam for Sickness	Routine Health Exam	< 1	1–2	> 2	Don't Remember	Total	No. of Cases
Total	85.8	2551	8.4	91.6	45.0	32.7	20.8	1.5	100.0	2,245
Strata										
Metro Tirana	91.4	900	4.9	95.1	42.7	38.2	18.3	0.8	100.0	823
Other Urban	88.5	807	8.0	92.0	44.8	31.0	23.0	1.2	100.0	720
Other Rural	83.0	844	9.6	90.4	45.8	32.0	20.4	1.9	100.0	702
Residence										
Urban	89.9	1522	7.2	92.8	44.1	33.5	21.5	1.0	100.0	1,390
Rural	83.0	1029	9.3	90.7	45.6	32.1	20.4	1.9	100.0	855
Age Group at Birth										
< 20	86.5	196	14.0	86.0	43.5	26.0	29.0	1.5	100.0	170
20–24	83.3	867	8.3	91.7	41.2	34.1	23.9	0.9	100.0	743
25–34	86.5	1350	7.4	92.6	47.3	32.1	18.5	2.1	100.0	1,200
35–44	94.1	138	8.3	91.7	48.5	38.6	12.6	0.3	100.0	132
Education Level										
Primary or Less	82.8	1228	9.4	90.6	42.9	33.6	22.0	1.5	100.0	1,032
Secondary Incomplete	85.7	179	8.1	91.9	52.8	30.9	12.1	4.2	100.0	159
Secondary Complete	90.1	880	6.4	93.6	45.7	32.1	21.0	1.2	100.0	807
Post-Secondary	92.8	264	8.9	91.1	50.7	29.2	19.6	0.5	100.0	247
Socioeconomic Index										
Low	82.8	1020	10.0	90.0	45.0	32.7	20.4	1.9	100.0	853
Medium	88.5	1222	7.5	92.5	43.8	33.0	21.8	1.4	100.0	1,107
High	88.8	309	4.2	95.8	52.3	30.2	17.4		100.0	285
Birth Order										
First	85.1	963	10.2	89.8	41.2	32.9	24.2	1.8	100.0	839
Second	85.6	964	6.8	93.2	46.5	31.6	19.7	2.1	100.0	856
Third or More	87.2	624	7.9	92.1	48.4	33.6	17.6	0.3	100.0	550
Place of Delivery										
District Maternity Hospital	85.6	1527	9.9	90.1	42.3	31.7	24.0	2.0	100.0	1,328
Tirana Maternity	91.8	744	5.1	94.9	41.0	40.4	18.5	0.2	100.0	685
Private Clinic/Hospital	**	**	**	**	**	**	**	**	**	16
Birth House/Health Center	89.6	154	3.4	96.6	61.0	31.6	6.5	0.8	100.0	136
At Home	70.1	107	5.0	95.0	70.7	22.5	6.5	0.4	100.0	80

Table 5.13Time Between Delivery and First Baby Clinic Visit by Selected Characteristics,
for Births in 1997–2002
Reproductive Health Survey: Albania 2002

** Percentages are not shown when base is less than 25 cases

ables

Table 5.14
Percentage of Babies With Birth Certificates Issued and Time Between
Delivery and Certificate Issued for Births in 1997–2002
Reproductive Health Survey: Albania 2002

Chapter 5

		h Birth ificate	lı				and Birth Cer ution of Week		
	X	No. of		4.0			Don't		No. of
Characteristics	Yes	Cases	< 1	1–2	3–4	> 4	Remember	Total	Cases
Total	92.6	2,551	19.9	49.4	16.9	10.8	3.0	100.0	2,361
Strata									
Metro Tirana	90.7	900	15.5	47.9	21.7	12.3	2.6	100.0	817
Other Urban	94.5	807	22.1	50.8	14.0	9.6	3.6	100.0	767
Other Rural	92.2	844	19.9	49.1	17.2	11.1	2.8	100.0	777
Residence									
Urban	93.3	1,522	20.8	49.7	15.9	10.2	3.3	100.0	1,414
Rural	92.1	1,029	19.2	49.3	17.5	11.2	2.8	100.0	947
Age Group at Birth									
< 20	88.8	196	19.5	47.0	12.5	15.6	5.3	100.0	166
20–24	92.5	867	20.3	46.9	16.6	12.8	3.3	100.0	798
25–34	93.0	1,350	20.4	49.9	18.0	9.3	2.4	100.0	1,263
35–44	95.5	138	14.5	62.3	15.3	5.5	2.4	100.0	134
Education Level									
Primary or Less	91.5	1,228	19.7	49.8	16.0	12.2	2.2	100.0	1,114
Secondary Incomplete	88.5	179	22.4	35.4	18.5	16.7	7.0	100.0	161
Secondary Complete	95.3	880	19.6	51.3	17.5	7.9	3.6	100.0	835
Post-Secondary	94.9	264	20.0	50.7	19.6	6.6	3.1	100.0	251
Socioeconomic Index									
Low	93.2	1,020	18.8	50.3	17.7	10.5	2.8	100.0	939
Medium	92.2	1,222	21.0	49.4	15.8	10.4	3.4	100.0	1,134
High	91.3	309	20.4	43.9	18.7	15.2	1.8	100.0	288
Birth Order									
First	90.1	963	**	**	**	**	**	**	**
Second	93.6	964	21.8	48.7	17.8	9.0	2.8	100.0	908
Third or More	94.9	624	19.7	52.5	16.2	8.8	2.8	100.0	591
Place of Delivery									
District Maternity Hospital	92.8	1,527	**	**	**	**	**	**	**
Tirana Maternity	91.3	744	11.9	48.5	23.8	10.5	5.2	100.0	676
Private Clinic/Hospital	**	**	**	**	**	**	**	**	17
Birth House/Health Center	94.4	154	17.4	58.5	15.1	7.5	1.5	100.0	145
At Home	91.3	107	22.2	42.1	15.3	16.7	3.8	100.0	95

** Percentages are not shown when base is less than 25 cases

	-			cteristic Health							
Characteristics	Post Partum Complication	Painful Uterus	Painful Urination	Bad Smelling Vaginal charge	Severe Bleeding	High Fever	Breast Infection	Infection of Surgical Wound	Faint/ Coma	Other	No. of Cases
Total	24.8	15.9	7.4	6.5	5.9	4.8	4.3	4.2	2.9	0.5	2,551
Strata											
Metro Tirana	32.0	19.1	8.2	8.1	8.6	6.5	5.9	6.6	2.8	0.9	900
Other Urban	23.5	14.0	6.7	5.6	5.5	4.1	4.3	3.6	2.7	0.3	807
Other Rural	23.7	16.1	7.5	6.5	5.3	4.7	3.9	3.9	3.1	0.5	844
Residence											
Urban	26.2	15.4	7.2	6.1	6.5	5.1	4.8	4.6	2.9	0.5	1,522
Rural	23.9	16.3	7.5	6.7	5.4	4.6	4.0	3.9	3.0	0.5	1,029
Age Group at Birth											
< 20	31.5	22.2	9.3	7.5	5.5	6.0	3.3	3.9	3.3	1.3	196
20–24	23.9	14.9	7.9	6.7	5.9	4.8	4.4	5.3	3.1	0.5	867
25–34	25.0	15.9	6.7	6.3	5.8	4.6	4.3	3.5	2.6	0.3	1,350
35–44	19.1	12.6	6.4	5.2	6.8	5.0	5.3	4.8	4.6	2.0	138
Education Level											
Primary or Less	24.8	15.8	7.8	6.7	6.6	5.4	4.5	4.2	3.2	0.6	1,228
Secondary Incomplete	27.7	20.6	5.1	5.6	3.0	3.9	4.8	1.0	3.0	0.7	179
Secondary Complete	24.1	15.9	7.4	6.8	4.7	3.3	3.8	4.9	2.6	0.3	880
Post-Secondary	25.4	12.6	5.9	4.4	7.9	6.9	4.7	4.4	2.1	1.2	264
Socioeconomic Index											
Low	23.7	15.6	7.3	6.7	5.8	5.1	4.5	3.9	3.1	0.6	1,020
Medium	25.3	16.1	7.0	6.3	5.9	4.3	4.0	4.1	2.4	0.4	1,222
High	29.7	16.7	9.6	6.7	6.0	5.8	5.1	6.8	5.2	1.1	309
Birth Order											
First	26.0	16.6	10.1	5.4	5.9	5.4	4.4	6.0	3.1	0.7	963
Second	23.3	14.5	5.7	6.8	4.4	3.9	4.7	3.3	2.3	0.2	964
Third or More	25.2	16.8	5.6	7.6	7.7	5.1	3.7	2.9	3.6	0.8	624
Pregnancy Complication	ons										
Yes	40.1	26.3	13.5	13.4	11.9	10.1	5.9	9.8	5.0	0.7	505
No	20.9	13.3	5.8	4.7	4.3	3.5	3.9	2.8	2.4	0.5	2,046
Prolonged Labor											
No	23.1	15.3	7.2	6.2	5.8	4.1	3.7	3.3	2.5	0.5	2,195
Yes	43.6	26.8	10.1	14.1	6.2	8.8	11.4	16.0	9.5	0.5	70
No Labor	35.8	18.5	7.9	6.5	6.0	9.6	7.0	9.1	6.2	0.6	233
Dk/Dr	31.2	16.6	6.9	6.6	6.1	9.0	10.3	6.0	0.8	0.0	53

Table 5.15 Percentage of Women Reporting Postpartum Complications, by Selected Characteristics, for Births in 1997–2002 Reproductive Health Survey: Albania 2002

Albania Reproductive Health Survey, 2002 Final Report

Table 5.16
Percent of Children Ever Breastfed and Percent Distribution of Initiation of Breastfeeding,
by Selected Characteristics, for Births in 1997–2002
Reproductive Health Survey: Albania 2002

Chapter 5

		dren Ever <u>eastfed</u>		Ini			ding After Bi on of Hours <u>)</u>		
Characteristics	%	No. of Cases	1 Hour or Less	2–23	24–47	48 Hours or More	Unknown	Total	No. of Cases
Total	93.2	2,551	15.4	51.8	8.8	23.2	0.9	100.0	2,381
Strata	00.2	2,001	1011	0110	0.0	2012	0.0	10010	2,001
Metro Tirana	93.5	900	22.5	47.0	7.0	22.5	1.0	100.0	839
Other Urban	92.9	807	14.7	50.9	9.2	24.8	0.5	100.0	753
Other Rural	93.3	844	13.9	53.5	9.0	22.6	1.0	100.0	789
Residence									
Urban	93.0	1,522	17.3	49.2	8.4	24.4	0.7	100.0	1,415
Rural	93.4	1,029	14.1	53.5	9.0	22.4	1.0	100.0	966
Age Group at Birth									
< 20	92.3	196	16.3	50.4	7.7	23.5	2.0	100.0	179
20–24	93.5	867	14.1	53.7	8.3	22.9	0.9	100.0	810
25–34	93.0	1,350	16.8	51.2	9.2	22.1	0.6	100.0	1,260
35–44	95.0	138	9.1	47.6	9.6	32.9	0.9	100.0	132
Education Level									
Primary or Less	93.3	1,228	14.4	54.6	7.2	23.0	0.8	100.0	1,143
Secondary Incomplete	90.7	179	18.4	49.0	6.0	25.6	1.0	100.0	166
Secondary Complete	93.1	880	16.1	48.1	12.6	22.1	1.1	100.0	820
Post-Secondary	95.6	264	17.2	47.2	7.7	27.6	0.2	100.0	252
Socioeconomic Index									
Low	93.9	1,020	15.1	52.6	8.2	22.7	1.4	100.0	957
Medium	93.0	1,222	15.0	52.1	9.1	23.3	0.5	100.0	1,142
High	90.8	309	19.1	44.6	10.6	25.7	0.0	100.0	282
Birth Order									
First	93.4	963	14.8	48.3	10.2	25.5	1.2	100.0	896
Second	93.5	964	17.2	53.6	7.6	21.0	0.6	100.0	902
Third or More	92.7	624	13.7	54.4	8.3	22.7	0.8	100.0	583
Type of Delivery									
Vaginal	93.9	2,182	16.8	55.6	7.9	18.8	0.9	100.0	2,054
Cesarean Section	88.3	369	4.7	23.4	15.4	55.7	0.8	100.0	327
Baby Weight at Birth									
< 2500 Grams	79.0	123	11.7	44.0	11.1	32.8	0.4	100.0	100
≥ 2500 Grams	94.0	2,414	15.6	52.1	8.7	22.7	0.9	100.0	2,270
Don't know	**	**	**	**	**	**	**	**	11

** Percentages are not shown when base is less than 25 cases

Maternal and Child Health

Table 5.17 Mean Duration of Breastfeeding in Months, by Type of Breastfeeding and Selected Characteristics, for Births in 1997–2002 Reproductive Health Survey: Albania, 2002

Characteristics Breastfeeding Breastfeeding Breastfeeding Total 2.8 4.9 14.3 Strata 2.7 4.8 15.2 Other Urban 2.5 4.9 13.8 Other Rural 2.9 4.9 14.6 Residence 14.1 Urban 2.4 4.7 14.1 Rural 3.0 5.0 14.7 Age Group at Birth 2 14.8 < 25 2.7 5.1 13.9 25-44 3.1 4.8 14.9 Education Level 2.0 4.1 14.8 Secondary Incomplete or Less 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.2 Birth Order 14.8 14.9 Edw 2.9 5.2 15.7 Medium/High 2.7 5.2 14.3 Second or More 2.6 4.8 14.8 <		Exclusive	Full	Any
Strata Intermediate Alternation Alternation Metro Tirana 2.7 4.8 15.2 Other Urban 2.5 4.9 13.8 Other Rural 2.9 4.9 14.6 Residence Urban 2.4 4.7 14.1 Rural 3.0 5.0 14.7 Age Group at Birth < 25 2.7 5.1 13.9 25-44 3.1 4.8 14.9 Education Level Secondary Incomplete or Less 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Uper of Delivery Vaginal <td< th=""><th></th><th></th><th></th><th></th></td<>				
Atto 2.7 4.8 15.2 Other Urban 2.5 4.9 13.8 Other Rural 2.9 4.9 14.6 Residence Urban 2.4 4.7 14.1 Rural 3.0 5.0 14.7 Age Group at Birth < 25 2.7 5.1 13.9 25-44 3.1 4.8 14.9 Education Level Secondary Incomplete or Less 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery Vaginal 2.7	Total	2.8	4.9	14.3
Other Urban 2.5 4.9 13.8 Other Rural 2.9 4.9 14.6 Residence Urban 2.4 4.7 14.1 Rural 3.0 5.0 14.7 Age Group at Birth - - - < 25	Strata			
Other Rural 2.9 4.9 14.6 Residence 14.6 Urban 2.4 4.7 14.1 Rural 3.0 5.0 14.7 Age Group at Birth 2.5 2.7 5.1 13.9 < 25 2.7 5.1 13.9 25 $25-44$ 3.1 4.8 14.9 Education Level $8condary Incomplete or Less$ 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index 2.9 5.2 15.7 Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order 13.5 5 14.8 First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery 14.8 14.8 Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** 1 Hour or Less 2.4 4.4 16.1 <				
Residence Virban 2.4 4.7 14.1 Rural 3.0 5.0 14.7 Age Group at Birth $<$ 25 2.7 5.1 13.9 25-44 3.1 4.8 14.9 Education Level $ <$				
Urban 2.4 4.7 14.1 Rural 3.0 5.0 14.7 Age Group at Birth . . . < 25	Other Rural	2.9	4.9	14.6
Rural 3.0 5.0 14.7 Age Group at Birth	Residence			
Age Group at Birth 2.7 5.1 13.9 25-44 3.1 4.8 14.9 Education Level Secondary Incomplete or Less 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding 1 Hour or Less 2.4 4.4 16.1 2-23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** S2500 Grams	Urban	2.4	4.7	14.1
< 25	Rural	3.0	5.0	14.7
25-44 3.1 4.8 14.9 Education Level	Age Group at Birth			
Education Level 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index 13.4 13.4 Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order 13.5 13.5 13.5 Second or More 2.6 4.8 14.8 Type of Delivery 2.7 5.2 14.3 Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** 1 Hour or Less 2.4 4.4 16.1 2-23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth ** ** ** < 2500 Grams	• •	2.7	5.1	13.9
Secondary Incomplete or Less 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index 13.4 13.4 Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order 13.5 13.5 13.5 First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery 14.8 14.8 Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding 1 14.4 16.1 2-23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Saby Weight at Birth ** ** ** < 2500 Grams	25–44	3.1	4.8	14.9
Secondary Incomplete or Less 3.0 5.2 14.8 Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index 13.4 13.4 Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order 13.5 13.5 13.5 First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery 14.8 14.8 Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding 1 14.4 16.1 2-23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Saby Weight at Birth ** ** ** < 2500 Grams	Education Level			
Secondary Complete or More 2.0 4.1 13.4 Socioeconomic Index 100 100 Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order 100 100 100 First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery 100 100 100 Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding 110 100 100 1 Hour or Less 2.4 4.4 16.1 2-23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Saby Weight at Birth 100 100 100 < 2500 Grams		3.0	5.2	14.8
Socioeconomic Index Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order	• •	2.0	4.1	13.4
Low 2.9 5.2 15.7 Medium/High 2.7 4.8 13.2 Birth Order 2.7 4.8 13.2 First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery 2.6 4.8 14.3 Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding 1 16.1 2-23 Hours 1 Hour or Less 2.4 4.4 16.1 2-23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth ** ** < 2500 Grams	Socioeconomic Index			
Medium/High 2.7 4.8 13.2 Birth Order		2.9	5.2	15.7
Birth Order First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding ** ** ** 1 Hour or Less 2.4 4.4 16.1 2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth ** ** ** < 2500 Grams	Medium/High			
First 3.1 4.9 13.5 Second or More 2.6 4.8 14.8 Type of Delivery Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** ** Initiation of Brestfeeding ** ** ** 1 Hour or Less 2.4 4.4 16.1 2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth ** ** ** < 2500 Grams	-			
Second or More 2.6 4.8 14.8 Type of Delivery Vaginal 2.7 5.2 14.3 Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding ** ** ** 1 Hour or Less 2.4 4.4 16.1 2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth ** ** ** < 2500 Grams ** ** ** ≥ 2500 Grams 3.0 5.2 14.7		3.1	4 9	13.5
Type of Delivery Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding ** ** ** 1 Hour or Less 2.4 4.4 16.1 2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth ** ** < 2500 Grams				
Vaginal 2.7 5.2 14.3 Cesarean Section ** ** ** Initiation of Brestfeeding 1 1 1 1 Hour or Less 2.4 4.4 16.1 2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth 2500 Grams ** ** ** ≥ 2500 Grams 3.0 5.2 14.7				
K** K** K** K** Initiation of Brestfeeding 1 1 1 1 Hour or Less 2.4 4.4 16.1 2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth < 2500 Grams		27	5.2	1/ 3
Initiation of Brestfeeding 1 Hour or Less 2.4 4.4 16.1 2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth 2500 Grams ** ** ** ≥ 2500 Grams 3.0 5.2 14.7	0			
1 Hour or Less2.44.416.12–23 Hours3.25.815.824 or More3.15.714.4Unknown******Baby Weight at Birth< 2500 Grams				
2–23 Hours 3.2 5.8 15.8 24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth 2500 Grams ** ** ≥ 2500 Grams 3.0 5.2 14.7	•	2.4	1 1	16 1
24 or More 3.1 5.7 14.4 Unknown ** ** ** Baby Weight at Birth - - - < 2500 Grams				
Unknown ** ** ** Baby Weight at Birth < 2500 Grams	= =0			
Baby Weight at Birth ** ** ** < 2500 Grams				
< 2500 Grams				
≥ 2500 Grams 3.0 5.2 14.7		**	**	**
		3.0		
	Don't know			

**Percentages are not shown when base is less than 25 cases

Table 5.18Infant and Child Mortality Rates (Infant and Child Deaths per 1,000 Live Births)by Selected Characterisitcs, for Children Born August 1992–July 2002Reproductive Health Survey: Albania 2002

		Infant Mo	rtality	- Child Mortality	Under-5	No. of
Characteristics	Total	Neonatal	Postneonatal	(1–4 years)	Mortality	Births
Total	26.2	11.9	14.3	5.8	31.9	4,823
Residence						
Urban	21.7	10.3	11.5	6.2	27.8	2,881
Rural	29.3	13.0	16.3	5.5	34.7	1,942
Age Group at Birth						
< 25	23.6	13.0	10.6	6.0	29.4	2,066
25–44	28.2	11.1	17.1	5.7	33.7	2,757
Education Level						
Secondary Incomplete or Less	33.4	12.8	20.6	7.4	40.6	2,576
Secondary Complete or Post-Secondary	14.7	10.4	4.3	3.3	18.0	2,247
Socioeconomic Index						
Low	31.4	11.6	19.8	5.7	36.9	1,949
Medium/High	21.4	12.1	9.2	5.9	27.2	2,874
Birth Order						
1	22.6	10.3	12.3	6.1	28.6	1,874
2	23.1	13.6	9.5	5.3	28.2	1,800
3+	33.9	11.9	22.0	5.9	39.7	1,148
Birth Interval *						
First Births	21.7	10.6	11.1	6.0	27.6	1,746
< 24	63.8	31.4	32.4	7.2	70.5	841
24 +	14.5	5.3	9.2	4.6	19.0	2,208

* Excludes 28 cases with birth interval undefined

CHAPTER 6 CONTRACEPTION AWARENESS AND KNOWLEDGE OF USE

Tn this chapter the level of contraceptive Lawareness and the knowledge of how to use contraceptive methods are analyzed. Men and women of reproductive age in Albania were also asked about their sources of information regarding contraception and their opinion on contraceptive effectiveness. The information described in this chapter can help family planning program planners assess the overall knowledge on contraception and make informed decisions on how and where to strengthen information, education and communication programs on contraception.

Contraceptive Awareness and Knowledge of Use

Almost all of the Albanian women were aware of at least one method of contraception (97%), with 90% aware of at least one modern method. The most frequently known modern methods were condoms (81%), pills (68%), and tubal ligation (68%). More than half of the women had never heard of other modern methods. For example, spermicides were recognized as a contraceptive method by only 5% of the women. In addition, Albanian women demonstrated a relatively high awareness of traditional methods (86%) with withdrawal being the most frequently known traditional method (85%). Only onefourth of the women were aware of periodic abstinence as a contraceptive method (Table 6.1A).

The level of awareness of any contraceptive method was highest among women who lived in urban areas other than Tirana, were between the age of 40 to 44 years, married, and more educated. Modern contraceptive awareness was almost universal among residents of Tirana and other urban areas. Not surprisingly, women from rural areas were overall less aware of any contraceptive method than their urban and Tirana residents' counterparts. Between Tirana and other urban areas, the level of condom and pill awareness were similar. However, in urban areas and Tirana awareness of tubal ligation (76% vs. 70% respectively) and injectables (43% vs 41% respectively) was slightly higher among residents of other urban areas. Awareness of emergency contraception, vasectomy, and spermicides was less than 11% and among rural residents was less than 6%.

Traditional methods were also more frequently recognized by women of urban residence. Among traditional methods, the largest gap in awareness was for periodic abstinence (33% vs. 19%).

Awareness of both modern and traditional methods among women was directly correlated with age (Table 6.1A). Almost all women aged 20 and older were aware of at least one contraceptive method. Awareness of modern methods had a weak inverse relationship with age for women aged 20 and older, while for traditional methods, level of awareness increased with the respondent's age. The best known methods for the youngest group were condoms (79%), pills (58%), and withdrawal (53%). However, among the oldest age group, withdrawal was the most recognized method (99%), followed by tubal ligation (79%), condoms (73%), and pills (64%). Only about one third of the 15-19 year olds have heard of tubal ligation as a contraceptive method.

Almost all women knew at least one modern method (87- 94%) regardless of their marital status. Withdrawal was the most widely known method among currently (100%) and previously married (99%) women, while never married women had mostly heard of condoms (82%). Currently married women were familiar with condoms as much as tubal ligation (80%), followed by pills (70%). However, awareness of modern methods was greatest among previously married women (94%), and tubal ligation and pills ranked second and third respectively after condoms.

Among the female respondents, the awareness of both modern and traditional methods was directly correlated with their level of education (Table 6.1A). The difference in modern method awareness between the least educated women and those with post secondary education ranges from 29% for condoms to 92% for spermicides. The smallest gap in the level of awareness by education level is for withdrawal. Overall, more educated women had slightly higher awareness of modern methods than traditional methods.

Awareness of contraceptive methods by men was almost universal (Table 6.1B) and slightly higher than overall women's awareness. About 99% of men in the survey knew at least one contraceptive method, mainly withdrawal and condoms (89% each). However, more than twothirds of men have not heard about other contraceptive methods. Among modern methods, pills were recognized by only one third of men, followed by tubal ligation and emergency contraception (15% and 10% respectively). Fewer men had heard of IUDs (8%), injectables (6%), vasectomy (5%), and spermicides (4%). Less than one- fourth of men knew about periodic abstinence as a contraceptive method.

By residence, the level of contraceptive awareness among men had little variation among Tirana residents and residents of other urban or rural areas (Table 6.1B). Urban residents (Tirana and other urban areas) were more aware of modern contraceptive methods than traditional methods (94% and 96% vs. 91%, respectively), while rural residents have heard of traditional methods more than modern ones (88% vs. 84%). Tirana residents had the highest level of awareness for all methods of contraception except for condoms and withdrawal, which had a slightly higher level of awareness among residents of other urban areas. Awareness of all modern methods was lowest for men in rural areas.

Contraceptive awareness (any method) was lowest, but still more than 90%, among 15-19 year old men (94%), however it was virtually universal among those aged 20 and older. Awareness of modern contraceptive methods had an inverse relationship with age after age 20, with awareness in the mid 90% for ages 25 to 34, mid 80% for ages 35 to 44, and 69% for ages 45 to 49. Nonetheless, this relationship was positive for traditional methods. Younger men (aged 15-19) were more aware of modern contraceptive methods than traditional ones (91% vs. 55%).


Never married men had greater awareness of modern contraceptive methods than currently married men (95% vs. 86%). Although virtually all currently married men have heard of at least one traditional contraceptive method, less than threefourths of never married men were aware of traditional methods. As shown in Table 6.1B there were only 14 male respondents who were previously married, thus percentages for this group were not calculated.

The awareness of both modern and traditional contraceptive methods was directly associated with men's educational level (Table 6.1B). The gap in the level of awareness between the most and the least educated men was 19 percentage points for condoms, the best known method. For other

modern methods the difference between post-secondary and primary or less educated men, was even greater.

Male and female respondents who have heard of at least one contraceptive method were asked whether they knew how to use the method. Among women, knowledge of how to use any method, modern or traditional was lower than the corresponding awareness of the method (59% vs. 90% for any modern and 80% vs. 86% for any traditional), (Tables 6.1A and 6.2A). The proportion of women who knew how to use specific methods was considerably lower than the corresponding awareness of methods. This gap was largest for the most widely known modern methods: condoms, pill, tubal ligation, injectables, and IUD (Figure 6.1). Although 81% of women

have heard of condoms, only slightly less than half of them stated that they knew how to use them (49%). The gap between awareness and knowledge of how to use the method was greatest for the pill (68% vs. 36%) and injectables (35% vs. 17%). The proportion of women who knew how tubal ligation and IUD protect against pregnancy was about 40% less than the proportion who have heard about them (68% vs. 40% and 24% vs. 14%, respectively). A relatively smaller gap between awareness of method and knowledge of its use was present for other modern methods (emergency contraception, vasectomy, and spermicides). The percentage of women who were aware of periodic abstinence and withdrawal were slightly different from the percentage who knew how to use them (85% vs. 79% and 25% vs. 20% respectively).




Among the female respondents, the gap between awareness of modern methods and knowledge of how to use them had a slight urban-rural variation, with a 10 percentage point gap among residents of Tirana and other urban areas vs. 13% among rural residents (Table 6.2A). Although 95% of urban residents had heard of at least one modern method, their knowledge of how to use them was substantially lower (70% and 66% among residents of Tirana and other urban areas respectively). Again, the difference between awareness of modern methods and knowledge of how to use them was greatest among rural residents (83% vs. 53%). The urban-rural variation in the knowledge of specific contraceptive methods was considerable. Knowledge of how to use condoms was to some extent lower among rural women (42%) than residents of urban areas (57% and 61% other urban and Tirana respectively). Women in rural areas had also considerably lower knowledge of the pill (29% v. 44% and 46%), injectables (13% vs. 22% and 23%), and IUD (10% vs. 19% and 20%) than other urban and Tirana residents. Women in the urban areas had 3-4 times greater knowledge about how emergency contraception and vasectomy prevent pregnancy than rural residents (12%-17% vs. 4% and 6%-8% vs. 2% respectively).

Regarding the age of respondents, among women, the difference between awareness of contraceptive methods and knowledge of how to use them was greatest among youngest women (88% vs. 57%) and smallest among women aged 35 and older (99%-100% vs. 97-98%). Although 86% of women aged 15-19 have heard of at least one modern contraceptive method, only about half of them knew how to use them. The knowledge of how to use specific contraceptive methods was highest among women aged 20-24 for condoms (56%); among women aged 30-34 for the pill (40%); among women aged 35-39 for tubal ligation (52%); and among women aged 25-34 for injectables (22%). The percentage of women who knew how to use withdrawal improved with increase in age of the respondent (from 39% for 15-19 year olds to 98% for 40-44 year olds). A similar increase for periodic abstinence (10% to 22%) is also observed.

Among currently and previously married women, there were no differences in contraceptive awareness and knowledge of use. However, never married women had a 32 percentage point gap between their contraceptive awareness and knowledge of use (89% vs. 57%). Knowledge of how to use modern methods was greatest among women who were previously married (75%), followed by married women (63%). Knowledge of how withdrawal works against pregnancy was universal among previously and currently married women (99%).

Women's education is directly associated with knowledge of how to use contraceptive methods. The gap in the knowledge of how to use modern methods between the least educated and the most educated women was greatest for condoms (41 percentage points) and the pills (42 percentage points).

Among the male respondents, knowledge of how to use modern contraceptive methods waslowerthantheir corresponding awareness (70% vs. 89%). However, this gap was only 2 percentage points for traditional methods (Table 6.2B). While 89% of men have heard of condoms, only 69% of them knew how to correctly use them. The proportion of men who knew how other modern contraceptive methods prevent pregnancy was relatively small, hence, the gap between knowledge of how to use and corresponding awareness was smaller: 21% vs. 33% for the pills; 11% vs. 15% for tubal ligation; and 7% vs. 10% for emergency contraception (Figure 6.2).



Knowledge of how contraceptive methods work had a slight urban-rural variation among the male respondents. The largest gap between knowledge of contraceptive use and the corresponding awareness was among rural residents (90% vs. 98%), followed by residents of Tirana (94% vs.100%) and other urban areas (94% vs. 99%). Although 60% of men in the rural areas knew how to use condoms, a considerably smaller percentage of them knew how the pills and tubal ligation prevents pregnancy (12% and 10%, respectively). The urbanrural variation for traditional methods was relatively small (89% vs. 86%).

With respect to age groups, there was only a slight gap in knowledge on how to use contraceptivemethods and the corresponding awareness for respondent men 25 years and older. Meanwhile, younger men (aged 15-19) had a huge gap between knowledge of how to use contraception and awareness of it (64% vs. 94%). While 91% of men aged 15-19 have heard of at least one contraceptive method, slightly more than half of them knew how to use them. Knowledge about how to use condoms was highest among men aged 20-24 (86%) and lowest among the oldest age group (51%). On the other hand, all the men in the oldest group stated that they knew how the withdrawal method works, while only half of the youngest respondents stated that they knew how to use withdrawal. One reason for this could be that 80% of adolescent men reported that they are not yet sexually experienced and therefore had not yet started to think about using contraceptives

Among men, all married men and 80% of never married men stated that they knew how to use at least one contraceptive method (Table 6.2B). The difference between awareness of modern contraceptive methods and knowledge of how to use them had only a slight variation according to the men's marital status. Never married men had greater knowledge of how to use all modern methods than married men, except for tubal ligation (7% vs. 14%). On the other hand, all married men knew how to use withdrawal while only 68% of never married men knew how it was used.

According to marital status, all men with post-secondary education knew how to use at least one contraceptive method. The difference between awareness of contraceptive methods and knowledge of how to use them among men with secondary and primary or less education was 9 and 6 percentage points, respectively.

Knowledge about Contraceptive Source

The Albania Reproductive Health Survey included questions about source of modern methods of contraception for both women and men. About three fourths of women knew where to get condoms; almost two thirds knew a source for the pills; more than half knew where to get tubal ligation; and 30% knew a source for injectables (Table 6.3A). However, only 21% and 10% of women knew where to get IUDs and emergency contraception respectively. Very few knew a source for spermicides or where vasectomies are performed.

Knowledge of a source for contraceptive methods was higher among residents of urban areas (91% Tirana and 92% other urban) than rural residents (76%). Knowledge about where to get contraceptive methods had little variation among different age groups, with the highest among women aged 35-39 (87%). Previously married women had the highest knowledge of contraceptive sources (87%) and women with the highest education had almost universal knowledge of where to get contraceptive methods, especially for condoms (97%) and pills (93%).

Knowledge of a source for contraceptive methods among men was highest for condoms (70%). A substantially smaller proportion of men knew where to get pills (26%), or where vasectomy procedure is performed (4%) (Table 6.3B). Among men, knowledge of contraceptive sources was highest among Tirana residents (82%); men aged 20-24 (89%); never married men (79%); and men with a post-secondary education (94%). The difference between the highest educated and the least educated men in regard to the knowledge of a source was 35% for condoms, 82% for the pills, 65% for tubal ligation, 88% for emergency contraception and IUD, 83% for vasectomy, and 85% for spermicides.

Most Important Source of Information about Contraception

Survey respondents were asked their most important source of information regarding specific contraceptive methods (Table 6.4A). Among women, the most important sources of information for condoms and pills were television or radio (64%, and 50% respectively), followed by friends (15% pills and 22% condoms). Friends were the most important source of information about IUDs (32%), injectables (46%), tubal ligation (43%) and periodic abstinence (60%). The second most important source of information about IUDs (25%) and injectables (16%) were television and radio; about tubal ligation was a physician (17%), and about periodic abstinence was a relative (13%). A partner or husband was the most important source of information for withdrawal (78%).

Among men, television or radio were mentioned by half of the men as the most important source of information for condoms and the pills, followed by friends for condoms (24%) and books/newspaper/ magazine for the pill (18%) (Table 6.4B). The same proportion of men (28%) named television or radio and a physician as the most important sources of information for tubal ligation. The most important source of information regarding injectables and IUDs were the media (37% and 26% respectively). The most important sources of information about withdrawal were friends and coworkers (72%); for periodic abstinence, the most important sources were their partner or wife and coworkers (40%).

Knowledge about Contraceptive Effectiveness

Tables 6.5A and 6.5B lists contraceptive methods according to their theoretical effectiveness as published by the United States Food and Drug Agency (USFDA) (2002)¹. Implants and vasectomy have the lowest failure rate (0.09 -0.1 pregnancies per 100 women) and therefore are the most effective methods. However, they are not widely available in Albania. The pills (combined oral contraceptives) have an expected failure rate of 0.1 pregnancies per 100 women if used correctly and consistently, but with typical use, the failure rate increases to 5 pregnancies per 100 women. The next effective methods are injectables, tubal ligation, and IUDs with failure rates of 0.3, 0.5 and 0.6 pregnancies per 100 women respectively when used correctly. Male condoms and other barrier methods have higher failure rates of 3-6 pregnancies per 100 women, and are less effective with the typical use (14-26 pregnancies per 100 women). Traditional methods are the least effective in preventing pregnancies.

Among women who have heard of modern methods, less than 20% thought that modern contraceptive methods were very effective. Condoms were the exception, with 18% of women considering them very effective (Table 6.5A). The proportion of women who were aware of a specific modern method and believed that they are very effective or effective was 57% for the pills, 51% for injectables, 72% for tubal ligation, 58% for IUDs, and 54% for condoms. On the other hand, 87% of women who have heard of withdrawal, the least effective method, believed that it is very effective or effective.

Among men, because of a very low awareness of modern contraceptive methods, the proportion that considered them as very effective or effective was small (Table 6.5B). After excluding men who were not aware of a specific contraceptive method, the proportion of men who believed certain methods were very effective or effective was 65% for the pills, 94% for tubal ligation, and 78% for condoms. Among men who have heard of periodic abstinence or withdrawal, 70% and 80%, respectively, considered them to be very effective or effective.

¹ http://www.fda.gov/fdac/features/1997/conceptbl.html

⁸ Contraception Awareness and Knowledge of Use Tables

Table 6.1 A
Percentage of Women Aged 15-44
Who Have Heard of Specific Methods of Contraception by Selected Characteristics
Reproductive Health Survey: Albania, 2002

		R	esidence)			A	ge			Marital Status				Education	1
Contraceptive Method	Total	Tirana	Other Urban	Rural	15-19	20-24	25-29	30-34	35-39	40-44	Married	Previously Married	Never Married	Primary or Less	Secondary	Post- Secondary
Any Method	96.5	97.5	98.2	95.2	88.0	97.0	99.3	99.3	98.6	99.6	99.9	99.3	89.4	94.5	98.5	100.0
Any Modern Method	89.6	95.3	94.9	85.2	85.5	92.0	92.7	91.4	90.6	86.9	90.9	94.0	86.9	83.5	96.2	99.8
Condoms	81.3	90.9	90.5	73.4	79.3	87.0	86.2	81.3	80.9	72.7	80.4	89.3	82.4	70.9	92.2	99.1
Pills	68.0	79.4	79.0	58.8	58.3	73.4	76.2	70.8	68.2	64.1	69.3	77.5	65.0	56.1	79.0	95.9
Tubal Ligation	67.6	69.8	75.6	62.5	36.2	59.6	77.9	81.7	83.0	79.0	80.3	82.8	41.2	60.3	73.8	86.9
Injectables	34.9	41.3	43.4	28.4	20.3	33.7	43.6	41.0	40.2	35.4	39.9	42.4	24.5	27.1	41.5	55.8
IUD	24.2	34.6	32.9	16.4	9.6	23.1	27.3	32.7	30.1	27.2	28.8	25.8	14.9	15.1	30.9	52.6
Emergency Contraception	10.7	20.8	15.2	5.3	8.6	14.2	10.9	13.4	9.5	7.9	10.3	10.6	11.5	4.9	12.6	40.3
Vasectomy	6.0	13.7	8.0	2.7	3.6	6.4	8.1	6.9	6.6	5.3	6.2	6.6	5.6	1.5	8.7	24.0
Spermicides	5.0	11.1	5.6	2.9	2.6	6.2	5.7	6.2	6.2	3.8	4.8	9.3	5.0	1.6	6.6	20.1
Any Traditional Method	86.0	86.8	90.5	83.3	55.0	82.9	94.5	97.4	98.3	99.4	99.8	99.3	57.8	83.9	86.8	95.7
Withdrawal	85.0	85.9	89.0	82.6	52.8	81.6	93.3	96.8	98.1	99.3	99.7	99.3	54.9	83.5	85.2	94.0
Periodic abstinence (rhythm method)	25.2	33.0	33.1	18.6	12.5	26.2	29.9	32.0	27.1	27.9	29.4	27.2	16.7	16.8	29.3	62.0
No. of Cases	5,697	2,108	1,816	1,773	1,094	936	946	1,067	958	696	3,965	88	1,644	2,519	2,483	695

Contraception Awareness and Knowledge of Use

		F	Residence	e		Age						Marital	Status*		Education	1
Contraceptive Method	Total	Tirana	Other Urban	Rural	15-19	20-24	25-29	30-34	35-39	40-44	45-49	Married	Never Married	Primary Or Less	Secondary	Post- Secondary
Any Method	98.6	99.6	99.3	97.8	93.6	99.9	100.0	100.0	99.3	99.4	100.0	99.7	96.9	97.2	99.8	100.0
Any Modern Method	89.4	94.1	95.7	83.9	91.0	99.7	95.5	95.6	87.6	83.4	68.8	85.9	94.9	81.4	96.3	99.8
Condoms	88.6	93.9	95.5	82.5	91.0	99.0	94.7	95.2	85.0	83.4	67.6	84.9	94.7	80.4	95.7	99.1
Pills	32.7	51.8	42.3	19.5	26.2	42.7	45.4	38.0	28.6	27.9	20.4	28.7	38.3	16.7	40.6	80.7
Tubal Ligation	14.8	18.5	18.3	11.2	6.0	8.0	16.6	18.9	22.2	20.1	15.0	18.5	9.1	10.4	15.8	33.4
Emergency Contraception	10.1	16.2	15.0	4.9	8.3	13.1	10.7	16.8	7.3	9.3	5.4	8.5	12.6	4.9	10.3	37.7
IUD	8.4	15.6	11.9	3.4	4.8	6.3	9.6	15.0	8.3	8.4	7.8	8.5	8.0	3.4	10.0	27.0
Injectables	6.4	11.0	10.3	2.3	4.9	5.4	7.1	10.6	6.1	7.1	4.0	5.8	7.3	3.3	6.4	23.2
Vasectomy	4.7	11.0	4.9	2.1	2.6	2.3	5.7	8.2	3.1	5.4	6.7	4.6	4.8	2.6	4.0	19.1
Spermicides	4.0	10.2	4.0	1.5	2.3	2.6	4.9	6.3	3.8	4.8	3.8	3.9	4.1	1.9	3.9	15.7
Any Traditional Method	89.4	90.6	91.2	87.9	55.2	88.6	98.0	100.0	99.1	99.4	100.0	99.7	73.6	88.5	88.9	96.8
Withdrawal	89.3	90.5	91.2	87.7	54.6	88.6	98.0	100.0	99.0	99.4	100.0	99.6	73.4	88.3	88.9	96.6
Periodic Abstinence (rhythm method)	22.5	28.7	24.6	18.9	12.2	23.2	30.1	33.7	21.8	22.9	17.0	23.6	20.7	16.7	25.1	41.6
No. of Cases	1,740	718	547	475	401	189	218	253	255	277	147	1,023	703	689	825	226

Table 6.1 BPercentage of Men Aged 15-49 Who Have Heard ofSpecific Methods of Contraception by Selected CharacteristicsReproductive Health Survey: Albania, 2002

*Excludes 14 previously married cases.

⁸ Contraception Awareness and Knowledge of Use Tables

eption

		R	esidenc	e			A	ge				Marital Statu	s	Education			
Contraceptive Method	Total	Tirana	Other Urban	Rural	15-19	20-24	25-19	30-34	35-39	40-44	Married	Previously Married	Never Married	Primary or Less	Secondary	Post- Secondary	
Any Method	85.4	87.6	88.4	83.2	57.2	80.6	94.6	95.5	97.3	97.9	99.5	99.3	56.6	83.2	86.9	93.9	
Any Modern Method	59.4	69.6	65.8	53.1	48.7	61.3	62.4	62.8	65.1	60.0	63.4	74.8	50.6	51.0	66.0	84.7	
Condoms	49.4	61.1	57.2	41.7	42.5	55.6	54.4	51.4	47.9	46.1	51.1	60.9	45.2	39.2	57.4	79.6	
Pills	36.0	46.0	44.4	28.6	27.9	38.5	38.7	39.7	38.8	35.2	37.9	53.0	31.2	27.0	41.9	68.5	
Tubal Ligation	40.2	38.5	46.2	37.4	20.6	35.4	43.1	48.1	51.6	49.7	47.5	64.9	24.0	34.9	44.1	56.7	
Injectables	17.4	22.8	21.8	13.4	9.0	15.7	21.9	22.0	20.9	17.7	20.4	27.8	10.7	12.4	20.4	36.4	
IUD	14.0	19.9	19.3	9.5	5.2	14.0	14.7	19.4	18.4	15.8	16.5	16.6	8.9	7.7	18.8	33.7	
Emergency Contraception	8.6	16.9	12.0	4.4	6.4	12.2	8.4	11.2	7.5	6.2	8.4	9.3	8.9	4.0	9.4	35.0	
Vasectomy	4.3	8.3	5.9	2.3	2.1	4.5	5.5	5.3	5.5	3.7	4.8	5.3	3.4	1.2	6.0	17.3	
Spermicides	3.6	7.4	3.8	2.3	1.4	4.5	4.2	4.5	4.7	2.8	3.6	8.6	3.1	1.2	4.4	15.1	
Any Traditional Method	79.5	80.8	83.4	77.0	40.4	72.4	90.4	94.9	96.2	97.6	99.2	99.3	39.2	78.4	79.5	86.7	
Withdrawal	78.9	79.8	83.0	76.4	38.9	71.7	89.6	94.4	96.2	97.6	99.2	99.3	37.4	77.9	78.9	85.7	
Periodic Abstinence (rhythm method)	19.7	25.8	24.8	15.1	9.5	20.0	23.6	24.9	22.1	21.5	23.5	21.2	11.9	13.4	22.4	48.3	
No. of Cases	5,697	2,108	1,816	1,773	1,094	936	946	1,067	958	696	3,965	88	1,644	2,519	2,483	695	

Contraception Awareness and Knowledge of Use

		F	Residenc	e			A	ge					Marital Status	6		Education	1
Contraceptive Method	Total	Tirana	Other Urban	Rural	15-19	20-24	25-29	30-34	35-39	40-44	45-49	Married	Previously Married	Never Married	Primary or Less	Secodary	Post- Seondary
Any Method	92.0	93.8	94.3	89.9	63.8	94.2	99.8	99.5	99.1	99.3	100.0	99.6	**	80.2	91.2	91.3	99.5
Any Modern Method	69.5	79.9	76.6	61.3	57.6	88.2	85.0	82.9	61.1	61.4	52.1	65.8	**	75.6	59.8	75.8	92.0
Condoms	68.7	79.6	75.8	60.1	57.4	86.2	85.0	82.0	59.8	60.6	50.8	64.8	**	74.8	59.1	74.6	91.6
Pills	20.8	37.7	25.2	11.5	13.1	28.2	34.1	26.1	17.7	16.7	11.2	17.9	**	25.1	10.2	24.3	60.6
Tubal Ligation	11.1	11.1	13.4	9.7	5.1	5.1	14.2	15.3	14.3	14.6	11.4	14.0	**	6.6	9.2	11.2	20.2
Emergency Contraception	7.4	13.7	9.8	3.6	5.4	8.1	9.3	14.2	5.5	6.0	4.1	6.4	**	9.0	3.5	7.6	27.7
IUD	6.8	12.3	10.0	2.7	4.0	5.5	9.0	12.1	5.0	6.1	6.7	6.4	**	7.2	2.9	7.7	22.5
Injectables	5.1	8.0	8.3	2.1	3.4	3.6	6.2	9.1	5.0	5.7	3.8	4.9	**	5.5	2.7	5.1	18.2
Vasectomy	3.6	7.5	4.3	1.7	2.3	2.1	3.9	6.3	2.6	4.4	4.5	3.5	**	3.9	2.1	3.1	14.3
Spermicides	3.5	8.2	3.7	1.5	2.0	2.6	4.0	5.4	3.5	4.2	3.2	3.3	**	3.7	1.7	3.7	11.6
Any Traditional Method	87.3	88.6	88.6	85.9	45.7	87.1	97.7	99.4	99.0	99.3	100.0	99.5	**	68.2	86.5	86.2	96.6
Withdrawal	87.1	88.5	88.6	85.7	45.2	87.1	97.7	99.4	98.8	99.3	100.0	99.5	**	68.0	86.2	86.2	96.4
Periodic Abstinence (rhythm method)	21.4	26.5	24.1	17.7	11.8	23.2	29.1	31.3	20.1	21.0	16.2	22.2	**	20.0	15.6	24.0	39.8
No. of Cases	1,740	718	547	475	401	189	218	253	255	277	147	1,023	14	703	689	825	226

Table 6.2 B Percentage of Men 15-49 Years Who Say They Know How Specific Methods of Contraception are Used by Selected Characteristics Reproductive Health Survey: Albania, 2002

** Percentages are not shown when base is less than 25 cases.

56

² Contraception Awareness and Knowledge of Use Tables

Table 6.3 A
Percentage of Women 15-44 Years Who Say They Know
Where to Get Specific Contraceptive Methods by Selected Characteristics
Reproductive Health Survey: Albania, 2002

		R	esidenc	е			Α	ge				Marital Status	5		Education	1
Contraceptive Method	Total	Tirana	Other Urban	Rural	15-19	20-24	25-29	30-34	35-39	40-44	Married	Previously Married	Never Married	Primary or Less	Secondary	Post- Secondary
Any Modern Method	83.3	90.9	92.0	76.4	77.4	85.1	86.3	84.5	87.3	81.1	85.6	87.3	78.5	74.6	92.4	98.7
Condoms	74.1	84.4	87.0	64.1	70.2	78.4	79.8	74.7	75.5	66.8	74.1	80.8	73.6	61.5	87.1	97.0
Pills	65.2	73.9	76.2	51.1	51.6	67.2	70.5	64.4	64.0	58.4	64.1	73.1	57.7	48.8	74.6	93.1
Tubal Ligation	58.8	59.5	68.1	53.6	30.2	50.3	66.5	72.3	75.7	68.9	70.8	77.5	34.0	51.8	64.7	78.3
Injectables	30.2	35.1	40.0	23.5	17.3	28.6	37.4	35.2	36.4	31.1	35.1	37.9	20.0	22.5	36.7	51.4
IUD	21.3	29.3	30.1	14.2	8.0	21.0	24.3	27.9	27.6	23.8	25.4	22.8	13.2	12.9	27.6	47.9
Emergency Contraception	10.2	19.6	14.6	5.0	8.0	13.3	10.5	12.9	9.0	7.6	9.8	10.0	10.8	4.6	11.9	38.9
Vasectomy	5.1	10.6	6.9	2.5	2.7	5.2	6.7	5.9	6.1	4.7	5.6	5.8	4.1	1.3	7.4	19.9
Spermicides	4.4	10.0	4.9	2.4	2.2	5.6	5.2	5.4	5.5	2.9	4.2	9.2	4.4	1.2	6.0	17.6
No. of Cases	5,697	2,108	1,816	1,773	1,094	936	946	1,067	958	696	3,965	88	1,644	2,519	2,483	695

		R	Residenc	e				Age					Marital Statu	S		Education	l
Contraceptive Method	Total	Tirana	Other Urban	Rural	15-19	20-24	25-29	30-34	35-39	40-44	45-49	Married	Previously Married	Never Married	Primary or Less	Secondary	Post- Secondary
Any Modern Method	70.8	82.2	80.0	60.8	64.2	88.9	86.1	82.4	64.2	59.3	49.4	65.8	**	78.5	60.2	77.9	93.9
Condoms	70.1	81.8	79.3	60.1	63.9	88.9	86.1	81.6	62.7	58.4	48.1	64.9	**	78.4	60.1	76.7	93.1
Pills	26.2	42.7	33.4	15.4	20.0	32.2	37.9	33.8	24.1	21.7	14.2	23.0	**	31.0	12.3	33.0	68.1
Tubal Ligation	12.2	12.2	15.6	10.3	5.0	6.2	13.9	17.1	17.6	17.0	11.5	15.6	**	7.1	9.3	12.7	26.1
Emergency Contraception	8.4	14.9	12.2	3.6	5.0	10.6	10.0	16.1	5.9	7.1	5.2	7.4	**	9.9	3.8	8.1	34.4
IUD	7.3	12.5	11.1	3.1	3.6	5.9	8.9	14.4	6.2	6.7	7.3	7.4	**	7.0	3.2	8.4	24.7
Injectables	5.5	8.5	9.0	2.3	4.0	4.1	6.3	10.0	5.0	6.3	3.2	5.0	**	6.4	3.0	5.1	20.7
Vasectomy	3.6	6.9	3.8	2.1	2.2	1.8	3.5	7.3	2.3	4.2	4.4	3.6	**	3.2	2.4	3.0	12.4
Spermicides	3.6	8.7	3.8	1.5	2.2	2.6	3.9	5.9	3.7	4.4	3.2	3.5	**	3.8	1.8	3.8	12.6
No. of Cases	1,740	718	547	475	401	189	218	253	255	277	147	1,023	14	703	689	825	226

Table 6.3 B

Percentage of Men 15-49 Years Who Say They Know Where to Get Specific Contraceptive Methods by Selected Characteristics

**Percentages are not shown when base is less than 25 cases.

Table 6.4 A

Percent Distribution of Most Important Source of Information about Contraception by Specific Method Among Women Aged 15–44 Who Have Heard About Specific Methods of Contraception **Reproductive Health Survey: Albania, 2002**

			Co	ontraceptive M	ethod		
Most Important Source of Information About Contraception	Condom	Pill	Tubal Ligation	Injectables	IUD	Withdrawal	Periodic Abstinence
Television or Radio	63.6	49.5	10.2	15.7	24.5	1.6	3.2
Friends	15.3	21.8	42.5	46.2	32.4	14.3	59.8
Pharmacist	6.0	5.7	0.0	1.7	1.0	0.0	0.3
Partner or Husband	4.6	0.0	0.7	0.3	1.0	77.7	2.8
Books/Newspaper/Magazine	2.0	4.9	2.2	4.6	6.6	0.6	7.0
Teacher	2.0	1.9	1.0	1.3	2.2	0.3	2.0
Nurse/Midwife/Community Health Worker	1.8	10.2	9.0	9.0	7.7	0.2	3.6
Coworker/Colleagues/Peers	2.5	2.0	2.5	3.6	3.6	1.9	4.1
Physician	1.0	1.1	16.8	10.2	15.5	0.5	2.8
Relative	1.0	2.4	12.6	6.9	4.6	2.2	13.3
Mother or Father	0.2	0.5	2.4	0.5	0.9	0.7	0.9
Other	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Don't Know	0.0	0.0	0.1	0.0	0.0	0.0	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
No. of Cases	4,901	4,178	4,023	2,179	1,643	4,985	1,653

Table 6.4 B

Percent Distribution of Most Important Source of Information about Contraception by Specific Method Among Men Aged 15–49 Who Have Heard About Specific Methods of Contraception Reproductive Health Survey: Albania, 2002

			Co	ontraceptive M	ethod		
Most Important Source of Information About Contraception	Condom	Pill	Tubal Ligation	Injectables	IUD	Withdrawal	Periodic Abstinence
Television or Radio	51.2	44.9	27.7	37.3	26.4	0.2	7.4
Friends	23.7	9.4	5.5	7.4	3.5	41.5	12.7
Pharmacist	1.6	2.0	0.3	1.6	0.3	0.0	0.0
Partner or Wife	0.1	3.5	5.6	4.6	9.0	14.5	25.9
Books/Newspaper/Magazine	6.3	17.9	14.9	15.7	24.3	1.3	8.9
Teacher	3.5	7.6	8.3	14.7	7.9	0.3	3.7
Nurse/Midwife/Community Health Worker	0.4	1.7	4.5	2.0	3.8	0.1	0.9
Coworker/Colleagues/Peers	10.6	7.2	4.3	9.1	14.7	30.2	24.2
Physician	1.6	5.6	28.0	7.6	9.0	0.2	1.4
Relative	0.5	0.6	0.2	0.0	0.3	9.9	14.5
Mother or Father	0.3	0.1	0.2	0.0	0.9	0.5	0.1
Other	0.2	0.0	0.2	0.0	0.0	0.5	0.2
Don't Know	0.1	0.1	0.3	0.0	0.0	0.7	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
No. of Cases	1,601	698	287	149	202	1,528	426

Table 6.5 A Percentage Distribution of Women 15-44 by Their Opinion About Contraceptive Effectiveness if the Method is Used Correctly and Consistently Reproductive Health Survey: Albania, 2002

Contraceptive Method*	Never Heard of Method	Heard of Method	No. of Cases	Very Effective	Effective	Some- what Effective	Not Very Effective	Do Not Know	Total	No. of Cases**
Pill	31.8	68.2	5697	9.5	47.7	9.8	2.9	30.1	100.0	4178
Injectables	64.6	35.4	5697	4.0	46.6	9.3	4.5	35.6	100.0	2179
Tubal Ligation	32.3	67.7	5697	12.7	59.5	4.6	1.9	21.3	100.0	4023
IUD	75.0	25.0	5697	6.0	52.0	8.4	3.2	30.4	100.0	1643
Condom	18.7	81.3	5697	18.3	36.0	17.2	1.6	26.9	100.0	4901
Calendar	74.3	25.7	5697	1.6	27.6	27.6	11.0	31.9	100.0	1653
Withdrawal	15.1	84.9	5697	69.3	17.2	6.9	1.6	4.9	100.0	4985

*Listed in descending order of contraceptive effectiveness when the method is used correctly and consistently (USFDA)

**Opinion about contraceptive effectiveness asked only of women who have heard of method

Table 6.5 BPercentage Distribution of Men 15-49 by Their Opinion AboutContraceptive Effectiveness if the Method is Used Correctly and ConsistentlyReproductive Health Survey: Albania, 2002

Contraceptive Method*	Never Heard of Method	Heard of Method	No. of Cases	Very Effective	Effective	Some- what Effective	Not Very Effective	Do Not Know	Total	No. of Cases***
Pill	67.3	32.7	1740	12.5	52.0	12.2	1.8	21.4	100.0	698
Injectables	93.6	6.4	1740	**	**	**	**	**	**	149
Tubal Ligation	85.2	14.8	1740	79.7	14.2	1.4	1.4	3.3	100.0	287
IUD	91.6	8.4	1740	**	**	**	**	**	**	202
Condom	11.4	88.6	1740	46.8	31.2	4.7	0.6	16.7	100.0	1601
Calendar	77.5	22.5	1740	13.3	56.6	23.0	4.0	3.1	100.0	426
Withdrawal	10.7	89.4	1740	42.4	37.7	16.2	1.9	1.9	100.0	1528

*Listed in descending order of contraceptive effectiveness when the method is used correctly and consistently (USFDA)

**Less than 10 percent of respondents heard of method

***Opinion about contraceptive effectiveness asked only of men who have heard of method

CHAPTER 7 CONTRACEPTIVE USE AND PREFERENCES

Family planning is one of the most important components of reproductive health and maternal and child health programs. Use of modern methods of contraception influences both fertility and abortion rates and is also known to improve infant and child survival through birth spacing by at least two-year intervals.

In this chapter we examine the use of family planning in the Albanian reproductive age population and describe the sociodemographic characteristics that are associated with its use. We group family planning methods into two broad categories, modern methods and traditional methods. Modern methods consist of the oral pill, the condom, the IUD, implants, injectables, vaginal methods, and female and male sterilization (i.e., tubal ligation and vasectomy). Periodic abstinence withdrawal (rhythm) and comprise traditional methods (V Zlidar et al, 2003).

Current Contraceptive Use

Table 7.1 presents data from recent demographic and reproductive health surveys conducted in Eastern Europe and the former Soviet Union republics. The Eastern European countries, including Albania, show a relatively high prevalence of contraceptive use among married women. However, for unmarried women, Albania departs from the Eastern European pattern and demonstrates exceptionally low contraceptive prevalence for previously married (5%) and never married (4%)women, similar to rates found in the Caucasus region. This reflects the fact that more than 90% of the unmarried female population reported that they were not sexually active at the time of the survey (see Chapter 8).

Since the majority of sexually active women in Albania are married and because, by convention, most international contraceptive prevalence data are reported for married women, most of the data in this chapter will be confined to married women and married men (V Zlidar et al., 2003; R Salem, 2004).

In Tables 7.2A and 7.2B, the reported current practice of contraception is relatively high and very similar for women and men, at 75% for married women and 77% for married men. However, these rates are almost entirely due to the use of the traditional method of withdrawal (67% and 73% for married women and men, respectively). Given the history and geographic location of Albania, it may not be surprising that withdrawal is the most prominent method of contraception used (G Santow, 1993; H Goldberg and A Toros, 1994; Y Tountas et al., 2004). Overall, married men report a lower level of modern contraceptive use than women - only 3% compared to the 8% reported by married women. Previously and never married women exhibit very low levels of use, presumably reflecting low levels of sexual activity, as mentioned above. Among males who are not currently married, contraceptive use is somewhat higher than among unmarried females (14% for never married men), again more likely a reflection of higher sexual activity among men in this category compared to women.

Only 8.0% of married women report using a modern method. There are two recent studies to which we can compare this result. A baseline survey, also in 2002, of basic health service utilization, expenditures and quality, in three districts (Berat, Kucova and Fier) yielded a modern contraceptive prevalence rate of 5.3% among married women (Abt, 2004). The upper limit of the 95% confidence interval is 6.4%, somewhat smaller than the lower limit of the RHS 95% confidence interval of 6.9%; the lower prevalence found in these three districts may be expected as the RHS includes metro Tirana with the highest geographic prevalence rate of 14.9%. The second survey was a UNICEF Multiple Indicator Cluster Survey (MICS) conducted in 2000. In that survey, a modern contraceptive prevalence rate of 15% was reported (95% CI = 13.5%, 16.5%), almost twice as high as the RHS result two vears later (UNICEF and INSTAT, 2000). A secondary analysis would be needed to determine if the difference is due to methodological differences in the two surveys or a difference between two time periods.

Tables 7.3 A and 7.3 B show contraceptive use controlling for selected sociodemographic characteristics among married women and men. For married women (Table 7.3A), the use of any method has a small association with residence, with women in Metro Tirana reporting higher contraceptive prevalence than rural women (81% vs. 72%). There is a more pronounced and expected positive association with age, number of living children, and educational level of the woman. The low use at ages 15-19 (49%) and among women with no children (29%) is likely reflecting higher proportions of married women in those categories who are pregnant or seeking to become pregnant (See Table 7.9A). When looking at use of traditional versus modern methods, again we observe low levels of use of modern contraception. Modern contraceptive use appears to have similar associations with age and number of living children as use of traditional methods. Additionally, strata demonstrates a strong

effect, with women in Metro Tirana three times as likely and women in urban areas twice as likely to use modern methods as women in rural areas (15%, 9% and 5%, respectively). There is no such effect of residence on traditional contraceptive use. Also, education and socioeconomic levels have a positive relationship with modern contraceptive use but little relation to use of traditional methods. Overall rates of modern use remain low in all categories of the demographic and socioeconomic characteristics shown. The highest percentages of married women using a modern method are 18% and 17% for women with a high socioeconomic level and with postsecondary education, respectively.

Among married men (Table 7.3B), there is no strong association between use of any method and stratum of residence. Men in Tirana report somewhat lower overall use than rural men or men in other urban areas. Furthermore, associations between use of any method and age, number of living children, and the man's educational level are much weaker than observed among the women or do not exist. Use of modern methods, however, is associated with residence, with 6% of urban men (7% in Tirana) versus less than 1% of rural men reporting modern contraceptive use, with educational level (increasing from 1% to 13% with increasing education), and with socioeconomic status (from 1% to 4% to 12% across the three SES categories). Only men living in Tirana (10%), men with one child (12%), men with a post-secondary education (17%), and men classified as high SES (14%) report at least 10% of their method use to be modern methods.

Table 7.4A shows the distribution of methods used by married women in greater detail. It is quickly evident that the principal modern method used by the few

Albanian women using modern methods is tubal ligation. Five percent of married Albanian women using contraception have been sterilized, and this percentage increases to 10% among women 40-44 and 8% among women with three or more children. The second most used modern method is the condom, whose highest prevalence is 10% among contracepting women with postsecondary education and 11% among women of high socioeconomic status.

Modern method use among married men is negligible at 3%. In Table 7.4B, we see that for these men the condom is by far the principal modern method used. Condom use is positively associated with urban residence, educational level and socioeconomic status. Among contraceptors, the highest percentage of condom use is 9% for married men residing in Tirana, 12% for those with postsecondary education, and 8% for men in the highest socioeconomic stratum.

While use of modern contraception increases with education, socioeconomic status and urbanization for both women and men, modern use is higher for women at each educational level and residence.

Source of Contraception

Because of the small numbers of married menusing modern methods of contraception, information on source of modern methods can only be shown for the married women (Table 7.5). The data show that almost all tubal ligations (99%) are obtained from government hospitals and clinic facilities and the vast majority of condoms (78%) are obtained from pharmacies. Fifty-eight percent of women who use the pill reported the pharmacy as their source, and a third reported family planning clinics (20%) and health posts (12%). The few women who use IUDs primarily obtain them from government hospitals (71%), with another 13% reporting family planning clinics as their source.

Among all women currently using modern contraceptives, approximately two-thirds (65%) reported they had received advice from a physician (Table 7.6). Physician advice is primarily present for women who are using tubal ligation or the IUD. Condom users are more likely to have obtained advice from their partner, and pill users obtained advice from physicians, nurses, midwives or pharmacists. Among women who did receive counseling from a physician or other health worker, only two out of five (41%) reported having received information about other contraceptive methods and about half (53%) reported having received information about possible side effects.

Satisfaction with and Preference for Current Method

The vast majority of users of contraception report they are satisfied with the method they use (data not shown). Almost 100% of married women and men using withdrawal or condoms indicate they are satisfied with their method. Women who have been surgically sterilized report the least satisfaction at 78%. Only 5% of women and 3% of men currently using withdrawal said they would prefer another method.

Women and men who were using traditional methods (predominantly withdrawal) were read a list of reasons why a person might prefer these traditional or "natural" methods over modern ones, and were asked to rank their importance on a three-point scale from not important to somewhat important to very important. Tables 7.7A and B show the percentage who indicated the given reasons to be somewhat or very important. In Table 7.7A, among the reasons provided, fear of side effects was by far the most important to the female respondents (84% reported somewhat or very important). Second among the given reasons was that the method used was the partner's preference (73%). And the third ranking reason was lack of knowledge of modern methods (64%). Almost half of the women said that access and cost (45%)each) were somewhat or very important factors. Religious beliefs and doctor's or other person's advice were significantly less important. The most important reasons (fear of side effects and partner's preference) sociodemographic varied little by However, access, cost characteristics. and lack of knowledge were substantially more important in rural compared to urban areas and among the least educated and the lowest socioeconomic level. There is no significant variation for each reason among the three main religious groups.

Among the male respondents (Table 7.7B), the most frequently ranked factor for deciding not to use a modern method was "partner's preference" at 87%. This is consistent with results of a recent study in Turkey (A Kulczycki, 2004). Second in frequency were lack of knowledge of modern methods (76%) and fear of side effects (73%). More than half of the males also ranked cost (69%), difficult access (64%), doctor's recommendation (61%), and religious beliefs (57%) as somewhat or very important. There is little association between sociodemographic characteristics and reasons for not using modern methods among the men. Cost, access and knowledge varied somewhat or not at all with residence and educational level in comparison to the women. On the other hand, high socioeconomic status was strongly associated with the five factors most frequently ranked as important, and rural residence was strongly associated with the ranking of religious beliefs as an important factor.

When asked about the effectiveness of their current method, almost three-quarters (73%) of women using traditional methods said the method they were using was more effective than a modern contraceptive method (Table 7.8A). Another 10% believed their method was equally effective as modern methods. Even traditional method users with postsecondary education or a high socioeconomic level believed in the effectiveness of their method. Only 10% of women in these sociodemographic categories thought their current traditional method to be less effective than a modern method.

Men also were not convinced that modern methods were more effective than traditional methods (Table 7.8B), although their responses did differ slightly from the women. Almost half (48%) of the men believed their current traditional method was more effective than modern methods, but another 42% said it was equally effective. While education had no significant effect on men's perception of the effectiveness of methods, as with the women, men in the highest socioeconomic category were most likely to think that modern methods were more effective (still at only 11%).

There is very low knowledge of the true failure rate for withdrawal compared with modern methods such as the IUD and oral contraceptives as shown by recent surveys conducted in nine countries of Eastern Europe and the Former Soviet Union (CDC and MACRO, 2004). Results from these surveys show that the 12 month failure rate for withdrawal use ranges from 12% to 30 %, and from 17% to 30% in the four countries where withdrawal is the method most used, compared with the 12 month failure rates for IUD and oral contraceptives varying between 1% and 16%.

Reasons for Not Using Contraception

Tables 7.9 (A & B) and 7.10 (A & B) provide information on married women and men who are not currently using contraception. In Table 7.9A & B, we can readily observe that the vast majority of married nonusers are not using because they are not sexually active (husbands or wives may be absent)--55% for both women and men. Another 36% of women and 22% of men report they (or their wives) are either trying to get pregnant, are currently pregnant or are postpartum. Among women, these are mostly 15-24 year olds (66%). Women in the oldest age group (35-44) also report hysterectomy or other sources of infecundity (16%), and men in the oldest age group often (16%) report dissatisfaction with contraception due to its interruption of lovemaking - likely with reference to the method of withdrawal.

Intention to Use Contraception among Nonusers

Only fecund married women and men are presented in Tables 7.10A & B and Table 11. Table 7.10A demonstrates that those women who may not be currently using due to an absent spouse or pregnancyrelated reasons are definitely thinking about using contraception in the future. It appears that the majority (80%) of nonusing married women want to practice family planning. There is little association with parity, except for the nulliparous women who report the least interest in practicing family planning; they are more likely to be 15-24 years of age and, as shown in Table 7.9A, were the group most likely to be trying to get pregnant. Thirtyone percent of nonusers with no children report they do not want to practice family planning. Men (Table 7.10B), on the other hand, are less likely than women to want to practice family planning in the future. Twenty percent of men do not want to use contraception and 37% are undecided. The percentage that does not want to use decreases with parity, while the undecided responses increase with parity. In Table 7.11, more than half (57%) of the married women currently not using contraception and who want more children also want to use a contraceptive method within the next 12 months and only 14% do not want to use a method. The women who do not want more children are more likely (69%) to want to use a method within the coming 12 months and less likely to not want to use a method.

Table 7.1

Percent of Women Currently Using Any Contraceptive Method by Marital Status Among Women Aged 15–44 Reproductive and Demographic Health Surveys (RHS and DHS) In Selected Eastern European and Former Soviet Union Countries Albania Reproductive Health Survey 2002, Final Report

			Marital Status		_
Region and Country	Total	Currently Married	Previously Married	Never Married	Number of Cases
Eastern Europe					
Albania, 2002	51	75	5	4	5,697
Czech Rep., 1993	59	69	46	38	4,497
Moldova, 1997	54	74	27	7	5,412
Romania, 1999	48	64	20	20	6,888
Russia, 1999*	59	73	42	29	6,004
Ukraine, 1999	54	68	35	22	7,128
Caucasus					
Armenia, 2000	38	61	2	0	5,624
Azerbaijan, 2001	32	55	2	0	7,668
Georgia, 1999	25	41	3	0	7,798
Central Asia					
Kazakhstan, 1999	45	62	40	8	4,267
Kyrgyz Rep., 1997	42	60	15	1	3,529
Turkmenistan, 2000	34	55	18	0	7,263
Uzbekistan, 1996	40	57	11	0	4,091

* Yekaterinburg, Perm, and Ivanovo, respectively (predominantly urban sample).

Source: Goldberg H et al., 1993, KIIS and CDC, 2001; MACRO International 1995-2001; Serbanescu et al., 1998, 2001, 2001, 2003; VCIOM and CDC, 1998, 2000.

Table 7.2 A **Current Use of Contraception Among Women** 15-44 Years of Age, by Marital Status (Percent Distribution) **Reproductive Health Survey: Albania, 2002**

			Marital Status	
Use of Contraception	Total	Married	Previously Married	Never Married
Currently Using	50.5	75.1	5.1	4.4
Modern Methods	5.6	8.0	2.7	1.0
Tubal Ligation	2.7	4.0	2.4	0.1
Condom	1.6	2.1	0.3	0.7
PIII	0.7	1.0	0.0	0.2
IUD	0.3	0.5	0.0	0.0
Injectables	0.3	0.4	0.0	0.0
Traditional Methods	44.8	67.1	2.4	3.4
Periodic Abstinence (rhythm)	0.0	0.0	0.0	0.0
Withdrawal	44.8	67.1	2.4	3.4
Not Current Using	49.5	24.9	94.9	95.6
Total	100.0	100.0	100.0	100.0
No. of Cases	5,697	3,965	88	1,644

Table 7.2 B Current Use of Contraception Among Men 15-49 Years of Age, by Marital Status (Percent Distribution) Reproductive Health Survey: Albania, 2002

			Marital Status	
Use of Contraception	Total	Married	Previously Married	Never Married
Currently Using	52.3	77.1	**	14.2
Modern Methods	4.2	3.3	**	5.4
Condom	3.6	2.4	**	5.4
Pill	0.2	0.3	**	0.0
IUD	0.2	0.3	**	0.0
Tubal Ligation	0.1	0.1	**	0.0
Injectables	0.1	0.2	**	0.0
Traditional Methods	48.2	73.9	**	8.8
Periodic Abstinence (Rhythm)	0.6	0.8	**	0.3
Withdrawal	47.6	73.1	**	8.5
Not Currently Using	47.7	22.9	**	85.8
Total	100.0	100.0	**	100.0
No. of Cases	1,740	1,023	14	703

**Percentages are not shown when base is less than 25 cases.

Table 7.3 A Percentage Currently Using Modern and Traditional Methods by Selected Characteristics Currently Married Women Aged 15-44 Years Reproductive Health Survey: Albania, 2002

		All Ma	Contrace	Contraceptors		
Characteristics	Any Method	Modern Method	Traditional Method	No. of Cases	Percentage Using a Modern	No. of Cases
Total	75.1	8.0	67.1	3,965	10.7	3,065
Strata						
Metro Tirana	81.3	14.9	66.4	1,438	18.3	1,177
Other Urban	77.2	9.4	67.7	1,308	12.2	1,015
Other Rural	72.2	5.1	67.0	1,219	7.1	873
Residence						
Urban	78.6	11.3	67.4	2,488	14.4	1,993
Rural	72.4	5.5	67.0	1,477	7.6	1,072
Age Group						
15-19	48.9	2.5	46.4	97	5.1	52
20-24	62.4	3.7	58.7	502	5.9	324
25-29	67.4	5.3	62.1	800	7.9	567
30-34	78.7	9.4	69.3	1,004	11.9	812
35-39	81.9	9.3	72.6	906	11.4	756
40-44	82.7	10.9	71.8	656	13.2	554
Living Children						
0	29.3	2.2	27.1	291	7.5	91
1	60.9	3.1	57.8	800	5.1	514
2	82.7	9.1	73.6	1,806	11.0	1,544
3	85.5	10.6	74.9	783	12.4	681
4 +	81.5	10.8	70.7	285	13.3	235
Education Level						
Primary or Less	71.0	5.8	65.2	1,821	8.2	1,310
Secondary Incomplete	72.3	9.5	62.7	237	13.1	179
Secondary Complete	81.3	9.4	72.0	1,487	11.6	1,230
Post-Secondary	81.2	17.2	64.0	420	21.2	346
Socioeconomic Index						
Low	76.0	6.2	69.8	1,429	8.2	1,093
Medium	73.1	7.9	65.2	2,011	10.8	1,528
High	82.1	17.8	64.3	525	21.7	444

		All M	Contraceptors				
Characteristics	Any Method	Modern	Traditional	No. of Cases	Percentage Using a Modern Method	No. of d Cases	
Total	77.1	3.3	73.9	1,023	4.3	797	
Strata							
Metro Tirana	72.2	7.3	64.9	421	10.1	314	
Other Urban	80.3	5.2	75.2	318	6.5	257	
Other Rural	77.2	0.6	76.6	284	0.8	226	
Residence							
Urban	78.1	6.3	71.8	670	8.1	526	
Rural	76.3	0.7	75.6	353	0.9	271	
Age Group							
20-24	67.7	0.0	67.7	32	**	21	
25-29	73.9	2.1	71.8	123	2.8	91	
30-34	82.2	5.6	76.6	215	6.8	176	
35-39	84.1	2.6	81.6	242	3.1	204	
40-44	76.7	3.0	73.7	270	3.9	205	
45-49	67.8	3.3	64.5	141	4.9	100	
Living Children							
0	45.9	2.7	43.2	105	5.9	51	
1	79.0	9.3	69.7	218	11.8	169	
2	83.0	2.3	80.8	464	2.8	393	
3	84.0	1.4	82.6	167	1.7	136	
4 +	68.7	0.5	68.3	69	0.7	48	
Education Level							
Primary or Less	75.0	0.7	74.3	408	0.9	314	
Secondary Incomplete	**	**	**	20	**	13	
Secondary Complete	80.6	4.4	76.2	458	5.5	363	
Post-Secondary	77.1	12.8	64.2	137	16.6	107	
Socioeconomic Index							
Low	77.7	1.2	76.5	403	1.5	319	
Medium	75.6	3.8	71.8	458	5.0	349	
High	80.6	11.5	69.1	162	14.3	129	

Table 7.3 BPercentage Currently Using Modern and Traditional Methods by Selected Characteristics
Currently Married Men Aged 15-49 Years
Reproductive Health Survey: Albania, 2002

** Percentages are not shown when base is less than 25 cases

ab es

Table 7.4 ACurrent Use of Specific Contraceptive Methods by Selected CharacteristicsCurrently Married Women Aged 15-44 YearsReproductive Health Survey: Albania, 2002

			Specific Contraceptive Method Use (Percent Distribution)								
Characteristics	Any Method	No. of Cases	Tubal Ligation	Condom	Pill	IUD	Injectables		Periodic Abstinence Rhythm	Total	No. of Cases
Total	75.1	3,965	5.3	2.8	1.3	0.6	0.5	89.3	0.0	100.0	3,065
Strata											
Metro Tirana	81.3	1,438	6.6	6.8	3.4	1.2	0.3	81.6	0.1	100.0	1,177
Other Urban	77.2	1,308	5.4	3.5	1.5	0.8	1.1	87.7	0.1	100.0	1,015
Other Rural	72.2	1,219	4.9	1.2	0.5	0.3	0.3	92.9	0.0	100.0	873
Residence											
Urban	78.6	2,488	5.8	4.7	2.0	0.9	0.8	85.6	0.1	100.0	1,993
Rural	72.4	1,477	5.0	1.3	0.7	0.3	0.3	92.5	0.0	100.0	1,072
Age Group											
15-19	48.9	97	0.0	2.6	0.8	0.0	1.8	94.9	0.0	100.0	52
20-24	62.4	502	0.7	4.0	0.2	0.9	0.1	94.0	0.0	100.0	324
25-29	67.4	800	1.9	2.9	2.1	0.6	0.3	92.1	0.0	100.0	567
30-34	78.7	1,004	5.5	3.6	1.8	0.5	0.5	88.1	0.0	100.0	812
35-39	81.9	906	6.0	2.4	1.4	0.9	0.6	88.5	0.2	100.0	756
40-44	82.7	656	9.5	2.0	0.6	0.3	0.7	86.8	0.0	100.0	554
Living Children											
0	29.3	291	0.0	7.4	0.0	0.0	0.0	92.6	0.0	100.0	91
1	60.9	800	1.2	2.7	1.1	0.0	0.1	94.9	0.0	100.0	514
2	82.7	1,806	4.6	3.5	2.0	0.7	0.2	88.9	0.1	100.0	1,544
3	85.5	783	8.3	1.6	0.7	0.8	1.0	87.6	0.0	100.0	681
4 +	81.5	285	8.2	2.2	0.6	0.7	1.5	86.7	0.0	100.0	235
Education Level											
Primary or Less	71.0	1,821	5.1	1.5	0.5	0.4	0.7	91.8	0.0	100.0	1,310
Secondary Incomplete	72.3	237	9.3	0.8	1.2	1.0	0.9	86.8	0.0	100.0	179
Secondary Complete	81.3	1,487	5.3	3.6	1.7	0.7	0.2	88.5	0.0	100.0	1,230
Post-Secondary	81.2	420	4.4	10.1	5.1	1.3	0.3	78.4	0.4	100.0	346
Socioeconomic Index											
Low	76.0	1,429	5.7	1.0	0.6	0.2	0.6	91.9	0.0	100.0	1,093
Medium	73.1	2,011	5.0	3.0	1.6	0.7	0.5	89.1	0.1	100.0	1,528
High	82.1	525	5.2	10.8	3.0	2.0	0.6	78.2	0.1	100.0	444

Table 7.4 BCurrent Use of Specific Contraceptive Methods by Selected Characteristics
Currently Married Men Aged 15-49 Years
Reproductive Health Survey: Albania, 2002

		-	Specific Contraceptive Method Use (Percent Distribution)								
Characteristics	Any Method	No. of Cases	Condom	Pill	ani	Injectables	Tubal Ligation	Withdrawal	Periodic Abstinence (Rhythm)	Total	No. of Cases
Total	77.1	1,023	3.1	0.3	0.4	0.3	0.1	94.8	1.0	100.0	797
Strata Metro Tirana Other Urban	72.2 80.3	421 318	8.7 3.6	0.5 0.7	0.3 1.3	0.0 0.8	0.6	88.5 91.8	1.4 1.8	100.0 100.0	314 257
Other Rural	77.2	284	0.8	0.0	0.0	0.0	0.0	98.8	0.4	100.0	226
Residence Urban Rural	78.1 76.3	670 353	5.6 0.9	0.7 0.0	1.0 0.0	0.5 0.0	0.2 0.0	90.2 98.7	1.7 0.4	100.0 100.0	526 271
Age Group 20-24	67.7	32	**	**	**	**	**	**	**	**	21
25-29	73.9	123	2.9	0.0	0.0	0.0	0.0	96.3	0.9	100.0	91
30-34	82.2	215	5.0	1.0	0.0	0.5	0.2	91.6	1.5	100.0	176
35-39	84.1	242	1.9	0.5	0.7	0.0	0.0	95.9	1.1	100.0	204
40-44	76.7	270	3.9	0.0	0.0	0.0	0.0	95.6	0.5	100.0	205
45-49	67.8	141	1.9	0.0	1.7	0.9	0.4	95.1	0.0	100.0	100
Living Children	45.9	105	5.8	0.0	0.0	0.0	0.0	90.7	3.4	100.0	51
1	79.0	218	9.2	1.4	0.6	0.6	0.0	87.2	1.1	100.0	169
2	83.0	464	1.6	0.1	0.8	0.0	0.3	96.2	1.1	100.0	393
3	84.0	167	1.0	0.0	0.0	0.7	0.0	97.8	0.5	100.0	136
4 +	68.7	69	0.7	0.0	0.0	0.0	0.0	99.3	0.0	100.0	48
Education Level Primary or Less	75.0	408	0.9	0.0	0.0	0.0	0.1	98.1	0.9	100.0	314
Secondary Incomplete	**	20	**	**	**	**	**	**	**	**	13
Secondary Complete	80.6	458	4.0	0.1	0.8	0.6	0.0	93.4	1.1	100.0	363
Post-Secondary	77.1	137	11.5	3.2	1.3	0.0	0.7	82.8	0.6	100.0	107
Socioeconomic Index											
Low	77.7	403	1.4	0.0	0.0	0.0	0.1	98.3	0.2	100.0	319
Medium	75.6	458	4.0	0.4	0.0	0.6	0.0	92.9	2.0	100.0	349
High	80.6	162	7.9	1.6	4.2	0.0	0.6	84.8	1.0	100.0	129

** Percentages are not shown when base is less than 25 cases

20

Ľ 2

Table 7.5Source of Supply for Last Modern Contraceptive Used Among Currently Married WomenAged 15-44 Years Currently Using a Modern Method by Specific Methods(Percent Distribution)Reproductive Health Survey: Albania, 2002

			Method Currently Using				
Source	Total	Tubal Ligation	Condom	Pill	IUD	Injectables	
Public Medical Sector	65.9	99.4	8.2	35.8	91.0	**	
Gov. Hospital-Maternity Ward	56.1	97.2	0.0	4.0	70.8	**	
Familiy Planning Clinic	5.7	0.0	4.0	19.8	13.4	**	
Health Post or Clinic	4.1	2.2	4.2	12.0	6.8	**	
Pharmacy	29.0	0.0	78.2	58.1	4.6	**	
Private Clinic or Office	0.5	0.0	0.0	1.0	4.4	**	
Partner/Husband	3.2	0.0	12.0	0.0	0.0	**	
Do Not Know	1.4	0.5	1.7	5.2	0.0	**	
TOTAL	100.0	100.0	100.0	100.0	100.0	**	
No. of Cases	410	169	130	67	28	16	

** Percentages are not shown when base is less than 25 cases.

Table 7.6 Percent Distribution of Type of Counseling by a Health Care Provider for Current Contraceptive Method Among Married Women Aged 15-44 Currently Using a Modern Method Reproductive Health Survey: Albania, 2002

		I			
Who Advised User	Total	Tubal Ligation	Condom	Pill	IUD
Physician	65.3	93.6	14.4	54.1	88.8
Partner	16.5	0.2	58.7	2.3	0.0
Nurse/Midwife	4.0	1.2	2.3	15.7	6.7
Friend	2.5	1.0	1.8	7.9	0.0
Pharmacist	2.0	0.0	2.7	10.5	0.0
Mother or Other Relative	1.9	0.0	1.9	2.1	4.5
No One	7.8	4.0	18.1	7.4	0.0
Total	100.0	100.0	100.0	100.0	100.0
No. of Cases	405	168	129	64	28
Type of Counseling by Physician, Nurse or Midw	vife				
General Information About Other Methods	40.7	25.6	**	67.9	77.4
Information About Possible Side Effects	52.5	44.9	**	68.0	73.3
No. of Cases	253	153	17	46	27

** Percentages are not shown when base is less than 25 cases

Table 7.7 A

Percentage of Contraceptive Users Who Stated that Selected Reasons Read to Them Were Very Important or Somewhat Important When Deciding To Use a Non-Supplied Method Instead of a Modern Method, by Selected Characteristics Women Aged 15-44 Currently Using Traditional Methods Reproductive Health Survey: Albania, 2002

Characteristics	Fear of or Experience With Side Effects	Partner Preference	Little Knowledge of Modern Methods	Difficult to Get a Modern Method	Cost of Modern Methods	Doctor Recomendation	Another Person's Advice	Religious Beliefs	No. of Cases
Total	84.1	73.2	64.2	45.4	44.7	33.0	23.4	21.5	2,739
Method Currently Using									
Rhythm Method	**	**	**	**	**	**	**	**	2
Withdrawal	84.2	73.2	64.1	45.5	44.7	33.0	23.4	21.5	2,737
Strata									
Metro Tirana	81.1	75.7	59.4	37.8	39.0	32.1	20.3	18.0	1,012
Other Urban	86.7	68.3	57.0	38.0	39.4	28.0	21.2	17.0	910
Other Rural	83.5	75.3	69.8	52.1	49.5	36.2	25.6	25.2	817
Residence									
Urban	85.0	70.6	57.6	37.7	38.8	28.4	20.0	16.5	1,750
Rural	83.4	75.2	69.4	51.7	49.4	36.7	26.0	25.5	989
Age Group									
15-24	85.1	74.0	63.3	45.4	43.9	31.0	23.8	19.8	407
25-34	83.7	73.8	64.2	45.7	44.6	31.1	22.2	19.6	1,216
35-44	84.2	72.4	64.4	45.2	45.1	35.3	24.3	23.7	1,116
Education Level									
Primary or Less	81.9	75.3	69.7	50.9	49.3	34.6	24.4	23.8	1,189
Secondary Incomplete	92.3	77.3	66.9	39.8	34.5	21.0	16.1	19.1	165
Secondary Complete	86.0	70.0	59.1	40.8	41.4	34.3	24.6	20.9	1,093
Post-Secondary	85.9	69.7	46.0	31.7	35.0	24.5	15.7	9.8	292
Socioeconomic Index									
Low	80.7	73.7	68.2	50.8	49.6	34.1	24.3	23.7	989
Medium	86.4	71.5	62.4	41.9	41.7	31.7	22.9	19.4	1,386
High	89.4	79.6	52.4	37.0	35.5	33.9	21.3	21.3	364
Religion									
Muslim	83.2	72.9	64.3	46.4	44.9	31.5	23.3	20.6	2,225
Orthodox	87.1	75.3	59.0	41.0	43.1	34.2	25.5	21.4	246
Catholic	89.4	73.9	69.3	44.8	46.4	42.6	22.4	27.6	240
Other/Undeclared	71.1	68.7	30.3	10.7	21.4	28.4	19.8	19.0	28

** Percentages are not shown when base is less than 25 cases

Contraceptive Use and Preferences

Table 7.7 B

Percentage of Contraceptive Users Who Stated That Selected Reasons Read to Them Were Very Important or Somewhat Important When Deciding To Use a Non-Supplied Method Instead of a Modern Method, by Selected Characteristics Men Aged 15-49 Currently Using Traditional Methods Reproductive Health Survey: Albania, 2002

Characteristics	Partner Preference	Little Knowledge of Modern Methods	Fear of or Experience With Side Effects	Cost of Modern Methods	Difficult to Get a Modern Method	Doctor Recommendations	Another Person's Advice	Religious Beliefs	Total	No. of Cases
Total	86.9	75.9	72.6	68.5	64.2	60.8	38.7	56.8	100.0	804
Method Currently Using Rhythm Method Withdrawal	** 86.8	** 75.9	** 72.5	** 68.5	** 64.2	** 60.6	** 38.9	** 56.7	** 100.0	13 791
Strata Metro Tirana Other Urban Other Rural	84.4 84.7 89.1	73.9 72.5 78.6	72.7 72.5 72.5	65.1 65.7 71.3	66.1 58.0 67.3	53.1 59.1 64.4	27.5 38.1 42.7	40.7 54.8 63.3	100.0 100.0 100.0	300 271 233
Residence Urban Rural	84.3 89.1	72.5 78.7	72.6 72.5	65.1 71.3	60.7 67.1	57.6 63.5	34.7 42.0	49.5 63.0	100.0 100.0	526 278
Age Group 15-24 25-34 35-49	90.3 91.2 84.1	82.0 79.0 73.5	79.5 77.0 69.3	84.5 71.2 65.3	71.2 73.1 58.4	59.3 67.8 57.0	42.5 41.6 36.6	62.2 61.7 53.5	100.0 100.0 100.0	43 277 484
Education Level Primary or Less Secondary Incomplete Secondary Complete Post-Secondary	84.7 ** 90.2 87.1	77.4 ** 76.6 65.5	70.8 ** 74.3 75.5	67.6 ** 69.6 66.7	62.2 ** 67.7 60.7	59.6 ** 62.4 60.3	39.6 ** 35.3 45.5	60.2 ** 53.6 51.3	100.0 ** 100.0 100.0	324 14 364 102
Socioeconomic Index Low Medium High	82.6 89.7 97.0	75.9 74.1 82.0	65.7 76.7 90.3	65.7 68.4 81.7	57.5 68.2 81.3	55.3 64.6 73.0	36.8 37.7 50.8	52.7 60.3 63.1	100.0 100.0 100.0	321 353 130
Religion Muslim Orthodox Catholic Other/Undeclared	87.3 82.6 82.3 **	75.1 74.7 83.1 **	71.3 73.3 85.0 **	66.9 75.3 74.3 **	62.9 67.4 69.8 **	59.2 63.5 71.8 **	38.3 39.3 31.6 **	54.7 63.2 67.9 **	100.0 100.0 100.0 **	675 80 32 17

** Percentages are not shown when base is less than 25 cases.

ables

Table 7.8 A Perceived Effectiveness of Traditional Methods Compared to Modern Methods by Selected Characteristics Women Aged 15-44 Currently Using Traditional Methods (Percent Distribution) Reproductive Health Survey: Albania, 2002

	Ті	aditional Met	hod vs Modern			
Characteristics	Current Method More Effective	About Equally Effective	Current Method/Less Effective	Do Not Know	Total	No. of Cases
Total	73.3	10.2	5.5	11.1	100.0	2,739
Strata						
Metro Tirana	72.8	13.2	7.8	6.2	100.0	1,012
Other Urban	72.4	11.7	5.7	10.2	100.0	910
Other Rural	73.9	8.3	4.6	13.1	100.0	817
Residence						
Urban	71.9	12.6	6.4	9.1	100.0	1,750
Rural	74.4	8.2	4.7	12.7	100.0	989
Age Group						
15-24	73.4	11.0	4.7	10.9	100.0	407
25-34	71.6	9.9	6.5	12.0	100.0	1,216
35-44	74.6	10.1	4.8	10.4	100.0	1,116
Education Level						
Primary or Less	74.4	7.4	4.8	13.4	100.0	1,189
Secondary Incomplete	74.3	15.3	3.3	7.1	100.0	165
Secondary Complete	72.3	12.0	5.8	9.9	100.0	1,093
Post-Secondary	68.7	17.5	10.5	3.3	100.0	292
Socioeconomic Index						
Low	73.2	7.6	4.7	14.5	100.0	989
Medium	74.8	11.2	5.3	8.6	100.0	1,386
High	65.2	17.5	10.2	7.1	100.0	364

Table 7.8 B Perceived Effectiveness of Traditional Methods Compared to Modern Methods by Selected Characteristics Men Aged 15-49 Currently Using Traditional Methods (Percent Distribution) Reproductive Health Survey: Albania, 2002

	Tra	Traditional Method vs. Modern						
Characteristics	Current Method More Effective	About Equally Effective	Current Method/ Less Effective	Do Not Know	Total	No. of Cases		
Total	47.7	41.5	4.6	6.1	100.0	804		
Strata								
Metro Tirana	48.7	32.8	7.1	11.4	100.0	300		
Other Urban	45.2	43.1	5.3	6.4	100.0	271		
Other Rural	49.0	43.4	3.4	4.2	100.0	233		
Residence								
Urban	46.5	39.5	6.0	7.9	100.0	526		
Rural	48.8	43.2	3.5	4.6	100.0	278		
Age Group								
15-24	21.7	68.3	7.4	2.5	100.0	43		
25-34	41.3	47.1	6.4	5.2	100.0	277		
35-49	54.0	35.7	3.4	7.0	100.0	484		
Education Level								
Primary or Less	50.3	38.8	3.6	7.3	100.0	324		
Secondary Incomplete	**	**	**	**	**	14		
Secondary Complete	48.4	41.3	5.9	4.4	100.0	364		
Post-Secondary	35.6	56.8	5.6	2.0	100.0	102		
Socioeconomic Index								
Low	52.5	35.8	3.0	8.6	100.0	321		
Medium	44.5	46.6	5.1	3.8	100.0	353		
High	36.9	49.7	10.6	2.8	100.0	130		

ab es

Table 7.9 A Most Commonly Cited Reasons for Not Currently Using Contraception Among Currently Married Women Aged 15-44 Years (Percent Distribution) **Reproductive Health Survey: Albania, 2002**

		Age Group			
Reason For Not Using Contraception	Total	15-24	25-34	35-44	
Not Sexually Active/No Partner	57.6	43.6	61.9	63.5	
Currently Pregnant	17.7	31.8	20.0	3.4	
Trying to Get Pregnant	12.4	17.6	10.9	10.0	
Postpartum/Breastfeeding	3.6	6.0	4.3	0.7	
Infecundity/Subfecundity	6.4	0.3	1.3	17.7	
Religious Reasons	0.8	0.0	0.3	1.9	
Respondent Does Not Want to Use Contraception	0.5	0.0	0.4	0.9	
Respondent Did Not Think About Using Contraception	0.4	0.0	0.2	0.9	
Other Reasons	0.7	0.7	0.8	0.9	
Total	100.0	100.0	100.0	100.0	
No. of Cases	1,063	282	482	299	

Table 7.9 B Most Commonly Cited Reasons for Not Currently Using Contraception by Age Group Among Currently Married Men Aged 15-49 Years (Percent Distribution) **Reproductive Health Survey: Albania, 2002**

			Age		
Reason Not Using Contraception	Total	15-24	25-34	35-49	
Not Sexually Active/No Partner	55.3	90.1	46.8	41.3	
Trying to Get Partner Pregnant	14.2	4.8	29.2	7.8	
Lovemaking Interrupted	7.1	0.0	1.2	16.2	
Partner Currently Pregnant	4.2	1.7	8.2	2.4	
Partner Postpartum/Breastfeeding	4.0	0.8	6.5	3.8	
Infecundity/Subfecundity	8.8	0.0	4.1	17.9	
Other Reasons	6.3	2.5	4.0	10.7	
Fotal	100.0	100.0	100.0	100.0	
No. of Cases	392	94	138	160	

Table 7.10 A Desire to Use Contraception in the Future by Number of Living Children Fecund Currently Married Women Aged 15-44 Years Who Are Not Using Contraception (Percent Distribution) Reproductive Health Survey: Albania, 2002

		Number of Living Children *				
Desire to Use Contraception	Total	0	1	2	3 +	
Want to Use a Method Within 12 Months	62.2	29.3	60.3	70.8	70.2	
Want to Use a Method Later	18.2	27.0	21.4	14.7	14.1	
Do Not Want to Use Contraception	10.7	30.7	10.0	6.6	6.7	
Undecided	9.0	13.1	8.3	8.0	9.1	
Total	100.0	100.0	100.0	100.0	100.0	
No. of Cases	838	110	262	298	168	

* Women who were pregnant at the time of the inteview are classified as having one more child than the actual number

Table 7.10 B Desire to Use Contraception in the Future by Number of Living Children Fecund Currently Married Men Aged 15-49 Years Who Are Not Using Contraception (Percent Distribution) Reproductive Health Survey: Albania, 2002

		Number of Living Children *				
Desire to Use Contraception	Total	0	1	2	3 +	
Want to Use a Method Within 12 Months	25.0	7.7	37.5	31.2	23.0	
Want to Use a Method Later	18.2	37.6	27.9	11.1	5.8	
Do Not Want to Use Contraception	19.8	45.8	22.1	11.0	8.9	
Undecided	37.0	8.9	12.6	46.8	62.3	
Total	100.0	100.0	100.0	100.0	100.0	
No. of Cases	202	40	41	70	51	

* Men whose partner was pregnant at the time of the interview are classified as having one more child than the actual number

Albania Reproductive Health Survey, 2002 Final Report

ables

Table 7.11 Desire to Use Contraception in the Future by Fertility Preferences Fecund Currently Married Women Aged 15-44 Years Who Are Not Using Contraception (Percent Distribution) Reproductive Health Survey: Albania, 2002

Chapter 7

		Desire for Aditional Children					
Desire to Use Contraception	Total	Want More Children	No More Children	Undecided			
Want to Use a Method Within 12 Months	62.2	56.5	69.0	66.1			
Want to Use a Method Later	18.2	20.9	15.1	15.3			
Do Not Want to Use Contraception	10.7	14.2	5.9	11.9			
Undecided	9.0	8.5	10.0	6.7			
Total	100.0	100.0	100.0	100.0			
No. of Cases	838	444	329	65			

CHAPTER 8

NEED FOR CONTRACEPTIVE SERVICES AND CONTRACEPTIVE COUNSELING

The unmet need for contraception is a very specific estimate that measures the gap between desired fertility and the contraceptive practices adopted to ensure that fertility preferences are met in any given population. The conventional definition of unmet need includes women currently married or in consensual unions who are fecund, currently sexually active, currently exposed to the risk of pregnancy, not wanting to become pregnant, and not using any form of pregnancy prevention (Bongaarts, 1991). In this report, the standard formulation of unmet need is extended to all women, not just those in union.

In addition to the unmet need for any family planning method, the Albania RHS 2002 estimated the need for modern contraception – an indicator used in other Eastern European surveys that expanded the definition to include users of nonsupplied methods in the category of unmet need. In countries with high use of nonsupplied methods, such as withdrawal, the standard definition of unmet need masks the real need for more effective contraception because these methods tend to have higher failure rates (CDC and MACRO, 2003). For these countries it is more useful to also estimate the need for modern contraception in addition to any contraceptive method, despite the small risk of overstating the unmet need in some cases where traditional methods are used more effectively.

Potential Demand and Unmet Need for Family Planning Services

The survey asked all women about their sexual, contraceptive and reproductive


behaviors, as well as their fecundity status and fertility preferences, allowing for an examination of contraceptive need among both married and unmarried respondents

Table 8.1 presents data on women in need of family planning, or at risk of an unintended pregnancy, in Albania by age group and marital status. Women who are currently pregnant or postpartum, currently seeking to become pregnant, sub-fecund, or who are not sexually active, are considered to not be in need of family planning services at the time of the survey. About one-half (49%) of women fall into this category, ranging from 80% of young adults to 36% of 25-44 year old women to only 23% of 35-44 year old women. This category excludes almost all previously married (95%) and never married women (96%) because of lack of sexual activity or lack of sexual experience, respectively. Only one-fourth (24%) of married women are classified as "not currently in need of family planning services". Thus, threefourths (75%) of married women are considered to have potential need for family planning services, of whom 8% are current users of modern methods and 67% report using traditional methods, principally withdrawal. As mentioned above, the questionnaire design of the survey instrument allows an estimation of women in need of modern contraception. As shown in Table 8.1, 68% percent of married women are in need of modern contraception. This includes well over half of all women from 25-44 years of age.

Table 8.2 shows need of any method and modern methods by selected characteristics for all women and currently married women. Focusing on the two-thirds of married women (68%) in need of a modern method, there is not much variation in need by geographic area of residence, education level, or socioeconomic status. Only married adolescents and women with no children have a substantially lower risk of an unintended pregnancy due the high proportion of these women wanting a pregnancy.

Unmet need among fecund married women is shown in Table 8.3 and Figure 8.1 by future fertility preferences, ie. whether they want to space an additional child or do not want any more children. Knowledge of reproductive intentions can assist providers to help couples choose the correct contraceptive method that will allow them to control if and when to have a pregnancy. Approximately two-thirds (69%) of fecund married women with a risk of an unintended pregnancy do not want any more children. This includes over 75% of women with 2 or more children or from 35-44 years of age. Only married women with no children or one child and from 15-29 years of age have a greater proportion of those at risk of an unintended pregnancy that want to have a child or additional children.

Table 8.4 summarizes the percentage of the potential demand for modern contraceptive methods "satisfied" for married women by selected characteristics. The first column showing potential demand comes from Table 8.2 and the second column listing the prevalence of modern method use comes from Table 7.3A. The potential demand "satisfied" is the second column divided by the first column. For all married women, only 12% of the potential demand (women at risk of an unintended pregnancy) is satisfied (using a modern contraceptive to prevent an unintended pregnancy). This percentage compares with 54% to 79% of potential demand satisfied in other



countries in Eastern Europe with a recent reproductive health survey. In the Caucasus region, both Armenia and Georgia have 31% of potential demand for modern contraceptive methods met and Azerbaijan has 18% met (CDC and MACRO, 2003). Potential demand satisfied is higher in urban areas and increases with age, educational attainment, socioeconomic status and number of living children. Only women in Metropolitan Tirana (22%), women with a post-secondary education (26%) and women classified as higher socioeconomic status (27%) have reached a level of 20% or higher.

Contraceptive Counseling

In Eastern Europe and the Former Soviet Union countries. most reproductive health services have been provided by physicians, and in some cases only by obstetricians, who traditionally have received little training in providing client-oriented counseling. Albanian The RHS included a series of questions designed to capture interactions between family planning providers and their clients. Specifically, women who had used a modern contraceptive method or had an abortion during the five years prior to interview were asked about the extent to which health providers provided basic information.

Women who had used at least one modern contraceptive method in the previous five years were asked who advised them to use their most recent modern method. If the advice came from a health care provider (i.e., physician, nurse, etc.) they were asked whether they received any information about other methods, including their comparative effectiveness and possible side effects associated with their use. As shown in Table 8.5, almost twothirds (63%) of women were advised by a health care provider (Ob/Gyn, general practitioner or nurse) to use their current or most recent modern method. However, for clinical methods, such as female sterilization and the IUD, 91% and 81%, respectively, were advised by an Ob/Gyn. Three-fourths of pill users (74%) were advised by a health care provider or a pharmacist. Condoms were usually suggested by the partner. About one-half of women (53%) received information about possible side effects, 41% received information about other methods and only 35% received counseling about method effectiveness. Although sample size is small, a higher proportion of women using the IUD received counseling.

Women who terminate their pregnancies in abortion and do not adopt an effective contraceptive method afterwards are probably at high risk for another unintended pregnancy and represent an important group whose family planning needs are not satisfied. A wide range of contraceptive methods, together with accurate information and/or referral for ongoing family planning care, should be made available and accessible to all women who have undergone abortions. As discussed in Chapter 4, women in the national household sample significantly underreported their experience with legally induced abortion. However, it may be instructive to look at the data for those women reporting at least one abortion in the past five years who were asked if they received any contraceptive advice either before or after the abortion procedure and whether they

received any contraceptive method or prescription for a method. As shown in Table 8.6 and Figure 8.2, only a minority of Albanian women who reported having had an abortion received contraceptive counseling before their abortion (18%), after their abortion (15%) or both before and after (5%). Only one in six women (16%) actually received a method or a prescription for a method. This result appears to confirm a missed opportunity to provide services to women who have just terminated an unintended pregnancy.




Table 8.1Potential Demand For Family Planning (FP) Services by Age Group And Marital Status
(Percent Distributions and Percentage)
Reproductive Health Survey: Albania, 2002

		A	ge Grou	ıp		Marital Status	
Demand for Family Planning Services	Total	15-24	25-34	35-44	Married	Previously Married	Never Married
Women Not Currently in Need of FP Services	48.8	80.0	36.0	22.5	23.6	94.9	95.5
Never Had Sexual Intercourse	30.1	67.7	11.2	2.8	0.0	0.0	91.8
Not Currently Sexually Active *	11.1	5.9	15.2	13.1	12.2	91.9	3.6
Currently Pregnant or Post-Partum	3.6	4.3	5.5	0.7	5.4	2.5	0.0
Seeking to Get Pregnant	2.2	2.0	3.2	1.2	3.2	0.5	0.1
Infecund/Subfecund ‡	1.8	0.1	0.9	4.7	2.8	0.0	0.0
Potential Demand for FP Services	50.5	19.0	63.0	76.7	75.1	5.1	4.4
Met Need							
Current Users of a Modern Method	5.6	1.5	6.6	9.6	8.0	2.7	1.0
Current Users of a Traditional Method	44.9	17.6	56.3	67.1	67.2	2.4	3.4
Unmet Need							
Non Users at Risk of Unintended Pregnancy	0.8	0.9	0.9	0.7	1.2	0.0	0.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Unmet Need For Modern Contraception [†]	45.7	18.5	57.2	67.8	68.4	2.4	3.4
No. of Cases	5,697	2,030	2,013	1,654	3,965	88	1,644

* Within the past month

† Includes nonusers at risk of unintended pregnancy and current users of traditional contraceptive methods

\$ Sterilization surgery for noncontraceptive reasons, medical conditions that preclude pregnancy, infertile partners, and menopause Note: The use of withdrawal with a relatively high failure rate may indicate not that fertility control was achieved but only that it was desired (HIMES, 1970). Thus, the unmet need for modern contraception is presented for countries in Eastern Europe where withdrawal is a predominant method

(CDC and Macro, 2004).

Table 8.2Percentage of Women Aged 15-44 Years in Need of Any or Modern Contraceptive Methods
by Selected Characteristics
Reproductive Health Survey: Albania, 2002

		All Women		М	arried Wome	en
Characteristics	Any Method	Modern Method	No. of Cases	Any Method	Modern Method	No. of Cases
Total	0.8	45.7	5,697	1.2	68.4	3,965
Strata						
Metro Tirana	1.0	45.4	2,108	1.5	67.8	1,438
Other Urban	0.7	48.3	1,816	1.1	68.8	1,308
Other Rural	0.8	44.4	1,773	1.3	68.3	1,219
Residence						
Urban	0.8	47.4	3,572	1.1	68.5	2,488
Rural	0.8	44.4	2,125	1.3	68.3	1,477
Age Group						
15-19	0.3	6.9	1,094	2.8	49.2	97
20-24	1.5	32.7	936	3.1	61.8	502
25-29	1.1	51.5	946	1.4	63.5	800
30-34	0.7	63.0	1,067	0.8	70.1	1,004
35-39	0.7	68.4	958	0.8	73.4	906
40-44	0.6	67.1	696	0.6	72.5	656
Living Children						
0	0.7	7.1	1,943	5.2	32.3	291
1	1.9	56.9	828	2.0	59.8	800
2	0.7	72.0	1,840	0.7	74.3	1,806
3 +	0.6	72.6	1,086	0.6	74.1	1,068
Education Level						
Primary or Less	0.9	45.1	2,519	1.4	66.6	1,821
Secondary Incomplete	0.1	24.3	653	0.3	63.1	237
Secondary Complete	0.6	55.6	1,830	0.8	72.8	1,487
Post-Secondary	1.6	42.8	695	2.5	66.5	420
Socioeconomic Index						
Low	0.8	48.1	1,940	1.3	71.1	1,429
Medium	0.8	43.4	2,985	1.3	66.5	2,011
High	0.6	47.2	772	0.9	65.3	525

Table 8.3Percentage of Unmet Need for Contraception among Fecund Married Women
of Reproductive Age by Future Fertility Preferences *
Reproductive Health Survey: Albania, 2002

		net Need Fo Contracept	-		et Need For Contracept			Need For of Total
Characteristics	Total	Spacing	Limiting	Total	Spacing	Limiting	Limiting Any Method	Limiting Modern Method
Total	1.3	1.0	0.3	70.3	22.0	48.4	23.1	68.8
Strata Metro Tirana	1.5	1.2	0.3	69.5	24.8	44.7	20.0	64.3
Other Urban	1.1	0.7	0.4	70.8	20.0	50.8	36.4	71.8
Other Rural	1.3	1.0	0.3	70.3	22.3	48.0	23.1	68.3
Residence Urban Rural	1.2 1.4	0.9 1.0	0.3 0.3	70.4 70.3	21.1 22.6	49.3 47.6	25.0 21.4	70.0 67.7
Age Group 15-19	2.8	2.8	0.0	49.2	47.2	2.0	0.0	4.1
20-24	3.1	2.8	0.3	62.2	51.1	11.1	9.7	17.8
25-29	1.5	1.0	0.4	64.3	36.3	28.0	26.7	43.5
30-34	0.8	0.4	0.4	70.8	20.8	50.1	50.0	70.8
35-39	0.8	0.7	0.1	76.4	8.0	68.4	12.5	89.5
40-44	0.7	0.2	0.5	77.6	2.2	75.4	71.4	97.2
Living Children	5.8	5.5	0.3	36.1	34.6	1.6	0.0	4.4
0 1	2.0	5.5 2.0	0.0	60.9	53.1	7.8	0.0	4.4 12.8
2	0.8	0.4	0.0	76.0	17.9	7.8 58.1	37.5	76.4
3+	0.6	0.4	0.5	75.8	7.0	68.9	83.3	90.9
Education Level								
Primary or Less	1.4	1.1	0.4	68.5	22.8	45.7	28.6	66.7
Secondary Incomplete	0.3	0.3	0.0	65.6	26.2	39.4	0.0	60.1
Secondary Complete	0.9	0.5	0.3	74.9	19.6	55.2	33.3	73.7
Post-Secondary	2.6	2.4	0.1	67.7	22.8	44.9	3.8	66.3
Socioeconomic Index Low	1.3	1.0	0.3	73.2	22.2	51.0	23.1	69.7
Medium	1.3	0.9	0.4	68.4	21.8	46.6	30.8	68.1
High	1.0	1.0	0.0	66.4	21.7	44.7	0.0	67.3

* Excludes 99 women classified as subfecund or infecund

Table 8.4
Percentage of Potential Demand for Modern Contraceptive Methods Satisfied
Married Women 15-44 Years of Age At Risk of an Unintended Pregnancy
By Selected Characteristics
Reproductive Health Survey: Albania, 2002

Characteristics	Total Potential Demand for Modern Methods	Percent Using Modern Methods	Percentage of Potential Demand for Modern Methods Satisfied
Total	68.4	8.0	11.7
Strata			
Metro Tirana	67.8	14.9	22.0
Other Urban	68.8	9.4	13.7
Other Rural	68.3	5.1	7.5
Residence			
Urban	68.5	11.3	16.5
Rural	68.3	5.5	8.1
Age Group			
15-19	49.2	2.5	5.1
20-24	61.8	3.7	6.0
25-29	63.5	5.3	8.3
30-34	70.1	9.4	13.4
35-39	73.4	9.3	12.7
40-44	72.5	10.9	15.0
Live Children			
0	32.3	2.2	6.8
1	59.8	3.1	5.2
2	74.3	9.1	12.2
3 +	74.1	10.7	14.4
Education Level			
Primary or Less	66.6	5.8	8.7
Secondary Incomplete	63.1	9.5	15.1
Secondary Complete	72.8	9.4	12.9
Post-Secondary	66.5	17.2	25.9
Socioeconomic Index			
Low	71.1	6.2	8.7
Medium	66.5	7.9	11.9
High	65.3	17.8	27.3

Table 8.5 Advise on Using Modern Methods by Method and Type of Counseling: Ever Users of Modern Methods Reproductive Health Survey: Albania, 2002

		Las	ceptive Metho stribution)	bd	
Advise Last Method	Total	Condom	Female Sterilization	Pill / Hormonal	IUD
OB/GYN	57.6	10.7	91.3	52.6	81.0
Partner	21.0	61.3	0.2	6.7	0.0
Nobody	8.6	18.8	3.8	5.2	0.0
Nurse/Midwife/Feldcher	3.5	1.6	1.6	10.3	5.8
Friend	2.7	1.2	1.0	7.8	7.3
Pharmacist	2.3	3.1	0.0	6.8	0.0
General Practitioner	2.3	1.1	2.1	4.7	2.0
Mother/Other Relative	2.0	2.1	0.0	5.9	3.9
Total	100.0	100.0	100.0	100.0	100.0
No. of Cases	509	193	175	110	31
% With Counseling	63.4	13.5	95.0	67.6	88.8

		Last Used Contraceptive Method (Percentage)						
Type of Counseling	Total	Condom	Female Sterilization	Pill / Hormonal	IUD			
General Information About Other Methods	40.9	**	26.1	61.5	78.9			
Information About Method Effectiveness	34.8	**	21.1	55.4	67.8			
Information About Possible Side Effects	53.4	**	45.4	67.5	70.7			
No. of Cases	287	22	160	76	29			

** Percentages are not shown when base is less than 25 cases.

ab			0
	ŀ	ľ	
		l	Ū

Table 8.6
Counseling and Contraception Offered
at the Time of Legally Performed Abortions
Reproductive Health Survey: Albania, 2002

	tha	entage Rep t MD Discu aception wit	ssed	Percent Distribution of Receipt of Contraceptive Method				
Characteristics	Yes, Before Abortion	Yes, After Abortion	Yes, Before And After	Received a Method	Received a Prescription	No Method or Prescription	Total	No. of Cases
Total	18.3	15.3	4.7	13.7	2.1	84.2	100.0	235
Strata								
Metro Tirana	15.1	26.0	5.4	15.4	5.4	79.2	100.0	109
Other Urban	21.1	14.4	6.1	12.3	1.9	85.7	100.0	92
Other Rural	16.8	9.3	2.3	14.3	0.0	85.7	100.0	34
Residence								
Urban	19.1	18.3	5.7	13.5	3.2	83.3	100.0	190
Rural	16.9	9.9	2.7	14.0	0.0	86.0	100.0	45
Mother's Age at Abortion	1							
15-24	9.0	20.5	1.1	9.2	1.1	89.7	100.0	50
25-34	20.0	14.1	5.7	16.4	2.6	81.0	100.0	147
35-44	22.2	13.9	5.0	8.9	1.3	89.8	100.0	38
Education Level								
Primary or Less Secondary Incomplete	16.5	15.6	5.3	16.4	1.1	82.6	100.0	114
or Higher	20.7	14.8	3.8	10.0	3.4	86.5	100.0	121
Socioeconomic Index								
Low	19.7	13.8	3.4	16.1	0.0	83.9	100.0	71
Medium	16.6	15.0	5.6	9.0	3.8	87.2	100.0	111
High	19.3	20.4	5.5	20.7	2.8	76.4	100.0	53
Where Abortion Performe	ed*							
Hospital / Maternity	21.8	17.1	6.4	16.5	1.7	81.8	100.0	166
Private Clinic / Office	9.7	14.5	0.0	6.3	4.3	89.4	100.0	62

* Excludes 7 cases that reported abortion performed in "other" location.

CHAPTER 9


OPINIONS ABOUT CONTRACEPTION AND ABORTION

Introduction

Tor five decades Albanians lived without access to modern contraception and abortion or even accurate information on the topics. Part of the purpose of the Albania RHS 2002 was to assess Albanians' interest in obtaining more information about contraception and their opinions about the advantages, disadvantages, and safety of known methods. Specifically, survey questions asked about desire contraceptive information. for more preferred source of such information, whether contraceptive information should be broadcast over television and radio, advantage/disadvantages of the intrauterine device (IUD) and oral contraceptives, and safety of selected methods of contraception and of abortion. These data on the opinions of the nation's reproductive age population can help direct efforts to meet the needs for accurate information about modern contraceptive methods in ways that suit the preferences of Albanian men and women

Desire for More Information about Contraception


Slightly more than two-thirds of Albanian women (68%) and nearly half of Albanian men (48%) wanted more information about contraception (Table 9.1). Desire for contraceptive information was not affected by survey strata or urban/rural residence. For both men and women, interest in more information about contraceptive methods declined as age and number of living children increased. Desire for information was nearly equal for women and men aged 15–19, 85% and 82%, respectively. Figure 9.1 shows that among women, interest was higher for those not currently using any contraception (75%) compared to those using either traditional or modern methods (58% and 62%), but among men, interest



was higher among both non-users (61%) and modern method users (73%) than those using traditional methods (32%). For women, interest in family planning information increased with socioeconomic status, but interest was higher among men of low and middle SES. Women with post-secondary education were more interested in contraceptive information, but education appeared to have no impact on interest among men. Women who are less than 25 years old, women who have no living children and women who have never been married are most likely to want more information about contraception (83%-85%). Men reporting a high level of interest are less than 20 years old (82%), but interest is also very high (71%-74%) among those who have never been married, have no living children, or are currently using a modern method of contraception (mostly condom).

Opinion on the Best source of Contraceptive Information

Among women who want more information about contraception, more than half (53%) think that radio or television would be the best source of that information (Table 9.2A and Figure 9.2). One in six women think a gynecologist would be the best source and 12% favor print media such as newspapers or brochures. Radio and television were most popular among rural women, divorced and widowed women, and women currently using condoms. Preference for receiving contraceptive information through mass electronic media increased with age and number of living children and decreased with increasing SES and educational level. Gynecologist as the best source of information was preferred most by women in metro Tirana, urban women, and women using modern contraceptive methods other than condoms and tubal ligation. Women



also were more likely to name a gynecologist as the best source of information as education and socioeconomic status increased. Printed information was most preferred by almost a third of women with post-secondary education (31%), nearly as many as named television and radio (37%). Younger women, never married, better educated women, and those with higher socioeconomic status and no children, showed a greater interest in family planning information in print form.

Most Albanian men who wanted more information about contraceptive methods thought the best sources to be radio or television (65%), print media (15%), or a health professional other than a gynecologist (12%) (Table 9.2B and Figure 9.2). Radio and TV were most favored by rural men, younger men, men with two living children, those not using a modern method of contraception, and those with less than a college education. Also, interest in receiving information over the airwaves was inversely related to socioeconomic status. Men most likely to report that a health professional would be the best source of contraceptive information are those aged 40-44, or with three or more living children. Such health professionals were generally more often named by men over the age of 24, currently married men, men with lower socioeconomic status, primary education or less, and those already using some method of contraception.

Opinion on Appropriateness of Broadcasting Contraceptive Information on Radio and Television

The Albania RHS 2002 asked all respondents whether information about contraception should be broadcast over television and radio. This question helps determine whether such broadcasts would be acceptable to the general population. Additional questions ask both men and women how much time they spend listening to radio or watching television, what time of day, and which stations and types of programs they favor. While these data are not included in this report, they are available for use in developing targeted IEC programs.

Nine out of ten (90%) women think contraceptive information should be broadcast (Table 9.3). There is little variation across characteristics, although women aged 20–24, of high SES or with post-secondary education are most likely to agree that this information should be available over the broadcast media (94%– 96%).

Approximately three-quarters of men (77%) think that information about contraception should be broadcast on radio or television (Table 9.3). Approval for broadcasting such information decreases as age and number of living children increases, and increases with SES and education. Currently married men are less likely than those still unmarried to think family planning information should be broadcast (69% vs. 89%), and men aged 45–49 or with 3 or more living children are least likely to accept such information on TV and the radio (58% and 57%, respectively).

Opinions Regarding the Advantages and Disadvantages of Pill and IUD Use

Women who have heard of the birth control pill were presented with a series of statements expressing advantages and disadvantages of using the Pill. Advantages presented were that the Pill is easy to use, easy to get, makes periods more regular, reduces menstrual cramps, protects against cancer, and decreases monthly blood loss. Potential disadvantages presented were that it can be stressful to remember to take the Pill everyday, pills make you gain weight, are too expensive, and are bad for circulation.

More than half of Albanian women who have heard of oral contraceptives think the Pill is easy to procure (68%) and easy to use (58%) (Table 9.4). Only one in five (19%) knows that the Pill can regulate the menstrual cycle and only one in ten knows that the Pill can reduce menstrual cramps, blood loss, and the risk of certain cancers. Agreement with these advantages was generally close between urban and rural women, although urban woman were somewhat more likely to know that pills can decrease blood loss. Women with education beyond secondary school were more aware of benefits from birth control pills than were women with primary education or less.

Approximately one-third of Albanian women agreed that the Pill can be stressful to remember to take (33%) or cause weight gain (32%), and a much smaller proportion thought that the Pill is too expensive (11%) or bad for circulation (8%). Urban women were somewhat more likely to agree that taking the Pill is stressful or causes weight gain, while women with higher education were much more likely to agree with these disadvantages.

Women were also presented with a similar set of statements regarding the advantages and disadvantages of IUD use. Statements about the possible advantages were that the IUD is Easy to use, relatively inexpensive, and decreases the risk of ectopic pregnancy. Possible disadvantages were that the IUD increases the risk of pelvic inflammatory disease (PID), may cause spotting between periods, may increase menstrual blood loss, and may increase painful menstruation. Slightly more than a third (37%) of women agreed that the IUD is easy to use (Table 9.5). Agreement with this statement was consistent across residence and education, ranging from 33% to 39%. Twenty two percent agreed that the IUD was relatively inexpensive, and a higher proportion of women with post-secondary education agreed (31%). Less than 15% of women, overall, incorrectly stated that the IUD decreases risk of ectopic pregnancy.

Agreement with the disadvantages was low as well. Almost a third of women (32%) agreed that IUD use increases the risk of PID, and that agreement increased with education. The percent of women agreeing with the other disadvantages (causes spotting between periods, increases menstrual blood loss, and increases painful menstruation) also increased with educational attainment. However, given the small sample size of women who have heard of the IUD, the differences for the education variable are not statistically significant. While knowledge of advantages and disadvantages is low, this may be a result of the low level of IUD use in the country.

Opinions on Risks to Women's Health Due to Use of Selected Birth Prevention Methods

All respondents were asked their opinion of the degree of risk (low, medium, or high) posed to a woman's health by selected methods of family planning, including the Pill, IUD, condom, tubal ligation, and abortion. Anywhere from a third of respondents to nearly 90% felt they did not know enough about a given method to assess its health risk (Table 9.6).

More than half of Albanian women surveyed did not know how great or small a risk the Pill, the IUD, or tubal ligation posed to a woman's health. Most women had opinions about condoms and abortion. The condom was generally believed to be low risk (47%) and abortion medium or high risk (23% and 40%, respectively).

More than two-thirds of Albanian men did not have enough knowledge to assess the risks the Pill, the IUD, or tubal ligation posed to a woman's health. Most men did report an opinion about the risk associated with condom and abortion. Sixty-four percent of Albanian men felt the condom posed a low degree of risk to a woman but 44% felt that abortion was of at least medium risk (18% medium, and 26% high).

In Tables 9.7 through 9.10, more detailed analyses are presented only for those birth prevention methods for which roughly half or more of respondents had an opinion. A look at women's perception of health risks from the contraceptive pill by selected characteristics shows that rural women. those under twenty years of age, with less than secondary education, or of low SES are the most likely to respond that they do not know what risks there might be (Table 9.7). Women more likely to have an opinion are those with a post-secondary education, of high SES, or those using a modern method of contraception other than tubal ligation. Most women with an opinion attribute a medium risk to the Pill, but previously married women and those currently using the Pill are more likely to say that the Pill poses a low risk to a woman's health.

Nearly half of Albanian women did not have enough knowledge to assess condoms' potential health risks. Women were more likely to have formed an opinion as their education and socioeconomic status increased, if they lived in metropolitan Tirana or other urban areas, and if they used a modern method of contraception other than tubal ligation (Table 9.8A). The vast majority of women with an opinion felt that the condom poses a low health risk to a woman. Condom users, especially, held the opinion that condoms pose a low risk (95%).

More than two-thirds of Albanian men had an opinion on condoms' health risks to women, although more than half of men 45–49 years of age or with three or more children did not know enough about condoms to form an opinion (Table 9.8B). As with the women, the vast majority of men who had an opinion rated condoms' health risks as low.

Slightly more than half of Albanian women surveyed did not know whether tubal ligation posed a health risk, but the proportion of women without an opinion decreased as age, educational level and SES increased (Table 9.9). Never married women, those with no living children, and those who are not using any form of contraception were much less likely to have an opinion, while most women who are surgically sterilized did have an opinion about the risks of the procedure. In general, few women with an opinion were inclined to say tubal ligation posed a high risk to a woman's health (8%). Most felt there was medium risk (27%), with about half as many assessing a low risk (13%).

Opinions on Risks to Women's Health Due to Abortion

About two-thirds of Albanian women had an opinion on the possible health risks posed to a woman by abortion, and nearly all of those attributed a medium or high risk to the procedure (Table 9.10A). Women were more likely to have an opinion on abortion risks as level of education and socioeconomic status increased, and if they lived in Metro Tirana or an urban area, or had ever had an abortion. Women under age twenty and those with no living children were less likely to have formed an opinion about the health risk from abortion. Most women with an opinion felt that abortion posed a high risk to a woman's health and roughly one-third attributed a medium risk. Almost half of women who reported ever having had an abortion perceived that the procedure posed a high risk to women's health.

Albanian men were less able to give an opinion on the risks that abortion might pose to women's health (44% of the total responded "don't know") (Table 9.10B). Men aged 15–19 were least likely to have an opinion (40%), while those with postsecondary education or high SES were more likely to have an opinion (68% and 70%, respectively), and users of modern contraception were the most likely to do so (89%). Generally, men felt that abortion was a medium- to high-risk procedure, with high risk accounting for one-fifth to nearly one-half of all men's responses (21%–49%) across characteristics.

Concluding Comments

These findings from the Albania RHS 2002 indicate that Albanians in general, both men and women, want and need to know

more about modern contraceptive methods so that they can decide which method or methods may be best for them as they plan their families. They want to receive this information by television and radio, and such broadcasts would be acceptable to most men and women. Women are less than knowledgeable of the benefits and disadvantages that come with using an IUD or the Pill, important facts that can affect a woman's decision to begin using these methods or to continue to do so when unexpected side effects occur. Albanian men and women express uncertainty about the health risks from using modern contraceptive methods; more information would likely reduce fear of such methods. Men and women alike are suspicious of health risks due to abortion, which would seem to make them more open to using effective modern methods of contraception if they have enough information to make informed decisions. By simply meeting the demand for contraceptive information, reproductive health programs may very well generate demand for modern contraceptive methods.

Characteristic	Women Who Want More Information %	No. of Cases	Men Who Want More Information %	No. of Cases
Total	68.1	5,697	47.6	1,740
Strata				
Metro Tirana	67.7	2,108	48.1	718
Other Urban	68.2	1,816	48.3	547
Other Rural	68.2	1,773	47.0	475
Residence				
Urban	68.5	3,572	47.9	1,155
Rural	67.8	2,125	47.3	585
Age Group				
15–19	85.0	1,094	82.3	401
20–24	83.0	936	66.6	189
25–29	71.9	946	57.5	218
30–34	62.9	1,067	41.1	253
35–39	56.0	958	31.3	255
40–44	41.9	696	20.4	277
45–49	N/A	N/A	16.1	147
Marital Status				
Currently Married	60.5	3,965	31.1	1,023
Previously Married	45.6	88	**	14
Never Married	84.7	1,644	73.5	703
Living Children				
0	83.6	1,943	71.0	815
1	68.6	828	40.7	221
2	60.4	1,840	31.1	468
3+	51.2	1,086	15.9	236
Education Level				
Primary or Less	66.5	2,519	47.0	689
Secondary	68.5	2,483	48.4	825
Post-Secondary	77.3	695	46.6	226
Socioeconomic Index				
Low	65.5	1,940	47.4	638
Medium	69.7	2,985	49.2	814
High	72.5	772	41.9	288
Current Method of Con	traception			
Modern	58.0	456	73.1	98
Traditional	61.8	2,739	32.4	804
Non-User	75.1	2,502	60.7	838

Table 9.1Percentage Who Want More Information about Contraception
by Selected CharacteristicsAmong Women Aged 15–44 and Men Aged 15–49Reproductive Health Survey: Albania, 2002

*Percentages are not shown when base is less than 25 cases.

Table

Percent Distribution of Women's Opinion on the Best Source of Contraceptive Information by Selected Characteristics Among Women 15–44 Who Want More Information About Contraception Reproductive Health Survey: Albania, 2002										
Characteristic	Radio/ TV	Gyne- colo- gist	Print Media	Other Health Pro- fessional	Relative/ Partner/ Boyfriend	Friend/ Cowork- er/ Contracep- tive User	Teacher	Other/ Don't Know	Total	No. of Cases
Total	52.8	16.3	11.7	8.8	4.1	2.8	1.4	2.1	100.0	3,885
Strata										
Metro Tirana	45.6	25.8	12.9	5.5	4.8	3.3	1.4	0.8	100.0	1,442
Other Urban	48.0	18.1	18.7	6.5	3.3	2.3	1.6	1.6	100.0	1,237
Other Rural	57.5	12.6	7.6	11.0	4.4	2.9	1.3	2.8	100.0	1,206
Residence										
Urban	47.1	21.2	16.7	5.9	3.7	2.7	1.5	1.2	100.0	2,464
Rural	57.1	12.6	7.9	10.9	4.4	2.9	1.3	2.8		1,421
Age Group										
15–19	49.3	11.1	13.7	5.7	9.0	4.0	4.8	2.4	100.0	923
20–24	52.6	16.9	12.6	8.4	4.5	2.9	0.3	1.8	100.0	769
25-29	50.9	18.6	13.2	9.7	3.1	2.2	0.2	2.0	100.0	663
30–34	54.6	18.4	9.2	12.2	0.8	2.7	0.1	2.1	100.0	671
35–39	56.8	18.7	8.7	12.1	0.4	1.7	0.0	1.5	100.0	545
40-44	58.0	18.9	9.8	6.9	1.5	2.0	0.0	3.0	100.0	314
Marital Status										
Currently Married	56.7	18.5	8.8	10.3	1.5	2.3	0.1	1.8	100.0	2,445
Previously Married	67.7	10.3	1.5	15.5	1.0	1.0	0.0	2.9	100.0	48
Never Married	46.7	13.4	16.2	6.4	7.9	3.7	3.2	2.6		1,392
Living Children				••••		•	0.2			.,
0	47.7	13.8	15.3	6.8	7.4	3.7	2.8	2.3	100.0	1,611
1	53.8	20.9	10.3	9.1	1.2	2.9	0.2	1.6	100.0	563
2	57.6	19.1	10.0	8.1	1.4	1.6	0.2	1.9		1,134
3+	59.1	15.6	5.1	14.8	1.1	2.0	0.0	2.3	100.0	577
Education Level	00.1	10.0	0.1	11.0		2.0	0.0	2.0	100.0	011
Primary or Less	58.2	13.2	6.8	10.7	3.9	3.3	1.0	2.9	100.0	1,649
Secondary	49.2	13.2 19.0	0.0 13.9	7.3	3.9 4.9	3.3 2.1	2.1	2.9 1.5	100.0	1,049
Post-Secondary	49.2 36.5	23.0	30.8	3.7	4.9	3.0	1.0	0.0	100.0	527
-		20.0	50.0	5.7	1.5	5.0	1.0	0.0	100.0	521
Socioeconomic Inde	x 56.9	13.8	7.3	11.4	4.4	2.3	1.2	2.6	100.0	1,241
Medium	51.0	16.5	14.3	7.7	3.9	3.1	1.4	2.0	100.0	2,090
High	43.8	26.4	14.5	2.8	4.2	3.3	1.4	0.7	100.0	2,090 554
Current Method of C				2.0		0.0		0.1		001
Tubal Ligation	54.8	17.4	9.8	16.8	0.7	0.6	0.0	0.0	100.0	70
Condom	60.7	21.6	9.0 10.0	4.8	0.7	2.4	0.0	0.0	100.0	123
Other Modern	43.8	21.0 36.4	10.0	4.8 9.0	0.0	0.0	0.0	0.0	100.0	94
Traditional	43.8 54.7	30.4 19.4	9.2	9.0 10.7	1.4	2.2	0.0	2.2	100.0	94 1,708
Non-User	54.7	13.2	9.2 13.7	7.2	6.5	3.4	2.4	2.2		1,890

Table 9.2 A ation ini

Opinions About Contraception and Abortion

Opinions About Contraception and Abortion

Table 9.2 B Percent Distribution of Men's Opinion on the Best Source of Contraceptive Information by Selected Characteristics Among Men Aged 15–49 Who Want More Information About Contraception										1
Among Men Aged 15–49 who want More Information About Contraception Reproductive Health Survey: Albania, 2002										
Characteristic	Radio/ TV	Print Media	Other Health Pro- fessional	Friend/ Cowork- er/ Contracep- tive User	Gyne- cologist	Teacher	Relative/ Partner/ Girlfriend	Other/ Don't Know	Total	No. of Cases
Total	65.4	15.2	11.6	3.2	2.6	1.2	0.3	0.5	100.0	863
Strata										
Metro Tirana	54.2	24.3	11.0	2.9	4.4	2.7	0.3	0.3	100.0	351
Other Urban	62.1	17.2	8.6	5.9	4.3	0.8	0.0	1.1	100.0	277
Other Rural	71.9	10.4	13.6	1.6	0.8	0.9	0.5	0.2	100.0	235
Residence	1110		1010		0.0	0.0	0.0	0.2	10010	200
	50.2	10.2	0.5	5.0	15	1.6	0.1	0.0	100.0	571
Urban	59.2	19.3	9.5	5.0	4.5	1.6	0.1	0.9	100.0	571
Rural	70.9	11.7	13.4	1.6	0.9	0.9	0.5	0.2	100.0	292
Age Group										
15–19	70.2	15.6	7.3	3.3	0.7	2.5	0.2	0.3	100.0	328
20–24	68.3	19.5	4.5	2.1	2.6	1.2	0.0	1.7	100.0	131
25–29	62.1	13.3	18.3	3.8	2.5	0.0	0.0	0.0	100.0	122
30–34	61.8	12.8	15.6	3.6	6.3	0.0	0.0	0.0	100.0	105
35–39	61.3	10.3	15.7	6.2	4.7	1.7	0.0	0.0	100.0	88
40–44	52.9	18.8	25.0	0.0	3.3	0.0	0.0	0.0	100.0	61
45–49	64.1	11.5	13.2	1.6	1.6	0.0	6.5	1.6	100.0	28
Marital Status										
Currently Married Previously	63.7	13.0	15.9	2.0	4.1	0.4	0.6	0.2	100.0	335
Married	**	**	**	**	**	**	**	**	**	6
Never Married	66.5	16.6	8.7	3.9	1.6	1.8	0.1	0.8	100.0	522
Living Children										
0	65.6	16.6	10.1	3.7	1.7	1.6	0.1	0.7	100.0	580
1	61.2	13.6	15.6	0.0	9.1	0.5	0.0	0.0	100.0	89
2	70.1	13.4	9.2	2.9	3.3	0.7	0.0	0.4	100.0	153
3+	57.3	7.7	28.1	2.8	0.0	0.0	4.1	0.0	100.0	41
Education Level										
Primary or Less	67.8	9.3	17.1	3.1	0.8	0.5	0.5	0.9	100.0	340
Secondary	65.9	16.9	7.5	3.8	4.0	1.8	0.0	0.0	100.0	419
Post-Secondary	50.3	39.4	1.7	0.5	5.0	2.4	0.0	0.7	100.0	104
Socieconomic Inde		00.4	1.7	0.0	0.0	2.4	0.0	0.7	100.0	104
		07	14.0	2.4	0.0	0.5	0.5	0.7	100.0	204
Low	71.1	8.7 10.7	14.2	3.4	0.9	0.5	0.5	0.7	100.0	321
Medium	60.7	19.7 27.2	10.1	3.6	3.5	2.1	0.1	0.3	100.0	414
High	58.5	27.2	5.2	0.0	7.1	1.1	0.3	0.7	100.0	128
Current Method of		•								
Modern	41.8	22.8	17.1	3.5	8.5	0.0	0.0	6.3	100.0	68
Traditional	66.8	12.9	15.1	2.2	2.5	0.5	0.0	0.0	100.0	273
Non-User	67.1	15.7	9.1	3.6	2.0	1.8	0.5	0.2	100.0	522

Tabl 0 0 0

ables

Table 9.3Percentage Who Think That Information about Contraception
Should be Broadcast on Radio or Television
by Selected CharacteristicsAmong Women Aged 15–44 and Men Aged 15–49
Reproductive Health Survey: Albania, 2002

Characteristic	Women	No. of Cases	Men	No. of Cases
Total	90.0	5,697	76.9	1,740
Strata				
Metro Tirana	91.1	2,108	82.5	718
Other Urban	91.6	1,816	79.5	547
Other Rural	88.9	1,773	73.1	475
Residence				
Urban	91.8	3,572	81.0	1,155
Rural	88.7	2,125	73.2	585
Age Group				
15–19	90.6	1,094	90.9	401
20–24	94.3	936	86.9	189
25–29	91.7	946	84.2	218
30–34	89.7	1,067	76.8	253
35–39	88.4	958	66.7	255
40–44	84.7	696	66.5	277
45–49	N/A	N/A	57.8	147
Marital Status				
Currently Married	89.5	3,965	68.9	1,023
Previously Married	83.4	88	**	14
Never Married	91.6	1,644	88.9	703
Living Children				
0	92.0	1,943	89.1	815
1	92.2	828	76.7	221
2	89.2	1,840	69.2	468
3+	86.6	1,086	56.8	236
Education Level				
Primary or Less	87.5	2,519	72.7	689
Secondary	92.3	2,483	80.1	825
Post-Secondary	96.2	695	83.7	226
Socioeconomic Index				
Low	86.8	1,940	71.5	638
Medium	92.0	2,985	79.6	814
High	94.9	772	90.0	288
Current Method of Contraception				
Modern	92.2	456	95.7	98
Traditional	89.8	2,739	70.9	804
Non-User	90.0	2,502	81.2	838

**Percentages are not shown when base is less than 25 cases.

Table 9.4 Percentage Who Agree with Selected Statements Concerning Possible Advantages and Disadvantages of Using the Pill, by Residence and Education Among Women Aged 15–44 Who Have Heard of the Pill Reproductive Health Survey: Albania 2002

		Resid	lence		vel	
Statements	Total	Urban	Rural	Primary Or Less	Secondary	Post- Secondary
Advantages						
Pills Are Easy to Procure	67.9	72.1	63.6	62.6	69.5	81.7
Pills Are Easy to Use	58.1	59.0	57.2	54.8	58.7	68.4
Pills Regulate Periods	18.9	21.4	16.5	13.1	20.2	36.6
Pills Decrease Menstrual Cramps	10.5	11.4	9.5	6.6	12.1	19.0
Pills Protect Against Cancer	10.5	11.2	9.9	7.2	12.8	14.4
Pills Decrease Blood Loss	9.7	12.4	7.0	5.9	11.2	19.0
Disadvantages						
Stressful to Remember to Take the Pill	32.9	35.8	30.0	28.4	34.7	43.4
Pills Make You Gain Weight	32.2	36.8	27.6	25.1	35.3	47.7
Pills Are Too Expensive	11.0	11.1	10.9	11.4	11.1	9.3
Pills Are Bad for Circulation	8.0	8.2	7.8	6.4	8.5	12.0
Number of Cases	4,178	2,887	1,291	1,475	2,031	672

Table 9.5

Percentage Who Agree with Selected Statements Concerning Possible Advantages and Disadvantages of Using the IUD, by Residence and Education Among Women Aged 15–44 Who Have Heard of the IUD Reproductive Health Survey: Albania 2002

		Residence		Education Level			
				Primary		Post-	
Statements	Total	Urban	Rural	Or Less	Secondary	Secondary	
Advantages							
IUD is Easy to Use	36.5	35.0	38.6	32.5	39.4	36.0	
IUD is Relatively Inexpensive	22.4	23.2	21.3	20.3	21.0	30.5	
IUD Decreases Risk of Ectopic Pregnancy	14.4	16.2	11.7	9.3	15.6	20.6	
Disadvantages							
Increases Risk of Pelvic Inflammantory Disease	31.6	30.5	33.2	27.1	32.6	37.5	
IUD May Cause Spotting Between Periods	21.2	20.0	23.0	20.8	20.0	25.3	
IUD May Increase Menstrual Blood Loss	15.3	14.0	17.2	14.3	14.9	18.0	
IUD May Increase Painful Menstruation	14.8	14.2	15.6	13.2	14.9	17.5	
Number of Cases	1,643	1,244	399	433	835	375	

Table 9.6

Percent Distribution of Opinion of Degree of Risk to a Woman's Health from Using Selected Birth Prevention Methods Among Women Aged 15-44 and Men Aged 15-49 Reproductive Health Survey: Albania, 2002

Birth Prevention Method	Low Risk	Medium Risk	High Risk	Don't Know	Total	No. of Cases
					Teta	
Women						
Pill	12.8	26.5	6.6	54.2	100.0	5,697
IUD	3.4	8.9	3.6	84.1	100.0	5,697
Condom	46.6	7.3	1.4	44.7	100.0	5,697
Tubal Ligation	12.7	26.8	7.6	52.9	100.0	5,697
Abortion	4.3	22.9	39.7	33.1	100.0	5,697
Men						
Pill	13.0	10.7	3.7	72.7	100.0	1,740
IUD	4.3	4.4	2.6	88.8	100.0	1,740
Condom	64.4	6.3	0.1	29.3	100.0	1,740
Tubal Ligation	3.4	7.8	7.3	81.6	100.0	1,740
Abortion	11.8	18.3	25.9	44.1	100.0	1,740

Table	9.7
-------	-----

Percent Distribution of Opinion of Degree of Risk to a Woman's Health From Using the Pill by Selected Characteristics Among Women Aged 15–44 Reproductive Health Survey: Albania, 2002

Characteristic	Low Risk	Medium Risk	High Risk	Don't Know	Total	No. of Cases
Total	12.8	26.5	6.6	54.2	100.0	5,697
Strata						
Metro Tirana	16.4	32.3	8.9	42.3	100.0	2,108
Other Urban	14.1	31.9	6.3	47.8	100.0	1,816
Other Rural	11.0	22.0	6.0	61.1	100.0	1,773
Residence						
Urban	14.9	32.4	7.2	45.5	100.0	3,572
Rural	11.1	22.1	6.1	60.7	100.0	2,125
Age						
15–19	8.8	23.7	3.7	63.8	100.0	1,094
20–24	15.5	22.8	8.6	53.1	100.0	936
25–29	12.8	29.8	8.6	48.7	100.0	946
30–34	15.8	27.9	6.5	49.8	100.0	1,067
35–39	12.4	27.6	6.2	53.8	100.0	958
40–44	12.4	28.6	6.6	52.5	100.0	696
Marital Status						
Currently Married	13.1	27.4	7.1	52.4	100.0	3,965
Previously Married	24.0	26.7	5.8	43.5	100.0	88
Never Married	11.3	24.9	5.6	58.3	100.0	1,644
Living Children						
0	11.2	25.6	5.6	57.7	100.0	1,943
1	17.0	24.2	7.7	51.1	100.0	828
2	14.0	30.5	7.0	48.4	100.0	1,840
3+	11.4	24.7	7.1	56.9	100.0	1,086
Education Level						
Primary or Less	10.0	19.7	5.7	64.7	100.0	2,519
Secondary	14.6	32.1	7.6	43.5	100.0	2,483
Post-Secondary	22.5	46.3	10.3	21.0	100.0	695
Socioeconomic Index						
Low	9.5	19.3	5.7	65.4	100.0	1,940
Medium	14.4	30.5	6.7	48.4	100.0	2,985
High	19.2	39.9	9.8	31.1	100.0	772
Current Method of Contraception	ı					
Tubal Ligation	12.2	34.4	5.5	47.9	100.0	173
Pill	78.7	19.6	0.0	1.7	100.0	74
Condom	11.7	53.7	11.6	23.0	100.0	165
Other Modern	22.9	39.0	13.8	24.4	100.0	44
Traditional	12.8	27.7	7.2	52.3	100.0	2,739
Non-User	11.7	24.1	5.9	58.3	100.0	2,502

Table 9.8 A

Percent Distribution of Opinion of Degree of Risk to a Woman's Health From Using the Condom by Selected Characteristics Among Women Aged 15–44 Reproductive Health Survey: Albania, 2002

Characteristic	Low Risk	Medium Risk	High Risk	Don't Know	Total	No. of Cases
Total	46.6	7.3	1.4	44.7	100.0	5,697
Strata						
Metro Tirana	61.0	7.1	0.8	31.1	100.0	2,108
Other Urban	52.6	8.2	1.0	38.2	100.0	1,816
Other Rural	39.1	6.8	1.8	52.3	100.0	1,773
Residence						
Urban	56.3	7.9	0.9	34.9	100.0	3,572
Rural	39.3	6.8	1.7	52.1	100.0	2,125
Age						
15–19	45.5	4.8	1.4	48.3	100.0	1,094
20–24	52.6	6.2	1.6	39.6	100.0	936
25–29	49.4	8.4	0.9	41.3	100.0	946
30–34	46.0	9.4	1.5	43.0	100.0	1,067
35–39	43.3	8.9	1.0	46.9	100.0	958
40–44	42.4	6.8	2.0	48.7	100.0	696
Marital Status						
Currently Married	45.7	8.3	1.5	44.5	100.0	3,965
Previously Married	47.9	8.9	2.3	40.8	100.0	88
Never Married	48.3	5.1	1.2	45.4	100.0	1,644
Living Children						
0	48.5	5.5	1.1	44.8	100.0	1,943
1	51.1	8.5	1.3	39.0	100.0	828
2	48.4	9.7	2.1	39.8	100.0	1,840
3+	38.4	6.5	1.1	53.9	100.0	1,086
Education Level						
Primary or Less	37.2	7.2	1.4	54.2	100.0	2,519
Secondary	53.6	7.4	1.3	37.8	100.0	2,483
Post-Secondary	76.3	7.4	2.0	14.3	100.0	695
Socioeconomic Index						
Low	35.3	6.6	1.8	56.2	100.0	1,940
Medium	52.4	7.7	1.1	38.7	100.0	2,985
High	69.1	7.9	1.1	21.9	100.0	772
Current Method of Contraception						
Tubal Ligation	52.3	4.1	0.0	43.6	100.0	173
Condom	95.2	0.4	0.4	4.1	100.0	163
Other Modern	61.1	15.0	0.0	23.8	100.0	120
Traditional	45.9	8.1	1.7	44.2	100.0	2,739
Non-User	44.9	6.7	1.3	47.1	100.0	2,502

Opinions About Contraception and Abortion

Characteristic	Low Risk	Medium Risk	High Risk	Don't Know	Total	No. of Cases
Total	64.4	6.3	0.1	29.2	100.0	1,740
Strata						
Metro Tirana	75.4	3.6	0.3	20.7	100.0	718
Other Urban	72.4	5.6	0.0	22.0	100.0	547
Other Rural	55.4	7.8	0.0	36.9	100.0	475
Residence						
Urban	73.8	4.8	0.1	21.2	100.0	1,155
Rural	56.2	7.6	0.0	36.2	100.0	585
Age						
15–19	63.2	5.8	0.0	31.0	100.0	401
20–24	83.5	3.9	0.0	12.6	100.0	189
25–29	72.7	8.5	0.0	18.8	100.0	218
30–34	74.0	7.5	0.0	18.5	100.0	253
35–39	59.3	5.5	0.0	35.2	100.0	255
40–44	55.2	8.0	0.0	36.8	100.0	277
45–49	38.7	5.1	0.5	55.7	100.0	147
Marital Status						
Currently Married	57.5	7.5	0.1	34.9	100.0	1,023
Previously Married	**	**	**	**	**	14
Never Married	75.2	4.5	0.0	20.3	100.0	703
Living Children						
0	74.9	5.5	0.0	19.6	100.0	815
1	73.5	6.8	0.0	19.7	100.0	221
2	58.4	8.3	0.2	33.1	100.0	468
3+	40.5	5.2	0.0	54.3	100.0	236
Education Level						
Primary or Less	55.8	6.4	0.0	37.9	100.0	689
Secondary	70.0	6.6	0.1	23.4	100.0	825
Post-Secondary	84.2	4.5	0.3	11.0	100.0	226
Socioeconomic Index						
Low	52.4	6.1	0.0	41.5	100.0	638
Medium	71.4	7.0	0.1	21.5	100.0	814
High	90.0	4.3	0.3	5.5	100.0	288
Current Method of Contraception						

Table 9.8 BPercent Distribution of Opinion of Degree of Risk to a Woman's Health From Using the Condom
by Selected Characteristics Among Men Aged 15–49Reproductive Health Survey: Albania, 2002

**Percentages are not shown when base is less than 25 cases.

Modern

Traditional

Non-user

1.2

7.0

6.1

98.3

58.3

67.6

0.0

0.1

0.0

0.6

34.6

26.3

100.0

100.0

100.0

98

804

838

Table 9.9
Percent Distribution of Opinion of Degree of Risk to a Woman's Health from Tubal Ligation
by Selected Characteristics Among Women Aged 15–44

Chapter 9

Reproductive Health Survey: Albania, 2002

Characteristic	Low Risk	Medium Risk	High Risk	Don't Know	Total	No. of Cases
Total	12.7	26.8	7.6	52.9	100.0	5,697
Strata						
Metro Tirana	16.4	23.7	9.4	50.5	100.0	2,108
Other Urban	12.1	29.6	8.9	49.4	100.0	1,816
Other Rural	12.0	26.2	6.3	55.5	100.0	1,773
Residence						
Urban	13.6	27.7	9.1	49.6	100.0	3,572
Rural	12.0	26.2	6.4	55.3	100.0	2,125
Age						
15–19	6.2	15.0	3.0	75.8	100.0	1,094
20–24	12.5	20.7	7.6	59.2	100.0	936
25–29	13.6	31.2	9.3	45.9	100.0	946
30–34	15.0	32.3	9.5	43.2	100.0	1,067
35–39	14.6	32.2	9.8	43.4	100.0	958
40–44	16.9	34.5	7.8	40.7	100.0	696
Marital Status						
Currently Married	15.2	32.5	9.2	43.1	100.0	3,965
Previously Married	12.7	33.6	13.8	39.9	100.0	88
Never Married	7.7	15.2	4.0	73.1	100.0	1,644
Living Children						
0	8.7	16.3	4.7	70.3	100.0	1,943
1	14.1	30.3	9.4	46.2	100.0	828
2	15.1	36.1	9.8	39.0	100.0	1,840
3+	15.8	31.5	8.8	44.0	100.0	1,086
Education Level						
Primary or Less	11.2	24.3	6.7	57.8	100.0	2,519
Secondary	13.3	28.7	8.1	49.9	100.0	2,483
Post-Secondary	19.9	34.6	11.5	34.0	100.0	695
Socioeconomic Index						
Low	11.4	22.9	6.4	59.3	100.0	1,940
Medium	13.2	29.5	7.4	49.8	100.0	2,985
High	16.7	31.0	14.2	38.1	100.0	772
Current Method of Contraception						
Tubal Ligation	47.6	32.0	5.0	15.4	100.0	173
Condom	8.7	36.1	14.1	41.1	100.0	163
Other Modern	22.9	29.6	13.2	34.4	100.0	120
Traditional	13.9	32.9	9.5	43.7	100.0	2,739
Non-User	9.6	20.7	5.6	64.1	100.0	2,502

Reproductive Health Survey: Albania, 2002									
Characteristic	Low Risk	Medium Risk	High Risk	Don't Know	Total	No. of Cases			
Total	4.3	22.9	39.7	33.1	100.0	5,697			
Strata									
Metro Tirana	5.7	23.0	46.2	25.1	100.0	2,108			
Other Urban	4.8	24.6	40.9	29.8	100.0	1,816			
Other Rural	3.6	22.0	37.2	37.2	100.0	1,773			
Residence									
Urban	5.3	24.0	43.4	27.4	100.0	3,572			
Rural	3.6	22.1	37.0	37.3	100.0	2,125			
Age									
15–19	3.1	21.1	31.1	44.8	100.0	1,094			
20–24	4.1	21.2	40.1	34.6	100.0	936			
25–29	4.4	25.3	43.8	26.5	100.0	946			
30–34	5.9	23.4	43.4	27.3	100.0	1,067			
35–39	3.6	24.7	42.5	29.3	100.0	958			
40–44	5.3	22.6	40.5	31.6	100.0	696			
Marital Status									
Currently Married	5.0	24.0	42.4	28.6	100.0	3,965			
Previously Married	4.0	19.4	44.7	31.9	100.0	88			
Never Married	3.0	21.0	34.0	41.9	100.0	1,644			
Living Children									
0	3.2	20.8	35.5	40.5	100.0	1,943			
1	5.0	25.9	42.4	26.7	100.0	828			
2	4.6	23.7	42.9	28.8	100.0	1,840			
3+	5.4	23.8	41.5	29.3	100.0	1,086			
Education Level									
Primary or Less	4.0	22.5	35.9	37.6	100.0	2,519			
Secondary	4.9	22.9	42.3	29.9	100.0	2,483			
Post-Secondary	3.8	25.8	53.0	17.5	100.0	695			
Socioeconomic Index									
Low	4.5	21.5	35.4	38.5	100.0	1,940			
Medium	3.9	24.0	41.8	30.3	100.0	2,985			
High	5.6	23.2	49.5	21.6	100.0	772			
Ever Had Abortion									
No	4.0	22.5	39.3	34.2	100.0	5,359			
Yes	11.0	31.6	48.1	9.3	100.0	338			

Table 9.10 A Percent Distribution of Opinion of Degree of Risk to a Woman's Health From Abortion by Selected Characteristics Among Women Aged 15–44 Reproductive Health Survey: Albania, 2002

Albania Reproductive Health Survey, 2002 Final Report

Table 9.10 B
Percent Distribution of Opinion of Degree of Risk to a Woman's Health from Abortion
by Selected Characteristics Among Men Aged 15–49
Reproductive Health Survey: Albania, 2002

Characteristic	Low Risk	Medium Risk	High Risk	Don't Know	Total	No. of Cases
Total	11.8	18.3	25.9	44.1	100.0	1,740
Strata						
Metro Tirana	8.6	20.2	26.1	45.1	100.0	718
Other Urban	12.6	18.1	29.6	39.7	100.0	547
Other Rural	12.5	17.7	23.5	46.3	100.0	475
Residence						
Urban	10.9	19.5	28.3	41.3	100.0	1,155
Rural	12.5	17.3	23.7	46.5	100.0	585
Age						
15–19	7.4	10.6	22.4	59.7	100.0	401
20–24	11.9	19.8	25.9	42.4	100.0	189
25–29	13.7	25.8	23.1	37.4	100.0	218
30–34	15.6	24.8	26.1	33.6	100.0	253
35–39	10.5	18.9	28.2	42.4	100.0	255
40–44	11.0	18.1	29.7	41.2	100.0	277
45–49	14.6	12.2	26.7	46.6	100.0	147
Marital Status						
Currently Married	12.5	19.2	25.9	42.4	100.0	1,023
Previously Married	**	**	**	**	**	14
Never Married	10.7	16.8	25.4	47.1	100.0	703
Living Children						
0	11.6	17.7	25.4	45.2	100.0	815
1	13.3	22.7	28.8	35.2	100.0	221
2	11.5	18.3	28.0	42.2	100.0	468
3+	11.5	16.8	22.0	49.6	100.0	236
Education Level						
Primary or Less	12.5	16.2	24.5	46.8	100.0	689
Secondary	10.4	19.3	26.7	43.6	100.0	825
Post-Secondary	14.5	24.7	29.1	31.7	100.0	226
Socioeconomic Index						
Low	12.3	17.8	20.9	49.0	100.0	638
Medium	10.3	17.9	29.6	42.2	100.0	814
High	15.3	22.0	32.9	29.8	100.0	288
Current Method of Contrac	eption					
Modern	14.9	26.0	48.5	10.6	100.0	98
Traditional	14.5	21.4	22.4	41.8	100.0	804
Non-User	8.8	14.5	27.4	49.3	100.0	838

**Percentages are not shown when base is less than 25 cases.

CHAPTER 10

REPRODUCTIVE HEALTH KNOWLEDGE AND ATTITUDES

In most of the Communist countries of LEastern Europe, as mentioned previously, choices for contraception were limited and fertility control was achieved chiefly through abortion (David, 1992). Communist Albania, like Romania, was even more restrictive than other countries of Eastern Europe, prohibiting both contraception and abortion in favor of a pronatalist national Post-Communist Albania policy. has enacted new laws changing the situation. Modern contraception was legalized in 1988 and abortion in 1992 (Islami et al, 1999). Reproductive health education and services in Albania are still evolving.

The ALRHS02 incorporated questions about both men's and women's opinions about ideal family size, basic knowledge of fertility, attitudes about abortion and unintended pregnancy, and perceptions about gender roles and norms. Responses to these questions can help target and shape information, education, and communications programs intended to improve reproductive health knowledge among Albanians of child bearing age.

Ideal Family Size

Respondents were asked "What do you think is the ideal number of children for a young family in Albania?" The question is intended to illuminate a cultural or social norm, rather than a personal desire for a certain number of children. Response categories ranged from 0 to "5 or more", and included nonnumeric responses such as "as many as God gives" and "as many as possible." The mean ideal number of children was calculated for women and men, excluding non-numeric responses (Table 10.1). Among women, the mean ideal number of children for a young family in Albania was 2.6 children. The ideal was higher outside of metro Tirana, in rural areas, and among currently married women. Ideal number of children increased with age and number of living children, and decreased as educational level and socioeconomic status increased.

In general, men thought the ideal number of children was somewhat lower (2.4). The same demographic patterns held true for men, although the mean ideal reported by men was always lower except among men and women with three or more children (3.1 for both).

Knowledge of the Menstrual Cycle

Men and women's knowledge of the menstrual cycle was explored with two questions. The first asked whether there are certain days during a woman's cycle when she is more likely to become pregnant. Respondents that answered yes were then asked when those days occur: just before her period, during her period, right after her period, or halfway between periods.

Fewer than one in five Albanian women (18%) correctly identified the time halfway between periods as a woman's most fertile time (Table 10.2A). One-fourth incorrectly thought that the week after menses was the most likely time for a woman to become pregnant (26%). Nearly half responded that they did not know if a woman is more likely to become pregnant at any particular point in her cycle (46%). Correct knowledge was more prevalent in urban areas, among ever married women, and increased with level of education and socioeconomic status.

Albanian men knew even less about the menstrual cycle. Three quarters of men surveyed responded that either they did not know or that a woman is equally likely to become pregnant at any point in the menstrual cycle (Table 10.2B). Just 11% knew that the time halfway between periods is when a woman is most fertile.

Knowledge of the Fertility Effect of Breastfeeding

The RHS questionnaire included a question asking if breastfeeding increases, decreases, or has no effect on a woman's chance of getting pregnant. Nearly half of the women (47%) responded that they did not know whether there was any effect (Table 10.3A). Only one in five women knew that breastfeeding reduced a woman's fertility (20%). Women outside metro Tirana, rural women, women of lower education and socioeconomic status were more likely to respond that they did not know and were less likely to correctly specify that breastfeeding lowers the chance of pregnancy. Unmarried women, women aged 15–19, or without children were most likely to have no knowledge of the effect of breastfeeding on fertility and least likely to know that pregnancy is less likely when a woman is breastfeeding.

Two-thirds of Albanian men (66%) surveyed reported not knowing if breastfeeding affected a woman's chance of becoming pregnant and approximately 20% knew that it lowers the chance (Table 10.3B). As with the women, knowledge was greater among urban men, and much lower among never married men, young men, men with less than complete secondary education, low socioeconomic status, and with no children.

Attitudes Toward Abortion

To explore whether Albanians, in the abstract, consider abortion to be an option for limiting fertility, the questionnaire asks whether respondents believe that a woman always has the right to decide about her pregnancy, including whether or not to have an abortion. Respondents who do not agree that abortion is always a woman's decision answer a question about whether abortion is acceptable under certain circumstances. Table 10.4A displays the percent distribution of surveyed women who agree that abortion is always a woman's decision, that abortion is acceptable under at least one of six selected circumstances, or never acceptable.

Nearly three quarters of Albanian women (72%) think that a woman always has the right to make decisions about her own pregnancy, including whether to have an abortion. Almost all other women believe that abortion is acceptable under some circumstances (26%). The opinion that abortion is a woman's decision does not vary much by demographic characteristics, although it is most common among women with postsecondary education (78%) and least prevalent among women of low socioeconomic status (69%).

Fewer Albanian men report that abortion is always a woman's personal decision (53%), but because most others believe it is acceptable in certain circumstances, the proportion that believe abortion is never acceptable is about equal to that of women (1%) (Table 10.4B). Acceptance of abortion varied among Albanian men by strata, residence and age; it was most common among men in urban areas other than Tirana (61%) and least likely among men aged 45 - 49 (44%). Acceptance of a woman's right to decide about her pregnancy increased somewhat with socioeconomic status (52% to 59%).

Respondents who did not think that a pregnant woman always has the right to decide about abortion were asked about the acceptability of abortion under specific circumstances: danger to a woman's life, danger to a woman's health, deformity of the fetus, pregnancy resulting from rape, pregnancy outside of marriage, and inability to support a/another child. Among both men and women, the selected circumstances can be grouped into three levels of acceptance.

Among women, being unmarried or unable to afford a child were the least acceptable reasons for abortion (38% and 22% respectively) (Table 10.5). Lack of money for a child was also found to be the situation with the greatest definitively negative response of abortion as opposed to "depends" or "don't know" (59%). The second level of acceptance among women was for situations in which the pregnancy was the result of a rape or it posed a danger to a woman's health (53% and 59%, respectively). About one in five respondents (19%) to these questions thought that abortion was not appropriate in these circumstances. The highest level of acceptance among women was for pregnancies in which the fetus had some sort of deformity (70%) or which threatened a woman's life (80%). One in ten women felt a woman must carry a pregnancy to term even if it might end her life and 19% felt she should do so even if the fetus was physically deformed.

Acceptance of abortion under specific circumstances shows some patterns when examined by women's characteristics (Table 10.6A). Women who are aged 15 - 19, never married, have no children, of lower

socioeconomic status, live in rural areas, or who have less education are in general less likely to accept abortion in any of the situations presented. However, in the first three cases shown, from one-half to threequarters of these women accepted abortion in these circumstances. With regard to urban/rural residence and socioeconomic status, the proportion was inverted when the circumstance in question was a pregnancy in a woman without a husband; in that case rural women were more likely to agree with abortion (42% vs. 32%) as were women in the lowest socioeconomic level (42% vs. 34% and 35%).

Albanian men were similar in their acceptance of abortion under the circumstances shown (Table 10.5). However, compared with female respondents, they were less likely to accept abortion as an option when a family can not afford a child or when a woman is unmarried, and most likely to disapprove outright in those circumstances. Albanian men also showed less acceptance when a fetus is malformed. Men placed the greatest value on a woman's health and life, agreeing that abortion was acceptable when her health or life were at risk (74% and 87%, respectively).

In looking at the situational acceptance of abortion among men by selected characteristics, one can see that, in general, approval was higher among men living in urban areas, currently married men, men with secondary complete education, and men with high socioeconomic status (Table 10.6B). Men who live outside of Tirana were less likely to accept abortion when a woman is unmarried or the family cannot support the child once it is born.

While most believe that abortion is a woman's personal decision, in the event of an unintended pregnancy, a large majority of Albanian women tend to think that a woman should have the baby and raise it herself (71%) and a significant minority think that she should have an abortion (21%) (Table 10.7A). Only 2% think a woman should have her baby adopted. Women most likely to think a woman should give birth and keep the baby are those in rural areas. "Keep the baby" decreases with higher levels of education and socioeconomic status.

Albanian men are much less likely to say that a woman should have and keep an unwanted baby (40%) (Table 10.7B). Adoption as a choice of action is generally more favored by men than women (10% vs. 2%). Men living in Tirana and in other urban areas, as well as men with postsecondary education and of high socioeconomic status, were most likely to think a woman should have an abortion. Men at the extremes of the reproductive ages were least likely to suggest a woman abort an unwanted pregnancy (19%). Reporting both that a woman should have an abortion and that she should have her baby adopted increased with education level. Increased socioeconomic status was associated with increased approval of abortion for an unwanted pregnancy. Older men aged 45 -49 were the most likely to expect a woman to give birth and to raise the child (58%).

Attitudes and Perceptions about Gender Norms

Both male and female respondents were asked whether they agreed with several statements reflecting gender norms and roles. Responses to these statements provide a limited snapshot of the cultural expectations regarding men and women and the roles they play in society as perceived by the respondents.

The vast majority of surveyed women agreed that housework is a woman's main job (87%) but that a husband should help

with household chores if his wife works (94%) (Table 10.8). At least three quarters of women agree that a woman should be a virgin at her marriage, that women need their husbands' permission to work outside the home, and that every person should marry. If a woman works, 70% of women believe she should give the money she earns to her husband, while two-thirds (66%) of women know a woman can become pregnant the first time she has sex. The least commonly held norms were that childcare is a woman's job (52%) and that men should have the final say in the family (42%). Agreement with all the statements, except those about a husband's help around the house and possible pregnancy at first sex, was greater among rural women and women with less education. With the exception of "husband helping with chores," agreement was inversely related to education attainment.

Generally, women in metropolitan Tirana are least likely to agree with these genderrelated statements and women in rural areas outside of Tirana are most likely to agree (Table 10.9A). Agreement tends to increase with age and with number of living children. On the whole, as socioeconomic status and level of education increase, agreement decreases. Currently married women are more likely to accept the norms and Orthodox Christian women are least likely to accept the norms.

In general, men most strongly agree that everyone should get married, that a woman's main job is housework, and that if a woman works her husband should help around the house (84% - 85%) (Table 10.8). More than half of Albanian men also agree that a woman needs her husband's permission to work; that, if she works, all of her money should go to her husband; that a woman should be a virgin until she marries, and that a woman can get pregnant the first time she has sex (64% - 71%). Slightly more than a third of Albanian men believe that childcare is a woman's job (36%) and that men have the final say in family matter (37%). As seen with the women, with the exception of a husband's responsibility to help with chores when his wife works and the belief that a woman can become pregnant at first sex, agreement with all statements was higher among rural men and decreased with higher educational attainment.

Belief that men should help working wives with household chores was nearly unaffected by men's demographic characteristics as was agreement that a woman can become pregnant the first time she has sex, with the exception of education noted above and the much lower agreement among men without any living children (Table 10.9B). Men from rural areas outside of metro Tirana were most likely to agree with nearly all the other gender norms. For the most part, more traditional norms were more popular with older men and agreement increased with number of living children. Marital status had no impact on whether men felt husbands should help their wives with chores, or have the final say in the family,

or that child care is a woman's job. For all other statements, higher proportions of currently married men agreed. Agreement with all the statements of gender norms decreased with socioeconomic status and level of education, although there was very little difference for beliefs that a woman can become pregnant at first sex or that a husband should help with household chores when his wife works. Muslim and Catholic men were very close in agreement, while Orthodox Christian men were, for the most part, less likely to agree with the selected gender norms. Otherwise, Orthodox men were more likely to agree that a woman should give her salary to her husband. Religion had no effect on attitudes toward helping with household chores.

When asked who should decide how many children a couple should have, most women reported that both the man and the woman should decide (96%) with little variation by urban/rural residence or education (Table 10.10). A majority of men also felt that the decision should be made jointly (89%), and that opinion increased with level of education (85% to 97%).

	Women 15-	-44	Men 15–49		
Characteristic	Mean Ideal Number of Children	No. of Cases*	Mean Ideal Number of Children	No. of Cases*	
Total	2.6	5,459	2.4	1,527	
Strata					
Metro Tirana	2.4	2,046	2.3	661	
Other Urban	2.6	1,750	2.3	472	
Other Rural	2.7	1,663	2.5	394	
Residence					
Urban	2.5	3,462	2.2	1,033	
Rural	2.7	1,997	2.5	494	
Age Group					
15–19	2.4	1,042	2.3	336	
20–24	2.5	907	2.2	166	
25–29	2.5	914	2.2	180	
30–34	2.7	1,025	2.3	232	
35–39	2.7	904	2.4	223	
40–44	2.9	667	2.5	257	
45–49	N/A	N/A	2.8	133	
Marital Status					
Currently Married	2.7	3,807	2.5	924	
Previously Married	2.4	82	**	12	
Never Married	2.4	1,570	2.3	591	
Number of Living Children					
0	2.4	1,851	2.2	686	
1	2.4	805	2.1	199	
2	2.5	1,792	2.3	445	
3 or More	3.1	1,011	3.1	197	
Educational Level					
Primary or Less	2.7	2,347	2.5	568	
Secondary	2.5	2,423	2.3	741	
Post-Secondary	2.3	689	2.2	218	
Socioeconomic Status					
Low	2.7	1,789	2.5	538	
Middle	2.6	2,909	2.3	727	
High	2.4	761	2.2	262	

Table 10.1 Mean Ideal Number of Children for a Young Family in Albania by Selected Characteristics Among Women Aged 15–44 and Men Aged 15–49 Reproductive Health Survey: Albania, 2002

*Excludes non-numeric responses: 4% of female and 12% of male respondents.

**Percentages are not shown when base is less than 25 cases.

Table 10.2 A
Percent Distribution of Women's Opinion
on the Most Likely Time in the Menstrual Cycle for a Woman to Become Pregnant
by Selected Characteristics Among Women 15–44
Reproductive Health Survey: Albania, 2002

	Just Before Menses	During	Just After Menses	Halfway Between		Do Not		No. of
Characteristic	Begins	Menses	Ends	Menses	Anytime	Know	Total	Cases
Total	3.4	1.6	25.6	17.7	5.7	46.0	100.0	5,697
Strata								
Metro Tirana	4.3	1.8	26.4	22.7	7.6	37.2	100.0	2,108
Other urban	3.4	1.4	29.3	22.4	4.7	38.8	100.0	1,816
Other rural	3.1	1.6	23.3	13.8	5.7	52.5	100.0	1,773
Residence								
Urban	3.9	1.6	28.7	23.2	5.8	36.8	100.0	3,572
Rural	3.0	1.6	23.2	13.6	5.6	53.0	100.0	2,125
Age Group								
15–19	2.5	0.8	10.3	11.2	3.3	71.9	100.0	1,094
20–24	2.9	0.9	20.5	19.1	5.4	51.2	100.0	936
25–29	2.3	1.5	35.7	15.4	7.2	37.9	100.0	946
30–34	4.1	1.9	29.3	22.2	5.9	36.6	100.0	1,067
35–39	3.6	2.0	33.7	18.7	5.9	36.2	100.0	958
40–44	5.3	2.8	29.9	22.2	7.4	32.3	100.0	696
Marital Status								
Currently Married	4.0	1.8	33.5	19.6	6.2	34.9	100.0	3,965
Previously Married	2.5	3.5	36.5	19.6	9.7	28.2	100.0	88
Never Married	2.2	1.1	9.2	13.9	4.3	69.2	100.0	1,644
Number of Living Cl	hildren							
0	2.3	1.2	11.6	15.1	4.4	65.4	100.0	1,943
1	4.0	1.3	37.6	19.1	5.2	32.9	100.0	828
2	3.0	1.9	33.1	22.3	6.9	32.8	100.0	1,840
3 or more	5.3	2.0	33.3	15.9	6.6	36.9	100.0	1,086
Educational Level								
Primary or Less	3.0	1.7	23.1	11.3	6.4	54.5	100.0	2,519
Secondary	3.8	1.5	28.7	21.4	4.7	39.9	100.0	2,483
Postsecondary	3.6	1.1	27.7	43.4	5.6	18.6	100.0	695
Socioeconomic Stat	tus							
Low	2.9	1.6	23.3	11.4	6.0	54.8	100.0	1,940
Middle	3.7	1.6	26.3	21.9	5.2	41.2	100.0	2,985
High	3.8	1.1	33.0	25.2	6.8	30.1	100.0	772

	In the Mens by Selec	strual Cy ted Chai	vcle for a W racteristics re Health Su	oman to E Among M	Become Pi len Aged 1	regnant 5–49	C	
Characteristic	Just Before Menses Begins	During Menses	Right After Menses Ends	Halfway Between Menses	Anytime	Do Not Know	Total	No. of Cases
Total	4.3	1.9	9.3	10.5	14.9	59.0	100.0	1,740
Strata								
Metro Tirana	3.9	2.0	10.9	13.2	20.2	49.9	100.0	718
Other urban	4.5	3.2	9.9	14.6	11.2	56.7	100.0	547
Other rural	4.4	1.1	8.4	7.0	15.0	64.0	100.0	475
Residence								
Urban	4.3	2.8	10.4	14.3	14.5	53.7	100.0	1,155
Rural	4.3	1.1	8.4	7.2	15.2	63.7	100.0	585
Age Group								
15–19	4.0	0.6	3.6	3.4	6.3	82.1	100.0	401
20–24	4.5	1.5	10.6	12.0	6.5	65.0	100.0	189
25–29	6.6	2.5	11.6	13.1	16.4	49.8	100.0	218
30–34	5.7	3.3	15.6	13.1	20.7	41.5	100.0	253
35–39	3.4	3.9	10.1	10.6	25.3	46.7	100.0	255
40–44	4.6	1.3	7.5	14.4	15.5	56.6	100.0	277
45–49	1.0	0.5	8.6	9.2	17.3	63.4	100.0	147
Marital Status								
Currently Married	4.4	2.3	10.7	11.2	20.3	51.2	100.0	1,023
Previously Married	**	**	**	**	**	**	**	14
Never Married	4.3	1.4	7.1	9.5	6.7	71.1	100.0	703
Number of Living Ch	ildren							
0	4.3	1.2	8.6	10.1	8.4	67.4	100.0	815
1	8.0	4.2	13.5	13.0	21.1	40.1	100.0	221
2	3.0	3.3	10.3	12.8	21.4	49.3	100.0	468
3 or more	3.8	0.3	7.2	6.9	18.2	63.6	100.0	236
Educational Level								
Primary or Less	2.8	1.5	8.2	5.4	14.7	67.4	100.0	689
Secondary	6.3	1.3	10.3	11.9	14.7	55.5	100.0	825
Postsecondary	3.4	7.1	10.8	31.1	16.9	30.8	100.0	226
Socioeconomic Statu	IS							
Low	2.8	1.8	6.4	5.1	17.0	66.9	100.0	638
Middle	5.2	1.3	11.8	13.0	13.0	55.7	100.0	814
High	7.8	4.8	12.5	24.5	13.0	37.5	100.0	288

Table 10.2 B

Percent Distribution of Men's Opinion on the Most Likely Time

**Percentages are not shown when base is less than 25 cases.

Table 10.3 A

Percent Distribution of Women's Opinion on the Likelihood of Pregnancy While Breastfeeding by Selected Characteristics Among Women Aged 15–44 Reproductive Health Survey: Albania 2002

Characteristic	Higher Risk	Lower Risk	Same Risk	Don't Know	Total	No. of Cases
Total	9.3	20.3	23.1	47.3	100.0	5,697
Strata						
Metro Tirana	9.4	23.8	23.0	43.7	100.0	2,108
Other urban	10.0	23.5	23.5	43.0	100.0	1,816
Other rural	8.9	17.6	22.8	50.6	100.0	1,773
Residence						
Urban	9.7	24.1	23.6	42.6	100.0	3,572
Rural	9.0	17.5	22.7	50.7	100.0	2,125
Age Group						
15–19	4.8	7.4	7.6	80.2	100.0	1,094
20–24	6.6	16.6	18.3	58.5	100.0	936
25–29	10.2	24.6	31.0	34.1	100.0	946
30–34	11.1	27.0	29.6	32.3	100.0	1,067
35–39	13.0	23.5	31.1	32.4	100.0	958
40–44	12.0	28.1	26.7	33.1	100.0	696
Marital Status						
Currently Married	11.6	25.8	29.5	33.1	100.0	3,965
Previously Married	11.3	38.5	27.9	22.4	100.0	88
Never Married	4.7	8.4	10.1	76.9	100.0	1,644
Number of Living Children						
0	4.9	10.2	10.3	74.6	100.0	1,943
1	10.0	24.8	30.9	34.4	100.0	828
2	12.0	29.8	30.0	28.3	100.0	1,840
3 or more	13.2	23.6	31.9	31.4	100.0	1,086
Educational Level						
Primary or Less	9.4	17.9	21.3	51.4	100.0	2,519
Secondary	9.4	22.0	24.4	44.2	100.0	2,483
Postsecondary	8.5	28.7	28.9	33.9	100.0	695
Socioeconomic Status						
Low	9.5	18.3	22.9	49.4	100.0	1,940
Middle	9.1	21.4	22.7	46.8	100.0	2,985
High	10.1	24.5	26.6	38.9	100.0	772

Characteristic	Higher Risk	Lower Risk	Same Risk	Don't Know	Total	No. of Cases
Total	1.1	19.3	13.8	65.8	100.0	1,740
Strata						
Metro Tirana	2.2	20.0	12.8	64.9	100.0	718
Other urban	1.3	22.4	18.9	57.4	100.0	547
Other rural	0.6	17.1	11.2	71.1	100.0	475
Residence						
Urban	1.7	21.8	16.3	60.2	100.0	1,155
Rural	0.6	17.1	11.7	70.7	100.0	585
Age Group						
15–19	0.4	6.1	7.4	86.1	100.0	401
20–24	1.1	14.7	11.5	72.8	100.0	189
25–29	0.4	23.5	16.0	60.2	100.0	218
30–34	2.1	26.0	20.7	51.3	100.0	253
35–39	1.1	21.6	18.7	58.6	100.0	255
40–44	2.3	26.2	12.4	59.2	100.0	277
45–49	0.6	22.6	12.3	64.5	100.0	147
Marital Status						
Currently Married	1.4	24.6	16.6	57.4	100.0	1,023
Previously Married	**	**	**	**	**	14
Never Married	0.7	11.1	9.5	78.7	100.0	703
Number of Living Children						
0	0.7	11.5	11.0	76.8	100.0	815
1	1.5	32.7	18.7	47.1	100.0	221
2	1.9	25.7	17.9	54.5	100.0	468
3 or more	0.9	21.5	12.0	65.6	100.0	236
Educational Level						
Primary or Less	0.7	15.0	11.7	72.6	100.0	689
Secondary	1.6	19.9	16.2	62.2	100.0	825
Postsecondary	1.1	38.9	13.5	46.6	100.0	226
Socioeconomic Status						
Low	0.8	15.7	12.7	70.7	100.0	638
Middle	1.1	20.9	13.4	64.6	100.0	814
High	2.4	28.6	20.0	49.0	100.0	288

Table 10.3 BPercent Distribution of Men's Opinion on the Likelihood of Pregnancy While Breastfeeding
by Selected Characteristics Among Men Aged 15–44
Reproductive Health Survey: Albania 2002

**Percentages are not shown when base is less than 25 cases.

Characteristic	Alucese	Under Certain	Never	Den't Know	Total	No. of
Characteristic	Always	Circumstances	Never	Don't Know	Total	Cases
Total	72.4	26.2	0.9	0.6	100.0	5,697
Strata						
Metro Tirana	73.1	25.6	0.9	0.3	100.0	2,108
Other urban	74.4	24.7	0.7	0.2	100.0	1,816
Other rural	71.1	27.2	0.9	0.8	100.0	1,773
Residence						
Urban	74.3	24.8	0.7	0.2	100.0	3,572
Rural	70.9	27.3	1.0	0.8	100.0	2,125
Age						
15–19	70.8	26.6	0.6	1.9	100.0	1,094
20–24	72.4	26.7	0.7	0.2	100.0	936
25–29	74.0	24.3	1.4	0.3	100.0	946
30–34	73.6	25.6	0.6	0.2	100.0	1,067
35–39	72.0	27.5	0.5	0.0	100.0	958
40–44	72.1	26.2	1.3	0.4	100.0	696
Marital Status						
Currently Married	72.8	26.2	0.7	0.3	100.0	3,965
Previously Married	77.0	23.0	0.0	0.0	100.0	88
Never Married	71.2	26.4	1.3	1.1	100.0	1,644
Number of Living Chil	dren					
0	71.6	26.3	1.2	1.0	100.0	1,943
1	72.9	26.2	0.4	0.5	100.0	828
2	72.7	25.9	1.1	0.3	100.0	1,840
3 or more	73.1	26.4	0.3	0.2	100.0	1,086
Educational Level						
Primary or Less	71.5	26.6	1.0	0.9	100.0	2,519
Secondary	72.5	26.7	0.6	0.1	100.0	2,483
Postsecondary	77.8	21.1	1.1	0.0	100.0	695
Socioeconomic Status	5					
Low	69.3	28.5	1.0	1.3	100.0	1,940
Middle	74.6	24.5	0.9	0.1	100.0	2,985
High	75.2	24.6	0.2	0.0	100.0	772

by Selected Characteristics Among Men Aged 15–49 Reproductive Health Survey: Albania, 2002									
Characteristic	Always	Under Certain Circumstances	Never	Don't Know	Total	No. of Cases			
Total	52.9	44.9	1.0	1.2	100.0	1,740			
Strata									
Metro Tirana	48.6	47.5	0.8	3.0	100.0	718			
Other urban	61.1	38.2	0.7	0.0	100.0	547			
Other rural	49.8	48.0	1.1	1.2	100.0	475			
Residence									
Urban	56.8	41.3	0.7	1.2	100.0	1,155			
Rural	49.5	48.1	1.2	1.2	100.0	585			
Age									
15–19	54.0	41.2	0.6	4.2	100.0	401			
20–24	52.5	46.6	0.0	0.8	100.0	189			
25–29	53.3	45.9	0.9	0.0	100.0	218			
30–34	56.1	43.2	0.3	0.3	100.0	253			
35–39	50.3	48.1	1.3	0.3	100.0	255			
40–44	58.8	39.5	1.1	0.6	100.0	277			
45–49	43.6	52.8	2.9	0.8	100.0	147			
Marital Status									
Currently Married	52.7	45.5	1.3	0.4	100.0	1,023			
Previously Married	**	**	**	**	**	14			
Never Married	53.3	44.0	0.4	2.3	100.0	703			
Number of Living Children									
0	53.0	44.3	0.7	2.0	100.0	815			
1	51.0	47.0	1.4	0.6	100.0	221			
2	55.5	42.9	1.0	0.6	100.0	468			
3 or more	50.5	48.0	1.3	0.2	100.0	236			
Educational Level									
Primary or Less	50.4	46.5	1.5	1.6	100.0	689			
Secondary	55.1	43.7	0.5	0.7	100.0	825			
Postsecondary	56.4	42.4	0.3	0.9	100.0	226			
Socioeconomic Status									
Low	51.9	45.4	1.5	1.2	100.0	638			
Middle	52.7	45.5	0.6	1.3	100.0	814			
High	58.5	40.8	0.1	0.6	100.0	288			

Table 10.4 B Percentage Distribution of Men's Agreement that Abortion Is a Woman's Personal Decision by Selected Characteristics Among Men Aged 15–49 Reproductive Health Survey: Albania, 2002

**Percentages are not shown when base is less than 25 cases.

a di

Table 10.5

Percent Distribution of Opinion Regarding Abortion Under Selected Circumstances by Selected Characteristics Among Women and Men of Reproductive Age Who Do Not Believe That Abortion Is Always a Woman's Personal Decision Reproductive Health Survey: Albania, 2002

	Women Aged 15–44								
		_							
Circumstances	Yes	No	Depends	Don't Know	Total	No. of Cases			
Woman's Life is Endangered	80.4	9.5	5.9	4.2	100.0	1,492			
Fetus is Malformed	70.1	18.6	5.0	6.4	100.0	1,492			
Woman's Health is Endangered	59.1	20.1	15.1	5.6	100.0	1,492			
Pregnancy is Result of Rape	53.1	20.2	16.9	9.8	100.0	1,492			
Woman is Unmarried	37.9	31.6	21.1	9.4	100.0	1,492			
Family Cannot Afford the Child	21.9	58.6	11.8	7.6	100.0	1,492			

	Men Aged 15–49								
		Acceptability of Abortion							
Circumstances	Yes	No	Depends	Don't Know	Total	No. of Cases			
Woman's Life is Endangered	86.5	4.0	6.4	3.2	100.0	818			
Woman's Health is Endangered	73.7	9.6	11.7	5.0	100.0	818			
Pregnancy is Result of Rape	59.0	11.6	20.2	9.2	100.0	818			
Fetus is Malformed	55.5	14.6	16.0	14.0	100.0	818			
Woman is Unmarried	17.9	46.1	22.5	13.5	100.0	818			
Family Cannot Afford the Child	13.9	45.4	22.6	18.0	100.0	818			

Albania Reproductive Health Survey, 2002 Final Report

Table 10.6 A Percentage Who Agree with the Acceptability of Abortion Under Selected Circumstances by Selected Characteristics Among Women Aged 15–44 Who Do Not Believe That Abortion is Always a Woman's Personal Decision Reproductive Health Survey: Albania, 2002

		Circumstances							
Characteristic	Woman's Life in Danger	Fetus Deformed	Woman's Health in Danger	Pregnancy Resulted From Rape	Woman Unmarried	Cannot Afford Child	No. of Cases		
Total	80.4	70.1	59.1	53.1	37.9	21.9	1,492		
Strata									
Metro Tirana	82.7	69.9	66.7	47.2	25.5	20.9	557		
Other urban	83.1	73.3	59.0	54.5	35.9	27.7	444		
Other rural	78.4	68.6	57.1	54.0	42.3	19.4	491		
Residence									
Urban	83.6	72.4	62.9	52.3	31.7	25.8	886		
Rural	78.2	68.6	56.6	53.6	42.0	19.3	606		
Age									
15–19	74.8	56.4	50.8	46.8	33.5	13.8	290		
20–24	81.0	69.2	61.9	55.6	44.0	20.9	244		
25–29	76.9	73.7	61.9	53.7	39.5	21.5	238		
30–34	87.8	74.5	63.0	56.5	40.3	28.2	284		
35–39	85.9	79.7	63.5	54.1	35.7	28.6	253		
40–44	78.1	73.2	57.2	54.6	36.2	22.1	183		
Marital Status									
Currently Married	83.8	75.0	62.1	55.3	39.4	24.6	1,049		
Previously Married	**	**	**	**	**	**	19		
Never Married	74.1	61.1	52.7	48.6	34.5	16.2	424		
Number of Living Children									
0	74.8	62.0	54.0	48.7	35.1	16.5	500		
1	85.7	77.4	63.7	59.8	43.8	22.0	211		
2	80.9	75.9	64.2	54.2	35.8	26.6	493		
3 or more	86.7	73.1	59.5	55.7	42.3	25.7	288		
Educational Level									
Primary or Less	76.0	66.1	56.5	52.4	41.8	18.6	700		
Secondary	85.4	75.4	55.2	60.8	32.9	26.2	641		
Postsecondary	88.8	72.9	71.3	47.0	33.9	25.2	151		
Socioeconomic Status									
Low	76.4	65.0	55.3	50.6	42.4	19.9	576		
Middle	82.9	74.7	61.7	55.0	33.9	23.6	746		
High	90.4	74.1	67.5	57.3	34.5	23.7	170		

**Percentages are not shown when base is less than 25 cases.

	Circumstances								
Characteristic	Woman's Life in Danger	Woman's Health in Danger	Pregnancy Resulted From Rape	Fetus Deformed	Woman Unmarried	Cannot Afford Child	No. of Cases		
Total	86.5	73.7	59.0	55.5	17.9	13.9	818		
Strata									
Metro Tirana	86.3	75.3	60.3	60.3	28.0	19.2	365		
Other urban	88.2	77.4	65.2	56.0	13.4	12.8	209		
Other rural	85.8	71.3	55.6	53.3	16.0	12.4	244		
Residence									
Urban	87.7	76.5	62.8	58.4	19.0	15.5	516		
Rural	85.6	71.6	56.1	53.3	17.1	12.8	302		
Age									
15–19	76.5	68.5	58.3	42.0	17.4	3.8	187		
20–24	88.4	74.8	57.9	56.3	12.9	14.9	89		
25–29	85.3	81.3	57.1	55.1	16.7	12.5	108		
30–34	90.4	74.7	59.6	60.4	19.2	19.1	112		
35–39	92.4	81.0	67.7	60.5	20.0	17.1	124		
40–44	88.7	78.6	61.1	69.0	21.7	23.0	126		
45–49	87.3	58.6	51.0	50.9	18.5	11.4	72		
Marital Status									
Currently Married	88.6	73.5	59.6	57.5	19.9	17.4	486		
Previously Married	**	**	**	**	**	**	7		
Never Married	83.1	73.9	57.7	51.9	15.0	8.7	325		
Number of Living Children									
0	84.2	75.0	58.3	53.0	16.5	9.6	382		
1	82.9	72.6	61.6	48.4	17.5	10.9	115		
2	90.1	75.4	62.7	60.0	19.0	20.1	214		
3 or more	89.9	69.1	54.3	60.1	20.4	18.4	107		
Educational Level									
Primary or Less	83.9	70.5	56.2	48.2	19.5	12.1	339		
Secondary	88.9	76.8	63.6	62.2	16.4	14.9	376		
Postsecondary	91.1	77.9	53.7	66.9	15.5	20.0	103		
Socioeconomic Status									
Low	83.5	69.9	57.4	51.3	16.5	14.0	299		
Middle	88.6	75.8	58.7	56.0	17.5	12.8	388		
High	92.8	83.9	68.6	74.6	27.3	18.8	131		

Table 10.6 B Percentage Who Agree with the Acceptability of Abortion Under Selected Circumstances by Selected Characteristics Among Men Aged 15–49 Who Do Not Believe That Abortion Is Always a Woman's Personal Decision Reproductive Health Survey: Albania, 2002

**Percentages are not shown when base is less than 25 cases.

Table 10.7 A

Percent Distribution of Women's Opinion on What a Woman Should Do If a Pregnancy is Unwanted by Selected Characteristics Among Women Aged 15–44 Reproductive Health Survey: Albania, 2002

	What Should A Woman Do If A Pregnancy is Unwanted									
			Give Birth and							
Characteristic	Give Birth and Keep the Baby	Have an Abortion	Have Baby Adopted	Don't Know	Total	No. of Cases				
Total	70.8	20.5	2.3	6.4	100.0	5,697				
	10.0	20.5	2.5	0.4	100.0	5,057				
Strata	05.7	04.4	0.0	0.0	100.0	0.400				
Metro Tirana	65.7	24.4	3.0	6.9	100.0	2,108				
Other urban	66.9	24.0	2.6	6.5	100.0	1,816				
Other rural	74.5	17.5	1.9	6.1	100.0	1,773				
Residence										
Urban	65.9	24.7	2.9	6.6	100.0	3,572				
Rural	74.6	17.4	1.8	6.2	100.0	2,125				
Age										
15–19	69.9	17.8	3.4	8.9	100.0	1,094				
20–24	72.8	18.1	2.7	6.4	100.0	936				
25–29	70.0	22.2	2.4	5.4	100.0	946				
30–34	69.2	24.3	1.3	5.1	100.0	1,067				
35–39	71.2	21.5	1.6	5.7	100.0	958				
40–44	72.2	20.3	1.8	5.6	100.0	696				
Marital Status										
Currently Married	71.4	21.9	1.6	5.0	100.0	3,965				
Previously Married	67.6	25.1	0.6	6.7	100.0	88				
Never Married	69.9	17.5	3.7	8.9	100.0	1,644				
Number of Living Children										
0	71.0	17.4	3.4	8.2	100.0	1,943				
1	69.6	24.7	1.8	3.9	100.0	828				
2	70.3	21.6	1.6	6.6	100.0	1,840				
3 or more	72.0	22.2	1.5	4.3	100.0	1,086				
Educational Level										
Primary or Less	73.8	18.1	1.5	6.6	100.0	2,519				
Secondary	68.0	22.6	2.9	6.2	100.0	2,483				
Postsecondary	64.4	26.1	4.3	5.2	100.0	695				
Socioeconomic Status										
Low	73.4	17.1	2.0	7.5	100.0	1,940				
Middle	70.7	21.4	2.4	5.6	100.0	2,985				
High	58.6	32.9	3.4	5.1	100.0	772				

Table 10.7 BPercent Distribution of Men's Opinion on What a Woman Should Do If aPregnancy is Unwanted by Selected Characteristics Among Men Aged 15–49Reproductive Health Survey: Albania, 2002

	What Should A Woman Do If A Pregnancy is Unwanted									
			Give Birth and							
Characteristic	Give Birth and Keep the Baby	Have an Abortion	Have Baby Adopted	Don't Know	Total	No. of Cases				
Total	40.2	23.4	9.6	26.8	100.0	1,740				
Strata										
Metro Tirana	37.7	29.0	9.5	23.7	100.0	718				
Other urban	39.1	28.7	11.1	21.2	100.0	547				
Other rural	41.8	18.1	8.7	31.3	100.0	475				
Residence										
Urban	38.6	29.0	10.6	21.8	100.0	1,155				
Rural	41.6	18.6	8.8	31.1	100.0	585				
Age										
15–19	32.3	19.0	10.5	38.1	100.0	401				
20–24	30.8	28.0	9.5	31.8	100.0	189				
25–29	41.6	21.6	9.5	27.3	100.0	218				
30–34	38.9	29.2	7.4	24.5	100.0	253				
35–39	41.4	21.4	13.2	24.0	100.0	255				
40–44	44.3	26.4	11.5	17.8	100.0	277				
45–49	58.2	19.0	4.3	18.6	100.0	147				
Marital Status										
Currently Married	45.4	22.5	9.5	22.6	100.0	1,023				
Previously Married	**	**	**	**	**	14				
Never Married	31.9	25.0	9.9	33.2	100.0	703				
Number of Living Children										
0	34.7	24.1	9.6	31.6	100.0	815				
1	45.9	21.0	6.2	26.9	100.0	221				
2	42.4	25.3	11.1	21.2	100.0	468				
3 or more	47.1	20.8	9.8	22.3	100.0	236				
Educational Level										
Primary or Less	41.8	20.2	6.0	32.0	100.0	689				
Secondary	39.1	24.7	12.4	23.8	100.0	825				
Postsecondary	36.5	35.1	15.4	13.1	100.0	226				
Socioeconomic Status										
Low	43.2	20.1	7.1	29.5	100.0	638				
Middle	37.5	24.4	12.4	25.7	100.0	814				
High	37.4	34.4	9.3	18.9	100.0	288				

**Percentages are not shown when base is less than 25 cases.

Table 10.8Percentage Who Agree with Statements on Gender Norms
by Residence and EducationAmong Women Aged 15–44 and Men Aged 15–49Reproductive Health Survey: Albania 2002

	Women Aged 15–44							
		Residence Educ			Education Le	ucation Level		
Statements	Total	Urban	Rural	Primary	Secondary	Post Secondary		
Husband Should Help with Chores if Wife Works	93.6	93.2	93.9	93.3	94.2	93.0		
Main Job of Woman is Housework	86.6	79.5	91.9	93.2	83.5	57.4		
Every Individual Should Get Married	78.9	70.5	85.1	85.5	74.9	53.3		
A Married Woman Needs Husband's Permission to Work	75.2	62.3	84.8	86.7	67.8	33.0		
A Woman Should Be A Virgin When She Marries	74.7	57.6	87.6	87.2	66.2	31.9		
If Woman Works, She Should Give Money to Husband	69.9	57.2	79.5	82.5	59.7	34.4		
A Woman Can Get Pregnant The First Time She Has Sex	65.5	65.7	65.4	64.5	66.6	67.0		
Child Care Is A Woman's Job	52.3	46.4	56.7	62.4	44.0	23.6		
Men Have Final Say in Family Matters	41.7	30.5	50.0	55.4	29.0	9.7		
Number of Cases	5,697	3,572	2,125	2,519	2,483	695		

	Men Aged 15–49							
		Residence			vel			
Statements	Total	Urban	Rural	Primary	Secondary	Post Secondary		
Husband Should Help with Chores if Wife Works	85.2	85.2	85.2	83.6	85.5	92.5		
Main Job of Woman is Housework	84.7	79.0	89.7	90.0	83.3	63.4		
Every Individual Should Get Married	84.2	77.5	90.0	88.0	83.1	68.1		
A Woman Can Get Pregnant The First Time She Has Sex	70.5	72.9	68.5	69.2	70.0	80.2		
A Woman Should Be A Virgin When She Marries	67.9	54.2	79.8	78.7	63.6	30.3		
If Woman Works, She Should Give Money to Husband	64.1	58.0	69.5	73.1	59.7	37.1		
A Married Woman Needs Husband's Permission to Work	64.1	56.8	70.5	71.8	60.2	42.0		
Men Have Final Say in Family Matters	37.0	30.3	42.7	50.0	26.9	14.5		
Child Care Is A Woman's Job	35.9	32.4	39.0	44.5	30.1	17.5		
Number of Cases	1,740	1,155	585	689	825	226		

Tables

Chapter 10
Table 10.9 A
Percentage Who Agree with Statements on Gender Norms
by Selected Characteristics Among Women Aged 15–44
Reproductive Health Survey: Albania, 2002

	Reproductive Health Survey: Albania, 2002									
Characteristic	Husband Should Help With Chores	Main Job of Woman is Housework	All People Should Marry	Need Husband's Permission To Work	Woman Should be Virgin at Marriage	lf Woman Works, All Money to Husband	Can Get Pregnant at First Sex	Child Care is a Woman's Job	Men Have Final Say in Family	No. of Cases
Total	93.6	86.6	78.9	75.2	74.7	69.9	65.5	52.3	41.7	5,697
Strata										
Metro Tirana	90.3	77.0	66.9	57.6	54.0	55.5	65.9	46.1	31.5	2,108
Other urban	94.8	81.9	73.0	66.8	62.2	60.0	65.3	46.6	31.2	1,816
Other rural	93.9	92.0	85.5	84.9	87.6	79.6	65.5	57.2	50.3	1,773
Residence										
Urban	93.2	79.5	70.5	62.3	57.6	57.2	65.7	46.4	30.5	3,572
Rural	93.9	91.9	85.1	84.8	87.6	79.5	65.4	56.7	50.0	2,125
Age										
15–19	89.9	80.3	72.4	71.2	72.7	63.9	41.8	52.0	40.4	1,094
20–24	93.4	85.0	75.1	76.3	70.2	69.1	58.2	49.6	40.0	936
25–29	94.1	87.1	80.9	75.7	71.2	72.1	70.8	55.3	44.0	946
30–34	96.4	88.3	79.0	76.1	75.3	74.0	75.1	53.3	44.6	1,067
35–39	95.2	89.2	83.8	75.7	78.6	71.7	76.5	54.0	41.5	958
40–44	94.0	92.2	84.9	77.4	81.7	71.0	79.9	49.9	40.1	696
Marital Status										
Currently Married Previously	94.7	90.7	83.5	79.3	77.9	74.3	76.4	55.1	44.8	3,965
Married	91.2	84.7	76.6	74.3	74.7	70.8	71.7	66.3	42.4	88
Never Married	91.7	78.5	69.9	67.0	68.5	61.3	43.4	45.9	35.3	1,644
Number of Living C	Children									
0	91.8	79.7	71.0	68.4	69.0	62.7	46.0	48.0	37.2	1,943
1	93.0	89.0	81.6	80.4	70.5	75.3	74.7	58.3	46.6	828
2	95.9	88.4	81.1	74.2	74.0	69.6	75.6	51.1	38.9	1,840
3 or more	94.3	94.7	87.9	84.8	87.9	79.6	81.0	57.6	49.8	1,086
Educational Level										
Primary or less	93.3	93.2	85.5	86.7	87.2	82.5	64.5	62.4	55.4	2,519
Secondary	94.2	83.5	74.9	67.8	66.2	59.7	66.6	44.0	29.0	2,483
Postsecondary	93.0	57.4	53.3	33.0	31.9	34.4	67.0	23.6	9.7	695
Socioeconomic Sta	atus									
Low	93.5	92.7	84.8	84.7	87.6	80.6	65.6	60.0	54.0	1,940
Middle	94.1	84.3	76.6	71.7	69.4	65.4	65.2	48.7	34.6	2,985
High	91.7	69.6	62.4	47.1	40.6	42.7	66.9	34.6	20.8	772
Religion*										
Muslim Orthodox	93.4	87.9	79.9	76.9	76.6	72.0	65.4	52.9	43.3	4,568
Christian	94.9	75.7	69.9	59.2	55.7	49.9	69.0	36.5	22.8	531
Catholic	94.1	86.7	79.6	77.2	77.0	72.4	62.9	61.4	45.2	530

* Excludes 68 women of other or undeclared religion.

Table 10.9 B Percentage Who Agree with Statements on Gender Norms by Selected Characteristics Among Men Aged 15–49 Reproductive Health Survey: Albania, 2002										
Characteristic	Husband Should Help With Chores	Main Job of Woman is Housework	All People 31	Can Get Pregnant at First	Woman Should AAA be Virgin at Marriage	If Woman Works, All Money to Husband	Need Husband's Permission To Work	Men Have Final Say in Family	Child Care Is A Woman's Job	No. of
Characteristic Total	85.2	<u>2 > 1</u> 84.7	84.2	70.5	<u> </u>	<u>= > 2 1</u> 64.1	<u> </u>	37.0	35.9	Cases 1,740
Strata	00.2	04.1	04.2	10.0	01.0	04.1	04.1	01.0	00.0	1,140
Metro Tirana	79.8	76.8	74.5	69.5	55.5	53.8	55.4	37.3	37.1	718
Other urban	79.8 87.4	80.5	74.5 79.9	75.4	55.5 54.7	53.8 59.7	61.1	27.8	31.0	547
Other rural	86.1	90.3	90.5	68.0	80.6	70.8	69.3	42.3	38.4	475
	00.1	90.5	90.5	00.0	00.0	70.0	09.5	42.3	30.4	475
Residence										
Urban	85.2	79.0	77.7	72.9	54.2	58.0	56.8	30.3	32.4	1,155
Rural	85.2	89.7	90.0	68.5	79.8	69.5	70.5	42.7	39.0	585
Age										
15–19	83.3	78.7	85.9	40.4	58.7	55.4	53.9	41.2	38.4	401
20–24	83.0	76.0	80.6	67.0	52.8	63.8	59.2	31.3	34.2	189
25–29	86.6	86.1	81.6	75.0	61.6	60.8	65.5	34.6	32.4	218
30–34	86.9	84.2	83.9	81.3	70.3	61.3	65.3	31.1	30.2	253
35–39	87.0	90.5	82.1	80.0	75.3	66.2	65.6	37.5	37.7	255
40–44	86.2	90.5	86.1	84.7	78.7	71.2	71.7	36.9	35.0	277
45–49	84.0	90.4	89.1	77.5	84.0	74.7	73.2	46.2	43.8	147
Marital Status										
Currently Married	86.1	88.9	86.8	79.4	78.7	68.5	69.8	38.0	36.5	1,023
Previously Married	**	**	**	**	**	**	**	**	**	14
Never Married	83.6	78.1	80.2	56.5	51.1	57.1	55.2	35.1	35.1	703
Number of Living Chil										
0	83.8	79.0	80.7	57.8	53.6	58.9	56.8	35.1	35.6	815
1	89.1	86.3	83.6	79.6	70.2	65.7	69.5	33.9	33.6	221
2	88.5	88.4	83.0 84.7	84.1	76.4	64.9	68.3	35.5 35.5	36.5	468
2 3 or more	81.9	93.1	92.4	78.1	90.4	75.1	73.5	45.6	37.5	236
	01.9	33.1	32.4	70.1	50.4	75.1	15.5	45.0	57.5	230
Educational Level					70 7	74.0	70.4	50.0		
Primary or Less	83.6	90.0	88.0	69.2	78.7	71.8	73.1	50.0	44.5	689
Secondary	85.5	83.3	83.1	70.0	63.6	60.2	59.7	26.9	30.1	825
Postsecondary	92.5	63.4	68.1	80.2	30.3	42.0	37.1	14.5	17.5	226
Socioeconomic Statu	S									
Low	85.4	92.0	88.3	69.4	79.6	72.0	69.2	47.9	44.6	638
Middle	85.4	80.5	82.4	71.0	60.3	59.1	61.3	29.3	30.3	814
High	83.7	68.9	72.8	73.8	46.0	49.3	52.8	18.7	19.7	288
Religion*										
Muslim	85.1	87.1	85.1	64.9	70.6	69.2	64.7	36.0	38.4	1,422
Orthodox Christian	85.4	68.3	73.4	56.2	43.1	77.6	57.3	32.6	23.2	181
Catholic	87.5	83.1	84.1	66.0	67.5	73.1	65.8	38.9	35.0	76

Table 10.9 B

**Percentages are not shown when base is less than 25 cases.

ables

	Women Aged 15–44								
		Resi	dence		Education Lev	vel			
Decision Maker	Total	Urban	Rural	Primary	Secondary	Post Secondary			
The Woman	0.7	1.0	0.4	0.6	0.6	1.6			
The Man	1.4	0.7	1.9	2.4	0.4	0.0			
Both	95.9	96.9	95.2	94.0	98.1	97.9			
The Mother-in-Law	0.0	0.0	0.0	0.0	0.0	0.0			
God	1.3	1.2	1.5	1.9	0.7	0.5			
Don't Know	0.7	0.2	1.0	1.1	0.2	0.0			
Total	100.0	100.0	100.0	100.0	100.0	100.0			
Number of Cases	5,697	3,572	2,125	2,519	2,483	695			

Chapter 10

		Resi	dence	Education Level					
Decision Maker	Total	Urban	Rural	Primary	Secondary	Post Secondary			
The Woman	0.3	0.4	0.2	0.0	0.6	0.3			
The Man	5.9	4.0	7.5	9.0	3.2	2.3			
Both	88.7	91.6	86.2	84.8	91.3	97.2			
The Mother-in-Law	1.5	0.7	2.3	0.9	2.6	0.0			
God	3.2	2.9	3.4	4.5	2.2	0.2			
Don't Know	0.4	0.5	0.4	0.7	0.1	0.0			
Total	100.0	100.0	100.0	100.0	100.0	100.0			
Number of Cases	1,740	1,155	585	689	825	226			

Men Aged 15-49

CHAPTER 11 HEALTH BEHAVIORS

espite a continuous reduction in the childhood mortality rate and the preservation of a relatively high life expectancy, the health of Albanians is being threatened by an increasing rate of chronic diseases, including cancers and cardiovascular diseases, which now are by far the most frequent health problems, accounting for about 75% of the deaths occurring in the country. In addition to demographic changes, these health problems are associated with an increase in risky behaviors, concurrent with dramatic political, economic and social changes in the last two decades. Among these risky behaviors, of significant importance are smoking and alcohol use. In Albania, as in many other post-communist societies of the former Soviet Union and other former eastern bloc countries in Europe, these new challenges are not being addressed appropriately by the health system, which often fails to provide adequate prevention services and treatment. Such inadequate services include hypertension screening and cervical cancer screening.

Tobacco is a potent human carcinogen that has been shown to be related to a significant number of cancers of the respiratory and digestive tracts, bladder, cervix and kidney. Cigarette smoking accounts for 87% of lung cancer deaths and 30% of all cancer deaths. Smoking is also considered a major risk factor for diseases of the circulatory system including heart attacks, strokes and blood clots of the legs and lungs. In addition, smoking contributes to increasing rates of chronic diseases of the lung such as emphysema and chronic bronchitis (Difranza and Lew, 1996).

There are data indicating that tobacco use is increasing in Albania. The transition toward a market economy and the arrival of international tobacco companies are some of the background factors related to the increase in tobacco use. In Albania, after several years of legislative debate, there are still no laws protecting the health of the population from cigarette smoking. Although unpublished data drawn from limited surveys indicate a high prevalence of smoking among men and a growing prevalence among women, no populationbased data existed until the Albania RHS. In these limited surveys, the prevalence of smoking is reported to be increasing rapidly, starting from a very low level in the late 1980's. The prevalence of smoking may be rising most rapidly among young women in Eastern Europe; a recent worldwide review estimated that among women of reproductive age who smoked, 40% were young adults (Jha et al., 2002). Smoking poses specific risks to women in addition to lung cancer and cardiovascular diseases: it may increase both the risk of cervical cancer and risks, for older women, associated with taking contraceptive pills. It also affects women's reproductive health by increasing the risk for early menopause, miscarriage, and low birth weight babies.


Cancer is a leading cause of death in women in most countries around the world. Among reproductive system cancers, breast and cervical cancer are the most common. In many developed countries screening systems are in place, contributing to better control and better survival rates. This is especially true in the case of cervical cancer. In developing countries most cases are detected at an advanced and incurable stage. A low awareness among the population, lack of information and mistrust towards possible treatments and a low priority for women's health issues adds to the lack of or inefficient screening services.

The Albania RHS provides information on the following specific health behaviors: health care seeking behavior, prevalence of routine gynecologic visits, cervical cancer screening, breast self-examination, fertility and pelvic inflammatory diseases and prevalence of tobacco and alcohol use. For some of these topics information is available for men too.

Use Of Health Care And Preventive Services

Patient's attitudes and behaviors regarding health care visits are important determinants of whether they receive routine counseling and screening, including hypertension screening and cervical and breast cancer screening. Recent data from Albania indicate a low use of health services by the population for both primary health care and secondary health care. Tables 11.1A and B and Figure 11.1 present the proportions of Albanian women and men who have visited a health facility during the past 12 months. All the data are shown by selected characteristics of respondents.

Slightly more than one in four women (26%) of reproductive age has sought services at a health facility during the year before the survey. This indicator includes all kinds of visits: cases of preventive services. illness. family planning, counseling and other health check ups. The proportion for men is much lower - almost the half of that of the women - indicating a less frequent use of health services by men of reproductive age. Women aged 25-34 years use the services more frequently than other age groups, especially when compared to those in the age group 15-19. This profile is influenced by an increase in the reproductive health related visits of women 25-34 years of age, as married



women use health services approximately twice as much as never married women. Pregnancies and caring for their children is a factor which increases the use of health services by mothers. Interestingly, with two or more children, the use of services slightly starts to decrease. Use of services is weakly related to socioeconomic status but does not have any strong relationship to education or employment status. Women in urban areas visit health facilities around 25% more frequently than women of reproductive age living in rural areas (Figure 11.1). Women classified in the highest socioeconomic category use the health services 50% more than those of the lowest category.

The men's profile in the use of health care appears to have more of an age influence; it continues to increase by five-year age groups beginning at 30-34 until ages 45-49. The distribution of the health services utilization rate also increases as the number of live children increases.

Table 11.2A shows the proportion of women who have received counseling for family planning methods during their visits to health facilities by type of facility. Only 16% of women who have used the health services during the 12 months previous to the survey received any counseling for family planning methods, representing only 4% of all women. In Tirana, women receive more counseling for family planning than in other areas. Women of higher socioeconomic status, higher education level and those employed also have higher rates of counseling.

Women who use private clinics for their needs are more likely to be exposed to family planning information and counseling (21%), compared to those who use public services (16%). Although the number of women in the sample using private clinics is small, the differences are more noteworthy for urban women, women 25-29, better educated women, high socioeconomic status and working women.

Although only 14% of men of reproductive age visited a health facility in the past 12 months, 83% received information or counseling regarding "control for pregnancy", which includes a partners maternal care, and 8% and 9% received information or counseling specifically on condoms and sexually transmitted infections, respectively (Table 11.2B). Counseling on condoms and STIs is highest for never married, 20-24 year old men, and those with no children.

The Albania RHS included questions about the reasons that may prevent women or men from getting medical advice or treatment for themselves. Questions on the specific problems shown in the Tables were asked of each respondent (Tables 11.3A and B). For both women and men the most important problem was financial: getting money needed for treatment prevented them from using health services (46% and 54%, respectively). Other factors for women included the need to go to the health services accompanied by someone (41%), and the distance to health facility (31%). For men, in addition to the distance (35%) was the fact that many among them did not know where to go (40%).

Prevalence Of Routine Gynecologic Visits Among Sexually Experienced Women

In the United States and Western Europe, it is generally recommended that women have a routine gynecologic examination every year after age 18, or even earlier if she has had sexual experience. In Albania, only 32% of sexually experienced women report that they have ever had a routine gynecologic examination (Table 11.4) and this represents the lowest rate reported among similar surveys carried out in countries of Eastern Europe and the Caucasus region. In these countries this proportion ranges from 58% to 93% (CDC and ORC MACRO, 2003).

When asked about routine gynecologic examinations in the last year, only 16% of sexually experienced women sampled in the Albania RHS answered that they have had such an exam. Again, this rate is the lowest among rates reported in similar surveys. In Eastern Europe, this proportion ranges from 65% to 70%, and in the Caucasus region, from 22% in Azerbaijan to 30% in Georgia (CDC and ORC MACRO, 2003).

Living in urban areas, and especially in Tirana, increases the likelihood of having routine gynecological examinations and having them more frequently than women in rural areas (20% versus 14% during the last year). Given sample sizes, there are no significant differences by age groups. The proportion of overall examinations among women with post secondary education is around 44%, while among women with primary education or less it is only 28%. Employment may be also positively associated with the indicator. Distribution of the proportion among married, unmarried, and previously married women most probably reflects their age distribution.

When asked about the most important reason why they have never had a routine gynecological examination (for those women with sexually experience only), most of the respondents answered that they were healthy or there is no need to receive a gynecological exam. A very low proportion (2%) mentioned problems related to the provider (Table 11.5). This reflects a very low awareness among women on the importance of cancer prevention and an information gap on the need for a routine gynecological examination. The low prevalence of routine gynecological exams could have a substantial negative effect on screening, counseling and health education. These findings raise the suggestion that much more effort must be made to modify general health beliefs and attitudes and to educate women about screening procedures and disease prevention. These efforts must be accompanied by other interventions aimed at increasing the access to and quality of services and better geographic coverage.

Breast Self Exam

According to registry data in Albania, breast cancer has increased from 13.8/100,000 population in 1994 to 20.6/100,000 in 2003 (IHP, 2004). It is generally accepted that secondary prevention for breast cancer has been proven to be effective. The aim of secondary prevention is to identify breast cancer, as early as possible, to increase the chances of survival after therapy. Examination of the breast may be as simple as a self examination but it also includes complicated procedures such as mammography, which for some populations might be not readily available or too expensive. A breast self exam is a very simple, self care procedure which can be performed by women by themselves and requires minimum training. Through a breast self exam, women can detect early changes in their breasts and then ask for further follow up by health specialists (Aubard Y et al., 2002; Rebentisch DP et al., 1995).

It was not the aim of the Albania RHS to evaluate any skills women might have

in applying a breast self-examination procedure. Questions were intended to explore the level of awareness about breast self exam and its prevalence. Only around half of the respondents have ever heard of the procedure and less than 10% have ever used it (Table 11.6).The level of awareness in the Albania RHS was higher in urban areas and among the employed; awareness levels also steadily increased with level of education. The rate was the lowest among women in the 15-19 age group.

Only 3.7% of women reported that they performed breast self-examination every month. Principal characteristics of these women were urban residency, higher level of education, employment, and ever having a routine gynecologic exam. The level of awareness and the utilization rate were always found to be higher among women who have received a routine gynecological examination. Nevertheless, more than 40% of these women (reporting at least one routine gynecological visit) were not aware about breast self examination, indicating a lack of counseling for this procedure by health professionals.

Cervical Cancer Screening

Risk factors for cervical cancer include early onset of sexual experience, a history of multiple sexual partners, smoking, HIV infection and infection with human papilloma virus. Every year, about 450, 000 new cases of cervical cancer are diagnosed all over the world (Parkin, et al, 1993). Age adjusted incidence rates range from 5 to 42 cases per 100,000 women with higher rates reported by developing countries. The cervical cancer proportional mortality is also much higher in developing countries. There are data from developed countries showing that while the frequency of cervical cancer in situ is increasing, cervical cancer mortality continues to decrease. This decline was mostly attributed to cervical screening programs (Miller AB, 1986).

Cervical cancer is the fourth cause of female mortality from cancers in Albania. Reported cervical cancer incidence is lower than other European countries accounting for 12.8% of deaths in 2000. However, there is an increasing trend of incidence from 3.1/100.000 in 1994 to 5.3/100,000 in 2003 (IPH, 2004).

Screening programs for cervical cancer are based on the use of the papanicolaou smear test (Pap test), a procedure which makes possible early identification of premalignant changes in the cells of the cervix and allows early intervention. Evaluation research has shown that performing pap test screening every three years reduces the risk of developing invasive cancer by more than 91% (Miller AB, 1986). Yearly screening increases this proportion only by 2 percentage points (93%), while the pap test screening performed every five years has a success rate of about 84%. In many countries health authorities have set up screening programs where sexually active women, or those who are at least 18 years old, are invited to have the test at least once in every three years. Among those who are older than 65 years old who have been regularly screened with normal results can continue the periodical tests at a lower frequency.

In Albania, the pap test procedure is offered in some gynecological-obstetrical centers and private clinics in Tirana, but there is no organized screening program in place. The Albania RHS produced an estimate of the extent of cervical screening in the general population of women of reproductive age. The questionnaire included a series of questions for all respondents regarding their pap test history: "have you ever had a cervical smear test (papanicolaou screening test)?"; "if yes, when did you have your last cervical smear test?"; and for those who never had screening, "what is the main reason you have never had a pap test?"

As shown in Table 11.7, among sexually experienced women, the percentage of those who reported ever having a pap test is very low (3%); among the lowest when compared to some other Eastern European countries. The result is similar to that reported by the above mentioned Knowledge, Attitude, Beliefs, and Practice (KABP) survey (2%). Since the prevalence of the pap smear test is generally very low, there is not enough variation to analyze characteristics of women with and without a test.

Routine gynecological visits can be considered opportunities to educate patients about healthy lifestyle choices and to promote appropriate screenings for preventable diseases such as cervical cancer. The Albania RHS found that having performed a routine gynecological examination is somewhat associated with pap test screening; nevertheless, only 8% of women, who have ever had a routine gynecological exam had a pap test too.

The large majority of women (70%) answered that the main reason why they have never had a pap test was because they have never heard of it (Table 11.8). The figure is very high and indicates that a serious information gap exists. This proportion is more than two times higher when compared to countries such as Romania, Moldova and Georgia, where the proportion of non-awareness ranges from 23% to 39%. Only in Azerbaijan is the proportion of women who never heard of the pap test slightly higher. Even 60% of women who have had a routine gynecologic exam have never heard of a pap smear.

Women of Tirana and, to some extent, those in other urban areas seem to be more informed about the pap test, compared to women living in rural areas. Only 48% in Tirana and 62% in other urban areas answered that they have never heard of the procedure. In rural areas this proportion increases to more than 80%. Education is also an important factor that influences the awareness of women about the pap test; only 33% of women with postsecondary education have never heard of it compared with more than 60% of other women. Employment is also positively associated with having heard of the pap test..

Apart from the 70% of women in the sample who have never heard of the test, another 13% answered that they were healthy and that they do not have gynecological problems. Another 10% of women answered that they have never been recommended to have the pap test by their physician.

These results clearly show the critical need for appropriate and sustained educational programs for general screening procedures in Albania. There is also a need for increasing the awareness of health professionals about the importance of screening for cervical cancer, and improving their practices when examining women of reproductive age.




Figure 11.2 is a stark demonstration of the need for preventive health behaviors among Albanian women.

Prevalence Of Selected Health Problems

In this section prevalence estimates are given for selected health problems (based on having been told by a physician that they have the problem) among women and men in Albania. For women these health problems include pelvic inflammatory disease (infection of the tubes of the uterus), anemia, high blood pressure, Pelvic and diabetes. inflammatory disease was investigated only among women 15-44 years who have had sexual experience. Diabetes and high blood pressure prevalence was studied among men 15-49 years of age. Apart from these health problems, men were also asked about two other conditions: hepatitis B and heart disease.

As mentioned above, to calculate the reported prevalence among the sample participants, only those who have been diagnosed by a doctor about the selected conditions were considered "positive". The question was formulated as follows: "has a doctor ever told you that you have had (selected health problem)?". This was the only way to standardize the answers of the respondents and to avoid maximizing the bias. On the other hand, the results extracted from the survey should be taken cautiously as they might be highly influenced by several factors, mainly the health seeking behavior of the different groups of the population. Furthermore, they do not reflect the current health status of the population. In general, the prevalence presented here are to be considered as minimums, as only the conditions already diagnosed and reported to the patient by a health professional are included. Generally, as seen in Tables 11.9A and B, the level

of self reporting of medical conditions in the sample was relatively low.

The prevalence of diagnosed pelvic inflammatory disease among sexually experienced women in Albania seems to be the lowest among countries having similar data. In selected countries of Eastern Europe and the Caucasus this prevalence ranges from 17% to 44% (CDC and ORC MACRO, 20034). In the Albania RHS it is around 11%.

Reported pelvic inflammatory disease is slightly higher in Tirana, it increases with age (until age 35-39 years), it is slightly lower among women with postsecondary education and it is, comparatively, very high among previously married women (about 7 times higher than never married women and almost 2 times higher than married women)

Diagnosed anemia has the highest prevalence among selected conditions included in the survey; almost 12% of women of reproductive health reported to have been told at least once by a doctor that they have this health problem. Nevertheless, this level remains lower than most of the other countries mentioned before. It is only higher than the prevalence reported by the survey carried out in Georgia. The distribution of self reported prevalence of anemia among selected categories of the Albanian population follows the same profile as the levels of pelvic inflammatory disease.

High blood pressure is the third condition most frequently reported (8.4%) and it is only slightly lower than that of Georgia, Romania and Moldova. Self reported high blood pressure in Albanian women increases with age; it is higher among previously married women and slightly higher among women living in rural areas; this later result might reflect the previously discussed distribution of high blood pressure during pregnancy.

The self reported high blood pressure among men of reproductive age is much lower then that of women and may be reflecting the lower utilization of health services by men, compared with women. Prevalence of heart disease among men 15-49 years of age is reported to be only 2% and it increases with age to reach a level of more than 8% among the age group 45-49.

The prevalence of self reported diagnosed diabetes in the total female sample is only 0.6%. Among men the level is 1.3%. The low prevalence does not allow the differences to be analyzed; however they are quite comparable to the prevalence produced in similar surveys in Romania, Moldova and Georgia (range 0.5%-1.1%).

Self-reported diagnosis of hepatitis B is at relatively high levels among men of reproductive age (6.1%). This level, as mentioned before, should be considered a minimum. In all the cases for women the most strongly associated factor associated with the rate of self reported conditions was the routine gynecological examination. For pelvic inflammatory diseases it was almost 20 times higher among those who had ever had a routine visit, for anemia the ratio was 2.7 times higher, and for high blood pressure 2.3 times higher.

Impaired Fecundity

The term "impaired fecundity" in this section refers to a couple's impaired ability to conceive or maintain pregnancy either because of a known medical condition or because of an absence of conception after at least 2 years of exposure to unprotected intercourse. The aim of the module in the questionnaire was to assess the levels of infertility in the country and document existing reproductive health services for women with impaired fecundity.

In Eastern Europe, infertility is often cited as a reproductive health problem, given dramatic declines in fertility, widespread use of abortion, increase in sexually transmitted infections, pelvic inflammatory diseases and deficient health infrastructure. There are no reliable data from Albania regarding this issue, apart from media reports responding to public interest and utilization of specialized services.

The questionnaire included questions for married women about receiving any medical help or using any infertility services. Table 11.10 presents three indicators: proportion of those currently married women who report that they have had impaired fecundity, the proportion of those with current impaired fecundity, and the proportion of those who had ever been to a health professional for help to become pregnant. The proportion of women who have been seeking medical help to become pregnant is slightly higher than that of women who are classified as ever having had impaired fecundity; 6.5% versus 4.7%. The proportion of current impaired fecundity is much lower; only 1.4%. The results of the Albania RHS are lower than those in the survey carried out in Azerbaijan (ever had impaired fecundity -10%), where the module of fecundity impairment was applied for the first time in 2001 (Serbanescu et al., 2003).

In Tirana, but not in other urban areas, the level of impaired fecundity seems to be higher than in rural areas. This is especially noted in the case of the third indicator – proportion of women that have ever been to a doctor to seek help to become pregnant. There is no clear trend observed in the rate of impaired fecundity related to age.

Number of lifetime partners might increase the likelihood of women seeking help from a health professional to become pregnant; women with a total number of two or more partners in their lives reported a rate almost two times higher compared to women with only one lifetime partner. But the numbers of women with two or more lifetime partners are too small to show any statistical significance.

A routine gynecological exam is the strongest factor associated with reported impaired fecundity; it is positively associated with ever having experienced impaired fecundity or ever seeking medical help for the problem more than 6 times. The risk of currently having impaired fecundity is even more strongly associated; it is more than 9 times higher among those who have never had a routine gynecological visit.

The questionnaire included a question about the diagnoses associated with fertility impairment that a doctor told the woman during her clinical visits for fecundity impairment (Table 11.11). Among married women in the sample who have ever sought help for increasing their chances of pregnancy, 42% of them were diagnosed by a doctor with ovulation problems. Three other types of problems diagnosed by the doctor were endometriosis (14%), sperm problems (12%) and blocked tubes (10%). Some 28% of women remained without diagnoses for their fecundity impairment. The numbers are too small to find statistical differences but there seems to be some possible trends. Ovulation problems are more frequent among women of older ages and endometriosis is particularly higher among the 20-24 year age group, while blocked tubes were reported more

frequently by younger women of ages 15-19 years old. Women of lower education status are more likely to have endometriosis (and to some extent blocked tubes) as a cause of their impaired fecundity, than women of higher education status. In this latter category, sperm problems were more frequently reported than in the lower education status category. Among working women, endometriosis and blocked tubes were found in much lower proportions than among not working women. However, working women reported sperm problems twice as frequently as women who do not work, as a diagnosis of their impaired fecundity. A routine gynecological exam was associated only with the "blocked tubes" diagnosis, and this diagnosis was reported three times more by those women who have had at least one routine gynecological visit.

Smoking And Alcohol Consumption

Smoking

As mentioned previously, the Albania RHS is the first population-based study to produce data on prevalence of cigarette smoking and alcohol. The questionnaire included several questions for estimating cigarette smoking prevalence among women and men of reproductive age in Albania.

Tables 11.12A and B show the status of cigarette smoking for selected respondent characteristics. Although the proportion of women who have ever smoked any quantities of tobacco, is around 16%, only 4% of them report having smoked at least 100 cigarettes in their lives and slightly less (3%) reported smoking during the last 30 days. The prevalence is much lower than that reported by similar surveys in Eastern European countries such as Romania, Czech Republic, Russia, Ukraine (rates ranging from 19 to 30%), but is similar to

the rates of the Caucasus region (1%-6%) (CDC and ORC MACRO, 2004).

The demonstrates a trend of increasing prevalence of cigarette use from rural areas to urban areas, Tirana having the highest rate. Rates of cigarette smoking seem to be highest among women of higher socioeconomic status, those with higher education, and those who work. Previously married women seem to be exposed to a higher risk too.

The proportion of men who currently smoke is much higher (46%) and there are 58% who have ever tried to smoke. These figures are comparable to those of most of the Eastern European countries. All the social categories of men have higher rates of cigarette smoking but some differences are either too small or inconsistent. It indicates that even if those men with higher socioeconomic and higher status do not smoke more than others, they still smoke at least as much as the lower social and education categories.

Smoking increases with age. There are slightly less than 10% of teenagers who currently smoke and more than one in four among them have tried at least once to start smoking.

Alcohol drinking

There are four percent of women of reproductive age in the RHS sample who can be classified as frequent drinkers (they drink alcohol beverages every day or almost every day). More than one in three women has had some quantities of alcohol during the past three months, but the majority of them drink only occasionally; 25% drink alcohol once or twice a week and 29% drink 2-3 times a month. Table 11.13A demonstrates a consistent tendency; women of urban areas, women with higher levels of education, higher socioeconomic status and those who work seem to drink more frequently. This profile indicates higher access of these categories to alcohol beverages and probably more frequent "social drinking" among them. Age does not seem to play an important role in drinking habits for women.

Among men, reported alcohol drinking is much more frequent than among women. More than 60% of men have used alcohol during the past three months and 41% of them drink every day or almost every day (Table 11.13B). All categories have high rates of drinking. Men 15-24 yeas of age seem to drink slightly less frequently than older age groups, but, almost one in five men of age 15-24 years is a frequent drinker. Working men seem to drink more frequently than those who do not work, indicating the same social phenomena observed among women. Other trends, nevertheless, are not as consistent as among women.

Table 11.1 A Percentage Visiting Any Health Facility* in the Past 12 Months, by Selected Characteristics Women Aged 15-44 Years Reproductive Health Survey: Albania 2002

	Visited Health Facility - Past 12 Months							
Characteristics	Yes	No	Total	No. of Cases				
Total	26.4	73.6	100.0	5,697				
Strata								
Metro Tirana	32.3	67.7	100.0	2,108				
Other Urban	28.4	71.6	100.0	1,816				
Other Rural	23.6	76.4	100.0	1,773				
Residence								
Urban	30.0	70.0	100.0	3,572				
Rural	23.8	76.2	100.0	2,125				
Age Group								
15-19	17.5	82.5	100.0	1,094				
20-24	25.8	74.2	100.0	936				
25-29	33.3	66.7	100.0	946				
30-34	31.3	68.7	100.0	1,067				
35-39	28.9	71.1	100.0	958				
40-44	24.8	75.2	100.0	696				
Marital Status								
Married	31.5	68.5	100.0	3,965				
Previously Married	29.7	70.3	100.0	88				
Never Married	16.2	83.8	100.0	1,644				
Live Children								
0	19.7	80.3	100.0	1,943				
1	36.9	63.1	100.0	828				
2	28.7	71.3	100.0	1,840				
3 +	29.0	71.0	100.0	1,086				
Education Level								
Primary or Less	25.6	74.4	100.0	2,519				
Secondary Incomplete	22.5	77.5	100.0	653				
Secondary Complete	27.9	72.1	100.0	1,830				
Post-Secondary	31.6	68.4	100.0	695				
Socioeconomic Index								
Low	23.2	76.8	100.0	1,940				
Medium	27.7	72.3	100.0	2,985				
High	35.4	64.6	100.0	772				
Employment								
Working	29.9	70.1	100.0	1,118				
Not Working	25.8	74.2	100.0	4,579				

* Includes preventive services, family planning, counseling and health check ups

Table 11.1 B

Visited Health Facility - Past 12 Months No. of Characteristics Yes No Total Cases Total 13.6 86.4 100.0 1,740 Strata Metro Tirana 11.2 88.8 100.0 718 Other Urban 19.0 81.0 100.0 547 Other Rural 11.3 88.7 100.0 475 Residence Urban 16.4 83.6 100.0 1,155 Rural 11.2 88.8 100.0 585 Age Group 15-19 8.5 91.5 100.0 401 20-24 9.1 90.9 100.0 189 25-29 7.0 93.0 100.0 218 30-34 11.7 88.3 100.0 253 35-39 13.1 86.9 255 100.0 40-44 27.0 73.0 100.0 277 45-49 21.7 78.3 100.0 147 **Marital Status** Married 16.6 83.4 100.0 1,023 ** ** ** **Previously Married** 14 **Never Married** 90.9 100.0 703 9.1 Live Children 0 9.7 90.3 100.0 815 1 10.7 89.3 100.0 221 2 17.1 82.9 100.0 468 3 + 20.5 79.5 100.0 236 Education Level 13.1 689 Primary or Less 86.9 100.0 Secondary Incomplete 8.3 91.7 100.0 199 Secondary Complete 15.4 84.6 100.0 626 Post-Secondary 14.5 85.5 100.0 226 Socioeconomic Index Low 15.2 84.8 100.0 638 Medium 11.2 88.8 100.0 814 High 16.0 84.0 100.0 288 Employment Working 15.8 84.2 100.0 913 Not Working 11.5 100.0 827 88.5

Percentage Visiting Any Health Facility* in the Past 12 Months, by Selected Characteristics, among Men Aged 15-49 Years **Reproductive Health Survey: Albania 2002**

* Includes preventive services, family planning, counseling and health check ups.

** Percentages are not shown when base is less than 25 cases.

Table 11.2 A Percentage of Women Who Visited a Health Facility in the Past 12 Months

	0
	6
1	

That Received Co	unselin	g For Far	nily Pla	nning M	ethods	wonting
by Type of Fa W		d Selecte ged 15-44		acteristi	cs,	
Reproduct				nia 2002		
	Anv	Facility		. Clinic/ spital	Private Clinic/ Hospital	
		No. of		No. of		No. of
Characteristics	%	Cases	%	Cases	%	Cases
Total	16.0	1,639	15.7	1,347	21.2	375
Strata						
Metro Tirana	20.3	693	19.0	524	22.9	219
Other Urban	15.6	521	14.0	441	24.4	101
Other Rural	14.6	425	15.6	382	16.3	55
Residence						
Urban	17.4	1,112	15.7	884	24.3	290
Rural	14.8	527	15.6	463	16.0	85
Age Group						
15-19	9.7	204	8.8	170	14.1	50
20-24	21.4	267	20.9	207	23.6	75
25-29	18.2	334	17.8	265	28.6	81
30-34	16.6	356	16.8	300	16.5	69
35-39	16.6	284	16.6	239	19.0	64
40-44	11.6	194	11.0	166	22.4	36
Marital Status						
Married	18.7	1,325	18.6	1,097	24.2	290
Previously Married	8.5	26	**	20	**	7
Never Married	6.6	288	5.1	230	12.6	78
Live Children						
0	10.7	422	9.2	321	17.7	131
1	23.9	326	24.7	266	26.0	80
2	18.3	570	17.7	466	27.1	130
3 +	13.7	321	14.2	294	8.5	34
Education Level						
Primary or Less	14.3	682	14.3	596	17.0	107
Secondary Incomplete	12.7	156	13.1	128	7.9	38
Secondary Complete	18.1	555	17.6	459	28.6	119
Post-Secondary	22.5	246	20.5	164	26.5	111
Socioeconomic Index						
Low	10.5	487	11.2	435	11.5	64
Medium	18.2	867	18.0	721	22.3	192
High	24.5	285	21.4	191	29.5	119
Employment						
Working	21.1	358	20.4	271	27.5	112
Not Working	15.0	1,281	14.8	1,076	18.8	263

** Percentages are not shown when base is less than 25 cases

Table 11.2 B

Percentage of Men Aged 15-49 Years Who Visited a Health Facility in Past 12 Months That Received information/Counseling on Selected Topics, By Selected Characteristics Reproductive Health Survey: Albania 2002

Characteristics	Condoms	STD	Control for Pregnancy	No. of Cases
Total	8.0	8.9	82.6	240
Strata				
Metro Tirana	7.5	8.3	80.7	79
Other Urban	9.2	12.9	87.6	103
Other Rural	7.0	5.1	78.3	58
Residence				
Urban	9.0	11.7	85.4	171
Rural	6.7	5.3	78.9	69
Age Group				
15-24	18.2	12.1	81.4	51
25-39	11.0	16.9	90.4	86
40-49	1.1	1.9	77.7	103
Marital Status				
Married	5.8	7.8	81.8	173
Previously Married	**	**	**	1
Never Married	14.3	11.4	84.7	66
Live Children				
0	20.6	19.5	84.3	82
1-2	3.2	6.4	86.6	108
3 +	0.0	0.0	74.6	50
Education Level				
Primary or Less	4.7	3.8	81.9	92
Secondary	8.0	9.8	82.6	115
Post-Secondary	23.4	29.7	85.8	33
Socioeconomic Index				
Low	6.2	7.0	75.6	101
Medium	10.9	11.8	87.4	99
High	7.2	9.0	98.6	40
Employment				
Working	7.3	7.9	83.1	146
Not Working	8.9	10.2	81.9	94

 ** Percentages are not shown when base is less than 25 cases

Chapter 11

Tables

Table 11.3 A
Percentage Agreeing that Selected Factors May Be a Major Problem
Preventing Women From Getting Medical Advice or Treatment for Themselves,
by Selected Characteristics, among Women Aged 15-44 Years
Reproductive Health Survey: Albania 2002

Characteristics	Getting Money Needed for Treatment	Not Wanting to Go Alone	Distance to the Health Facility	Concern There May Not Be a Female Health Provider	Having to Take Transport	Knowing Where to Go	Getting Permission to Go	No. of Cases
Total	46.3	41.3	30.9	27.1	22.6	20.0	19.2	5,697
Strata								
Metro Tirana	34.8	29.2	13.8	16.1	8.5	15.0	13.2	2,108
Other Urban	36.9	29.4	9.4	18.3	5.0	12.2	10.3	1,816
Other Rural	54.7	51.3	47.6	35.1	36.2	25.6	25.8	1,773
Residence								
Urban	35.9	28.8	9.8	17.5	5.5	12.8	10.7	3,572
Rural	54.1	50.6	46.7	34.2	35.3	25.4	25.6	2,125
Age Group								
15-19	45.0	60.5	33.0	46.9	25.3	27.5	30.6	1,094
20-24	44.8	46.8	33.5	30.3	22.1	20.7	20.1	936
25-29	46.1	42.2	29.3	24.9	21.1	17.8	17.8	946
30-34	43.8	31.5	27.6	18.8	19.0	17.2	15.4	1,067
35-39	48.6	29.6	28.3	15.6	19.5	15.5	13.3	958
40-44	50.1	29.5	33.1	18.6	27.6	18.6	13.8	696
Marital Status								
Married	46.6	33.9	29.3	19.5	21.0	17.2	16.7	3,965
Previously Married	50.2	36.5	31.6	19.4	18.5	18.1	4.9	88
Never Married	45.4	56.2	34.1	42.7	25.8	25.6	25.1	1,644
Live Children								
0	44.3	54.1	32.8	40.8	24.5	24.6	24.4	1,943
1	42.6	38.7	23.9	20.0	15.9	15.3	16.9	828
2	42.7	29.8	25.1	16.2	16.6	15.8	13.3	1,840
3 +	56.1	35.0	38.8	21.0	30.3	19.9	18.8	1,086
Education Level								
Primary or Less	55.9	50.1	43.9	33.8	32.8	25.9	26.2	2,519
Secondary Incomplete	38.6	41.3	18.2	28.2	10.9	17.4	17.3	653
Secondary Complete	38.4	30.4	17.6	18.0	12.3	13.5	11.0	1,830
Post-Secondary	18.7	19.8	6.3	12.0	4.3	6.1	2.8	695
Socioeconomic Index								
Low	64.5	50.9	50.5	36.1	39.0	27.0	28.5	1,940
Medium	35.2	35.4	18.5	21.3	11.8	15.5	13.3	2,985
High	19.0	27.2	5.6	15.6	3.1	11.1	7.3	772
Employment								
Working	29.9	23.3	13.5	12.9	9.9	13.3	8.8	1,118
Not Working	49.2	44.5	34.1	29.6	24.8	21.2	21.1	4,579
Religion								
Muslim	47.3	41.1	32.5	26.5	24.0	19.3	20.3	4,568
Orthodox	34.1	32.2	17.0	19.6	10.5	14.7	8.4	531
Catholic	49.1	50.1	31.6	37.2	22.2	28.9	20.0	530
Other/Undeclared	22.7	22.6	9.9	17.2	4.7	10.2	8.0	68

Table 11.3 B

Percentage Agreeing that Selected Factors May Be a Major Problem Preventing Men From Getting Medical Advice or Treatment for Themselves, by Selected Characteristics, among Men Aged 15-49 Years Reproductive Health Survey: Albania 2002

	Getting Money				Concern There	
Characteristics	Needed for Treatment	Knowing Where to Go	Distance to the Health Facility	Having to Take Transport	May not Be a Male Health Provider	No. of Cases
Total	54.1	40.4	34.8	30.1	20.7	1,740
Strata						
Metro Tirana	35.6	27.7	12.9	11.2	16.4	718
Other Urban	45.5	28.4	11.0	8.2	11.9	547
Other Rural	66.4	52.5	57.6	50.6	27.5	475
Residence						
Urban	40.6	27.1	10.4	8.2	12.9	1,155
Rural	65.7	52.0	56.0	49.2	27.4	585
Age Group						
15-19	61.9	49.4	36.7	34.0	24.2	401
20-24	48.7	38.5	37.2	31.2	18.0	189
25-29	45.7	35.4	34.2	27.1	15.1	218
30-34	47.6	33.8	29.6	24.3	17.9	253
35-39	59.2	44.0	38.5	35.2	29.4	255
40-44	52.6	37.7	33.6	26.5	17.2	277
45-49	60.8	40.7	32.7	31.1	21.6	147
Marital Status						
Married	55.3	40.3	36.0	30.1	21.5	1,023
Previously Married	**	**	**	**	**	14
Never Married	52.3	40.3	33.0	30.2	19.1	703
Live Children						
0	50.5	39.3	32.6	28.8	18.7	815
1	47.4	35.6	30.1	24.5	18.8	221
2	54.0	41.0	29.9	24.6	21.4	468
3+	67.3	45.4	50.1	44.5	25.8	236
Education Level						
Primary or Less	63.8	49.7	48.8	41.9	28.3	689
Secondary Incomplete	52.5	42.6	24.1	21.4	15.4	199
Secondary Complete	47.1	32.8	24.8	21.3	15.2	626
Post-Secondary	29.5	17.3	8.2	9.0	5.1	226
Socioeconomic Index			0.2		•••	
Low	65.0	47.2	48.9	43.5	27.7	638
Medium	49.4	38.8	26.8	21.9	16.1	814
High	23.9	16.4	4.2	3.0	7.3	288
Employment						
Working	44.9	29.3	25.8	22.1	14.2	913
Not Working	62.9	29.3 51.1	43.5	37.9	26.8	827
-	02.0	01.1	-0.0	01.0	20.0	021
Religion	57 A	<i>ЛЛ</i> 4	20 4	22 0	20.7	1 400
Muslim	57.4	44.1 15.1	39.4	33.8	22.7	1,422
Orthodox	36.9	15.1	6.5 15 0	8.6	5.6	181
Catholic	31.6	28.0	15.9	11.7	9.1	76
Other/Undeclared ** Percentages are not show	<u>41.7</u>	27.4	15.8	16.3	24.6	61

Table 11.4

Time since Last Routine Gynecologic Examination by Selected Characteristics, among Women Aged 15-44 Years Who Have Ever Had Sexual Intercourse **Reproductive Health Survey: Albania 2002**

	Time of	Last Routine	Gynecologi	c Examinat	ion (Perce	ent Distrib	ution)
Characteristics	Within Past Year	Within 1-3 Years	> 3 Yrs. Ago	Never Had	Don't Know	Total	No. of Cases
Total	16.4	9.2	5.6	68.5	0.4	100.0	4,258
Strata							
Metro Tirana	22.5	10.5	4.7	61.3	1.0	100.0	1,618
Other Urban	17.7	11.1	7.0	64.0	0.3	100.0	1,379
Other Rural	13.7	7.6	5.0	73.4	0.3	100.0	1,261
Residence							
Urban	19.7	11.0	6.2	62.6	0.6	100.0	2,734
Rural	13.7	7.7	5.1	73.3	0.3	100.0	1,524
Age Group							
15-19	19.3	1.9	1.1	77.8	0.0	100.0	165
20-24	17.0	10.6	1.1	70.8	0.4	100.0	583
25-29	17.9	10.3	3.3	68.3	0.2	100.0	864
30-34	17.6	9.5	5.4	66.9	0.6	100.0	1,026
35-39	17.2	8.7	7.0	66.9	0.3	100.0	940
40-44	11.9	8.8	10.0	68.7	0.6	100.0	680
Marital Status							
Married	16.3	9.1	5.7	68.6	0.4	100.0	3,965
Previously Married	13.1	14.6	8.2	63.2	1.0	100.0	88
Never Married	20.8	6.1	1.1	71.1	0.9	100.0	205
Education Level							
Primary or Less	15.0	7.7	4.8	72.1	0.4	100.0	1,894
Secondary Incomplete	22.9	10.0	4.7	62.2	0.1	100.0	272
Secondary Complete	16.0	10.0	6.8	66.7	0.5	100.0	1,568
Post-Secondary	22.0	14.9	6.4	56.3	0.3	100.0	524
Employment							
Working	19.4	13.0	7.0	59.9	0.8	100.0	957
Not Working	15.7	8.3	5.2	70.4	0.3	100.0	3,301

Table 11.5

Most Important Reason That Women Have Never Had a Routine Gynecologic Exam, by Selected Characteristics, among Women Aged 15-44 Years Who Have Ever Had Sexual Intercourse Reproductive Health Survey: Albania 2002

	Most Important	(Percent	Percent Distribution)					
Characteristics	She is Healthy and Has Not Had Gynecologic Problems	Do Not Need To Get a Gynecologic Exam	Respondent Problems *	Provider Problems **	Cannot Afford the Cost	Don't Know	Total	No. of Cases
Total	48.4	41.9	6.0	2.0	1.4	0.3	100.0	2,791
Strata								
Metro Tirana	47.4	44.8	4.5	1.8	1.3	0.1	100.0	996
Other Urban	51.1	42.1	5.0	1.0	0.7	0.1	100.0	878
Other Rural	47.3	41.1	6.9	2.6	1.7	0.5	100.0	917
Residence								
Urban	49.3	43.6	4.9	1.2	0.8	0.1	100.0	1,691
Rural	47.8	40.8	6.7	2.6	1.8	0.4	100.0	1,100
Age Group								
15-19	36.3	57.5	0.5	4.3	1.4	0.0	100.0	124
20-24	49.2	42.1	4.7	3.8	0.0	0.1	100.0	400
25-29	48.5	44.9	4.1	0.9	1.2	0.4	100.0	558
30-34	47.5	42.3	6.7	1.4	1.8	0.2	100.0	660
35-39	48.6	41.1	6.2	2.5	1.2	0.5	100.0	617
40-44	51.2	36.2	8.8	1.5	2.1	0.3	100.0	432
Marital Status								
Married	48.8	41.4	6.1	1.9	1.4	0.3	100.0	2,598
Previously Married	40.5	42.9	10.0	5.8	0.7	0.0	100.0	57
Never Married	43.9	53.7	0.2	2.2	0.0	0.0	100.0	136
Education Level								
Primary or Less	46.7	40.8	7.1	2.8	2.1	0.5	100.0	1,331
Secondary Incomplete	48.0	46.6	4.5	0.5	0.5	0.0	100.0	170
Secondary Complete	50.8	43.0	4.7	0.9	0.5	0.0	100.0	1,008
Post-Secondary	52.3	42.6	3.3	1.5	0.2	0.0	100.0	282
Employment								
Working	49.7	42.2	4.7	2.6	0.8	0.0	100.0	567
Not Working	48.2	41.9	6.2	1.9	1.5	0.4	100.0	2,224

* Includes no time to go for exam, she forgets about it, does not like exam, and she never thought about it.

** Difficult to get appointment, does not like facility, is embarrassed to have a gynecologic exam, does not know where to go, and doctor never recommended

Tables

Table 11.6 Percentage with Awareness of Breast Self-Examination (BSE) and Frequency of BSE, by Selected Characteristics, among Women Aged 15-44 Years Who Have Ever Had Sexual Intercourse

Reproductive Health Survey: Albania 2002

	Awareness Frequency of BSE (Percent Distribution)						
Characteristics	Total	Every Month	Every 2-5 Months	1-2 Times Per Year or Less	Never	Total	No. of Cases
Total	51.9	3.7	1.4	3.3	91.6	100.0	4,258
Strata							
Metro Tirana	61.8	5.7	2.2	5.6	86.5	100.0	1,618
Other Urban	63.1	5.1	1.6	4.2	89.1	100.0	1,379
Other Rural	42.2	2.2	1.1	2.0	94.7	100.0	1,261
Residence							
Urban	63.4	5.5	1.9	4.9	87.7	100.0	2,734
Rural	42.5	2.2	1.0	2.0	94.8	100.0	1,524
Age Group							
15-19	31.6	2.5	0.3	1.8	95.4	100.0	165
20-24	43.8	2.6	0.6	1.8	95.0	100.0	583
25-29	51.6	5.9	1.2	4.1	88.7	100.0	864
30-34	56.8	3.7	1.6	4.5	90.3	100.0	1,026
35-39	56.8	3.4	1.5	3.2	91.9	100.0	940
40-44	51.5	2.9	2.0	2.9	92.2	100.0	680
Marital Status							
Married	51.7	3.5	1.4	3.2	91.9	100.0	3,965
Previously Married	57.7	2.0	0.7	5.7	91.6	100.0	88
Never Married	53.1	8.8	1.8	4.7	84.7	100.0	205
Education Level							
Primary or Less	39.7	1.5	0.9	1.8	95.8	100.0	1,894
Secondary Incomplete	57.8	3.9	1.1	0.7	94.2	100.0	272
Secondary Complete	64.8	5.0	1.7	4.8	88.5	100.0	1,568
Post-Secondary	78.7	13.0	3.6	9.6	73.8	100.0	524
Employment							
Working	71.3	7.6	2.6	6.7	83.1	100.0	957
Not Working	47.6	2.8	1.1	2.6	93.5	100.0	3,301
Routine Gynecologic Exam	ı						
Ever	57.6	5.1	2.6	5.2	87.1	100.0	1,467
Never	49.3	3.0	0.9	2.4	93.6	100.0	2,791

	Free		ervical Cance cent Distribu		ing Test	
Characteristics	Within Past Year	1-3 Years Ago	> 3 Years Ago	Never Had	Total	No. of Cases
Total	1.5	1.2	0.5	96.8	100.0	4,258
Strata						
Metro Tirana	3.3	1.8	1.0	93.9	100.0	1,618
Other Urban	2.1	1.6	0.5	95.9	100.0	1,379
Other Rural	0.6	0.8	0.3	98.2	100.0	1,261
Residence						
Urban	2.6	1.7	0.7	95.1	100.0	2,734
Rural	0.7	0.8	0.3	98.2	100.0	1,524
Age Group						
15-19	0.3	0.0	0.0	99.7	100.0	165
20-24	1.5	1.5	0.2	96.8	100.0	583
25-29	1.5	0.6	0.3	97.6	100.0	864
30-34	1.8	1.0	0.6	96.6	100.0	1,026
35-39	1.3	2.1	0.8	95.8	100.0	940
40-44	1.7	1.2	0.4	96.7	100.0	680
Marital Status						
Married	1.5	1.2	0.4	96.9	100.0	3,965
Previously Married	3.7	1.8	2.6	91.9	100.0	88
Never Married	1.0	0.8	0.9	97.2	100.0	205
Education Level						
Primary or Less	0.9	0.7	0.2	98.1	100.0	1,894
Secondary Incomplete	1.3	0.3	0.6	97.8	100.0	272
Secondary Complete	1.9	1.6	0.6	95.9	100.0	1,568
Post-Secondary	3.8	3.7	1.3	91.2	100.0	524
Employment						
Working	2.8	2.0	1.2	94.1	100.0	957
Not Working	1.2	1.1	0.3	97.4	100.0	3,301
Routine Gynecologic Exam						
Ever	3.7	2.9	1.3	92.1	100.0	1,467
Never	0.5	0.5	0.1	98.9	100.0	2,791

Table 11.7 Frequency of Cervical Cancer Screening by Selected Characteristics, among Women Aged 15-44 Years Who Have Ever Had Sexual Intercourse Reproductive Health Survey: Albania 2002

ables

Main Reason that Women Have Never Had a Pap Smear, by Selected Characteristics, among Women Aged 15-44 Years Who Have Ever Had Sexual Intercourse **Reproductive Health Survey: Albania 2002**

Chapter 11

		Reason Never Had a Pap Smear (Percent Distribution)									
Characteristics	Never Heard of It	She is Healthy and Has Not Had Gynecologic Problems	Doctor Never Recom- mended It	She Does Not Feel Test is Necessary		Not Sexually Active	Don't Know	Total	No. of Cases		
Total	69.8	12.8	9.4	4.3	2.9	0.6	0.3	100.0	4,074		
Strata											
Metro Tirana	47.9	24.5	15.2	7.6	3.4	0.9	0.5	100.0	1,517		
Other Urban	62.4	14.7	12.5	6.0	4.1	0.1	0.1	100.0	1,318		
Other Rural	80.6	8.2	5.9	2.3	2.1	0.7	0.3	100.0	1,239		
Residence											
Urban	56.8	6.8	13.3	6.8	4.0	0.4	0.2	100.0	2,580		
Rural	80.0	2.2	6.3	2.2	2.0	0.7	0.3	100.0	1,494		
Age Group											
15-19	88.5	3.8	3.4	3.7	0.6	0.0	0.0	100.0	163		
20-24	71.8	13.0	8.0	4.4	2.3	0.2	0.3	100.0	562		
25-29	71.3	11.7	8.5	3.5	4.5	0.5	0.0	100.0	835		
30-34	68.5	13.9	10.5	3.3	3.1	0.5	0.1	100.0	981		
35-39	65.8	12.3	11.3	5.6	3.5	0.8	0.7	100.0	889		
40-44	68.5	15.1	9.4	4.5	1.4	0.7	0.4	100.0	644		
Marital Status											
Married	69.9	12.5	9.6	4.2	3.0	0.5	0.3	100.0	3,794		
Previously Married	70.2	13.4	8.0	5.1	2.0	0.6	0.6	100.0	83		
Never Married	67.1	19.2	5.1	5.2	2.3	1.2	0.0	100.0	197		
Education Level											
Primary or Less	79.1	8.9	6.4	2.6	2.1	0.6	0.3	100.0	1,849		
Secondary Incomplete	68.5	15.0	7.2	6.1	2.4	0.7	0.0	100.0	264		
Secondary Complete	62.7	15.2	12.1	5.4	3.8	0.5	0.2	100.0	1,485		
Post-Secondary	33.4	29.2	21.4	10.0	5.3	0.3	0.5	100.0	476		
Employment											
Working	52.1	18.9	16.7	7.6	3.7	0.7	0.3	100.0	882		
Not Working	73.6	11.5	7.8	3.5	2.7	0.5	0.3	100.0	3,192		
Routine Gynecologic E	xam										
Ever	60.2	10.2	16.2	7.1	4.4	1.2	0.7	100.0	1,321		
Never	73.9	14.0	6.5	3.1	2.3	0.3	0.1	100.0	2,753		



Table 11.9 A Percentage of Women Aged 15-44 Years Who Have Been Told by a Doctor That They Have Selected Health Problems, by Selected Characteristics **Reproductive Health Survey: Albania 2002**

		PID **		High Blood		No. of	
Characteristics	% No. of Cases*		Anemia	Pressure	Diabetes	Cases	
Total	10.8	4,258	11.8	8.4	0.6	5,697	
Strata							
Metro Tirana	14.2	1,618	15.7	8.6	1.1	2,108	
Other Urban	9.6	1,379	12.7	6.9	0.6	1,816	
Other Rural	10.4	1,261	10.2	9.2	0.4	1,773	
Residence							
Urban	11.2	2,734	13.9	7.6	0.8	3,572	
Rural	10.5	1,524	10.3	9.1	0.4	2,125	
Age Group							
15-19	4.5	165	3.9	2.1		1,094	
20-24	8.3	583	10.2	3.8	0.1	936	
25-29	10.4	864	15.4	8.4	0.8	946	
30-34	10.9	1,026	14.7	10.7	0.4	1,067	
35-39	14.3	940	16.7	12.3	1.3	958	
40-44	10.2	680	13.0	16.3	1.1	696	
Marital Status							
Married	10.8	3,965	15.7	11.5		3,965	
Previously Married	19.0	88	22.1	15.9	3.0	88	
Never Married	2.9	205	3.4	1.9	0.2	1,644	
Education Level							
Primary or Less	10.8	1,894	10.1	9.0		2,519	
Secondary Incomplete	13.1	272	10.1	5.4	0.5	653	
Secondary Complete	11.0	1,568	14.0	9.2	0.9	1,830	
Post-Secondary	8.0	524	17.8	5.7	1.1	695	
Employment							
Working	11.9	957	15.5	8.9		1,118	
Not Working	10.5	3,301	11.2	8.4	0.5	4,579	
Routine Gynecologic Exam							
Ever	30.8	1,467	22.8	15.0	1.4	1,513	
Never	1.6	2,791	8.5	6.5	0.3	4,184	

* Restricted to women aged 15-44 years who have ever had sexual intercourse

** 19% hospitalized with 84% spending 6 or more nights in hospital

Table 11.9 B
Percentage of Men Aged 15-49 Years Who Have Been Told by a Doctor
That They Have Selected Health Problems, by Selected Characteristics
Reproductive Health Survey: Albania 2002

Characteristics	Hepatitis B	High Blood Pressure	Heart Disease	Diabetes	No. of Cases
Total	6.1	3.7	2.0	1.3	1,740
Strata					, -
Metro Tirana	5.1	2.3	1.5	1.3	718
Other Urban	7.5	5.0	2.8	1.8	547
Other Rural	5.7	3.5	1.7	1.0	475
Residence					
Urban	6.6	4.0	2.5	1.5	1,155
Rural	5.7	3.4	1.6	1.2	585
Age Group					
15-19	6.1	0.0	0.4	0.3	401
20-24	3.8	0.0	0.2	0.0	189
25-29	7.1	0.4	0.2	0.0	218
30-34	8.1	3.2	0.0	1.9	253
35-39	8.5	2.1	2.8	1.6	255
40-44	5.6	9.0	3.9	2.9	277
45-49	3.1	14.5	8.1	3.2	147
Marital Status					
Married	5.9	6.0	3.1	2.0	1,023
Previously Married	**	**	**	**	14
Never Married	6.4	0.3	0.3	0.2	703
Education Level					
Primary or Less	7.0	4.3	2.8	1.2	689
Secondary Incomplete	5.3	1.0	0.3	1.1	199
Secondary Complete	5.2	3.8	1.4	1.6	626
Post-Secondary	5.5	2.9	1.2	1.0	226
Employment					
Working	7.0	5.1	2.2	2.2	913
Not Working	5.2	2.4	1.8	0.5	827
Number of Partners					
0	5.0	0.0	0.4	0.2	446
1	3.5	5.9	3.5	1.5	448
2+	6.6	3.7	1.4	1.5	702
Don't Know	11.6	2.1	0.0	1.4	42
No Response	19.1	9.0	5.6	3.8	102

** Percentages are not shown when base is less than 25 cases

Tables

Table 11.10Percentage of Currently Married Women Aged 15-44 Years Who Reported Fecundity Impairment,by Selected CharacteristicsReproductive Health Survey: Albania 2002

Chapter 11

Characteristics	Ever Had Impaired Fecundity	Current Impaired Fecundity	Ever Been to Doctor for Help to Become Pregnant	No. of Cases
Total	4.7	1.4	6.5	3,965
Strata				
Metro Tirana	6.0	2.3	9.3	1,438
Other Urban	3.6	1.7	5.3	1,308
Other Rural	4.9	1.0	6.3	1,219
Residence				
Urban	4.3	1.9	6.6	2,488
Rural	4.9	1.0	6.3	1,477
Age Group				
15-19	4.4	1.7	5.2	97
20-24	5.3	2.3	7.5	502
25-29	6.6	1.6	8.5	800
30-34	3.6	1.5	4.6	1,004
35-39	4.4	1.5	6.7	906
40-44	3.9	0.4	5.8	656
Education Level				
Primary or Less	5.6	1.4	7.3	1,821
Secondary Incomplete	3.1	1.2	6.2	237
Secondary Complete	3.5	1.4	5.1	1,487
Post-Secondary	4.2	1.5	6.7	420
Employment				
Working	4.3	1.4	6.9	859
Not Working	4.7	1.4	6.4	3,106
Number of Partners *				
1	4.6	1.4	6.4	3,904
2+	6.1	1.8	11.0	60
Routine Gynecologic Exam				
Ever	11.1	3.7	15.0	1,367
Never	1.7	0.4	2.5	2,598

* Excludes one case that did not respond

Table 11.11 Percentage of Currently Married Women Aged 15-44 Years Who Reported Fecundity Impairment by Type of Problem and Selected Characteristics Reproductive Health Survey: Albania 2002

		r Had y Service	Type of Problem							
Chavastavistica	0/	No. of	Ovulation Drableme	Endo-	Sperm Broklama	Blocked	No	No. of		
Characteristics	%	Cases	Problems		Problems	Tubes	Diagnosis	Cases		
Total	6.5	3,965	42.0	13.5	12.3	10.0	27.9	271		
Strata										
Metro Tirana	9.3	1,438	32.3	13.0	15.5	9.1	36.1	131		
Other Urban	5.3	1,308	51.1	5.1	11.6	7.6	32.2	72		
Other Rural	6.3	1,219	41.7	17.8	11.3	11.5	22.4	68		
Residence										
Urban	6.6	2,488	42.5	7.2	14.0	8.6	35.0	183		
Rural	6.3	1,477	41.5	18.6	10.9	11.1	22.2	88		
Age Group										
15-19	5.2	97	**	**	**	**	**	5		
20-24	7.5	502	30.3	8.6	9.6	23.0	28.5	38		
25-29	8.5	800	32.9	29.2	16.6	2.5	21.7	57		
30-34	4.6	1,004	47.6	15.5	15.0	6.7	23.2	62		
35-39	6.7	906	50.2	7.4	8.8	12.7	34.1	62		
40-44	5.8	656	43.9	4.3	11.4	10.3	33.5	47		
Education Level										
Secondary Incomplete or Less	7.2	2,058	42.0	16.2	10.2	10.6	25.5	145		
Secondary Complete or More	5.4	1,907	41.9	7.9	16.7	8.7	32.9	126		
Employment										
Working	6.9	859	35.9	4.0	20.9	4.8	38.5	68		
Not Working	6.4	3,106	43.4	15.6	10.3	11.2	25.5	203		
Routine Gynecologic Exam										
Ever	15.0	1,367	41.2	13.3	12.8	12.1	26.2	203		
Never	2.5	2,598	44.1	14.0	11.0	4.2	32.6	68		

** Percentages are not shown when base is less than 25 cases

Table 11.12 A Percentage of Women Aged 15-44 Who Have Ever Smoked and Who Currently Smoke by Selected Characteristics Reproductive Health Survey: Albania 2002											
		Cigarette L	lse								
Characteristics	Ever Smoked Cigarettes	Ever Smoked 100 + Cigarettes	Currently Smoke	No. of Cases							
Total	16.1	3.6	3.0	5,697							
Strata											
Metro Tirana	28.4	9.2	7.9	2,108							
Other Urban	20.3	5.0	4.1	1,816							
Other Rural	10.3	1.3	0.9	1,773							
Residence											
Urban	23.8	6.8	5.8	3,572							
Rural	10.4	1.3	0.9	2,125							
Age Group				,							
15-19	18.7	1.9	1.7	1,094							
20-24	22.0	4.0	3.1	936							
25-29	16.9	3.7	2.9	946							
30-34	13.2	4.0	3.2	1,067							
35-39	12.8	4.4	3.8	958							
40-44	11.4	4.5	3.8	696							
Marital Status											
Married	13.1	3.4	2.6	3,965							
Previously Married	28.6	16.3	13.9	88							
Never Married	21.3	3.4	3.0	1,644							
Education Level											
Primary or Less	11.6	1.9	1.6	2,519							
Secondary Incomplete	24.1	4.7	4.0	653							
Secondary Complete	16.6	4.7	3.8	1,830							
Post-Secondary	34.1	10.3	8.2	695							
Socioeconomic Index											
Low	10.6	1.7	1.4	1,940							
Medium	17.9	4.1	3.4	2,985							
High	33.2	10.8	8.7	772							
Employment											
Working	22.5	8.5	7.4	1,118							
e e				*							

Chapter 11

Characteristics	Ever Smoked Cigarettes	Ever Smoked 100 + Cigarettes	Currently Smoke	No. of Cases
Total	57.6	49.2	46.3	1,740
Strata				·
Metro Tirana	65.6	55.5	50.4	718
Other Urban	56.8	47.9	45.5	547
Other Rural	54.9	47.4	45.2	475
Residence				
Urban	60.0	50.7	47.2	1,155
Rural	55.5	47.8	45.5	585
Age Group				
15-19	25.1	9.8	9.6	401
20-24	56.3	43.2	42.4	189
25-29	63.5	55.4	53.1	218
30-34	65.0	58.9	55.4	253
35-39	65.8	60.0	57.2	255
40-44	67.0	62.5	56.7	277
45-49	75.1	72.9	66.6	147
Marital Status				
Married	66.6	61.2	57.1	1,023
Previously Married	**	**	**	14
Never Married	43.2	30.2	29.2	703
Education Level				
Primary or Less	58.2	51.6	49.9	689
Secondary Incomplete	26.6	12.6	12.6	199
Secondary Complete	63.7	55.2	50.9	626
Post-Secondary	61.4	48.5	41.9	226
Socioeconomic Index				
Low	55.3	48.0	46.4	638
Medium	58.6	49.7	46.1	814
High	63.9	52.5	46.8	288
Employment				
Working	64.3	57.6	53.3	913
Not Working	51.1	41.1	39.6	827

Table 11.12 BPercentage of Men Aged 15-49 Who Have Ever Smokedand Who Currently Smoke, by Selected CharacteristicsReproductive Health Survey: Albania 2002

** Percentages are not shown when base is less than 25 cases

Table 11.13 A Percentage of Women Aged 15-44 Who Report Having a Drink Containing Alcohol in past 3 Months, and Frequency of Drinking, by Selected Characteristics **Reproductive Health Survey: Albania 2002**

	Frequency of Drinking Alcohol (Percent Distribution)											
Characteristics	Drink in Past 3 Months	No. of Cases	Every Day	Almost Every Day	1-2 Times a Week	2-3 Times a Month	Once a Month	1-2 Times in Three Months	Only a Few Sips	Don't Know	Total	No. of Cases
Total	35.5	5,697	1.5	3.3	25.3	29.3	16.6	6.8	16.6	0.7	100.0	2,612
Strata												
Metro Tirana	44.4	2,108	2.8	5.1	30.4	22.2	15.7	5.5	15.9	2.4	100.0	1,101
Other Urban	40.5	1,816	1.0	3.8	28.4	30.9	13.2	5.4	16.8	0.5	100.0	872
Other Rural	30.2	1,773	1.4	2.2	20.8	31.2	19.4	8.3	16.7	0.0	100.0	639
Residence												
Urban	42.7	3,572	1.7	4.4	29.5	27.3	14.2	5.4	16.4	1.2	100.0	1,849
Rural	30.1	2,125	1.4	2.2	20.9	31.4	19.1	8.2	16.8	0.1	100.0	763
Age Group												
15-24	35.4	2,030	0.8	2.2	23.3	29.9	16.1	8.2	18.7	0.9	100.0	961
25-34	34.6	2,013	1.4	4.6	25.5	31.5	17.4	4.6	14.9	0.2	100.0	871
35-44	36.4	1,654	2.7	3.5	27.6	26.5	16.5	7.0	15.5	0.8	100.0	780
Marital Status												
Married	34.8	3,965	1.9	4.0	27.8	28.7	16.4	5.6	14.9	0.7	100.0	1,751
Previously Married	34.9	88	2.5	1.3	20.7	23.3	18.1	14.4	17.5	2.2	100.0	37
Never Married	36.9	1,644	0.8	2.1	21.1	30.8	16.9	8.4	19.5	0.4	100.0	824
Education Level												
Primary or Less	29.9	2,519	0.9	2.6	20.7	32.6	19.3	7.5	15.9	0.5	100.0	905
Secondary Incomplete	40.9	653	2.7	1.8	25.0	28.9	15.4	8.1	17.6	0.6	100.0	330
Secondary Complete	40.0	1,830	1.8	3.8	27.6	27.9	14.8	6.2	17.1	0.8	100.0	939
Post-Secondary	50.3	695	1.8	6.0	37.3	20.8	12.0	4.2	17.1	0.8	100.0	438
Socioeconomic Index												
Low	26.7	1,940	1.2	1.5	19.5	32.8	21.0	7.9	15.8	0.4	100.0	646
Medium	40.5	2,985	1.5	3.2	27.1	28.4	15.8	6.9	16.4	0.6	100.0	1,462
High	49.8	772	2.6	8.2	32.1	24.3	8.5	3.2	19.4	1.7	100.0	504
Employment												
Working	45.0	1,118	2.8	5.4	33.2	21.6	12.7	5.2	18.6	0.5	100.0	631
Not Working	33.7	4,579	1.2	2.8	23.3	31.2	17.6	7.2	16.1	0.7	100.0	1,981

Table 11.13 B

Percentage of Men Aged 15-49 Who Report Having a Drink Containing Alcohol in Past 3 Months, and Frequency of Drinking, by Selected Characteristics Reproductive Health Survey: Albania 2002

	ast				F			nking Alco tribution)	ohol			
<u>Characteristics</u>	Drinking in Past 3 Months	No. of Cases	Every Day	Almost Every Day	1-2 Times a Week	2-3 Times a Month	Once a Month	1-2 Times in Three Months	Only Few Sips	Don't Know	Total	No. of Cases
Total	60.8	1,740	13.6	27.5	37.3	12.6	2.8	0.5	5.0	0.6	100.0	1,123
Strata												
Metro Tirana	69.3	718	16.9	27.3	35.5	11.6	2.4	0.9	4.8	0.7	100.0	507
Other Urban	60.0	547	10.0	29.8	38.0	13.2	4.2	0.9	3.3	0.6	100.0	330
Other Rural	57.8	475	14.3	26.2	37.8	12.7	2.1	0.2	6.2	0.6	100.0	286
Residence												
Urban	63.8	1,155	13.1	28.5	37.2	12.2	3.7	0.7	4.0	0.6	100.0	768
Rural	58.1	585	14.2	26.5	37.4	13.0	1.9	0.4	6.0	0.6	100.0	355
Age Group												
15-24	43.2	590	4.8	14.0	43.0	23.5	4.6	1.3	7.5	1.3	100.0	266
25-34	73.2	471	12.3	31.0	40.7	9.7	3.2	0.5	2.5	0.1	100.0	356
35-49	67.1	679	19.5	32.4	31.8	8.7	1.4	0.2	5.4	0.6	100.0	501
Marital Status												
Married	68.5	1,023	17.6	31.8	34.7	9.1	1.6	0.3	4.5	0.3	100.0	756
Previously Married	**	14	**	**	**	**	**	**	**	**	**	9
Never Married	48.6	703	4.6	18.2	43.5	20.2	4.8	1.1	6.3	1.3	100.0	358
Education Level												
Primary or Less	58.6	689	13.0	28.3	38.2	11.4	2.3	0.5	5.3	1.0	100.0	419
Secondary Incomplete	35.8	199	8.6	9.2	30.8	34.1	3.1	3.2	11.0	0.0	100.0	78
Secondary Complete	69.0	626	16.0	29.4	35.0	11.0	3.4	0.3	4.5	0.4	100.0	467
Post-Secondary	65.5	226	10.3	26.0	46.1	12.8	2.4	0.0	2.5	0.0	100.0	159
Socioeconomic Index												
Low	57.1	638	14.7	27.5	33.1	13.4	3.1	0.5	6.4	1.4	100.0	389
Medium	62.0	814	12.0	26.7	40.8	12.5	2.5	0.7	4.7	0.0	100.0	524
High	72.3	288	15.8	30.0	40.9	10.1	2.3	0.0	0.9	0.0	100.0	210
Employment												
Working	71.6	913	17.1	33.4	32.7	9.7	3.5	0.2	2.6	0.8	100.0	677
Not Working	50.3	827	9.2	19.9	43.2	16.4	1.8	1.0	8.2	0.3	100.0	446

 ** Percentages are not shown when base is less than 25 cases

CHAPTER 12

FAMILY LIFE AND SEX EDUCATION

Introduction

↑oncerns about teenage sexuality, ✓ pregnancy and sexual health have been increasing worldwide in recent years. Prevention programs designed to reduce the rate of adolescent pregnancy and sexually transmitted infections require a multifaceted approach, and school-based sex education is one important component of a broad effort. A number of studies have demonstrated that high-quality sex education programs can lead to higher levels of abstinence, later initiation of sexual activity, increased use of contraception and fewer sexual partners (Dawson, 1986; Kirby, 1999; Kirby et al., 1994). Health education interventions are widely seen as appropriate strategies for promoting young people's sexual health, particularly when information among young people about sexuality, reproduction, contraception, and sexually transmitted infections is lacking. Family life or sexual education has been part of the school curriculum in many countries, although teaching about contraceptive methods is sometimes omitted.

Health education, including family life and sex education, became part of the Albanian school curricula only in 1994 starting in the fifth class of elementary schools. Only nine hours in each of the eight classes are dedicated to sex education while another six hours are for HIV/AIDS and sexually transmitted diseases. Still there is a need to improve the family life education curricula and extend it further to high schools. Extracurricular sessions applied by teachers, NGOs or health promoting schools' network are important but are still spontaneous and do not fulfill the current standardized needs of such education.

Opinions about Sex Education in Schools

Table 12.1A shows that over 90% of Albanian women of reproductive age agree that sex education, including human reproduction, contraception and sexually transmitted infections, should be taught in school. There is little variation by various socio-demographic characteristics, with the percentage in agreement dropping only to 88% for women with 3 or more living children. Nevertheless, there are some patterns worth noting. The largest differentials are observed for educational levels, with a positive association between educational level and agreement that sex education should be taught in school (from 89% to 99% agreement). Socioeconomic status also shows a positive relationship, although not as strong as that for education. Urban women, younger women, and never married women are somewhat more likely to favor sex education in school than their rural, older and married counterparts.

Among men 15-49 years of age (Table 12.1B), there is also a high percentage in favor of sex education in school, although not as high as that of women (84% vs. 92%). Similar relationships with sociodemographic variables are observed for men, but with stronger associations. Thus, while 90% of urban men favor sex education, only 79% of rural men are in agreement. Agreement by age ranges from 91% among 15-24 year olds to 76% for 35-49 year olds. Ninety-one percent of never married men agree to sex education in school compared to 80% of married men, and the positive associations of educational level and socioeconomic status with favoring sex education are even stronger than observed for women.

When asked the best age to begin courses in school on human reproduction and contraception, the majority of women who favored the teaching of sex education in school reported ages 14-15 (Table 12.2A and Figure 12.1). This was the preferred age group among women in all categories of the socio-demographic control variables for both human reproduction and contraception. The second preferred age groups for receiving these topics in school varied depending on the topic. For the topic of human reproduction, most women preferred the younger ages (13 years or younger) over the older ages (16 years or older), with the exception of women living in rural areas, having 3 or more children, primary or lower educational level and low socioeconomic status. On the other hand, with respect to the teaching of contraception, the women who did not choose ages 14–15 were more likely to prefer older ages (16 years or older). The exceptions to this pattern were urban women, more highly educated women and those of high socioeconomic status.


Compared to women, men who favored the teaching of sex education in school appeared to be somewhat more conservative than women in terms of the ages at which they felt young persons should receive courses on human reproduction and contraception. Overall, men reported a preference for ages 16 and over as the best age to receive information on human reproduction and contraception in school (44% and 47%, respectively) (Table 12.2B and Figure 12.2). Those men who expressed a preference for ages 15 or younger were those living in urban areas (57% and 55%), the youngest (64% and 58% of 15-24 year olds), never married (66% and 60%), those with no living children (64% and 59%), having post-secondary education (65% and 62%), and having high socioeconomic status (72%) and 69%).


Tables 12.3A and B present the preferred ages for young people to receive lessons in school on sexually transmitted infections as reported by women and men of reproductive age who favor sex education in school. The preferred ages and the patterns for various socio-demographic categories are similar to those observed for contraception. The majority of women prefer this topic be taught to young people at 14-15 years of age (54%) (Table 12.3A). The second overall preferred age group is 16 years of age or older (25%), with the exceptions noted earlier among urban women (24%), women with postsecondary education (32%) and women with high socioeconomic status (29%). Overall, men (Table 12.3B) prefer the older ages of 16 years or more (46%), with the younger ages of 14–15 preferred only by 15–24 year old men (53%), the never married (53%), the childless (51%), those with post-secondary education (44%), and those with high socioeconomic status (50%).

Tables 12.4 through 12.6 present the actual experiences of young adults in Albania with regard to education on matters related to sexuality.

Discussions with Parents on Family Life Education Topics

In Table 12.4A are the percentages of women 15–24 years of age who had discussed family life education topics with a parent before the age of 18. Two-thirds of young women (64%) reported they had discussed any topic with a parent. However, only 21% reported discussing how pregnancy occurs, 15% reported discussing HIV/AIDS, and 8% reported discussing methods of contraception. These percentages varied markedly by certain socio-demographic characteristics. Urban women were more likely to have discussed a topic with a parent than rural women, and the probability of having discussed a topic increased with educational level and





socioeconomic status of the young woman. There were no pronounced aged differences in the likelihood of having discussed a topic. And, with the exception of how pregnancy occurs, there were no significant differences between sexually experienced and nonsexually experienced young women in whether or not they had discussed a family life education topic with a parent before age 18.

Young men report much less communication with parents on family life topics compared to women (Table 12.4B). Only 11% of 15-24 year old men said they had discussed a family life education topic with a parent before age 18. Less than 10% had discussed HIV/AIDS or how pregnancy occurs, and only 2% had discussed methods of contraception. As with young women, differences are observed socio-demographic characteristics, by with urban, highly educated and high socioeconomic status men more likely to have discussed family life education topics with a parent. Differences in this likelihood by age or by sexual experience are small.

Sex Education in School

Tables 12.5A and B demonstrate that Albanian young adults are more likely to receive education on sexual topics in school than in the home. Seventy-seven percent of young women had received some type of sex education in school before the age of 18 (Table 12.5A). The majority had received information on the female and male reproductive systems (67% and 63%, respectively), the menstrual cycle (66%), how pregnancy occurs (57%) and HIV/AIDS (53%). But only one-fourth had received information on contraceptive methods (24%) or sexually transmitted infections other than HIV/AIDS (24%). Differences according to socio-demographic categories are also observed. Residence in urban areas, younger age, higher educational level and higher socioeconomic status all provide greater access to sex education in school.

Young men report lower levels of sex education in school than the young women (Table 12.5B). Sixty-four percent report having received any topic of sex education in school before the age of 18. As with the young women, contraceptive methods and sexually transmitted infections other than HIV/AIDS were topics not often covered in their sex education lectures (30% and 24%, respectively). Men also report low coverage of the menstrual cycle, with only 23% of men receiving such information. As with young women, residence in urban areas, higher educational level and higher socioeconomic status seem to provide greater access to sex education in school. Although sexually experienced young men reported sex education in school in slightly higher proportions than non-sexually experienced young men, the differences are not statistically significant.

Table 12.6A shows the age by which selected topics have been received in school reported by young adult women who have had sex education in school. With the exception of contraception, most young women had received lessons on pregnancy-related topics before the age of 16. In fact, according to the magnitude of increase in the cumulative percentages, it appears that all indicated topics were most likely to be received in school at age 14. Table 12.6B shows a similar pattern for young adult males, that is, all topics were most likely to have been presented in school when the young men were 14 years of age.

Sources of Information on Sexual Matters

When young adults who had received sex education in school were asked what was the most important source for them with regard to information on sexual matters, approximately half of the women and men responded their teachers to be the most important source (49% and 53%, respectively) (Tables 12.7A and B). The second most frequently reported source was radio and TV, followed closely by books, newspapers and magazines, with 24% of women and 21% of men saying radio and TV and 10% of women and 19% of men reporting the print media were the most important source to them. When sociodemographic characteristics are controlled, the three most reported sources remain in the same rank order for almost all of the categories for both young adult women and young adult men.

Tables 12.8A and B present the most important sources for sexual information reported by young adults who did not have sex education in school. For these young people, radio and TV is reported most frequently - 44% for women and 53% for men. Co-workers and peers are the second most frequently reported important source, with 15% of young women and 20% of young men offering this response. Finally, the third and fourth most cited sources for women was their parents (11%) and other relatives (9%), although these sources were infrequent among men (2% and 0.2%, respectively). Again, the rank order of the most cited sources for both women and men does not vary greatly across sociodemographic categories.

The Impact of Sexuality Education on Knowledge about Pregnancy Issues

Questions were included in the ALRHS-02 to assess respondents' knowledge of pregnancy and contraception issues. Tables 12.9A and B show the answers given by young adult women and men to several of these questions.

Knowledge of the most fertile time in a woman's cycle is an important measure of a woman's ability to assess the risk of pregnancy during unprotected intercourse, and consequently to prevent unintended pregnancy. Only 15% of women 15–24 years of age correctly identified halfway between periods to be the most likely time for a woman to become pregnant (Table 12.9A). Sixtyseven percent simply said they did not know the answer. However, there does seem to be some impact of having discussed the topic with parents or in school. Young women who discussed the topic with a parent were more than twice as likely as those who did not to know the correct answer (21% vs. 8%), and women who received the topic in school were also more than twice as likely to know the correct answer compared to those who did not have the topic in school (19% vs. 7%). Similarly, only 12% of young women knew that breastfeeding can decrease the chance of pregnancy. This percentage increased for women who had discussed pregnancy with a parent (19%) compared to those who had not (9%). Learning about pregnancy in school, on the other hand, had little effect on knowledge of the correct answer to this question -13% compared to 10% for those who did not learn about pregnancy in school. Finally, when asked if it is possible to get pregnant at first intercourse, only 49% of young adult women agreed. Both having discussed pregnancy with a parent and having learned about pregnancy in school appear to have an effect on this knowledge. Sixty-three percent of young women who had discussed pregnancy with a parent knew that it is possible to get pregnant at first sexual intercourse, whereas only 45% of those who did not discuss pregnancy with a parent knew this answer. Having received a

lesson in school about pregnancy also had an effect on women's knowledge of this question. Fifty-three percent of women who learned about pregnancy in school knew the correct answer compared to 44% of those who were not taught about pregnancy in school.

Young men's knowledge on pregnancy matters is shown in Table 12.9B. Only 10% of young adult men knew that breastfeeding can reduce the likelihood of becoming pregnant. Having discussed pregnancy with a parent appears to have a pronounced effect on young men's correct answer to this question. Thirty percent of men who had discussed pregnancy with a parent knew that breastfeeding can decrease the chance of pregnancy compared to only 9% of those who did not discuss the topic of pregnancy with a parent. The effect of having been taught about pregnancy in school is minimal in relation to the effect of breastfeeding. About one-half (52%) of the 15–24 year old men knew a woman could become pregnant at first sexual intercourse. Again, we observe a large effect of having discussed pregnancy with a parent— 70% vs. 51% for those who had not discussed pregnancy with a parent. Having learned about pregnancy in school also appears to have been effective, but to a lesser extent.




Table 12.1 A

Percentage of Women Aged 15-44 Who Agree Certain Sex Education Topics Be Taught in School, By Selected Characteristics **Reproductive Health Survey: Albania 2002**

Characteristic	Any Topic	Human Reproduction	Contraception	Sexually Transmitted Infections	No. of Cases
Total	92.3	92.3	92.0	92.1	5,697
Strata					
Metro Tirana	94.0	93.9	93.6	93.8	2,108
Other Urban	94.3	94.3	94.2	94.3	1,816
Other Rural	90.8	90.8	90.3	90.4	1,773
Residence					
Urban	94.2	94.2	94.1	94.2	3,572
Rural	90.9	90.9	90.4	90.6	2,125
Age Group					
15-24	94.3	94.3	93.9	94.0	2,030
25-34	92.1	92.1	91.9	91.9	2,013
35-44	90.1	90.1	89.8	90.0	1,654
Marital Status					
Married	91.4	91.4	91.0	91.1	3,965
Previously Married	90.6	90.6	90.6	90.6	88
Never Married	94.4	94.4	94.1	94.2	1,644
Living Children					
0	94.3	94.3	94.0	94.2	1,943
1	91.8	91.8	91.4	91.2	828
2	93.1	93.1	93.0	93.0	1,840
3 +	88.3	88.3	87.7	88.0	1,086
Education Level					
Primary or Less	89.1	89.1	88.6	88.8	2,519
Secondary Incomplete	95.8	95.8	95.5	95.6	653
Secondary Complete	95.5	95.5	95.4	95.4	1,830
Post-Secondary	98.5	98.5	98.3	98.5	695
Socioeconomic Index					
Low	89.8	89.7	89.4	89.4	1,940
Medium	93.9	93.9	93.6	93.8	2,985
High	95.9	95.9	95.9	95.9	772

Table 12.1 B	
Percentage of Men Aged 15-49 Who Agree Certain Sex Education T	opics
Be Taught in School, By Selected Characteristics	
Reproductive Health Survey: Albania 2002	
Sexual	ly

Characteristics	Any Topic	Human Reproduction	Contraception	Transmitted Infections	No. of Cases
Total	84.3	84.3	84.3	84.3	1,740
Strata					ŗ
Metro Tirana	89.7	89.7	89.7	89.7	718
Other Urban	88.8	88.8	88.8	88.8	547
Other Rural	79.5	79.5	79.5	79.5	475
Residence					
Urban	89.8	89.8	89.8	89.8	1,155
Rural	79.3	79.3	79.3	79.3	585
Age Group					
15-24	90.9	90.9	90.9	90.9	590
25-34	87.6	87.6	87.6	87.6	471
35-49	76.4	76.4	76.4	76.4	679
Marital Status					
Married	79.8	79.8	79.8	79.8	1,023
Previously Married	**	**	**	**	14
Never Married	91.1	91.1	91.1	91.1	703
Living Children					
0	91.4	91.4	91.4	91.4	815
1	79.5	79.5	79.5	79.5	221
2	83.9	83.9	83.9	83.9	468
3 +	69.8	69.8	69.8	69.8	236
Education Level					
Primary or Less	72.5	72.5	72.5	72.5	689
Secondary Incomplete	95.4	95.4	95.4	95.4	199
Secondary Complete	94.6	94.6	94.6	94.6	626
Post-Secondary	97.6	97.6	97.6	97.6	226
Socioeconomic Index					
Low	76.7	76.7	76.7	76.7	638
Medium	89.6	89.6	89.6	89.6	814
High	96.4	96.4	96.4	96.4	288

** Percentages are not shown when base is less than 25 cases.

Table 12.2 A

Percent Distribution of Best Age to Start School-Based Courses on Human Reproduction and Contraception, as Reported by Women Aged 15-44 Who Are in Favor of Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

		Н	uman R	leproduc Don't	tion	No. of			Contra	aception Don't	1	No. of
Characteristic	≤ 13	14-15	16 +	Know	Total	Cases	≤13	14-15	16 +	Know	Total	Cases
Total	24.4	55.7	17.8	2.2	100.0	5,300	18.6	53.7	24.2	3.5	100.0	5,283
Strata												
Metro Tirana	34.2	51.7	12.1	2.0	100.0	1,979	24.9	52.2	19.3	3.5	100.0	1,973
Other Urban	28.3	56.1	14.2	1.4	100.0	1,718	22.6	56.7	18.7	2.0	100.0	1,716
Other Rural	19.3	56.6	21.5	2.6	100.0	1,603	14.5	52.5	28.7	4.3	100.0	1,594
Residence												
Urban	30.8	54.4	13.5	1.4	100.0	3,367	24.2	55.3	18.5	2.0	100.0	3,362
Rural	19.4	56.7	21.1	2.8	100.0	1,933	14.3	52.6	28.5	4.6	100.0	1,921
Age Group												
15-24	21.3	60.6	15.7	2.4	100.0	1,918	15.4	60.0	21.1	3.5	100.0	1,910
25-34	27.6	52.7	17.2	2.4	100.0	1,875	22.2	50.9	23.6	3.3	100.0	1,872
35-44	25.1	52.3	21.0	1.6	100.0	1,507	19.2	48.5	28.8	3.5	100.0	1,501
Marital Status												
Married	26.2	52.4	19.2	2.2	100.0	3,659	20.2	50.1	26.1	3.6	100.0	3,647
Previously Married	21.2	52.0	21.3	5.6	100.0	81	15.3	47.6	27.5	9.6	100.0	81
Never Married	21.2	62.1	14.8	1.9	100.0	1,560	15.7	61.1	20.3	2.9	100.0	1,555
Living Children												
0	21.6	61.6	14.6	2.2	100.0	1,840	15.8	60.7	20.2	3.2	100.0	1,834
1	28.2	51.7	18.9	1.2	100.0	770	23.4	51.0	23.1	2.5	100.0	768
2	28.9	51.2	17.1	2.8	100.0	1,725	21.5	50.3	24.3	3.9	100.0	1,724
3 +	21.4	53.1	23.7	1.8	100.0	965	17.2	47.1	31.8	3.9	100.0	957
Education Level												
Primary or Less	18.5	57.1	20.7	3.6	100.0	2,240	13.1	53.8	27.5	5.6	100.0	2,227
Secondary Incomplete	24.8	57.4	16.8	1.0	100.0	627	19.9	56.3	21.7	2.1	100.0	626
Secondary Complete	29.8	53.8	15.8	0.6	100.0	1,749	24.2	52.2	22.5	1.1	100.0	1,747
Post-Secondary	41.4	51.0	7.6	0.0	100.0	684	31.8	55.3	12.7	0.2	100.0	683
Socioeconomic Index												
Low	18.6	55.5	22.0	3.8	100.0	1,739	14.1	52.1	28.6	5.3	100.0	1,732
Medium	26.7	56.3	15.8	1.1	100.0	2,809	20.6	54.6	22.4	2.3	100.0	2,799
High	38.6	52.5	8.6	0.3	100.0	752	28.4	56.5	13.6	1.5	100.0	752

Table 12.2 B

Percent Distribution of Best Age to Start School-Based Courses on Human Reproduction and Contraception, as Reported by Men Aged 15-49 Who Are in Favor of Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

	Human Reproduction						Contraception					
Characteristic	≤ 13	14-15	16 +	Don't Know	Total	No. of Cases	≤ 13	14-15	16 +	Don't Know	Total	No. of Cases
Total	11.7	39.7	44.1	4.5	100.0	1,515	10.8	37.6	47.1	4.5	100.0	1,515
Strata												
Metro Tirana	19.2	41.2	37.1	2.5	100.0	645	17.5	40.0	40.0	2.5	100.0	645
Other Urban	10.9	43.9	41.3	3.9	100.0	490	10.3	42.2	43.6	3.9	100.0	490
Other Rural	8.9	36.3	49.1	5.7	100.0	380	8.2	33.4	52.7	5.7	100.0	380
Residence												
Urban	14.4	42.8	39.5	3.3	100.0	1,047	13.3	41.2	42.1	3.3	100.0	1,047
Rural	9.1	36.7	48.6	5.6	100.0	468	8.4	34.0	52.1	5.6	100.0	468
Age Group												
15-24	8.9	55.2	33.3	2.6	100.0	536	8.5	49.7	39.2	2.6	100.0	536
25-34	11.3	36.9	46.3	5.6	100.0	419	9.9	36.8	47.8	5.6	100.0	419
35-49	14.8	26.4	53.2	5.6	100.0	560	13.8	26.1	54.6	5.6	100.0	560
Marital Status												
Married	13.0	28.1	53.2	5.8	100.0	857	11.7	28.0	54.5	5.8	100.0	857
Previously Married	**	**	**	**	**	**	**	**	**	**	**	12
Never Married	9.9	55.7	32.0	2.4	100.0	646	9.5	50.7	37.4	2.4	100.0	646
Living Children												
0	11.3	52.8	33.1	2.8	100.0	750	10.3	48.7	38.2	2.8	100.0	750
1	5.9	44.3	43.5	6.2	100.0	186	5.2	42.6	46.0	6.2	100.0	186
2	12.2	27.0	55.9	4.9	100.0	404	11.4	27.1	56.6	4.9	100.0	404
3 +	16.4	14.6	61.0	8.0	100.0	175	15.3	14.9	61.8	8.0	100.0	175
Education Level												
Primary or Less	7.1	39.0	45.8	8.0	100.0	523	6.4	36.7	48.9	8.0	100.0	523
Secondary Incomplete	14.1	46.9	38.4	0.6	100.0	186	13.7	42.5	43.1	0.6	100.0	186
Secondary Complete	13.4	37.4	46.3	2.9	100.0	585	12.3	36.0	48.8	2.9	100.0	585
Post-Secondary	21.4	43.9	34.7	0.0	100.0	221	20.0	42.2	37.8	0.0	100.0	221
Socioeconomic Index												
Low	8.2	35.9	49.3	6.6	100.0	500	7.6	32.9	52.9	6.6	100.0	500
Medium	12.4	40.4	44.1	3.2	100.0	737	11.4	39.0	46.4	3.2	100.0	737
High	21.5	50.8	25.9	1.8	100.0	278	20.0	49.1	29.1	1.8	100.0	278

** Percentages are not shown when base is less than 25 cases.

Table 12.3 A

Percent Distribution of Best Age to Start School-Based Courses on Sexually Transmitted Infections, as Reported by Women 15-44 Who are in Favor of Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

	Sexually Transmitted Infections							
				Don't		No. of		
Characteristic	≤ 13	14-15	16 +	Know	Total	Cases		
Total	18.0	53.6	24.8	3.7	100.0	5,291		
Strata								
Metro Tirana	25.1	53.1	18.8	3.0	100.0	1,977		
Other Urban	22.4	56.3	19.3	2.0	100.0	1,718		
Other Rural	13.4	52.2	29.6	4.8	100.0	1,596		
Residence								
Urban	24.0	55.2	18.8	1.9	100.0	3,366		
Rural	13.3	52.3	29.4	5.0	100.0	1,925		
Age Group								
15-24	14.7	60.6	21.0	3.7	100.0	1,913		
25-34	21.0	51.0	24.6	3.4	100.0	1,873		
35-44	19.2	47.2	29.8	3.9	100.0	1,505		
Marital Status								
Married	19.7	49.5	27.0	3.8	100.0	3,652		
Previously Married	16.9	46.2	24.7	12.2	100.0	81		
Never Married	14.9	61.8	20.4	2.9	100.0	1,558		
Living Children								
0	14.9	61.6	20.3	3.2	100.0	1,837		
1	22.7	51.3	23.6	2.4	100.0	767		
2	21.4	49.7	25.0	3.9	100.0	1,725		
3 +	16.5	45.4	33.2	5.0	100.0	962		
Education Level								
Primary or Less	12.5	53.2	28.2	6.1	100.0	2,232		
Secondary Incomplete	19.9	56.5	22.1	1.6	100.0	627		
Secondary Complete	23.2	52.5	23.2	1.2	100.0	1,748		
Post-Secondary	31.6	55.8	12.6	0.0	100.0	684		
Socioeconomic Index								
Low	13.1	51.6	29.2	6.2	100.0	1,734		
Medium	20.0	54.7	23.1	2.2	100.0	2,805		
High	29.3	56.7	13.4	0.6	100.0	752		

Tabl	e 1	2.3	В
Ian	6	2.0	

Percent Distribution of Best Age to Start School-Based Courses on Sexually Transmitted Infections, as Reported by Men 15-49 Who are in Favor of Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

	Sexually Transmitted Infections								
				Don't		No. of			
Characteristic	≤ 13	14-15	16 +	Know	Total	Cases			
Total	11.1	38.5	45.9	4.5	100.0	1,515			
Strata									
Metro Tirana	18.9	39.3	39.6	2.3	100.0	645			
Other Urban	10.1	43.0	42.7	4.2	100.0	490			
Other Rural	8.3	35.1	50.8	5.7	100.0	380			
Residence									
Urban	13.5	41.7	41.3	3.5	100.0	1,047			
Rural	8.8	35.3	50.4	5.5	100.0	468			
Age Group									
15-24	9.2	52.5	35.7	2.6	100.0	536			
25-34	9.8	36.6	48.1	5.6	100.0	419			
35-49	14.1	26.0	54.3	5.6	100.0	560			
Marital Status									
Married	11.9	28.0	54.3	5.8	100.0	857			
Previously Married	**	**	**	**	**	12			
Never Married	9.9	52.9	34.7	2.5	100.0	646			
Living Children									
0	10.7	50.6	35.8	2.9	100.0	750			
1	5.4	42.4	45.9	6.2	100.0	186			
2	11.9	27.0	56.3	4.9	100.0	404			
3 +	15.3	15.0	61.6	8.0	100.0	175			
Education Level									
Primary or Less	6.6	36.6	48.7	8.1	100.0	523			
Secondary Incomplete	13.4	50.0	36.0	0.6	100.0	186			
Secondary Complete	12.9	36.1	48.1	2.9	100.0	585			
Post-Secondary	20.3	43.8	35.9	0.0	100.0	221			
Socioeconomic Index									
Low	7.9	33.2	52.3	6.6	100.0	500			
Medium	11.7	40.4	44.7	3.2	100.0	737			
High	20.5	50.2	27.5	1.8	100.0	278			

** Percentages are not shown when base is less than 25 cases.

Table 12.4 APercentage of Young Adult Women Aged 15-24Who Discussed The Indicated Family Life Education TopicsWith a Parent Before They reached Age 18, by Selected CharacteristicsReproductive Health Survey: Albania 2002

Characteristic	Any Topic	Menstrual Cycle	Abstinence Before Marriage	How Pregnancy Occurs	HIV/ AIDS	Other Sexually Transmitted Diseases	Methods of Contraception	No. of Cases
Total	64.4	50.8	44.5	21.4	14.7	7.9	7.6	2,030
Strata								
Metro Tirana	69.9	56.4	40.9	26.5	25.6	14.7	11.4	746
Other Urban	70.2	61.0	48.8	28.3	20.0	13.2	12.5	583
Other Rural	60.5	44.8	43.5	17.0	9.5	3.8	4.4	701
Residence								
Urban	72.0	61.6	46.6	28.7	23.0	14.3	12.5	1,193
Rural	59.7	44.1	43.2	16.8	9.5	3.9	4.5	837
Age Group								
15-17	63.9	53.9	44.3	22.5	16.7	9.8	8.9	658
18-19	68.0	51.1	49.5	19.7	15.3	6.6	7.6	436
15-19	65.5	52.7	46.4	21.4	16.2	8.5	8.4	1,094
20-24	63.0	48.4	42.1	21.3	12.9	7.1	6.6	936
Education Level								
Primary or Less	58.2	43.2	40.9	15.9	8.7	3.6	4.2	1,013
Secondary Incomplete	72.4	63.4	50.4	29.5	25.5	16.1	12.4	446
Secondary Complete	71.4	54.9	50.5	25.9	17.8	9.7	11.7	332
Post-Secondary	80.4	72.2	46.8	36.4	29.8	18.3	13.8	239
Socioeconomic Index								
Low	55.6	40.9	39.8	13.0	6.3	2.5	3.9	649
Medium	69.5	56.3	48.0	25.9	19.1	10.9	9.4	1,106
High	76.8	65.9	46.3	35.2	28.9	16.4	14.5	275
Sexual Experience								
Yes	65.0	51.6	42.7	25.4	13.7	7.1	6.6	748
No	64.2	50.4	45.4	19.5	15.2	8.3	8.0	1,282

Table 12.4 BPercentage of Young Adult Men Aged 15-24Who Discussed The Indicated Family Life Education TopicsWith a Parent Before They Reached Age 18, by Selected CharacteristicsReproductive Health Survey: Albania 2002

Characteristic	Any Topic	HIV/AIDS	How Pregnancy Occurs	Methods of Contraception	Other Sexually Transmitted Diseases	No. of Cases
Total	11.1	8.7	5.4	2.1	2.1	590
Strata						
Metro Tirana	20.4	18.3	7.2	5.8	5.2	234
Other Urban	17.4	13.3	8.7	2.2	3.4	177
Other Rural	4.9	3.2	3.2	0.9	0.3	179
Residence						
Urban	19.6	16.2	8.4	3.8	4.6	367
Rural	5.1	3.3	3.3	0.9	0.3	223
Age Group						
15-17	13.1	11.6	5.4	3.2	1.9	282
18-19	11.5	6.8	7.0	2.0	1.3	119
15-19	12.7	10.2	5.8	2.8	1.7	401
20-24	9.1	6.8	4.8	1.2	2.5	189
Education Level						
Primary or Less	5.2	3.1	3.0	0.8	0.2	264
Secondary Incomplete	21.9	19.0	8.7	4.4	3.7	178
Secondary Complete	9.0	6.4	5.4	1.1	3.2	106
Post-Secondary	28.6	25.8	13.5	8.0	8.0	42
Socioeconomic Index						
Low	8.6	5.5	4.6	0.8		222
Medium	10.5	8.6	5.4	2.6	3.1	282
High	28.1	26.3	10.1	6.6	7.8	86
Sexual Experience						
Yes	13.7	9.8	7.3	2.9	3.4	157
No	10.1	8.3	4.6	1.8	1.5	433

Table 12.5 A Percentage of Young Adult Women Aged 15-24 Who Were Taught Indicated Family Life Education Topics in School Before They Reached Age 18, by Selected Characteristics Reproductive Health Survey: Albania 2002

Characteristic	Any Topic	Female Reproductive System	Menstrual Cycle	Male Reproductive System	How Pregnancy Occurs	HIV/ AIDS	Contraceptive Methods	Other Sexually Transmitted Diseases	No. of Cases
Total	76.8	66.9	65.9	63.0	57.2	52.5	24.1	23.5	2,030
Strata									
Metro Tirana	79.4	69.2	69.4	64.6	61.7	61.1	35.3	35.1	746
Other Urban	87.8	80.1	76.0	77.6	72.0	68.5	36.4	37.1	583
Other Rural	71.2	60.5	60.5	56.1	49.5	43.2	15.7	14.5	701
Residence									
Urban	86.4	77.7	75.2	74.8	69.7	68.2	37.5	38.2	1,193
Rural	70.8	60.2	60.1	55.7	49.4	42.8	15.7	14.4	837
Age Group									
15-17	83.5	74.4	73.4	70.6	63.0	64.5	29.0	28.8	658
18-19	80.3	69.3	69.1	65.8	61.1	56.9	26.1	25.5	436
15-19	82.2	72.4	71.7	68.6	62.2	61.4	27.8	27.5	1,094
20-24	70.2	60.1	58.7	56.1	51.0	41.5	19.4	18.7	936
Education Level									
Primary or Less	64.9	52.2	53.2	47.9	41.8	35.8	11.7	9.9	1,013
Secondary Incomplete	95.4	90.4	84.2	85.3	79.2	78.5	43.7	44.4	446
Secondary Complete	91.9	83.9	82.4	81.7	76.4	73.1	36.7	41.7	332
Post-Scondary	96.7	92.9	89.6	91.9	88.2	81.7	49.3	45.4	239
Socioeconomic Index									
Low	67.6	55.7	55.7	51.7	46.9	41.2	15.4	12.2	649
Medium	81.7	72.7	71.7	69.0	62.0	57.4	28.4	29.4	1,106
High	92.9	86.8	80.4	82.5	79.2	79.1	40.5	43.7	275

Table 12.5 B Percentage of Young Adult Men Aged 15-24 Who Were Taught Indicated Family Life Education Topics in School Before They Reached Age 18, by Selected Characteristics Reproductive Health Survey: Albania 2002

Characteristics	Any Topic	Female Reproductive System	Menstrual Cycle	Male Reproductive System	How Pregnancy Occurs	HIV/ AIDS	Contraceptive Methods	Other Sexually Transmitted Diseases	No. of Cases
Total	63.7	42.9	23.2	44.3	44.7	56.5	29.8	24.2	590
Strata									
Metro Tirana	67.4	44.6	24.9	45.8	47.2	61.4	32.9	26.4	234
Other Urban	75.1	53.6	33.0	53.6	54.8	67.1	36.3	32.0	177
Other Rural	56.9	37.1	17.8	39.1	38.7	49.5	25.6	19.6	179
Residence									
Urban	72.9	50.1	30.8	50.8	52.4	66.2	36.4	30.9	367
Rural	57.2	37.9	17.9	39.6	39.1	49.6	25.2	19.5	223
Age Group									
15-17	63.3	42.4	22.3	44.7	42.8	58.4	31.4	25.8	282
18-19	68.0	47.4	22.1	47.4	49.2	58.1	34.1	21.7	119
15-19	64.6	43.8	22.2	45.5	44.6	58.3	32.2	24.6	401
20-24	62.6	41.8	24.5	42.7	44.7	54.1	26.8	23.7	189
Education Level									
Primary or Less	47.5	30.8	15.5	30.8	29.5	38.7	17.0	15.9	264
Secondary Incomplete	80.8	57.2	27.1	61.3	60.5	76.3	41.8	29.8	178
Secondary Complete	77.6	47.6	28.4	49.0	55.9	71.9	40.4	29.1	106
Post-Secondary	91.9	77.8	58.1	79.0	76.5	82.4	59.0	58.3	42
Socioeconomic Index									
Low	54.8	34.1	14.8	35.8	35.5	44.9	20.7	15.1	222
Medium	69.0	47.7	26.7	48.6	48.3	64.6	34.1	28.5	282
High	85.6	66.7	51.6	68.0	75.7	77.9	57.9	52.0	86
Sexual Experience									
Yes	66.4	46.0	27.8	46.7	49.3	57.9	31.4	28.8	157
No	62.6	41.7	21.4	43.2	42.7	55.9	29.2	22.3	433

Tables

Table 12.6 A Percentage of Young Adult Women Aged 15-24 Who Were Taught in School About Indicated Family Life Education Topics by Specific Ages

Reproductive Health Survey: Albania 2002

Chapter 12

	Perce	By Age	- No of			
Торіс	<14	<15	<16	<17	<18	No. of Cases
Female Reproductive System	10.6	54.4	60.1	65.9	66.9	2,030
The Menstrual Cycle	12.9	56.7	61.8	65.2	65.9	2,030
Male Reproductive System	9.2	51.4	56.4	61.9	63.0	2,030
How Pregnancy Occurs	6.8	45.0	50.2	56.1	57.2	2,030
HIV/AIDS	3.6	35.8	41.9	49.6	52.5	2,030
Contraceptive Methods	1.4	13.4	16.8	21.9	24.1	2,030
Other Sexually Transmitted Diseases	1.2	13.4	17.2	21.2	23.5	2,030

Table 12.6 B Percentage of Young Adult Men Aged 15-24 Who Were Taught in School About Indicated Family Life Education Topics by Specific Ages Reproductive Health Survey: Albania 2002

	Percent Who Have Taken Course By Age								
Торіс	<14	<15	<16	<17	<18	No. of Cases			
HIV/AIDS	8.7	34.6	44.4	52.9	56.5	590			
Male Reproductive System	8.1	28	36.6	43	44.3	590			
Female Reproductive System	7.8	27.1	35.4	41.7	42.9	590			
How Pregnancy Occurs	7.6	26.2	37.6	43.6	44.7	590			
The Menstrual Cycle	4.2	15.6	19.4	21.4	23.2	590			
Contraceptive Methods	2.9	14.9	23.1	27.4	29.8	590			
Other Sexually Transmitted Diseases	2.4	14.7	20.1	23.2	24.2	590			

Table 12.7 A Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Women 15-24 Who Had Received Any Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

		Most In	Most Important Source of Information Related to Sexual Matters									
Characteristic	Teacher	Radio/ TV	Books/ Newspapers/ Magazines	Co-Worker/ Peer	Mother/ Father	Other Relative	Boyfriend/ Friends	Partner/ Husband	Doctor/Nurse/ Midwife	Don't Know	Total	No. of Cases
Total	48.8	23.7	9.9	7.6	3.9	2.0	3.0	0.3	0.5	0.4	100.0	1,599
Strata												
Metro Tirana	45.6	19.7	12.7	7.2	8.0	1.6	4.0	0.1	0.8	0.3	100.0	599
Other Urban	46.1	20.2	12.4	8.7	4.8	2.8	4.0	0.5	0.4	0.0	100.0	511
Other Rural	51.2	26.8	7.6	7.1	2.3	1.7	2.2	0.2	0.4	0.6	100.0	489
Residence												
Urban	45.6	20.0	12.6	8.3	5.9	2.4	4.1	0.4	0.6	0.1	100.0	1,027
Rural	51.2	26.5	7.8	7.0	2.5	1.7	2.2	0.1	0.4	0.5	100.0	572
Age Group												
15-17	55.7	22.9	7.8	5.1	4.1	1.8	1.9	0.0	0.6	0.0	100.0	563
18-19	46.4	21.4	15.2	7.1	4.2	2.2	2.8	0.3	0.2	0.3	100.0	357
15-19	52.0	22.3	10.7	5.9	4.1	2.0	2.3	0.1	0.5	0.1	100.0	920
20-24	44.1	25.8	8.7	10.0	3.7	2.1	4.1	0.5	0.4	0.7	100.0	679
Education Level												
Primary or Less	46.9	28.0	6.1	9.4	2.7	2.6	2.8	0.4	0.5	0.5	100.0	631
Secondary Incomplete	54.3	21.2	9.3	5.2	5.6	1.9	2.1	0.0	0.4	0.1	100.0	431
Secondary Complete	47.5	19.3	16.5	5.8	4.3	0.6	4.8	0.3	0.3	0.6	100.0	304
Post-Secondary	47.9	15.3	18.9	6.5	6.1	1.6	2.9	0.0	0.6	0.0	100.0	233
Socioeconomic Index												
Low	50.7	26.6	7.4	7.0	2.4	1.3	3.0	0.3	0.6	0.7	100.0	422
Medium	47.6	23.2	10.9	8.0	4.0	2.7	2.8	0.3	0.4	0.2	100.0	915
High	48.4	15.8	13.1	7.5	9.3	0.7	4.4	0.0	0.8	0.0	100.0	262
Sexual Experience												
Yes	39.5	27.4	7.3	10.1	5.0	3.1	5.7	0.9	0.5	0.4	100.0	524
No	52.4	22.3	10.9	6.6	3.5	1.6	2.0	0.0	0.5	0.3	100.0	1,075

Table 12.7 B

Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Men 15-24 Who Had Received Any Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

	Most Important Source of Sexual Information Related to Sexual Matters										
Characteristic	Teacher	Radio/TV	Books/ Newspapers, Magazines	Co-Worker/ Peer	Girlfriend/ Friends	Mother/ Father	Doctor	Other Relative	Don't Know	Total	No. of Cases
Total	53.3	21.4	17.8	3.8	1.6	1.1	0.7	0.1	0.3	100.0	400
Strata											
Metro Tirana	46.1	22.1	20.4	5.7	1.9	2.1	1.2	0.6	0.0	100.0	161
Other Urban	61.6	18.2	13.6	3.8	0.8	1.2	0.0	0.0	0.8	100.0	137
Other Rural	50.8	23.2	19.5	3.0	2.0	0.5	1.0	0.0	0.0	100.0	102
Residence											
Urban	56.2	18.4	16.5	4.9	1.3	1.7	0.5	0.0	0.5	100.0	272
Rural	50.8	24.0	19.0	2.8	1.8	0.5	1.0	0.2	0.0	100.0	128
Age Group											
15-17	61.0	15.8	16.5	2.8	2.0	0.6	1.2	0.0	0.0	100.0	186
18-19	52.2	21.7	15.8	8.1	0.4	1.8	0.0	0.0	0.0	100.0	86
15-19	58.4	17.6	16.3	4.4	1.5	1.0	0.9	0.0	0.0	100.0	272
20-24	46.6	26.4	19.8	3.0	1.7	1.1	0.5	0.3	0.6	100.0	128
Education Level											
Primary or Less	44.0	24.2	20.9	6.1	1.5	1.0	1.3	0.3	0.7	100.0	126
Secondary or Greater	59.1	19.6	15.9	2.4	1.6	1.1	0.4	0.0	0.0	100.0	274
Socioeconomic Index											
Low	45.1	29.7	16.8	5.5	1.3	0.0	1.2	0.3	0.0	100.0	122
Medium	60.6	16.5	17.0	2.3	2.1	1.1	0.0	0.0	0.5	100.0	207
High	52.3	12.3	24.3	4.0	0.5	4.7	1.9	0.0	0.0	100.0	71
Sexual Experience											
Yes	50.2	26.3	14.5	2.9	1.2	3.0	0.7	0.4	0.8	100.0	111
No	54.7	19.2	19.2	4.2	1.7	0.2	0.7	0.0	0.0	100.0	289

Table 12.8 A

Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Women 15-24 Who Never Received Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

	Most Important Source of Information Related to Sexual Matters											
Characteristic	Radio/ TV	Co-Worker/ Peer	Books/ Newspapers/ Magazines	Mother/ Father	Other Relative	Boyfriend/ Friends	Partner/ Husband	Doctor	Teacher	Don't Know	Total	No. of Cases
Total	43.9	15.1	5.9	10.7	9.0	4.3	2.4	0.4	2.0	6.3	100.0	431
Strata												
Tirana	49.3	12.1	6.8	15.3	7.6	1.5	4.5	0.5	0.0	2.4	100.0	147
Other Urban	41.7	20.0	9.1	6.7	6.9	1.7	5.1	2.1	2.8	4.0	100.0	72
Rural	43.3	14.8	5.1	10.6	9.6	5.4	1.5	0.0	2.3	7.5	100.0	212
Residence												
Urban	43.3	15.6	9.1	12.2	6.3	1.9	5.5	1.3	1.7	3.0	100.0	166
Rural	44.0	15.0	4.9	10.3	9.7	5.1	1.5	0.1	2.1	7.2	100.0	265
Age Group												
15-17	40.2	10.6	6.0	12.1	11.2	2.2	0.0	0.0	5.2	12.3	100.0	95
18-19	50.2	16.8	6.5	6.3	8.1	3.8	3.4	0.0	3.4	1.5	100.0	79
15-19	44.7	13.4	6.2	9.5	9.8	3.0	1.5	0.0	4.4	6.5	100.0	174
20-24	43.3	16.5	5.6	11.6	8.3	5.4	3.0	0.6	0.3	5.4	100.0	257
Education Level												
Primary or Less	43.8	14.9	5.8	11.3	8.5	4.4	2.7	0.1	1.6	7.0	100.0	382
Secondary or Greater	44.5	17.7	6.5	5.7	12.9	4.1	0.0	2.9	5.7	0.0	100.0	49
Socioeconomic Index												
Low	43.6	13.9	6.6	9.4	11.1	5.6	1.0	0.0	2.6	6.1	100.0	227
Medium	44.5	17.0	4.8	12.8	5.8	2.8	3.6	0.9	1.4	6.4	100.0	191
High	**	**	**	**	**	**	**	**	**	**	**	13

** Percentages are not shown when base is less than 25 cases.

Table 12.8 B

Percent Distribution of The Most Important Source of Information Related to Sexual Matters, Reported by Young Adult Men 15-24 Who Never Received Sex Education in School, by Selected Characteristics Reproductive Health Survey: Albania 2002

	Mos	st Import	tant Source	of Inf	ormation	Related	d to Se	xual Mat	ters		
Characteristic	Radio/TV	Co-Worker/ Peer	Books/ Newspapers, Magazines	Teacher	Mother/ Father	Girlfriend/ Friends	Doctor	Other Relative	Don't Know	Total	No. of Cases
Total	52.5	19.8	4.7	2.4	2.0	1.6	1.3	0.2	15.6	100.0	190
Strata											
Metro Tirana	52.7	14.4	0.0	4.8	2.6	5.5	7.9	0.0	12.1	100.0	73
Other Urban	34.8	38.7	4.8	1.3	1.3	1.3	0.0	1.3	16.4	100.0	40
Other Rural	57.5	15.7	5.8	2.0	2.0	0.7	0.0	0.0	16.2	100.0	77
Residence											
Urban	43.8	26.5	2.9	3.3	2.2	3.3	3.5	0.8	13.7	100.0	95
Rural	56.5	16.8	5.5	1.9	1.9	0.8	0.3	0.0	16.4	100.0	95
Age Group											
15-17	61.8	14.3	3.2	3.2	3.2	1.3	0.0	0.6	12.4	100.0	96
18-19	45.3	34.0	0.0	1.7	4.8	1.7	0.8	0.0	11.7	100.0	33
15-19	57.6	19.4	2.4	2.8	3.6	1.4	0.2	0.4	12.2	100.0	129
20-24	46.3	20.3	7.4	1.8	0.0	1.8	2.7	0.0	19.7	100.0	61
Education Level											
Primary or Less	54.4	21.9	1.8	1.8	2.2	0.9	1.2	0.3	15.5	100.0	138
Secondary or Greater	47.0	13.7	12.9	4.0	1.4	3.7	1.5	0.0	15.8	100.0	52
Socioeconomic Index											
Low	46.9	25.6	0.0	2.6	2.2	0.7	0.0	0.4	21.6	100.0	100
Medium	60.2	12.6	12.3	2.1	1.5	3.1	0.5	0.0	7.7	100.0	75
High	**	**	**	**	**	**	**	**	**	**	15
Sexual Experience											
Yes	49.6	19.9	3.3	2.9	1.6	2.9	3.7	0.0	16.1	100.0	46
No	53.6	19.8	5.2	2.1	2.1	1.1	0.4	0.3	15.4	100.0	144

** Percentages are not shown when base is less than 25 cases.

Table 12.9 A

Knowledge of Young Adult Women Aged 15-24, on Selected Reproductive Health Issues by Whether or Not Specific Topics Were Discussed With a Parent or Taught in School (Percent Distribution) Reproductive Health Survey: Albania 2002

			Menstrual h Parents	Taught in School About Menstrual Cycl	
Most Likely to Become Pregnant During Menstrual Cycle	Total	Yes	No	Yes	No
Just Before Her Periods Starts	2.7	2.8	2.6	3.4	1.3
During Her Periods	0.8	1.2	0.5	1.0	0.6
Right After Period Ends	14.8	15.7	14.0	13.7	17.1
Halfway Between Periods	14.7	20.9	8.3	18.7	7.0
Do Not Know	66.9	59.5	74.6	63.2	74.1
Total	100.0	100.0	100.0	100.0	100.0
No. of Cases	2,030	1,109	921	1,389	641

		Pregnancy	sed How Occurs With ents	Taught in S How Pregna	
Breastfeeding Affect on Getting Pregnant	Total	Yes	No	Yes	No
Increases The Chance	5.6	7.1	5.2	4.7	6.8
Decreases The Chance	11.5	19.2	9.4	12.7	9.8
Has No Effect	12.4	10.0	13.1	12.7	12.0
Do Not Know	70.5	63.7	72.3	69.9	71.3
Total	100.0	100.0	100.0	100.0	100.0
No. of Cases	2,030	510	1,520	1,225	805

		Pregnancy	sed How Occurs With ents	Taught in School Abou How Pregnancy Occurs		
Possibile to Get Pregnant at First Intercourse	Total	Yes	No	Yes	No	
Agree	49.1	63.0	45.4	52.8	44.1	
Disagree	21.1	17.8	22.0	20.1	22.4	
Do Not Know	29.8	19.2	32.7	27.0	33.5	
Total	100.0	100.0	100.0	100.0	100.0	
No. of Cases	2,030	510	1,520	1,225	805	

Table 12.9 B Knowledge of Young Adult Men Aged 15-24, on Selected Reproductive Health Issues by Whether or Not Specific Topics Were Discussed With a Parent or Taught in School (Percent Distribution) Reproductive Health Survey: Albania 2002

			ow Pregnancy ith Parents	Abou	In School it How cy Occurs
Breastfeeding Affect on Getting Pregnant	Total	Yes	No	Yes	No
Increases the Chance	0.7	1.3	0.7	0.8	0.6
Decreases the Chance	9.8	29.6	8.7	12.6	7.6
Has No Effect	9.2	8.6	9.2	16.0	3.6
Do Not Know	80.3	60.5	81.4	70.6	88.1
Total	100.0	100.0	100.0	100.0	100.0
No. of Cases	590	39	551	282	308
			ow Pregnancy ith Parents	Abou	in School It How cy Occurs
Possible to Get Pregnant at First Intercourse	Total	Yes	No	Yes	No
Agree	51.9	69.6	50.9	57.0	47.9
Disagree	19.4	4.1	20.3	23.4	16.3
Do Not Know	28.6	26.3	28.7	19.7	35.8
Total	100.0	100.0	100.0	100.0	100.0

282

308

551

590

39

No. of Cases

CHAPTER 13


YOUNG ADULT SEXUAL AND CONTRACEPTIVE EXPERIENCE

Sexual Experience

module included The young adult questions on age and partner at first sexual intercourse, as well as use of contraception for those with sexual experience. The first sexual relation was classified as premarital or marital as has been done in other countries of Eastern Europe (CDC and MACRO, 2003). This classification was obtained from three specific questions in the survey instrument: the dates of the first sexual relation and first marriage (or consensual union), if ever married, and the relationship to the partner at the time of this first sexual experience. If the first partner was reported to be a husband or wife, the dates of first sexual experience and first marriage (or consensual union) were compared to determine if the first sexual experience was marital or pre-marital. If the first sexual experience occurred at least

one month prior to the date of marriage, this was classified as pre-marital.

As shown in Table 13.1A and Figure 13.1, 32% of young adult women reported sexual experience, of which slightly less than half (14% of the total) are classified as having premarital sex. Sexual experience varies by residence, with slightly higher proportions of urban women having had sexual experience. Premarital sex, in particular, is reported by a greater proportion of urban women (23%) than rural women (8%). The sexual experience rate increases with age and is eight times higher for older young adults 22-24 years of age (66%) than for adolescents 15-17 years of age (8.5%), although, among sexually experienced young women, adolescents are more likely to have had premarital sex compared with marital sex.



Among 15-17 year olds, 5% have had premarital sex and only 3% marital sex; whereas for 22-24 year olds, 23% have had premarital sex while 43% report first sex after marriage. For both educational level and socioeconomic status, there is a positive association with premarital first intercourse. For example, among the least educated, 25% of women had first intercourse after marriage and 11% had premarital first intercourse, and among those with postsecondary education, only 7% had had postmarital first intercourse while 28% had premarital first intercourse. A similar pattern is observed for socioeconomic status, with marital first intercourse declining from 19.5% to 12.5% and premarital first intercourse increasing from 8% to 32% with increasing socioeconomic level.

Twenty-nine percent of the young adult men reported sexual experience (Table 13.1B and Figure 13.1). The great majority were classified as premarital (27%)compared to 3% marital). Premarital sexual experience rates tend to be higher for urban men and increase with age, educational attainment and socioeconomic status. The variation in sexual experience rates among young adult men is almost entirely due to premarital sexual experience. Thus, 40% of young men in Metro Tirana, 61% of all 22-24 year olds, 79% of those with postsecondary education, and 47% of young men with a high socioeconomic level have had premarital sexual intercourse.

Table 13.2 shows young adult female sexual experience rates for Albania in the context of other reproductive health and demographic health surveys for countries in its region. It appears that Albania's rates have more in common with countries in the Caucasus region than Eastern Europe in terms of the overall level of sexual experience. With regard to rates of premarital sexual experience, however, Albania resembles neither Eastern Europe nor the Caucasus. Albania's rates of premarital sex (7% of 15–19 year olds and 21% of 20–24 year olds) are substantially lower than Eastern Europe rates but substantially higher than those for Azerbaijan and Georgia.

First Sexual Intercourse

The relationship to their first partner is shown for females in Table 13.3A and for males in Table 13.3B. As previously noted, 58% of sexually experienced young adult women were married at the time of their first sexual encounter (Table 13.3A). Another 35% reported their partner to be their fiancée. A much higher proportion of women who live in rural areas (74%) had their first sexual experience with their husband than did their counterparts in urban areas (37%). Marital sex is inversely related to educational attainment and socioeconomic status. Most premarital sex is reported to have been with a fiancée or boyfriend. There is no evidence of casual sex at first sexual experience among the women. However, post-secondary education is the only group to report most first partners to be a "boyfriend" rather than their "fiancée."

Most of the men indicated that their first partner was a "girlfriend" (43%) (Table 13.3B). Only 9% of young men reported their first sexual experience to be with their "wife/consensual partner." Overall, the young men were more likely to report first sex with weaker relationships than the young women. Thirty-two percent of young men reported first sex with a "friend" or "lover" and 7% with an "acquaintance" compared to only 0.2% of young women reporting first sex with a "friend" and none with an "acquaintance." The young men show the same direction of association as the young women between premarital relationship and the selected sociodemographic variables, although at much higher levels of premarital sex. Only 1% of sexually experienced young men reported first sex with a prostitute.

Tables 13.4A and 13.4B show the percentages that have had sexual intercourse by a given age for young women and men, respectively. We can see that a very low percentage (1%) of young women and men have had sex before the age of 15. Percentages having had sex by specific ages are quite similar for women and men. Thirteen percent of young adult women and 12% of young adult men have had sexual intercourse before age 18, and 30% of women and 27% of men have had sex before age 22.

Among both women and men, those living in urban areas and having highest socioeconomic status are more likely to have a young age at first sex than young adults in rural areas or with low SES. Women whose first sexual experience was premarital also report a younger age at first sex than those who were married at first sex. With respect to educational level, however, women and men have distinctly different patterns. The least educated women have younger ages at first sex compared to better educated women, but among men it is the most educated that have the youngest ages at first sex.


Tables 13.5A and B present the age differences between young adults and their sexual partners at first sexual experience. Approximately 2% of the young adult women reported their first partner to be younger (Table 13.5A). Three-quarters of the women (73%) report a wide age difference of five years or more between themselves and their first partner. There are not strong effects of residence or marital status at first sex on the age difference at first sex. However, the largest age difference appears to be negatively associated with age at first sex.

Among the young adult men, again the evident norm is that the men tend to be older than their female partner (Table 13.5B). Fifty-six percent report having been older than their first partner. However, another 13% report having had a first partner older than themselves. These older partners among males suggests that young men are sometimes initiated into sex with older women, a result found in other countries for which male data are available (E Ferraz et al., 1992; L Morris et al., 1995; R Tejada, J Herold and L Morris, 1997). The younger the age at first sex, the more likely the man had an older first partner.

InTable13.6A, four-fifths(81%) of sexually experienced young women reported that they or their partner used a contraceptive method at first sexual relationship; 74% of those with marital sex and 91% of those with premarital sex. Figure 13.2 shows that 77% of all women reported use of withdrawal at first sexual intercourse and only 4% used another method. Thus, among contraceptors, almost all women who reported using a method said that their partner used withdrawal at first sexual experience (96%) (Table 13.6A). Use of withdrawal does not show an association with residence or age at first sex. In all groups except those with a post-secondary education, more than 91% of users used withdrawal. Condom use, while at low levels, tends to be higher for premarital intercourse and to be positively related to educational attainment and socioeconomic status.

At first sexual experience, 85% of young adult men report using contraception, closely divided between condoms (37%) and withdrawal (47%) (Figure 13.2). Among contraceptive users, 56% of men used withdrawal at first intercourse, while 43% used condoms (Table 13.6). The relative weight of condoms vs. withdrawal varies by several sociodemographic characteristics. Overall contraceptive use at first sex tended to be higher in rural areas than in urban areas among

the men. This can be attributed to the much higher use of withdrawal in rural areas (67%) compared to urban areas (42%), although again interpreted with caution due to small sample size. At the same time, condom use was only 33% of users in rural areas compared to 57% in urban areas. Condoms are more likely to be used at first sex by urban and younger men, more educated men and young men from higher socioeconomic groups. The patterns for withdrawal are the reverse of those for condoms for these variables.



Reasons for not using contraception at first sex are shown for young adult women in Table 13.7. Three-quarters of young women reported that they did not use contraception because they wanted to become pregnant. For women who were married at first sexual experience, this proportion reaches 85%, compared with 23% for women whose first sexual experience was premarital. Only 4% reported they did not know about contraception and 4% said they thought contraceptives harmful. were that Although the sample size is especially low for women who had premarital sex, they reported a high frequency of "fear of side effects" (18%) and "sexual intercourse was unexpected" (17%). Reasons for not using contraception cannot be calculated for the young adult men, because too few men (n=29) reported no contraception at first sex.

Current Sexual Activity

First sexual experience does not always reflect current sexual activity, especially among unmarried young adults. For all reproductive age women, particularly the married, the sexually active are conventionally defined as those having had sex in the month prior to the survey. Unmarried young adults, however, are known to have more sporadic sexual consequently encounters. and the sexually active are variously defined as having had sexual intercourse in the last 30 days, the last two months or the last three months prior to interview. In Table 13.8, we can see the differences in sexual activity by marital status for young adult women and men. While all of the married women have had sexual experience, only 7 percent of 15-24 year old divorced, widowed, and never married women report having had

sexual experience (Table 13.8). The age difference in sexual experience reflects higher proportions married among 20-24 year olds. The recency of sexual activity by marital status shows that two-thirds (65%) of married women had sexual intercourse in the last month, and of those who did not have recent sexual intercourse the majority were pregnant or postpartum (19%). Among sexually experienced unmarried women, however, only about half had intercourse in the last 30 days, none report being pregnant or postpartum, and half reported the last sexual experience more than a month ago.

The pattern of current sexual activity for young men is similar to that of young women (Table 13.8). All married 15–24 year olds report having sexual experience, with 92% having had sex in the last 30 days, and 5% abstaining due to their wife's pregnancy. At the same time, 25% of widowed, divorced and never married men reported having had sex, with 17% reporting last sex in the last 30 days and 7% reporting last sexual intercourse more than one month ago.

Recent contraceptive use among young adult women is exceptionally high, although there is very low use of modern methods. Table 13.9A shows that 76% of sexually experienced women 15-24 years of age used contraception at last intercourse. When marital status is controlled, the percentage jumps to 98% for unmarried women (Table 13.9 and Figure 13.2). This high level of contraceptive use is due almost exclusively to the withdrawal method, which was reported to have been used by 69% of married women and 87% of unmarried women at last sexual intercourse. Unmarried women have a

higher rate of use of modern methods compared to married women (11% vs. 3%, respectively), with these rates reflecting use of the condom. Two percent of married young adult women and 9% of the unmarried reported using a condom at last sexual experience.

In contrast to the young adult women, only 41% of sexually experienced young adult men report having used contraception at last intercourse (Table 13.9B). The breakdown by marital status is 68% among the married and 34% among the unmarried. The small number of cases in the denominator of these rates, however, would produce a large margin of error for the young men and thus give us less confidence in the percentages shown. However, it is clear that withdrawal is the major method and condom the only modern method. There is no use of modern methods reported by the married, and modern method use among the unmarried men is due entirely to the condom. Indeed, condom rates are at similar levels to withdrawal among the unmarried young adult men (15% and 18%, respectively) (Figure 13.2).

When nonusers of contraception were asked why they currently were not using, 44% of the sexually experienced young women responded that they were not sexually active – that is, approximately one-third of the married and almost all of the unmarried (Table 13.10). The remainder of the married young adult women said they were not using for pregnancy related reasons, either they were pregnant, trying to get pregnant, or they were postpartum. Ninety percent of the nonusing young adult men also reported no current sexual activity as their reason for not using contraception. Seven percent gave a pregnancy-related reason (data not shown).

Tables 13.11A and B show the number of sexual partners reported by sexually experienced young adults. Among the women, 11% reported no sexual partner in the last three months and 88% reported only one partner in that time period. Unmarried women are more likely to report no sexual partner compared to married women. There is little variation by educational level. Moreover, very few of the women report more than one sexual partner in their lifetime, less than 1% of married women and 15% of sexually experienced unmarried women. A larger proportion of the young men (compared to the women) reported no sexual partner in the last three months, that is, 34%. Very few men reported more than one sexual partner in the last three months (5%), and there was no variation in this percentage by educational level. A greater proportion of young men than voung women had more than one sexual partner in their lifetime. Fifty-two percent of married men and 77 percent of unmarried men reported two or more lifetime sexual partners.

Attitudes toward Condom Use

Sexually experienced young adults were asked a series of questions meant to measure attitudes towards condom use. Table 13.12 shows the results for this series of attitudinal questions for women and men. Positive attitudes were more prevalent among persons who had used condoms in the past compared to never users of condoms. Almost 100% of young female and male ever-users agreed with the statement, "Using condoms with a new partner is a smart idea." Two-thirds of females and males who had ever used condoms agreed with the statement, "It is easy to discuss using a condom with a prospective partner." The item, "Women should ask their partners to use condoms," received 73% agreement from the women but only 46% from men, and "Condoms diminish sexual pleasure was agreed to by 57% of women and 77% of men. Less than 50% of male and female ever-users agreed to such statements as "Using condoms is not necessary if you know your partner," "It's embarrassing to ask for condoms in family planning clinics or pharmacies," and "People who use condoms sleep around a lot."

Never-users of condoms presented a different pattern of response to the condom attitude questions than everusers. Only around 50% of both females and males agreed with the statement, "Using condoms with a new partner is a smart idea." One-third or fewer of both sexes agreed with "Women should ask their partners to use condoms," and "It is easy to discuss using a condom with a prospective partner." Moreover, half of the women and 57% of men believe "Using condoms is not necessary if you know your partner." The young male ever-users are more likely to have negative attitudes towards condoms than the young female ever-users, as is demonstrated by their agreement with, "People who use condoms sleep around a lot" and "It is embarrassing to ask for condoms in family planning clinics or pharmacies."

It is important to point out that very few young adults, whether ever-users or never-users or male or female, agreed with the statement, "The same condom can be used more than once." Another result that deserves attention is the high proportion of never-users who responded "don't know" to the condom attitude questions.

In Tables 13.13A & B we have the percentage of sexually experienced young adults who agreed with statements about personal reactions to a partner requesting the use of condoms for sexual intercourse. Two-thirds of the women and 84% of the men agreed that they would feel safe from pregnancy. As for safety from STDs or HIV/AIDS, 62% of women and 74% of men felt agreement with that statement. These positive reactions were more common for women in urban areas, with higher levels of education and who had ever used condoms. Among men, among the selected variables, only ever use of condoms appeared to have an effect on positive attitudes. For both young women and men, there appeared to be no age effect on positive responses.

The negative items for this question were, [would you feel] "Insulted or angry," "Like you had done something wrong," and "Suspicious that he/she may sleep around." Less than one-third of both women and men agreed with the negative items. Among women, rural residence and never use of condoms tended to increase the percentage agreement with negative attitudes. There was no clear age or education effect. Among men, however, agreement with negative attitudes was greater for rural than urban residents, the youngest men, the less educated men, and never users of condoms.

Finally, Table 13.14 shows the percentages of young adults who have ever talked with a partner about using

condoms. Twenty-four percent of the sexually experienced young adult women and 58% of the young men responded affirmatively to the question. Talking with a partner about condom use is positively associated with urban residence, educational level and past condom use, but appears to have no relationship with age among 15–24 year olds of both sexes. The strong association with educational level is shown in Figure 13.3. Twenty-seven percent of all young adult men said they had tried to obtain a condom in the last 12 months, and all of those who tried were successful in obtaining condoms (data not shown). Among the 81 young men who had obtained condoms, 51% said they have had condoms in their possession almost all or all of the time in the last 12 months (data not shown). The brands of condoms reported to be most often obtained were "For You" (47%), "Love Plus" (19%), "For You More" (13%), and "Durex" (1%).




Table 13.1 A Percent Distribution of Reported Sexual Experience by Marital Status at Time of First Sexual Experience by Selected Characteristics Among Young Women Aged 15–24 Reproductive Health Survey: Albania, 2002

	No Sexual –	Sexual E	Experience		
Characteristics	Experience	Marital	Premarital	Total	No. of Cases
Total	67.7	18.7	13.6	100.0	2,030
Strata					
Metro Tirana	61.2	17.1	21.7	100.0	746
Other Urban	66.1	12.6	21.3	100.0	583
Other Rural	70.1	21.8	8.1	100.0	701
Residence					
Urban	64.2	13.1	22.7	100.0	1,193
Rural	69.8	22.2	8.0	100.0	837
Current Age					
15-17	91.5	3.4	5.1	100.0	658
18-19	78.3	10.9	10.8	100.0	436
20-21	59.0	21.6	19.5	100.0	393
22-24	33.7	43.4	22.9	100.0	543
Age Group					
15-19	86.2	6.5	7.4	100.0	1,094
20-24	44.8	33.8	21.4	100.0	936
Education Level					
Primary or Less	63.5	25.2	11.3	100.0	1,013
Secondary	75.8	9.7	14.5	100.0	778
Post-Secondary	65.2	6.8	28.0	100.0	239
Socioeconomic Index					
Low	72.8	19.5	7.7	100.0	649
Medium	65.5	19.0	15.5	100.0	1,106
High	55.7	12.5	31.8	100.0	275

Young Adult Sexual and Contraceptive Experience

Table 13.1 B Percent Distribution of Reported Sexual Experience by Marital Status at Time of First Experience by Selected Characteristics, Among Young Men Aged 15–24 Reproductive Health Survey: Albania, 2002

		Sexual I	Experience	_	
Characteristics	No Sexual [—] Experience	Marital	Premarital	Total	No. of Cases
Total	70.7	2.7	26.6	100.0	590
Strata					
Metro Tirana	55.9	4.3	39.8	100.0	234
Other Urban	70.5	0.6	28.9	100.0	177
Other Rural	75.8	3.2	21.0	100.0	179
Residence					
Urban	65.6	1.4	33.0	100.0	367
Rural	74.3	3.6	22.1	100.0	223
Current Age					
15-17	97.8	0.0	2.2	100.0	282
18-19	88.9	0.0	11.1	100.0	119
20-21	49.2	3.7	47.1	100.0	74
22-24	31.3	8.1	60.6	100.0	115
Age Group					
15-19	95.2	0.0	4.8	100.0	401
20-24	38.7	6.3	55.1	100.0	189
Education Label					
Primary or Less	76.0	4.2	19.8	100.0	264
Secondary	71.0	1.4	27.6	100.0	284
Post-Secondary	21.4	0.0	78.6	100.0	42
Socioeconomic Index					
Low	82.5	3.1	14.3	100.0	222
Medium	61.5	2.8	35.7	100.0	282
High	52.8	0.0	47.2	100.0	86

Table 13.2

Percent Distribution of Reported Sexual Experience by Marital Status at Time of First Sexual Experience by Current Age Among Young Women Aged 15–24 Reproductive and Demographic Health Surveys (RHS and DHS) In Selected Eastern European and Former Soviet Union Countries Albania Reproductive Health Survey 2002, Final Report

	Reporte	ed Sexual Experie	ence		No. of
Region and Country	No Sexual Experience	After Marriage	Before Marriage	Total	Cases
Eastern Europe					
Albania, 2002					
15–19	86	7	7	100	1,094
20–24	45	34	21	100	936
Total	68	19	14	100	2,030
Czech Rep., 1993					
15–19	46	†	54	100	646
20–24	2	1	97	100	737
Total	27	t	73	100	1,383
Moldova, 1997					
15–19	79	6	14	100	747
20–24	17	43	40	100	910
Total	50	23	26	100	1,657
Romania, 1999					
15–19	74	4	22	100	924
20–24	22	20	58	100	1,239
Total	47	13	41	100	2,163
Russia, 1999*					
15–19	51	†	49	100	748
20–24	8	5	87	100	1,058
Total	25	3	71	100	1,806
Ukraine, 1999					
15–19	68	3	30	100	1,079
20–24	12	15	73	100	1,151
Total	40	9	51	100	2,230
Caucasus					
Azerbaijan, 2001					
15–19	90	10	†	100	1,207
20–24	53	45	3	100	1,207
Total	74	25	1	100	2,414
Georgia, 1999					
15–19	84	15	†	100	1,142
20–24	47	50	2	100	1,246
Total	67	31	1	100	2,388

* Yekaterinburg, Perm, and Ivanovo, respectively (predominantly urban sample).

† Less than 0.5%.

Source: Goldberg H et al., 1993, KIIS and CDC, 2001; MACRO International 1995-2001; Serbanescu et al., 1998, 2001, 2001, 2003; VCIOM and CDC, 1998, 2000.

Young Adult Sexual and Contraceptive Experience

Table 13.3 A Percent Distribution of Relationship to Partner at First Sexual Intercourse Among Sexually Experienced Young Women Aged 15–24 Reproductive Health Survey: Albania, 2002

	Relation	ship at Firs	st Sexual	Intercour	se	_	
Characteristics	Husband/ Consensual Partner	Fiancee	Boy- friend	Friend	Rape	- Total	No. of Cases
Total	57.9	34.5	7.4	0.2	0.1	100.0	747*
Strata							
Metro Tirana	44.2	33.3	21.6	0.3	0.6	100.0	313
Other Urban	37.1	51.2	11.3	0.4	0.0	100.0	203
Other Rural	72.9	26.5	0.6	0.0	0.0	100.0	231
Residence							
Urban	36.6	46.2	16.6	0.4	0.3	100.0	462
Rural	73.5	25.9	0.6	0.0	0.0	100.0	285
Age 1st Sex							
< 18	52.8	39.6	7.1	0.3	0.3	100.0	305
18-19	57.6	31.6	10.7	0.2	0.0	100.0	254
20-24	65.7	30.2	4.1	0.0	0.0	100.0	188
Marital Status at 1st Sex							
Premarital	0.0	81.9	17.5	0.4	0.3	100.0	371
Marital	100.0	0.0	0.0	0.0	0.0	100.0	376
Education Level							
Primary or Less	69.0	29.6	1.4	0.1	0.0	100.0	434
Secondary	40.0	48.7	10.7	0.4	0.2	100.0	218
Post-Secondary	19.6	31.6	47.8	0.0	0.9	100.0	95
Socioeconomic Index							
Low	71.6	28.0	0.4	0.0	0.0	100.0	222
Medium	55.1	36.6	8.0	0.2	0.1	100.0	406
High	28.3	44.4	26.1	0.4	0.7	100.0	119

* Excludes one case with missing information on age at first sexual intercourse

Table 13.3 BPercent Distribution of Relationship to Partner at First Sexual Intercourse
Among Sexually Experienced Young Men Aged 15–24
Reproductive Health Survey: Albania, 2002

	Relationship at First Sexual Intercourse								_	
Characteristics	Wife/ Consensual Partner	Fiancee	Girlfriend	Lover	Friend	Just Met Acquaintance	Prostitute	Unknown	Total	No. of Cases
Total	9.3	1.7	42.7	18.7	13.6	6.9	1.1	6.0	100.0	156
Strata										
Metro Tirana	9.9	2.7	42.0	17.6	11.5	11.9	0.0	4.5	100.0	81
Other Urban	2.0	0.0	50.9	13.3	21.5	2.0	2.0	8.2	100.0	40
Other Rural	13.4	2.2	38.1	22.5	10.1	6.7	1.2	5.6	100.0	35
Residence										
Urban	4.2	0.5	49.3	14.9	17.5	5.8	1.1	6.6	100.0	106
Rural	14.2	2.9	36.5	22.2	9.9	7.9	1.1	5.5	100.0	50
Age 1st Sex										
< 18	2.5	0.6	46.3	22.6	11.8	5.0	1.3	9.8	100.0	73
18-19	7.9	0.0	40.1	13.1	25.8	6.2	0.0	7.0	100.0	44
20-24	19.9	5.0	40.4	18.7	4.2	10.0	1.8	0.0	100.0	39
Marital Status at 1st Sex										
Premarital	0.0	1.9	47.1	20.6	15.0	7.6	1.2	6.6	100.0	142
Marital	**	**	**	**	**	**	**	**	**	14
Education Level										
Primary or Less	17.5	1.2	32.5	18.6	16.4	4.5	1.3	8.0	100.0	54
Secondary	4.7	3.0	50.2	15.7	10.4	9.6	1.3	5.0	100.0	73
Post-Secondary	0.0	0.0	49.7	26.4	14.9	5.5	0.0	3.5	100.0	29
Socioeconomic Index										
Low	18.0	6.2	37.3	15.4	14.2	5.4	0.0	3.6	100.0	38
Medium	7.3	0.0	44.3	18.2	13.6	7.8	1.9	6.9	100.0	83
High	0.0	0.0	46.8	27.1	12.6	6.4	0.0	7.2	100.0	35

* Exclude one case with missing information on age at first sexual intercourse

**Percentages are not shown when base is less than 25 cases

Table 13.4 A Percent of Women Aged 15-24 Who Had Their First Sexual Intercourse Before Selected Ages, by Various Characteristics Reproductive Health Survey: Albania 2002

	Age at F	irst Sex	ual Inte	rcourse	_		
Characteristics	<15	<18	<20	<22	Has Had Intercourse	Never Had Intercourse	No. of Cases
Total	1.3	13.3	23.4	30.1	32.3	67.7	2029*
Strata							
Metro Tirana	2.4	14.2	29.7	36.3	38.8	61.2	745
Other Urban	1.4	16.1	26.7	31.2	33.9	66.1	583
Other Rural	1.1	11.8	20.2	28.0	29.9	70.1	701
Residence							
Urban	1.7	15.6	28.2	33.2	35.8	64.2	1,192
Rural	1.1	11.8	20.4	28.2	30.2	69.8	837
Current Age							
15-19	1.2	(10.8)	(13.8)	NA	13.8	86.2	1,094
20-24	1.6	16.3	35.1	(50.2)	55.2	44.8	935
Marital Status at 1st Sex							
Premarital Sex	4.2	46.0	77.4	93.4	100.0	0.0	371
Marital Sex	4.1	37.5	68.6	93.0	100.0	0.0	376
Education Level							
Primary or Less	2.1	15.5	26.6	34.2	36.5	63.5	1,013
Secondary	0.3	11.2	18.7	22.9	24.2	75.8	778
Post-Secondary	0.2	4.7	18.2	29.5	34.6	65.4	238
Socioeconomic Index							
Low	1.4	10.7	19.2	25.1	27.2	72.8	649
Medium	1.4	14.4	25.1	32.3	34.5	65.5	1,106
High	0.8	19.2	34.0	41.5	44.2	55.8	274

* Excludes one case with missing information on age at first sexual intercourse

() Time exposed partially truncated because not all cases have exposure throughout the period of analysis NA. Not applicable

Table 13.4 B Percent of Men Aged 15-24 Who Had Their First Sexual Intercourse Before Selected Ages, by Various Characteristics Reproductive Health Survey: Albania 2002

		Age at I	First Sex	ual Interc	ourse	_ Never		
Characteristics	<15	<18	<20	<22	Has Had Intercourse	Had Intercourse	No. of Cases	
Total	1.0	11.9	20.4	27.0	29.3	70.7	589*	
Strata								
Metro Tirana	3.9	24.0	35.2	41.5	43.9	56.1	233	
Other Urban	0.0	9.4	23.5	28.3	29.5	70.5	177	
Other Rural	0.5	9.1	14.0	21.5	24.2	75.8	179	
Residence								
Urban	1.4	14.8	28.4	33.3	34.3	65.7	366	
Rural	0.8	9.8	14.8	22.6	25.7	74.3	223	
Age Group								
15-19	0.6	(4.5)	(4.8)	NA	4.8	95.2	401	
20-24	1.5	21.5	40.9	(56.1)	61.3	38.7	188	
Education Level								
Primary or Less	1.4	7.2	15.5	22.5	24.0	76.0	264	
Secondary	0.6	15.9	22.8	26.1	28.9	71.1	283	
Post-Secondary	0.7	23.1	46.8	73.1	78.6	21.4	42	
Socioeconomic Index								
Low	0.0	3.7	7.4	15.4	17.5	82.5	222	
Medium	1.1	17.6	29.5	35.6	38.5	61.5	282	
High	6.1	27.5	45.1	46.8	46.8	53.2	85	

* Excludes one case with missing information on age at first sexual intercourse

()Time exposed partially truncated because not all cases have exposure throughout the period of analysis

N.A. Not applicable

Table 13.5 A

Percent Distribution of Age Difference Between Partners at First Sexual Intercourse Among Sexually Experienced Young Women Aged 15–24 by Residence, Age at first Sex and Marital Status at First Sex Reproductive Health Survey: Albania, 2002

	Age Difference Between Partners at First Sexual Intercourse								
Characteristics	Partner Same Age or Younger	Partner 1-4 Years Older	Partner 5 or More Years Older	Total	No. of Cases				
Total	1.9	25.2	72.9	100.0	744*				
Residence									
Urban	1.3	26.9	71.9	100.0	459				
Rural	2.3	24.0	73.7	100.0	285				
Age 1st Sex									
< 18	0.6	12.7	86.7	100.0	302				
18-19	0.1	29.8	70.1	100.0	254				
20-24	5.6	38.6	55.7	100.0	188				
Marital Status at 1st Sex									
Premarital	1.2	26.1	72.7	100.0	368				
Marital	2.3	24.6	73.1	100.0	376				

* Excludes one case with missing information on partner's age and 3 cases in which the first sex was rape

Young Adult Sexual and Contraceptive Experience 248

Table 13.5 B

Percent Distribution of Age Difference Between Partners at First Sexual Intercourse Among Sexually Experienced Young Men Aged 15–24 by Residence, Age at first Sex and Marital Status at First Sex Reproductive Health Survey: Albania, 2002

	Age Diff	Age Difference Between Partners at First Sexual Intercourse					
Characteristics	Partner Younger	Partner Same Age	Partner Older	Don't Know	Total	No. of Cases	
Total	56.1	28.0	12.5	3.5	100.0	156*	
Residence							
Urban	53.0	32.3	11.7	3.0	100.0	106	
Rural	59.0	23.9	13.2	3.9	100.0	50	
Age 1st Sex							
< 18	31.4	43.9	19.8	4.8	100.0	73	
18-19	60.5	24.4	14.3	0.8	100.0	44	
20-24	85.1	10.0	0.8	4.2	100.0	39	
Marital Status at 1st Sex							
Premarital	52.6	30.6	13.3	3.4	100.0	134	
Marital	**	**	**	**	**	22	

* Excludes one cases with missing information on partner's age

**Percentages are not shown when base is less than 25 cases

Table 13.6 A

Percent Who Used Contraception at First Sexual Intercourse and Percent Distribution of Method Used Among Sexually Experienced Young Women Aged 15–24 Reproductive Health Survey: Albania, 2002

	Used a l	Nethod		Туре	of Method		
		No. of			Other		No. of
Characteristics	Percent	Cases	Withdrawal	Condom	Modern	Total	Cases
Total	81.3	744*	95.8	3.4	0.8	100.0	597*
Strata							
Metro Tirana	78.5	310	92.0	6.7	1.3	100.0	243
Other Urban	87.9	203	96.1	3.4	0.5	100.0	176
Other Rural	78.9	231	96.8	2.3	0.9	100.0	178
Residence							
Urban	85.7	459	94.5	4.7	0.8	100.0	384
Rural	78.1	285	96.8	2.3	0.9	100.0	213
Age 1st Sex							
<18	83.3	302	95.4	4.3	0.3	100.0	245
18-19	82.0	254	96.5	3.2	0.3	100.0	212
20-24	77.7	188	95.4	2.3	2.3	100.0	140
Marital Status at 1st Sex							
Premarital	91.3	368	92.8	6.4	0.8	100.0	326
Marital	74.1	376	98.4	0.7	0.9	100.0	271
Education Level							
Primary or Less	77.7	434	99.4	0.6	0.0	100.0	324
Secondary	89.2	217	91.5	8.0	0.5	100.0	193
Post-Secondary	87.3	93	81.8	9.9	8.3	100.0	80
Socioeconomic Index							
Low	72.8	222	100.0	0.0	0.0	100.0	157
Medium	84.7	405	94.3	4.6	1.1	100.0	335
High	91.3	117	91.7	6.3	2.0	100.0	105

* Excludes three women who reported first intercourse to have been rape.

	Used a	Method	Туре	of Method	ł		
		No. of			Other		No. of
Characteristics	Percent	Cases	Withdrawal	Condom	Modern	Total	Cases
Total	85.1	155*	56.2	43.1	0.7	100.0	124*
Strata							
Metro Tirana	69.3	81	35.2	64.8	0.0	100.0	58
Other Urban	80.2	39	46.7	50.7	2.6	100.0	32
Other Rural	97.8	35	69.9	30.1	0.0	100.0	34
Residence							
Urban	77.0	105	41.8	56.7	1.5	100.0	81
Rural	92.7	50	67.4	32.6	0.0	100.0	43
Age 1st Sex							
< 18	81.7	72	49.4	50.6	0.0	100.0	56
18-19	87.3	44	50.0	47.8	2.2	100.0	37
20-24	87.6	39	70.7	29.3	0.0	100.0	31
Marital Status at 1st Sex							
Premarital	88.5	133	50.2	49.0	0.7	100.0	111
Marital	**	22	**	**	**	**	13
Education Level							
Primary or Less	84.0	54	73.7	26.3	0.0	100.0	41
Secondary	83.9	72	43.1	55.3	1.6	100.0	59
Post-Secondary	**	**	**	**	**	**	24
Socioeconomic Index							
Low	86.5	38	75.6	24.4	0.0	100.0	29
Medium	84.3	83	48.8	50.1	1.1	100.0	65
High	85.9	34	46.8	53.2	0.0	100.0	30

Table 13.6 BPercent Who Used Contraception at First Sexual Intercourse AndPercent Distribution of Method Used Among Sexual Experienced Young Men Aged 15–24Reproductive Health Survey: Albania, 2002

* Excludes two cases with missing information

** Percentages are not shown when base is less than 25 cases

Table 13.7

Percent Distribution of Most Commonly Cited Reasons for Not Using Contraception at First Sexual Intercourse Among Sexually Experienced Young Women Aged 15-24 by Marital Status at First Sexual Intercourse Reproductive Health Survey: Albania, 2002

	Marital S	Status at First I	ntercourse
Main Reason for Not Using Contraception	Total	Married	Not Married
She Wanted to Get Pregnant	75.3	84.8	23.0
She Did Not Think About Using a Method	4.7	4.4	6.6
She Did Not Know About Contraception	4.2	4.2	4.6
Fear Of Side Effects	4.2	1.6	18.1
Partner Against	2.9	2.3	6.3
Sexual Intercourse Was Unexpected	3.3	0.8	17.0
Thought it Was Safe	1.7	0.0	11.1
Other Reasons	3.7	1.9	13.3
Total	100.0	100.0	100.0
No. of Cases	126	93	33

Table 13.8 Percent Distribution of Current Sexual Activity Status by Sex, Current Marital Status and Age Group Among Young Adults Aged 15–24 Reproductive Health Survey: Albania, 2002

			Women				
		Marita	al Status	Age Group			
	Total	Married	Not Married	15-19	20-24		
Never Had Intercourse	67.7	0.0	93.1	86.2	44.8		
Ever Had Intercourse	32.3	100.0	6.9	13.8	55.2		
Within the Last Month	20.2	65.2	3.2	7.7	35.7		
1-3 Months Ago	4.7	11.8	2.1	2.5	7.5		
Over 3 Months Ago But Within Last Year	1.2	2.6	0.7	0.6	1.9		
One Year or Longer	1.1	1.6	0.9	0.4	1.9		
Currently Pregnant or Postpartum	5.1	18.8	0.0	2.7	8.2		
Total	100.0	100.0	100.0	100.0	100.0		
No. of Cases	2,030	599	1,431	1,094	936		

	-	Marital Status		Age Group	
	Total	Married	Not Married	15-19	20-24
Never Had Intercourse	70.7	0.0	75.1	95.2	38.7
Ever Had Intercourse	29.3	100.0	24.9	4.8	61.3
Within the Last Month	21.8	92.3	17.4	2.8	46.6
1-3 Months Ago	0.5	2.7	0.3	0.6	0.4
Over 3 Months Ago But Within Last Year	4.3	0.0	4.5	0.9	8.7
One Year or Longer	2.5	0.0	2.6	0.5	5.0
Partner Currently Pregnant	0.3	5.0	0.0	0.0	0.7
Total	100.0	100.0	100.0	100.0	100.0
No. of Cases	590	33	557	401	189

Men

Table 13.9 A

Percent Who Used Contraception at Last Sexual Intercourse and Percent Distribution of Methods Used by Current Marital Status and Age Group Among Sexually Experienced Young Women Aged 15–24 Reproductive Health Survey: Albania, 2002

				Marital Status						
	Total			Married			Not Married			
	Total	15-19	20-24	Total	15-19	20-24	Total	15-19	20-24	
Percent Using	76.0	74.5	76.5	72.0	63.4	74.1	98.0	98.5	97.5	
Modern Methods	4.4	5.5	4.0	3.1	4.0	2.9	11.3	8.7	13.7	
Traditional Methods	71.6	69.0	72.5	68.9	59.4	71.1	86.7	89.8	83.8	
Methods Used										
Pill	0.5	0.3	0.6	0.2	0.4	0.1	2.1	0.0	4.1	
IUD	0.4	0.0	0.5	0.4	0.0	0.5	0.0	0.0	0.0	
Condom	3.1	4.4	2.6	2.0	2.7	1.8	8.9	8.2	9.6	
Tubal Ligation	0.2	0.0	0.3	0.3	0.0	0.4	0.0	0.0	0.0	
Injectables	0.2	0.8	0.1	0.2	0.9	0.1	0.2	0.5	0.0	
Withdrawal	71.6	69.0	72.5	68.9	59.4	71.1	86.7	89.8	83.8	
Not Using	24.0	25.5	23.5	28.0	36.6	25.9	2.0	1.5	2.5	
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	
No. of Cases	747*	164	583	599	97	502	148	67	81	

* Excludes one case with missing information

Table 13.9 BPercent Who Used Contraception at Last Sexual Intercourse and PercentDistribution of Methods Used by Current Marital Status and Age GroupAmong Sexually Experienced Young Men Aged 15–24Reproductive Health Survey: Albania, 2002

				Marital Status						
		Total			Married			Not Married		
	Total	15-19	20-24	Total	15-19	20-24	Total	15-19	20-24	
Percent Using	41.0	25.5	42.5	67.7	**	67.7	34.4	25.5	35.6	
Modern Methods	11.8	15.2	11.5	0.0	**	0.0	14.7	15.2	14.7	
Traditional Methods	29.1	10.3	31.1	67.7	**	67.7	19.7	10.3	20.9	
Methods Used										
Condom	11.8	15.2	11.5	0.0	**	0.0	14.7	15.2	14.7	
Periodic Abstinence (Rhythm)	2.1	0.0	2.3	5.1	**	5.1	1.4	0.0	1.6	
Withdrawal	27.0	10.3	28.7	62.6	**	62.6	18.3	10.3	19.4	
Not Using	59.0	74.5	57.5	32.3	**	32.3	65.6	74.5	64.4	
Total	100.0	100.0	100.0	100.0	**	100.0	100.0	100.0	100.0	
No. of Cases	157	31	126	32	0	32	125	31	94	

** Percentages are not shown when base is less than 25 cases

Table 13.10Percent Distribution of Main Reason Not Currently Using Contraceptionby Sex and Current Marital Status Among Sexually Active Young Adults Aged 15–24Reproductive Health Survey: Albania, 2002

	Total	Wo	men	_	Men		
Main Reason for Not Using Contraception		Married	Not Married	Total	Married	Not Married	
Not Sexually Active/No Partner	43.6	32.2	97.9	90.1	**	98.1	
Currently Pregnant	31.8	38.5	0.0	1.7	**	0.0	
Trying to Get Pregnant	17.6	21.3	0.0	4.8	**	0.0	
Postpartum/Breastfeeding	6.0	7.3	0.0	0.8	**	0.0	
Lovemaking Interrupted	0.5	0.6	0.0	0.0	**	0.0	
Believes She is Subfecund	0.3	0.1	1.0	0.0	**	0.0	
Did Not Think About Using Contraception	0.0	0.0	0.0	1.3	**	1.0	
Other	0.2	0.0	1.0	1.2	**	0.9	
Total	100.0	100.0	100.0	100.0	**	100.0	
No. of Cases	282	223	59	94	11	83	

**Percentages are not shown when base is less than 25 cases.

	S
l	
l	9
	6

Table 13.11 A Percent Distribution of Number of Sexual Partners in Last Three Months and in Lifetime by Current Marital Status and Education Level Among Sexually Experienced Young Women Aged 15–24 Reproductive Health Survey: Albania, 2002

		Marital	Status		Education Level			
Number of Sexual Partners	Total	Married	Not Married	Primary or Less	Secondary	Post- Secondary		
Last Three Months								
None	11.4	9.4	22.2	11.1	11.0	14.4		
One	88.2	90.2	77.3	88.4	88.6	85.6		
Two or More	0.4	0.4	0.5	0.5	0.3	0.0		
Lifetime								
One	97.3	99.5	85.4	98.9	95.5	89.9		
Two or More	2.7	0.5	14.6	1.0	4.5	10.1		
Total	100.0	100.0	100.0	100.0	100.0	100.0		
No. of Cases	748	599	149	434	218	96		

Table 13.11 B Percent Distribution of Number of Sexual Partners in Last Three Months And in Lifetime by Marital Status and Education Level Among Sexually Experienced Young Men Aged 15–24 Reproductive Health Survey: Albania, 2002

		Current Marital Status			Education Lev	vel
Number of Sexual Partners	Total	Married	Not Married	Primary or Less Secondary		Post- Secondary
Last Three Months						
None	33.8	5.1	40.8	25.9	40.4	37.1
One	61.1	92.4	53.4	69.3	54.6	56.6
Two or More	5.1	2.5	5.8	4.8	5.0	6.2
Lifetime						
One	28.3	48.5	23.4	37.6	21.5	22.3
Two or More	71.8	51.5	76.7	62.4	78.5	77.7
Total	100.0	100.0	100.0	100.0	100.0	100.0
No. of Cases	157	32	125	54	74	29

Table 13.12Percent Agreeing With Statements About Condoms and Condom Use, by Sex and
Condom Experience for Sexually Experienced Young Adults Aged 15-24 Years
Reproductive Health Survey: Albania, 2002

				Women								
		Ever Use	r (n=13	9)		Never Use	er (n=60)9)				
Statement	Agree	Disagree	Do Not Know	Refused to Answer	Agree	Disagree	Do Not Know	Refused to Answer				
Smart Idea to Use Condom With New Partner	94.6	2.5	2.9	0.0	58.0	10.4	30.7	0.8				
Women Should Ask Partner to Use a Condom	72.9	20.1	7.0	0.0	23.5	36.7	38.7	1.1				
Easy to Discuss Condom Use With Partner	68.7	20.9	9.6	0.8	33.4	17.4	47.4	1.9				
Condoms Diminish Sexual Enjoyment	57.2	39.9	2.9	0.0	16.2	6.0	75.8	2.1				
Condoms Not Necessary if you Know Partner	45.1	52.4	1.7	0.8	48.6	19.4	31.2	0.8				
Embarrassed to Ask for Condoms At Store	25.8	67.5	6.8	0.0	31.1	27.1	40.3	1.5				
People Who Use Condoms are Promiscuous	17.5	69.9	12.6	0.0	28.1	21.4	49.0	1.4				
The Same Condom Can be Used More Than Once	6.0	93.2	0.8	0.0	4.0	51.6	42.8	1.6				

	Ever User (n=117)					Never Us	er (n=4	0)
Statement	Agree	Disagree	Do Not Know	Refused to Answer	Agree	Disagree	Do Not Know	Refused to Answer
Smart Idea to Use Condom With New Partner	99.2	0.0	0.8	0.0	48.8	18.0	33.2	0.0
Women Should Ask Partner to Use a Condom	45.7	42.2	12.1	0.0	31.4	39.0	28.8	0.7
Easy to Discuss Condom Use With Partner	68.5	19.4	12.1	0.0	17.6	21.7	58.3	2.4
Condoms Diminish Sexual Enjoyment	77.0	21.4	1.6	0.0	43.4	3.1	51.9	1.7
Condoms Not Necessary if You Know Partner	36.3	62.2	1.5	0.0	57.1	17.9	25.0	0.0
Embarrassed to Ask for Condoms at Store	24.2	72.0	3.8	0.0	44.5	9.0	46.5	0.0
People Who Use Condoms are Promiscuous	33.9	53.0	13.1	0.0	49.3	6.8	43.9	0.0
The Same Condom Can be Used More Than Once	2.5	96.7	0.8	0.0	6.1	72.2	21.6	0.0

Men

Table 13.13 A Percent Agreeing With Hypothetical Responses to a Partner Requesting Condom Use During Sex, by Selected Characteristics, Among Sexually Experienced Young Women 15-24 Years of Age Reproductive Health Survey: Albania, 2002

Characteristics	Feels Safe From Getting Pregnant	Feels Safe From Getting STD/HIV/ AIDS	Would be Insulted or Angry	Would Feel Like She Had Done Something Wrong	Suspicious That He May "Sleep Around"	No. of Cases
Total	66.7	61.6	30.1	22.6	18.2	748
Residence						
Urban	73.2	68.4	27.1	19.0	18.2	463
Rural	61.8	56.6	32.3	25.3	18.1	285
Age Group						
15-19	65.7	61.0	31.1	23.2	19.5	165
20-24	66.9	61.7	29.7	22.4	17.8	583
Education Level						
Primary or Less	58.5	54.0	32.9	24.8	18.8	434
Secondary	81.7	76.1	26.5	23.2	18.8	218
Post-Secondary	88.7	80.0	17.6	2.4	10.6	96
Know How Condom is Used						
Yes	78.4	69.8	31.5	25.9	21.0	477
No	49.1	49.2	27.9	17.7	13.9	271
Condom Use						
Ever User	87.2	83.1	8.9	10.1	11.6	139
Never User	63.8	58.5	33.0	24.4	19.1	609

Table 13.13 B Percent Agreeing With Hypothetical Responses to a Partner Requesting Condom Use During Sex, by Selected Characteristics, Among Sexually Experienced Young Men 15-24 Years of Age Reproductive Health Survey: Albania, 2002

Characteristics	Feels Safe From Getting Her Pregnant	Feels Safe From Getting STD/ HIV/AIDS	Would be Insulted or Angry	Would Feel Like He Had Done Something Wrong	Suspicious That She May "Sleep Around"	No. of Cases
Total	83.6	74.4	20.6	19.0	32.8	157
Residence						
Urban	81.0	75.6	10.8	7.1	27.5	107
Rural	86.2	73.0	30.0	30.4	37.7	50
Age Group						
15-19	79.6	75.2	7.4	5.9	17.9	31
20-24	84.1	74.4	21.9	20.4	34.3	126
Education Level						
Primary or Less	83.8	64.3	31.4	30.2	39.0	54
Secondary	84.0	79.2	15.7	13.8	31.3	74
Post-Secondary	82.4	88.6	5.1	3.5	20.3	29
Know How Condo	om is Used					
Yes	83.9	77.9	20.6	18.8	32.9	144
No	**	**	**	**	**	13
Condom Use						
Ever User	89.3	79.7	9.3	11.2	27.9	117
Never User	72.2	63.8	43.7	35.1	42.8	40

 ** Percentages are not shown when base is less than 25 cases

Table 13.14 Percent Who Have Ever Talked to a Partner About Using Condoms, by Sex and Selected Characteristics for Sexually Experienced Young Adults 15-24 Years of Age Reproductive Health Survey: Albania, 2002

		Women						
Characteristics		No. of Cases						
Total	24.4	748	57.7	156*				
Residence								
Urban	32.1	463	79.2	106				
Rural	18.8	285	37.4	50				
Age Group								
15-19	26.0	165	51.6	31				
20-24	23.9	583	58.3	125				
Education Level								
Primary or Less	15.4	434	39.6	53				
Secondary	39.9	218	68.7	74				
Post-Secondary	52.1	96	75.4	29				
Condom Use								
Ever User	89.0	139	84.1	117				
Never User	15.4	609	3.0	39				
Withdrawal Use								
Ever User	24.0	698	55.6	139				
Never User	31.0	50	**	17				

* Excludes one case with missing information

** Percentages are not shown when base is less than 25 cases

CHAPTER 14

KNOWLEDGE AND EXPERIENCE OF SEXUALLY TRANSMITTED INFECTIONS AND KNOWLEDGE OF HIV/AIDS TRANSMISSION AND PREVENTION

Countries of South Eastern Europe have not been challenged by a substantial HIV/AIDS epidemic, or other sexually transmitted infections (STI), compared with countries of Eastern Europe and Central Asia (EE/CA) which continue to have expanding epidemics. It appears that infection rates are growing faster in EE/ CA than in any other region of the world and injecting drug use has been identified as the main force driving the increase in infection rates. Also, sexual transmission of HIV infection is increasing, especially between injecting drug users and their partners (UNAIDS, 2004). The number of infected women has been increasing from one-in-four cases in 2001 to one-in-three in 2003 and they account for an increasing share of newly diagnosed cases (UNAIDS and WHO 2003).

As of the end of 2004, Albania is still considered a low HIV prevalence country. Since the first detected HIV-infected case, found in 1993 through routine blood bank HIV screening, the number of identified cases has increased to 132, of which 88 cases were reported in the last three years. It has been estimated that HIV prevalence in Albania does not exceed 0.1% which follows the low prevalence pattern of South Eastern European Countries (MoH, 2003; IPH,2004) The majority (about 80%) of HIV infections and AIDS cases are believed to have acquired their HIV infection outside of Albania and over 90% of them were sexually transmitted. However, during the last two years there has been an increase of cases infected within the country.

Most of the cases belong to the 30-40 year old age group but a trend toward younger ages has been observed during the last three years with the number of cases in the age group of 20-30 year olds increasing. Also, a feminization of the epidemic has been observed since 2000 with the number of women infected with HIV/AIDS increasing (IPH, 2003; IPH, 2004). Sexual HIV transmission is predominant among married women in Albania, and mother to child transmission has been identified during the last two years resulting in the infection being present among children (IPH, 2004).

During the decades of Albania's isolation from the 1960s until the early 1990s, STIs such as syphilis or gonorrhea were virtually eliminated. By the end of 2003, 138 cases of syphilis were identified in Albania with women accounting for 48% of the cases; 45% had primary syphilis (IPH, 2004). A concomitant HIV infection was observed in 4.5% of the cases in the last three years (IPH, unpublished data).

There are no accurate data available on other sexually transmitted diseases in Albania. But, an increase of gonorrhea cases has been detected through routine hospital based surveillance and a laboratory surveillance of gonorrhea is in the process of being established. Recent studies show the presence of herpes viruses among women of child bearing age. A study performed by the STI lab in the Institute of Public Health documented vaginal discharge and the presence of Chlamydia trachomatis in 21% of 527 women studied in Tirana (IPH, 2003). There is a concern that a hidden epidemic might occur among drug users, men having sex with men and female sex workers working within the country.

A National AIDS Program (NAP) and plans for prevention/control of HIV/AIDS/STI were developed during the late 1980s with the help of the Global Programme on AIDS of the World Health Organization (GPA/ WHO). During the long period of isolation, and as a result of a rigid prevention/control measures that included police investigations and follow-up, as well as enforced screening and treatment of STIs, there was a virtual elimination of STIs and visible commercial sex activities. With the re-opening of Albania to population movements both within and out of the country in the early 1990s, STI cases began to reappear and international agencies also gradually re-entered to provide economic and technical assistance to social and health programs in Albania. A law on HIV/AIDS prevention was passed in the parliament in 2002 and the national strategy with the objective to keep Albania a low prevalence country was approved by the government by the end of 2003. This strategy underlined the importance of prevention activities and especially information and education programs (MoH, 2003).

Often, prevention programs do not reach people who most need them, especially women and young people. Increased access to treatment will offer better opportunities to strengthen prevention and volunteer testing. The current voluntary testing and counseling services are poor and the uptake of existing service is still low due to stigma and fear of discrimination (MoH, 2003; UNAIDS, 2004).

HIV-risk behavior surveillance data are needed by the HIV/AIDS/STI prevention and control program, including: specific patterns and prevalence of persons who routinely have unprotected sex with multiple and concurrent sex partners, and specific patterns and prevalence of persons who routinely share their injecting equipment with other IDU. Also, among high-risk groups, it will be important to evaluate how many sex partners persons have had during the past month and/or year, and for each of these sex encounters, if a condom was used. There are major concerns that HIV risk behaviors have been steadily increasing in the country without a substantial increase of knowledge and education that leads to disease prevention.

As a low prevalence country it is important to be aware of the level of correct knowledge about transmission and prevention among different population groups, especially among women and men of reproductive age, and identify factors that influence the misconceptions related to HIV transmission or other STIs. Detailed information about the level of awareness, source of information, and knowledge related to HIV/ AIDS and also exposure to HIV/AIDS/STI testing and perceived risks for HIV/AIDS/ STI were collected through a module of the questionnaire devoted to this topic


Awareness of STIs and Knowledge of Symptoms Related to STIs

When asked about general awareness of sexually transmitted diseases (STIs), Albanian women showed a very low level of knowledge (Table 14.1A). Overall, HIV/ AIDS is the only STI that most of the women (96%) have heard of. Only 57% of women had heard of Yeast infection, and for other STIs the level of awareness was even lower (4%-14%).

With the exception of HIV and Yeast infection, overall awareness was higher for women living in Tirana, and other urban areas. Rural women were in general less aware of STIs. Marital status does not

seem to be related to the level of awareness of HIV as an STI or other STIs with the exception of Yeast infection. Yeast infection is the only disease that never married women are least aware of (43%) compared to previously married (70%) and currently married women (63%). In contrast, for all other diseases, never married women have higher awareness or about the same level of awareness when compared with currently or previously married women. Although in general, there is a tendency for women with fewer children to have slightly greater awareness of STIs, the number of living children does not show a clear pattern of association between level of awareness and STIs. In contrast, the level of education of women is clearly associated with the level of awareness. Higher educated women are more likely to have heard of specific STIs. HIV awareness is high, independent of the number of lifetime partners, but women with two or more lifetime partners were consistently more likely to have heard of other STIs. However with only 113 women in this category, 95% confidence limits range from 6 to 11 percentage points.

Overall, the percentage of men that have heard of HIV is very similar to that of women (96%). Regarding all other STIs, similar to women, the general level of awareness of Albanian men is low (Figure 14.1). The biggest differences between men and women can be found in the percentages of those who have heard of syphilis (22% of men vs. 14% of women) and gonorrhea (14% vs. 7%), and the 12% of men who have heard of Yeast infection (Table 14.1B), compared with 57% of women. Knowledge of remaining STIs was very low among men, around 4%. By residence, rural men were less likely than urban men, including Tirana residents, to have heard of STIs. Similarly to what was found for Albanian women, there is no clear pattern of association between age and awareness of specific STIs. Despite this fact, its clear from Table 14.1B that men



between the ages of 45-49 are somewhat less likely to be aware of HIV/AIDS (87%), compared with all other age groups where awareness is greater than 93%. Because the number of men previously married in the sample is very low, marital status comparison will be made only between never married and currently married men in this chapter. Awareness among never married and currently married men for HIV/AIDS and all other STIs are not statistically different. Among men, there is no clear relationship between HIV knowledge and number of living children. Not surprisingly, knowledge of STIs increases with education. However, there is no significant difference for HIV awareness by educational attainment as all groups have at least 93% awareness. With exception of HIV and yeast infection, men with 2 or more lifetime partners are more likely to be aware of STIs than their counterparts with fewer lifetime partners.

Table 14.2A and B show knowledge of symptoms associated with STIs other than HIV for women and men, respectively. Among women, 59% have heard of at least one STI other than HIV/AIDS, compared to only 27% of the men. Among all women 15-44 years of age, 41% are not aware of any STIs, and 19% do not know any symptoms that can be associated with STIs. Only 12% of the women knew one symptom associated with STIs, and 28% recognized two or more symptoms. According geographic to strata. knowledge of STI symptoms is greater in Tirana where 47% of the residents know at least one symptom related to an STI. Younger women (15-19) are less aware of any STI symptom than others. By age 25-29 women showed more awareness of STI related symptoms, from 43% to 48% of women aged 25-44 knowing at least one symptom.

Previously married and currently married women are more aware of STI symptoms than never married women (45%, 33%) and 19%, respectively, know at least two symptoms). Level of education and number of lifetime partners are directly related to knowledge of STI related symptoms. Women with higher education are twice as likely as their primary or less educated counterparts to know at least one STI related symptom. Exposure to multiple partners also increases the likelihood of knowing at least one STI related symptom. Thus, more than half of the women with two or more lifetime partners at the time of the survey know at least one symptom related to STIs compared to 45% of women with only one partner and 27% of the women without sexual experience.

Among men, knowledge of symptoms of STIs is even lower than among women. The vast majority of men (74%) are not aware of any STI, and only 20% know at least one STI related symptom (Table 14.2B). As expected, rural residents are less aware of STIs (84%) compared with Tirana (64%) and residents of other urban areas (63%). No clear relationship can be found between STI awareness and age or marital status. For all age groups the vast majority of men are not aware of any STI. Similar to women, education seems to be an important determinant of STI awareness. Only 29% of the men with post-secondary schooling were not aware of any STI compared to 68% of men with secondary education and 87% with primary or less education. While having either zero or one partner seems to have the same impact on STI awareness (only 18% and 17%, respectively, have heard of at least one STI), those with two or more partners are more likely to be aware of STIs (37%). Results for men who don't remember how many partners they have

had are similar to those reporting two or more partners.

The prevalence rate of syphilis is very similar to that of HIV, and in a blind study conducted by the IPH, 4.5 % of the patients with syphilis tested positive for HIV and 11% of persons living with HIV/AIDS (PLWHA) were positive for syphilis. Also, Syphilis or other STIs were seen in 20% of HIV/AIDS cases (IPH, unpublished data). This fact shows the important role that STIs play in the transmission of HIV/AIDS and the need to increase STI awareness among women and men in Albania through culturally appropriate programs.

Most Important Source of Information and Messages About STIs

Tables 14.3A and B show the most important source of information about STIs for women and men, respectively, for those who have heard of at least one STI. Overall, among women, mass media is the most important source of information (84%). Women of all ages reported mass media as their most important source of information. However, among adolescents (15-19), 20% reported having received STI information from schools. By marital status, mass media is also the most important source of information for STIs, but 15% of never married women reported receiving information from schools. These are probably younger women still in school. Similar results are shown according to lifetime partners, with 16% of women with zero partners (probably the youngest women and still in school) receiving STI information via schools.

Mass media is also the most important source of information on STIs among men (66%), but significantly lower (p<0.01)

than for women (84%). The second most important source of information on STIs for men is the doctor (10%), followed by books (8%), and friends (7%). By strata, STI information via mass media ranges from 63% to 70%. Similarly to the finding for women, an important percentage of younger men (15-19) get their STI information from school (23%). When comparing age groups among men, major differences are found in the second most important source of STI information. Whereas for the 20-24 and 25-29 year olds the second most important source is books (14% and 11%, respectively), for 30-34 it is friends (12%), for 35-39 it is the doctor (15%), for 40-44 it is friends (15%), and for 45-49 it is again the doctor (17%). Married men reported the second most important source to be the doctor (12%), while never married men reported books (13%). Regardless of their level of education, men in Albania reported mass media as their most important source of STI information. However, the second most important source differs by education. Primary and secondary educated men have as their second source of information the doctor (16% and 10%, respectively), while post-secondary educated have books (23%). According to number of lifetime partners, men with 2 or more partners reported their second source of information to be the doctor (16%), compared to men without sexual experience that get their information (second most important) from school (20%).


Respondents were asked if, in the past six months, they have seen or heard any public announcements or messages on television or radio about HIV/AIDS or other STIs (Tables 14.4A and B). Fifty seven percent of women and 63 percent of men said that they saw an announcement or message about HIV/AIDS on television in the past six months. Only 6% and 23%, respectively, heard a message about HIV/AIDS on the radio. Almost half of women (40%) and approximately one third of men (34%) did not see or hear a message on television nor radio (Figure 14.2). For women, seeing messages on TV increased as educational attainment increased and was higher for women living in Tirana or other urban areas. Otherwise, there is no significant variation associated with the characteristics shown in the Table. For men, reception of messages about HIV/AIDS on TV was higher for men living in Tirana, those with more than a primary education and men with fewer than two or more lifetime partners. Less than one percent of both women and men heard messages on the radio only.

Fewer women and men saw or heard messages about other STIs. Only 8% of

women saw messages on TV and less than 2% heard messages on the radio. For men, the corresponding figures are 19% and 13%, respectively. Ninety-two percent of women and 81% of men did not see or hear any message or did not remember any message about STIs on either TV or the radio. Those who did were more likely to live in Tirana and/or have a post-secondary education.

Media, and prevention and education messages released by them, are potentially important tools in educating the population about HIV/AIDS prevention. A need for training media specialists on HIV/AIDS/STI prevention messages is very much needed as well as the integration of primary care services into HIV/AIDS/STI prevention initiatives.

Self-Reported Diagnosis, Testing and Treatment of STIs



Tables 14.5A and B present the percentage of women and men, respectively, that said they have been diagnosed with any STI during their lifetime, by selected characteristics. Overall, the most prevalent reproductive tract infection was yeast infection, but much higher for women than for men (8% and 1% respectively).

Among women, diagnosed STIs such as syphilis, gonorrhea, genital herpes, trichomoniasis, genital warts and chlamydia were extremely low (less than 1%). A diagnosis of Yeast infection (YI) was reported to a greater extent by women living in Tirana (12%), women between the ages of 25 and 39 (11% - 13%), previously married women (15%), and women with one or more living children (11% - 13%).Women with children were between 4 and 5 times more likely to have been diagnosed with a YI. Diagnosed YI also increases with number of lifetime partners.

Among sexually experienced women (Table14.6A), overall awareness of HIV/ AIDS was high (96%) followed by awareness of YI (63%) and syphilis (13%). Levels of awareness of all other STIs were less than 10%. The most common reproductive tract infection tested was YI (14%); of those tested, 82% were confirmed (11.5%) and all diagnosed cases were treated. Among sexually experienced men (Table 14.6B), 96% were aware of any HIV/AIDS, followed by syphilis (24%), and gonorrhea (16%). Awareness of other STIs and yeast infection was less than 15%. Less than 1% of men were ever tested for HIV. The most commonly tested STI among men was for genital warts (4%); in all cases, the diagnosis was not confirmed.

Policies to establish the syndromic approach for surveillance or case management of persons with STIs in Albania, according to the National HIV/AIDS Prevention Strategy (MoH, 2003), should take into account the survey data of self reported diagnosis and testing.

Perceived Risk of STIs

Tables 14.7A and B show the perceived risk of acquiring an STI among men and women, respectively, who have heard of at least one STI other than AIDS. When asked about the potential risk of acquiring an STI, the vast majority of Albanian women consider themselves at no risk (80%) or at low risk (14%). Only 2% consider themselves to be at high risk or medium risk. Socio-demographic characteristics did not strongly influence the perceived risk of acquiring an STI. However, not surprisingly, perceived risk increased with the number of lifetime partners. About 9% of women with 2 or more lifetime partners perceived themselves at high or medium risk of acquiring STIs with another 24% saying that they had a low risk.

Male responses regarding perceived risk are even more troublesome (Table 14.7B). The vast majority of males (97%) said they have no risk or do not know how to assess their risk of acquiring an STI. The low rate of perceived risks of contracting STIs among women and men in Albania makes them more vulnerable in a situation where rates for both HIV/AIDS and other STIs are increasing and the main route of transmission in the country is sexual transmission.

Knowledge of HIV/AIDS

Even though the vast majority of women have heard of HIV/AIDS (96%), as shown in Table 14.1A, only 56% believe that HIV/AIDS can be asymptomatic, showing significant lack of understanding of the disease (Table 14.8A). Furthermore, only 17% of reproductive age women know where to get an HIV test and less than 1% have ever been tested. Not surprisingly, rural women are less likely to know that HIV can be asymptomatic (49%), compared to their urban counterparts (63% for both Tirana and other urban). Tirana women are more likely to know where an HIV test can be obtained (32%). Across age groups, belief that HIV can be asymptomatic ranges from 50% to 61%, and 15% - 19% know where to get an HIV test. Among women, the belief that HIV can be asymptomatic decreases with number of living children (data not shown). Women with more children (4 or more) were less likely to understand the disease, and also less likely to know where to be tested. Education is highly associated with awareness, overall understanding of the disease and testing locations. More educated women had universal awareness of HIV; 85% of women with post-secondary education know that HIV can be asymptomatic; and almost half of them know where to be tested. Regarding the number of lifetime partners, sample size is very low but, women with 2 or more lifetime partners better understand the disease and are more likely to know where to be tested.

Although HIV awareness is almost universal among men, rural men are less likely to have heard of HIV (Table 14.8B), less likely to know that HIV can be asymptomatic and also to know where HIV tests are provided. Tirana residents were most likely to have been tested for HIV (4%). Slightly over half of men 20-34 years of age (52%-53%) know that HIV infection can be asymptomatic. Overall, only 45% of men know that HIV can be asymptomatic. By age group, only from 21% (45-49) to 44% (25-29) know where HIV tests are provided. Men aged 20-24 were the ones with highest percentage of tested men (3%). Married men are less likely to know that HIV can be asymptomatic (40% vs. 52%). However, they are equally knowledgeable of where to be tested (33%).

Similar to women, education is associated with increased knowledge of HIV infection, where to be tested, and having sought out the HIV test. Men with two or more lifetime partners are more likely to know that HIV infection can be asymptomatic, and know where to be tested than men with no sexual experience (56% vs. 43% and 41% vs. 25% respectively).

Evidence based and culturally appropriate programs that increase the level of knowledge about HIV, as well as culturally appropriate information, education and communication programs for women and men living in rural areas, should be designed and implemented to further prevent the spread of HIV/AIDS among the general population.

Knowledge of HIV/AIDS Transmission

To assess the knowledge of HIV/AIDS transmission, respondents were asked to agree or disagree with statements on how the AIDS virus can be transmitted. Not knowing about a particular mode of transmission or a "no" response to a documented transmission mechanism was considered as lack of knowledge about a particular transmission mechanism. The least known transmission route was MTCT/breastfeeding among both women (24%) and men (55%) of reproductive age (Tables 14.9A and B, respectively). Among women, MTCT/during pregnancy

or delivery (18%), unprotected homosexual intercourse (16%), blood transfusion and use of non-sterile needles (both with 10%) followed. Rural women, women previously married, less educated women and women without children (not shown in Table) were overall less likely to know important ways of HIV/AIDS transmission.

Among men (Table 14.9B), transmission through MTCT delivery/pregnancy (37%), unprotected homosexual intercourse (36%), and unprotected heterosexual intercourse (17%) followed as the least known methods of HIV transmission. Similar to women, rural men, less educated men and men with less than two lifetime partners were less likely to know important ways of HIV transmission. However, contrary to women, the level of knowledge of HIV transmission was similar for currently married and never married men.

Respondents were classified as correctly rejecting a misconception if they answered "no" to an incorrect mechanism of transmission. The percentages of women who correctly rejected misconceptions about HIV transmission are presented in Table 14.10A. None of the misconceptions presented have been scientifically documented as common modes of HIV transmission. Overall, most of the misconceptions were rejected by more than a third of the women. Exceptions are mosquito bites (25%) and dental/surgical procedures (4%). However, even though most of the misconceptions were rejected by more than a third of the women, the results show that from one-third to two-thirds do have misconceptions about HIV transmission.

Albanian men have overall fewer misconceptions than Albanian women (Table 14.10B). Men were more likely than women to correctly reject all the misconceptions about HIV transmission presented in the Table. Overall, misconceptions about HIV transmission are higher among men with primary or less education. Similar to women, the most common misconception was HIV transmission trough dental or surgical procedures. This misconception may be associated with the fact that receiving infected blood products or medical treatment with infected instruments can transmit HIV. Other surveys in Eastern European countries have shown similar results about this particular misconception (Serbanescu et al, 1998, 2001, and 2003).

Tables 14.11A and B (women and men, respectively) display the **UNAIDS** indicator 2 by respondents' background characteristics. UNAIDS indicator 2 represents: the percentage of respondents with correct knowledge that HIV can be asymptomatic, and is not spread by two most common misconceptions in the specific country; in this case, HIV is not spread by mosquito bites or through medical procedures. This indicator gives the overall knowledge about HIV/AIDS in the population. It is estimated using all respondents, not just those who have heard of HIV/AIDS. As mentioned before, 56% of Albanian women of reproductive age know that HIV can be asymptomatic, only 25% know that HIV is not spread by mosquito bites, and only 4% know that HIV is not spread by dental or surgical procedures. Overall, less than 1% of Albanian women correctly answered all three questions.

Among men, the level of UNAIDS knowledge indicator 2 is also extremely low (1%). These low levels of knowledge for indicator 2 for men and women are a reflection of the high level of misconceptions related to transmission mechanisms. To increase the overall level of knowledge,

AIDS education campaigns will have to focus on transmission misconceptions, especially those pertaining to the spread of disease through dental or surgical procedures and mosquito bites.

Knowledge of HIV/AIDS Prevention

A two-part question was asked to assess knowledge of HIV prevention. Respondents were asked about what a person can do to reduce the risk of HIV infection. Individuals who spontaneously answered a correct prevention mechanism were coded "yes (spontaneous)". In part two, the respondents were asked about the means of transmission that they did not answer spontaneously ("probed").

More than a third of the women (36%)spontaneously mentioned "stay faithful to one partner" as a possible means of HIV prevention (Table 14.12A). When probed, an additional one-half (54%) of the women agreed with faithfulness as a means of HIV prevention. Condom use was spontaneously mentioned as preventive measure against HIV infection by 31% of the women, but increased to 75% when probed (44%). Somewhat surprising was the fact that 21% of the women did not mention condom use at all. The characteristics of these women are shown in Table 14.13A. This is important as low condom use is observed in the country. Less than one-fourth of Albanian women spontaneously mentioned most of the other possible means of HIV transmission. However, percentages these increase significantly when probed (i.e avoid sex with prostitutes 15% vs. 76%; limit number of sexual partners 12% vs.78%; avoid sex with bisexual 10% vs. 79%). Nearly half of the women did not mention "abstinence" as a means to prevent HIV (49%), and even when probed only 37% of the women happen to have mentioned it in addition to the 10% who spontaneously mentioned "abstinence".

Among men (Table 14.12B), the most frequent spontaneous report of means to prevent HIV was also condom use (38%). When probed, another 39% of the men mentioned condom use and 19% did not mention condom use. Slightly over one-third of men (35%) spontaneously mentioned "avoid sex with prostitutes" and another 57% agreed when probed. Abstinence was spontaneously mentioned by only 5% of the men as a mean to prevent HIV, and even though a little more than half (56%) mentioned it when probed, 35% of the men did not agree with it at all. Possible means of HIV transmission such as partner limitation, avoiding sharing sharp objects, avoid sex with bisexuals, donate blood and HIV testing were less frequently mentioned spontaneously (less than 20%), but when probed the vast majority of men agreed with these means as ways to prevent HIV transmission.

Looking at background characteristics of women who know possible means of HIV prevention(spontaneously and after probing), rural women, and women with primary or less education were overall less likely to know the possible means of HIV prevention (Table 14.13A). Among women, the least known way of preventing HIV is abstinence (46%). Surprisingly, abstinence as a way of preventing HIV infection is not significantly associated with residence, has no clear pattern of association with age, and women with post-secondary education are less likely to agree with it. Moreover, currently married women (45%) and women with one lifetime partner (45%) were somewhat less likely to know about abstinence as a way to prevent HIV.

Among men, abstinence is also the least known way considered to prevent HIV (Table 14.13B), but at a much higher rate than women (61% vs. 46% respectively). No clear pattern between can be identified between respondents characteristics and knowledge of abstinence to prevent HIV transmission. Never married and currently married men showed similar levels of knowledge on abstinence and HIV prevention. Men with two or more lifetime partners were more likely to mention "avoid risky sex" and "use condoms".

Tables 14.14A and B (women and men, respectively) present results from the UNAIDS knowledge indicator 1 and the variables that form this indicator: percent of women who believe HIV can be prevented by being monogamous, limiting the number of sexual partners, and using condoms. Around 90% of the women mentioned all three ways of preventing HIV. Below the average knowledge for this indicator 1 were rural women (87%); women 15-19 years old (87%); never married women (88%), women with primary or less education (84%), and women without a partner (87%).

UNAIDS knowledge indicator 1 was much lower for men (69%). According to background characteristics, rural men (63%), men aged 35 and older (58% to 62%), currently married men (67%), men with primary or less education (62%), and men with one lifetime partner (58%) were less likely to be classified as "in compliance" with the UNAIDS knowledge indicator 1.

The data presented here show again the need to increase the level of education toward appropriate prevention measures among men and especially among rural men and women, young women and less educated men and women, taking into account the high rate of migration of these groups and their potential risk to contract HIV/AIDS.

Beliefs About the Risk of HIV/AIDS and Self-Perceived Risk of HIV/AIDS

Respondents were asked to rate their self-

perceived risk of contracting HIV/AIDS. We assume that their responses are based on their acquired knowledge about HIV transmission, prevention, and risk factors. Overall, only 3% of Albanian women did not know how to assess their risk, and 4%, who never heard of HIV/AIDS, were not asked this question (Table 14.15A). The vast majority of women perceive themselves at no risk (79%) or little risk (11%). Moderate and great risk ("high risk") was reported by only 3% of the women. Differences by residence, age, and marital status are negligible. However, women with postsecondary education, and women with two or more lifetime partners were slightly more likely to assess themselves at moderate or great risk than their counterparts. It is important to note that these two groups were also the ones that consistently showed more understanding of HIV/AIDS, and therefore their assessment might be more accurate than the other groups.

Among men (Table 14.15B), even though the vast majority knew how to assess their risk of HIV infection, they were almost two and a half times more likely than women not to know how to assess their risk. Similar to women, most of the men considered themselves at no (79%) or little risk of HIV infection (8%). Of the 2% of the men who considered themselves at moderate or great risk of HIV infection, the only group to reach 5% was men between the ages of 20-24 (5%).

Tables 14.16A and B (women and men, respectively)presentthepercent distribution of the risk factors for contracting HIV among those who have heard of the disease and think that they have any risk of contracting HIV. Among women, medical/ dental treatment was universally reported (91%) as the main possible risk factor of contracting HIV. Men on the other hand, even though medical /dental risk was also very high (54%), also mentioned unprotected sex with a casual partner (16%), and use of IV drugs/shared needles and "many sexual partners/ trade sex for money" (both with 7% each) as other risk factors for their contracting HIV.

These results are somewhat troublesome. It is well established that HIV transmission is primarily transmitted trough individual risk behaviors. The popular notion that medical/dental treatment is a significant mechanism of HIV transmission can have negative implications for health care service utilization. Education programs should address this misconception and emphasize the distinction between HIV transmission through contaminated blood products and the very low probability of transmission through medical/dental treatment or procedures.

The 79% of women and men that assessed themselves at no risk of contracting HIV (of those who have heard of HIV) were asked about the main factor that protects them against HIV infection. Results are presented in Tables 14.17A for women and 14.17B for men. Among women, the most important protective factor was monogamy (41%), followed by abstaining from sex (32%), and trust in the partner (23%). The same pattern was seen for men with monogamy (48%), abstaining from sex (28%) and trust in partner (13%), mentioned in that order. Men mentioned condom use at a higher level (8.4%) compared to women (0.8%) but, still, 8% is a very low percentage.

In conclusion, this study shows the gaps that exist in knowledge and awareness about HIV/AIDS/STI transmission and prevention among different groups of the reproductive age population in Albania. Other studies also have shown the lack of accurate knowledge and awareness on HIV/AIDS and other STIs among young mobile populations or other high risk groups such as drug users or female sex workers (UNICEF, 2002). It is important to develop and build culturally appropriate information, education and communication programs for those living in rural areas and the less educated. Special attention must be given to married and young women in rural areas where the lack of knowledge increases their vulnerability toward HIV/AIDS. Appropriate integrated information, education and communication interventions might prevent further spread of the infection among these groups.

ai l	
<u> </u>	
•	
Š	
6	
5	
- 1	
9	
5	
S	
2	
Į.	
5	
4	
0	
9	
60	
U.	
•	
9	
U	
(1) (4)	
S S	
TIS a	
STIS a	
STIS a	
j	
JO I	
OF	
ce of 2	
nce of S	
ence of S	
ence of S	
erience of S	
rience of S	
erience of S	
erience of S	
xperience of 3	
d Experience of S	
nd Experience of S	
and Experience of S	
and Experience of S	
se and Experience of S	
e and Experience of S	
edge and Experience of S	
ledge and Experience of S	
vledge and Experience of S	
ledge and Experience of S	
owledge and Experience of S	
owledge and Experience of S	

Table 14.1 A
Percentage of Women Aged 15-44 Who Have Heard of Specified
Sexually Transmitted Infections by Selected Characteristics
Reproductive Health Survey: Albania, 2002

Characteristic	HIV/ AIDS	Yeast Infection	Syphilis	Gonorhea	Genital Hepes	Trichomonas	Genital Warts	Chlamydia	No. of Cases
Total	95.7	56.6	13.6	6.8	5.5	4.3	4.3	3.9	5,697
Strata									
Tirana	97.1	61.8	26.5	14.4	12.4	9.2	8.7	8.3	2,108
Other Urban	98.1	65.2	21.7	11.6	8.3	6.4	6.8	6.7	1,816
Rural	94.0	50.3	5.5	2.0	2.0	1.7	1.7	1.1	1,773
Age Group									
15–19	95.7	36.8	13.0	6.7	3.9	2.8	1.8	3.3	1,094
20–24	97.1	55.5	13.8	8.6	6.4	4.4	5.8	5.1	936
25–29	97.2	63.6	12.8	6.1	4.4	3.4	3.5	2.9	946
30–34	94.1	62.4	16.0	7.9	7.3	6.0	6.3	4.9	1,067
35–39	96.1	65.3	13.3	5.6	5.5	4.7	4.1	3.6	958
40–44	93.9	62.9	12.8	6.1	6.1	5.1	5.3	3.7	696
Marital Status									
Currently Married	95.5	62.8	12.5	5.8	5.2	4.4	4.5	3.5	3,965
Previously Married	96.0	70.2	13.9	4.7	5.3	2.7	2.6	2.7	88
Never Married	96.1	43.2	15.7	9.1	6.2	4.2	4.1	4.7	1,644
No. of Living Children									
0	96.4	45.3	16.2	8.8	6.0	4.3	4.6	4.8	1,943
1	96.0	62.2	13.8	7.1	5.7	4.4	3.7	4.0	828
2	96.9	66.2	15.0	7.5	6.6	5.8	5.6	4.5	1,840
3	94.9	64.7	9.2	3.2	3.4	3.1	3.0	2.0	795
4+	89.1	52.7	4.1	0.9	3.0	1.3	1.9	0.6	291
Education									
Primary or Less	92.8	47.9	3.6	1.0	1.4	0.8	0.8	0.7	2,519
Secondary	98.9	62.7	19.0	9.0	7.0	5.6	5.5	4.8	2,483
Post-Secondary	99.8	85.2	3.6	35.9	26.4	21.6	21.9	21.3	695
Lifetime No. of Partner	s								
0	95.8	41.2	13.9	7.9	5.1	3.5	3.5	4.0	1,439
1	95.7	63.1	13.1	6.2	5.6	4.6	4.6	3.7	4,140
2+	95.5	66.1	29.1	12.7	10.9	8.2	7.3	9.1	113
Don't Know	*	*	*	*	*	*	*	*	5

* Percentages are not shown when base is less than 25 cases.

		0	hea	Ę	nonas		dia		
Characteristic	HIV/ AIDS	Syphilis	Gonorrhea	Yeast Infection	Trichomonas	Genital Herpes	Chlamydia	Genital Warts	No. of Cases
Total	96.1	21.8	14.2	11.7	4.4	3.8	3.7	3.5	1,740
Strata									
Tirana	98.4	29.7	19.6	18.2	8.6	8.5	8.1	7.4	718
Urban	98.2	30.9	21.1	19.2	7.0	5.1	5.9	4.4	547
Rural	93.9	13.3	8.0	4.8	1.2	1.1	0.6	1.6	475
Age Group									
15–19	98.4	18.0	9.8	9.1	4.1	2.3	2.5	2.3	401
20–24	99.9	21.7	17.4	11.7	3.6	2.7	3.3	3.4	189
25–29	96.6	25.2	16.8	12.7	4.3	3.1	4.5	4.3	218
30–34	99.5	23.3	17.9	13.9	7.0	5.6	5.4	6.5	253
35–39	93.9	19.7	13.9	12.7	6.0	5.1	5.1	4.2	255
40–44	95.3	21.5	10.0	11.9	3.1	3.9	2.3	2.3	277
45-49	86.7	25.3	15.5	11.0	2.9	4.1	2.7	1.8	147
Marital Status									
Currently Married	94.9	20.2	13.5	11.8	3.8	3.6	3.4	3.0	1,023
Previously Married	*	*	*	*	*	*	*	*	14*
Never Married	98.2	24.2	15.3	11.6	5.3	3.9	4.0	4.3	703
No. of Living Children									
0	98.5	24.5	15.5	12.0	5.5	3.9	4.5	4.5	815
1	98.7	20.6	18.5	15.1	5.7	3.8	4.5	4.8	221
2	95.7	23.6	14.5	13.2	3.7	4.3	3.6	3.1	468
3	93.6	14.6	7.3	9.2	2.4	3.0	1.7	1.6	167
4+	80.7	12.7	8.8	2.6	0.3	2.0	0.3	0.0	69
Education									
Primary or Less	92.6	10.3	5.2	5.1	1.4	0.6	0.5	1.2	689
Secondary	99.3	24.9	17.2	14.1	4.4	3.8	4.3	3.5	825
Post-Secondary	99.8	68.4	48.2	36.4	20.5	20.7	18.0	16.5	226
Lifetime No. of Partners**									
0	98.0	14.5	8.4	7.5	3.4	1.8	2.0	1.8	446
1	93.5	14.4	9.2	6.0	4.1	2.8	3.2	2.0	448
2+	98.4	29.6	18.1	17.0	5.4	5.1	5.3	5.8	702
Don't Know	89.2	35.8	33.4	22.1	4.4	7.3	2.9	4.1	144

Table 14.1 BPercentage of Men Aged 15–49 Who Have Heard of SpecifiedSexually Transmitted Infections by Selected CharacteristicsReproductive Health Survey: Albania, 2002

* Percentages are not shown when base is less than 25 cases.

Tables

Table 14.2 A Percentage of Women Aged 15–44 With Knowledge of Symptoms Associated With STIs, Other Than HIV/AIDS, in a Woman by Selected Characteristics Reproductive Health Survey: Albania, 2002

		d of at						
	one S	east TI other IIV/AIDS	Knowled					
Characteristic	%	N	No Awareness of STIs	No	<u>Distribution</u> One Symptom	, Two or More Symptoms	- Total	No. of Cases
Total	58.8	5,697	41.2	18.8	11.7	28.3	100.0	5,697
Strata								
Tirana	65.6	2,108	34.4	19.0	9.9	36.8	100.0	2,108
Urban	67.9	1,816	32.1	24.9	12.1	30.9	100.0	1,816
Rural	52.0	1,773	48.0	15.5	12.0	24.5	100.0	1,773
Age Group								
15–19	42.2	1,094	57.8	17.5	11.1	13.6	100.0	1,094
20–24	56.6	936	43.4	17.8	12.0	26.8	100.0	936
25–29	65.1	946	34.9	22.3	11.1	31.8	100.0	946
30–34	64.2	1,067	35.8	16.7	11.7	35.8	100.0	1,067
35–39	66.9	958	33.1	18.7	12.2	36.0	100.0	958
40–44	64.3	696	35.7	20.6	12.4	31.3	100.0	696
Marital Status								
Currently Married	64.3	3,965	35.7	19.4	12.2	32.7	100.0	3,965
Previously Married	71.8	88	28.2	12.8	14.5	44.6	100.0	88
Never Married	47.2	1,644	52.8	18.1	10.5	18.6	100.0	1,644
Education								
Primary or Less	49.3	2,519	50.7	15.3	11.5	22.6	100.0	2,519
Secondary	66.0	2,483	34.0	23.0	12.0	31.1	100.0	2,483
Post-Secondary	88.5	695	11.5	23.2	12.1	53.2	100.0	695
Lifetime No. of Partne	rs							
0	45.1	1,439	54.9	17.6	10.4	17.0	100.0	1,439
1	64.7	4,140	35.4	19.5	12.2	33.0	100.0	4,140
2+	69.9	113	30.1	13.8	16.9	39.3	100.0	113
Don't Know	*	5*	*	*	*	*	*	5

* Percentages are not shown when base is less than 25 cases.

	Le One S	d of at east TI Other HV/AIDS	Knowle					
Characteristic	%	N	No Awareness of STIs	No	<u>Distribution</u> One Symptom	Two or More Symptoms	- Total	No. of Cases
Total	26.5	1,740	73.5	6.6	4.6	15.2	100.0	1,740
Strata								
Tirana	35.8	718	64.2	5.7	6.8	23.3	100.0	718
Urban	37.3	547	62.7	9.6	6.6	21.1	100.0	547
Rural	16.5	475	83.5	5.3	2.6	8.6	100.0	475
Age Group								
15–19	22.7	401	77.3	6.7	4.4	11.6	100.0	401
20–24	26.0	189	74.0	7.4	4.2	14.4	100.0	189
25–29	30.2	218	69.8	5.3	5.2	19.8	100.0	218
30–34	29.5	253	70.5	4.1	5.0	20.4	100.0	253
35–39	24.7	255	75.3	5.8	4.3	14.6	100.0	255
40–44	26.5	277	73.5	9.7	5.2	11.6	100.0	277
45-49	27.8	147	72.2	7.3	4.3	16.2	100.0	147
Marital Status								
Currently Married	25.2	1,023	74.8	6.1	4.4	14.8	100.0	1,023
Previously Married	*	14*	*	*	*	*	*	14
Never Married	28.5	703	71.5	7.6	4.9	16.0	100.0	703
Education								
Primary or Less	13.5	689	86.5	2.7	2.8	8.0	100.0	689
Secondary	31.8	825	68.2	10.4	5.5	15.9	100.0	825
Post-Secondary	71.2	226	28.8	9.7	10.4	51.1	100.0	226
Lifetime No. of Partn	ers							
0	18.2	446	81.8	4.5	3.5	10.1	100.0	446
1	16.6	448	83.4	2.8	3.6	10.2	100.0	448
2+	36.5	702	63.5	10.3	5.6	20.6	100.0	702
Don't Know	43.1	144	56.9	10.5	7.9	24.7	100.0	144

Table 14.2 B Percentage of Men Aged 15–49 With Knowledge of Symptoms Associated With STIs, Other Than HIV/AIDS, in a Man by Selected Characteristics Reproductive Health Survey: Albania, 2002

* Percentage not shown when base is less than 25 cases.

2

Table	14.3 A
-------	--------

Most Important Source of Information About Sexually Transmitted Infections All Women Aged 15–44 Who Have Heard of at Least One STI by Selected Characteristics Reproductive Health Survey: Albania, 2002 (Percent Distribution)

	Most Important Source of Information About STIs										
Characteristic	Mass Media	Friends	Doctor	A Parent or Relative	Books	School	Partner	Other	Do Not Remember	Total	No. of Cases
Total	84.2	1.0	2.6	1.3	2.9	4.4	3.4	0.0	0.2	100.0	3,596
Strata											
Tirana	75.4	1.9	3.9	1.9	5.2	7.7	4.0	0.0	0.1	100.0	1,417
Urban	84.17	1.5	1.1	1.5	3.7	5.5	2.4	0.0	0.1	100.0	1,244
Rural	87.5	0.3	3.1	0.9	1.5	2.5	4.0	0.0	0.3	100.0	935
Age Group											
15–19	66.7	0.5	0.4	3.9	4.6	20.1	3.9	0.0	0.1	100.0	512
20–24	82.0	0.8	1.6	1.2	4.3	6.6	3.4	0.0	0.0	100.0	578
25–29	89.9	0.6	2.6	0.3	2.0	0.6	3.8	0.0	0.3	100.0	646
30–34	87.5	1.2	4.3	0.4	3.0	0.7	2.5	0.0	0.4	100.0	722
35–39	88.6	1.3	2.6	0.9	2.1	0.4	3.9	0.0	0.2	100.0	664
40–44	88.2	1.5	3.7	1.4	1.8	0.1	3.1	0.0	0.2	100.0	474
Marital Status											
Currently Married	88.5	1.0	3.1	1.3	1.9	0.8	3.1	0.0	0.3	100.0	2,659
Previously Married	91.2	1.8	0.7	0.7	0.7	0.0	5.0	0.0	0.0	100.0	61
Never Married	71.7	0.8	1.3	1.4	5.9	14.7	4.1	0.0	0.0	100.0	876
Education											
Primary or Less	88.0	0.5	2.5	1.2	1.0	1.6	4.9	0.0	0.2	100.0	1,275
Secondary	82.8	1.1	2.5	1.5	3.2	6.4	2.3	0.0	0.3	100.0	1,702
Post-Secondary	74.6	2.6	3.1	1.1	9.2	7.8	1.7	0.0	0.0	100.0	619
Lifetime No. of Partn	ers										
0	71.5	0.7	1.1	1.5	5.5	15.6	4.2	0.0	0.0	100.0	713
1	88.4	1.0	2.9	1.2	2.0	1.0	3.2	0.0	0.2	100.0	2,796
2+	71.0	4.2	6.7	1.3	9.6	3.8	3.4	0.0	0.0	100.0	83
Don't Know	*	*	*	*	*	*	*	*	*	*	4

* Percentage not shown when base is less than 25 cases.

Table 14.3 B
Most Important Source of Information on Sexually Transmitted Infections
All Men Aged 15–49 Who Have Heard of at Least One STI by Selected Characteristics
Reproductive Health Survey: Albania, 2002
(Percent Distribution)

	Most Important Source of Information About STIs										
Characteristic	Mass Media	Friends	Doctor	A Parent or Relative	Books	School	Partner	Other	Do Not Remember	Total	No. of Cases
Total	66.4	7.1	10.1	1.4	7.7	4.8	0.1	0.0	2.4	100.0	545
Strata											
Tirana	67.6	4.6	11.7	0.9	9.3	4.5	0.3	0.0	1.0	100.0	264
Urban	62.8	8.9	9.4	1.8	8.2	5.2	0.0	0.0	3.7	100.0	203
Rural	70.2	6.8	9.5	1.3	5.6	4.9	0.0	0.0	1.8	100.0	78
Age Group											
15–19	59.8	0.7	1.6	4.1	9.8	22.8	0.0	0.0	1.2	100.0	100
20–24	63.7	4.2	7.4	3.5	13.6	4.1	0.0	0.0	3.5	100.0	60
25–29	69.0	7.2	10.8	0.0	10.7	2.3	0.0	0.0	0.0	100.0	76
30–34	69.8	11.8	9.6	0.0	5.5	1.3	0.0	0.0	1.9	100.0	86
35–39	70.8	5.3	15.0	0.0	3.8	0.0	0.7	0.0	4.3	100.0	76
40–44	67.5	15.4	11.5	0.0	4.4	0.6	0.0	0.0	0.6	100.0	92
45-49	64.7	5.0	17.3	2.1	4.8	0.0	0.0	0.0	6.2	100.0	55
Marital Status											
Currently Married	69.6	8.3	12.2	0.4	4.2	1.8	0.2	0.0	3.4	100.0	313
Previously Married	*	*	*	*	*	*	*	*	*	*	6
Never Married	61.7	5.3	7.4	2.8	12.6	9.3	0.0	0.0	1.0	100.0	226
Education											
Primary or Less	62.7	9.1	16.0	1.7	5.0	2.0	0.0	0.0	3.5	100.0	99
Secondary	70.8	6.2	9.9	1.1	1.6	8.1	0.0	0.0	2.4	100.0	281
Post-Secondary	60.9	7.0	4.5	1.9	23.1	1.2	0.4	0.0	1.2	100.0	165
Lifetime No. of Partne	ers										
0	62.4	0.7	2.1	3.3	10.0	20.4	0.0	0.0	1.2	100.0	99
1	78.2	4.4	3.9	1.6	8.0	0.0	0.0	0.0	3.9	100.0	101
2+	59.6	11.9	16.3	1.1	8.0	2.4	0.2	0.0	0.6	100.0	277
Don't Know	82.0	0.0	4.1	0.0	3.1	1.6	0.0	0.0	9.2	100.0	68

* Percentage not shown when base is less than 25 cases.

Tables

Percentage of Women of Reproductive Age Who Had Received Radio And Television Messages About HIV/AIDS and Other STIs During The Past Six Months By Selected Characteristics

Reproductive Health Survey: Albania, 2002

	HIV/AIDS							HER S	Tl's			
Characteristic	TV	Radio And TV	Radio	Neither Radio Nor TV	Do Not Know	TV	Radio And TV	Radio	Neither Radio Nor TV	Do Not Know	Total	No. of Cases
Total	51.5	5.5	0.3	39.5	3.3	6.2	1.3	0.1	86.4	6.0	100.0	5,697
Strata												
Tirana	55.4	7.8	0.5	33.5	2.9	11.7	2.8	0.6	78.6	6.3	100.0	2,108
Urban	55.9	7.4	0.3	33.3	3.2	10.1	2.0	0.1	80.7	7.0	100.0	1,816
Rural	48.1	3.7	0.2	44.6	3.4	2.4	0.5	0.0	91.7	5.4	100.0	1,773
Age Group												
15–19	48.9	6.7	0.3	41.7	2.6	3.5	1.2	0.2	90.3	4.8	100.0	1,094
20–24	51.0	7.3	0.4	38.4	2.9	7.5	1.5	0.2	85.4	5.5	100.0	936
25–29	51.2	4.2	0.4	40.5	3.8	5.7	1.1	0.1	86.9	6.3	100.0	946
30–34	52.5	4.5	0.4	39.4	3.2	8.5	1.1	0.2	83.2	7.0	100.0	1,067
35–39	54.5	4.5	0.3	36.6	4.1	6.9	1.6	0.2	84.6	6.7	100.0	958
40–44	52.1	5.0	0.0	39.6	3.3	5.7	1.5	0.1	86.5	6.2	100.0	696
Marital Status												
Currently Married	51.4	4.7	0.2	40.5	3.2	6.2	1.2	0.1	86.4	6.1	100.0	3,965
Previously Married	57.9	2.6	0.7	32.9	5.9	9.9	1.3	0.7	69.7	18.4	100.0	88
Never Married	51.5	7.1	0.4	37.9	3.1	6.0	1.5	0.3	87.3	5.0	100.0	1,644
Education												
Primary or Less	46.7	4.2	0.2	45.8	3.2	2.7	0.4	0.0	91.0	5.9	100.0	2,519
Secondary	56.4	6.1	0.3	33.6	3.6	8.4	2.3	0.2	83.1	6.0	100.0	2,483
Post-Secondary	61.1	10.9	0.3	25.2	2.6	19.0	3.2	0.9	70.7	6.1	100.0	695
Lifetime No. of Partne	rs											
0	50.7	7.2	0.4	38.4	3.3	5.1	1.5	0.3	88.1	5.0	100.0	1,439
1	51.8	4.7	0.2	40.0	3.3	6.4	1.3	0.1	85.8	6.4	100.0	4,140
2+	55.5	3.6	0.9	40.0	0.0	15.5	0.0	0.9	77.3	6.4	100.0	113
Don't Know	*	*	*	*	*	*	*	*	*	*	*	5

* Percentage not shown when base is less than 25 cases.

Percentage of Men of Reproductive Age Who Had Received Radio And Television Messages About HIV/AIDS and Other STIs During The Past Six Months By Selected Characteristics Reproductive Health Survey: Albania, 2002												
					alth Su	irvey						
			HIV/AIC				0	THER				
Characteristic	TV	Radio And TV	Radio	Neither Radio Nor TV	Do Not Know	TV	Radio And TV	Radio	Neither Radio Nor TV	Do Not Know	Total	No. of Cases
Total	44.9	18.2	0.5	34.3	2.1	6.6	12.5	0.3	71.2	9.4	100.0	1,740
Strata												
Tirana	45.3	23.2	0.5	28.4	2.6	10.8	13.5	0.4	62.6	12.7	100.0	718
Urban	41.6	17.4	1.1	38.6	1.2	5.9	12.7	0.2	73.5	7.8	100.0	547
Rural	46.7	12.5	0.4	38.3	2.1	1.5	11.5	0.2	80.9	6.0	100.0	475
Age Group												
15–19	49.9	18.1	0.0	28.7	3.2	4.1	11.9	0.7	75.0	8.3	100.0	401
20–24	53.1	7.3	0.3	37.5	1.9	3.8	6.3	0.0	82.2	7.7	100.0	189
25–29	37.7	24.1	0.3	36.0	1.9	5.7	16.7	0.0	70.5	7.1	100.0	218
30–34	48.8	17.8	1.2	31.3	0.9	7.0	14.7	0.8	69.1	8.5	100.0	253
35–39	44.0	16.5	1.0	36.7	1.7	4.5	14.0	0.0	72.7	8.8	100.0	255
40–44	40.6	14.4	1.0	42.1	1.9	3.8	11.6	0.0	78.8	5.8	100.0	277
45-49	36.2	13.9	0.5	47.2	2.2	3.8	10.8	0.0	76.8	8.6	100.0	147
Marital Status												
Currently Married	41.4	16.5	0.9	39.6	1.7	4.2	13.7	0.1	75.2	6.9	100.0	1,023
Previously Married	*	*	*	*	*	*	*	*	*	*	*	14
Never Married	50.7	15.4	0.3	31.2	2.5	5.2	10.1	0.5	74.8	9.4	100.0	703
Education												
Primary or Less	45.4	13.4	0.5	38.8	1.9	1.7	11.8	0.2	79.4	6.9	100.0	689
Secondary	44.6	18.2	0.9	34.6	1.7	5.2	12.1	0.2	73.7	8.7	100.0	825
Post-Secondary	43.5	20.5	0.0	32.8	3.2	18.0	15.0	0.2	58.0	8.9	100.0	226
Lifetime No. of Partn	ers											
0	53.8	15.9	0.0	28.1	2.2	3.5	10.9	0.5	78.1	7.0	100.0	446
1	49.9	19.0	0.8	28.9	1.5	3.2	15.2	0.1	75.8	5.7	100.0	448
2+	38.5	14.5	0.8	44.6	1.6	5.7	11.0	0.2	74.2	9.0	100.0	702
Don't Know	28.5	13.1	0.9	52.4	5.1	8.8	11.4	0.0	66.9	12.9	100.0	42

Table 14.4 B fМ 4 D . \A/I . .

* Percentage not shown when base is less than 25 cases.

Tables

Table 14.5 A Percentage of Women Aged 15-44 Years with a Diagnosis of Specified Sexually Transmitted Infections by Selected Characteristics Reproductive Health Survey: Albania, 2002

	Yeast			Genital		Genital		No. of
Characteristic	Infection	Syphilis	Gonorrhea	Herpes	Trichomoniasis	Warts	Chlamydia	Cases
Total	8.3	0.0	0.0	0.0	0.1	0.1	0.0	5,697
Strata								
Tirana	11.5	0.0	0.0	0.2	0.2	0.1	0.0	2,108
Urban	9.0	0.0	0.0	0.0	0.1	0.1	0.0	1,816
Rural	7.1	0.0	0.0	0.0	0.1	0.1	0.0	1,773
Age Group								
15–19	1.2	0.0	0.0	0.0	0.0	0.0	0.0	1,094
20–24	6.3	0.0	0.0	0.0	0.0	0.0	0.0	936
25–29	11.3	0.0	0.0	0.2	0.1	0.1	0.0	946
30–34	11.3	0.0	0.0	0.0	0.1	0.3	0.0	1,067
35–39	13.3	0.0	0.0	0.1	0.2	0.0	0.0	958
40–44	9.4	0.0	0.0	0.1	0.1	0.1	0.1	696
Marital Status								
Currently Married	11.7	0.0	0.0	0.1	0.1	0.1	0.0	3,965
Previously Married	15.2	0.0	0.0	0.0	0.0	0.0	0.0	88
Never Married	1.2	0.0	0.0	0.0	0.0	0.0	0.0	1,644
No. of Living Children	n							
0	2.7	0.0	0.0	0.0	0.0	0.0	0.0	1,943
1	12.9	0.0	0.0	0.0	0.3	0.0	0.0	828
2	11.5	0.0	0.0	0.2	0.1	0.3	0.1	1,840
3	10.9	0.0	0.0	0.0	0.2	0.0	0.0	795
4+	12.0	0.0	0.0	0.0	0.0	0.0	0.0	291
Education								
Primary or Less	7.3	0.0	0.0	0.1	0.0	0.1	0.1	2,519
Secondary	9.4	0.0	0.0	0.0	0.1	0.0	0.0	2,483
Post-Secondary	10.2	0.0	0.0	0.0	0.2	0.0	0.0	695
Lifetime No. of Partne	ers							
0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	1,439
1	11.5	0.0	0.0	0.1	0.1	0.1	0.0	4,140
2+	17.3	0.0	0.0	0.0	0.9	0.0	0.0	113
Don't Know	**	**	**	**	**	**	**	5

* Respondents were not asked about the results of HIV testing.

 * *Percentage not shown when base is less than 25 cases.

	Yeast			Genital		Genital		No. of
Characteristic	Infection	Syphilis	Gonorrhea		Trichomoniasis	Warts	Chlamydia	
Total	1.2	0.0	0.0	0.0	0.0	0.0	0.1	1,740
Strata								
Tirana	1.5	0.0	0.0	0.1	0.0	0.0	0.0	718
Urban	2.3	0.0	0.0	0.0	0.0	0.0	0.0	547
Rural	0.4	0.0	0.0	0.0	0.0	0.0	0.0	475
Age Group								
15–19	0.0	0.0	0.0	0.0	0.0	0.0	0.0	401
20–24	0.7	0.0	0.0	0.0	0.0	0.0	0.0	189
25–29	1.9	0.0	0.0	0.0	0.0	0.0	0.0	218
30–34	1.7	0.0	0.0	0.0	0.0	0.0	0.8	253
35–39	1.3	0.0	0.0	0.0	0.0	0.0	0.0	255
40–44	0.6	0.0	0.0	0.1	0.0	0.0	0.0	277
45-49	2.9	0.0	0.0	0.0	0.0	0.0	0.0	147
Marital Status								
Currently Married	1.7	0.0	0.0	0.0	0.0	0.0	0.2	1,023
Previously Married	**	**	**	**	**	**	**	14
Never Married	0.4	0.0	0.0	0.0	0.0	0.0	0.0	703
No. of Living Children	l							
0	0.9	0.0	0.0	0.0	0.0	0.0	0.2	815
1	2.7	0.0	0.0	0.2	0.0	0.0	0.0	221
2	1.3	0.0	0.0	0.0	0.0	0.0	0.0	468
3	1.0	0.0	0.0	0.0	0.0	0.0	0.0	167
4+	0.0	0.0	0.0	0.0	0.0	0.0	0.0	69
Education								
Primary or Less	0.6	0.0	0.0	0.0	0.0	0.0	0.2	689
Secondary	2.0	0.0	0.0	0.0	0.0	0.0	0.0	825
Post-Secondary	0.2	0.0	0.0	0.0	0.0	0.0	0.0	226
Lifetime No. of Partne	rs							
0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	446
1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	448
2+	2.0	0.0	0.0	0.1	0.0	0.0	0.3	702
Don't Know * Respondents were not as	5.2 sked about the r	0.0 esults of HIV	0.0	0.0	0.0	0.0	0.0	42

Table 14.5 B Percentage Men Aged 15-49 With a Diagnosis of Specified Sexually Transmitted Infections by Selected Characteristics Reproductive Health Survey: Albania, 2002

* Respondents were not asked about the results of HIV testing.

* *Percentage not shown when base is less than 25 cases.

ß

Table 14.6 A Level of Awareness, Testing, Diagnosis, And Treatment For STIs Among Women Aged 15-44 Years Who Have Ever Had Sexual Intercourse by Specific Sexually Transmitted Infections Reproductive Health Survey: Albania, 2002

Specific STIs	Awareness of The STI	Testing for The STI	Diagnosis of The STI	Treatment For The STI	Number of Caes
HIV/AIDS	95.7	0.3	*	*	4,087
Yeast Infection	62.9	14.0	11.5	11.5	4,087
Syphilis	13.4	0.1	0.0	0.0	4,087
Gonorrhea	6.3	0.1	0.0	0.0	4,087
Genital Herpes	5.7	0.2	0.1	0.1	4,087
Genital Warts	4.6	0.1	0.1	0.1	4,087
Trichomonas	4.5	0.3	0.1	0.1	4,087
Chlamydia	3.8	0.1	0.0	0.0	4,087

* Respondents were not asked about the results of HIV testing.

Table 14.6 B Level of Awareness, Testing, Diagnosis, And Treatment For STIs Among Men Aged 15-49 Years Who Have Ever Had Sexual Intercourse by Specific Sexually Transmitted Infections Reproductive Health Survey: Albania, 2002

Specific STIs	Awareness of The STI	Testing For The STI	Diagnosis of The STI	Treatment For The STI	Number of Cases
HIV/AIDS	95.5	0.9	*	*	1,296
Syphilis	24.2	0.2	0.0	0.0	1,296
Gonorrhea	16.1	0.2	0.0	0.0	1,296
Yeast Infection	13.1	1.7	1.5	1.4	1,296
Trichomonas	4.8	0.1	0.0	0.0	1,296
Genital Herpes	4.4	0.0	0.0	0.0	1,296
Chlamydia	4.2	0.2	0.1	0.1	1,296
Genital Warts	4.1	4.1	0.0	0.0	1,296

* Respondents were not asked about the results of HIV testing.

Table 14.7 A	
Percent Distribution of Women Aged 15-44 Who Have Heard	
of at Least One Sexually Transmitted Infection	
By Self-Perceived Risk of Acquiring an STI by Selected Characteristics	
Reproductive Health Survey: Albania, 2002	
Self-Perceived Risk	

Characteristic	High or Medium risk	Low Risk	No Risk	Do Not Know	Total	No. of Cases
Total	2.2	13.7	79.7	4.3	100.0	3596
Strata						
Tirana	3.6	14.7	77.6	4.1	100.0	1417
Urban	2.6	13.6	79.9	4.0	100.0	1244
Rural	1.5	13.4	80.4	4.7	100.0	935
Age Group						
15–19	1.0	7.2	88.2	3.6	100.0	512
20–24	2.1	13.1	79.1	5.7	100.0	578
25–29	2.1	18.6	74.8	4.5	100.0	646
30–34	3.4	12.9	79.8	4.0	100.0	722
35–39	1.7	13.6	79.6	5.1	100.0	664
40–44	3.0	16.1	77.9	3.1	100.0	474
Marital Status						
Currently Married	2.8	15.3	77.5	4.4	100.0	2659
Previously Married	0.0	5.8	90.6	3.7	100.0	61
Never Married	1.1	10.1	84.6	4.2	100.0	876
Education						
Primary or Less	2.0	11.8	80.4	5.8	100.0	1275
Secondary	2.0	15.5	79.0	3.5	100.0	1702
Post-Secondary	4.0	14.3	79.7	2.1	100.0	619
Lifetime No. of Partners						
0	0.7	7.9	87.3	4.0	100.0	713
1	2.6	15.3	77.7	4.5	100.0	2796
2+	9.0	24.0	65.0	2.1	100.0	83
Don't Know	*	*	*	*	*	4

* Percentage not shown when base is less than 25 cases.

Tabl	e	14.	7	В

Percent Distribution of Men Aged 15-49 Who Have Heard of at Least One Sexually Transmitted Infection By Self-Perceived Risk of Acquiring an STI by Selected Characteristics Reproductive Health Survey: Albania, 2002

	Self-Perceived Risk				_	
Characteristic	High or Medium Risk	Low Risk	No Risk	Do Not Know	Total	No. of Cases
Total	1.7	1.8	88.3	8.2	100.0	545
Strata						
Tirana	1.5	1.7	82.7	14.1	100.0	264
Urban	1.1	1.3	91.0	6.6	100.0	203
Rural	2.8	2.5	89.5	5.2	100.0	78
Age Group						
15–19	3.2	2.3	88.6	6.0	100.0	100
20–24	0.0	1.7	82.8	15.5	100.0	60
25–29	1.4	6.6	84.8	7.2	100.0	76
30–34	3.7	0.6	80.5	15.2	100.0	86
35–39	0.7	0.0	92.5	6.8	100.0	76
40–44	0.0	0.7	95.6	3.7	100.0	92
45-49	2.2	0.0	95.9	1.9	100.0	55
Marital Status						
Currently Married	2.0	0.2	90.2	7.6	100.0	313
Previously Married	*	*	*	*	*	6
Never Married	1.3	4.0	85.9	8.9	100.0	226
Education						
Primary or Less	1.9	3.7	86.9	7.6	100.0	99
Secondary	2.1	1.6	88.2	8.1	100.0	281
Post-Secondary	0.8	0.4	90.0	8.9	100.0	165
Lifetime No. of Partners						
0	3.1	1.2	91.8	3.9	100.0	99
1	0.5	0.0	96.7	2.8	100.0	101
2+	1.4	3.1	84.2	11.3	100.0	277
Don't Know	3.0	0.0	86.8	10.1	100.0	68

* Percentage not shown when base is less than 25 cases.

lables

Table 14.8 A
Percentage of Women Aged 15–44 Who Have Heard of HIV/AIDS,
Who Believe HIV/AIDS Infection Can Be Asymptomatic,
and Who Know Where HIV Testing Is Provided by Selected Characteristics
Reproductive Health Survey: Albania, 2002

Characteristic	Have Heard of HIV/AIDS	Believe That HIV/AIDS Infection Can Be Asymptomatic	Know Where HIV Tests Are Provided	Have Been Tested For HIV	No. of Cases
Total	95.7	55.5	16.9	0.2	5,697
Strata					
Tirana	97.1	62.9	32.3	0.6	2,108
Other Urban	98.1	62.8	19.3	0.2	1,816
Rural	94.0	49.4	11.1	0.1	1,773
Age Group					
15–19	95.7	55.2	15.9	0.0	1,094
20–24	97.1	60.7	17.2	0.1	936
25–29	97.2	56.7	17.1	0.2	946
30–34	94.1	55.8	18.7	0.3	1,067
35–39	96.1	53.6	15.2	0.2	958
40–44	93.9	50.4	17.7	0.3	696
Marital Status					
Currently Married	95.5	53.9	16.3	0.2	3,965
Previously Married	96.0	58.9	20.5	0.5	88
Never Married	96.1	58.5	17.8	0.1	1,644
Education					
Primary or Less	92.8	45.1	9.0	0.0	2,519
Secondary	98.9	64.1	21.4	0.3	2,483
Post-Secondary	99.8	84.7	48.1	0.7	695
Lifetime No. of Partners					
0	95.8	57.3	16.2	0.1	1,439
1	95.7	54.6	16.8	0.2	4,140
2+	95.5	60.0	32.7	0.0	113
Don't Know	*	*	*	*	5

* Percentage not shown when base is less than 25 cases.

Who Believe HIV/AIDS Infection Can be Asymptomatic, and Who Know Where HIV Testing is Provided by Selected Characteristics Reproductive Health Survey: Albania, 2002							
Characteristic	Have Heard of HIV/AIDS	Believe That HIV/AIDS Infection Can be Asymptomatic	Know Where HIV Tests Are Provided	Have Been Tested For HIV	No. of Cases		
Total	96.1	44.7	33.2	0.8	1,740		
Strata							
Tirana	98.4	50.6	46.0	3.7	718		
Other Urban	98.2	62.2	40.7	0.2	547		
Rural	93.9	31.9	23.6	0.0	475		
Age Group							
15–19	98.4	44.4	26.0	0.1	401		
20–24	99.9	53.1	34.4	2.5	189		
25–29	96.6	53.2	43.9	0.9	218		
30–34	99.5	52.1	41.6	0.2	253		
35–39	93.9	39.9	31.4	0.5	255		
40–44	95.3	40.3	34.5	0.5	277		
45-49	86.7	27.1	21.3	1.0	147		
Marital Status							
Currently Married	94.9	40.0	33.2	0.5	1,023		
Previously Married	*	*	*	*	14		
Never Married	98.2	52.0	33.2	1.1	703		
Education							
Primary or Less	92.6	29.7	20.5	0.4	689		
Secondary	99.3	53.7	39.5	0.8	825		
Post-Secondary	99.8	82.7	70.9	3.0	226		
Lifetime No. of Partne	ers						
0	98.0	43.2	24.6	0.4	446		
1	93.5	32.3	29.5	0.3	448		
2+	98.4	55.8	40.9	1.5	702		
Don't Know	89.3	43.2	36.7	0.7	42		

Table 14.8 B Percentage of Men Aged 15-49 Who Have Heard of HIV/AIDS,

*Percentage not shown when base is less than 25 cases.

Albania Reproductive Health Survey, 2002 Final Report

S

Table 14.9 A Percentage of Women Aged 15-44 Who Do Not Know Principle Mechanisms of HIV Transmission by Selected Characteristics Reproductive Health Survey: Albania, 2002

Characteristic	MTCT Breastfeeding	MTCT Pregnancy or Delivery	Unprotected Homosexual Intercourse	Non- Sterile Needles	Blood Transfusion	Unprotected Heterosexual Intercourse	No. of Cases
Total	23.9	17.5	15.8	10.0	10.2	5.3	5,697
Strata							
Tirana	25.7	14.1	13.5	7.6	7.1	5.5	2,108
Urban	21.2	13.2	12.6	6.1	7.1	5.9	1,816
Rural	24.8	20.8	18.2	12.8	12.8	4.9	1,773
Age Group							
15–19	28.7	20.7	20.4	10.5	10.4	5.7	1,094
20–24	26.2	19.8	14.5	9.4	10.1	5.0	936
25–29	21.2	13.4	15.2	11.6	10.7	5.2	946
30–34	22.5	18.0	15.0	10.3	9.6	5.6	1,067
35–39	21.1	16.7	14.5	8.8	11.2	5.0	958
40–44	21.5	15.3	13.7	8.8	9	5.1	696
Marital Status							
Currently Married	22.2	16.7	15.2	10.4	10.7	5.4	3,965
Previously Married	31.8	22.5	11.1	9.8	13.0	4.1	88
Never Married	26.7	18.8	17.3	9.1	8.9	5.2	1,644
Education							
Primary or Less	26.3	22.3	19.2	14.3	15.4	6.6	2,519
Secondary	21.0	13.3	13.0	5.5	4.9	3.9	2,483
Post-Secondary	20.9	5.7	6.3	1.7	0.4	3.3	695
Lifetime No. of Partr	iers						
0	26.4	19.0	17.3	9.2	9.0	5.4	1,439
1	22.7	17.1	15.3	10.4	10.8	5.3	4,140
2+	24.9	8.7	7.2	6.9	7.1	3.4	113
Don't Know	*	*	*	*	*	*	5

* Percentage not shown when base is less than 25 cases.

Characteristic	MTCT Breastfeeding	MTCT Pregnancy or Delivery	Unprotected Homosexual Intercourse	Non-Sterile Needles	Blood Transfusion	Unprotected Heterosexual Intercourse	No. of Cases
Total	55.2	37.4	35.7	14.0	8.0	16.5	1,740
Strata							
Tirana	53.6	32.1	29.4	6.8	3.6	18.8	718
Urban	52.8	27.9	33	12.2	6.2	14.1	547
Rural	57.1	45.1	39.7	17.8	10.8	17.0	475
Age Group							
15–19	62.8	48.0	47.1	16.0	5.8	21.4	401
20–24	54.3	34.7	33.8	12.9	6.4	19.8	189
25–29	52.4	34.2	22.6	9.6	4.5	9.5	218
30–34	53.9	32.7	27.2	8.0	4.2	15.2	253
35–39	56.0	37.0	38.7	12.3	10.2	19.0	255
40–44	50.4	32.2	40.6	15.5	10.1	15.5	277
45-49	53.3	39.3	34.8	24.2	16.7	12.0	147
Marital Status							
Currently Married	54.2	36.9	35.8	14.9	9.9	15.5	1,023
Previously Married	*	*	*	*	*	*	14
Never Married	56.7	38.1	36.1	12.6	5.1	18.2	703
Education							
Primary or Less	55.3	43.5	41.9	18.1	11.0	17.9	689
Secondary	56.4	35.6	32.5	11.6	5.9	16.4	825
Post-Secondary	48.5	13.2	17.4	2.7	1.8	9.7	226
Lifetime No. of Partne	rs						
0	63.7	47.8	43.8	15.8	6.2	23.5	446
1	63.7	46.5	39.2	15.6	12.7	22.8	448
2+ Don't Know	41.5 61.4	25.0 28.9	30.9 19.5	10.3 19.2	5.8 5.7	8.4 9.3	702 42

Table 14.9 B Percentage of Men Aged 15-49 Who Do Not Know Principle Mechanisms of HIV Transmission by Selected Characteristics Reproductive Health Survey: Albania, 2002

 * Percentage not shown when base is less than 25 cases.

2

Table 14.10 A Percentage of Women Aged 15-44 Who Correctly Reject Misconceptions About HIV Transmission by Selected Characteristics Reproductive Health Survey: Albania, 2002

	Shaking		Public		Eating	Mosquito	Dental/	Never Heard of	No. of
Characteristic	Hands	Manicure	Toilet	Kissing	Utensils	Bites	Surgical	HIV/AIDS	Cases
Total	66.3	66.1	44.2	41.6	38.3	24.8	4.0	4.3	5,697
Strata									
Tirana	73.2	72.4	49.7	50.3	47.3	26.5	3.4	3.0	2,108
Urban	73.1	73.6	50.2	51.2	45.4	28.9	3.0	1.9	1,816
Rural	60.7	60.2	39.3	33.9	31.5	22.1	4.8	6.0	1,773
Age Group									
15–19	72.7	72.2	50.8	47.4	42.6	28.5	6.3	4.3	1,094
20–24	74.9	72.3	48.3	44.5	40.1	26.9	4.6	2.9	936
25–29	65.6	66.2	40.2	38.5	37.9	24.6	3.8	2.9	946
30–34	60.8	60.9	44.4	41.3	38.4	23.6	2.9	5.9	1,067
35–39	60.7	61.2	38.5	36.6	35.1	19.3	3.7	3.9	958
40–44	60.2	61.1	40.0	38.8	33.7	24.5	2.0	6.1	696
Marital Status									
Currently Married	62.3	62.6	40.9	38.3	36.0	23.0	3.4	4.5	3,965
Previously Married	62.1	59.0	41.5	40.5	26.7	25.0	2.3	4.3	88
Never Married	74.7	73.5	50.9	48.2	43.4	28.3	5.5	4.0	1,644
Education									
Primary or Less	57.2	56.2	37.0	32.6	28.3	21.2	4.9	7.2	2,519
Secondary	74.3	75.6	49.3	48.3	46.6	27.5	3.5	1.1	2,483
Post-Secondary	90.4	87.4	68.5	70.1	65.5	36.3	0.8	0.3	695
Lifetime No. of Parti	ners								
0	73.6	72.9	51.2	46.7	42.6	28.3	5.7	4.2	1,439
1	63.1	63.0	41.1	39.3	36.2	23.3	3.4	4.3	4,140
2+	70.1	70.5	45.9	45.0	48.7	23.4	1.8	4.4	113
Don't Know	*	*	*	*	*	*	*	*	5

* Percentage not shown when base is less than 25 cases.

Table 14.10 B Percentage of Men Aged 15-49 Who Correctly Reject Misconception About HIV Transmission by Selected Characteristics Reproductive Health Survey: Albania, 2002

Characteristic	Shaking Hands	Public Toilet	Manicure	Kissing	Eating Utensils	Mosquito Bites	Dental/ Surgical	Never Heard of HIV/AIDS	No. of Cases
Total	87.8	77.4	61.9	56.9	41.5	35.0	13.3	3.9	1,740
Strata									
Tirana	88.1	72.1	61.5	53.8	51.6	31.9	9.4	1.6	718
Urban	88.9	83.2	67.0	63.9	47.1	38.0	16.1	1.8	547
Rural	87.0	76.0	58.8	53.9	33.8	34.3	13.1	6.1	475
Age Group									
15–19	88.6	71.9	61.1	52.8	43.8	30.6	13.8	1.6	401
20–24	92.2	83.3	66.0	66.0	47.6	40.0	17.5	0.1	189
25–29	89.9	84.6	65.0	58.8	47.0	39.7	13.0	3.4	218
30–34	86.5	79.2	61.6	53.0	44.7	35.4	9.0	0.5	253
35–39	83.4	77.8	66.1	56.8	35.9	33.3	13.2	6.1	255
40–44	83.5	70.4	55.2	54.6	36.1	31.9	12.2	4.7	277
45-49	90.9	76.5	57.3	57.5	31.3	35.3	14.1	13.3	147
Marital Status									
Currently Married	86.0	76.2	61.4	55.0	37.3	33.5	13.4	5.1	1,023
Previously Married	*	*	*	*	*	*	*	*	14
Never Married	90.4	79.1	62.2	59.8	47.0	37.2	13.3	1.8	703
Education									
Primary or Less	82.9	71.7	55.6	50.2	32.3	29.9	14.2	7.4	689
Secondary	91.4	80.2	66.8	60.6	46.3	38.6	13.8	0.7	825
Post-Secondary	95.4	92.9	69.6	72.4	64.2	43.3	6.4	0.2	226
Lifetime No. of Part	ners								
0	90.3	76.4	62.6	55.4	45.1	37.4	15.5	2.0	446
1	86.7	78.4	55.4	49.9	33.0	37.9	15.3	6.5	448
2+	85.8	78.8	64.7	63.2	44.0	31.1	12.1	1.6	702
Don't Know	94.3	69.1	70.4	57.1	49.8	35.4	3.9	10.7	42

* Percentage not shown when base is less than 25 cases.

ß

Table 14.11 A Percentage of Women Aged 15-44 Who Know HIV Infection Can Be Asymptomatic, And is Not Spread by Dental Treatment or Mosquito Bite, by Selected Characteristics Reproductive Health Survey: Albania, 2002

	-	Know That H	IV is Not Spread by:	-	
Characteristic	Know HIV Can Be Asymptomatic	Mosquito Bites	Dental/ Surgical Treatment	UNAIDS Knowledge Indicator 2**	No. of Cases
Total	55.5	24.8	4.0	0.8	5,697
Strata					
Tirana	62.9	26.5	3.4	0.8	2,108
Urban	62.8	28.9	3.0	0.5	1,816
Rural	49.4	22.1	4.8	0.9	1,773
Age Group					
15–19	55.2	28.5	6.3	1.5	1,094
20–24	60.7	26.9	4.6	1.0	936
25–29	56.7	24.6	3.8	0.7	946
30–34	55.8	23.6	2.9	0.6	1,067
35–39	53.6	19.3	3.7	0.6	958
40–44	50.4	24.5	2.0	0.0	696
Marital Status					
Currently Married	53.9	23.0	3.4	0.5	3,965
Previously Married	58.9	25.0	2.3	0.0	88
Never Married	58.5	28.3	5.5	1.3	1,644
Education					
Primary or Less	45.1	21.2	4.9	0.8	2,519
Secondary	64.1	27.5	3.5	1.0	2,483
Post-Secondary	84.7	36.3	0.8	0.1	695
Lifetime No. of Partners					
0	57.3	28.3	5.7	1.3	1,439
1	54.6	23.3	3.4	0.6	4,140
2+	60.0	23.4	1.8	0.0	113
Don't Know	*	*	*	*	5

* Percentage not shown when base is less than 25 cases.

** Indicator 2 represents the percentage of all women with correct knowledge that HIV could be asymptomatic, and is not spread by

mosquito bites or through medical treatment.

Know That HIV is Not Spread by: UNAIDS Know HIV Can Mosquito Dental/ Knowledge No. of Characteristic **Be Asymptomatic** Bites Surgical Treatment Indicator 2** Cases Total 44.7 33.6 12.8 1.1 1,740 Strata 9.3 Tirana 50.6 31.4 1.7 718 Urban 62.2 37.3 15.8 1.6 547 Rural 31.9 32.3 12.4 0.5 475 Age Group 15 - 1944.4 30.1 13.5 0.7 401 20-24 53.1 40.0 17.5 2.7 189 25-29 53.2 38.4 12.5 1.4 218 30-34 52.1 35.2 8.9 253 0.8 35-39 39.9 31.2 12.4 0.5 255 40-44 40.3 30.5 11.7 0.2 277 45-49 27.1 30.7 12.3 1.6 147 **Marital Status Currently Married** 40.0 31.8 12.8 0.9 1,023 * * * * **Previously Married** 14 Never Married 52.0 36.5 703 13.0 1.4 Education Primary or Less 29.7 27.7 13.1 1.3 689 38.3 Secondary 53.7 13.8 1.1 825 Post-Secondary 82.7 43.3 6.4 0.5 226 Lifetime No. of Partners 0 43.2 36.6 15.2 1.4 446 32.3 35.4 1 14.3 0.7 448 2+ 55.8 30.6 11.9 702 1.3 Don't Know 43.2 31.7 3.5 0.7 42

Table 14.11 B Percentage of Men Aged 15-49 Who Know HIV Infection Can Be Asymptomatic, And is Not Spread by Medical Treatment or Mosquito Bite, by Selected Characteristics Reproductive Health Survey: Albania, 2002

* Percentage not shown when base is less than 25 cases.

** Indicator 2 represents the percentage of all men with correct knowledge that HIV could be asymptomatic, and is not spread by mosquito bites or through medical treatment.

Table 14.12 APercent Distribution of Women Aged 15-44 Who Mentioned Possible Means
of Preventing HIV/AIDS Spontaneously And After Probing
Reproductive Health Survey: Albania, 2002

	Mention	ed			
Possible Means	Spontaneously	Probed	Did Not Mention	Have Not Heard of HIV/AIDS	Total
Stay Faithful to One Partner	35.8	53.9	6.0	4.3	100.0
Use Condoms	30.5	44.2	21.0	4.3	100.0
Sterilize Needles And Syringes	24.1	62.9	8.7	4.3	100.0
Avoid Blood Transfusions	21.6	61.8	12.3	4.3	100.0
Avoid Sex With Prostitutes	14.7	75.6	5.4	4.3	100.0
Limit Number of Sexual Partners	11.6	77.9	6.2	4.3	100.0
Avoid Sharing Razors or Needles	11.4	72.3	12.0	4.3	100.0
Avoid Sex With Bisexuals	10.3	79.3	6.1	4.3	100.0
Abstain From Sex	9.5	36.9	49.3	4.3	100.0
Ask Partner to Be Tested For HIV	9.2	69.9	16.6	4.3	100.0
Do Not Donate Blood	8.8	74.8	12.1	4.3	100.0

Table 14.12 BPercent Distribution of Men Aged 15-49 Who Mentioned Possible Meansof Preventing HIV/AIDS Spontaneously and After ProbingReproductive Health Survey: Albania, 2002

	Mentione	d			
Possible Means	Spontaneously	Probed	Did Not Mention	Have Not Heard HIV/AIDS	Total
Use Condoms	38.3	38.5	19.3	3.9	100.0
Avoid Sex With Prostitutes	34.9	56.9	4.3	3.9	100.0
Stay Faithful to One Partner	18.5	60.8	16.8	3.9	100.0
Avoid Blood Transfusions	13.3	68.6	14.2	3.9	100.0
Sterilize Needles And Syringes	12.3	74.6	9.2	3.9	100.0
Limit Number of Sexual Partners	5.6	77.2	13.3	3.9	100.0
Abstain From Sex	5.3	55.5	35.3	3.9	100.0
Avoid Sharing Razors or Needles	3.9	78.1	14.1	3.9	100.0
Avoid Sex With Bisexuals	3.4	71.6	21.1	3.9	100.0
Do Not Donate Blood	1.9	65.9	28.3	3.9	100.0
Ask Partner to Be Tested For HIV	1.8	74.6	19.7	3.9	100.0

Table 14.13 A Percent of Women Aged 15-44 Who Know Possible Means of Preventing HIV/AIDS Transmission Spontaneously and After Probing, by Selected Characteristics Reproductive Health Survey: Albania, 2002

Characteristic	Avoid Risky Sex	Monogamy Limit Number of Partners	Avoid Blood Donation/ Transfusion	Sterilize Needles/ Avoid Sharing Needles	Ask Partner to Be Tested	Use Condoms	Abstain From Sex	No. of Cases
Total	87.1	85.9	84.6	83.8	79.1	74.7	46.4	5,697
Strata								
Tirana	88.9	86.4	87.0	87.6	83.2	86.2	47.0	2,108
Urban	90.4	89.0	89.5	88.7	84.0	82.0	47.0	1,816
Rural	84.8	84.1	81.3	80.0	75.3	67.4	45.9	1,773
Age Group								
15–19	85.9	84.5	83.5	82.2	76.6	73.4	46.9	1,094
20–24	87.9	85.9	86.5	85.7	79.7	78.5	49.4	936
25–29	90.3	89.9	87.1	85.2	83.4	79.0	46.7	946
30–34	85.8	85.8	84.4	82.0	77.7	74.7	44.9	1,067
35–39	86.9	85.6	82.8	84.7	78.0	73.2	44.4	958
40–44	86.4	84.4	83.7	83.2	75.8	69.6	45.7	696
Marital Status								
Currently Married	87.0	86.4	84.2	83.4	78.2	73.8	45.0	3,965
Previously Married	90.0	87.0	89.2	81.2	80.3	79.3	54.2	88
Never Married	87.1	84.9	85.2	84.7	80.8	76.3	48.7	1,644
Education								
Primary or Less	82.4	81.5	78.9	77.0	72.7	64.8	44.5	2,519
Secondary	92.4	91.2	90.3	90.7	85.8	84.4	50.3	2,483
Post-Secondary	94.0	90.5	95.8	96.1	90.1	95.4	41.0	695
Lifetime No. of Partn	ers							
0	86.8	84.4	84.5	84.0	79.8	75.2	49.4	1,439
1	87.2	86.6	84.6	83.6	78.7	74.3	45.0	4,140
2+	89.0	84.9	89.0	86.4	82.7	84.4	50.5	113
Don't Know	*	*	*	*	*	*	*	5

* Percentage not shown when base is less than 25 cases.

		Reproduct	ive Health S	urvey: Albar	nia, 2002			
Characteristic	Avoid Risky Sex	Sterilize Needles/ Avoid Sharing Needles	Avoid Blood Donation/ Transfusion	Monogamy Limit Number of Partners	Use Condoms	Ask Partner to Be Tested	Abstain From Sex	No. of Cases
Total	77.0	80.9	70.7	79.5	76.8	76.4	60.8	1,740
Strata								
Tirana	80.3	80.5	69.5	75.4	87.0	69.7	59.6	718
Urban	81.5	85.3	76.5	83.7	83.3	80.0	63.7	547
Rural	73.1	78.5	67.7	78.6	69.0	76.9	59.6	475
Age Group								
15–19	73.2	81.2	69.9	75.6	84.2	73.6	61.8	401
20–24	75.0	82.7	74.1	71.3	85.1	74.0	54.6	189
25–29	80.0	86.0	77.2	84.7	86.2	82.4	65.3	218
30–34	85.7	88.3	77.6	85.6	84.7	82.7	64.2	253
35–39	78.6	79.2	66.7	79.7	65.8	78.8	56.9	255
40–44	77.7	79.2	68.9	82.3	65.9	74.3	63.1	277
45-49	69.5	67.8	59.2	79.0	61.0	69.2	59.8	147
Marital Status								
Currently Married	77.4	79.5	68.9	82.4	70.9	77.2	61.8	1,023
Previously Married	*	*	*	*	*	*	*	14
Never Married	76.3	83.2	73.4	75.4	86.3	75.2	59.5	703
Education								
Primary or Less	73.8	75.9	65.6	78.0	68.0	74.4	60.3	689
Secondary	78.7	84.5	74.3	80.8	83.4	78.1	60.8	825
Post-Secondary	86.3	91.1	80.7	81.3	93.1	79.0	64.0	226
Lifetime No. of Partn	ners							
0	74.8	80.5	71.6	76.1	82.3	74.0	63.2	446
1	70.6	80.7	63.0	82.5	61.2	77.1	67.5	448
2+	82.8	81.6	73.7	81.5	86.2	76.9	55.6	702
Don't Know	80.6	80.0	82.9	68.5	74.3	79.0	52.8	42

Table 14.13 BPercent of Men Aged 15-49 Who Know Possible Means of Preventing HIV/AIDS TransmissionSpontaneously and After Probing, by Selected CharacteristicsReproductive Health Survey: Albania, 2002

 * Percentage not shown when base is less than 25 cases.

20

0		
D		
5		
-		
-		
D)		
5		
D		
-		
-		
U		
2		
4		
2		
-		
-		
-		
5		
5		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
3		
REALED THE PLACE TO		
analy nue cite to a		
analy nue cite to a		
analy nurver of the structure of the str		
analy on a la and who we are a		
Relice of a LIS alla Milowieus		
analy on a la and who we are a		
Relice of a LIS alla Milowieus		
helielice of alla vilomied		
Relice of a LIS alla Milowieus		
helielice of alla vilomied		
alla Experience of 2115 alla Milowied		
alla Experience of 2115 and Milowied		
alla Experience of 3113 and Miowied		
alla Experience of 2115 and Milowied		
alla Experience of 2115 and Milowied		
alla Experience of 2115 and Milowied		
alla Experience of 2115 and Milowied		
alla Experience of 2115 and Milowied		
alla Experience of 2115 and Milowied		
alla Experience of 2115 and Milowied		
alla Experience of 2115 and Milowied		

Table 14.14a Percent of Women Aged 15-44 Who Believe HIV Can Be Prevented By Limiting Number of Sexual Partners, Being Monogomous, And Using Condoms, by Selected Characteristics Reproductive Health Survey: Albania, 2002

Characteristic	Monogomy	Limit Number of Sexual Partners	Condoms	UNAIDS Knowledge Indicator 1*	No. of Cases
Total	89.6	89.5	79.0	72.6	5,697
Strata					
Tirana	91.1	92.4	89.2	84.3	2,108
Urban	92.6	92.9	83.9	79.6	1,816
Rural	87.6	86.8	73.4	65.4	1,773
Age Group					
15–19	87.3	86.6	77.8	70.9	1,094
20–24	90.4	90.6	81.4	76.3	936
25–29	92.2	92.5	81.9	77.1	946
30–34	89.3	88.6	80.6	72.6	1,067
35–39	90.6	90.3	77.0	70.9	958
40–44	88.7	89.1	75.6	67.9	696
Marital Status					
Currently Married	90.4	90.1	78.3	72.1	3,965
Previously Married	92.8	92.2	83.6	76.9	88
Never Married	87.8	88.0	80.2	73.3	1,644
Education					
Primary or Less	85.5	84.3	72.0	62.5	2,519
Secondary	94.4	95.0	85.5	82.4	2,483
Post-Secondary	94.7	97.9	95.6	93.8	695
Lifetime No. of Partne	rs				
0	87.7	87.4	79.4	72.2	1,439
1	90.5	90.3	78.6	72.5	4,140
2+	85.2	91.4	88.8	81.1	113
Don't Know	**	**	**	**	5**

* Indicator 1 represents the percentage of all women who identify monogomy, condom use and partner limitation as prevention measures against HIV.

** Percentage not shown when base is less than 25 cases.

Table 14.14 B

Percent of Men Aged 15-49 Who Believe HIV Can Be Prevented By Limiting Number of Sexual Partners, Being Monogamous, And Using Condoms, by Selected Characteristics Reproductive Health Survey: Albania, 2002

				UNAIDS	
Characteristic	Monogamy	Limit Number of Sexual Partners	Condoms	Knowledge Indicator 1*	No. of Cases
Total	79.2	82.8	80.7	68.8	1,740
Strata					
Tirana	80.3	81.4	88.6	76.0	718
Urban	83.0	86.3	85.2	74.5	547
Rural	76.6	81.2	75.0	62.6	475
Age Group					
15–19	73.0	77.3	85.8	71.2	401
20–24	69.3	75.7	85.3	67.8	189
25–29	81.3	86.8	89.6	81.1	218
30–34	85.2	89.2	85.2	78.4	253
35–39	80.0	85.6	71.9	61.4	255
40–44	86.1	85.4	70.7	61.8	277
45-49	83.2	81.8	74.2	58.3	147
Marital Status					
Currently Married	84.2	85.6	76.0	66.7	1,023
Previously Married	**	**	**	**	14
Never Married	71.5	78.4	88.2	72.3	703
Education					
Primary or Less	78.9	80.6	75.4	62.4	689
Secondary	79.4	84.4	84.1	73.3	825
Post-Secondary	80.6	86.5	93.3	82.3	226
Lifetime No. of Partne	ers				
0	74.1	78.4	84.3	71.4	446
1	88.1	83.3	67.7	58.4	448
2+	76.8	86.2	87.9	76.4	702
Don't Know	73.1	77.9	85.2	64.1	42

* Indicator 1 represents the percentage of all men who identify monogamy, condom use and partner limitation as prevention measures against HIV.

 ** Percentage not shown when base is less than 25 cases.

Tables

	Table 14.15 A Percent Distribution of Women Aged 15-44 by Self-Perceived Risk of Contracting HIV/AIDS by Selected Characteristics Reproductive Health Survey: Albania, 2002									
Characteristic	Great Risk	Moderate Risk	Little Risk	No Risk	Don't Know	Never Heard of HIV/AIDS	Total	No. of Cases		
Total	1.2	1.8	11.4	78.5	2.9	4.3	100.0	5,697		
Strata										
Tirana	1.3	2.5	10.8	80.0	2.5	3.0	100.0	2,108		
Urban	1.0	2.1	11.8	80.6	2.7	1.9	100.0	1,816		
Rural	1.2	1.4	11.3	77.0	3.1	6.0	100.0	1,773		
Age Group										
15–19	1.3	1.5	10.4	79.5	3.0	4.3	100.0	1,094		
20–24	1.4	2.1	10.2	80.9	2.5	2.9	100.0	936		
25–29	1.5	1.1	10.3	81.4	2.9	2.9	100.0	946		
30–34	1.5	2.3	11.9	74.6	3.9	5.9	100.0	1,067		
35–39	0.8	2.0	13.6	77.1	2.7	3.9	100.0	958		
40–44	0.6	1.6	12.2	77.2	2.4	6.1	100.0	696		
Marital Status										
Currently Married	1.3	1.8	11.7	77.4	3.3	4.5	100.0	3,965		
Previously Married	0.0	4.2	9.3	79.7	2.5	4.3	100.0	88		
Never Married	1.1	1.5	10.7	80.7	2.1	4.0	100.0	1,644		
Education										
Primary or Less	1.1	1.6	10.0	76.2	3.9	7.2	100.0	2,519		
Secondary	1.2	1.6	12.2	82.0	1.8	1.1	100.0	2,483		
Post-Secondary	1.4	3.1	15.9	77.9	1.4	0.3	100.0	695		
Lifetime No. of Partne	ers									
0	1.2	1.4	10.6	80.7	1.9	4.2	100.0	1,439		
1	1.2	1.9	11.3	78.0	3.3	4.3	100.0	4,140		
2+	2.3	2.7	24.8	64.2	1.6	4.4	100.0	113		
Don't Know	*	*	*	*	*	*	*	5		

* Percentage not shown when base is less than 25 cases.

Table 14.15 B Percent Distribution of Men Aged 15-49 by Self-Perceived Risk of Contracting HIV/AIDS by Selected Characteristics Reproductive Health Survey: Albania, 2002								
Characteristic	Great Risk	Moderate Risk	Little Risk	No Risk	Don't Know	Never Heard of HIV/AIDS	Total	No. of Cases
Total	0.7	1.2	8.4	78.8	7.0	3.9	100.0	1,740
Strata								
Tirana	0.2	0.8	14.6	75.9	6.8	1.6	100.0	718
Urban	0.9	0.4	9.1	80.9	6.9	1.8	100.0	547
Rural	0.8	1.9	5.4	78.7	7.2	6.0	100.0	475
Age Group								
15–19	0.0	0.6	5.9	87.1	4.8	1.6	100.0	401
20–24	1.4	3.6	13.6	72.8	8.5	0.2	100.0	189
25–29	1.5	1.9	9.1	78.8	5.3	3.4	100.0	218
30–34	0.4	1.8	11.2	78.0	8.1	0.5	100.0	253
35–39	1.0	0.7	7.8	78.0	6.4	6.1	100.0	255
40–44	0.6	0.2	7.6	75.3	11.6	4.7	100.0	277
45-49	0.0	0.0	3.0	78.8	4.9	13.3	100.0	147
Marital Status								
Currently Married	0.6	0.6	7.4	78.4	7.9	5.1	100.0	1,023
Previously Married	*	*	*	*	*	*	*	14
Never Married	0.8	2.2	10.0	79.7	5.6	1.8	100.0	703
Education								
Primary or Less	1.1	0.8	4.9	78.6	7.2	7.4	100.0	689
Secondary	0.3	1.8	9.4	80.5	7.3	0.7	100.0	825
Post-Secondary	0.6	1.0	21.4	71.5	5.2	0.3	100.0	226
Lifetime No. of Partne	ers							
0	0.4	2.4	4.1	87.6	3.5	2.0	100.0	446
1	0.5	0.0	0.6	88.2	4.2	6.5	100.0	448
2+	1.2	1.7	16.6	70.9	8.0	1.6	100.0	702
Don't Know	0.0	0.0	11.6	52.5	25.1	10.8	100.0	42

* Percentage not shown when base is less than 25 cases.

Tables

Table 14.16 A Opinion About The Main Risk Factor of Contracting HIV/AIDS Among Women 15-44 Who Have Heard About HIV/AIDS And Believe They Have Any Risk of Contracting HIV/AIDS Reproductive Health Survey: Albania, 2002

Main Risk Factors	Percent Distribution
Medical/Dental Treatment	90.8
Does Not Trust Partner	4.9
Received Blood Transfusion	2.0
Manicure/ Haircut	0.7
Used IV Drugs/ Shared Needles	0.4
Unprotected Sex With Casual Partner	0.2
Many Sexual Partners/ Trade Sex For Money	0.0
Other	0.4
Don't Know	0.7
Total	100.0
Total Number of Cases	827

Table 14.16 B Opinion About The Main Risk Factor of Contracting HIV/AIDS Among Men 15-44 Who Have Heard About HIV/AIDS And Believe They Have Any Risk of Contracting HIV/AIDS Reproductive Health Survey: Albania, 2002

Main Risk Factors	Percent Distribution
Medical/Dental Treatment	54.4
Unprotected Sex With Casual Partner	16.2
Used IV Drugs/ Shared Needles	7.6
Many Sexual Partners/ Trade Sex For Money	7.5
Manicure/ Haircut	4.8
Does Not Trust Partner	3.8
Received Blood Transfusion	1.1
Other	0.0
Don't Know	4.6
Total	100.0
Total Number of Cases	203

bles

Table 14.17 A

Opinion About The Main Factor That Protects From Contracting HIV/AIDS Among Women Aged 15-44 Who Have Heard of HIV/AIDS And Believe That They Have No Risk of Contracting HIV/AIDS, by Selected Characteristics Reproductive Health Survey: Albania, 2002 (Percent Distribution)

Characteristic	Monogamy	Not Sexually Active	Trust Partner	Do Not Share Needles	Use Condoms	Do Not Get/Need Transfusion	Other	Don't Know	Total	No. of Cases
Total	40.5	31.5	23.0	0.8	0.8	0.6	0.0	2.8	100.0	4,515
Strata										
Tirana	45.6	29.8	20.1	0.5	1.7	0.9	0.0	1.4	100.0	1,690
Urban	42.6	28.6	24.3	0.5	1.0	0.7	0.2	2.1	100.0	1,458
Rural	37.8	33.7	23.1	1.1	0.4	0.4	0.0	3.5	100.0	1,367
Age Group										
15–19	8.6	81.0	5.7	1.5	0.2	0.5	0.0	2.5	100.0	875
20–24	33.8	43.8	17.3	0.3	1.1	0.8	0.0	2.9	100.0	766
25–29	45.7	18.5	30.1	1.2	1.4	0.4	0.0	2.7	100.0	760
30–34	54.3	10.0	30.5	0.6	1.2	0.5	0.4	2.5	100.0	813
35–39	58.3	6.1	29.6	0.2	0.6	0.5	0.0	4.7	100.0	744
40–44	56.4	8.3	32.2	1.0	0.4	0.6	0.0	1.1	100.0	557
Marital Status										
Currently Married	60.6	0.4	34.3	0.5	0.8	0.4	0.0	3.0	100.0	3,123
Previously Married	5.8	82.6	1.7	6.6	0.8	1.7	0.0	0.8	100.0	72
Never Married	4.4	87.6	2.8	1.1	0.7	0.7	0.1	2.6	100.0	1,320
Education										
Primary or Less	40.9	31.1	22.6	0.7	0.2	0.3	0.0	4.2	100.0	1,947
Secondary	39.6	32.2	24.2	1.1	0.8	0.8	0.0	1.3	100.0	2,047
Post-Secondary	42.2	31.3	19.5	0.9	4.2	0.9	0.4	0.6	100.0	521
Lifetime No. of Partne	ers									
0	1.1	94.3	0.1	1.2	0.0	0.7	0.1	2.5	100.0	1,166
1	58.3	3.3	33.2	0.7	1.1	0.5	0.0	2.9	100.0	3,276
2+	47.1	14.3	32.9	0.0	4.3	1.4	0.0	0.0	100.0	70
Don't Know	*	*	*	*	*	*	*	*	*	3

* Percentage not shown when base is less than 25 cases.

Table 14.17 B

Opinion About The Main Factor That Protects From Contracting HIV/AIDS Among Men Aged 15-49 Who Have Heard of HIV/AIDS And Believe That They Have No Risk of Contracting HIV/AIDS, by Selected Characteristics Reproductive Health Survey: Albania, 2002 (Percent Distribution)

Characteristic	Monogamy	Not Sexually Active	Trust Partner	Use Condoms	Do Not Get/Need Transfusion	Do Not Share Needles	Total	No. of Cases
Total	47.6	26.7	12.8	8.4	3.1	1.4	100.0	1,371
Strata								
Tirana	50.3	21.3	10.7	14.1	1.7	1.9	100.0	553
Urban	42.6	24.2	17.5	11.2	3.6	0.9	100.0	439
Rural	49.6	30.1	10.7	4.7	3.3	1.6	100.0	379
Age Group								
15–19	0.6	90.6	0.6	1.5	5.0	1.6	100.0	341
20–24	19.9	43.9	6.1	22.9	7.2	0.0	100.0	135
25–29	52.0	7.1	19.4	17.7	1.8	2.0	100.0	171
30–34	64.9	3.1	18.4	8.8	2.9	1.9	100.0	195
35–39	73.8	1.2	17.9	5.4	0.0	1.7	100.0	202
40–44	74.2	0.8	18.0	3.3	1.9	1.8	100.0	211
45-49	76.9	2.3	16.0	2.7	1.4	0.7	100.0	116
Marital Status								
Currently Married	74.3	1.3	18.1	3.5	1.4	1.4	100.0	801
Previously Married	*	*	*	*	*	*	*	11
Never Married	7.5	65.1	4.8	15.6	5.5	1.5	100.0	559
Education								
Primary or Less	47.9	30.8	12.6	4.1	2.6	2.0	100.0	545
Secondary	47.0	25.2	12.3	11.0	3.6	0.9	100.0	666
Post-Secondary	49.7	9.2	15.9	20.7	3.5	1.0	100.0	160
Lifetime No. of Partner	rs							
0	1.0	93.1	0.0	0.1	4.7	1.1	100.0	390
1	89.8	0.7	3.9	2.3	2.5	0.8	100.0	401
2+	40.7	2.5	32.1	19.9	2.6	2.2	100.0	502
Don't Know	65.7	3.1	10.9	17.2	1.0	2.1	100.0	78

* Percentage not shown when base is less than 25 cases.

Chapter 15

VIOLENCE AGAINST WOMEN

Introduction

Physical and sexual violence against women is increasingly recognized as a global problem that can affect physical and mental health. The adoption of Resolution WHA49.25 by the World Health Organization drew international attention to the potential consequences of violence on the impact on the delivery of health care services (WHO, 1987). The resolution also called on member nations to establish data systems that could document the dimensions of the problem of violence.

Violence against women, also known as "gender-based violence," encompasses a wide variety of acts and behaviors, including verbal, physical, and sexual violence, but also includes restriction of access to food, health care, or economic assets, female genital cutting, and other forms of violence across the life cvcle. Women in all cultures experience violence, and although some indicators such as poverty and lack of education are often associated with higher levels of violence, experience of violence is reported by women of all socioeconomic and educational levels. Often cited as potential contributors to the cultural acceptability of violence against women are gender norms and stereotypes, women's economic dependence on men, lack of legislation or loose enforcement of existing laws, and lack of preventive activities that support the development of alternatives to physical violence in resolving conflict or anger (WHO, 2001).

The data presented in this report represent the first national population based data available on the issue of violence against women in Albania. The lack of data on domestic violence at the national level in Albania has hampered the development of program services and strategies (Haxhiymeri et al., 2000). Reliable data on violence in general and specifically about intimate partner violence are very scarce. Police departments, legal offices and health care centers seldom record such type of data and the Ministry of Public Order lacks accurate statistics on the ratio between different types of crimes. Both the Ministry of Public Order and the Ministry of Justice categorize domestic crime simply as crimes of one person against another. There are no separate entries and no further break down of statistics to identify intimate partner violence (Refleksione, 2000; Haxhiymeri et al., 2000). Mortality data on fatalities are not accurate and information by geographic areas or age groups is incomplete.

Since the late 1990's, Refleksione. Albanian women's an association. and other researchers have conducted several investigations of the issue of violence against women in Albania on a limited scale. One study, conducted in selected districts of Albania, found that 39% of female participants had experienced physical abuse and 25% of them experienced emotional abuse (Miria, 1996). In another community study, sexual abuse was reported by 9% to 23% of women (Miria et al., 2000). Another local study found that 46% of women living in rural areas compared with 36% of women living in urban areas experienced physical abuse by their partners while 28% of women living in the rural areas reported sexual abuse compared to 36% of women living the urban areas (Kaci et al., 1996). However, this was not a population-based study. Some other qualitative studies have examined the characteristics of victims of domestic violence in Albania (Van Hook et al., 2000; Co-PLAN, 2001; Baban et al., 2003). A women's center based in Tirana since the late 1990s has monitored the press as one of its main activities. The monitoring process of 7 newspapers in 2001 made possible the review of 1,130 articles and 207 (18.3%) were about domestic violence, while in 2002 the monitoring process reviewed 1,244 articles with domestic violence accounting for 9.9% of them (Gjermeni et al., 2003).

Domestic violence is still being treated as a private problem and not a societal problem. Gender based violence is still not considered as a health problem and health workers are not trained on this issue. Community based work to address the problem of domestic violence is almost not existing.

Limited services for victims of abuse exist, generally sponsored by NGOs or other civil society organizations. These include counseling centers and women's shelters mainly in Tirana. Training programs designed to raise awareness among the police, social workers, and the communities are very rare (Refleksione, 2000; Haxhiymeri et al., 2000). The services that exist function in the absence of adequate legal protections for victims of violence. Only 5% of cases of domestic violence asked for legal protection according to the women's legal office (Mecaj, 1997).

The Albanian civil code does not address the issue of domestic violence and the Albanian criminal code offers little support to victims, who must file a complaint and prepare their own case; but, often, cases of domestic violence are not treated as criminal cases. When cases are withdrawn, no legal provisions exist to pursue prosecution. Activists have pointed out that revisions of the penal code and civil procedures in Albania are necessary in order to adequately address the manner in which domestic cases are addressed (Van Hook et al., 2000).

Data on violence against women can be of critical use in drawing local and national attention to the problem and the need to strengthen services and preventive efforts to address it. In several countries, survey data on violence have been instrumental in the establishment or reinforcement of laws that protect women and children. The data also can be used to educate providers of health care services, law enforcement or legal services, and social services. Moreover, national level data can be an important key to understanding the dimensions of the problems of intimate partner violence (IPV) and sexual violence in Albania. They can be used to identify individual characteristics and risk factors for violence, including the association of violence and key indicators of women's reproductive health or women's health in general.

Scientific investigation of the dimensions of violence against women is relatively new, and questions on violence against women in population-based surveys only became common in the mid-1990s. Collection of data in Albania followed

guidelines set forth by the World Health Organization regarding the ethical implementation of survey research on violence against women (World Organization, 2001). WHO Health guidelines recommend the adoption of methodological strategies to minimize underreporting of violence by survey participants. Similar to other national, population-based surveys in Eastern Europe (Serbanescu et al., 1995, 1998, 2001, 2003) and in other parts of the world (Kishor, 2004), the Albania Reproductive Health Survey used a modified Conflict Tactics Scale (Straus and Gelles, 1979) to collect data on specific acts of verbal, physical, and sexual violence. Research has indicated that asking people about specific acts (e.g., slapping, throwing an object, hitting with a fist) can help avoid underreporting due to cultural differences in what is considered an "abusive" or "violent" act.

The WHO guidelines include training for interviewers and other field staff on the topic of violence against women. In addition to an in-depth orientation to the topic, field staff also was trained in specific techniques for the suspension of questions regarding violence in cases where absolute privacy is not possible during the interview, and measures to be followed when a participant disclosed abuse.

Despite these and other efforts to minimize underreporting, researchers have generally assumed that due to factors such as stigma, fear of reprisals, or cultural norms, violence data are underreported in population-based surveys. The degree of underestimation in survey research is thought to vary across countries and within countries due to culture and religion, and likely depends on the cultural acceptability of discussing or reporting violence (Kishor S and Johnson K, 2004.) The data in this report, therefore, should be interpreted with caution as they might reflect a lower estimate of the true prevalence of violence in Albania.

In Albania, the Reproductive Health Survey focused on three distinct aspects of violence against women: 1) history of violence in childhood; 2) verbal, physical or sexual violence inflicted by current or former intimate partners, here termed "intimate partner violence (IPV)", also often referred to as "domestic violence"; and 3) women's experience of forced sexual intercourse, or sexual coercion. In addition to asking reproductive age women about their experience of these three types of violence, Albanian men were also asked about their history of violence during childhood, and their perpetration of verbal, physical, or sexual violence against a current or former intimate partner. The male data from this survey documents the male perpetration of domestic violence in Albania for the first time.


Comparison of Violence across Countries of Eastern Europe

The questions asked of women in Albania are comparable to questions that have been asked in reproductive health surveys in a number of other countries of Eastern Europe and the Former Soviet Union (FSU). Culturally and legally, Albania shares with other countries of the region some common characteristics regarding the legal status of women and gender norms (CDC and ORC Macro, 2003). Figure 15.1 shows substantial variation in reports of lifetime and past year physical violence according to country. Reported lifetime abuse ranges from 29% in Romania to 5% in the Republic of Georgia. Albanian women reported prevalence of lifetime violence on the lower end of that range, at 8%. Similarly, reported physical violence during the past year ranged from a high of 10% in Romania, to a low of 2% in Georgia. Again, Albanian women reported prevalence of physical violence during the past year just above that reported by Georgian women (5%).

History of Witnessing or Experiencing Abuse during Childhood

Having witnessed violence in the home during childhood is one of the most noted risk factors for violence as an adult (Hoteling and Sugarman, 1986). In Albania, both women and men were asked whether, during their childhood or adolescence, they ever saw or heard their parents or step-parents physically abuse each other. Respondents were also asked whether, as a child, they were ever beaten or physically mistreated by anyone in their family.

Among Albanian women aged 15–44, 12% reported having witnessed parental abuse, and 27% reported having received physical abuse themselves during childhood (Table 15.1A). Prevalence of both indicators was greater among residents of rural areas compared to urban residents. Prevalence of both having witnessed abuse and having been physically abused was greater among women with four or more children, lower education, lower socioeconomic status,




and among those who did not work for wages. For example, 20% of women with a post-secondary education reported having been physically abused as a child, compared to 31% of women with an education level of primary or less.

Albanian men aged 15-49 reported significantly higher levels of both indicators compared to women (Table 15.1B and Figure 15.2). Overall, 47% of men reported having witnessed parental violence compared to 12% of women, and 67% of men reported having been beaten or physically mistreated themselves as children, compared to 27% of women. While it might be expected that having experienced physical abuse would vary by sex, the difference in reports of men and women of having witnessed parental abuse, is quite striking and suggests potential underreporting on the part of women. Also, according to Albanian tradition, it is

said that young men should endure more hardship than young women.

Similar to women, among Albanian men variations in the two indicators by individual characteristics revealed higher prevalence both of having witnessed abuse and having experienced physical abuse among rural residents compared to urban (Table 15.1B). Among men, there was no significant difference by age. As with women, men who reported a higher number of living children, lower education, and lower socioeconomic status also reported higher levels of both indicators of violence during childhood. For example, among men with primary education or less, 52% reported having witnessed abuse and 72% reported having experienced abuse, compared with 27% and 44%, respectively, among men with post-secondary education. Employment



status was not associated with either indicator for men.

The data indicate that having witnessed violence in the home is highly associated with having been abused for both men and women. Among women who witnessed physical violence between their parents, 81% experienced physical abuse themselves as a child, compared to only 19% among women who reported that they did not witness physical abuse between their parents. For men, a similar strong association was found between having witnessed parental abuse and having experienced physical abuse as a child. Whereas only 10% of men who did not witness physical violence between parents experienced physical abuse themselves as a child, 90% of those who witnessed parental physical violence reported that they themselves were also physically abused.

Women's Experience of Intimate Partner Violence

Intimate partner violence (IPV) constitutes one aspect of gender-based violence, and is defined as verbal, physical, or sexual violence that occurs between current or former intimate partners, including husbands and wives, members of a consensual union, or in casual intimate relationships. In Albania, the definition of intimate partner violence was restricted to that occurring between current or former married partners or members of a consensual union with cohabitation. As noted in the introduction. IPV in Albania was measured using a modified Conflict Tactics Scale (CTS) that asks about specific acts or behaviors. In Albania, the modified CTS inquired about specific acts of verbal violence or abuse, physical violence, and coercive sex as outlined in Figure 15.3.

Type of Violence From Intimate Partner	CTS Element					
Verbal	• Insulted you or swore at you?					
	• Threatened to hurt you or someone you care about?					
Physical	• Pushed you, shook you, shoved you, or threw something at you?					
	• Slapped you or twisted your arm?					
	• Hit you with his fist or with something else?					
	• Threatened you with a knife or weapon?					
	• Kicked you, choked you, or beat you up?					
Sexual	• Physically forced you to have sexual relations even though you did not want to?					

Figure 15.3 Questions asked using modified Conflict Tactics Scale (CTS)


For each element, women were asked whether the type of violence had ever occurred. Respondents who answered that they had experienced a specific type of violence were then asked when their most recent experience had happened.

For data analysis, responses were combined and categorized as shown in Figure 15.4 into verbal, physical, or sexual violence. Responses were also categorized into two timeframes: lifetime experience of verbal, physical, or sexual violence; and experience of the three types of violence within the past year, as an estimate of the current magnitude of the problem.

Table 15.2A shows the reported prevalence of violence occurring in the women's lifetime and in the past year to Albanian women aged 15–44 who were ever married or in union. Overall,

30% of Albanian women reported ever having experienced verbal abuse from an intimate partner, and 23% reported having experienced verbal abuse during the past year. Lifetime physical violence was reported by 8% of women, with 5% reporting physical violence during the past year. Finally, 3% of Albanian women reported ever having been forced to have unwanted sex by a husband or partner, with 2% reporting unwanted sex during the past year.

Table 15.2A also shows data on violence during women's lifetime and during the past year disaggregated by selected individual characteristics. Little variation was found according to place of residence. When women's current age was taken into account, little variation was found in lifetime prevalence of the three types of violence. For violence during the past year, reports of verbal abuse were lowest



among women aged 20-24 (18%) compared to women aged 15-19 and 30-34 (25% and 25%, respectively). Reported physical violence in the past year was highest among 15–19 year olds (11%) compared to other age groups. No variation by age was found in reported sexual violence during the past year. Previously married women reported significantly higher percentages of lifetime violence compared to currently married women (see also Figure 15.4). For example, whereas 31% of previously married women reported physical violence at some time in their lives by a husband or partner, this was reported by 8% of women who were currently married or in union (Figure 15.4). Likewise, 19% of women previously married or in union reported ever having experienced forced sex by a husband or partner compared to 2% of currently married women. This pattern was not seen for violence during the past year; in fact, a greater proportion of currently married women reported verbal abuse during the past year compared to previously married women. Presumably, this pattern is affected by exposure to the risk of violence, with previously married women having less recent exposure than currently married women as they have been separated or divorced in the past year. Also, as the culture of silence surrounds intimate partner violence, a sense of freedom of expression might be experienced by divorced or separated women.

Reported violence in the two time frames varied somewhat by other characteristics of Albanian women. Higher parity was associated with greater proportions of lifetime and current verbal abuse; lower levels of education were associated with greater prevalence of lifetime and past year verbal and physical violence; and lower socioeconomic status was significantly associated with higher prevalence of lifetime and past year verbal abuse. Little variation was found according to whether women were currently employed. This might be explained by the results of one study in Albania showing that the financial contribution of women does not increase or decrease their status inside the family (Baban et al., 2003)

Table 15.3A shows women's reports of individual acts of physical violence from a current or past husband or partner in their lifetime and in the past year. Acts of physical violence are classified into "moderate" and "severe" forms of physical violence. Overall, women reported moderate forms of physical violence more commonly than severe forms, regardless of women's characteristics. Slapping or twisting the arm was the single most commonly reported act of physical violence. Among the severe forms of physical violence, hitting with a fist or with something else and kicking, choking, or beating were reported more commonly than threatening with a knife or weapon. Previously married women reported the highest prevalence of all acts of lifetime physical violence. No individual characteristics stood out as strongly associated with individual types of violence in the past year.


Male Infliction of Intimate Partner Violence

Albanian men also responded to the same elements of the CTS, but were asked about their experience of having inflicted the different types of violence on a current or former wife or partner. While evidence suggests that women also direct physical violence at men, most research has shown that men inflict violence against women far more often than the reverse (WHO, 2001). Table 15.2B shows that overall, 33% of men reported ever having committed verbal abuse against an intimate partner, 14% reported have inflicted physical abuse, and 1% admitted having forced unwanted sex. During the past year, 19% of men reported inflicting verbal abuse, 5% physical violence, and 0.3% sexual violence against a current or former partner.

As shown in Figure 15.5, women's reports of violence received and men's reports of violence inflicted were similar. Reports of lifetime verbal abuse did not differ significantly, and were only slightly higher among women than men in the past year. Women did report a lower proportion of lifetime physical violence (8%) compared to the proportion of men who reported having inflicted violence (14%), However, the two sexes reported the same level of physical violence received and inflicted during the past year. Finally, slightly higher proportions of women reported forced sex both in their lifetime and in the past year compared to reported infliction of forced sex by men. It is important to note when interpreting these data that the female and male samples were independent and

therefore do not represent couple-level data. Nevertheless, the notable finding that a greater proportion of men reported infliction of physical violence compared to the proportion of women who reported receiving physical violence during their lifetime could point to reluctance on the part of women to report such incidents to a survey interviewer.

Table 15.2B shows that, among Albanian men, having perpetrated violence was not strongly associated with residence, similar to women. Men's reports of having inflicted verbal abuse or physical violence in their lifetime increased with men's age, as did recent physical violence (with the exception of men aged 20-24 likely due to small sample size in this age group (n=33). Men's reports of ever having perpetrated verbal and physical violence and of infliction of violence during the past year increased with parity and was inversely related to education and socioeconomic status, a pattern similar to that reported by women.



In Table 15.3B, men's reports of specific acts of physical violence, categorized into "moderate" and "severe" violence, are similar to the responses of women with the exception of a greater proportion of "moderate" violence reported to have been inflicted by men at some time in their lives, compared to that reported by women. Whereas 5% of women reported that a partner had pushed, shaken, shoved or thrown something at her, this act was reported by 9% of men. Likewise, 8% of women reported having been slapped or having had her arm twisted, compared to a reported 13% of men saying they had inflicted this type of physical violence. Women's and men's reports of severe physical violence during over lifetime and of specific acts of physical violence during the past year did not differ significantly.

Regarding the association of selected characteristics of men with infliction of specific acts of physical violence, rural residents reported somewhat greater lifetime levels of having pushed, shook, or thrown something at a partner compared to urban residents. Also among men, reports of having inflicted moderate violence over the lifetime increased with age, parity, lower educational level, and low socioeconomic status. For past year violence, reported infliction of moderate violence was slightly higher among men with higher parity, lower education, and low socioeconomic status. These associations were not found for severe acts of violence.

Characteristics of Physical or Sexual Violence Experienced by Women in the Past Year

Women who experienced physical or sexual violence during the past year were asked how many episodes of violence had occurred during the past year, and if they had any physical injuries as a result of the violence.

This data is presented by type of abuse in Table 15.4. From one-third (36%) to onehalf (52%) of women suffering physical or sexual abuse in the past year reported that the abuse happened at least four times. Another one-fifth (20%) to almost onethird (28%) said they did not remember how many times the abuse occurred which may indicate multiple episodes. Among women who reported physical violence in the past year, 44% reported swelling, bruises, cuts or other physical injury.

Women who had experienced abuse during the past year were asked whether they talked to anyone about the violence (Table 15.5). Just over half the abused women (54%) responded that they did not talk to anyone. A greater proportion of residents of rural areas reported that they did not talk to anyone compared to urban residents, particularly those in Metro Tirana. Also less likely to have talked to someone were women with incomplete secondary education or less, and those with low socioeconomic status. Whereas 47% of women with medium or high socioeconomic status reported not talking to someone, this was true of 61% of women with low socioeconomic status.

Women who reported having talked to someone about the physical or sexual violence abuse identified most frequently family members or friends as the person(s) to whom they turned (Table 15.6), with 52% responding that they talked to their mother, 39% stating that they talked to their mother, 39% stating that they talked to a member of their husband's family, and 32% reporting that they turned to another relative. Only 10% reported that they went to the police, 8% talked to a doctor, and 3% talked to a legal advisor. When the data were stratified by individual characteristics, denominators were too small to detect significant differences. Generally, rural residents and older women reported with greater frequency that they talked to their mother or another family member compared to urban and younger residents, who reported more often talking to a friend.

Women who reported that they did not talk to a doctor/medical worker, the police, or a legal advisor about the violence were asked the main reason why they did not (Table 15.7). The most common responses included one third of the women (33%) who reported that they felt it would be of no use or would not do any good and 28% who reported that violence is normal and there was no need for her to complain. Other responses included that the woman did not know where to seek help (9%), fear of divorce or ending the relationship (7%), and fear of more beatings or other retaliation (7%). Reasons given by residence were similar but rural women were somewhat more likely to say that "they did not know where to seek help." The same was true of women classified "low socioeconomic status." The as results presented here reiterate the need of improving legislation and strengthening the training of police, legal advisers and health care workers. Also information. education, communication activities on IPV should be promoted.

Women's Attitudes and Beliefs about the Effects of Violence

Women who had ever experienced physical or sexual violence were asked questions about the violence and how they believed it affected them or their children. First, women were asked to tell the interviewer about the circumstances surrounding the violence. Situations commonly identified by women, shown in Table 15.8, included the partner being drunk (42%), financial troubles (41%) or when the husband/ partner was unemployed (37%); when there were "family problems" (21%) or when the husband/partner was jealous (32%).

Women who had ever experienced physical or sexual violence were also asked whether they believed the incidents affected their health. Almost half the women (47%) (data not shown) agreed that they had experienced some health effect of violence, with commonly cited affects being sleep disorders, depression, low self-esteem, and physical injuries. Physically or sexually abused women who had living children were asked if they thought the children were affected by the violence. Almost one third (31%) responded that they did think the children were affected. Effects on children most commonly cited by women included having witnessed the violence, living in fear, and decreased learning abilities of the children (data not shown).

All women who experienced any form of abuse were also asked whether they would want to know the hotline number in Metro Tirana where they could ask advice about domestic violence. Overall, 12% wanted to know the hotline number, with some differences by residence, age group, education, and socioeconomic status of the woman (Table 15.9).

All women were also asked their attitudes about situations in which it might be justifiable for a man to hit or beat his wife (Table 15.10). Most common agreement in urban areas was if she asked if he had other girlfriends (47%), she neglects the children (23%) and she dresses too sexy (18%). In rural areas, the top three areas of agreement were the same but at a higher level; asking if he has other girlfriends (69%), she neglects the children (42%), she looks too "sexy" (34%) or she goes out without telling him (34%). The same pattern was seen by level of education. The data presented here show that IPV should be considered as an important public health problem in Albania.

Prevalence of Forced Sexual Intercourse

Every Albanian woman who participated in the Reproductive Health Survey was asked about her experience of forced sexual intercourse against the her will at any time in her life, her age the first time this occurred, and her relationship to the person who forced the unwanted sex. For these questions, forced sexual intercourse was defined in the questionnaire as including vaginal, anal, and/or oral penetration. In examining the data resulting from these questions, it is important to keep in mind the probability of underreporting due to the sensitivity of the subject and the potential shame and social stigma associated with reporting sexual violence in a standardized interview. Although it is not possible to estimate the amount of underreporting on these questions, it is possible to assert that the data resulting from these questions represents a low-end estimation of the true prevalence of sexual violence among Albanian women. Another reason for underreporting is that violence and sex within marriage are almost never discussed within Albanian culture and sometimes sexual intercourse is used to decrease violence or the man's anger. Also, women may feel that sexual intercourse is

to fulfill their duties within a marriage. Women in Albania generally do not have sexual autonomy and negotiation skills do not exist (Baban et al., 2003; Lesko et al., 2003). Questions to evaluate the power of negotiation were not included in this survey, but should be in future surveys.

A total of 2% of Albanian women reported ever having experienced forced intercourse against their will. In comparison to other countries of the region, this percentage is quite low (data not shown). Since the numerator was so small when the data were disaggregated by characteristics of the women, no significant differences were found. Previously married women did report a higher prevalence (16%), as did women with four or more children (5%).

The data indicate that almost three quarters of women who experienced forced sex reported that they were older than 19 years of age at first forced sex (73%), compared to 18% of women who reported that forced sex occurred for the first time at ages 19 and under. Almost 10% did not reveal their age. Finally, over 80% of women reported that the perpetrator of forced sex was a husband or partner, 9% reported that the perpetrator was an ex-husband or partner, and 3% stated that the perpetrator was a boyfriend. It is notable that no women reported forced sexual intercourse by a casual partner or stranger.



Table 15.1 A Percentage of Women Aged 15-44 Who Witnessed or **Experienced Parental Abuse by Selected Characteristics Reproductive Health Survey: Albania 2002**

Characteristics	Witnessed Abuse	Experienced Abuse	No. of Cases*
Total	11.5	27.3	5,686
Strata			·
Metro Tirana	7.7	22.0	2,105
Other Urban	9.9	25.2	1,811
Other Rural	13.6	29.9	1,770
Residence			
Urban	9.3	24.4	3,565
Rural	13.2	29.4	2,121
Age Group			
15–19	10.6	28.9	1,091
20–24	14.3	28.3	934
25–29	10.5	27.3	946
30–34	9.8	26.4	1,065
35–39	12.3	26.6	956
40–44	12.0	25.4	694
Marital Status			
Married/Previously Married	12.2	26.7	4,046
Never Married	10.2	28.4	1,640
Living Children			
0	11.3	28.2	1,938
1	10.0	23.8	828
2	11.3	26.1	1,838
3	11.8	27.7	793
4 +	15.5	32.2	289
Education Level			
Primary or Less	14.1	31.2	2,514
Secondary Incomplete	7.7	23.1	651
Secondary Complete	9.4	23.3	1,826
Post-Secondary	6.8	20.1	695
Socioeconomic Index			
Low	14.7	31.1	1,933
Medium	9.6	24.6	2,982
High	7.0	23.5	771
Employment			
Working	8.4	20.4	1,116
Not Working	12.1	28.5	4,570

* Excluding 9 cases who did not live with 2 parents and 2 cases without information

Table 15.1 B Percentage of Men Aged 15–49 Who Witnessed or Experienced Parental Abuse by Selected Characteristics Reproductive Health Survey: Albania 2002

Characteristics	Witnessed Abuse	Experienced Abuse	No. of Cases*
Total	47.4	66.8	1,739
Strata			
Metro Tirana	37.3	58.7	717
Other Urban	47.8	62.5	547
Other Rural	51.2	72.5	475
Residence			
Urban	43.2	60.3	1,154
Rural	51.1	72.5	585
Age Group			
15–19	47.5	68.3	401
20–24	45.1	65.4	188
25–29	44.0	68.0	218
30–34	45.9	63.1	253
35–39	49.7	70.2	255
40–44	48.1	64.0	277
45–49	52.3	68.3	147
Marital Status			
Married	49.1	67.4	1,023
Previously Married	**	**	14
Never Married	44.9	65.7	702
Live Children			
0	44.8	66.6	814
1	45.0	62.8	221
2	47.7	64.3	468
3	54.3	69.1	167
4 +	57.2	81.0	69
Education Level			
Primary or Less	52.3	71.6	689
Secondary Incomplete	45.5	68.6	199
Secondary Complete	46.6	65.6	625
Post-Secondary	26.5	43.8	226
Socioeconomic Index			
Low	51.5	69.4	638
Medium	46.4	66.7	814
High	33.1	55.3	287
Employment			
Working	46.2	67.3	912
Not Working * Excludes 1 case who did not live wi	48.6	66.3	827

** Percentages are not shown when base is less than 25 cases

ables

Chapter	1	5
---------	---	---

Table 15.2 A Percentage Who Reported Intimate Partner Violence (IPV) in Their Lifetime and Percentage Who Reported Intimate Partner Violence in The Last Year, by Type of Abuse and Selected Characteristics, among Women Aged 15–44 Ever Married or in Union Reproductive Health Survey: Albania 2002

	Lifetime IPV IPV During The Last Year								
	Verbal	Physical	Sexual	No. of	Verbal	Physical	Sexual	No. of	
Characteristics	Abuse	Abuse	Abuse	Cases	Abuse	Abuse		Cases *	
Total	30.4	8.2	2.9	4,049	22.8	4.5	1.5	4,049	
Strata									
Metro Tirana	30.4	8.8	3.2	1,476	20.9	4.6	1.6	1,476	
Other Urban	33.0	8.8	2.7	1,331	24.8	4.1	1.2	1,331	
Other Rural	28.9	7.7	2.9	1,242	22.1	4.7	1.7	1,242	
Residence									
Urban	32.8	9.0	2.9	2,545	24.0	4.2	1.4	2,545	
Rural	28.5	7.6	2.8	1,504	21.8	4.7	1.7	1,504	
Age Group									
15–19	25.0	10.6	2.7	98	25.0	10.6	2.7	98	
20–24	20.5	6.0	1.6	505	18.0	4.2	1.0	505	
25–29	30.5	10.5	3.4	822	23.7	6.4	2.5	822	
30–34	35.0	9.3	3.5	1,017	25.4	4.6	1.9	1,017	
35–39	31.8	7.7	2.7	928	22.2	3.3	0.9	928	
40–44	30.9	6.7	2.7	679	22.3	3.2	1.1	679	
Marital Status									
Married	29.9	7.5	2.4	3,963	23.2	4.5	1.5	3,963	
Previously Married	45.8	31.2	18.9	86	7.1	4.0	3.5	86	
Living Children									
0	18.8	6.9	2.7	299	17.0	3.7	1.3	299	
1	28.1	8.9	2.7	828	20.6	5.3	1.2	828	
2	30.2	7.4	2.7	1,837	22.3	4.2	1.3	1,837	
3	34.0	9.6	2.5	794	25.2	4.5	1.6	794	
4 +	35.7	8.2	4.8	291	27.2	4.6	3.1	291	
Education Level									
Primary or Less	31.4	9.0	3.1	1,863	24.0	5.4	2.0	1,863	
Secondary Incomplete	31.8	13.7	6.8	246	22.6	6.7	3.3	246	
Secondary Complete	29.8	6.5	2.0	1,514	21.9	2.8	0.6	1,514	
Post-Secondary	24.2	5.4	2.0	426	16.7	2.7	0.7	426	
Socioeconomic Index									
Low	32.3	8.6	2.7	1,458	24.7	5.1	1.7	1,458	
Medium	29.7	8.3	3.2	2,063	22.1	4.2	1.6	2,063	
High	24.4	6.0	2.0	528	16.5	2.5	0.6	528	
Employment							5.0		
Working	29.5	8.4	4.5	890	19.8	3.5	1.6	890	
Not Working	30.6	8.2	2.5	3,159	23.4	4.7	1.5	3,159	

* Data are missing for 4 ever married women who were not asked about violence

Table 15.2 B

Percentage of Men Who Reported Perpetration of Intimate Partner Violence (IPV) in Their Lifetime And Percentage Who Perpetrated Intimate Partner Violence in The Last Year by Type of Abuse and Selected Characteristics, among Men Aged 15–49 Ever Married or in Union Reproductive Health Survey: Albania 2002

	Lifetime IPV				IPV During The Last Year			
Characteristics	Verbal Abuse	Physical Abuse	Sexual Abuse	No. of Cases	Verbal Abuse	Physical Abuse	Sexual Abuse	No. of Cases
Total	32.5	14.2	1.0	1,037	18.9	4.8	0.3	1,037
Strata								
Metro Tirana	30.6	16.9	1.9	430	15.1	6.9	1.4	430
Other Urban	35.2	13.3	0.9	321	20.8	4.7	0.0	321
Other Rural	31.6	13.7	0.6	286	19.4	4.1	0.0	286
Residence								
Urban	33.5	14.7	1.4	682	18.7	5.5	0.6	682
Rural	31.6	13.8	0.6	355	19.1	4.2	0.0	355
Age Group *								
20–24	21.7	13.5	6.1	33	20.5	13.5	4.9	33
25–29	16.1	7.1	1.7	128	11.5	2.0	0.6	128
30–34	29.3	9.7	0.4	216	20.3	3.4	0.2	216
35–39	33.8	11.7	0.8	243	21.6	4.2	0.0	243
40–44	39.0	17.2	0.8	273	21.5	4.4	0.0	273
45–49	39.9	23.5	0.6	144	16.3	7.9	0.0	144
Living Children								
0	11.8	5.0	1.1	112	8.4	2.2	0.7	112
1	28.3	9.2	1.8	221	19.3	5.1	0.8	221
2	37.7	14.4	1.0	468	22.8	5.3	0.1	468
3	36.0	19.5	0.4	167	15.8	2.6	0.0	167
4 +	34.2	22.3	0.0	69	19.7	9.1	0.0	69
Education Level								
Secondary Incomplete or Less	35.0	16.5	1.5	433	21.5	6.5	0.5	433
Secondary Complete or More	29.9	11.9	0.4	604	16.3	3.0	0.0	604
Socioeconomic Index								
Low	36.4	16.1	1.1	405	21.8	5.9	0.1	405
Medium	28.9	13.1	0.6	464	17.7	4.3	0.2	464
High	27.5	9.5	1.9	168	9.7	1.5	1.5	168
Employment								
Working	33.1	14.7	0.5	710	18.7	4.0	0.3	710
Not Working	31.5	13.4	1.7	327	19.3	6.0	0.2	327

* No ever-married men ages 15-19 were in the sample

Tables

Tables

Table 15.3 A	
Percentage Who Reported Lifetime Physical Violence and Recent	
Violence by Severity of Violent Act by Selected Characteristics,	
among Women Aged 15–44 Ever Married or in Union	
Reproductive Health Survey: Albania 2002	

	Lif	etime Ph	ysical Vio	lence		Physical	Violence	During TI	ne Last Ye	ar	
	Moderat	е		Severe		Moderate Severe					
Characteristics	Pushed You, Shook You, Shoved You, or Threw Something at You	Slapped You or Twisted Your Arm	Hit You With a Fist or With Something Else	Kicked You, Choked You or Beat You up	Threatened You With a Knife or Other Weapon	Pushed You, Shook You, Shoved You, or Threw Something at You	Slapped You or Twisted Your Arm	Hit You With a Fist or With Something Else	Kicked You, Choked You or Beat You up	Threatened You With a Knife or Other Weapon	No. of Cases*
Total	4.6	7.8	2.3	1.8	0.6	2.5	4.2	0.9	0.8	0.3	4,049
Strata											
Metro Tirana	4.2	8.6	2.4	1.8	0.3	2.2	4.1	1.2	1.2	0.2	1,476
Other Urban	5.1	8.2	2.7	2.0	0.8	2.5	3.8	1.1	0.9	0.1	1,331
Other Rural	4.5	7.3	2.0	1.6	0.5	2.6	4.5	0.7	0.6	0.3	1,242
Residence											
Urban	4.8	8.5	2.7	2.0	0.6	2.3	3.8	1.1	1.0	0.2	2,545
Rural	4.4	7.2	2.0	1.6	0.5	2.6	4.5	0.8	0.6	0.3	1,504
Age Group											
15–19	4.7	10.6	0.4	0.4	0.0	4.7	10.6	0.4	0.4	0.0	98
20–24	3.0	6.0	2.6	1.3	0.5	1.8	4.2	1.3	0.8	0.1	505
25–29	6.8	10.0	3.2	2.4	0.8	4.8	6.1	1.4	1.5	0.5	822
30–34	4.6	8.3	2.5	2.4	0.9	2.4	4.3	1.2	1.1	0.5	1,017
35–39	5.3	7.2	2.3	1.8	0.3	2.2	2.8	0.7	0.5	0.3	928
40–44	2.9	6.6	1.4	1.0	0.4	1.0	3.1	0.3	0.3	0.0	679
Marital Status											
Married	3.9	7.0	1.8	1.2	0.4	2.5	4.2	0.8	0.7	0.2	3,963
Previously Married	25.9	31.2	17.2	19.0	7.7	4.0	4.0	4.0	3.4	2.6	86
Living Children											
0	5.5	6.2	2.6	1.1	0.3	2.7	3.4	0.5	0.4	0.0	299
1	5.6	8.4	3.1	2.9	1.1	3.1	5.1	1.4	0.9	0.6	828
2	4.5	6.9	2.3	1.5	0.5	2.6	3.8	1.0	0.9	0.1	1,837
3	4.0	9.3	1.6	1.2	0.4	1.7	4.4	0.5	0.6	0.2	794
4 +	3.8	7.7	1.9	2.2	0.5	2.5	4.2	1.0	1.1	0.5	291
Education Level											
Primary or Less	4.8	8.5	2.6	1.8	0.7	2.8	5.2	0.9	0.7	0.3	1,863
Secondary Incomplete	8.7	13.0	2.1	3.9	0.6	2.8	6.7	1.0	1.5	0.6	246
Secondary Complete	3.8	6.2	1.9	1.5	0.5	1.9	2.5	0.9	0.8	0.2	1,514
Post-Secondary	3.3	5.1	1.8	1.3	0.0	2.5	2.3	1.3	0.9	0.0	426
Socioeconomic Index	(
Low	4.5	8.1	2.1	1.6	0.7	2.8	4.8	0.6	0.7	0.2	1,458
Medium	4.9	7.8	2.7	2.1	0.6	2.3	4.0	1.3	0.9	0.4	2,063
High	3.2	5.5	1.2	0.9	0.0	1.8	2.2	0.6	0.6	0.0	528
Employment											
Working	5.8	7.9	3.0	2.4	0.5	2.2	3.3	1.1	0.9	0.2	890
Not Working	4.3	7.7	2.2	1.6	0.6	2.6	4.4	0.9	0.8	0.3	3,159

* Data are missing for 4 ever married women who were not asked about violence

	am				49 Ever M th Surve						
			Physical					nce During	g the La	ist Year	
	Moder			Sever	e	Moder	ate		Sever	е	
Characteristics	Pushed, Shook, or Shoved, or Threw Something at Partner	Slapped or Twisted Partner Arm	Hit With a First or With Some- thing Else	Kicked, Choked or Beat Partner	Threatened Partner With a Knife or Other Weapon	Pushed, Shook, or Shoved, or Threw Something at Partner	Slapped or Twisted Partner Arm	Hit With a First or With Some- thing Else	Kicked, Choked or Beat Partner	Threatened Partner With a Knife or Other Weapon	No. of Cases
Total	9.0	13.1	3.6	1.4	0.1	2.7	4.6	0.4	0.6	0.0	1,037
Strata											
Metro Tirana	7.5	14.6	3.5	0.8	0.4	2.5	6.7	1.4	0.4	0.0	430
Other Urban	6.1	12.7	3.3	0.3	0.0	2.0	4.7	0.0	0.0	0.0	321
Other Rural	11.2	12.7	3.9	2.3	0.0	3.1	3.7	0.3	1.1	0.0	286
Residence											
Urban	6.6	13.6	3.4	0.5	0.2	2.1	5.5	0.6	0.2	0.0	682
Rural	11.0	12.6	3.8	2.1	0.0	3.2	3.8	0.3	1.0	0.0	355
Age Group *											
20–24	6.8	7.7	4.5	2.1	0.0	3.3	4.2	2.2	0.2	0.0	161
25–29	5.3	9.7	2.3	1.1	0.1	1.4	3.8	0.2	0.5	0.0	459
30–34	13.5	18.7	4.6	1.4	0.1	3.7	5.4	0	0.9	0.0	417
Marital Status											
Married	9.0	13.1	3.6	1.4	0.1	2.7	4.6	0.4	0.6	0.0	1,023
Previously Married	3.7	12.1	3.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	14
Living Children											
0	3.0	4.6	0.4	0.0	0.0	1.5	2.2	0.0	0.0	0.0	112
1	4.1	8.6	3.7	1.8	0.0	1.7	5.1	1.1	1.0	0.0	221
2	7.5	13.1	2.9	1.4	0.2	2.6	5.3	0.5	0.6	0.0	468
3	15.6	18.7	8.2	2.5	0.0	2.3	2.6	0.2	1.0	0.0	167
4 +	17.1	19.3	1.2	0.0	0.0	6.8	6.9	0.0	0.0	0.0	69
Education Level											
Primary or Less	10.2	15.0	3.6	2.2	0.1	3.8	6.1	0.9	1.2	0.0	433
Secondary Incomplete	7.7	11.1	3.6	0.6	0.1	1.5	3.0	0.0	0.0	0.0	604
Socioeconomic Index	Σ.										
Low	10.7	14.6	3.5	2.4	0.0	3.8	5.4	0.2	1.3	0.0	405
Medium	8.2	12.7	3.5	0.5	0.2	1.6	4.3	0.5	0.0	0.0	464
High	3.5	7.3	4.6	0.0	0.0	1.5	1.5	1.5	0.0	0.0	168
Employment											
Working	9.2	13.8	4.9	1.2	0.1	1.9	4.0	0.6	0.6	0.0	710
Not Working	8.6	12.0	1.5	1.7	0.0	4.0	5.4	0.1	0.7	0.0	327

Table 15.3 B Percentage Who Reported Inflicting Lifetime Physical Violence And Recent Violence by Severity of Violent Act and Selected Characteristics, among Men Aged 15–49 Ever Married or in Union

* No ever-married men ages 15-19 were in the sample

** Percentages are not shown when base is less than 25 cases

ables

Table 15.4

Percent Distribution of Number of Incidents of Physical Abuse by Type of Abuse and Percentage Resulting in Physical Injuries Among Women Who Were Physically Abused During The Past Year Women Aged 15-44 Ever Married or in Union **Reproductive Health Survey: Albania 2002**

		Num	ber of l	ncident	s Past Year			% With	With	
Type of Abuse	1	2	3	4+	Don't Remember	Refused	Total	Physical Injuries	No. of Cases	
Pushed, Shook, Shoved or Threw Something at you	3.5	9.1	11.7	52.4	20.3	3.0	100.0	33.4	101	
Slapped You or Twisted Your Arm	10.0	16.7	4.5	35.9	22.6	10.2	100.0	19.4	172	
Hit You	8.7	9.4	4.1	48.0	27.8	2.0	100.0	56.3	43	
Threatened You With a Knife or Other Weapon	**	**	**	**	**	**	**	**	8	
Kicked You, Choked You or Beat You up	11.7	7.9	5.6	44.0	22.8	8.0	100.0	73.5	38	
Forced you to Have Sexual Relations	3.3	19.5	10.0	43.2	22.0	1.9	100.0	24.3	65	

** Percentages are not shown when base is less than 25 cases.

Table 15.5 Women Who Were Physically Abused by an Intimate Partner During The Past Year Who Talked to Anyone About This Incident of Violence by Selected Characteristics Reproductive Health Survey: Albania 2002

	Talked to Anyone About Violence (Percent Distribution)					
Characteristics	Yes	No	Total	No. of Cases		
Total	46.2	53.8	100.0	200		
Strata						
Metro Tirana	56.3	43.7	100.0	71		
Other Urban	48.2	51.8	100.0	62		
Other Rural	42.4	57.6	100.0	67		
Residence						
Urban	51.5	48.5	100.0	119		
Rural	42.4	57.6	100.0	81		
Age Group						
15–24	54.3	45.7	100.0	40		
25–34	43.9	56.1	100.0	101		
35+	45.1	54.9	100.0	59		
Education Level						
Secondary Incomplete or Less	42.0	58.0	100.0	143		
Secondary Complete or More	59.2	40.8	100.0	57		
Socioeconomic Index						
Low	39.5	60.5	100.0	93		
Medium/High	52.9	47.1	100.0	107		

Table 15.6 Percentage of Women Who Were Physically Abused by an Intimate Partner During The Past Year Who Discussed the Abuse With Other Persons by Type of Person by Selected Characteristics Reproductive Health Survey: Albania 2002

		Husband's	Other				Doctor/ Health	1	Legal	No. of
Characteristics	Mother	Family	Relative	Friend	Police	Neighbor	Provider	Children	Adviser	Cases
Total	51.9	38.9	31.6	25.7	9.5	8.5	8.1	6.4	2.7	97
Strata										
Metro Tirana	37.5	23.8	38.8	20.1	2.2	9.6	11.4	5.6	0.0	39
Other Urban	50.4	38.1	31.3	34.1	2.9	0.0	9.6	6.5	3.5	30
Other Rural	57.9	44.8	29.2	23.0	15.9	13.0	6.2	6.6	3.1	28
Residence										
Urban	45.3	32.4	34.3	29.4	2.8	3.9	10.9	6.5	2.3	63
Rural	57.4	44.4	29.4	22.6	15.2	12.4	5.9	6.3	3.0	34
Age Group										
15–29	43.2	43.1	22.6	24.9	14.1	10.7	7.4	0.0	3.6	46
30–44	59.0	35.5	39.1	26.4	5.7	6.8	8.7	11.6	1.9	51
Education Level										
Secondary Incomplete or Less	52.6	41.5	29.4	18.8	7.3	7.1	5.1	4.0	0.0	63
Secondary Complete or More	50.3	33.0	36.6	41.2	14.4	11.8	15.0	11.8	8.6	34
Socioeconomic Index										
Low	46.3	61.0	31.2	18.4	7.4	3.6	8.4	7.9	3.8	39
Medium/High	56.1	22.1	32.0	31.3	11.1	12.2	8.0	5.2	1.8	58

Table 15.7 Percentage Distribution of Women Who Were Ever Physically Abused by an Intimate Partner Who Did not Report the Abuse to a Health Provider, Police, or Lawyer by reasons for Not Reporting The Abuse by Residence and Socio-Economic Index Reproductive Health Survey: Albania 2002

		Resid	lence	Socioeco	onomic Index
Main Reason no Legal/Medical Help	Total	Urban	Rural	Low	Medium/High
No Use Would Not do Any Good	32.5	31.5	33.3	30.2	35.0
Violence is Normal /No Need to Complain	27.7	30.0	26.1	27.7	27.7
Did not Know Where to Seek Help	9.0	5.5	11.6	13.4	4.2
Afraid of Divorce	7.1	6.8	7.4	6.5	7.8
Afraid of More Beatings	6.5	6.2	6.7	8.4	4.3
Bring Bad Name to Family	4.7	5.2	4.3	4.5	4.9
Embarrassed	3.8	5.2	2.8	3.7	3.9
Thought Would Not be Taken Seriously	2.8	4.4	1.7	1.2	4.6
Thought she Would be Blamed	1.1	2.6	0.0	0.0	2.3
Afraid of Losing the Children	1.0	0.0	1.8	2.0	0.0
Don't Know/Refused	3.8	2.7	4.5	2.3	5.3
Total	100.0	100.0	100.0	100.0	100.0
No. of Cases	189	113	76	91	98

Tables ا

Violence Against Women

Table 15.8 Situations That Make Partner Violent as Reported by Women Who Were Physically Abused by an Intimate Partner During The Past Year by Residence Reproductive Health Survey: Albania 2002

		Reside	ence
Characteristics	Total	Urban	Rural
Husband/Partner is Drunk	42.0	42.6	41.4
Family Has Financial Troubles	40.5	38.7	42.1
Husband/Partner is Unemployed	36.7	35.8	37.6
Husband/Partner is Jealous	32.3	33.0	31.7
There Are Family Problems	21.4	24.7	18.3
She Does Not Look After Children	20.1	15.4	24.4
Husband/Partner Acts According to Kanun	12.4	10.8	13.8
Husband/Partner Does Not Have Food at Home	9.8	10.0	9.7
She is Unemployed/Has no Income	8.9	6.1	11.4
Husband/Partner Cannot Get Alcohol/Drugs	3.8	4.3	3.3
She is Pregnant	2.6	2.7	2.6
Other—Partner Violence	0.6	0.5	0.8
No. of Cases	352	241	111

Table 15.9

Percentage of Women Aged 15–44 Ever Married or in Union Who Ever Experienced Any Form of Abuse Who Want to Know of The Tirana Hotline Number Where They Can Ask Questions About Violence Against Women Reproductive Health Survey: Albania 2002

	Percentage Who Want to Know Tirana Hotline Number					
Characteristics	Yes	No	Total	No. of Cases		
Total	12.1	87.9	100.0	1,275		
Strata						
Metro Tirana	15.2	84.8	100.0	451		
Other Urban	15.3	84.7	100.0	447		
Other Rural	9.1	90.9	100.0	377		
Residence						
Urban	15.3	84.7	100.0	841		
Rural	9.3	90.7	100.0	434		
Age Grouped						
15–29	16.4	83.6	100.0	397		
30–44	10.3	89.7	100.0	878		
Education Level						
Secondary Incomplete or Less	10.3	89.7	100.0	688		
Secondary Complete or More	15.4	84.6	100.0	587		
Socioeconomic Index						
Low	10.5	89.5	100.0	480		
Medium/High	13.6	86.4	100.0	795		

Table 15.10 Percentage of Women Age 15-44, Who Stated That a Man Has a Right to Hit or Beat His Wife Under The Following Situations by Residence and Education **Reproductive Health Survey: Albania 2002**

		Resid	dence	Level of E	ducation
Situation	Total	Urban	Rural	Secondary Incomplete or Less	Secondary Complete or More
She Asks Him if He Has Other Girlfriends	59.8	47.4	69.2	67.8	45.6
She Neglects The Children	33.5	22.5	41.7	40.2	21.4
She Dresses Too "Sexy or Depends Too Much on Her Looks"	27.5	18.4	34.3	33.8	16.2
She Goes Out Without Telling Him	25.8	14.3	34.4	33.1	12.8
She Argues With Him	22.2	13.7	28.6	28.6	10.8
He is Not Happy With Her Care of Household	15.2	8.5	20.2	19.9	6.7
He Finds Out That She Has Been Unfaithful	11.6	6.5	15.4	15.3	5.1
She Refuses to Have Sex	11.4	6.2	15.2	14.6	5.6
No. of Cases	5,697	3,572	2,125	3,172	2,525

Glossary

Abortion: Purposeful termination of an intrauterine pregnancy with the intention of avoiding a live birth.

Abortion-to-live birth ratio: The number of abortions to women aged 15-44 years during a specified period of time divided by the number of live births to women aged 15-44 years during the same time period.

Age-specific fertility rate (ASFR): The number of births to women within a specific age group during a specified period of time per 1,000 women aged 15-44; traditionally given for 5-year age groups.

Age-specific abortion rate (ASAR): The number of abortions to women within a specific age group during a specified period of time per 1,000 women aged 15-44; traditionally given for 5-year age groups.

Acquired Immune Deficiency Syndrome (AIDS): Disease caused by the human immunodeficiency virus (HIV), which disrupts the immune system and leads to death from opportunistic infections.

Below replacement fertility: Total fertility rate below that which is necessary to maintain the current size of a population, generally less than 2.1 births per woman.

CDC: Centers for Disease Control and Prevention.

Child mortality rate: Deaths that occur between exact ages 1 and 5, during a specified time period per 1,000 children aged 1 up to but not including 5 years of age during the same time period.

Contraceptive effectiveness: The reduction in pregnancy rate due to the use of a method of contraception compared with the pregnancy rate expected by chance if not using contraception. Contraceptive failure is the inverse of contraceptive effectiveness.

Contraceptive prevalence rate (CPR): Percentage of currently married and inunion women who are using some method of pregnancy prevention (modern or traditional).

Districts (Rrethet): Albania is divided into 36 districts.

Fecund: Physically capable of becoming pregnant.

General fertility rate (GFR): The number of live births to women of all ages during a specified period of time per 1,000 women aged 15-44 years.

General abortion rate (GAR): The number of abortions to women of all ages during a specified period of time per 1,000 women aged 15-44 years.

Human Immunodeficiency Virus (HIV): Virus which causes AIDS, spread by sexual contact with an infected person, use of needles or blood products contaminated with the virus, and from mother to child during pregnancy, delivery or breastfeeding.

Household: one or more persons who share a dwelling and expenses.

Infant mortality rate (IMR): Deaths that occur dying in the first year of life (from birth to, but not including 1 year of age), during a specified time period per 1,000 live births during the same time period.

Infecund: Physically incapable of becoming pregnant.

INSTAT: National Institute of Statistics, Rruga Leke Dukagjini 5, Tirana, Albania

Intimate partner violence (IPV): Any act of physical, sexual or psychological harm, including threats of such acts, coercion or arbitrary deprivations of liberty, whether occurring in public or private life, perpetrated against a woman by her male partner.

Institute of Public Health (IPH): Institute of Public Health, Rruga Alexander Moisiu 80, Tirana, Albania.

Kotelchuck Index: Index to measure the adequacy of prenatal care utilization combining the time of initiation of prenatal care with the number of visits.

Low birth weight (LBW): Infant weight at birth of less than 2,500 grams.

Maternal mortality ratio (MMR): Number of women who die of any cause related to or aggravated by pregnancy or its management during a specified period of time per 100,000 live births during the same time period.

Method mix: Percent distribution of methods used by contracepting women, adding up to 100%.

Mistimed pregnancy: An unintended pregnancy that occurs when the woman wants a child at some point in the future.

Modern methods of contraception: Pregnancy prevention by supplied or surgical means including condoms, diaphragms, cervical caps, spermicides, intrauterine devices (IUDs), oral contraceptives, contraception, injectables, emergency patches, implants, male and female sterilization; generally more effective than traditional methods of contraception in preventing pregnancy.

Mother-to-child transmission (MTCT): Refers to ways in which HIV is passed from an HIV-infected mother to her child. These include transmission during pregnancy, during delivery, and through breastfeeding.

Neonatal mortality rate (NNMR): Deaths that occur within the first month of life (from birth to, but not including, 28 days) during a specified time period per 1,000 live births during the same time period.

Permanent method of contraception: Male or female sterilization, also called surgical contraception, which provides permanent and highly effective pregnancy prevention.

Post-neonatal mortality rate (PNMR): The difference between neonatal and infant mortality rates, interpreted as deaths that occur during the post-neonatal period (from 28 days to, but not including, 1 year of age) during a specified time period per 1,000 live births during the same time period.

Rate of natural increase: The birth rate minus the death rate, implying the annual rate of population growth without regard to migration.

Reproductive Health Survey (RHS): Population-based survey of women aged 15–44, and sometimes also men aged 15– 49, performed with technical assistance from CDC.

Replacement level fertility: The total fertility rate necessary to maintain the current size of a population, generally 2.1 births per woman.

Reproductive age: Age range during which most women are assumed to be capable of bearing children and contributing significantly to the TFR and CPR; in Reproductive Health Surveys considered to be 15–44 years of age. Sexually transmitted infection (STI): Infection spread by sexual contact.

Sub-fecund: Having a diminished capacity of becoming pregnant.

Total fertility rate (TFR): The average number of children that a woman would have during her childbearing years if she passed through those years experiencing the observed age-specific fertility rates (ASFRs); the sum of the ASFRs.

Total abortion rate (TAR): The average number of abortions that a woman would have over the course of her life if she experienced the observed age-specific abortion rates (ASARs); the sum of the ASARs.

Traditional methods of contraception: Pregnancy prevention by means such as periodic abstinence or withdrawal, often considered natural, but generally less effective than modern methods.

Under five mortality rate (Under5MR): Deaths that occur before 5 years of age during a specified period of time per 1,000 live births during the same time period. UNICEF: United Nations Children's Fund

Unintended pregnancy: A pregnancy is classified as unintended if the woman stated that "just before she got pregnant with that pregnancy" she did not want to have a(another) baby then or at any time in the future" (unwanted pregnancy) or if she wanted to get pregnant at a later time (mistimed pregnancy).

Unmet need for contraception: In the RHS, the percentage of women who are currently sexually active, fecund, not wanting to become pregnant and not currently using any method of contraception.

Unwanted pregnancy: A pregnancy is classified as unwanted if the woman stated that "just before she got pregnant with that pregnancy she did not want to have a(another) baby then or at any time in the future."

Women in union: Women who are currently married or living with a man in a consensual, unregistered union.

Young Adult Reproductive Health Survey (YARHS): Population-based survey of young people aged 15–24 conducted with technical assistance from CDC.

Golssary

References

Abt Associates, 2004. Primary Health Care Reform in Albania: Baseline Survey of Basic Health Service Utilization, Expenditures and Quality. 156 pp.; Bethesda, MD. USA.

ACOG committee opinion, 1995. Recommendations on frequency of Pap test screening. Number 152. Committee on Gynecologic Practice. American College of Obstetricians and Gynecologists. International Journal of Gynecology and Obstetrics, 49:210-211.

AFPA, IFAW and UNFPA, 2002. Albania. NGO shadow report. Convention on the elimination of all forms of discrimination against women (CEDAW), Tirana.

Alan Guttmacher Institute, 1998. Into a New World-Young Women's Sexual and Reproductive Lives. 56 p. New York NY, USA

Augard Y, Genet D, Euraud JL, Clavene P, Tubiana-Mathieu N, Philippe HJ, 2002. Impact of Screening on Breast Cancer Detection. Retrospective Comparative Study of Two Periods Ten Years Apart. European Journal of Gynecological Oncology; 23(1):37-41.

Baban et al., 2003. Domestic violence in Albania. An explanatory study. UNICEF.

Bongaarts J, 1978. A framework for analyzing the Proximate Determinants of Fertility. Population and Development Review; 17:293-313.

Bongaarts J, 1991. The KAP-Gap and the Unmet Need for Contraception. Population and Development Review; 17:293–313.

Cates W, Jr., 1982. Legal Abortion: The Public Health Record. Science; 215(4540):1586-1590. Centers for Disease Control and Prevention [CDC] and ORC/Macro, 2003. Reproductive, Maternal and Child Health in Eastern Europe and Eurasia: A Comparative Report. Atlanta, GA (USA) and Calverton, MD (USA).

Co PLAN, Center for Habitat Development, 2001. Gender survey in the Bathore urban area of the Kamza Municipality, Tirana, Albania.

Coles CD, 1993. Impact of Prenatal Alcohol Exposure on the Newborn and the Child. Clinical Obstetrics and Gynecology; 36:255-266.

Council of Europe, 1998. The Demographic Characteristics of National Minorities in Certain European States. Volume 1. Council of Europe Publishing. Strasbourg, Germany.

Czech Statistical Office, Factum, non Fabula, WHO Collaborating Center for Perinatal Medicine/Institute for the Care of Mother and Child, Prague, and Centers for Disease Control and Prevention, 1995. 1993 Czech Republic Reproductive Health Survey Final Report. Prague, CZ and Atlanta, GA (USA).

David HP, 1992. Abortion in Europe, 1920– 91: A Public Health Perspective. Studies in Family Planning; 23(1):1–22.

Dawson DA, 1986. The Effects of Sex Education on Adolescent Behavior. Family Planning Perspectives; 18 (4):162–170.

Dehne KL, Khodakevich L, Hamers FF, Schwartzander B, 1999. The HIV/AIDS Epidemic in Eastern Europe: Recent Patterns and Trends and Their Implications for Policy-Making. AIDS; 13:741–749. DiFranza JR, Lew RA, 1996. Morbidity and Mortality Associated with the Use of Tobacco Products by Other People. Pediatrics; 97: 560-568.

Duran A, Emslie K, 2000. Health System Strategy:2000-2010 for Albania. Ministry of Health, Tirana, Albania.

Edleson, JL, 1999. Children's Witnessing of Adult Domestic Violence. Journal of Interpersonal Violence; 14(8):839-870.

Fairbank A and Gaumer G, 2004. Organization and Financing of Primary Health Care in Albania. Paper presented at the American Public Health Association Annual Meeting. Washington D.C., USA.

Ferraz E, Ferreira I, Soares M, Morris L, 1992. Pesquisa sobre saúde reprodutiva e sexualidade do jovem: Rio de Janeiro, Curitiba e Recife em 1989-1990. Rio de Janeiro Brasil; BEMFAM and Centers for Disease Control.

Gjepali S, Progri V, Lezo M, Qirjaku S, Zaze V, Ikonomi M, Bizhga V, Rama S, 1999. Albanian National Women Report 1999. United Nations Development Programme, Tirana, Albania.

Gjipali et al., 1994. The Albanian women: Hesitation and perspectives. European Network for Women Studies, Bucharest.

Gjermeni E, Bregu M, 2003. Monitoring media on domestic violence (2001-2002). Publication of Women's Center (www. women-center.org.al), Tirana.

Goldberg H, Sullivan JM, Serbanescu F, Sharmanov A, 1999. Survey Findings on Actual and Intended Fertility Prevention in Eastern Europe and the Former Soviet Union. Paper presented at the 1999 Annual Meeting of the Population Association of America. Goldberg HI and Toros A, 1994. The use of Traditional Methods of Contraception among Turkish Couples. Studies in Family Planning; 25(2):122-128.

Greene ME, Rasekh Z, Amen K, Chaya N, Dye J, 2002. In This Generation: Sexual and Reproductive Health Policies for a Youthful World. Population Action International. Washington, D.C.

Grunseit A, Kippax S, 1997. Impact of HIV and Sexual Health Education on the Sexual Behavior of Young People: a Review Update. Joint United Nations Programme on HIV/AIDS (UNAIDS). Geneva, Switzerland.

Gryboski K, Hall R, Vian T, 2004. Informed Payments in the Public Sector in Albania. Paper presented at the American Public Health Association Annual Meeting. Washington, D.C., USA.

Haggerty PA, Rutstein SO, 1999. Breastfeeding and Complementary Infant Feeding, and the Postpartum Effects of Breastfeeding. DHS Comparative Studies No. 30. Macro International. Calverton, MD.

Hamers FF, Downs A, 2003. HIV in Central and Eastern Europe. Lancet; 361(9362): 1035–1044.

Haxhiymeri E et al., 2000. Mapping of existing information on domestic violence in Albania. UNICEF. Tirana, Albania

Henshaw SK, Singh S and Haas T, 1999. Recent trends in abortion rates worldwide, International Family Planning Perspectives, 25(1):4448.

Hill K, Abouzahr C, Wardlaw T, 2001. Estimates of Maternal Mortality for 1995. Bulletin of the World Health Organization; 79(3):182-193. Himes NE, 1970. Medical History of Contraception. New York, Schoken Books.

Hobcraft JN, McDonald JW, Rutstein SO, 1985. Demographic Determinants of Infant and Early Child Mortality: A Comparative Analysis. Population Studies; 39(3):363-386.

Hotaling G, Sugarman DB, 1986. An Analysis of Risk Markers in Husband to Wife Violence: The Current State of Knowledge. Violence and Victims. 1(2):101-124.

Hotchkiss DR, Hutchinson P, Berruti A, 2004. Poverty, Out of Pocket Payments and Utilization of Primary Health Care: Evidence from Albania. Paper presented at the American Public Health Association Annual Meeting. Washington, D.C., USA.

Instituti i Statistikës (INSTAT), 2002. The Population of Albania in 2001: Main Results of the Population and Housing Census. INSTAT, Tirana, Albania.

Instituti i Statistikës (INSTAT), 2003. Aborti Ne Shqipeeri. Available at www.instat.gov. al/graphics/doc/tabelat/sh n/shn01.html. Last visited 11/17/2003. INSTAT, Tirana, Albania.

International Centre for Migration and Health, 2000. Reproductive and Family Health Related Isuues in Albania. 85 pp.; Geneva, Switzerland.

IPH, 2003. Bulletin of Institute of Public Health (in Albanian). Tirana, Albania.

IPH, 2004. Bulletin of Institute of Public Health (in Albanian). Tirana, Albania.

Islami D, Kallajahi F, Gliozheni O, Reproductive Health in Albania. Available at www.gfmer.ch/Endo/Reprod_Health_ Eastern_Europe/albania/ALNANIA_ ISLAMI.html. Last updated August 13, 2003. Geneva Foundation for Medical Education and Research.

Jha P, Ranson MK, Nguyen SN, Yach D, 2002. Estimates of Global and Regional Smoking Prevalence in 1995, by Age and Sex. American Journal of Public Health; 92:1002-1006.

Joint United Nations Programme on AIDS [UNAIDS], 2001. Regional HIV/AIDS StatisticsandFeatures,Endof2001.Available at www.unaids.org/worldaidsday/2001/ EPIgr aphics2001/EPIgraphic2_en.gif. Last updated December, 2001.

Kaçi B, 1996. Violence against women-A national problem (in Albanian). Tirana.

Kirby, D, 1999. Reducing Adolescent Pregnancy: Approaches that work. Contemporary Pediatrics; 16 (1): 83–94.

Kirby D, Short L, Collins J, Rugg D, Kolbe L, Howard M, Miller B, Sonenstein F, Zabing LS, 1994. School-Based Programs to Reduce Sexual Risk Behaviors: A Review of Their Effectiveness. Public Health Reports; 109:339–359

Kish L., 1967. Cluster Sampling and Subsampling. In: Survey Sampling. New York: John Wiley and Sons.

Kishor S and Johnson K, 2004. Profiling Domestic Violence—A Multi-Country Study. ORC MACRO, Calverton, MD-USA. Kolbo JR, Blakely EH, Engleman D, 1996. Children Who Witness Domestic Violence: A Review of Empirical Literature. Journal of Interpersonal Violence; 11(2):281-293.

Kotelchuck M, 1994. An Evaluation of the Ressner Adequacy of Prenatal Care index and a Purposed Adequacy of Prenatal Care Utilization Index. American Journal of Public Health; 84: 1414-1420.

Krug EG, Dahlberg LL, Mercy JA et al., 2002. World Report on Violence and Health. World Health Organization, Geneva, Switerland.

Kulczycki A, 2004. The Determinants of Withdrawal use in Turkey: A Husband's Imposition or a Woman's Choice? Soc. Sci. Med. 59(5): 1019-1033.

Labbok MH and Krasovek K, 1990. Toward Consistency in Breastfeeding Definitions. Studies in Family Planning; 21: 226-230.

Latthe PM and Shafi MI, 2001. Screening for Gynecological Conditions. Current Obstetrics and Gynecology; 11(1):31-37.

Le TN and Verma VK., 1997. An Analysis of Sample Designs and Sampling Errors of the Demographic and Health Surveys. DHS Analytical Reports No. 3. Calverton, Maryland: MACRO International Inc.

Lesko et al., 2003. The girls and trafficking (in Albanian). Vlora. Tirana, Albania.

Macaluso A, Ronfani L, Bacci A, Cattaneo A, 1999. Safe Motherhood Needs Assessment in Albania. WHO Regional Office for Europe and WHO Collaborating Centre, Trieste, Italy. McCauley AP and Salter C, 1995. Meeting the Needs of Young Adults. Population Reports, Series J, No. 41. Johns Hopkins Population Information Program, Baltimore, MD.

Meçaj V, 2000. Protection of women rights in Albania – Current analysis (in Albanian), Tirana 1997 in "Study about contribution of women movement in Albania 1990-1998," Tirana. (Studim mbi kntributin e shkruar the levizjes se gruas ne Shqiperi 1990-1998, Tirane).

Miller AB, 1986. Screening for Cancer: Issues and Future Directions. Journal of Chronic Diseases; 39:1067-1077.

Ministry of Finance, 2002. National Strategy for Social and Economic Development (NSSED). Tirana, Albania.

Ministry of Health (MOH), 2003. Albanian Health Promotion and Public Health Strategy. Tirana, Albania.

Ministry of Health (MOH), 2003. The National Strategy of Prevention and Control of HIV/AIDS in Albania (2004-2010). Tirana, Albania.

Ministry of Health (MOH), 2004. Albania Health Care System Strategy. Tirana, Albania.

Minnesota Advocates for Human Rights, 1996. Domestic Violence in Albania. Minneapolis, Minnesota.

Miria S, 1996. Violence against women and psychosocial taboo (in Albanian). Refleksione. Tirana, Albania.

Miria et al., 2000. Survey on violence against women and trafficking in rural areas of Lezha District (in Albanian). Reflecksione. Tirana, Albania. Morris L, 2000. History and Current Status of Reproductive Health Surveys at CDC. American Journal of Preventive Medicine. 19(1S):31-34.

Morris L, Sedivy V, Friedman J and McFarlane C, 1995. Sexual Behavior and Contraceptive Use Among Young Adults in Jamaica: Results from 1993 Contraceptive Prevalence Survey. National Family Planning Board, Kingston, Jamaica.

National Committee of Women and Family and UNICEF, 2000. Mapping of Existing Information on Domestic Violence in Albania. Tirana, Albania.

National Equity Committee, 2002. First National Report on Convention on the elimination of all forms of discrimination against women (CEDAW), Tirana, Albania.

Neza A, 2000. Welcome to Albania. ILAR, Tirana, Albania.

O'Campo P, Gielen AC, Fadan RR, Kass N, 1994. Verbal Abuse and Physical Violence Among a Cohort of Low-Income Pregnant Women. Women's Health Issues; 4(1):29-37.

Parkin DM, Adair L, Akin JS et al., 1993. Estimates of the Worldwide Incidence of Eighteen Major Cancers in 1985. International Journal of Cancer, 54: 594-606.

Piha T, Besselink E, Lopez AD, 1993. Tobacco or Health. World Health Statistics Quarterly; 46(3):188-194.

Population Reference Bureau (PRB). 1999 and 2000. 1998 and 1999 World Population Data Sheet. Population Reference Bureau, Washington D.C. Population Reference Bureau (PRB), 2002a. Women of the World Data Sheet. Population Reference Bureau, Washington D.C.

Population Reference Bureau (PRB), 2002b. 2001 World Population Data Sheet. Population Reference Bureau, Washington D.C.

Rahman A, Katzive L, Henshaw SL, 1998. A Global Review of Laws on Induced Abortion, 1985-1997. International Family Planning Perspectives; 24(2):56-64.

Rebentisch DP, Rebentish HD, Thomas K, Karat S, Jadhav AJ, 1995. A Proven and Highly Cost-Effective Method of Early Detection of Breast Cancer for Developing Countries. Radiotherapy and Oncology; 37:246-248.

Refleksione, 2000. Study about contribution of women movement in Albania 1990-1998 (in Albanian), Tirana. (Studim mbi kontributin e shkruar the levizjes se gruas ne Shqiperi 1990-1998, Tirane).

Riedner G, Dehne KL, and Gromyko A, 2000. Recent Declines in Reported Syphilis Rates in Eastern Europe and Central Asia: Are the Epidemics Over? Sexually Transmitted Infections; 76(5):363–365.

Robey B, Rutstein SO, Morris L, Blackburn R, 1992. The Reproductive Revolution: New Survey Findings. Population Reports, Series M, No. 11. Johns Hopkins Population Information Program, Baltimore, MD.

Roudi-Fahimi F, 2004. Islam and Family Planning. Population Reference Bureau Policy Brief. Washington D.C. USA. Salem R, 2004. Men's Surveys: New Findings, Population Reports, Series M, No. 18. Baltimore USA. Johns Hopkins School of Public Health.

Santow G, 1993. Coitus Interruptus in the Twentieth Century. Population and Development Review; 19(4):767-792.

Serbanescu F, 1998. Reproductive Health Issues in Eastern Europe and the Newly Independent States: Country Cases. Paper presented at the Regional UN Population Meeting. Budapest, Hungary 7-9 December.

Shapo L, Gilmore A, Coker R, 2003. Prevalence and Determinants of Smoking in Tirana City, Albania: A Population-based Survey. Public Health, 117:228-236.

Song LY, Singer MI, Anglin TM, 1998. Violence Exposure and Emotional Trauma as Contributors to Adolescents' Violence Behaviors. Archives of Pediatrics and Adolescent Medicine; 152:531-536.

Strauss MA, Gelles RJ, 1979. Measuring Intrafamily Conflict and Violence: The Conflict Tactics Scale. Journal of Marriage and the Family, 41:75-88.

Tejada R, Herold J, Morris L, 1997. Encuesta Nacional de Jóvenes: Republica Dominicana, 1992. Instituto do Estudios de Población y Desarrollo, Santo Domingo, D.R.

Tountas Y, Dimitrakaki C, Antoniou A et al., 2004. Attitudes and Behaviors Toward Contraception Among Greek Women During Reproductive Age: A Country-Wide Survey. Eur. J. Obstet. Gynecol. Reprod. Biology. 116(2): 190-195. United Nations (UN), 2003. World Population (Wall Chart). Population Division Publication (ST/ESA/SER. A/224). UN, New York, NY.

United Nations Economic Commission for Europe. 1997-2001. Fertility and Family Planning Surveys in Countries of the Economic Commission for Europe, Country Reports.

UNAIDS, 2004. Executive Summary. 2004 Report on the global AIDS epidemic. Last updated June, 2004.

UNAIDS, 2001. Regional HIV/AIDS statistics and features, end of 2001. Available at www.unaids.org/worldaidsday/2001/ EPIgraphics2001/EPIgraphic2_en.gif. Last updated December, 2001.

UNAIDS and WHO, 2003. AIDS epidemic update. Geneva, Switzerland.

UNDP, 2004. Albanian National Report. On progress toward achieving the Millenium Development Goals, Tirana, Albania.

UNICEF, 1998. Situation Analysis: Children's and Women's Rights in Albania.

UNICEF-Albania, 2000. Assessment of Social and Economic Conditions of Districts in Albania. UNICEF-Albania, Tirana, Albania.

UNICEF and INSTAT, 2000. Albania Multiple Indicator Survey Report. Tirana, Albania.

UNICEF et al., 2002. Rapid Assessment and Response on HIV/AIDS Among Especially Vulnerable Young People in Albania. Tirana, Albania. U.S. Department of Health and Human Services, 2001. Women and Smoking, A Report of the Surgeon General—2001. Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. Atlanta, GA.

Van Hook et al. Responding to gender violence in Albania. A partnership effort. International Social Work 43(3):351-363.

Weisman CS, Macconnaon DS, Henderson JT, Shortridge E, Orso CL, 2002. Contraceptive Counseling in Managed Care: Preventing Unintended Pregnancy in Adults. Women's Health Issues; 12(2):79-95.

Westoff CF, 1976. The Decline of Unplanned Births in the United States. Science; 191:38–41.

Westoff CF, 2000. The Substitution of Contraception for Abortion in Kazakhstan in the 1990s. DHS Analytical Studies No. 1. ORC Macro. Calverton, MD.

Westoff CF, Sharmonov AT, Sullivan JM, Croft T, 1998. Replacement of Abortion by Contraception in Three Central Asian Republics. The Policy Project and Macro International. Calverton, MD.

Westoff CF, Sullivan JM, Newby HA, Themme AR, 2002. Contraception-Abortion Connections in Armenia. DHS Analytical Studies No.6. ORC Macro. Calverton, MD.

World Bank, 1997. Albania: Growing Out of Poverty. Report No.15698-ALB. World Bank, Albania. World Health Organization (WHO), 1987. Violence Against Women: A Public Health Priority. WHO/FRH/WHD/97.8, Geneva, Switzerland.

World Health Organization (WHO), 1991. Indicators for Assessing Breast-feeding Practices. WHO/CDD/SER/91.14, Geneva, Switzerland.

World Health Organization (WHO), 1993. International classification of diseases and related health problems, tenth revision. Geneva, Switzerland: WHO.

World Health Organization (WHO), 2001. Ethical Recommendations for Investigations on Domestic Violence against Women. WHO/ FCH/GWH/01.1, Geneva, Switzerland.

World Health Organization (WHO), 2003. Country Health Indicators. Available at http:// www3.who.int/whosis/country/indica tors. cfm?country=alb&language=en.

World Health Organization (WHO), 2003. European Health for All Database. Available at www.euro.who.int/HFADB. Last visited April 2003. WHO Regional Office for Europe. Copenhagen, Denmark.

Zlidar VM, Gardner R, Rutstein SO, Morris L, Goldberg HI, 2003. New Survey Findings: The Reproductive Revolution Continues. Population Reports, Series M, No. 17. Johns Hopkins Bloomberg School of Public Health, Population Information Program. Baltimore, MD.

References

APPENDIX A

SAMPLING ERROR ESTIMATES

The estimates for a sample survey are affected by two types of errors: nonsampling error and sampling error.

Non-sampling error is the result of mistakes made in carrying out data collection and data processing, including the failure to locate and interview the right household, errors in the way questions are asked or understood, and data entry errors. Although intensive quality-control efforts were made during the implementation of the Albania Reproductive Health Survey (ALRHS-02) to minimize this type of error, non-sampling errors are impossible to avoid altogether and difficult to evaluate statistically.

Sampling error is a measure of the variability between an estimate and the true value of the population parameter being estimated, which can be attributed to the fact that a sample rather than a complete enumeration was used to produce it. In other words, sampling error is the difference between the real value for any variable measured in a survey and the value estimated by the survey. The samples selected are only one of the many probability samples that could have been selected from the female population aged 15–44 and the male population 15-49 years of age using the same sample

design and projected sample size. Each of these samples would have yielded slightly different results compared with results from the actual sample selected.

Because the statistics presented here are based on a sample, they may differ by chance variation from the statistics that would result if all women 15-44 years of age or men 15-49 years of age in Albania would have been interviewed. Sampling error is usually measured in terms of the variance and standard error (square root of the variance) for a particular statistic (mean, proportion, or ratio). The standard error (SE) can be used to calculate confidence intervals (CI) of the estimates within which we can say with a given level of certainty that the true value of population parameter lies. For example, for any given statistic calculated from the survey sample, there is a 95 percent probability that the true value of that statistic will lie within a range of plus or minus two SE of the survey estimate. The chances are about 68 out of 100 (about two out of three) that a sample estimate would fall within plus or minus one SE of a statistic obtained from a complete census count or survey of the complete population.

The estimated sampling errors for 95% confidence intervals (1.96 x SE) for selected proportions and sample sizes

are shown in Table A.1. The estimates in Table A.1 can be used to estimate 95% confidence intervals for the estimated proportions shown for each sample size. The sampling error estimates include an average design effect of 1.6, needed because the ALRHS-02 did not employ a simple random sample but included clusters of elements in the second stage of the sample selection.

Table A.1 Sampling Error Estimates (Expressed in Percentage Points) for 95% Confidence Intervals for Selected Estimated Proportions and Sample Sizes on Which the Proportions Are Based Assuming a Design Effect of 1.6

			Estimated Pr	<u>oportions (Pi</u>)	
Sample Size	0.05/0.95	0.10/0.90	0.20/0.80	0.30/0.70	0.40/0.60	0.50/0.50
25	0.108	0.149	0.198	0.227	0.243	0.248
50	0.076	0.105	0.140	0.161	0.172	0.175
100	0.054	0.074	0.099	0.114	0.121	0.124
200	0.038	0.053	0.070	0.080	0.086	0.088
400	0.027	0.037	0.050	0.057	0.061	0.062
800	0.019	0.026	0.035	0.040	0.043	0.044
1000	0.017	0.024	0.031	0.036	0.038	0.039
1500	0.014	0.019	0.026	0.029	0.031	0.032
2000	0.012	0.017	0.022	0.025	0.027	0.028
3000	0.011	0.014	0.020	0.021	0.022	0.023
4000	0.008	0.012	0.016	0.018	0.019	0.020
5000	0.008	0.011	0.014	0.016	0.017	0.018

The selection of clusters is generally characterized by some homogeneity that generally tends to increase the variance of the sample. Thus, the variance in the sample for the ALRHS-02 is greater than a simple random sample would be due to the effect of clustering. The design effect represents the ratio of the two variance estimates: the variance of the complex design using clusters, divided by the variance of a simple random sample using the same sample size (Kish L, 1967). For more details regarding design effects for specific reproductive health variables, the reader is referred to the Le and Verma report, which studied demographic and health surveys in 48 countries (Le TN and Verma JK, 1997). The pattern of variation of design effects is shown to be consistent across countries and variables. Variation among surveys is high but less so among variables. Urban -rural and regional differentials in design effects are small, which can be attributed to the fact that similar sample designs and cluster sizes were used across domains within each country. At the country level, the overall design effect, averaged over all variables and countries, is about 1.5 (we used 1.6 in Table A.1 to be slightly more conservative).

To obtain the 95% CI for proportions or sample sizes not shown in the table, one may interpolate. For example, for a sample size of 200 and a point estimate of 25% (midway between 0.20/0.80 and 0.30/0.70), the 95% CI would be plus or minus 7.5 percentage points; for a sample size of 300 (midway between 200 and 400) and an estimate of 20%, the 95% CI would be plus or minus 6.0 percentage points.

Specific to this report, one could use the table for a quick estimate of the 95% confidence interval for the percentage of women with births in the period from 1997 to 2002 that did not have prenatal care.

According to Table 5.1 in this report, 19.1% of 2,551 births did not have prenatal care. Let us round off the percent with no prenatal care to 20% and the number of births to 2500. Using the table, we look down the column for 0.20 and interpolate halfway between sample sizes of 2000 (0.022) and 3000 (0.020) and estimate the confidence

interval to be 0.021, or roughly between a lower limit of 17.0% and an upper limit of 21.2%. The actual confidence interval is 0.019 or 1.9 percentage points, which yields a lower limit of 17.2% and an upper limit of 21.0%..

Differences between estimates discussed in this report were found to be statistically significant at the five percent level using a two-tailed normal deviate test (p=0.05). This means that in repeated samples of the same type and size, a difference as large as the one observed would occur in only 5% of samples if there were, in fact, no differences between the percentage in the population.

The relative standard error of a statistic (also called "coefficient of variation") is the ratio of the standard error (SE) for that statistic to the value of the statistic. It is usually expressed as a percent of the estimate. Estimates with a relative standard error of 30% or more are generally viewed as unreliable by themselves, but they may be combined with other estimates to make comparisons of greater precision. For example, an estimate of 20% based on a sample size of only 50 observations yields a SE of 7% (one half the 95% confidence interval shown in Table A.1). The relative standard error would be 35% (the ratio of the SE of 7% to the estimate of 20%), too large for the estimate to be reliable.

APPENDIX B

PARTICIPANT INSTITUTIONS

Institute of Public Health	Silvia Bino, Survey Director Alban Ylli Manuela Murthi Eduard Kakarriqi Alba Merdani Erieta Dobi Bukurie Gega Dritan Roshi Xhemile Miho Albana Ahmeti Vilson Topulli
Ministry of Health	Leonard Solis, Minister of Health Petrit Vasili Saemira Pino Nedime Ceka Donika Beba Miranda Hysi Nurie Caushi
University Hospital	Gjergji Theodhosi
Institute of Statistics	Milva Ekonomi Bashkim Ramizi Ermiona Kali Abedin Cici Ermira Galanxhi Margarita Caci Myhedin Llagami
USAID/Albania	Rita Hudson Harry Birnholz Pamela Wyville-Staples Zhaneta Shatri
UNFPA	Katy Shroff Manuella Bello Albana Gribo Blendi Cano

UNICEF	Roberto Migliorini	
	Lenin Guzman	
	Judith Leveille	
	Mariana Bukli	
	Alketa Zazo	
Centers for Disease Control	Alicia Ruiz	
and Prevention, Division of	Danielle B. Jackson	
Reproductive Health, Atlanta,	Fernando Carlosama	
GA, USA	Florina Serbanescu	
	Howard Goldberg	
	Joan Herold	
	Leo Morris	
	Paul Stupp	
	Ranee Seither	
	Rose Pecoraro	
	Shin Kim	
	Soloman Iyasu	
	Sonia Bowens	
	Stephen McCracken	
	Wyndy Amerson	

Participant Institutions

Valerie Haynes Jackson

Persons Involved In The 2002 Albanian Reproductive Health Survey Field Operations And Supervision

Female Survey Teams

Field Work Coordinator: Dritan Dushaj

	Team I	Team II
Supervisor	Rudina Baboci	Altina Peshkatari
Interviewers:	Ndricime Bulku	Areta Mecaj
	Ines Kristo	Daniela Shuke
	Elona Nikolli	Aferdita Alizoti
	Rozeta Gjela	Majlinda Skenderi

	Team III	Team IV
Supervisor	Oriola Halili	Odeta Hoxhaj
Interviewers:	Brixhilda Gjeka	Esmeralda Dosti
	Olsa Refugjati	Lumnie Doda
	Eliona Mucaj	Suzana Osmani
	Valbona Dervishi	Aida Daci

Supervisor Interviewers: Team V

Enkeleida Latifi Albana Bregu Enkelejda Ngjelina Jonida Naska Monika Hasani

Back-ups:

Nekida Jolldachi Eriketa Luzi Anila Buci Julia Mitrushi

Male Survey Teams

Field Work Coordinator: Euglen Hoxha

Team VI.

Supervisor: Ilirjan Bashllari

Team VII. Supervisor: Agim Giikopulli

Interviewers:

Dritan Mema Bledar Pine Arti Tarja Selim Kacidhe Lulezim Bardhi Emiliano Lazo Arjan Boci Roland Myhypi

Back-up

Klodian Vasia

Drivers

Vladimir Jace Alket Simani Arjan Zenuni Fatmir Koci Bashkim Kaci Mehmet Becolli Maksim Mulellari Olsi Abazi

2002 ALBANIA REPRODUCTIVE HEALTH SURVEY

FEMALE QUESTIONNAIRE

Hello. I'm from the National Institute of Public Health. We are doing a national survey about the health of women and men in Albania. The purpose of the survey is to collect information that will help us to plan reproductive health services in Albania.			
I would like to ask you about your health and where you obtain health services. All of the information you give us will be confidential. The interview is completely voluntary and if we should come to any question that you don't want to answer, just let me know and we'll go on to the next question. The interview will take about 35-40 minutes. I would like to start now, is that OK?			
SIGNATURE OF THE INTERVIEWER	DAY MONTH		
MARK IF THE WOMAN AGREES TO BE INTERVIEWED	1. YES \rightarrow CONTINUE		
	2. NO \rightarrow END OF INTERVIEW		
TIME STARTED:: ID NU	J MBER		

I. BACKGROUND CHARACTERISTICS

100. In what month and year were you born?

MONTH _____ YEAR _____

98 DON'T KNOW

101. How old are you (at last birthday)? ____YEARS OLD 98 DON'T KNOW

MAKE SURE THAT AGE AND DATE OF BIRTH CORRESPOND

102. What is the highest level of education you completed, not counting the current grade you are in?

000. NEVER ATTENDED (NO FORMAL EDUCATION)

1. GENERAL SCHOOL	1 2 3 4 5 6 7 8	88
2. THEORETICAL HIGH SCHOOL/ GIMNAZIUM	1 2 3 4	88
3. PROFESSIONAL(VOCATIONAL) SCHOOL	1 2 3 4 5	88
4. TECHNICAL SCHOOL (POSTSECONDARY)	1 2 3	88
5. UNIVERSITY/FACULTY	1 2 3 4 5+	88
6. POST UNIVERSITY/POSTGRADUATE STUDIES	1 2 3 4 5+	88
888. DON'T REMEMBER/ DON'T KNOW		

IF Q102 = 2 OR Q102=3 CONTINUE; ELSE GO TO Q104

- 103. Do you have a high school diploma?
 - 1. YES

2. NO

- 104. Do you currently work outside of the home (at least 20 hours per week)?
 - 1. YES \rightarrow GO TO Q106
 - 2. YES, BUT ON MATERNITY/PREGNANCY LEAVE \rightarrow GO TO Q106
 - 3. NO

- 105. What is the main reason that you are not working at this time?
 - 1. ATTENDING SCHOOL
 - 2. I NTERNAL DISPLACEMENT
 - 3. LOOKING FOR WORK
 - 4. LAID OFF
 - 5. DOES NOT NEED/WANT/LIKE TO WORK
 - 6. MEDICAL LEAVE
 - 7. MATERNITY LEAVE
 - 8. INABILITY TO FIND/AFFORD CHILD CARE
 - 9. HOMEMAKER
 - 10. PERMANENT DISABILITY
 - 11. HUSBAND DOES NOT ALLOW HER TO WORK
 - 12. PARENTS DO NOT ALLOW HER TO WORK
 - 13. ODD JOBS (<20 HOURS PER WEEK)
 - 14. TEMPORARY BACK FROM WORK ABROAD
 - 20. OTHER (SPECIFY)
- 106. I would like to ask you some questions about where you have lived. For most of the time until you were 12 years old, did you live in a town, or in a village?
 - 1. TOWN
 - 2. VILLAGE
- 107. In what month and year did you start to live continuously in ______ (NAME THE PLACE OF RESIDENCE)?

MONTH YEAR 00. ALWAYS, SINCE BIRTH \rightarrow GO TO 108 88. DON'T REMEMBER

- 107A. Just before you moved here (CURRENT PLACE OF RESIDENCE), did you live in a town, a village, or outside Albania ?
 - 1. A town (URBAN AREA),
 - 2. A village (RURAL AREA), or
 - 3. Outside Albania?
- 108. Are you currently married, not married but living with someone, separated, divorced, widowed, or have you never been married ?
 - 1. MARRIED \rightarrow GO TO Q111
 - 2. NOT MARRIED BUT LIVING WITH A PARTNER \rightarrow GO TO Q111
 - 3. SEPARATED \rightarrow GO TO Q111
 - 4. DIVORCED \rightarrow GO TO Q111
 - 5. WIDOWED \rightarrow GO TO Q111
 - 6. NEVER MARRIED
- 109. Have you ever lived with a boyfriend or partner ? (LIVING TOGETHER MEANS HAVING A SEXUAL RELATIONSHIP WHILE SHARING THE SAME USUAL ADDRESS.)
 - 1. YES \rightarrow GO TO Q111

2. NO

- 110. If you could choose exactly the number of children to have in your whole life, how many would that be?

 ______CHILDREN
 22. AS MANY AS GOD GIVES
 - 33. AS MANY AS HUSBAND WOULD WANT
 - 88. NOT SURE/DON'T REMEMBER

GO TO Q124

Female Questionnaire

	TIMES	<u>9. KE</u>	$FUSAL \rightarrow GO$			
TIMES	112. In what month and year did you begin living with your (first, second, third, or fourth) husband/partner?	13. How old was your I, II, III, IV husband/ partner when you started to live together?	114. What was the highest grade in school that your I,II,III,IV husband/ partner completed when you got married/started to live together ?	115. What is your current union relationship with your I, II, III, IV, husband/ partner, are you still in the relationship or how did the relationship end?	116. In what month and year did your union with your I,II,III,IV, .husband/ partner end?	117. IF:
I	MTH _ YR 88. DON'T KNOW/ REF	^{AGE} 88. DK	0. NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married → Q117 Living with partner >Q117 Separated Divorced Widowed 	MTH YR 88. DON'T KNOW/ REF	Q111=1 GO TO Q118; ELSE CONTINUE
п	MTH _ YR 88. DON'T KNOW/ REF	^{AGE} 88. DK	0.NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married → Q117 Living with partner→Q117 Separated Divorced Widowed 	MTH YR 88. don't know/ REF	Q111=2 GO TO Q118; ELSE CONTINUE
Ш	MTH YR 88. DON'T KNOW/ REF	– – AGE 88. DK	0.NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married → Q117 Living with partner→Q117 Separated Divorced Widowed 	MTH YR 88. DON'T KNOW/ REF	Q111=3 GO TO Q118 ELSE CONTINUE
IV	MTH _ YR 88. Don't Know/ Ref	AGE 88. DK	0.NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married → Q117 Living with partner→Q117 Separated Divorced Widowed 	MTH YR 88. DON'T KNOW/ REF	GO TO Q118

111.	How many times hav	e you been married or lived with a man as husband and wife?	•
	TIMES	9 REFUSAL \rightarrow GO TO O118	

118. When you first got married/living together as husband and wife did you wish to have any children?1. YES

2. NO \rightarrow GO TO Q120

8. NOT SURE- \rightarrow GO TO Q120

119. How many children did you wish to have when you first got married?

____ CHILDREN

- 22. AS MANY AS GOD GIVES
- 33. AS MANY AS HUSBAND WANTS
- 88. NOT SURE/DON'T REMEMBER

120. How many children did your husband wish to have when you first got married?

- ____ CHILDREN
- 22. AS MANY AS GOD GIVES
- 33. AS MANY AS RESPONDENT WANTS
- 77. NEVER DISSCUSSED
- 88. NOT SURE/DON'T REMEMBER

IF CURRENTLY SEPARATED, DIVORCED OR WIDOWED (Q115_LAST=3,4,5) GO TO Q124

- 121. Is your husband currently employed (either in Albania or abroad)?
 - 1. YES
 - 2. NO
 - 8. DK/REF
- 122. Since January 1997 has your (current) husband ever worked abroad/outside Albania?
 - 1. YES
 - 2. NO ----- \rightarrow GO TO Q124
 - 8. NUK E DI/REF \rightarrow GO TO Q124

	Appendix C				
122A.	Since January 1997, how many times did h	e work abroad?	TIMES		
122B.	Since January 1997, how many months tota	l did he work abroad	!? 		
	MONTHS				
123.	When was the last time when he worked out TRIP ?	tside Albania (MON	TH AND YEAR OF BEGINNING THE		
	MONTH	YEAR 88.	DO NOT REMEMBER		
123A.	In what month and year did he return from TRIP ?				
	MONTH	YEAR 77. 88.	STILL ABROAD/OUTSIDE ALBANIA DO NOT REMEMBER		
124.	More or less how many hours a day do you	listen to the radio?			
	;		\rightarrow GO TO Q128 TE ACCESS TO RADIO \rightarrow GO TO Q128 Y		
125.	What stations do you most often listen to? (READ LIST)	PROBE FOR MOR	E THAN ONE STATION, DO NOT		
		MENTIONED	NOT MENTIONED		
	A. TOP ALBANIA	1	2		
	B. RADIO TIRANA	1	2		
	C. CLUB FM	1	2		
	D. RASH	1	2		
	D. RASH E. STINET	1	2 2		
	E. STINET E. STINET	1	2		
	F. GOLD MUSIC	1	2		
	G. ITALIAN STATIONS	1	2		
	H. VOICE OF AMERICA	1	2		
	I. BBC	1	2		
	J. RADIO PLANET	1	2		
	K. OTHER	1	2		
126.	What types of programs do you most often l		OR MORE THAN ONE PROGRAM,		
	DO NOT READ LIST)				
		MENTIONED	NOT MENTIONED		
	A.NEWS	1	2		
	B.SPORTS	1	2		
	C.MUSIC D.PLAYS/DRAMAS	1	2 2		
	E.CHURCH/RELIGIOUS PROGRAMS	1 ' 1	2		
	F.WOMEN'S PROGRAMS	1	2		
	G.HEALTH PROGRAMS	1	2		
	H.POLITICAL EVENTS	1	2		
	I.OTHER	1	2		
127.	What times do you most often listen to the r READ LIST)	adio? (PROBE FOF	R MORE THAN ONE TIME, DO NOT		
		ENTIONED N	NOT MENTIONED		
	A. 6-8 AM	1	2		
	B. 8-10 AM	1	2		
	C. 10AM-NOON	1	2		
	D. NOON-2 PM	1	2		
	E. 2-4 PM	1	2		
	F. 4-6 PM	1	2		
	G. 6-8 PM	1	2		
	H. 8-10 PM	1	2		
	I. AFTER 10 PM	1	2		
	J. NO REGULAR TIMES	1	2		

128. More or less how many hours a day do you spend watching television?

HOURS A DAY	00. NEVER \rightarrow GO TO Q132
	55. DOES NOT HAVE ACCESS TO TV \rightarrow GO TO Q132
	66. WHEN THE HOUSEHOLD HAS ELECTRICITY
	77. NOT EVERY DAY
	88. DON'T KNOW

129. What channels do you most often watch? (PROBE FOR MORE THAN ONE CHANNEL, DO NOT READ LIST)

	MENTIONED	NOT MENTIONED
A. TVSH	1	2
B. TVKLAN	1	2
C. TVA	1	2
D. TVKOHA	1	2
E. VISION PLUS	1	2
F. SHIJAK TV	1	2
G. TELENORBA	1	2
H. TVALBA	1	2
I. CALVIN	1	2
J. TOP CHANNEL	1	2
K. FOREIGN CHANNELS (e.g. CNN, BBC, EURONEWS, RAI)	1	2
L. NESER TV	1	2
M. OTHER	1	2

130. What types of programs do you most often watch? (PROBE FOR MORE THAN ONE PROGRAM, DO NOT READ LIST)

	MENTIONED	NOT MENTIONED
A. NEWS	1	2
B. COMERCIALS	1	2
C. ENTERTAINMENT PROGRAMS	1	2
D. SERIALS/MOVIES	1	2
E. SPORTS	1	2
F. MUSIC PROGRAMS, VIDEO CLIPS	1	2
G. PLAYS/DRAMAS	1	2
H. CHILDREN'S PROGRAMS	1	2
I. CHURCH/RELIGIOUS PROGRAMS	1	2
J. WOMEN'S PROGRAMS	1	2
K. HEALTH PROGRAMS	1	2
L. POLITICAL EVENTS	1	2
M. OTHER	1	2

131. What times do you most often watch television? (CIRCLE ALL MENTIONED, DO NOT READ LIST)

what times do you most often waten ter		initioned, bo not ki	
	MENTIONED	NOT MENTIONED	
A. 6-8 AM	1	2	
B. 8-10 AM	1	2	
C. 10AM-NOON	1	2	
D. NOON-2 PM	1	2	
E. 2-4 PM	1	2	
F. 4-6 PM	1	2	
G. 6-8 PM	1	2	
H. 8-10 PM	1	2	
I. AFTER 10 PM	1	2	
J. NO REGULAR TIMES	1	2	

- 132. How often do you read a newspaper?
 - 1. DAILY/NEARLY EVERY DAY
 - 2. ABOUT 3-4 TIMES PER WEEK
 - 3. ONCE OR TWICE PER WEEK
 - 4. LESS THAN ONCE PER WEEK
 - 5. NEVER/ALMOST NEVER

II. SEX EDUCATION

The next set of questions is about sex education.

201. Do you think schools should teach courses about human reproduction, contraception, and prevention of sexually transmitted diseases?

1. YES 2. NO \rightarrow GO TO 203 8. DK 9. NR \rightarrow GO TO 203

202. At what year of age should schools begin to teach about? (READ A-C)

A. Human Reproduction?	 77. SHOULD NOT BE TAUGHT IN SCHOOL.
B. Contraception?	 88. DK
C. STD's	 99. NR

GO TO BOX 2-I

203. Now I want to read some reasons for which one may oppose sex education in school. Please tell me if you agree or don't agree. (**READ A-D**)

	AGREE	DISAGREE	<u>DK</u>	<u>NR</u>
A. Sex education will give adolescents the idea to begin sex earlier	1	2	8	9
B. Sex education should be taught only in the house	1	2	8	9
C. Sex education goes against my religious beliefs	1	2	8	9
D. Teachers do not have enough training to teach such courses	1	2	8	9

BOX 2-I IF RESPONDENT IS 15-24 YEARS OF AGE CONTINUE; IF SHE IS 25-44 YEARS GO TO SECTION III

204. Before you were 18 years old, did a parent ever talked to you about.....(READ A-F)

	<u>YES</u>	<u>NO</u>	<u>DK/DR</u>	<u>REF</u>
A. Menstrual Cycle?	1	2	8	9
B. How Pregnancy Occurs?	1	2	8	9
C. Not Having Sexual Intercourse Before Marriage?	1	2	8	9
D. Methods of Contraception?	1	2	8	9
E. HIV/AIDS?	1	2	8	9
F. Other Sexually Transmitted Diseases?	1	2	8	9

READ EACH QUESTION 205-207 FROM THE TABLE FOR EACH TOPIC OF SEX EDUCATION:

TOPIC	205. Before you were 18 years old, have you ever been taught at school about.? (READ A-G)	when you first were taught at	207. Who taught you at school about?
A. Menstrual Cycle	$1 \text{ YES } \rightarrow \text{GO TO Q206}$ $2 \text{ NO } \rightarrow \text{ GO TO Q205}_B$ $8 \text{ DK } \rightarrow \text{ GO TO Q205}_B$ $9 \text{ NR } \rightarrow \text{ GO TO Q205}_B$		1 TEACHER 2 DOCTOR/NURSE 3 VOLUNTEER 7 OTHER 8 DON'T REMEMBER
B. Female Reproductive System	$1 \text{ YES } \rightarrow \text{ GO TO Q206}$ $2 \text{ NO } \rightarrow \text{ GO TO Q205_C}$ $8 \text{ DR } \rightarrow \text{ GO TO Q205_C}$ $9 \text{ NR } \rightarrow \text{ GO TO Q205_C}$		1 TEACHER 2 DOCTOR/NURSE 3 VOLUNTEER 7 OTHER 8 DON'T REMEMBER
C. Male Reproductive System	$1 \text{ YES } \rightarrow \text{ GO TO Q206}$ $2 \text{ NO } \rightarrow \text{ GO TO Q205_D}$ $8 \text{ DR } \rightarrow \text{ GO TO Q205_D}$ $9 \text{ NR } \rightarrow \text{ GO TO Q205_D}$		1 TEACHER 2 DOCTOR/NURSE 3 VOLUNTEER 7 OTHER 8 DON'T REMEMBER
D. How Pregnancy Occurs	$\begin{array}{rcl} 1 \text{ YES } & \rightarrow & \textbf{GO TO Q206} \\ 2 \text{ NO } & \rightarrow & \textbf{GO TO Q205}_E \\ 8 \text{ DR } & \rightarrow & \textbf{GO TO Q205}_E \\ 9 \text{ NR } & \rightarrow & \textbf{GO TO Q205}_E \end{array}$		1 TEACHER 2 DOCTOR/NURSE 3 VOLUNTEER 7 OTHER 8 DON'T REMEMBER
E. Contraceptive Methods	$\begin{array}{rcl} 1 \text{ YES } & \rightarrow & \textbf{GO TO Q206} \\ 2 \text{ NO } & \rightarrow & \textbf{GO TO Q205}_F \\ 8 \text{ DR } & \rightarrow & \textbf{GO TO Q205}_F \\ 9 \text{ NR } & \rightarrow & \textbf{GO TO Q205}_F \end{array}$		1 TEACHER 2 DOCTOR/NURSE 3 VOLUNTEER 7 OTHER 8 DON'T REMEMBER
F. HIV/AIDS	$1 \text{ YES } \rightarrow \text{ GO TO Q206}$ $2 \text{ NO } \rightarrow \text{ GO TO Q205_G}$ $8 \text{ DR } \rightarrow \text{ GO TO Q205_G}$ $9 \text{ NR } \rightarrow \text{ GO TO Q205_G}$		1 TEACHER 2 DOCTOR/NURSE 3 VOLUNTEER 7 OTHER 8 DON'T REMEMBER
G. Other Sexually Transmitted Diseases	$\begin{array}{rrrr} 1 & \text{YES} & \rightarrow & \textbf{GO TO Q206} \\ 2 & \text{NO} & \rightarrow & \textbf{GO TO Q208} \\ 8 & \text{DR} & \rightarrow & \textbf{GO TO Q208} \\ 9 & \text{NR} & \rightarrow & \textbf{GO TO Q208} \end{array}$		1 TEACHER 2 DOCTOR/NURSE 3 VOLUNTEER 7 OTHER 8 DON'T REMEMBER

208. In your opinion, who or what was the most important source of information you have had about topics related to sexual matters?

1. MOTHER	10. NURSE, MIDWIFE
2. FATHER	11. TEACHER
3. RELATIVE	12. PHARMACIST
4. BOYFRIEND	13. BOOKS
5. FRIENDS	14. NEWSPAPERS, MAGAZINES, BROCHURES, FLYERS
6. CO-WORKER	15. RADIO
7. COLLEAGUES, PEER	16. TV
8. PARTNER/HUSBAND	20. OTHER (SPECIFY):
9. DOCTOR	88. DON'T REMEMBER

III. FERTILITY/PREGNANCY

- 300. Are you currently pregnant?
 - 1. YES
 - 2. NO \rightarrow **GO TO Q305**
 - 3. NOT SURE \rightarrow **GO TO Q305**
- 301. How many months pregnant are you now? _____ MONTHS
- 302. Just before you get pregnant, did you want to get pregnant then, did you want to get pregnant later, or did you not want to get pregnant then or any time in the future?
 - WANTED TO GET PREGNANT THEN
 WANTED TO GET PREGNANT LATER
 DID NOT WANT THE PREGNANCY THEN OR ANY TIME IN THE FUTURE
 NOT SURE
- 303. Is this your first pregnancy?
 - 1. YES
 - 2. NO \rightarrow **GO TO Q307**
 - 8. NOT SURE
- 304. Have you ever had a stillbirth, ectopic pregnancy, miscarriage, or an induced abortion?
 - 1. YES \rightarrow GO TO PREGNANCY HISTORY, PAGE 9 2. NO \rightarrow GO TO MODULE IV, PAGE 20
- 305. Have you ever been pregnant?
 - 1. YES \rightarrow GO TO Q307
 - 2. NO
 - 3. NOT SURE
 - 4. NEVER HAD SEX \rightarrow GO TO MODULE IV, PAGE 20
- 306. Have you ever had a stillbirth, ectopic pregnancy, miscarriage, or an induced abortion?
 - 1. YES \rightarrow GO TO PREGNANCY HISTORY, PAGE 9
 - 2. NO \rightarrow GO TO MODULE IV, PAGE 20
- 307. Have you ever had any live-born children?
 - 1. YES

2. NO → GO TO PREGNANCY HISTORY, PAGE 9

308. How many living children do you have, including those who do not live with you?

_____ CHILDREN

- 309. Have you ever had a child born alive who later died, including those who may have died in the first hours or days after birth?
 - 1. YES
 - 2. NO \rightarrow **GO TO PREGNANCY HISTORY, PAGE 9**
- 310.
 How many children died?
 _____ CHILDREN
- 311. So altogether you had a total of ____ (Q308+Q310) live births?
 - 1. YES
 - 2. NO \rightarrow CHECK Q308 AND Q310 AND MAKE CHANGES IF NECESSARY

PREGNANCY HISTORY

Now I would like to talk to you about all your pregnancies (not counting the current one). Please, make sure you include all pregnancies, it doesn't matter when they happened or how they ended, whether in a live birth, an abortion, a miscarriage, or a stillbirth. Starting with your most recent pregnancy, please give me the following information:

<u>#</u>	312 How did that pregnancy end?	did that pregnancy end? (month & year)		315 Was the baby a boy or a girl?	1	LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS;	318 IF Q313B < 1997→GO TO NEXT PREGNANCY Just before you get pregnant, did you want to get pregnant then, did you want to get pregnant later, or did you not want to get pregnant then or any time in the future?
	 MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH 		1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES→ Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
-	2. MULTIPLE LIVE BIRTH	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES →Q318 2. NO	3 YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	 MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCAPPLACE 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES→ Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK 999. NP/REE	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	 MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCAPPLACE 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES →Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK 9999. NP/REE	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE

<u>#</u>	312	313	314	315	316	317	318
		did that pregnancy end? (month & year)	How many weeks or months had you been pregnant when that pregnancy ended?	Was the baby a boy or a girl?		DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS;	IF Q313B <1997→GO TO <u>NEXT PREGNANCY</u> Just before you get pregnant, did you want to get pregnant then, did you want to get pregnant later, or did you not want to get pregnant then or any time in the future?
	2. MULTIPLE LIVE BIRTH 3. MULTIPLE (LB WITH SB) 4. STILLBIRTH (SINGLE) 5. MULTIPLE STILLBIRTH	98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES →Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK 999. NP/REE	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	 STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH 	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES →Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK 900. NB /BEE	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	2. MULTIPLE LIVE BIRTH	BYR 98. DK/NR	OR	1. BOY 2. GIRL 3. BOTH	1. YES→ Q318 2. NO	OR 2 MTHS OR 3YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	2. MULTIPLE LIVE BIRTH	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318		1. YES →Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK	1. WANTED TO GET PREG. THEN 2. WANTED TO GET PREG. LATER 3. DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE 8. NOT SURE
	2. MULTIPLE LIVE BIRTH	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318		1. YES →Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK 900. NB /BEE	1. WANTED TO GET PREG. THEN 2. WANTED TO GET PREG. LATER 3. DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE 8. NOT SURE

#	312 How did that pregnancy end?	did that pregnancy end? (month & year)		315 Was the baby a boy or a girl?	316 Is the child still alive?	LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS;	318 IF Q313B < 1997→GO TO <u>NEXT PREGNANCY</u> Just before you get pregnant, did you want to get pregnant then, did you want to get pregnant later, or did you not want to get pregnant then or any time in the future?
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES →Q318 2. NO	3 YEARS 888. DK 999. NR/REE	1. WANTED TO GET PREG. THEN 2. WANTED TO GET PREG. LATER 3. DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE 8. NOT SURE
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES →Q318 2. NO	3 YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES→ Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	2. MULTIPLE LIVE BIRTH	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES →Q318 2. NO	OR 2 MTHS OR 3 YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE
	2. MULTIPLE LIVE BIRTH	BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO Q318	1. BOY 2. GIRL 3. BOTH	1. YES→ Q318 2. NO	3 YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE

#	312 How did that pregnancy end?	313 When did that pregnancy end? (month & year)	314 How many weeks or months had you been pregnant when that pregnancy ended?	315 Was the baby a boy or a girl?	316 Is the child still alive?	317 How old was the child when he died? (RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YRS.)
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES →GO TO THE NEXT PG. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES→ GO TO THE NEXT PG. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	A MTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES→ GO TO THE NEXT PG. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES →GO TO THE NEXT PG. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF
	2. MULTIPLE LIVE BIRTH	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES→ GO TO THE NEXT PG. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF

			1	1	1	
#	312 How did that pregnancy end?	313 When did that pregnancy end? (month & year)	314 How many weeks or months had you been pregnant when that pregnancy ended?	315 Was the baby a boy or a girl?	316 Is the child still alive?	317 How old was the child when he died? (RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YRS.)
<u>20</u>	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES →GO TO The Next Pg. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES →GO TO The Next pg. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES →GO TO The Next Pg. 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK 999. NR/REF
	 LIVE BIRTH (SINGLE) MULTIPLE LIVE BIRTH MULTIPLE (LB WITH SB) STILLBIRTH (SINGLE) MULTIPLE STILLBIRTH MISCARRIAGE INDUCED ABORTION MOLAR PREGNANCY ECTOPIC PREGNANCY 	AMTH BYR 98. DK/NR	1 WEEKS OR 2 MONTHS 888. DK 999. NR/REF IF Q312>3 GO TO NEXT LINE	1. BOY 2. GIRL 3. BOTH	1. YES→ GO TO THE BOX 3-1 2. NO	1DAYS OR 2MTHS OR 3YEARS 888. DK 999. NR/REF

BOX 3-1

• THE FOLLOWING QUESTIONS ARE ONLY FOR PREGNANCIES ENDED BETWEEN 1997–2002 IF THE RESPONDENT HAD AT LEAST ONE LIVE BIRTH, STILLBIRTH OR AN INDUCED ABORTION (Q312=1–5 or 7) THAT ENDED BETWEEN 1997–2002 THEN <u>CONTINUE WITH Q319</u> ON THE NEXT PAGE;

• IF SHE HAD ONLY MISCARRIAGE(S) OR MOLAR PREGNANCY OR ECTOPIC PREGNANCY(IES) (312=6,8,9), <u>GO TO MODULE IV, PAGE 21 AFTER WRITING "0" IN Q319 AND Q338</u>

• IF SHE DID NOT HAVE ANY PREGNANCY ENDED IN 1997-2001 (CHECK Q313B), GO TO MODULE IV, PAGE 21 AFTER WRITING "0" IN Q319 AND Q338

319. HOW MANY INDUCED ABORTIONS DID THE RESPONDENT HAVE BETWEEN JANUARY 1997 AND THE DATE OF THE INTERVIEW (SEE PAGE 9) INDUCED ABORTIONS (IF NO INDUCED ABORTIONS GO TO 0338)

INDUCED ABORTIONS (IF NO INDUCED ABORTIONS GO TO Q338)						
319A. COPY LINE #. FROM PG. TABLE PAGE 19-13	LAST ABORTION	NEXT TO LAST AB. — —	SECOND TO LAST AB. 	THIRD TO LAST AB. — —		
320 . What was the principal reason that you decided to have this abortion?	 PREGNANCY WAS LIFE OR HEALTH THREATENING RISK OF BIRTH DEFECTS SOCIOECONOMIC REASONS RESPONDENT DID NOT WANT (ANYMORE) CHILDREN SPACING NEXT PREGNANCY PARTNER DID NOT WANT (ANY) CHILDREN DID NOT HAVE A PARTNER OTHER 	 PREGNANCY WAS LIFE OR HEALTH THREATENING RISK OF BIRTH DEFECTS SOCIOECONOMIC REASONS RESPONDENT DID NOT WANT (ANYMORE) CHILDREN SPACING NEXT PREGNANCY PARTNER DID NOT WANT (ANY) CHILDREN DID NOT HAVE A PARTNER OTHER 	 PREGNANCY WAS LIFE OR HEALTH THREATENING RISK OF BIRTH DEFECTS SOCIOECONOMIC REASONS RESPONDENT DID NOT WANT (ANYMORE) CHILDREN SPACING NEXT PREGNANCY PARTNER DID NOT WANT (ANY) CHILDREN DID NOT HAVE A PARTNER OTHER 	 PREGNANCY WAS LIFE OR HEALTH THREATENING RISK OF BIRTH DEFECTS SOCIOECONOMIC REASONS RESPONDENT DID NOT WANT (ANYMORE) CHILDREN SPACING NEXT PREGNANCY PARTNER DID NOT WANT (ANY) CHILDREN DID NOT HAVE A PARTNER OTHER 		
320A. What was the at- titude of the child's father toward you having that abortion?	1. FAVORED 2. OPPOSED 3. NEUTRAL	 FAVORED OPPOSED NEUTRAL DID NOT KNOW ABOUT IT DO NOT REMEMBER 	 FAVORED FAVORED OPPOSED NEUTRAL DID NOT KNOW ABOUT IT DO NOT REMEMBER 	 FAVORED OPPOSED NEUTRAL DID NOT KNOW ABOUT IT DO NOT REMEMBER 		
321. When you got pregnant with this baby, were you using any method of contracep- tion?	 YES NO → GO TO Q323 DK/NR → GO TOQ323 	 YES NO → GO TO Q323 DK/NR → GO TOQ323 	 YES NO → GO TO Q323 DK/NR → GO TOQ323 	 YES NO → GO TO Q323 DK/NR → GO TOQ323 		
322. What method of contracep- tion was that?	1. PILL9. INJECTABLES2. IUD11. CALENDAR3. CONDOM12. WITHDRAWAL4. CONDOM+SP13. CAL+WDR5. CONDM+TRD20. OTHER6. SPERMICIDE88. DR	1. PILL 9. INJECTABLES 2. IUD 11. CALENDAR 3. CONDOM 12. WITHDRAWAL 4. CONDOM+SP 4. CONDOM+TRD 20. OTHER 6. SPERMICIDE 88. DR	1. PILL 9. INJECTABLES 2. IUD 11. CALENDAR 3. CONDOM 12. WITHDRAWAL 4. CONDOM+SP 13. CAL+WDR 5. CONDM+TRD 20. OTHER 6. SPERMICIDE 88. DR	1. PILL9. INJECTABLES2. IUD11. CALENDAR3. CONDOM12.WITHDRAWAL4. CONDOM+SP13. CAL+WDR5. CONDM+TRD20. OTHER6. SPERMICIDE88. DR		
323. Before this abortion, did you have an ultrasound exam of the pregnancy?	 YES NO → GO TO Q325 DK/NR → GO TOQ325 	 YES NO → GO TO Q325 DK/NR → GO TOQ325 	 YES NO → GO TO Q325 DK/NR → GO TOQ325 	 YES NO → GO TO Q325 DK/NR → GO TOQ325 		
324. Did you find out the gender of the baby after the ultrasound?	8. DON'T KNOW/ DR	1. YES 2. NO 8. DON'T KNOW/ DR	1. YES 2. NO 8. DON'T KNOW/ DR	1. YES 2. NO 8. DON'T KNOW/ DR		
325. Where was that abortion performed?	 HOSPITAL/ MATERNITY PRIVATE CLINIC/OFFICE RESPONDENT'S HOME DOCTOR'S HOME AT HOME AND HOSP. OTHER 	 HOSPITAL/ MATERNITY PRIVATE CLINIC/OFFICE RESPONDENT'S HOME DOCTOR'S HOME AT HOME AND HOSP. OTHER 	 HOSPITAL/ MATERNITY PRIVATE CLINIC/OFFICE RESPONDENT'S HOME DOCTOR'S HOME AT HOME AND HOSP. OTHER 	 HOSPITAL/ MATERNITY PRIVATE CLINIC/OFFICE RESPONDENT'S HOME DOCTOR'S HOME AT HOME AND HOSP. OTHER OP/GYN 		
326. Who performed that abortion?	1. OB/GYN 2. OTHER PHYSICIAN 3. NURSE/MIDWIFE 4. LAY PERSON 5. SELF-INDUCED 8. DON'T KNOW/ DR	 OB/GYN OTHER PHYSICIAN NURSE/MIDWIFE LAY PERSON SELF-INDUCED DON'T KNOW/ DR 	1. OB/GYN 2. OTHER PHYSICIAN 3. NURSE/MIDWIFE 4. LAY PERSON 5. SELF-INDUCED 8. DON'T KNOW/ DR	1. OB/GYN 2. OTHER PHYSICIAN 3. NURSE/MIDWIFE 4. LAY PERSON 5. SELF-INDUCED 8. DON'T KNOW/ DR		
327. What method was used?	1. D&C 2. VACUUM ASPIRATION 3. OXITOCIN 4. CATHETER 7. OTHER	 D&C VACUUM ASPIRATION OXITOCIN CATHETER OTHER DON'T KNOW/ DR 	1. D&C 2. VACUUM ASPIRATION 3. OXITOCIN 4. CATHETER 7. OTHER	1. D&C 2. VACUUM ASPIRATION 3. OXITOCIN 4. CATHETER 7. OTHER		

CONTINUE ON NEXT PAGE

	LAST ABORTION	NEXT TO LAST AB.	SECOND TO LAST AB.	THIRD TO LAST AB.
328. How much did you pay for that abortion, including gifts or money given to the doctor?	LEKI 0 0 0 0 0 0 NO CHARGE 7 7 7 7 7 ONLY GIFTS 8 8 8 88 DK	LEKI 0 0 0 0 0 0 NO CHARGE 7 7 7 7 7 ONLY GIFTS 8 8 8 88 DK	LEKI 0 0 0 0 0 0 NO CHARGE 7 7 7 7 7 ONLY GIFTS 8 8 8 88 DK	LEKI 0 0 0 0 0 0 NO CHARGE 7 7 7 7 7 ONLY GIFTS 8 8 8 88 DK
329. Did you have any local or intravenous anesthesia for that abortion? By local anesthesia we mean an injection in the uterus opening.	1. LOCAL (UTERINE CERVIX) 2. INTRAVENOUS 3. NEITHER LOCAL NOR IV 8. DK/DR	1. LOCAL (UTERINE CERVIX) 2. INTRAVENOUS 3. NEITHER LOCAL NOR IV 8. DK/DR	1. LOCAL (UTERINE CERVIX) 2. INTRAVENOUS 3. NEITHER LOCAL NOR IV 8. DK/DR	1. LOCAL (UTERINE CERVIX) 2. INTRAVENOUS 3. NEITHER LOCAL NOR IV 8. DK/DR
330. Did you take any antibiotics after that abortion?	1. YES 2. NO 8. NOT REMEMBER	1. YES 2. NO 8. NOT REMEMBER	1. YES 2. NO 8. NOT REMEMBER	1. YES 2. NO 8. NOT REMEMBER
331. Within 30 days after that abortion did you have any health problems as a result of that abortion?	 YES NO → GO TO Q333 	 YES NO → GO TO Q333 	 YES NO → GO TO Q333 	 YES NO → GO TO Q333
332. Did you have one of the following problems: (READ A-F)	A. Perforation <u>YES NO</u> B. Severe Bleeding 1 2 C. Fever >38 °C 1 2 D. Purulent Discharge 1 2 E. Belly Pain 1 2 F. Other 1 2	A. PerforationYESNOB. Severe Bleeding12C. Fever >38 °C12D. Purulent Discharge12E. Belly Pain12F. Other12	A. PerforationYESNOB. Severe Bleeding12C. Fever >38 °C12D. Purulent Discharge12E. Belly Pain12F. Other12	A. PerforationYESNOB. Severe Bleeding12C. Fever >38 °C12D. Purulent Discharge12E. Belly Pain12F. Other12
333. How many nights did you spend in the hospital after that abortion (+re- admissions during the first month)?	NIGHTS 88 DK	NIGHTS 88 DK	NIGHTS 88 DK	NIGHTS 88 DK
334. Did you have any related health problems more than 6 months later as a result of that abortion?	1. YES 2. NO \rightarrow Q336 3. NOT YET 6 MTH. \rightarrow Q336 8. DON'T REMEMBER \rightarrow Q336	 YES NO → Q336 NOT YET 6 MTH. → Q336 DON'T REMEMBER → Q336 	 YES NO → Q336 NOT YET 6 MTH. → Q336 DON'T REMEMBER → Q336 	 YES NO → Q336 NOT YET 6 MTH. → Q336 DON'T REMEMBER → Q336
335.What was the most important health problem?	BELLY PAIN STERILITY INFECTION LACK OF MENSES IRREGULAR BLEEDING MORE PAINFUL PERIODS OTHER	1. BELLY PAIN 2. STERILITY 3. INFECTION 4. LACK OF MENSES 5. IRREGULAR BLEEDING 6. MORE PAINFUL PERIODS 7. OTHER	1. BELLY PAIN 2. STERILITY 3. INFECTION 4. LACK OF MENSES 5. IRREGULAR BLEEDING 6. MORE PAINFUL PERIODS 7. OTHER	1. BELLY PAIN 2. STERILITY 3. INFECTION 4. LACK OF MENSES 5. IRREGULAR BLEEDING 6. MORE PAINFUL PERIODS 7. OTHER
336. Either before or after the most recent abortion, did a doctor talk to you about contraception?	 YES, BEFORE ABORTION YES, AFTER ABORTION YES, BEFORE & AFTER NO → GO TO Q337A 	 YES, BEFORE ABORTION YES, AFTER ABORTION YES, BEFORE & AFTER NO → GO TO Q337A 	 YES, BEFORE ABORTION YES, AFTER ABORTION YES, BEFORE & AFTER NO → GO TO Q337A 	 YES, BEFORE ABORTION YES, AFTER ABORTION YES, BEFORE & AFTER NO → GO TO Q337A
337. After that abortion, did you receive a method of contraception or prescription for a method?	 RECEIVED A METHOD RECEIVED PRESCRIPTION NO METHOD OR PRESCR. DON'T REMEMBER 	 RECEIVED A METHOD RECEIVED PRESCRIPTION NO METHOD OR PRESCR. DON'T REMEMBER 	 RECEIVED A METHOD RECEIVED PRESCRIPTION NO METHOD OR PRESCR. DON'T REMEMBER 	 RECEIVED A METHOD RECEIVED PRESCRIPTION NO METHOD OR PRESCR. DON'T REMEMBER
337A. After that abortion, did a doctor or nurse refer you to a Family Planning cabinet?	1. YES 2. NO 8. DON'T REMEMBER	1. YES 2. NO 8. DON'T REMEMBER	1. YES 2. NO 8. DON'T REMEMBER	1. YES 2. NO 8. DON'T REMEMBER

HOW MANY BIRTHS HAS THE RESPONDENT HAD BETWEEN JANUARY 1997 AND PRESENT (SEE PG. 11-15) 1. LIVE BIRTHS 2. STILLBIRTHS (IF NO LIVE BIRTH OR STILLBIRTH GO TO MODULE IV PAGE 21) 338.

2. STILLBI	<u>KIHS (</u>	IF NO LIVE BIRTH OF	K STILLBIRTH GC	D TO MODULE IV PAC	3E 21)		
339. COPY LINE #. FROM PREGNANCY TABLE PAGE 9-13	LAST BIRTH		NEXT TO	LAST BIRTH	SECOND	SECOND TO LAST BIRTH	
340. During the 6 mths before you found out you were pregnant, how many cigarettes did you smoke a day, on average?	1. 1-4 (JUST A FEV 2. 5-10 CIGARETT 3. 11 + (MORE TH	V) ES (OR ½ PACK)	1. 1-4 (JUST A FEV 2. 5-10 CIGARETT 3. 11 + (MORE TH	0. NONE → GO TOQ342 1. 1-4 (JUST A FEW) 2. 5-10 CIGARETTES (OR ½ PACK) 3. 11 + (MORE THAN ½ PACK) 8. DON'T REMEMBER → GO TOQ342		0. NONE → GO TOQ342 1. 1-4 (JUST A FEW) 2. 5-10 CIGARETTES (OR ½ PACK) 3. 11 + + (MORE THAN ½ PACK) 8. DON'T REMEMBER → GO TOQ342	
341.On the average, how many cigarettes did you smoke per day after you found out that you were pregnant?		ES (OR ½ PACK) HAN ½ PACK)	0. NONE 1. 1-4 (JUST A FEV 2. 5-10 CIGARETT 3. 11 + (MORE TH 8. DON'T REMEN	TES (OR ½ PACK) AN ½ PACK)	0. NONE 1. 1-4 (JUST A FI 2. 5-10 CIGARET 3. 11 + + (MORE 8. DON'T REME	TES (OR ½ PACK) THAN ½ PACK)	
342. How many times per week did you drink alcoholic beverages during that pregnancy?	 4 TIMES OR M 1-3 TIMES LESS THAN OR NEVER 	ORE /ALMOST DAILY	 4 TIMES OR M 1-3 TIMES LESS THAN OR NEVER 	ORE /ALMOST DAILY NCE PER WEEK	2. 1-3 TIMES	MORE/ALMOST DAILY	
343. When you got pregnant with this baby, were you using any method of contracep- tion?		Q345 IBER → GO TO Q345	 YES NO → GO TO DON'T REMEN 	Q345 ABER \rightarrow GO TO Q345	1. YES 2. NO \rightarrow GO TO 8. DON'T REME	0 Q345 EMBER \rightarrow GO TO Q345	
344. What method of contraception was that?	1. PILL 2. IUD 3. CONDOM 4. CONDOM+SP 5. CONDM+TRAD 6. SPERMICIDES	9. INJECTABLES 11. CALENDAR 12. WITHDRAWAL 13. CAL+WDR 20. OTHER 88. DR	1. PILL 2. IUD 3. CONDOM 4. CONDOM+SP 5.CONDM+TRAD 6. SPERMICIDES	9. INJECTABLES 11. CALENDAR 12. WITHDRAWAL 13. CAL+WDR 20. OTHER 88. DR	1. Pill 2. IUD 3. CONDOM 4. CONDOM+SP 5. CONDM+TRAD 6. SPERMI- CIDES	20. OTHER	
345. How many weeks or months pregnant were you when you learned that you were pregnant that time?	1WEEKS OR 2MONTHS 888 DK/DR		1		1		
346. During that preg- nancy, did you have any prenatal care visits?		Q355 1BER → GO TO Q355	 YES NO → GO TO Q355 DON'T REMEMBER → GO TO Q355 		 YES NO → GO TO Q355 DON'T REMEMBER → GO TO Q355 		
347. How many weeks or months pregnant were you at the time of your first prenatal care visit?	1 WE 2 MO	EKS OR NTHS 888 DK/DR	1 WEEKS OR 2 MONTHS 888 DK/DR			EEKS OR 10nths 888 DK/DR	
348 How many prena- tal visits did you have during that pregnancy?	VISITS	88. DK 99. REF	VISITS	88. DK 99. REF	VISITS	88. DK 99. REF	
349. Where did you receive most of the prenatal care visits?	1. HEALTH POST ("AMBULANTZA") 2. RURAL/URBAN HEALTH CENTER 3. POLICLINIC 4. DISTRICT MATERNITY/HOSPITAL 5. TIRANA MATERNITY 6. PRIVATE CLINIC 7. AT HOME 8. OTHER		 HEALTH POST ("AMBULANTZA") RURAL/URBAN HEALTH CENTER POLICLINIC DISTRICT MATERNITY/HOSPITAL TIRANA MATERNITY PRIVATE CLINIC AT HOME OTHER 		2. RURAL/URBA 3. POLICLINIC		
350. Who provided most of the prenatal care?	 GENERAL PRA OB/GYN NURSE/MIDWI OTHER 		 GENERAL PRA OB/GYN NURSE/MIDWI OTHER 		 GENERAL PR OB/GYN NURSE/MIDW OTHER 		
351. During those vis- its, did you receive any information about: (READ A-H):	A. Nutrition B. Smoking during I C. Drinking Alcohol D. Breastfeeding E. Delivery F. Contraception G. Warning Signs of H. Postnatal Care	during Pg. 1 2 1 2 1 2 1 2 Pg Complic 1 2	A. Nutrition B. Smoking during I C. Drinking Alcohol D. Breastfeeding E. Delivery F. Contraception G. Warning Signs of H. Postnatal Care	Pregnancy 1 2 I during Pg. 1 2 I 2 1 2 I 2 1 2 FPg Complic 1 2 2	A. Nutrition B. Smoking during C. Drinking Alcoh D. Breastfeeding E. Delivery F. Contraception G. Warning Signs H. Postnatal Care	ol during Pg. 1 2 1 2 1 2 1 2 0f Pg Complic 1 2	

	LACT DIDTH	NEVT TO LAST DIDTH	SECOND TO LAST BIDTH
352 . During this preg-	LAST BIRTH <u>YES NO</u>	NEXT TO LAST BIRTH <u>YES NO</u>	SECOND TO LAST BIRTH YES NO
nancy, were any of the	<u>TES NO</u>		<u>115 NO</u>
following done at least once:	A. WEIGHT12B. HEIGHT12	A. WEIGHT12B. HEIGHT12	A. WEIGHT 1 2 B. HEIGHT 1 2
A. Were you eighed?	C. URINE SAMPLE 1 2	C. URINE SAMPLE 1 2	C. URINE SAMPLE 1 2
B. Was your height measured? C. Did you give a urine sample?	D. BLOOD SAMPLE 1 2	D. BLOOD SAMPLE 1 2	D. BLOOD SAMPLE 1 2
D. Did you give a blood sample?			
353 . During those visits, did you have your blood		 YES NO → GO Q355 	 YES NO → GO Q355
pressure measured?	2. NO \rightarrow GO Q355 8. DON'T REMEMBER \rightarrow GO TO Q355	2. NO \rightarrow GO Q355 8. DON'T REMEMBER \rightarrow GO Q355	2. NO \rightarrow GO Q355 8. DON'T REMEMBER \rightarrow GO Q355
354. During those visits,	1. YES	1. YES	1. YES
were you ever told that you have high blood	2. NO	2. NO	2. NO
pressure?	8. DON'T REMEMBER	8. DON'T REMEMBER	8. DON'T REMEMBER
355. Did you have an	1. YES	1. YES	1. YES
ultrasound (US) exam during that pregnancy?	2. NO \rightarrow GO TO Q357 8. DON'T REMEMBER \rightarrow GO TO Q357	2. NO \rightarrow GO TO Q357 8. DON'T REMEMBER \rightarrow GO TO Q357	2. NO \rightarrow GO TO Q357 8. DON'T REMEMBER \rightarrow GO TO Q357
356 . How many weeks	1 WEEKS OR	1WEEKS OR	$1. _ WEEKS OR$
or months pregnant were			I WEEKS OK
you at the time of your first US?	2 MONTHS 888 DK/DR	2 MONTHS 888 DK/DR	2 MONTHS 888 DK/DR
357. During this preg-	1. YES	1. YES	1. YES
nancy, were you given an injection in the arm	2. NO 8. DON'T REMEMBER	2. NO 8. DON'T REMEMBER	2. NO 8. DON'T REMEMBER
to prevent the baby from	. DOW I REMEMBER	. DOI'T REMEMBER	0. DOW I REMEMBER
getting tetanus (convul- sions after birth)?			
358. During this preg-	1. YES	1. YES	1. YES
nancy, have you taken any iron supplements	2. NO \rightarrow GO TO Q360 8. DON'T REMEMBER \rightarrow GO TO Q360	2. NO \rightarrow GO TO Q360 8. DON'T REMEMBER \rightarrow GO TO Q360	2. NO \rightarrow GO TO Q360 8. DON'T REMEMBER \rightarrow CO TO Q360
(iron tablets, injection or	8. DON I REMEMBER \rightarrow GO IO Q300	8. DON I REMEMBER \rightarrow GO IO Q300	8. DON 1 REMEMBER \rightarrow GO 10 Q300
iron syrup) ? 358A. In what week	1 WEEKS OR	1 WEEKS OR	1 WEEKS OR
or month of pregnancy did you start taking iron supplements?	2 MONTHS 888 DK/DR	2 MONTHS 888 DK/DR	2 MONTHS 888 DK/DR
359 . How often did you	1. EVERY DAY	1. EVERY DAY	1. EVERY DAY
take iron supplements?	2. SEVERAL TIMES PER WEEK	2. SEVERAL TIMES PER WEEK	2. SEVERAL TIMES PER WEEK
	 ONCE A WEEK LESS THAN ONCE A WEEK 	 ONCE A WEEK LESS THAN ONCE A WEEK 	 ONCE A WEEK LESS THAN ONCE A WEEK
	8. DK/DR	8. DK/DR	8. DK/DR
360. During that	1. YES	1. YES	1. YES
pregnancy, did you have any complications	2. NO \rightarrow GO TO Q364 8. DON'T REMEMBER \rightarrow GO TO Q364	2. NO \rightarrow GO TO Q364 8. DON'T REMEMBER \rightarrow GO TO Q364	2. NO \rightarrow GO TO Q364 8. DON'T REMEMBER \rightarrow GO TO Q364
that required medical attention?	8. DOIN 1 REMEMBER \rightarrow GO 10 Q304	0 = 0 +	0 = 0 +
361. What complica-	YES NO	YES NO	YES NO
tions did you have? Did you have:	A. Weak Cervix 1 2		A. Weak Cervix 1 2
you nave.	B. Bleeding During First 6 Mth12C. Bleeding at 6 Mths or More12	B. Bleeding During First 6 Mth12C. Bleeding at 6 Mths or More12	B. Bleeding During First 6 Mth12C. Bleeding at 6 Mths or More12
(READ EACH CON-	D. High BP Related to Preg. 1 2		D. High BP Related to Preg. 1 2
DITION FROM A-K)	E. Diabetes Related to Preg. 1 2	E. Diabetes Related to Preg. 1 2	E. Diabetes Related to Preg. 1 2
	F. Water Retention or Edema12G. Anemia Related to Preg.12		F. Water Retention or Edema12G. Anemia Related to Preg.12
		e	H. Urinary Tract Infection 1 2
	I. Risk of Preterm Delivery 1 2 J. Rh Isoimmunization 1 2		I. Risk of Preterm Delivery 1 2 J. Rh Isoimmunization 1 2
			J. Rh Isoimmunization12K. Other12
362 . Not including the			
delivery, how many times were you hospital-	TIMES 00. NEVER HOSP.	TIMES 00. NEVER HOSP.	TIMES 00. NEVER HOSP.
ized for pregnancy	88. DK/DR IF "00" GO TO Q364	88. DK/DR IF "00" GO TO Q364	88. DK/DR IF "00" GO TO Q364
complications?			
363. Altogether, how	NIGHTS 85. 85+ NIGHTS	NIGHTS 85. 85+ NIGHTS	NIGHTS 85. 85+ NIGHTS
many nights were you in the hospital for these	88. DK/DR	88. DK/DR	88. DK/DR
complications?			
	1. DISTRICT MATERNITY HOSPITAL	1. DISTRICT MATERNITY HOSPITAL	1. DISTRICT MATERNITY HOSPITAL
364 . Where did you give birth to this baby?	2. TIRANA MATERNITY 3. PRIVATE CLINIC/HOSPITAL	2. TIRANA MATERNITY 3. PRIVATE CLINIC/HOSPITAL	2. TIRANA MATERNITY 3. PRIVATE CLINIC/HOSPITAL
[4. BIRTH HOUSE/HEALTH CENTER	4. BIRTH HOUSE/HEALTH CENTER	4. BIRTH HOUSE/HEALTH CENTER
	5. AT HOME \rightarrow Q370	5. AT HOME \rightarrow Q370	5. AT HOME \rightarrow Q370
	6. ON THE WAY TO HOSP. \rightarrow Q366	6. ON THE WAY TO HOSP. \rightarrow Q366	6. ON THE WAY TO HOSP. \rightarrow Q366
365 . How many hours	HOURS 85. 85+ HOURS	HOURS 85. 85+ HOURS	HOURS 85. 85+ HOURS
before delivery were	88. DK/DR	88. DK/DR	88. DK/DR
you admitted to the place where you gave			
birth?			

Appendix C

		Appendix C	
	LAST BIRTH	NEXT TO LAST BIRTH	SECOND TO LAST BIRTH
366. How many nights were you in that place after delivery?	NIGHTS	NIGHTS	NIGHTS
367. Where was your nusband or partner at he time of delivery, was he: READ 1- 4)	 In the Delivery Room, At the Hospital/clinic, At Home or with friends At Work Travelling DK/DR 	 In the Delivery Room, At the Hospital/clinic, At Home or with friends At Work Travelling DK/DR 	 In the Delivery Room, At the Hospital/clinic, At Home or with friends At Work Travelling DK/DR
668. Was that baby born by vaginal delivery, orceps, or C-section?	1. VAGINAL DELIVERY \rightarrow GO TO Q370 2. FORCEPS ("APARAT") \rightarrow GO TO Q370 3. CESAREAN SECTION	1. VAGINAL DELIVERY→ GO TO Q370 2. FORCEPS ("APARAT")→ GO TO Q370 3. CESAREAN SECTION	1. VAGINAL DELIVERY→ GO TO Q370 2. FORCEPS ("APARAT")→ GO TO Q370 3. CESAREAN SECTION
369. Do you know what was the most mportant reason that you had to deliver by resarean section ?	BABY TOO BIG (CPD) MALPRESENTATION BABY STARTED TO SUFFER PROLONGED LABOR/FAILED INDUCTION OBSTETRIC HEMORRHAGE PREVIOUS C- SECTION ON REQUEST OON REQUEST DON'T KNOW	BABY TOO BIG (CPD) MALPRESENTATION BABY STARTED TO SUFFER PROLONGED LABOR/FAILED INDUCTION OBSTETRIC HEMORRHAGE PREVIOUS C- SECTION ON REQUEST OON THER B. DON'T KNOW	 BABY TOO BIG (CPD) MALPRESENTATION BABY STARTED TO SUFFER PROLONGED LABOR/FAILED INDUCT. OBSTETRIC HEMORRHAGE PREVIOUS C- SECTION ON REQUEST OTHER BON'T KNOW
70. How long had you been in labor with hat pregnancy (regular contractions 5' apart)	HOURS 77. C-SECTION BEFORE LABOR 88. DK/DR	HOURS 77. C-SECTION BEFORE LABOR 88. DK/DR	HOURS 77. C-SECTION BEFORE LABOR 88. DK/DR
71. Who attended the lelivery of that child?	 PHYSICIAN NURSE/MIDWIFE TRADITIONAL BIRTH ATTENDANT OTHER UNATTENDED 	 PHYSICIAN NURSE/MIDWIFE TRADITIONAL BIRTH ATTENDANT OTHER UNATTENDED 	 PHYSICIAN NURSE/MIDWIFE TRADITIONAL BIRTH ATTEN- DANT OTHER UNATTENDED
72. How much did he baby weigh t birth?	GRAMS → GO TO Q374 8888 DON'T KNOW	GRAMS → GO TO Q374 8888 DON'T KNOW	GRAMS → GO TO Q374 8888 DON'T KNOW
73. Do you know if he baby weighed less han 2500 g or was considered too small?	1. YES, WAS LESS THAN 2500g 2. NO, WAS MORE THAN 2500g 8. DK/DR	1. YES, WAS LESS THAN 2500g 2. NO, WAS MORE THAN 2500g 8. DK/DR	1. YES, WAS LESS THAN 2500g 2. NO, WAS MORE THAN 2500g 3. DK/DR
874. During the first 6 veeks after birth, did you have any of the fol- owing complications: READ A-I)	YES NOA. Severe Bleeding1B. Bad-smelling Vaginal Discharge1C. Infection of Surgical Wound1D. Faint/coma1E. High Fever (39-40c)1F. Painful Urination1G. Painful Uterus (pelvic pain)1H. Breast Infection1I. Other12	YES NOA. Severe Bleeding12B. Bad-smelling Vaginal Discharge12C. Infection of Surgical Wound12D. Faint/coma12E. High Fever (39-40c)12F. Painful Urination12G. Painful Uterus (pelvic pain)12H. Breast Infection12I. Other12	YESNOA. Severe Bleeding12B. Bad-smelling Discharge12C. Infection of Surgical Wound12D. Faint/coma12E. High Fever (39-40c)12F. Painfu Urination12G. Painful Uterus (pelvic pain)12H. Breast Infection12I. Other12
875. After leaving the nospital (DO NOT READ IF HOME DELIVERY) did you nave any postdelivery sheck-ups?	 YES NO → GO TO Q378 DO NOT REMEMBER → GO TO Q378 	 YES NO → GO TO Q378 DON'T REMEMBER → GO TO Q378 	 YES NO → GO TO Q378 DO NOT REMEMBER→GO TO Q378
76. How many days or weeks after the deliv- ry did the first check ake place?	1 DAYS 000. SAME DAY OR 88. DON'T REMEMBER 2. WEEKS	1 DAYS 000. SAME DAY OR 88. DON'T REMEMBER 2. WEEKS	1 DAYS 000. SAME DAY OR 88. DON'T REMEMBER 2. WEEKS
77. During those isit(s) did you receive nformation about: READ A-F)	YES NOA. Breastfeeding1B. Breast Care1C. Child Care1D. Immunization1E. Nutrition1F. Contraception1	YES NOA. Breastfeeding12B. Breast Care12C. Child Care12D. Immunization12E. Nutrition12F. Contraception12	YES NOA. Breastfeeding1B. Breast Care1C. Child Care1D. Immunization1E. Nutrition1F. Contraception1
78. For how many nonths after birth did you not have a period?	MONTHS 88. DK/DR 77. NOT YET	MONTHS 88. DK/DR	MONTHS 88. DK/DR
379. How many months after birth did you re- sume sexual relations?	MONTHS 88. DK/DR 77. NOT YET	MONTHS 88. DK/DR	MONTHS 88. DK/DR

	LAST BIRTH IF STILLBIRTH → NEXT BIRTH	NEXT TO LAST BIRTH IF STILLBIRTH → NEXT BIRTH	SECOND TO LAST BIRTH IF STILLBIRTH → MODULE IV
380. After leaving the hospital (DO NOT READ IF HOME DE- LIVERY) did a health professional check on the baby's health?	 YES NO → GO TO Q382 NO, BABY DIED- → GO TO Q382 DO NOT REMEMBER → GO TO Q382 	1. YES 2. NO \rightarrow GO TO Q382 3. NO, BABY DIED- \rightarrow GO TO Q382 8. DO NOT REMEMBER \rightarrow GO TO Q382	 YES NO → GO TO Q382 NO, BABY DIED- → GO TO Q382 DO NOT REMEMBER → GO TO Q382
381. How many days or weeks after delivery did the first health check take place?	1 DAYS OR 2 WEEKS 000. SAME DAY 888. DO NOT REMEMBER	1 DAYS OR 2 WEEKS 000. SAME DAY 888. DO NOT REMEMBER	1 DAYS OR 2 WEEKS 000. SAME DAY 888. DO NOT REMEMBER
381A. Was the health check because the baby was sick or was a rou- tine health exam?	 HEALTH CHECK FOR SICKNESS ROUTINE HEALTH CHECK DO NOT REMEMBER 	 HEALTH CHECK FOR SICKNESS ROUTINE HEALTH CHECK DO NOT REMEMBER 	1. HEALTH CHECK FOR SICKNESS 2. ROUTINE HEALTH CHECK 8. DO NOT REMEMBER
382. How many days or weeks after the delivery did you register the baby at the city/village council?	1 DAYS OR 2 WEEKS 000. NOT REGISTERED YET 777. BABY DIED AND NOT REGISTERED 888. DO NOT REMEMBER	1 DAYS OR 2 WEEKS 000. NOT REGISTERED YET 777. BABY DIED AND NOT REGISTERED 888. DO NOT REMEMBER	1 DAYS OR 2 WEEKS 000. NOT REGISTERED YET 777. BABY DIED AND NOT REGISTERED 888. DO NOT REMEMBER
383. Did you breast- feed?	1. YES 2. NO → GO TO Q387 3. NO, INFANT DIED → NEXT BIRTH	1. YES 2. No → Go to Q387 3. No, infant died → Next Birth	1. YES 2. NO \rightarrow GO TO Q387 3. NO, INFANT DIED \rightarrow MODULE IV
384. How long after birth did you start breastfeeding?	1HOURS 777. LESS THAN 1HR OR 0R 2DAYS 888. DON'T REMEMB.	1HOURS 777. LESS THAN 1HR OR 2DAYS 888. DON'T RE- MEMB.	1 HOURS 777. LESS THAN 1HR OR 2 DAYS 888. DON'T REMEMB.
385. Are you still breastfeeding?	1. YES \rightarrow GO TO Q387 2. NO 3. NO, INFANT DIED \rightarrow NEXT BIRTH		
386. How old was the baby when you stopped breastfeeding?	1 DAYS OR 2 WEEKS OR 888. DK/DR 3 MTHS	1 DAYS OR 2 WEEKS OR 888. DK/DR 3 MTHS	1 DAYS OR 2 WEEKS OR 888. DK/DR 3 MTHS
387. How old was the baby when you gave him/her water or other liquids?	 WEEKS OR 888. DK/DR 	1DAYS OR 777. NOT YET 2WEEKS OR 888. DK/DR 3MTHS 888. DK/DR	1. DAYS_OR 777. NOT YET 2. WEEKS OR 888. DK/DR 3. MTHS
388. How old was the baby when you started feeding with formula or other milk?	1DAYS_OR 777. NOT YET 2WEEKS OR 888. DK/DR 3MTHS 888. DK/DR	1. DAYS_OR 777. NOT YET 2. WEEKS OR 888. DK/DR 3. MTHS	1. DAYS_OR 777. NOT YET 2. WEEKS OR 888. DK/DR 3. MTHS
389. How old was the baby when you started feeding with solid or semi-solid food?	1 WEEKS OR777. NOT YET2 MTHS888. DK/DRIF STILL BREASTFEEDING \rightarrow GO TO Q391	1WEEKS OR 777. NOT YET 2MTHS 888. DK/DR	1WEEKS OR 777. NOT YET 2MTHS 888. DK/DR
390. Why did you stop breastfeeding this baby? FOR WOMEN WHO DID NOT BREAST- FEED (Q383=2) ASK: Why did you not breast- feed this baby?	MOTHER WEAK/ILL CHILD WEAK/ILL CHILD DIED NIPPLE/BREAST PROBLEMS NOT ENOUGH MILK MOTHER WORKING CHILD REFUSED BECAME PREGNANT WEANING AGE/AGE TO STOP PREFERRED BOTTLE-FEEDING OTHER	MOTHER WEAK/ILL CHILD WEAK/ILL CHILD DIED NIPPLE/BREAST PROBLEMS NOT ENOUGH MILK MOTHER WORKING CHILD REFUSED BECAME PREGNANT WEANING AGE/AGE TO STOP PREFERRED BOTTLE-FEEDING OTHER 8. DK/DR	MOTHER WEAK/ILL CHILD WEAK/ILL CHILD DIED NIPPLE/BREAST PROBLEMS NOT ENOUGH MILK MOTHER WORKING CHILD REFUSED BECAME PREGNANT WEANING AGE/AGE TO STOP PREFERRED BOTTLE-FEEDING OTHER
	NEXT BIRTH	NEXT BIRTH	GO TO MODULE IV

MODULE IV: FAMILY PLANNING KNOWLEDGE/ SEXUAL EXPERIENCE

400. Have you ever	401. Do you know	402. Have you ever	403. Do you know	404. What was the most important source
heard of it?	how to use it?	used it?	where to get it?	of information about this method (SEE CODES BELOW)
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \; \mathrm{YES} \to \mathrm{Q402}$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow Q404$	
$2 \text{ NO} \rightarrow \text{B}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{ Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow \text{Q402}$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$	
$2 \text{ NO} \rightarrow \text{C}$	$2 \text{ NO} \rightarrow Q402$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow \text{Q402}$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$	
$2 \text{ NO} \rightarrow \text{D}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{ Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow \text{Q402}$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$	
$2 \text{ NO} \rightarrow \text{E}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{ Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow Q402$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$	
$2 \text{ NO} \rightarrow F$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{ Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow Q402$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$	
$2 \text{ NO} \rightarrow \text{G}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{ Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow Q402$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$	
$2 \ \mathrm{NO} \to \mathrm{H}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{ Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow Q402$	$1 \text{ YES} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$	
$2 \text{ NO} \rightarrow \text{I}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q403}$	$2 \text{ NO} \rightarrow \text{ Q404}$	
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow \text{Q402}$	$1 \text{ YES} \rightarrow \text{Q404}$		
$2 \ \mathrm{NO} \to \mathrm{J}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q404}$		
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow \text{Q402}$	$1 \text{ YES} \rightarrow \text{Q404}$		
$2 \ \mathrm{NO} \to \mathrm{K}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q404}$		
$1 \text{ YES} \rightarrow \text{Q401}$	$1 \text{ YES} \rightarrow \text{Q402}$	$1 \text{ YES} \rightarrow \text{Q404}$		
$2 \text{ NO} \rightarrow \text{Q405}$	$2 \text{ NO} \rightarrow \text{Q402}$	$2 \text{ NO} \rightarrow \text{ Q404}$		
	heard of it? 1 YES \rightarrow Q401 2 NO \rightarrow B 1 YES \rightarrow Q401 2 NO \rightarrow C 1 YES \rightarrow Q401 2 NO \rightarrow D 1 YES \rightarrow Q401 2 NO \rightarrow E 1 YES \rightarrow Q401 2 NO \rightarrow F 1 YES \rightarrow Q401 2 NO \rightarrow H 1 YES \rightarrow Q401 2 NO \rightarrow H 1 YES \rightarrow Q401 2 NO \rightarrow J 1 YES \rightarrow Q401 2 NO \rightarrow K	heard of it?how to use it? $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $2 ext{NO} ext{B}$ $2 ext{NO} ext{Q402}$ $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $2 ext{NO} ext{C}$ $2 ext{NO} ext{Q402}$ $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $2 ext{NO} ext{D}$ $2 ext{NO} ext{Q402}$ $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $2 ext{NO} ext{D}$ $2 ext{NO} ext{Q402}$ $2 ext{NO} ext{Q401}$ $1 ext{YES} ext{Q402}$ $2 ext{NO} ext{Q402}$ $1 ext{YES} ext{Q402}$ $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $2 ext{NO} ext{Q402}$ $2 ext{NO} ext{Q402}$ $1 ext{YES} ext{Q401}$ $1 ext{YES} ext{Q402}$ $2 ext{NO} ext{Q402}$ ext{Q402}$ <	heard of it?how to use it?used it? $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $2 NO \rightarrow B$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $2 NO \rightarrow C$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $2 NO \rightarrow C$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow D$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow D$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow E$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow E$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow E$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow F$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow G$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $2 NO \rightarrow G$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $2 NO \rightarrow H$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q404$ $2 NO \rightarrow J$ $2 NO \rightarrow Q402$ $1 YES \rightarrow Q404$ $2 NO \rightarrow J$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q404$ $2 NO \rightarrow K$ $2 NO \rightarrow Q402$ $1 YES \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q404$ $1 YES \rightarrow Q404$ $2 NO \rightarrow K$ $2 NO \rightarrow Q402$ $1 YES \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q404$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q404$ $1 YES \rightarrow Q404$ $2 NO \rightarrow Q404$	heard of it?how to use it?used it?where to get it? $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow B$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow C$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow D$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $2 NO \rightarrow D$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow E$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $2 NO \rightarrow E$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow F$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $2 NO \rightarrow G$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow H$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow H$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q403$ $1 YES \rightarrow Q404$ $2 NO \rightarrow J$ $2 NO \rightarrow Q402$ $2 NO \rightarrow Q403$ $2 NO \rightarrow Q404$ $1 YES \rightarrow Q401$ $1 YES \rightarrow Q402$ $1 YES \rightarrow Q404$ $2 NO \rightarrow Q404$

For each of the following methods of preventing pregnancy, please tell me:

CODES FOR Q404 (DO NOT READ)

- 1. MOTHER 2. FATHER 3. RELATIVE 4. BOYFRIEND 5. FRIENDS 6. CO-WORKER 7. COLLEAGUES, PEER 16. RADIO 8.PARTNER/HUSBAND 9. DOCTOR
- 10. NURSE, MIDWIFE, FELDCHER
 - 11. COMMUNITY HEALTH WORKER
 - 12. TEACHER
 - 13. PHARMACIST
 - 14. BOOKS
 - 15 NEWSPAPERS, MAGAZINES, BROCHURE, FLYERS

 - 17. TV
 - 20. OTHER (SPECIFY):
 - 88. DON'T REMEMBER

405. Looking at this CARD, please tell me which do you think is the most effective contraceptive method? (SHOW CARD A)

- 1. The Pill
- 2. IUD
- 3. Condom
- 6. Foams/jelly/creams/Foamy Tablets
- 7. Tubal Ligation
- 8. Emergency Hormonal Contraception ("Morning After Pill")
- 9. Injectables (Depo-Provera)
- 10. Vasectomy (Male Sterilization)
- 11. Rhythm Method
- 12. Withdrawal
- 77. NONE OF THEM
- 88. DON'T KNOW/NOT SURE
- 406. How would you rate each of the following methods with regard to effectiveness at preventing pregnancy? Would you say tha _________ is very effective, effective, somewhat effective, not very effective or not at all effective? (INTERVIEWER: ASK THE QUESTION FOR EACH OF THE METHODS LISTED BELOW, UNLESS RESPONDENT HAS SAID IN Q400 THAT SHE NEVER HEARD OF THAT SPECIFIC METHOD; MARK "9" FOR THOSE CASES WITHOUT ASKING)

	Very <u>Effective</u>	<u>Effective</u>	Somewhat <u>Effective</u>	Not Very <u>Effective</u>	Not <u>Effective</u>	DO NOT <u>KNOW</u>	NEVER <u>HEARD</u>
1. The Pill	1	2	3	4	5	8	9
2. Iud	1	2	3	4	5	8	9
3. Condom	1	2	3	4	5	8	9
7. Tubal Ligation	1	2	3	4	5	8	9
9. Injectables	1	2	3	4	5	8	9
11. Calendar	1	2	3	4	5	8	9
12. Withdrawal	1	2	3	4	5	8	9

BOX 4-I

IF RESPONDENT IS 15-24 YEARS OF AGE CONTINUE; IF SHE IS 25-44 YEARS GO TO Q420 PAGE 24

408. How old were you when you had your first menstruation _____AGE

00. NOT YET 88. DON'T REMEMBER 99. REFUSE TO ANSWER

409. Did you know what menstruation was at that time?

- 1. YES 2. NO
- 8 NO SR410.

410.	Now I have some questions about your first sexual intercourse. When did you have sexual intercourse for the first time - in what month and year was that? (PROBE: Can you tell me what year that was?)								
	88. D0	EVER HAD INTERCOURSE → GO TO Q601, PG 36 On't Remember EFUSE TO ANSWER							
411.	How old were you at that time? Y	EARS 88. DON'T REMEMBER							
412.	At the time you first had sexual intercourse, what	was your relationship to that man?							
	 HUSBAND, CONSENSUAL PARTNER FIANCEE BOYFRIEND FRIEND LOVER ACQUAINTANCE 	7. JUST MET 8. RELATIVE 9. RAPE \rightarrow GO TO Q421 10 INCEST \rightarrow GO TO Q421 20. OTHER(SPECIFY) 88. DO NOT REMEMBER/REF							
413.	How long were you and your first partner dating	when you first had sexual relations?							
	1DAYS OR 2WEEKS OR 000=FIRST TIME WE MET 888=DON'T REMEMBER 999=NO RESPONSE 777=OTHER	3MONTHS OR 4YEARS							
414.	How old was your first partner?	YEARS 88. DK/DR							
415.	Before you had sex for the first time, did you and 1. YES 2. NO 8. DON'T REMEMBER	your partner ever talk about using contraception?							
416.	At the time you had first sexual intercourse, did y 1. YES 2. NO \rightarrow GO TO Q419 8. DK/DO NOT REMEMBER \rightarrow GO TO Q421 9. REF \rightarrow GO TO Q421								
417.	 Which contraceptive method did you or your part 1 THE PILL 2 IUD 3 CONDOM 6 FOAM/JELLY/CREAM/VAGINAL FILMS 8 EMERGENCY HORMONAL CONTRACEP' 9 INJECTABLES 10 OTHER MODERN METHODS 11 CALENDAR METHOD 12 WITHDRAWAL 19 DOUCHE 20 OTHER:	TION							

418. Who made the decision to use contraception at that time? (**READ 1-3**)

- 1. You
- 2. Your partner
- 3. Both you and your partner
- 8. DON'T REMEMBER

GO TO Q421

- 419. What was the main reason for not using a contraceptive method at that time?
 - 1. SEX WAS NOT EXPECTED
 - 2. THOUGHT IT WAS A SAFE TIME OF THE MONTH
 - 3. DID NOT KNOW WHERE TO GET A METHOD//DIFFICULT TO GET/NOT AVAILABLE
 - 4. RESPONDENT WAS AGAINST IT
 - 5. PARTNER WAS AGAINST IT
 - 6. DID NOT KNOW ABOUT CONTRACEPTION
 - 7. WANTED TO GET PREGNANT
 - 8. THOUGHT THAT CONTRACEPTIVE METHODS ARE HARMFUL
 - 9. DID NOT THINK ABOUT USING A METHOD/NEGLIGENCE
 - 10. RESPONDENT AFRAID OF PARTNER'S REACTION
 - 11. TOO DRUNK (PARTNER OR RESPONDENT)
 - 12. RESPONDENT WAS TOO EMBARRASSED TO USE A METHOD
 - 20. OTHER (SPECIFY)
 - 88. DON'T REMEMBER/DON'T KNOW

GO TO Q421

420. How old were you at the time of your first sexual intercourse?

_____YEARS

00. NEVER HAD INTERCOURSE- \rightarrow GO TO Q601 PAGE 36 88. DK/DR

- 421. During the past 30 days (past month) have you had sexual intercourse?
 - 1. YES 2. NO \rightarrow **GO TO Q436** 9. REF \rightarrow **GO TO Q436**
- 422. How often have you had sexual intercourse during the past 30 days (**READ 1-5**)?
 - 1. Every day
 - 2. 3-5 times per week,
 - 3. 1-2 times per week,
 - 4. 2-3 times per month, or
 - 5. Only once
 - 9. REF

GO TO Q445

436. During the past 3 months, have you had sexual intercourse? 1. YES 2. NO \rightarrow GO TO Q453 9. REF \rightarrow **GO TO Q453** 445. During the past 3 months, with how many different men have you had intercourse? Please count every sexual partner, even those you had sex with only once. PARTNERS 88. DK 99. NR When was the last time you had sexual intercourse? 453. ____YEAR MONTH 88. DK/DR 99. REFUSE TO ANSWER 454. At the time of your last intercourse, what was your relationship with that man? 1. HUSBAND, CONSENSUAL PARTNER 7. JUST MET 2. FIANCEE 8. RELATIVE 3. BOYFRIEND 9. RAPE \rightarrow GO TO Q460 10..INCEST \rightarrow GO TO Q460 4. FRIEND 20. OTHER(SPECIFY) 5. LOVER 88. DO NOT REMEMBER/REF 6. ACQUAINTANCE 455. At the time you had your last sexual intercourse, did you or your partner use any contraceptive method? 1. YES 2. NO \rightarrow **GO TO Q460** 9. DK/REF \rightarrow GO TO Q460 456. Which contraceptive method did you or your partner use at the last intercourse? 1. THE PILL 2. IUD 3. CONDOM 4. CONDOM +SPERMICIDE 5. CONDOM +WITHDRAWAL/CALENDAR 6. FOAM/JELLY/CREAMS/C-FILMS 7. FEMALE STERILIZATION 8. EMERGENCY HORMONAL CONTRACEPTION 9. INJECTABLES(DEPO PROVERA) 10. OTHER MODERN METHODS 11. CALENDAR 12. WITHDRAWAL 13. WITHDRAWAL AND CALENDAR 20. OTHER TRADITIONAL METHODS 88. NOT SURE 460. Counting all the men you had sexual intercourse with, even only once, how many partners have you had sexual intercourse with in your life? PARTNERS 85. 85+ PARTNERS 88. DK

99. NR

V. CURRENT AND PAST CONTRACEPTIVE USE

- 501. RECORD WHETHER RESPONDENT REPORTED HAVING USED ANY METHOD (ANY Q402=1 AT PG.20)
 1 NEVER USED (NO Q402=1)
 2 EVER USED (ANY Q402=1)- → GO TO Q503
- 502. So, you said that you or any of your partners have never used any method to prevent pregnancy?
 1. NEVER USED → GO TO Q515, PAGE 27
 2. EVER USED → CORRECT Q402 THEN CONTINUE
- 503. Are you (or your partner) currently using (in the last 30 days) any method or doing anything to prevent pregnancy?
 - 1. YES
 - 2. NO \rightarrow GO TO Q515 PAGE 27
- 504. What method are you currently using?
- 1. THE PILL
- 2. IUD 3. CONDOM \rightarrow **GO TO Q506**
- 4. CONDOM +SPERMICIDE \rightarrow GO TO Q506
- 5. CONDOM +WITHDRAWAL/CALENDAR \rightarrow GO TO Q506
- 6. FOAM/JELLY/CREAMS/C-FILMS
- 7. FEMALE STERILIZATION
- 8. EMERGENCY HORMONAL CONTRACEPTION
- 9. INJECTABLES(DEPO PROVERA)
- 10. OTHER MODERN METHODS
- 11. CALENDAR
- 12. WITHDRAWAL
- 13. WITHDRAWAL AND CALENDAR
- 20. OTHER TRADITIONAL METHODS
- 88. NOT SURE
- 505. In the last 30 days, did you and your partner ever use a condom in addition to the method you are using? 1. YES
 - 2. NO

IF Q504=1,2,7, 9, 10, 11, or 88 GO TO Q507

- 506. In the last 30 days how often did you/your partner use this method (METHOD MENTIONED IN Q504) (READ 1-4)?
 - 1. Always, at each sexual intercourse,
 - 2. almost always,
 - 3. Sometimes,
 - 4. Only once
 - 9. REF
- 507. What was the most important reason for choosing this method?
 - 1. DOCTOR RECOMMENDED
 - 2. AFFORDABLE COST
 - 3. VERY EFFECTIVE
 - 4. VERY SAFE (FEW SIDE EFFECTS)
 - 5. SAW ADS (TV, RADIO, PRESS, BROCHURES)
 - 6. EASY TO USE
 - 7. PARTNER PREFERS IT
 - 8. KNOWS SOMEBODY WHO USES IT
 - 9. CURIOSITY/WANTED TO TRY IT
 - 10. ALLOWS SPONTANEITY DURING INTERCOURSE
 - 20. OTHER_
 - 88. DO NOT KNOW/ DO NOT REMEMBER

BOX 5-I

IF Q504 = 1-10, OR 88 GO TO Q510; IF SHE USES NATURAL METHODS (Q504 =11-20), CONTINUE

Appendix C

508.	Please tell me whether each of the followin	g reasons was very important, somewhat important, or not
	important at all in your decision to use	(CODE FROM Q504 FOR TRADITIONAL METHOD)
	instead of a modern method:	

	Very <u>Important</u>	Somewhat <u>Important</u>	Not <u>Important</u>	Not <u>Sure</u>
A. Difficult to get a modern method	1	2	3	8
B. Cost of these modern methods	1	2	3	8
C. Little knowledge of modern methods	1	2	3	8
D. Fear of or experience with side effects	1	2	3	8
E. Husband/Partner choice	1	2	3	8
F. Religious beliefs	1	2	3	8
G. Doctor's recommendation	1	2	3	8
H. Another person's advice	1	2	3	8

509. How effective at preventing pregnancy do you think _____ (CODE FROM Q504 FOR TRADITIONAL METHOD) is compared to modern methods, like the pill or the IUD? (READ 1-3)

- 1. Current method more effective
- 2. About equally effective
- 3. Current method less effective
- 8. DON'T KNOW/NOT SURE

510. Do you have any problems or concerns with using your current method?

- 1. YES
 - 2. NO \rightarrow GO TO Q512
- 511. What is the most important problem?
 - 1. SIDE EFFECTS
 - 2. HEALTH CONCERNS
 - 3. ACCESS/AVAILABILITY
 - 4. COST
 - 5. SOMETIMES FORGET TO USE
 - 6. SOMETIMES DIFFICULT/INCONVENIENT TO USE
 - 7. HUSBAND/PARTNER DISAPPROVES
 - 8. LESS EFFECTIVE METHOD/GOT PREGNANT WHILE USING IT
 - 9. DEEPLY UNSATISFIED WITH THE METHOD
 - 0. OTHER
- 512. Would you prefer to use a different method of family planning from the one you are currently using?1. YES
 - 2. NO \rightarrow GO TO BOX 5-II
- 513. What method would you prefer to use (OTHER THAN THE METHOD SPECIFIED IN Q504)?
 - 1. THE PILL
 - 2. IUD
 - 3. CONDOM
 - 4. CONDOM +SPERMICIDE
 - 5. CONDOM +WITHDRAWAL/CALENDAR-
 - 6. FOAM/JELLY/CREAMS/C-FILMS
 - 7. FEMALE STERILIZATION
 - 8. EMERGENCY HORMONAL CONTRACEPTION
 - 9. INJECTABLES (DEPO PROVERA)
 - 10. OTHER MODERN METHODS____
 - 11. CALENDAR
 - 12. WITHDRAWAL
 - 13. WITHDRAWAL AND CALENDAR
 - 20. OTHER TRADITIONAL METHODS____
 - 88. DO NOT NOW/NO SR514.

- 514. What is the most important reason that you do not use that method?
 - 1. DOCTOR WILL NOT PRESCRIBE IT
 - 2. COST
 - 3. NOT AVAILABLE/UNRELIABLE SUPPLIES/DIFFICULT TO OBTAIN
 - 4. TOO FAR AWAY
 - 5. DO NOT KNOW HOW/WHERE TO OBTAIN IT
 - 6. HUSBAND/PARTNER OBJECTS TO IT
 - 7. RELIGIOUS REASONS
 - 8. FEAR OF SIDE EFFECTS
 - 9. HAS NOT YET MADE UP HER MIND
 - 10. DIFFICULT TO USE
 - 11. FEAR OF SURGICAL PROCEDURE (IUD, TL, NORPLANT)
 - 20. OTHER_____ 88. DON'T KNOW
- BOX 5-II GO TO Q 521 PAGE 28
- 515. What is the main reason that you or your partner are not currently using a contraceptive method?
 - 1. DOES NOT CURRENTLY HAVE A PARTNER/ NOT SEXUALLY ACTIVE IN THE LAST MONTH
 - 2. TRYING TO GET PREGNANT
 - 3. POSTPARTUM/ BREASTFEEDING
 - 4. CURRENTLY PREGNANT
 - 5. HYSTERECTOMY/MENOPAUSE \rightarrow GO TO Q523
 - 6. DOCTOR SAID SHE OR HER PARTNER CANNOT HAVE CHILDREN \rightarrow GO TOQ523
 - 7. SHE/COUPLE TRIED TO GET PREGNANT FOR AT LEAST 2 YEARS AND DIDN'T SUCCEED \rightarrow Q523
 - 8. FEAR OF SIDE EFFECTS
 - 9. LOVEMAKING WOULD BE INTERRUPTED
 - 10. RESPONDENT DID NOT THINK ABOUT USING CONTRACEPTION
 - 11. COST, CANNOT AFFORD BIRTH CONTROL
 - 12. BIRTH CONTROL IS THE PARTNER'S RESPONSIBILITY
 - 13. BIRTH CONTROL IS NOT (VERY) EFFECTIVE
 - 14. RESPONDENT DOES NOT WANT TO USE A METHOD
 - 15. PARTNER OBJECTS TO USING METHOD
 - 16. OBJECTS DUE TO RELIGIOUS REASONS
 - 17. DOES NOT KNOW WHERE TO GET METHOD
 - 18. RESPONDENT DOES NOT KNOW HOW TO USE BIRTH CONTROL METHODS
 - 19. RESPONDENT DOES NOT THINK THAT SHE CAN GET PREGNANT
 - 20. RESPONDENT USES DOUCHING
 - 77. OTHER (SPECIFY)
 - 88. DK/REF
- 516. Do you think that you will use a contraceptive method during the next 12 months (ADD:OTHER THAN DOUCHING IF Q515=20)?
 - 1. YES \rightarrow GO TO Q518
 - 2. NO
 - 8. NOT SURE
- 517. Do you think that you will use a contraceptive method any time in the future?
 - 1. YES
 - 2. NO \rightarrow **GO TO Q521**
 - 8. NOT SURE \rightarrow GO TO Q521
- 518. What method would you want to use most?
 - 1. THE PILL
 - 2. IUD
 - 3. CONDOM
 - 4. CONDOM +SPERMICIDE
 - 5. CONDOM +WITHDRAWAL/CALENDAR-
 - 6. FOAM/JELLY/CREAMS/C-FILMS
 - 7. FEMALE STERILIZATION
 - 8. EMERGENCY HORMONAL CONTRACEPTION
 - 9. INJECTABLES(DEPO PROVERA)
 - 10. OTHER MODERN METHODS_____
 - 11. CALENDAR \rightarrow GO TO Q521
 - 12. WITHDRAWAL \rightarrow GO TO Q521
 - 13. WITHDRAWAL + CALENDAR \rightarrow GO TO Q521
 - 20. OTHER \rightarrow GO TO Q521
 - 88. NOT SURE \rightarrow GO TO Q521

- 519.
 On average, how much are you willing to pay for contraception, per month?

 _____HUNDRED LEKE
 85. 85 HUNDRED LEKE OR MORE

 98.
 NOT SURE/DON'T KNOW

 520.
 Where would you want to get your contraceptive method?

 1.
 HEALTH POST

 2.
 HEALTH CENTER

 3.
 POLICLINIC

 4.
 FAMILY PLANNING CLINIC
 - 5. GOV HOSPITAL-MATERNITY WARD
 - 6. PRIVATE CLINIC OR OFFICE
 - 7. NGO

STORE/ KIOSK
 PARTNER/HUSBAND
 FRIEND
 RELATIVE
 OTHER (SPECIFY): _
 BON'T KNOW

- 521. During the last year, how often did you talk about contraception with your husband/ partner? 1. NEVER \rightarrow GO TO Q523
 - 2. ONE OR TWO TIMES
 - 3. THREE TIMES OR MORE
 - 4. RESPONDENT HAD NO PARTNER DURING THE LAST YEAR \rightarrow GO TO Q523
- 522. Generally, does your husband/ partner agree or disagree with the use of contraceptive methods?
 - 1. AGREES
 - 2. DISAGREES
 - 3. NEITHER AGREES NOR DISAGREES
 - 8. NOT SURE/DON'T KNOW
- 523. Some people use condoms for reasons other than birth control, for instance because they are concerned about getting diseases that can result from sexual intercourse. Have you ever used condoms for: (**READ 1-4**)
 - 1. Birth Control Only \rightarrow **GO TO BOX 5-III**
 - 2. Disease Prevention Only \rightarrow GO TO BOX 5-III
 - 3. Both, or \rightarrow **GO TO BOX 5-III**
 - 4. You Never Used a Condom?
 - 5. USED CONDOM OUT OF CURIOSITY
 - 8. NOT SURE/ DO NOT REMEMBER
- 524. Why have you and your partner(s) never used condoms?
 - 1. PREVENTING PREGNANCY IS WOMAN'S RESPONSIBILITY
 - 2. PARTNER(S) OBJECTED TO USE CONDOMS
 - 3. HAVE ONLY ONE PARTNER
 - 4. THEY ARE ONLY FOR USE WITH PROSTITUTES
 - 5. THEY ARE ONLY FOR EXTRAMARITAL RELATIONS
 - 6. CONDOMS DIMINISH PLEASURE/SPONTANEITY
 - 7. CONDOMS ARE LESS EFFECTIVE IN PREVENTING PREGNANCY
 - 8. CONDOMS ARE TOO DIFFICULT TO USE
 - 9. LOVEMAKING WOULD BE INTERRUPTED
 - 10. CONDOM USE IS TOO MESSY
 - 11. COST
 - 12. SHE HAS NEVER THOUGHT ABOUT IT
 - 13. IT IS EMBARRASSING TO BUY CONDOMS
 - 14. PREFERS OTHER CONTRACEPTIVE METHODS
 - 20. OTHER___
 - 88. DON'T KNOW

BOX 5-III

FIRST COMPLETE COLUMN 1 AND 4 (SEE PREGNANCY HISTORY PG.9-13 AND MARITAL HISTORY PAGE 3). USING INFO FROM THE COMPLETED COLUMNS, ASK THE MONTH-BY-MONTH CONTRACEPTIVE HISTORY STARTING WITH THE CURRENT MONTH AND GOING BACK TO JANUARY 1997.

 \rightarrow IF RESPONDENT HAS USED ANY CONTRACEPTIVE METHOD SINCE JANUARY 1997, FILL IN ALL FOUR COLUMNS OF THE CALENDAR

→ IF NO METHOD HAS BEEN USED SINCE JANUARY 1997, WRITE "0" AT THE BEGINNING AND THE END OF THE 2ND COLUMN THEN GO TO Q551, PAGE 33

<u>COLUMN 1</u>	DATE	1	2	3	4	DATE	1	2	3	4
PREGNANCY OUTCOME	1997					2000				
1. PREGNANT THAT MONTH	1 Jan					1 Jan				\square
2. LIVE BIRTH	2 Feb					2 Feb				\square
4. STILLBIRTH	3 Mar					3 Mar				
6. MISCARRIAGE	4 Apr					4 Apr				
7. INDUCED ABORTION	5 May					5 May				
8. MOLAR PREGNANCY	6 Jun					6 Jun				
9. ECTOPIC PREGNANCY	7 Jul					7 Jul				
COLUMN 2	8 Aug					8 Aug				
METHOD USED	9 Sep		İ			9 Sep				
0. NO METHOD	10 Oct					10 Oct				
1. PILL	11 Nov					11 Nov				
2. IUD	12 Dec					12 Dec				
3. CONDOM		199	98				2001			
4. CONDOM+SPERMICIDES	1 Jan					1 Jan				
5. CONDOM+CAL./WITHDRAWAL	2 Feb					2 Feb				
6. SPERMICIDES	3 Mar					3 Mar				
7. TUBAL LIGATION	4 Apr					4 Apr				
8. EMERGENCY HORM. CONTRACEPTION	5 May					5 May				
9. INJECTABLES (E.G. DEPO-PROVERA)	6 Jun					6 Jun				
10. OTHER MODERN METHODS	7 Jul					7 Jul				
11. CALENDAR	8 Aug					8 Aug				
12. WITHDRAWAL	9 Sep					9 Sep				
13. WITHDRAWAL +CALENDAR	10 Oct					10 Oct				
20. OTHER TRADITIONAL MET	11 Nov					11 Nov				
88. DO NOT REMEMBER	12 Dec					12 Dec				
<u>COLUMN 3</u>		199	99			2002				
REASON STOPPED USING A METHOD	1 Jan					1 Jan				
1. GOT PREGNANT WHILE USING	2 Feb					2 Feb				
2. WANTED TO GET PREGNANT	3 Mar					3 Mar				
3. HUSBAND OBJECTED	4 Apr					4 Apr				
4. SIDE EFFECTS	5 May					5 May				
5. HEALTH CONCERNS	6 Jun					6 Jun				
6. STOPPED TO "REST THE BODY"	7 Jul					7 Jul				
7. PHYSICIAN DECISION	8 Aug					8 Aug				
8. SUPPLY/AVAILABILITY	9 Sep					9 Sep				
9. DIFFICULT/INCONVENIENT TO USE	10 Oct					10 Oct				
10. MARRIAGE/RELATIONSHIP ENDED	11 Nov					11 Nov				
11. WANTED TO TRY OTHER METHOD	12 Dec					12 Dec				

- WANTED TO TRY OTHER METHOD
 SPORADIC SEXUAL ACTIVITY
- 12. SPORADIC SEXUAL ACTIVIT 13. SHE NEGLECTED TO USE
- 20. OTHER

COLUMN 4 (MARITAL STATUS)

- 0. NOT MARRIED/NOT IN UNION
- 1. MARRIED/IN UNION

IF SHE DID NOT USE A METHOD IN JANUARY 1997 (COLUMN 2_JAN 1997=0) GO TO Q527

526. You said that in January of 1997 you were using ____ (WRITE CODE # FOR THE MET. USED IN COLUMN 2_JAN 1997).

When did you start using that method?

MONTH _____ YEAR ____ ___

88. DK/DO NOT REMEMBER

527. LAST CONTRACEPTIVE METHOD USED (COPY THE METHOD FROM THE CONTRACEPTIVE CALENDAR):

1. THE PILL 9. DEPO-PROVERA 2. IUD **10. OTHER MODERN METHOD** 3. CONDOM 11. CALENDAR \rightarrow GO TO Q536 4. CONDOM +SPERMICIDES 12. WITHDRAWAL \rightarrow GO TO Q536 5. CONDOM +WITHDRAWAL/CALENDAR 13. WITHDRAWAL+CALENDAR \rightarrow GO TO Q536 6. FOAM/JELLY/CREAMS 20. OTHER TRADITIONAL MET. \rightarrow GO TO Q536 7. FEMALE STERILIZATION 88. DO NOT REMEMBER \rightarrow GO TO Q536 8. EMERGENCY HORMONAL CONTRACEP-TION 528. The next following questions concern the last contraceptive method you have used. Where did you get that method? 1. HEALTH POST 8. PHARMACY 2. HEALTH CENTER 9. OPEN MARKET 3. POLICLINIC 10. STORE/ KIOSK 4. FAMILY PLANNING CLINIC 11. PARTNER/HUSBAND 5. GOV HOSPITAL-MATERNITY WARD 12. FRIEND 6. PRIVATE CLINIC OR OFFICE 13. RELATIVE 7. NGO 20. OTHER (SPECIFY): 88. DON'T KNOW 529 Do (Did) you pay for this method? 1. YES 2. NO \rightarrow GO TO Q531 3. PARTNER GETS THE METHOD \rightarrow GO TO Q531 530. On average, how much did you pay for contraception, per month? HUNDRED LEKE 85. 85 HUNDRED LEKE OR MORE 98. NOT SURE/DON'T KNOW 531. At the time you started using the last contraceptive method, who advised you about how to use that method? 1. OB/GYN 6. OTHER RELATIVE \rightarrow GO TO Q536 2. GENERAL PRACTITIONER 7. FRIEND- \rightarrow GO TO O536 3. NURSE/MIDWIFE/FELDCHER 8. PARTNER \rightarrow GO TO Q536 4. PHARMACIST \rightarrow GO TO Q536 9. NOBODY \rightarrow GO TO Q536 5. MOTHER \rightarrow GO TO Q536 20. OTHER \rightarrow GO TO Q536 532. When you received the information concerning use of the method, did the health provider tell you about other contraceptive methods?

1. YES

2. NO \rightarrow GO TO Q534

533. Did the health provider explain how effective your method is compared to other contraceptive methods?

1. YES 2. NO

534. Did the health provider explain the possible side effects of your method?

1. YES

2. NO

536. OBSERVE THE CALENDAR AND RECORD IF RESPONDENT HAS USED PILLS OR IUD AT ANY TIME DURING THE PAST FIVE YEARS:

ONLY PILLS
 PILL AND IUD
 ONLY IUD → GO TO Q543
 NEITHER PILL NOR IUD (OTHER MODERN OR TRAD. METHODS) → GO TO BOX 5-V, PG. 33

537. OBSERVE THE CALENDAR AND VERIFY IN WHAT MONTH AND YEAR RESPONDENT STARTED TO TAKE PILLS MOST RECENTLY (PAST OR CURRENT USERS). You said you most recent started taking pills in:

____MONTH _____YEAR

88. DO NOT REMEMBER

538. What brand of pills did you use most recently? (SHOW CARD B; ASK TO SEE PACKAGE, IF SHE IS CURRENTLY USING PILLS)

 BISECURIN CILEST 	 MARVELON MERCILON 	 SIGORAL MICROLUT
3. CYCLO-PROGYNOVA	10. MICROGYNON	20. OTHER
4. DIANE-35	11. NEOGYNON	88. DO NOT KNOW
5. FEMODEN	12. OVIDON	
6. FERTILAN	13. POSTINOR	
7. GYNOFEN 35	14. RIGEVIDON	

539. When you started taking pills, how long did your physician tell you that you could take them? (Q539 REFERS TO THE LAST INTERVAL OF USE, INCLUDING CURRENT USE)

MONTHS	00. NEVER TALKED TO A DOCTOR ABOUT IT
	44. THREE OR MORE YEARS (36 MONTHS OR MORE)
	55. AS LONG AS RESPONDENT WANTED/INDEFINITELY
	66. DID NOT SAY HOW LONG
	77. OTHER (SPECIFY)
	88. DON'T REMEMBER

540. At any time during the last usage of pills have you had any health problems or side effects that you think are related to using pills?

1. YES

2. NO \rightarrow GO TO BOX IV

541. What kind of problem or side effect have you had? (CIRCLE ALL MENTIONED, DO NOT READ LIST)

MENTIONED	NOT MENTIONED
A. HEADACHES OR DIZZINES 1	2
B. BLURRED VISION, SEEING FLASHING LIGHTS 1	2
C. WEIGHT GAIN 1	2
D. NAUSEA 1	2
E. BREAST TENDERNESS 1	2
F. BLEEDING/SPOTTING BETWEEN MENSTRUAL PERIODS 1	2
G. MOOD CHANGES (LESS INTEREST IN SEX, DEPRESSION) 1	2
H. OTHER (SPECIFY) 1	2

- 542. Was this problem serious enough that you went to a doctor or clinic about it?
 - 1. YES
 - 2. NO

BOX IV

IF RESPONDENT HAS USED ONLY PILLS (Q536=1) THEN GO TO BOX 5-V PAGE 33; ELSE CONTINUE

88. DO NOT REMEMBER

543. OBSERVE THE CALENDAR AND VERIFY IN WHAT MONTH AND YEAR RESPONDENT STARTED TO USE THE LAST (OR CURRENT) IUD. You said you had an IUD inserted in....

____YEAR

MONTH

- 544. Now, I want you to think back at the time when you had inserted your (last) IUD. Where did you have the IUD inserted?
 - 1. HEALTH CENTER
 - 2. POLICLINIC
 - 3. FAMILY PLANNING CLINIC
 - 4. GOV HOSPITAL-MATERNITY WARD
 - 5. PRIVATE CLINIC OR OFFICE
 - 6. MARIE STOPES
- 545. After the IUD was inserted, when did the physician tell you to come back for a routine check-up?

- 546. When the IUD was inserted, did the physician tell you how to check that the IUD is in place?
 - 1. YES
 - 2. NO
 - 8. DON'T REMEMBER
- 547. Did the physician tell you how long the IUD could be left in place?
 - 1. YES
 - 2. NO
 - 8. DON'T REMEMBER
- 548. Thinking back to the first year after you had this IUD inserted, did you have any health problems or side effects that you think are related to your IUD?
 - 1. YES
 - 2. NO \rightarrow GO TO BOX 5-V ON NEXT PAGE

549. What kind of problem or side effect did you have? ? (CIRCLE ALL MENTIONED, DO NOT READ LIST)

	MENTIONED	NOT MENTIONED
A. ABDOMINAL CRAMPING	1	2
B. HEAVY BLEEDING DURING MENSTRUAL PERIO	DDS 1	2
C. SPOTTING/BLEEDING BETWEEN PERIODS	1	2
D. INFECTION/DISCHARGE/PID	1	2
E. PARTNER'S COMPLAINS ABOUT THE STRINGS	1	2
F. EXPULSION	1	2
G. OTHER (SPECIFY)	1	2

- 550. Did you see a doctor for this(ese) problem(s)?
 - 1. YES
 - 2. NO

BOX 5-V

IF ANY CONTRACEPTIVE METHOD WAS USED IN THE LAST MONTH (LAST CELL IN COLUMN 2 >"0") THEN GO TO Q553; ELSE CONTINUE

- 551. Do you think you are physically able to get pregnant at the present time?
 - 1. YES \rightarrow GO TO Q553
 - 2. NO
 - 3. NOT SURE
 - 4. CURRENTLY PREGNANT \rightarrow GO TO Q553
- 552. What is the main reason why you think you cannot get pregnant?
 - 1. RESPONDENT DOES NOT HAVE A PARTNER/ IS NOT SEXUALLY ACTIVE
 - 2. CURRENTLY BREAST-FEEDING /POSTPARTUM
 - 3. PELVIC IINFLAMMATORY DISEASE (PID)
 - 4. ENDOCRINE DYSFUNCTION
 - 5. HYSTERECTOMY (SURGICAL REMOVAL OF UTERUS) \rightarrow GO TO Q558 PAGE 34
 - 6. PREMENOPAUSE/ MENOPAUSE \rightarrow GO TO Q558 PAGE 34
 - 7. OVARIAN CYSTS/ OVARIAN DYSFUNCTION \rightarrow GO TO Q557 PG. 34
 - 8. RESPONDENT HAD BOTH TUBES REMOVED OR OBSTRUCTED \rightarrow GO TO Q557 PG. 34
 - 9. HAS TRIED TO GET PREGNANT IN THE PAST 2 YEARS AND DID NOT SUCCEED \rightarrow GO TO Q557 PG. 34
 - 10. PARTNER HAD A MEDICAL OPERATION AND CANNOT HAVE CHILDREN \rightarrow GO TO Q557 PG. 34
 - 11. PARTNER IS INFERTILE \rightarrow GO TO Q557 PG. 34
 - 12. CURRENTLY USES A METHOD (GO BACK TO Q504 AND CORRECT IT)
 - 20. OTHER (SPECIFY)
 - 88. DO NOT KNOW
 - 99. REFUSE TO ANSWER
- 553. Looking to the future, do you yourself intend to have (a/another) baby at some time (IF CURRENTLY PREGNANT ADD "...after this pregnancy"?)

WANTS A BABY
 DOES NOT WANT A BABY → GO TO Q555
 RESPONDENT WANTS A BABY BUT PARTNER DISAGREES
 RESPONDENT DOES NOT WANT A BABY BUT PARTNER WANTS → GO TO Q555
 DK → GO TO Q555

554. When do you, yourself, actually want to get pregnant (again)...(READ 1-4)

Right away, (DO NOT READ IF THE WOMAN IS ALREADY PREGNANT)
 Within the next 12 months,
 Within 1-2 years,
 or after 2 years?
 AFTER SHE MARRIES
 WHEN GOD WANTS

8. DK

- 555. IF Q553 =1,3, OR 8 BEGIN WITH: "After having all the children you want,...") Do you think you would be interested in having an operation to prevent you from having any more children?
 - 1. YES \rightarrow GO TO Q558
 - 2. NO
 - 3. ALREADY STERILIZED \rightarrow GO TO Q558
 - 8. NOT SURE
- 556. What is the most important reason you wouldn't be interested in such a procedure?
 - 1. HEALTH RISKS/FEAR OF SIDE EFFECTS
 - 2. FEAR OF OPERATION
 - 3. DOESN'T KNOW ENOUGH ABOUT /NEVER HEARD OF STERILIZATION
 - 4. MIGHT WANT ANOTHER CHILD
 - 5. COST
 - 6. DOES NOT HAVE A PARTNER/NOT SEXUALLY ACTIVE
 - 7. AGE TOO YOUNG OR TOO OLD (APPROACHING MENOPAUSE)
 - 8. HAVEN'T THOUGHT ABOUT IT
 - 9. NOT CULTURALLY ACCEPTABLE
 - 10. RELIGIOUS REASONS
 - 11. PREFERS (OR USES) OTHER CONTRACEPTIVE METHODS
 - 12. CANNOT GET PREGNANT (INFERTILITY, MEDICAL REASONS)
 - 20. OTHER_
- 88. DON'T KNOW

GO TO Q558

- 557. Looking to the future, do you yourself intend to seek any medical help to have a(nother) baby ?
 - 1. YES
 - 2. NO
 - 3. RESPONDENT WANTS A BABY BUT PARTNER DISAGREES
 - 4. RESPONDENT DOES NOT WANT A BABY BUT PARTNER WANTS
 - 8. DK
- 558. The next questions are about any infertility services you may have ever received. This includes any medical help to become pregnant that you or your husband(s)/partner(s) may have received. Have you or your husband(s)/partner(s) ever been to a doctor or other medical care provider to talk about ways to help you become pregnant?

YES
 NO → GO TO Q571
 DK/NR → GO TO Q571

565. Thinking back to your or your partner's first visit when you sought medical help for becoming pregnant, in what month and year was that visit (IF PARTNER 1ST VISIT PRECEEDED HERS, RECORD THAT DATE)

____ MONTH

YEAR

88. DK/NOT REMEMBER

566.	When you first went for medical help in (MONTH/YEAR), how many months or years had you (and your husband/part- ner) been trying to become pregnant?									
	1	MONTHS	OR	2	YEARS		888. DK/DR			
567.							ome pregnant were yo ND CODE ALL THA		u	
	B. Bloc	ked tubes?						1	<u>NO</u> 2 2	
	D Seme	en or sperm pro	blems (l	ow count	, poor motility, v	aricocele) '	s fixes to other places	1	2 2 2	
568.	1. YES	5		re you (a	nd your (husband	l/partner)) j	pursuing medical help	to become pregna	nt?	
569.		\rightarrow GO TO Q5' the past 12 more		w many v	isits have you or	your husba	and/partner made to a	doctor to help you	to get preg-	
		VISITS	8	8. DK/DR	Ł					
570.	In what	t month and yea	r was y	our (most	recent/last) visit	for help to	become pregnant?			
	A	MONTH	[В	l	_YEAR	88. DK/NOT REME	MBER		
571.	Have you ever been treated for an infection in your fallopian tubes, womb, or ovaries, also called a pelvic infection, pelvic inflammatory disease, or P.I.D.? (IF DON'T KNOW, PROBE: This is a female infection that sometimes causes abdominal pain or lower stomach cramps.) NOTE: INFECTIONS OF THE VAGINA ALONE, ENDOMETRIOSIS, PELVIC TUMORS, AND CYSTS DO NOT COUNT AS PELVIC INFECTIONS									
		\rightarrow GO TO MC NR \rightarrow GO TO								
572.	Were ye	ou having any s	ympton	ns, such a	s pain, discharge	, or bleedin	g, that caused you to	go for treatment?		
	1. YES 2. NO	5								
573.	Please t that?	try to remember	when y	vou first re	eceived treatmen	t for a pelv	ic infection or P.I.D.	In what month and	year was	
		MONTH		-		YEAR	88. DK/NC	T REMEMBER		
574.	In what	In what month and year did you last receive treatment for a pelvic infection or P.I.D.?								
		MONTH		-		YEAR	88. DK/NC	T REMEMBER		
575.	Altoget	Altogether, how many different times have you been hospitalized one night or longer for a pelvic infection?								
	2. ONO 3. 2-3 4. 4 TI		E	ULE VI						
576.				you spend	d in the hospital	for a pelvic	infection or P.I.D.?		NIGHTS	

VI. WOMEN'S HEALTH

Now I would like to ask you some questions about your health.

601. In the past 12 months, have you visited any health facility for care for yourself, including obtaining preventive services, such as family planning counselling or health check-ups?

1. YES

2. NO \rightarrow GO TO Q604

8. DK/DO NOT REMEMBER \rightarrow GO TO Q604

602. Where did you receive these services (CIRCLE ALL MENTIONED AND PROBE "Any Other Place")?

MENTIONED	NOT MENTIONED
A. GOVT. HEALTH CLINIC OR HOSPITAL 1	2
B. PRIVATE HEALTH CLINIC OR HOSPITAL 1	2
C. INTERNATIONAL RELIEF ORGANIZATION (e.g. Red Cross) 1	2
D. LOCAL NGO (e.g. Albanian Family Planning) 1	2
E MARIE STOPES 1	2
F. OTHER 1	2

603. During your visit in the past 12 months at the health facility, did a doctor or medical provider talk to you about family planning methods?

- 1. YES
- 2. NO
- 8. DK
- 604. Many different factors can prevent women from getting medical advice or treatment for themselves. When you want to get medical advice or treatment, is each of the following a big problem or not (**READ A G**)?

	BIG PI	ROBLEM	NOT A BIG PROBLEM
1	A. Knowing where to go	1	2
]	B. Getting permission to go	1	2
(C. Getting money needed for treatment	1	2
]	D. The distance to the health facility	1	2
]	E. Having to take transport	1	2
]	F. Not wanting to go alone	1	2
(G. Concern that there may not be a female health provider	1	2

- 605. Have you ever had a routine gynecologic exam (PHYSICAL EXAMINATION OF EXTERNAL AND INTERNAL GENITAL AREA FOR DIAGNOSTIC OTHER THAN PREGNANCY)?
 - 1. YES \rightarrow **GO TO Q607** 2. NO 8. NR
- 606. What is the most important reason that you have never had a routine gynecologic exam?
 - 1. DOES NOT NEED TO GO TO GYNECOLOGIC EXAM
 - 2. SHE IS HEALTHY AND HAS NOT GYNECOLOGIC PROBLEMS
 - 3. THERE IS NOT TIME TO GO FOR EXAM
 - 4. SHE FORGETS ABOUT IT
 - 5. SHE DOES NOT LIKE GYNECOLOGIC EXAM
 - 6. IT IS DIFFICULT TO GET APPOINTMENT
 - 7. DOES NOT LIKE PLACE/FACILITY
 - 8. DOES NOT LIKE THE STAFF
 - 9. WAITING TIME IS TOO LONG
 - 10. DOCTOR DID NOT RECOMMEND
 - 11. SHE IS EMBARRASSED TO HAVE GYNECOLOGIC EXAM
 - 12. NEVER THOUGHT ABOUT IT
 - 13. DOES NOT KNOW WHERE TO GO FOR SUCH AN EXAM
 - 14. CANNOT AFFORD THE COST
 - 15. NEVER HAD SEXUAL INTERCOURSE (VIRGIN)
 - 20. OTHER _
 - 88. DK/NOT RESPONSE

GO TO Q608

Female Questionnaire

- 607. When was your last routine gynecologic exam (not pregnancy related)? (READ 1-4)
 - During the past 12 months
 1-2 years ago (12-23 MTH)
 2-3 years ago (24-35 MTH)
 3 or more years ago
 DK/DR
- 608. Have you ever had a cervical smear (a test that takes a sample of cells from the cervix, or opening to the uterus to detect cancer), also called Papanicolau test?
 - 1. YES \rightarrow GO TO Q610

2. NO

- 8. DK
- 9. REF
- 609. What is the main reason you have never had a Pap smear?
 - 1. NEVER HEARD OF IT
 - 2. DOCTOR HAS NOT RECOMMENDED IT
 - 3. SHE IS HEALTHY AND HAS NO GYNECOLOGIC PROBLEMS
 - 4. SHE DOES NOT FEEL TEST IS NECESSARY
 - 5. DOES NOT HAVE TIME TO GO FOR A TEST/ SHE FORGETS ABOUT IT
 - 6. NEVER THOUGHT OF IT
 - 7. SHE IS AFRAID OF THE RESULTS
 - 8. SHE IS AFRAID IT COULD BE PAINFUL
 - 9. TOO EMBARRASSED TO GET THE TEST OR A PELVIC EXAM.
 - 10. SHE HAD NO PARTNER/ NOT SEXUALLY ACTIVE
 - 11. NEVER HAD SEXUAL INTERCOURSE
 - 20. OTHER (SPECIFY):___
 - 88. DON'T KNOW
 - 99. REFUSE TO ANSWER

GO TO Q611

- 10. When did you have your last Pap smear? Was it...(READ 1-4)
 - 1. within the last year, (O TO 11 MONTHS AGO)
 - 2. 1 to 2 years ago, (12 TO 23 MONTHS AGO)
 - 3. 2-3 years ago, (24 to 35 MONTHS AGO)
 - 4. more than 3 years ago? (36+MONTHS AGO)
 - 8. DON'T KNOW
- 611. Have you heard about breast self-examinations?
 - 1. YES
 - 2. NO \rightarrow GO TO Q614
- 612. Do you ever do breast self-examinations?
 - 1. YES 2. NO \rightarrow **GO TO Q614**
- 613. How often do you do it, on average?

ONCE A MONTH/AFTER EACH MENSTRUATION
 EVERY 2-5 MONTHS
 EVERY 6-11 MONTHS
 ONCE PER YEAR OR LESS

614. Have you ever tried cigarette smoking, even one or two puffs?

1. YES 2. NO → **GO TO 621**

615. How old were you when you smoked a cigarette for the first time?

___YEARS 88. DO NOT REMEMBER 99. REFUSE TO ANSWER

616. Have you smoked at least 100 cigarettes in your entire life? (PROBE: 100 cigarettes is about 5 packs)

1. YES 2. NO \rightarrow GO TO 621 8. DK \rightarrow GO TO 621 9. REF \rightarrow GO TO 621

617. How old were you when you first started smoking fairly regularly?

YEARS	00. NEVER SMOKED REGULARLY
	88. DO NOT REMEMBER
	99. REFUSE TO ANSWER

618. During the last 30 days, did you smoke cigarettes: (READ 1-4)

Every Day
 Almost Every Day
 Some Days
 Not at All in the last 30 days → GO TO Q620
 REF → GO TO Q620

619. During the last 30 days, on the days you smoked, how many cigarettes did you smoke per day?

_____YEAR

- 1. 1 CIGARETTE PER DAY
- 2. 2-5 CIGARETTES PER DAY
- 3. 6-10 CIGARETTES PER DAY
- 4. 11-19 CIGARETTES PER DAY
- 5. 20 OR MORE CIGARETTES PER DAY

GO TO Q621

620. In what month and year did you last smoke cigarettes at all? (PROBE FOR SEASON IF MONTH IS UNKNOWN)

____ MONTH ____

88. DO NOT REMEMBER 99. REF

- 621. Now, I will ask you about some medical conditions that you may have had. Has a doctor or other medical care provider ever told you that you had Diabetes or "high sugar"?
 - 1. YES
 - 2. NO \rightarrow **GO TO Q623**
 - 8. NOT SURE \rightarrow **GO TO Q623**
 - 9. REFUSAL \rightarrow GO TO Q623

622. Were you ever told you had diabetes when you were not pregnant?

- 1. YES
- 2. NO
- 3. NEVER BEEN PREGNANT

623. Has a doctor or other medical care provider ever told you that you had Anemia, or "thin blood"?

- 1. YES
- 2. NO \rightarrow GO TO Q625
- 8. NOT SURE \rightarrow GO TO Q625
- 9. REFUSAL \rightarrow GO TO Q625
- 624. Were you ever told you had anemia or "thin blood" when you were not pregnant?
 - 1. YES
 - 2. NO
 - 3. NEVER BEEN PREGNANT
- 625. Has a doctor or other medical care provider ever told you that you had Hypertension or High Blood Pressure?
 1. YES
 2. NO. CO TO O(27)
 - 2. NO \rightarrow GO TO Q627
 - 8. NOT SURE \rightarrow **GO TO Q627**
 - 9. REFUSAL \rightarrow GO TO Q627
- 626. Were you ever told you had Hypertension or High Blood Pressure when you were not pregnant?
 - 1. YES
 - 2. NO
 - 3. NEVER BEEN PREGNANT
- 627. In the past 12 months have you had any of the following symptoms?

	<u>YES</u>	<u>NO</u>	NOT SURE
A. Vaginal discharge with a bad smell	1	2	8
B. Itching or burning in the genital area	1	2	8
C. Burning pain on urination	1	2	8
D. Pain during sexual intercourse	1	2	8
E. Sore, ulcer or warts in genital area	1	2	8
F. Swelling in genital area	1	2	8

IF ALL Q627_A—Q627_F >1 (NO SYMPTOMS IN THE PAST 12 MONTHS) GO TO Q631; ELSE CONTINUE

628. Did you have any treatment for this(ese) condition(s)?

- 1. YES
- 2. NO \rightarrow GO TO Q630
- 8. NOT SURE \rightarrow GO TO Q630

629. Who treated you?

- 1. OB/GYN
- 2. FAMILY DOCTOR/GP
- 3. DERMATOLOG
- 4. INFECTIOUS DISEASES DOCTOR
- 5. NURSE/MIDWIFE
- 6. PHARMACIST

- 7. PARTNER
- 8. FRIEND/RELATIVE
- 9. SELF-TREATMENT
- 20. OTHER:
- 99. REFUSE TO ANSWER

GO TO Q631

- 630. What was the main reason you did not seek treatment?
 - 1. SERVICES TOO FAR AWAY / INACCESSIBLE
 - 2. DON'T KNOW WHERE TO GO FOR SERVICES
 - 3. CANNOT AFFORD SERVICES OR TREATMENT
 - 4. AFRAID OF KNOWING THE RESULTS
 - 5. IT IS EMBARRASSING
 - 6. DID NOT THINK WAS AN STD
 - 7. SYMPTOM(S) DISAPPEARED
 - 8. OTHER (SPECIFY)_____
 - 9. REFUSE
- 631. In the past 3 months, have you had a drink containing alcohol, that is a beer, a glass of wine, a cocktail, a shot of liqueur, vodka, or whiskey?
 - 1. YES
 - 2. NO \rightarrow **GO TO MODULE VII**
 - 8. NOT SURE/DO NOT REMEMBER \rightarrow **GO TO MODULE VII**
 - 9. REFUSE \rightarrow **GO TO MODULE VII**
- 632. In the past 3 months, on the days that you drank alcohol, how many drinks did you usually have?
 - _____ # OF DRINKS

00. NO DRINKS/ONLY FEW SIPS → **GO TO MODULE VII** 88. NOT SURE/DO NOT REMEMBER → **GO TO MODULE VII** 99. REFUSE → **GO TO MODULE VII**

- 633. In the past 3 months, how often did you drink that amount (PROBE: per day, week, or month)?
 - 1. EVERYDAY
 - 2. ALMOST EVERY DAY
 - 3. 1-2 TIMES A WEEK
 - 4. 2-3 TIMES A MONTH
 - 5. ONCE A MONTH
 - 6. 1-2 TIMES IN THREE MONTHS
- 634. In the past 3 months, have there been days when you had more than usual (# FROM Q632) drinks?
 - 1. YES

2. NO \rightarrow GO TO MODULE VII 8. NOT SURE/DO NOT REMEMBER \rightarrow GO TO MODULE VII 9. REFUSE \rightarrow GO TO MODULE VII

635. In the past 3 months, how many drinks did you have on the days that you drank more than usual (# FROM Q632)? (CHECK IF # FROM Q635># FROM Q632)

```
_____ # OF DRINKS
```

88. NOT SURE/DO NOT REMEMBER \rightarrow **GO TO MODULE VII** 99. REFUSE \rightarrow **GO TO MODULE VII**

636. How often did you drink that amount?

1. EVERYDAY

- 2. ALMOST EVERY DAY
- 3. 1-2 TIMES A WEEK
- 4. 2-3 TIMES A MONTH
- 5. ONCE A MONTH
- 6. 1-2 TIMES IN THREE MONTHS

VII REPRODUCTIVE HEALTH KNOWLEDGE/ATTITUDES

700. What do you think is the ideal number of children for a young family in Albania?

	0 CHILDREN		3-4 CHILDREN
	1 CHILD 1-2 CHILDREN		4 CHILDREN 5 OR MORE
	2 CHILDREN		AS MANY AS GOD GIVES
4.	2-3 CHILDREN	77.	AS MANY AS POSSIBLE
5.	3 CHILDREN	88.	DON'T KNOW

701. During a woman's menstrual cycle, are there certain days when she is more likely to become pregnant if she has sexual relations?

1. YES

2. NO \rightarrow GO TO 702

8. DO NOT KNOW \rightarrow GO TO 702

- 701A. When is it most likely for a woman to become pregnant, just before her period begins, during her period, right after her period has ended, or halfway between two periods?
 - 1 Just before her period starts
 - 2 During her period
 - 3 Right after period ends
 - 4 Halfway between her periods
 - 8 DON'T KNOW
- 702. Do you think that breastfeeding increases, decreases or has no effect on a woman's chance to get pregnant?
 - 1. INCREASES THE CHANCE
 - 2. DECREASES THE CHANCE
 - 3. HAS NO EFFECT
 - 8. DO NOT KNOW
- 703. Do you think that a woman always has the right to decide about her pregnancy, including whether or not to have an abortion?

1. YES \rightarrow GO TO Q705

- 2. NO
- 3. DO NOT KNOW
- 704. Under which of the following conditions is it all right for a woman to have an abortion (**READ A-F**)?

	YES	<u>NO</u>	DEPENDS	<u>DK</u>
A. Her life is endangered by the pregnancy	1	2	3	8
B. The fetus has a physical deformity	1	2	3	8
C. The pregnancy has resulted from rape	1	2	3	8
D. Her health is endangered by the pregnancy	1	2	3	8
E. She is unmarried	1	2	3	8
F. The couple cannot afford to have a(nother) child	1	2	3	8

- 705. If a woman had an unwanted pregnancy what should she do? (**READ 1-3**):
 - 1. Have the baby and keep it
 - 2. Have the baby and give it up for adoption
 - 3. Have an abortion
 - 8. DON'T KNOW

706. I would like to know if you are in agreement with the following statements (**READ A-I**):

	AGREE	DISAGREE	<u>DK</u>
A. A woman can become pregnant the first time she has sexual intercourse	1	2	8
B. All people should get married	1	2	8
C. A woman should be a virgin when she marries	1	2	8
D. The main job for a woman is to take care of the home and cook for her family	1	2	8
E. A married woman needs her husband's permission to work outside the home	1	2	8
F. If a woman works, she should give her money to her husband	1	2	8
G. If a woman works, her husband should help her with the household chores	1	2	8
H. The men in the family should have the final say in all family matters	1	2	8
I. Child care is a woman job	1	2	8

707. Who do you think should decide how many children a couple should have (**READ 1-5**)?

- 1. The woman,
- 2. The man,
- 3. Both
- 4. Mother in law, or
- 5. God?
- 8 DON'T KNOW
- 708. How would you rank each of the following birth control methods (SHOW CARD C) with regard to their risk of developing health problems; please tell me if the risk is low, medium, or high:

	Low <u>Risk</u>	Medium <u>Risk</u>	High <u>Risk</u>	<u>DK</u>
A. Pill	1	2	3	8
B. IUD	1	2	3	8
C. Condom	1	2	3	8
D. Tubal Ligation	1	2	3	8
E. Injectables	1	2	3	8
F. Abortion on Request	1	2	3	8

BOX 7-I

IF Q400_A=2 ON PAGE 20 (NEVER HEARD OF PILLS), GO TO BOX 7-II BELOW

710. Please tell me if you agree or disagree with the following statements about birth control pills (READ A-J):

	AGREE	DISAGREE	<u>DK</u>
A. Pills are easy to use	1	2	8
B. Pills are easy to get	1	2	8
C. Pills are too expensive	1	2	8
D. It is stressful to remember to take the pill every day	1	2	8
E. Pills protect against some gynecologic cancers	1	2	8
F. Pills may make you gain weight	1	2	8
G. Pills make women's periods more regular	1	2	8
H. Pills decrease blood loss during menstruation	1	2	8
I. Pills decrease menstrual cramps and pain	1	2	8
J. Pills are bad for blood circulation	1	2	8

BOX 7-II

IF Q400_B=2 ON PAGE 20 (NEVER HEARD ABOUT IUD), GO TO Q712

711. Please tell me if you agree or disagree with the following statements about IUDs (READ A-G:

	AGREE	DISAGREE	<u>DK</u>
A. IUD is easy to use	1	2	8
B. IUD increases the risk of pelvic inflammatory disease	1	2	8
C. IUD is a relatively inexpensive contraceptive method	1	2	8
D. IUD may cause spotting between periods	1	2	8
E. IUD may increase the blood loss during menses	1	2	8
F. IUD increases menstrual pains	1	2	8
G. IUD decreases the risk of ectopic pregnancy	1	2	8

712. Do you want to have more information about contraceptive methods?

YES
 NO → GO TO Q714
 DON'T KNOW → GO TO Q714

713. Who do you think would be the best source of information about contraceptive methods?

- MOTHER
 OTHER RELATIVE
 BOYFRIEND
 HUSBAND, PARTNER
 SOMEBODY WHO USES CONTRACEPTION
 CO-WORKER
 FRIEND, COLLEAGUE, PEER
 GYNECOLOGIST
 GENERAL PRACTITIONER
- 10. NURSE, MIDWIFE 11. TEACHER 12. PHARMACIST 13. BOOKS 14 NEWSPAPERS, MAGAZINES, BROCHURES 15. RADIO \rightarrow GO TO Q715 16. TV- \rightarrow GO TO Q715 20. OTHER (SPECIFY): ______ 88. DON'T REMEMBER

714. Do you think that information about contraception should be broadcast on radio or television?

- 1. YES 2. NO
- 8. DO NOT KNOW
- 715. Some people use condoms to keep from getting sexual transmitted diseases. How effective do you think a properly used condom is for this purpose? (**READ 1-3**)
 - 1. Very Effective
 - 2. Somewhat effective
 - 3. Not effective
 - 8. DON'T KNOW

BOX 7-IV

IF RESPONDENT IS 15-24 YEARS OF AGE CONTINUE; IF SHE IS 25-44 YEARS GO TO SECTION VIII

- 718. Have you ever talked to a partner about him using a condom?
 - YES
 NO
 NEVER HAD A SEXUAL PARTNER → GO TO Q721
 DON'T REMEMBER
 REFUSE
- 719. Have you ever asked a partner to use a condom?

1. YES 2. NO \rightarrow **GO TO Q721** 8. DON'T REMEMBER \rightarrow **GO TO Q721** 9. REFUSE \rightarrow **GO TO Q721**

720. Has any of the following ever happened because you asked a partner to wear a condom....... (READ A-F) (ANY OF THESE INCIDENTS COULD HAVE HAPPENED MORE THAN ONCE, WITH THE SAME PARTNER OR DIFFERENT PARTNERS)

	YES	<u>NO</u>	<u>DK</u>	<u>REF</u>
A. Did a partner refuse to wear a condom?	1	2	8	9
B. Did a partner refuse to have sexual intercourse with you?	1	2	8	9
C. Did a partner threaten to break up with you?	1	2	8	9
D. Did a partner yell at you or threaten to hurt you?	1	2	8	9
E. Did a partner make you have sex anyway without a condom?	1	2	8	9
F. Did a partner physically hurt you?	1	2	8	9

721. If your partner/husband would want to use a condom when having sex with you, would you feel: (READ A-E)

	AGREE	DISAGREE	<u>DK</u>	<u>REF</u>
A. Insulted or angry?	1	2	8	9
B. Safe from getting pregnant?	1	2	8	9
C. Like you had done something wrong?	1	2	8	9
D. Safe from getting STD or HIV/AIDS?	1	2	8	9
E. Suspicious that he may sleep around?	1	2	8	9

722. Please indicate whether you agree or disagree with the following statements about condoms (READ A-H):

	AGREE	DISAGREE	<u>DK</u>	<u>REF</u>
A. Using condoms with a new partner is a smart idea	1	2	8	9
B. Using condoms is not necessary if you know your partner	1	2	8	9
C. Women should ask their partners to use condoms	1	2	8	9
D. It is easy to discuss using a condom with a prospective partner	1	2	8	9
E. Condoms diminish sexual enjoyment	1	2	8	9
F. Same condoms can be used more than once	1	2	8	9
G. People who use condoms sleep around a lot	1	2	8	9
H. It is embarrassing to ask for condoms in FP clinics or pharmacies	1	2	8	9

VIII. SOCIOECONOMIC CHARACTERISTICS

800. Please tell me whether this household or any member of it has the following items: (**READ A-I**):

	YES	NO
A. Flush Toilet	1	2
B. Heating System	1	2
C. Refrigerator	1	2
D. TV	1	2
E. Working Automobile	1	2
F. VCR	1	2
G. Household phone	1	2
H. Cellular phone	1	2
I. Vacation home (villa)	1	2
J. Air Conditioner	1	2
K. Gas or Electric Stove	1	2
L. Computer	1	2
M. Satellite Antenna	1	2
Does your family have access to a garden whe 1. YES 2. NO	ere you grov	w your own vegetables?
What is the main source of drinking water for	members o	f your household?
1. PIPED WATER (PIPED INTO RES	IDENCE O	$(R YARD) \rightarrow GO TO Q804$
2. PIPED WATER (PUBLIC TAP)		
3. WELL WATER (RESIDENCE OR Y	$(ARD) \rightarrow 0$	GO TO Q804
4. PUBLIC WELL	,	-

- 5. SURFACE WATER (SPRING, RIVER, POND, LAKE)
- 6. RAIN WATER \rightarrow **GO TO Q804**
- 7. BOTTLED WATER \rightarrow **GO TO Q804**

803.	How long does it	t take to go to the	water source, ge	et water and come back	, in minutes?		MINUTES
------	------------------	---------------------	------------------	------------------------	---------------	--	---------

- 804. Which of the following describes your living arrangements. Do you live: (READ 1-5)
 - 1. In your privately owned flat or house
 - 2. In rented space (room, flat or house)
 - 3. With your immediate family (NO RENT)
 - 4. With other relatives (NO RENT)
 - 5. With friends (NO RENT)
 - 7. OTHER (SPECIFY):

801.

802.

805. How many rooms are occupied by you and your family (not including bathrooms and kitchen): _____ ROOMS

 806.
 How many hours per day do you have electricity?
 HOURS
 88 DK/NOT SURE

807. What is your ethnic background?

1. ALBANIAN	5. ALBANIAN FROM KOSOVO	
2. GREEK	6. MIXED ETHNICITY (SPECIFY)	
3. GYPSY (ROMA)	7. OTHER (SPECIFY):	
4. MACEDONIAN	9. REFUSED/NOT STATED	
What is your religion?		

1. MUSLIM.

808.

- 2. ORTHODOX
- 3. CATHOLIC
- 4. PROTESTANT (BAPTIST, LUTHERAN, PENTECOSTAL, ETC)
- 7. OTHER (SPECIFY):_____
- 8. NO RELIGION \rightarrow **GO TO Q900**
- 9. UNDECLARED \rightarrow **GO TO Q900**
- 809. About how often do you usually attend religious services? (READ 1-5)
 - 1. At least once a week
 - 2. At least once a month, but less than once a week
 - 3. Less than once a month
 - 4. Only on holidays, or
 - 5. Never

IX-A. AIDS/STDs

The next set of questions are about sexually transmitted infections Including HIV/ AIDS. For each of the following conditions please tell me if:

CONDITION	900. Have you ever heard of it?	901. Have you ever been tested for?	902. Have you ever been told that you have?	903. Did you take any treatment for?	904. Who treated you for? (SEE CODES BELOW)
A. Syphilis	$1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{B}$	1. YES 2. NO \rightarrow B 8. DK \rightarrow B	1. YES 2. NO \rightarrow B 8. DK/DR \rightarrow B	$1. YES$ $2. NO \rightarrow B$ $8. DK/DR \rightarrow B$	
B. Gonorrhea	$1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{C}$	1. YES 2. NO \rightarrow C 8. DK \rightarrow C	1. YES 2. NO \rightarrow C 8. DK/DR \rightarrow C	1. YES 2. NO \rightarrow C 8. DK/DR \rightarrow C	
C. Chlamydia	$1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{D}$	1. YES 2. NO \rightarrow D 8. DK \rightarrow D	1. YES 2. NO \rightarrow D 8. DK/DR \rightarrow D	1. YES 2. NO \rightarrow D 8. DK/DR \rightarrow D	
D. Yeast Infection	$1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{E}$	1. YES 2. NO \rightarrow E 8. DK \rightarrow E	1. YES 2. NO \rightarrow E 8. DK/DR \rightarrow E	1. YES 2. NO \rightarrow E 8. DK/DR \rightarrow E	
E. Genital Herpes	$\begin{array}{c} 1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{F} \end{array}$	1. YES 2. NO \rightarrow F 8. DK \rightarrow F	1. YES 2. NO \rightarrow F 8. DK/DR \rightarrow F	1. YES 2. NO \rightarrow F 8. DK/DR \rightarrow F	
F. Genital Warts	$1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{G}$	1. YES 2. NO \rightarrow G 8. DK \rightarrow G	1. YES 2. NO \rightarrow G 8. DK/DR \rightarrow G	1. YES 2. NO \rightarrow G 8. DK/DR \rightarrow G	
G. Trichomoniasis	$\begin{array}{c} 1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{H} \end{array}$	1. YES 2. NO \rightarrow H 8. DK \rightarrow H	1. YES 2. NO \rightarrow H 8. DK/DR \rightarrow H	$\begin{array}{c} 1. \text{ YES} \\ 2. \text{ NO} \rightarrow \text{H} \\ 8. \text{ DK/DR} \rightarrow \text{H} \end{array}$	
H. HIV/AIDS	1. YES 2. NO → Q905	1. YES \rightarrow Q905 2. NO \rightarrow Q905 8. DK \rightarrow Q905			

CODES FOR Q904:

- 1. OB/GYN
- 2. FAMILY DOCTOR/GP
- 3. DERMATOLOG
- 4. INFECTIOUS DISEASES DOCTOR
- 5. NURSE/MIDWIFE
- 6. PHARMACIST

- 7. PARTNER
- 8. FRIEND/RELATIVE
- 9. SELF-TREATMENT
- 20. OTHER:____
 - 99. REFUSE TO ANSWER

905. If a woman has a sexually transmitted disease, what symptoms might she have? (RECORD ALL MENTIONED, DO NOT READ LIST)

	<u>YES</u>	<u>NO</u>
A. ABDOMINAL PAIN	1	2
B. VAGINAL DISCHARGE	1	2
C. FOUL SMELLING DISCHARGE	1	2
D. BURNING PAIN ON URINATION	1	2
E. REDNESS/INFLAMMATION IN GENITAL AREA	1	2
F. SWELLING IN GENITAL AREA	1	2
G. GENITAL SORES/ULCERS OR WARTS	1	2
H. GENITAL ITCHING	1	2
I. WEIGHT LOSS		1
J. HARD TO GET PREGNANT/HAVE A CHILD	1	2

906. Do you know a place where you could get an HIV/AIDS test?

- 1. YES
- 2. NO
- 907. In general, what has been your most important source of information about STDs including AIDS? (Where or from whom have you learned the most about STDs?)

11. FAMILY DOCTOR, GP
12. NURSE, MIDWIFE, FELDCHER
13. TEACHER
14. PHARMACIST
15. SPECIALITY BOOKS
16. NEWSPAPERS, MAGAZINES, BROCHURES, FLYERS
17. RADIO
18. TV
20. OTHER (SPECIFY):
77. NEVER HEARD OF ANY STDs (Q900_AQ900_H=2)
99. DR/REF.

908. In the past 6 months, have you seen or heard any public announcements or ads on television or radio about: (READ A-D, PROBE FOR BOTH)

				DO NOT
<u>YES, RADIO</u>	<u>YES, TV</u>	<u>YES, BOTH</u>	<u>NO</u>	<u>REMEMBER</u>
1	2	3	4	8
1	2	3	4	8
1	2	3	4	8
1	2	3	4	8
	<u>YES, RADIO</u> 1 1 1 1	YES, RADIO YES, TV 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2	YES, RADIOYES, TVYES, BOTH123123123123123	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$

IF Q900_H =2 (NEVER HEARD OF HIV/AIDS) GO TO Q914; ELSE CONTINUE

909. Do you think that a person can be infected with the HIV virus but have no symptoms of disease?

- 1. YES
- 2. NO
- 8. DK

910. Please tell me whether you think that the AIDS virus can be transmitted in the following ways? (READ A-M)

	<u>YES</u>	<u>NO</u>	<u>DK</u>
A. Through blood transfusion	1	2	8
B. Using public toilets	1	2	8
C. Through kissing	1	2	8
D. Through unprotected sexual intercourse between a man and a woman	1	2	8
E. Through unprotected sexual intercourse between men	1	2	8
F. By shaking hands	1	2	8
G. Using non-sterile syringes or needles	1	2	8
H. Through mosquito bites	1	2	8
I. Sharing plates, forks, or glasses with someone who has HIV/AIDS	1	2	8
J. From a woman who has the AIDS virus to her baby during pregnancy/delivery	1	2	8
K. From a mother to her child through breast milk	1	2	8
L. Getting a manicure, pedicure or haircut	1	2	8
M. Having dental or surgical treatment	1	2	8

911. What can a person do to reduce the risk of getting AIDS? PROBED

	SPONTANEOUS	VEC	PROBED	ЪV
	<u>YES</u>	<u>YES</u>	<u>NO</u>	<u>DK</u>
A. USE CONDOMS	1	3	4	8
B. ABSTAIN FROM SEX	1	3	4	8
C. HAVE ONLY ONE PARTNER/STAY FAITHFUL TO ONE PARTN	IER 1	3	4	8
D. LIMIT NUMBER OF SEXUAL PARTNERS	1	3	4	8
E. AVOID SEX WITH PROSTITUTES	1	3	4	8
F. AVOID SEX WITH BISEXUALS	1	3	4	8
G. DO NOT DONATE BLOOD	1	3	4	8
H. AVOID TRANSFUSIONS	1	3	4	8
I. ASK PARTNER TO GET BLOOD TESTED FOR AIDS	1	3	4	8
J. STERILIZE NEEDLES AND SYRINGES	1	3	4	8
K. DO NOT SHARE RAZORS/BLADES, NEEDLES OR SYRINGES	5 1	3	4	8
L. OTHER (SPECIFY)	1	3	4	8

912. How much of a risk do you think you personally have of getting HIV/AIDS? Would you say you are at (READ 1-4):

- 1. Great risk,
- 2. Moderate Risk,
- 3. Little risk, or
- 4. No risk at all \rightarrow GO TO Q913A
- 8. DON'T KNOW \rightarrow GO TO Q913B

913. Why do you think you have any risk of getting AIDS?

- 1. RECEIVED BLOOD TRANSFUSIONS/BLOOD PRODUCTS
- 2. HAD MANY SEXUAL PARTNERS/ TRADED SEX FOR MONEY
- 3. HAD UNPROTECTED INTERCOURSE WITH CASUAL PARTNER(S)
- 4. USED IV DRUGS/SHARED NEEDLES
- 5. PARTNER HAD/HAS SEX WITH OTHER WOMEN
- 6. SHE MAY GET INFECTED WHILE RECEIVING MEDICAL OR DENTAL TREATMENT
- 7. SHE MAY GET INFECTED GETTING A MANICURE, PEDICURE, OR HAIRCUT
- 8. OTHER (SPECIFY)_____
- 9. DK/REF

GO TO Q913B

- 913A Why do you think you have no risk of getting AIDS?
 - 1. ONLY ONE PARTNER
 - 2. NO SEXUAL RELATIONS
 - 3. USES CONDOMS
 - 4. CONFIDENCE IN PARTNER
 - 5. DOES NOT GET/NEED TRANSFUSIONS
 - 6. DOES NOT SHARE NEEDLES
 - 7. OTHER (SPECIFY)
 - 9. DK/REF

913B How much of a risk do you think you personally have of getting other STD? Would you say you are at (**READ 1-4**):

- 1. Great risk,
- 2. Moderate Risk,
- 3. Little risk, or
- 4. No risk at all
- 8. DON'T KNOW

913C Is there a cure for AIDS?

1. Yes Common 2. No-Not common

9.

Don't Know

Female Questionnaire

IX-B VIOLENCE

- 914. Thinking back to your childhood and adolescence, did you ever see or hear your parents or step-parents physically abuse each other?
 - 1. YES
 - 2. NO
 - 3. DID NOT LIVE WITH 2 PARENTS \rightarrow GO TO Q916
 - 8. DR/REF

915. As a child, have you ever being beaten or physically mistreated in any way by anyone in your family?

- 1. YES
- 2. NO
- 8. DR/REF
- 916. THE INTERVIEWER SHOULD GO BACK TO PAGE 3 AND RECORD HOW MANY TIMES THE RESPONDENT LIVED WITH A MEN AS HUSBAND AND WIFE (SEE Q111): ______TIMES

IF Q916=0 GO TO Q936; IF Q916>0 CONTINUE

The next set of questions is about violence and physical abuse that may have happened between you and a partner or ex-partner. When we say a partner we mean a husband, ex-husband, as well as any other man you have been living with as husband and wife.

918. Please tell me	if any of your part-	919. When was the last time when	920. During the last year, how many			
ners or ex-partners ever (READ A-H):		(A-H) happened to you?	times did			
• • • •			(A-H) happen to you?			
A. Insulted you, or		1. WITHIN THE LAST YEAR \rightarrow Q920				
swore at you?	2. NO \rightarrow Q918_B	2. 1-3 YEARS AGO \rightarrow Q918_B	66. ALMOST DAILY			
	8. DK \rightarrow Q918 B	3. 4-5 YEARS AGO \rightarrow Q918 B	TIMES 77. WEEKLY			
	9. REF \rightarrow Q918 B	4. 5 YEARS AGO OR MORE \rightarrow Q918 B	88. DON'T REMEMB.			
			99. REFUSES			
B. Threatened	1. YES → Q919	1. WITHIN THE LAST YEAR \rightarrow Q920	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			
to hurt you or	2. NO \rightarrow Q918 C	2. 1-3 YEARS AGO \rightarrow Q918_C	66. ALMOST DAILY			
	8. DK \rightarrow Q918 C	3. 4-5 YEARS AGO \rightarrow Q918 C	TIMES 77. WEEKLY			
about?	9. REF \rightarrow Q918_C	4. 5 YEARS AGO OR MORE \rightarrow Q918_C	88. DON'T REMEMB.			
			99. REFUSES			
C. Pushed you,	1. YES → Q919	1. WITHIN THE LAST YEAR \rightarrow Q920				
	2. NO \rightarrow Q918_D	2. 1-3 YEARS AGO \rightarrow Q918_D	66. ALMOST DAILY			
you, or threw	8. DK \rightarrow Q918_D	3. 4-5 YEARS AGO \rightarrow Q918_D	TIMES 77. WEEKLY			
something at you?	9. REF \rightarrow Q918_D	4. 5 YEARS AGO OR MORE \rightarrow Q918 D	88. DON'T REMEMB.			
			99. REFUSES			
D. Slapped you	1. YES → Q919	1. WITHIN THE LAST YEAR \rightarrow Q920				
or twisted your	2. NO \rightarrow Q918 E	2. 1-3 YEARS AGO \rightarrow Q918_E	66. ALMOST DAILY			
arm?	8. DK \rightarrow Q918 E	3. 4-5 YEARS AGO \rightarrow Q918 E	TIMES 77. WEEKLY			
d1111.	9. REF \rightarrow Q918 E	4. 5 YEARS AGO OR MORE \rightarrow Q918 E	88. DON'T REMEMB.			
	$9. \text{ KEP} \rightarrow \text{Q910}_{\text{E}}$	$ 4.51 \text{ EARS ACC OR MORE} \rightarrow 0.16 \text{ E}$				
F II'4 . '41.1'	1 VEG 0010		99. REFUSES			
	1. YES \rightarrow Q919	1. WITHIN THE LAST YEAR \rightarrow Q920				
	2. NO \rightarrow Q918_F	2. 1-3 YEARS AGO \rightarrow Q918_F	66. ALMOST DAILY			
thing else?	8. DK \rightarrow Q918_F	3. 4-5 YEARS AGO \rightarrow Q918_F	TIMES 77. WEEKLY			
	9. REF \rightarrow Q918_F	4. 5 YEARS AGO OR MORE \rightarrow Q918_F	88. DON'T REMEMB.			
			99. REFUSES			
F. Threatened you	1. YES \rightarrow Q919	1. WITHIN THE LAST YEAR \rightarrow Q920				
with a knife or	2. NO \rightarrow Q918_G	2. 1-3 YEARS AGO \rightarrow Q918_G	66. ALMOST DAILY			
other weapon?	8. DK $\rightarrow \mathbf{Q918}^{\mathbf{G}}$	3. 4-5 YEARS AGO \rightarrow Q918 G	TIMES 77. WEEKLY			
		4. 5 YEARS AGO OR MORE \rightarrow Q918_G	88. DON'T REMEMB.			
			99. REFUSES			
G. Kicked you,	1. YES → Q919	1. WITHIN THE LAST YEAR \rightarrow Q920				
	$1. 1123 \rightarrow Q919$ 2. NO $\rightarrow Q918_H$	2. 1-3 YEARS AGO \rightarrow Q918 H	66. ALMOST DAILY			
you up?	8. DK \rightarrow Q918_H	3. 4-5 YEARS AGO \rightarrow Q918_H	TIMES 77. WEEKLY			
	$ 9. \text{ KEF} \rightarrow \mathbf{Q918}_{\mathbf{H}}$	4. 5 YEARS AGO OR MORE \rightarrow Q918_H	88. DON'T REMEMB.			
			99. REFUSES			
H. Physically forced		1. WITHIN THE LAST YEAR \rightarrow Q920				
you to have	2. NO \rightarrow BOX 9-I	2. 1-3 YEARS AGO → BOX 9-I	66. ALMOST DAILY			
sexual relations	8. DK → BOX 9-I	3. 4-5 YEARS AGO \rightarrow BOX9-I	TIMES 77. WEEKLY			
even though you	9. REF \rightarrow BOX 9-I	4. 5 YEARS AGO OR MORE \rightarrow BOX 9-I	88. DON'T REMEMB.			
did not want to?			99. REFUSES			
and not many tot		1	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			

B	X 9-I	
	ALL Q918_AQ918_H >1 (NEVER EXPERIENCED ANY TYPE OR ABUSE) GO TO Q936; ELSE CONTIN	UE

II.	
III.	
IV	

BOX 9-II

< IF ANY PHYSICAL VIOLENCE TOOK PLACE DURING THE LAST YEAR (ANY Q919_C--Q919_H=1), CONTINUE; < IF ANY F PHYSICAL VIOLENCE TOOK PLACE MORE THAN ONE YEAR AGO (ANY Q919_C--Q919_H>1) GO TO Q925; < IF R. SUFFERED ONLY VERBAL VIOLENCE (Q918_C--Q918_H>1) THEN GO TO Q935

922. In the past 12 months, did you have any swelling, bruises, cuts, or other physical injuries as a result of this/these incident(s)?

1. YES

- 2. NO \rightarrow GO TO Q925
- 8. DON'T REMEMBER \rightarrow GO TO Q925
- 923. In the past 12 months, did you see a doctor, or other medical care provider for medical treatment of these injuries?

YES
 NO → GO TO Q925
 DO NOT REMEMBER → GO TO Q925

924. Did this(these) injury(ies) require hospitalization?

1. YES

- 2. NO
- 8. DO NOT REMEMBER
- 925. During or after a violent incident, does your partner want to have sexual relations with you sometimes, always, or never?
 - 1. YES, SOMETIMES
 - 2. YES, ALWAYS
 - 3. NO, NEVER
 - 8. DO NOT REMEMBER/REFUSE
- 926. Do you believe that violence incidents affected your health?
 - 1. YES
 - 2. NO \rightarrow GO TO Q928
 - 8. DO NOT KNOW \rightarrow **GO TO Q928**
- 927. What health problems did you have that you think were related to violence? (**READ A—K**)

	YES	<u>NO</u>	DR/REF
A. Injuries?	1	2	8
B. Chronic pains?	1	2	8
C. General deterioration of health?	1	2	8
D. Depression?	1	2	8
E. Sleep disorders?	1	2	8
F. Suicide attempt(s)?	1	2	8
G. Low self-esteem?	1	2	8
H. Unplanned pregnancy?	1	2	8
I. Miscarriage or other pregnancy complications?	1	2	8
J. Gynecological disorders?	1	2	8
K. Anything else?	1	2	8

928. (READ ONLY IF RESPONDENT HAS EVER HAD A LIVING CHILD) Would you say that this violence has affected your child(ren)?

- 1. YES
- 2. NO \rightarrow **GO TO Q930**

3. NO CHILDREN \rightarrow **GO TO Q930**

- 8. NOT SURE \rightarrow **GO TO Q930**
- 929. Please tell me if your child(ren) were affected in the following ways (**READ A—F**):

	<u>YES</u>	<u>NO</u>
A. Witnessed violence	1	2
B. Children living in fear	1	2
C. Children injured too	1	2
D. Children left home, live with relative	1	2
E. Decreased Learning abilities	1	2
F. Anything else (specify)		1

- 930. Did you talk to anyone about this(these) incidents of violence?
 - 1. YES
 - 2. NO \rightarrow GO TO Q932

931. Who did you talk with? (MARK ALL MENTIONED AND PROBE FOR ANYONE ELSE)?

	MENTIONED	NOT MENTIONED
A. YOUR MOTHER	1	2
B. OTHER RELATIVE	1	2
C. HUSBAND'S (PARTNER'S) FAMILY	1	2
D. CHILDREN	1	2
E. FRIEND	1	2
F. NEIGHBOR	1	2
G. DOCTOR/HEALTH PROVIDER/SOCIAL WORKER	1	2
H. POLICE	1	2
I. LEGAL ADVISER	1	2

IF Q931_G, Q931_H, AND Q931_I=1 GO TO Q933; ELSE CONTINUE

- 932. What is the main reason you have never sought any medical or legal help?
 - 1. DID NOT KNOW WHERE TO SEEK HELP
 - 2. NO USE/WOULD NOT DO ANY GOOD
 - 3. EMBARRASSED
 - 4. AFRAID OF MORE BEATINGS/BEING PUNISHED
 - 5. AFRAID OF DIVORCE/END OF RELATIONSHIP
 - 6. AFRAID OF LOOSING THE CHILDREN
 - 7. THOUGHT WOULD NOT BE TAKEN SERIOUSLY/NOT BELIEVED/LAUGHED AT
 - 8. VIOLENCE IS NORMAL/NO NEED TO COMPLAIN
 - 9. THOUGHT SHE WOULD BE BLAMED
 - 10. BRING BAD NAME TO FAMILY
 - 20. OTHER_
 - 88. DK/REF

933. Did anyone intervene or try to stop the violence? (MARK ALL MENTIONED AND PROBE FOR ANYONE ELSE)?

	MENTIONED	NOT MENTIONED
A. YOUR MOTHER	1	2
B. OTHER RELATIVE	1	2
C. HUSBAND'S (PARTNER'S) FAMILY	1	2
D. CHILDREN	1	2
E. FRIEND	1	2
F. NEIGHBOR	1	2
G. DOCTOR/HEALTH PROVIDER/SOCIAL WORKER	1	2
H. POLICE	1	2
I. LEGAL ADVISER	1	2

934. Could you tell me a little more about what usually happens when your partner is/was violent. Are there any particular situations that make him violent? (CIRCLE ALL THAT APPLY PROBING "ANY OTHER...")

		··)
	MENTIONED	NOT MENTIONED
A. WHEN DRUNK	1	2
B. WHEN SHE DOES NOT LOOK AFTER CHILDREN	1	2
C. WHEN THE FAMILY HAS MONEY TROUBLES	1	2
D. WHEN HE IS UNEMPLOYED	1	2
E. WHEN SHE IS UNEMPLOYED/HAS NO INCOME	1	2
F. WHEN THERE ARE FAMILY PROBLEMS/MOTHER-IN-LAW PROBLEM	AS 1	2
G. WHEN HE IS JEALOUS	1	2
H. WHEN SHE IS PREGNANT	1	2
I. WHEN HE CANNOT GET ALCOHOL/DRUGS	1	2
J. WHEN HE DOES NOT HAVE FOOD AT HOME (DINNER NOT READY)) 1	2
K. WHEN HE ACTS ACCORDING TO THE "KANUN"	1	2
L. OTHER	1	2

935. In Tirana there is a hotline number where a woman can ask advice about domestic violence. Do you want to know the hotline number?

1. YES, NUMBER GIVEN (TELEPHONE NUMBER: 233 408)

- 2. NO
- 936. Sometimes a husband is annoyed or angered by things that his wife does. In your opinion, is a husband justified in hitting or beating his wife in the following situations (**READ A—H**):

	YES	<u>NO</u>	<u>DK</u>
A. If she goes out without telling him?	1	2	8
B. If she neglects the children?	1	2	8
C. If she argues with him?	1	2	8
D. If she refuses to have sex with him?	1	2	8
E. If he is not happy with her household work or food provisions?	1	2	8
F. If she asks him whether he has other girlfriends?	1	2	8
G. If he finds out that she has been unfaithful?	1	2	8
H. If she dresses too sexy or spends too much on her "look"?	1	2	8

- 937. At any time in your life, have you ever been forced by a man to have sexual intercourse against your will? (For this question, sexual intercourse includes vaginal, anal or oral penetration)
 - 1. YES

2. NO \rightarrow END OF INTERVIEW, GO TO Q940

8. DON'T REMEMBER → END OF INTERVIEW, GO TO Q940

938. How old were you the first time you were forced by a man to have sexual intercourse against your will?

_____ AGE 88. DON'T REMEMBER

939. At that time, what was your relationship with the person(s) who forced you to have sexual intercourse?

- 1. STRANGER
- 2. ACQUAINTANCE
- 3. FRIEND
- 4. DATE
- 5. BOYFRIEND
- 6. HUSBAND OR PARTNER
- 7. EX-HUSBAND OR EX-PARTNER
- 8. FATHER OR STEP-FATHER
- 9. OTHER RELATIVE (SPECIFY_____)
- 10. CO-WORKER
- 11. TEACHER
- 77. OTHER (SPECIFY_____
- 98. DON'T REMEMBER/REF

940. THANK THE WOMAN FOR GIVING HER TIME AND RECORD THE TIME THE INTERVIEW ENDED :

TIME INTERVIEW ENDED _____: _____:

ALBANIA REPRODUCTIVE HEALTH SURVEY 2002 MALE QUESTIONNAIRE

of me	For the National Institute of Public Health en and women in Albania. The purpose of the survey is to colle a services in Albania.							
be con answe	Id like to ask you about your health and where you obtain healt nfidential. The interview is completely voluntary and if we sho er, just let me know and we'll go on to the next question. The i uld like to start now, is that OK?	ould co	me to an	y ques	tion th	at you	ı don't	
SIGN	NATURE OF THE INTERVIEWER I	DAY		M	ONTH	- 		
	RK IF THE MAN AGREES TO BE INTERVIEWED		1. YE 2. NC	$\mathbf{S} \rightarrow \mathbf{C}$	CONT	INUE		EW
TIME	STARTED:: ID) NUM	IBER					
<u>I. BAC</u>	CKGROUND CHARACTERISTICS							
100.	In what month and year were you born?							
	MONTH YEAR			98 E	ON'T	KNO	W	
101.	How old are you (at last birthday)?YEARS OLD			98.	DON"	T KNO	WC	
	MAKE SURE THAT AGE AND DATE OF	BIRT	H COR	RESP	OND			
102.	What is the highest level of education you completed, not co	ounting	the curr	ent gra	ade yo	u are i	n?	
	000. NEVER ATTENDED (NO FORMAL EDUCATION)							
	1. GENERAL SCHOOL		1	2	3	4	5	88
	2. THEORETICAL HIGH SCHOOL/ GIMNAZIUM		1	2	3	4		88
	3. PROFESSIONAL (VOCATIONAL) SCHOOL		1	2	3	4	5	88
	4. TECHNICAL SCHOOL (POSTSECONDARY)		1	2	3	4	5	88
	5. UNIVERSITY		1	2	3	4	5+	88
	6. POST UNIVERSITY/POSTGRADUATE STUDIES		1	2	3	4	5+	88
	888. DON'T REMEMBER/ DON'T KNOW							
	IF Q102 = 2 OR 3 CONTINUE; EI	LSE G	о то q	104				
103.	Do you have a high school diploma?							
	1. YES 2. NO							

- 104. Do you currently work outside of the home (at least 20 hours per week)?
 - 1. YES \rightarrow GO TO Q106
 - 3. NO

105. What is the main reason that you are not working at this time?

- 1. ATTENDING SCHOOL
- 2. I NTERNAL DISPLACEMENT
- 3. LOOKING FOR WORK
- 4. LAID OFF
- 5. DOES NOT NEED/WANT/LIKE TO WORK
- 6. MEDICAL LEAVE
- 7. CARING FOR CHILDREN
- 8. INABILITY TO FIND/AFFORD CHILD CARE
- 9. HOMEMAKER
- 10. PERMANENT DISABILITY
- 11. PARTNER/SPOUSE DOES NOT ALLOW RESPONDENT TO WORK
- 12. PARENTS DO NOT ALLOW RESPONDENT TO WORK
- 13. ODD JOBS (<20 HOURS PER WEEK)
- 14. TEMPORARILY BACK FROM WORK ABROAD
- 20. OTHER (SPECIFY)___
- 106. I would like to ask you about where you have lived. For most of the time until you were 12 years old, did you live in a town, or in a village?
 - 1. TOWN
 - 2. VILLAGE
- 107. In what month and year did you start to live continuously in _____ (NAME THE PLACE OF RESIDENCE)?

MONTH____YEAR ____

00. ALWAYS, SINCE BIRTH \rightarrow **GO TO Q108** 88. DON'T REMEMBER

- 107A. Just before you moved here (CURRENT PLACE OF RESIDENCE), did you live in a town, a village, or outside Albania ?
 - 1. A town (URBAN AREA),
 - 2. A village (RURAL AREA), or
 - 3. Outside Albania?
- 108. Are you currently married, not married but living with someone, separated, divorced, widowed, or have you never been married ?
 - 1. MARRIED \rightarrow GO TO Q111
 - 2. NOT MARRIED BUT LIVING WITH A PARTNER \rightarrow GO TO Q111
 - 3. SEPARATED \rightarrow GO TO Q111
 - 4. DIVORCED \rightarrow GO TO Q111
 - 5. WIDOWED \rightarrow GO TO Q111
 - 6. NEVER MARRIED
- 109. Have you ever lived with a girlfriend or partner? (LIVING TOGETHER MEANS HAVING A SEXUAL RELATIONSHIP WHILE SHARING THE SAME USUAL RESIDENCE.)

1. YES \rightarrow **GO TO Q111** 2. NO

110. If you could choose exactly the number of children to have in your whole life, how many would that be?

____ CHILDREN

22. AS MANY AS GOD GIVES
33. AS MANY AS WIFE WOULD WANT
88. NOT SURE/DON'T REMEMBER
GO TO Q122

111. How many times have you been married or lived with a woman as husband and wife?

	TIMES	9. REFU	$JSAL \to \mathbf{GO} \ \mathbf{TO} \ \mathbf{Q118}$			
TIMES	112. In what month and year did you begin living with your (first, second, third, or fourth) wife / partner?	113. How old was your I, II, III, IV wife / partner when you started to live together?	114. What was the highest level in school that your I,II,III,IV wife / partner attended when you got married / started to live together ?	115. What is your current union relationship with your I, II, III, IV, wife / partner, are you still in the relationship or how did the relationship end?	116. In what month and year did your union with your I,II,III,IV, .wife / partner end?	117. IF:
I	MTH YR 88. DON'T KNOW/REF.	AGE 88. DK	0. NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married →Q117 Living with partner→ Q117 Separated Divorced Widowed 	MTH YR 88. DON'T KNOW / REF.	Q111=1 GO TO Q118 ELSE CON- TINUE
п	MTH YR 88. DON'T KNOW/REF.	AGE 88. DK	0.NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married → Q117 Living with partner→ Q117 Separated Divorced Widowed 	MTH YR 88. DON'T KNOW / REF.	Q111=2 GO TO Q118 ELSE CON- TINUE
ш	MTH YR 88. DON'T KNOW/REF.	AGE 88. DK	0.NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married →Q117 Living with partner→ Q117 Separated Divorced Widowed 	MTH YR 88. DON'T KNOW / REF.	Q111=3 GO TO Q118 ELSE CON- TINUE
IV	MTH YR 88. DON'T KNOW/REF.	AGE 88. DK	0.NEVER ATTENDED 1. PRIMARY (1-8) 2. SECONDARY (9-12) 3. VOCATIONAL 4. TECHNICAL SCH. 5. UNIVERSITY 8. UNKNOWN	 Married →Q117 Living with partner→ Q117 Separated Divorced Widowed 	MTH YR 88. DON'T KNOW / REF.	GO TO Q118

118. When you first got married/started living together as husband and wife did you wish to have any children?

1. YES

- 2. NO \rightarrow GO TO Q120
- 8. NOT SURE \rightarrow GO TO Q120

119. How many children did you wish to have when you first got married?

CHILDREN

22. AS MANY AS GOD GIVES33. AS MANY AS WIFE WANTS88. NOT SURE/DON'T REMEMBER

120. How many children did your wife/partner wish to have when you first got married?

____ CHILDREN

- 22. AS MANY AS GOD GIVES33. AS MANY AS RESPONDENT WANTS
- 77. NEVER DISCUSSED NUMBER OF CHILDREN
- 88. NOT SURE/DON'T REMEMBER

IF CURRENTLY SEPARATED, DIVORCED OR WIDOWED (Q115_LAST=3,4,5) GO TO Q122; ELSE CONTINUE

121. Is your wife currently employed?

- 1. YES
- 2. NO
- 8. DON'T KNOW
- 122. Since 1991, have you worked outside Albania?
 - 1. YES
 - 2. NO \rightarrow GO TO Q124
 - 8. DON'T KNOW \rightarrow GO TO Q124

		Ap	opendix D					
122A.]	How many times have you worked ou	tside Albania s	ince 1991?					
	TIMES							
123.	When was the last time you worked	outside Albani	ia ? (BEGINNING	THE TRIP – MONT	TH AND YEAR)			
	MONTH			/IBER				
		99	. REFUSAL					
123A.	A. In what month and year did you return from the last trip? (END OF TRIP – MONTH AND YEAR)							
	MONTH	YEAR 88	. DON'T REMEN	/BER				
			. REFUSAL					
124.	More or less how many hours a day	do vou listen	to the radio?					
	HOURS A DAY	00. NEVER	\rightarrow GO TO Q128					
				SS TO RADIO \rightarrow G	GO TO Q128			
		77. NOT EV 88. DON'T						
125.	What stations do you most often list	en to? (DDOE	RE EOD MODE 1	THAN ONE STATI	ON DO NOT DEAD LIST)			
125.	what stations do you most often his		DE FOR MORE I	NOT	on, bo not kead hist)			
			MENTIONED	MENTIONED				
	A. TOP ALBANIA B. RADIO TIRANA		1	2 2				
	C. CLUB FM		1	2				
	D. RASH		1	2				
	E. STINET F. GOLD MUSIC		1	2 2				
	G. EUROPE PLUS		1	2				
	H. ITALIAN STATIONS		1	2				
	I. BBC		1	2 2				
	J. OTHER		1	Z				
126.	What types of programs do you mos READ LIST)	t often listen to	o? (PROBE FOR	MORE THAN ON	E PROGRAM, DO NOT			
	KEAD LIST)			NOT				
			MENTIONED	MENTIONED				
	A. NEWS B. SPORTS		1 1	2 2				
	C. MUSIC		1	2				
	D. PLAYS/DRAMAS		1	2				
	E. CHURCH/RELIGIOUS PROGI F. WOMEN'S PROGRAMS	RAMS	1	2 2				
	G. HEALTH PROGRAMS		1	2				
	H. POLITICAL EVENTS		1	2				
	I. OTHER		1	2				
127.	What times do you most often listen	to the radio? ((PROBE FOR M	ORE THAN ONE	ΓΙΜΕ, DO NOT READ			
	LIST)			NOT				
			MENTIONED	NOT <u>MENTIONED</u>				
	A. 6 - 8 AM		1	2				
	B. 8-10 AM		1	2				
	C. 10AM-NOON		1	2				
	D. NOON-2 PM		1	2				
	E. 2 - 4 PM		1	2				
	F. 4 - 6 PM		1	2				
	G. 6 - 8 PM		1	2				
	H. 8-10 PM		1	2				
	I. AFTER 10 PM		1	2				
	J. NO REGULAR TIMES		1	2				

128. More or less how many hours a day do you spend watching television?

HOURS A DAY $\,$ 00. NEVER \rightarrow GO TO Q132 $\,$

55. DOES NOT HAVE ACCESS TO TV \rightarrow GO TO Q132

66. WHEN THE HOUSEHOLD HAS ELECTRICITY

- 77. NOT EVERY DAY
- 88. DON'T KNOW

129. What channels do you most often watch? (PROBE FOR MORE THAN ONE CHANNEL, DO NOT READ LIST)

	METIONED	NOT METIONED
A. TVSH	1	2
B. TVKLAN	1	2
C. TVA	1	2
D. TVKOHA	1	2
E. VISION PLUS	1	2
F. SHIJAK TV	1	2
G. TELENORBA	1	2
H. TVALBA	1	2
I. CALVIN	1	2
J. FOREIGN CHANNELS (e.g. CNN, BBC, EURONEWS, RAI)	1	2
K. OTHER	1	2

130. What types of programs do you most often watch? (PROBE FOR MORE THAN ONE PROGRAM, DO NOT READ LIST)

	MENTIONED	NOT METIONED
A. NEWS	1	2
B. COMERCIALS	1	2
C. ENTERTAINMENT PROGRAMS	1	2
D. SERIALS/MOVIES	1	2
E. SPORTS	1	2
F. MUSIC PROGRAMS, VIDEO CLIPS	1	2
G. PLAYS/DRAMAS	1	2
H. CHILDREN'S PROGRAMS	1	2
I. CHURCH/RELIGIOUS PROGRAMS	1	2
J. WOMEN'S PROGRAMS	1	2
K. HEALTH PROGRAMS	1	2
L. POLITICAL EVENTS	1	2
M. OTHER	1	2

131. What times do you most often watch television? (CIRCLE ALL MENTIONED, DO NOT READ LIST)

	MENTIONED	NOT METIONED
A. 6 - 8 AM	1	2
B. 8-10 AM	1	2
C. 10AM-NOON	1	2
D. NOON-2 PM	1	2
E. 2 - 4 PM	1	2
F. 4 - 6 PM	1	2
G. 6 - 8 PM	1	2
H. 8-10 PM	1	2
I. AFTER 10 PM	1	2
J. NO REGULAR TIMES	1	2

132. How often do you read a newspaper?

- 1. DAILY/NEARLY EVERY DAY
- 2. ABOUT 3-4 TIMES PER WEEK
- 3. ONCE OR TWICE PER WEEK
- 4. LESS THAN ONCE PER WEEK
- 5. NEVER/ALMOST NEVER

II. SEX EDUCATION

The next set of questions is about sex education.

201. Do you think schools should teach courses about human reproduction, contraception, and prevention of sexually transmitted diseases (STDs)?

1. YES 2. NO \rightarrow GO TO Q203 8. DK 9. NR \rightarrow GO TO Q203

202. At what year of age should schools begin to teach about? (READ A-C)

A. Human Reproduction?	 77. SHOULD NOT BE TAUGHT IN SCHOOL
B. Contraception?	 88. DK
C. Sexually Transmitted Diseases (STDs)	 99. NR

GO TO BOX 2-I

203. Now I want to read some reasons for which one may oppose sex education in school. Please tell me if you agree or don't agree. (**READ A-D**)

	<u>AGREE</u>	DISAGREE	<u>DK</u>	<u>NR</u>
A. Sex education will give adolescents the idea to begin sex earlier	1	2	8	9
B. Sex education should be taught only in the house	1	2	8	9
C. Sex education goes against my religious beliefs	1	2	8	9
D. Teachers do not have enough training to teach such courses	1	2	8	9

BOX 2-I

IF RESPONDENT IS 15-24 YEARS OF AGE CONTINUE; IF HE IS 25-49 YEARS GO TO SECTION III

204. Before you were 18 years old, did a parent ever talked to you about.....(READ A-F)

	<u>YES</u>	<u>NO</u>	<u>DK/DR</u>	<u>REF</u>
B. How Pregnancy Occurs?	1	2	8	9
D. Contraceptive Methods? (How to avoid getting a woman pregnant?)	1	2	8	9
E. HIV/AIDS?	1	2	8	9
F. Other Sexually Transmitted Diseases?	1	2	8	9

TOPIC	205. Before you were 18 years	206. How old were you
	old, have you ever been taught about (READ A-G) while at school?	when you first were taught about(READ A-G) at school?
A. Menstrual Cycle	$1 \text{ YES} \rightarrow \text{GO TO Q206}$ $2 \text{ NO} \rightarrow \text{GO TO Q205}_B$ $8 \text{ DK} \rightarrow \text{GO TO Q205}_B$ $9 \text{ NR} \rightarrow \text{GO TO Q205}_B$	
B. Female Reproductive System	$1 \text{ YES} \rightarrow \text{GO TO Q206}$ $2 \text{ NO} \rightarrow \text{GO TO Q205_C}$ $8 \text{ DR} \rightarrow \text{GO TO Q205_C}$ $9 \text{ NR} \rightarrow \text{GO TO Q205_C}$	
C. Male Reproductive System	$1 \text{ YES } \rightarrow \text{GO TO } \text{Q206}$ $2 \text{ NO } \rightarrow \text{GO TO } \text{Q205_D}$ $8 \text{ DR } \rightarrow \text{GO TO } \text{Q205_D}$ $9 \text{ NR } \rightarrow \text{GO TO } \text{Q205_D}$	
D. How Pregnancy Occurs	$1 \text{ YES } \rightarrow \text{GO TO } \text{Q206}$ $2 \text{ NO } \rightarrow \text{GO TO } \text{Q205}_{\text{E}}$ $8 \text{ DR } \rightarrow \text{GO TO } \text{Q205}_{\text{E}}$ $9 \text{ NR } \rightarrow \text{GO TO } \text{Q205}_{\text{E}}$	
E. Contraceptive Methods	$1 \text{ YES } \rightarrow \text{GO TO } \text{Q206}$ $2 \text{ NO } \rightarrow \text{GO TO } \text{Q205}_{\text{F}}$ $8 \text{ DR } \rightarrow \text{GO TO } \text{Q205}_{\text{F}}$ $9 \text{ NR } \rightarrow \text{GO TO } \text{Q205}_{\text{F}}$	
F. HIV/AIDS	$1 \text{ YES } \rightarrow \text{GO TO } \text{Q206}$ $2 \text{ NO } \rightarrow \text{GO TO } \text{Q205}_{\text{G}}$ $8 \text{ DR } \rightarrow \text{GO TO } \text{Q205}_{\text{G}}$ $9 \text{ NR } \rightarrow \text{GO TO } \text{Q205}_{\text{G}}$	
G. Other Sexually Transmitted Diseases	$\begin{array}{c} 1 \text{ YES } \rightarrow \textbf{GO TO Q206} \\ 2 \text{ NO } \rightarrow \textbf{GO TO Q208} \\ 8 \text{ DR } \rightarrow \textbf{GO TO Q208} \\ 9 \text{ NR } \rightarrow \textbf{GO TO Q208} \end{array}$	

READ EACH QUESTION 205-207 FROM THE TABLE FOR EACH TOPIC OF SEX EDUCATION:

208. In your opinion, who or what was the most important source of information you have had about topics related to sexual matters?

MOTHER
 FATHER
 RELATIVE
 GIRLFRIEND
 FRIENDS
 CO-WORKER
 COLLEAGUES, PEER
 PARTNER/WIFE
 DOCTOR

10. NURSE, MIDWIFE

- 11. TEACHER
- 12. PHARMACIST
- 13. BOOKS
- 14. NEWSPAPERS, MAGAZINES, BROCHURES, FLYERS
- 15. RADIO
- 16. TV
- 20. OTHER (SPECIFY):
- 88. DON'T REMEMBER

III. FERTILITY/PREGNANCY

305.	Have you ever made a wo	man pregnant?		
	1. YES \rightarrow GO TO Q307 2. NO 3. NOT SURE 4. NEVER HAD SEX \rightarrow	GO TO MODULE IV	, PAGE 12	
306.	Has a female partner/wife	ever had a stillbirth, ect	opic pregnancy, miscarriage, or a	n induced abortion?
	1. YES \rightarrow GO TO Q312 2. NO \rightarrow GO TO MOD			
307.	Have you ever had any liv	e-born children?		
	 YES NO → GO TO Q312 			
308.	How many living children	do you have, including	those who do not live with you?	
	CHILDREN			
309.	Have you ever had a child	born alive who later die	d or died in the first hours or day	s after birth?
	1. YES 2. NO \rightarrow GO TO Q312			
310.	How many children died?	CHILD	REN	
311.	So altogether you had a to	tal of (Q308+Q	310) live births?	
	 YES NO → CHECK Q308 	AND Q310 AND MAK	E CHANGES IF NECESSARY	7
312.			a got a woman pregnant. Please to non. How did that first pregnancy	
	 LIVE BIRTH (SINGL STILLBIRTH (SINGL SPONTANEOUS ABO INDUCED ABORTIO ECTOPIC PREGNAN MOLAR PREGNANC 	E OR MULTIPLE) ORTION (MISCARRIA) N CY	GE)	
	10. CURRENTLY PREGN 88. DON'T KNOW	VANT (FIRST TIME) –	→ GO TO Q315	
313.	In what month and year di	d that first pregnancy en	d?	
	MONTH	YEAR	88. DON'T RE	MEMBER
	IF Q313_Y =8888 (I	OOESN'T REMEMBE	R YEAR) CONTINUE; OTHE	RWISE GO TO Q315

314. How old were you at that time? ____YEARS 88. DON'T REMEMBER

- 315. Thinking back to the time when your partner got pregnant, with that first pregnancy, did you want her to get pregnant or not?
 - 1. YES
 - 2. NO
 - 8. NOT SURE
- 316. How old was your partner when she got pregnant with that first pregnancy?

AGE ____ 88 DON'T KNOW

317. What was your relationship to her when she first got pregnant?

1. WIFE/PARTNER	6. JUST MET
2. FIANCEÉ	7 CASUAL SEXUAL PARTNER
3. GIRLFRIEND	8. RELATIVE
4. FRIEND	11. SHE WAS A PROSTITUTE
5. ACQUAINTANCE	20. OTHER (SPECIFY)
	88. DON'T REMEMBER

IF Q312=10 DO NOT ASK Q318 BUT CIRCLE="1"

- 318. As far as you know, is any woman currently pregnant by you?
 - 1. YES
 - 2. NO \rightarrow **GO TO BOX 3-I**
 - 3. NOT SURE \rightarrow **GO TO BOX 3-I**
- 319. How many months pregnant is the woman?

_____MONTHS 88. DOES NOT KNOW 99. REFUSED / NO RESPONSE

IF Q312=10 DO NOT ASK Q320 BUT WRITE THE ANSWER FROM Q315

320. Thinking back to the time when she got pregnant, did you want her to get pregnant or not?

- 1. YES
- 2. NO
- 3. NOT SURE

BOX 3-I

IF Q311=0 (NEVER HAD A LIVE BIRTHS) GO TO Q327; OTHERWISE CONTINUE WITH HISTORY OF ALL LIVE BIRTHS

HISTORY OF ALL LIVE BIRTHS

Now I would like to talk to you about all live births you and your partner(s) have ever had. Please, make sure you include all births you have fathered regardless of which partner was the mother of the child. Let's start with your most recent birth, please give me the following information:

WRITE ## OF LIVE BIRTHS _____ (FROM Q311)

#						326
	321	322	323	324	325	IF Q321 < 1997→GO TO NEXT LIVE- BIRTH
	When was that live birth? (month & year)	How many weeks or months had your partner been pregnant when she gave birth?	Was the baby a boy or a girl?	Is the child still alive?	How old was the child when he/she died? (RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YRS.)	Just before you got pregnant, did you want to get pregnant then, did you want to get pregnant later, or did you not want to get pregnant then or any time in the future?
1 LAST LIVE BIRTH	MTH YR 98. DK/NR	1 WEEKS OR 2 Months 888. DK 999. NR/REF	1. BOY 2. GIRL 3. BOTH	1. YES → Q326 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS 888. DK	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SUBE
					999. NR/REF	8. NOT SURE
2	MTH YR	1 WEEKS OR 2. MONTHS	1. BOY		1 DAYS OR 2. MTHS OR	1. WANTED TO GET PREG. THEN 2. WANTED TO GET PREG. LATER
	98. DK/NR	888. DK 999. NR/REF	2. GIRL 3. BOTH	1. YES → Q326 2. NO	3YEARS 888. DK 999. NR/REF	3. DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE 8. NOT SURE
3	MTH YR 98. DK/NR	1 WEEKS OR 2 Months 888. DK 999. NR/REF	1. BOY 2. GIRL 3. BOTH	1. YES → Q326 2. NO	1 DAYS OR 2 MTHS OR 3 YEARS	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE
					888. DK 999. NR/REF	8. NOT SURE
4		1 WEEKS			1 DAYS OR 2 MTHS OR	
	MTH YR 98. DK/NR	OR 2 MONTHS 888. DK 999. NR/REF	1. BOY 2. GIRL 3. BOTH	1. YES → Q326 2. NO	3 YEARS 888. DK 999. NR/REF	 WANTED TO GET PREG. THEN WANTED TO GET PREG. LATER DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE NOT SURE

#	321	322	323	324	325	326
					How old was the child when he/she died?	IF Q321 < 1997 →GO TO NEXT LIVE BIRTH
	When was that live birth? (month & year)	How many weeks or months had your partner been pregnant when she gave birth?	Was the baby a boy or a girl?		(RECORD DAYS IF LESS THAN 1 MONTH; MONTHS IF LESS THAN TWO YEARS; OR YRS.)	Just before she got pregnant, did you want to get pregnant then, did you want to get pregnant later, or did you not want to get pregnant then or any time in the future?
<u>5</u>		1 WEEKS			1 DAYS OR	1. WANTED TO GET PREG. THEN
	MTH	OR	1. BOY		2 MTHS_OR	2. WANTED TO GET PREG. LATER
	YR	2 MONTHS 888. DK	2. GIRL 3. BOTH	1. YES \rightarrow Q326 2. NO	3 YEARS	3. DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE
	98. DK/NR	999. NR/REF	5. 60111	2. NO	888. DK 999. NR/REF	8. NOT SURE
<u>6</u>	MTH	1 WEEKS			1 DAYS OR	1. WANTED TO GET PREG. THEN
	YR	OR 2 MONTHS	1. BOY		2 MTHS OR	2. WANTED TO GET PREG. LATER
	98. DK/NR	2 MONTHS 888. DK 999. NR/REF	2. GIRL 3. BOTH	1. YES \rightarrow Q326 2. NO	3 YEARS	3. DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE
					888. DK 999. NR/REF	8. NOT SURE
Z	MTH	1 WEEKS OR			1 DAYS OR	1. WANTED TO GET PREG. THEN
	YR	2MONTHS	1. BOY		2 MTHS OR	2. WANTED TO GET PREG. LATER
	98. DK/NR	888. DK 999. NR/REF	2. GIRL 3. BOTH	1. YES \rightarrow Q326 2. NO	3 YEARS	3. DID NOT WANT PREGNANCY THEN OR ANY TIME IN THE FUTURE
					888. DK 999. NR/REF	8. NOT SURE

VERIFY THAT SUM OF LIVE BIRTHS = Q311, IF NOT, CORRECT HISTORY OF LIVE BIRTH TABLE VERIFY THAT THE LAST ROW IN THE TABLE CORRESPONDS TO THE FIRST LIVE BIRTH ASK: "Was this your first child?"

- 327. Has your wife/partner ever had a spontaneous or induced abortion (therapeutic, induced, or planned) in the last five years?
 - SPONTANEOUS ABORTION, → CONTINUE WITH Q328
 INDUCED ABORTION, OR → GO TO Q329
 BOTH, SPONTANEOUS AND INDUCED ABORTION? → CONTINUE WITH Q328
 NONE (IN THE LAST 5 YEARS) → GO TO MODULI IV
- 328. How many spontaneous abortions in the last five years? _____ SPONTANEOUS ABORTIONS
- 329. How many induced, therapeutic or planned abortions in the last five years?

_ ___ INDUCED ABORTIONS

MODULE IV: FAMILY PLANNING KNOWLEDGE/ SEXUAL EXPERIENCE

For each of the following methods of preventing pregnancy, please tell me:

метнор	400. Have you ever heard of it?	401 . Do you know how to use it?	402 . Have you ever used it?	403. Do you know where to get it?	404. What was the most important source of information about this method (SEE CODES BELOW)
A. The Pill (Oral Contraceptives)	$\begin{array}{l} 1 \text{ YES} \rightarrow \text{Q401} \\ 2 \text{ NO-} \rightarrow \text{B} \end{array}$	$\begin{array}{l} 1 \text{ YES} \rightarrow \text{Q402} \\ 2 \text{ NO} \rightarrow \text{Q402} \end{array}$	$\begin{array}{l} 1 \text{ YES} \rightarrow \text{Q403} \\ 2 \text{ NO} \rightarrow \text{Q403} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$	
B. IUD (Spirali)	$1 \text{ YES} \rightarrow \text{Q401}$ $2 \text{ NO-} \rightarrow \text{C}$	$1 \text{ YES} \rightarrow \text{Q402}$ $2 \text{ NO} \rightarrow \text{Q402}$	$\begin{array}{l} 1 \text{ YES} \rightarrow \text{Q403} \\ 2 \text{ NO} \rightarrow \text{Q403} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$	
C. Condoms (Prezervativ)	$1 \text{ YES} \rightarrow \text{Q401}$ $2 \text{ NO-} \rightarrow \text{D}$	$1 \text{ YES} \rightarrow \text{Q402}$ $2 \text{ NO} \rightarrow \text{Q402}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q403} \\ 2 \text{ NO} \rightarrow \text{Q403} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$	
D. Foam/Jelly/ Cream/Foamy Tablets (Spermicides)	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q401} \\ 2 \text{ NO-} \rightarrow \text{E} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q402} \\ 2 \text{ NO} \rightarrow \text{Q402} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q403} \\ 2 \text{ NO} \rightarrow \text{Q403} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$	
E. Tubal Ligation	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q401} \\ 2 \text{ NO-} \rightarrow \text{F} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q402} \\ 2 \text{ NO} \rightarrow \text{Q402} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q403} \\ 2 \text{ NO} \rightarrow \text{Q403} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$	
F. Vasectomy (Male Sterilization)	$1 \text{ YES} \rightarrow \text{Q401}$ $2 \text{ NO-} \rightarrow \text{G}$	$1 \text{ YES} \rightarrow \text{Q402}$ $2 \text{ NO} \rightarrow \text{Q402}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q403} \\ 2 \text{ NO} \rightarrow \text{Q403} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$	
G. Injectables (e.g. Depo-Provera)	$\begin{array}{l} 1 \text{ YES} \rightarrow \text{Q401} \\ 2 \text{ NO-} \rightarrow \text{H} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q402} \\ 2 \text{ NO} \rightarrow \text{Q402} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q403} \\ 2 \text{ NO} \rightarrow \text{Q403} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$	
H. Emergency Hormonal Contraception ("Morning After Pill"; Postinor)	$1 \text{ YES} \rightarrow \text{Q401}$ $2 \text{ NO-} \rightarrow \text{I}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q402} \\ 2 \text{ NO} \rightarrow \text{Q402} \end{array}$	$1 \text{ YES} \rightarrow \text{Q403}$ $2 \text{ NO} \rightarrow \text{Q403}$	$1 \text{ YES} \rightarrow \text{Q404}$ $2 \text{ NO} \rightarrow \text{Q404}$	
I. Rhythm/Calendar Method	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q401} \\ 2 \text{ NO-} \rightarrow \text{J} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q402} \\ 2 \text{ NO} \rightarrow \text{Q402} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$		
J. Withdrawal (Coitus Interruptus)	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q401} \\ 2 \text{ NO-} \rightarrow \text{K} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q402} \\ 2 \text{ NO-} \rightarrow \text{Q402} \end{array}$	$\begin{array}{c} 1 \text{ YES} \rightarrow \text{Q404} \\ 2 \text{ NO} \rightarrow \text{Q404} \end{array}$		
K. Other contraceptive methods (SPECIFY):	$1 \text{ YES} \rightarrow \text{Q401}$ $2 \text{ NO-} \rightarrow \text{Q405}$	$1 \text{ YES} \rightarrow \text{Q402}$ $2 \text{ NO-} \rightarrow \text{Q402}$	$1 \text{ YES} \rightarrow \text{Q404}$ $2 \text{ NO} \rightarrow \text{Q404}$		

CODES FOR Q404 (DO NOT READ)

- 1. MOTHER
- 2. FATHER
- 3. RELATIVE
- 4. BOYFRIEND
- 5. FRIENDS
- 6. CO-WORKER
- 7. COLLEAGUES, PEER
- 8. PARTNER/HUSBAND
- 9. DOCTOR

- 10. NURSE, MIDWIFE, FELDCHER
- 11. COMMUNITY HEALTH WORKER
- 12. TEACHER
- 13. PHARMACIST
- 14. BOOKS
- 15 NEWSPAPERS, MAGAZINES, BROCHURE, FLYERS
- 16. RADIO
- 17. TV
- 20. OTHER (SPECIFY):
- 88. DON'T REMEMBER

405. Looking at this CARD, please tell me which do you think is the most effective contraceptive method? (SHOW CARD A)

- 1. The Pill
- 2. IUD
- 3. Condom
- 6. Foams/jelly/creams/Foamy Tablets
- 7. Tubal Ligation
- 8. Emergency Hormonal Contraception ("Morning After Pill")
- 9. Injectables (Depo-Provera)
- 10. Vasectomy (Male Sterilization)
- 11. Rhythm Method
- 12. Withdrawal
- 77. NONE OF THEM
- ') 88. DON'T KNOW/NOT SURE
- 406. How would you rate each of the following methods with regard to effectiveness at preventing pregnancy? Would you say that ______ is very effective, effective, somewhat effective, not very effective or not at all effective?

(INTERVIEWER: ASK THE QUESTION FOR EACH OF THE METHODS LISTED BELOW, UNLESS RESPON-DENT HAS SAID IN Q400 THAT HE NEVER HEARD OF THAT SPECIFIC METHOD; MARK "9" FOR THOSE CASES WITHOUT ASKING)

		Very <u>Effective</u>	Effective	Somewhat Effective	Not Very <u>Effective</u>	Do Not <u>Know</u>	Never <u>Heard</u>
1.	The pill	1	2	3	4	8	9
2.	IUD	1	2	3	4	8	9
3.	Condom	1	2	3	4	8	9
7.	Tubal ligation	1	2	3	4	8	9
9.	Injectables	1	2	3	4	8	9
11.	Calendar	1	2	3	4	8	9
12.	Withdrawal	1	2	3	4	8	9

BOX 4-I

414.

IF RESPONDENT IS 15-24 YEARS OF AGE CONTINUE; IF HE IS 25-49 YEARS GO TO Q420, Pg. 14

410. Now I have some questions about your first sexual intercourse. When did you have sexual intercourse for the first time - in what month and year was that? (PROBE: Can you tell me what year that was?)

MONTH	YEAR	00. NEVER HAD INTERCOURSE \rightarrow GO TO Q601, P26
		88. DON'T REMEMBER
		99. REFUSE TO ANSWER

411. How old were you at that time? _____ YEARS 88. DON'T REMEMBER

412. At the time you first had sexual intercourse, what was your relationship to that woman?

1. WIFE	8. RELATIVE
2. FIANCEE	9. RAPE \rightarrow GO TO Q421
3. GIRLFRIEND	10. INCEST \rightarrow GO TO Q421
4. FRIEND	11. PROSTITUTE- \rightarrow GO TO Q415
5. LOVER	20. OTHER (SPECIFY)
6. ACQUAINTANCE	88. DO NOT REMEMBER
7. JUST MET \rightarrow GO TO Q414	99. REFUSAL

413. How long were you and your first partner dating when you first had sexual relations?

 1. _____ DAYS OR 2. _____ WEEKS OR 3. _____ MONTHS OR 4. _ ____YEARS

 000=FIRST TIME WE MET

 888=DON'T REMEMBER

 999=NO RESPONSE/REFUSAL

 777=OTHER______

 How old was your first partner?

- 415. Before you had sex for the first time, did you and your partner ever talk about using contraception?
 - 1. YES
 - 2. NO
 - 8. DON'T REMEMBER
- 416. At the time you had first sexual intercourse, did you or your partner use any contraceptive method?

1 YES 2. NO \rightarrow GO TO Q419 8. DK/DO NOT REMEMBER \rightarrow GO TO Q421 9. REF \rightarrow **GO TO Q421**

- 417. Which contraceptive method did you or your partner use at the first intercourse?
 - 1. THE PILL
 - 2. IUD
 - 3. CONDOM
 - 6. FOAM/JELLY/CREAM/VAGINAL FILMS
 - 8. EMERGENCY HORMONAL CONTRACEPTION
 - 9. INJECTABLES
 - 10. OTHER MODERN METHODS
 - 11. CALENDAR METHOD
 - 12. WITHDRAWAL
 - 19. DOUCHE
 - 20. OTHER:
 - 88. DON'T KNOW/DON'T REMEMBER
- 418. Who made the decision to use contraception at that time? (READ 1-3)
 - 1. You
 - 2. Your partner
 - 3. Both you and your partner
 - 8. DON'T REMEMBER

GO TO Q421

- 419. What was the main reason for not using a contraceptive method at that time?
 - 1. SEX WAS NOT EXPECTED
 - 2. THOUGHT IT WAS A SAFE TIME OF THE MONTH
 - 3. DID NOT KNOW WHERE TO GET A METHOD//DIFFICULT TO GET/NOT AVAILABLE
 - 4. RESPONDENT WAS AGAINST IT
 - 5. PARTNER WAS AGAINST IT
 - 6. DID NOT KNOW ABOUT CONTRACEPTION
 - 7. WANTED TO GET PREGNANT
 - 8. THOUGHT THAT CONTRACEPTIVE METHODS ARE HARMFUL
 - 9. DID NOT THINK ABOUT USING A METHOD/NEGLIGENCE
 - 10. RESPONDENT AFRAID OF PARTNER'S REACTION
 - 11. TOO DRUNK (PARTNER OR RESPONDENT)
 - 12. RESPONDENT WAS TOO EMBARRASSED TO USE A METHOD
 - 20. OTHER (SPECIFY)
 - 88. DON'T REMEMBER/DON'T KNOW

GO TO Q421

420. How old were you at the time of your first sexual intercourse?

YEARS

00. NEVER HAD INTERCOURSE-- \rightarrow GO TO Q601, PG26 88. DK/DR

- 421. During the past 30 days (past month) have you had sexual intercourse?
 - 1. YES
 - 2. NO \rightarrow **GO TO Q436, PG.17** 9. REF \rightarrow GO TO Q436, PG.17

422. How many times have you had sexual intercourse during the past 30 days (READ 1-5)?

- 1. Every day,
- 2. 3-5 times per week,
- 3. 1-2 times per week,
- 4. 2-3 times per month, or
- 5. Only once
- 9. REFUSED

423. At the time of your last sexual intercourse, what was your relationship with that woman?

1. WIFE \rightarrow GO TO Q425	7. JUST MET \rightarrow GO TO Q425
2. FIANCEE	8. RELATIVE \rightarrow GO TO Q425
3. GIRLFRIEND	11. PROSTITUTE \rightarrow GO TO Q425
4. FRIEND	20. OTHER (SPECIFY)
5. LOVER	88. DO NOT REMEMBER
6. ACQUAINTANCE	99. REFUSAL

424. How long were you and your last partner dating when you last had sexual relations?

1DAYS OR 2	WEEKS OR	3	MONTHS OR	4	YEARS
000=FIRST TIME WE MET					
888=DON'T REMEMBER					
999=NO RESPONSE					
777=OTHER					
At the time or your last sexual	intercourse, did you or	your partner u	ise any contracept	ive method?	

1. YES 2. NO \rightarrow **GO TO Q428** 9. DR/ REF \rightarrow **GO TO Q428**

- 426. Which method did you use?
 - 1. THE PILL
 - 2. IUD

425.

- 3. CONDOM \rightarrow GO TO Q431
- 4. SPERMICIDE (FOAM/JELLY/CR1EAM/VAGINAL FILMS)
- 5. CONDOM+SPERMICIDE \rightarrow GO TO Q431
- 6. CONDOM +CALENDAR / WITHDRAWAL \rightarrow GO TO Q431
- 7. TUBAL LIGATION
- 8. EMERGENCY HORMONAL CONTRACEPTION / MORNING AFTER PILL
- 9. INJECTABLES (EXAMPLE: DEPO-PROVERA)
- 10. OTHER MODERN METHOD (SPECIFY)
- 11. CALENDAR METHOD
- 12. WITHDRAWAL \rightarrow GO TO Q427A
- 13. WITHDRAWAL+CALENDAR \rightarrow GO TO Q427A
- 20. OTHER NATURAL METHOD (SPECIFY)
- 88. DON'T KNOW/DON'T REMEMBER

427. At the time or your last sexual intercourse, did you or your partner use a condom in addition to your current method?

- 1. YES \rightarrow GO TO BOX 4-1
- 2. NO

9. DON'T REMEMBER / REFUSE

- 427A. During the last 30 days did you use a condom?
 - 1. YES
 - 2. NO

9. DOES NOT REMEMBER / REFUSAL

BOX 4-1

IF Q426=1, 2, 7, 9, 10, 11 GO TO Q432

- 427B. During the last 30 days did you or your partner use that current method: (READ 1-4)
 - 1. At every sexual intercourse,
 - 2. At almost all acts of sexual intercourse,
 - 3. From time to time, or
 - 4. Only once?
 - 9. REFUSAL

GO TO Q432

- 428. In the last 30 days, did you or your partner use any contraceptive method to avoid pregnancy?
 - 1. YES

2. NO \rightarrow **GO TO Q445, PAGE 18**

- 429. Which contraceptive method did you use?
 - 1. THE PILL
 - 2. IUD
 - 3. CONDOM \rightarrow GO TO Q431
 - 4. SPERMICIDE (FOAM/JELLY/CR1EAM/VAGINAL FILMS)
 - 5. CONDOM+SPERMICIDE \rightarrow GO TO Q431
 - 6. CONDOM +CALENDAR / WITHDRAWAL \rightarrow GO TO Q431
 - 7. TUBAL LIGATION
 - 8. EMERGENCY HORMONAL CONTRACEPTION / MORNING AFTER PILL
 - 9. INJECTABLES (EXAMPLE: DEPO-PROVERA)
 - 10. OTHER MODERN METHOD (SPECIFY)
 - 11. CALENDAR METHOD
 - 12. WITHDRAWAL
 - 13. WITHDRAWAL+CALENDAR
 - 20. OTHER NATURAL METHOD (SPECIFY)
 - 88. DON'T KNOW
- 430. In the last 30 days, did you or your partner use a condom?
 - 1. YES
 - 2. NO
 - 9. REFUSAL

IF Q429=1, 2, 7, 9, 10, 11, GO TO Q432

431. During the last 30 days did you or your partner use that current method: (READ OPTIONS 1-4)

- 1. At every sexual intercourse?
- 2. At almost all acts of sexual intercourse, or
- 3. From time to time, or
- 4. Only once?
- 9. REFUSAL

432. Why did you choose that current method? (REFERS TO METHOD CITED IN Q426 OR Q429)

- 1. DOCTOR RECOMMENDED
- 2. COST
- 3. VERY EFFECTIVE
- 4. NO SIDED EFFECTS (FEW SIDE EFFECTS)
- 5. SAW ADS (TV, RADIO, PRESS, BROCHURES)
- 6. EASY TO USE
- 7. PARTNER PREFERS IT
- 8. KNOWS SOMEBODY WHO USES IT
- 9. CURIOSITY / WANTED TO TRY IT
- 10. ALLOWS SPONTANEITY DURING INTERCOURSE
- 11. TRADITIONAL METHOD / CULTURALLY ACCEPTABLE
- 20. OTHER___
- 88 DON'T KNOW

433.	ou or your partner used. Where did you or time you had sex?					
	1. HEALTH POST		9. OPEN MARKET			
	2. HEALTH CENTER		DRE/ KIOSK			
	3. POLICLINIC		TNER/WIFE			
	4. FAMILY PLANNING CLINIC	12. FRI				
	5. GOV HOSPITAL-MATERNITY WARD	13. REL				
	6. PRIVATE CLINIC OR OFFICE		IER (SPECIFY):			
	7. NGO		FURAL METHOD \rightarrow GO TO Q445			
	8. PHARMACY	88. DOI	N'T KNOW			
434.	Do (Did) you pay for this method?					
	1. YES					
	2. NO \rightarrow GO TO Q445 3. PARTNER GETS THE METHOD \rightarrow GO TO (Q445				
435.	On average, how much did you pay for the contract	ceptive metho	od, per month?			
	HUNDRED LEKI		85 HUNDRED LEKI OR MORE NOT SURE/DON'T KNOW			
	G	O TO Q445				
436.	During the past 3 months, have you had sexual int	ercourse?				
	1. YES					
	2. NO → GO TO Q453, PAGE 19 9. REF → GO TO Q453, PAGE 19					
437.	At the time of your last sexual intercourse, what w					
	1. WIFE \rightarrow GO TO Q439		T MET → GO TO Q439			
	2. FIANCEE		$ATIVE \rightarrow GO TO Q439$			
	3. GIRLFRIEND		$OSTITUTE \rightarrow GO TO Q439$			
	4. FRIEND 5. LOVER	20. 01F	HER (SPECIFY) NOT REMEMBER			
	6. ACQUAINTANCE	99. REF				
120						
438.	How long were you and your last partner dating w 1DAYS OR 2WEEKS	-				
	000=FIRST TIME WE MET	on n				
	888=DON'T REMEMBER					
	999=NO RESPONSE					
	777=OTHER					
439.	At your last sexual intercourse, did you or your pa	rtner use any	v contraceptive method to avoid pregnancy?			
	1. YES					
	2. NO \rightarrow GO TO Q445					
	8. DON'T REMEMBER \rightarrow GO TO Q445 9. REFUSED \rightarrow GO TO Q445					
440.	Which contraceptive method did you use?					
	1. THE PILL		9. INJECTABLES (EXAMPLE: DEPO-PROVERA)			
	2. IUD		10. OTHER MODERN METHOD (SPECIFY)			
	 CONDOM → GO TO Q441A SPERMICIDE (FOAM/JELLY/CR1EAM/VAGINA 	I FILMS)	11. CALENDAR METHOD 12. WITHDRAWAL \rightarrow GO TO Q445			
	4. SPERMICIDE (FOAM/JELLY/CRTEAM/VAGINA 5. CONDOM+SPERMICIDE \rightarrow GO TO Q441A	L I ILIVIS)	12. WITHDRAWAL \rightarrow GO TO Q445 13. WITHDRAWAL+CALENDAR \rightarrow GO TO Q445			
	6. CONDOM +CALENDAR / WITHDRAWAL \rightarrow GO	O TO Q441A				
	7. TUBAL LIGATION		88. DON'T KNOW			
	8. EMERGENCY HORMONAL CONTRACEPTION AFTER PILL	/ MORNING				
	AFTERTILL					

- 441. At your last sexual intercourse, did you or your partner use a condom in addition to your current method?
 - 1. YES 2. NO \rightarrow **GO TO Q442** 8. DON'T REMEMBER \rightarrow GO TO Q442 9. REFUSED \rightarrow GO TO Q442
- 441A. During the last 3 months how often did you use a condom when you had sexual relations with that woman? Did you use a condom: (READ OPTIONS 1-3)
 - 1. At every sexual intercourse? 2. At almost all acts of sexual intercourse, or 3. From time to time? 8. DON'T KNOW 9. REFUSAL
- 442. Where did you or your partner get the method of contraception you used the last time you had sex?

1. HEALTH POST	9. OPEN MARKET
2. HEALTH CENTER	10. STORE/ KIOSK
3. POLICLINIC	11. PARTNER/WIFE
4. FAMILY PLANNING CLINIC	12. FRIEND
5. GOV HOSPITAL-MATERNITY WARD	13. RELATIVE
6. PRIVATE CLINIC OR OFFICE	20. OTHER (SPECIFY):
7. NGO	77. NATURAL METHOD \rightarrow GO TO Q445
8. PHARMACY	88. DON'T KNOW
Do (Did) you pay for this method?	
1 YES	
2 NO \rightarrow GO TO Q445	

444. On average, how much did you pay for the contraceptive method, per month?

3 PARTNER GETS THE METHOD \rightarrow GO TO Q445

 HUNDRED	LEKI

85. 85 HUNDRED LEKI OR MORE 88. NOT SURE/DON'T KNOW

445. During the past 3 months, with how many different women have you had intercourse?

PARTNERS

443.

447.

85. 85+ WOMEN 88. DON'T KNOW / NOT SURE 99. NO RESPONSE

IF Q445>1 CONTINUE; IF Q445=1, 88 OR 99 GO TO Q460, PAGE 20

446. You told me that in the last 3 months you had sexual relations with more than 1 woman. What was the relationship with the next-to-last woman when you last had sexual intercourse with her?

1. WIFE → GO TO Q449	7. JUST MET → GO TO 0449
2. FIANCEE	8. RELATIVE \rightarrow GO TO Q449
3. GIRLFRIEND	11. PROSTITUTE \rightarrow GO TO Q449
	20. OTHER (SPECIFY)
	88. DO NOT REMEMBER
6. ACQUAINTANCE	99. REFUSAL
How long had you and that partner been together whe	en you last had sexual relations?
1DAYS OR 2WEEKS OR	. 3 MONTHS OR 4 YEARS
000=FIRST TIME WE MET	

888=DON'T REMEMBER 999=NO RESPONSE 777=OTHER

- 449. Did you and that next-to-last partner use any contraceptive method to avoid pregnancy the last time you had sexual relations?
 - 1. YES
 - 2. NO \rightarrow **GO TO Q460**
 - 9. DON'T REMEMBER / REFUSE \rightarrow GO TO Q460
- 450. Which contraceptive method did you use?
 - 1. THE PILL
 - 2. IUD
 - 3. CONDOM \rightarrow GO TO Q452
 - 4. SPERMICIDE (FOAM/JELLY/CR1EAM/VAGINAL FILMS)
 - 5. CONDOM+SPERMICIDE \rightarrow GO TO Q452
 - 6. CONDOM +CALENDAR / WITHDRAWAL \rightarrow GO TO Q452
 - 7. TUBAL LIGATION
 - 8. EMERGENCY HORMONAL CONTRACEPTION / MORNING AFTER PILL
 - 9. INJECTABLES (EXAMPLE: DEPO-PROVERA)
 - 10. OTHER MODERN METHOD (SPECIFY)
 - 11. CALENDAR METHOD
 - 12. WITHDRAWAL \rightarrow **GO TO Q460**
 - 13. WITHDRAWAL+CALENDAR \rightarrow GO TO Q460
 - 20. OTHER NATURAL METHOD (SPECIFY)
 - 88. DON'T KNOW
- 451. At your last sexual intercourse with this (next-to-last) partner, did you or your partner use a condom in addition to your current method?

1. YES 2. NO \rightarrow **GO TO Q460** 8. DON'T REMEMBER \rightarrow **GO TO Q460** 9. REFUSE \rightarrow **GO TO Q460**

452. During the last 3 months how often did you use a condom when you had sexual relations with that (next-to-last) woman? Did you use a condom: (READ OPTIONS 1-4)

1. At every sexual intercourse?

- 2. At almost all acts of sexual intercourse,
- 3. From time to time, or
- 4. Only once?
- 9. DON'T KNOW / REFUSAL

GO TO Q460

453. When was the last time you had sexual intercourse?

____MONTH _____YEAR 88. DK 99. NR

454. At the time of your last intercourse, what was your relationship with your partner?

1. WIFE	8. RELATIVE
2. FIANCEE	9. RAPE \rightarrow GO TO Q460
3. GIRLFRIEND	10. INCEST \rightarrow GO TO Q460
4. FRIEND	11. PROSTITUTE
5. LOVER	20. OTHER (SPECIFY)
6. ACQUAINTANCE	88. DO NOT REMEMBER
7. JUST MET	99. REFUSAL

455. At the time you had your last sexual intercourse, did you or your partner use any contraceptive method?

- 1. YES
- 2. NO \rightarrow GO TO Q460
- 9. DK/REF \rightarrow GO TO Q460

456.	Which contraceptive method did you or your partner use at the last intercourse?			
	 THE PILL IUD CONDOM CONDOM +SPERMICIDE CONDOM + WITHDRAWAL/CALENDAR FOAM/JELLY/CREAM/VAGINAL FILMS/DIAFRAGM TUBAL LIGATION EMERGENCY HORMONAL CONTRACEPTION 	 9. INJECTABLES (EXAMPLE: DEPO-PROVERA) 10. OTHER MODERN METHOD (SPECIFY)		
457.	Where did you or your partner get the method of cont	raception you used the last time?		
	 HEALTH POST HEALTH CENTER POLICLINIC FAMILY PLANNING CLINIC GOV HOSPITAL-MATERNITY WARD PRIVATE CLINIC OR OFFICE NGO PHARMACY 	 9. OPEN MARKET 10. STORE/ KIOSK 11. PARTNER/WIFE 12. FRIEND 13. RELATIVE 20. OTHER (SPECIFY):		
458.	 Do (Did) you pay for this method? 1 YES 2 NO → GO TO Q460 			
	3 PARTNER GETS THE METHOD \rightarrow GO TO Q46	0		
459.	On average, how much did you pay for the contracept	ive method, per month?		
	HUNDRED LEKI	85. 85 HUNDRED LEKI OR MORE88. NOT SURE/DON'T KNOW		
460.	How many women have you had sexual intercourse with in the last 12 months? (even those you had intercourse with only once).			
	85. 85 OR MO WOMEN 88. DON'T KN 99. NO RESPO	NOW / DON'T REMEMBER		
460A.	Counting all the women you had sexual intercourse w women have you had sexual intercourse with in your	ith, even those you had intercourse with only once, how many ife?		
	85. 85 OR MO WOMEN 88. DON'T KN 99. NO RESPO	NOW / DON'T REMEMBER		
461.	Have you ever had sexual relations with a prostitute?			
	 YES NO → GO TO MODULE V DON'T REMEMBER / REFUSE → GO TO MOD 	ULE V		
462.	When was the last time you had sexual relations with a prostitute? Did you have relations: (READ 1-4)?			
	 In the last month? 2-3 months ago In the last year, but not in the last month, or More than a year ago DON'T REMEMBER / REFUSE 			
463.	Did you use a condom the last time you had sexual relations with a prostitute?			
	1. YES 2. NO			

9 DON'T REMEMBER / REFUSE

V. CURRENT AND PAST CONTRACEPTIVE USE

503A. VERIFY WHETHER RESPONDENT REPORTED CURRENTLY (LAST 30 DAYS) USING ANY METHOD (SEE Q425 AND Q428: PAGE 15 AND 16). CIRCLE "1" OR "2".

1. DID NOT USE A METHOD IN THE PAST 30 DAYS (Q425=2 AND Q428=2 OR BLANK) CONTINUE

2. USED A METHOD IN THE LAST 30 DAYS (Q425=1 or Q428=1 \rightarrow GO to Box 5-1

- 503. So, you said that you or any of your partners are not currently (in the last 30 days) using any method to prevent pregnancy?
 - 1. CURRENTLY USING \rightarrow **CORRECT Q425 OR Q428, AND THEN CONTINUE** 2. NOT CURRENTLY USING \rightarrow **GO TO Q515**

BOX 5-I

IFCURRENTLY USES TRADITIONAL METHOD (SEE Q426 AND Q429) CONTINUE, ELSE GO TO Q510

508. Please tell me whether each of the following reasons was very important, somewhat important, or not important at all in your decision to use ______ (WRITE NAME OF TRADITIONAL METHOD) instead of a modern method:

	VERY <u>IMPORTANT</u>	SOMEWHAT <u>IMPORTANT</u>	NOT <u>IMPORTANT</u>	DON'T <u>KNOW</u>
A. Difficult to get a modern method	1	2	3	8
B. Cost of these modern methods	1	2	3	8
C. Little knowledge of modern methods	1	2	3	8
D. Fear of or experience with side effects	1	2	3	8
E. Husband/Partner choice	1	2	3	8
F. Religious beliefs	1	2	3	8
G. Doctor's recommendation	1	2	3	8
H. Another person's advice	1	2	3	8

- 509. How effective at preventing pregnancy do you think (**TRADITIONAL METHOD NAMED**) is compared to modern methods, like the pill or the IUD? (**READ 1-3**)
 - 1. Current method more effective
 - 2. About equally effective
 - 3. Current method less effective
 - 8. DON'T KNOW/NOT SURE

510. Do you have any problems or concerns with using your current method?

1. YES

2. NO \rightarrow GO TO Q512

- 511. What is the most important problem?
 - 1. SIDE EFFECTS
 - 2. HEALTH CONCERNS
 - 3. ACCESS/AVAILABILITY
 - 4. COST
 - 5. SOMETIMES FORGET TO USE
 - 6. SOMETIMES DIFFICULT/INCONVENIENT TO USE
 - 7. WIFE / PARTNER DISAPPROVES
 - 8. LESS EFFECTIVE METHOD/GOT PREGNANT WHILE USING IT
 - 9. DEEPLY UNSATISFIED WITH THE METHOD
 - 10. OTHER_____

- 512. Would you prefer to use a different method of family planning from the one you are currently using?
 - 1. YES
 - 2. NO \rightarrow GO TO 521
- 513. What method would you prefer to use (OTHER THAN THE METHOD ALREADY SPECIFIED)?
 - 1. THE PILL
 - 2. IUD
 - 3. CONDOM
 - 4. CONDOM +SPERMICIDE
 - 5. CONDOM +WITHDRAWAL/CALENDAR-
 - 6. FOAM/JELLY/CREAMS/C-FILMS
 - 7. FEMALE STERILIZATION
 - 8. EMERGENCY HORMONAL CONTRACEPTION
 - 9. INJECTABLES (DEPO PROVERA)
 - 10. OTHER MODERN METHODS_____
 - 11. CALENDAR
 - 12. WITHDRAWAL
 - 13. WITHDRAWAL AND CALENDAR
 - 20. OTHER NATURAL METHODS_
 - 88. DO NOT KNOW/NOT SURE
- 514. What is the most important reason that you do not use that method?
 - 1. DOCTOR WILL NOT PRESCRIBE IT
 - 2. COST
 - 3. NOT AVAILABLE/UNRELIABLE SUPPLIES/DIFFICULT TO OBTAIN
 - 4. TOO FAR AWAY
 - 5. DO NOT KNOW HOW/WHERE TO OBTAIN IT
 - 6. WIFE/PARTNER OBJECTS TO IT
 - 7. RELIGIOUS REASONS
 - 8. FEAR OF SIDE EFFECTS
 - 9. HAS NOT YET MADE UP HIS/THEIR MIND
 - 10. DIFFICULT TO USE
 - 11. FEAR OF SURGICAL PROCEDURE (IUD, TL, NORPLANT)
 - 20. OTHER___
 - 88. DON'T KNOW

BOX 5-II

GO TO Q 521

- 515. What is the main reason that you or your partner are not currently using a contraceptive method?
 - 1. DOES NOT CURRENTLY HAVE A PARTNER/ NOT SEXUALLY ACTIVE IN THE LAST MONTH
 - 2. TRYING TO GET PREGNANT
 - 3. PARTNER/WIFE POSTPARTUM/ BREASTFEEDING
 - 4. PARTNER/WIFE CURRENTLY PREGNANT
 - 5. PARTNER/WIFE HYSTERECTOMY/MENOPAUSE \rightarrow GO TO Q523
 - 6. DOCTOR SAID HE OR HIS WIFE/PARTNER CANNOT HAVE CHILDREN \rightarrow GO TO Q523
 - 7. COUPLE TRIED TO GET PREGNANT FOR AT LEAST 2 YEARS AND DIDN'T SUCCEED \rightarrow GO TO Q523
 - 8. FEAR OF SIDE EFFECTS
 - 9. LOVEMAKING WOULD BE INTERRUPTED
 - 10. RESPONDENT DID NOT THINK ABOUT USING CONTRACEPTION
 - 11. COST, CANNOT AFFORD BIRTH CONTROL
 - 12. BIRTH CONTROL IS THE PARTNER'S RESPONSIBILITY
 - 13. BIRTH CONTROL IS NOT (VERY) EFFECTIVE
 - 14. RESPONDENT DOES NOT WANT TO USE A METHOD
 - 15. PARTNER OBJECTS TO USING METHOD
 - 16. OBJECTS DUE TO RELIGIOUS REASONS
 - 17. DOES NOT KNOW WHERE TO GET METHOD
 - 18. RESPONDENT DOES NOT KNOW HOW TO USE BIRTH CONTROL METHODS
 - 19. RESPONDENT DOES NOT THINK THAT HE AND PARTNER CAN GET PREGNANT
 - 20. RESPONDENT HAS NO TIME TO GO TO A FP CLINIC/ OR BUY CONDOMS
 - 21. RESPONDENT'S PARTNER USES DOUCHING
 - 77. OTHER (SPECIFY)
 - 88. DON'T KNOW

516. Do you think that you will use a contraceptive method during the next 12 months (ADD:OTHER THAN DOUCHING IF Q515=21)? (YOU OR YOUR PARTNER)

1. YES \rightarrow **GO TO Q518** 2. NO 8. NOT SURE

- 517. Do you think that you will use a contraceptive method any time in the future?
 - 1. YES

2. NO \rightarrow GO TO Q521

8. NOT SURE \rightarrow GO TO Q521

- 518. What method would you want to use most?
 - 1. THE PILL
 - 2. IUD
 - 3. CONDOM
 - 4. CONDOM +SPERMICIDE
 - 5. CONDOM +WITHDRAWAL/CALENDAR-
 - 6. FOAM/JELLY/CREAMS/C-FILMS
 - 7. FEMALE STERILIZATION
 - 8. EMERGENCY HORMONAL CONTRACEPTION
 - 9. INJECTABLES(DEPO PROVERA)
 - 10. OTHER MODERN METHODS
 - 11. CALENDAR \rightarrow GO TO Q521
 - 12. WITHDRAWAL \rightarrow GO TO Q521
 - 13. WITHDRAWAL + CALENDAR \rightarrow GO TO Q521
 - 20. OTHER \rightarrow GO TO Q521
 - 88. NOT SURE \rightarrow GO TO Q521

519. On average, how much are you willing to pay for contraception, per month? (YOU OR YOUR PARTNER)

_____ HUNDRED LEKI

85. 85 HUNDRED LEKI OR MORE88. NOT SURE/DON'T KNOW

520. Where would you want to get your contraceptive method? (YOU OR YOUR PARTNER)

- 1. HEALTH POST
- 2. HEALTH CENTER
- 3. POLICLINIC
- 4. FAMILY PLANNING CLINIC
- 5. GOV HOSPITAL-MATERNITY WARD
- 6. PRIVATE CLINIC OR OFFICE
- 7. NGO

88. NOI SUKE/DON I KNOW

8. PHARMACY
 9. OPEN MARKET
 10. STORE/ KIOSK
 11. PARTNER/WIFE
 12. FRIEND
 13. RELATIVE
 20. OTHER (SPECIFY): ______
 88. DON'T KNOW

521. During the last year, how often did you talk about contraception with your wife / partner?

- 1. NEVER \rightarrow GO TO Q523
- 2. ONE OR TWO TIMES
- 3. THREE TIMES OR MORE
- 4. RESPONDENT HAD NO PARTNER DURING THE LAST YEAR \rightarrow GO TO Q523
- 522. Generally, does your wife / partner agree or disagree with the use of contraceptive methods?
 - 1. AGREES
 - 2. DISAGREES
 - 3. NEITHER AGREES NOR DISAGREES
 - 8. NOT SURE/DON'T KNOW

- 523. Some people use condoms for reasons other than birth control, for instance because they are concerned about getting diseases that can result from sexual intercourse. Have you ever used condoms for: (**READ 1-4**)
 - 1. Birth Control Only \rightarrow **GO TO BOX 5-III**
 - 2. Disease Prevention Only \rightarrow **GO TO BOX 5-III**
 - 3. Both, or \rightarrow **GO TO BOX 5-III**
 - 4. You Never Used a Condom?
 - 5. USED CONDOM OUT OF CURIOSITY
 - 8. NOT SURE/ DO NOT REMEMBER
- 524. Why have you and your partner(s) never used condoms?
 - 1. PREVENTING PREGNANCY IS WOMAN'S RESPONSIBILITY
 - 2. PARTNER(S) OBJECTED TO USE CONDOMS
 - 3. HAVE ONLY ONE PARTNER
 - 4. THEY ARE ONLY FOR USE WITH PROSTITUTES
 - 5. THEY ARE ONLY FOR EXTRAMARITAL RELATIONS
 - 6. CONDOMS DIMINISH PLEASURE/SPONTANEITY
 - 7. CONDOMS ARE LESS EFFECTIVE IN PREVENTING PREGNANCY
 - 8. CONDOMS ARE TOO DIFFICULT TO USE
 - 9. LOVEMAKING WOULD BE INTERRUPTED
 - 10. CONDOM USE IS TOO MESSY
 - 11. COST
 - 12. HE HAS NEVER THOUGHT ABOUT IT
 - 13. IT IS EMBARRASSING TO BUY CONDOMS
 - 14. PREFERS OTHER CONTRACEPTIVE METHODS
 - 20. OTHER
 - 88. DON'T KNOW

BOX 5-III

VERIFY Q425 AND 428 (PAGES 15 & 16); IF RESPONDENT CURRENTLY USING A METHOD GO TO Q553

- 551. Do you think that you can get your partner pregnant now?
 - 1. YES
 - 2. NO
 - 3. NOT SURE
 - 4. PARTNER IS PREGNANT NOW
- 552. What is the reason why you think you cannot get your partner pregnant now?
 - 1. DOES NOT CURRENTLY HAVE A PARTNER / NOT SEXUALLY ACTIVE
 - 2. PARTNER IS POSTPARTUM / BREASTFEEDING
 - 3. GENITAL INFECTION (ANEXITA, ENDOMETRITIS)
 - 4. SYSTEMATIC OR ENDOCRINAL DISEASE
 - 5. HYSTERECTOMY (OPERATION TO REMOVE THE UTERUS) \rightarrow GO TO MODULE VI
 - 6. MENOPAUSAL \rightarrow GO TO MODULE VI
 - 7. OVARIAN CYST OR DYSFUNCTION \rightarrow **GO TO MODULE VI**
 - 8. OBSTRUCTED OR MISSING FALLOPIAN TUBES → GO TO MODULE VI
 - 9. NO PREGNANCY FOR 2 YEARS DESPITE NO CONTRACEPTION → GO TO MODULE VI
 - 10. RESPONDENT HAD OPERATION SO CANNOT HAVE KIDS \rightarrow **GO TO MODULE VI**
 - 11. RESPONDENT STERILE → GO TO MODULE VI
 - 12. USING A CONTRACEPTIVE METHOD → VERIFY Q425 & Q428 (P. 15-16), CORRECT IF NECESSARY
 - 20. OTHER (SPECIFY)_
 - 88. DON'T KNOW

553. Looking to the future, do you yourself intend to have (a/another) baby at some time (IF CURRENTLY PREGNANT ADD "...after this pregnancy"?)

WANTS A BABY
 DOES NOT WANT A BABY → GO TO Q555
 RESPONDENT WANTS A BABY BUT PARTNER DISAGREES
 RESPONDENT DOES NOT WANT A BABY BUT PARTNER WANTS → GO TO Q555
 DK → GO TO Q555

554. When do you, yourself, actually want to get your wife/partner pregnant (again)...(READ 1-4)

Right away, (DO NOT READ IF THE HIS WIFE / PARTNER IS ALREADY PREGNANT)
 Within the next 12 months,
 Within 1-2 years,
 or after 2 years?
 AFTER HE MARRIES
 WHEN GOD WANTS
 DON'T KNOW

- 555. IF Q553 =1, 3, OR 8 BEGIN WITH: "After having all the children you want,...") Do you think you would be interested in your partner having an operation (tubal ligation) to prevent having any more children?
 - 1. YES \rightarrow GO TO MODULI VI

2. NO

- 3. WIFE / PARTNER IS ALREADY STERILIZED \rightarrow GO TO MODULI VI
- 4. HE HAS HAD A VASECTOMY \rightarrow **GO TO MODULI VI**
- 8. NOT SURE
- 556. What is the most important reason you wouldn't be interested in her having such a procedure?
 - 1. HEALTH RISKS/FEAR OF SIDE EFFECTS
 - 2. FEAR OF OPERATION
 - 3. DOESN'T KNOW ENOUGH ABOUT /NEVER HEARD OF STERILIZATION
 - 4. MIGHT WANT ANOTHER CHILD

5. COST

- 6. DOES NOT HAVE A PARTNER/NOT SEXUALLY ACTIVE
- 7. AGE TOO YOUNG OR TOO OLD (APPROACHING MENOPAUSE)
- 8. HAVEN'T THOUGHT ABOUT IT
- 9. NOT CULTURALLY ACCEPTABLE
- 10. RELIGIOUS REASONS
- 11. PREFERS (OR USES) OTHER CONTRACEPTIVE METHODS
- 12. CANNOT GET PREGNANT (INFERTILITY, MEDICAL REASONS)
- 20. OTHER___
- 88. DON'T KNOW

VI. MEN'S HEALTH

Now I would like to ask you some questions about your health.

601. In the past 12 months, have you visited any health facility for care for yourself, including obtaining preventive services, such as family planning counseling or health check-ups?

1. YES

- 2. NO \rightarrow GO TO Q604
- 8. DK/DO NOT REMEMBER \rightarrow GO TO Q604
- 603. During your visit in the past 12 months at the health facility, did a doctor or medical provider talk to you about

			DON'T KNOW
	<u>YES</u>	<u>NO</u>	DON'T REM.
A. Condoms	1	2	8
B. STDs (Sexually Transmitted Diseases)	1	2	8
C. Control for pregnancy	1	2	8

604. Many different factors can prevent us from getting medical advice or treatment. When you want to get medical advice or treatment, are any of the following major problems or not (READ A -G)?

	BIG <u>PROBLEM</u>	NOT A BIG <u>PROBLEM</u>
A. Knowing where to go	1	2
C. Getting money needed for treatment	1	2
D. The distance to the health facility	1	2
E. Having to take transport	1	2
G. Concern that there may not be a male health provider	1	2

- 614. Have you ever tried cigarette smoking, even one or two puffs?
 - 1. YES 2. NO \rightarrow GO TO 621
- 615. How old were you when you smoked a cigarette for the first time?

YEARS 88. DK 99.NR

616. Have you smoked at least 100 cigarettes in your entire life? (PROBE: 100 cigarettes is about 5 packs)

1. YES 2. NO \rightarrow GO TO 621 8. DK \rightarrow GO TO 621 9. REFUSAL \rightarrow GO TO 621

617. How old were you when you first started smoking fairly regularly?

YEARS	00. NEVER SMOKED REGULARLY
	88. DO NOT REMEMBER
	99. REFUSE

- 618. During the last 30 days, did you smoke cigarettes: (READ 1-4)
 - 1. Every Day 2. Almost Every Day 3. Some Days 4. Not at All in the last 30 days \rightarrow GO TO Q620
 - 9. REFUSAL \rightarrow GO TO Q620

619. During the last 30 days, on the days you smoked, how many cigarettes did you smoke per day?

- 1. 1 CIGARETTE PER DAY
- 2. 2-5 CIGARETTES PER DAY
- 3. 6-10 CIGARETTES PER DAY
- 4. 11-19 CIGARETTES PER DAY
- 5. 20 OR MORE CIGARETTES PER DAY

GO TO Q621

620. In what month and year did you last smoke cigarettes at all? (PROBE FOR SEASON IF MONTH IS UNKNOWN)

	MONTH

_____YEAR 88. DK 99. REF

- 621. Now, I will ask you about some medical conditions that you may have had. Has a doctor or other medical care provider ever told you that you had Diabetes or "high sugar"?
 - 1. YES
 - 2. NO
 - 8. NOT SURE
 - 9. REFUSAL
- 622. Has a doctor or other medical care provider ever told you that you had heart disease?
 - 1. YES
 - 2. NO
 - 8. NOT SURE
 - 9. REFUSAL
- 625. Has a doctor or other medical care provider ever told you that you had Hypertension or High Blood Pressure?
 - 1. YES
 - 2. NO
 - 8. NOT SURE
 - 9. REFUSAL
- 626. Has a doctor or other medical care provider ever told you that you had Hepatitis B?
 - 1. YES
 - 2. NO
 - 8. NOT SURE
 - 9. REFUSAL

627. In the past 12 months have you had any of the following symptoms?

	<u>YES</u>	<u>NO</u>	<u>DON'T KNOW</u>
A. Discharge from the penis	1	2	8
B. Itching or burning in the genital area	1	2	8
C. Burning pain on urination	1	2	8
D. Pain during sexual intercourse	1	2	8
E. Sore, ulcer or warts in genital area	1	2	8
F. Swelling in genital area	1	2	8

IF ALL Q627 (A-F) > 1 (NO SYMPTOMS IN THE PAST 12 MONTHS) GO TO Q631; ELSE CONTINUE

- 628. Did you have any treatment for this(ese) condition(s)?
 - 1. YES
 - 2. NO \rightarrow GO TO Q630
 - 8. NOT SURE \rightarrow **GO TO Q630**

- 629. Who treated you?
 - 1. OB/GYN
 - 2. FAMILY DOCTOR/GENRAL PRACTIONER
 - 3. DERMATOLOGIST
 - 4. INFECTIOUS DISEASES DOCTOR
 - 5. NURSE/MIDWIFE
 - 6. PHARMACIST

- PARTNER
 FRIEND/RELATIVE
 SELF-TREATMENT
 OTHER:
 REFUSE TO ANSWER
- GO TO Q631
- 630. What was the main reason you did not seek treatment?
 - 1. SERVICES TOO FAR AWAY / INACCESSIBLE
 - 2. DON'T KNOW WHERE TO GO FOR SERVICES
 - 3. CANNOT AFFORD SERVICES OR TREATMENT
 - 4. AFRAID OF KNOWING THE RESULTS
 - 5. IT IS EMBARRASSING
 - 6. DID NOT THINK WAS A STD
 - 7. SYMPTOM(S) DISAPPEARED
 - 8. OTHER (SPECIFY)
 - 9. REFUSE
- 631. In the past 3 months, have you had a drink containing alcohol, that is raki, beer, wine, a cocktail, shot of liqueur, vodka, or whiskey?
 - 1. YES
 - 2. NO \rightarrow **GO TO MODULE VII**
 - 8. NOT SURE/DO NOT REMEMBER \rightarrow GO TO MODULE VII
 - 9. REFUSE \rightarrow GO TO MODULE VII

OF DRINKS

632. In the past 3 months, on the days that you drank alcohol, how many drinks did you usually have?

00. NO DRINKS/ONLY FEW SIPS → GO TO MODULE VII 88. NOT SURE/DO NOT REMEMBER → GO TO MODULE VII 99. REFUSE → GO TO MODULE VII

- 633. In the past 3 months, how often did you drink that amount (**PROBE:** per day, week, or month)?
 - 1. EVERYDAY
 - 2. ALMOST EVERY DAY
 - 3. 1-2 TIMES A WEEK
 - 4. 2-3 TIMES A MONTH
 - 5. ONCE A MONTH
 - 6. 1-2 TIMES IN THREE MONTHS
- 634. In the past 3 months, have there been days when you had more than usual (# FROM Q632) drinks?
 - 1. YES

2. NO \rightarrow GO TO MODULE VII 8. NOT SURE/DO NOT REMEMBER \rightarrow GO TO MODULE VII 9. REFUSE \rightarrow GO TO MODULE VII

635. In the past 3 months, how many drinks did you have on the days that you drank more than usual (# FROM Q632)? (CHECK IF # FROM Q635 > # FROM Q632)

 $= _ \# \text{ OF DRINKS}$ $88. \text{ NOT SURE/DO NOT REMEMBER} \rightarrow \text{ GO TO MODULE VII}$ $99. \text{ REFUSE} \rightarrow \text{ GO TO MODULE VII}$

VII REPRODUCTIVE HEALTH KNOWLEDGE/ATTITUDES

700. What do you think is the ideal number of children for a young family in Albania?

0. 0 CHILDREN	6. 3-4 CHILDREN
1. 1 CHILD	7. 4 CHILDREN
2. 1-2 CHILDREN	8. 5 OR MORE
3. 2 CHILDREN	9. AS MANY AS GOD GIVES
4. 2-3 CHILDREN	77. AS MANY AS POSSIBLE
5. 3 CHILDREN	88. DON'T KNOW

- 701. During a woman's menstrual cycle, are there certain days when she is more likely to become pregnant if she has sexual relations?
 - 1. YES
 - 2. NO \rightarrow GO TO 702
 - 8. DO NOT KNOW \rightarrow **GO TO 702**
- 701A. When is it most likely for a woman to become pregnant, just before her period begins, during her period, right after her period has ended, or halfway between two periods?
 - 1 Just before her period starts,
 - 2 During her period,
 - 3 Right after period ends,
 - 4 Halfway between her periods,
 - 8 DON'T KNOW
- 702. Do you think that breastfeeding increases, decreases or has no effect on a woman's chance to get pregnant?
 - 1. INCREASES THE CHANCE
 - 2. DECREASES THE CHANCE
 - 3. HAS NO EFFECT
 - 8. DO NOT KNOW
- 703. Do you think that a woman always has the right to decide about her pregnancy, including whether or not to have an abortion?

1. YES \rightarrow GO TO Q705

2. NO

704. Under which of the following conditions is it all right for a woman to have an abortion (**READ A-F**)?

	<u>YES</u>	<u>NO</u>	DEPENDS	<u>DK</u>
A. Her life is endangered by the pregnancy	1	2	3	8
B. The fetus has a physical deformity	1	2	3	8
C. The pregnancy was the result of rape	1	2	3	8
D. Her health is endangered by the pregnancy	1	2	3	8
E. She is unmarried	1	2	3	8
F. The couple cannot afford to have a (nother) child	1	2	3	8

- 705. If a woman had an unwanted pregnancy what should she do? (**READ 1-3**):
 - 1. Have the baby and keep it
 - 2. Have the baby and give it up for adoption
 - 3. Have an abortion
 - 8. DON'T KNOW

706. I would like to know if you are in agreement with the following statements (**READ A-I**):

	<u>AGREE</u>	DISAGREE	<u>DK</u>
A. A woman can become pregnant the first time she has sexual intercourse	1	2	8
B. All people should get married	1	2	8
C. A woman should be a virgin when she marries	1	2	8
D. The main job for a woman is to take care of the home and cook for her family	1	2	8
E. A married woman needs her husband's permission to work outside the home	1	2	8
F. If a woman works, she should give her money to her husband	1	2	8
G. If a woman works, her husband should help her with the household chores	1	2	8
H. The men in the family should have the final say in all family matters	1	2	8
I. Child care is a woman's job	1	2	8

707. Who do you think should decide how many children a couple should have (READ 1-5)?

- 1. The woman,
- 2. The man,
- 3. Both
- 4. Mother in law, or
- 5. God?
- 8 DON'T KNOW
- 708. How would you rank each of the following birth control methods (SHOW CARD C) with regard to their risk of developing health problems; please tell me if the risk is low, medium, or high:

	LOW <u>RISK</u>	MEDIUM <u>RISK</u>	HIGH <u>RISK</u>	<u>DK</u>
A. Pill	1	2	3	8
B. IUD	1	2	3	8
C. Condom	1	2	3	8
D. Tubal Ligation	1	2	3	8
E. Injectable (Depo-Provera)	1	2	3	8
F. Abortion on Request	1	2	3	8

712. Do you want to have more information about contraceptive methods?

- 1. YES
- 2. NO \rightarrow GO TO Q714
- 8. DON'T KNOW \rightarrow GO TO Q714

713. Who do you think would be the best source of information about contraceptive methods?

1. MOTHER	10. NURSE, MIDWIFE
2. OTHER RELATIVE	11. TEACHER
3. GIRLFRIEND	12. PHARMACIST
4. WIFE, PARTNER	13. BOOKS
5. SOMEBODY WHO USES CONTRACEPTION	14 NEWSPAPERS, MAGAZINES, BROCHURES
6. CO-WORKER	15. RADIO \rightarrow GO TO Q715
7. FRIEND, COLLEAGUE, PEER	16. TV → GO TO Q715
8. GYNECOLOGIST	20. OTHER (SPECIFY):
9. GENERAL PRACTITIONER	88. DON'T REMEMBER

714. Do you think that information about contraception should be broadcast on radio or television?

- 1. YES
- 2. NO
- 8. DO NOT KNOW

- 715. Some people use condoms to keep from getting sexual transmitted diseases. How effective do you think a properly used condom is for this purpose? (**READ 1-4**)
 - 1. Very Effective,
 - 2. Somewhat effective,
 - 3. Not effective?
 - 8. DON'T KNOW
- 716. In the last 12 months have you tried to obtain condoms?

YES
 NO → GO TO Q718
 NEVER HAD SEXUAL PARTNER → GO TO Q721
 DON'T KNOW WHERE TO GET THEM → GO TO Q718

716A. In the last 12 months have you succeeded in obtaining condoms?

1. YES 2. NO → **GO TO Q718**

717. In the past 12 months, how often did you have condoms in your possession?

ALL THE TIME
 ALMOST ALL THE TIME
 SOME OF THE TIME
 NEVER
 DON'T REMEMBER

717A. In the last 12 months which brand of condoms did you use most frequently?

1. DUREX	10. SILCO
2. KAMASUTRA	11. PROTEX
3. ROMED	12. AROMA
4. UNIDUS	13. MASCULAN
5. LOVEPLUS	14. TAHITI
6. LIFESTYLES	15. "FOR YOU"
7. MANIX	16. "FOR YOUR MORE"
8. TENTATION	20. OTHER
9. PLAYBOY	88. DON'T KNOW

GO TO Q718

717B. Why couldn't you succeed in obtaining condoms?

- 718. Have you ever talked with a partner about you using a condom?
 - 1. YES
 - 2. NO
 - 3. NEVER HAD A SEXUAL PARTNER \rightarrow GO TO Q721
 - 8. DON'T REMEMBER
 - 9. REFUSE

719. Have you ever suggested to any partner that you use a condom?

1. YES

2. NO \rightarrow **GO TO Q721** 8. DON'T REMEMBER \rightarrow **GO TO Q721**

9. REFUSE \rightarrow GO TO Q721

720. Has any of the following ever happened because you did not want to use a condom......(READ A-F) (ANY OF THESE INCIDENTS COULD HAVE HAPPENED MORE THAN ONCE, WITH THE SAME PARTNER OR DIFFERENT PARTNERS)

	<u>YES</u>	<u>NO</u>	<u>DK</u>	<u>REF</u>
A. You refused to use a condom?	1	2	8	9
B. You refused to have sexual intercourse with her?	1	2	8	9
C. You threaten to break up with her?	1	2	8	9
D. You shouted at and threatened her?	1	2	8	9
E. You made her have sex anyway without a condom?	1	2	8	9
F. You hit her?	1	2	8	9

721. If your partner/wife wanted you to use a condom when having sex with her, would you feel: (READ A-E)

	AGREE	DISAGREE	<u>DK</u>	<u>REF</u>
A. Angry or insulted?	1	2	8	9
B. Safe from getting pregnant?	1	2	8	9
C. Like you had done something wrong?	1	2	8	9
D. Safe from getting STDs / AIDS?	1	2	8	9
E. Suspicious that she might be sleeping around / promiscuous?	1	2	8	9

722. Please indicate whether you agree or disagree with the following statements about condoms (READ A-H):

	AGREE	DISAGREE	<u>DK</u>	<u>REF</u>
A. Using condoms with a new partner is a smart idea	1	2	8	9
B. Using condoms is not necessary if you know your partner	1	2	8	9
C. Women should ask their partners to use condoms	1	2	8	9
D. It is easy to discuss using a condom with a prospective partner	1	2	8	9
E. Condoms diminish sexual enjoyment	1	2	8	9
F. Same condoms can be used more than once	1	2	8	9
G. People who use condoms sleep around a lot / promiscuous	1	2	8	9
H. It is embarrassing to ask for condoms in FP clinics or pharmacies	1	2	8	9

723. As far as you know, do any of your friends use condoms?

1. YES

2. NO

8. DON'T KNOW

724.

What are your friends' opinions about condoms?

GOOD OPINION, FAVOR USE
 BAD OPINION, DISLIKE
 OPINIONS VARY - MIXED
 DON'T KNOW

VIII. SOCIOECONOMIC CHARACTERISTICS

800. Please tell me whether this household or any member of it has the following items: (READ A-I):

	<u>YES</u>	<u>NO</u>
A. Flush Toilet	1	2
B. Heating System	1	2
C. Refrigerator	1	2
D. TV	1	2
E. Working Automobile	1	2
F. VCR	1	2
G. Household Phone	1	2
H. Cellular Phone	1	2
I. Vacation home(villa)	1	2
J. Air Conditioner	1	2
K. Gas or Electric Stove	1	2
L. Personal Computer with Internet Access	1	2
M. Satellite Antenna	1	2

- 801. Does your family have a garden where you grow your own vegetables?
 - 1. YES
 - 2. NO
- 802. What is the main source of drinking water for members of your household?
 - 1. PIPED WATER (PIPED INTO RESIDENCE OR YARD \rightarrow GO TO Q804
 - 2. PIPED WATER (PUBLIC TAP)
 - 3. WELL WATER (RESIDENCE OR YARD) \rightarrow GO TO Q804
 - 4. PUBLIC WELL
 - 5. SURFACE WATER (SPRING, RIVER, POND, LAKE)
 - 6. RAIN WATER \rightarrow GO TO Q804
 - 7. BOTTLED WATER \rightarrow GO TO Q804
- 803. How long does it take to go to the water source, get water and come back, in minutes? _____ MINUTES
- 804. Which of the following describes your living arrangements. Do you live: (READ 1-5)
 - 1. In your privately owned flat or house,
 - 2. In rented space (room, flat or house),
 - 3. With your immediate family (NO RENT),
 - 4. With other relatives (NO RENT), or
 - 5. With friends (NO RENT)?
 - 7. OTHER (SPECIFY):

805. How many rooms are occupied by you and your family (excl. bathrooms and kitchen):

806. How many hours per day do you have electricity?

807. What is your ethnic background?

 ALBANIAN GREEK GYPSY (ROMA) MAKEDONIAN 	 ALBANIAN FROM KOSOVO MIXED ETHNICITY (SPECIFY) OTHER (SPECIFY): REFUSED/NOT STATED
What is your religion?	
1. MUSLIM	7. OTHER (SPECIFY):
2. ORTHODOX	8. NO RELIGION \rightarrow GO TO Q900

809. About how often do you usually attend religious services? (READ 1-5)

4. PROTESTANT (BAPTIST, LUTHERAN, PENTECOSTAL, ETC)

1. At least once a week

3. CATHOLIC

- 2. At least once a month, but less than once a week
- 3. Less than once a month
- 4. Only on religious holidays or special events, or
- 5. Never

808.

HOURS

9. UNDECLARED \rightarrow GO TO Q900

IX-A. AIDS/STDs

The next set of questions are about sexually transmitted infections Including HIV/ AIDS. For each of the following conditions please tell me if:

	900.	901.	902.	903.	904.
CONDITION	Have you ever	Have you ever	Have you ever	Did you take any	Who treated you
CONDITION	heard of it?	been tested for?	been told that you	treatment for?	for? (SEE
	ficulty of it.		have?	freutinent for	CODES BELOW)
A. Syphilis	1. YES	1. YES	1. YES	1. YES	CODES BEEO (II)
rt. Syphilis	$2. \text{ NO} \rightarrow \mathbf{B}$	2. NO \rightarrow B	2. NO \rightarrow B	2. NO \rightarrow B	
	$2.10 \rightarrow \mathbf{D}$	8. DK \rightarrow B	8. DK/DR \rightarrow B	8. DK/DR \rightarrow B	
B. Gonorrhea	1. YES	1. YES	1. YES	1. YES	
B. Gonornica	1.123 $2. \text{NO} \rightarrow \text{C}$	1. TES 2. NO \rightarrow C	1. TES 2. NO \rightarrow C	1. TES 2. NO \rightarrow C	
	2. NO \rightarrow C		2. NO \rightarrow C 8. DK/DR \rightarrow C	2. NO \rightarrow C 8. DK/DR \rightarrow C	
C. Chlamadia	1. YES	8. DK \rightarrow C 1. YES	$\frac{\partial DK}{DK} \rightarrow C$	8. $DK/DK \rightarrow C$ 1. YES	
C. Chlamydia					
	2. NO \rightarrow D	2. NO \rightarrow D	2. NO \rightarrow D	2. NO \rightarrow D	
		8. DK \rightarrow D	8. DK/DR \rightarrow D	8. DK/DR \rightarrow D	
D. Yeast Infection	1. YES	1. YES	1. YES	1. YES	
	2. NO $\rightarrow \mathbf{E}$	2. NO $\rightarrow \mathbf{E}$	2. NO \rightarrow E	2. NO $\rightarrow \mathbf{E}$	
		8. DK \rightarrow E	8. DK/DR \rightarrow E	8. DK/DR \rightarrow E	
E. Genital Herpes	1. YES	1. YES	1. YES	1. YES	
	2. NO \rightarrow F	2. NO \rightarrow F	2. NO \rightarrow F	2. NO \rightarrow F	
		8. DK \rightarrow F	8. DK/DR \rightarrow F	8. DK/DR \rightarrow F	
F. Genital Warts	1. YES	1. YES	1. YES	1. YES	
	2. NO \rightarrow G	2. NO \rightarrow G	2. NO \rightarrow G	2. NO \rightarrow G	
		8. DK \rightarrow G	8. DK/DR \rightarrow G	8. DK/DR \rightarrow G	
G. Trichomoniasis	1. YES	1. YES	1. YES	1. YES	
	2. NO \rightarrow H	2. NO \rightarrow H	2. NO \rightarrow H	2. NO \rightarrow H	
		8. DK \rightarrow H	8. DK/DR \rightarrow H	8. DK/DR \rightarrow H	
H. HIV / AIDS	1. YES	1. YES → Q905			
	2. NO \rightarrow Q905	2. NO \rightarrow Q905			
		8. DK \rightarrow Q905			

CODES FOR Q904:

- 1. OB/GYN
- 2. FAMILY DOCTOR/GP
- 3. DERMATOLOG
- 4. INFECTIOUS DISEASES DOCTOR
- 5. NURSE/MIDWIFE
- 6. PHARMACIST

- 7. PARTNER
- 8. FRIEND/RELATIVE
- 9. SELF-TREATMENT
- 20. OTHER:_____
- 99. REFUSE TO ANSWER

905. If a man has a sexually transmitted disease, what symptoms might he have? (RECORD ALL MENTIONED, DO NOT READ LIST)

	MENTIONED	NOT MENTIONED
A. ABDOMINAL PAIN	1	2
B. DISCHARGE FROM PENIS	1	2
C. FOUL SMELLING DISCHARGE	1	2
D. BURNING PAIN ON URINATION	1	2
E. REDNESS/INFLAMMATION IN GENITAL AREA	1	2
F. SWELLING IN GENITAL AREA	1	2
G. GENITAL SORES/ULCERS OR WARTS	1	2
H. GENITAL ITCHING	1	2
I. WEIGHT LOSS	1	2
J. HARD TO GET PREGNANT/HAVE A CHILD	1	2

906. Do you know a place where you could get an HIV/AIDS test?

- 1. YES
- 2. NO
- 907. In general, what has been your most important source of information about STDs including AIDS? (Where or from whom have you learned the most about STDs?)

1. MOTHER	11. FAMILY DOCTOR, GP
2. FATHER	12. NURSE, MIDWIFE, FELDCHER
3. OTHER RELATIVE	13. TEACHER
4. GIRLFRIEND	14. PHARMACIST
5. WIFE, PARTNER	15. SPECIALITY BOOKS
6. SOMEBODY WHO HAD STDs	16. NEWSPAPERS, MAGAZINES, BROCHURES
7. FRIENDS COLLEAGUES, PEERS	17. RADIO
8. OB/GYN DOCTOR	18. TV
9. DERMATOLOGIST	20. OTHER (SPECIFY):
10. INFECTIOUS DISEASES DOCTOR	77. NEVER HEARD OF ANY STDs $(Q900_A - H = 2)$

99. DR/REF.

908. In the past 6 months, have you seen or heard any public announcements or ads on television or radio about: (READ A-D, PROBE FOR BOTH)

				<u>DO NOT</u>
<u>YES, RADIO</u>	<u>YES, TV</u>	<u>YES, BOTH</u>	<u>NO</u>	<u>REMEMBER</u>
1	2	3	4	8
1	2	3	4	8
1	2	3	4	8
1	2	3	4	8
	YES, RADIO 1 1 1 1	YES, RADIO YES, TV 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2	YES, RADIOYES, TVYES, BOTH123123123123123	YES, RADIOYES, TVYES, BOTHNO12341234123412341234

IF Q900_H =2 (NEVER HEARD OF HIV/AIDS) GO TO Q914; ELSE CONTINUE

909. Do you think that a person can be infected with the HIV virus but have no symptoms of disease?

- 1. YES
- 2. NO
- 8. DK

910. Please tell me whether you think that the AIDS virus can be transmitted in the following ways? (READ A-M)

	<u>YES</u>	<u>NO</u>	<u>DK</u>
A. Through blood transfusion	1	2	8
B. Using public toilets	1	2	8
C. Through kissing	1	2	8
D. Through unprotected sexual intercourse between a man and a woman	1	2	8
E. Through unprotected sexual intercourse between men	1	2	8
F. By shaking hands	1	2	8
G. Using non-sterile syringes or needles	1	2	8
H. Through mosquito bites	1	2	8
I. Sharing plates, forks, or glasses with someone who has HIV/AIDS	1	2	8
J. From a woman who has the AIDS virus to her baby during pregnancy/delivery	1	2	8
K. From a mother to her child through breast milk	1	2	8
L. Getting a manicure, pedicure or haircut	1	2	8
M. Having dental or surgical treatment	1	2	8

Appendix D

911. What can a person do to reduce the risk of getting AIDS?

	SPONTANEOUS	PRO	<u>BED</u>	
	<u>YES</u>	<u>YES</u>	<u>NO</u>	<u>DK</u>
A. USE CONDOMS	1	3	4	8
B. ABSTAIN FROM SEX	1	3	4	8
C. HAVE ONLY ONE PARTNE/STAY FAITHFUL TO ONE PARTNE	R 1	3	4	8
D. LIMIT NUMBER OF SEXUAL PARTNERS	1	3	4	8
E. AVOID SEX WITH PROSTITUTES	1	3	4	8
F. AVOID SEX WITH BISEXUALS	1	3	4	8
G. DO NOT DONATE BLOOD	1	3	4	8
H. AVOID TRANSFUSIONS	1	3	4	8
I. ASK PARTNER TO GET BLOOD TESTED FOR AIDS	1	3	4	8
J. STERILIZE NEEDLES AND SYRINGES	1	3	4	8
K. DO NOT SHARE RAZORS/BLADES, NEEDLES OR SYRINGES	1	3	4	8
L. OTHER (SPECIFY)	1	3	4	8

912. How much of a risk do you think you personally have of getting HIV/AIDS? Would you say you are at (READ 1-4):

- 1. Great risk,
- 2. Moderate Risk,
- 3. Little risk, or
- 4. No risk at all \rightarrow **GO TO Q913A**
- 8. DON'T KNOW \rightarrow GO TO Q913B

913. Why do you think you have any risk of getting AIDS?

- 1. RECEIVED BLOOD TRANSFUSIONS/BLOOD PRODUCTS
- 2. HAD MANY SEXUAL PARTNERS/ TRADED SEX FOR MONEY
- 3. HAD UNPROTECTED INTERCOURSE WITH CASUAL PARTNER(S)
- 4. USED IV DRUGS/SHARED NEEDLES
- 5. PARTNER HAD/HAS SEX WITH OTHER WOMEN
- 6. HE MAY GET INFECTED WHILE RECEIVING MEDICAL OR DENTAL TREATMENT
- 7. HE MAY GET INFECTED GETTING A HAIRCUT
- 8. OTHER (SPECIFY)
- 9. DK/REF

GO TO Q913B

- 913A. Why do you think you have no risk of getting AIDS?
 - 1. ONLY ONE PARTNER
 - 2. NO SEXUAL RELATIONS
 - 3. USES CONDOMS
 - 4. CONFIDENCE IN PARTNER
 - 5. DOES NOT GET/NEED TRANSFUSIONS
 - 6. DOES NOT SHARE NEEDLES
 - 7. OTHER (SPECIFY)
 - 9. DK/REF
- 913B How much of a risk do you think you personally have of getting other STD? Would you say you are at **(READ 1-4):**
 - 1. Great risk,
 - 2. Moderate Risk,
 - 3. Little risk, or
 - 4. No risk at all
 - 8. DON'T KNOW

913C. In your opinion, it is common for unmarried man to have multiple sexual partners?

- 1. YES COMMON
- 2. NO NOT COMMON
- 8. DR/REF

913D. In your opinion, it is common for married man to have multiple sexual partners?

- 1. YES COMMON
- 2. NO NOT COMMON
- 8. DR/REF

913E. In your opinion, it is common for unmarried man to have sexual relations with prostitutes?

- 1. YES COMMON
- 2. NO NOT COMMON
- 8. DR/REF

913F. In your opinion, it is common for married man to have sexual relations with prostitutes?

- 1. YES COMMON
- 2. NO NOT COMMON
- 8. DR/REF

913G. As far as you know, is there is a cure for AIDS?

- 1. YES
- 2. NO
- 8. DR/REF

IX-B VIOLENCE

914. Thinking back to your childhood and adolescence, did you ever see or hear your parents or step-parents physically abuse each other?

- 1. YES
- 2. NO
- 3. DID NOT LIVE WITH BOTH PARENTS \rightarrow GO TO Q916
- 8. DR/REF

915. As a child, have you ever being beaten or physically mistreated in any way by anyone in your family?

- 1. YES
- 2. NO
- 8. DR/REF

916. THE INTERVIEWER SHOULD GO BACK TO PAGE 3 AND RECORD HOW MANY TIMES THIS MAN HAS EVER LIVED WITH A WOMAN AS HUSBAND AND WIFE (SEE Q111):

TIMES -----→ IF "0" GO TO Q936 ; ELSE > 0 CONTINUE

The next set of questions is about violence and physical abuse that may have happened between you and a partner or ex-partner. When we say a partner we mean a wife, ex-wife, as well as any other woman you have lived with as husband and wife.

918. Please tell me if (REAL	you have ever) A-H):	919. When was the last time you (READ A-H) your partner?	920. During the last year, how many times did you (A-H) your partner?
at partner?	1. YES \rightarrow Q919 2. NO- \rightarrow Q918_B 8. DK- \rightarrow Q918_B 9. REF \rightarrow Q918_B	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow Q918_B 3. 4-5 YEARS AGO \rightarrow Q918_B 4. 5 YEARS AGO OR MORE \rightarrow Q918_B	66. ALMOST DAILY 77. WEEKLY TIMES 88. DON'T REMEMBER 99. REFUSES
B. Threatened to hurt her or anyone she cares about?	$1. \text{ YES} \rightarrow \text{ Q919}$ $2. \text{ NO-} \rightarrow \text{ Q918}_{C}$ $8. \text{ DK-} \rightarrow \text{ Q918}_{C}$ $9. \text{ REF} \rightarrow \text{ Q918}_{C}$	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow Q918_C 3. 4-5 YEARS AGO \rightarrow Q918_C 4. 5 YEARS AGO OR MORE \rightarrow Q918_C	66. ALMOST DAILY 77. WEEKLY — TIMES 88. DON'T REMEMBER 99. REFUSES
C. Pushed, shook, shoved her, or threw something at your partner?	$1. \text{ YES} \rightarrow \text{ Q919}$ $2. \text{ NO-} \rightarrow \text{ Q918}_D$ $8. \text{ DK-} \rightarrow \text{ Q918}_D$ $9. \text{ REF} \rightarrow \text{ Q918}_D$	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow Q918_D 3. 4-5 YEARS AGO \rightarrow Q918_D 4. 5 YEARS AGO OR MORE \rightarrow Q918_D	66. ALMOST DAILY 77. WEEKLY TIMES 88. DON'T REMEMBER 99. REFUSES
D. Slapped or twisted your partner's arm?	$1. \text{ YES} \rightarrow \text{ Q919}$ $2. \text{ NO-} \rightarrow \text{ Q918}_{E}$ $8. \text{ DK-} \rightarrow \text{ Q918}_{E}$ $9. \text{ REF} \rightarrow \text{ Q918}_{E}$	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow Q918_E 3. 4-5 YEARS AGO \rightarrow Q918_E 4. 5 YEARS AGO OR MORE- \rightarrow Q918_E	66. ALMOST DAILY 77. WEEKLY TIMES 88. DON'T REMEMBER 99. REFUSES
E. Hit her with your fist or with some- thing else?	$1. \text{ YES} \rightarrow \text{ Q919}$ $2. \text{ NO-} \rightarrow \text{ Q918}_{\text{F}}$ $8. \text{ DK-} \rightarrow \text{ Q918}_{\text{F}}$ $9. \text{ REF} \rightarrow \text{ Q918}_{\text{F}}$	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow Q918_F 3. 4-5 YEARS AGO \rightarrow Q918_F 4. 5 YEARS AGO OR MORE \rightarrow Q918_F	66. ALMOST DAILY 77. WEEKLY TIMES 88. DON'T REMEMBER 99. REFUSES
F. Threatened her with a knife or other weapon?	$1. \text{ YES} \rightarrow \text{ Q919}$ $2. \text{ NO-} \rightarrow \text{ Q918}_G$ $8. \text{ DK-} \rightarrow \text{ Q918}_G$ $9. \text{ REF} \rightarrow \text{ Q918}_G$	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow Q918_G 3. 4-5 YEARS AGO \rightarrow Q918_G 4. 5 YEARS AGO OR MORE \rightarrow Q918_G	66. ALMOST DAILY 77. WEEKLY — TIMES 88. DON'T REMEMBER 99. REFUSES
G. Kicked, choked or beaten your part- ner?	$1. \text{ YES} \rightarrow \text{ Q919}$ $2. \text{ NO-} \rightarrow \text{ Q918}_{H}$ $8. \text{ DK-} \rightarrow \text{ Q918}_{H}$ $9. \text{ REF} \rightarrow \text{ Q918}_{H}$	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow Q918_H 3. 4-5 YEARS AGO \rightarrow Q918_H 4. 5 YEARS AGO OR MORE \rightarrow Q918_H	66. ALMOST DAILY 77. WEEKLY TIMES 88. DON'T REMEMBER 99. REFUSES
H. Physically forced her to have sexual relations even though she did not want to?	$\begin{array}{l} 1. \text{ YES} \rightarrow \text{ Q919} \\ 2. \text{ NO-} \rightarrow \text{ BOX 9-I} \\ 8. \text{ DK-} \rightarrow \text{ BOX 9-I} \\ 9. \text{ REF} \rightarrow \text{ BOX 9-I} \end{array}$	1. WITHIN THE LAST YEAR \rightarrow Q920 2. 1-3 YEARS AGO \rightarrow BOX 9-I 3. 4-5 YEARS AGO \rightarrow BOX9-I 4. 5 YEARS AGO OR MORE \rightarrow BOX 9-I	66. ALMOST DAILY 77. WEEKLY TIMES 88. DON'T REMEMBER 99. REFUSES

BOX 9-I

IF ALL Q918 (A-H) >1 (NEVER CAUSED ANY TYPE OF ABUSE) GO TO Q936; ELSE CONTINUE

921. You told me before that you lived with _____ partners (RECORD THE NUMBER OF PARTNERS FROM Q916). Which of these partners has physically you abused as you have just mentioned? MARK THE PARTNER(S) NUMBER FROM THE UNION TABLE AT PAGE 3 (ALLOW FOR MULTIPLE RESPONSES):

I.	 _	
II.		
III.		
IV.		

BOX 9-II

IF ANY PHYSICAL VIOLENCE TOOK PLACE DURING THE LAST YEAR (ANY Q919_C - H = 1) CONTINUE; IF ANY PHYSICAL VIOLENCE TOOK PLACE MORE THAN ONE YEAR AGO (ANY Q919_C - H > 1) GO TO Q925; IF RESPONDENT INFLICTED ONLY VERBAL VIOLENCE (Q918_C - H > 1) THEN GO TO Q936

- 922. In the past 12 months, did you cause your partner any swelling, bruises, cuts, or other physical injuries as a result of this/these incident(s)?
 - 1. YES
 - 2. NO \rightarrow GO TO Q925
 - 8. DON'T REMEMBER \rightarrow GO TO Q925
- 923. In the past 12 months, did your partner see a doctor, or other medical care provider for medical treatment of these injuries?
 - 1. YES 2. NO \rightarrow GO TO Q925 8. DO NOT REMEMBER \rightarrow GO TO Q925
- 924. Did this(these) injury(ies) require hospitalization?
 - 1. YES
 - 2. NO
 - 8. DO NOT REMEMBER
- 925. During or after a violent incident, did you want to have sexual relations with your partner sometimes, always, or never?
 - YES, SOMETIMES
 YES, ALWAYS
 NO, NEVER
 DO NOT REMEMBER/REFUSE
 REFUSED
- 926. Do you believe that violence incidents affected her health?
 - YES
 NO
 DO NOT KNOW
 REFUSED

IF RESPONDENT HAS NEVER HAD CHILD(REN), GO TO 930; OTHERWISE CONTINUE

- 928. Would you say that this violence has affected your child(ren)?
 - 1. YES
 - 2. NO \rightarrow GO TO Q930
 - 8. NOT SURE \rightarrow GO TO Q930

929. Please tell me if your child(ren) were affected in the following ways (READ A—F):

		YES	<u>NO</u>
А.	Witnessed violence	1	2
В.	Children living in fear	1	2
С.	Children injured too	1	2
D.	Children left home, live with relative	1	2
E.	Decreased Learning abilities	1	2
F.	Other (specify)	1	2

- 930. Did she talk to anyone about this(these) incidents of violence?
 - 1. YES
 - 2. NO \rightarrow GO TO Q932

931. Who did she talk with? (MARK ALL MENTIONED AND PROBE FOR ANYONE ELSE)?

	MENTIONED	NOT MENTIONED
A. HER MOTHER	1	2
B. OTHER RELATIVE	1	2
C. YOUR (RESPONENT'S) FAMILY	1	2
D. CHILDREN	1	2
E. FRIEND	1	2
F. NEIGHBOR	1	2
G. DOCTOR/HEALTH PROVIDER/SOCIAL WORKER	1	2
H. POLICE	1	2
I. LEGAL ADVISER	1	2

IF Q931_G , H, OR I = 1, GO TO Q933; ELSE CONTINUE

932. What is the main reason your partner have never sought any medical or legal help?

- 1. DID NOT KNOW WHERE TO SEEK HELP
- 2. NO USE/WOULD NOT DO ANY GOOD
- 3. EMBARRASSED
- 4. AFRAID OF MORE BEATINGS/BEING PUNISHED
- 5. AFRAID OF DIVORCE/END OF RELATIONSHIP
- 6. AFRAID OF LOOSING THE CHILDREN
- 7. THOUGHT WOULD NOT BE TAKEN SERIOUSLY/NOT BELIEVED/LAUGHED AT
- 8. VIOLENCE IS NORMAL/NO NEED TO COMPLAIN
- 9. THOUGHT SHE WOULD BE BLAMED
- 10. BRING BAD NAME TO FAMILY
- 20. OTHER
- 88. DK/REF

933. Did anyone intervene or try to stop the violence? (MARK ALL MENTIONED AND PROBE FOR ANYONE ELSE)?

	MENTIONED	NOT MENTIONED
A. YOUR MOTHER	1	2
B. OTHER RELATIVE	1	2
C. WIFE'S (PARTNER'S) FAMILY	1	2
D. CHILDREN	1	2
E. FRIEND	1	2
F. NEIGHBOR	1	2
G. DOCTOR/HEALTH PROVIDER/SOCIAL WORKER	1	2
H. POLICE	1	2
I. LEGAL ADVISER		

934. Could you tell me a little more about what usually happens when you are/were violent. Are there any particular situations that make you violent? (CIRCLE ALL THAT APPLY PROBING "ANY OTHER...")

]	MENTIONED	NOT MENTIONED
A. WHEN DRUNK	1	2
B. WHEN SHE DOES NOT LOOK AFTER CHILDREN	1	2
C. WHEN THE FAMILY HAS MONEY TROUBLES	1	2
D. WHEN YOU ARE UNEMPLOYED	1	2
E. WHEN SHE IS UNEMPLOYED/HAS NO INCOME	1	2
F. WHEN THERE ARE FAMILY PROBLEMS/MOTHER-IN-LAW PROBL	EMS 1	2
G. WHEN YOU ARE JEALOUS	1	2
H. WHEN SHE IS PREGNANT	1	2
I. WHEN HE CANNOT GET ALCOHOL/DRUGS	1	2
J. WHEN HE DOES NOT HAVE FOOD AT HOME (DINNER NOT READ	Y) 1	2
K. WHEN HE ACTS ACCORDING TO THE "KANUN"	1	2
L. OTHER	1	2

936. Sometimes a husband is annoyed or angered by things that his wife does. In your opinion, is a husband justified in hitting or beating his wife in the following situations (READ A—H):

	YES	NO	<u>DK</u>
A. If she goes out without telling him?	1	2	8
B. If she neglects the children?	1	2	8
C. If she argues with him?	1	2	8
D. If she refuses to have sex with him?	1	2	8
E. If he is not happy with her household work or food provisions?	1	2	8
F. If she asks him whether he has other girlfriends?	1	2	8
G. If he finds out that she has been unfaithful?	1	2	8
H. If she dresses too sexy or spends too much on her "look"?	1	2	8

940. THANK THE MAN FOR GIVING HIS TIME AND RECORD THE TIME THE INTERVIEW ENDED :

TIME INTERVIEW ENDED _____: _____: