

Influenza (Flu)

Weekly U.S. Influenza Surveillance Report

FLUVIEW

A Weekly Influenza Surveillance Report Prepared by the Influenza Division



Key Updates for Week 46, ending November 16, 2019

According to this week's FluView report, seasonal influenza activity in the United States continues to increase but the amount of activity and the predominant influenza virus varies by region.

Viruses

Clinical Lab

7.3% of respiratory specimens tested by clinical laboratories were positive for influenza viruses. This is higher than the previous week.

Public Health Lab

Nationally, B/Victoria viruses are most common followed by A(H1N1)pdm09 and A(H3N2) viruses. The predominant virus varies by region and age group.

Virus Characterization


Genetic and antigenic characterization and antiviral susceptibility of viruses collected in the U.S. this season is now being reported.

Illness

Outpatient Illness: ILINet


2.5% of visits to health care providers were for influenza-like illness (ILI). ILI was above the national baseline of 2.4% for the first time. Four of 10 regions were at or above their baselines.

Outpatient Illness: ILI Activity Map



The number of jurisdictions experiencing high ILI activity increased from 1 last week to 8 this week; however 44 jurisdictions experienced minimal or low ILI activity.

Geographic Spread



The number of jurisdictions reporting regional or widespread activity increased from 10 last week to 15 this week.

Severe Disease

Hospitalizations

The overall hospitalization rate for the season is 1.4 per 100,000. This is similar to what has been seen at this time during other recent seasons.

P&I Mortality

5.2% of deaths were attributed to pneumonia and influenza (P&I). This is below the epidemic threshold of 6.2%.

Pediatric Deaths

One new influenza-associated pediatric death occurring during the 2019-2020 season was reported to CDC this week. The total for the season is 4.

All data are preliminary and may change as more reports are received.

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods](#) page.

Additional information on the current and previous influenza seasons for each surveillance component are available on [FluView Interactive](#)

Key Messages from CDC


- Nationally, ILI activity has been at or above baseline for two weeks; however, the amount of influenza activity across the country varies with the south and parts of the west seeing elevated activity while other parts of the country are still seeing low activity.
- There is significant cocirculation of influenza A(H3N2), A(H1N1)pdm09 and B/Victoria viruses with the predominant virus varying by region and age group.
- The flu season is just getting started. It's not too late to get vaccinated. Flu vaccination is always the best way to prevent flu and its potentially serious complications.

U.S. Virologic Surveillance

Clinical Laboratories

The results of tests performed by clinical laboratories nationwide are summarized below. Data from clinical laboratories (the percentage of specimens tested that are positive for influenza) are used to monitor whether influenza activity is increasing or decreasing.

	Week 46	Data Cumulative since September 29, 2019 (week 40)
No. of specimens tested	24,405	158,740
No. of positive specimens (%)	1,786 (7.3%)	5,299 (3.3%)
<i>Positive specimens by type</i>		
Influenza A	470 (26.3%)	1,681 (31.7%)
Influenza B	1,316 (73.7%)	3,618 (68.3%)



Public Health Laboratories

The results of tests performed by public health laboratories nationwide are summarized below. Data from public health laboratories are used to monitor the proportion of circulating viruses that belong to each influenza subtype/lineage.


	Week 46	Data Cumulative since September 29, 2019 (week 40)
No. of specimens tested	779	8,105
No. of positive specimens	241	1,676
<i>Positive specimens by type/subtype</i>		
Influenza A	87 (36.1%)	811 (48.4%)
(H1N1)pdm09	54 (71.1%)	337 (45.7%)
H3N2	22 (28.9%)	401 (54.3%)
Subtyping not performed	11	73
Influenza B	154 (63.9%)	865 (51.6%)
Yamagata lineage	3 (2.8%)	26 (4.0%)
Victoria lineage	106 (97.2%)	630 (96.0%)

Lineage not performed

45

209

Nationally influenza B/Victoria viruses have been reported more frequently than other influenza viruses this season; followed by A(H1N1)pdm09 and A(H3N2) viruses, which are also circulating in significant numbers. The predominant virus varies by region and the proportion of influenza B/Victoria viruses is increasing in some regions. Regional and state level data about circulating influenza viruses can be found on [FluView Interactive](#). The predominant virus also varies by age group. Nationally, influenza B/Victoria viruses are the most commonly reported influenza viruses among children age 0-4 years (48% of reported viruses) and 5-24 years (56% of reported viruses), while A(H3N2) viruses are the most commonly reported influenza viruses among persons 65 years of age and older (70% of reported viruses). Among adults aged 25-64 years, approximately equal proportions of influenza A(H1N1)pdm09, A(H3N2) and B/Victoria viruses (33%, 24% and 31%, respectively) have been reported. Additional age data can be found on [FluView Interactive](#).

[View Chart Data](#) | [View Full Screen](#)

Additional virologic surveillance information for current and past seasons:

[Surveillance Methods](#) | [FluView Interactive: National, Regional, and State Data](#) or [Age Data](#)

Influenza Virus Characterization

CDC performs [genetic](#) and [antigenic](#) characterization of U.S. viruses submitted from state and local health laboratories using Right Size Roadmap submission guidance. These data are used to compare how similar the currently circulating influenza viruses are to the reference viruses used for developing new influenza vaccines and to monitor evolutionary changes that continually occur in circulating influenza. CDC also tests susceptibility of influenza viruses to antiviral medications including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the PA endonuclease inhibitor baloxavir.

CDC **genetically characterized** 174 influenza viruses collected in the U.S. from September 29, 2019 to November 16, 2019.

Virus Subtype or Lineage	Genetic Characterization				
	Total No. of Subtype/Lineage Tested	Clade	Number (% of subtype/lineage tested)	Subclade	Number (% of subtype/lineage tested)
A/H1	35				
		6B.1A	35 (100%)		
A/H3	77				
		3C.2a	77 (100%)	2a1	77 (100%)
				2a2	0
				2a3	0
				2a4	0
		3C.3a	0	3a	0
B/Victoria	58				
		V1A	58 (100%)	V1A	0
				V1A.1	10 (17.2%)
				V1A.3	48 (82.8%)
B/Yamagata	4				
		Y3	4 (100%)		

CDC **antigenically characterizes** a subset of influenza viruses by [hemagglutination inhibition \(HI\)](#) or neutralization based Focus Reduction assays (FRA). Antigenic drift is evaluated by comparing antigenic properties of cell-propagated reference viruses representing currently recommended vaccine components with those of cell-propagated circulating viruses. CDC antigenically characterized 13 influenza viruses collected in the U.S. from September 29, 2019 to November 16, 2019.

Influenza A Viruses

- **A (H1N1)pdm09:** Two A(H1N1)pdm09 viruses were antigenically characterized by HI with ferret antisera, and all were antigenically similar (reacting at titers that were within 4-fold of the homologous virus titer) to cell-propagated A/Brisbane/02/2018-like reference viruses representing the A(H1N1)pdm09 component for the 2019-20 Northern Hemisphere influenza vaccines.
- **A (H3N2):** Antigenic characterization is pending.

Influenza B Viruses

- **B/Victoria:** 11 B/Victoria lineage viruses, including viruses from both co-circulating sub-clades, were antigenically characterized by HI with ferret antisera, and seven (63.6%) were antigenically similar to cell-propagated B/Colorado/06/2017-like reference viruses representing the B/Victoria component for the 2019-20 Northern Hemisphere influenza vaccines.
- **B/Yamagata:** Antigenic characterization is pending.


CDC assesses **susceptibility of influenza viruses to the antiviral medications** oseltamivir, zanamivir, peramivir, and baloxavir using next generation sequence analysis supplemented by laboratory assays. One hundred and sixty-nine viruses collected in the U.S. from September 29, 2019 to November 16, 2019 were tested for antiviral susceptibility as follows:

Antiviral Medication			Total Viruses	A/H1	A/H3	B/Victoria	B/Yamagata
Neuraminidase Inhibitors	Oseltamivir	Viruses Tested	169	35	73	57	4
		Reduced Inhibition	0	0	0	0	0
		Highly Reduced Inhibition	1 (0.6%)	1 (2.9%)	0	0	0
	Peramivir	Viruses Tested	169	35	73	57	4
		Reduced Inhibition	0	0	0	0	0
		Highly Reduced Inhibition	1 (0.6%)	1 (2.9%)	0	0	0
	Zanamivir	Viruses Tested	169	35	73	57	4
		Reduced Inhibition	0	0	0	0	0
		Highly Reduced Inhibition	0	0	0	0	0
PA Endonuclease Inhibitor	Baloxavir	Viruses Tested	169	35	74	56	4
		Reduced Susceptibility	0	0	0	0	0

Outpatient Illness Surveillance

ILINet

Nationwide during week 46, 2.5% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). This percentage is above the national baseline of 2.4%.



[View Chart Data](#) | [View Full Screen](#)

On a regional level, the percentage of outpatient visits for ILI ranged from 1.4% to 4.4% during week 46. Region 3 (Delaware, the District of Columbia, Maryland, Pennsylvania, Virginia, and West Virginia), Region 4 (Alabama, Florida, Georgia, Kentucky, Mississippi, North Carolina, South Carolina, and Tennessee), Region 6 (Arkansas, Louisiana, New Mexico, Oklahoma, and Texas), and Region 9 (Arizona, California, Hawaii, and Nevada) reported a percentage of outpatient visits for ILI which is equal to or above their region-specific baselines. Regions 1, 2, 5, 7, 8, and 10 were below their region-specific baselines.

ILI Activity Map


Data collected in ILINet are used to produce a measure of **ILI activity*** by state.

During week 46, the following ILI activity levels were experienced:

- High – Puerto Rico and seven states (Alabama, Arkansas, Georgia, Mississippi, Nevada, South Carolina, and Texas)
- Low – New York City and 12 states (Alaska, Arizona, California, Colorado, Connecticut, Hawaii, Maryland, New Jersey, New Mexico, Oklahoma, Tennessee, and Virginia)
- Minimal – the District of Columbia and 30 states (Delaware, Florida, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, New Hampshire, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Utah, Vermont, Washington, West Virginia, Wisconsin, and Wyoming)
- Data were insufficient to calculate an ILI activity level from the U.S. Virgin Islands and one state (Louisiana).

A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet



*Data collected in ILINet may disproportionately represent certain populations within a state, and therefore, may not accurately depict the full picture of influenza activity for the whole state. Differences in the data presented here by CDC and independently by some state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.

Additional information about medically attended visits for ILI for current and past seasons:

[Surveillance Methods](#) | FluView Interactive: [National](#), [Regional](#), and [State Data](#) or [ILI Activity Map](#)

Geographic Spread of Influenza as Assessed by State and Territorial Epidemiologists


The influenza activity reported by state and territorial epidemiologists indicates geographic spread of influenza viruses but does not measure the severity of influenza activity.

During week 46 the following influenza activity was reported:

- Widespread – six states (Alabama, California, Louisiana, Massachusetts, Nevada and South Carolina)

- Regional – nine states (Arizona, Connecticut, Florida, Georgia, Hawaii, New Mexico, Oregon, Tennessee and Texas)
- Local – Puerto Rico and 23 states (Arkansas, Colorado, Delaware, Illinois, Indiana, Iowa, Kentucky, Maryland, Michigan, Minnesota, Mississippi, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oklahoma, Virginia, Washington, West Virginia, Wisconsin and Wyoming)
- Sporadic – the District of Columbia, the U.S. Virgin Islands and 11 states (Alaska, Idaho, Kansas, Maine, Missouri, Montana, Ohio, Pennsylvania, South Dakota, Utah and Vermont)
- No activity – one state (Rhode Island)
- Guam did not report.

A Weekly Influenza Surveillance Report Prepared by the Influenza Division
Weekly Influenza Activity Estimates Reported by State and Territorial Epidemiologists*



Download Image

Download Data

Most Recent Flu Activity data in XML Format (<https://www.cdc.gov/flu/weekly/fluereport.xml>) | View Full Screen (<http://gis.cdc.gov/grasp/fluview/FluView8.html>)

***This map indicates geographic spread and does not measure the severity of influenza activity.**


Additional geographic spread surveillance information for current and past seasons:

[Surveillance Methods](#) | [FluView Interactive](#)

Influenza-Associated Hospitalizations

The Influenza Hospitalization Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-related hospitalizations in select counties in the Emerging Infections Program (EIP) states and Influenza Hospitalization Surveillance Project (IHSP) states.

A total of 393 laboratory-confirmed influenza-associated hospitalizations were reported by FluSurv-NET sites between October 1, 2019 and November 16, 2019. The overall hospitalization rate was 1.4 per 100,000 population. The highest rate of hospitalization was among adults aged ≥ 65 (3.6 per 100,000 population), followed by children aged 0-4 (2.2 per 100,000 population) and adults aged 50-64 (1.4 per 100,000 population). Rates are similar to what has been seen at this time during other recent seasons. Among 393 hospitalizations, 246(62.6%) were associated with influenza A virus, 142 (36.1%) with influenza B virus, 3 (0.8%) with influenza A virus and influenza B virus co-infection, and 2 (0.5%) with influenza virus for which the type was not determined. Among those with influenza A subtype information, 28 (50.0%) were A(H1N1)pdm09 virus and 28 (50.0%) were A(H3N2).




[View Full Screen](#)

Additional hospitalization surveillance information for current and past seasons and additional age groups:
[Surveillance Methods](#) | [FluView Interactive](#)

Pneumonia and Influenza (P&I) Mortality Surveillance

Based on National Center for Health Statistics (NCHS) mortality surveillance data available on November 21, 2019, 5.2% of the deaths occurring during the week ending November 9, 2019 (week 45) were due to P&I. This percentage is below the epidemic threshold of 6.2% for week 45.

Pneumonia and Influenza Mortality from
the National Center for Health Statistics Mortality Surveillance System
Data through the week ending November 9, 2019, as of November 21, 2019




[View Chart Data](#) | [View Full Screen](#)

Additional pneumonia and influenza mortality surveillance information for current and past seasons:
[Surveillance Methods](#) | [FluView Interactive](#)

Influenza-Associated Pediatric Mortality

One influenza-associated pediatric death was reported to CDC during week 46. The death was associated with an influenza A (H1N1)pdm09 virus and occurred during week 45 (the week ending November 9, 2019).

A total of four influenza-associated pediatric deaths occurring during the 2019-2020 season have been reported to CDC.



[View Full Screen](#)

Additional pediatric mortality surveillance information for current and past seasons:
[Surveillance Methods](#) | [FluView Interactive](#)

Additional National and International Influenza Surveillance Information

FluView Interactive: FluView includes enhanced web-based interactive applications that can provide dynamic visuals of the influenza data collected and analyzed by CDC. These FluView Interactive applications allow people to create customized, visual interpretations of influenza data, as well as make comparisons across flu seasons, regions, age groups and a variety of other demographics. To access these tools, visit <http://www.cdc.gov/flu/weekly/fluviewinteractive.htm>

National Institute for Occupational Safety and Health: Monthly surveillance data on the prevalence of health-related workplace absenteeism among full-time workers in the United States are available from NIOSH at <https://www.cdc.gov/niosh/topics/absences/default.html>

U.S. State and local influenza surveillance: Select a jurisdiction below to access the latest local influenza information

Alabama	Alaska	Arizona	Arkansas	California
Colorado	Connecticut	Delaware	District of Columbia	Florida
Georgia	Hawaii	Idaho	Illinois	Indiana
Iowa	Kansas	Kentucky	Louisiana	Maine
Maryland	Massachusetts	Michigan	Minnesota	Mississippi

Missouri	Montana	Nebraska	Nevada	New Hampshire
New Jersey	New Mexico	New York	North Carolina	North Dakota
Ohio	Oklahoma	Oregon	Pennsylvania	Rhode Island
South Carolina	South Dakota	Tennessee	Texas	Utah
Vermont	Virginia	Washington	West Virginia	Wisconsin
Wyoming	New York City	Puerto Rico	Virgin Islands	

World Health Organization: Additional influenza surveillance information from participating WHO member nations is available through [FluNet](#) and the [Global Epidemiology Reports](#).

WHO Collaborating Centers for Influenza located in [Australia](#), [China](#), [Japan](#), the [United Kingdom](#), and the [United States](#) (CDC in Atlanta, Georgia).

Europe: For the most recent influenza surveillance information from Europe, please see WHO/Europe and the European Centre for Disease Prevention and Control at <http://www.flunewseurope.org/>.

Public Health Agency of Canada: The most up-to-date influenza information from Canada is available at <http://www.phac-aspc.gc.ca/fluwatch/>

Public Health England: The most up-to-date influenza information from the United Kingdom is available at <https://www.gov.uk/government/statistics/weekly-national-flu-reports>

Any links provided to non-Federal organizations are provided solely as a service to our users. These links do not constitute an endorsement of these organizations or their programs by CDC or the Federal Government, and none should be inferred. CDC is not responsible for the content of the individual organization web pages found at these links.

An overview of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component, is available at: <http://www.cdc.gov/flu/weekly/overview.htm>.