

HHS Public Access

Author manuscript *Inj Prev.* Author manuscript; available in PMC 2019 November 29.

Published in final edited form as:

Inj Prev. 2016 February ; 22(1): 52-58. doi:10.1136/injuryprev-2014-041434.

Helmets for Kids: evaluation of a school based helmet intervention in Cambodia

David J. Ederer^{1,2}, Truong Van Bui³, Erin M. Parker¹, Douglas R. Roehler^{1,2,4}, Mirjam Sidik³, Michael J. Florian³, Pagna Kim³, Sophal Sim³, Michael F. Ballesteros⁵

¹Division of Unintentional Injury Prevention, National Center for Injury Prevention and Control, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

²McNeal Professional Services, Kennesaw, Georgia, USA 3Asia Injury Prevention Foundation, Phnom Penh, Cambodia

³Asia Injury Prevention Foundation, Phnom Penh, Cambodia

⁴Department of Health Behavior and Health Education, University of Michigan School of Public Health, Ann Arbor, Michigan, USA

⁵Division of Global Health Protection, Center for Global Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

Abstract

Objective—This paper analyses helmet use before and after implementing Helmets for Kids, a school based helmet distribution and road safety programme in Cambodia.

Methods—Nine intervention schools (with a total of 6721 students) and four control schools (with a total of 3031 students) were selected using purposive sampling to target schools where students were at high risk of road traffic injury. Eligible schools included those where at least 50% of students commute to school on bicycles or motorcycles, were located on a national road (high traffic density), had few or no street signs nearby, were located in an area with a history of crash injuries and were in a province where other Cambodia Helmet Vaccine Initiative activities occur. Programme's effectiveness at each school was measured through preintervention and postintervention roadside helmet observations of students as they arrived or left school. Research assistants conducted observations 1–2 weeks preintervention, 1–2 weeks postintervention, 10–12 weeks postintervention and at the end of the school year (3–4 months postintervention).

Disclaimer

Correspondence to: David Ederer, Division of Unintentional Injury Prevention, National Center for Injury Prevention and Control, Centers for Disease Control and Prevention, 4770 Buford Highway NE, MS 62, Atlanta, GA 30341, USA; dederer@cdc.gov. Contributors

All authors reviewed the manuscript and provided revisions several times. DJE, TVB and EMP were primarily responsible for drafting, revisions and analysis. MS, PK and SS were primarily responsible for implementing the programme and data collection. MFB, DRR and MJF conceptualised and designed the evaluation, and helped interpret the data. All authors approved the submission for publication.

Twitter Follow Douglas Roehler at @droehler

The findings and conclusions in this report are those of the authors, and do not necessarily represent the official position of the CDC. Competing interests

Asia Injury Prevention Foundation authors are reporting on evaluation results of a programme they developed as part of the Cambodia Helmet Vaccine Initiative.

Results—In intervention schools, observed student helmet use increased from an average of 0.46% at 1–2 weeks preintervention to an average of 87.9% at 1–2 weeks postintervention, 83.5% at 10–12 weeks postintervention and 86.5% at 3–4 months postintervention, coinciding with the end of the school year. Increased helmet use was observed in children commuting on bicycle or motorcycle, which showed similar patterns of helmet use. Helmet use remained between 0.35% and 0.70% in control schools throughout the study period.

Conclusions—School based helmet use programmes that combine helmet provision and road safety education might increase helmet use among children.

BACKGROUND—RTCs are among the world's foremost public health problems with 1.24 million deaths and 20–50 million non fatal injuries per year. Road crashes are the eighth leading cause of all deaths globally and the leading cause of death for people aged 15–29 years. [1] Without appropriate and prompt interventions, by 2030, road traffic fatalities are expected to increase to nearly 1.9 million deaths annually. [2, 3] Over 90% of road traffic deaths occur in low income and middle income countries, and fatality risk is expected to rise as low income countries motorise. [1, 4]

RTCs are a growing public health problem in Cambodia. According to Cambodia's national Road Crash and Victim Information System, road traffic fatalities increased from 9.3 per 100 000 population in 2006 to 13.1 per 100 000 population in 2012. [5,6] The number of registered vehicles in Cambodia also increased more than 200% between 2006 and 2012, while the population increased 7.7%. [5, 6]

In 2012, 1966 of the 15 615 road crash injuries in Cambodia were fatal; of these, 156 were among children under 15 years old. [5] Furthermore, thousands of people sustain severe injuries requiring hospitalisation, resulting in lost income and educational opportunities, and sometimes permanent dis ability. The estimated economic cost of RTCs in Cambodia in 2012 was US\$329 million. [5]

In Cambodia, motorcycles are the primary means of transportation. Motorcycles comprise over 80% of Cambodia's registered motor vehicles, and account for 67% of road traffic deaths annually.[1] From 2006 to 2012, the motorcyclist fatality rate increased by 28%, from 6.99 per 100 000 to 8.96 per 100 000. [5, 6] Many motorcycle related fatalities could be prevented if motorcycle drivers and passengers in Cambodia wore helmets consistently and correctly. Motorcycle helmets have been shown to reduce the risk of death by 42% and head injury by 69%. [7] Cambodian law mandates helmet use for motorcycle drivers, but not passengers, including child passengers, contributing to low rates of helmet use. [8] One can legally drive a motorcycle in Cambodia at age 16, although this age restriction is not always observed in practice. Previous studies have shown that additional barriers to motorcycle helmet use include lack of access to high quality and low cost helmets, a belief that crash risk depends on trip length and helmet comfort. [8–10]

Although little data are available on injuries and deaths among bicyclists in Cambodia, bicyclists are also at high risk for injury and death. Like motorcyclists, bicyclists do not benefit from the protective shell of a car or from seat belts. Globally, few roads are designed with bicyclists' safety in mind, and in Cambodia, bicyclists share the same road space with

cars and motorcycles. [11] Bicycle helmets have been shown to reduce the risk of head injuries by 63%–88%; therefore, increasing helmet use among bicyclists could prevent many head injuries in Cambodia. [12]

School based and community based road safety programmes promoting bicycle helmet use have shown some success in high income countries in Europe, North America and Australia. [13–32] These interventions typically included one or more of the following: mandatory helmet policies, education campaigns and free or subsidised helmets. [13–32]

Many school based bicycle helmet programmes report increase in helmet use, decrease in head injuries and increase in road safety knowledge, but few have followed up several months after the intervention to measure long term effectiveness. [19] A meta analysis of non legislative efforts indicates that community based interventions providing free bicycle helmets are more likely to increase helmet use than interventions providing subsidised helmets and school based programmes.[19] Several analyses of bicycle helmet promotion interventions and policies have used helmet observations to measure effectiveness. [13, 17–30] However, none of these studies evaluated helmet promotion programmes in low income and middle income countries where helmet use is low, and the burden of road traffic injuries is highest, particularly among bicyclists and motorcyclists.

There is little research on school based motorcycle helmet programmes. However, a school based motorcycle helmet promotion programme in Greece showed a significant improvement in helmet and road safety knowledge after the programme was implemented. [32] Self reported helmet related attitudes and practices increased for students at public and vocational schools, but not at private schools. [32]

In 2010, the Asia Injury Prevention Foundation, together with government, private sector and nongovernmental organisations, launched the Cambodia Helmet Vaccine Initiative to promote motorcycle helmet wearing in Cambodia. The Cambodia Helmet Vaccine Initiative is part of the Global Helmet Vaccine Initiative, a multicountry effort to increase helmet use. The Helmets for Kids (HFK) programme, which is now a component of the Global Helmet Vaccine Initiative, was launched in Vietnam in 2000, and was first implemented in Cambodia in 2006. The school based programme aims to increase helmet use and road safety knowledge of students and teachers at target intervention schools. With assistance from the CDC, Asia Injury Prevention Foundation began collecting evaluation data for HFK in Cambodia in the 2011–2012 and 2012–2013 school years. The evaluation data included school based helmet observations and surveys of road safety knowledge and attitudes among students. This paper describes the HFK programme's effect on student helmet use while travelling to and from participating Cambodian schools on motorcycles or bicycles.

METHODS

Nine primary schools were selected for the HFK intervention over 2 years: five in school year 2011–2012 and four in 2012–2013. Four control schools were added in the 2012–2013 school year to determine whether the observed change in helmet use could be attributed, in

part, to other changes in the environment, such as other Cambodia Helmet Vaccine Initiative activities (see table 1 for details of control and intervention schools).

The selection of schools for the HFK intervention was meant to ensure that the programme was implemented at schools where children were at high risk for road traffic injury. To determine high risk schools, Asia Injury Prevention Foundation consulted with school officials to collect information on all primary schools located within the Cambodia Helmet Vaccine Initiative target areas. Based on this information, schools were purposively selected for the intervention using the following criteria: (1) at least 50% of students travelled to school on motorcycles or bicycles (2) school must be located on a national road (highdensity traffic area), (3) school must be in an area with no or very few traffic signs and (4) school must be in a location identified as high risk based on police reports and a history of students and teachers being involved in crashes. In the second year, the four schools selected for the HFK intervention were matched 1:1 with control schools selected on the same criteria, and matched on size and urban/rural status.

All selected schools were primary schools with students in grades 1–5. The number of students ranged from 241 to 1883 at intervention schools and from 328 to 1132 at control schools. The road user mix of students commuting to each school varied from 53% to 85% bicyclists/motorcyclists in intervention schools and from 58% to 92% at control schools. The majority of students at intervention and control schools commuted on bicycle (see table 1).

HFK Programme

HFK provides free, high quality tropical helmets to primary school students. Tropical helmets were developed to meet the needs of motorcyclists in hot climates: they are well ventilated and light weight. [33, 34] The donated helmets are produced in Vietnam to meet Standard QCVN 2 2008:B. [35] The helmets are appropriate for use on motorcycles and bicycles, and are fitted to each student. HFK educates students and teachers on traffic safety, and asks parents to encourage their children to wear helmets (http://asiainjury.org/our reach/cambodia/). Further, HFK attempts to create an environment that encourages helmet use through social norms and school policy.

At each intervention school, a handover ceremony occurred with each helmet donation. The handover ceremony served as the beginning of the programme activities. The ceremonies emphasised the importance of helmets and commitment to the HFK programme.

The road safety education programme included 1–2 h of in school training and extracurricular activities facilitated by Asia Injury Prevention Foundation staff to reinforce students' road safety knowledge. The curriculum and activities were the same across all schools. The road safety curriculum emphasised the importance of wearing helmets, how to fit and properly wear a helmet (emphasising the need to strap the helmet) and how to care for helmets. Extracurricular activities included road safety games and quizzes that reinforced HFK messages. Murals with road safety messages were painted on the walls and fences at schools.

Among the nine intervention schools, two received the intervention in two consecutive school years. This included road safety education programmes over 2 years, a helmet donation to all students in 2011–2012 and a 'topoff' donation in the second year (2012–2013) to new students and students who had damaged or lost their helmets.

Overall, 7031 helmets were distributed, including 310 in the 'topoff' donations. Students at control schools did not receive helmets, and did not take part in any intervention activities. However, both intervention and control schools were located in provinces where provincial helmet promotion occurred. Therefore, students in control schools may have been exposed to Cambodia Helmet Vaccine Initiative television commercials, billboards or radio ads promoting helmet use.

HFK evaluation

Trained research assistants videotaped either as students arrived to or departed from school to document helmet use. Helmet observations were conducted at both intervention and control schools. Control schools were not added until the second year of the programme (2012–2013). At each school, research assistants conducted helmet observations four times per school year: 1–2 weeks before helmet distribution (pre), 1–2 weeks after the helmet distribution (post I), 10–12 weeks after helmet distribution (post II) and at the end of the school year (post III). The school year ended 3–4 months after the intervention at each school. Schools were not told in advance on which day helmet observations would occur, but knew helmet observations would occur four times throughout the school year. Schools were coded from I1 to I4 for intervention commencing school year 2012–2013, from I5 to I9 for intervention commencing 2011–2012 and from C1 to C4 for control schools.

Cameras were placed in the same position and at the same time of the day for each set of observations at a school. Individual students could not be identified in the recordings. Research assistants conducted the video observations and visited the schools in advance to identify the best camera location to capture students arriving and leaving school. Observations were typically recorded at the school's main gates under comparable weather conditions.

Data captured for each student observed on a bicycle or motorcycle included: wearing helmet (yes or no) and type of vehicle (motorcycle or bicycle). Students riding motorcycles were passengers, while students riding bicycles were drivers. Coders also attempted to capture whether or not the helmet was buckled (yes or no); however, this information was difficult to observe and less reliable than whether a student was wearing a helmet at all. A single observer watched each video three times and averaged the results.

The primary purpose of the helmet observation data was to determine the effectiveness of the programme at each school; the analytical approach stems from this purpose. We employed a basic pre–post design, and included controls in the second year to capture potential province level effects. For each school, the percentage of students wearing helmets is reported at each observation point in the initial year of the intervention (pre, post I, post II and post III), along with the difference in helmet use between the first and last observation point. This difference is used to determine programme effectiveness for each school. As a

summary measure, average helmet wearing percentages, weighted by the number of observations, are also presented for both intervention and control schools. Difference in weighted average helmet use between intervention and control schools is also reported as a summary measure. A two sample t test was conducted to compare weighted average helmet use at intervention and control schools at each observation period.

Schools I5 and I8 received the intervention in consecutive years; for these schools, data from both years of the intervention are presented graphically to show the extent to which helmet use was sustained across school years. The protocol was determined to satisfy the requirements of the institutional review boards of the CDC and the National Center for Injury Prevention and Control.

RESULTS

Before helmet distribution (pre), the helmet wearing rate among students was negligible: overall, only 0.46% (range 0.0%-3.30%) of the 2478 students observed commuting to and from intervention schools by motorcycle or bicycle were wearing a helmet. Similarly, baseline helmet wearing rates among the 607 students observed at the four control schools averaged 0.35% (range 0.0%-0.7%).

Observed helmet use at intervention schools increased by at least 70 percentage points at each intervention school by the end of the school year, as shown in table 2.

The observed helmet use rate at intervention schools, weighted by the number of observations per school, averaged 87.9% (range 58.9%–100.0%) at post I, 83.5% (range 52.7%–98.0%) at post II and 86.5% (range 70.7%–100.0%) at post III. This increase was consistent for both bicycle and motorcycle helmet use, as shown in figure 1. Of students wearing helmets, approximately 99% of helmets were strapped.

In table 3, helmet wearing rates at control schools did not change much over the course of the study period, with a weighted average of 0.33% at post I (range 0.0%-4.9%), 0.70% at post II (range 0.0%-3.0%) and 0.38% at post III (range 0.0%-2.1%).

Weighted average helmet use preintervention was not significantly different between intervention and control schools (t (10)=1.15, p=0.277). However, the weighted average helmet use of intervention and control schools was significantly different at post I (t (10)=306.17, p<0.00001), post II (t (10)=265.16, p<0.00001) and post III (t (10)=270.08, p<0.00001).

In the schools that received the intervention over two consecutive years (I5 and I8), observed helmet use remained above 50% over the course of the intervention (figure 2). An additional 310 helmets were distributed to students who were not enrolled at I5 and I8 schools in the first year and to students who had damaged or lost their helmets. Prior to the handover event for the second year, the helmet wearing rates were 86.3% and 59.9% at schools I5 and I8, and continued to remain high throughout the study period, as shown in figure 2.

DISCUSSION

HFK helped raise helmet wearing rates at intervention schools. Although the ability to observe whether helmets were buckled was difficult, nearly all (approximately 99%) students had buckled helmets, suggesting that students understood how to properly wear a helmet. In contrast, observed helmet use at the four control schools did not change significantly during the study period. These findings are similar to the implementation of HFK in Vietnam where helmet wearing rates increased from 25% before intervention to an average of 94% after helmet distribution and education (results presented at conference). [36] Similar programmes in high income countries have also increased helmet use. [17] Cambodia's road traffic injury burden is high, and the majority of road users are motorcyclists or bicyclists. HFK, and programmes like it, may reduce childhood head injuries and deaths. Further research is needed to determine whether helmet education and provision programmes can be widely applied in low income and middle income countries where bicycles are widely used.

Previous studies have shown that children in Cambodia and elsewhere are strongly influenced by the helmet use behaviour of their peers; therefore, programmes such as HFK might have a role in building a broader road safety culture in families and communities where HFK is implemented. [18, 21, 30, 36, 37] The Cambodia Helmet Vaccine Initiative also implements population based interventions to encourage helmet use, such as behaviour change communication campaigns. Messages from the community based campaign and HFK are meant to be complementary to each other. The increase in helmet wearing rates might not have been as high without the community based campaign.

However, results from control schools suggest that public awareness campaigns alone are not enough to increase child helmet use. Only one control school (C1) had helmet use above 1% at any observation point. This school had a high proportion of students commuting to school by motorcycle. Helmet use may have been influenced by other local activities focused on promoting motorcycle passenger helmet use. School I1 also had a high rate of students commuting as motorcycle passengers, and was the only intervention school with recorded helmet use in the 'pre' observation period (3.30%).

Of note, one school (I9) showed a markedly lower rate of helmet use than other intervention schools. Qualitative investigations found that teachers in this school spent less time on road safety education activities, in part due to a lack of commitment from the school principal. Although the aim was to have a helmet for each child, this school did not receive enough helmets for each student. This school demonstrates the potential challenges associated with implementing HFK (eg, lack of buy in from administration, limited resources), but suggests that comprehensive school based helmet programmes can make a significant difference in helmet use even when implementation is compromised.

This study has limitations. First, research assistants did not conduct helmet observations after the end of the school year to determine how long helmet use was maintained. However, schools I5 and I8 implemented HFK over 2 years. These schools maintained helmet use rates of 86.3% and 59.9%, respectively, in the weeks before the second year of activities

began (figure 2). Although the proportion of helmeted students decreased during the school break, helmet use among students remained high beyond the initial intervention school year. This shows promise for long term increases in helmet use. Helmet use was only measured when students arrived at or left school. Further research is needed to determine whether or not students exposed to the programme use helmets outside the school setting and in the years after the intervention. It would also be beneficial to determine whether or not schools receiving HFK continue to teach safe road behaviours, and encourage students to consistently wear a helmet after the programme ends. Last, the pre–post study design is limited. Determining the magnitude of the change in helmet use is not feasible with the current study design. The findings might not be generalisable to all schools in Cambodia as schools were selected using purposive sampling as described above. The intervention was implemented at schools where students are at high risk of crashes, and have an interest in road safety. Schools interested in participating in the programme may be more likely to have increased helmet use.

In future evaluations, it would be useful to implement more rigorous evaluation such as a randomised controlled trial. Further evaluations may also compare helmet use in schools that receive the full HFK intervention against control schools with educational activities only and control schools that receive only helmet provision without educational activities. This would help quantify which components increase helmet use most and by how much. Future evaluations could also test the importance of providing helmets for free compared with subsidised helmets. Evidence from high income countries suggests that free helmet programmes are more likely to increase helmet use than programmes offering discounted helmets. [17]

This study contributes new evidence for programmes that aim to increase child helmet use in low income and middle income countries, and specifically for school based programmes that combine helmet provision and education. Evaluation data from Vietnam have demonstrated HFK's ability to increase helmet use, [36] but this is the first time that control schools have been included for an HFK programme. This study is also important given the current increasing burden of road traffic injury in low income and middle income countries, and the pressing need for effective interventions to decrease motor vehicle injuries and deaths and increase road safety.

Acknowledgements

The authors would like to thank David Sleet for his expertise, guidance and mentorship while evaluating the programme.

Funding

This work was conducted as part of the Cambodia Helmet Vaccine Initiative, funded by cooperative agreement 1U50DP001117 01 between the CDC and the International Union for Health Promotion and Education.

REFERENCES

 World Health Organization. Global status report on road safety 2013: supporting a decade of action [Internet]. Geneva: World Health Organization, 2013 http://www.who.int/iris/bitstream/ 10665/78256/1/9789241564564_eng.pdf (accessed Jul 2014).

- 2. World Health Organization. Decade of Action for Road Safety 2011–2020: saving millions of lives. http://www.who.int/violence_injury_prevention/publications/road_traffic/ saving_millions_lives_en.pdf?ua=1 (accessed Jul 2014).
- 3. World Health Organization. Projections of mortality and causes of death, 2015 and 2030. http://www.who.int/healthinfo/global_burden_disease/projections/en/ (accessed Jul 2014).
- 4. Kopits E, Cropper M. Traffic fatalities and economic growth. Accid Anal Prev 2005;37:169–78. [PubMed: 15607288]
- Cambodia National Road Safety Committee Road Crash and Victim Information System 2012 Annual Report: Road crashes and casualties in Cambodia [Internet]. Phnom Penh, Cambodia http:// www.roadsafetycambodia.info (accessed Jul 2013).
- Central Intelligence Agency. The world fact book. Cambodia, 2014 Updated 10 July 2014, (accessed 15 Jul 2014).
- 7. Liu B, Ivers R, Norton R, et al. Helmets for preventing injury in motorcycle riders. Cochrane Database Syst Rev 2008;(1):CD004333. [PubMed: 18254047]
- Bachani AM, Branching C, Ear C, et al. Trends in prevalence, knowledge, attitudes, and practices of helmet use in Cambodia: results from a two year study. Injury 2013;44:S31–7. [PubMed: 24377776]
- Roehler D, Sann S, Kim P, et al. Motorcycle helmet attitudes, behaviours and beliefs among Cambodians. Int J Inj Contr Saf Promot 2013;20:179–83. [PubMed: 23324068]
- 10. Bachani A, Tran N, Sann S, et al. Helmet use among motorcyclists in Cambodia: a survey of use, knowledge, attitudes, and practices. Traffic Inj Prev 2012;13(Supp 1):31–6.
- 11. Wegman F, Zhang F, Dijkstra A. How to make more cycling good for road safety? Accid Anal Prev 2012;44:19–29. [PubMed: 22062332]
- Thompson D, Rivara F, Thompson R. Helmets for preventing head and facial injuries in bicyclists. Cochrane Database Syst Rev 2000;(2):CD001855. [PubMed: 10796827]
- Spinks A, Turner C, McClure R, et al. Community based programmes to promote use of bicycle helmets in children aged 0–14 years: a systematic review. Int J Inj Contr Saf Promot 2005;12:131– 42. [PubMed: 16335431]
- 14. Lachapelle U, Noland R, Von Hagen L. Teaching children about bicycle safety: an evaluation of the New Jersey Bike School program. Accid Anal Prev 2013;52:237–49. [PubMed: 23414911]
- 15. Hooshmand J, Hotz G, Neilson V, et al. BikeSafe: Evaluating a bicycle safety program for middle school aged children. Accid Anal Prev 2014;66:182–6. [PubMed: 24560440]
- Ekman DS, Ekman R. Twenty five years of bicycle helmet promotion for children in Skaraborg District, Sweden. Int J Inj Contr Saf Promot 2012;19:213–7. [PubMed: 22827535]
- Kendrick D, Royal S. Cycle helmet ownership and use; a cluster randomised controlled trial in primary school children in deprived areas. Arch Dis Child 2004;89:330–5. [PubMed: 15033841]
- Buckley L, Sheehan M, Chapman R. Bicycle helmet wearing among adolescents. Trans Res Rec 2009;2140:173–81.
- Royal S, Kendrick D, Coleman T. Promoting bicycle helmet wearing by children using non legislative interventions: systematic review and meta analysis. Inj Prev 2007;13:162–7. [PubMed: 17567970]
- 20. Cote T, Sacks J, Kresnow M, et al. Bicycle helmet use among Maryland children: effect of legislation and education. Pediatrics 1992;89:1216–20. [PubMed: 1594379]
- 21. Rivara F, Thompson D, Thompson R, et al. The Seattle children's bicycle helmet campaign: changes in helmet use and head injury admissions. Pediatrics 1994;93:567–9. [PubMed: 8134210]
- 22. Rourke L Bicycle helmet use among school children. Impact of a community education program and a cycling fatality. Can Fam Physician 1994;40:1116. [PubMed: 8019188]
- Liller K, Smorynski A, McDermott R, et al. The MORE HEALTH bicycle safety project. J Sch Health 1995;65:87–90. [PubMed: 7609469]
- 24. Parkin P, Hu X, Spence L, et al. Evaluation of a subsidy program to increase bicycle helmet use by children of low income families. Pediatrics 1995;96:283–7. [PubMed: 7630685]
- Rouzier P, Alto W. Evolution of a successful community bicycle helmet campaign. J Am Board Fam Pract 1995;8:283–7. [PubMed: 7572292]

- 26. Farley C, Haddad S, Brown B. The effects of a 4 year program promoting bicycle helmet use among children in Quebec. Am J Public Health 1996;86:46–51. [PubMed: 8561241]
- 27. Logan P, Leadbetter S, Gibson R, et al. Evaluation of a Bicycle helmet giveaway program—Texas, 1995. Pediatrics 1998;101:578–82. [PubMed: 9521937]
- 28. Gilchrist J, Schieber R, Leadbetter S, et al. Police enforcement as part of a comprehensive bicycle helmet program. Pediatrics 2000;106:6–9. [PubMed: 10878141]
- Wesson D, Spence L, Hu X, et al. Trends in bicycling related head injuries in children after implementation of a community based bike helmet campaign. J Pediatr Surg 2000;35:688–9. [PubMed: 10813324]
- 30. Hall M, Cross D, Howat P, et al. Evaluation of a school based peer leader bicycle helmet intervention. Inj Control Saf Promot 2004;11:165–74. [PubMed: 15764103]
- 31. Kirsch S, Pullen N. Evaluation of a school-based education program to promote bicycle safety. Health Promot Prac 2003;4:138–45.
- 32. Germeni E, Lionis C, Kalampoki V, et al. Evaluating the impact of a school-based helmet promotion program on eligible adolescent drivers: different audiences, different needs? Health Educ Res2010;25:865–76.
- Peden M, Scurfield R, Sleet D, et al. World report on road traffic injury prevention Geneva: World Health Organization, 2004.
- Passmore JW, Nguyen LH, Nguyen NP, et al. The formulation and implementation of a national helmet law: a case study from Viet Nam. Bull World Health Organ 2010;88:783–7. [PubMed: 20931064]
- 35. QUY CHUẨN KỸTHUẬT QUỐC GIA VỀMŨBẢO HIEMCHO NGƯỜI I MÔ TÔ, XE MÁY. VietNam QCVN 2:2008/BKHCN [National technical regulation on protective helmets for motorcycle and moped users. Vietnam Standard QCVN 2:2008/BKHCN]. http://qcvn.gov.vn/ FileUpload/Documents/Mu%20Bao%20Hiem/Van%20Ban/QCVN%2002_2008_BKHCN.pdf
- 36. Brondum L, Truong T, Dinh K. Helmets for Kids programme increases helmet use among students [abstract]. Inj Prev 2012;18(Suppl 1):A112.
- De Jong M, Brijs T, Brijs K, et al. Sociological Analysis of Road Safety Situation In Cambodia. 2011 https://doclib.uhasselt.be/dspace/handle/1942/12070 (accessed Jan 2014).

What is already known about this topic?

- Vulnerable road users, including motorcyclists and bicyclists, suffer a disproportionate burden of road traffic injuries and deaths in low income and middle income countries.
- Helmet use is low among Cambodian children.
- Motorcycle and bicycle helmet use reduces the risk of head injury in children.

What this study adds?

- School based helmet programmes that include helmet distribution and education can increase child helmet use during school commutes.
- Helmet use increases can be sustained throughout the school year.

Ederer et al.



Figure 1.

Average helmet use among students commuting on bicycle or motorcycle at each observation period in the nine Helmets for Kids intervention schools by vehicle type

Ederer et al.





Figure 2. Observed helmet use in two Cambodian schools (15 and 18) that received Helmets for Kids over two consecutive years.

The postintervention II observation for I5 coincided with the end of the school year. No further observations were made. School I5 did not conduct postintervention III helmet observations because the school ended classes early in 2011–2012 due to illness. The post-topoff II helmet observation for I8 coincided with the end of the school year. School I8 did not conduct the post-topoff III helmet observation because helmets were distributed late.

Table 1

Characteristics of Helmets for Kids intervention and control schools in Cambodia

School year implemented	School ID	Number of students	Number of teachers		Students commuting to school by bicycle or motorcycle (bicycle; motorcycle)	Rural/urban status
2012-2013	I1	304		20	85% (5%; 80%)	Urban
2012-2013	I2	1206		35	80% (70%; 10%)	Rural
2012-2013	13	946		45	60% (50%; 10%)	Urban
2012-2013	I4	603		19	73% (63%; 10%)	Rural
2011-2012	15	800		28	67% (51%; 16%)	Rural
2011-2012	I6	263		8	59% (48%; 11%)	Rural
2011-2012	I7	241		8	53% (39%; 14%)	Rural
2011-2012	18	475		25	80% (74%; 6%)	Rural
2011-2012	19	1883		50	69% (43%; 26%)	Urban
Control schools	School ID	Number of students	Number of teachers		Students commuting to school by bicycle or motorcycle (bicycle; motorcycle)	Rural/urban status
2012-2013	C1	328		20	80% (6%; 74%)	Urban
2012-2013	C2	1132		29	62% (54%; 8%)	Rural
2012-2013	C3	950		28	92% (69%; 23%)	Urban
2012-2013	C4	621		19	58% (50%; 8%)	Rural

Table 2

Observed helmet use among students commuting on bicycles or motorcycles at each observation period in Helmets for Kids intervention schools in the first year of intervention in Cambodia

School ID and school year intervention	No of student in school	Pre (1–2 weeks before intervention)		Post I (1–2 weeks after intervention)		Post II (10–12 weeks after intervention)		Post III (3–4 months after intervention)		Difference between Pre and Post III
commenced	N=6721	n (obs)	% helmet	n (obs)	obs) % n (obs) % n (obs) helmet helmet	% helmet	Percentage points			
I1 (2012–13)	304	60	3.3	61	75.4	81	77.8	44	85.7	82.4
I2 (2012–13)	1206	500	0.2	589	94.5	512	97.5	473	95.3	95.1
I3 (2012–13)	946	306	0.0	293	90.1	219	82.6	323	80.5	80.5
I4 (2012–13)	603	224	0.0	255	91.8	192	84.4	140	82.9	82.9
I5 (2011–12)	800	272	0.0	323	96.9	284	97.5	NA [*]	NA [*]	97.5 *
I6 (2011–12)	263	146	0.0	130	99.2	110	94.5	80	100.0	100.0
I7 (2011–12)	241	59	0.0	103	100.0	61	90.2	36	100.0	100.0
I8 (2011–12)	475	282	0.0	166	98.8	216	97.2	229	93.9	93.9
I9 (2011–12)	1883	775	0.0	419	58.9	495	52.7	317	70.7	70.7
Weighted Average**			0.46		87.9		83.5		86.5	

n=total number of students observed; %=percent of students observed wearing a helmet.

^{*}NA: end of year helmet use could not be observed at school I5 because it closed unexpectedly early due to a viral illness among the students. Difference in per cent helmeted between post II and pre is reported.

^{*t*}Each school's reported helmet use is weighted by the number of observations when calculating weighted average.

Table 3

Observed helmet use among students commuting on bicycles or motorcycles at each observation period in the four control schools in Cambodia, 2012–2013

School ID and school-year	No of students in	Pre (1–2 weeks before intervention)		Post I (1–2 weeks after intervention)		Post II (10–12 weeks after intervention)		Post III (3–4 months after intervention)	
commenced	school N=3031	n (obs)	% helmet	n (obs)	% helmet	n (obs)	% helmet	n (obs)	% helmet
C1 (2012–13)	328	73	0.0%	82	4.9%	70	3.0%	95	2.1%
C2 (2012–13)	1132	303	0.7%	268	0.4%	230	0.0%	261	0.0%
C3 (2012–13)	950	62	0.0%	90	0.0%	88	0.0%	62	0.0%
C4 (2012–13)	621	169	0.0%	284	0.0%	253	0.0%	106	0.0%
Weighted Average*			0.35%		0.70%		0.33%		0.38%

n=total number of students observed; %=percent of students observed wearing a helmet.

* Each school's reported helmet use is weighted by the number of observations when calculating weighted average.