



# HHS Public Access

Author manuscript

*Prehosp Emerg Care*. Author manuscript; available in PMC 2020 November 01.

Published in final edited form as:

*Prehosp Emerg Care*. 2020 ; 24(1): 8–14. doi:10.1080/10903127.2019.1597954.

## How Well Do Ems Providers Predict Intracranial Hemorrhage in Head-Injured Older Adults?

**Simson Hon, BS, Samuel D. Gaona, BS**

Department of Emergency Medicine, UC Davis School of Medicine, Sacramento, California

**Mark Faul, PhD, MA**

Disease Control and Prevention, Atlanta, Georgia

**James F. Holmes, MD, MPH, Daniel K. Nishijima, MD, MAS**

Department of Emergency Medicine, UC Davis School of Medicine, Sacramento, California

**Sacramento County Prehospital Research Consortium**

### Abstract

**Objective:** To evaluate the accuracy of emergency medical services (EMS) provider judgment for traumatic intracranial hemorrhage (tICH) in older patients following head trauma in the field. We also compared EMS provider judgment with other sets of field triage criteria.

**Methods:** This was a prospective observational cohort study conducted with five EMS agencies and 11 hospitals in Northern California. Patients 55 years and older who experienced blunt head trauma were transported by EMS between August 1, 2015 and September 30, 2016, and received an initial cranial computed tomography (CT) imaging, were eligible. EMS providers were asked, “What is your suspicion for the patient having intracranial hemorrhage (bleeding in the brain)?” Responses were recorded as ordinal categories (<1%, 1–5%, >5–10%, >10–50%, or >50%) and the incidences of tICH were recorded for each category. The accuracy of EMS provider judgment was compared to other sets of triage criteria, including current field triage criteria, current field triage criteria plus multivariate logistical regression risk factors, and actual transport.

**Results:** Among the 673 patients enrolled, 319 (47.0%) were male and the median age was 75 years (interquartile range 64–85). Seventy-six (11.3%) patients had tICH on initial cranial CT imaging. The increase in EMS provider judgment correlated with an increase in the incidence of tICH. EMS provider judgment had a sensitivity of 77.6% (95% CI 67.1–85.5%) and a specificity of 41.5% (37.7–45.5%) when using a threshold of 1% or higher suspicion for tICH. Current field triage criteria (Steps 1–3) was poorly sensitive (26.3%, 95% CI 17.7–37.2%) in identifying tICH and current field trial criteria plus multivariate logistical regression risk factors was sensitive (97.4%, 95% CI 90.9–99.3%) but poorly specific (12.9%, 95% CI 10.4–15.8%). Actual transport was comparable to EMS provider judgment (sensitivity 71.1%, 95% CI 60.0–80.0%; specificity 35.3%, 95% CI 31.6–38.3%).

Address correspondence to Daniel Nishijima, MD, MAS, Department of Emergency Medicine, University of California, Davis School of Medicine, 4150 V. St, PSSB 2100, Sacramento, CA 95817. [dnishijima@ucdavis.edu](mailto:dnishijima@ucdavis.edu).

Author Contribution: J.F. Holmes and D.K. Nishijima were responsible for conception of this article; S.D. Gaona, M. Faul, J.F. Holmes, and D.K. Nishijima were responsible for the performance; S. Hon and D.K. Nishijima were responsible for the analysis of the research; and all authors were responsible for writing the manuscript.

**Conclusions:** As EMS provider judgment for tICH increased, the incidence for tICH also increased. EMS provider judgment, using a threshold of 1% or higher suspicion for tICH, was more accurate than current field triage criteria, with and without additional risk factors included.

### Keywords

emergency medical services; head injuries; closed

---

## Background

Approximately 2.8 million people suffer from a traumatic brain injury (TBI) per year in the United States (1). TBI contributes to 30% of all injury-related deaths with an estimated cost of \$60 billion annually (1, 2). In addition, more than 50% of those who experience TBI will also develop some form of tICH (3). The presence of tICH after TBI confers an increased risk of morbidity and mortality (4–6).

EMS providers are often the first to evaluate trauma patients and must make an initial assessment that includes deciding where to transport the patient (7). This initial decision carries significant implications, because severely injured patients who receive trauma center care experience a 25% reduction in mortality versus those treated at non-trauma centers (8). Unfortunately, EMS providers often have to make decisions based on the limited information that can be immediately observed, rather than data that is precisely measured (9). Several studies have measured the effectiveness of EMS provider judgment as a way to triage trauma patients, but the results have been mixed (10, 11). Major issues with using EMS provider judgment for prehospital trauma triage include over-triaging, low specificity, and variable accuracy across multiple sites (11–14). Despite these challenges, EMS providers play an important role in identifying seriously-injured patients in need of trauma center care (11, 15–18). It is important to further explore how well EMS providers predict the immediate needs of trauma patients, particularly in the context of TBI and possible tICH.

Our objective for this study was to determine the accuracy of EMS provider judgment to predict the presence of tICH, as identified on initial cranial CT scan, in older adults with head trauma. We also compared EMS provider judgment with other sets of field triage criteria.

## Methods

### Study Design and Setting

This was a prospective observational cohort study conducted in Sacramento County, California, involving 5 EMS agencies and 11 hospitals (one level I trauma center, 3 level II trauma centers, and 7 non-trauma centers). We obtained Institutional Review Board approval at all sites.

### Study Participants

Inclusion and exclusion criteria regarding participants for this study has been outlined previously (19, 20). Briefly, our participants were patients 55 years and older who

experienced head trauma and were transported by EMS between August 1, 2015 and September 30, 2016. Participants were excluded if they experienced penetrating head trauma, inter-facility transport, or did not undergo cranial CT imaging at their index ED visit. In addition, patients who did not consent to a follow-up telephone call or those who did not have a reliable means for follow-up were also excluded. EMS providers were given standardized data collection forms that included information pertaining to demographics, anticoagulant usage, and other clinical variables.

### Study Protocol and Measurements

The data collection method has been described previously (19, 20). In summary, a standardized prehospital patient care report (PCR) form was completed for each patient by participating EMS agencies. Information collected included the patient's demographics, vital signs, medical history, and history of present illness (e.g., vomiting, headache, loss of consciousness, amnesia, seizure after head injury), mechanism of injury, initial GCS score, and EMS provider judgment for tICH. EMS providers were asked "What is your suspicion for the patient having intracranial hemorrhage (bleeding in the brain)?" Responses were recorded as ordinal categories (<1%, 1–5%, >5–10%, >10–50%, or >50%).

Each patient's EMS record was then linked to hospital electronic medical records (EMR) using patient name, date of birth, and date of EMS transport. A trained research coordinator then summarized the data from the PCR and EMR. This included patient name, date of birth, mechanism of injury, detail regarding EMS transport, initial GCS score, history of vomiting, headache, loss of consciousness, amnesia, or seizure after head injury, EMS suspicion for tICH, antiplatelet and anticoagulant use, and initial cranial CT results. All tests, including cranial CT imaging, were ordered at the discretion of the treating physician.

### Outcome Measure

The primary outcome measured was the presence of tICH on initial cranial CT imaging. tICH included subarachnoid hemorrhage, subdural hemorrhage, epidural hemorrhage, intraparenchymal hemorrhage/contusion, or intraventricular hemorrhage.

### Primary Analysis

We formatted and de-identified the data and recoded the variables using STATA 14.0 statistical software (STATA Corp, College Station, TX). Descriptive statistics were used to characterize the study population. Non-normal interval data were reported with medians and quartiles 1 (Q1) and 3 (Q3). The incidence of tICH was stratified by EMS provider judgment ordinal categories (<1%, 1–5%, >5–10%, >10–50%, or >50%).

We compared the sensitivity and specificity of EMS provider judgment to identify tICH to other sets of triage criteria, including current field triage criteria, current field triage criteria plus additional risk factors identified using a multivariate logistical regression model, and actual transport. EMS provider judgment was stratified as <1% judgment for tICH vs. 1% or higher judgment for tICH. Current field triage criteria included physiologic criteria (Step 1), anatomic criteria (Step 2), and mechanism of injury criteria (Step 3) (7).

Multivariate logistical regression risk factors were identified using a regression model that included a parsimonious set of covariates, comprised of age 85 years or older (ideal cut-point based on receiver operating curve), loss of consciousness or amnesia, history of vomiting, abnormal initial EMS GCS score (dichotomized where abnormal equals GCS score less than 15), evidence of trauma above the clavicles, mechanism of injury other than a fall from standing height or less, any anticoagulant or antiplatelet use, and the presence of any Step 1 to 3 field triage criteria. Anticoagulant or antiplatelet use included the use of warfarin, direct oral anticoagulants (dabigatran, rivaroxaban, apixaban, or edoxaban), aspirin, and other antiplatelet medications (clopidogrel, ticlopidine, prasugrel, dipyridamole, cilostazol, or ticagrelor). To identify the ideal cut-point for age, we explored various age cut-points using a receiver operating curve for nonparametric data and chose the cut-point that maximized sensitivity and specificity (Youden Index) (21). Significance was defined as a  $p$  value  $<0.05$  and the results of the multivariate logistic regression model were presented as adjusted OR with 95% CIs. We used complete-case analysis to handle missing data due to the infrequency of missing data ( $<1\%$ ) and model fit was evaluated using the c-statistic (22, 23).

### Sensitivity Analysis

To evaluate for potential selection bias, we compared age, male sex, EMS GCS score, Injury Severity Score (ISS), and incidence of tICH between those who did and did not have an EMS provider judgment recorded. Because a higher risk threshold for EMS provider judgment might be tolerated, we also evaluated the sensitivity and specificity of EMS provider judgment stratified as 5% or lower judgment for tICH vs.  $>5\%$  judgment for tICH.

## Results

### Characteristics of the Patients

Our initial cohort included 1,304 patients, of which 673 patients (51.6%) had an EMS provider judgment for tICH documented. These 673 patients had a median age of 75 years (interquartile range 64–85) and 319 (47.0%) were male. Most (72.8%) of these patients had a GCS score of 15 and the majority (73.8%) had experienced a fall from standing height or less. Following emergency department evaluation, most patients (57.9%) were later discharged home. The complete description of patients included in this study is documented in Table 1. There were no differences in age, male sex, GCS score, ISS, and the presence of tICH in patients with and without an EMS provider judgment recorded (Table 2).

### Main Results

Overall, 76 (11.3%) patients enrolled in the study had a tICH on initial cranial CT imaging. EMS providers most commonly had a low suspicion for tICH and rated 265 patients (39.4%) as  $<1\%$  risk and 201 patients (29.9%) as 1–5% risk. In general, as EMS provider judgment for tICH increased, the incidence of tICH also increased (Table 3).

Adjusted analysis demonstrated a history of vomiting (OR 4.70, 95% CI 1.58–13.97), evidence of trauma above the clavicle (OR 2.41, 95% CI 1.09–5.32), an abnormal initial EMS GCS score (OR 2.23, 95% CI 1.26–3.96), Step 1 to 3 field triage criteria (OR 2.04,

95% CI 1.05 to 3.98), and loss of consciousness or amnesia (OR 2.02, 95% CI 1.14 to 3.59) as predictive of the incidence of tICH on initial cranial CT (Table 4).

Using a threshold of 1% or higher suspicion for tICH, EMS provider judgment had a sensitivity of 77.6% (95% CI 67.1–85.5%) and a specificity of 41.5% (95% CI 37.7–45.5%). Using a threshold of >5% suspicion for tICH, EMS provider judgment had a sensitivity of 53.9% (95% CI 42.8–64.7%) and a specificity of 72.2% (95% CI 68.5–75.6%).

Steps 1 to 3 of the field triage criteria was poorly sensitive (26.3%, 95% CI 17.7–37.2%) in identifying tICH and Steps 1 to 3 of the field triage criteria plus multivariate logistical regression risk factors was sensitive (97.4%, 95% CI 90.9–99.3%) but poorly specific (12.9%, 95% CI 10.4–15.8%). Actual transport had similar accuracy as EMS provider judgment (sensitivity 71.1%, 95% CI 60.0–80.0%; specificity 35.3%, 95% CI 31.6–38.3%) (Table 5). The c-statistic for the logistic regression model was 0.73, which indicates an overall good model fit.

## Discussion

Our study demonstrated that in general, as EMS provider judgment for tICH increased, the incidence for tICH also increased. Most patients were considered in the two lowest risk categories (<1% and 1 to 5%), however, the incidence of tICH in these categories was higher than predicted by EMS providers (6.4% in the <1% risk category and 9.0% in the 1 to 5% risk category). This suggests that many of these patients were well-appearing but ultimately had a tICH on initial cranial CT. This is consistent with prior literature demonstrating the difficulty in accurately identifying tICH in this patient population (24).

EMS provider judgment, using a threshold of 1% or higher suspicion, demonstrated similar accuracy compared to actual transport. Current field triage criteria using Step 1 to 3 criteria was not sensitive while the addition of the additional risk factors included in the multivariate model proved to be highly sensitive but not specific, limiting its utility as triage criteria. All sets of triage criteria failed to meet the American College of Surgeons – Committee on Trauma (ACS-COT) benchmark of a sensitivity 95% and a specificity 50% for field age criteria (20, 25).

Several studies have been conducted to evaluate EMS prehospital triage capabilities, such as determining patient ambulance usage, severity of injuries to specific body regions, and hospital admissions (10, 12, 13). While some studies support the use of EMS provider judgment in prehospital triage, others have cautioned against such usage (10–18).

While none of the sets of triage criteria met the ACS-COT benchmark for sensitivity and specificity, there are questions as to whether this benchmark can be realistically achieved (25). In particular, this patient population of older adults with head trauma are difficult to risk stratify. They often have minor, low energy mechanisms of injury and have minimal evidence of trauma but are at increased risk for tICH compared to younger patients due to anatomical differences, comorbid conditions, and frequent use of anticoagulant or antiplatelet agents. Prior decision rules have included older age as a risk factor for clinically

important TBI (26–29). However, in practice, injured older adults are also under-triaged to trauma centers more frequently than younger injured patients (30–32).

The results of our study have potential clinical and policy implications. Our results support the most recent field triage guidelines that include special considerations criteria (Step 4) (7). These criteria highlight the higher risk for potentially serious injuries in older adults despite minor mechanisms of injury. Our results also support the field triage guidelines recommendation to consider EMS provider judgment in the decision to transport injured patients to a trauma center. Particularly in well-appearing older adults with head trauma, EMS provider judgment may be more accurate to predict trauma center need than any specific set of physiologic, anatomical, or mechanistic criteria. Additional research is needed to determine the best possible triage criteria to use in evaluating these patients.

Our results should be evaluated in the context of some limitations. First, our study was conducted in a single EMS system and as such, the results may not be applicable in other EMS settings. Second, we set a low threshold for EMS provider judgment (1% suspicion) to suggest transport to a trauma center. We conducted a sensitivity analysis to evaluate a higher threshold (>5% suspicion) and the sensitivity to predict tICH decreased while the specificity increased. These, and the other, ordinal cutoff values were sensible, but were not based on any scientific evidence. Third, almost half of the subjects did not have an EMS provider judgment reported and this could lead to potential selection bias. However, there were no differences in age, sex, EMS GCS score, and the incidence of tICH in subjects with and without an EMS provider judgment reported. Fourth, EMS providers may not accurately identify anticoagulants or antiplatelet use (33). Knowledge of anticoagulant or antiplatelet use may influence EMS provider judgment. Finally, EMS providers typically do not have the medical experience to assess and diagnose tICH and, thus, this may limit the accuracy of their clinical judgment.

## Conclusions

In summary, our results suggest that increasing EMS provider judgment for tICH is consistent with an increasing incidence of tICH in head injured older adults. EMS provider judgment, using a threshold of 1% or higher suspicion for tICH, was more accurate than current field triage criteria, with and without additional risk factors included.

## Acknowledgments

The findings and conclusion of this research are those of the authors and do not represent the official views of the US Department of Health and Human Services (DHHS) and the CDC. The inclusion of individuals, programs, or organizations in this article does not constitute endorsement by the US federal government, DHHS, or CDC. D.K. Nishijima has previously served as a grant reviewer for Pfizer.

This work was funded by the Centers for Disease Control and Prevention (CDC), grant number U01CE002177. D.K. Nishijima was supported through Mentored Clinical Research Training Program Award (Grant Number UL1TR000002 and linked award KL2TR000134) from the National Center for Advancing Translational Sciences (NCATS), a component of the National Institutes of Health (NIH), and the NIH Roadmap for Medical Research.



## References

1. Taylor CA, Bell JM, Breiding MJ, Xu L. Traumatic brain injury-related emergency department visits, hospitalizations, and deaths—United States, 2007 and 2013. *MMWR Surveill Summ.* 2017;66(9):1–16. doi:10.15585/mmwr.ss6609a1.
2. Ma VY, Chan L, Carruthers KJ. Incidence, prevalence, costs, and impact on disability of common conditions requiring rehabilitation in the United States: stroke, spinal cord injury, traumatic brain injury, multiple sclerosis, osteoarthritis, rheumatoid arthritis, limb loss, and back pain. *Arch Phys Med Rehabil.* 2014;95(5):986–95.e1. doi:10.1016/j.apmr.2013.10.032. [PubMed: 24462839]
3. Roberts I, Yates D, Sandercock P, Farrell B, Wasserberg J, Lomas G, Cottingham R, Svoboda P, Brayley N, Mazairac G, et al. Effect of intravenous corticosteroids on death within 14 days in 10008 adults with clinically significant head injury (MRC CRASH trial): randomised placebo-controlled trial. *Lancet.* 2004;364(9442):1321–8. doi:10.1016/S0140-6736(04)17188-2. [PubMed: 15474134]
4. Perel P, Roberts I, Bouamra O, Woodford M, Mooney J, Lecky F. Intracranial bleeding in patients with traumatic brain injury: a prognostic study. *BMC Emerg Med.* 2009;9:15. [PubMed: 19650902]
5. Sacco S, Marini C, Toni D, Olivieri L, Carolei A. Incidence and 10-year survival of intracerebral hemorrhage in a population-based registry. *Stroke.* 2009;40(2):394–9. doi:10.1161/STROKEAHA.108.523209. [PubMed: 19038914]
6. Togha M, Bakhtavar K. Factors associated with in-hospital mortality following intracerebral hemorrhage: a three-year study in Tehran, Iran. *BMC Neurol.* 2004;4:9. [PubMed: 15193159]
7. Sasser SM, Hunt RC, Faul M, Sugerman D, Pearson WS, Dulski T, Wald MM, Jurkovich GJ, Newgard CD, Lerner EB, et al. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage, 2011. *MMWR Recomm Rep.* 2012;61(RR-1):1–20.
8. MacKenzie EJ, Rivara FP, Jurkovich GJ, Nathens AB, Frey KP, Egleston BL, Salkever DS, Scharfstein DO. A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med.* 2006;354(4):366–78. doi:10.1056/NEJMsa052049. [PubMed: 16436768]
9. Jones CM, Cushman JT, Lerner EB, Fisher SG, Seplaki CL, Veazie PJ, Wasserman EB, Dozier A, Shah MN. Prehospital trauma triage decision-making: a model of what happens between the 9-1-1 call and the hospital. *Prehosp Emerg Care.* 2016;20(1):6–14. doi:10.3109/10903127.2015.1025157. [PubMed: 26017368]
10. Lavoie A, Emond M, Moore L, Camden S, Liberman M. Evaluation of the prehospital index, presence of high-velocity impact and judgment of emergency medical technicians as criteria for trauma triage. *CJEM.* 2010;12(2):111–8. doi:10.1017/S1481803500012136. [PubMed: 20219158]
11. Brown LH, Hubble MW, Cone DC, Millin MG, Schwartz B, Patterson PD, Greenberg B, Richards ME. Paramedic determinations of medical necessity: a meta-analysis. *Prehosp Emerg Care.* 2009;13(4):516–27. doi:10.1080/10903120903144809. [PubMed: 19731166]
12. Levine SD, Colwell CB, Pons PT, Gravitz C, Haukoos JS, McVane KE. How well do paramedics predict admission to the hospital? a prospective study. *J Emerg Med.* 2006;31(1): 1–5. doi:10.1016/j.jemermed.2005.08.007. [PubMed: 16798145]
13. Mulholland SA, Cameron PA, Gabbe BJ, Williamson OD, Young K, Smith KL, Bernard SA. Prehospital prediction of the severity of blunt anatomic injury. *J Trauma.* 2008;64(3): 754–60. doi:10.1097/01.ta.0000244384.85267.c5. [PubMed: 18332820]
14. Mulholland SA, Gabbe BJ, Cameron P, Victorian State Trauma Outcomes Registry and Monitoring Group (VSTORM). Is paramedic judgement useful in prehospital trauma triage? *Injury.* 2005;36(11):1298–305. doi:10.1016/j.injury.2005.07.010. [PubMed: 16214474]
15. Emerman CL, Shade B, Kubincanek J. A comparison of EMT judgment and prehospital trauma triage instruments. *J Trauma.* 1991;31(10):1369–75. [PubMed: 1942145]
16. Fries GR, McCalla G, Levitt MA, Cordova R. A prospective comparison of paramedic judgment and the trauma triage rule in the prehospital setting. *Ann Emerg Med.* 1994;24(5): 885–9. doi:10.1016/S0196-0644(94)70207-1. [PubMed: 7978562]
17. Newgard CD, Kampp M, Nelson M, Holmes JF, Zive D, Rea T, Bulger EM, Liao M, Sherck J, Hsia RY, et al. Deciphering the use and predictive value of “emergency medical services provider judgment” in out-of-hospital trauma triage: a multisite, mixed methods assessment. *J Trauma Acute Care Surg.* 2012;72(5):1239–48. doi:10.1097/TA.0b013e3182468b51. [PubMed: 22673250]

18. Simmons E, Hedges JR, Irwin L, Maassberg W, Kirkwood HA, Jr. Paramedic injury severity perception can aid trauma triage. *Ann Emerg Med.* 1995;26(4):461–8. [PubMed: 7574129]
19. Chenoweth JA, Gaona SD, Faul M, Holmes JF, Nishijima DK, Sacramento County Prehospital Research Consortium. Incidence of delayed intracranial hemorrhage in older patients after blunt head trauma. *JAMA Surg.* 2018;153(6): 570–5. [PubMed: 29450470]
20. Nishijima DK, Gaona SD, Waechter T, Maloney R, Blitz A, Elms AR, Farrales RD, Montoya J, Bair T, Howard C, et al. The incidence of traumatic intracranial hemorrhage in head-injured older adults transported by EMS with and without anticoagulant or antiplatelet use. *J Neurotrauma.* 2018;35(5):750–9.
21. Ruopp MD, Perkins NJ, Whitcomb BW, Schisterman EF. Youden index and optimal cut-point estimated from observations affected by a lower limit of detection. *Biom J.* 2008; 50(3):419–30. doi:10.1002/bimj.200710415. [PubMed: 18435502]
22. Haukoos JS, Newgard CD. Advanced statistics: missing data in clinical research—part 1: an introduction and conceptual framework. *Acad Emerg Med.* 2007;14(7):662–8. doi:10.1111/j.1553-2712.2007.tb01855.x. [PubMed: 17538078]
23. Hosmer DW, Lemeshow S. *Applied logistic regression.* 2nd ed. New York: Wiley; 2000.
24. Nishijima DK, Offerman SR, Ballard DW, Vinson DR, Chettipally UK, Rauchwerger AS, Reed ME, Holmes JF; Clinical Research in Emergency Services and Treatment (CREST) network. Risk of traumatic intracranial hemorrhage in patients with head injury and preinjury warfarin or clopidogrel use. *Acad Emerg Med.* 2013;20(2):140–5. doi:10.1111/acem.12074. [PubMed: 23406072]
25. Newgard CD, Hsia RY, Mann NC, Schmidt T, Sahni R, Bulger EM, Wang NE, Holmes JF, Fleischman R, Zive D, et al. The trade-offs in field trauma triage: a multiregion assessment of accuracy metrics and volume shifts associated with different triage strategies. *J Trauma Acute Care Surg.* 2013;74(5):1298–306. doi:10.1097/TA.0b013e31828b7848. [PubMed: 23609282]
26. Haydel MJ, Preston CA, Mills TJ, Luber S, Blaudeau E, DeBlieux PM. Indications for computed tomography in patients with minor head injury. *N Engl J Med.* 2000;343(2): 100–5. doi:10.1056/NEJM200007133430204. [PubMed: 10891517]
27. Smits M, Dippel DW, de Haan GG, Dekker HM, Vos PE, Kool DR, Nederkoorn PJ, Hofman PA, Twijnstra A, Tanghe HL, et al. External validation of the Canadian CT head rule and the New Orleans Criteria for CT scanning in patients with minor head injury. *JAMA.* 2005;294(12):1519–25. doi:10.1001/jama.294.12.1519. [PubMed: 16189365]
28. Stiell IG, Clement CM, Rowe BH, Schull MJ, Brison R, Cass D, Eisenhauer MA, McKnight RD, Bandiera G, Holroyd B, et al. Comparison of the Canadian CT head rule and the New Orleans Criteria in patients with minor head injury. *JAMA.* 2005;294(12):1511–8. doi:10.1001/jama.294.12.1511. [PubMed: 16189364]
29. Stiell IG, Wells GA, Vandemheen K, Clement C, Lesiuk H, Laupacis A, McKnight RD, Verbeek R, Brison R, Cass D, et al. The Canadian CT head rule for patients with minor head injury. *Lancet.* 2001;357(9266):1391–6. [PubMed: 11356436]
30. Chang DC, Bass RR, Cornwell EE, MacKenzie EJ. Undertriage of elderly trauma patients to state-designated trauma centers. *Arch Surg.* 2008;143(8):776–81. [PubMed: 18711038]
31. Faul M, Xu L, Sasser SM. Hospitalized traumatic brain injury: low trauma center utilization and high interfacility transfers among older adults. *Prehosp Emerg Care.* 2016; 20(5):594–600. doi: 10.3109/10903127.2016.1149651. [PubMed: 26986195]
32. Garwe T, Stewart K, Stoner J, Newgard CD, Scott M, Zhang Y, Cathey T, Sacra J, Albrecht RM. Out-of-hospital and inter-hospital under-triage to designated tertiary trauma centers among injured older adults: a 10-year statewide geospatial-adjusted analysis. *Prehosp Emerg Care.* 2017;21(6): 734–43. doi:10.1080/10903127.2017.1332123. [PubMed: 28661712]
33. Nishijima DK, Gaona S, Waechter T, Maloney R, Bair T, Blitz A, Elms AR, Farrales RD, Howard C, Montoya J, et al. Do EMS providers accurately ascertain anticoagulant and anti-platelet use in older adults with head trauma? *Prehosp Emerg Care.* 2017;21(2):209–15. doi: 10.1080/10903127.2016.1218985. [PubMed: 27636529]



**TABLE 1.**Characteristics of the patient population,  $n = 673$ 

Characteristic	N (%)
Age, median (Q1, Q3)	75 (64, 85)
Male sex	319 (47)
Race <sup>*</sup>	
• White	484 (71.9)
• Black	50 (7.4)
• Asian	59 (8.8)
• American Indian/Alaskan Native	6 (0.9)
• Pacific Islander/Native Hawaiian	8 (1.2)
• Other	65 (9.7)
• Unknown	1 (0.1)
Ethnicity <sup>†</sup>	
• Hispanic	57 (8.5)
• Not Hispanic	603 (89.6)
EMS provider was a paramedic	664 (98.7)
Initial prehospital Glasgow Coma Scale (GCS) score <sup>‡</sup>	
• GCS score 15	490 (72.8)
• GCS score 14	125 (18.6)
• GCS score 13	16 (2.4)
• GCS score < 13	37 (5.5)
Mechanism of injury	
• Direct blow to head	39 (5.8)
• Fall from greater than standing height	30 (4.5)
• Fall from standing height or less	497 (73.8)
• Motor vehicle collision >35 miles per hour	26 (3.9)
• Motor vehicle collision ≤ 35 miles per hour	24 (3.6)
• Auto versus pedestrian/bicyclist	17 (2.5)
• Other mechanism of injury	27 (4.0)
• Unknown mechanism	13 (1.9)
Trauma above the clavicles	533 (79.2)
History of vomiting	18 (2.7)
History of headache	41 (6.1)
History of loss of consciousness or amnesia	192 (28.5)
Anticoagulant/antiplatelet medication use	
• Warfarin alone	66 (9.8)
• Direct oral anticoagulant alone	28 (4.2)
• Aspirin alone	115 (17.1)
• Other antiplatelet alone	30 (4.5)
• More than one anticoagulant or antiplatelet medication	32 (4.8)
• None	402 (59.7)

Characteristic	N (%)
International normalized ratio, median (Q1, Q3) <sup>§</sup>	2.3 (1.8, 2.8)
Platelet count, median (Q1, Q3)	215 (173.3, 262)
ED disposition	
• Discharged home	390 (57.9)
• Admitted to observation unit	19 (2.8)
• Admitted to the floor	161 (23.9)
• Admitted to the intensive care unit	73 (10.8)
• Death in the ED	2 (0.3)
• Operating room	8 (1.2)
• Transferred to another hospital	11 (1.6)
• Left against medical advice	6 (0.9)
• Other	3 (0.4)
Hospital length of stay, median (Q1, Q3) <sup>e</sup>	3 (2, 6)
Injury severity score, median (Q1, Q3) <sup>e</sup>	0 (0, 1)
Isolated head injury <sup>#</sup>	635 (94.5)

Abbreviations: ED = *emergency department*; EMS = *emergency medical services*; Q1 = *first quartile*; Q3 = *third quartile*.

\* May have more than one race.

<sup>†</sup> Missing in 13 patients.

<sup>‡</sup> Missing in five patients.

<sup>§</sup> In patients taking warfarin.

<sup>e</sup> Calculated in admitted patients only.

<sup>#</sup> If Abbreviated Injury Scale score for all non-head body regions is less than 3.

**TABLE 2.**

Comparison of characteristics and outcomes in patients with and without EMS clinical impression provided

Characteristic	EMS gestalt, <i>n</i> = 673	No EMS gestalt, <i>n</i> = 474
Age, median (IQR)	75 (64 to 85)	72 (62 to 84)
Male sex, <i>n</i> (%)	319 (47.4)	227 (47.9)
EMS GCS score, median (IQR)	15 (14 to 15)	15 (15 to 15)
ISS, median (IQR)	6 (4 to 14)	5 (2 to 12)
HCH, <i>n</i> (%)	76 (11.3)	36 (7.6)

Abbreviations: EMS = *emergency medical services*; IQR = *interquartile range*; GCS = *Glasgow Coma Score*; ISS = *injury severity score*; tICH = *traumatic intracranial hemorrhage*.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**TABLE 3.**

Incidence of traumatic intracranial hemorrhage by EMS provider judgment

EMS provider judgment*	Traumatic intracranial hemorrhage, <i>n</i> (%)
<1%	17/265 (6.4)
1–5%	18/201 (9.0)
>5–10%	7/92 (7.6)
>10–50%	23/90 (25.6)
>50%	11/25 (44.0)

\* EMS providers were asked, “What is your suspicion for the patient having intracranial hemorrhage (bleeding in the brain)?” (choices <1%, 1–5%, >5–10%, >10–50%, >50%).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**TABLE 4.**

Adjusted analysis to predict the incidence of traumatic intracranial hemorrhage on initial cranial CT scan,  $n = 668$  \*

Variable	OR (95% CI)
History of vomiting	4.70 (1.58 to 13.97)
Evidence of trauma above the clavicles	2.41 (1.09 to 5.32)
Abnormal EMS GCS score, initial	2.23 (1.26 to 3.96)
Step 1 to 3 criteria	2.04 (1.05 to 3.98)
Loss of consciousness or amnesia	2.02 (1.14 to 3.59)
Any anticoagulant or antiplatelet use	1.61 (0.94 to 2.75)
Age 85 years or older	1.43 (0.78 to 2.60)
Mechanism of injury other than a fall from standing height or less	1.37 (0.74 to 2.55)

Abbreviations: CT = *computed tomography*; EMS = *emergency medical services*; GCS = *Glasgow Coma Scale*; OR = *odds ratio*; CI = *confidence interval*.

\* Five patients were not included in the regression model due to missing data

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**TABLE 5.** Test characteristics for various combinations of triage criteria to identify traumatic intracranial hemorrhage ( $n = 76$ )

Triage criteria	Sensitivity		Specificity	
	<i>n</i>	% (95% CI)	<i>n</i>	% (95% CI)
EMS provider judgment*	59/76	77.6% (67.1–85.5%)	248/597	41.5% (37.7–45.5%)
Step 1–3 criteria	20/76	26.3% (17.7–37.2%)	527/597	88.3% (85.4–90.6%)
Step 1–3 criteria and multivariate logistic regression risk factors <sup>†</sup>	74/76	97.4% (90.9–99.3%)	77/597	12.9% (10.4–15.8%)
Actual transport	54/76	71.1% (60.0–80.0%)	211/597	35.3% (31.6–39.3%)

\* If EMS provider judgment was <1% suspicion for the patient having intracranial hemorrhage.

<sup>†</sup> Loss of consciousness or amnesia OR abnormal initial EMS GCS score OR history of vomiting OR evidence of trauma above the clavicles.