

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Hypertension in Relation to Dioxins and Polychlorinated Biphenyls from the Anniston Community Health Survey Follow-Up

Marian Pavuk, Tara C. Serio, Caroline Cusack, Matt Cave, Paula F. Rosenbaum, and Linda S. Birnbaum

Table of Contents

Table S1. Logistic regression models of dioxin TEQs, PCB Groups (lipid adjusted) and hypertension.

Table S2. Logistic regression models of summary lipid adjusted substituted TEQ quartiles and hypertension.

Table S3. Linear Regression models assessing independent effects of wet weight total dioxin TEQ and \sum non-dioxin-like PCBs on selected cytokines among individuals with hypertension in ACHS II (n=263).

Table S4. Linear Regression Parameter Estimates of Cytokines vs. Chemical Groups (Whole Weight TEQs) among individuals without hypertension (n=75).

Table S5. Linear regression models of dioxin TEQs, PCB groups, and systolic and diastolic blood pressure.

Table S6. Hazard Ratios (95% CIs) of Hypertension and TEQ tertiles (whole weight).

Figure S1. Linear regression of systolic and diastolic blood pressure on log₁₀ sum of 35 PCBs' concentration without adjustment of hypertension risk factors. The participants in the model are not on antihypertensive medication (n=124).

Figure S2. Linear regression of systolic and diastolic blood pressure on log₁₀ total dioxin TEQ concentration without adjustment of hypertension risk factors. The participants in the model are not on antihypertensive medication (n=124).