

Supplementary table 1 : comparison of outcomes and main clinical features in the original data ('base') and the cross-validation ('partition') over 10 iterations

Comorbidities	partition_subtype_1	base_subtype_1	partition_subtype_2	base_subtype_2	partition_subtype_3	base_subtype_3	partition_subtype_4	base_subtype_4	partition_subtype_5	base_subtype_5
Cirrhosis	0.4 +/- 0.1	0.3	2.5 +/- 3.0	2	0.3 +/- 0.2	0.3	15.0 +/- 3.5	17.2	8.9 +/- 1.9	9.8
HCC	0.2 +/- 0.1	0.2	1.6 +/- 2.3	1.3	0.2 +/- 0.1	0.2	13.9 +/- 3.3	16.3	6.0 +/- 1.6	6.8
CVD	11.2 +/- 0.9	13.5	44.6 +/- 11.5	29.6	19.1 +/- 6.9	5.8	23.7 +/- 3.5	27	32.0 +/- 3.3	33.9
MI	1.6 +/- 0.1	1.7	8.6 +/- 2.6	7.1	2.0 +/- 0.7	0.6	5.6 +/- 1.2	6.5	8.8 +/- 1.6	9.8
CKD	4.9 +/- 0.4	5.9	29.8 +/- 8.9	16.8	8.6 +/- 3.7	2.9	6.7 +/- 1.1	6.1	20.1 +/- 3.3	23.3
Dead	0.3 +/- 0.0	0.3	5.6 +/- 5.5	1.8	0.2 +/- 0.1	0.1	5.0 +/- 1.4	5.1	35.9 +/- 8.8	35.8
Obesity	53.9 +/- 1.1	56.2	57.1 +/- 6.2	54.4	58.7 +/- 5.4	50.1	48.0 +/- 2.8	46.2	44.1 +/- 2.2	43.4
Diabetes	28.4 +/- 1.9	31.8	66.0 +/- 14.1	48.2	52.3 +/- 18.3	27.2	42.2 +/- 4.8	45.7	46.5 +/- 5.2	48
Hypertension	50.5 +/- 2.5	55.9	83.8 +/- 13.1	70.4	67.1 +/- 16.1	39	59.9 +/- 2.9	62.9	66.0 +/- 3.0	63.7
Elevated ALT	38.2 +/- 4.1	45.7	60.2 +/- 4.2	57.7	43.4 +/- 16.1	13.7	40.5 +/- 2.0	37.4	54.2 +/- 3.0	52.8
Low platelets	8.4 +/- 1.0	9.8	36.5 +/- 13.4	30.1	13.0 +/- 4.3	3.6	68.1 +/- 12.1	78.1	76.9 +/- 5.0	79.1
Nb patients	8721 +/- 704	8665	252 +/- 140	548	1709 +/- 720	2857	970 +/- 385	851	309 +/- 118.0	369