



Published in final edited form as:

*J Parasitol.* 2019 August ; 105(4): 571–575.

## FIRST REPORT OF THE INTRODUCTION OF AN EXOTIC TICK, *AMBLYOMMA COELEBS* (ACARI: IXODIDAE), FEEDING ON A HUMAN TRAVELER RETURNING TO THE UNITED STATES FROM CENTRAL AMERICA

Goudarz Molaei<sup>1,2</sup>, Sandor E. Karpthy<sup>3</sup>, Theodore G. Andreadis<sup>1,2</sup>

<sup>1</sup>Center for Vector Biology and Zoonotic Diseases, Connecticut Agricultural Experiment Station, 123 Huntington Street, New Haven, Connecticut 06511.

<sup>2</sup>Department of Epidemiology of Microbial Diseases, Yale School of Public Health, 60 College Street, P.O. Box 208034, New Haven, Connecticut 06520-8034.

<sup>3</sup>Rickettsial Zoonoses Branch, Division of Vector-Borne Diseases, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention (CDC), Atlanta, Georgia 30329.

### Abstract

Introduction of ticks into the United States that can carry disease-causing pathogens to humans, companion animals, and wildlife has accelerated in recent years, mostly due to globalization, frequency of travel, and a rise in legal and illegal animal trades. We hereby report for the first time introduction of a live fully engorged *Amblyomma coelebs* feeding on a human into the United States from Central America. *Amblyomma coelebs* is geographically distributed in the Neotropical region and reaches the southern states of Mexico. This species is capable of transmitting a number of pathogens of public health and veterinary importance including spotted fever group rickettsiae, raising concern that *A. coelebs*, if it became established in the United States, might also be able to carry these pathogens. Considering the risks of exotic ticks as vectors of numerous pathogens and their potential to establish new populations under conducive climatic and habitat conditions, rigorous inspection practices of imported livestock and pet animals at ports of entry are vital. It is also important for travelers and practitioners to develop a heightened awareness of the public health risks associated with the unintended importation of exotic ticks and the potential such parasites have for breaching United States biosecurity defenses.

### Keywords

Exotic Tick; *Amblyomma coelebs*; Spotted Fever Group; Rickettsiae; Central America; United States

---

Introduction of ticks into the United States that can carry disease-causing pathogens to humans, companion animals, and wildlife has accelerated in recent years, mostly due to

globalization, frequency of travel, and rise in legal and illegal animal trades (Keirans and Durden, 2001; BurrIDGE, 2011). Ticks are vectors of a broad range of pathogens with global distributions, including bacteria, protozoa, filarial parasites, and arboviruses (Molaei et al., 2018), and are responsible for approximately 95% of the reported vector-borne diseases in the United States every year (Eisen et al., 2017).

Of the more than 140 exotic tick species introduced into the United States, 63 are reported to readily feed on humans and 23 are known to transmit pathogens of public health and veterinary importance (Keirans and Durden, 2001; BurrIDGE, 2011). Exotic ticks have been reported to be infected with a number of human and veterinary disease-causing pathogens in the United States, including *Ehrlichia ruminantium* (formerly known as *Cowdria ruminantium*) infection (heartwater), a lethal disease of cattle, sheep, goats, and deer, in *Amblyomma sparsum* collected from tortoises imported into Florida from Africa (BurrIDGE et al., 2000a, 2000b; BurrIDGE, 2001), and *Rickettsia* sp. in *Amblyomma exornatum* in a reptile facility in Alabama (Reeves et al., 2006).

The recent introduction and establishment of the Asian longhorned tick, *Haemaphysalis longicornis*, into the United States in 2017 (Rainey et al., 2018) highlights the enduring risk and consequences associated with invasive tick species of medical and/or veterinary importance. Reports of established populations of the Asian longhorned tick are now increasing rapidly along the Eastern Seaboard and westward, with it being reported from Arkansas, Connecticut, Kentucky, Maryland, New Jersey, New York, North Carolina, Pennsylvania, Tennessee, Virginia, and West Virginia (Beard et al., 2018; [https://www.aphis.usda.gov/animal\\_health/animal\\_diseases/tick/downloads/longhorned-tick-sitrep.pdf](https://www.aphis.usda.gov/animal_health/animal_diseases/tick/downloads/longhorned-tick-sitrep.pdf)).

A 66-yr-old male resident of North Haven, Connecticut, with a history of recent leisure travel to Costa Rica and Panama discovered a live tick attached to his back (Fig. 1). The resident's Central American trip included visits to Coiba National Park in Panama and to the Osa Conservation area, Piedras Blancas National Park, Corcovado National Park, and Wildlife Refuge in Costa Rica. The tick was removed by a dermatologist and submitted to the Connecticut Agricultural Experiment Station-Tick Testing Laboratory (CAES-TTL) on 23 March 2018 for species identification, engorgement status, and pathogen testing.

The specimen measured 5 mm in length and 4 mm in width and was identified as a fully engorged *Amblyomma* nymph upon initial morphological identification. This was followed by capturing dorsal and ventral images of the specimen (Fig. 2) using an AXIO Scope.A1 (Zeiss, Göttingen, Germany) with an attached RT3 camera system (SPOT Imaging, Sterling Heights, Michigan). Capturing higher resolution scanning electron microscopy (SEM) images was not feasible due to the risk of exposing the fully engorged tick specimen to the SEM vacuum, sample surface charging, and beam damage.

For genetic identification, DNA was extracted from the tarsi and tibiae of the specimen using the DNeasy Blood & Tissue extraction kit (Qiagen, Valencia, California) according to the manufacturer's recommendations. The mitochondrial 16S region of the extracted DNA was amplified by PCR using the method described by Ushijima et al. (2003). The PCR

product was then purified using a QIAquick PCR Purification Kit (Qiagen) and sequenced in both directions directly in cycle-sequencing reactions (Keck Sequencing Facility, New Haven, Connecticut) with a 3730xl DNA Analyzer (Applied Biosystems, Foster City, California). Sequences were annotated with ChromasPro (Technelysium Pty Ltd., Tewantin, Queensland, Australia) and identified by comparison to the GenBank DNA sequence database ([https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&PAGE\\_TYPE=BlastSearch&LINK\\_LOC=blasthome](https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&PAGE_TYPE=BlastSearch&LINK_LOC=blasthome)). The annotated sequence was deposited into the NCBI GenBank (GenBank accession ). The identity of the specimen was determined by morphological features and genetic analysis as *Amblyomma coelebs* Neumann.

The specimen was submitted on 30 March 2018 at ambient temperature to the Rickettsial Zoonoses Branch, Division of Vector-Borne Diseases, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention (CDC), Atlanta, Georgia, to screen for *Rickettsia* spp. pathogens including those common to Central and South America. The tick was macerated using a sterile scalpel and DNA extracted using a DNeasy Blood & Tissue extraction kit (Qiagen) according to the manufacturer's instructions. The DNA was examined using a pan-rickettsia qPCR assay, PanR8, as previously published (Kato et al., 2013), and the results did not indicate the presence of rickettsial pathogens. Both a positive control and a no template negative control were run along with the sample.

We hereby report for the first time introduction of a live fully engorged *A. coelebs* feeding on a human in the United States. Like other *Amblyomma* ticks, *A. coelebs* is a 3-host parasite (M. L. Levin, available at <https://www.merckvetmanual.com/integumentary-system/ticks/Amblyomma-spp.>) and is geographically distributed in the Neotropics (Onofrio et al., 2006). Adults of this species frequently feed on tapirs (*Tapirus terrestris*) as their usual hosts (Labruna and Guglielmone, 2009), but they have also been reported parasitizing a wide range of other hosts including humans, equines (Beldomenico et al., 2003), cattle, peccaries, South American foxes, wild felids, opossums, armadillos, ring-tailed cats, and various large rodents, and larvae and nymphs may also attach to birds (Ogrzewalska et al., 2010; Lopes et al., 2016; Nava et al., 2017). Reports also indicate that nymphs might be more generalists and exploit a larger host range that includes carnivores, marsupials, rodents, birds, and occasionally humans (Beldomenico et al., 2003; Ogrzewalska et al., 2009, 2010; Garcia et al., 2015; Sponchiado et al., 2015; Aguirre et al., 2018; Ito et al., 2018).

Given the wide host range and geographical distribution, it is surprising there has been no documented introduction of *A. coelebs* into the United States. The National Veterinary Services Laboratories, United States Department of Agriculture, has no record of an introduction, nor do any published reports exist of this species being introduced into the country. However, Keirens and Durden (2001) have reported on introduction of this species on seed imported into Louisiana, and Burridge (2011) has also made reference to this introduction.

Infection of *A. coelebs* with spotted fever group rickettsiae (SFGR), *Rickettsia Amblyommatis* (formerly known as '*Rickettsia amblyommii*' and later '*Candidatus*

*Rickettsia amblyommii*'), has been reported from French Guyana and from the southeastern, midwestern, and western Amazon, Brazil (Parola et al., 2007; Martins et al., 2014; Silveira et al., 2015; Witter et al., 2016). As likely the most prevalent and widely distributed SFGR species in the Americas (Karpathy et al., 2016), *R. Amblyommatis* has also been reported from several other tick species of the genus *Amblyomma*, most commonly from *Amblyomma americanum* (Labruna et al., 2004, 2007; Fitak et al., 2014; Hermance et al., 2014; Castro et al., 2015). In laboratory analysis, the WB-8-2<sup>T</sup> strain of *R. Amblyommatis* has been shown to be non-pathogenic to guinea pigs (*Cavia porcellus*) and has produced mild and transient infections in the tunica vaginalis of male meadow voles (*Microtus pennsylvanicus*) when inoculated with densely infected cell culture suspensions (Burgdorfer et al., 1981). Serological evidence suggests that humans develop a robust immune response to this organism (A. Medina and colleagues, unpubl. data; Apperson et al., 2008), and it may be associated with disease manifestations in some patients (Delisle et al., 2016). The inability of *R. Amblyommatis* to cause disease in guinea pigs (Burgdorfer et al., 1981), coupled with a lack of epidemiological evidence of human infections, suggest that this rickettsia species is not pathogenic to humans. However, its high prevalence in ticks may complicate the diagnosis and surveillance of other SFGR infections in humans (Karpathy et al., 2016).

Considering the risks of exotic ticks as vectors of numerous pathogens and their greater potential to rapidly establish new populations under conducive climatic condition and habitats and the availability of competent hosts (Ogden et al., 2008; Wu et al., 2016), rigorous inspection practices of imported livestock and pet animals at ports of entry are vital. Continuous monitoring of the infection status of parasitizing ticks and evaluating their competency for transmission of the local tick-borne pathogens could also prove important to protecting human and veterinary health. It is also important for travelers and practitioners to develop a heightened awareness of the public health risks associated with the unintended importation of exotic ticks and the potential such parasites have for breaching United States biosecurity defenses.

We are grateful to Jack L. Schlater, DVM, National Veterinary Services Laboratories, United States Department of Agriculture, for his contribution to the earlier version of this paper and for invaluable information on the biology and status of *Amblyomma coelebs* in the United States. We thank Dr. Gale E. Ridge and Katherine Dugas, CAES–Department of Entomology, for the light microscopy images, and the staff members at the CAES-TTL: Alex Diaz for technical assistance, and Mallery Breban for image editing and illustrations. The CAES-TTL is funded by the State of Connecticut. This publication was supported in part by cooperative agreement no. U01 CK000509, funded by the Centers for Disease Control and Prevention. The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

## LITERATURE CITED


- Aguirre AAR, Rodrigues VDS, Costa IND, Garcia MV, Higa LOS, Medeiros JF, and Andreotti R. 2018 Biological parameters of *Amblyomma coelebs* Neumann, 1906 (Acari: Ixodidae) under experimental conditions. *Revista Brasileira de Parasitologia* 27: 81–86.

- Apperson CS, Engber B, Nicholson WL, Mead DG, Engel J, Yabsley MJ, Dail K, Johnson J, and Watson DW. 2008 Tick-borne diseases in North Carolina: Is 'Rickettsia amblyommii' a possible cause of rickettsiosis reported as Rocky Mountain spotted fever? *Vector-Borne and Zoonotic Diseases* 8: 597–606. [PubMed: 18447622]
- Beard CB, Occi J, Bonilla DL, Egizi AM, Fonseca DM, Mertins JW, Backenson BP, Bajwa WI, Barbarin AM, Bertone MA, et al. 2018 Multistate infestation with the exotic disease-vector tick *Haemaphysalis longicornis*—United States, August 2017–September 2018. *MMWR (Morbidity and Mortality Weekly Report)* 67: 1310–1313. doi:10.15585/mmwr.mm6747a3. [PubMed: 30496158]
- Beldomenico PM, Baldi CJ, Antoniazzi LR, Orduna GM, Mastropaolo M, Macedo AC, Ruiz MF, Orcellet VM, Peralta JL, Venzal JM, et al. 2003 Ixodid ticks (Acari: Ixodidae) present at Parque Nacional El Rey, Argentina. *Neotropical Entomology* 32: 273–277.
- Burgdorfer W, Hayes SF, and Thomas LA. 1981 A new spotted fever group rickettsia from the lone star tick, *Amblyomma americanum* *In* *Rickettsiae and Rickettsial Diseases*, Burgdorfer W and Anacker RL (eds.). Academic Press, New York, New York, p. 595–602.
- Burridge MJ 2001 Ticks (Acari: Ixodidae) spread by the international trade in reptiles and their potential roles in dissemination of diseases. *Bulletin of Entomological Research* 91: 3–23. [PubMed: 11354992]
- Burridge MJ 2011 Non-native and invasive ticks: Threats to human and animal health in the United States. University Press of Florida, Gainesville, Florida, 292 p.
- Burridge MJ, Simmons LA, and Allan SA. 2000a Introduction of potential heartwater vectors and other exotic ticks into Florida on imported reptiles. *Journal of Parasitology* 86: 700–704. [PubMed: 10958443]
- Burridge MJ, Simmons LA, Simbi BH, Peter TF, and Mahan SM. 2000b Evidence of Cowdria ruminantium infection (heartwater) in *Amblyomma sparsum* ticks found on tortoises imported into Florida. *Journal of Parasitology* 86: 1135–1136. [PubMed: 11128494]
- Castro D, M. A, García S GG., Dzul-Rosado K, Aguilar A, Castillo J, Gabster A, Trejos D, Zavala-Castro J, and Bermúdez C SE. 2015 Questing *Amblyomma mixtum* and *Haemaphysalis juxtakochi* (Acari: Ixodidae) infected with *Candidatus 'Rickettsia amblyommii'* from the natural environment in Panama Canal Basin, Panama. *Tropical Medicine and Health* 43: 217–222. [PubMed: 26865823]
- Delisle J, Mendell NL, Stull-Lane A, Bloch KC, Bouyer DH, and Moncayo AC. 2016 Human infections by multiple spotted fever group Rickettsiae in Tennessee. *American Journal of Tropical Medicine and Hygiene* 94: 1212–1217. [PubMed: 27022147]
- Eisen RJ, Kugeler KJ, Eisen L, Beard CB, and Paddock CD. 2017 Tick-borne zoonoses in the United States: Persistent and emerging threats to human health. *ILAR (Institute for Laboratory Animal Research) Journal* 23: 1–17.
- Fitak RR, Kelly DJ, Daniels MK, Jiang J, Richards AL, and Fuerst PA. 2014 The prevalence of rickettsial and ehrlichial organisms in *Amblyomma americanum* ticks collected from Ohio and surrounding areas between 2000 and 2010. *Ticks and Tick-Borne Diseases* 5: 797–800. [PubMed: 25108789]
- Garcia MV, Matias J, Aguirre Ade A, Csordas BG, Szabo MP, and Andreotti R. 2015 Successful feeding of *Amblyomma coelebs* (Acari: Ixodidae) nymphs on humans in Brazil: Skin reactions to parasitism. *Journal of Medical Entomology* 52: 117–119. [PubMed: 26336294]
- Hernance M, Dos Santos RI, Heinze D, Hausser N, Bouyer DH, and Thangamani S. 2014 Detection of *Rickettsia amblyommii* in ticks collected from Missouri, USA. *Emerging Microbes and Infections* 3: e34. doi:10.1038/emi.2014.31. [PubMed: 26038740]
- Ito K, Taniguchi H, Ohtaki N, Ando S, and Kawabata H. 2018 A first case of tick bite by *Amblyomma coelebs* in Japan. *Journal of Dermatology* 45: 243–244. [PubMed: 28326613]
- Karpathy SE, Slater KS, Goldsmith CS, Nicholson WL, and Paddock CD. 2016 *Rickettsia amblyommatis* sp. nov., a spotted fever group Rickettsia associated with multiple species of *Amblyomma* ticks in North, Central and South America. *International Journal of Systematic and Evolutionary Microbiology* 66: 5236–5243. [PubMed: 27638476]

- Kato CY, Chung IH, Robinson LK, Austin AL, Dasch GA, and Massung RF. 2013 Assessment of real-time PCR assay for detection of *Rickettsia* spp. and *Rickettsia rickettsii* in banked clinical samples. *Journal of Clinical Microbiology* 51: 314–317. [PubMed: 23135935]
- Keirans JE, and Durden LA. 2001 Invasion: Exotic ticks (Acari: Ixodidae) imported into the United States. A review and new records. *Journal of Medical Entomology* 38: 850–861. [PubMed: 11761384]
- Labruna MB, and Guglielmone AA. 2009 Ticks of New World tapirs. *Tapir conservation. Newsletter of the IUCN/SSC, Tapir Specialist Group* 18/1, 21–28.
- Labruna MB, Pacheco RC, Nava S, Brandão PE, Richtzenhain LJ, and Guglielmone AA. 2007 Infection by *Rickettsia bellii* and *Candidatus 'Rickettsia amblyommii'* in *Amblyomma neumannii* ticks from Argentina. *Microbial Ecology* 54: 126–133. [PubMed: 17264991]
- Labruna MB, Whitworth T, Bouyer DH, McBride JW, Camargo LMA, Camargo EP, Popov V, and Walker DH. 2004 *Rickettsia bellii* and *Rickettsia amblyommii* in *Amblyomma* ticks from the state of Rondônia, Western Amazon, Brazil. *Journal of Medical Entomology* 41: 1073–1081. [PubMed: 15605647]
- Lopes MG, Junior JM, Foster RJ, Harmsen BJ, Sanchez E, Martins TF, Quigley H, Marcili A, and Labruna MB. 2016 Ticks and rickettsiae from wildlife in Belize, Central America. *Parasites & Vectors* 9: 62. doi:10.1186/s13071-016-1348-1. [PubMed: 26831147]
- Martins TF, Labruna MB, Mangold AJ, Cafrune MM, Guglielmone AA, and Nava S. 2014 Taxonomic key to nymphs of the genus *Amblyomma* (Acari: Ixodidae) in Argentina, with description and redescription of the nymphal stage of four *Amblyomma* species. *Ticks and Tick-Borne Diseases* 5: 753–770. [PubMed: 25113984]
- Molaei G, Andreadis TG, Anderson JF, and Stafford III KC. 2018 An exotic hitchhiker: A case report of importation into Connecticut from Africa of the human parasitizing tick, *Hyalomma truncatum* (Acari: Ixodidae). *Journal of Parasitology* 104: 302–305. [PubMed: 29412044]
- Nava S, Venzal J, González Acuña D, Martins T, and Guglielmone A. 2017 Ticks of the Southern Cone of America: Diagnosis, distribution and hosts with taxonomy, ecology and sanitary importance. Elsevier, London, U.K., 372 p.
- Ogden NH, Lindsay LR, Hanincová K, Barker IK, Bigras-Poulin M, Charron DF, Heagy A, Francis CM, O'Callaghan CJ, Schwartz I, et al. 2008 Role of migratory birds in introduction and range expansion of *Ixodes scapularis* ticks and of *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in Canada. *Applied and Environmental Microbiology* 74: 1780–1790. [PubMed: 18245258]
- Ogrzewalska M, Pacheco RC, Uezu A, Richtzenhain LJ, Ferreira F, and Labruna MB. 2009 Ticks (Acari: Ixodidae) infesting birds in an Atlantic rain forest region of Brazil. *Journal of Medical Entomology* 46: 1225–1229. [PubMed: 19769058]
- Ogrzewalska M, Uezu A, and Labruna MB. 2010 Ticks (Acari: Ixodidae) infesting wild birds in the eastern Amazon, northern Brazil, with notes on rickettsial infection in ticks. *Parasitology Research* 106: 809–816. [PubMed: 20140452]
- Onofrio VC, Venzal JM, Pinter A, and Szabó MPJ. 2006 Família Ixodidae: Características gerais, comentários e chave para gêneros In *Carrapatos de Importância Médico-Veterinária da Região Neotropical*, Barros-Battesti DM, Arzua M, and Bechara GH (eds.). Vox/ICTTD-3/Butantan, São Paulo, Brazil, p. 29–39.
- Parola P, Matsumoto K, Socolovschi C, Parzy D, and Raoult D. 2007 A tick-borne rickettsia of the spotted-fever group, similar to *Rickettsia amblyommii*, in French Guyana. *Annals of Tropical Medicine and Parasitology* 101: 185–188. [PubMed: 17316505]
- Rainey T, Occi JL, Robbins RG, and Egizi A. 2018 Discovery of *Haemaphysalis longicornis* (Ixodida: Ixodidae) parasitizing a sheep in New Jersey, United States. *Journal of Medical Entomology* 55: 757–759. [PubMed: 29471482]
- Reeves WK, Durden LA, and Dasch GA. 2006 A spotted fever group *Rickettsia* from an exotic tick species, *Amblyomma exornatum* (Acari: Ixodidae), in a reptile breeding facility in the United States. *Journal of Medical Entomology* 43: 1099–1101. [PubMed: 17017252]
- Silveira I, Martins TF, Olegário MM, Peterka C, Guedes E, Ferreira F, and Labruna MB. 2015 Rickettsial infection in animals, humans and ticks in Brazil. *Zoonoses and Public Health* 62: 525–533. [PubMed: 25643912]




- Sponchiado J, Melo GL, Martins TF, Krawczak FS, Labruna MB, and Cáceres NC. 2015 Association patterns of ticks (Acari: Ixodida: Ixodidae, Argasidae) of small mammals in Cerrado fragments, western Brazil. *Experimental and Applied Acarology* 65: 389–401. [PubMed: 25633262]
- Ushijima Y, Oliver JH, Keirans JE, Tsurumi M, Kawabata H, Watanabe H, and Fukunaga M. 2003 Mitochondrial sequence variation in *Carios capensis* (Neumann), a parasite of seabirds, collected on Torishima Island in Japan. *Journal of Parasitology* 89: 196–198. [PubMed: 12659332]
- Witter R, Martins TF, Campos AK, melo ALT, Corrêa SHR, Morgado TO, Wolf RW, May-Júnior JA, SInkoc AL, Strüssmann C, et al. 2016 Rickettsial infection in ticks (Acari: Ixodidae) of wild animals in midwestern Brazil. *Ticks and Tick-Borne Diseases* 7: 415–423. [PubMed: 26775021]
- Wu X, Rost G, and Zou X. 2016 Impact of spring bird migration on the range expansion of *Ixodes scapularis* tick populations. *Bulletin of Mathematical Biology* 78: 138–168. [PubMed: 26688012]



**Figure 1.**  
(A) *Amblyomma coelebs* engorged on the back of the patient. (B) Tick bite lesion on the patient's back after removing the tick. Color version available online.





**Figure 2.** Light microscopy images of the engorged *Amblyomma coelebs*. (A) Dorsal, (B) ventral, (C) ventral close-up of anus, (D) dorsal close-up of capitulum, and (E) ventral close-up of capitulum. Color version available online.