



National Center for Health Statistics



National Vital Statistics System

Provisional Death Counts for Coronavirus Disease (COVID-19)

i Note: Provisional death counts are based on death certificate data received and coded by the National Center for Health Statistics as of April 30, 2020. Death counts are delayed and may differ from other published sources (see Technical Notes). Counts will be updated periodically. Additional information will be added to this site as available.

- **This page has daily updates of national provisional counts for deaths involving COVID-19 in the United States by week, by age, by sex, by place of death and by jurisdiction of residence.**
- **For weekly state-specific updates by demographic and geographic characteristics, including race and Hispanic origin, visit our [Weekly Updates](#) page.**

The provisional counts for coronavirus disease (COVID-19) deaths are based on a current flow of mortality data in the National Vital Statistics System. National provisional counts include deaths occurring within the 50 states and the District of Columbia that have been received and coded as of the date specified. It is important to note that it can take several weeks for death records to be submitted to National Center for Health Statistics (NCHS), processed, coded, and tabulated. Therefore, the data shown on this page may be incomplete, and will likely not include all deaths that occurred during a given time period, especially for the more recent time periods. Death counts for earlier weeks are continually revised and may increase or decrease as new and updated death certificate data are received from the states by NCHS. COVID-19 death counts shown here may differ from other published sources, as data currently are lagged by an average of 1–2 weeks.

The provisional data presented on this page include the weekly provisional count of deaths in the United States due to COVID-19, deaths from all causes and percent of expected deaths (i.e., number of deaths received over number of deaths expected based on data from previous years), pneumonia deaths (excluding pneumonia deaths involving influenza), pneumonia deaths involving COVID-19, influenza deaths, and deaths involving pneumonia, influenza, or COVID-19; (a) by week ending date, (b) by age at death, (c) by sex, (d) by place of death, and (e) by specific jurisdictions. Future updates to this release may include additional detail such as demographic characteristics, additional causes of death (e.g., acute respiratory distress syndrome or other comorbidities), or estimates based on models that account for reporting delays to generate more accurate predicted provisional counts.

Pneumonia and influenza deaths are included to provide context for understanding the completeness of COVID-19 mortality data and related trends. Deaths due to COVID-19 may be misclassified as pneumonia or influenza deaths in the absence of positive test results, and these conditions may appear on death certificates as a comorbid condition. Thus, increases in pneumonia or influenza deaths may be an indicator of excess COVID-19-related mortality. Additionally, estimates of completeness for influenza or pneumonia deaths may provide context for understanding the lag in

reporting for COVID-19 deaths, as it is anticipated that these causes would have similar delays in reporting, processing, and coding. However, it is possible that reporting of COVID-19 mortality may be slower or faster than for other causes of death, and that the delay may change over time. Analyses to better understand and quantify reporting delays for COVID-19 deaths and related causes are underway. The list of causes provided in these tables may expand in future releases as more data are received, and other potentially comorbid conditions are determined.

Download Datasets

- [CSV Format](#) 
- [Data.CDC.gov \(Export to CSV, JSON, XLS, XML\)](#)^[?]

Table 1. Deaths involving coronavirus disease 2019 (COVID-19), pneumonia, and influenza reported to NCHS by week ending date, United States. Week ending 2/1/2020 to 4/25/2020.*

Data as of April 30, 2020

Week ending date in which the death occurred	COVID-19 Deaths (U07.1) ¹	Deaths from All Causes	Percent of Expected Deaths ²	Pneumonia Deaths (J12.0–J18.9) ³	Deaths with Pneumonia and COVID-19 (J12.0–J18.9 and U07.1) ³	Influenza Deaths (J09–J11) ⁴	Deaths with Pneumonia, Influenza, or COVID-19 (U07.1 or J09–J18.9) ⁵
Total Deaths	34,521	713,386	97	62,725	15,301	5,821	86,973
2/1/2020	0	57,263	97	3,687	0	469	4,156
2/8/2020	0	57,613	97	3,671	0	494	4,165
2/15/2020	0	56,871	97	3,692	0	516	4,208
2/22/2020	0	56,795	98	3,556	0	536	4,092
2/29/2020	5	56,831	99	3,620	3	619	4,241
3/7/2020	25	56,190	97	3,719	14	583	4,312
3/14/2020	49	53,940	95	3,674	25	578	4,275
3/21/2020	487	53,924	95	4,138	224	495	4,890
3/28/2020	2,659	57,299	103	5,598	1,220	407	7,397
4/4/2020	7,932	63,955	115	8,643	3,813	429	12,978
4/11/2020	11,864	65,676	119	9,580	5,281	427	16,272
4/18/2020	9,333	52,642	98	6,928	3,874	210	12,434
4/25/2020	2,167	24,387	45	2,219	847	58	3,553

NOTE: Number of deaths reported in this table are the total number of deaths received and coded as of the date of analysis and do not represent all deaths that occurred in that period. The United States population, based on 2018 postcensal estimates from the U.S. Census Bureau, is 327,167,434 (9).

*Data during this period are incomplete because of the lag in time between when the death occurred and when the death certificate is completed, submitted to NCHS and processed for reporting purposes. This delay can range from 1 week to 8 weeks or more, depending on the jurisdiction, age, and cause of death.

¹Deaths with confirmed or presumed COVID-19, coded to ICD-10 code U07.1

²Percent of expected deaths is the number of deaths for all causes for this week in 2020 compared to the average number across the same week in 2017–2019. Previous analyses of 2015–2016 provisional data completeness have found that completeness is lower in the first few weeks following the date of death (<25%), and then increases over time such that data are generally at least 75% complete within 8 weeks of when the death occurred (8).

³Pneumonia death counts exclude pneumonia deaths involving influenza.

⁴Influenza death counts include deaths with pneumonia or COVID-19 also listed as a cause of death.

⁵Deaths with confirmed or presumed COVID-19, Pneumonia, or Influenza, coded to ICD-10 codes U07.1 or J09–J18.9.

Table 2. Deaths involving coronavirus disease 2019 (COVID-19), pneumonia, and influenza reported to NCHS by age group, United States. Week ending 2/1/2020 to 4/25/2020.*

Data as of April 30, 2020

Age group	COVID-19 Deaths (U07.1) ¹	Deaths from All Causes	Pneumonia Deaths (J12.0–J18.9) ²	Deaths with Pneumonia and COVID-19 (J12.0–J18.9 and U07.1) ²	Influenza Deaths (J09–J11) ³	Population ⁴
All ages	34,521	713,386	62,725	15,301	5,821	327,167,434
Under 1 year	4	3,699	34	1	11	3,848,208
1–4 years	2	712	33	2	30	15,962,067
5–14 years	3	1,063	37	0	40	41,075,169
15–24 years	39	6,334	133	16	41	42,970,800
25–34 years	263	13,430	444	109	130	45,697,774
35–44 years	647	19,374	1,003	255	206	41,277,888
45–54 years	1,776	37,546	2,596	728	511	41,631,699
55–64 years	4,347	90,338	7,594	1,902	1,076	42,272,636
65–74 years	7,419	140,332	13,037	3,256	1,267	30,492,316
75–84 years	9,416	176,370	17,346	4,295	1,301	15,394,374
85 years and over	10,605	224,188	20,468	4,737	1,208	6,544,503

NOTE: Number of deaths reported in this table are the total number of deaths received and coded as of the date of analysis and do not represent all deaths that occurred in that period.

*Data during this period are incomplete because of the lag in time between when the death occurred and when the death certificate is completed, submitted to NCHS and processed for reporting purposes. This delay can range from 1 week to 8 weeks or more, depending on the jurisdiction, age, and cause of death.

¹Deaths with confirmed or presumed COVID-19, coded to ICD-10 code U07.1.

²Pneumonia death counts exclude pneumonia deaths involving influenza.

³Influenza death counts include deaths with pneumonia or COVID-19 also listed as a cause of death.

⁴Population is based on 2018 postcensal estimates from the U.S. Census Bureau (9)

Table 3. Deaths involving coronavirus disease 2019 (COVID-19), pneumonia, and influenza reported to NCHS by sex, United States. Week ending 2/1/2020 to 4/25/2020.*

Data as of April 30, 2020

Sex	COVID-19 Deaths (U07.1) ¹ ⬇	Deaths from All Causes ⬇	Pneumonia Deaths (J12.0-J18.9) ² ⬇	Deaths with Pneumonia and COVID-19 (J12.0-J18.9 and U07.1) ² ⬇	Influenza Deaths (J09-J11) ³ ⬇
Total deaths	34,521	713,386	62,725	15,301	5,821
Male	19,581	368,275	33,635	8,785	2,989
Female	14,940	345,085	29,089	6,515	2,832
Unknown	1	25	1	1	0

NOTE: Number of deaths reported in this table are the total number of deaths received and coded as of the date of analysis and do not represent all deaths that occurred in that period.

*Data during this period are incomplete because of the lag in time between when the death occurred and when the death certificate is completed, submitted to NCHS and processed for reporting purposes. This delay can range from 1 week to 8 weeks or more, depending on the jurisdiction, age, and cause of death.

¹Deaths with confirmed or presumed COVID-19, coded to ICD-10 code U07.1.

²Pneumonia death counts exclude pneumonia deaths involving influenza.

³Influenza death counts include deaths with pneumonia or COVID-19 also listed as a cause of death.

Table 4. Deaths involving coronavirus disease 2019 (COVID-19), pneumonia, and influenza reported to NCHS by place of death, United States. Week ending 2/1/2020 to 4/25/2020.*

Data as of April 30, 2020

Place of death	COVID-19 Deaths (U07.1) ¹ ⬇	Deaths from All Causes ⬇	Pneumonia Deaths (J12.0-J18.9) ² ⬇	Deaths with Pneumonia and COVID-19 (J12.0-J18.9 and U07.1) ² ⬇	Influenza Deaths (J09-J11) ³ ⬇
Total	34,521	713,386	62,725	15,301	5,821
Healthcare setting, inpatient	23,441	205,899	43,022	12,092	3,600
Healthcare setting, outpatient or emergency room	1,648	42,360	2,418	619	191
Healthcare setting, dead on arrival	44	1,884	49	12	11
Decedent's home	2,415	230,062	4,634	338	1,201
Hospice facility	439	49,755	3,089	216	289
Nursing home/long term care facility	6,160	138,583	8,509	1,924	449

Other	369	44,579	983	97	80
Place of death unknown	5	264	21	3	0

NOTE: Number of deaths reported in this table are the total number of deaths received and coded as of the date of analysis and do not represent all deaths that occurred in that period.

*Data during this period are incomplete because of the lag in time between when the death occurred and when the death certificate is completed, submitted to NCHS and processed for reporting purposes. This delay can range from 1 week to 8 weeks or more, depending on the jurisdiction, age, and cause of death.

¹Deaths with confirmed or presumed COVID-19, coded to ICD-10 code U07.1.

²Pneumonia death counts exclude pneumonia deaths involving influenza.

³Influenza death counts include deaths with pneumonia or COVID-19 also listed as a cause of death.

Table 5. Deaths involving coronavirus disease 2019 (COVID-19), pneumonia, and influenza reported to NCHS by jurisdiction of occurrence, United States. Week ending 2/1/2020 to 4/25/2020.*

Data as of April 30, 2020

Jurisdiction of Occurrence	COVID-19 Deaths (U07.1) ¹	Deaths from All Causes	Percent of Expected Deaths ²	Pneumonia Deaths (J12.0–J18.9) ³	Deaths with Pneumonia and COVID-19 (J12.0–J18.9 and U07.1) ³	Influenza Deaths (J09–J11) ⁴
United States ⁵	34,521	713,386	97	62,725	15,301	5,821
Alabama	153	12,427	91	769	43	83
Alaska	-	856	79	41	-	-
Arizona	194	15,859	101	1,115	102	105
Arkansas	26	7,774	94	519	-	67
California	1,111	69,341	98	5,947	643	552
Colorado	511	10,721	104	950	311	91
Connecticut	16	1,277	15	95	-	18
Delaware	53	2,034	85	123	24	14
District of Columbia	58	1,452	92	165	58	-
Florida	784	54,148	99	3,905	423	281
Georgia	460	19,739	91	1,306	213	94
Hawaii	-	2,830	94	185	-	18
Idaho	44	3,553	96	195	13	24
Illinois	977	28,696	104	2,529	558	170
Indiana	493	16,502	95	1,526	264	121
Iowa	78	7,447	94	531	20	80
Kansas	82	6,559	94	448	35	85
Kentucky	105	10,794	86	959	60	85

Louisiana	779	11,174	95	825	329	61
Maine	39	3,813	100	318	11	30
Maryland	631	13,661	105	1,233	268	110
Massachusetts	1,872	17,627	112	2,036	693	150
Michigan	1,746	26,798	106	2,580	838	221
Minnesota	169	11,351	99	832	60	113
Mississippi	186	8,064	99	737	90	51
Missouri	181	15,085	89	956	65	169
Montana	-	2,248	84	132	-	32
Nebraska	21	3,971	90	306	-	27
Nevada	134	6,431	96	534	106	36
New Hampshire	54	3,193	99	223	18	29
New Jersey	4,147	25,480	131	3,815	2,100	108
New Mexico	36	4,113	85	291	22	25
New York ⁶	4,800	32,000	122	5,080	2,480	188
New York City	10,978	30,711	219	6,076	4,019	850
North Carolina	0	10,727	43	639	0	127
North Dakota	-	1,546	85	140	-	18
Ohio	171	26,464	82	1,502	74	225
Oklahoma	114	8,762	84	782	43	90
Oregon	70	8,392	89	446	30	59
Pennsylvania	1,388	28,073	78	2,164	553	176
Rhode Island	55	2,235	82	124	13	23
South Carolina	150	12,807	101	786	56	91
South Dakota	-	1,898	89	149	-	21
Tennessee	127	18,292	95	1,364	62	119
Texas	347	48,574	93	3,567	135	310
Utah	28	4,675	96	277	11	38
Vermont	42	1,570	104	108	10	14
Virginia	326	17,624	99	981	103	102
Washington	527	14,139	95	1,175	281	98
West Virginia	11	5,022	85	364	-	57
Wisconsin	209	13,718	100	786	32	145
Wyoming	-	1,139	99	89	-	-
Puerto Rico	63	5,158	0	730	31	28

NOTE: Number of deaths reported in this table are the total number of deaths received and coded as of the date of analysis and do not represent all deaths that occurred in that period. Data cells with counts greater than zero and less than 10 have been suppressed in accordance with NCHS confidentiality standards.

*Data during this period are incomplete because of the lag in time between when the death occurred and when the death certificate is completed, submitted to NCHS and processed for reporting purposes. This delay can range from 1 week to 8 weeks or more, depending on the jurisdiction, age, and cause of death.

¹Deaths with confirmed or presumed COVID-19, coded to ICD-10 code U07.1.

²Percent of expected deaths is the number of deaths for all causes for this week in 2020 compared to the average number across the same week in 2017-2019.

³Pneumonia death counts exclude pneumonia deaths involving influenza.

⁴Influenza death counts include deaths with pneumonia or COVID-19 also listed as a cause of death.

⁵United States death count includes the 50 states, plus the District of Columbia and New York City.

⁶Excludes New York City.

Technical Notes

Comparing data in this report to other sources

Provisional death counts in this report will not match counts in other sources, such as media reports or numbers from county health departments. Death data, once received and processed by National Center for Health Statistics (NCHS), are tabulated by the state or jurisdiction in which the death occurred. Death counts are not tabulated by the decedent's state of residence. COVID-19 deaths may also be classified or defined differently in various reporting and surveillance systems. Death counts in this report include laboratory confirmed COVID-19 deaths and clinically confirmed COVID-19 deaths. This includes deaths where COVID-19 is listed as a "presumed" or "probable" cause. Some local and state health departments only report laboratory-confirmed COVID-19 deaths. This may partly account for differences between NCHS reported death counts and death counts reported in other sources. Provisional counts reported here track approximately 1-2 weeks behind other published data sources on the number of COVID-19 deaths in the U.S. (1,2,3).

Nature and sources of data

Provisional death counts are based on death records received and processed by NCHS as of a specified cutoff date. National provisional counts include deaths occurring within the 50 states and the District of Columbia. NCHS receives the death records from state vital registration offices through the Vital Statistics Cooperative Program. Provisional data are based on available records that meet certain data quality criteria at the time of analysis and may not include all deaths that occurred during a given time period especially for more recent periods. Estimates of completeness are provided. Therefore, they should not be considered comparable with final data and are subject to change.

Cause-of-death classification and definition of deaths

Mortality statistics are compiled in accordance with World Health Organization (WHO) regulations specifying that WHO member nations classify and code causes of death with the current revision of the International Statistical Classification of Diseases and Related Health Problems (ICD). ICD provides the basic guidance used in virtually all countries to code and classify causes of death. It provides not only disease, injury, and poisoning categories but also the rules used to select the single underlying cause of death for tabulation from the several diagnoses that may be reported on a single death certificate, as well as definitions, tabulation lists, the format of the death certificate, and regulations on use of the classification. Causes of death for data presented in this report were coded according to ICD guidelines described in annual issues of Part 2a of the NCHS Instruction Manual (4).

Coronavirus disease deaths are identified using the ICD-10 code U07.1. Deaths are coded to U07.1 when coronavirus disease 2019 or COVID-19 are reported as a cause that contributed to death on the death certificate. These can include laboratory confirmed cases, as well as cases without laboratory confirmation. If the certifier suspects COVID-19 or

determines it was likely (e.g., the circumstances were compelling within a reasonable degree of certainty), they can report COVID-19 as “probable” or “presumed” on the death certificate (5, 6).

Pneumonia deaths are identified using underlying cause-of-death codes from the 10th Revision of ICD (ICD-10): J12–J18, excluding deaths that involve influenza (J09–J11). Influenza deaths are identified from the ICD-10 codes J09–J11, and include deaths with pneumonia or COVID-19 listed as a contributing cause of death.

Estimated completeness of data

Provisional data are incomplete, and the level of completeness varies by jurisdiction, week, decedent’s age, and cause of death. Until data for a calendar year are finalized, typically in December of the following year, completeness of provisional data cannot be determined. However, completeness can be estimated in a variety of ways. Surveillance systems that rely on weekly monitoring of provisional mortality data, such as CDC’s FluView Interactive mortality surveillance (7), estimate completeness by comparing the count of deaths in a given week of the current year to the average count of deaths in that same week of the previous 3 years. These estimates can be generated for specific causes of death, jurisdictions, and age groups, and updated on a weekly or daily basis. For the purposes of COVID-19 surveillance, completeness is approximated by comparing the provisional number of deaths received to the number of expected deaths based on prior years data. Percent of expected deaths provided in this data release are based on the total count of deaths in the most recent weeks of the current year, compared with an average across the same weeks of the three previous years (i.e., 2017–2019). These estimates of completeness are calculated by week, jurisdiction of occurrence, and age group.

It is important to note that the true levels of completeness are unknown, and the estimates provided here are only a proxy. In cases where mortality rates are increasing rapidly, particularly when excess deaths due to a novel cause are occurring, values for completeness for recent weeks may exceed 100% even when NCHS has yet to receive all available data. Conversely, if the number of deaths was elevated in prior years due to a severe flu season, for example, estimated completeness in the most recent weeks may be lower than the true value. To avoid relying too heavily on comparisons to a single week of a single prior year, estimates of completeness included in this release are based on the average counts in a given week across 3 prior years (e.g., the 12th week of 2017, 2018, and 2019).

Percent of expected deaths provided in this release are shown to provide context for interpreting provisional counts of COVID-19 deaths and deaths due to related causes. Where estimated values are high (e.g., greater than 100%), this suggests that mortality is higher in 2020 relative to the same weeks of prior years. Where estimated values of completeness are low, this could indicate that data are incomplete due to delayed reporting, or that mortality is lower in 2020 compared with prior years, or some combination of these factors.

Delays in reporting

Provisional counts of deaths are underestimated relative to final counts. This is due to the many steps involved in reporting death certificate data. When a death occurs, a certifier (e.g. physician, medical examiner or coroner) will complete the death certificate with the underlying cause of death and any contributing causes of death. In some cases, laboratory tests or autopsy results may be required to determine the cause of death. Completed death certificates are sent to the state vital records office and then to NCHS for cause of death coding. At NCHS, about 80% of deaths are automatically processed and coded within seconds, but 20% of deaths need to be manually coded, or coded by a person. Deaths involving certain conditions such as influenza and pneumonia are more likely to require manual coding than other causes of death. Furthermore, all deaths with COVID-19 are manually coded. Death certificates are typically manually coded within 7 days of receipt, although the coding delay can grow if there is a large increase in the number of deaths. As a result, underestimation of the number of deaths may be greater for certain causes of death than others.

Previous analyses of provisional data completeness from 2015 suggested that mortality data is approximately 27% complete within 2 weeks, 54% complete within 4 weeks, and at least 75% complete within 8 weeks of when the death occurred (8). Pneumonia deaths are 26% complete within 2 weeks, 52% complete within 4 weeks, and 72% complete within 8 weeks (unpublished). Data timeliness has improved in recent years, and current timeliness is likely higher than published rates.

Place of Death

Place of death noted on the death certificate is determined by where the death was pronounced and on the physical location where the death occurred (10). Healthcare setting includes hospitals, clinics, medical facilities, or other licensed institutions providing diagnostic and therapeutic services by medical staff. Decedent's home includes independent living units such as private homes, apartments, bungalows, and cottages. Hospice facility refers to a licensed institution providing hospice care (e.g., palliative and supportive care for the dying), but not to hospice care that might be provided in other settings, such as a patient's home. Nursing home/long-term care facility refers to a facility that is not a hospital but provides patient care beyond custodial care, such as a nursing home, skilled nursing facility, a long-term care facility, convalescent care facility, intermediate care facility, or residential care facility. Other includes such locations as a licensed ambulatory/surgical center, birthing center, physician's office, prison ward, public building, worksite, outdoor area, orphanage, or facilities offering housing and custodial care but not patient care (e.g., board and care home, group home, custodial care facility, foster home).

Comparing deaths from different states

Death counts should not be compared across states. Data timeliness varies by state. Some states report deaths on a daily basis, while other states report deaths weekly or monthly. Furthermore, health departments and state vital record offices may be affected by COVID-19 related response activities, which could further delay death certificate reporting. Currently, 63% of U.S. deaths are reported within 10 days of the date of death, but there is variation within states. Twenty states report over 75% of deaths within the first 10 days, while three states report fewer than 1% of deaths within 10 days.

Why are pneumonia and influenza deaths included in this report?

Pneumonia and influenza deaths are included to provide context for understanding the completeness of COVID-19 mortality data and related trends. Deaths due to COVID-19 may be misclassified as pneumonia or influenza deaths in the absence of positive test results, and pneumonia or influenza may appear on death certificates as a comorbid condition. Additionally, COVID-19 symptoms can be similar to influenza-like illness, thus deaths may be misclassified as influenza. Thus, increases in pneumonia and influenza deaths may be an indicator of excess COVID-19-related mortality. Additionally, estimates of completeness for pneumonia and influenza deaths may provide context for understanding the lag in reporting for COVID-19 deaths, as it is anticipated that these causes would have similar delays in reporting, processing, and coding.




Source

NCHS, National Vital Statistics System. Estimates are based on provisional data.

References

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis*. 2020. Available from: [https://doi.org/10.1016/S1473-3099\(20\)30120-1](https://doi.org/10.1016/S1473-3099(20)30120-1). [↗](#)
2. Wu J, McCann A, Collins K, Harris R, Huang J, Almukhtar S. Coronavirus in the U.S.: Latest map and case count. *New York Times*. <https://www.nytimes.com/interactive/2020/us/coronavirus-us-cases.html>. [↗](#)
3. National Center for Immunization and Respiratory Diseases (NCIRD), Division of Viral Diseases. Cases in the US.

Centers for Disease Control and Prevention. 2020.

4. National Vital Statistics System. Instructions for classifying the underlying cause of death. In: NCHS instruction manual; Part 2a. Published annually.
5. National Center for Health Statistics. Guidance for certifying deaths due to COVID-19. Hyattsville, MD. 2020. Available from: <https://www.cdc.gov/nchs/data/nvss/vsrg/vsrg03-508.pdf>. 
6. National Center for Health Statistics. New ICD code introduced for COVID-19 deaths. Hyattsville, MD. 2020. Available from: <https://www.cdc.gov/nchs/data/nvss/coronavirus/Alert-2-New-ICD-code-introduced-for-COVID-19-deaths.pdf> 
7. National Center for Immunization and Respiratory Diseases (NCIRD). CDC's FluView Interactive. Centers for Disease Control and Prevention. Available from: <https://www.cdc.gov/flu/weekly/index.htm>.
8. Spencer MR, Ahmad F. Timeliness of death certificate data for mortality surveillance and provisional estimates. National Center for Health Statistics. 2016.
9. U.S. Census Bureau. Annual estimates of the resident population by single year of age and sex for the United States: April 1, 2010 to July 1, 2018. Available from: https://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=PEP_2018_PEPSYASEXN&prodType=table. 
10. Centers for Disease Control and Prevention. Medical Examiners' and Coroners' handbook on death registration and Fetal death reporting. Hyattsville, Maryland: DHHS Publication; 2003. PHS2003-1110

Page last reviewed: April 30, 2020

Content source: [National Center for Health Statistics](#)