



Published in final edited form as:

Travel Med Infect Dis. 2019 ; 32: 101504. doi:10.1016/j.tmaid.2019.101504.

International mass gatherings and travel-associated illness: A GeoSentinel cross-sectional, observational study

A full list of authors and affiliations appears at the end of the article.

Abstract

Background—Travelers to international mass gatherings may be exposed to conditions which increase their risk of acquiring infectious diseases. Most existing data come from single clinical sites seeing returning travelers, or relate to single events.

Methods—Investigators evaluated ill travelers returning from a mass gathering, and presenting to a GeoSentinel site between August 2015 and April 2019, and collected data on the nature of the event and the relation between final diagnoses and the mass gathering.

Results—Of 296 ill travelers, 51% were female and the median age was 54 years (range: 1–88). Over 82% returned from a religious mass gathering, most frequently Umrah or Hajj. Only 3% returned from the Olympics in Brazil or South Korea. Other mass gatherings included other sporting events, cultural or entertainment events, and conferences. Respiratory diseases accounted for almost 80% of all diagnoses, with vaccine preventable illnesses such as influenza and pneumonia accounting for 26% and 20% of all diagnoses respectively. This was followed by gastrointestinal illnesses, accounting for 4.5%. Sixty-three percent of travelers reported having a pre-travel encounter with a healthcare provider.

Conclusions—Despite this surveillance being limited to patients presenting to GeoSentinel sites, our findings highlight the importance of respiratory diseases at mass gatherings, the need for pre-travel consultations before mass gatherings, and consideration of vaccination against influenza and pneumococcal disease.

Keywords

Surveillance; Mass gatherings; Olympics; Respiratory

1. Introduction

Attendance at an international mass gathering (MG) may expose travelers to health risks related to crowded conditions, population movement, and inadequate sanitation [1,2]. According to the World Health Organization, an event can be classified as a MG if the number of people attending is sufficient to strain the planning and response resources of the community, state, or nation hosting the event [2]. However, much of the available literature describes mass gatherings as those exceeding 25,000 persons. We describe demographic

*Corresponding author. VITROME, Institut Hospitalo-Universitaire Méditerranée Infection, 19-21 Boulevard Jean Moulin, 13385, Marseille Cedex 05, France. philippe.gautret@ap-hm.fr (P. Gautret).

characteristics and diagnoses among travelers who attended a MG and presented with a travel-associated illness to a GeoSentinel site.

2. Methods

Records of ill travelers attending a MG between August 17, 2015, and April 30, 2019, were collected by GeoSentinel, a global clinicianbased surveillance network that monitors travel-related illnesses among international travelers and migrants [3]. GeoSentinel was established in 1995 as a collaboration between the Centers for Disease Control and Prevention and the International Society of Travel Medicine. It consists of 68 clinical sites in 28 countries. GeoSentinel records ill persons' visits to network sites; well travelers are not captured. Attendance at a MG is routinely recorded. During the study period, investigators were directed to enter supplemental details on the nature and location of the MG and to evaluate whether the diagnosis was likely associated with MG attendance. Records were excluded if the mass gathering was likely to have < 25,000 attendees or if data were missing regarding the type of MG. GeoSentinel's data collection protocol has been reviewed by a human subjects advisor at CDC's National Center for Emerging and Zoonotic Infectious Diseases and is classified as public health surveillance and not human subjects research. Additional ethics clearance was obtained by participating sites as required by their respective institutions or national regulations.

3. Results

A total of 327 records of ill travelers attending a MG during international travel had surveys completed by the investigator providing their care. Thirty-one records were excluded. Of 296 ill travelers included, 151 (51.0%) were female and the median age was 54 years (range: 1–88; IQR: 39–64). Purposes of MGs were religious (243 cases; 82.1%), cultural (e.g., music, dance, carnival) (19 cases; 6.4%), the World Scout Jamboree (17 cases; 5.7%), major sport events (13 cases; 4.4%), or a large conference (4 cases; 1.4%). The top three specific MGs were Umrah or Hajj in Saudi Arabia with 241 cases (81.4%), including 87 at Umrah and 154 at Hajj, followed by the World Scout Jamboree in Japan with 17 cases (5.7%), and the Olympics in Brazil and South Korea with 9 cases (3.1%) (Table 1). Ill travelers who acquired illness at Umrah or Hajj were older (median age: 57 years [range: 1–88; IQR: 47–66]) than those at the Olympics (median age: 36 years [range: 19–55; IQR: 31–46]) or the World Scout Jamboree (median age: 18 years [range: 14–52; IQR: 17–20]). Sixty-four percent of ill travelers who attended Umrah or Hajj were hospitalized because of their illness; one traveler who attended the World Scout Jamboree and no travelers who attended the Olympics were hospitalized.

Overall, 130 of 206 (63%) ill travelers with information available reported having a pre-travel encounter with a healthcare provider. The majority (260 of 296, 87.8%) of acquired illnesses were directly associated with MG attendance, while the relation between the illness and the MG was not ascertainable for 25 travelers (8.4%), and the illness was travel-related but not linked to MG attendance for 11 travelers (3.7%). Only three of nine (33.3%) diagnoses among travelers who attended the Olympics were associated with attending the MG.

A total of 303 diagnoses were reported among the 260 ill travelers whose illness was associated with MG attendance (Table 2). Respiratory diseases were the most frequently reported disease category with 236 diagnoses (77.9%), followed by gastrointestinal diseases (14 diagnoses; 4.6%). Diagnoses related to attendance at the three most common MGs — Umrah or Hajj, World Scout Jamboree, and Olympics — are presented in Table 1.

4. Discussion

GeoSentinel sites evaluated 260 pilgrims with Umrah- or Hajj-related illnesses among an estimated 32 million foreign MG attendees over the study period [4], 17 Scouts among 33,628 attendees [5] and 9 Olympic spectators among 7 million attendees [6,7]. It should be noted, however, that GeoSentinel collects data only on ill travelers presenting to a network site and some GeoSentinel sites may care for more MG attendees than others, which may not be representative of all travelers attending a MG. In particular, the Scouts reported to GeoSentinel were from a single site in Sweden and were identified because of an international alert following a meningococcal outbreak (W ST11 serotype) among six Scottish and Swedish nationals who attended this event [8].

Ill MG attendees seen at a GeoSentinel site most frequently attended Umrah or Hajj, likely due to the large number of travelers to these pilgrimages. These findings are consistent with previously published literature demonstrating that outbreaks are not frequently reported during or after MGs other than Umrah and Hajj pilgrimages, although they sometimes occur at Muslim, Christian, and Hindu religious events, sports events, and large-scale open-air festivals [9,10]. Its size, international recognition, unique multinational component, and yearly recurrence, likely account for the preponderance of Umrah and Hajj among international MGs responsible for outbreaks. Our data are also consistent with previous reports regarding the older age of Umrah and Hajj travelers [11], and the younger age of Scout Jamboree travelers. The finding that almost three-quarters of ill travelers who attended Umrah or Hajj were hospitalized is likely due to a recruitment bias, given that ill travelers seen at GeoSentinel sites may be sicker because of the specialized infectious disease or tertiary care nature of these sites; or to initial clinical suspicion for Middle East respiratory syndrome, resulting in hospitalization and isolation pending testing. The paucity of reported trauma is also likely a reflection of the infectious disease specialization of most GeoSentinel network sites, and of the fact that trauma usually requires immediate attention and is most likely to occur during travel [12].

The predominance of respiratory tract infections reported in this analysis, including 4 cases of pneumococcal disease with one death, corroborates results obtained in Saudi hospitals of pneumonia diagnoses among Umrah and Hajj attendees [13] as well as the high number of influenza virus infections observed among both patients in Saudi hospitals and patients hospitalized on returning to their home countries from Umrah and Hajj [14]. Crowded conditions with close proximity of large numbers of attendees at Umrah and Hajj are a likely explanation for the frequency of respiratory infections among pilgrims, given the transmissibility of these infections. Our results confirm the importance of influenza vaccination for Umrah and Hajj travelers [15]. Although the availability of influenza vaccine may be limited depending on the time of year Hajj occurs [16], immunization with expired

influenza vaccine from the recently ended influenza season may have few associated adverse events [17]. By contrast, most illnesses among travelers attending the Olympics were linked to travel, but not to attending the Olympics, and these illnesses' were mild, which may be due to the physical separation of various sporting events at the Olympics, the propensity to hold the Olympics in high income countries, the relatively short travel duration of attendees and participants, and the relatively young age of participants. This is supported by the rarity of documented outbreaks during Olympic Games between 1984 and 2014 [9,18].

Our identification of respiratory tract infections, especially pneumonia, among Umrah and Hajj attendees suggests the need for additional research to document responsible pathogens. Such data may have important consequences regarding vaccine recommendations before travel. In particular, it may be possible to validate that the influenza serotypes found are representative of strains circulating during the prior northern winter. These data also may provide justification for recommending pneumococcal vaccination for some high-risk travelers [19–21]. However, one-third of all travelers in this report did not attend a pre-travel consultation, and this will likely be a barrier to implementation of such recommendations. Many countries have a staging area where Umrah and Hajj attendees gather before departure, or have pre-Hajj classes at local mosques. Leaders there could raise awareness of recommended (but not required) vaccines and personal protective measures such as carrying and routinely using hand sanitizer may lead to higher level of protective behaviors of traveling pilgrims [22]. Also, primary care physicians should inquire about planned travel to mass gatherings and vaccinate travelers against meningococcus, influenza, and pneumococcus, as appropriate.

Given its broad international catchment, GeoSentinel plays a role in identifying emerging infectious diseases with epidemic potential, thus contributing to efforts to create enhanced international multidisciplinary surveillance of MG-associated illnesses, as recently recommended by experts [23]. Since our surveillance was limited to patients presenting to GeoSentinel sites, to better understand travel-associated illnesses acquired at MGs improved global surveillance mechanisms are needed.

Authors

Philippe Gautret^{a,*}, Kristina M. Angelo^b, Hilmir Asgeirsson^c, Alexandre Duvignaud^d, Perry J.J. van Genderen^e, Emmanuel Bottieau^f, Lin H. Chen^g, Salim Parker^h, Bradley A. Connor^{i,j}, Elizabeth D. Barnett^k, Michael Libman^l, Davidson H. Hamer^{m,n}, GeoSentinel Network,

Emilie Javelle^o, Marc Mendelson^p, Hedvig Glans^q, Eric Caumes^r, Poh Lian Lim^s, Jonathan Alpern^t, William Stauffer^t, Andrea Boggild^u, Kevin Kain^u, Karin Leder^v, Joseph Torresi^v, Katherine Plewes^w, Wayne Ghesquiere^w, Denis Malvy^x, Francois Chappuis^y, Giles Eperon^y, Sarah Borwein^z, Johnnie Yates^{aa}, Brian Ward^{ab}, Jesse Waggoner^{ac}, Henry Wu^{ac}, Israel Molina^{ad}, Dan Bourque^{ae}, Kunjana Mavunda^{af}, Mugen Ujiie^{ag}, Satoshi Katsuna^{ag}, Carsten Larsen^{ah}, Christian Wejse^{ah}, Noreen Hynes^{ai}, Paul Kelly^{aj}, Susan Kuhn^{ak}, Cecilia Perret Perez^{al}, Thomas Weitzel^{am}, Sabine Jordan^{an}, Mauro Saio^{ao}, Michael Beadsworth^{ap}, Nicholas Beeching^{aq}, Mogens Jensenius^{aq}, Susan Anderson^{ar}, Patricia Schlagenhauf^{as}, Rainer Weber^{as}

Affiliations

^aAix Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France and IHU-Méditerranée Infection, Marseille, France

^bDivision of Global Migration and Quarantine, Centers for Disease Control and Prevention, GA, USA

^cDepartment of Infectious Diseases, Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden

^dDepartment of Infectious Diseases and Tropical Medicine, Division of Tropical Medicine and Clinical International Health, CHU Bordeaux, Bordeaux, France

^eDepartment of Medical Microbiology and Infectious Diseases, University Hospital Erasmus Medical Centre, Rotterdam, the Netherlands

^fInstitute of Tropical Medicine, Antwerp, Belgium

^gDivision of Infectious Diseases and Travel Medicine, Mount Auburn Hospital and Harvard Medical School, MA, USA

^hDivision of Infectious Diseases & HIV Medicine, Department of Medicine, University of Cape Town, Cape Town, South Africa

ⁱWeill Cornell Medical College, New York, USA

^jNew York Center for Travel and Tropical Medicine, New York, USA

^kDepartment of Pediatrics, Boston University School of Medicine, MA, USA

^lJ.D. MacLean Centre for Tropical Diseases, McGill University, Montreal, Canada

^mDepartment of Global Health, Boston University School of Public Health, MA, USA

ⁿSection of Infectious Diseases, Department of Medicine, Boston University School of Medicine, MA, USA

^oIHU Méditerranée Infection, Marseille, France

^pDivision of Infectious Diseases & HIV Medicine - Groote Schuur Hospital, Cape Town, South Africa

^qClinic of Infectious Diseases, Karolinska University Hospital, Stockholm, Sweden

^rService des Maladies Infectieuses et Tropicales, Hôpital Pitié-Salpêtrière, Sorbonne Université, Paris, France

^sTravellers' Health & Vaccine Center, Singapore

^tHealthPartners Center for International Health - Regions Hospital, Minneapolis, MN, USA

^uCentre for Travel and Tropical Medicine - Toronto General Hospital, Toronto, Canada

- ^vVictorian Infectious Diseases Service, Royal Melbourne Hospital, Melbourne, Australia
- ^wVancouver General Hospital and Tropical Medicine Expert Group BC, Vancouver, Canada
- ^xSanté-Voyages et Médecine Tropicale, Hôpital Saint-André, Bordeaux, France
- ^yDivision of Tropical and Humanitarian Medicine, Geneva University Hospitals, Geneva, Switzerland
- ^zTravelSafe Medical Centre - Central Health Medical Practice, Hong Kong, China
- ^{aa}Kaiser Permanente Honolulu Clinic, Honolulu, HI, USA
- ^{ab}McGill University Centre for Tropical Diseases - Montreal General Hospital, Montreal, Canada
- ^{ac}Emory TravelWell Center, Atlanta, GA, USA
- ^{ad}Antiga Escola de Enfermeria, Hospital Universitari Vall d'Hebron, Barcelona, Spain
- ^{ae}Travel Medicine Center, Mt. Auburn Hospital, Cambridge, MA, USA
- ^{af}International Travel Clinic, Miami, FL, USA
- ^{ag}Travel Clinic - National Center for Global Health and Medicine, Toyko, Japan
- ^{ah}Department of Infectious Disease - Aarhus University Hospital, Aarhus, Denmark
- ^{ai}John Hopkins Univ. School of Medicine - Geographic Medicine Center - Division of Infectious Diseases, Baltimore, MD, USA
- ^{aj}Tropical Medicine Clinic - Jacobi Medical Center, Bronx, NY, USA
- ^{ak}Alberta Children's Hospital, Calgary, Canada
- ^{al}Centro de Enfermedades - Tropicales y Salud del Viajero - Pontificia Universidad Católica de Chile, Santiago, Chile
- ^{am}Travel Medicine Program, Clínica Alemana, Universidad del Desarrollo, Santiago, Chile
- ^{an}University Medical Centre Hamburg Eppendorf, Department of Medicine, Division of Infectious Diseases and Tropical Medicine, Bernhard-Nocht-Kilnik, Hamburg, Germany
- ^{ao}Nairobi Hospital, Nairobi, Kenya
- ^{ap}Liverpool School of Tropical Medicine, Liverpool, United Kingdom
- ^{aq}Department of Infectious Diseases - Oslo University Hospital, Ullevål, Oslo, Norway
- ^{ar}Travel Medicine & Emergency/Urgent Care - Palo Alto Medical Foundation, Palo Alto, CA, USA

^{as}University of Zürich Centre for Travel Medicine, WHO Collaborating Centre for Travellers' Health, Zurich, Switzerland

Acknowledgments

Funding

GeoSentinel, the Global Surveillance Network of the International Society of Travel Medicine (ISTM), is supported by a Cooperative Agreement (U50CK00189) from the Centers for Disease Control and Prevention (CDC), as well as funding from the ISTM and the Public Health Agency of Canada. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of CDC.

References

- [1]. Abubakar I, Gautret P, Brunette G, Blumberg L, Johnson D, Pomeroy G, et al. Global perspectives for prevention of infectious diseases associated with mass gatherings. *Lancet Infect Dis* 2012 Jan;12(1):66–74. [PubMed: 22192131]
- [2]. World Health Organization. Public health for mass gatherings: key considerations Available at: https://www.who.int/ihr/publications/WHO_HSE_GCR_2015.5/en/, Accessed date: 20 May 2019.
- [3]. Harvey K, Esposito DH, Han PKozarsky P, Freedman DO, Plier DA, et al. Surveillance for travel-related disease —GeoSentinel surveillance system, United States, 1997–2011. *MMWR Surveill Summ* 2013;62:1–23.
- [4]. General Authority for Statistics. Kingdom of Saudi Arabia Available at: <http://www.stats.gov.sa/en/28>, Accessed date: 20 May 2019.
- [5]. Japan 2015. 23rd World Scout jamboree Available at: https://db0nus869y26v.cloudfront.net/en/23rd_World_Scout_Jamboree, Accessed date: 20 May 2019.
- [6]. International Olympic Committee. Marketing report Rio; 2016 Available at: <https://stillmed.olympic.org/media/Document%20Library/OlympicOrg/Games/Summer-Games/Games-Rio-2016-Olympic-Games/Media-Guide-for-Rio-2016/IOC-Marketing-Report-Rio-2016.pdf>, Accessed date: 20 May 2019.
- [7]. Panja T Olympic Organizers say tickets are sold, but where are the people? *The New York Times*. 15 February 2018 Available at: <https://www.nytimes.com/2018/02/15/sports/olympics/olympics-venues-empty.html>, Accessed date: 20 May 2019.
- [8]. Kanai M, Kamiya H, Smith-Palmer A, Takahashi H, Hachisu Y, Fukusumi M, et al. Meningococcal disease outbreak related to the World Scout jamboree in Japan. *Western Pac Surveill Response J* 2015;8(2):25–30. 2017 May 8.
- [9]. Gautret P, Steffen R. Communicable diseases as health risks at mass gatherings other than Hajj: what is the evidence? *Int J Infect Dis* 2016 Jun;47:46–52. [PubMed: 26987476]
- [10]. Hoang VT, Gautret P. Infectious diseases and mass gatherings. *Curr Infect Dis Rep* 2018 Aug 28;20(11):44. [PubMed: 30155747]
- [11]. Memish ZA, Zumla A, Alhakeem RF, Assiri A, Turkestani A, Al Harby KD, et al. Hajj: infectious disease surveillance and control. *Lancet* 2014;383:2073–82. [PubMed: 24857703]
- [12]. Khan ID, Khan SA, Asima B, Hussaini SB, Zakiuddin M, Faisal FA. Morbidity and mortality amongst Indian Hajj pilgrims: a 3-year experience of Indian Hajj medical mission in mass-gathering medicine. *J Infect Public Heal* 2018;11:165–70.
- [13]. Benkouiten S, Al-Tawfiq JA, Memish ZA, Albarrak A, Gautret P. Clinical respiratory infections and pneumonia during the Hajj pilgrimage: a systematic review. *Trav Med Infect Dis* 2019 Mar - Apr;28:15–26.
- [14]. Gautret P, Benkouiten S, Al-Tawfiq JA, Memish ZA. Hajj-associated viral respiratory infections: a systematic review. *Trav Med Infect Dis* 2016 Mar-Apr;14(2):92–109.
- [15]. Al-Tawfiq JA, Gautret P, Memish ZA. Expected immunizations and health protection for Hajj and Umrah 2018 -An overview. *Trav Med Infect Dis* 2017 Sep;19:2–7.

- [16]. Alfelali M, Alqahtani AS, Barasheed O, Booy R, Rashid H. Mandating influenza vaccine for Hajj pilgrims. *Lancet Infect Dis* 2016 Jun;16(6):633–4. [PubMed: 27301914]
- [17]. Hesse EM, Hibbs BF, Cano MV. Notes from the Field: administration of expired injectable influenza vaccines reported to the vaccine adverse event reporting system — United States, July 2018–March 2019. *MMWR Morb Mortal Wkly Rep* 2019;68(23):529–30. [PubMed: 31194725]
- [18]. Zieliński A. Enhanced surveillance at mass gatherings. *Przegl Epidemiol* 2009;63(4):477–85. [PubMed: 20120944]
- [19]. Alfelali M, Khandaker G, Booy R, Rashid H. Mismatching between circulating strains and vaccine strains of influenza: effect on Hajj pilgrims from both hemispheres. *Hum Vaccines Immunother* 2016 Mar 3;12(3):709–15.
- [20]. Yezli S, van der Linden M, Booy R, AlOtaibi B. Pneumococcal disease during Hajj and Umrah: research agenda for evidence-based vaccination policy for these events. *Trav Med Infect Dis* 2019 May - Jun;29:8–15.
- [21]. Goeijenbier M, van Genderen P, Ward BJ, Wilder-Smith A, Steffen R, Osterhaus AD. Travellers and influenza: risks and prevention. *J Travel Med* 2017 Jan;24(1):tav078. [PubMed: 28077609]
- [22]. Alqahtani AS, Heywood AE, Rashid H. Preparing Australian pilgrims for the Hajj 2018. *J Travel Med* 2018;1–2:tay068.
- [23]. Memish ZA, Steffen R, White P, Dar O, Azhar EI, Sharma A. Mass gathering medicine: public health issues arising from mass gathering religious and sporting events. *Lancet* 2019 May 18;393:2073–84. [PubMed: 31106753]

Mass gathering characteristics, ill traveler demographics, and most frequent illnesses acquired at Umrah and Hajj, the 23rd Scout Jamboree, and the Olympics (n = 296).

Table 1

	Umrah or Hajj	23rd World Scout Jamboree	Olympics
Location	Mecca, Saudi Arabia	Yamagushi, Japan	Rio de Janeiro, Brazil Pyeongchang, South Korea
Dates	Hajj: 21–26 September 2015 10–15 September 2016 30 August–4 September 2017 19–29 August 2018 Umrah: At any time during the year, but preferentially during Ramadan	28 July to 8 August 2015	5–21 August 2016 9–25 February 2018
Attendance [4–7]	Hajj 2015: 1,384,941 foreign attendees Hajj 2016: 1,325,372 foreign attendees Hajj 2017: 1,752,014 foreign attendees Hajj 2018: 1,758,722 foreign attendees Umrah 2017: 6,532,074 foreign attendees in 2017 (data not available in other years)	33,628 attendees	> 6,000,000 entrances > 1,000,000 entrances
Number of GeoSentinel cases	241	17	Rio: 8 Pyeongchang: 1
Patient age, median (years)	57 (range: 1–88; IQR: 47–66)	18 (range: 14–52; IQR: 17–20)	36 (range: 19–55; IQR: 31–46)
Female gender (%)	51.0	47.1	22.2
Pre-travel encounter (%)	67.1 ^a	Not documented	71.4
Inpatient (%)	63.9	5.9	0
Association of illness(es) with MG (%)	96.7	100	33.3 ^b
Most frequent diagnoses associated with MG (% of patients)	Pneumonia (28.6) Confirmed influenza (22.7) Acute bronchitis (21.9) Influenza-like illness (9.9) Upper respiratory tract infection (7.3)	Upper respiratory tract infection (29.4) Viral syndrome without rash (29.4) Influenza-like illness (17.7) Pharyngitis (17.7) Skin infection (5.9)	Viral syndrome without rash (2 patients) Viral syndrome (1 case) Acute bronchitis (1 case)
Death	1	0	0

^aOf 173 records for which information was available.

^bOf 7 records for which information was available.

Table 2

Diagnoses (N = 303) among 260 ill travelers with diseases associated with attendance at an international mass gathering.

Diagnosis	Number of Diagnoses ^a	% of Ill Travelers
Respiratory		
Pneumonia ^b	67	25.7
Influenza ^c	53	20.3
Acute bronchitis	52	19.9
Influenza-like illness	27	10.4
Upper respiratory tract infection, unspecified	22	8.4
Acute exacerbation of chronic respiratory disease	5	1.9
Pharyngitis	4	1.5
Cough, no etiology	2	0.8
Enterovirus 68 infection	1	0.4
Pleural effusion	1	0.4
Pulmonary embolism	1	0.4
Pulmonary tuberculosis	1	0.4
Gastrointestinal		
Acute gastroenteritis	6	2.3
Acute diarrhea	4	1.5
<i>Campylobacter</i> infection	1	0.4
Nausea and vomiting, no etiology	1	0.4
Non-pathogenic protozoa	1	0.4
<i>Salmonella</i> infection	1	0.4
Ear, nose, and throat		
Acute sinusitis	3	1.1
Acute otitis media	2	0.8
Tonsillitis	1	0.4
Dermatologic		
Skin and soft tissue infection	7	2.7
Insect bite	1	0.4
Urogenital		
Pyelonephritis ^d	2	0.8
Acute renal failure	1	0.4
Cystitis	1	0.4
Vaginitis	1	0.4
Neurologic		
Viral meningitis	1	0.4
Trauma		
Injury	1	0.4
Strain	1	0.4

Diagnosis	Number of Diagnoses ^a	% of Ill Travelers
Blisters	1	0.4
Febrile systemic illness		
Viral syndrome without rash	9	3.4
Sepsis or bacteremia	3	1.1
Viral syndrome	1	0.4
Miscellaneous		
Antibiotic resistant organism ^e	5	1.9
Dehydration	2	0.8
Fatigue	2	0.8
Myalgia	2	0.8
Adverse drug event ^d	1	0.4
Dizziness	1	0.4
Heart disease	1	0.4
Osteomyelitis	1	0.4
Thrombophlebitis	1	0.4
Death ^f	1	0.4

^aAll diagnoses were confirmed according to pre-established GeoSentinel Network criteria for each diagnosis, unless otherwise specified.

^bIncluding *Streptococcus pneumoniae* infection (4 cases).

^cInfluenza A (38 cases) and influenza B (15 cases).

^dOne case was probable.

^eOne organism was specified (*Salmonella* spp.).

^fDirectly related to complications from pneumococcal pneumonia.