

Coronavirus Disease 2019 (COVID-19)

When to wear gloves

For the general public, CDC recommends wearing gloves at these times.



When cleaning

When you are routinely cleaning and disinfecting your home.

- Follow precautions listed on the disinfectant product label, which may include-
 - wearing gloves (reusable or disposable) and
 - having good ventilation by turning on a fan or opening a window to get fresh air into the room you're cleaning.
- Wash your hands after you have removed the gloves.



When caring for someone who is sick

If you are providing care to someone who is sick at home or in another non-healthcare setting

- Use disposable gloves when cleaning and disinfecting the area around the person who is sick or other surfaces that may be frequently touched in the home.
- Use disposable gloves when touching or having contact with blood, stool, or body fluids, such as saliva, mucus, vomit, and urine.
- After using disposable gloves, throw them out in a lined trash can. Do not disinfect or reuse the gloves.
- Wash your hands after you have removed the gloves.



When gloves aren't needed

Wearing gloves outside of these instances (for example, when using a shopping cart or using an ATM) will not necessarily protect you from getting COVID-19 and may still lead to the spread of germs. The best way to protect yourself from germs when running errands and after going out is to regularly wash your hands with soap and water for 20 seconds or use hand sanitizer with at least 60% alcohol.



Other ways to protect yourself

COVID-19 is a respiratory virus and is mainly spread through droplets created when a person who is infected coughs, sneezes, or talks. You can protect yourself by keeping social distance (at least 6 feet) from others and washing your hands with soap and water for 20 seconds (or using a hand sanitizer with at least 60% alcohol) at key times, and practicing everyday preventive actions.



Gloves in the workplace

Guidelines and recommendations for glove use in healthcare and work settings will differ from recommendations for the general public.

Page last reviewed: May 9, 2020

Content source: National Center for Immunization and Respiratory Diseases (NCIRD), Division of Viral Diseases