Acute Flaccid Myelitis

AFM Cases and Outbreaks

As of August 31, 2020, there have been 21 confirmed cases in 2020. There are 44 reports of patients under investigation (PUIs) for 2020. Two patients with confirmed AFM died in the acute phase of their illness, one in 2017 and one in 2020. We have also learned of deaths in cases confirmed in previous years.


There were **46 confirmed cases in 2019** out of 142 PUIs. CDC, state, and local health departments are still investigating some of the PUIs.

There have been **638 confirmed cases** since CDC began tracking AFM in August of 2014. CDC has been thoroughly investigating cases since that time. We have seen increases in AFM cases, mostly in young children, every two years.

Annual AFM Cases by State


Annual maps represent only cases for whom information was sent to and confirmed by CDC as of August 31, 2020. Patients under investigation are still being classified, and the case counts are subject to change. Cases of AFM have occurred in 49 states and the District of Columbia.

2020 –




Thus far in 2020, there have been 21 confirmed cases in 13 states and the District of Columbia.

2019




2018




In 2018, there were 238 total confirmed cases in 42 states. One of the confirmed cases is a foreign resident (based on the country of usual residence) and therefore not included in the state map.






In 2016, there were 153 total confirmed cases in 39 states and the District of Columbia.



2014




In 2014, there were 120 total confirmed cases in 34 states between the months of August and December. These case counts include only pediatric patients younger than 22 years of age.

We defer to the states to release additional information on cases as they choose.

Cases by Month, 2014-2020

Most patients had onset of AFM between August and November, with increases in AFM cases every two years since 2014. Many viruses commonly circulate at this same time of year, including enteroviruses, which are likely responsible for the increase in cases in peak years.

Confirmed AFM cases by CDC



Data table —

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec	Total
2014								21	51	24	15	9	120
2015	2	2	1	0	1	0	2	3	1	4	2	4	22
2016	1	0	6	1	7	9	13	33	43	27	9	4	153
2017	2	5	5	4	2	4	3	1	4	0	3	5	38
2018	0	4	1	2	3	8	8	30	88	66	21	7	238
2019	2	4	4	3	5	2	5	4	6	7	2	2	46
2020	7	3	5	2	1	2	1						21

Confirmed cases of AFM by month of illness onset. Case Counts are subject to change.

- ^ Confirmed AFM cases by CDC from August 2014 through August 31, 2020. Case counts are subject to change.
- * The data shown from August 2014 to July 2015 are based on the AFM investigation case definition: onset of acute limb weakness on or after August 1, 2014, and a magnetic resonance image (MRI) showing a spinal cord lesion largely restricted to gray matter in a patient age ≤21 years.
- † The data shown from August 2015 to present are based on the AFM case definition adopted by the Council of State and Territorial Epidemiologists (CSTE): acute onset of focal or flaccid limb weakness and an MRI showing spinal cord lesion largely restricted to gray matter and spanning one or more spinal segments, regardless of age.

For more information, visit the Case Definitions page.

It is currently difficult to interpret trends of the AFM data. Collecting information about PUIs for AFM is relatively new. There may initially be more variability in the AFM data from year to year, making it difficult to interpret or compare case counts between years.







For Parents of Children with AFM

Page last reviewed: September 2, 2020

Content source: National Center for Immunization and Respiratory Diseases, Division of Viral Diseases